

A Theory of Higher-Order Subtyping with Type Intervals

Extended Version Including Supplementary Material

SANDRO STUCKI, Chalmers University of Technology, Sweden

PAOLO G. GIARRUSSO, Bedrock Systems Inc., Germany

The calculus of Dependent Object Types (DOT) has enabled a more principled and robust implementation of Scala, but its support for type-level computation has proven insufficient. As a remedy, we propose F^ω , a rigorous theoretical foundation for Scala's higher-kinded types. F^ω extends $F^\omega_{<}$ with *interval kinds*, which afford a unified treatment of important type- and kind-level abstraction mechanisms found in Scala, such as bounded quantification, bounded operator abstractions, translucent type definitions and first-class subtyping constraints. The result is a flexible and general theory of higher-order subtyping. We prove type and kind safety of F^ω , as well as weak normalization of types and undecidability of subtyping. All our proofs are mechanized in Agda using a fully syntactic approach based on hereditary substitution.

CCS Concepts: • **Theory of computation** → **Type theory; Type structures; Operational semantics; Program verification.**

Additional Key Words and Phrases: Scala, higher-kinded types, subtyping, type intervals, bounded polymorphism, bounded type operators, singleton kinds, dependent kinds, hereditary substitution

ACM Reference Format:

Sandro Stucki and Paolo G. Giarrusso. 2021. A Theory of Higher-Order Subtyping with Type Intervals: Extended Version Including Supplementary Material. *Proc. ACM Program. Lang.* 5, ICFP, Article 69 (August 2021), 73 pages. <https://doi.org/10.1145/3473574>

1 INTRODUCTION

Modern statically typed programming languages provide powerful type-level abstraction facilities such as parametric polymorphism, subtyping, generic datatypes, and varying degrees of type-level computation. When several of these features are present in the same language, new and more expressive combinations arise, such as (F1) bounded quantification, (F2) bounded type operators, and (F3) translucent type definitions. Such mechanisms further increase the expressivity of the language, but also the complexity of its type system and, ultimately, its implementation.

A case in point is the Scala programming language. Scala is a multi-paradigm language that integrates functional and object-oriented concepts. It features all of the aforementioned type-level constructs and many more. While programmers enjoy the expressivity of Scala's type system, the language developers have struggled for years to manage its complexity – tracing down elusive compiler bugs and soundness issues in a seemingly ad-hoc fashion. The development of the calculus of Dependent Object Types (DOT) [Amin et al. 2016] marked a turning point. By providing a solid theoretical foundation for a large part of the Scala language, DOT inspired a complete redesign of the Scala compiler as well as a substantial redesign of the language itself – giving rise to Scala 3 [Dotty Team 2020]. But despite initial hopes to the contrary, DOT has proven insufficient to express Scala's *higher-kinded* (HK) types, which are used pervasively throughout the Scala standard library [Moors et al. 2008b]. The lack of a proper theory of HK types severely complicates

Authors' addresses: Sandro Stucki, Computer Science and Engineering, Chalmers University of Technology, Gothenburg, Sweden, sandros@chalmers.se; Paolo G. Giarrusso, Bedrock Systems Inc., Berlin, Germany, p.giarrusso@gmail.com.

© 2021 Copyright held by the owner/author(s).

This is the author's version of the work. It is posted here for your personal use. Not for redistribution. The definitive Version of Record was published in *Proceedings of the ACM on Programming Languages*, <https://doi.org/10.1145/3473574>.

their implementation in Scala 3 [Odersky et al. 2016]. To address this problem, we introduce F^ω , a rigorous theoretical foundation for Scala's HK types with a machine-checked safety proof.

We start by illustrating the use of HK types through an example.

```
type Ordering[A] = (A, A) => Boolean
class SortedView[A, B >: A](xs: List[A], ord: Ordering[B]) {
  def foldLeft[C](z: C, op: (C, A) => C): C = //...
  def concat[C >: A <: B](ys: List[C]): SortedView[C, B] = //...
  // definitions of further operations such as 'map', 'flatMap', etc.
}
```

The example is inspired by the definition of the class `SeqView.Sorted` from the standard library.¹ It is heavily simplified for conciseness and to avoid the use of Scala idioms irrelevant to this paper.

The `SortedView` class allows one to iterate over the elements of an underlying list `xs` in increasing order (via `foldLeft`) without modifying the original list. The second constructor argument `ord` provides the comparison operation used for sorting. Note that `ord` compares data of type `B`, which is declared a supertype of the element type `A` via the annotation `B >: A`. Thus we may use a comparison function for a less precise type to sort the elements in the view.²

In Scala, parametrized classes like `SortedView` are first-class type operators. Hence, `SortedView` is an instance of a lower-bounded type operator (F2). The operator `Ordering[A]` is just a convenient type alias for the function type `(A, A) => Boolean` of binary operators. It is a *transparent* type definition (F3), in that instances of `Ordering[A]` can be replaced by its definition anywhere in the program. The method `concat` illustrates the use of lower- and upper-bounded polymorphism (F1) to combine views on lists of different but related element types. Views are covariant: if `A` is a subtype of `C` then a view `v` on an `A`-list is also a view on a `C`-list and can be extended as such using `v.concat(ys)`. The upper bound on `C` ensures that the ordering remains applicable. This pattern of using lower-bounds to implement operations on covariant data structures is common in the Scala standard library.

Although bounded quantification (F1) and bounded operators (F2) may seem conceptually different from translucent type definitions (F3), all three are closely related. A type alias declaration of the form `X = A` (such as `Ordering` above) declares that the type `X` has type `A` as upper and lower bound. Because subtyping in Scala is antisymmetric, this effectively identifies `X` with `A`. In general, a type declaration of the form `X >: A <: B`, introduces an abstract type `X` that is bounded by `A` from below and by `B` from above. In other words, the declaration `X >: A <: B` specifies a *type interval* in which `X` must be contained. Type aliases take the form of *singleton intervals* `X >: A <: A`, where the lower and upper bounds coincide.

In our example, the types `Ordering` and `SortedView` can be seen as abstract types. We can write their declarations as³

```
type Ordering[A] >: (A, A) => Boolean <: (A, A) => Boolean
type SortedView[A, B >: A] <: {
  def foldLeft[C](z: C, op: (C, A) => C): C
  def concat[C >: A <: B](ys: List[C]): SortedView[C, B] /* ... */ }

```

¹See [https://www.scala-lang.org/api/2.13.6/scala/collection/SeqView\\$\\$Sorted.html](https://www.scala-lang.org/api/2.13.6/scala/collection/SeqView$$Sorted.html)

²Expert readers may notice that `Ordering` is contravariant in its argument, hence `SortedView` could simply take a parameter of type `Ordering[A]` and still accept instances of `Ordering[B]`. But this simplification does not extend to the Scala standard library, where `Ordering` is invariant due to additional methods; hence the lower bound is necessary. For readability, we use the simplified version of `Ordering` but keep the bound in the definition of `SeqView`.

³This code is accepted by the Scala 2.13.6 compiler. Ironically, it is rejected by Scala 3, which is built on DOT.

This version is closer to how type definitions are represented in DOT. The class `SortedView` is now represented as an abstract type declaration with only an upper bound. This means that its interface remains exposed – any instance of `SortedView[A, B]` can be up-cast to a record with fields `foldLeft`, `concat`, etc. But not every record (or class) containing those fields is automatically an instance of the type `SortedView[A, B]`. This ensures that `SortedView` continues to behave like a nominal type, as Scala classes are supposed to.⁴ Unlike `SortedView`, the abstract type `Ordering` is both upper- and lower-bounded.

Intervals can also encode abstract types missing a bound. Scala features a pair of extremal types `Any` and `Nothing`: the maximal type `Any` is a supertype of every other type; the minimal type `Nothing` is a subtype of every other type. Abstract types X with only an upper or lower bound A thus inhabit the degenerate intervals $X >: \text{Nothing} <: A$ and $X >: A <: \text{Any}$, respectively.

This suggests a uniform treatment of **F1–F3** through type intervals, and indeed, this is essentially how bounded quantification (**F1**) and type definitions (**F3**) are modeled in DOT. Unfortunately, DOT lacks intrinsics for higher-order type computation, such as type operator abstractions and applications, preventing it from encoding simple HK type definitions such as the identity operator `type Id[X] = X` [Odersky et al. 2016]. Traditionally, **F1–F3** have been studied through orthogonal extensions of Girard’s higher-order polymorphic λ -calculus F_ω [1972]. Bounded higher-order subtyping (**F1** and **F2**) has been formalized in variants of $F_{<}^\omega$: [Compagnoni and Goguen 2003; Pierce and Steffen 1997], translucent type definitions (**F3**) through singleton kinds [Stone and Harper 2000]. The treatment of **F1** and **F3** in DOT suggest a different, unified approach to studying **F1–F3**: via a formal theory of higher-order subtyping with type intervals, which we pursue with $F_{<}^\omega$.

Our goal in doing so is twofold. First, we want to establish a theoretical foundation for Scala’s HK types, with full support for type-level computations (including **F1–F3**). A principled theoretical understanding of HK types is crucial because potential safety issues are hard to identify and fix by “trial and error” alone, especially when they arise from feature interactions. Second, we want to study the concept of type intervals in its own right. Despite their apparent simplicity, adding type intervals to $F_{<}^\omega$ leads to a surprisingly rich theory of higher-order subtyping that goes beyond previous treatments of **F1–F3**. That is because type intervals encode first-class subtyping constraints, or *type inequations*, similar to extensional identity types in Martin-Löf type theory (MLTT).

In DOT, type intervals are baked into abstract type members and therefore tied to the use of *path-dependent types*; in $F_{<}^\omega$, we break this bond. A DOT type member declaration $\{X: A .. B\}$ roughly corresponds to a Scala-style type member declaration `class C { type X >: A <: B }`. It combines two separate type-system features: the declaration of an abstract type member X in a record or class, and the declaration of subtyping constraints on X via the bounds A and B . These two features are independent. For example, the Agda programming language features unbounded abstract type members via record types while the $F_{<}^\omega$ calculus features subtyping constraints via bounded quantification but no type members. The notion of path-dependent types in DOT and Scala is intimately linked to abstract type members. Given an instance $z: \{X: A .. B\}$, the type expression $z.X$ denotes the type value assigned to X in z . The type $z.X$ is *path-dependent* because it depends on the term-level expression z (the “path” to X). In DOT (but not Scala), type members are also used to model bounded quantification.

In $F_{<}^\omega$, we deliberately separate the notion of type intervals from that of abstract type members (and path-dependent types) and drop the latter. This simplifies the theory and allows us to study the power of type intervals in the context of higher-order subtyping: for bounded quantification, bounded operator abstraction and translucent type definitions – all of which are independent of path-dependent types, yet commonly used when working with Scala’s HK types.

⁴For details on how Scala classes may be encoded in DOT, we refer the reader to Amin [2016, Chap. 2].

We leave the development of a combined theory of HK and path-dependent types for future work, and focus here on the theoretical and practical insights afforded by the novel combination of *higher-order subtyping with type intervals*. Concretely, we make the following contributions.

- (1) We propose *type intervals* as a unifying concept for expressing bounded quantification, bounded operator abstractions, and translucent type definitions. Going beyond the status quo, we show that type intervals are expressive enough to also cover less familiar constructs, such as lower-bounded operator abstractions and first-class inequations (§2).
- (2) We introduce F^{ω} – an extension of F_{ω} with *interval kinds* – as a formal calculus of higher-order subtyping with type intervals (§3). F^{ω} is the first formalization of Scala’s higher-kinded types with a rigorous, machine-checked type safety proof. As such, it provides a theoretical foundation for several important features of Scala-like type systems.
- (3) We establish important metatheoretic properties of our theory: kind safety (§3), weak normalization of types (§4), type safety (§5), and undecidability of subtyping (§6). The metatheoretic proofs are complicated substantially by the interaction of advanced type system features such as dependent kinds, subtyping and subkinding, and (in)equality reflection. As others have recognized [Aspinall and Compagnoni 2001; Yang and Oliveira 2017; Zwanenburg 1999], the combination of dependent types (or kinds) and subtyping poses a particular challenge in metatheoretic developments. The usefulness of our proof techniques thus extends beyond the scope of F^{ω} to other systems combining dependent types and subtyping.
- (8) The metatheoretic development is entirely syntactic (it involves no model constructions) and has been fully mechanized using the Agda proof assistant [Norell 2007]. The main technical device is a purely syntactic, bottom-up normalization procedure based on a novel variant of *hereditary substitution* that computes the $\beta\eta$ -normal forms of types and kinds (§4).

We outline our proof strategy in §3.6, review related work in §7 and give concluding remarks in §8.

Because of space constraints, we omit most proofs and many details of the metatheory from the paper and focus instead on the big picture: the design and expressiveness of F^{ω} as well as the many challenges involved in proving its type safety and our strategies for addressing them. However, the complete metatheory, including full proofs of all lemmas and theorems stated in the paper, has been mechanized in Agda, and the source code is freely available as an artifact [Stucki and Giarrusso 2021]. An overview of the Agda formalization, establishing the connection to the theory presented in the paper, is included in the appendix, along with detailed human-readable descriptions of metatheoretic results that have been omitted from the paper. Yet more details can be found in the first author’s PhD dissertation [Stucki 2017].

2 SUBTYPING WITH TYPE INTERVALS

Before we define our formal theory of type intervals, let us illustrate the core ideas in a bit more detail. Consider the following Scala type definitions.

```
abstract class Bounded[B, F[_ <: B]] { def apply[X <: B]: F[X] }
type All[F[_]] = Bounded[Any, F]
```

The class `Bounded` and the type alias `All` are Scala encodings of the bounded and unbounded universal quantifiers found in F_{\leq} [Curien and Ghelli 1992]. We chose this example for its brevity and because it exemplifies the type-level mechanisms found in more realistic definitions, such as those given in §1. In particular, it features bounded quantification (of $X <: B$ in `apply`), a bounded operator (the parameter `F[_ <: B]` of `Bounded`) and a transparent type alias (`All`).

We want to translate these two definitions into a typed λ -calculus. The challenge is to find a type system that is expressive enough to do so. The example involves type-level computations, so our

first candidate is Girard's higher-order polymorphic λ -calculus F_ω [1972], but it lacks even basic support for subtyping. Our next candidate is F_{\leq}^ω , which extends F_ω with higher-order subtyping and bounded quantification. But most variants of F_{\leq}^ω lack support for bounded operators [cf. Pierce and Steffen 1997; Pierce 2002]. Thankfully, Compagnoni and Goguen have developed $\mathcal{F}_{\leq}^\omega$, a variant of F_{\leq}^ω with bounded operators [2003]. $\mathcal{F}_{\leq}^\omega$ has four type variable binders:

$\lambda X \leq A : K . t$ term-level type abstraction $\forall X \leq A : K . B$ type-level bounded quantifier
 $\lambda X \leq A : K . B$ type-level type abstraction $(X \leq A : K) \rightarrow J$ kind-level dependent arrow

The type A in a binding $X \leq A : K$ is called the *upper bound* of X , and must be of kind K . To represent unconstrained bindings, $\mathcal{F}_{\leq}^\omega$ features a *top* type \top , which is a supertype of every other type (like `Any` in Scala). The bound \top in $X \leq \top : *$ is thus trivially satisfied and can be omitted.

Since operator abstractions carry bounds in $\mathcal{F}_{\leq}^\omega$, so must arrow kinds. This makes arrow kinds type-dependent, which substantially complicates the meta theory of $\mathcal{F}_{\leq}^\omega$ when compared to other variants of F_{\leq}^ω . As usual, we abbreviate $(X : J) \rightarrow K$ to $J \rightarrow K$ when X does not occur freely in K .

A possible translation of `Bounded` and `All` to $\mathcal{F}_{\leq}^\omega$ is

$Bounded := \lambda B : * . \lambda F : (X \leq B : *) \rightarrow * . \forall X \leq B : * . F X$ $All := Bounded \top$

The named, parametrized class `Bounded[B, F[<: B]]` has been replaced by a pair of nested anonymous operator abstractions taking arguments $B : *$ and $F : (X \leq B : *) \rightarrow *$; the signature `apply[X <: B] : F[X]` of the method `apply` by the bounded universal $\forall X \leq B : * . F X$.

The declared kinds of the variables B, F and X in the definition of *Bounded* indicate what sort of type they represent: B and X are proper types, while F is a unary bounded operator. This makes *Bounded* itself a higher-order type operator of kind $* \rightarrow ((X \leq B : *) \rightarrow *) \rightarrow *$. For example, we obtain the type of the polymorphic identity function by applying *Bounded* as follows:

$Bounded \top (\lambda X : * . X \rightarrow X) = (\lambda B : * . \lambda F : (X \leq B : *) \rightarrow * . \forall X \leq B : * . F X) \top (\lambda X : * . X \rightarrow X)$
 $= \forall X : * . X \rightarrow X.$

The translation of the type alias `All[F[_]]` is then just the partial application $All = Bounded \top$.

The above definitions of *Bounded* and *All* are meta-definitions, i.e. they are just convenient shorthands for the type expressions $\lambda B : * . \lambda F : (X \leq B : *) \rightarrow * . \forall X \leq B : * . F X$ and $Bounded \top$. But we can also give object-level definitions of *Bounded* and *All* in $\mathcal{F}_{\leq}^\omega$, using standard syntactic sugar for let-binding type and term variables:

let $X : K = A$ **in** $t := (\lambda X : K . t) A,$ **let** $x : B = s$ **in** $t := (\lambda x : B . t) s.$

We can use *Bounded* as an abstract type operator in a term t by let-binding it to a type variable:

let $Bounded : (B : *) \rightarrow ((X \leq B : *) \rightarrow *) \rightarrow * = \lambda B : * . \lambda F : (X \leq B : *) \rightarrow * . \forall X \leq A : * . F X$ **in** $t.$

This definition is *opaque*, i.e. the term t sees the signature of *Bounded*, but not its definition. Consider

let $Bounded : (B : *) \rightarrow ((X \leq B : *) \rightarrow *) \rightarrow * = \dots$ **in**
let $x : \forall X : * . X \rightarrow X$ $= \lambda X : * . \lambda z : X . z$ **in** $— OK$
let $y : Bounded \top (\lambda X : * . X \rightarrow X)$ $= \lambda X : * . \lambda z : X . z$ **in** \dots $— type error$

The third definition does not type check because $Bounded \neq \lambda B : * . \lambda F : (X \leq B : *) \rightarrow * . \forall X : * . F X$ as types, despite the binding. Indeed, $\mathcal{F}_{\leq}^\omega$ cannot express *transparent* type definitions.

Furthermore, $\mathcal{F}_{\leq}^\omega$ also lacks support for *lower-bounded* definitions. As discussed in §1, these have important applications e.g. in the Scala standard library. Both transparent and lower-bounded definitions, and all the features of $\mathcal{F}_{\leq}^\omega$, can be uniformly expressed using *interval kinds*.

2.1 Intervals and Singletons

As the name implies, an interval kind $A .. B$ is inhabited by a range of proper types C , namely those that are supertypes $C \geq A$ of its lower bound A and subtypes $C \leq B$ of its upper bound B . Hence, kinding statements of the form $C : A .. B$ are equivalent to pairs of subtyping statements $A \leq C$ and $C \leq B$. We make this equivalence formal in the next section.

Since every proper type is a subtype of \top , intervals of the form $A .. \top$ are effectively unconstrained from above, and can thus be used to encode *lower-bounded* definitions. Similarly, upper-bounded definitions can be expressed using intervals of the form $\perp .. A$ where the lower bound is the minimum or *bottom* type \perp , our equivalent of Scala's `Nothing` type. For example, we recover F_{\leq} -style bounded quantifiers $\forall X \leq B. A$ as $\forall X : \perp .. B. A$. Interval kinds of the form $A .. A$, where the lower and upper bounds coincide, are called *singleton kinds* or simply *singletons*. Given $B : A .. A$, we have both $A \leq B$ and $B \leq A$, which, assuming an antisymmetric subtyping relation, implies $A = B$. Singleton kinds can thus encode transparent definitions and have been studied for that purpose by [Stone and Harper \[2000\]](#). We adopt their notation $S(A) = A .. A$ for the singleton containing just A .

Using interval kinds, we refine our definitions of `Bounded` and `All` to make them transparent.

$$\begin{aligned} \text{let } \text{Bounded} &: (B : *) \rightarrow (F : \perp .. B \rightarrow *) \rightarrow S(\forall X : \perp .. B. F X) \\ &= \lambda B : *. \lambda F : \perp .. B \rightarrow *. \forall X : \perp .. B. F X \quad \mathbf{in} \\ \text{let } \text{All} &: (F : * \rightarrow *) \rightarrow S(\text{Bounded } \top F) = \text{Bounded } \top \quad \mathbf{in} \dots \end{aligned}$$

The signature of `Bounded` tells us that, when we apply it to suitable type arguments B and F , the result is both a subtype and a supertype of $\forall X : \perp .. B. F X$. In other words, we have $\text{Bounded } B F = \forall X : \perp .. B. F X$ in the body of the `let`-binding. Similarly, we have $\text{All } F = \text{Bounded } \top F$, as desired.

Interval kinds $A .. B$ are only well-formed if A and B are proper types, i.e. of kind $*$. To express all of the binders found in $\mathcal{F}_{\leq}^{\omega}$, we need a way to encode bindings of the form $X \leq A : K$, for arbitrary kinds K . As we will see in §3, this is indeed possible because F^{ω} can encode *higher-order interval kinds* $A .._K B$ for arbitrary K using its other kind- and type-level constructs.

2.2 First-Class Inequalities

Instances $C : A .. B$ of an interval kind $A .. B$ represent types bounded by A and B respectively. But they also represent proofs that $A \leq C$ and $C \leq B$, and – by transitivity of subtyping – that $A \leq B$. In other words, the inhabitants of interval kinds $A .. B$ represent first-class *type inequations* $A \leq B$. Similarly, higher-order intervals represent type operator inequations. Interval kinds thus provide us with a mechanism for *(in)equality reflection*, i.e. a way to extend the subtyping relation via assumptions made at the term- or type-level (via type abstractions).

Among other things, this allows us to postulate type operators with associated subtyping rules through type variable bindings. We will see an example of this in §6; other examples are intersection types or equi-recursive types and their associated subtyping theories (see [Appendix E](#) for a detailed example). This is possible because we do not impose any *consistency constraints* on the bounds of intervals. That is, an interval kind $A .. B$ is well-formed, irrespective of whether $A \leq B$ is actually provable or not. If we can prove that $A \leq B$, we say that the bounds of $A .. B$ are *consistent*.

Having both (in)equality reflection and inconsistent bounds makes F^{ω} very expressive, but breaks subject reduction of open terms and decidability of (sub)typing. This is common in type theories with equality reflection (e.g. extensional MLTT [[Nordström et al. 1990](#)]) because they allow the reflection of absurd assumptions. For example, in a context where $Z : \top .. \perp$, we have $\top \leq Z \leq \perp$,

x, y, z, \dots	Term variable	X, Y, Z, \dots	Type variable
$s, t ::= x \mid \lambda x:A. t \mid s t \mid \lambda X:K. t \mid t A$			Term
$u, v ::= \lambda x:A. t \mid \lambda X:K. t$			Value
$A, B, C ::= X \mid \top \mid \perp \mid A \rightarrow B \mid \forall X:K. A \mid \lambda X:K. A \mid A B$			Type
$\Gamma, \Delta ::= \emptyset \mid \Gamma, x:A \mid \Gamma, X:K$			Typing context
$J, K, L ::= A..B \mid (X:J) \rightarrow K$	Kind	$j, k, l ::= * \mid j \rightarrow k$	Shape (simple kind)

Fig. 1. Syntax of F^ω .

i.e. the subtyping relation becomes trivial, and we can type non-terminating and stuck terms.

$$\begin{aligned}
 (\lambda x:\top. x x) (\lambda x:\top. x x) : \top & \quad - \textit{non-terminating, but } \top \leq Z \leq \perp \leq \top \rightarrow \top \\
 (\lambda X:*. \lambda x:X. x) (\lambda x:\top. x) : \top & \quad - \textit{stuck, but } \forall X:*. X \rightarrow X \leq \top \leq Z \leq \perp \leq \top \rightarrow \top
 \end{aligned}$$

It is therefore unsafe to reduce terms under absurd assumptions in general. Note that these examples do not break type safety of F^ω overall though. The absurd assumption $Z : \top.. \perp$ can never be instantiated, and hence reduction of *closed* terms remains perfectly safe. We discuss this point in more detail at the end of §3. The use of inconsistent bounds to prove undecidability of subtyping is more subtle; we return to it in §6.

Seeing the trouble inconsistent bounds can cause, one may wonder why we do not just enforce consistency of interval bounds statically. There are several reasons.

- *Statically enforcing consistent bounds in Scala is hard.* While we could statically enforce consistent bounds in F^ω , Amin et al. [2014] have shown that this would not extend to systems closer to Scala; a detailed explanation is given by Amin [2016, Sec. 3.4.4, 4.2.3].
- *Inconsistent bounds are useful.* Intervals with unconstrained bounds are useful, e.g. to encode generalized algebraic datatypes (GADTs) via first-class inequality constraints [Cretin and Rémy 2014; Parreaux et al. 2019]. Consistency of such constraints cannot be established when a GADT is defined, only when it is instantiated.
- *Decidability of subtyping could be recovered.* Even if we enforced consistent bounds in F^ω , subtyping would likely remain undecidable because F^ω , like full F_\leq and all variants of the DOT calculus, use a strong subtyping rule for universals that is a known source of undecidability [Pierce 1992]. Recent work by Hu and Lhoták [2019] suggests a novel approach for algorithmic subtyping that handles both inconsistent bounds and strong subtyping for universals. Whether or not their approach can be generalized to higher-order subtyping is a question we leave for future work.

3 THE DECLARATIVE SYSTEM

In this section, we introduce F^ω – our formal theory of higher-order subtyping with type intervals. We present its syntax and its type system, and establish some basic metatheoretic properties – just enough to show that subject reduction holds for well-kinded open types. Finally, we discuss the challenges involved in proving type safety, and outline our strategy for doing so.

3.1 Syntax

The syntax of F^ω is given in Fig. 1. The syntax of terms and types is identical to that of F_ω except for the extremal type constants \top and \perp . The *top type* \top is the maximal proper type: any other proper type is a subtype of \top . Dually, the *bottom type* \perp is the minimal proper type. Following Pierce [2002], λs carry domain annotations. This will become important in §3.2, §4.3 and §4.4.

Kind constants	Higher-order type intervals	$\boxed{A .._K B}$ $\boxed{*_K}$
$* := \perp .. \top$	$A .._{A' .. B'} B := A .. B$	$*_{A .. B} := \perp .. \top$
$\emptyset := \top .. \perp$	$A .._{(X:J) \rightarrow K} B := (X:J) \rightarrow A X .._K B X$ for $X \notin \text{fv}(A) \cup \text{fv}(B)$	$*_{(X:J) \rightarrow K} := (X:J) \rightarrow *_K$
Higher-order extrema	$\boxed{\top_K}$ $\boxed{\perp_K}$	Type erasure $\boxed{ K }$
$\top_{A .. B} := \top$	$\perp_{A .. B} := \perp$	$ A .. B := *$
$\top_{(X:J) \rightarrow K} := \lambda X:J. \top_K$	$\perp_{(X:J) \rightarrow K} := \lambda X:J. \perp_K$	$ (X:J) \rightarrow K := J \rightarrow K $
Bounded quantification and type operators		
$\forall X \leq A:K. B := \forall X:(\perp_K) .._K A. B$	$(X \leq A:J) \rightarrow K := (X:(\perp_J) .._J A) \rightarrow K$	
$\lambda X \leq A:K. t := \lambda X:(\perp_K) .._K A. t$	$\lambda X \leq A:K. B := \lambda X:(\perp_K) .._K A. B$	

Fig. 2. Syntactic shorthands and encodings

The main differences between F^ω and other variants of F_ω are reflected in its kind language. First, the usual kind of proper types $*$ is replaced by the *interval kind* former $A .. B$. The interval $A .. B$ is inhabited by exactly those proper types that are supertypes of A and subtypes of B . The degenerate interval $\perp .. \top$ spans *all* proper types. Hence we use $*$ as a shorthand for $\perp .. \top$. Second, most variants of F_ω have a *simple* kind language (as described by the non-terminal k in Fig. 1). In contrast, F^ω has a *dependent* kind language. The arrow kind $(X:J) \rightarrow K$ acts as a binder for the type variable X which may appear freely in the codomain K . Dependent kinds play an important role when modeling *bounded type operators*. For example, consider a binary type operator of kind $(X:*) \rightarrow (Y:\perp .. X) \rightarrow *$. The upper-bounded kind $\perp .. X$ of Y ensures that the operator can only be applied to types A, B if B is a subtype of A . This idea goes back to [Compagnoni and Goguen's \$\mathcal{F}_{\leq}^\omega\$](#) , which features both upper-bounded type operators and dependent arrow kinds [2003].

We abbreviate $\perp .. \top$ by $*$ and $(X:J) \rightarrow K$ by $J \rightarrow K$ when X is not free in K . This allows us to treat *simple kinds* or *shapes* k as a subset of (dependent) kinds K . In the opposite direction, we define an *erasure* map $|K|$ which forgets any dependencies in K (see Fig. 2). Given a kind K , we say K has *shape* $|K|$. Unlike kinds, shapes are stable under substitution, i.e. $|K[A/X]| \equiv |K|$.

Following [Barendregt \[1992\]](#), we identify expressions e (terms, types and kinds), up to α -*equivalence* and assume that the names of bound and free variables are distinct. We write $e \equiv e'$ to stress that e and e' are α -equivalent. The set of free variables of e is denoted by $\text{fv}(e)$, and we write $e[t/x]$ and $e[A/X]$ for capture-avoiding term and type substitutions in e , respectively. We require that the variables bound in a typing context Γ be distinct so that we may think of Γ as a finite map and use function notation, such as $\text{dom}(\Gamma)$, $\Gamma(x)$, $\Gamma(X)$. We write (Γ, Δ) for the concatenation of two contexts Γ and Δ with disjoint domains, and we often omit the empty context \emptyset , writing e.g. $\Gamma = x:A, Y:K$ instead of $\Gamma = \emptyset, x:A, Y:K$.

3.1.1 Encodings. Together with the extremal types \top and \perp , interval kinds allow us to express *bounded quantification* and *bounded operators* over proper types. For example, the F_{\leq} -style universal type $\forall X \leq A. B$ can be expressed as $\forall X:\perp .. A. B$ in F^ω . To extend this principle to *higher-order* bounded quantification and type operators, we define encodings for higher-order interval kinds and extremal types via type abstraction and dependent kinds in Fig. 2. The encoding of *higher-order maxima* \top_K is standard [cf. [Compagnoni and Goguen 2003](#); [Pierce 2002](#)]; that of *higher-order*

$\text{minima } \perp_K$ follows the same principle. The encoding of *higher-order interval kinds* $A .._K B$ resembles that of higher-order *singleton kinds* given by Stone and Harper [2000]. Indeed, singleton kinds are just interval kinds where the upper and lower bounds coincide. Encodings of higher-order F_{\leq}^{ω} -style bounded operators and universal quantifiers are also given in Fig. 2.

3.1.2 Structural Operational Semantics. For computations in terms, we adopt the standard call-by-value (CBV) semantics given by Pierce [2002, Fig. 30-1], writing $t \longrightarrow_v^* t'$ when the term t CBV-reduces in one or more steps to t' . For types and kinds, we define the *one-step β -reduction* relation \longrightarrow_{β} as the compatible closure of β -contraction of type operators w.r.t. all the type and kind formers. We write $\longrightarrow_{\beta}^*$ for its reflexive, transitive closure, *β -reduction*.

3.2 Declarative Typing and Kinding

The static semantics of F_{\leq}^{ω} are summarized in Figs. 3 and 4. We refer to this set of judgments as the *declarative* system, as opposed to the *canonical* system introduced in §5. We sometimes write $\Gamma \vdash \mathcal{J}$ to denote an arbitrary judgment of the declarative system. Throughout the paper, we silently assume that judgments are *well-scoped*, i.e. if $\Gamma \vdash \mathcal{J}$ then $\text{fv}(\mathcal{J}) \subseteq \text{dom}(\Gamma)$. We now discuss each of the judgments, emphasizing novel rules.

3.2.1 Context and Kind Formation. The rules for context formation $\Gamma \text{ ctx}$ are standard. They ensure that the type and kind annotations of all bindings are well-formed. The rules of the remaining judgments are set up so that they can only be derived in well-formed contexts.

Our kind formation judgment $\Gamma \vdash K \text{ kd}$ ensures that (a) all types appearing in K are well-kinded, and (b) that the bounds of intervals are proper types (not λ s), forbidding for instance $\perp .. \lambda X : K. A$. The formation rule **WF-DARR** for dependent arrows is standard. An interval $A .. B$ is well-formed if A and B are proper types. As discussed in §2.2, we choose not to enforce $A \leq B$; e.g. the empty kind $\emptyset = \top .. \perp$ is well-formed. We say that the bounds of an interval $A .. B$ are *consistent* in Γ if $\Gamma \vdash A \leq B : *$ and *inconsistent* otherwise. If A and B are closed types and $\not\vdash A \leq B : *$ in the empty context, we say that the bounds of $A .. B$ are *absurd*. For example, the bounds of $*$ are always consistent while those of $\top .. \perp$ are absurd.⁵ The bounds of $X .. Y$ are inconsistent in $\Gamma = X : *, Y : *$ but not absurd because X and Y are open types.

3.2.2 Kinding and Typing. The kinding rules **K-VAR**, **K-TOP**, **K-BOT**, **K-ARR** and **K-ALL** are all standard. The rule **K-ALL** resembles that found in F_{ω} : no bound annotations are needed because the bounds of X are internalized in the kind K . Similarly, **K-ABS** and **K-APP**, resemble those in F_{ω} more than those in F_{\leq}^{ω} . The rules **K-SING** and **K-SUB** are used to adjust the kind of a type: **K-SUB** is the kind-level analog of **T-SUB** (subsumption); **K-SING** resembles Stone and Harper's singleton introduction rule [2000]. Note that **K-SING** only *narrows* the kind of a type whereas **K-SUB** only *widens* it. The premise of **K-SING** may look a bit surprising: why use $\Gamma \vdash A : B .. C$ instead of $\Gamma \vdash A : *$? This extra flexibility is necessary to prove that types inhabiting intervals are proper types, i.e. that $\Gamma \vdash A : B .. C$ implies $\Gamma \vdash A : *$. Thus the relaxed premise justifies itself.

The typing rules are again entirely standard, with the possible exception of some additional context and kind formation premises that would be redundant in variants of F_{ω} with simple kinds.

3.2.3 Subkinding and Subtyping. F_{\leq}^{ω} features both subtyping and *subkinding*. The use of subkinding in F_{\leq}^{ω} is both natural and essential. If a type is contained in an interval $A .. B$, then one naturally expects it to also be contained in a *wider* interval $A' .. B'$ where $A' \leq A$ and $B \leq B'$. This is captured in the rule **SK-INTV**. Subkinding is also essential. It is thanks to **SK-INTV** that we can express bounded polymorphism and bounded type operators in F_{\leq}^{ω} . Consider e.g. a polymorphic

⁵See Lemma 5.4 in §5.2.

Context formation $\Gamma \text{ ctx}$

$$\frac{}{\emptyset \text{ ctx}} \text{ (C-EMPTY)} \quad \frac{\Gamma \text{ ctx} \quad \Gamma \vdash K \text{ kd}}{\Gamma, X:K \text{ ctx}} \text{ (C-TMBIND)} \quad \frac{\Gamma \text{ ctx} \quad \Gamma \vdash A : *}{\Gamma, x:A \text{ ctx}} \text{ (C-TPBIND)}$$

Kind formation $\Gamma \vdash K \text{ kd}$

$$\frac{\Gamma \vdash A : * \quad \Gamma \vdash B : *}{\Gamma \vdash A .. B \text{ kd}} \text{ (WF-INTV)} \quad \frac{\Gamma \vdash J \text{ kd} \quad \Gamma, X:J \vdash K \text{ kd}}{\Gamma \vdash (X:J) \rightarrow K \text{ kd}} \text{ (WF-DARR)}$$

Kinding $\Gamma \vdash A : K$

$$\frac{\Gamma \text{ ctx} \quad \Gamma(X) = K}{\Gamma \vdash X : K} \text{ (K-VAR)} \quad \frac{\Gamma \text{ ctx}}{\Gamma \vdash \top : *} \text{ (K-TOP)} \quad \frac{\Gamma \text{ ctx}}{\Gamma \vdash \perp : *} \text{ (K-BOT)}$$

$$\frac{\Gamma \vdash A : * \quad \Gamma \vdash B : *}{\Gamma \vdash A \rightarrow B : *} \text{ (K-ARR)} \quad \frac{\Gamma \vdash K \text{ kd} \quad \Gamma, X:K \vdash A : *}{\Gamma \vdash \forall X:K. A : *} \text{ (K-ALL)}$$

$$\frac{\Gamma \vdash J \text{ kd} \quad \Gamma, X:J \vdash A : K \quad \Gamma, X:J \vdash K \text{ kd}}{\Gamma \vdash \lambda X:J. A : (X:J) \rightarrow K} \text{ (K-ABS)} \quad \frac{\Gamma \vdash A : (X:J) \rightarrow K \quad \Gamma \vdash B : J \quad \Gamma, X:J \vdash K \text{ kd} \quad \Gamma \vdash K[B/X] \text{ kd}}{\Gamma \vdash AB : K[B/X]} \text{ (K-APP)}$$

$$\frac{\Gamma \vdash A : B .. C}{\Gamma \vdash A : A .. A} \text{ (K-SING)} \quad \frac{\Gamma \vdash A : J \quad \Gamma \vdash J \leq K}{\Gamma \vdash A : K} \text{ (K-SUB)}$$

Typing $\Gamma \vdash t : A$

$$\frac{\Gamma \text{ ctx} \quad \Gamma(x) = A}{\Gamma \vdash x : A} \text{ (T-VAR)} \quad \frac{\Gamma \vdash K \text{ kd} \quad \Gamma, X:K \vdash t : A}{\Gamma \vdash \lambda X:K. t : \forall X:K. A} \text{ (T-TABS)}$$

$$\frac{\Gamma \vdash A : * \quad \Gamma \vdash B : * \quad \Gamma, x:A \vdash t : B}{\Gamma \vdash \lambda x:A. t : A \rightarrow B} \text{ (T-ABS)} \quad \frac{\Gamma \vdash t : \forall X:K. A \quad \Gamma \vdash B : K}{\Gamma \vdash tB : A[B/X]} \text{ (T-TAPP)}$$

$$\frac{\Gamma \vdash s : A \rightarrow B \quad \Gamma \vdash t : A}{\Gamma \vdash st : B} \text{ (T-APP)} \quad \frac{\Gamma \vdash t : A \quad \Gamma \vdash A \leq B : *}{\Gamma \vdash t : B} \text{ (T-SUB)}$$

Fig. 3. Declarative presentation of F_{ω}^{ω} – part 1. Premises in gray are validity conditions (see §3.3).

term $t : \forall X:\perp .. A. B$ and a type argument C such that $C \leq A : *$. We can apply t to C because C has kind $C .. C$ (by **K-SING**) which in turn is a subkind of $\perp .. A$ (by **SK-INTV**). The rule **SK-DARR** lifts the interval containment order through dependent arrow kinds. It resembles **Aspinall and Compagnoni's** subtyping rule ($s\text{-}\pi$) for dependent product types **Aspinall and Compagnoni** [2001].

Subtyping judgments $\Gamma \vdash A \leq B : K$ are indexed by the common kind K in which A and B are related. Note that two types may be related in some kinds but not others. For example, the extremal types \top and \perp are related as proper types, i.e. $\perp \leq \top : *$, but not as inhabitants of

Subkinding

$$\boxed{\Gamma \vdash J \leq K}$$

$$\frac{\Gamma \vdash A_2 \leq A_1 : * \quad \Gamma \vdash B_1 \leq B_2 : *}{\Gamma \vdash A_1 .. B_1 \leq A_2 .. B_2} \text{ (SK-INTV)} \quad \frac{\Gamma \vdash (X:J_1) \rightarrow K_1 \text{ kd} \quad \Gamma \vdash J_2 \leq J_1 \quad \Gamma, X:J_2 \vdash K_1 \leq K_2}{\Gamma \vdash (X:J_1) \rightarrow K_1 \leq (X:J_2) \rightarrow K_2} \text{ (SK-DAARR)}$$

Subtyping

$$\boxed{\Gamma \vdash A \leq B : K}$$

$$\frac{\Gamma \vdash A : K}{\Gamma \vdash A \leq A : K} \text{ (ST-REFL)} \quad \frac{\Gamma \vdash A \leq B : K \quad \Gamma \vdash B \leq C : K}{\Gamma \vdash A \leq C : K} \text{ (ST-TRANS)}$$

$$\frac{\Gamma \vdash A : B .. C}{\Gamma \vdash A \leq \top : *} \text{ (ST-TOP)} \quad \frac{\Gamma \vdash A : B .. C}{\Gamma \vdash \perp \leq A : *} \text{ (ST-BOT)}$$

$$\frac{\Gamma, X:J \vdash A : K \quad \Gamma \vdash B : J \quad \Gamma \vdash A[B/X] : K[B/X] \quad \Gamma, X:J \vdash K \text{ kd} \quad \Gamma \vdash K[B/X] \text{ kd}}{\Gamma \vdash (\lambda X:J. A) B \leq A[B/X] : K[B/X]} \text{ (ST-}\beta_1) \quad \frac{\Gamma, X:J \vdash A : K \quad \Gamma \vdash B : J \quad \Gamma \vdash A[B/X] : K[B/X] \quad \Gamma, X:J \vdash K \text{ kd} \quad \Gamma \vdash K[B/X] \text{ kd}}{\Gamma \vdash A[B/X] \leq (\lambda X:J. A) B : K[B/X]} \text{ (ST-}\beta_2)$$

$$\frac{\Gamma \vdash A : (X:J) \rightarrow K \quad X \notin \text{fv}(A)}{\Gamma \vdash \lambda X:J. A X \leq A : (X:J) \rightarrow K} \text{ (ST-}\eta_1) \quad \frac{\Gamma \vdash A : (X:J) \rightarrow K \quad X \notin \text{fv}(A)}{\Gamma \vdash A \leq \lambda X:J. A X : (X:J) \rightarrow K} \text{ (ST-}\eta_2)$$

$$\frac{\Gamma \vdash A_2 \leq A_1 : * \quad \Gamma \vdash B_1 \leq B_2 : *}{\Gamma \vdash A_1 \rightarrow B_1 \leq A_2 \rightarrow B_2 : *} \text{ (ST-ARR)} \quad \frac{\Gamma \vdash \forall X:K_1. A_1 : * \quad \Gamma \vdash K_2 \leq K_1 \quad \Gamma, X:K_2 \vdash A_1 \leq A_2 : *}{\Gamma \vdash \forall X:K_1. A_1 \leq \forall X:K_2. A_2 : *} \text{ (ST-ALL)}$$

$$\frac{\Gamma \vdash \lambda X:J_1. A_1 : (X:J) \rightarrow K \quad \Gamma \vdash \lambda X:J_2. A_2 : (X:J) \rightarrow K \quad \Gamma, X:J \vdash A_1 \leq A_2 : K}{\Gamma \vdash \lambda X:J_1. A_1 \leq \lambda X:J_2. A_2 : (X:J) \rightarrow K} \text{ (ST-ABS)} \quad \frac{\Gamma \vdash A_1 \leq A_2 : (X:J) \rightarrow K \quad \Gamma \vdash B_1 = B_2 : J \quad \Gamma \vdash B_1 : J \quad \Gamma, X:J \vdash K \text{ kd} \quad \Gamma \vdash K[B_1/X] \text{ kd}}{\Gamma \vdash A_1 B_1 \leq A_2 B_2 : K[B_1/X]} \text{ (ST-APP)}$$

$$\frac{\Gamma \vdash A : B_1 .. B_2}{\Gamma \vdash B_1 \leq A : *} \text{ (ST-BND}_1) \quad \frac{\Gamma \vdash A : B_1 .. B_2}{\Gamma \vdash A \leq B_2 : *} \text{ (ST-BND}_2)$$

$$\frac{\Gamma \vdash A_1 \leq A_2 : B .. C}{\Gamma \vdash A_1 \leq A_2 : A_1 .. A_2} \text{ (ST-INTV)} \quad \frac{\Gamma \vdash A_1 \leq A_2 : J \quad \Gamma \vdash J \leq K}{\Gamma \vdash A_1 \leq A_2 : K} \text{ (ST-SUB)}$$

Kind equality

$$\boxed{\Gamma \vdash J = K}$$

Type equality

$$\boxed{\Gamma \vdash A = B : K}$$

$$\frac{\Gamma \vdash J \leq K \quad \Gamma \vdash K \leq J}{\Gamma \vdash J = K} \text{ (SK-ANTI-SYM)} \quad \frac{\Gamma \vdash A \leq B : K \quad \Gamma \vdash B \leq A : K}{\Gamma \vdash A = B : K} \text{ (ST-ANTI-SYM)}$$

Fig. 4. Declarative presentation of F^{ω} – part 2. Premises in gray are validity conditions (see §3.3).

their respective singleton kinds $\perp .. \perp$ and $\top .. \top$. For a given context Γ and kind K , the subtyping relation $\Gamma \vdash A \leq B : K$ is a preorder, as witnessed by the rules **ST-REFL** and **ST-TRANS**. Note that there are no such rules for subkinding, but it is easy to prove them admissible. In §5, we will see that some (but not all) instances of **ST-REFL** and **ST-TRANS** can be eliminated too.

The rules **ST-TOP** and **ST-BOT** establish \top and \perp as the maximum and minimum proper types w.r.t. subtyping. The premise $\Gamma \vdash A : B .. C$ ensures that the extremal types are only related to other proper types. As for **K-SING**, the kind $B .. C$ in the premise allows us to prove that types inhabiting intervals are proper types:

$$\frac{\text{(K-SING)} \frac{\Gamma \vdash A : B .. C}{\Gamma \vdash A : A .. A} \quad \text{(ST-BOT)} \frac{\Gamma \vdash A : B .. C}{\Gamma \vdash \perp \leq A : *} \quad \text{(ST-TOP)} \frac{\Gamma \vdash A : B .. C}{\Gamma \vdash A \leq \top : *} \quad \text{(SK-INTV)} \frac{\Gamma \vdash A : B .. C}{\Gamma \vdash A \leq \top : *}}{\Gamma \vdash A .. A \leq \perp .. \top} \text{(K-SUB)}$$

This derivation would not be possible if the rules **K-SING**, **ST-BOT** or **ST-TOP** had premise $\Gamma \vdash A : *$.

The rules **ST- β_1** and **ST- β_2** correspond to β -contraction and expansion, respectively. Two separate rules are needed because subtyping is not symmetric. We could have combined them into a single type equality rule but that would have complicated the definition of type equality. Similarly, the rules **ST- η_1** and **ST- η_2** relate η -convertible types. The rule for universals resembles **SK-DARR**.

Most variants of F_{\leq}^{ω} separate subtyping of type operator applications into a subtyping rule that only compares the heads of applications, and a congruence rule for type equality w.r.t. application. Here we fuse these two rules into a single subtyping rule **ST-APP**. Since we do not track the variance of type operators, the arguments must be equal types. Because arrow kinds are dependent, either B_1 or B_2 must be substituted for X in K in the conclusion. Both are equally suitable; we pick B_1 .

The rule for subtyping operator abstractions, **ST-ABS**, is maybe the most unusual when compared to other variants of F_{ω} since it allows abstractions to be subtypes even if their domain annotations J_1 and J_2 are not subkinds. Other systems adopt weaker versions of this rule where $J \equiv J_2$ or even $J \equiv J_1 \equiv J_2$. But such rules are not suitable for a theory featuring both subkinding and η -equality. Let $A : * \rightarrow *$ be an operator and $B : *$ a type in Γ . By **K-SUB**, **ST-SUB**, **ST-ETA_{1,2}** and antisymmetry,

$$\Gamma \vdash \lambda X : \perp .. B. AX = A = \lambda X : B .. \top. AX : (X : B .. B) \rightarrow *$$

i.e. the two η -expansions of A are equal as types, despite having distinct domain annotations. Because the η -rules allow such equations, we adopt a compatible subtyping rule for abstractions. The first two premises of **ST-ABS** ensure that both abstractions – irrespective of their domain annotations – inhabit the common arrow kind $(X : J) \rightarrow K$; the remaining premise ensures that the bodies of the two abstractions are pointwise subtypes assuming the common domain $X : J$. Note that systems without domain annotations avoid such complications [cf. [Abel 2008](#)].

So far, we have seen how a type interval $A .. B$ can be formed using **WF-INTV**, and introduced using **K-SING**, but we have yet to see how the bounds A and B of the interval can be put to use. Type intervals are “eliminated” by turning them into subtyping judgments via a pair of *bound projection* rules **ST-BND₁** and **ST-BND₂**. Given a type $A : B .. C$, the rules **ST-BND₁** and **ST-BND₂** assert that B and C are indeed lower and upper bounds, respectively, of A . When A is a variable, we may use rule **ST-BND₂** to derive judgments of the form $\Gamma \vdash X \leq C$, similar to those obtained using the variable subtyping rule from F_{\leq} . More generally, the bound projection rules allow us to *reflect* any well-formed assumption $\Gamma(X) = A .. B$ – consistent or not – into a corresponding subtyping judgment $\Gamma \vdash A \leq B : *$. We discuss the ramifications this has for type safety in §3.5.

As for kinding judgments, there are two subtyping rules that allow us to adjust the kinds of subtyping judgments, **ST-SUB** and **ST-INTV**. The former is the analog of **K-SUB** for subtyping, whereas the latter is the subtyping counterpart of the interval introduction rule **K-SING**: if A and B are subtypes in some interval $C .. D$, then surely they are still subtypes in the interval $A .. B$ bounded

by those very same types. Indeed, $A..B$ is the smallest interval in which the two types are related. The **ST-INTV** rule plays an important role in the proof of subject reduction for types and kinds (**Theorem 3.3**) because it allows us to relate β -equal types inhabiting singleton kinds.

3.2.4 Kind and Type Equality. The kind and type equality judgments are each generated by exactly one rule: **SK-ANTI-SYM** for kind equality and **ST-ANTI-SYM** for type equality. In most variants of F_ω with subtyping, the subtyping relation is not defined to be antisymmetric. Instead antisymmetry may or may not be an admissible property that has to be proven [cf. **Compagnoni and Goguen 1999**]. In F_ω^ω , antisymmetry is not an admissible property, however. To see this, consider the context $\Gamma = X:A..A$ for some proper type A . Then $\Gamma \vdash X : A..A$, and we can derive that X and A are mutual subtypes using **ST-BND₁** and **ST-BND₂**. But without antisymmetry we have no way to derive $\Gamma \vdash X = A : *$. Faced with this issue, we could have chosen to add a *singleton reflection* rule for deriving $\Gamma \vdash X = A : *$ from $\Gamma \vdash X : A..A$ directly, such as the one due to **Stone and Harper [2000]**. Interestingly, antisymmetry for proper types is derivable from **Stone and Harper's** rule and other rules about type intervals. We conjecture that antisymmetry of subtyping under arbitrary kinds would also have been admissible in such a system, albeit at the cost of a more complicated type equality judgment. We prefer the simpler judgment with an explicit antisymmetry rule.

3.3 Basic Metatheoretic Properties

With the dynamics and statics in place, we can begin our work on the metatheory of F_ω^ω . Our system enjoys the usual basic metatheoretic properties, such as preservation of all the judgments under weakening, substitution and narrowing of contexts, as well as admissibility of the missing order-theoretic and congruence rules for subkinding, kind equality and type equality. Although these properties constitute the foundation on which we build the remainder of our metatheory, they are also entirely standard, and little insight is gained by spelling them out in detail. We therefore relegate them to **Appendix B**. There, the reader will also find a collection of admissible rules that justify the encodings of the higher-order extrema and interval kinds given in **§3.1.1**. These include formation, subtyping and subkinding rules for the encoded kinds, and typing rules for introducing and eliminating the more familiar forms of bounded universals. Unlike the other admissible rules, they are not important for the remainder of the metatheoretic development.

There are two standard properties of the declarative system that are exceptional in that their proofs are not routine inductions, namely *validity* of the various judgments and *functionality* of substitutions. Roughly, a judgment is valid if all its parts are well-formed.

Lemma 3.1 (validity).

(kinding validity) If $\Gamma \vdash A : K$, then Γ ctx and $\Gamma \vdash K$ kd.

(typing validity) If $\Gamma \vdash t : A$, then Γ ctx and $\Gamma \vdash A : *$.

(kind (in)equation validity) If $\Gamma \vdash J = K$ or $\Gamma \vdash J \leq K$, then $\Gamma \vdash J$ kd and $\Gamma \vdash K$ kd.

(type (in)equation validity) If $\Gamma \vdash A = B : K$ or $\Gamma \vdash A \leq B : K$, then $\Gamma \vdash A : K$ and $\Gamma \vdash B : K$.

The validity lemma provides a “sanity check” for the static semantics, but it also plays a crucial role in the proofs of other important properties, such as subject reduction or soundness of type normalization. Unfortunately, it is harder to prove than one might expect. The proofs of kinding, subkinding and subtyping validity require the following *functionality* lemma for the case of **ST-APP**.

Lemma 3.2 (functionality). Let $\Gamma \vdash A_1 = A_2 : K$.

(1) If $\Gamma, X:K \vdash J$ kd, then $\Gamma \vdash J[A_1/X] = J[A_2/X]$.

(2) If $\Gamma, X:K \vdash B : J$, then $\Gamma \vdash B[A_1/X] = B[A_2/X] : J[A_1/X]$.

The proof of functionality, in turn, depends on kinding and subtyping validity. Proving the two statements by simultaneous induction is not enough to resolve the circular dependency because the proof of functionality would require us to apply the IH to derivations obtained via validity. Since these are not generally sub-derivations of the relevant premise, the induction does not go through.

Instead, we follow Harper and Pfenning [2005] and establish validity by “temporarily extending” certain rules of the declarative system with additional premises, which we call *validity conditions*, shown in gray in Figs. 3 and 4. We then prove functionality and validity for the extended system, show that the two systems are equivalent and the validity conditions are redundant after all, and obtain Lemma 3.1 for the original system. For details see Appendix B.3.

3.4 Subject Reduction for Well-Kinded Types

For most versions of F_ω , subject reduction for types is easy to prove because types are simply-kinded. In F_ω^ω , the proof is complicated by the presence of type-dependent kinds and subkinding. However these complications are minor since there are only two *shapes* of kinds – intervals and arrows – with exactly one subkinding rule per shape. Hence subkinding is easy to invert.

To prove subject reduction, we show that β -reduction steps can be lifted to type and kind equality.

Theorem 3.3.

- (1) If $\Gamma \vdash J$ kd and $J \longrightarrow_\beta K$, then $\Gamma \vdash J = K$.
- (2) If $\Gamma \vdash A : K$ and $A \longrightarrow_\beta B$, then $\Gamma \vdash A = B : K$.

Subject reduction for kinds and types then follows immediately from Theorem 3.3 and validity.

Corollary 3.4 (subject reduction for kinding).

- (1) If $\Gamma \vdash J$ kd and $J \longrightarrow_\beta^* K$, then $\Gamma \vdash K$ kd.
- (2) If $\Gamma \vdash A : K$ and $A \longrightarrow_\beta^* B$, then $\Gamma \vdash B : K$.

3.5 The Long Road to Type Safety

After establishing subject reduction for well-kinded types, we prove type safety via *progress* and *preservation* (aka subject reduction) [Wright and Felleisen 1994]. But as we show in this section, we must first weaken the statement of preservation for it to hold in F_ω^ω .

Preservation typically applies to all open terms:

Proposition 3.1 (preservation). *If $\Gamma \vdash t : A$ and $t \longrightarrow_v t'$, then $\Gamma \vdash t' : A$.*

However, in F_ω^ω this statement fails because reduction of open terms is unsafe. The culprit are type variable bindings with absurd bounds. Consider the following example. Let v be the polymorphic identity function $v = \lambda X : *. \lambda x : X. x$ which is of type $A = \forall X : *. X \rightarrow X$. In F_ω^ω , closed universals are not subtypes of closed arrows;⁶ hence v cannot be applied to itself. For the same reason, the term application $t = (\lambda x : A. x) v v$ is ill-typed as a closed term. Yet, t is well-typed in the context $\Gamma = X : (A \rightarrow A) .. (A \rightarrow A \rightarrow A)$ because we can use **ST-BND₁** and **ST-BND₂** to derive $\Gamma \vdash A \rightarrow A \leq X \leq A \rightarrow A \rightarrow A : *$ and subsumption to derive $\Gamma \vdash \lambda x : A. x : A \rightarrow A \rightarrow A : *$. Note that the bounds of X are proper types, so its kind is well-formed, as is the context Γ . But since $\not\leq A \leq A \rightarrow A : *$, the bounds of the interval are absurd.

Next, consider what happens when t takes a reduction step.

$$(\lambda x : A. x) v v \longrightarrow_v (x[v/x]) v \equiv v v.$$

According to preservation, the application $v v$ should have type A , but instead it is ill-typed, even in Γ . The assumption $X : (A \rightarrow A) .. (A \rightarrow A \rightarrow A)$ is useless here, since v does not have type

⁶See Lemma 5.4 in §5.2.

$A \rightarrow A$. Not only is $v v$ ill-typed, it is also stuck. Hence $v v$ is neither a value nor can it be reduced further – type safety clearly does not hold in Γ .

But all is not lost. Type safety still holds for closed terms, as does a weaker form of preservation.

Proposition 3.2 (preservation – weak version). *If $\vdash t : A$ and $t \rightarrow_v t'$, then $\vdash t' : A$.*

Throughout the next two sections, we will work our way towards a proof of this proposition.

3.6 Challenges and Proof Strategy

To conclude the section, let us briefly explore the challenges involved in proving weak preservation and our strategy to address them. The complexity of the subtyping relation throws a spanner in the works when we try to prove weak preservation for cases where β -contractions occur. To prove these cases, one normally starts by showing that the following rules are admissible:

$$\frac{\Gamma \vdash A_1 \rightarrow B_1 \leq A_2 \rightarrow B_2 : *}{\Gamma \vdash A_2 \leq A_1 : * \quad \Gamma \vdash B_1 \leq B_2 : *} \quad \frac{\Gamma \vdash \forall X:K_1. A_1 \leq \forall X:K_2. A_2 : *}{\Gamma \vdash K_2 \leq K_1 \quad \Gamma, X:K_2 \vdash A_1 \leq A_2 : *}$$

These properties are generally known as *inversion of subtyping*, and are closely related to the Π -injectivity property, which is a well-known source of complexity in dependent type theories. There are several features of subtyping that severely complicate the proof of subtyping inversion.

- (1) The rules for β and η -conversion, together with transitivity, may change the shapes of related types in the middle of a subtyping derivation, e.g. from a type former to a type application.

$$\Gamma \vdash A_1 \rightarrow A_2 \leq (\lambda X:*. X \rightarrow A_2) A_1 \leq \dots \leq (\lambda X:*. X \rightarrow B_2) B_1 \leq B_1 \rightarrow B_2 : *$$

- (2) The subsumption rule **ST-SUB** may change the kinds of related types at any time.
- (3) As outlined above, we can derive judgments of the form $\Gamma \vdash A \leq X \leq B : *$ where A and B need not be of the same shape, from absurd assumptions in Γ . For example

$$\Gamma \vdash A \rightarrow B \leq X \leq \forall X:K. C : *$$

We address these points as follows. First, we eliminate uses of the $\beta\eta$ rules (1) by adopting an alternative presentation of subtyping which we dub *canonical subtyping* (§5). Canonical subtyping only relates types in η -long β -normal form (§4), so there is no need for $\beta\eta$ rules. This approach works in variants of F_ω with η -conversion [cf. Abel and Rodriguez 2008], whereas rewriting-based proofs of Π -injectivity [e.g. Adams 2006; Barendregt 1992] do not generalize readily to our setting.

The canonical presentation of subtyping also restricts the placement of subsumption (2) to certain strategic positions, just as in algorithmic or bidirectional subtyping (§5).

Finally, we avoid issues caused by absurd bounds (3) by proving subtyping inversion only in the empty context, i.e. only for closed types. That suffices for proving weak preservation and, as the above example illustrates, it is the best we can do in a system with inequality reflection (§5.2).

The core challenges of the proof of subtyping inversion thus consists in showing that well-formed kinds and well-kinded types have normal forms (§4), so that the canonical and declarative versions of subtyping can be proven equivalent (§5). We address these challenges in the next two sections.

4 NORMALIZATION OF TYPES

As discussed in the previous section, we cannot prove inversion of subtyping directly, because $\beta\eta$ -convertible types and kinds differ in their syntactic structure. We address this problem in two steps: (1) in this section, we show that types and kinds in F_ω^ω can be reduced to $\beta\eta$ -normal form via a *bottom-up normalization* procedure based on *hereditary substitution*; (2) in next section, we give a canonical presentation of subtyping that only relates types in normal form.

4.1 Syntax

We begin by introducing an alternative syntax for types that is better suited to our definition of hereditary substitution. The key difference is that the type arguments of applications are grouped together in sequences called *spines*. Hence, we refer to this presentation of types as *spine form*.

$$D, E ::= F \mathbf{E} \quad F, G ::= X \mid \top \mid \perp \mid D \rightarrow E \mid \forall X:K. E \mid \lambda X:K. E \quad \mathbf{D}, \mathbf{E} ::= \epsilon \mid D, \mathbf{E}$$

Applications form a separate syntactic category, *eliminations* E , and consist of a *head* F and a spine \mathbf{E} . A head is any type former that is not an application. We adopt vector notation for spines, writing \mathbf{E} for the sequence $\mathbf{E} = E_1, E_2, \dots, E_n$ and (\mathbf{D}, \mathbf{E}) for the concatenation of \mathbf{D} and \mathbf{E} .

The two representations of types are isomorphic, so we mix them freely, knowing that explicit conversions can always be inserted where necessary.

4.2 Hereditary Substitution in Raw Types

In the §5, we will introduce a system of canonical judgments defined directly on normal forms. Since kinds in F^ω are dependent, some of the kinding and subtyping rules involve substitutions in kinds, e.g. **K-APP** or **ST- $\beta_{1,2}$** . Unfortunately, substitutions do not preserve normal forms because substituting an operator abstraction for the head of a neutral type introduces a new redex. For example $(Y A)[\lambda X:K. B/Y] \equiv (\lambda X:K. B) A$ is not a normal form, even if $Y A$ and $\lambda X:K. B$ are. To define a canonical counterpart of e.g. **K-APP** directly on normal kinds and types, we need a variant of substitution that immediately eliminates the β -redexes it introduces. This type of substitution operation is known as *hereditary substitution* [Watkins et al. 2004]. The challenge in defining a hereditary substitution function is, of course, ensuring its totality.

Our definition of hereditary substitution is given in Fig. 5. Hereditary substitution is defined mutually with *reducing application* of eliminations by recursion on the structure of the shape k . The definitions of the three hereditary substitution functions proceed by inner recursion on the structure of the parameters K , D and \mathbf{D} , respectively. Note that there are some recursive calls where no parameter decreases, but one can check (and we have done so) that at least one of the relevant parameters decreases strictly along every cycle in the call graph. Hence the five functions remain structurally recursive, ensuring their totality. Note the crucial use of spine forms: $D \cdot^k \mathbf{E}$ simultaneously unwinds \mathbf{E} and k using the fact that \mathbf{E} matches the right-associative structure of k .

Our presentation of hereditary substitution differs from others in the literature. Like Keller and Altenkirch [2010], we define hereditary substitution by structural recursion and mutually with reducing application. But their definition is based on an intrinsically typed representation, which does not readily generalize to a system with dependent types (or kinds). Instead, like Abel and Rodriguez [2008] we define hereditary substitution directly on raw (i.e. unkinded) types, so our definition contains degenerate cases; unlike Abel and Rodriguez's, our definition is structurally recursive hence easier to mechanize. Our approach of defining hereditary substitution by recursion on shapes rather than (dependent) kinds was inspired by Harper and Licata's formalization of Canonical LF [2007]. However, they define hereditary substitution as an inductive relation, thus they avoid degenerate cases but must establish functionality and termination separately.

Because the essential difference between ordinary and hereditary substitution is that the latter reduces newly created β -redexes, the results of the two operations are β -convertible.

Lemma 4.1. *Let E be an elimination, X a type variable and k a shape, then*

- (1) $K[E/X] \longrightarrow_\beta^* K[E/X^k]$ for any kind K ;
- (2) $D[E/X] \longrightarrow_\beta^* D[E/X^k]$ for any type D ;
- (3) $E D \longrightarrow_\beta^* E \cdot^k D$ for any type D ;
- (4) $E \mathbf{D} \longrightarrow_\beta^* E \cdot^k \mathbf{D}$ for any spine \mathbf{D} .

It is an immediate consequence of Lemma 4.1 and subject reduction that ordinary and hereditary substitutions produce judgmentally equal results.

Hereditary substitution $D[E/X^k]$

$$(Y \mathbf{D})[E/X^k] = \begin{cases} E.^k (\mathbf{D}[E/X^k]) & \text{if } Y = X, \\ Y (\mathbf{D}[E/X^k]) & \text{otherwise,} \end{cases}$$

$$(\top \mathbf{D})[E/X^k] = \top (\mathbf{D}[E/X^k])$$

$$(\perp \mathbf{D})[E/X^k] = \perp (\mathbf{D}[E/X^k])$$

$$((D_1 \rightarrow D_2) \mathbf{D})[E/X^k] = (D_1[E/X^k] \rightarrow D_2[E/X^k]) (\mathbf{D}[E/X^k])$$

$$((\forall Y:K. D') \mathbf{D})[E/X^k] = (\forall Y:K[E/X^k]. D'[E/X^k]) (\mathbf{D}[E/X^k]) \quad \text{for } Y \neq X, Y \notin \text{fv}(E),$$

$$((\lambda Y:K. D') \mathbf{D})[E/X^k] = (\lambda Y:K[E/X^k]. D'[E/X^k]) (\mathbf{D}[E/X^k]) \quad \text{for } Y \neq X, Y \notin \text{fv}(E).$$

$$\epsilon[E/X^k] = \epsilon$$

 $\mathbf{D}[E/X^k]$

$$(D', \mathbf{D})[E/X^k] = (D'[E/X^k], (\mathbf{D}[E/X^k]))$$

$$(D_1 .. D_2)[E/X^k] = D_1[E/X^k] .. D_2[E/X^k]$$

 $K[E/X^k]$

$$((Y:J) \rightarrow K)[E/X^k] = (Y:J[E/X^k]) \rightarrow K[E/X^k] \quad \text{for } Y \neq X, Y \notin \text{fv}(E).$$

Reducing application $D.^k E$ $D.^k \mathbf{E}$

$$D.^* E = DE$$

$$D.^k \epsilon = D$$

$$D.^{k \rightarrow l} E = \begin{cases} D'[E/X^k] & \text{if } D = \lambda X:J. D', \\ DE & \text{otherwise.} \end{cases}$$

$$D.^k (E, \mathbf{E}) = \begin{cases} (D.^{k_1 \rightarrow k_2} E).^k_2 \mathbf{E} & \text{if } k = k_1 \rightarrow k_2, \\ D(E, \mathbf{E}) & \text{otherwise.} \end{cases}$$

Fig. 5. Hereditary substitution and reducing application

Corollary 4.2 (soundness of hereditary substitution). *Let $\Gamma \vdash A : K$, then*

(1) *if $\Gamma, X:K, \Delta \vdash J \text{ kd}$, then $\Gamma, \Delta[A/X] \vdash J[A/X] = J[A/X]^{|K|}$;*

(2) *if $\Gamma, X:K, \Delta \vdash B : J$, then $\Gamma, \Delta[A/X] \vdash B[A/X] = B[A/X]^{|K|} : J[A/X]$.*

4.3 Normalization of Raw Types

Based on hereditary substitution, we define a bottom-up *normalization* function nf on kinds and types. It is a straightforward extension of the normalization function by [Abel and Rodriguez \[2008\]](#) to dependent kinds. The function nf is defined directly on raw types and kinds and relies on a separate function for η -expanding variables. The definition of both functions is given in [Fig. 6](#).

The η -expansion $\eta_K(A)$ of a type A of kind K is defined by recursion on the structure of K . It is used in the definition of nf to expand type variables. Note that $\eta_K(A)$ immediately expands newly introduced argument variables to produce η -long forms. Normalization $\text{nf}_\Gamma(A)$ and $\text{nf}_\Gamma(K)$ of raw types and kinds in context Γ are defined by mutual recursion on A and K , respectively. The case of applications uses hereditary substitution to eliminate β -redexes. Note the crucial use of domain-annotations: in order to hereditarily substitute a type argument in the body of an operator abstraction $\lambda X:K. E$, we need to guess its shape, or equivalently, the shape of X . Since the normalization function is defined directly on raw, unkinded types, the only way to obtain this information is from the kind annotation K in the abstraction.

The context parameter Γ of nf is used to look up the declared kinds of variables, which drive their η -expansion. To ensure that the resulting η -expansions are normal, the context Γ must itself

η -expansion

$$\eta_{D_1..D_2}(A) = A \quad \eta_{(X:J) \rightarrow K}(A) = \lambda X:J. \eta_K(A(\eta_J(X))) \quad \text{for } X \notin \text{fv}(A).$$

 $\eta_K(A)$ **Normalization**

$$\begin{aligned} \text{nf}_\Gamma(X) &= \eta_{\Gamma(X)}(X) && \text{if } X \in \text{dom}(\Gamma), \\ &X && \text{otherwise,} \\ \text{nf}_\Gamma(A) &= A && \text{for } A \in \{\perp, \top\}, \\ \text{nf}_\Gamma(A \rightarrow B) &= \text{nf}_\Gamma(A) \rightarrow \text{nf}_\Gamma(B) \\ \text{nf}_\Gamma(\forall X:K. A) &= \forall X:K'. \text{nf}_{\Gamma, X:K'}(A) && \text{where } K' = \text{nf}_\Gamma(K), \\ \text{nf}_\Gamma(\lambda X:K. A) &= \lambda X:K'. \text{nf}_{\Gamma, X:K'}(A) && \text{where } K' = \text{nf}_\Gamma(K), \\ \text{nf}_\Gamma(AB) &= E[\text{nf}_\Gamma(B)/X^{|K|}] && \text{if } \text{nf}_\Gamma(A) = \lambda X:K. E, \\ &(\text{nf}_\Gamma(A)) (\text{nf}_\Gamma(B)) && \text{otherwise.} \end{aligned}$$

 $\text{nf}_\Gamma(K)$

$$\begin{aligned} \text{nf}_\Gamma(A..B) &= \text{nf}_\Gamma(A) .. \text{nf}_\Gamma(B) \\ \text{nf}_\Gamma((X:J) \rightarrow K) &= (X:J') \rightarrow \text{nf}_{\Gamma, X:J'}(K) \quad \text{where } J' = \text{nf}_\Gamma(J). \end{aligned}$$

Fig. 6. Normalization of types and kinds

be normal. We therefore extend normalization pointwise to contexts, defining $\text{nf}(\Gamma)$ as

$$\text{nf}(\emptyset) = \emptyset \quad \text{nf}(\Gamma, x:A) = \text{nf}(\Gamma), x:\text{nf}_{\text{nf}(\Gamma)}(A) \quad \text{nf}(\Gamma, X:K) = \text{nf}(\Gamma), X:\text{nf}_{\text{nf}(\Gamma)}(K)$$

Since nf is a total function defined directly on raw types and kinds, it necessarily contains degenerate cases, i.e. the resulting types need not be β -normal. For example, the case of applications relies on the domain annotations K of operator abstractions $\lambda X:K. A$ in head position to be truthful. The ill-kinded type $\Omega = (\lambda X:*. X X)(\lambda X:*. X X)$ will result in $\text{nf}(\Omega) \equiv (X X)[\lambda X:*. X X/X^*] \equiv (\lambda X:*. X X) \cdot^* (\lambda X:*. X X) \equiv \Omega$. However, for well-kinded types $\Gamma \vdash A : K$, the type $\text{nf}_\Gamma(A)$ is guaranteed to be an η -long β -normal form, as we will see in §5 (cf. Lemma 5.3).

Furthermore, η -expansion and normalization are *sound*, i.e. they do not alter the meaning of types and kinds. In particular, well-kinded types and well-formed kinds are judgmentally equal to their normalized counterparts. The proof relies on soundness of hereditary substitutions.

Lemma 4.3 (soundness of normalization).

- (1) If $\Gamma \vdash K$ kd, then $\Gamma \vdash K = \text{nf}_{\text{nf}(\Gamma)}(K)$. (2) If $\Gamma \vdash A : K$, then $\Gamma \vdash A = \text{nf}_{\text{nf}(\Gamma)}(A) : K$.

4.4 Commutativity of Normalization and Hereditary Substitution

We are now almost ready to introduce the canonical presentation of F^ω . Our final task in this section is to establish a series of *commutativity properties* of hereditary substitution and normalization. They say, roughly, that the order of these operations can be switched without changing the result. We require these properties to prove equivalence of the canonical and declarative systems. For example, to prove that **ST-BETA**₁ is admissible in the canonical system, we must show that

$$\Gamma \vdash \text{nf}_\Gamma(A)[\text{nf}_\Gamma(B)/X^{|J|}] = \text{nf}_\Gamma(A[B/X]) : \text{nf}_\Gamma(K[B/X]).$$

Since normalization involves hereditary substitutions, we must further show that these preserve the canonical judgments. The case for applications involves kind equations of the form

$$\Gamma \vdash K[A/Y^k][B/X^j] = K[B/X^j][A[B/X^j]/Y^k].$$

Unfortunately, the commutativity properties do not hold for arbitrary raw types and kinds. The reasons are twofold. First, our definition of hereditary substitution contains degenerate cases for ill-kinded inputs, which can cause inconsistencies when we commute hereditary substitutions. For example, it's easy to verify that for $B = \lambda Z: * \rightarrow *. Z$

$$(XA)[Y/X^*][B/Y^{**}] \equiv A \neq BA \equiv (XA)[B/Y^{**}][Y[B/Y^{**}]/X^*].$$

Second, normalization involves η -expansion and, as we have seen in §3.2, η -expansions of the same type variable can differ syntactically in their domain annotations.

We address the two problems separately. For the former, we adopt the approach taken by [Abel and Rodriguez \[2008\]](#), namely to prove commutativity of hereditary substitutions only for well-kinded normal forms. To apply their technique, we first need to show that hereditary substitutions preserve kinding (of normal forms). This is easy in their setting, which is simply kinded, but challenging in ours because our kinding rules involve substitutions in dependent kinds. A direct proof that hereditary substitutions preserve kinding would require the very commutativity lemmas we are trying to establish. We circumvent this issue by relaxing our requirements: for $e[V/Y^k]$ to be non-degenerate, the normal form V need not actually be well-kinded; it only needs to have *shape* k . Using this insight, we prove commutativity of hereditary substitutions in 4 steps.

- (1) We define a *simple kinding judgment* that assign shapes (rather than kinds) to normal forms.
- (2) We show that hereditary substitution preserves simple kinding. Because shapes have no type dependencies, the proof does not require any commutativity lemmas.
- (3) We show that hereditary substitutions in simply kinded normal forms commute.
- (4) We show that simple kinding is sound: every type of kind K has a normal form of shape $|K|$.

We refer the reader to [Appendix C](#) for details.

It remains to show that normalization commutes with substitution. To work around the issue of incongruous domain annotations, we introduce an auxiliary equivalence $A \approx B$ on types and kinds, called *weak equality*, that identifies operator abstractions up to the *shape* of their domain annotations, i.e. $\lambda X: J. A \approx \lambda X: K. A$ when $|J| \equiv |K|$. Substitution then *weakly commutes* with normalization of well-formed kinds and well-kinded types.

Lemma 4.4. *Let $\Gamma \vdash A : J$ and $V = \text{nf}_{\text{nf}(\Gamma)}(A)$, then*

- (1) *if $\Gamma, X: J, \Delta \vdash K \text{ kd}$, then $\text{nf}_{\text{nf}(\Gamma, \Delta[A/X])}(K[A/X]) \approx (\text{nf}_{\text{nf}(\Gamma, X: J, \Delta)}(K))[V/X^{|J|}]$;*
- (2) *if $\Gamma, X: J, \Delta \vdash B : K$, then $\text{nf}_{\text{nf}(\Gamma, \Delta[A/X])}(B[A/X]) \approx (\text{nf}_{\text{nf}(\Gamma, X: J, \Delta)}(B))[V/X^{|J|}]$;*

The proof uses commutativity of hereditary substitutions and a few helper lemmas, e.g. that substitutions weakly commute with η -expansion. The full proof is given in [Appendix C.2.3](#).

5 THE CANONICAL SYSTEM

Having characterized the normal forms of kinds and types in the previous section, we now present our *canonical system* of judgments directly defined on normal forms, and summarize its most important metatheoretic properties: the hereditary substitution lemma, equivalence w.r.t. the declarative system, and inversion of subtyping. We conclude the section by revisiting the type safety proof outlined in §3.

The rules for canonical kinding, subtyping and spine equality are given in [Figs. 7](#) and [8](#). The canonical system also contains judgments for kind and context formation, subkinding, and type and kind equality, but the rules of those judgments are analogous to their declarative counterparts,

Canonical kinding of variables

$$\boxed{\Gamma \vdash_{\text{var}} X : K}$$

$$\frac{\Gamma \text{ ctx} \quad \Gamma(X) = K}{\Gamma \vdash_{\text{var}} X : K} \quad (\text{CV-VAR})$$

$$\frac{\Gamma \vdash_{\text{var}} X : J \quad \Gamma \vdash J \leq K \quad \Gamma \vdash K \text{ kd}}{\Gamma \vdash_{\text{var}} X : K} \quad (\text{CV-SUB})$$

Spine kinding

$$\boxed{\Gamma \vdash J \Rightarrow \mathbf{V} \Rightarrow K}$$

$$\frac{}{\Gamma \vdash K \Rightarrow \epsilon \Rightarrow K} \quad (\text{CK-EMPTY}) \quad \frac{\Gamma \vdash U \Leftarrow J \quad \Gamma \vdash J \text{ kd} \quad \Gamma \vdash K[U/X^{|J|}] \Rightarrow \mathbf{V} \Rightarrow L}{\Gamma \vdash (X:J) \rightarrow K \Rightarrow U, \mathbf{V} \Rightarrow L} \quad (\text{CK-CONS})$$

Kinding of neutral types

$$\boxed{\Gamma \vdash_{\text{ne}} N : K}$$

Kinding checking

$$\boxed{\Gamma \vdash V \Leftarrow K}$$

$$\frac{\Gamma \vdash_{\text{var}} X : J \quad \Gamma \vdash J \Rightarrow \mathbf{V} \Rightarrow K}{\Gamma \vdash_{\text{ne}} X \mathbf{V} : K} \quad (\text{CK-NE})$$

$$\frac{\Gamma \vdash V \Rightarrow J \quad \Gamma \vdash J \leq K}{\Gamma \vdash V \Leftarrow K} \quad (\text{CK-SUB})$$

Kind synthesis for normal types

$$\boxed{\Gamma \vdash V \Rightarrow K}$$

$$\frac{\Gamma \text{ ctx}}{\Gamma \vdash \top \Rightarrow \top .. \top} \quad (\text{CK-TOP})$$

$$\frac{\Gamma \text{ ctx}}{\Gamma \vdash \perp \Rightarrow \perp .. \perp} \quad (\text{CK-BOT})$$

$$\frac{\Gamma \vdash U \Rightarrow U .. U \quad \Gamma \vdash V \Rightarrow V .. V}{\Gamma \vdash U \rightarrow V \Rightarrow (U \rightarrow V) .. (U \rightarrow V)} \quad (\text{CK-ARR}) \quad \frac{\Gamma \vdash K \text{ kd} \quad \Gamma, X:K \vdash V \Rightarrow V .. V}{\Gamma \vdash \forall X:K. V \Rightarrow (\forall X:K. V) .. (\forall X:K. V)} \quad (\text{CK-ALL})$$

$$\frac{\Gamma \vdash J \text{ kd} \quad \Gamma, X:J \vdash V \Rightarrow K}{\Gamma \vdash \lambda X:J. V \Rightarrow (X:J) \rightarrow K} \quad (\text{CK-ABS})$$

$$\frac{\Gamma \vdash_{\text{ne}} N : U .. V}{\Gamma \vdash N \Rightarrow N .. N} \quad (\text{CK-SING})$$

Fig. 7. Canonical presentation of F_{ne}^{ω} – part 1

so we omit them. We use the following naming conventions to distinguish normal forms: U, V, W denote normal types; M, N denote neutral types. No special notation is used for normal kinds.

The judgments for kinding, subtyping and spine equality are bidirectional. Double arrows are used to indicate whether a kind is an input ($K \Leftarrow$ or $\Leftarrow K$) or an output ($\Rightarrow K$) of the judgment. The contexts and subjects of judgments are always considered inputs. The rules for kind synthesis are similar to those for declarative kinding, except that the synthesized kinds are more precise, that there is no subsumption rule, and that kinding of variables and applications has been combined into a single rule **CK-SING** for kinding neutral proper types. A quick inspection of the rules reveals that all synthesized kinds are singletons. In particular, $\Gamma \vdash V \Rightarrow V .. V$ for proper types V . The kind checking judgment $\Gamma \vdash V \Leftarrow K$ has only one inference rule: the subsumption rule **CK-SUB**.

The canonical kinding judgments $\Gamma \vdash_{\text{var}} X : K$ for variables and $\Gamma \vdash_{\text{ne}} N : K$ for neutral types are not directed because of the presence of the subsumption rule **CV-SUB**. While this rule is not actually necessary for variable kinding, it considerably simplifies the metatheory. Without it, the proof of context narrowing would circularly depend on at least three other lemmas – transitivity of subkinding, functionality and the hereditary substitution lemma – all of which use somewhat idiosyncratic, possibly incompatible induction strategies.

Subtyping of proper types

$$\boxed{\Gamma \vdash U \leq V}$$

$$\frac{\Gamma \vdash V \Rightarrow V .. V}{\Gamma \vdash V \leq \top} \quad (\text{CST-Top})$$

$$\frac{\Gamma \vdash V \Rightarrow V .. V}{\Gamma \vdash \perp \leq V} \quad (\text{CST-Bot})$$

$$\frac{\Gamma \vdash U \leq V \quad \Gamma \vdash V \leq W}{\Gamma \vdash U \leq W} \quad (\text{CST-TRANS})$$

$$\frac{\Gamma \vdash U_2 \leq U_1 \quad \Gamma \vdash V_1 \leq V_2}{\Gamma \vdash U_1 \rightarrow V_1 \leq U_2 \rightarrow V_2} \quad (\text{CST-ARR})$$

$$\frac{\Gamma \vdash_{\text{var}} X : K \quad \Gamma \vdash K \Rightarrow \mathbf{V}_1 = \mathbf{V}_2 \Rightarrow U .. W}{\Gamma \vdash X \mathbf{V}_1 \leq X \mathbf{V}_2} \quad (\text{CST-NE})$$

$$\frac{\Gamma \vdash \forall X : K_1. V_1 \Rightarrow \forall X : K_1. V_1 .. \forall X : K_1. V_1 \quad \Gamma \vdash K_2 \leq K_1 \quad \Gamma, X : K_2 \vdash V_1 \leq V_2}{\Gamma \vdash \forall X : K_1. V_1 \leq \forall X : K_2. V_2} \quad (\text{CST-ALL})$$

$$\frac{\Gamma \vdash_{\text{ne}} N : V_1 .. V_2}{\Gamma \vdash V_1 \leq N} \quad (\text{CST-BND}_1)$$

$$\frac{\Gamma \vdash_{\text{ne}} N : V_1 .. V_2}{\Gamma \vdash N \leq V_2} \quad (\text{CST-BND}_2)$$

Checked subtyping

$$\boxed{\Gamma \vdash U \leq V \Leftarrow K}$$

$$\frac{\Gamma \vdash V_1 \Leftarrow U .. W \quad \Gamma \vdash V_2 \Leftarrow U .. W \quad \Gamma \vdash V_1 \leq V_2}{\Gamma \vdash V_1 \leq V_2 \Leftarrow U .. W} \quad (\text{CST-INTV})$$

$$\frac{\Gamma \vdash \lambda X : J_1. V_1 \Leftarrow (X : J) \rightarrow K \quad \Gamma \vdash \lambda X : J_2. V_2 \Leftarrow (X : J) \rightarrow K \quad \Gamma, X : J \vdash V_1 \leq V_2 \Leftarrow K}{\Gamma \vdash \lambda X : J_1. V_1 \leq \lambda X : J_2. V_2 \Leftarrow (X : J) \rightarrow K} \quad (\text{CST-ABS})$$

Spine equality

$$\boxed{\Gamma \vdash J \Rightarrow \mathbf{U} = \mathbf{V} \Rightarrow K}$$

$$\frac{}{\Gamma \vdash K \Rightarrow \epsilon = \epsilon \Rightarrow K} \quad (\text{SPeq-EMPTY})$$

$$\frac{\Gamma \vdash U_1 = U_2 \Leftarrow J \quad \Gamma \vdash K[U_1/X^{U_1}] \Rightarrow \mathbf{V}_1 = \mathbf{V}_2 \Rightarrow L}{\Gamma \vdash (X : J) \rightarrow K \Rightarrow U_1, \mathbf{V}_1 = U_2, \mathbf{V}_2 \Rightarrow L} \quad (\text{SPeq-CONS})$$

Fig. 8. Canonical presentation of F_{\leq}^{ω} – part 2

Canonical spine kinding $\Gamma \vdash J \Rightarrow \mathbf{V} \Rightarrow K$ differs from the other judgments in that it features both an input kind J and an output kind K . When an operator of shape J is applied to the spine \mathbf{V} , the resulting type is of shape K – as exemplified by the rule **CK-NE**. In **CK-CONS**, the head U of the spine U, \mathbf{V} is hereditarily substituted for X in the codomain K of the overall input kind $(X : J) \rightarrow K$ to obtain the input kind $K[U/X^{U_1}]$ for kinding the tail \mathbf{V} of the spine. The use of hereditary (rather than ordinary) substitution ensures that the resulting kind remains normal.

The rules for canonical subtyping are divided into two judgments: *subtyping of proper types* $\Gamma \vdash U \leq V$ and *checked subtyping* $\Gamma \vdash U \leq V \Leftarrow K$. The separate judgment for proper subtyping $\Gamma \vdash U \leq V$ simplifies the metatheory by disentangling subtyping and kinding. It resembles the subtyping judgment of F_{\leq} . Notable differences are the two bound projection rules **CST-BND₁** and **CST-BND₂**, which replace the variable subtyping rule, and the rule **CST-NE** for subtyping neutrals. The most interesting of these is **CST-NE**. It says that two neutral types $X \mathbf{V}_1$ and $X \mathbf{V}_2$ headed by a common type variable X are subtypes if they have canonically equal spines. Importantly, the spines \mathbf{V}_1 and \mathbf{V}_2 need not be syntactically equal. In F_{\leq}^{ω} , normal forms may be judgmentally equal yet differ syntactically, e.g. because they have different domain annotations (via **CST-ABS**), because one

of the types is declared as a type alias of the other (via a singleton kind), or because one of the types can be proven to alias the other due to inconsistent bounds: $\Gamma, X: U .. V, Y: V .. U, \Delta \vdash U = V$. The last example illustrates that there is no easy way for the normalization function nf to resolve such equations. In systems without dependent kinds and equality reflection, judgmentally equal types have syntactically equal normal forms, so that **CST-NE** can be omitted [see e.g. [Abel and Rodriguez 2008](#)]; in systems with singleton kinds (but no type intervals), type aliases can be resolved during normalization [see e.g. [Stone and Harper 2000](#)]. Neither of these apply in F^ω . Unfortunately, the presence of **CST-NE** complicates the metatheory of the canonical system (cf. §5.1.1).

The checked subtyping judgment $\Gamma \vdash U \leq V \Leftarrow K$ is kind-directed. The kind K determines whether U and V are compared as proper types (**CST-INTV**) or type operators (**CST-ABS**). The rule **CST-INTV** checks that the types V_1 and V_2 have the expected kind $U .. W$ and are proper subtypes. Because normal types are η -long, the only normal types of arrow kind are operator abstractions. They are compared using rule **CST-ABS**, exactly as in the declarative system. The rules of the *spine equality* judgment $\Gamma \vdash J \Rightarrow U = V \Rightarrow K$ resemble those of spine kinding.

5.1 Metatheoretic Properties of the Canonical System

It is easy to show that the canonical system is sound w.r.t. the declarative one, i.e. that normal forms related by the canonical judgments are also related by the declarative counterparts. To avoid confusion, we mark canonical judgments with the subscript “c” and declarative ones with “d” in the following soundness lemma. The full statement of the lemma has 13 parts, one for each canonical judgment; we only show the most important ones.

Lemma 5.1 (soundness of the canonical rules – excerpt).

- (1) If $\Gamma \vdash_c V \Rightarrow K$ or $\Gamma \vdash_c V \Leftarrow K$, then $\Gamma \vdash_d V : K$.
- (2) If $\Gamma \vdash_c U \leq V$, then $\Gamma \vdash_d U \leq V : *$.
- (3) If $\Gamma \vdash_c U \leq V \Leftarrow K$, then $\Gamma \vdash_d U \leq V : K$.

Many of the basic properties of the declarative system – weakening, context narrowing, admissible order-theoretic rules, many validity properties, and the commutativity lemmas from the previous section – also hold in the canonical system, and their proofs carry over with minor modifications. Full statements and proofs are given in [Appendix D.1](#). Notable exceptions are the substitution and functionality lemmas. These do not hold because ordinary substitutions do not preserve normal forms. Instead, we need to establish analogous lemmas for hereditary substitutions.

5.1.1 The Hereditary Substitution Lemma. The most important metatheoretic property of the canonical system is the *hereditary substitution lemma*, which states, roughly, that canonical judgments are preserved by hereditary substitutions of canonically well-kinded types. It is key to proving completeness w.r.t. the declarative system, and thus to our overall goal of establishing type safety. Proving and even stating the hereditary substitution lemma is challenging. The full statement of the lemma has 24 separate parts, all of which have to be proven simultaneously. The large number of canonical judgments is one reason for the complexity of the lemma. But the foremost reason is that the proof of the hereditary substitution lemma circularly depends on *functionality of the canonical judgments*. This circular dependency is caused by the subtyping rule **CST-NE**.

To illustrate this, assume we are given two normal forms V and W such that $\Gamma \vdash V = W \Leftarrow *$, but $V \not\equiv W$ syntactically. We have seen examples of such normal forms V and W earlier. Assume further that there is some X with $\Gamma(X) = * \rightarrow *$ and consider what happens when we hereditarily substitute the operator $\lambda Y:*. U$ for X in the judgment $\Gamma \vdash X V \leq X W$ obtained using **CST-NE**. We would like to show that hereditary substitution preserves this inequation, i.e. that

$$\Gamma \vdash (XV)[\lambda Y:*. U/X^{**}] \equiv U[V/Y^*] \leq U[W/Y^*] \equiv (XW)[\lambda Y:*. U/X^{**}],$$

which requires functionality. The example illustrates a second point, namely that, in order to prove that hereditary substitutions preserve canonical kinding and subtyping, we need to prove that kinding and subtyping of reducing applications is admissible. Our hereditary substitution lemma must cover all of these properties, leading to the aforementioned grand total of 24 parts. We give a small excerpt here, illustrating some of the properties just discussed.

Lemma 5.2 (hereditary substitution – excerpt). *Assume $\Gamma \vdash U_1 = U_2 \Leftarrow K$.*

(1) *Hereditary substitution preserves kind checking:*

if $\Gamma, X:K, \Delta \vdash V \Leftarrow J$, then $\Gamma, \Delta[U_1/X^{|K|}] \vdash V[U_1/X^{|K|}] \Leftarrow J[U_1/X^{|K|}]$.

(2) *Functionality/monotonicity of hereditary substitution:*

if $\Gamma, X:K, \Delta \vdash V \Leftarrow J$ and $\Gamma, X:K, \Delta \vdash J \text{ kd}$, then

$$\Gamma, \Delta[U_1/X^{|K|}] \vdash V[U_1/X^{|K|}] \leq V[U_2/X^{|K|}] \Leftarrow J[U_1/X^{|K|}].$$

(3) *Hereditary substitution preserves checked subtyping:*

if $\Gamma, X:K, \Delta \vdash V_1 \leq V_2 \Leftarrow J$ and $\Gamma, X:K, \Delta \vdash J \text{ kd}$, then

$$\Gamma, \Delta[U_1/X^{|K|}] \vdash V_1[U_1/X^{|K|}] \leq V_2[U_2/X^{|K|}] \Leftarrow J[U_1/X^{|K|}].$$

(4) *Canonical equality of reducing applications is admissible:*

if $\Gamma \vdash V_1 = V_2 \Leftarrow (X:K) \rightarrow J$, then $\Gamma \vdash V_1 \cdot^{|K| \rightarrow |J|} U_1 = V_2 \cdot^{|K| \rightarrow |J|} U_2 \Leftarrow J[U_1/X^{|K|}]$.

The structure of the proof mirrors that of the recursive definition of hereditary substitution itself. All 24 parts are proven simultaneously by induction in the structure of the shape $|K|$. Most parts proceed by an inner induction on the derivations of the judgments into which U_1 and U_2 are being substituted. The cases involving the rules [CK-CONS](#) and [SP-EQ-CONS](#), rely on commutativity of hereditary substitutions in kinds. Details are given in [Appendix D.2](#).

Just as for the declarative system, the proof of functionality enables us to prove some additional validity properties, which, in turn, are necessary to prove completeness of the canonical system.

5.1.2 Completeness of Canonical Kinding. In [§4](#), we saw that every declaratively well-kinded type has a judgmentally equal $\beta\eta$ -normal form ([Lemmas 4.3](#)). To establish equivalence of the canonical and declarative systems, it remains to show that the normal forms of types related by a declarative judgment are also canonically related. The full statement of the completeness lemma has 11 parts, one for each declarative judgment plus three auxiliary parts for dealing with hereditary substitutions and $\beta\eta$ -conversions. The most important ones are the following, where we again use the subscripts “c” and “d” to distinguish canonical judgments from declarative ones.

Lemma 5.3 (completeness of the canonical rules – excerpt).

(1) *If $\Gamma \vdash_d A : K$, then $\text{nf}(\Gamma) \vdash_c \text{nf}(A) \Leftarrow \text{nf}(K)$.*

(2) *If $\Gamma \vdash_d A \leq B : K$, then $\text{nf}(\Gamma) \vdash_c \text{nf}(A) \leq \text{nf}(B) \Leftarrow \text{nf}(K)$.*

The completeness proof relies on [Lemma 5.2](#) to show that the normal forms of applications are canonically well-kinded, and on the weak commutativity properties established in [§4.4](#) to show that β - and η -conversions are admissible in the canonical system. In addition, the proof relies on the validity conditions discussed in [§3.3](#). The full statement and proof of the completeness lemma are given in [Appendix D.3](#).

5.2 Inversion of Subtyping and Type Safety

As we saw in [§3](#), reductions in open terms are unsafe because variable bindings with inconsistent bounds can inject arbitrary inequations into the subtyping relation. Under such assumptions, subtyping cannot be inverted in any meaningful way. We therefore prove inversion of subtyping only in the empty context, following the approach by [Rompf and Amin \[2016\]](#):

- (1) We introduce a helper judgment $\vdash_{\text{tf}} U \leq V$, which states that U is a proper subtype of V in the empty context. It is obtained from the canonical subtyping judgment $\Gamma \vdash U \leq V$ by fixing $\Gamma = \emptyset$ and removing **CST-TRANS** and any rules involving free variables (**CST-NE**, **CST-BND₁** and **CST-BND₂**). Soundness of $\vdash_{\text{tf}} U \leq V$ w.r.t. canonical subtyping is immediate.
- (2) We prove that transitivity is admissible in $\vdash_{\text{tf}} U \leq V$. This is straightforward since there are no $\beta\eta$ -conversion rules or bound projection rules that get in the way.
- (3) Because transitivity is admissible, it is straightforward to establish completeness, and thus equivalence of $\vdash_{\text{tf}} U \leq V$ w.r.t. canonical subtyping.
- (4) Inversion of the canonical subtyping relation in the empty context then follows immediately by inspection of the rules for $\vdash_{\text{tf}} U \leq V$ and equivalence of the two judgments.
- (5) Inversion of top-level declarative subtyping follows by equivalence of canonical and declarative subtyping and soundness of normalization.

Lemma 5.4 (inversion of top-level subtyping). *Let $\emptyset \vdash A_1 \leq A_2 : *$.*

- (1) *If $A_1 = B_1 \rightarrow C_1$ and $A_2 = B_2 \rightarrow C_2$, then $\emptyset \vdash B_2 \leq B_1 : *$ and $\emptyset \vdash C_1 \leq C_2 : *$.*
- (2) *If $A_1 = \forall X:K_1. B_1$ and $A_2 = \forall X:K_2. B_2$, then $\emptyset \vdash K_2 \leq K_1$ and $X:K_2 \vdash B_1 \leq B_2 : *$.*
- (3) *$\emptyset \not\vdash \top \leq \perp$, and $\emptyset \not\vdash A \rightarrow B \leq \forall X:K. C$, and $\emptyset \not\vdash \forall X:K. A \leq B \rightarrow C$.*

With subtyping inversion in place, we are finally ready to prove type safety of F^{ω} .

Theorem 5.5 (type safety). *Well-typed terms do not get stuck.*

- (progress) *If $\vdash t : A$, then either t is a value, or $t \rightarrow_{\nu} t'$ for some term t' .*
 (weak preservation) *If $\vdash t : A$ and $t \rightarrow_{\nu} t'$, then $\vdash t' : A$.*

The proofs are standard. Details are given in [Appendix D.4](#).

6 UNDECIDABILITY OF SUBTYPING

In this section, we prove that subtyping for F^{ω} is undecidable. The culprit is equality reflection via the bound projection rules **ST-BND₁** and **ST-BND₂** (cf. §2.2). Following [Castellan et al. \[2015\]](#), we prove undecidability of subtyping by reduction from convertibility of SK combinator terms. The *SK combinator calculus* has only three term formers $t ::= S \mid K \mid st$ and two equational axioms: $Sst =_{\text{SK}} su(tu)$ and $Kst =_{\text{SK}} s$. Yet SK is Turing-complete, and convertibility of SK terms is undecidable. We embed SK terms and equations into F^{ω} via the following declarations:

$$\begin{aligned} \Gamma_{\text{SK}} &:= S : *, \quad K : *, \quad \odot : * \rightarrow * \rightarrow *, \\ S_r &: (X:*) \rightarrow (Y:*) \rightarrow (Z:*) \rightarrow S \odot X \odot Y \odot Z .. X \odot Z \odot (Y \odot Z), \\ S_e &: (X:*) \rightarrow (Y:*) \rightarrow (Z:*) \rightarrow X \odot Z \odot (Y \odot Z) .. S \odot X \odot Y \odot Z, \\ K_r &: (X:*) \rightarrow (Y:*) \rightarrow K \odot X \odot Y .. X, \quad K_e : (X:*) \rightarrow (Y:*) \rightarrow X .. K \odot X \odot Y. \\ \llbracket S \rrbracket &:= S \quad \llbracket K \rrbracket := K \quad \llbracket st \rrbracket := \llbracket s \rrbracket \odot \llbracket t \rrbracket. \end{aligned}$$

The map $\llbracket - \rrbracket$ encodes SK terms as types under Γ_{SK} and induces a reduction from SK convertibility $s =_{\text{SK}} t$ to subtyping $\Gamma_{\text{SK}} \vdash \llbracket s \rrbracket \leq \llbracket t \rrbracket : *$, which can be used to prove undecidability of subtyping.

Theorem 6.1 (undecidability). *Let s, t be SK terms. Then $\Gamma_{\text{SK}} \vdash \llbracket s \rrbracket \leq \llbracket t \rrbracket : *$ iff $s =_{\text{SK}} t$.*

It is easy to verify the “if” direction. For example, the contraction law for K corresponds to the inequation $\Gamma_{\text{SK}} \vdash K \odot \llbracket s \rrbracket \odot \llbracket t \rrbracket \leq K_r \llbracket s \rrbracket \llbracket t \rrbracket \leq \llbracket s \rrbracket : *$. The “only if” direction is more challenging. The complexity of subtyping derivations precludes a direct decoding into $=_{\text{SK}}$ for many of the same reasons that a direct proof of subtyping inversion is unfeasible. In addition, types such as

\top or $\forall X:K. A$ that are not encodings of SK terms can appear as intermediate expression along a subtyping derivation. For example, note the fleeting appearance of \top in the following.

$$\Gamma_{\text{SK}} \vdash \llbracket S \rrbracket \equiv S \leq K \odot S \odot (K \odot \top \odot S) \leq K \odot S \odot \top \leq S \equiv \llbracket S \rrbracket : *.$$

To eliminate such spurious appearances of undecodable types, our proof takes a detour through four auxiliary judgment forms.

- (1) We first prove undecidability of canonical subtyping to eliminate instances of $\beta\eta$ -conversions. Undecidability of declarative subtyping follows by equivalence of the two judgments.
- (2) We define a *reduced* version of the canonical system from which we exclude any rules (and judgment forms) that are not relevant to the embedding shown above. For example, any judgments involving higher-order operators are excluded, as are the rules **CST-ARR** and **CST-ALL** and the variable subsumption rule **CV-SUB**. The reduced system still contains **CST-BOT** and **CST-TOP** since we cannot rule out intermediate occurrences of these rules a-priori. We show that canonical subtyping derivations under Γ_{SK} can be translated into reduced ones.
- (3) We extend the SK term syntax with \perp and \top and define an order \leq_{SK} on extended terms. The term order is an asymmetric version of \leq_{SK} which features the rules $\perp \leq_{\text{SK}} t$ and $t \leq_{\text{SK}} \top$. Thanks to these, reduced canonical subtyping derivations can be directly decoded into \leq_{SK} .
- (4) We introduce a pair of *parallel reduction relations* \Rightarrow_{\leq} and \Rightarrow_{\geq} on the extended syntax. These contain the rules $\perp \Rightarrow_{\leq} t$ and $\top \Rightarrow_{\geq} t$ for eliminating occurrences of \perp and \top , along with the usual contraction rules for SK terms. Crucially, the reduction rules can eliminate but never introduce instances of \perp and \top . Hence, if s is a pure SK term, $s \Rightarrow t$ implies $s =_{\text{SK}} t$. The parallel reductions enjoy a confluence property w.r.t. the term order: If $s \leq_{\text{SK}} t$, then there is a u such that $s \Rightarrow_{\leq}^* u \Leftarrow_{\leq}^* t$. Via confluence, $s \leq_{\text{SK}} t$ implies $s =_{\text{SK}} t$ for pure s and t .

Thus we have established a chain of implications from which the result follows.

$$\Gamma_{\text{SK}} \vdash \llbracket s \rrbracket \leq \llbracket t \rrbracket : * \implies \Gamma_{\text{SK}} \vdash_{\text{red}} \llbracket s \rrbracket \leq \llbracket t \rrbracket \implies s \leq_{\text{SK}} t \implies s \Rightarrow_{\leq}^* u \Leftarrow_{\leq}^* t \implies s =_{\text{SK}} t.$$

For full details, we refer the intrepid reader to the mechanized proof of [Theorem 6.1](#), which is given in the `F0megaInt.Undecidable` module of our Agda formalization [[Stucki and Giarrusso 2021](#)].

7 RELATED WORK

Bounded quantification has been studied extensively through F_{\leq} , a variant of System F with bounded quantification, which comes in two flavors: the *Kernel* variant $F_{<}$. [[Cardelli et al. 1991](#)] based on [Cardelli and Wegner’s Kernel Fun \[1985\]](#) has decidable subtyping, while *Full* F_{\leq} [[Curien and Ghelli 1992](#)] features a more expressive subtyping rule for bounded universal quantifiers that renders subtyping undecidable [[Pierce 1992](#)]. Recently, [Hu and Lhoták \[2019\]](#) have shown that the $D_{<}$ calculus – a simplified variant of DOT that uses an expressive \forall -subtyping rule – suffers from the same decidability issue as Full F_{\leq} . For compatibility with DOT, and knowing that subtyping in F^{ω} is undecidable either way, we also adopt the more expressive rule. The metatheory developed in [§§3–6](#) is largely unaffected by this choice.

An extension of [Girard’s \$F_{\omega}\$ \[1972\]](#) with higher-order subtyping and bounded quantification was first proposed by [Cardelli \[1990\]](#) under the name $F_{<}^{\omega}$. Basic meta theoretic properties of $F_{<}^{\omega}$ were established by [Pierce and Steffen \[1997\]](#), [Compagnoni \[1995\]](#), and [Compagnoni and Goguen \[1999\]](#). An extension with bounded operator abstractions ($\mathcal{F}_{\leq}^{\omega}$) has been developed by [Compagnoni and Goguen \[2003\]](#). More recently, [Abel and Rodriguez \[2008\]](#) developed a variant of $F_{<}^{\omega}$, where types are identified up to $\beta\eta$ -equality and proved its decidability using hereditary substitution. Their work inspired the syntactic approach taken in this paper.

Many of the ideas in $F_{<}^{\omega}$ go back to early work by [Cardelli \[1988\]](#) on *power types*. Though very expressive, power types render the type language non-normalizing, and in a later work [Cardelli](#)

and Longo [1991] replaced them with the better behaved *power kinds*. Power kinds can be directly expressed in F^ω as interval kinds that are bounded by \perp from below: $P(A) = \perp .. A$. Crary [1997] developed an extension of F_ω with power kinds as a general calculus for higher-order subtyping. His representations of higher-order bounded quantifiers and operators closely resemble ours.

The notion of *translucency* was introduced by Harper and Lillibridge [1994] in the setting of ML-style modules with sharing constraints. They proposed *translucent sums* as a uniform way of representing translucent type definitions. Stone and Harper [2000] later proposed *singleton kinds* as an alternative mechanism for representing type definitions with sharing constraints. Interval kinds are closely related, conceptually and formally, to Stone and Harper’s singleton kinds.

The safety of (in)consistent subtyping constraints in F_\leq -like systems has been studied in depth by Cretin and Rémy [2014] and Scherer and Rémy [2015]. They formalize two distinct types of subtyping coercions: *coherent* coercions can be erased (i.e. used implicitly) while *incoherent* coercions are introduced and eliminated explicitly. Reductions is allowed (and safe) only under coherent abstractions. It is unclear if their results extend to \mathcal{F}_\leq^ω -like systems such as ours.

Hereditary substitution is due to Watkins et al. [2004] and has been used to prove weak normalization of a variety of systems. A particularly illuminating example is provided by Keller and Altenkirch [2010] who use it to implement a normalization function for STLC in Agda. Other examples are the work by Abel and Rodriguez [2008] on F_\leq^ω and the presentation of Canonical LF by Harper and Licata [2007], both of which inspired the metatheoretic development in this paper. Hereditary substitution has also been used to mechanize the equational theory of singleton kinds [Crary 2009] and the semantics of the SML language [Lee et al. 2007] in Twelf.

F^ω belongs to a long line of calculi developed to model Scala’s type system. One of the first to support complex type operators and a form of interval kinds was *Scalina* [Moors et al. 2008b]. Type and kind safety of Scalina was never established but it inspired an extension of Scala’s type system with HK types, including higher-order bounded polymorphism, type operators and type definitions [Moors et al. 2008a]. More recently, Amin et al. [2016] introduced the calculus of *Dependent Object Types (DOT)* as a theoretical foundation for Scala and a core calculus for the *Scala 3* compiler [Dotty Team 2020]. Many variants of DOT have been developed, differing in expressiveness and presentation; most come with mechanized type safety proofs [see e.g. Amin 2016; Giarrusso et al. 2020; Rapoport and Lhoták 2019; Rompf and Amin 2016]. Central to all is the notion of *abstract type members*. Because type members can have lower and upper bounds, they provide a form of type intervals. In F^ω , we separate the concept of type intervals from that of abstract type members via interval kinds. DOT admits encodings of some type operators, but none of the DOT calculi developed so far can express general HK types as supported by Scala. But the development of the *Scala 3* compiler has shown the need for a principled theory of HK types [Odersky et al. 2016]. In this paper, we have proposed such a theory.

In the future, we wish to extend our work on HK types with existing work on DOT by recombining abstract type members with interval kinds. A type member definition would then be of the form $\{X : K\}$ where K may be a higher-order interval kind. We expect this to cause new feature interactions, some of which may be problematic. A sketch of such an extension, including a brief discussion of potential issues, can be found in the first author’s dissertation [Stucki 2017, Ch. 6].

Another direction for future work is to adapt the techniques developed by Hu and Lhoták [2019] for algorithmic subtyping in the DOT-like calculus $D_{<}$ to our system. Hu and Lhoták address the problems caused by inconsistent bounds in $D_{<}$ by replacing the general subtyping transitivity rule with a specialized rule that combines transitivity and bound projection. This isolates the problematic use of inequality reflection in a single rule. They then show that one can obtain a decidable system by removing this rule and weakening the rule for subtyping universals. Not only is this an elegant solution, it also closely reflects the strategy implemented in the *Scala* compiler.

We do believe that a variant of [Hu and Lhoták](#)'s strategy could be applied to our system. However, we expect transitivity elimination to be considerably more challenging in our system than in $D_{<}$, as one would expect in a dependently kinded setting.

8 CONCLUSIONS

We have described F^ω , a formal theory of higher-order subtyping with type intervals. In F^ω , type intervals are represented through interval kinds. We showed how interval kinds can be used to encode bounded universal quantification, bounded type operators and singleton kinds. We illustrated the use of interval kinds to abstract over and reflect type inequations and discussed the problems that arise when the corresponding intervals have inconsistent bounds.

We established basic metatheoretic properties of F^ω . We proved subject reduction in its full generality on the type level, and in a restricted form on the term level. We showed that types and kinds are weakly normalizing by defining a bottom-up normalization procedure on raw kinds and types and proving its soundness. We gave an alternative, canonical presentation of the kind and type level of F^ω , defined directly on $\beta\eta$ -normal forms. We showed that hereditary substitutions preserve canonical judgments and used this result to establish equivalence of the declarative and canonical presentations. We showed that canonical and, by equivalence, declarative subtyping can be inverted in the empty context. Based on these results, we established type safety of F^ω . We concluded our metatheoretic development by showing that subtyping is undecidable in F^ω . The metatheory has been fully mechanized in Agda.

Our goal in developing F^ω was twofold: study the theory of type intervals for higher-order subtyping, and develop a foundation for Scala's higher-kinded types. We believe that F^ω fulfills this goal and constitutes an important step toward a full formalization of Scala's expressive type system. During the development of F^ω , we discovered a number of minor flaws in Scala 3 (listed in [Appendix F](#)). None of these issues constitute critical bugs – in particular, they do not break type safety. But they do illustrate that subtyping in Scala 3 is slightly weaker than necessary, suggesting that there is room for improvement. Indeed, we hope that F^ω will serve as a blueprint for a more principled implementation of higher-order subtyping in future versions of Scala.

ACKNOWLEDGMENTS

We owe special thanks to Guillaume Martres for many discussions about this work, for his patience in answering our questions about the Scala 3 type checker, and for his striking ability to produce counterexamples to type system drafts. For insightful discussions and feedback on earlier versions of this work we thank Andreas Abel, Nada Amin, Jesper Cockx, Martin Odersky and François Pottier. We thank the anonymous reviewers for their helpful comments and suggestions. This paper is based upon work supported by the European Research Council (ERC) under Grant 587327 DOPPLER and by the Swedish Research Council (VR) under Grants 2015-04154 PolUser and 2018-04230 Perspex.

A OVERVIEW OF THE AGDA MECHANIZATION

The following is a list of the modules included in the Agda mechanization, with short descriptions of their purpose. The list is organized in blocks which correspond roughly to the sections of the paper where the metatheoretic definitions and properties corresponding to the contents of the module are (first) described. For more information, see the *README.md* and *Correspondence.agda* files included in our artifact [[Stucki and Giarrusso 2021](#)].

A.1 The Declarative System

Syntax of raw (i.e. untyped) terms along with support for untyped substitutions.

- `F0megaInt.Syntax`

Variants of β -reduction/equivalence and properties thereof.

- `F0megaInt.Reduction.Cbv`
- `F0megaInt.Reduction.Full`

Declarative typing, kinding, subtyping, etc. along with corresponding substitution lemmas.

- `F0megaInt.Typing`

An alternative presentation of kinding and subtyping that is better suited for proving functionality and validity lemmas, and a proof that the two presentations are equivalent.

- `F0megaInt.Kinding.Declarative`
- `F0megaInt.Kinding.Declarative.Validity`
- `F0megaInt.Kinding.Declarative.Equivalence`

Encodings and properties of higher-order extremal types, interval kinds and bounded quantifiers.

- `F0megaInt.Typing.Encodings`

A.2 Normalization of Types

Hereditary substitutions and normalization of raw types and kinds.

- `F0megaInt.Syntax.SingleVariableSubstitution`
- `F0megaInt.Syntax.HereditarySubstitution`
- `F0megaInt.Syntax.Normalization`

Weak equality of raw terms (up to kind annotations).

- `F0megaInt.Syntax.WeakEquality`

Soundness of normalization w.r.t. to declarative kinding.

- `F0megaInt.Kinding.Declarative.Normalization`

Simple kinding of types, and hereditary substitution lemmas; lemmas about η -expansion of simply kinded types and kinds.

- `F0megaInt.Kinding.Simple`
- `F0megaInt.Kinding.Simple.EtaExpansion`

Normalization and simultaneous simplification of declaratively kinded types.

- `F0megaInt.Kinding.Simple.Normalization`

A.3 The Canonical System

Canonical kinding of types along with (hereditary) substitution, validity and inversion lemmas for canonical kinding and subtyping.

- `F0megaInt.Kinding.Canonical`
- `F0megaInt.Kinding.Canonical.HereditarySubstitution`
- `F0megaInt.Kinding.Canonical.Validity`
- `F0megaInt.Kinding.Canonical.Inversion`

Lifting of weak (untyped) kind and type equality to canonical kind and type equality.

- `F0megaInt.Kinding.Canonical.WeakEquality`

Equivalence of canonical and declarative kinding.

- `F0megaInt.Kinding.Canonical.Equivalence`

Generation of typing and inversion of declarative subtyping in the empty context.

- `F0megaInt.Typing.Inversion`

Type safety (preservation and progress).

- `F0megaInt.Typing.Preservation`
- `F0megaInt.Typing.Progress`

A.4 Undecidability of Subtyping

A reduced variant of the canonical system.

- `F0megaInt.Kinding.Canonical.Reduced`

Setup for the undecidability proof: syntax and lemmas for the SK combinator calculus, and support for encoding/decoding SK terms and equality proofs into types and subtyping derivations.

- `F0megaInt.Undecidable.SK`
- `F0megaInt.Undecidable.Encoding`
- `F0megaInt.Undecidable.Decoding`

Undecidability of subtyping

- `F0megaInt.Undecidable`

A.5 Auxiliary Modules Providing Generic Functionality

Generic support for typing contexts over abstract bindings.

- `Data.Context`
- `Data.Context.WellFormed`
- `Data.Context.Properties`

Extra lemmas that are derivable in the substitution framework of the Agda standard library, as well as support for binary (term) relations lifted to substitutions, typed substitutions, and typed relations lifted to substitutions.

- `Data.Fin.Substitution.Extra`
- `Data.Fin.Substitution.ExtraLemmas`
- `Data.Fin.Substitution.Relation`
- `Data.Fin.Substitution.Typed`
- `Data.Fin.Substitution.TypedRelation`

Support for generic reduction relations, and relational reasoning for transitive relations.

- `Relation.Binary.Reduction`
- `Relation.Binary.TransReasoning`

B BASIC METATHEORY AND ADMISSIBLE RULES OF THE DECLARATIVE SYSTEM

We establish basic meta-theoretic properties of the declarative system and introduce a number of admissible rules, many of which are used in proofs later on. In particular, we establish a series of standard lemmas stating that the declarative judgments are preserved under common operations on contexts. Next, we show that the usual order-theoretic rules for subkinding and type and kind equality are admissible, as are congruence rules (w.r.t. to all the type formers) for type equality. We further introduce admissible typing, kinding, subtyping and subkinding rules for the encoded higher-order extremal types, interval kinds and bounded quantifiers described in §3 of the paper, and discuss alternative encodings. Finally, we state and prove a number of standard validity properties for the various judgments.

The contents of this section are based on Chapter 3 of the first author's PhD dissertation. We refer the interested reader to the dissertation for the full details [Stucki 2017, Ch. 3].

B.1 Basic Metatheoretic Properties

We start our metatheoretic development by showing that judgments can only be derived in well-formed contexts.

Lemma B.1 (context validity). *If $\Gamma \vdash \mathcal{J}$ for any of the judgments defined above, then Γ ctx.*

PROOF. By (simultaneous) induction on the derivations of the various judgments. The cases for context formation judgments and rules that contain Γ ctx as a premise are trivial. For other rules, the result follows by applying the IH to any of the premises that do not extend the context. There is always at least one such premise. \square

Next, we establish a series of standard lemmas stating that the declarative judgments are preserved under common operations on contexts, namely, addition and narrowing of bindings, and substitutions.

Lemma B.2 (weakening). *If $\Gamma, \Delta \vdash \mathcal{J}$, then*

- (1) *for any type A and $x \notin \text{dom}(\Gamma, \Delta)$, if $\Gamma \vdash A : *$, then $\Gamma, x:A, \Delta \vdash \mathcal{J}$;*
- (2) *for any kind K and $X \notin \text{dom}(\Gamma, \Delta)$, if $\Gamma \vdash K$ kd, then $\Gamma, X:K, \Delta \vdash \mathcal{J}$.*

Corollary B.3 (iterated weakening). *If Γ, Δ ctx and $\Gamma \vdash \mathcal{J}$, then $\Gamma, \Delta \vdash \mathcal{J}$.*

Lemma B.4 (substitution).

- (1) *If $\Gamma \vdash t : A$ and $\Gamma, x:A, \Delta \vdash \mathcal{J}$, then $\Gamma, \Delta \vdash \mathcal{J}[t/x]$.*
- (2) *If $\Gamma \vdash A : K$ and $\Gamma, X:K, \Delta \vdash \mathcal{J}$, then $\Gamma, \Delta[A/X] \vdash \mathcal{J}[A/X]$.*

Lemma B.5 (context narrowing).

- (1) *If $\Gamma \vdash A : *$, $\Gamma \vdash A \leq B : *$ and $\Gamma, x:B, \Delta \vdash \mathcal{J}$, then $\Gamma, x:A, \Delta \vdash \mathcal{J}$.*
- (2) *If $\Gamma \vdash J$ kd, $\Gamma \vdash J \leq K$ and $\Gamma, X:K, \Delta \vdash \mathcal{J}$, then $\Gamma, X:J, \Delta \vdash \mathcal{J}$.*

The context narrowing lemma is a bit weaker than one might expect; the premises $\Gamma \vdash A : *$ and $\Gamma \vdash J$ kd seem redundant. Surely, if $\Gamma \vdash A \leq B : *$ then A and B ought to be proper types. This property – called *subtyping validity* – does indeed hold, but we are not yet ready to prove it. Indeed, one of the prerequisites is the context narrowing lemma itself.

Lemmas B.2, B.4 and **B.5** are proven in that order, each by simultaneous induction on the derivations of the various judgments. The proofs of **Lemmas B.4** and **B.5** rely on **Corollary B.3** for the variable cases **T-VAR** and **K-VAR**. All three proofs are entirely standard, so we only present an excerpt from the proof of **Lemma B.4** to illustrate the basic strategy. In it, we make use of the following helper lemma about substitutions.

Lemma B.6 (substitutions commute). *Let e be some arbitrary expression, A, B types and X, Y distinct type variables such that $X \notin \text{fv}(B)$. Then $e[A/X][B/Y] \equiv e[B/Y][A[B/Y]/X]$*

PROOF. By induction on the structure of e . \square

PROOF OF **LEMMA B.4**. The two parts are proven separately, each by induction on the derivation of the second premise ($\Gamma, x:A, \Delta \vdash \mathcal{J}$ for the first part, $\Gamma, X:K, \Delta \vdash \mathcal{J}$ for the second). For the context formation judgment, the proofs proceed by a local induction on the structure of Δ . We show the cases for **K-VAR** and **ST- β_1** for the second part of the lemma. The other cases are similar.

- *Case K-VAR.* \mathcal{J} is $X : J$ and we have $\Gamma, X:K, \Delta$ ctx and $(\Gamma, X:K, \Delta)(Y) = J$. By the IH, we get $\Gamma, \Delta[A/X]$ ctx. We distinguish two sub-cases based on whether $Y = X$.
 - *Sub-case $Y = X$.* We want to show that $\Gamma, \Delta[A/X] \vdash A : K[A/X]$. By well-scopedness, X does not occur freely in K , so we have $K[A/X] \equiv K$. The desired result follows from applying iterated weakening (**Corollary B.3**) to the premise $\Gamma \vdash A : K$.

Another consequence of these rules is that we can treat well-typed terms and well-kinded types up to type and kind equality, respectively.

Corollary B.10 (conversion). *The following are admissible.*

$$\frac{\Gamma \vdash A : J \quad \Gamma \vdash J = K}{\Gamma \vdash A : K} \quad (\text{K-CONV}) \qquad \frac{\Gamma \vdash t : A \quad \Gamma \vdash A = B : *}{\Gamma \vdash t : B} \quad (\text{T-CONV})$$

$$\frac{\Gamma \vdash A \leq B : J \quad \Gamma \vdash J = K}{\Gamma \vdash A \leq B : K} \quad (\text{ST-CONV}) \qquad \frac{\Gamma \vdash A = B : J \quad \Gamma \vdash J = K}{\Gamma \vdash A = B : K} \quad (\text{TEQ-CONV})$$

In light of their order-theoretic properties, we often call kind and type equality judgments *equations* and subkinding and subtyping judgments *inequations*. We sometimes use equational reasoning notation in proofs, i.e. we write

$$\Gamma \vdash A_1 = A_2 = \dots = A_i \leq A_{i+1} \leq \dots = A_n \equiv A_{n+1} \equiv \dots = A_m : K$$

to denote chains of (in)equations where the use of the corresponding transitivity rules is left implicit. In so doing, we may freely mix the relations \leq , $=$ and \equiv provided that they are defined on the same sort (i.e. kinds or types). Such chains are always interpreted as judgments of the weakest relation they contain.

B.3 Validity of the Declarative System

In this section, we state and prove a number of *validity properties* for the various judgments defined in Figs. 3 and 4 of the paper (page 10). Roughly, we say that a judgment is valid if all its parts are well-formed. For example, *subkinding validity* states that, if $\Gamma \vdash J \leq K$, then both J and K are actually well-formed kinds. We saw another example earlier: *context validity* (Lemma B.1) states that the context Γ of any judgment $\Gamma \vdash \mathcal{J}$ is well-formed. Here is a summary of the validity properties that remain to be proven.

Lemma B.11 (validity). *The judgments defined in Figs. 3 and 4 enjoy the following validity properties.*

- (kinding validity) *If $\Gamma \vdash A : K$, then $\Gamma \vdash K$ kd.*
- (typing validity) *If $\Gamma \vdash t : A$, then $\Gamma \vdash A : *$.*
- (subkinding validity) *If $\Gamma \vdash J \leq K$, then $\Gamma \vdash J$ kd and $\Gamma \vdash K$ kd.*
- (subtyping validity) *If $\Gamma \vdash A \leq B : K$, then $\Gamma \vdash A : K$ and $\Gamma \vdash B : K$.*
- (kind equation validity) *If $\Gamma \vdash J = K$, then $\Gamma \vdash J$ kd and $\Gamma \vdash K$ kd.*
- (type equation validity) *If $\Gamma \vdash A = B : K$, then $\Gamma \vdash A : K$ and $\Gamma \vdash B : K$.*

These validity properties provide a “sanity check” for the static semantics developed in this section, but they also play a crucial role in the proofs of other important properties, such as subject reduction, soundness of type normalization and and type safety.

Unfortunately, the validity properties are harder to prove than one might expect. The proofs of kinding, subkinding and subtyping validity require the following *functionality* lemma for the case of **ST-APP**.

Lemma B.12 (functionality). *Let $\Gamma \vdash A_1 = A_2 : K$.*

- (1) *If $\Gamma, X : K, \Delta \vdash J$ kd, then $\Gamma, \Delta[A_1/X] \vdash J[A_1/X] = J[A_2/X]$.*
- (2) *If $\Gamma, X : K, \Delta \vdash B : J$, then $\Gamma, \Delta[A_1/X] \vdash B[A_1/X] = B[A_2/X] : J[A_1/X]$.*

But a naive attempt at proving this lemma directly leads to a circular dependency on kinding and subtyping validity, in a way that is not easily resolved. In particular, it is not sufficient to simply prove the two statements simultaneously.

It is instructive to play through the critical cases encountered when attempting to prove [Lemmas B.11](#) and [B.12](#) directly to see where things go wrong and to better understand the solution described in the next section. We start with subtyping validity, attempting a proof by induction on subtyping derivations. For the case of the application rule [ST-APP](#), we are given $\Gamma \vdash A_1 \leq A_2 : (X:J) \rightarrow K$ and $\Gamma \vdash B_1 = B_2 : J$, and we would like to show that $\Gamma \vdash A_1 B_1 : K[B_1/X]$ and $\Gamma \vdash A_2 B_2 : K[B_1/X]$. We can already spot the source of trouble: the type B_1 that is being substituted for X in the kind of the second type application differs from the argument type B_2 . By the IH, we get $\Gamma \vdash A_2 : (X:J) \rightarrow K$ and $\Gamma \vdash B_2 : J$, and applying [K-APP](#) we obtain $\Gamma \vdash A_2 B_2 : K[B_2/X]$ but, as expected, the kinds do not match up. If we could show that $\Gamma \vdash K[B_2/X] = K[B_1/X]$, then by [K-CONV](#), we would be done. Enter functionality of kind formation.

For the functionality lemma, we attempt a proof by simultaneous induction on kind formation and kinding derivations and consider the case where the current kinding derivation ends in an instance of the operator abstraction rule [K-ABS](#). In addition to the premise $\Gamma \vdash B_1 = B_2 : J$, we are given derivations for $\Gamma, X:J \vdash K_1$ kd and $\Gamma, X:J, Y:K_1 \vdash A : K_2$, and we want to show that

$$\Gamma \vdash (\lambda Y:K_1. A)[B_1/X] = (\lambda Y:K_1. A)[B_2/X] : ((Y:K_1) \rightarrow K_2)[B_1/X].$$

To do so, we would like to use the rule [ST-ABS](#) together with [ST-ANTI-SYM](#) but we first need to establish the right-hand validity of the above equation, i.e. that

$$\Gamma \vdash (\lambda Y:K_1. A)[B_2/X] : ((Y:K_1) \rightarrow K_2)[B_1/X].$$

Clearly, [Lemma B.11](#) would be helpful here: equation validity would give us $\Gamma \vdash B_2 : J$, from which we could obtain $\Gamma \vdash (\lambda Y:K_1. A)[B_2/X] : ((Y:K_1) \rightarrow K_2)[B_2/X]$ by [Lemma B.4](#). This is almost what we need. Again, we face a mismatch in kinds that could, in principle, be remedied by using functionality of kind formation together with [K-CONV](#). Concretely, we would like to invoke the IH to derive $\Gamma \vdash ((Y:K_1) \rightarrow K_2)[B_1/X] = ((Y:K_1) \rightarrow K_2)[B_2/X]$. Alas, we do not have a suitable sub-derivation to do so. Although $\Gamma \vdash (Y:K_1) \rightarrow K_2$ kd follows from kinding validity, we cannot apply the IH to this result because it is not a sub-derivation of our overall premise. Indeed, none of the sub-derivations we are given are sufficient to derive the required kind equation.

Note that we could finish the proof of this case if only (1) the rule [K-ABS](#) had an additional premise $\Gamma, X:J, Y:K_1 \vdash K_2$ kd and (2) the IH was a bit stronger, so that we could use it to derive

$$\Gamma, Y:K_1[B_1/X] \vdash K_2[B_1/X] = K_2[B_2/X].$$

This, together with a similar use of the IH on the first premise of [ST-ABS](#) and some uses of [Lemma B.4](#), [ST-REFL-TEQ](#) and [SK-DARR](#), would be enough to derive $\Gamma \vdash ((Y:K_1) \rightarrow K_2)[B_2/X] \leq ((Y:K_1) \rightarrow K_2)[B_1/X]$, which we could then put to use with [K-SUB](#). Indeed, these are the basic ideas that will allow us to resolve the circular dependency between [Lemma B.11](#) and [Lemma B.12](#). We start by addressing point (1).

B.3.1 The Extended System. We define a pair of *extended* kinding and subtyping judgments where some rules have been endowed with additional premises. These are precisely the premises highlighted in gray in [Figs. 3](#) and [4](#). We call these extra premises *validity conditions*. Crucially, the validity conditions of an extended rule are redundant in the sense that they follow (more or less) directly from the remaining premises of the rule via [Lemma B.11](#). For example, the extended rule [K-ABS](#) carries the extra premise $\Gamma, X:J \vdash K$ kd which follows directly from applying kinding validity to the rule's second premise $\Gamma, X:J \vdash A : K$. Thanks to this invariant, the two sets of rules are in fact equivalent – every derivation of an extended kinding or subtyping judgment has a corresponding derivation that uses only original rules, and vice-versa. We give a formal equivalence proof in [§B.3.2](#).

Since kinding and subkinding are defined mutually with all the other judgments of the declarative system (except typing), the extension indirectly affects those judgments as well. We call the entire set of extended judgments the *extended (declarative) system*, as opposed to the *original (declarative) system*. We will sometimes distinguish the two systems by writing $\Gamma \vdash_d \mathcal{J}$ for judgments of the original system and $\Gamma \vdash_e \mathcal{J}$ for those of the extended system. Since the two systems are equivalent, this distinction only matters in a few key situations – notably the development in the remainder of this section. When we refer to the “declarative system” in the following sections, we will always mean the original declarative system, unless otherwise noted.

The idea of extending a set of inference rules with redundant premises in order to simplify metatheoretic proofs is not new. For example, Harper and Pfenning [2005] use similar premises to establish validity properties for the typing judgments of a variant of LF. Furthermore, some readers will have noticed that a few of the original declarative rules already carry redundant premises. For example, the first premise $\Gamma \vdash J \text{ kd}$ of the rule **K-ABS** could easily be reconstructed from its second premise $\Gamma, X:J \vdash A : K$ via context validity (Lemma B.1). The rules **WF-DARR**, **K-ALL**, **T-ABS**, and **T-TABS** carry similar validity conditions. We include these premises primarily because their presence simplifies the proof of the substitution lemma (Lemma B.4). Context validity, on the other hand, remains easily provable without them.

To prove the validity properties stated in Lemma B.11 for both the original and extended systems, we use the following strategy:

- (1) prove that the validity properties hold for the extended judgments;
- (2) prove that the two systems are equivalent, i.e. that
 - (a) the extended rules are *sound* w.r.t. to the original ones – we can drop the validity conditions without affecting the conclusions of any derivations – and that
 - (b) the extended rules are *complete* w.r.t. to the original ones – the additional validity conditions follow from the remaining premises of the extended rules via the validity properties proved in step 1;
- (3) prove that the validity properties hold for the original system via the equivalence – convert original derivations to extended derivations (via completeness), derive the property in question, convert the conclusion back (via soundness).

Before we continue, we should point out that some of the validity conditions of the extended system are not actually necessary for the proof of Lemmas B.11 and B.12 – some even complicate the proofs. However, we will face a similar cyclic dependency later on when attempting to prove the equivalence of the (original) declarative system and the *canonical system* of judgments introduced in §5. Rather than introducing yet another extension to the declarative system later, we opt for a combined system containing all of the extra conditions.

We start the development set out above by noting that all the basic metatheoretic properties established in §B.1 still hold for the extended system. The proofs carry over with minor adjustments to deal with the additional premises. Next, we prove a variant of the functionality lemma discussed above but using the extended kinding and subtyping rules. To do so, we need the following auxiliary definition of *context equality* and an associated corollary of context weakening.

$$\frac{}{\emptyset = \emptyset \text{ ctx}} \quad \frac{\Gamma = \Delta \text{ ctx} \quad \Gamma \vdash J = K}{\Gamma, X:J = \Delta, X:K \text{ ctx}} \quad \frac{\Gamma = \Delta \text{ ctx} \quad \Gamma \vdash A = B : *}{\Gamma, x:A = \Delta, x:B \text{ ctx}}$$

Corollary B.13 (context conversion). *If $\Gamma \text{ ctx}$ and $\Gamma = \Delta \text{ ctx}$ and $\Delta \vdash \mathcal{J}$, then $\Gamma \vdash \mathcal{J}$.*

Context equality $\Gamma = \Delta \text{ ctx}$ is simply the pointwise lifting of type and kind equality to contexts. It allows us to relate the bindings appearing in two syntactically different contexts Γ and Δ , as illustrated in our extended functionality lemma.

Lemma B.14 (functionality – extended version). *Substitutions of equal types in well-formed expressions result in well-formed equations. Let Γ, Δ, Σ be contexts, K a kind and A_1, A_2 types, such that $\Gamma \vdash A_1 : K$ and $\Gamma \vdash A_2 : K$, the context Γ, Σ is well-formed and the following equations hold:*

$$\Gamma \vdash A_1 = A_2 : K \quad \Gamma, \Sigma = \Gamma, \Delta[A_1/X] \text{ ctx} \quad \Gamma, \Sigma = \Gamma, \Delta[A_2/X] \text{ ctx}.$$

(1) *If $\Gamma, X:K, \Delta \vdash J \text{ kd}$, then $\Gamma, \Sigma \vdash J[A_1/X] = J[A_2/X]$.*

(2) *If $\Gamma, X:K, \Delta \vdash B : J$, then $\Gamma, \Sigma \vdash B[A_1/X] = B[A_2/X] : J[A_1/X]$.*

Compared to [Lemma B.12](#), the lemma has been strengthened – so that it is applicable to any type variable binding in a context, not just the last one – and simultaneously weakened – by adding extra conditions on A_1, A_2 and the target context Γ, Σ . The latter are effectively validity conditions ensuring that the proof of the lemma does not depend on [Lemma B.11](#). The separate target context Σ is used to symmetrize the treatment of context extensions, which is helpful when dealing with kind annotations in contravariant positions.

PROOF. The two parts are proven simultaneously, by induction on extended kind formation and kinding derivations, respectively. The proof of the first part is relatively straightforward, while the proof of the second part deserves some attention. We present a few key cases, the others are similar.

- **Case *K-VAR*.** We have $B = Y, J = \Gamma(Y)$ and $\Gamma, X:K, \Delta \text{ ctx}$. We distinguish two cases based on Y .
 - **Sub-case $Y = X$.** We have $J = K \equiv K[A_1/X]$ since $X \notin \text{fv}(K)$. By iterated weakening, we get $\Gamma, \Sigma \vdash A_1 = A_2 : K[A_1/X]$ and we are done.
 - **Sub-case $Y \neq X$.** We have $(\Gamma, \Delta[A_1/X])(Y) \equiv J[A_1/X]$, either because $Y \in \text{dom}(\Gamma)$ and $X \notin \text{fv}(J)$, or because $Y \in \text{dom}(\Delta)$ and $(\Delta[A_1/X])(Y) = J[A_1/X]$. Furthermore, since $\Gamma, \Sigma = \Gamma, \Delta[A_1/X] \text{ ctx}$ we have

$$\Gamma, \Sigma \vdash (\Gamma, \Sigma)(Y) = (\Gamma, \Delta[A_1/X])(Y) \equiv J[A_1/X].$$

We conclude by [K-VAR](#), [K-CONV](#), and [TEQ-REFL](#).

- **Case *K-ALL*.** We have $B = \forall Y:J_1. B_1$ and $J = *$ for some kind J_1 and type B_1 , as well as $\Gamma, X:K, \Delta \vdash J_1 \text{ kd}$ and $\Gamma, X:K, \Delta, Y:J_1 \vdash B_1 : *$. We want to show that $(\forall Y:J_1. B_1)[A_1/X]$ and $(\forall Y:J_1. B_1)[A_2/X]$ are mutual subtypes. To do so, we first prove that

$$\begin{aligned} \Gamma, \Sigma \vdash (\forall Y:J_1. B_1)[A_1/X] & : * & \Gamma, \Sigma \vdash (\forall Y:J_1. B_1)[A_2/X] & : * \\ \Gamma, \Sigma \vdash J_1[A_2/X] \leq J_1[A_1/X] & & \Gamma, \Sigma \vdash J_1[A_1/X] \leq J_1[A_2/X] & \\ \Gamma, \Sigma, X:J_1[A_2/X] \vdash B_1[A_1/X] \leq B_1[A_2/X] & : * & & \\ \Gamma, \Sigma, X:J_1[A_1/X] \vdash B_1[A_2/X] \leq B_1[A_1/X] & : * & & \end{aligned}$$

then apply [ST-ALL](#) twice, and conclude with [ST-ANTI-SYM](#). The two kinding judgments follow from the premises by the substitution lemma ([Lemma B.4](#)), the two subkinding judgments by the IH. The last two subtyping judgments require some extra work.

Note that the additional type variable bindings in the two judgments differ syntactically, so we will have to use the IH twice, with different target contexts. In each case we need to show that the kind of the additional binding is well-formed and equal to $J_1[A_1/X]$ and $J_1[A_2/X]$, respectively. Concretely, we need to show that

$$\begin{aligned} \Gamma, \Sigma \vdash J_1[A_2/X] = J_1[A_1/X] \text{ kd} & & \Gamma, \Sigma \vdash J_1[A_2/X] \text{ kd} & \\ \Gamma, \Sigma \vdash J_1[A_2/X] = J_1[A_2/X] \text{ kd} & & & \end{aligned}$$

for the first invocation of the IH, and three analogous statements for the second. The first equation follows from the two subkinding judgments above via [ST-ANTI-SYM](#). The context

formation judgment and the first equation follow from the substitution lemma and **TEQ-REFL**. This is sufficient to apply the IH and obtain the first of the two remaining subtyping judgments via **ST-REFL-TEQ**. The proof of the second one is similar.

- *Case K-APP*. We have $B = B_1 B_2$ and $J = J_2[B_2/Y]$ for some B_1, B_2, J_2 , as well as

$$\begin{array}{ll}
 \Gamma \vdash A_1 : K, & \Gamma \vdash A_2 : K \\
 \Gamma \vdash A_1 = A_2 : K, & \Gamma, \Sigma \text{ ctx} \\
 \Gamma, \Sigma = \Gamma, \Delta[A_1/X] \text{ ctx}, & \Gamma, \Sigma = \Gamma, \Delta[A_2/X] \text{ ctx}, \\
 \Gamma, X:K, \Delta \vdash B_1 : (Y:J_1) \rightarrow J_2, & \Gamma, X:K, \Delta \vdash B_2 : J_1, \\
 \Gamma, X:K, \Delta, Y:J_1 \vdash J_2 \text{ kd}, & \Gamma, X:K, \Delta \vdash J_2[B_2/Y] \text{ kd}
 \end{array}$$

for some J_1 . We want to establish that that $(B_1 B_2)[A_1/X]$ and $(B_1 B_2)[A_2/X]$ are mutual subtypes in $J_2[B_2/Y][A_1/X]$, i.e. that

$$\Gamma, \Sigma \vdash (B_1 B_2)[A_1/X] \leq (B_1 B_2)[A_2/X] : J_2[B_2/Y][A_1/X], \quad \text{and} \quad (1)$$

$$\Gamma, \Sigma \vdash (B_1 B_2)[A_2/X] \leq (B_1 B_2)[A_1/X] : J_2[B_2/Y][A_1/X]. \quad (2)$$

The first half is fairly straightforward. Applying the IH to the first two premises of **K-APP** yields corresponding equations, the first of which we turn into an inequation via **ST-REFL-TEQ**.

$$\begin{array}{l}
 \Gamma, \Sigma \vdash B_1[A_1/X] \leq B_1[A_2/X] : ((Y:J_1) \rightarrow J_2)[A_1/X], \\
 \Gamma, \Sigma \vdash B_2[A_1/X] = B_2[A_2/X] : J_1[A_1/X].
 \end{array}$$

In order to apply **ST-APP** we also need to derive the following validity conditions:

$$\begin{array}{ll}
 \Gamma, \Sigma \vdash B_2[A_1/X] : J_1[A_1/X], & \Gamma, \Sigma, Y:J_1[A_1/X] \vdash J_2[A_1/X] \text{ kd}, \\
 \Gamma, \Sigma \vdash J_2[A_1/X][B_2[A_1/X]/Y] \text{ kd}.
 \end{array}$$

All three follow from premises of **K-APP** and the substitution lemma (**Lemma B.4**), followed by a use of **Corollary B.13** to adjust the contexts. Adjusting the context of $\Gamma, \Delta[A_1/X], Y:J_1[A_1/X] \vdash J_2[A_1/X] \text{ kd}$, requires a bit more work because we need to prove that the kind of the extra binding $Y:J_1[A_1/X]$ is well-formed. To do so, we first invoke context validity (**Lemma B.1**) on the second validity condition of **K-APP**, which gives us $\Gamma, X:K, \Delta \vdash J_1 \text{ kd}$. From this, we derive the desired well-formedness proof via the substitution lemma. By **ST-APP** and **Lemma B.6**, we arrive at (1).

We have to work a bit harder to prove (2). Again, we want to apply **ST-APP**, and again, the first two premises follow from the IH – this time followed by a use of **TEQ-SYM** to adjust the direction – and **ST-REFL-TEQ** to turn the first equation into a subtyping statement. The validity conditions are

$$\begin{array}{ll}
 \Gamma, \Sigma \vdash B_2[A_2/X] : J_1[A_1/X], & \Gamma, \Sigma, Y:J_1[A_1/X] \vdash J_2[A_1/X] \text{ kd}, \\
 \Gamma, \Sigma \vdash J_2[A_1/X][B_2[A_2/X]/Y] \text{ kd}.
 \end{array}$$

We have already established the second condition; the third one follows from applying the substitution lemma to the first two. So it remains to prove the first.

We start by deriving $\Gamma, \Sigma \vdash B_2[A_2/X] : J_1[A_2/X]$ via the substitution lemma and context conversion. Next, we would like to use the IH to show that $\Gamma, \Sigma \vdash J_1[A_2/X] = J_1[A_1/X]$ in order to adjust the kind of the previous judgment via **K-CONV**. But to do so, we need to find a derivation of $\Gamma, X:K, \Delta \vdash J_1 \text{ kd}$ that is a strict sub-derivation of our current instance of **K-APP**.

Fortunately, this is always possible, thanks to the validity condition $\Gamma, X:K, \Delta, Y:J_1 \vdash J_2 \text{ kd}$. Since the contexts of kind formation judgments are always well-formed themselves (see **Lemma B.1**),

it suffices to traverse the derivation tree of this judgment upwards along kind formation and kinding rules until one arrives at a “leaf” – an instance of **K-VAR**, **K-TOP** or **K-BOT** – which holds a well-formedness derivation for the current context. That context formation derivation, in turn, contains a sub-derivation of the desired kind formation judgment. Readers who are skeptical of this somewhat informal argument are encouraged to state and prove a helper lemma that combines the IH with the “lookup procedure” just described. The lemma is proven simultaneously with the main lemma, by induction on kind formation and kinding derivations.

With all the validity conditions in place, we apply **ST-APP** to obtain

$$\Gamma, \Sigma \vdash (B_1 B_2)[A_2/X] \leq (B_1 B_2)[A_1/X] : J_2[A_1/X][B_2[A_2/X]/Y].$$

To complete the proof of 2, we use **ST-CONV** and

$$\begin{aligned} \Gamma, \Sigma \vdash J_2[A_1/X][B_2[A_2/X]/Y] & \\ &= J_2[A_2/X][B_2[A_2/X]/Y] && \text{(by Lemma B.4)} \\ &\equiv J_2[B_2/Y][A_2/X] && \text{(by Lemma B.6)} \\ &= J_2[B_2/Y][A_1/X] && \text{(by the IH)} \end{aligned}$$

using our earlier result $\Gamma, \Sigma \vdash J_1[A_2/X] = J_1[A_1/X]$ in the first step and the final validity condition of **K-APP** in the last. \square

We are now ready to prove **Lemma B.11** in the extended system, simultaneously with the following lemma.

Lemma B.15. *Subtypes inhabiting interval kinds are proper subtypes. If $\Gamma \vdash A \leq B : C .. D$, then also $\Gamma \vdash A \leq B : *$.*

PROOF OF LEMMA B.11 AND LEMMA B.15 – EXTENDED VERSION. All the validity properties are proven simultaneously with **Lemma B.15**, by induction on the derivations of the respective premises. The proof is now mostly routine, thanks to the validity conditions. The only interesting cases are those of **ST-APP**, where we use the functionality lemma to adjust the kind of the right-hand validity proof, and **ST-INTV**, where we use **Lemma B.15**. The proof of **Lemma B.15** uses subtyping validity in turn. \square

Corollary B.16. *Equal types in intervals are equal as proper types. If $\Gamma \vdash A = B : C .. D$, then also $\Gamma \vdash A = B : *$.*

B.3.2 Equivalence. The next and final step in our program for proving **Lemma B.11** is to establish the equivalence of the two declarative systems.

Lemma B.17. *The original and extended declarative systems are equivalent: $\Gamma \vdash_d \mathcal{J}$ iff $\Gamma \vdash_e \mathcal{J}$.*

Thanks to this equivalence, all the validity properties laid out in **Lemma B.11** also hold for the original judgments of the declarative system. Our original functionality lemma (**Lemma B.12**) and the following strengthened version of **Lemma B.5** follow as corollaries of validity and **Lemmas B.14** and **B.5**, respectively.

Corollary B.18 (context narrowing – strong version).

- (1) *If $\Gamma \vdash A \leq B : *$ and $\Gamma, x:B, \Delta \vdash \mathcal{J}$, then $\Gamma, x:A, \Delta \vdash \mathcal{J}$.*
- (2) *If $\Gamma \vdash J \leq K$ and $\Gamma, X:K, \Delta \vdash \mathcal{J}$, then $\Gamma, X:J, \Delta \vdash \mathcal{J}$,*

B.4 Admissible Congruence Rules for Type and Kind Equality

Thanks to the validity properties established in §B.3, we are able to prove a number of admissible *congruence rules* for kind and type equality. These follow the same structure as the corresponding subkinding and subtyping rules but are generally a bit simpler. First, we no longer need to pay attention to the variance (or polarity) of constructor arguments because equality is symmetric. Second, the left-hand validity conditions present in the rules **SK-DARR** and **ST-ALL** become redundant in the corresponding equality rules because the kind annotations in the left- and right-hand sides are convertible. Finally, thanks to symmetry, only one rule is needed for β -conversion, and likewise for η -conversion.

Lemma B.19. *Kind equality is a congruence with respect to the interval and dependent arrow kind formers, i.e. the following kind equality rules are admissible.*

$$\frac{\Gamma \vdash A_1 = A_2 : * \quad \Gamma \vdash B_1 = B_2 : *}{\Gamma \vdash A_1 .. B_1 = A_2 .. B_2 : *} \text{ (KEQ-INTV)} \quad \frac{\Gamma \vdash J_1 = J_2 \quad \Gamma, X:J_1 \vdash K_1 = K_2}{\Gamma \vdash (X:J_1) \rightarrow K_1 = (X:J_2) \rightarrow K_2} \text{ (KEQ-DARR)}$$

Lemma B.20. *Type equality is a congruence with respect to the various type formers and includes β and η -conversion, i.e. the following type equality rules are admissible.*

$$\frac{\Gamma \vdash K_1 = K_2 \quad \Gamma, X:K_1 \vdash A_1 = A_2 : *}{\Gamma \vdash \forall X:K_1. A_1 = \forall X:K_2. A_2 : *} \text{ (TEQ-ALL)} \quad \frac{\Gamma \vdash \lambda X:J_1. A_1 : (X:J) \rightarrow K \quad \Gamma \vdash \lambda X:J_2. A_2 : (X:J) \rightarrow K \quad \Gamma, X:J \vdash A_1 = A_2 : K}{\Gamma \vdash \lambda X:J_1. A_1 = \lambda X:J_2. A_2 : (X:J) \rightarrow K} \text{ (TEQ-ABS)}$$

$$\frac{\Gamma \vdash A_1 = A_2 : * \quad \Gamma \vdash B_1 = B_2 : *}{\Gamma \vdash A_1 \rightarrow B_1 = A_2 \rightarrow B_2 : *} \text{ (TEQ-ARR)} \quad \frac{\Gamma \vdash A_1 = A_2 : (X:J) \rightarrow K \quad \Gamma \vdash B_1 = B_2 : J}{\Gamma \vdash A_1 B_1 = A_2 B_2 : K[B_1/X]} \text{ (TEQ-APP)}$$

$$\frac{\Gamma, X:J \vdash A : K \quad \Gamma \vdash B : J}{\Gamma \vdash (\lambda X:J. A) B = A[B/X] : K[B/X]} \text{ (TEQ-}\beta\text{)} \quad \frac{\Gamma \vdash A : (X:J) \rightarrow K \quad X \notin \text{fv}(A)}{\Gamma \vdash \lambda X:J. AX = A : (X:J) \rightarrow K} \text{ (TEQ-}\eta\text{)}$$

$$\frac{\Gamma \vdash A_1 = A_2 : * \quad \Gamma \vdash B_1 = B_2 : *}{\Gamma \vdash A_1 \rightarrow B_1 = A_2 \rightarrow B_2 : *} \text{ (TEQ-ARR)} \quad \frac{\Gamma \vdash A_1 = A_2 : B .. C}{\Gamma \vdash A_1 = A_2 : A_1 .. A_1} \text{ (TEQ-SING)}$$

PROOF. The admissibility proofs of the above rules all follow the same basic pattern. We want to show that the left- and right-hand sides of the conclusions are mutual subkinds or subtypes, respectively. To do so, we employ the respective subkinding and subtyping rules, adjusting the kinds of additional bindings and subtyping judgments using context narrowing (**Corollary B.18**), subsumption **ST-SUB**, conversion **ST-CONV** and functionality (**Lemma 3.2**) where necessary. When additional validity properties are required, **Lemma 3.1** delivers the required well-formedness or well-kindedness proofs.

For example, the proof of **TEQ-SING** proceeds as follows. We note that the premise must have been derived using **ST-ANTISYM**, hence we have $\Gamma \vdash A_1 \leq A_2 : B .. C$ and $\Gamma \vdash A_2 \leq A_1 : B .. C$. By **ST-INTV**, **Lemma B.15** and subtyping validity

$$\begin{array}{lll} \Gamma \vdash A_1 \leq A_2 : A_1 .. A_2 & (1a) & \Gamma \vdash A_1 \leq A_2 : * & (1b) & \Gamma \vdash A_1 : * & (1c) \\ \Gamma \vdash A_2 \leq A_1 : A_2 .. A_1 & (2a) & \Gamma \vdash A_2 \leq A_1 : * & (2b) & \Gamma \vdash A_2 : * & (2c) \end{array}$$

From (1a), (1c) and (2b) we derive

$$\frac{\frac{\frac{\Gamma \vdash A_1 : *}{\Gamma \vdash A_1 \leq A_1 : *} \text{(ST-REFL)}}{\Gamma \vdash A_1 \leq A_2 : A_1 .. A_2} \quad \frac{\Gamma \vdash A_2 \leq A_1 : *}{\Gamma \vdash A_1 .. A_2 \leq A_1 .. A_1} \text{(SK-INTV)}}{\Gamma \vdash A_1 \leq A_2 : A_1 .. A_1} \text{(ST-SUB)}$$

and similarly $\Gamma \vdash A_2 \leq A_1 : A_1 .. A_1$ from (2a), (2b) and (1c). We conclude with **ST-ANTI-SYM**. \square

Note that the rule **TEQ-SING** (and by transitivity, the rule **ST-INTV**) plays an important role in the proof of subject reduction for types (**Theorem 3.3**). It is used in the case for **K-SING**, where it allows us to relate β -equal types inhabiting singleton kinds.

B.5 Admissible Rules for Higher-Order Extrema and Intervals

In this section, we state and prove admissible rules that justify the encodings of higher-order extremal types and interval kinds given in §3.1.1 of the paper (page 8). Many of these rules are straightforward generalizations of the corresponding rules for the types \top , \perp and for proper type intervals $A .. B$. The remaining rules and lemmas mostly deal with the family of kinds $*_K$, which plays a crucial role in the other encodings and the proofs of their respective properties.

We start by stating and proving a formation rule for $*_K$.

Lemma B.21. *The kind $*_K$ is well-formed whenever K is, i.e. the following is admissible.*

$$\frac{\Gamma \vdash K \text{ kd}}{\Gamma \vdash *_K \text{ kd}} \text{(WF-KMAX)}$$

PROOF. By induction on the structure of K . The base case uses **K-BOT**, **K-TOP** and **WF-INTV** to show that $*$ is well-formed. \square

The kind $*_K$ is a widened version of K , i.e. the latter is always a subkind of the former. As a consequence, any type of kind K is also of kind $*_K$.

Lemma B.22. *Any well-formed kind K is a subkind of $*_K$.*

$$\frac{\Gamma \vdash K \text{ kd}}{\Gamma \vdash K \leq *_K} \text{(SK-KMAX)}$$

PROOF. By straightforward induction on the structure of K . \square

Corollary B.23. *If $\Gamma \vdash A : K$, then also $\Gamma \vdash A : *_K$.*

The following two lemmas introduce admissible kinding rules for the higher-order extremal types, and prove that \perp_K and \top_K are in fact extrema in $*_K$, i.e. they are the least and greatest inhabitants of $*_K$, respectively.

Lemma B.24. *Higher-order extremal types are well-formed if their index kind is.*

$$\frac{\Gamma \vdash K \text{ kd}}{\Gamma \vdash \top_K : *_K} \text{(K-TMAX)} \quad \frac{\Gamma \vdash K \text{ kd}}{\Gamma \vdash \perp_K : *_K} \text{(K-TMIN)}$$

PROOF. Separately, by induction on the structure of K . The cases for dependent arrow kinds use **WF-KMAX**. \square

Lemma B.25. *The types \top_K and \perp_K are the maximal and minimal elements of $*_K$, respectively.*

$$\frac{\Gamma \vdash A : K}{\Gamma \vdash A \leq \top_K : *_K} \text{(ST-TMAX)} \quad \frac{\Gamma \vdash A : K}{\Gamma \vdash \perp_K \leq A : *_K} \text{(ST-TMIN)}$$

PROOF. Separately, by induction on the structure of K . [Corollary B.23](#) is used to adjust the kind of the premises where necessary. In the inductive step, we use [ST-Abs](#) and the η -rules [ST- \$\eta_{1,2}\$](#) . For example, for $K = (X:K_1) \rightarrow K_2$ we have

$$\begin{aligned} \Gamma \vdash A &\leq \lambda X:K_1. AX && \text{for } X \notin \text{fv}(A) && \text{(by ST-}\eta_2\text{)} \\ &\leq \lambda X:K_1. \top_{K_2} && && \text{(by the IH and ST-Abs)} \\ &\equiv \top_{(X:K_1) \rightarrow K_2} : *_{(X:K_1) \rightarrow K_2}. && && \text{(by definition)} \end{aligned}$$

□

Having generalized the properties of the extremal types to their higher-order counterparts, we now turn to interval kinds. We start with an admissible formation rule for higher-order intervals.

Lemma B.26. *Higher-order interval kinds are well-formed if their bounds are.*

$$\frac{\Gamma \vdash A : K \quad \Gamma \vdash B : K}{\Gamma \vdash A .._K B \text{ kd}} \text{ (WF-HoINTV)}$$

PROOF. By induction on the structure of K . The inductive step uses kinding validity, [K-Var](#) and [K-App](#) to expand the bounds. □

The subkinding rule [SK-INTV](#) for proper type intervals also generalizes straightforwardly to intervals over arbitrary type operators.

Lemma B.27. *Higher-order interval kinds are widened in accordance with their bounds.*

$$\frac{\Gamma \vdash A_2 \leq A_1 : K \quad \Gamma \vdash B_1 \leq B_2 : K}{\Gamma \vdash A_1 .._K B_1 \leq A_2 .._K B_2} \text{ (SK-HoINTV)}$$

PROOF. By induction on the structure of K . The inductive step uses subtyping validity, [K-Var](#), [TEQ-REFL](#) and [ST-APP](#) to expand the bounds, and [WF-HoINTV](#) to establish well-formedness of the left-hand side. □

Next, we would like to prove an admissible higher-order singleton introduction rule that generalizes [K-SING](#). Ideally, we would like to show that any well-kinded type $\Gamma \vdash A : K$ inhabits its corresponding singleton kind $S(A : K) = A .._K A$. This is not necessarily true, however. Consider the case of an operator variable X with declared type $\Gamma(X) = * \rightarrow *$. The singleton kind corresponding to X is $S(X : * \rightarrow *) = (Y:*) \rightarrow X Y .. X Y$, so we would like to prove that $\Gamma \vdash X : (Y:*) \rightarrow X Y .. X Y$. Which kinding rules could we use to adjust the kind of X to the desired singleton kind? Since [K-SING](#) can only be applied to proper types, our only option is to use the subsumption rule [K-SUB](#). But unfortunately, the declared kind $* \rightarrow *$ of X is a *strict supertype* of the singleton kind $(Y:*) \rightarrow X Y .. X Y$, so this cannot work.

We can, however, assign the desired singleton kind to the η -expansion of X , i.e. to $\lambda Y:*. X Y$. Unlike X , the application $X Y$ in the body of the η -expansion is a proper type, so we can use [K-SING](#) to narrow its kind. The full derivation is

$$\frac{\Gamma \vdash * \text{ kd} \quad \frac{\Gamma, Y:*\vdash X Y : *}{\Gamma, Y:*\vdash X Y : *} \text{ (K-SING)}}{\Gamma \vdash * \text{ kd} \quad \Gamma, Y:*\vdash X Y : X Y .. X Y} \text{ (K-Abs)}$$

$$\frac{\Gamma, Y:*\vdash X : * \rightarrow * \quad \Gamma, Y:*\vdash Y : *}{\Gamma, Y:*\vdash X Y : *} \text{ (K-App)}$$

$$\frac{\Gamma \vdash * \text{ kd} \quad \Gamma, Y:*\vdash X Y : X Y .. X Y}{\Gamma \vdash \lambda Y:*. X Y : (Y:*) \rightarrow X Y .. X Y}$$

This principle generalizes to arbitrary well-kinded types: the η -expansion of a well-kinded type $\Gamma \vdash A : K$ always inhabits the corresponding singleton kind $S(A : K)$.

Given a type A , we define the *weak η -expansion* $\bar{\eta}_K(A)$ of A as $\bar{\eta}_{B..C}(A) = A$ and $\bar{\eta}_{(X:J) \rightarrow K}(A) = \lambda X:J. \bar{\eta}_K(A X)$ where, as usual, we assume that $X \notin \text{fv}(A)$. We call this expansion “weak” because the argument X in the definition $\lambda X:J. \bar{\eta}_K(A X)$ of the arrow case is not η -expanded further. This means that the result is not η -long. This is sufficient for the purpose of this section; we will define a stronger version in the next section.

As expected, a type of kind K is equal to its weak η -expansion in K .

Lemma B.28. *Weak η -expansion is sound, i.e. if $\Gamma \vdash A : K$, then $\Gamma \vdash A = \bar{\eta}_K(A) : K$.*

PROOF. By induction on the structure of K , using **TEQ- η** and **TEQ-Abs** in the inductive case. \square

Lemma B.29. *The η -expansions of type operators inhabit their higher-order singleton intervals.*

$$\frac{\Gamma \vdash A : K}{\Gamma \vdash \bar{\eta}_K(A) : S(A : K)} \text{ (K-HoSING)}$$

PROOF. By induction on the structure of K . The base case follows from **K-SING**, the inductive step from the usual combination of kinding validity, **K-VAR**, **K-APP** and **WF-HoINTV**. \square

Corollary B.30. *If $\Gamma \vdash B_1 \leq A : K$ and $\Gamma \vdash A \leq B_2 : K$, then $\Gamma \vdash \bar{\eta}_K(A) : B_1 .._K B_2$.*

Having found ways to form, widen and populate higher-order intervals, we still need a way to put their bounds to use. To this end, we introduce two *higher-order bound projection* rules, which generalize the corresponding rules **ST-BND₁** and **ST-BND₂** for proper type intervals.

Lemma B.31 (higher-order bound projection). *Inhabitants of a higher-order interval are supertypes of its lower bound and subtypes of its upper bound.*

$$\frac{\Gamma \vdash A : K \quad \Gamma \vdash B_1 : K}{\Gamma \vdash A : B_1 .._K B_2} \text{ (ST-HoBND}_1\text{)} \qquad \frac{\Gamma \vdash A : K \quad \Gamma \vdash B_2 : K}{\Gamma \vdash A : B_1 .._K B_2} \text{ (ST-HoBND}_2\text{)}$$

These rules are a bit weaker than one might expect. In particular, the additional premises $\Gamma \vdash A : K$, $\Gamma \vdash B_1 : K$ and $\Gamma \vdash B_2 : K$, might seem redundant. They are necessary because we cannot, in general, invert well-formedness judgments about higher-order intervals. That is, $\Gamma \vdash B_1 .._K B_2$ kd does *not* imply $\Gamma \vdash B_1 : K$ and $\Gamma \vdash B_2 : K$, nor does $\Gamma \vdash A : B_1 .._K B_2$ imply $\Gamma \vdash A : K$. To see this, consider the kind $K = \perp .._\emptyset \top$, where $\emptyset = \top .. \perp$ is the empty interval (note the absurd bounds). The kind K is well-formed and inhabited by both \perp and \top , yet clearly \perp, \top are not inhabitants of \emptyset . Note that the formation rule **WF-HoINTV** for higher-order intervals is not to blame: although K is well-formed, we cannot prove this fact using **WF-HoINTV**. There are simply more well-formed higher-order intervals than can be derived using **WF-HoINTV**.

PROOF OF **LEMMA B.31**. Separately, by induction on the structure of K . In the base case, we use the interval projection rules **ST-BND_{1,2}** as well as **ST-INTV** and **ST-SUB** to adjust the kinds of the resulting inequations. In the inductive step, we use **ST-Abs** and the η -rules **ST- $\eta_{1,2}$** . For example, for the left-hand case and $K = (X:K_1) \rightarrow K_2$ we have

$$\begin{aligned} \Gamma \vdash B_1 &\leq \lambda X:K_1. B_1 X && \text{(by ST-}\eta_2\text{)} \\ &\leq \lambda X:K_1. A X && \text{(by the IH and ST-Abs)} \\ &\leq A && : (X:K_1) \rightarrow K_2. \text{(by ST-}\eta_1\text{)} \end{aligned}$$

\square

Thanks to the admissible kinding and subtyping rules for higher-order intervals and extrema, we can now easily derive judgments for forming, introducing or eliminating bounded universal quantifiers over arbitrary type operators.

For example, well-formedness of the higher-order universal quantifier $\forall X \leq A : K . B$ can be derived as

$$\begin{array}{c}
 \text{(kinding validity)} \frac{\Gamma \vdash A : K}{\Gamma \vdash K \text{ kd}} \\
 \text{(K-TMIN)} \frac{\Gamma \vdash A : K}{\Gamma \vdash \perp_K : *K} \quad \text{(Corollary B.23)} \frac{\Gamma \vdash A : K}{\Gamma \vdash A : *K} \\
 \text{(WF-HOINTV)} \frac{\Gamma \vdash \perp_K : *K \quad \Gamma \vdash A : *K}{\Gamma \vdash \perp_K \dots *K A \text{ kd}} \\
 \text{(Lemma B.32)} \frac{\Gamma \vdash \perp_K \dots *K A \text{ kd}}{\Gamma \vdash \perp_K \dots K A \text{ kd}} \\
 \text{(K-ALL)} \frac{\Gamma \vdash \perp_K \dots K A \text{ kd} \quad \Gamma, X : \perp_K \dots K A \vdash B : *}{\Gamma \vdash \forall X \leq A : K . B : *}
 \end{array}$$

The derivation uses the following lemma for simplifying interval kinds; its proof is by structural induction on the index K .

Lemma B.32. *Let A, B be types and K a kind. Then $A \dots *K B \equiv A \dots K B$*

Similar derivations exist for the introduction and elimination rules.

Corollary B.33 (bounded quantification). *The following rules for the formation, introduction and elimination of bounded universal quantifiers are admissible.*

$$\begin{array}{c}
 \frac{\Gamma \vdash A : K \quad \Gamma, X : \perp_K \dots K A \vdash B : *}{\Gamma \vdash \forall X \leq A : K . B : *} \text{(K-ALLBND)} \\
 \frac{\Gamma \vdash A : K \quad \Gamma, X : \perp_K \dots K A \vdash t : B}{\Gamma \vdash \lambda X \leq A : K . t : \forall X \leq A : K . B} \text{(T-TABSND)} \\
 \frac{\Gamma \vdash t : \forall X \leq A : K . B \quad \Gamma \vdash C \leq A : K}{\Gamma \vdash t (\bar{\eta}_K(C)) : B[\bar{\eta}_K(C)/X]} \text{(T-TAPPBND)}
 \end{array}$$

Similar rules for the formation, abstraction and elimination of bounded operators are also admissible.

Note that we need to η -expand the type argument C in the elimination rule **T-TAPPBND** before it can be applied to t . This is because C has kind K , while the polymorphic expression t expects an argument of kind $\perp_K \dots K C$. As discussed earlier, C is not guaranteed to inhabit that kind but its η -expansion is – via **K-HOSING** and a subsequent widening of its kind from $C \dots K C$ to $\perp_K \dots K C$.

There is an alternative encoding of higher-order bounded quantification (and bounded type operators) that separates the declaration of type variables from that of the subtyping constraints imposed by their bounds, at the cost of using an auxiliary type variable with potentially inconsistent bounds. Assume a partition of the set of type variable names into two distinct sets of *operator names* denoted by X_n, Y_n, \dots and *constraint names* denoted by X_c, Y_c, \dots . We may then encode an upper-bounded type variable binding $X \leq A : K$ as a pair of bindings $X_n : K, X_c : X_n \dots K A$, separating the declaration of the operator name X from the subtyping constraint $X \leq A$. For example, the encoding of bounded universal quantifiers according to this scheme would be $\forall X \leq A : K . B = \forall X_n : K . \forall X_c : X_n \dots K A . B$ where $X_c \notin \text{fv}(B)$.

The advantage of this encoding is a cleaner separation between the uses of bounded variable bindings in kinding and subtyping. Whenever we want to refer to the original type variable X or its kind, we simply use X_n . When we require a proof of the fact that $X \leq A$ we obtain one from X_c via **ST-HOBIND₁**, **ST-HOBIND₂**, and **ST-TRANS**. The same is true when we instantiate type parameters. For example, a type application $t C$, where t has type $\forall X \leq A : K . B$ and $C \leq A$, is now desugared to $t C (\bar{\eta}_K(C))$, i.e. only the second type argument, which corresponds to the constraint

parameter X_c , needs to be η -expanded, while the argument C for the parameter X_n can be left as is. Since X_c does not occur freely in the codomain B of the desugared universal type, the overall type of the desugared application is just $B[C/X_n]$. A clear drawback of this encoding is the necessary duplication of bindings and the corresponding introduction and elimination forms (abstraction, application). In addition, the kind $X_n \dots_K A$ of the constraint X_c has inconsistent bounds in general, which can be problematic.

There is an obvious alternative definition for the family of kinds $*_K$, namely $*_K = (\perp_K) \dots_K (\top_K)$. The original definition, given in Fig. 2 of the paper (page 8), has the advantage of being independent of the definition of the higher-order extrema \top_K and \perp_K . This allowed us to prove properties such as **WF-KMAX** and **Lemma B.22** admissible without appealing to any of the properties of higher-order extrema, and thereby avoid some cyclic dependencies in the proofs of the latter. The alternative definition, on the other hand, seems more intuitive. To conclude the section, we show that the two definitions are equal for well-formed kinds K .

Lemma B.34. *The kind $*_K$ is equal to the higher-order interval bounded by \perp_K and \top_K . If $\Gamma \vdash K$ kd, then $\Gamma \vdash *_K = (\perp_K) \dots_K (\top_K)$.*

PROOF OF LEMMA B.34. By induction on the structure of K . The base case is immediate. In the inductive step, we use **SK-HOINTV** and the β -rule **TEQ- β** .

Let $K = (X:K_1) \rightarrow K_2$. We want to show that

$$\Gamma \vdash (X:K_1) \rightarrow *_K = (Y:K_1) \rightarrow (\perp_{(X:K_1) \rightarrow K_2} Y) \dots_{K_2} (\top_{(X:K_1) \rightarrow K_2} Y)$$

for some Y that does not occur freely in $\text{fv}(\perp_{(X:K_1) \rightarrow K_2})$ or $\text{fv}(\top_{(X:K_1) \rightarrow K_2})$. By the IH, we have $\Gamma \vdash *_K = (\perp_{K_2}) \dots_{K_2} (\top_{K_2})$ but we need to adjust the bounds of the right-hand interval. We use the following equation for the lower bound, and an similar one for the upper bound.

$$\begin{aligned} \Gamma \vdash \perp_{(X:K_1) \rightarrow K_2} Y &\equiv \lambda X:K_1. \perp_{K_2} Y && \text{(by definition)} \\ &= \perp_{K_2} [Y/X] && \text{(by K-TMIN and TEQ-}\beta\text{)} \\ &\equiv \perp_{K_2} && : *_K. \quad (\alpha\text{-renaming}) \end{aligned}$$

By **SK-REFL-KEQ**, **SK-HOINTV** and **SK-ANTI-SYM** we obtain

$$\Gamma \vdash (\perp_{K_2}) \dots_{*_K_2} (\top_{K_2}) = (\perp_{(X:K_1) \rightarrow K_2} Y) \dots_{*_K_2} (\top_{(X:K_1) \rightarrow K_2} Y)$$

which we re-index using **Lemma B.32**. We conclude by **KEQ-REFL** and **KEQ-DARR**. \square

C SIMPLE KINDING OF NORMAL TYPES

This section introduces a system of *simplified kinding* judgments which provide a syntactic characterization of normal types and allow us to establish important properties about hereditary substitutions and normal forms, notably a pair of commutativity lemmas (**Lemmas C.11** and **C.21**) that play an important role in the development of §5 in the paper.

The contents of this section are based on Chapter 4 of the first author's PhD dissertation. We refer the interested reader to the dissertation for the full details [**Stucki 2017**, Ch. 4].

C.1 Preliminaries

Before we can introduce the simplified system, we need to introduce a few auxiliary definitions and lemmas that were omitted from §4 in the paper.

We start with the definition of *shape contexts* – the type of context used in our simplified kinding judgments. Shape contexts γ, δ are best thought of as typing contexts consisting exclusively of type

variable bindings $X:k$ with shape annotations k (as opposed to full kind annotations K). Their grammar is defined as follows.

$$\gamma, \delta ::= \emptyset \mid \gamma, X:k \quad (\text{Shape context})$$

As for full contexts, we assume that the variables bound in a shape context are all distinct. We write $\text{dom}(\gamma)$ for the set of variables bound in γ and (γ, δ) for the concatenation of two shape contexts with disjoint domains.

C.1.1 Weak Equality. Recall that *weak equality* is an equivalence $A \approx B$ on types and kinds that identifies operator abstractions up to the shape of their domain annotations.

$$\frac{|J| \equiv |K| \quad A \approx B}{\lambda X:J. A \approx \lambda X:K. B} \text{ (WEQ-ABS)}$$

Formally, weak equality is defined as the smallest congruence (w.r.t. to all the type and kind formers) that includes both syntactic equality and the above rule **WEQ-ABS**. It is easy to verify that weak equality is a congruence w.r.t. ordinary substitution. In the following sections, we will show that weak equality is also a congruence w.r.t. hereditary substitution, η -expansion and normalization, and that weakly equal kinds $K_1 \approx K_2$ have equal shapes $|K_1| \equiv |K_2|$.

C.1.2 Properties of Hereditary Substitution. Since it is defined pointwise on spines, hereditary substitution commutes with spine concatenation.

Lemma C.1. $(\mathbf{D}_1, \mathbf{D}_2)[E/X^k] \equiv (\mathbf{D}_1[E/X^k], \mathbf{D}_2[E/X^k])$.

Just as for ordinary substitution, shapes are stable under hereditary substitution.

Lemma C.2 (stability of shapes under hereditary substitution). $|J[A/X^k]| \equiv |J|$.

PROOF. By straightforward induction on the structure of J . \square

As a consequence, weak equality is a congruence w.r.t. hereditary substitution and reducing application.

Lemma C.3. *Weak equality is a congruence w.r.t. hereditary substitution and reducing application.*

Let $E_1 \approx E_2$,

- (1) if $K_1 \approx K_2$, then $K_1[E_1/X^k] \approx K_2[E_2/X^k]$;
- (2) if $D_1 \approx D_2$, then $D_1[E_1/X^k] \approx D_2[E_2/X^k]$;
- (3) if $\mathbf{D}_1 \approx \mathbf{D}_2$, then $\mathbf{D}_1[E_1/X^k] \approx \mathbf{D}_2[E_2/X^k]$;
- (4) if $D_1 \approx D_2$, then $E_1 \cdot^k D_1 \approx E_2 \cdot^k D_2$;
- (5) if $\mathbf{D}_1 \approx \mathbf{D}_2$, then $E_1 \cdot^k \mathbf{D}_1 \approx E_2 \cdot^k \mathbf{D}_2$.

PROOF. The structure of the proof mirrors that of the recursive definitions of hereditary substitution and reducing application. All five parts are proven simultaneously, by induction on the structure of k . Parts 1–3 proceed by an inner induction on the derivations of $K_1 \approx K_2$, $D_1 \approx D_2$ and $\mathbf{D}_1 \approx \mathbf{D}_2$, respectively. Parts 4 and 5 proceed by a case analysis on the final rules used to derive $E_1 \approx E_2$ and $\mathbf{D}_1 \approx \mathbf{D}_2$, respectively. Part 2 proceeds by a case analysis on the final rule used to derive $F_1 \approx F_2$, where F_1 and F_2 are the heads, respectively, of $E_1 = F_1 \mathbf{D}_1$ and $E_2 = F_2 \mathbf{D}_2$. In the case for **WEQ-ABS**, we use stability of kind simplification under hereditary substitution (**Lemma C.2**). In the variable case, we use the IH twice: first for part 3 to derive $\mathbf{D}_1[E_1/X^k] \approx \mathbf{D}_2[E_2/X^k]$, then for part 5, to derive $E_1 \cdot^k (\mathbf{D}_1[E_1/X^k]) \approx E_2 \cdot^k (\mathbf{D}_2[E_2/X^k])$. In the second instance, k does not decrease nor is $\mathbf{D}_1[E_1/X^k] \approx \mathbf{D}_2[E_2/X^k]$ a sub-derivation of the current premise. This use of the IH is nevertheless justified because any subsequent use of the IH for part 2 in the proof of part 5 must occur after the use of the IH for part 4, at which point k has necessarily decreased. \square

C.1.3 Properties of η -Expansion and Normalization. Unsurprisingly, shapes are stable under normalization.

Lemma C.4 (stability of shapes under normalization). $|\text{nf}_\Gamma(K)| \equiv |K|$.

PROOF. By straightforward induction on the structure of K . \square

Extending kind simplification pointwise to contexts, we define $|\Gamma|$ as

$$|\emptyset| = \emptyset \qquad |\Gamma, x:A| = |\Gamma| \qquad |\Gamma, X:K| = |\Gamma|, X:|K|$$

It is easy to see that context lookup commutes with simplification, i.e. $|\Gamma|(X) = |\Gamma(X)|$, and that simplified contexts are stable under (hereditary) substitution and normalization, i.e.

$$|\Gamma[A/X]| = |\Gamma| \qquad |\Gamma[A/X^k]| = |\Gamma| \qquad |\text{nf}(\Gamma)| = |\Gamma|$$

The following two lemmas show that η -expansion and normalization preserve weak equality. Importantly, this is true even if the corresponding kinds and contexts, respectively, are not themselves weakly equal but have equal shapes – a much weaker requirement.

Lemma C.5. *Weak equality is preserved by η -expansion along kinds of equal shape. If $|J| \equiv |K|$ and $D \approx E$, then $\eta_J(D) \approx \eta_K(E)$.*

PROOF. By induction on the structure of J and case analysis on the final rule used to derive $D \approx E$. \square

Lemma C.6. *Kinds and types normalize weakly equally in contexts that simplify equally. Let Γ and Δ be contexts such that $|\Gamma| \equiv |\Delta|$. Then*

- (1) $\text{nf}_\Gamma(K) \approx \text{nf}_\Delta(K)$ for any kind K , and
- (2) $\text{nf}_\Gamma(A) \approx \text{nf}_\Delta(A)$ for any type A .

PROOF. Simultaneously, by induction on the structure of K and A , respectively. In the type variable case $A = X$ we use [Lemma C.5](#); in the operator application case $A = A_1 A_2$ we use [Lemma C.3.2](#); in the cases for dependent operator kinds, universal types and operator abstraction, we use [Lemma C.18](#). \square

C.2 The Simplified System

The function nf assigns to each raw type A in a given context Γ a unique type $\text{nf}_\Gamma(A)$. But as we have seen, the type $\text{nf}_\Gamma(A)$ may not be $\beta\eta$ -normal if A is ill-kinded. In this section, we prove the converse: whenever A is well-kinded in Γ , the type $\text{nf}_{\text{nf}(\Gamma)}(A)$ is a η -long β -normal form. To do so, we first introduce a set of *simplified kinding* judgments. Roughly, a simplified kinding judgment $\gamma \vdash E : k$ establishes that the type E is a normal form of *shape* k in the *shape context* γ . Given $\gamma \vdash E : k$, we say that E is a *well-shaped* or *simply (well-)kinded normal form*, or just that E is *simply kinded*. As we are about to show, every well-kinded type $\Gamma \vdash A : K$ has a well-shaped normal form $|\Gamma| \vdash E : |K|$, namely $E = \text{nf}_{\text{nf}(\Gamma)}(A)$ (see [Lemma C.14](#) below).

It is important to note that the converse is not true: not every simply kinded type is well-kinded. Because kind simplification forgets dependencies, there are necessarily some ill-kinded types that are considered simply well-kinded according to the judgments we are about to introduce. However, every simply kinded type is guaranteed to be an η -long β -normal form and, as we will see in this section, simple kinding is preserved by operations such as hereditary substitution and η -expansion. Hence simple kinding allows us to prove important properties about these operations on $\beta\eta$ -normal forms without subjecting ourselves to the complexity of fully dependent kinds.

Simplified well-formedness of kinds

$$\frac{\gamma \vdash U : * \quad \gamma \vdash V : *}{\gamma \vdash U .. V \text{ kds}}$$

(SWF-INTV)

$$\frac{\gamma \vdash J \text{ kds} \quad \gamma, X : |J| \vdash K \text{ kds}}{\gamma \vdash (X : J) \rightarrow K \text{ kds}}$$

(SWF-DARR)

$$\boxed{\gamma \vdash K \text{ kds}}$$

Kinding of neutral types

$$\frac{\gamma(X) = j \quad \gamma \vdash j : \mathbf{V} : k}{\gamma \vdash_{\text{ne}} X \mathbf{V} : k}$$

(SK-VARAPP)

$$\boxed{\gamma \vdash_{\text{ne}} N : k}$$

Simple spine kinding

$$\frac{}{\gamma \vdash k : \epsilon : k}$$

(SK-EMPTY)

$$\frac{\gamma \vdash U : j \quad \gamma \vdash k : \mathbf{V} : l}{\gamma \vdash j \rightarrow k : U, \mathbf{V} : l}$$

(SK-CONS)

$$\boxed{\gamma \vdash j : \mathbf{V} : k}$$

Simple kinding of normal types

$$\frac{}{\gamma \vdash \top : *}$$

(SK-TOP)

$$\frac{}{\gamma \vdash \perp : *}$$

(SK-BOT)

$$\frac{\gamma \vdash U : * \quad \gamma \vdash V : *}{\gamma \vdash U \rightarrow V : *}$$

(SK-ARR)

$$\frac{\gamma \vdash K \text{ kds} \quad \gamma, X : |K| \vdash V : *}{\gamma \vdash \forall X : K. V : *}$$

(SK-ALL)

$$\frac{\gamma \vdash J \text{ kds} \quad \gamma, X : |J| \vdash V : k}{\gamma \vdash \lambda X : J. V : |J| \rightarrow k}$$

(SK-ABS)

$$\frac{\gamma \vdash_{\text{ne}} N : *}{\gamma \vdash N : *}$$

(SK-NE)

$$\boxed{\gamma \vdash V : k}$$

Fig. 9. Simplified kinding

To enhance readability, we use the following naming conventions for normal forms: the metavariables U, V, W denote normal types, while M and N denote neutral types.⁷ No special notation is used for normal kinds.

Judgments. Fig. 9 defines the following judgments by mutual induction.

$\gamma \vdash K \text{ kds}$	the kind K is simply well-formed and normal in γ
$\gamma \vdash V : k$	the type V is a normal form of shape k in γ
$\gamma \vdash_{\text{ne}} N : k$	the type N is a neutral form of shape k in γ
$\gamma \vdash j : \mathbf{V} : k$	applying an operator of shape j to the normal spine \mathbf{V} yields a type of shape k in γ .

The judgments for simple kind formation and kinding follow the syntactic structure of normal kinds and types. A type V is a $\beta\eta$ -normal form $\gamma \vdash V : k$ of shape k if it is either a proper type introduced by one of the basic type formers applied to normal arguments (rules **SK-TOP**, **SK-BOT**, **SK-ARR**, and **SK-ALL**), an operator abstraction with a normal body (rule **SK-ABS**), or a simply kinded neutral type (rule **SK-NE**). Simply kinded neutral forms $\gamma \vdash_{\text{ne}} N : k$ are eliminations headed by an abstract type operator, i.e. a type variable X , which is applied to a spine of normal types \mathbf{V}

⁷This is just notation. We do not consider normal forms a separate syntactic category, e.g. the letters U, V, W are metavariables denoting types (typically in elimination form) rather than non-terminals in some grammar of normal forms.

(rule **SK-VARAPP**). Finally, a simply well-formed normal kind $\gamma \vdash K$ kds is either a type interval bounded by normal types (rule **SWF-INTV**) or a dependent arrow with normal domain and codomain (rule **SWF-DARR**). Note that type operator abstractions are the only normal forms of arrow shape. This ensures that normal types are always η -long.

The simple spine kinding judgment $\gamma \vdash j : \mathbf{V} : k$ is different from the other judgment forms in that it is a quaternary rather than a ternary relation. The shapes j and k should be read as inputs and outputs, respectively, of such judgments: when a type of shape j is applied to the spine \mathbf{V} (the subject of the judgment), the resulting type is of shape k – as exemplified by the rule **SK-VARAPP**.

There is no formation judgment for shape contexts γ since such contexts only contain bindings assigning shapes to type variables, and there is no such thing as an ill-formed simple kind.

Because kinding is simplified, there is no notion of subkinding or kind equality, and hence no need for a subsumption rule. As a consequence, the simple kind formation and kinding rules are syntax-directed. Another important property of simplified kinding is that none of the rules involve substitutions in kinds. This substantially simplifies the proofs of key lemmas about hereditary substitutions, in particular that of **Lemma C.10** which states that hereditary substitutions preserve simple kinding (and thus normal forms). It is also important in establishing admissibility of the following simple rules about spines and neutral types.

Lemma C.7. *The following simple kinding rules for spine concatenation and application of neutrals are admissible.*

$$\frac{\gamma \vdash j : \mathbf{U} : k \quad \gamma \vdash k : \mathbf{V} : l}{\gamma \vdash j : \mathbf{U}, \mathbf{V} : l} \text{ (SK-CONCAT)} \quad \frac{\gamma \vdash j : \mathbf{U} : k \rightarrow l \quad \gamma \vdash V : k}{\gamma \vdash j : \mathbf{U}, V : l} \text{ (SK-SNOC)}$$

$$\frac{\gamma \vdash_{\text{ne}} N : j \rightarrow k \quad \gamma \vdash V : j}{\gamma \vdash_{\text{ne}} NV : k} \text{ (SK-NEAPP)}$$

PROOF. The proofs are done separately for each of the three rules in the order the rules are listed. The proof for **SK-CONCAT** is by induction on the derivation of the first premise. The rule **SK-CONS** is derivable from **SK-SNOC** as a special case where $\mathbf{V} = V, \epsilon$, using **SK-EMPTY** and **SK-CONS**. The proof of **SK-NEAPP** starts with a case analysis on the final rule used to derive $\Gamma \vdash_{\text{ne}} N : j \rightarrow k$. The only rule for deriving such judgments is **SK-VARAPP**, hence N must be of the form $N = X U$ with $\gamma(X) = l$ and $\gamma \vdash l : \mathbf{U} : j \rightarrow k$. We conclude by **SK-SNOC** and **SK-VARAPP**. \square

C.2.1 Simply-Kinded Hereditary Substitution. Before we can prove that hereditary substitutions preserve simple kinding, we first need to establish the usual weakening properties for simple kind formation and kinding.

Lemma C.8 (weakening). *A simple judgment remains true if its context is extended by an additional binding. Let γ, δ be shape contexts, k a shape and $X \notin \text{dom}(\gamma, \delta)$. If $\gamma, \delta \vdash \mathcal{J}$ for any of the simple judgments defined above, then $\gamma, X : k, \delta \vdash \mathcal{J}$.*

PROOF. Simultaneously for all four judgments, by induction on the derivation of $\gamma, \delta \vdash \mathcal{J}$. \square

Corollary C.9 (Iterated weakening). *Given a pair γ, δ of disjoint shape contexts, if $\gamma \vdash \mathcal{J}$, then $\gamma, \delta \vdash \mathcal{J}$.*

Lemma C.10 (hereditary substitution). *Hereditary substitutions and reducing applications preserve the shapes of types and simple well-formedness of kinds. Let γ, δ be shape contexts and X such that $X \notin \text{dom}(\gamma, \delta)$. Assume further that $\gamma \vdash V : k$ for some V and k . Then*

- (1) if $\gamma, X : k, \delta \vdash J$ kds, then $\gamma, \delta \vdash J[V/X^k]$ kds;
- (2) if $\gamma, X : k, \delta \vdash U : j$, then $\gamma, \delta \vdash U[V/X^k] : j$;

- (3) if $\gamma, X:k, \delta \vdash_{\text{ne}} N : j$, then $\gamma, \delta \vdash N[V/X^k] : j$ as a normal form;
- (4) if $\gamma, X:k, \delta \vdash j : \mathbf{U} : l$, then $\gamma, \delta \vdash j : \mathbf{U}[V/X^k] : l$;
- (5) if $k = k_1 \rightarrow k_2$ and $\gamma \vdash U : k_1$, then $\gamma \vdash V.^{k_1 \rightarrow k_2} U : k_2$;
- (6) if $\gamma \vdash k : \mathbf{U} : j$, then $\gamma \vdash V.^k \mathbf{U} : j$.

Note that hereditary substitutions preserve the shapes of neutral types but not neutrality itself.

PROOF. All six parts are proven simultaneously by induction on the structure of k . Parts 1–4 proceed by an inner induction on the simple formation or kinding derivations for J, U, N and \mathbf{U} , respectively. Parts 5 and 6 proceed by a case analysis on the final rules used to derive $\gamma \vdash V : k_1 \rightarrow k_2$ and $\gamma \vdash k : \mathbf{V} : j$, respectively; for part 5, the only applicable rule is **SK-Abs**. For part 3, in the case for **SK-VARAPP** when $N = X \mathbf{U}$, we use iterated weakening (**Corollary C.9**) and the IH (for 4), respectively, to obtain $\gamma, \delta \vdash V : k$ and $\gamma, \delta \vdash k : \mathbf{U}[V/X^k] : j$. To conclude the case, we apply the IH again (for 6). In this second use of the IH, k does not decrease nor is $\gamma, \delta \vdash k : \mathbf{U}[V/X^k] : j$ a strict sub-derivation of the current premise. However, in order to use the IH for part 3 again from within the proof of part 6, we must go through part 5, at which point k necessarily decreases. Again, the structure of the proof mirrors that of the mutually recursive definitions of hereditary substitution and reducing application. \square

Thanks to **Lemma C.10**, we can now prove the following commutativity lemma about hereditary substitutions, which will play an important role in the proof of **Lemma C.21** below and in §5 of the paper.

Lemma C.11 (commutativity of hereditary substitutions). *Hereditary substitutions of simply kinded types commute; hereditary substitutions of simply kinded types commute with simply kinded reducing applications. Let $\gamma_1 \vdash U : j$ and $\gamma_1, X:j, \gamma_2 \vdash V : k$. Then*

- (1) if $\gamma_1, X:j, \gamma_2, Y:k, \gamma_3 \vdash J$ kds, then

$$J[V/Y^k][U/X^j] \equiv J[U/X^j][V[U/X^j]/Y^k];$$

- (2) if $\gamma_1, X:j, \gamma_2, Y:k, \gamma_3 \vdash W : l$, then

$$W[V/Y^k][U/X^j] \equiv W[U/X^j][V[U/X^j]/Y^k];$$

- (3) if $\gamma_1, X:j, \gamma_2, Y:k, \gamma_3 \vdash_{\text{ne}} N : l$, then

$$N[V/Y^k][U/X^j] \equiv N[U/X^j][V[U/X^j]/Y^k];$$

- (4) if $\gamma_1, X:j, \gamma_2, Y:k, \gamma_3 \vdash l_1 : \mathbf{W} : l_2$, then

$$\mathbf{W}[V/Y^k][U/X^j] \equiv \mathbf{W}[U/X^j][V[U/X^j]/Y^k];$$

- (5) if $k = k_1 \rightarrow k_2$ and $\gamma_1, X:j, \gamma_2 \vdash W : k_1$, then

$$(V.^{k_1 \rightarrow k_2} W)[U/X^j] \equiv (V[U/X^j]).^{k_1 \rightarrow k_2} (W[U/X^j]);$$

- (6) if $\gamma_1, X:j, \gamma_2 \vdash k : \mathbf{W} : l$, then $(V.^k \mathbf{W})[U/X^j] \equiv (V[U/X^j]).^k (\mathbf{W}[U/X^j])$.

PROOF. All six parts are proven simultaneously by simultaneous induction on the structures of j and k . Simultaneous structural induction on j and k means roughly that it is sufficient for either one of j or k to decrease in an induction step. More formally, denote by \sqsubseteq the sub-expression order on shapes, then the simultaneous induction order $<$ on unordered pairs $\{j, k\}$ of shapes is defined as $\{j_1, j_2\} < \{k_1, k_2\}$ if $j_1 \sqsubset k_1$ and $j_2 \sqsubseteq k_2$. Importantly, $<$ is defined over *unordered* pairs which allows us to exchange j and k in an induction step. Parts 1–4 proceed by an inner induction on the simple formation or kinding derivations for J, W, N and \mathbf{W} , respectively.

As usual, the interesting cases are those for part 3, when $N = Y \mathbf{W}$ and $N = X \mathbf{W}$.

- Case *SK-VARAPP*, $N = Y \mathbf{W}$. We have $\gamma_1, X:j, \gamma_2, Y:k, \gamma_3 \vdash k : \mathbf{W} : l$. By [Lemma C.10.4](#), we obtain $\gamma_1, X:j, \gamma_2, \gamma_3 \vdash k : \mathbf{W}[V/Y^k] : l$, and hence we have

$$\begin{aligned}
(Y \mathbf{W})[V/Y^k][U/X^j] & \equiv (V.^k (\mathbf{W}[V/Y^k]))[U/X^j] && \text{(by definition)} \\
& \equiv (V[U/X^j]).^k (\mathbf{W}[V/Y^k][U/X^j]) && \text{(by the IH for 6)} \\
& \equiv (V[U/X^j]).^k (\mathbf{W}[U/X^j][V[U/X^j]/Y^k]) && \text{(by the IH for 4)} \\
& \equiv (Y \mathbf{W})[U/X^j][V[U/X^j]/Y^k]. && \text{(by definition)}
\end{aligned}$$

- Case *SK-VARAPP*, $N = X \mathbf{W}$. We have $\gamma_1, X:j, \gamma_2, Y:k, \gamma_3 \vdash j : \mathbf{W} : l$. By [Lemma C.10.4](#), we obtain $\gamma_1, X:j, \gamma_2, \gamma_3 \vdash j : \mathbf{W}[V/Y^k] : l$, and hence we have

$$\begin{aligned}
(X \mathbf{W})[V/Y^k][U/X^j] & \equiv (X (\mathbf{W}[V/Y^k]))[U/X^j] && \text{(by definition)} \\
& \equiv U.^j (\mathbf{W}[V/Y^k][U/X^j]) && \text{(by definition)} \\
& \equiv U.^j (\mathbf{W}[U/X^j][V[U/X^j]/Y^k]) && \text{(by the IH for 4)} \\
& \equiv (U[V[U/X^j]/Y^k]).^j (\mathbf{W}[U/X^j][V[U/X^j]/Y^k]) && \text{(as } Y \notin \text{fv}(U)\text{)} \\
& \equiv (U.^j (\mathbf{W}[U/X^j]))[V[U/X^j]/Y^k] && \text{(by the IH for 6)} \\
& \equiv (X \mathbf{W})[U/X^j][V[U/X^j]/Y^k]. && \text{(by definition)}
\end{aligned}$$

Note that, in the second case, we switched the roles of the shape j and k when invoking the IH for part 6. \square

C.2.2 Simplification and Normalization of Kinding. Thanks to [Lemma 4.3](#) we know that the definition of the normalization function nf is sound, i.e. that well-formed kinds $\Gamma \vdash K \text{ kd}$ and well-kinded types $\Gamma \vdash A : K$ are convertible with $\text{nf}_{\text{nf}(\Gamma)}(K)$ and $\text{nf}_{\text{nf}(\Gamma)}(A)$, respectively. But we have yet to establish that $\text{nf}_{\text{nf}(\Gamma)}(K)$ and $\text{nf}_{\text{nf}(\Gamma)}(A)$ are actually normal forms. In this section, we prove a more general result, namely that, whenever $\Gamma \vdash K \text{ kd}$ and $\Gamma \vdash A : K$, it follows that $\text{nf}_{\text{nf}(\Gamma)}(K)$ is a simply well-formed normal kind and $\text{nf}_{\text{nf}(\Gamma)}(A)$ is a simply well-kinded normal type.

As a first step, we show that the shapes of variables and, more generally, of neutral types are preserved by η -expansion.

Lemma C.12. *η -expansion preserves the shapes of neutral types. Assume $\gamma \vdash K \text{ kds}$ and $\gamma \vdash_{\text{ne}} N : |K|$. Then $\gamma \vdash \eta_K(N) : |K|$.*

PROOF. By induction on the structure of K . The case fore $K = (X:K_1) \rightarrow K_2$ proceeds by case analysis on the final rules used to derive $\gamma \vdash (X:K_1) \rightarrow K_2 \text{ kds}$ and $\gamma \vdash_{\text{ne}} N : |K_1| \rightarrow |K_2|$ and uses the weakening lemma ([Lemma C.8](#)) as well as *SK-NEAPP*. \square

Next, we require a syntactic notion of normal contexts. We define the *simple context formation* judgment $\Gamma \text{ ctxs}$ as the pointwise lifting of simple kind formation and kinding to bindings:

$$\frac{}{\emptyset \text{ ctxs}} \quad \frac{\Gamma \text{ ctxs} \quad |\Gamma| \vdash K \text{ kds}}{\Gamma, X:K \text{ ctxs}} \quad \frac{\Gamma \text{ ctxs} \quad |\Gamma| \vdash V : *}{\Gamma, x:V \text{ ctxs}}$$

Since simple kind formation and kinding is defined on normal kinds and types, a simply well-formed context Γ is also normal. Conversely, if we lookup the declared kind or type of a variable in a simply well-formed context, the result is guaranteed to be a normal form.

Lemma C.13. *The declared kinds and types of variables in a simply well-formed context Γ are simply well-formed and well-kinded, respectively, in $|\Gamma|$, i.e*

$$\frac{\Gamma, X:K, \Delta \text{ ctxs}}{|\Gamma, X:K, \Delta| \vdash K \text{ kds}} \text{ (SC-TpLOOKUP)} \qquad \frac{\Gamma, x:V, \Delta \text{ ctxs}}{|\Gamma, x:V, \Delta| \vdash V : * } \text{ (SC-TmLOOKUP)}$$

PROOF. Both parts are proven separately by structural induction on Δ and case analysis on the final rule used to derive the premise. In the inductive case, we use the weakening lemma for simple kind formation. \square

With [Lemmas C.12](#) and [C.13](#) at hand, it is easy to show that nf does indeed produce normal forms.

Lemma C.14 (normalization and simplification). *Well-formed kinds and well-kinded types have simply well-formed and simply kinded normal forms, respectively.*

- (1) If $\Gamma \vdash K \text{ kd}$, then $|\text{nf}(\Gamma)| \vdash \text{nf}_{\text{nf}(\Gamma)}(K) \text{ kds}$.
- (2) If $\Gamma \vdash A : K$, then $|\text{nf}(\Gamma)| \vdash \text{nf}_{\text{nf}(\Gamma)}(A) : |K|$.
- (3) If $\Gamma \text{ ctx}$, then $\text{nf}(\Gamma) \text{ ctxs}$.

The proof uses the following helper lemma about the shapes of subkinds, which is proven by straightforward induction on subkinding derivations.

Lemma C.15. *Subkinds have equal shapes. If $\Gamma \vdash J \leq K$, then $|J| \equiv |K|$.*

PROOF OF [LEMMA C.14](#). Simultaneously by induction on declarative kind formation, kinding, and context formation derivations. The only interesting cases are [K-SUB](#) (where we use [Lemma C.15](#)), [K-VAR](#) and [K-APP](#). In the case for [K-VAR](#), where $A = X$, we use the IH for part 3 and [SC-TpLOOKUP](#) to obtain $|\text{nf}(\Gamma)| \vdash K \text{ kds}$ for $K = \text{nf}(\Gamma)(X)$, and we conclude by [Lemma C.12](#). In the case for [K-APP](#), where $A = A_1 A_2$, we start by applying the IH to obtain $|\text{nf}(\Gamma)| \vdash \text{nf}_{\text{nf}(\Gamma)}(A_1) : |K_1| \rightarrow |K_2|$, and $|\text{nf}(\Gamma)| \vdash \text{nf}_{\text{nf}(\Gamma)}(A_2) : |K_1|$. The first of these judgments must be derived using [SK-ABS](#) because that is the only simple kinding rule assigning an arrow kind to a type. Hence $\text{nf}_{\text{nf}(\Gamma)}(A_1) = \lambda X:J. V$ for some J and V such that $|J| \equiv |K_1|$ and $|\text{nf}(\Gamma), X:|K_1| \vdash V : |K_2|$. We conclude by the hereditary substitution lemma for normal types ([Lemma C.10.2](#)) and [Lemma C.2](#). \square

C.2.3 Commutativity of Normalization and Substitution. Our final task in this section is to establish another commutativity property that will play a crucial role in proving equality of declarative and canonical subtyping: the fact that normalization commutes with substitution.

In the past few sections, we have seen that well-formed kinds and well-kinded types have normal forms ([Lemma C.14](#)) and that these normal forms are convertible to the kinds and types they were computed from ([Lemma 4.3](#)). By validity, context conversion and kind conversion, this means that every declarative subtyping judgment $\Gamma \vdash A \leq B : K$ has an associated judgment $\text{nf}(\Gamma) \vdash \text{nf}(A) \leq \text{nf}(B) : \text{nf}(K)$ relating the normal forms of the original expressions.

In [§5](#) of the paper, we describe a system of canonical rules for deriving such judgments which are defined directly on normal forms – similar to the simple kinding and kind formation judgments introduced in this section. The proof of equivalence of the two systems requires one canonical rule – possibly a derivable or admissible one – for every declarative rule. But some of the declarative rules, such as the subtyping rules [ST- \$\beta_{1,2}\$](#) for β -conversions, or the kinding rule [K-APP](#) for applications, involve substitutions, which do not preserve normal forms. To see why this is a problem, consider the declarative rule [K-APP](#):

$$\frac{\Gamma \vdash A : (X:J) \rightarrow K \quad \Gamma \vdash B : J}{\Gamma \vdash AB : K[B/X]}$$

By soundness of normalization (Lemma 4.3), type equation validity (Lemma 3.1), and context conversion (Corollary B.13), we know that the following is also admissible:

$$\frac{\text{nf}(\Gamma) \vdash U : (X:J') \rightarrow K' \quad \text{nf}(\Gamma) \vdash V : J'}{\text{nf}(\Gamma) \vdash \text{nf}(AB) : \text{nf}(K[B/X])}$$

where $U = \text{nf}(A)$, $V = \text{nf}(B)$, $J' = \text{nf}(J)$ and $K' = \text{nf}(K)$. By Lemma C.14, we know that U and V are simply well-kinded normal types, and that J' and K' are simply well-formed normal kinds. For our canonical application rule, we would like to express the type $\text{nf}(AB)$ and the kind $\text{nf}(K[B/X])$ in the conclusion directly using U, V, J' and K' . This is relatively straightforward for the application $\text{nf}(AB)$ because we know that U must be an operator abstraction $U = \lambda X:L. W$; after all, U has shape $|J| \rightarrow |K|$ and normal forms are η -long. We also know that $|L| \equiv |J| \equiv |J'|$ (see the proof of Lemma C.14 for details). Hence $\text{nf}(AB) \equiv W[V/X^{|J'|}]$ by definition of nf , and we are done.

Things are more complicated for the normal kind $\text{nf}(K[B/X])$. The definition of the normalization function nf does not tell us anything immediately useful about substitutions. Indeed, we know that substitutions do not preserve normal forms, e.g. $(YV)[\lambda X:J'. W/Y]$ is not a normal form, even if YV and $\lambda X:J'. W$ are. However, Corollary 4.2.1 tells us that substitutions in kinds are judgmentally equal to hereditary substitutions, i.e. $\Gamma \vdash K[B/X] = K[B/X^{|J'|}]$, and Lemma C.10.1 tells us that hereditary substitutions preserve normal forms, all of which suggests that $\text{nf}(K[B/X])$ should be equal to $K'[V/X^{|J'|}]$. This is indeed the case; one can show that $\Gamma \vdash \text{nf}(K[B/X]) = K'[V/X^{|J'|}]$. But there is a caveat: the two normal forms are not *syntactically* equal, i.e. $\text{nf}(K[B/X]) \not\equiv K'[V/X^{|J'|}]$. Similarly, $\text{nf}(A[B/X]) \not\equiv \text{nf}(A)[\text{nf}(B)/X^k]$ for types A and B in general.

This fact is best illustrated through the case of type variables, i.e. when $A = X$ and we have $\text{nf}_\Gamma(X[B/X]) \equiv \text{nf}_\Gamma(B)$ and $(\text{nf}_\Gamma(X))[\text{nf}(B)/X^k] \equiv (\eta_{\Gamma(X)}(X))[\text{nf}(B)/X^k]$. We would like to show that $(\eta_K(X))[V/X^k]$ is syntactically equal to V at least when all the involved types and kinds are well-kinded and well-formed, i.e. when $\Gamma \vdash X : K$, $\Gamma \vdash V : K$ and $k = |K|$. But this is not the case. The culprit is a mismatch of kind annotations in operator abstractions, as illustrated by the following counterexample.

Let $J_1 = \top .. \top$ and $J_2 = *$ so that $\Gamma \vdash J_1 \leq J_2$ for any context Γ . Let $U = \top$, $V = \lambda X:J_2. U$ and $K = (X:J_1) \rightarrow *$ so that $\Gamma \vdash V : (X:J_2) \rightarrow *$, $\Gamma \vdash (X:J_2) \rightarrow * \leq K$ and hence $\Gamma \vdash V : K$. Then

$$\begin{aligned} \eta_K(Y) &\equiv \lambda Z:J_1. YZ \\ \eta_K(Y)[V/Y^{|K|}] &\equiv (\lambda Z:J_1. YZ)[V/Y^{|K|}] \\ &\equiv \lambda Z:J_1. (YZ)[V/Y^{|K|}] && \text{(because } Y \notin \text{fv}(J_1)) \\ &\equiv \lambda Z:J_1. V \cdot^{|K|} Z \\ &\equiv \lambda Z:J_1. \lambda X:J_2. U \cdot^{|J_1| \rightarrow *} Z \\ &\equiv \lambda Z:J_1. U[Z/X^{|J_1|}] \\ &\equiv \lambda X:J_1. U && \text{(as } X, Z \notin \text{fv}(U)) \\ &\not\equiv \lambda X:J_2. U \equiv V. && \text{(because } J_1 \not\equiv J_2) \end{aligned}$$

So we are forced to conclude that $\eta_K(Y)[V/Y^{|K|}] \not\equiv V$ in general. The problem, as illustrated by this example, is that the domain annotation J_2 of the type operator abstraction $V = \lambda X:J_2. U$ is not necessarily preserved by the hereditary substitution. It is replaced by the domain J_1 of the declared kind $K = (X:J_1) \rightarrow U$ of Y , which need not be *syntactically* equal to J_1 .

However, we do have $\Gamma \vdash J_1 \leq J_2$ and thus $|J_1| \equiv |J_2|$. The solution, therefore, is to be more lenient when comparing domain annotations in operator abstractions: the *weak* equation $\eta_K(Y)[V/Y^{|K|}] \approx$

V does hold. In fact, it holds for any simply well-formed kind K and simply well-kinded type U , as the following lemma shows.

Lemma C.16.

(1) Let $\gamma, X: j, \delta \vdash K$ kds, $\gamma \vdash U : j$ and $\gamma, X: j, \delta \vdash_{\text{ne}} X \mathbf{V} : |K|$. Then

$$(\eta_K(X \mathbf{V})) [U/X^j] \approx (X \mathbf{V}) [U/X^j].$$

Let $\gamma \vdash K$ kds, then

(2) if $\gamma, X: |K|, \delta \vdash J$ kds, then $J[\eta_K(X)/X^{|K|}] \approx J$;

(3) if $\gamma, X: |K|, \delta \vdash V : j$, then $V[\eta_K(X)/X^{|K|}] \approx V$;

(4) if $\gamma, X: |K|, \delta \vdash j : \mathbf{V} : l$, then $\mathbf{V}[\eta_K(X)/X^{|K|}] \approx \mathbf{V}$;

(5) if $K = (X:K_1) \rightarrow K_2$, $\gamma \vdash_{\text{ne}} N : |K|$ and $\gamma \vdash V : |K_1|$, then $\eta_K(N) \cdot^{|K|} V \approx \eta_{K_2[V/X^{|K_1|}]}(N V)$;

(6) if $\gamma \vdash_{\text{ne}} N : |K|$ and $\gamma \vdash |K| : \mathbf{V} : *$, then $\eta_K(N) \cdot^{|K|} \mathbf{V} \approx N \mathbf{V}$.

Corollary C.17. If $\gamma, X: j, \delta \vdash K$ kds and $\gamma \vdash U : j$, then $(\eta_K(X)) [U/X^j] \approx U$.

It is in this lemma that we see the true usefulness of weak equality. While syntactic equality is too strict for this particular commutativity property, using judgmental type and kind equality would have forced us to formulate its premises in terms of declarative kinding. This would have resulted in a weaker lemma with a more complicated proof. In §D.2.2 we show that weak equations can be converted into judgmental ones provided the related types or kinds are well-kinded or well-formed, respectively. Hence, weak equality affords us a relatively straightforward proof of this lemma (and the next) with a minimal overhead in complexity.

In the proof of Lemma C.16, we employ the following helper lemmas. They are proven separately by three easy inductions: the first on the derivation of $K_1 \approx K_2$, the second on the structure of the kind K , and the third on the derivation of $\gamma \vdash j : \mathbf{V} : k$.

Lemma C.18. Weakly equal kinds have equal shapes. If $K_1 \approx K_2$, then $|K_1| \equiv |K_2|$.

Lemma C.19. Let $X \neq Y$, then $(\eta_K(X \mathbf{D})) [E/Y^j] \equiv \eta_{K[E/Y^j]}(X (\mathbf{D}[E/Y^j]))$ for any K, \mathbf{D}, j and E .

Lemma C.20. Let $\gamma \vdash U : j, \gamma \vdash j : \mathbf{V} : k$ and $\gamma \vdash k : \mathbf{W} : l$, then $U \cdot^j (\mathbf{V}, \mathbf{W}) \equiv (U \cdot^j \mathbf{V}) \cdot^k \mathbf{W}$.

PROOF OF LEMMA C.16. All six parts are proven simultaneously by induction on the structure of K . Parts 2–4 proceed by an inner induction on the simple formation and kinding derivations for J, V and \mathbf{V} , respectively. We show a few key cases, the remainder of the proof is routine.

- *Part 1*, $K = (Y:K_1) \rightarrow K_2$. By inspection of the formation and kinding rules, we must have $\gamma, X: j, \delta \vdash K_1$ kds, $\gamma, X: j, \delta, Y: |K_1| \vdash K_2$ kds and $\gamma, X: j, \delta \vdash j : \mathbf{V} : |K|$. By Lemma C.10 we have

$$\gamma, \delta \vdash (X \mathbf{V}) [U/X^j] : |K_1| \rightarrow |K_2| \quad \text{and} \quad \gamma, \delta \vdash K_1 [U/X^j] \text{ kds.}$$

The final kinding rule used to derive the first of these judgments must be **SK-Abs** since that is the only rule assigning arrow shapes to normal types. Therefore, the following must hold for some J and W :

$$(X \mathbf{V}) [U/X^j] = \lambda Y: J. W \tag{3}$$

$$|K_1| = |J| \tag{4}$$

$$\gamma, \delta, Y: |J| \vdash W : |K_2|. \tag{5}$$

By weakening ([Lemma C.8](#)), [Lemma C.12](#) and [SK-NEAPP](#) we also have

$$\gamma, X:j, \delta, Y:|K_1| \vdash_{\text{ne}} X \mathbf{V}(\eta_{K_1}(Y)) : |K_2|$$

and hence

$$\begin{aligned} & (\eta_{K_2}(X \mathbf{V}(\eta_{K_1}(Y))))[U/X^j] \\ & \approx (X \mathbf{V}(\eta_{K_1}(Y)))[U/X^j] && \text{(by IH for 1)} \\ & \equiv U \cdot^j ((\mathbf{V}, (\eta_{K_1}(Y)))[U/X^j]) && \text{(by definition)} \\ & \equiv (U \cdot^j (\mathbf{V}[U/X^j])) \cdot^{|K_1|} ((\eta_{K_1}(Y)))[U/X^j] && \text{(by Lemmas C.1 and C.20)} \\ & \equiv (U \cdot^j (\mathbf{V}[U/X^j])) \cdot^{|K_1|} (\eta_{K_1}[U/X^j](Y)) && \text{(by Lemma C.19)} \\ & \equiv ((X \mathbf{V})[U/X^j]) \cdot^{|K_1| \mapsto |K_2|} (\eta_{K_1}[U/X^j](Y)) && \text{(by definition)} \\ & \equiv (\lambda Y:J. W) \cdot^{|K_1| \mapsto |K_2|} (\eta_{K_1}[U/X^j](Y)) && \text{(by (3))} \\ & \equiv W[(\eta_{K_1}[U/X^j](Y))/Y^{|K_1|}] && \text{(by definition)} \\ & \approx W && \text{(by Lemma C.18, (5) and IH for 3)} \end{aligned}$$

We conclude that

$$\begin{aligned} & (\eta_K(X \mathbf{V}))[U/X^j] \\ & \equiv \lambda Y:K_1[U/X^j]. (\eta_{K_2}(X \mathbf{V}(\eta_{K_1}(Y))))[U/X^j] && \text{(by definition)} \\ & \approx \lambda Y:J. W && \text{(by Lemma C.18, (4), the above and WEQ-Abs)} \\ & \equiv (X \mathbf{V})[U/X^j]. && \text{(by (3))} \end{aligned}$$

- *Part 3, case SK-NE.* The rule [SK-NE](#) has only one premise which must have been derived using [SK-VARAPP](#), so $V = N = Y \mathbf{V}$ and we have $\gamma, X:|K|, \delta \vdash j : \mathbf{V} : *$ with $(\gamma, X:|K|, \delta)(Y) = j$. We distinguish two cases: $Y = X$ and $Y \neq X$ but consider only the first case here; the second case is simpler. Since $Y = X$, we have $j = |K|$, and

$$\begin{aligned} & V[\eta_K(X)/X^{|K|}] \\ & \equiv \eta_K(X) \cdot^{|K|} (\mathbf{V}[\eta_K(X)/X^{|K|}]) && \text{(by definition)} \\ & \approx \eta_K(X) \cdot^{|K|} \mathbf{V} && \text{(by reflexivity of } \approx, \text{ IH for 4 and Lemma C.3.5)} \\ & \approx X \mathbf{V}. && \text{(by SK-VARAPP and the IH for 6)} \end{aligned}$$

- *Part 5, $K = (X:K_1) \rightarrow K_2$.* By inspection of the formation and kinding rules, we must have $\gamma \vdash K_1$ kds and $N = Y \mathbf{U}$ with $\gamma \vdash \gamma(Y) : \mathbf{U} : |K|$.

$$\begin{aligned} \eta_K(N) \cdot^{|K|} V & \equiv \lambda X:K_1. \eta_{K_2}(Y \mathbf{U}(\eta_{K_1}(X))) \cdot^{|K|} V && \text{(by definition)} \\ & \equiv \eta_{K_2}(Y \mathbf{U}(\eta_{K_1}(X)))[V/X^{|K_1|}] && \text{(by definition)} \\ & \equiv \eta_{K_2[V/X^{|K_1|}]}(Y((\mathbf{U}, (\eta_{K_1}(X)))[V/X^{|K_1|}])) && \text{(by Lemma C.19)} \\ & \equiv \eta_{K_2[V/X^{|K_1|}]}(Y \mathbf{U}((\eta_{K_1}(X)))[V/X^{|K_1|}])) && \text{(as } X \notin \text{fv}(\mathbf{U})) \\ & \approx \eta_{K_2[V/X^{|K_1|}]}(Y \mathbf{U} V). && \text{(by IH for 1 and Lemma C.5)} \end{aligned}$$

The use of the IH in the last step corresponds to [Corollary C.17](#). \square

With [Lemma C.16](#) in place, we are ready to prove that normalization weakly commutes with substitution. In the following, $\Gamma \approx \Delta$ denotes the pointwise lifting of weak equality to contexts.

Lemma C.21. *Substitution weakly commutes with normalization of well-formed kinds and well-kinded types. Let $\Gamma \vdash A : J$ and $V = \text{nf}_{\text{nf}(\Gamma)}(A)$, then*

- (1) *if $\Gamma, X : J, \Delta \vdash K \text{ kd}$, then $\text{nf}_{\text{nf}(\Gamma, \Delta[A/X])}(K[A/X]) \approx (\text{nf}_{\text{nf}(\Gamma, X : J, \Delta)}(K))[V/X]^{|J|}$;*
- (2) *if $\Gamma, X : J, \Delta \vdash B : K$, then $\text{nf}_{\text{nf}(\Gamma, \Delta[A/X])}(B[A/X]) \approx (\text{nf}_{\text{nf}(\Gamma, X : J, \Delta)}(B))[V/X]^{|J|}$;*

PROOF. Simultaneously by induction on declarative kind formation and kinding derivations. In the case for **K-VAR** where $B = Y$, we use **Corollary C.17** if $Y = X$. Otherwise, let $\Sigma = \Gamma, X : J, \Delta$ and $\Sigma' = \Gamma, \Delta[A/X]$. By stability of simplified contexts under normalization and hereditary substitution, and by commutation of simplification and context lookup, we have

$$\begin{aligned} |\text{nf}(\Sigma')(Y)| &\equiv |\text{nf}(\Sigma')|(Y) \equiv |\Sigma'| (Y) \\ &\equiv (|\Gamma|, |\Delta[A/X]|)(Y) \equiv (|\Gamma|, |\Delta|)(Y) \equiv (|\Gamma|, X : |J|, |\Delta|)(Y) \equiv |\Sigma|(Y) \\ &\equiv |\text{nf}(\Sigma)|(Y) \equiv |\text{nf}(\Sigma)(Y)| \equiv |\text{nf}_{\Sigma}(\Sigma(Y))| \equiv |(\text{nf}_{\Sigma}(\Sigma(Y)))[V/X]^{|J|}| \end{aligned}$$

Hence, by **Lemma C.5** and **Lemma C.19** we have

$$\eta_{\text{nf}(\Sigma')(Y)}(Y) \approx \eta_{(\text{nf}_{\Sigma}(\Sigma(Y)))[V/X]^{|J|}}(Y) \equiv (\eta_{\text{nf}_{\Sigma}(\Sigma(Y))}(Y))[V/X]^{|J|}.$$

In the case for **K-APP**, we use **Lemma C.3.2** and **Lemma C.11.2**. \square

The very last lemma of this section will be used in our equivalence proof in **§D.3** to show that subtyping rules for η -conversion of normal operators are admissible in canonical kinding.

Lemma C.22. *If $\Gamma \vdash A : (X : J) \rightarrow K$ with $X \notin \text{fv}(A)$, then $\text{nf}_{\text{nf}(\Gamma)}(\lambda X : J. A X) \approx \text{nf}_{\text{nf}(\Gamma)}(A)$.*

PROOF. The proof uses **Lemma C.14** to obtain $|\text{nf}(\Gamma)| \vdash \text{nf}_{\text{nf}(\Gamma)}(A) : |J| \rightarrow |K|$ and proceeds by case analysis on the final rule used to derive this simple kinding judgment; the only applicable rule is **SK-Abs**. The remainder of the proof uses equational reasoning very similar to that used in the proof of **Lemma C.16.1**. \square

D PROPERTIES OF THE CANONICAL SYSTEM

This section states and proves some basic yet important metatheoretic properties of the canonical system that have been omitted from **§5** of the paper. As a first step, we prove soundness of the canonical system in **§D.1**. Next, we state and prove a *hereditary substitution lemma* in **§D.2**, which establishes that canonical judgments are preserved by hereditary substitutions. This lemma is a key ingredient in proving completeness of the canonical system. We prove completeness in **§D.3**, after showing that canonical subtyping can be inverted at the top-level in **§D.4**. **Fig. 10** shows the canonical judgments that have been omitted from the paper: context and kind formation, subkinding, and kind and type equality.

The contents of this section are based on Chapter 5 of the first author's PhD dissertation. We refer the interested reader to the dissertation for the full details [**Stucki 2017**, Ch. 5].

D.1 Soundness and Basic Properties

Before we establish any other metatheoretic properties of the canonical system, let us prove its soundness with respect to the declarative presentation. To avoid confusion, we mark canonical judgments with the subscript “c” and declarative ones with “d” in the following lemma.

Lemma D.1 (soundness of the canonical rules).

- (1) *If $\Gamma \text{ ctx}_c$, then $\Gamma \text{ ctx}_d$.*
- (2) *If $\Gamma \vdash_c K \text{ kd}$, then $\Gamma \vdash_d K \text{ kd}$.*
- (3) *If $\Gamma \vdash_c J \leq K$, then $\Gamma \vdash_d J \leq K$.*
- (4) *If $\Gamma \vdash_c J = K$, then $\Gamma \vdash_d J = K$.*

Context formation $\boxed{\Gamma \text{ ctx}}$

$$\frac{}{\emptyset \text{ ctx}} \text{ (CC-EMPTY)} \quad \frac{\Gamma \text{ ctx} \quad \Gamma \vdash K \text{ kd}}{\Gamma, X:K \text{ ctx}} \text{ (CC-TMBIND)} \quad \frac{\Gamma \text{ ctx} \quad \Gamma \vdash V \Rightarrow V .. V}{\Gamma, x:V \text{ ctx}} \text{ (CC-TPBIND)}$$

Kind formation $\boxed{\Gamma \vdash K \text{ kd}}$

$$\frac{\Gamma \vdash U \Rightarrow U .. U \quad \Gamma \vdash V \Rightarrow V .. V}{\Gamma \vdash U .. V \text{ kd}} \text{ (CWF-INTV)} \quad \frac{\Gamma \vdash J \text{ kd} \quad \Gamma, X:J \vdash K \text{ kd}}{\Gamma \vdash (X:J) \rightarrow K \text{ kd}} \text{ (CWF-DARR)}$$

Subkinding $\boxed{\Gamma \vdash J \leq K}$

$$\frac{\Gamma \vdash U_2 \leq U_1 \quad \Gamma \vdash V_1 \leq V_2}{\Gamma \vdash U_1 .. V_1 \leq U_2 .. V_2} \text{ (CSK-INTV)} \quad \frac{\Gamma \vdash (X:J_1) \rightarrow K_1 \text{ kd} \quad \Gamma \vdash J_2 \leq J_1 \quad \Gamma, X:J_2 \vdash K_1 \leq K_2}{\Gamma \vdash (X:J_1) \rightarrow K_1 \leq (X:J_2) \rightarrow K_2} \text{ (CSK-DARR)}$$

Kind equality $\boxed{\Gamma \vdash J = K}$ **Type equality** $\boxed{\Gamma \vdash U = V \Leftarrow K}$

$$\frac{\Gamma \vdash J \text{ kd} \quad \Gamma \vdash K \text{ kd} \quad \Gamma \vdash J \leq K \quad \Gamma \vdash K \leq J}{\Gamma \vdash J = K} \text{ (CSK-ANTISYM)}$$

$$\frac{\Gamma \vdash K \text{ kd} \quad \Gamma \vdash U \leq V \Leftarrow K \quad \Gamma \vdash V \leq U \Leftarrow K}{\Gamma \vdash U = V \Leftarrow K} \text{ (CST-ANTISYM)}$$

Fig. 10. Canonical presentation of F^{ω} – supplement

- (5) If $\Gamma \vdash_{\text{var}} X : K$, then $\Gamma \vdash_{\text{d}} X : K$.
- (6) If $\Gamma \vdash_{\text{ne}} N : K$, then $\Gamma \vdash_{\text{d}} N : K$.
- (7) If $\Gamma \vdash_{\text{c}} V \Rightarrow K$, then $\Gamma \vdash_{\text{d}} V : K$.
- (8) If $\Gamma \vdash_{\text{c}} V \Leftarrow K$, then $\Gamma \vdash_{\text{d}} V : K$.
- (9) If $\Gamma \vdash_{\text{c}} U \leq V$, then $\Gamma \vdash_{\text{d}} U \leq V : *$.
- (10) If $\Gamma \vdash_{\text{c}} U \leq V : K$, then $\Gamma \vdash_{\text{d}} U \leq V : K$.
- (11) If $\Gamma \vdash_{\text{c}} U = V : K$, then $\Gamma \vdash_{\text{d}} U \leq V : K$.
- (12) If $\Gamma \vdash_{\text{d}} A : J$ and $\Gamma \vdash_{\text{c}} J \Rightarrow \mathbf{V} \Rightarrow K$, then $\Gamma \vdash_{\text{d}} A \mathbf{V} : K$.
- (13) If $\Gamma \vdash_{\text{d}} A \leq B : J$ and $\Gamma \vdash_{\text{c}} J \Rightarrow \mathbf{U} = \mathbf{V} \Rightarrow K$, then $\Gamma \vdash_{\text{d}} A \mathbf{U} \leq B \mathbf{V} : K$.

The proof uses the following two helper lemmas. The first is a derived rule, the second is proven by easy case analysis of the rules for kind synthesis.

Lemma D.2. *If $\Gamma \vdash A : B .. C$, then also $\Gamma \vdash A : *$.*

Lemma D.3. *If $\Gamma \vdash U \Rightarrow V .. W$, then $V \equiv U$ and $W \equiv U$.*

PROOF OF LEMMA D.1. By induction on derivations of the various canonical judgments; for parts 1–11, on the derivation of the first premise, for parts 12 and 13, on that of the second premise. Most cases are straightforward. In cases involving synthesized kinding of proper types, we use **K-SING** and **Lemmas D.2, B.15** and **D.3** to adjust kinds where necessary. In cases involving hereditary substitutions, i.e. those for **CK-CONS** and **SP-EQ-CONS**, we use soundness of hereditary substitution in kinds (**Corollary 4.2.1**). \square

As for the declarative system, the contexts of most canonical judgments are well-formed. There are two exceptions: kinding and equality of spines. The rules **CK-EMPTY** and **SpEQ-EMPTY** for empty spines offer no guarantee that the enclosing context is well-formed. This is not a problem in practice, since well-kinded spines only appear in judgments about neutral types, the heads of which must be kinded in a well-formed context.

Lemma D.4 (context validity). *Assume $\Gamma \vdash \mathcal{J}$ for any canonical judgment except spine kinding or equality. Then Γ ctx.*

PROOF. By simultaneous induction on the derivations of the various judgments. \square

We can prove a few more validity properties at this point. First, since synthesized kinds are singletons, validity of synthesized kinding judgments follows by **CWF-INTV** for proper types and by an easy induction for type operators. Second, validity of kind equality as well as the checked subtyping and type equality judgments follows immediately from the validity conditions included in the rules **CSK-ANTI-SYM**, **CST-INTV**, **CST-ABS** and **CST-ANTI-SYM**.

Lemma D.5 (canonical validity – part 1).

(synthesized kinding validity) *If $\Gamma \vdash V \Rightarrow K$, then $\Gamma \vdash K$ kd.*

(checked subtyping validity) *If $\Gamma \vdash U \leq V \Leftarrow K$, then $\Gamma \vdash U \Leftarrow K$ and $\Gamma \vdash V \Leftarrow K$.*

(kind equation validity) *If $\Gamma \vdash J = K$, then $\Gamma \vdash J$ kd and $\Gamma \vdash K$ kd.*

(type equation validity) *If $\Gamma \vdash U = V \Leftarrow K$, then $\Gamma \vdash U \Leftarrow K$, $\Gamma \vdash V \Leftarrow K$ and $\Gamma \vdash K$ kd.*

We prove the remaining validity properties of the canonical system in §D.2.1, once we have shown that hereditary substitutions preserve well-formedness of kinds.

Before we can do so, we need to establish the usual weakening and context narrowing lemmas for canonical typing.

Lemma D.6 (weakening). *Assume $\Gamma, \Delta \vdash \mathcal{J}$ for any of the canonical judgments.*

(1) *If $\Gamma \vdash A \Rightarrow A..A$, then $\Gamma, x:A, \Delta \vdash \mathcal{J}$.*

(2) *If $\Gamma \vdash K$ kd, then $\Gamma, X:K, \Delta \vdash \mathcal{J}$.*

Corollary D.7 (iterated weakening). *If Γ, Δ ctx and $\Gamma \vdash \mathcal{J}$, then $\Gamma, \Delta \vdash \mathcal{J}$.*

Lemma D.8 (context narrowing – weak version).

(1) *If $\Gamma \vdash A \Rightarrow A..A$, $\Gamma \vdash A \leq B : *$ and $\Gamma, x:B, \Delta \vdash \mathcal{J}$, then $\Gamma, x:A, \Delta \vdash \mathcal{J}$.*

(2) *If $\Gamma \vdash J$ kd, $\Gamma \vdash J \leq K$ and $\Gamma, X:K, \Delta \vdash \mathcal{J}$, then $\Gamma, X:J, \Delta \vdash \mathcal{J}$.*

The proofs of both lemmas are routine inductions on the derivations of the respective judgments. The proof of context narrowing is only easy thanks to the rule **CV-SUB**. Without this rule, the canonical kinding judgments for variables and neutral types would become synthesis judgments and context narrowing would no longer hold in its present form for these two judgments. To see this, consider the variable kinding judgment $\Gamma, X:K, \Delta \vdash_{\text{var}} X : K$, which, after eliminating **CV-SUB**, could only be derived using **CV-VAR**. If we were to narrow the context by changing the declared kind K of X to some J such that $\Gamma \vdash J \leq K$, then the synthesized kind of X would necessarily change to J too.

For neutral kinding, we would be in a similar situation as the new synthesized kind of the head would have to be propagated through the kinding derivation of the spine. Along the way, the new kind J would have to be unraveled by repeatedly separating $J = (X:J_1) \rightarrow J_2$ into J_1 , J_2 and hereditarily substituting the next element of the spine into the codomain J_2 . To prove context narrowing for the remainder of the canonical judgments, we would have to maintain the

invariant that the new synthesized kind of a neutral type is a subtype of the original one, i.e. if $\Gamma, X:K_1, \Delta \vdash_{\text{var}} N \Rightarrow K_2$ and $\Gamma \vdash J_1 \leq K_1$ then $\Gamma, X:J_1, \Delta \vdash_{\text{var}} N \Rightarrow J_2$ and $\Gamma \vdash J_2 \leq K_2$. Because spine kinding involves hereditary substitution, a proof involving this invariant would require a hereditary substitution lemma for subtyping, i.e. a proof that hereditary substitutions preserve subtyping. We give such a proof in §D.2 and, as we will see there, it makes crucial use of context narrowing itself. By allowing us to establish Lemma D.8 independently, the rule CV-SUB thus simplifies the proof of an otherwise rather complicated lemma (see Lemma D.17 for details).

D.1.1 Order-Theoretic Properties. Having established context narrowing, we can prove the usual order-theoretic properties of canonical subkinding, subtyping, as well as kind and type equality. We start by stating and proving the various reflexivity properties which, unlike those for the declarative relations, have to be proven simultaneously for the canonical variants.

Lemma D.9. *The following reflexivity rules are all admissible.*

$$\begin{array}{c}
\frac{\Gamma \vdash K \text{ kd}}{\Gamma \vdash K \leq K} \quad (\text{CSK-REFL}) \qquad \frac{\Gamma \vdash K \text{ kd}}{\Gamma \vdash K = K} \quad (\text{CKEQ-REFL}) \\
\\
\frac{\Gamma \vdash U \Rightarrow V \dots W}{\Gamma \vdash U \leq U} \quad (\text{CST-REFLSYN}) \qquad \frac{\Gamma \vdash V \Leftarrow K \quad \Gamma \vdash K \text{ kd}}{\Gamma \vdash V \leq V \Leftarrow K} \quad (\text{CST-REFLCK}) \\
\\
\frac{\Gamma \vdash V \Rightarrow J \quad \Gamma \vdash J \leq K \quad \Gamma \vdash K \text{ kd}}{\Gamma \vdash V \leq V \Leftarrow K} \quad (\text{CST-REFLSUB}) \\
\\
\frac{\Gamma \vdash J \Rightarrow \mathbf{V} \Rightarrow K}{\Gamma \vdash J \Rightarrow \mathbf{V} = \mathbf{V} \Rightarrow K} \quad (\text{SPEQ-REFL}) \qquad \frac{\Gamma \vdash V \Leftarrow K \quad \Gamma \vdash K \text{ kd}}{\Gamma \vdash V = V \Leftarrow K} \quad (\text{CTEQ-REFL})
\end{array}$$

PROOF. The proof is by mutual induction in the structure of the kinds and types being related to themselves, and then by case-analysis on the final rules of the corresponding kind formation and kinding derivations. The proof for CST-REFLSUB proceeds by an inner induction on the derivation of $\Gamma \vdash J \leq K$. In the case for CSK-DARR where $V = \lambda X:J_1.U$ and we have $\Gamma \vdash K_1 \leq J_1$, $\Gamma, X:K_1 \vdash J_2 \leq K_2$ and $\Gamma, X:J_1 \vdash U \Rightarrow J_2$, we use context narrowing to adjust the declared kind of X from J_1 to K_1 in $\Gamma, X:J_1 \vdash U \Rightarrow J_2$ before applying the IH to obtain $\Gamma, X:K_1 \vdash U \leq U \Leftarrow K_2$. \square

Transitivity of the various relations and symmetry of the equalities are more easily established, thanks to the rule CST-TRANS on the one hand, and to the structure of equality on the other.

Lemma D.10. *Canonical subkinding, subtyping, kind and type equality are transitive.*

$$\begin{array}{c}
\frac{\Gamma \vdash J \leq K \quad \Gamma \vdash K \leq L}{\Gamma \vdash J \leq L} \quad (\text{CSK-TRANS}) \qquad \frac{\Gamma \vdash J = K \quad \Gamma \vdash K = L}{\Gamma \vdash J = L} \quad (\text{CKEQ-TRANS}) \\
\\
\frac{\Gamma \vdash U \leq V \Leftarrow K \quad \Gamma \vdash V \leq W \Leftarrow K}{\Gamma \vdash U \leq W \Leftarrow K} \quad (\text{CST-TRANSCk}) \qquad \frac{\Gamma \vdash U = V \Leftarrow K \quad \Gamma \vdash V = W \Leftarrow K}{\Gamma \vdash U = W \Leftarrow K} \quad (\text{CTEQ-TRANS})
\end{array}$$

PROOF. The proof of **CSK-TRANS** is by induction on the structure of K and case analysis on the final rules used to derive the premises. In the case for $K = (X:K_1) \rightarrow K_2$ we use context narrowing. The proof of **CST-TRANS** is by induction on the derivation of the first premise. The proofs of **CKEQ-TRANS** and **CTEQ-TRANS** are by inspection of the equality rules and use **CSK-TRANS** and **CST-TRANS**, respectively. \square

Lemma D.11. *Canonical kind and type equality are symmetric.*

$$\frac{\Gamma \vdash J = K}{\Gamma \vdash K = J} \quad (\text{CKEQ-SYM}) \qquad \frac{\Gamma \vdash U = V \Leftarrow K}{\Gamma \vdash V = U \Leftarrow K} \quad (\text{CTEQ-SYM})$$

PROOF. By inspection of the equality rules, **CSK-ANTI-SYM** and **CST-ANTI-SYM**. \square

Thanks to context narrowing and subkinding transitivity, we can prove admissibility of the following subsumption rules for the three checked judgments.

Lemma D.12. *Kind subsumption is admissible in the checked judgments.*

$$\frac{\Gamma \vdash U \Leftarrow J \quad \Gamma \vdash J \leq K}{\Gamma \vdash U \Leftarrow K} \quad (\text{CK-SUBCK}) \qquad \frac{\Gamma \vdash U \leq V \Leftarrow J \quad \Gamma \vdash J \leq K}{\Gamma \vdash U \leq V \Leftarrow K} \quad (\text{CST-SUBCK})$$

$$\frac{\Gamma \vdash U = V \Leftarrow J \quad \Gamma \vdash J \leq K}{\Gamma \vdash U = V \Leftarrow K} \quad (\text{CTEQ-SUBCK})$$

PROOF. Admissibility is proven separately for the three rules, in the order they are listed. The proof of **CK-SUBCK** is by inspection of the kind checking rules and uses **CSK-TRANS**. The proof of **CST-SUBCK** is by induction on the derivation of the second premise $\Gamma \vdash J \leq K$ and uses **CK-SUBCK** as well as context narrowing for the inductive step. The proof of **CTEQ-SUBCK** is by inspection of the equality rules and uses **CST-SUBCK**. \square

Kind subsumption subsumes kind conversion thanks to the first of the following three rules, all of which follow immediately by inspection of the equality and checked subtyping rules.

$$\frac{\Gamma \vdash J = K}{\Gamma \vdash J \leq K} \quad (\text{CSK-REFL-KEQ}) \qquad \frac{\Gamma \vdash U = V \Leftarrow K}{\Gamma \vdash U \leq V \Leftarrow K} \quad (\text{CST-REFL-TEQ})$$

$$\frac{\Gamma \vdash U = V \Leftarrow W \dots W'}{\Gamma \vdash U \leq V} \quad (\text{CST-REFL-TEQ}')$$

D.1.2 Canonical Replacements for Declarative Rules. Our completeness proof for the canonical system relies on the fact that the normal form $\text{nf}(A)$ of any declaratively well-kinded type $\Gamma \vdash A : K$ kind checks against the normal form $\text{nf}(K)$ of the corresponding kind K , i.e. that we have $\text{nf}(\Gamma) \vdash \text{nf}(A) \Leftarrow \text{nf}(K)$. To simplify the proof of this fact, we establish a set of admissible kind checking rules below that mirror the corresponding declarative rules.

We begin by proving that normal forms with synthesized or checked interval kinds also check against $*$.

Lemma D.13. *Types inhabiting interval kinds are proper types. If $\Gamma \vdash U \Rightarrow V \dots W$ or $\Gamma \vdash U \Leftarrow V \dots W$, then also $\Gamma \vdash U \Leftarrow *$.*

PROOF. If $\Gamma \vdash U \Rightarrow V \dots W$, then by **Lemma D.3**, $\Gamma \vdash U \Rightarrow U \dots U$. From this we derive the result by **CST-BOT**, **CST-TOP**, **CSK-INTV** and **CK-SUB**. If $\Gamma \vdash U \Leftarrow V \dots W$, then this must have been derived using **CK-SUB** and hence $\Gamma \vdash U \Rightarrow K$ and $\Gamma \vdash K \leq V \dots W$. By inspection of the subkinding rules, $K = V' \dots W'$ for some V' and W' . The result then follows by the first part of the lemma. \square

The second part of the proof follows a pattern that is quite typical for proofs in the remainder of this section. Thanks to the division of kinding into kind synthesis and checking, and thanks to the simplicity of both kind checking and subkinding derivations, we can often prove properties of kind checking judgments $\Gamma \vdash U \Leftarrow K$ by appealing to similar properties of kind synthesis judgments $\Gamma \vdash U \Rightarrow K'$ where K and K' have the same shape, i.e. where K and K' are either both intervals or both arrows. Two more examples of this pattern appear in the following lemma, where the admissibility proofs of the rules **CST-CkBND₁** and **CST-CkBND₂**, which have kind checking judgments as their premises, appeal to instances of **CST-SYNBND₁** and **CST-SYNBND₂**, respectively, which have similar kind synthesis judgments as their premises.

Lemma D.14. *All of the following are admissible.*

$$\begin{array}{c}
\frac{\Gamma \vdash U \Leftarrow V .. W}{\Gamma \vdash U \Rightarrow U .. U} \text{ (CK-SING')} \quad \frac{\Gamma \vdash V \Rightarrow K}{\Gamma \vdash V \Leftarrow K} \text{ (CK-SYNCK)} \quad \frac{\Gamma \vdash_{ne} N : U .. V}{\Gamma \vdash N \Leftarrow U .. V} \text{ (CK-NECK)} \\
\\
\frac{\Gamma \vdash U \Leftarrow * \quad \Gamma \vdash V \Leftarrow *}{\Gamma \vdash U \rightarrow V \Leftarrow *} \text{ (CK-ARR')} \quad \frac{\Gamma \vdash K \text{ kd} \quad \Gamma, X:K \vdash V \Leftarrow *}{\Gamma \vdash \forall X:K. V \Leftarrow *} \text{ (CK-ALL')} \\
\\
\frac{\Gamma \vdash J \text{ kd} \quad \Gamma, X:J \vdash V \Leftarrow K}{\Gamma \vdash \lambda X:J. V \Leftarrow (X:J) \rightarrow K} \text{ (CK-ABS')} \quad \frac{\Gamma \vdash V_1 \leq V_2 \Leftarrow U .. W}{\Gamma \vdash V_1 \leq V_2 \Leftarrow V_1 .. V_2} \text{ (CST-INTV')} \\
\\
\frac{\Gamma \vdash U \Rightarrow V_1 .. V_2}{\Gamma \vdash V_1 \leq U} \text{ (CST-SYNBND}_1\text{)} \quad \frac{\Gamma \vdash U \Rightarrow V_1 .. V_2}{\Gamma \vdash U \leq V_2} \text{ (CST-SYNBND}_2\text{)} \\
\\
\frac{\Gamma \vdash U \Leftarrow V_1 .. V_2}{\Gamma \vdash V_1 \leq U} \text{ (CST-CkBND}_1\text{)} \quad \frac{\Gamma \vdash U \Leftarrow V_1 .. V_2}{\Gamma \vdash U \leq V_2} \text{ (CST-CkBND}_2\text{)} \\
\\
\frac{\Gamma \vdash J \Rightarrow U \Rightarrow (X:K) \rightarrow L \quad \Gamma \vdash V \Leftarrow K \quad \Gamma \vdash K \text{ kd}}{\Gamma \vdash J \Rightarrow U, V \Rightarrow L[V/X^{|K|}]} \text{ (CK-SNOC)} \\
\\
\frac{\Gamma \vdash J \Rightarrow U_1 = U_2 \Rightarrow (X:K) \rightarrow L \quad \Gamma \vdash V_1 = V_2 \Leftarrow K}{\Gamma \vdash J \Rightarrow U_1, V_1 = U_2, V_2 \Rightarrow L[V_1/X^{|K|}]} \text{ (SPEQ-SNOC)}
\end{array}$$

PROOF. Some of the rules are derivable others are admissible; most of the proofs are straightforward, so we omit the details. The proofs of the alternate type formation rules use **CK-SING'** and **Lemma D.13**. The proofs of the alternate projection rules **CST-SYNBND₁** and **CST-SYNBND₂** use **Lemma D.3** and reflexivity; those of **CST-CkBND₁** and **CST-CkBND₂** are by inspection of kind checking and subkinding and use **CST-SYNBND₁** and **CST-SYNBND₂**, respectively. The proofs of the last two rules are by induction on the derivations of the respective first premises. \square

As in the declarative system, we define *canonical context equality* $\Gamma = \Delta \text{ ctx}$ as the pointwise lifting of canonical type and kind equality to context bindings:

$$\frac{}{\emptyset = \emptyset \text{ ctx}} \quad \frac{\Gamma = \Delta \text{ ctx} \quad \Gamma \vdash J = K}{\Gamma, X:J = \Delta, X:K \text{ ctx}} \quad \frac{\Gamma = \Delta \text{ ctx} \quad \Gamma \vdash U = V \Leftarrow W .. W'}{\Gamma, x:U = \Delta, x:V \text{ ctx}}$$

D.1.3 Simplification of Canonical Kinding. In §C.2.2 of the previous section, we showed that every well-formed kind and well-kinded type has a simply well-formed or simply well-kinded normal form, respectively (see Lemma C.14). Canonically well-formed kinds and canonically well-kinded types are already in normal form, but we can still simplify their kind formation and kinding derivations, as the following pair of lemma shows. In the statement of the second lemma, we use the subscript “nes” to mark simple kinding judgments for neutral types and “ne” to mark their canonical counterparts.

Lemma D.15. *Canonical subkinds and equal kinds simplify equally. If $\Gamma \vdash J \leq K$ or $\Gamma \vdash J = K$ then $|J| \equiv |K|$.*

PROOF. Separately, by induction on subkinding and kind equality derivations, respectively. \square

Lemma D.16 (simplification). *Well-formed kinds and well-kinded normal forms, neutrals and spines are also simply well-formed and well-kinded, respectively.*

- (1) *If $\Gamma \vdash_{\text{var}} X : K$, then $|\Gamma| \vdash_{\text{nes}} X : |K|$.*
- (2) *If $\Gamma \vdash K \text{ kd}$, then $|\Gamma| \vdash K \text{ kds}$.*
- (3) *If $\Gamma \vdash_{\text{ne}} N : K$, then $|\Gamma| \vdash_{\text{nes}} N : |K|$.*
- (4) *If $\Gamma \vdash V \Rightarrow K$, then $|\Gamma| \vdash V : |K|$.*
- (5) *If $\Gamma \vdash V \Leftarrow K$, then $|\Gamma| \vdash V : |K|$.*
- (6) *If $\Gamma \vdash J \Rightarrow \mathbf{V} \Rightarrow K$, then $|\Gamma| \vdash |J| : \mathbf{V} : |K|$.*

PROOF. Part 1 is proven separately, parts 2–6 are proven simultaneously, all by induction on the derivations of the respective premises. The cases of CV-SUB and CK-SUB use of Lemma D.15. \square

Thanks to Lemma D.16, properties of simply kinded normal forms still hold for canonically kinded normal forms. For example, by Lemmas D.16 and C.11, hereditary substitutions in canonically kinded types commute.

D.2 The Hereditary Substitution Lemma

We have arrived at the core of the technical development of this section: the proof of the *hereditary substitution lemma*. The hereditary substitution lemma states, roughly, that canonical judgments are preserved by hereditary substitutions of canonically well-kinded types. Just as the ordinary substitution lemma for the declarative system (Lemma B.4) played a key role in the proofs of several metatheoretic properties in §3 of the paper, the hereditary substitution lemma is key to proving important metatheoretic properties of the canonical system. But unlike that of its ordinary counterpart, the proof of the hereditary substitution lemma is rather challenging. This is reflected already in the statement of the lemma, which features 24 separate parts, all of which have to be proven simultaneously (see Lemma D.17 below).

One reason for the large number of parts is simply that there are more judgment forms in the canonical system than there are in the declarative system. But the foremost reason is that the proof of the hereditary substitution lemma circularly depends on *functionality of the canonical judgments*, i.e. on the fact that hereditarily substituting canonically equal types in normal forms yields canonically equal normal forms. This also renders the proof more challenging since both properties have to be established at the same time.

The main source of complexity is the subtyping rule CST-NE. It is because of this rule that we have to prove the hereditary substitution and functionality lemmas simultaneously.

To illustrate this, consider the neutral types $N = X \mathbf{V}$ and $M = X \mathbf{W}$ with $\mathbf{V} = V_1, V_2$, $\mathbf{W} = W_1, W_2$ such that $X \notin \text{fv}(\mathbf{V}) \cup \text{fv}(\mathbf{W})$, and assume some Γ, Δ and U such that

$$\Gamma, \Delta \vdash \lambda Y_1 : J_1. \lambda Y_2 : J_2. U \Leftarrow J \quad \text{and} \quad \Gamma, X : J, \Delta \vdash J \Rightarrow \mathbf{V} = \mathbf{W} \Rightarrow *$$

for $J = (Y_1:J_1) \rightarrow (Y_2:J_2) \rightarrow J_3$. Then, by **CST-NE**, we have $\Gamma, X:J, \Delta \vdash N \leq M$.

We would like to hereditarily substitute $U' = \lambda Y_1:J_1. \lambda Y_2:J_2. U$ for X in N and M and show that the resulting types remain subtypes, i.e. that

$$\Gamma, \Delta[U'/X^{|J|}] \vdash N[U'/X^{|J|}] \leq M[U'/X^{|J|}].$$

By the definition of hereditary substitution, we have

$$\begin{aligned} N[U'/X^{|J|}] &\equiv (X \mathbf{V})[U'/X^{|J|}] \equiv U' \cdot^{|J|} (\mathbf{V}[U'/X^{|J|}]) \\ &\equiv U' \cdot^{|J|} V_1 \cdot^{|J_2| \rightarrow |J_3|} V_2 \\ &\equiv ((\lambda Y_2:J_2. U)[V_1/Y_1^{|J_1|}]) \cdot^{|J_2| \rightarrow |J_3|} V_2 \\ &\equiv (\lambda Y_2:J_2[V_1/Y_1^{|J_1|}]. U[V_1/Y_1^{|J_1|}]) \cdot^{|J_2| \rightarrow |J_3|} V_2 \\ &\equiv V'[V_2/Y_2^{|J_2|}] \end{aligned}$$

where $V' = U[V_1/Y_1^{|J_1|}]$. Similarly, $M[U'/X^{|J|}] \equiv W'[W_2/Y_2^{|J_2|}]$ for $W' = U[W_1/Y_1^{|J_1|}]$. Hence, we need to show that

$$\begin{aligned} \Gamma, \Delta[U'/X^{|J|}] \vdash U[V_1/Y_1^{|J_1|}] &\leq U[W_1/Y_1^{|J_1|}] \quad \text{and} \\ \Gamma, \Delta[U'/X^{|J|}] \vdash V'[V_2/Y_2^{|J_2|}] &\leq W'[W_2/Y_2^{|J_2|}]. \end{aligned}$$

Since they belong to equal spines, V_1 and W_1 are judgmentally equal as types, and so are V_2 and W_2 . But in general, neither of these pairs of types are syntactically equal, i.e. $V_1 \not\equiv W_1$, $V_2 \not\equiv W_2$ and $V' \not\equiv W'$. To establish the above inequations, we therefore need to show that simultaneous hereditary substitutions of judgmentally equal types preserve inequations.

The example illustrates a second point, namely that, in order to prove that hereditary substitutions preserve canonical kinding and subtyping, we need to prove that kinding and subtyping of reducing applications is admissible. Our hereditary substitution lemma must cover all of these properties, leading to the aforementioned grand total of 24 parts.

Lemma D.17 (hereditary substitution). *Hereditary substitutions of canonically kind-checked types preserve the canonical judgments; substitutions of canonically equal types in canonically well-formed and well-kinded expressions result in canonical equations; substitutions of canonically equal types preserve canonical (in)equations; kinding and subtyping of reducing applications is admissible. Assume that the following equations hold*

$$\Gamma \vdash U_1 = U_2 \Leftarrow K \quad \Gamma, \Sigma = \Gamma, \Delta[U_1/X^{|K|}] \text{ ctx} \quad \Gamma, \Sigma = \Gamma, \Delta[U_2/X^{|K|}] \text{ ctx}$$

for given $\Gamma, \Delta, \Sigma, U_1, U_2, K$ and $X \notin \text{dom}(\Gamma, \Delta, \Sigma)$.

- (1) If $\Gamma, X:K, \Delta \vdash J \text{ kd}$, then $\Gamma, \Sigma \vdash J[U_1/X^{|K|}] \text{ kd}$.
- (2) If $\Gamma, X:K, \Delta \vdash_{\text{var}} X : J$, then $\Gamma, \Sigma \vdash U_1 \Leftarrow J[U_1/X^{|K|}]$.
- (3) If $\Gamma, X:K, \Delta \vdash_{\text{var}} Y : J$ and $Y \neq X$, then $\Gamma, \Sigma \vdash_{\text{var}} Y : J[U_1/X^{|K|}]$.
- (4) If $\Gamma, X:K, \Delta \vdash_{\text{ne}} N : V .. W$, then $\Gamma, \Sigma \vdash N[U_1/X^{|K|}] \Leftarrow (V .. W)[U_1/X^{|K|}]$.
- (5) If $|\Gamma| \vdash J \text{ kds}$ and $\Gamma, X:K, \Delta \vdash J \Rightarrow \mathbf{V} \Rightarrow L$, then

$$\Gamma, \Sigma \vdash J[U_1/X^{|K|}] \Rightarrow \mathbf{V}[U_1/X^{|K|}] \Rightarrow L[U_1/X^{|K|}].$$

- (6) If $\Gamma, X:K, \Delta \vdash V \Rightarrow J$, then $\Gamma, \Sigma \vdash V[U_1/X^{|K|}] \Rightarrow J[U_1/X^{|K|}]$.
- (7) If $\Gamma, X:K, \Delta \vdash V \Leftarrow J$, then $\Gamma, \Sigma \vdash V[U_1/X^{|K|}] \Leftarrow J[U_1/X^{|K|}]$.
- (8) If $\Gamma, X:K, \Delta \vdash J \text{ kd}$, then $\Gamma, \Sigma \vdash J[U_1/X^{|K|}] \leq J[U_2/X^{|K|}]$.
- (9) If $\Gamma, X:K, \Delta \vdash J \text{ kd}$, then $\Gamma, \Sigma \vdash J[U_1/X^{|K|}] = J[U_2/X^{|K|}]$.
- (10) If $\Gamma, X:K, \Delta \vdash_{\text{var}} X : J$, then $\Gamma, \Sigma \vdash U_1 = U_2 \Leftarrow J[U_1/X^{|K|}]$.
- (11) If $\Gamma, X:K, \Delta \vdash_{\text{ne}} N : V .. W$, then $\Gamma, \Sigma \vdash N[U_1/X^{|K|}] \leq N[U_2/X^{|K|}]$.

(12) If $|\Gamma| \vdash J$ kds and $\Gamma, X:K, \Delta \vdash J \Rightarrow \mathbf{V} \Rightarrow L$, then

$$\Gamma, \Sigma \vdash J[U_1/X^{|\mathbf{K}|}] \Rightarrow \mathbf{V}[U_1/X^{|\mathbf{K}|}] = \mathbf{V}[U_2/X^{|\mathbf{K}|}] \Rightarrow L[U_1/X^{|\mathbf{K}|}].$$

(13) If $\Gamma, X:K, \Delta \vdash V \Rightarrow W .. W'$, then $\Gamma, \Sigma \vdash V[U_1/X^{|\mathbf{K}|}] \leq V[U_2/X^{|\mathbf{K}|}]$.

(14) If $\Gamma, X:K, \Delta \vdash V \Rightarrow J$ and $\Gamma, X:K, \Delta \vdash J$ kd, then

$$\Gamma, \Sigma \vdash V[U_1/X^{|\mathbf{K}|}] \leq V[U_2/X^{|\mathbf{K}|}] \Leftarrow J[U_1/X^{|\mathbf{K}|}].$$

(15) If $\Gamma, X:K, \Delta \vdash V \Leftarrow J$ and $\Gamma, X:K, \Delta \vdash J$ kd, then

$$\Gamma, \Sigma \vdash V[U_1/X^{|\mathbf{K}|}] \leq V[U_2/X^{|\mathbf{K}|}] \Leftarrow J[U_1/X^{|\mathbf{K}|}].$$

(16) If $\Gamma, X:K, \Delta \vdash J_1 \leq J_2$, then $\Gamma, \Sigma \vdash J_1[U_1/X^{|\mathbf{K}|}] \leq J_2[U_2/X^{|\mathbf{K}|}]$.

(17) If $|\Gamma| \vdash J$ kds and $\Gamma, X:K, \Delta \vdash J \Rightarrow \mathbf{V}_1 = \mathbf{V}_2 \Rightarrow L$, then

$$\Gamma, \Sigma \vdash J[U_1/X^{|\mathbf{K}|}] \Rightarrow \mathbf{V}_1[U_1/X^{|\mathbf{K}|}] = \mathbf{V}_2[U_2/X^{|\mathbf{K}|}] \Rightarrow L[U_1/X^{|\mathbf{K}|}].$$

(18) If $\Gamma, X:K, \Delta \vdash V_1 \leq V_2$, then $\Gamma, \Sigma \vdash V_1[U_1/X^{|\mathbf{K}|}] \leq V_2[U_2/X^{|\mathbf{K}|}]$.

(19) If $\Gamma, X:K, \Delta \vdash V_1 \leq V_2 \Leftarrow J$ and $\Gamma, X:K, \Delta \vdash J$ kd, then

$$\Gamma, \Sigma \vdash V_1[U_1/X^{|\mathbf{K}|}] \leq V_2[U_2/X^{|\mathbf{K}|}] \Leftarrow J[U_1/X^{|\mathbf{K}|}].$$

(20) If $\Gamma, X:K, \Delta \vdash V_1 = V_2 \Leftarrow J$, then $\Gamma, \Sigma \vdash V_1[U_1/X^{|\mathbf{K}|}] = V_2[U_2/X^{|\mathbf{K}|}] \Leftarrow J[U_1/X^{|\mathbf{K}|}]$.

(21) If $\Gamma \vdash K \Rightarrow \mathbf{V} \Rightarrow J$, then $\Gamma \vdash U_1 \cdot^{|\mathbf{K}|} \mathbf{V} \Leftarrow J$.

(22) If $K = (X:K_1) \rightarrow K_2$, $\Gamma \vdash V \Leftarrow K_1$ and $\Gamma \vdash K_1$ kd, then $\Gamma \vdash U_1 \cdot^{|\mathbf{K}|} V \Leftarrow K_2[V/X^{|\mathbf{K}_1|}]$.

(23) If $\Gamma \vdash K \Rightarrow \mathbf{V}_1 = \mathbf{V}_2 \Rightarrow J$, then $\Gamma \vdash U_1 \cdot^{|\mathbf{K}|} \mathbf{V}_1 = U_2 \cdot^{|\mathbf{K}|} \mathbf{V}_2 \Leftarrow J$.

(24) If $K = (X:K_1) \rightarrow K_2$ and $\Gamma \vdash V_1 = V_2 \Leftarrow K_1$, then

$$\Gamma \vdash U_1 \cdot^{|\mathbf{K}|} V_1 = U_1 \cdot^{|\mathbf{K}|} V_2 \Leftarrow K_2[V_1/X^{|\mathbf{K}_1|}].$$

PROOF. As for the proof of [Lemma C.10](#), the structure of the proof mirrors that of the recursive definition of hereditary substitution itself. All [24](#) parts are proven simultaneously by induction in the structure of the simple kind $|\mathbf{K}|$. Parts [1–20](#) proceed by an inner induction on the respective formation, kinding, subkinding, subtyping or equality derivations of the expressions in which U_1 and U_2 are being substituted for X . Parts [21–24](#) proceed by a case analysis on the final rule used to derive $\Gamma \vdash K \Rightarrow \mathbf{V} \Rightarrow J$, $\Gamma \vdash U_1 \Leftarrow K$, $\Gamma \vdash K \Rightarrow \mathbf{V}_1 = \mathbf{V}_2 \Rightarrow J$ and $\Gamma \vdash U_1 = U_2 \Leftarrow K$, respectively.

The proofs of parts [1–7](#) are similar to that of the declarative substitution lemma ([Lemma B.4](#)), while those of parts [8–20](#) resemble the proof of the extended functionality lemma ([Lemma B.14](#)). In cases like [CWF-DARR](#) or [CK-ALL](#), where the context is extended by an additional binding, we use the IH together with context narrowing ([Lemma D.8](#)) to maintain the invariants $\Gamma, \Sigma = \Gamma, \Delta[U_1/X^{|\mathbf{K}|}]$ ctx and $\Gamma, \Sigma = \Gamma, \Delta[U_2/X^{|\mathbf{K}|}]$ ctx.

The cases where the proofs of parts [1–20](#) differ most substantially from those of [Lemma B.4](#) and [Lemma B.14](#) are parts [4](#), [11](#) and the case for [CST-NE](#) of part [18](#), which deal with neutral types. There, we proceed by case distinction on $X = Y$, where Y is the head of the corresponding neutral types. If $X = Y$, then we proceed using either part [21](#), or part [23](#) followed by [CST-REFL-TEQ'](#). If $X \neq Y$, then we use part [3](#) and proceed with either part [5](#) followed by [CK-NECK](#), or with parts [12](#) or [17](#) followed by [CST-NE](#). In the cases for [CST-BND₁](#) and [CST-BND₂](#), we use part [4](#) followed by [CST-CKBND₁](#) or [CST-CKBND₂](#).

In the cases for [CK-CONS](#) and [SP-EQ-CONS](#) of parts [5](#), [12](#) and [17](#), respectively, where $J = (Y:J_1) \rightarrow J_2$, we use [Lemma C.11.1](#) to show that hereditary substitutions in kinds commute, i.e. that

$$J_2[V_1/Y^{|\mathbf{L}_1|}][U_1/X^{|\mathbf{K}|}] \equiv J_2[U_1/X^{|\mathbf{K}|}][V_1[U_1/X^{|\mathbf{K}|}]/Y^{|\mathbf{L}_1|}].$$

The necessary simple kinding derivations are provided by case analysis of the final rule used to derive the premise $|\Gamma| \vdash J$ kds and [Lemma D.16.5](#).

The proofs of parts [22](#) and [24](#) resemble that of [Lemma C.10.5](#) but are complicated slightly by the presence of subkinding. We show the proof of part [22](#), that of part [24](#) is similar. We have $K = (X:K_1) \rightarrow K_2$, $\Gamma \vdash U_1 \Leftarrow K$, $\Gamma \vdash V \Leftarrow K_1$ and $\Gamma \vdash K_1$ kd, and we want to show that $\Gamma \vdash U_1 \cdot^{|\mathcal{K}|} V \Leftarrow K_2[V/X^{|\mathcal{K}_1|}]$. By inspection of the kind checking and subkinding rules, we must have $U_1 = \lambda X:J_1. U$ such that $\Gamma, X:J_1 \vdash U \Rightarrow J_2$, $\Gamma \vdash K_1 \leq J_1$ and $\Gamma, X:K_1 \vdash J_2 \leq K_2$, and by the definition of reducing application, $U_1 \cdot^{|\mathcal{K}|} V \equiv U[V/X^{|\mathcal{K}_1|}]$. Using context narrowing and the IH for part [7](#), we obtain $\Gamma \vdash U[V/X^{|\mathcal{K}_1|}] \Rightarrow J_2[V/X^{|\mathcal{K}_1|}]$. By [TEQ-REFL](#) and the IH for part [16](#), we have $\Gamma \vdash J_2[V/X^{|\mathcal{K}_1|}] \leq K_2[V/X^{|\mathcal{K}_1|}]$. We conclude by [CK-SUB](#). \square

D.2.1 Validity. With the hereditary substitution lemma in place, we can now prove the remaining validity properties of the canonical judgments. The most intricate cases are those for spine kinding and equality, which is where we use [Lemma D.17](#).

Lemma D.18 (canonical validity – part 2).

- (*spine kinding validity*) If $\Gamma \vdash J$ kd and $\Gamma \vdash J \Rightarrow \mathbf{U} \Rightarrow K$, then $\Gamma \vdash K$ kd.
- (*spine equation validity*) If $\Gamma \vdash J_1 \leq J_2$ and $\Gamma \vdash J_2 \Rightarrow \mathbf{U} = \mathbf{V} \Rightarrow K_2$, then $\Gamma \vdash J_2 \Rightarrow \mathbf{U} \Rightarrow K_2$, $\Gamma \vdash J_1 \Rightarrow \mathbf{V} \Rightarrow K_1$ and $\Gamma \vdash K_1 \leq K_2$ for some K_1 .
- (*neutral kinding validity*) If $\Gamma \vdash N : K$, then $\Gamma \vdash K$ kd.
- (*subkinding validity*) If $\Gamma \vdash J \leq K$, then $\Gamma \vdash J$ kd and $\Gamma \vdash K$ kd.
- (*proper subtyping validity*) If $\Gamma \vdash U \leq V$, then $\Gamma \vdash U \Rightarrow U \dots U$ and $\Gamma \vdash V \Rightarrow V \dots V$.
- (*checked kinding validity*) If $\Gamma \vdash V \Leftarrow K$, then $\Gamma \vdash K$ kd.

PROOF. Subkinding and proper subtyping validity are proven simultaneously, the remaining parts are proven separately, in the order they are listed. All parts are proven by induction on derivations of the judgments they are named after: spine kinding and equation validity are proven by induction on their respective second premises, the remaining parts on their respective first premises. In inductive steps of the proofs of spine kinding and equation validity, we use the hereditary substitution lemmas to derive suitable first premises for applying the IH. The proof of spine equation validity relies on checked equation validity from [Lemma D.18](#). The proof of neutral kinding validity relies on spine kinding validity. In the proof of proper subtyping validity, we use neutral kinding validity in the cases for the bound projection rules [CST-BND_{1,2}](#). The proof of checked kinding validity relies on proper subtyping validity. \square

D.2.2 Lifting of Weak Equality to Canonical Equality. In [§C.2.2](#) of the previous section, we established a number of weak commutativity properties (see [Lemmas C.16](#) and [C.21](#)). Among others, we showed that normalization weakly commutes with hereditary substitution. But up until now, we do not have any effective means to put these properties to use – we have yet to establish a relationship between weak equality and the equality judgments of the declarative and canonical systems.

To remedy this situation, we prove that a weak equation $U \approx V$ can be lifted to canonical equation $\Gamma \vdash U = V \Leftarrow K$, provided the left- and right-hand sides U, V are well-kinded, i.e. $\Gamma \vdash U \Leftarrow K$ and $\Gamma \vdash V \Leftarrow K$. Similarly, we show that weakly equal kinds are canonically equal if they are well-formed.

Lemma D.19. *Weakly equal canonically well-formed kinds and well-kinded types are canonically equal.*

- (1) If $\Gamma \vdash J$ kd, $\Gamma \vdash K$ kd and $J \approx K$, then $\Gamma \vdash J \leq K$.
- (2) If $\Gamma \vdash U \Rightarrow U \dots U$, $\Gamma \vdash V \Rightarrow V \dots V$ and $U \approx V$, then $\Gamma \vdash U \leq V$.
- (3) If $\Gamma \vdash K$ kd, $\Gamma \vdash U \Leftarrow K$, $\Gamma \vdash V \Leftarrow K$ and $U \approx V$, then $\Gamma \vdash U \leq V \Leftarrow K$.

- (4) If $\Gamma \vdash J \text{ kd}$, $\Gamma \vdash J \leq K_1$, $\Gamma \vdash J \leq K_2$, $\Gamma \vdash K_1 \Rightarrow \mathbf{U}_1 \Rightarrow V_1 .. W_1$, $\Gamma \vdash K_2 \Rightarrow \mathbf{U}_2 \Rightarrow V_2 .. W_2$, and $\mathbf{U}_1 \approx \mathbf{U}_2$, then $\Gamma \vdash J \Rightarrow \mathbf{U}_1 = \mathbf{U}_2 \Rightarrow V .. W$ for some V and W .
- (5) If $\Gamma \vdash J \text{ kd}$, $\Gamma \vdash K \text{ kd}$ and $J \approx K$, then $\Gamma \vdash J = K$.
- (6) If $\Gamma \vdash K \text{ kd}$, $\Gamma \vdash U \Leftarrow K$, $\Gamma \vdash V \Leftarrow K$ and $U \approx V$, then $\Gamma \vdash U = V \Leftarrow K$.

PROOF. Simultaneously for all 6 parts by simultaneous induction on the corresponding pairs of kinds, types or spines being related, then by case analysis on the final rules used to derive the corresponding formation, kinding and weak equality judgments. In the proof of part 5, we apply the IH for part 1 directly to the equations $J \approx K$ and $K \approx J$, where the latter is derived using symmetry of weak equality. Neither J nor K decrease in this step, but the proof of part 5 does not make any further use of the IH and could therefore be inlined in the proofs of the other parts. The proof of part 5 is similar.

The proofs of the remaining parts are largely routine. The most interesting case is the inductive one in part 4, where we have $\mathbf{U}_1 = (U_1, \mathbf{U}'_1)$, $\mathbf{U}_2 = (U_2, \mathbf{U}'_2)$, $K_1 = (X:K_{11}) \rightarrow K_{12}$, $K_2 = (X:K_{21}) \rightarrow K_{22}$, $U_1 \approx U_2$ and $\mathbf{U}'_1 \approx \mathbf{U}'_2$. Analyzing the derivations of the remaining premises, we have

$$\Gamma \vdash K_{11} \leq J_1 \quad \Gamma, X:K_{11} \vdash J_2 \leq K_{12} \quad \Gamma \vdash K_{21} \leq J_1 \quad \Gamma, X:K_{21} \vdash J_2 \leq K_{22}$$

such that $J = (X:J_1) \rightarrow J_2$, as well as

$$\begin{array}{ll} \Gamma \vdash U_1 \Leftarrow K_{11} & \Gamma \vdash K_{12}[U_1/X^{|K_{11}|}] \Rightarrow \mathbf{U}'_1 \Rightarrow V_1 .. W_1 \\ \Gamma \vdash U_2 \Leftarrow K_{21} & \Gamma \vdash K_{22}[U_2/X^{|K_{21}|}] \Rightarrow \mathbf{U}'_2 \Rightarrow V_2 .. W_2 \end{array}$$

We use **CK-SUBCK** and the IH for part 6 to derive $\Gamma \vdash U_1 = U_2 \Leftarrow J_1$, then we use hereditary substitution (**Lemmas D.17.16** and **D.17.8**) and **Lemma D.15** to derive

$$\begin{array}{ll} \Gamma \vdash J_2[U_1/X^{|J_1|}] \leq & K_{12}[U_1/X^{|K_{11}|}] \\ \Gamma \vdash J_2[U_1/X^{|J_1|}] \leq J_2[U_2/X^{|J_1|}] \leq & K_{22}[U_2/X^{|K_{21}|}] \end{array}$$

We conclude the case by the IH for part 4 and **SPeQ-CONS**. \square

D.3 Completeness of Canonical Kinding

In the previous section, we saw that every declaratively well-formed kind or well-kinded type has a judgmentally equal $\beta\eta$ -normal form (**Lemmas 4.3** and **C.14**). In this section, we prove that every declarative judgment has a canonical counterpart where the expressions related by the original judgment have been normalized. Roughly, whenever $\Gamma \vdash_d \mathcal{J}$ holds, we also have $\text{nf}(\Gamma) \vdash_c \text{nf}_{\text{nf}(\Gamma)}(\mathcal{J})$. Since normalization does not change the meaning of an expression, this result establishes completeness of the canonical system w.r.t. to the declarative one.

There are several judgments for kinding types in the canonical system, but only one in the declarative system. To establish completeness, we show that, if a type A is of kind K according to declarative kinding, then the normal form $\text{nf}(A)$ kind checks against the normal form $\text{nf}(K)$, i.e. if $\Gamma \vdash_d A : K$, then $\text{nf}(\Gamma) \vdash_c \text{nf}(A) \Leftarrow \text{nf}(K)$.

When A is a variable $A = X$, the normal form $\text{nf}(A)$ is its η -expansion $\text{nf}(A) = \eta_{\text{nf}(K)}(X)$ and we use the following lemma to prove that it kind checks against $\text{nf}(K)$.

Lemma D.20 (η -expansion). *η -expansion preserves the canonical kinds of neutral types. If $\Gamma \vdash_{\text{ne}} N : K$, then $\Gamma \vdash \eta_K(N) \Leftarrow K$.*

Instead of proving the lemma directly, we first prove the following helper lemma.

Lemma D.21.

- (1) If $\Gamma, X:J \vdash K \text{ kd}$ and $\Gamma, X:J \vdash \eta_J(X) \Leftarrow J$, then $\Gamma, X:J \vdash K[\eta_J(X)/X^{|J|}] = K$.
- (2) If $\Gamma \vdash_{\text{ne}} N : J$ and $\Gamma \vdash J \leq K$, then $\Gamma \vdash \eta_K(N) \Leftarrow K$.

The first part says that hereditary substitutions of η -expanded variables in kinds vanish, while the second part is a strengthened version of [Lemma D.20](#).

PROOF. The two parts are proven separately. For the first part, we use simplification of canonical kinding ([Lemma D.16.2](#)) and [Lemma C.16.2](#) to derive $K[\eta_J(X)/X^{|J|}] \approx K$. By weakening and the hereditary substitution lemma ([Lemma D.17.1](#)), we have $\Gamma, X:J \vdash K[\eta_J(X)/X^{|J|}]$ kd. The conclusion of the first part then follows by [Lemma D.19.5](#).

The proof of the second part is by induction on the structure of K and case analysis on the final rule used to derive $\Gamma \vdash J \leq K$. In the base case, we use **CK-NECK** and **CK-SUBCK**. In the inductive case, we have $K = (X:K_1) \rightarrow K_2$, $J = (X:J_1) \rightarrow J_2$ such that $\Gamma \vdash K_1 \leq J_1$ and $\Gamma, X:K_1 \vdash J_2 \leq K_2$. By subkinding validity, we further have $\Gamma \vdash K_1$ kd and $\Gamma, X:K_1 \vdash K_2$ kd. By weakening, the IH and **CSK-REFL**, we obtain first $\Gamma, X:K_1 \vdash \eta_{K_1}(X) \Leftarrow K_1$, then $\Gamma, X:K_1 \vdash \eta_{K_1}(X) \Leftarrow J_1$ by weakening and **CK-SUBCK**. By inspection of neutral kinding, we know that $N = Y \mathbf{V}$ for some Y and \mathbf{V} and that $\Gamma \vdash_{\text{var}} Y : L$ and $\Gamma \vdash L \Rightarrow \mathbf{V} \Rightarrow (X:J_1) \rightarrow J_2$. We use weakening, **CK-SNOC** and **CK-NE** to derive $\Gamma, X:K_1 \vdash_{\text{ne}} Y \mathbf{V} (\eta_{K_1}(X)) : J_2[\eta_{K_1}(X)/X^{|J_1|}]$.

Now we see why it was necessary to strengthen the IH: the body of the η -expansion of N has kind $J_2[\eta_{K_1}(X)/X^{|J_1|}]$ rather than K_2 as required by [Lemma D.20](#). In order to apply the IH, we show that

$$\begin{aligned} \Gamma, X:K_1 \vdash J_2[\eta_{K_1}(X)/X^{|J_1|}] &\equiv J_2[\eta_{K_1}(X)/X^{|K_1|}] && \text{(by Lemma D.15)} \\ &\leq K_2[\eta_{K_1}(X)/X^{|K_1|}] && \text{(by Lemma D.17.16)} \\ &= K_2. && \text{(by part 1)} \end{aligned}$$

We conclude the case by the IH and **CK-ABS'**. \square

[Lemma D.20](#) as well as a strengthened version of [Lemma D.21.1](#) now follow as corollaries.

Corollary D.22. *If $\Gamma, X:J \vdash K$ kd, then $\Gamma, X:J \vdash K[\eta_J(X)/X^{|J|}] = K$.*

To establish completeness of the canonical system w.r.t. the declarative system, we show that every declarative judgment derived using the *extended* declarative rules, rather than the original ones, has a canonical counterpart. The proof makes crucial use of the validity conditions present in the extended rules. To avoid confusion, we again mark canonical judgments with the subscript “c” and extended declarative ones with “e”. To enhance readability, we omit the subscript $\text{nf}(\Gamma)$, writing e.g. $\text{nf}(A)$ instead of $\text{nf}_{\text{nf}(\Gamma)}(A)$.

Lemma D.23 (completeness of the canonical rules – extended version).

- (1) If $\Gamma \text{ ctx}_e$, then $\text{nf}(\Gamma) \text{ ctx}_c$.
- (2) If $\Gamma \vdash_e K$ kd, then $\text{nf}(\Gamma) \vdash_c \text{nf}(K)$ kd.
- (3) If $\Gamma \vdash_e A : K$, then $\text{nf}(\Gamma) \vdash_c \text{nf}(A) \Leftarrow \text{nf}(K)$.
- (4) If $\Gamma \vdash_e J \leq K$, then $\text{nf}(\Gamma) \vdash_c \text{nf}(J) \leq \text{nf}(K)$.
- (5) If $\Gamma \vdash_e A \leq B : C..D$, then $\text{nf}(\Gamma) \vdash_c \text{nf}(A) \leq \text{nf}(B)$.
- (6) If $\Gamma \vdash_e A \leq B : K$, then $\text{nf}(\Gamma) \vdash_c \text{nf}(A) \leq \text{nf}(B) \Leftarrow \text{nf}(K)$.
- (7) If $\Gamma \vdash_e J = K$, then $\text{nf}(\Gamma) \vdash_c \text{nf}(J) = \text{nf}(K)$.
- (8) If $\Gamma \vdash_e A = B : K$, then $\text{nf}(\Gamma) \vdash_c \text{nf}(A) = \text{nf}(B) \Leftarrow \text{nf}(K)$.
- (9) If $\Gamma \vdash_e A : (X:J) \rightarrow K$ and $X \notin \text{fv}(A)$, then

$$\text{nf}(\Gamma) \vdash_c \text{nf}(\lambda X:J. AX) = \text{nf}(A) \Leftarrow \text{nf}((X:J) \rightarrow K).$$

- (10) If $\Gamma, X:J \vdash_e K$ kd, $\Gamma \vdash_e A : J$ and $\Gamma \vdash_e K[A/X]$ kd, then

$$\text{nf}(\Gamma) \vdash_c \text{nf}(K)[\text{nf}(A)/X^{\text{nf}(J)}] = \text{nf}(K[A/X]).$$

(11) If $\Gamma, X:J \vdash_e A : K$, $\Gamma \vdash_e B : J$, $\Gamma, X:J \vdash_e K \text{ kd}$, $\Gamma \vdash_e A[B/X] : K[B/X]$
and $\Gamma \vdash_e K[B/X] \text{ kd}$, then

$$\text{nf}(\Gamma) \vdash_c \text{nf}((\lambda X:J. A) B) = \text{nf}(A[B/X]) \Leftarrow \text{nf}(K[B/X]).$$

Equivalent statements w.r.t. the original declarative rules follow by equivalence of the original and extended declarative systems.

PROOF. All parts are proven simultaneously, by induction on the derivations of the respective premises, except for parts 9–11, which are helper lemmas that apply the IH directly to all of their premises but could be inlined in the proofs of the other parts.

Thanks to the admissible rules introduced in Lemma D.14, the proofs of parts 1–3 are straightforward, except for the cases of **K-VAR**, where we use Lemma D.20, and **K-APP**, where we use the hereditary substitution lemma to normalize $A = A_1 A_2$ if $\text{nf}(A_1)$ is an abstraction, and the IH for part 10 together with **CK-SUBCK** to adjust the kind of the result. The validity conditions of **K-APP** are crucial in this last step.

Parts 5 and 8 follow almost immediately from part 6, part 7 from part 4.

The proofs of parts 9–11 all follow the same pattern. First, we use the IH to normalize the premises and establish validity of the left- and right-hand sides of the respective equations. Then we use Lemma C.22, Lemma C.21.1 and Lemma C.21.2, respectively, to derive weak versions of these equations, and Lemma D.19 to turn them into canonical equations.

The remaining parts 4 and 6 are the most difficult to prove. The cases of the extended β and η -conversion rules are covered by parts 9 and 11 thanks to the validity conditions in the extended rules. The case of **ST-APP** is similar to that of **K-APP** – again the validity conditions are crucial. Some of the remaining cases are covered by the admissible rules introduced in §§D.1.1 and D.1.2. The challenging cases are those where one of the premises of the corresponding rule extends the contexts, i.e. those of **CSK-DARR**, **CST-ALL** and **CST-ABS**. We show the case for **CSK-DARR** here, the other two are similar.

We are given $\Gamma \vdash_e (X:J_1) \rightarrow J_2 \text{ kd}$, $\Gamma \vdash_e K_1 \leq J_1$ and $\Gamma, X:K_1 \vdash_e J_2 \leq K_2$ with $J = (X:J_1) \rightarrow J_2$ and $K = (X:K_1) \rightarrow K_2$. We start by applying the IH to all the premises and analyze the first of the resulting derivations to obtain

$$\begin{array}{ll} \text{nf}(\Gamma) \vdash_c \text{nf}(J_1) \text{ kd} & \text{nf}(\Gamma, X:\text{nf}(J_1)) \vdash_c \text{nf}_{\text{nf}(\Gamma, X:J_1)}(J_2) \text{ kd} \\ \text{nf}(\Gamma) \vdash_c \text{nf}(K_1) \leq \text{nf}(J_1) & \text{nf}(\Gamma, X:\text{nf}(K_1)) \vdash_c \text{nf}_{\text{nf}(\Gamma, X:K_1)}(J_2) \leq \text{nf}_{\text{nf}(\Gamma, X:K_1)}(K_2). \end{array}$$

Note the different contexts $\text{nf}(\Gamma, X:J_1)$ and $\text{nf}(\Gamma, X:K_1)$ used to normalize J_2 in the second and fourth of these judgments, respectively. This leads to a syntactic difference in the resulting normal forms, i.e. we have $\text{nf}_{\text{nf}(\Gamma, X:J_1)}(J_2) \neq \text{nf}_{\text{nf}(\Gamma, X:K_1)}(J_2)$. In order to apply **CSK-DARR**, we need to resolve this difference.

We notice that $|\text{nf}(\Gamma, X:J_1)| \equiv |\text{nf}(\Gamma, X:K_1)|$ by Lemma D.15. Hence, by Lemma C.6, we have $\text{nf}_{\text{nf}(\Gamma, X:J_1)}(J_2) \approx \text{nf}_{\text{nf}(\Gamma, X:K_1)}(J_2)$. Using context narrowing and subkinding validity, we derive

$$\text{nf}(\Gamma, X:\text{nf}(K_1)) \vdash \text{nf}_{\text{nf}(\Gamma, X:J_1)}(J_2) \text{ kd} \quad \text{and} \quad \text{nf}(\Gamma, X:\text{nf}(K_1)) \vdash \text{nf}_{\text{nf}(\Gamma, X:K_1)}(J_2) \text{ kd},$$

from which we obtain, by Lemma D.19.1,

$$\text{nf}(\Gamma, X:\text{nf}(K_1)) \vdash_c \text{nf}_{\text{nf}(\Gamma, X:J_1)}(J_2) = \text{nf}_{\text{nf}(\Gamma, X:K_1)}(J_2) \leq \text{nf}_{\text{nf}(\Gamma, X:K_1)}(K_2)$$

We conclude the case by **CWF-DARR** and **CSK-DARR**. \square

Transitivity-free subtyping of closed proper types

$$\boxed{\vdash_{\text{tf}} U \leq V}$$

$$\begin{array}{c} \frac{\emptyset \vdash V \Rightarrow V .. V}{\vdash_{\text{tf}} V \leq \top} \quad (\text{TfST-Top}) \qquad \frac{\emptyset \vdash V \Rightarrow V .. V}{\vdash_{\text{tf}} \perp \leq V} \quad (\text{TfST-Bot}) \\ \\ \frac{\emptyset \vdash U_2 \leq U_1 \quad \emptyset \vdash V_1 \leq V_2}{\vdash_{\text{tf}} U_1 \rightarrow V_1 \leq U_2 \rightarrow V_2} \quad (\text{TfST-ARR}) \qquad \frac{\emptyset \vdash \forall X:K_1. V_1 \Rightarrow \forall X:K_1. V_1 .. \forall X:K_1. V_1 \quad \emptyset \vdash K_2 \leq K_1 \quad X:K_2 \vdash V_1 \leq V_2}{\vdash_{\text{tf}} \forall X:K_1. V_1 \leq \forall X:K_2. V_2} \quad (\text{TfST-ALL}) \end{array}$$

Fig. 11. Top-level transitivity-free canonical subtyping

D.4 Inversion of Subtyping

As we saw in §3 of the paper, preservation of types under CBV reduction does not hold in arbitrary contexts. The culprit are type variable bindings with inconsistent bounds. Such bindings can inject arbitrary inequations into the subtyping relation and hence break putative properties that hold for subtyping of closed types. For example, the absurd assumption $X : \top .. \perp$ trivializes the subtyping relation under any context in which it appears. To see this, consider the following derivation, where $\Gamma = X : \top .. \perp$.

$$\frac{\frac{\vdots \quad \frac{\Gamma \vdash_{\text{ne}} X : \top .. \perp}{\Gamma \vdash \top \leq X} \quad \frac{\vdots \quad \frac{\Gamma \vdash_{\text{ne}} X : \top .. \perp}{\Gamma \vdash X \leq \perp}}{\Gamma \vdash \top \leq \perp} \quad \vdots}{\Gamma \vdash U \rightarrow V \leq \top} \quad \frac{\Gamma \vdash \top \leq \perp}{\Gamma \vdash U \rightarrow V \leq \perp} \quad \frac{\vdots}{\Gamma \vdash \perp \leq \forall X:K. W}}{\Gamma \vdash U \rightarrow V \leq \forall X:K. W} \quad (\text{CST-TRANS})$$

Under such conditions, subtyping cannot be inverted in any meaningful way. We therefore consider inversion of canonical subtyping only in the empty context, following the approach taken by Rompf and Amin in their type safety proof for DOT [Rompf and Amin 2016]

As a first step we show that any top-level uses of the transitivity rule **CST-TRANS** can be eliminated. To do so, we introduce a helper judgment $\vdash_{\text{tf}} U \leq V$, which states that U is a proper subtype of V in the empty context (see Fig. 11). It is easy to see that this judgment is sound w.r.t. canonical subtyping in the empty context (the proof is by routine induction on subtyping derivations).

Lemma D.24 (soundness of top-level subtyping). *If $\vdash_{\text{tf}} U \leq V$, then $\emptyset \vdash U \leq V$.*

Crucially, the inference rules for the judgment $\vdash_{\text{tf}} U \leq V$ do not include a transitivity rule, but the following variant of that rule is admissible.

Lemma D.25 (top-level transitivity elimination). *The following is admissible.*

$$\frac{\emptyset \vdash U \leq V \quad \vdash_{\text{tf}} V \leq W}{\vdash_{\text{tf}} U \leq W} \quad (\text{TfST-TRANS})$$

PROOF. The proof is by induction on the derivation of the first premise and case analysis of the final rule used to derive the second. In the case of **CST-TRANS**, we use the IH twice. In the case of **CST-Bot** where $\emptyset \vdash \perp \leq U$ and $\vdash_{\text{tf}} U \leq V$, we use **Lemma D.24** and validity of canonical typing to derive $\emptyset \vdash V \Rightarrow V .. V$ and conclude with **TfST-Bot**. Similarly, in cases where the second premise was derived using **TfST-Top**, we use validity of canonical subtyping and **TfST-Top**. \square

Thanks to **TFST-TRANS**, it is straightforward to establish completeness, and thus equivalence of the judgments $\emptyset \vdash U \leq V$ and $\vdash_{\text{tf}} U \leq V$.

Lemma D.26 (equivalence of top-level subtyping). *The two versions of canonical subtyping are equivalent in the empty context: $\emptyset \vdash U \leq V$ iff $\vdash_{\text{tf}} U \leq V$.*

PROOF. We have already proven soundness (\Leftarrow). Completeness (\Rightarrow) is by induction on the derivations of $\emptyset \vdash U \leq V$ and uses **TFST-TRANS** in the case of **CST-TRANS**. \square

Inversion of the canonical subtyping relation in the empty context now follows immediately by inspection of the transitivity-free subtyping rules and **Lemma D.26**. We only state the relevant cases.

Corollary D.27 (inversion of canonical subtyping – embedding). *Let $\emptyset \vdash U_1 \leq U_2$.*

- (1) *If $U_1 = V_1 \rightarrow W_1$ and $U_2 = V_2 \rightarrow W_2$, then $\emptyset \vdash V_2 \leq V_1$ and $\emptyset \vdash W_1 \leq W_2$.*
- (2) *If $U_1 = \forall X:K_1. V_1$ and $U_2 = \forall X:K_2. V_2$, then $\emptyset \vdash K_2 \leq K_1$ and $X:K_2 \vdash V_1 \leq V_2$.*

Corollary D.28 (inversion of canonical subtyping – contradiction). *For any U, V, W and K ,*

- (1) $\emptyset \not\vdash \top \leq \perp$,
- (2) $\emptyset \not\vdash U \rightarrow V \leq \forall X:K. W$, and
- (3) $\emptyset \not\vdash \forall X:K. U \leq V \rightarrow W$.

A bit more work is needed to also prove the declarative version of subtyping inversion. Again, we only state the relevant cases.

Lemma D.29 (inversion of declarative subtyping – embedding). *Let $\emptyset \vdash A_1 \leq A_2 : *$.*

- (1) *If $A_1 = B_1 \rightarrow C_1$ and $A_2 = B_2 \rightarrow C_2$, then $\emptyset \vdash B_2 \leq B_1 : *$ and $\emptyset \vdash C_1 \leq C_2 : *$.*
- (2) *If $A_1 = \forall X:K_1. B_1$ and $A_2 = \forall X:K_2. B_2$, then $\emptyset \vdash K_2 \leq K_1$ and $X:K_2 \vdash B_1 \leq B_2 : *$.*

The proof makes use of the following generation lemma for well-kinded arrow and universal types, which is proven by induction on kinding derivations.

Lemma D.30 (generation of kinding for arrows and universals). *The following are admissible.*

$$\frac{\Gamma \vdash A \rightarrow B : *}{\Gamma \vdash A : * \quad \Gamma \vdash B : *} \qquad \frac{\Gamma \vdash \forall X:K. A : *}{\Gamma \vdash K \text{ kd} \quad \Gamma, X:K \vdash A : *}$$

PROOF OF LEMMA D.29. By completeness of canonical subtyping and soundness of normalization. We show only the first part, the second is analogous. Assume $\emptyset \vdash B_1 \rightarrow C_1 \leq B_2 \rightarrow C_2 : *$. Then by validity of declarative subtyping (**Lemma 3.1**), generation of kinding for arrow types, soundness of normalization (**Lemma 4.3**) and completeness of canonical subtyping, we have

$$\begin{aligned} \emptyset \vdash_{\text{d}} B_1 &= \text{nf}_{\text{nf}(\Gamma)}(B_1) : * & \emptyset \vdash_{\text{d}} C_1 &= \text{nf}_{\text{nf}(\Gamma)}(C_1) : * \\ \emptyset \vdash_{\text{d}} B_2 &= \text{nf}_{\text{nf}(\Gamma)}(B_2) : * & \emptyset \vdash_{\text{d}} C_2 &= \text{nf}_{\text{nf}(\Gamma)}(C_2) : * \\ \emptyset \vdash_{\text{c}} \text{nf}_{\text{nf}(\Gamma)}(B_1) &\rightarrow \text{nf}_{\text{nf}(\Gamma)}(C_1) \leq \text{nf}_{\text{nf}(\Gamma)}(B_2) &\rightarrow \text{nf}_{\text{nf}(\Gamma)}(C_2) \end{aligned}$$

By inversion and soundness of canonical subtyping, it follows that

$$\begin{aligned} \emptyset \vdash_{\text{d}} B_2 &= \text{nf}_{\text{nf}(\Gamma)}(B_2) \leq \text{nf}_{\text{nf}(\Gamma)}(B_1) = B_1 : * \quad \text{and} \\ \emptyset \vdash_{\text{d}} C_1 &= \text{nf}_{\text{nf}(\Gamma)}(C_1) \leq \text{nf}_{\text{nf}(\Gamma)}(C_2) = C_2 : *. \end{aligned} \quad \square$$

We also prove a declarative counterpart of **Corollary D.28**, which is used in the proof of the progress theorem below.

Lemma D.31 (inversion of declarative subtyping – contradiction). *For any A, B, C and K ,*

- (1) $\emptyset \not\leq \top \leq \perp$,
- (2) $\emptyset \not\leq A \rightarrow B \leq \forall X:K. C$, and
- (3) $\emptyset \not\leq \forall X:K. A \leq B \rightarrow C$.

PROOF. By completeness of canonical subtyping, then by contradiction using [Corollary D.28](#). \square

D.4.1 Type Safety Revisited. We are finally ready to prove type safety of F^ω . The proof of weak preservation requires a standard generation lemma for term and type abstractions.

Lemma D.32 (generation of typing for term and type abstraction).

- (1) If $\Gamma \vdash \lambda x:A. t : B$, then $\Gamma, x:A \vdash t : C$ and $\Gamma \vdash A \rightarrow C \leq B : *$ for some C .
- (2) If $\Gamma \vdash \lambda X:K. t : A$, then $\Gamma, X:K \vdash t : B$ and $\Gamma \vdash \forall X:K. B \leq A : *$ for some B .

Recall that weak preservation ([Prop. 3.2](#)) states that CBV reduction preserves the types of closed terms, i.e. if $\vdash t : A$ and $t \rightarrow_v t'$, then $\vdash t' : A$.

PROOF OF [PROP. 3.2](#). The proof is by induction on typing derivations and case analysis on CBV reduction rules. The interesting cases are those where β -contractions occur. We describe the case of [T-APP](#). The corresponding case for [T-TAPP](#) is similar. We have $t = (\lambda x:B. s) v$ with $\vdash \lambda x:B. s : C \rightarrow A$ and $\vdash v : C$ for some B and C . By generation of term abstractions ([Lemma D.32.1](#)), $x:B \vdash s : D$ and $\vdash B \rightarrow D \leq C \rightarrow A : *$ for some D . By inversion of subtyping ([Lemma D.29.2](#)), we have $\vdash C \leq B : *$ and $\vdash D \leq A : *$ and hence $\vdash v : B$ and $x:B \vdash s : A$ by subsumption. To conclude the proof we need to show that $\vdash s[v/x] : A$, which follows from the substitution lemma ([Lemma B.4](#)). \square

This establishes the first half of type safety. For the second half, progress, we first need to prove a standard canonical forms lemma.

Lemma D.33 (canonical forms). *Let v be a closed, well-typed value.*

- (1) If $\emptyset \vdash v \in A \rightarrow B$, then $v = \lambda x:C. t$ for some C and t .
- (2) If $\emptyset \vdash v \in \forall X:K. A$, then $v = \lambda X:J. t$ for some J and t .

PROOF. Separately for the two parts; each by case analysis, first on v , then on the final typing rule used to derive the respective premise. Since the only values are abstractions, the relevant sub-cases are [T-ABS](#), [T-TABS](#) and [T-SUB](#). The sub-cases for [T-ABS](#) and [T-TABS](#) are immediate. In the sub-cases for [T-SUB](#), we first use the generation lemma for abstractions ([Lemma D.32](#)), then dismiss impossible sub-cases using [Lemma D.31](#). \square

Thanks to the canonical forms lemma, the proof of the progress theorem is now entirely standard.

Theorem D.34 (progress). *If $\vdash t : A$, then either t is a value, or $t \rightarrow_v t'$ for some term t' .*

PROOF. By routine induction on typing derivations. The cases for [T-APP](#) and [T-TAPP](#) use the canonical forms lemma ([Lemma D.33](#)). \square

E ENCODING CUSTOM SUBTYPING THEORIES

Disclaimer. *The following examples have not been mechanized in Agda.*

Recall that bindings of the form $X : A .. B$ represent first-class *type inequations* $A \leq B$ in F^ω because, in a context containing such a binding, we have $A \leq X$ and $X \leq B$, and hence – by transitivity of subtyping – $A \leq B$. Interval kinds thus provide us with a mechanism for (*in*)equality reflection, i.e. a way to extend the subtyping relation via assumptions made at the term- or type-level (via type abstractions). Among other things, this allows us to postulate type operators with associated subtyping rules through type variable bindings. For example, we may postulate intersection types

$A \wedge B$ by assuming an abstract binary type operator $\bar{\wedge} : * \rightarrow * \rightarrow *$ and the usual typing rules for intersections as abstract type inequations:

$$Y_{\leq L} : (X_1, X_2 : *) \rightarrow (\bar{\wedge} X_1 X_2) .. X_1, \quad Y_{\leq R} : (X_1, X_2 : *) \rightarrow (\bar{\wedge} X_1 X_2) .. X_2,$$

$$Y_{\leq \wedge} : (Z, X_1, X_2 : *) \rightarrow Z .. X_1 \rightarrow Z .. X_2 \rightarrow Z .. (\bar{\wedge} X_1 X_2).$$

where we abbreviated dependent arrow kinds $(X_1 : J) \rightarrow (X_2 : J) \rightarrow \dots \rightarrow (X_n : J) \rightarrow K$ with multiple parameters as $(X_1, X_2, \dots, X_n : J) \rightarrow K$ for readability. The two variables $Y_{\leq L}$ and $Y_{\leq R}$ represent, respectively, the left- and right-hand projection rules for intersections ($A \wedge B \leq A$ and $A \wedge B \leq B$). The assumption $Y_{\leq \wedge}$ encodes the fact that intersections are greatest lower bounds, i.e. that $A \leq B \wedge C$ when $A \leq B$ and $A \leq C$. To see how this last “rule” can be put to work, let A, B and C be proper types and assume that $A \leq B$ and $A \leq C$. Then we also have $A : A .. B$ and $A : A .. C$, and hence $Y_{\leq \wedge} A B C A A : A .. (\bar{\wedge} B C)$, from which we conclude $A \leq \bar{\wedge} B C$.

We can also postulate recursive inequations. For example, the following bindings encode an equi-recursive type constructor $\mu : (* \rightarrow *) \rightarrow *$ that, when applied to a unary operator A , represents the fixpoint μA of A .

$$\bar{\mu} : (* \rightarrow *) \rightarrow *,$$

$$Y_{\leq L} : (X : * \rightarrow *) \rightarrow (\bar{\mu} X) .. (X (\bar{\mu} X)), \quad Y_{\leq R} : (X : * \rightarrow *) \rightarrow (X (\bar{\mu} X)) .. (\bar{\mu} X).$$

Together, the assumptions $Y_{\leq L}$ and $Y_{\leq R}$ say that $\mu A = A (\mu A)$, i.e. that μA is a fixpoint of A .

The above examples are only possible because we do not impose any *consistency constraints* on the bounds of intervals. That is, an interval kind $A .. B$ is well-formed, irrespective of whether $A \leq B$ is actually provable or not. For example, the signature of the abstract intersection operator $\bar{\wedge}$ above does not tell us anything about how the type application $\bar{\wedge} A B$ is related to its first argument A , nor does the abstract left projection inequality $Y_{\leq L}$ impose any constraints on its parameters X_1 and X_2 . It is therefore impossible to say anything about the relationship of the bounds $\bar{\wedge} X_1 X_2$ and X_1 of the codomain of $Y_{\leq L}$, other than that they are both proper types. We can certainly not prove that $\bar{\wedge} X_1 X_2 \leq X_1$ in general.

F REPORTED SCALA 3 ISSUES

During the development of F^ω we discovered and reported the following issues to the Scala 3 bug tracker.

- <https://github.com/lampepfl/dotty/issues/2887>
- <https://github.com/lampepfl/dotty/issues/6320>
- <https://github.com/lampepfl/dotty/issues/6499>
- <https://github.com/lampepfl/dotty/issues/9691>
- <https://github.com/lampepfl/dotty/issues/9695>
- <https://github.com/lampepfl/dotty/issues/9697>

REFERENCES

- Andreas Abel. 2008. Polarized Subtyping for Sized Types. *Mathematical Structures in Computer Science* 18 (10 2008), 797–822. Issue Special Issue 05. <https://doi.org/10.1017/S0960129508006853>
- Andreas Abel and Dulma Rodriguez. 2008. Syntactic Metatheory of Higher-Order Subtyping. In *Proceedings of the 22nd International Workshop on Computer Science Logic (CSL 2008), 17th Annual Conference of the EACSL, Bertinoro, Italy (LNCS, Vol. 5213)*, Michael Kaminski and Simone Martini (Eds.). Springer, Berlin, Heidelberg, 446–460. https://doi.org/10.1007/978-3-540-87531-4_32
- Robin Adams. 2006. Pure type systems with judgemental equality. *Journal of Functional Programming* 16, 2 (2006), 219–246. <https://doi.org/10.1017/S0956796805005770>

- Nada Amin. 2016. *Dependent Object Types*. Ph.D. Dissertation. School of Computer and Communication Sciences, École polytechnique fédérale de Lausanne, Lausanne, Switzerland. <https://doi.org/10.5075/epfl-thesis-7156> EPFL thesis no. 7156.
- Nada Amin, Samuel Grütter, Martin Odersky, Tiark Rompf, and Sandro Stucki. 2016. The Essence of Dependent Object Types. In *A List of Successes That Can Change the World: Essays Dedicated to Philip Wadler on the Occasion of His 60th Birthday*, Sam Lindley, Conor McBride, Phil Trinder, and Don Sannella (Eds.). LNCS, Vol. 9600. Springer International Publishing, Cham, 249–272. https://doi.org/10.1007/978-3-319-30936-1_14
- Nada Amin, Tiark Rompf, and Martin Odersky. 2014. Foundations of Path-dependent Types. In *Proceedings of the 2014 ACM International Conference on Object Oriented Programming Systems Languages & Applications (OOPSLA 2014), Portland, Oregon, USA*. ACM, New York, NY, USA, 233–249. <https://doi.org/10.1145/2660193.2660216>
- David Aspinall and Adriana Compagnoni. 2001. Subtyping dependent types. *Theoretical Computer Science* 266, 1-2 (2001), 273–309. [https://doi.org/10.1016/S0304-3975\(00\)00175-4](https://doi.org/10.1016/S0304-3975(00)00175-4)
- Hendrik P. Barendregt. 1992. Lambda Calculi with Types. In *Handbook of Logic in Computer Science*, Samson Abramsky, Dov M. Gabbay, and Thomas S. E Maibaum (Eds.). Vol. 2. Oxford University Press, Oxford, UK, Chapter 2, 117–309.
- Luca Cardelli. 1988. Structural Subtyping and the Notion of Power Type. In *Proceedings of the 15th ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages (POPL 1988), San Diego, California, USA*. ACM, New York, NY, USA, 70–79. <https://doi.org/10.1145/73560.73566>
- Luca Cardelli. 1990. Notes about F_{\leq}^{ω} . (October 1990). Unpublished manuscript.
- Luca Cardelli and Giuseppe Longo. 1991. A Semantic Basis for Quest. *Journal of Functional Programming* 1, 4 (1991), 417–458. <https://doi.org/10.1017/S0956796800000198>
- Luca Cardelli, Simone Martini, John C. Mitchell, and Andre Scedrov. 1991. An extension of system F with subtyping. In *Proceedings of the International Conference on Theoretical Aspects of Computer Software (TACS 1991), Sendai, Japan*, Takayasu Ito and Albert R. Meyer (Eds.). Springer, Berlin, Heidelberg, 750–770. https://doi.org/10.1007/3-540-54415-1_73
- Luca Cardelli and Peter Wegner. 1985. On Understanding Types, Data Abstraction, and Polymorphism. *Comput. Surveys* 17, 4 (Dec. 1985), 471–523. <https://doi.org/10.1145/6041.6042>
- Simon Castellan, Pierre Clairambault, and Peter Dybjer. 2015. Undecidability of Equality in the Free Locally Cartesian Closed Category. In *13th International Conference on Typed Lambda Calculi and Applications, (TLCA 2015), Warsaw, Poland (LIPIcs, Vol. 38)*, Thorsten Altenkirch (Ed.). Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl, Germany, 138–152. <https://doi.org/10.4230/LIPIcs.TLCA.2015.138>
- Adriana Compagnoni and Healfdene Goguen. 1999. Anti-Symmetry of Higher-Order Subtyping. In *Proceedings of the 13th International Workshop on Computer Science Logic (CSL 1999), 8th Annual Conference of the EACSL Madrid, Spain (LNCS, Vol. 1683)*, Jörg Flum and Mario Rodríguez-Artalejo (Eds.). Springer, Berlin, Heidelberg, 420–438. https://doi.org/10.1007/3-540-48168-0_30
- Adriana Compagnoni and Healfdene Goguen. 2003. Typed operational semantics for higher-order subtyping. *Information and Computation* 184, 2 (2003), 242–297. [https://doi.org/10.1016/S0890-5401\(03\)00062-2](https://doi.org/10.1016/S0890-5401(03)00062-2)
- Adriana B. Compagnoni. 1995. Decidability of higher-order subtyping with intersection types. In *Computer Science Logic, 8th International Workshop, (CSL 1994), Kazimierz, Poland, September 25–30, 1994, Selected Papers (LNCS, Vol. 933)*, Leszek Pacholski and Jerzy Tiuryn (Eds.). Springer, Berlin, Heidelberg, 46–60. <https://doi.org/10.1007/BFb0022246>
- Karl Cray. 1997. Foundations for the Implementation of Higher-order Subtyping. In *Proceedings of the Second ACM SIGPLAN International Conference on Functional Programming (ICFP 1997), Amsterdam, The Netherlands*. ACM, New York, NY, USA, 125–135. <https://doi.org/10.1145/258948.258961>
- Karl Cray. 2009. A Syntactic Account of Singleton Types via Hereditary Substitution. In *Proceedings of the Fourth International Workshop on Logical Frameworks and Meta-Languages, Theory and Practice (LFMTP 2009), Montreal, Quebec, Canada*. ACM, New York, NY, USA, 21–29. <https://doi.org/10.1145/1577824.1577829>
- Julien Cretin and Didier Rémy. 2014. System F with Coercion Constraints. In *Proceedings of the Joint Meeting of the Twenty-Third EACSL Annual Conference on Computer Science Logic (CSL 2014) and the Twenty-Ninth Annual ACM/IEEE Symposium on Logic in Computer Science (LICS 2014), Vienna, Austria*. ACM, New York, NY, USA, Article 34, 10 pages. <https://doi.org/10.1145/2603088.2603128>
- Pierre-Louis Curien and Giorgio Ghelli. 1992. Coherence of Subsumption, Minimum Typing and Type-checking in F_{\leq} . *Mathematical Structures in Computer Science* 2, 1 (March 1992), 55–91. <https://doi.org/10.1017/S0960129500001134>
- The Dotty Team. 2020. Scala 3 – A next-generation compiler for Scala – <http://dotty.epfl.ch>. Source code available from <https://github.com/lampepfl/dotty>.
- Paolo G. Giarrusso, Léo Stefanescu, Amin Timany, Lars Birkedal, and Robbert Krebbers. 2020. Scala Step-by-Step: Soundness for DOT with Step-Indexed Logical Relations in Iris. *PACMPL* 4, ICFP, Article 114 (Aug. 2020), 29 pages. <https://doi.org/10.1145/3408996>
- Jean-Yves Girard. 1972. *Interprétation fonctionnelle et élimination des coupures de l'arithmétique d'ordre supérieur*. Ph.D. Dissertation. Université Paris VII.

- Robert Harper and Daniel R. Licata. 2007. Mechanizing Metatheory in a Logical Framework. *Journal of Functional Programming* 17, 4-5 (July 2007), 613–673. <https://doi.org/10.1017/S0956796807006430>
- Robert Harper and Mark Lillibridge. 1994. A Type-theoretic Approach to Higher-order Modules with Sharing. In *Proceedings of the 21st ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages (POPL 1994)*, Portland, Oregon, USA. ACM, New York, NY, USA, 123–137. <https://doi.org/10.1145/174675.176927>
- Robert Harper and Frank Pfenning. 2005. On Equivalence and Canonical Forms in the LF Type Theory. *ACM Transactions on Computational Logic* 6, 1 (Jan. 2005), 61–101. <https://doi.org/10.1145/1042038.1042041>
- Jason Z. S. Hu and Ondřej Lhoták. 2019. Undecidability of $D_{<}$: And Its Decidable Fragments. *PACMPL* 4, POPL, Article 9 (Dec. 2019), 30 pages. <https://doi.org/10.1145/3371077>
- Chantal Keller and Thorsten Altenkirch. 2010. Hereditary Substitutions for Simple Types, Formalized. In *Proceedings of the Third ACM SIGPLAN Workshop on Mathematically Structured Functional Programming (MSFP 2010)*, Baltimore, Maryland, USA. ACM, New York, NY, USA, 3–10. <https://doi.org/10.1145/1863597.1863601>
- Daniel K. Lee, Karl Cray, and Robert Harper. 2007. Towards a Mechanized Metatheory of Standard ML. In *Proceedings of the 34th Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages (POPL 2007)*, Nice, France. ACM, New York, NY, USA, 173–184. <https://doi.org/10.1145/1190216.1190245>
- Adriaan Moors, Frank Piessens, and Martin Odersky. 2008a. Generics of a Higher Kind. In *Proceedings of the 23rd ACM SIGPLAN Conference on Object-oriented Programming Systems Languages and Applications (OOPSLA 2008)*, Nashville, TN, USA. ACM, New York, NY, USA, 423–438. <https://doi.org/10.1145/1449764.1449798>
- Adriaan Moors, Frank Piessens, and Martin Odersky. 2008b. Safe type-level abstraction in Scala. In *Proceedings of the International Workshop on Foundations of Object-Oriented Languages (FOOL 2008)*, San Francisco, CA, USA. 1–13. <https://www.cs.cmu.edu/~aldrich/FOOL/fool08/moors.pdf>
- Bengt Nordström, Kent Petersson, and Jan M. Smith. 1990. *Programming in Martin-Löf's type theory*. Vol. 200. Oxford University Press, Oxford, UK.
- Ulf Norell. 2007. *Towards a practical programming language based on dependent type theory*. Ph.D. Dissertation. Department of Computer Science and Engineering, Chalmers University of Technology, Göteborg, Sweden.
- Martin Odersky, Guillaume Martres, and Dmitry Petrashko. 2016. Implementing Higher-kinded Types in Dotty. In *Proceedings of the 7th ACM SIGPLAN Symposium on Scala (SCALA@SPLASH 2016)*, Amsterdam, Netherlands. ACM, New York, NY, USA, 51–60. <https://doi.org/10.1145/2998392.2998400>
- Lionel Parreaux, Aleksander Boruch-Gruszecki, and Paolo G. Giarrusso. 2019. Towards Improved GADT Reasoning in Scala. In *Proceedings of the Tenth ACM SIGPLAN Symposium on Scala (Scala 2019)*, London, United Kingdom. ACM, New York, NY, USA, 12–16. <https://doi.org/10.1145/3337932.3338813>
- Benjamin Pierce and Martin Steffen. 1997. Higher-order subtyping. *Theoretical Computer Science* 176, 1–2 (1997), 235–282. [https://doi.org/10.1016/S0304-3975\(96\)00096-5](https://doi.org/10.1016/S0304-3975(96)00096-5)
- Benjamin C. Pierce. 1992. Bounded Quantification is Undecidable. In *Proceedings of the 19th ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages (POPL 1992)*, Albuquerque, NM, USA. Association for Computing Machinery, New York, NY, USA, 305–315. <https://doi.org/10.1145/143165.143228>
- Benjamin C. Pierce. 2002. *Types and programming languages*. MIT Press, Cambridge, MA, USA.
- Marianna Rapoport and Ondřej Lhoták. 2019. A Path to DOT: Formalizing Fully Path-Dependent Types. *PACMPL* 3, OOPSLA, Article 145 (Oct. 2019), 29 pages. <https://doi.org/10.1145/3360571>
- Tiark Ropmf and Nada Amin. 2016. Type Soundness for Dependent Object Types (DOT). In *Proceedings of the 2016 ACM SIGPLAN International Conference on Object-Oriented Programming, Systems, Languages, and Applications (OOPSLA 2016)*, Amsterdam, Netherlands. ACM, New York, NY, USA, 624–641. <https://doi.org/10.1145/2983990.2984008>
- Gabriel Scherer and Didier Rémy. 2015. Full Reduction in the Face of Absurdity. In *Proceedings of the 24th European Symposium on Programming on Programming Languages and Systems (ESOP 2015)*, Held as Part of the European Joint Conferences on Theory and Practice of Software (ETAPS 2015), London, UK (LNCS, Vol. 9032), Jan Vitek (Ed.). Springer, Berlin, Heidelberg, 685–709. https://doi.org/10.1007/978-3-662-46669-8_28
- Christopher A. Stone and Robert Harper. 2000. Deciding Type Equivalence in a Language with Singleton Kinds. In *Proceedings of the 27th ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages (POPL 2000)*, Boston, MA, USA. ACM, New York, NY, USA, 214–227. <https://doi.org/10.1145/325694.325724>
- Sandro Stucki. 2017. *Higher-Order Subtyping with Type Intervals*. Ph.D. Dissertation. School of Computer and Communication Sciences, École polytechnique fédérale de Lausanne, Lausanne, Switzerland. <https://doi.org/10.5075/epfl-thesis-8014>
- Sandro Stucki and Paolo G. Giarrusso. 2021. *A Theory of Higher-order Subtyping with Type Intervals – Agda Formalization*. Zenodo. <https://doi.org/10.5281/zenodo.4775731>
- Kevin Watkins, Iliano Cervesato, Frank Pfenning, and David Walker. 2004. A Concurrent Logical Framework: The Propositional Fragment. In *International Workshop on Types for Proofs and Programs (TYPES 2003)*, Torino, Italy, April 30–May 4, 2003, Revised Selected Papers (LNCS, Vol. 3085), Stefano Berardi, Mario Coppo, and Ferruccio Damiani (Eds.). Springer, Berlin, Heidelberg, 355–377. https://doi.org/10.1007/978-3-540-24849-1_23

- Andrew K. Wright and Matthias Felleisen. 1994. A Syntactic Approach to Type Soundness. *Information and Computation* 115, 1 (Nov. 1994), 38–94. <https://doi.org/10.1006/inco.1994.1093>
- Yanpeng Yang and Bruno C. d. S. Oliveira. 2017. Unifying Typing and Subtyping. *PACMPL* 1, OOPSLA, Article 47 (Oct. 2017), 26 pages. <https://doi.org/10.1145/3133871>
- Jan Zwanenburg. 1999. Pure Type Systems with Subtyping. In *Proceedings of the 4th International Conference on Typed Lambda Calculi and Applications (TLCA 1999), L'Aquila, Italy*, Jean-Yves Girard (Ed.). LNCS, Vol. 1581. Springer, Berlin, Heidelberg, 381–396. https://doi.org/10.1007/3-540-48959-2_27