*R***-EQUIVALENCE ON REDUCTIVE GROUP SCHEMES**

PHILIPPE GILLE AND ANASTASIA STAVROVA

ABSTRACT. Let A be an equicharacteristic henselian regular local ring. Let k and K be the residue field and the fraction field of A. We show that for any reductive group scheme \mathfrak{G} over A there is a canonical isomorphism of Manin's R-equivalence class groups $\mathfrak{G}_K(K)/R \cong \mathfrak{G}_k(k)/R$. Our proof is based on extending the notion of R-equivalence from algebraic varieties over fields to schemes over commutative rings, and showing that the two canonical homomorphisms $\mathfrak{G}(A)/R \to \mathfrak{G}_k(k)/R$ and $\mathfrak{G}(A)/R \to \mathfrak{G}_K(K)/R$ are isomorphisms. If \mathfrak{G} is a torus or an isotropic simply connected semisimple group, the first isomorphism in fact holds without the assumption that A is regular, and the second one without the assumption that A is negative, if X is a connected smooth scheme over a field k, and \mathfrak{G} is a reductive X-group scheme belonging to one of the two classes mentioned above, then \mathfrak{G} being retract rational at the generic point of X implies that all fibers $\mathfrak{G}_x, x \in X$, are retract rational.

<u>Keywords</u>: reductive group scheme, algebraic torus, R-equivalence, A^1 -equivalence, Whitehead group, non-stable K_1 -functor.

MSC 2020: 20G15, 20G35, 19B99

Contents

1. Introduction	2
2. <i>R</i> -equivalence for schemes	5
2.1. Definition	5
2.2. Elementary properties	6
2.3. Retract rationality and <i>R</i> -equivalence	8
3. <i>R</i> -equivalence for reductive groups	11
3.1. <i>R</i> -equivalence as a birational invariant	11
3.2. The case of tori	13
3.3. Parabolic reduction	15
4. \mathbf{A}^{1} -equivalence and non-stable K_{1} -functors	16
4.1. \mathbf{A}^{1} -equivalence	16
4.2. Patching pairs and \mathbf{A}^1 -equivalence	17
4.3. Non stable K_1 -functor	20
4.4. Comparison of $K_1^{\mathfrak{G}}$, \mathbf{A}^1 -equivalence and <i>R</i> -equivalence	20
5. Passage to the field of fractions	22

Date: February 18, 2022.

The second author is supported by the Russian Science Foundation grant 19-71-30002.

PHILIPPE GILLE AND ANASTASIA STAVROVA

6. The case of simply connected semisimple isotropic groups	26
6.1. Coincidence of equivalence relations	26
6.2. The retract rational case	27
7. Behaviour for henselian pairs	31
7.1. The torus case	32
7.2. A generalization	32
7.3. The semisimple case	33
8. Specialization for R -equivalence	34
8.1. The case of tori	34
8.2. Reduction to the anisotropic case	35
8.3. The lifting map	36
8.4. The case of DVRs	37
8.5. Specialization in the equicharacteristic case	39
9. Appendices	41
9.1. The big Bruhat cell is a principal open subscheme.	41
9.2. Colliot-Thélène and Ojanguren method for functors in pointed sets.	41
9.3. Fields of representants for henselian regular local rings	43
References	43

1. INTRODUCTION

Yu. Manin [40, §14] introduced the notion of R-equivalence for points of algebraic varieties over a field. This notion has been used extensively in the study of reductive algebraic groups, e.g. [14, 15, 23, 2]. In the present paper, we propose a generalized definition of R-equivalence that is applicable to arbitrary schemes over an affine base and allows to extend several of the above-mentioned results to reductive group schemes in the sense of [64].

Among reductive groups, two classes play a fundamental role, the tori and the semisimple simply connected isotropic groups. In these two cases the *R*-equivalence class group G(k)/R of a reductive group *G* over a field *k* is already known to coincide with the value of a certain functor defined on the category of all commutative *k*-algebras, and even on all commutative rings *B* such that *G* is defined over *B*.

Namely, if G = T is a k-torus and

$$(1.1) 1 \to F \to P \to T \to 1$$

is a flasque resolution of T, then T(k)/R coincides with the first Galois (or étale) cohomology group $H^1_{\acute{e}t}(k,F)$ [14], and $H^1_{\acute{e}t}(-,F)$ is the functor of the above kind.

If G is a simply connected absolutely almost simple k-group having a proper parabolic k-subgroup, then G(k)/R coincides with the Whitehead group of G, which is the subject of the Kneser–Tits problem, and with the group of \mathbf{A}^1 -equivalence classes of k-points. Recall that the Whitehead group of G over k is defined as the quotient

 $\mathbf{2}$

of G(k) by the subgroup generated by the k-points of the unipotent radicals of all proper parabolic k-subgroups of G. In the setting of reductive groups over rings, the Whitehead group is also called a non-stable K_1 -functor, which is defined as follows.

Let *B* be a ring. If \mathfrak{G} is a reductive *B*-group scheme equipped with a parabolic *B*-subgroup \mathfrak{P} of unipotent radical $R_u(\mathfrak{P})$, we define the elementary subgroup $E^{\mathfrak{P}}(B)$ of $\mathfrak{G}(B)$ to be the subgroup generated by the $R_u(\mathfrak{P})(B)$ and $R_u(\mathfrak{P}^-)(B)$ where $\mathfrak{P}^$ is an opposite *B*-parabolic to \mathfrak{P} . We define the non stable K_1 -functor $K_1^{\mathfrak{G},\mathfrak{P}}(B) = \mathfrak{G}(B)/E^{\mathfrak{P}}(B)$ called also the Whitehead coset. We say that \mathfrak{G} has *B*-rank $\geq n$, if every normal semisimple *B*-subgroup of \mathfrak{G} contains $(\mathbf{G}_{m,B})^n$. If *B* is semilocal and \mathfrak{P} is minimal, or if the *B*-rank of \mathfrak{G} is ≥ 2 and \mathfrak{P} is strictly proper (i.e. \mathfrak{P} intersects properly every semisimple normal subgroup of *G*), then $E^{\mathfrak{P}}(B)$ is a normal subgroup independent of the specific choice of \mathfrak{P} [64, Exp. XXVI], [54].

A related, more universal construction is the 1st Karoubi-Villamayor K-functor, or the group of \mathbf{A}^1 -equivalence classes, denoted here by $\mathfrak{G}(B)/A^1$ where $A^1\mathfrak{G}(B)$ consists in the (normal) subgroup of $\mathfrak{G}(B)$ generated by the elements $g(0)g^{-1}(1)$ for g running over $\mathfrak{G}(B[t])$.

If G is semisimple simply connected over a field k and equipped with a strictly proper parabolic k-subgroup P, we know that the natural maps

$$K_1^{G,P}(k) \to G(k)/A^1 \to G(k)/R$$

are bijective [23]. In the present paper, we investigate to which extent such a result holds over the ring B, especially in the semilocal case and in the regular case.

Our first task is the extension of the notion of R-equivalence for rational points of algebraic varieties to integral points of a B-scheme in such a way that it is functorial with respect to ring homomorphisms. This is the matter of section 2; an advantage of R-equivalence is the nice functoriality with respect to fibrations.

In the subsequent sections we study the properties of R-equivalence on reductive group schemes. For tori over regular rings, the Colliot-Thélène and Sansuc computation of R-equivalence extend verbatim to the ring setting, see §3.2.

For non-toral reductive groups we obtain several results under the assumption that B is an equicharacteristic semilocal regular domain. Namely, we show in Theorem 6.5 that for a semisimple simply connected B-group \mathfrak{G} of B-rank ≥ 2 , the maps

$$K_1^{\mathfrak{G},\mathfrak{P}}(B) \to \mathfrak{G}(B)/A^1 \to \mathfrak{G}(B)/R$$

are isomorphisms; if the *B*-rank of \mathfrak{G} is only ≥ 1 , then the second map is an isomorphism (Theorem 6.2). In particular, this provides several new cases where $E^{\mathfrak{P}}(B) = \mathfrak{G}(B)$ holds (Cor. 6.9).

Let K be the fraction field of B. Another main result is the surjectivity of the map $\mathfrak{G}(B)/R \to \mathfrak{G}(K)/R$, assuming either that \mathfrak{G} is a reductive group of B-rank ≥ 1 , or that \mathfrak{G} has no parabolic subgroups over the residue fields of B (Theorem 5.4). If \mathfrak{G} is simply connected semisimple of B-rank ≥ 1 , then this map is an isomorphism (Theorem 6.2). This statement was previously known (for $\mathfrak{G}(B)/A^1$ instead of $\mathfrak{G}(B)/R$)

in the case where \mathfrak{G} is defined over an infinite perfect subfield of B and is of classical type, see [3, Corollary 4.3.6] and [46, Example 2.3].

As a corollary, we conclude that if \mathfrak{G} is a *B*-torus or a simply connected semisimple *B*-group of *B*-rank ≥ 1 , then \mathfrak{G} is retract rational over *B* if and only if \mathfrak{G}_K is retract rational over *K* (Proposition 3.5 and Theorem 6.8). In particular, if *X* is a connected smooth scheme over a field *k*, and \mathfrak{G} is a reductive *X*-group scheme belonging to one of the two classes mentioned above, then \mathfrak{G} being retract rational at the generic point of *X* implies that all fibers \mathfrak{G}_x , $x \in X$, are retract rational. This is reminiscent of the recent results on the rationality of fibers of smooth proper schemes over smooth curves [37, 49].

The assumption that B is equicharacteristic arises from the fact that we use a geometric construction developped by I. Panin for the proof of the Serre–Grothendieck conjecture for equicharacteristic semilocal regular rings [50, Theorem 2.5] (see also [52, 20]). Recently, K. Česnavičius partially generalized this construction to semilocal regular rings which are essentially smooth over a discrete valuation ring and proved the Serre–Grothendieck conjecture for quasi-split reductive groups over such rings in the unramified case [10]. We expect that in the future this approach will yield a similar extension of our results.

Another motivation for the present work was to deal with the specialization problem for *R*-equivalence [11, 6.1], [36], [22]. Let *A* be a henselian local domain of residue field *k* and fraction field *K*. Let \mathfrak{G} be a reductive *A*-group scheme and denote by $G = G \times_A k$ its closed fiber. We address the questions whether there exists a natural specialization homomorphism $\mathfrak{G}(K)/R \to G(k)/R$ and a lifting map $G(k)/R \to \mathfrak{G}(K)/R$. It makes sense to approach these questions using the generalized *R*-equivalence for \mathfrak{G} , since we may investigate whether the maps in the diagram

$$G(k)/R \longleftarrow \mathfrak{G}(A)/R \longrightarrow \mathfrak{G}(K)/R$$

are injective/surjective/bijective. In general, the only apriori evidence is the surjectivity of $\mathfrak{G}(A)/R \to G(k)/R$ which follows from the surjectivity of $\mathfrak{G}(A) \to G(k)$ (Hensel's lemma). We prove that if \mathfrak{G} is a torus or a simply connected semisimple group scheme equipped with a strictly proper parabolic A-subgroup, then the map $\mathfrak{G}(A)/R \to G(k)/R$ is an isomorphism (Proposition 8.1 and Theorem 8.6). In the case where A is a henselian regular local ring containing a field k_0 , we prove that for any reductive \mathfrak{G} there are two isomorphisms

$$G(k)/R \xleftarrow{\sim} \mathfrak{G}(A)/R \xrightarrow{\sim} G(K)/R$$

and in particular there is a well-defined specialization (resp. lifting) homomorphism (Theorem 8.14). Note that the recent results on the local-global principles over semiglobal fields [12] crucially use the existense of an (idependently constructed) specialization map for two-dimensional rings. Acknowledgments. We thank K. Cesnavičius for valuable comments in particular about fields of representants for henselian rings. We thank D. Izquierdo for useful conversations.

Notations and conventions.

We use mainly the terminology and notation of Grothendieck-Dieudonné [28, §9.4 and 9.6], which agrees with that of Demazure-Grothendieck used in [64, Exp. I.4]

Let S be a scheme and let \mathcal{E} be a quasi-coherent sheaf over S. For each morphism $f: T \to S$, we denote by $\mathcal{E}_T = f^*(\mathcal{E})$ the inverse image of \mathcal{E} by the morphism f. Recall that the S-scheme $\mathbf{V}(\mathcal{E}) = \operatorname{Spec}(\operatorname{Sym}^{\bullet}(\mathcal{E}))$ is affine over S and represents the S-functor $T \mapsto \operatorname{Hom}_{\mathcal{O}_T}(\mathcal{E}_T, \mathcal{O}_T)$ [28, 9.4.9].

We assume now that \mathcal{E} is locally free of finite rank and denote by \mathcal{E}^{\vee} its dual. In this case the affine *S*-scheme $\mathbf{V}(\mathcal{E})$ is of finite presentation (ibid, 9.4.11); also the *S*functor $T \mapsto H^0(T, \mathcal{E}_{(T)}) = \operatorname{Hom}_{\mathcal{O}_T}(\mathcal{O}_T, \mathcal{E}_T)$ is representable by the affine *S*-scheme $\mathbf{V}(\mathcal{E}^{\vee})$ which is also denoted by $\mathbf{W}(\mathcal{E})$ [64, I.4.6].

For scheme morphisms $Y \to X \to S$, we denote by $\prod_{X/S} (Y/X)$ the S-functor defined

by

$$\left(\prod_{X/S} (Y/X)\right)(T) = Y(X \times_S T)$$

for each S-scheme T. Recall that if $\prod_{X/S} (Y/X)$ is representable by an S-scheme, this

scheme is called the Weil restriction of Y to S.

If \mathfrak{G} is a S-group scheme locally of finite presentation, we denote by $H^1(S, \mathfrak{G})$ the set of isomorphism classes of sheaf \mathfrak{G} -torsors for the fppf topology.

2. R-Equivalence for schemes

2.1. **Definition.** Let *B* be a ring (unital, commutative). We denote by Σ the multiplicative subset of polynomials $P \in B[T]$ satisfying $P(0), P(1) \in B^{\times}$. Note that evaluation at 0 (and 1) extend from B[t] to the localization $B[t]_{\Sigma}$.

Let \mathcal{F} be a *B*-functor in sets. We say that two points $x_0, x_1 \in \mathcal{F}(B)$ are <u>directly</u> <u>*R*-equivalent</u> if there exists $x \in \mathcal{F}(B[t]_{\Sigma})$ such that $x_0 = x(0)$ and $x_1 = x(1)$. The *R*-equivalence on $\mathcal{F}(B)$ is the equivalence relation generated by this elementary relation.

Remarks 2.1. (a) If B is a field, then $B[t]_{\Sigma}$ is the semilocalization of B[t] at 0 and 1 so that the definition agrees with the classical definition.

(b) If B is a semilocal ring with maximal ideals $\mathfrak{m}_1, \ldots, \mathfrak{m}_r$, then $B[t]_{\Sigma}$ is the semilocalization of B[t] at the maximal ideals $\mathfrak{m}_1 B[t] + tB[t]$, $\mathfrak{m}_1 B[t] + (t-1)B[t]$, ..., $\mathfrak{m}_r B[t] + tB[t]$, $\mathfrak{m}_r B[t] + (t-1)B[t]$. In particular $B[t]_{\Sigma}$ is a semilocal ring.

(c) The most important case is for the *B*-functor of points $h_{\mathfrak{X}}$ of a *B*-scheme \mathfrak{X} . In this case we write $\mathfrak{X}(B)/R$ for $h_{\mathfrak{X}}(B)/R$.

(d) If the *B*-functor \mathcal{F} is locally of finite presentation (that is commutes with filtered direct limits), then two points $x_0, x_1 \in \mathcal{F}(B)$ are directly *R*-equivalent if there exists a polynomial $P \in B[t]$ and $x \in \mathcal{F}(B[t, \frac{1}{P}])$ such that $P(0), P(1) \in B^{\times}$ and $x_0 = x(0)$ and $x_1 = x(1)$. This applies in particular to the case of $h_{\mathfrak{X}}$ for a *B*-scheme \mathfrak{X} locally of finite presentation.

The important thing is the functoriality. If $B \to C$ is a morphism of rings, then the map $\mathcal{F}(B) \to \mathcal{F}(C)$ induces a map $\mathcal{F}(B)/R \to \mathcal{F}(C)/R$. We have also a product compatibility $(\mathfrak{X} \times_B \mathfrak{Y})(B)/R \xrightarrow{\sim} \mathfrak{X}(B)/R \times \mathfrak{Y}(B)/R$ for *B*-schemes $\mathfrak{X}, \mathfrak{Y}$.

If \mathfrak{G} is a *B*-group scheme (and more generally a *B*-functor in groups), then the *R*-equivalence is compatible with left/right translations by $\mathfrak{G}(B)$, also the subset $R\mathfrak{G}(B)$ of elements of $\mathfrak{G}(B)$ which are *R*-equivalent to 1 is a normal subgroup. It follows that the set $\mathfrak{G}(B)/R \cong \mathfrak{G}(B)/R\mathfrak{G}(B)$ is equipped with a natural group structure.

2.2. Elementary properties. We start with the homotopy property.

Lemma 2.2. Let \mathcal{F} be a *B*-functor.

(1) The map $\mathcal{F}(B)/R \to \mathcal{F}(B[u])/R$ is bijective.

(2) Assume that \mathcal{F} is a *B*-functor in groups. Then two points of $\mathcal{F}(B)$ which are *R*-equivalent are directly *R*-equivalent.

Proof. (1) The specialization at 0 provides a splitting of $B \to B[u]$, so that the map $\mathcal{F}(B)/R \to \mathcal{F}(B[u])/R$ is split injective. It is then enough to establish the surjectivity. Let $f \in \mathcal{F}(B[u])$. We put $x(u,t) = f(ut) \in \mathcal{F}(B[u,t])$ so that $x(u,0) = f(0)_{B[u]}$ and x(u,1) = f. In other words, f is directly R-equivalent to $f(0)_{B[u]}$ and we conclude that the map is surjective.

(2) We put $\mathcal{B} = B[t]$ and are given two elements $f, f' \in \mathcal{F}(B)$ which are *R*-equivalent. By induction on the length of the chain connecting f and f', we can assume that there exists $f_1 \in \mathcal{F}(B)$ which is directly *R*-equivalent to f and $f' = f_2$. Also by translation we can assume that f = 1. There exists $g(t), h(t) \in \mathcal{F}(\mathcal{B})$ such that g(0) = 1, $g(1) = f_1^1 = h(0)$ and $h(1) = f_2$. We put $f(t) = g(t)^{-1}h(1-t) \in \mathcal{F}(\mathcal{B})$. Then f(0) = 1 and $f(1) = f_2$ as desired.

Lemma 2.3. Let \mathcal{F} be a *B*-functor locally of finite presentation and consider a direct limit $B_{\infty} = \underline{\lim}_{\lambda \in \Lambda} B_{\lambda}$ of *B*-rings. Then the map $\underline{\lim}_{\lambda \in \Lambda} \mathcal{F}(B_{\lambda})/R \to \mathcal{F}(B_{\infty})/R$ is bijective.

Lemma 2.4. Let C be a locally free B-algebra of degree d. Let \mathcal{E} be a C-functor and consider the B-functor $\mathcal{F} = \prod_{C/B} \mathcal{E}$ defined by $\mathcal{F}(B') = \mathcal{E}(C \otimes_B B')$ for each B-algebra B'. Then the morphism $\mathcal{F}(B)/R \to \mathcal{E}(C)/R$ is an isomorphism.

Proof. We distinguish the multiplicative subsets Σ_B and Σ_C . The map $B[t]_{\Sigma_B} \otimes_B C \to C[t]_{\Sigma_C}$ induces a map $\mathcal{F}(B[t]_{\Sigma}) = \mathcal{E}(B[t]_{\Sigma_B} \otimes_B C) \to \mathcal{E}(C[t]_{\Sigma_C})$. We get then a morphism $\mathcal{F}(B)/R \to \mathcal{E}(C)/R$. We claim that $B[t]_{\Sigma_B} \otimes_B C \to C[t]_{\Sigma_C}$ is an isomorphism.

Since C is locally free over B of degree d, we can consider the norm map $N: C \to B$ as defined in [65, Tag 0BD2, 31.17.6]. It is well-known that there exists a polynomial map $N': C \to B$ such that N(c) = c N'(c) for each $c \in C$. Given $Q(t) \in C[t]$ such that $Q(1), Q(0) \in C^{\times}$, we have that $P(T) = N_{C/B}(Q(T))$ belongs to Σ so that Q(t) divides P(T). It follows that we have an isomorphism $C[t]_{\Sigma_B} \to C[t]_{\Sigma_C}$. Since $B[t]_{\Sigma} \otimes_{B[t]} C[t] \xrightarrow{\sim} C[t]_{\Sigma_B}$ [65, Tag 00DK, 9.11.15], we conclude that the map $B[t]_{\Sigma_B} \otimes_B C \to C[t]_{\Sigma_C}$ is an isomorphism. As counterpart we get that the map $\mathcal{F}(B)/R \to \mathcal{E}(C)/R$ is an isomorphism. \Box

Lemma 2.5. Let \mathfrak{X} be a *B*-scheme.

(1) Assume that $\mathfrak{X} = \operatorname{Spec}(B[\mathfrak{X}])$ is affine and let $\mathfrak{U} = \mathfrak{X}_f$ be a principal open subset of \mathfrak{X} where $f \in B[\mathfrak{X}]$. If two points $x_0, x_1 \in \mathfrak{U}(B)$ are directly *R*-equivalent in $\mathfrak{X}(B)$, then they are directly *R*-equivalent in $\mathfrak{U}(B)$.

(2) Assume that B is semilocal. Let \mathfrak{U} be an open B-subscheme of \mathfrak{X} . If two points $x_0, x_1 \in \mathfrak{X}(B)$ are directly R-equivalent in $\mathfrak{X}(B)$, then they are directly R-equivalent in $\mathfrak{U}(B)$.

(3) Let \mathfrak{G} be a *B*-group scheme and let \mathfrak{U} be an open *B*-subscheme of \mathfrak{G} . If \mathfrak{U} is a principal open subset or if *B* is semilocal, then the map $\mathfrak{U}(B)/R \to \mathfrak{G}(B)/R$ is injective.

Note that (3) was known in the field case under an assumption of unirationality [14, Prop. 11].

Proof. (1) Let $x_1, x_2 \in \mathfrak{U}(B)$ and let $x(t) \in \mathfrak{X}(B[t]_{\Sigma})$ such that $x(0) = x_0$ and $x(1) = x_1$. We consider the polynomial $P(t) = f(x(t)) \in B[t]_{\Sigma}$. Since $P(0) = f(x(0)) \in B^{\times}$ and $P(1) = f(x(1)) \in B^{\times}$, it follows that $P \in \Sigma$ hence $x(t) \in \mathfrak{U}(B[t]_{\Sigma})$. Thus x_0 and x_1 are directly R-equivalent in $\mathfrak{U}(B)$.

(2) Let $x(t) \in \mathfrak{X}(B[t]_{\Sigma})$ such that $x(0) = x_1$ and $x(1) = x_1$. Since $B[t]_{\Sigma}$ is a semilocal ring and the closed points of $\text{Spec}(B[t]_{\Sigma})$ map to points of \mathfrak{U} , it follows that $x(t) \in \mathfrak{U}(B[t]_{\Sigma})$.

(3) This follows from the fact that two points of $\mathfrak{G}(B)$ are *R*-equivalent if and only they are directly *R*-equivalent according to Lemma 2.2.(2).

Lemma 2.6. (1) Let \mathcal{L} be a finitely generated locally free B-module and consider the associated vector group scheme $\mathbf{W}(\mathcal{L})$. Let $\mathfrak{U} \subset \mathbf{W}(\mathcal{L})$ be an open subset of the affine space $\mathbf{W}(\mathcal{L})$. We assume that \mathfrak{U} is a principal open subset or that B is semilocal. Then any two points of $\mathfrak{U}(B)$ are directly R-equivalent. In particular if $\mathfrak{U}(B) \neq \emptyset$, we have $\mathfrak{U}(B)/R = \bullet$.

(2) Let \mathfrak{G} be an affine B-scheme of finite presentation such that $H^1(B, \mathfrak{G}) = 1$, $H^1(B[t]_{\Sigma}, \mathfrak{G}) = 1$ and $\mathfrak{G}(B)/R = 1$. Let $f : \mathfrak{Y} \to \mathfrak{X}$ be a morphism of B-schemes which is a \mathfrak{G} -torsor. Then the map $\mathfrak{Y}(B)/R \to \mathfrak{X}(B)/R$ is bijective.

(3) In (2), assume that \mathfrak{G} arises by successive extensions of vector group schemes. Then the map $\mathfrak{Y}(B)/R \to \mathfrak{X}(B)/R$ is bijective. (4) In (2), assume that \mathfrak{G} is a split B-torus and that $\operatorname{Pic}(B) = \operatorname{Pic}(B[t]_{\Sigma}) = 0$. Then the map $\mathfrak{Y}(B)/R \to \mathfrak{X}(B)/R$ is bijective.

(5) Assume that B is semilocal and that \mathfrak{T} is a quasitrivial B-torus and let $f: \mathfrak{Y} \to \mathfrak{X}$ be a morphism of B-schemes which is a \mathfrak{T} -torsor. Then the map $\mathfrak{Y}(B)/R \to \mathfrak{X}(B)/R$ is bijective.

Proof. (1) According to Lemma 2.5.(3), it is enough to show that two points of $\mathbf{W}(\mathcal{L})(B) = \mathcal{L}$ are *R*-equivalent. Let $x_0, x_1 \in \mathcal{L}$ and consider $x(t) = (1-t)x_0 + tx_1 \in \mathcal{L}$ $\mathcal{L} \otimes_B B[t] \subset \mathcal{L} \otimes_B B[t]_{\Sigma} = \mathbf{W}(\mathcal{L})(B[t]_{\Sigma})$. Since $x(0) = x_0$ and $x(1) = x_1$, we conclude that x_0 and x_1 are directly *R*-equivalent.

(2) Since $H^1(B, \mathfrak{G}) = 0$, it follows that the map $\mathfrak{Y}(B) \to \mathfrak{X}(B)$ is surjective and a fortiori the map $\mathfrak{Y}(B)/R \to \mathfrak{X}(B)/R$ is onto. For the injectivity, it is enough to prove that two points $y_0, y_1 \in \mathfrak{Y}(B)$ such that their images $x_0, x_1 \in \mathfrak{X}(B)$ are directly *R*-equivalent are *R*-equivalent. Our assumption is that there exists $x(t) \in \mathfrak{X}(B[t]_{\Sigma})$ such that $x(0) = x_0$ and $x(1) = x_1$. Since $H^1(B[t]_{\Sigma}, \mathfrak{G}) = 1$ by assumption, we can lift x(t) to some element $y(t) \in \mathfrak{Y}(B[t]_{\Sigma})$. Then $y_0 = y(0).g_0$ and $y_1 = y(1).g_1$ for (unique) elements g_0, g_1 of $\mathfrak{G}(B)$. By (1), g_0 and g_1 are *R*-equivalent to 1 which enables us to conclude that y_0 and y_1 are *R*-equivalent.

(3) By induction we can assume that \mathfrak{G} is a vector group scheme. In this case $H^1(C, \mathfrak{G}) = 1$ for each *B*-ring *C* and $\mathbf{G}(C)/R = 1$ according to (1). Hence part (2) of the statement applies.

(4) By induction by the rank we can assume that $\mathfrak{G} = \mathbb{G}_m$. We have $H^1(B, \mathbb{G}_m) =$ $\operatorname{Pic}(B) = 0$ and similarly for $H^1(B[t]_{\Sigma}, \mathbb{G}_m)$. Finally we have $\mathbb{G}_m(B)/R = 1$ according to (1) so that part (2) of the statement applies.

(5) follows from similar vanishing properties.

2.3. Retract rationality and *R*-equivalence. It is well-known that over a field, there is a close relation between retract rationality of algebraic varieties and the triviality of their *R*-equivalence class groups; see e.g. the survey [11]. We extend Saltman's definition [60] of retract rationality over fields to the setting of pointed B-schemes.

Definition 2.7. Let (\mathfrak{X}, x) be a pointed B-scheme. We say that (\mathfrak{X}, x) is

- (1) <u>B-rational</u> if (\mathfrak{X}, x) admits an open B-subscheme (\mathfrak{U}, x) such that (\mathfrak{U}, x) is *B*-isomorphic to an affine open subscheme of $(\mathbf{A}_{B}^{N}, 0)$.
- (2) stably B-rational if (\mathfrak{X}, x) admits an open B-subscheme (\mathfrak{U}, x) such that $(\mathfrak{U} \times_B \mathbf{A}_B^d, (x, 0))$ is B-rational for some $d \ge 0$.
- (3) retract *B*-rational if (\mathfrak{X}, x) admits an open *B*-subscheme (\mathfrak{U}, x) such that (\mathfrak{U}, x) is a \overline{B} -retract of an open subset of some $(\mathbf{A}_{B}^{N}, 0)$.

Lemma 2.8. Assume that B is semilocal with residue fields $\kappa_1, \ldots, \kappa_c$. Let (\mathfrak{U}, x) be a pointed B-scheme which is a retract of an open subset $(\mathfrak{V}, 0)$ of some $(\mathbf{A}_B^N, 0)$.

(1) There is an affine open (\mathfrak{U}', x) of (\mathfrak{U}, x) which is a retract of an affine open subset $(\mathfrak{V}', 0)$ of some $(\mathbf{A}_B^N, 0)$.

- (2) We have $\mathfrak{U}(B)/R = \bullet$.
- (3) The map $\mathfrak{U}(B) \to \mathfrak{U}(\kappa_1) \times \cdots \times \mathfrak{U}(\kappa_c)$ is onto.

Proof. (1) Since the map $\mathfrak{U} \to \mathfrak{V}$ is a closed immersion, it is enough to deal with the case of $(\mathfrak{V}, 0)$. According to [65, Tag 01ZU], there exists an open affine subset \mathfrak{V}' of \mathfrak{V} containing $0_{\kappa_1}, \ldots, 0_{\kappa_c}$. Then the zero section $\operatorname{Spec}(B) \to \mathfrak{V}$ factorizes through \mathfrak{V}' , so that $(\mathfrak{V}', 0)$ does the job.

(2) We have $\mathfrak{V}(B) \neq \emptyset$ and $\mathfrak{V}(B)/R = \bullet$ according to Lemma 2.6.(1). On the other hand, the functoriality of *R*-equivalence provides a section of the map $\mathfrak{U}(B)/R \to \mathfrak{V}(B)/R$. We conclude that $\mathfrak{U}(B)/R = 1$.

(3) Once again we can work with an open subset \mathfrak{V} of \mathbf{A}_B^N . Let $(v_i)_{i=1,\dots,c}$ be an element of $\mathfrak{V}(\kappa_1) \times \cdots \times \mathfrak{V}(\kappa_c)$. There exists $f \in B[t_1,\dots,t_n]$ such that $\mathfrak{V}_f = \mathbf{A}_{B,f}^N \subseteq \mathfrak{V}$ and $v_i \in \mathfrak{V}_f(\kappa_i)$ for $i = 1, \dots, c$. We can deal with \mathfrak{V}_f and observe that

$$\mathfrak{V}_f(B) = \left\{ b \in B^N \ | \ f(b) \in B^{\times} \right\}$$

maps onto $\prod_{i=1,\ldots,c} \mathfrak{V}_f(\kappa_i).$

Definition 2.9. We say that a B-scheme \mathfrak{X} satisfies the <u>lifting property</u> if for each semilocal B-ring C, the map

$$\mathfrak{X}(C) \to \prod_{\mathfrak{m}\in \max(C)} \mathfrak{X}(C/\mathfrak{m})$$

is onto, where $\max(C)$ denotes the maximal spectrum of C.

We extend Saltman's criterion of retract rationality [60, th. 3.9].

Proposition 2.10. We assume that B is semilocal with residue fields $\kappa_1, \ldots, \kappa_c$. Let (\mathfrak{X}, x) be a pointed affine finitely presented integral B-scheme. Then the following assertions are equivalent:

(i) (\mathfrak{X}, x) is retract *B*-rational;

(ii) (\mathfrak{X}, x) admits an open B-subscheme (\mathfrak{V}, x) which satisfies the lifting property.

Remarks 2.11. (a) Note that the assumption $X(B) \neq \emptyset$ implies that B is an integral ring.

(b) Assume that B is an integral ring of field of fractions K. Let \mathfrak{Y} be a flat affine B-scheme such that \mathfrak{Y}_K is integral. Then $B[\mathfrak{Y}]$ injects in $K[\mathfrak{Y}]$ so that \mathfrak{Y} is integral. In particular, if \mathfrak{G} is a smooth affine B-group scheme such that \mathfrak{G}_K is connected, then \mathfrak{G} is integral.

Proof. Let $\mathfrak{m}_1, \ldots, \mathfrak{m}_c$ be the maximal ideals of B and put $\kappa_i = B/\mathfrak{m}_i$ for $i = 1, \ldots, c$.

 $(i) \Longrightarrow (ii)$. By definition (\mathfrak{X}, x) admits an open *B*-subscheme (\mathfrak{U}, x) such that (\mathfrak{U}, x) is a *B*-retract of an open subset of some \mathbf{A}_B^N . We take $\mathfrak{V} = \mathfrak{U}$, it satisfies the lifting property according to Lemma 2.8.(3).

 $(ii) \implies (i)$. Up to replacing \mathfrak{V} by \mathfrak{X} , we can assume that \mathfrak{X} satisfies the lifting property. Furthermore an argument as in Lemma 2.8.(1) permits to assume that \mathfrak{X} is affine. We denote by $x_i \in \mathfrak{X}(\kappa_i)$ the image of x. We write $B[\mathfrak{X}] = B[t_1, \ldots, t_N]/\mathcal{P}$ for a prime ideal \mathcal{P} of $B[t_1, \ldots, t_N]$. We denote by $\eta : \operatorname{Spec}(\kappa(\mathfrak{X})) \to \mathbf{A}_B^N$ the generic point of \mathfrak{X} . We consider the semilocalization C of $B[t_1, \ldots, t_N]$ at the points η, x_1, \ldots, x_c of \mathbf{A}_B^N . Our assumption implies that the map

$$\mathfrak{X}(C) \to \mathfrak{X}(\kappa(\mathfrak{X})) \times \mathfrak{X}(\kappa_1) \times \cdots \times \mathfrak{X}(\kappa_c)$$

is onto. Let $y \in \mathfrak{X}(C)$ be a lifting of (η, x_1, \ldots, x_c) . Then y extends to a principal neighborhood $B[t_1, \ldots, t_N]_f$ of (η, x_1, \ldots, x_c) , i.e. there is a B-map $\phi : (\mathbf{A}_B^N)_f \to \mathfrak{X}$ which satisfies $\phi(\eta) = \eta$ and $\phi(x_i) = x_i$ for $i = 1, \ldots, c$. The composite $\mathfrak{X}_f \to (\mathbf{A}_B^N)_f \xrightarrow{\phi} \mathfrak{X}$ fixes the points η, x_1, \ldots, x_c . There exists then a function $g \in B[t_1, \ldots, t_N]$ such that $g(\eta) \in \kappa(\mathfrak{X})^{\times}, g(x_i) \in \kappa_i^{\times}$ for $i = 1, \ldots, c$ and the restriction

$$\mathfrak{X}_{fg} \to (\mathbf{A}_B^N)_{fg} \xrightarrow{\phi} \mathfrak{X}$$

is the canonical map. Thus the open subset (\mathfrak{X}_{fg}, x) of (\mathfrak{X}, x) is a *B*-retract of the principal open subscheme of $(\mathbf{A}_B^N)_{fg}$ of \mathbf{A}_B^N .

Remarks 2.12. (a) Under the assumptions of the proposition, it follows that the retract rationality property is of birational nature (with respect to our base point). Furthermore inspection of the proof shows that if (\mathfrak{X}, x) is *B*-retract rational, we can take \mathfrak{V} to be a principal open subset of \mathfrak{X} and it is a *B*-retract of a principal open subset of \mathfrak{A}_B^N .

(b) The direct implication $(i) \Longrightarrow (ii)$ does not require \mathfrak{X} to be integral.

Example 2.13. Let *B* be a semilocal ring having infinite residue fields $\kappa_1, \ldots, \kappa_c$. Let \mathfrak{G} be a reductive *B*-group scheme and let \mathfrak{T} be a maximal *B*-torus of \mathfrak{G} (such a torus exists according to Grothendieck's theorem [64, XIV.3.20 and footnote]). Let $\mathfrak{X} = \mathfrak{G}/\mathfrak{N}_{\mathfrak{G}}(\mathfrak{T})$ be its *B*-scheme of maximal tori. We claim that \mathfrak{X} satisfies the lifting property so that, if *B* is integral, (\mathfrak{X}, \bullet) is retract rational over *B* according to Proposition 2.10. It is enough to show that the map $\mathfrak{X}(B) \to \prod_{i=1,\ldots,c} \mathfrak{X}(\kappa_i)$ is onto. Let T_i be a maximal κ_i -torus of \mathfrak{G}_{κ_i} for $i = 1, \ldots, c$. Since κ_i is infinite, there exists $X_i \in \operatorname{Lie}(\mathcal{T}_i)(\kappa_i) \subset \operatorname{Lie}(\mathfrak{G})(\kappa_i)$ such that $T_i = C_{\mathfrak{G}\kappa_i}(X_i)$ [64, XIV.5.1]. We pick a lift $X \in \operatorname{Lie}(\mathfrak{G})(R)$ of the X'_i s. Then $\mathfrak{T} = C_{\mathfrak{G}}(X)$ is a maximal *B*-torus of \mathfrak{G} which lift the T_i 's. By inspection of the argument we can actually assume only that $\sharp_{\kappa_i} \ge \dim_{\kappa_i}(\mathfrak{G}_{\kappa_i})$ by using [4, Thm. 1].

Proposition 2.14. We assume that B is semilocal. Let \mathfrak{G} be a B-group scheme with connected geometric fibers such that

(i) $(\mathfrak{G}, 1)$ is retract *B*-rational;

(ii) $\mathfrak{G}(\kappa)$ is dense in \mathfrak{G}_{κ} for each residue field κ of a maximal ideal of B. Then $\mathfrak{G}(B)/R = 1$.

Note that (ii) is satisfied if \mathfrak{G} is reductive and if B has infinite residue fields.

Proof. Let $\mathfrak{m}_1, \ldots, \mathfrak{m}_c$ be the maximal ideals of B and put $\kappa_i = B/\mathfrak{m}_i$ for $i = 1, \ldots, c$. The algebraic groups \mathfrak{G}_{κ_i} are then retract rational and $\mathfrak{G}(\kappa_i)$ is dense in \mathfrak{G}_{κ_i} for $i = 1, \ldots, c$.

Let $(\mathfrak{U}, 1)$ be an open subset of $(\mathfrak{G}, 1)$ which is a *B*-retract of some open of \mathbf{A}_B^N . Since $\mathfrak{U}(B)$ maps onto $\mathfrak{U}(\kappa_1) \times \cdots \times \mathfrak{U}(\kappa_c)$ (Lemma 2.8.(3)) and since $\mathfrak{U}(\kappa_i)$ is dense in $\mathfrak{G}(\kappa_i)$ by assumption, it follows that $\mathfrak{U}(B)$ is *B*-dense in \mathfrak{G} . In particular, there exist $u_1, \ldots, u_s \in \mathfrak{U}(B)$ such that $\mathfrak{G} = u_1 \mathfrak{U} \cup \cdots \cup u_s \mathfrak{U}$ so that $\mathfrak{U}(B)$ generates $\mathfrak{G}(B)$ as a group. Lemma 2.8.(2) shows that $\mathfrak{U}(B)/R = 1$. We conclude that $\mathfrak{G}(B)/R = 1$. \Box

Remark 2.15. Proposition 2.14 applies to quasitrivial tori so that it is coherent with Lemma 2.4.

3. R-Equivalence for reductive groups

3.1. *R*-equivalence as a birational invariant. The following statement generalizes [14, Prop. 11].

Proposition 3.1. Assume that B is semilocal with infinite residue fields. Let \mathfrak{G} be a B-group scheme and let $f : (\mathfrak{V}, v_0) \to (\mathfrak{G}, 1)$ be a B-morphism of pointed B-schemes such that (\mathfrak{V}, v) is an open subset of some $(\mathbf{A}_B^n, 0)$ and such that $f_{B/\mathfrak{m}}$ is dominant for each maximal ideal \mathfrak{m} of B. Let $(\mathfrak{U}, 1)$ be an open neighborhood of $(\mathfrak{G}, 1)$.

(1) We have $f(\mathfrak{V}(B)) \cdot \mathfrak{U}(B) = \mathfrak{G}(B)$.

(2) The map $\mathfrak{U}(B)/R \to \mathfrak{G}(B)/R$ is bijective.

Proof. Let $\mathfrak{m}_1, \ldots, \mathfrak{m}_c$ be the maximal ideals of B.

(1) From the proof of [14, Prop. 11], we have $f(\mathfrak{V}(B/\mathfrak{m}_i)) \cdot \mathfrak{U}(B/\mathfrak{m}_i) = \mathfrak{G}(B/\mathfrak{m}_i)$ for $i = 1, \ldots, c$. We are given $g \in \mathfrak{G}(B)$ and denote by g_i its reduction to $\mathfrak{G}(B/\mathfrak{m}_i)$ for $i = 1, \ldots, c$, then $g_i = f(v_i) u_i$ for some $v_i \in \mathfrak{V}(B/\mathfrak{m}_i)$ and $u_i \in \mathfrak{U}(B/\mathfrak{m}_i)$. Let $v \in \mathfrak{V}(B)$ be a common lift of the elements v_i , then $f(v)^{-1}g \in \mathfrak{G}(B)$ belongs to $\mathfrak{U}(B)$. (2) The surjectivity follows from (1) and the fact $\mathfrak{V}(B)/R = 1$ established in Lemma 2.6.(1). On the other hand, the injectivity has been proven in Lemma 2.5.(3).

We say that a *B*-group scheme *G* is *B*-linear, if for some $N \ge 1$ there is a closed embedding of *B*-group schemes $G \to \operatorname{GL}_{N,B}$.

Lemma 3.2. Assume that B is semilocal with infinite residue fields $\kappa_1, \ldots, \kappa_c$. Let \mathfrak{G} be a reductive B-group scheme.

(1) There exist maximal B-tori $\mathfrak{T}_1, \ldots, \mathfrak{T}_n$ of \mathfrak{G} such that the product map $\psi : \mathfrak{T}_1 \times \cdots \times \mathfrak{T}_n \to \mathfrak{G}$ satisfies the following property: ψ_{κ_j} is smooth at the origin for each $j = 1, \ldots, c$. Furthermore, the submodules $\operatorname{Lie}(\mathfrak{T}_i)(B)$ together generate $\operatorname{Lie}(\mathfrak{G})(B)$ as a B-module.

(2) Assume furthermore that \mathfrak{G} is B-linear. Then there exists a quasi-trivial Btorus \mathfrak{Q} and a B-morphism of pointed B-schemes $f : (\mathfrak{Q}, 1) \to (\mathfrak{G}, 1)$ such that f_{κ_j} is smooth at the origin for each $j = 1, \ldots, c$.

Proof. (1) We start with the case of an infinite field k and of a reductive k-group G. We know that G(k) is Zariski dense in G. Let T be a maximal k-torus of G and let $1 = g_1, g_2, \ldots, g_n$ be elements of G(k) such that Lie(G) is generated by the $g_i \text{Lie}(T)(k)$'s. We consider the map of B-schemes

 $\gamma:\mathfrak{T}^n \longrightarrow \mathfrak{G}$

 $(t_1,\ldots,t_n) \qquad \mapsto \qquad {}^{g_1}t_1\ldots {}^{g_n}t_n.$

Its differential at 1 is

 $d\gamma_{1_k}: \mathrm{Lie}(T)(k)^n \longrightarrow \mathrm{Lie}(G)(k)$

 $(X_1,\ldots,X_n) \qquad \mapsto \qquad {}^{g_1}X_1 + \cdots + {}^{g_n}X_n$

which is onto by construction. We put $T_i = {}^{g_i}T$ for i = 1, ..., n and observe that the product map $\psi : T_1 \times_k \cdots \times_k T_n \to G$ is smooth at 1. In this construction we are free to add more factors.

In the general case, we fix *n* large enough and maximal κ_j -tori $T_{1,j}, \ldots, T_{n,j}$ such that the product map $\psi_i : T_{1,j} \times_{\kappa_j} \cdots \times_{\kappa_j} T_{n,j} \to \mathfrak{G}_{\kappa_j}$ is smooth at 1 for $j = 1, \ldots, c$. Example 2.13 shows that there exists a maximal *B*-torus \mathfrak{T}_i which lifts the $T_{i,j}$'s for $i = 1, \ldots, n$. Then the product map $\psi : \mathfrak{T}_1 \times_B \cdots \times_k \mathfrak{T}_n \to \mathfrak{G}$ satisfies the desired requirements. Nakayama's lemma implies that the $\operatorname{Lie}(\mathfrak{T}_i)(B)$'s generate $\operatorname{Lie}(\mathfrak{G})(B)$ as a *B*-module.

(2) We assume that \mathfrak{G} is linear so that the \mathfrak{T}_i 's are isotrivial according to [27, Cor. 5.1]. Then by [15, Prop. 1.3] there exist flasque resolutions $1 \to \mathfrak{S}_i \to \mathfrak{Q}_i \xrightarrow{q_i} \mathfrak{T}_i \to 1$ of \mathfrak{T}_i where \mathfrak{Q}_i is a quasi-trivial *B*-torus and \mathfrak{S}_i is a flasque *B*-torus for i = 1, ..., n. We consider the map

 $f: \mathfrak{Q}_1 \times_B \cdots \times_B \mathfrak{Q}_n \quad \to \quad \mathfrak{G}$

 $(v_1,\ldots,v_n) \qquad \mapsto \qquad q_1(v_1)\ldots q_n(v_n).$

Since the q_i 's are smooth, $f = \psi \circ (q_1, \ldots, q_n)$ satisfies the desired requirements. \Box

Putting together Lemma 3.2 and Proposition 3.1 leads to the following fact.

Corollary 3.3. Assume that B is semilocal with infinite residue fields. Let \mathfrak{G} be a B-linear reductive B-group scheme.

(1) Let $f : (\mathfrak{Q}, 1) \to (\mathfrak{G}, 1)$ be the morphism constructed in Lemma 3.2.(2). Then $f(\mathfrak{Q}(B)) \cdot \mathfrak{U}(B) = \mathfrak{G}(B)$.

(2) Let $(\mathfrak{U}, 1)$ be an arbitrary open subset of $(\mathfrak{G}, 1)$. The map $\mathfrak{U}(B)/R \to \mathfrak{G}(B)/R$ is bijective.

3.2. The case of tori. Let B be a commutative ring such that the connected components of Spec(B) are open (e.g. B is Noetherian or semilocal). Let \mathfrak{T} be an isotrivial B-torus. According to [15, Prop. 1.3], there exists a flasque resolution

$$1 \to \mathfrak{S} \to \mathfrak{Q} \xrightarrow{\pi} \mathfrak{T} \to 1,$$

that is an exact sequence of B-tori where \mathfrak{Q} is a quasitrivial B-torus and \mathfrak{S} is a flasque B-torus. The following statement generalizes the corresponding result over fields due to Colliot-Thélène and Sansuc [14, Thm. 3.1].

Proposition 3.4. Assume additionally that B is a regular integral domain. We have $\pi(\mathfrak{Q}(B)) = R\mathfrak{T}(B)$ and the characteristic map $\mathfrak{T}(B) \to H^1(B, \mathfrak{S})$ induces an isomorphism

$$\mathfrak{T}(B)/R \xrightarrow{\sim} \ker (H^1(B,\mathfrak{S}) \to H^1(B,\mathfrak{Q})).$$

In particular, if B is a regular semilocal domain, we have an isomorphism $\mathfrak{T}(B)/R \xrightarrow{\sim} H^1(B,\mathfrak{S})$.

Proof. According to Lemma 2.6.(1) we have $\mathbb{G}_m(B)/R = 1$. Then Lemma 2.4 shows that $R\mathfrak{Q}(B) = \mathfrak{Q}(B)$. Hence the inclusion $\pi(\mathfrak{Q}(B)) \subseteq R\mathfrak{T}(B)$. For the converse, it is enough to show that a point $x \in \mathfrak{T}(B)$ which is directly *R*-equivalent to 1 belongs to $\pi(\mathfrak{Q}(R))$. By definition, there exists a polynomial $P \in B[t]$ such that $P(0), P(1) \in B^{\times}$ and $x(t) \in \mathfrak{T}(B[t, 1/P])$ satisfying x(0) = 1 and x(1) = x. We consider the obstruction $\delta(x(t)) \in H^1(B[t, 1/P], \mathfrak{S})$. Since \mathfrak{S} is flasque and *B* is a regular domain, the map

$$H^1(B,\mathfrak{S}) \to H^1(B[t,1/P],\mathfrak{S})$$

is onto by [15, Cor. 2.6]. It follows that $\delta(x(t)) = \delta(x)(0) = \delta(x(0)) = 1$ so that x(t) belongs to the image of $\pi : \mathfrak{Q}(B[t, 1/P]) \to \mathfrak{T}(B[t, 1/P])$. Thus x = x(1) belongs to $\pi(\mathfrak{Q}(R))$. as desired.

Using the above result, we extend Colliot-Thélène and Sansuc's criterion of retract rationality, see [15, Prop. 7.4] and [44, Prop. 3.3].

Proposition 3.5. Let B be a semilocal ring and let \mathfrak{T} be an isotrivial B-torus. Let $1 \to \mathfrak{S} \to \mathfrak{Q} \xrightarrow{\pi} \mathfrak{T} \to 1$ be a flasque resolution.

(1) We consider the following assertions:

(i) \mathfrak{S} is an invertible B-torus (i.e. a direct summand of a quasitrivial B-torus);

(ii) there exists an open subset $(\mathfrak{U}, 1)$ of $(\mathfrak{T}, 1)$ such that $\pi^{-1}(\mathfrak{U}) \cong \mathfrak{S} \times_B \mathfrak{U}$;

(iii) the pointed B-scheme $(\mathfrak{T}, 1)$ is retract rational;

(iv) \mathfrak{T} is R-trivial on semilocal rings, that is $\mathfrak{T}(C)/R = 1$ for each semilocal B-ring C;

(iv') \mathfrak{T} is R-trivial on fields, that is $\mathfrak{T}(F)/R = 1$ for each B-field F.

(v) \mathfrak{T} satisfies the lifting property.

Then we have the implications $(i) \implies (ii) \implies (iii) \implies (iv) \implies (iv') \implies (v)$. Furthermore if B is integral, we have the equivalences $(iii) \iff (iv) \iff (iv') \iff (v)$.

(2) We assume furthermore that B is a normal domain of fraction field K. We consider the following assertions:

(vi) \mathfrak{S}_K is an invertible K-torus;

(vii) \mathfrak{T}_K is R-trivial on semilocal rings, that is, $\mathfrak{T}(A)/R = 1$ for each semilocal K-ring A;

(vii') \mathfrak{T}_K is R-trivial on fields, that is, $\mathfrak{T}(F)/R = 1$ for each K-field F;

(viii) the pointed K-scheme $(\mathfrak{T}_K, 1)$ is retract rational.

Then the assertions (i), (ii), (iii), (iv), (iv), (v), (v), (vi), (vii), (vii') and (viii) are equivalent.

Proof. Let $\mathfrak{m}_1, \ldots, \mathfrak{m}_c$ be the maximal ideals of B and put $\kappa_i = B/\mathfrak{m}_i$ for $i = 1, \ldots, c$.

(1) (i) \Longrightarrow (ii). Let C be the semilocal ring of \mathfrak{T} at the points $1_{\kappa_1}, \ldots, 1_{\kappa_c}$ of \mathfrak{T} . Since \mathfrak{S} is invertible, we have $H^1(C, \mathfrak{S}) = 1$. In particular the \mathfrak{S} -torsor $\pi : \mathfrak{Q} \to \mathfrak{T}$ admits a splitting $s : \operatorname{Spec}(C) \to \mathfrak{Q}$. It follows that there exists an open neighborhood ($\mathfrak{U}, 1$) of \mathfrak{T} such that the \mathfrak{S} -torsor $\pi : \mathfrak{Q} \to \mathfrak{T}$ admits a splitting $s : \mathfrak{U} \to \mathfrak{Q}$.

 $(ii) \Longrightarrow (iii)$. We are given an open neighborhood $(\mathfrak{U}, 1)$ of $(\mathfrak{T}, 1)$ such that $\pi^{-1}(\mathfrak{U}) \cong \mathfrak{S} \times_B \mathfrak{U}$. Thus $(\mathfrak{U}, 1)$ is a retract of $\pi^{-1}(\mathfrak{U})$ which is open in some affine *B*-space, so that $(\mathfrak{T}, 1)$ is retract rational.


 $(iii) \Longrightarrow (iv)$. Let $\kappa_1, \ldots, \kappa_c$ the residue fields of the maximal ideals of B. If all κ_i 's are infinite, Proposition 2.14 shows that \mathfrak{T} is R-trivial. For the general case, it is enough to show that $\mathfrak{T}(B)/R = 1$. We may assume that $\kappa_1, \ldots, \kappa_b$ are finite fields and that $\kappa_{b+1}, \ldots, \kappa_c$ are infinite.

The maps $\mathfrak{Q}(\kappa_i) \to \mathfrak{T}(\kappa_i)$ are onto for $i = 1, \ldots, b$ (a finite field is of cohomological dimension 1) so that there exists a finite subset Θ of $\mathfrak{Q}(B)$ mapping onto $\prod_{i=1,\ldots,b} \mathfrak{T}(\kappa_i)$.

Let $(\mathfrak{U}, 1)$ be an open subset of $(\mathfrak{G}, 1)$ which is a *B*-retract of some open of $(\mathbf{A}_B^N, 0)$. Reasoning as in the proof of Proposition 2.14, we observe that $\mathfrak{T} = \mathfrak{U}(\mathfrak{U}(B)\pi(\Theta))$ so that $\mathfrak{T}(B)/R = 1$.

 $(iv) \Longrightarrow (iv')$. Obvious.

 $(iv') \Longrightarrow (v)$. We assume that \mathfrak{T} is *R*-trivial on fields. It enough to show that $\mathfrak{T}(B)$ maps onto $\mathfrak{T}(\kappa_1) \times \cdots \times \mathfrak{T}(\kappa_c)$. We consider the commutative diagram



The left vertical map is onto since \mathfrak{Q} satisfies the lifting property and the bottom horizontal map is onto by Proposition 3.4, since $\mathfrak{T}(\kappa_i)/R = 1$. Thus the right vertical map is onto.

Finally, if B is integral, then \mathfrak{T} is an integral scheme according to Remark 2.11.(b). Proposition 2.10 shows that (v) implies (iii).

(2) Let B' be a Galois connected cover of B which splits \mathfrak{T} and \mathfrak{S} . Let Γ be its Galois group. Then B' is a normal ring and its fraction field K' is a Galois extension of K of group Γ . According to Lemma [14, Lemme 2, (vi)], (i) (resp. (vi)) is equivalent to saying that the Γ -module $\mathfrak{S}(B')$ (resp. $\mathfrak{S}(K')$) is invertible. Since $\mathfrak{S}(B') = \mathfrak{S}(K')$ we get the equivalence $(i) \iff (vi)$. The statement over fields [15, Prop. 7.4] provides the equivalences $(vi) \iff (vii') \iff (viii)$. Taking into account the first part of the Proposition and the obvious implications, we have the following picture

Thus the assertions (i), (ii), (ii), (iv), (iv), (v), (v), (vi), (vii), (vii') and (viii) are equivalent.

3.3. **Parabolic reduction.** Let *B* be a ring and let \mathfrak{G} be a reductive *B*-group scheme. Let \mathfrak{P} be a parabolic *B*-subgroup of \mathfrak{G} together with an opposite parabolic *B*-subgroup \mathfrak{P}^- . We know that $\mathfrak{L} = \mathfrak{P} \times_{\mathfrak{G}} \mathfrak{P}^-$ is a Levi subgroup of \mathfrak{P} . We consider the big Bruhat cell

$$\Omega := R_u(\mathfrak{P}^-) \times_B \mathfrak{L} \times R_u(\mathfrak{P}^-) \subseteq \mathfrak{G}$$

Lemma 3.6. We have

$$\mathfrak{L}(B)/R \xrightarrow{\sim} \mathfrak{P}(B)/R \hookrightarrow \mathfrak{G}(B)/R$$

Proof. The left isomorphism follows from Lemma 2.6.(3). According to Lemma 9.1, the big cell Ω is a principal open subset of \mathfrak{G} so $\Omega(B)/R$ injects in $\mathfrak{G}(B)/R$ according to Lemma 2.5.(1). Since $R_u(\mathfrak{P})$ and $R_u(\mathfrak{P}^-)$ are extensions of vector group schemes, the map $\Omega(B)/R \to \mathfrak{P}(B)/R$ is bijective by Lemma 2.6.(3), hence $\mathfrak{P}(B)/R$ injects into $\mathfrak{G}(B)/R$. According to [24, Th. 7.3.1], there exists a homomorphism $\lambda : \mathbb{G}_{m,B} \to \mathfrak{G}$ such that $\mathfrak{P} = \mathfrak{P}_{\mathfrak{G}}(\lambda)$ and $\mathfrak{L} = \mathfrak{Z}_{\mathfrak{G}}(\lambda)$.

Lemma 3.7. We assume that there exists a central split B-subtorus \mathfrak{S} of \mathfrak{L} which factorizes λ . Assume that $\mathfrak{G}(B)$ is generated by $\mathfrak{P}(B)$ and $\mathfrak{P}^-(B)$ and that $\operatorname{Pic}(B) = \operatorname{Pic}(B[t]_{\Sigma}) = 0$. Then we have isomorphisms

$$\mathfrak{G}(B)/R \xleftarrow{\sim} \mathfrak{L}(B)/R \xrightarrow{\sim} (\mathfrak{L}/\mathfrak{S})(B)/R.$$

Proof. The assumption implies that $\mathfrak{L}(B)$ generates $\mathfrak{G}(B)/R$ so that the injective map $\mathfrak{L}(B)/R \to \mathfrak{G}(B)/R$ is onto hence an isomorphism. The right handside homomorphism follows of Lemma 2.6.(4).

Corollary 3.8. We assume that B is semilocal connected. Let \mathfrak{S} be the central maximal split B-subtorus \mathfrak{S} of \mathfrak{L} (as defined in [64, XXVI.7.1]). Then we have isomorphisms

$$\mathfrak{G}(B)/R \xleftarrow{\sim} \mathfrak{L}(B)/R \xrightarrow{\sim} (\mathfrak{L}/\mathfrak{S})(B)/R.$$

Proof. The second condition of Lemma 3.7 is satisfied [64, XXVI.5.2]. Also we have $\operatorname{Pic}(B) = \operatorname{Pic}(B[t]_{\Sigma}) = 0$ since B and $B[t]_{\Sigma}$ are semilocal rings. Thus Lemma 3.7 applies.

4. \mathbf{A}^1 -EQUIVALENCE AND NON-STABLE K_1 -FUNCTORS

4.1. A¹-equivalence. Let *B* be an arbitrary (unital, commutative) ring. Let \mathcal{F} be a *B*-functor in sets. We say that two points $x_0, x_1 \in \mathcal{F}(B)$ are directly A¹-equivalent if there exists $x \in \mathcal{F}(B[t])$ such that $x_0 = x(0)$ and $x_1 = x(1)$. The (naive) A¹-equivalence on $\mathcal{F}(B)$ is the equivalence relation generated by this relation.

Let G be a B-group scheme. We denote the equivalence class of $1 \in G(B)$ by $A^1G(B)$ and the group of \mathbf{A}^1 -equivalence classes by

$$G(B)/A^1 = G(B)/A^1G(B).$$

This group is functorial in B, and the functor $G(-)/A^1$ on the category of B-schemes is sometimes called the 1st Karoubi-Villamayor K-theory functor corresponding to G, and denoted by $KV_1^G(B)$ [34, 3].

Clearly, for any ring B we have a canonical surjection

$$G(B)/A^1 \longrightarrow G(B)/R.$$

The analog of Lemma 2.2 is true for \mathbf{A}^1 -equivalence. In particular, two points $g_0, g_1 \in G(B)$ are \mathbf{A}^1 -equivalent if and only if they are directly \mathbf{A}^1 -equivalent.

4.2. Patching pairs and A¹-equivalence. Let $R \to R'$ be a morphism of rings and let $f \in R$.

We say that that $(R \to R', f)$ is a *patching pair* if R' is flat over R and $R/fR \xrightarrow{\sim} R'/fR$. The other equivalent terminology is to say that

$$(4.1) \qquad \qquad \begin{array}{c} R \longrightarrow R_f \\ \downarrow \qquad \qquad \downarrow \\ R' \longrightarrow R'_f \end{array}$$

is a patching diagram. In this case, there is an equivalence of categories between the category of R-modules and the category of glueing data (M', M_1, α_1) where M' is an R'-module, M_1 an R_f -module and $\alpha_1 : M' \otimes_{R'} R'_f \xrightarrow{\sim} M_1 \otimes_{R_f} R'_f$ [65, Tag 05ES]. Note that this notion of a patching diagram is less restrictive than the one used by Colliot-Thélène and Ojanguren in [13, §1].

Examples 4.1. (a) (Zariski patching) Let $g \in R$ such R = fR + gR. Then $(R \to R_g, f)$ is a patching pair.

(b) Assume that R is noetherian. If $\widehat{R} = \varprojlim R/f^n R$, then $(R \to \widehat{R}, f)$ is a patching pair according to [65, Tags 00MB, 05GG].

(c) Assume that $R = k[[x_1, \ldots, x_n]]$ is a ring of formal power series over a field and let h be a monic Weierstrass polynomial of R[x] of degree ≥ 1 . Then (R[x], R[[x]], h) is a patching pair, see [5, page 803].

We recall that $(R \to R', f)$ is a glueing pair if $R/f^n R \xrightarrow{\sim} R'/f^n R'$ for each $n \ge 1$ and if the sequence

$$(4.2) 0 \to R \to R_f \oplus R' \xrightarrow{\gamma} R'_f \to 0$$

is exact where $\gamma(x, y) = x - y$ [65, Tag 01FQ].

Examples 4.2. (a) A patching pair is a glueing pair, we have $R/f^n R \xrightarrow{\sim} R'/f^n R'$ for all $n \ge 1$ [65, Tag 05E9] and the complex (4.2) is exact at 0, R and $R_f \oplus R'$ [65, Tag 05EK]. Since the map γ is surjective, the complex is exact.

(b) If f is a non zero divisor in R and $\widehat{R} = \varprojlim R/f^n R$, then $(R \to \widehat{R}, f)$ is a glueing pair [65, Tag 0BNS].

If $(R \to R', f)$ is a glueing pair, the Beauville-Laszlo theorem provides an equivalence of categories between the category of flat *R*-modules and the category of glueing data (M', M_1, α_1) where M' is a flat R'-module, M_1 a flat R_f -module and $\alpha_1 : M' \otimes_{R'} R'_f \xrightarrow{\sim} M_1 \otimes_{R_f} R'_f$ [65, Tags 0BP2, 0BP7 and 0BNX]. In particular we can patch torsors under an affine flat *R*-group scheme *G* in this setting, this means that the base change induces an equivalence from the category of *G*-torsors to that of triples (T, T', ι) where *T* is a *G*-torsor over $\operatorname{Spec}(R_f), T'$ a *G*-torsor over $\operatorname{Spec}(R')$ and $\iota: T \times_{R_f} R'_f \xrightarrow{\sim} T' \times_{R'} R'_f$ an isomorphism of *G*-torsors over $\operatorname{Spec}(R'_f)$, see [6, lemma 2.2.10]. This is a generalization of [13, proposition 2.6]. More specifically, there is an exact sequence of pointed sets

(4.3)
$$1 \to G(R') \setminus G(R'_f) / G(R_f) \to H^1(R,G) \to H^1(R',G) \times H^1(R_f,G).$$

This sequence can be used to relate the \mathbf{A}^1 -equivalence on G with local triviality of G-torsors.

Lemma 4.3. Let G be a flat B-linear B-group scheme. Let $h \in B$.

(1) Let $(B \to A, h)$ be a glueing pair and assume that

(4.4)
$$\ker(H^1(B[x], G) \to H^1(B_h[x], G)) = 1.$$

Then we have $A^1G(A_h) = A^1G(A) A^1G(B_h)$ and the map

(4.5)
$$\ker \left(G(B)/A^1 \to G(B_h)/A^1 \right) \to \ker \left(G(A)/A^1 \to G(A_h)/A^1 \right)$$

is surjective.

(2) Assume that h is a non zero divisor in B. Let $\widehat{B} = \varprojlim_{n \ge 0} B/h^{n+1}B$ be the completion. Then we have the inclusion $A^1G(\widehat{B}_h) \subseteq A^1G(\widehat{B})A^1(B_h)$ and the map

(4.6)
$$\ker \left(G(B)/A^1 \to G(B_h)/A^1 \right) \to \ker \left(G(\widehat{B})/A^1 \to G(\widehat{B}_h)/A^1 \right)$$

is surjective. Assuming furthermore that $G(\widehat{B}_h) = G(\widehat{B}) A^1 G(\widehat{B}_h)$, we have $G(B_h) = G(B) A^1(B_h)$.

Proof. (1) Since $(B[t] \rightarrow A[t], h)$ is a glueing pair, we have an exact sequence of pointed sets

$$1 \to G(B_h[x]) \setminus G(A_h[x]) / G(A[x]) \to H^1(B[x], G) \to H^1(B_h[x], G) \times H^1(A[x], G).$$

Our assumption provides a decomposition $G(A_h[x]) = G(A[x]) G(B_h[x])$, and a fortiori a decomposition $G(A_h) = G(A) G(B_h)$. Let $x \in A^1 G(A_h)$. Then there exists $g \in G(A[x]_h)$ such that g(0) = 1 and g(1) = x. We can decompose then $g = g_1 g_2$ with $g_1 \in G(A[x]), g_2 \in G(B[x]_h)$. Since $1 = g_1(0)g_2(0)$ we can assume that $g_1(0) = 1$ and $g_2(0) = 1$. It follows that $x \in A^1 G(A) A^1 G(B_h)$. This establishes the equality $A^1 G(A_h) = A^1 G(A) A^1 G(B_h)$.

For showing the surjectivity of the map (4.5), we are given $[x] \in G(A)/A^1$ and $[y] \in G(B_h)/A^1$ such that $x = y \in G(A_h)/A^1$. The preceding identity allows us to assume that $x = y \in G(A_h)$. Since (B, A, h) is a glueing pair, x, y define a point $g \in G(B)$.

(2) This the special case $A = \hat{B}$. The last fact is a straightforward consequence. \Box

The condition (4.4) in Lemma 4.3 is not easy to check in general. Later on we will discuss a case where it is known to hold as a corollary of the work of Panin on the Serre–Grothendieck conjecture [50, 51]. However, Moser obtained the following unconditional result in the special case of Example 4.1 (a).

Lemma 4.4. (Moser, [48, lemma 3.5.5], see also [3, lemma 3.2.2]) Let G be a finitely presented B-group scheme which is B-linear.

(1) Let $f_0, f_1 \in B$ such that $Bf_0 + Bf_1 = B$. Let $g \in G(B_{f_0f_1}[T])$ be an element such that g(0) = 1. Then there exists a decomposition $g = h_0^{-1} h_1$ with $h_i \in G(B_{f_i}[T])$ and $h_i(0) = 1$ for i = 0, 1.

(2) The sequence of pointed sets

$$G(B)/A^1 \longrightarrow G(B_{f_0})/A^1 \times G(B_{f_1})/A^1 \Longrightarrow G(B_{f_0f_1})/A^1$$

is exact at the middle term.

Proof. (1) The original reference does the case B noetherian and the general case holds by the usual noetherian approximation trick.

(2) Let $[g_0] \in G(B_{f_0})/A^1$ and let $[g_1] \in G(B_{f_1})/A^1$ such that $[g_0] = [g_1] \in G(B_{f_0f_1})/A^1$. Then there exists $g \in G(B_{f_0f_1}[T])$ such that $g_0 g_1^{-1} = g(1) \in G(B_{f_0f_1}[T])$ and g(0) = 1. By (1) we write $g = h_0^{-1} h_1$ with $h_i \in G(B_{f_i}[T])$ and $h_i(0) = 1$ for i = 0, 1 so that $g_0 g_1^{-1} = h_0^{-1}(1) h_1(1)$. Since $[h_i(1)g_i] = [g_i] \in G(B_{f_0})/A^1$, we can replace g_i by $h_i(1)g_i$ and deal then with the case $g_0 = g_1 \in G(B_{f_0f_1})$. This defines an unique element $m \in G(B)$ such that $[m] = [g_i] \in G(B_{f_i})/A^1$.

Remark 4.5. By induction we get the following generalization. Let $f_1, \ldots, f_c \in B$ such that $Bf_1 + \cdots + Bf_c = B$ and put $f = f_1 \ldots f_c$. Let $g \in G(B_f[T])$ be an element such that g(0) = 1. Then there exists a decomposition $g = h_1 \ldots h_c$ with $h_i \in G(B_{f_i}[T])$ and $h_i(0) = 1$ for $i = 1, \ldots, c$. It follows that the image of $G(B)/A^1$ in $\prod_{i=1,\ldots,c} G(B_{f_i})/A^1$ consists of elements having same image in $G(B_f)/A^1$.

Since Lemma 4.4 does not presuppose any results about G-torsors, Moser was able to use it to establish a local-global principle for torsors [48, 3.5.1] generalizing Quillen's local-global principle for finitely presented modules [56, Theorem 1]. In our context, we combine Lemma 4.4 with a theorem of Colliot-Thélène and Ojanguren to obtain the following result.

Proposition 4.6. Let k be an infinite field and let G be an affine k-algebraic group. Let A be the local ring at a prime ideal of a polynomial algebra $k[t_1, \ldots, t_d]$. Then the homomorphism

$$G(A)/A^1 \to G(k(t_1,\ldots,t_d))/A^1$$

is injective.

Proof. Our plan is to use Colliot-Thélène and Ojanguren method [13, §1] as abstracted in the appendix 9.2. We consider the k-functor in groups $B \mapsto F(B) = G(B)/A^1$. The claim follows from Proposition 9.3 once properties $\mathbf{P_1}$, $\mathbf{P_2}$ and $\mathbf{P'_3}$ are checked for the *k*-functor *F*. The property $\mathbf{P_1}$ is clear, since *G* is finitely presented over *k*.

Let *L* be a *k*-field and let $d \ge 0$ be an integer. We have $F(L) = F(L[t_1, \ldots, t_d])$, and F(L) injects in $F(L(t_1, \ldots, t_d))$, since every polynomial over *L* has an invertible value. Property **P**₂ is established. On the other hand Lemma 4.4.(2) establishes the surjectivity of the map

$$\ker \left(G(B)/A^1 \to G(B_{f_0})/A^1 \right) \to \ker \left(G(B_{f_1})/A^1 \to G(B_{f_0f_1})/A^1 \right)$$

for $B = Bf_0 + Bf_1$ so that Zariski patching property $\mathbf{P'_3}$ holds for the functor F. \Box

Remark 4.7. The extension to the finite field case is established in Corollary 5.5.

4.3. Non stable K_1 -functor. Let \mathfrak{G} be a reductive group scheme over our base ring *B*. Let \mathfrak{P} be a strictly proper parabolic subgroup of \mathfrak{G} . Let \mathfrak{P}^- be an opposite *B*-parabolic subgroup scheme of \mathfrak{G} , and denote by $E_{\mathfrak{P}}(B)$ the subgroup of $\mathfrak{G}(B)$ generated by $R_u(\mathfrak{P})(B)$ and $R_u(\mathfrak{P}^-)(B)$ (it does not depend on the choice of $\mathfrak{P}^$ by [64, XXVI.1.8]). We consider the Whitehead coset

$$K_1^{\mathfrak{G},\mathfrak{P}}(B) = \mathfrak{G}(B)/E_{\mathfrak{P}}(B).$$

As a functor on the category of commutative *B*-algebras, $K_1^{\mathfrak{G},\mathfrak{P}}(-)$ is also called the non-stable (or unstable) K_1 -functor associated to \mathfrak{G} and \mathfrak{P}.

Recall that if B is semilocal, then the functor $C \mapsto E_{\mathfrak{P}}(C)$ on the category of commutative B-algebras C does not depend on the choice of a strictly proper parabolic B-subgroup \mathfrak{P} , see [64, XXVI.5] and [62, th. 2.1.(1)]. In particular, in this case $E_{\mathfrak{P}}(B)$ is a normal subgroup of $\mathfrak{G}(B)$. For an arbitrary ring B, the same holds if \mathfrak{G} satisfies the condition (E) below, see [54]. In these two cases we will occasionally write $K_1^{\mathfrak{G}}(C)$ instead of $K_1^{\mathfrak{G},\mathfrak{P}}(C)$, omitting the specific strictly proper parabolic B-subgroup.

Condition (E). For any maximal ideal \mathfrak{m} of B, all irreducible components of the relative root system of $\mathfrak{G}_{B_{\mathfrak{m}}}$ in the sense of [64, XXVI.7] are of rank at least 2.

Note that the condition (E) is satisfied if \mathfrak{G} has B-rank ≥ 2 , since in this case all $\mathfrak{G}_{B_{\mathfrak{m}}}$ also have $B_{\mathfrak{m}}$ -rank ≥ 2 .

Since the radicals $R_u(\mathfrak{P})$ and $R_u(\mathfrak{P}^-)$ are successive extensions of vector group schemes [64, XXVI.2.1], Lemma 2.6.(1) implies that $E_{\mathfrak{P}}(B) \subseteq A^1\mathfrak{G}(B) \subseteq \mathfrak{G}(B)$. We get then surjective maps

$$K_1^{\mathfrak{G},\mathfrak{P}}(B) \longrightarrow \mathfrak{G}(B)/A^1 \longrightarrow \mathfrak{G}(B)/R.$$

4.4. Comparison of $K_1^{\mathfrak{G}}$, \mathbf{A}^1 -equivalence and R-equivalence.

Lemma 4.8. We consider the following assertions:

- (i) The map $K_1^{\mathfrak{G},\mathfrak{P}}(B) \to K_1^{\mathfrak{G},\mathfrak{P}}(B[u])$ is bijective;
- (*ii*) $\mathfrak{G}(B[u]) = \mathfrak{G}(B) E_{\mathfrak{P}}(B[u]);$
- (iii) The map $K_1^{\mathfrak{G},\mathfrak{P}}(B) \to G(B)/A^1$ is bijective.

Then we have the implications $(i) \iff (ii) \implies (iii)$. Furthermore if (iii) holds, we have that $E_{\mathfrak{P}}(B) = A^1\mathfrak{G}(B)$; in particular $E_{\mathfrak{P}}(B)$ is a normal subgroup of $\mathfrak{G}(B)$ which does not depend of \mathfrak{P} .

Proof. $(i) \iff (ii)$. The map $K_1^{\mathfrak{G},\mathfrak{P}}(B) \to K_1^{\mathfrak{G},\mathfrak{P}}(B[u])$ is always injective, since it has a left inverse induced by $u \mapsto 0$. Clearly, this map is surjective, if and only if we have the decomposition $\mathfrak{G}(B[u]) = \mathfrak{G}(B) E_{\mathfrak{P}}(B[u])$.

(ii) \implies (iii). The map $K_1^{\mathfrak{G},\mathfrak{P}}(B) \to G(B)/A^1$ is surjective. Let $g_0, g_1 \in \mathfrak{G}(B)$ mapping to the same element of $G(B)/A^1$. There exists $g(t) \in \mathfrak{G}(B[t])$ such that $g(0) = g_0$ and $g(1) = g_1$. Our assumption implies that g(t) = gh(t) with $g \in \mathfrak{G}(B)$ and $h(t) \in E_{\mathfrak{P}}(B[u])$. It follows that $g_i = gh(i)$ for i = 0, 1 with $h(i) \in E_{\mathfrak{P}}(B)$. We get that $g_0 = gh(0) = (gh(1))(h(1)^{-1}h(0)) \in g_1 E_{\mathfrak{P}}(B)$. Thus g_0, g_1 have same image in $K_1^{\mathfrak{G},\mathfrak{P}}(B)$.

Remarks 4.9. (a) Assume that \mathfrak{G} satisfies condition (*E*). In this case, homotopy invariance reduces to the case of the ring $B_{\mathfrak{m}}$ for each maximal ideal \mathfrak{m} of *B* according to a generalization of the Suslin local-global principle [54, lemma 17].

(b) If B is a regular ring containing a field k, and \mathfrak{G} satisfies (E), then we know that $K_1^{\mathfrak{G}}(B) \xrightarrow{\sim} K_1^{\mathfrak{G}}(B[u])$ by [63, th. 1.1].

(c) Let us provide a counterexample to $K_1^{\mathfrak{G}}(B) \xrightarrow{\sim} K_1^{\mathfrak{G}}(B[u])$ in the non-regular case. Given a field k (of characteristic zero), we consider the domain $B = k[x^2, x^3] \subset$ k[x]. We claim that $K_1^{\mathrm{SL}_n}(B) \subsetneq K_1^{\mathrm{SL}_n}(B[u])$ for n >> 0 so that $1 = K_1^{\mathrm{SL}_n}(B_{\mathfrak{m}}) \subsetneq$ $K_1^{\mathrm{SL}_n}(B_{\mathfrak{m}}[u])$ for some maximal ideal of B. For n >> 0, we have $K_1^{\mathrm{SL}_n}(B) = SK_1(B)$ and $K_1^{\mathrm{SL}_n}(B[u]) = \mathrm{SK}_1(B[u])$. Inspection of the proof of Krusemeyer's computation of $\mathrm{SK}_1(B)$ [38, prop. 12.1] provides functorial maps $\Omega_A^1 \to \mathrm{SK}_1(A \otimes_k B)$ for a k-algebra A. We get then commutative diagram of maps

$$\begin{array}{c} \Omega_{k}^{1} & \xrightarrow{\sim} \operatorname{SK}_{1}(B) \\ \downarrow & \downarrow \\ \Omega_{k[u]}^{1} & \longrightarrow \operatorname{SK}_{1}(B[u]) \\ \downarrow & \downarrow \\ \Omega_{k(u)}^{1} & \xrightarrow{\sim} \operatorname{SK}_{1}(B_{k(u)}) \end{array}$$

where the top and the bottom horizontal maps are isomorphisms [38, prop. 12.1]. Since $\Omega_k^1 \subseteq \Omega_{k[u]}^1$, a diagram chase yields that $\mathrm{SK}_1(B) \subseteq \mathrm{SK}_1(B[u])$. Since $K_1^{\mathrm{SL}_n}(B_m) =$ 1, this example also shows that the condition (*iii*) of Lemma 4.8 does not imply (*i*). (d) In case of regular rings, the condition (*iii*) of Lemma 4.8 may hold while (*i*) does not, if \mathfrak{G} does not satisfy (E). Let k be a field. Let \mathfrak{P} be the standard parabolic subgroup of SL_2 consisting of upper triangular matrices. Then one has $\mathrm{SL}_2(k[x]) =$ $E_{\mathfrak{P}}(k[x])$. Consequently, $K_1^{\mathrm{SL}_2,\mathfrak{P}}(k[x]) = 1$, and hence $\mathrm{SL}_2(k[x])/A^1 = 1$, so (*iii*) holds. On the other hand, $K_1^{\mathrm{SL}_2,\mathfrak{P}}(k[x,u]) \neq 1$ [16], so (*i*) does not hold.

Lemma 4.10. We consider the following assertions:

- (i) The map $K_1^{\mathfrak{G},\mathfrak{P}}(B) \to K_1^{\mathfrak{G},\mathfrak{P}}(B[u]_{\Sigma})$ is bijective;
- (ii) $\mathfrak{G}(B[u]_{\Sigma}) = \mathfrak{G}(B) E_{\mathfrak{P}}(B[u]_{\Sigma});$
- (iii) The map $K_1^{\mathfrak{G},\mathfrak{P}}(B) \to G(B)/R$ is bijective.

Then we have the implications (i) \iff (ii) \implies (iii). Furthermore if (iii) holds, we have that $E_{\mathfrak{P}}(B) = R\mathfrak{G}(B)$; in particular $E_{\mathfrak{P}}(B)$ is a normal subgroup of $\mathfrak{G}(B)$ which does not depend of \mathfrak{P} .

Proof. This is similar with that of Lemma 4.8

5. Passage to the field of fractions

Lemma 5.1. Let B be a regular ring containing a field, and let G be a reductive group over B having a strictly proper parabolic B-subgroup. Let $f \in B[x]$ be a monic polynomial. Then the natural map of étale cohomology sets $H^1_{\acute{e}t}(B[x], G) \to H^1_{\acute{e}t}(B[x]_f, G)$ has trivial kernel.

Proof. Clearly, we can assume that B is a domain. Let K be the field of fractions of B. By [63, Lemma 5.4] for any maximal ideal m of B the map $H^1_{\acute{e}t}(B_m[x], G) \rightarrow$ $H^1_{\acute{e}t}(K[x], G)$ has trivial kernel. Furthermore, the map $H^1_{\acute{e}t}(K[x], G) \rightarrow H^1_{\acute{e}t}(K(x), G)$ has trivial kernel by [13, Proposition 2.2]. Then for any monic polynomial f the map $H^1_{\acute{e}t}(B_m[x], G) \rightarrow H^1_{\acute{e}t}(B_m[x]_f, G)$ has trivial kernel. Since B is regular, by [69, Corollary 3.2] G is B-linear. Then the claim holds by [63, Lemma 4.2].

In the extreme opposite case we have the following fact.

Lemma 5.2. Let B be a Noetherian commutative ring, and let G be a B-linear reductive B-group. We assume that $G_{B/m}$ is anisotropic for each maximal ideal m of B. Let $f \in B[x]$ be a monic polynomial. Then the natural map of étale cohomology sets $H^1_{\acute{e}t}(B[x], G) \to H^1_{\acute{e}t}(B[x]_f, G)$ has trivial kernel.

Proof. Assume first that B is semilocal. Let $\xi = [E] \in H^1_{\acute{e}t}(B[x], G)$ be an element of the kernel. We extend E to a G-bundle \hat{E} on \mathbb{P}^1_B by patching it to the trivial G-bundle over $\mathbb{P}^1_B \setminus \{f = 0\}$. We denote by $\hat{\xi}$ its class; since f is monic, we have $\hat{\xi}|_{\infty} = *$.

Let m_1, \ldots, m_c be the maximal ideals of B and put $k_i = B/m_i$. Since G_{k_i} is anisotropic, then by [21, Th. 3.8 (b)] $\hat{\xi}_{k_i}$ is trivial. Next we apply [10, lemma 5.2.1] and get that $\hat{\xi}$ belongs to the image of $H^1_{\acute{e}t}(B,G) \to H^1_{\acute{e}t}(\mathbb{P}^1_B,G)$. Since $\hat{\xi}|_{\infty} = *$, we conclude that $\hat{\xi} = *$. Thus E is a trivial G-torsor over B[x].

If B is not necessarily semilocal, the claim reduces to the maximal localizations of B by applying the local-global principle [63, Lemma 4.2]. \Box

Remarks 5.3. (a) The rigidity property for \mathbb{P}^1_B -torsors under reductive groups was proved in [57, Th. 1] and [52, Prop. 9.6] under the assumption that B is semilocal and contains a field (i.e. is equicharacteristic). Tsybyshev [71, Theorem 1] was able to prove it assuming only that B is reduced and $\operatorname{Pic}(B) = 0$. Česnavičius [10] observed that one can remove the condition that B contains a field by using Alper's theorem stating that GL_N/G is affine for any B [1, cor. 9.7.7]. The idea to use [21, Th. 3.8 (b)] for anisotropic groups appeared in [20, p. 178] and in [19, th. 1 and remark 2.1.(iii) on the anisotropic case]. Fedorov also introduced the use of affine Grassmannians to treat the case of not necessarily semilocal B and anisotropic G [19, Theorem 5].

(b) Let G_0 the underlying Chevalley *B*-group scheme of *G*. The condition of linearity on *G* is satisfied if the $Out(G_0)_S$ -torsor $Isomext(G_0, G)$ is isotrivial, see [42, prop. 3.2]; this reference provides then a representation such that GL_n/G is affine, so there is no need to appeal to Alper's result in this case. This includes the semisimple case and the case when *B* is a normal ring due to Thomason [69, Corollary 3.2].

(c) The claim of Lemma 5.2 does not hold if G is anisotropic over B and isotropic over B/m, even if B is regular local and G is simply connected [19, Corollary 2.3].

Theorem 5.4. Let B be a regular semilocal domain that contains a field k, and let K be the fraction field of B. Let \mathfrak{G} be a reductive B-group scheme.

(1) Assume that either \mathfrak{G} contains a strictly proper parabolic B-subgroup, or \mathfrak{G} is anisotropic over B/m for all maximal ideals m of B. Then the map

$$\mathfrak{G}(B)/R \to \mathfrak{G}(K)/R$$

is surjective.

(2) Assume that \mathfrak{G} contains a strictly proper parabolic B-subgroup. Then the map

$$\mathfrak{G}(B)/A^1 \to \mathfrak{G}(K)/A^1$$

is injective.

Proof. Clearly, we can assume that k is a finite field or \mathbb{Q} without loss of generality. Then the embedding $k \to B$ is geometrically regular, since k is perfect [43, (28.M), (28.N)]. Then by Popescu's theorem [55, 67] B is a filtered direct limit of smooth k-algebras. Since the group scheme \mathfrak{G} is finitely presented over B, and the functors $\mathfrak{G}(-)/R$ and $\mathfrak{G}(-)/A^1$ commute with filtered direct limits, we can assume that \mathfrak{G} is defined over a smooth k-domain C, and $B = C_S$ is a localization of C at a set S that is a union of a finite set of prime ideals p_i of C. Moreover, since parabolic subgroups of \mathfrak{G} are also finitely presented, depending on the assumption on \mathfrak{G} we can secure that \mathfrak{G} contains a strictly proper parabolic subgroup over C, or \mathfrak{G} is anisotropic over $C_{p_i}/p_i C_{p_i}$ for all p_i 's.

(1) We need to show that $\mathfrak{G}(B)/R \to \mathfrak{G}(K)/R$ is surjective, where K is the fraction field of B and C. Clearly, it is enough to show the same for the localization of C at the complement of the union of maximal ideals $m_i \supseteq p_i$ (note that if \mathfrak{G} is anisotropic over $C_{p_i}/p_i C_{p_i}$, then it is automatically anisotropic over C/m_i). Hence we can assume that *B* is a localization of *C* at a union of a finite set of maximal ideals. On top of that, in order to show that $\mathfrak{G}(B)/R \to \mathfrak{G}(K)/R$ is surjective, it is enough to show that for any $f \in \bigcap_i m_i$ and any $g \in \mathfrak{G}(C_f)$ the image of g in $\mathfrak{G}(K)$ belongs to $\mathfrak{G}(B) \cdot R\mathfrak{G}(K)$.

We apply Panin's theorem [50, th. 2.5]. This provides a monic polynomial $h \in B[t]$, an inclusion of rings $B[t] \subset A$, a homomorphism $\phi : A \to B$ and a commutative diagram

such that

(i) the left hand square is a elementary distinguished Nisnevich square in the category of smooth B-schemes in the sense of [47, 3.1.3];

- (ii) the composite $C \xrightarrow{u} A \xrightarrow{\phi} B$ is the canonical localization homomorphism;
- (iii) the map $B[t] \to A \xrightarrow{\phi} B$ is the evaluation at 0;
- (iv) $h(1) \in B^{\times}$;
- (v) there is an A-group scheme isomorphism $\Phi : \mathfrak{G}_B \times_B A \xrightarrow{\sim} \mathfrak{G} \times_C^u A$.

By inspection of the construction A is finite étale over B[t] and $h(t) = N_{A/B[t]}(u(f)) = u(f)a$ with $a \in A$. Property (4) of [52, theorem 3.4] states that the map $\phi : A \to B$ extends to a map $A_a \to B$, so that $\phi(g) \in B^{\times}$. We compute

$$h(0) = \phi(h) \quad [\text{property } (iii)]$$

= $\phi(u(f)) \phi(a)$
= $f \phi(a) \quad [\text{property } (ii)]$

it follows that h(0) is a non-zero element of B. In particular ϕ extends to a map $\phi_h : A_h \to B_{h(0)}$.

Since $(B[t] \to A, h)$ is a glueing pair, we have an exact sequence of pointed sets

$$1 \to \mathfrak{G}(B[t]_h) \setminus \mathfrak{G}(A_h) / \mathfrak{G}(A) \to H^1(B[t], \mathfrak{G}) \to H^1(B[t]_h, \mathfrak{G}) \times H^1(A, \mathfrak{G}).$$

Our assumptions on \mathfrak{G} imply that the map $H^1(B[t], \mathfrak{G}) \to H^1(B[t]_h, \mathfrak{G})$ has trivial kernel. Indeed, if \mathfrak{G} contains a strictly proper parabolic subgroup over B, this follows from Lemma 5.1. If \mathfrak{G} is anisotropic modulo every maximal ideal of B, then the same follows from Lemma 5.2, taking into account that B is regular and hence by [69, Corollary 3.2] G is B-linear. Therefore we have $\mathfrak{G}(A_h) = \mathfrak{G}(B[t]_h)\mathfrak{G}(A)$.

Set $\tilde{g} = \Phi^{-1}(v_*(g)) \in \mathfrak{G}(A_h)$. Then $\tilde{g} = b \cdot a$, where $b \in \mathfrak{G}(B[t]_h)$ and $a \in \mathfrak{G}(A)$. Note that by (iii) we have $\phi(h) = h(0)$. We have $\phi_h(\tilde{g}) = \phi_h(v(g)) = g \in \mathfrak{G}(B_{h(0)})$ by (ii). It follows that $g = \phi_h(b) \cdot \phi_h(a)$. Clearly we have $\phi_h(a) \in \mathfrak{G}(B) \subseteq \mathfrak{G}(B_{h(0)})$. We claim that $\phi_h(b) \in \mathfrak{G}(B) \cdot R\mathfrak{G}(B_{h(0)})$. Indeed, we have $\phi_h(b) = b|_{t=0}$ by (iii), and since $h(1) \in B^{\times}$, we have $b|_{t=1} \in \mathfrak{G}(B)$. Then the image of b in $\mathfrak{G}(B_{h(0)}[t]_h)$ provides an R-equivalence between $\phi_h(b)$ and an element of $\mathfrak{G}(B)$. Summing up, the image of $g \in \mathfrak{G}(C_f)$ under the composition $\mathfrak{G}(C_f) \xrightarrow{v} \mathfrak{G}(A_h) \xrightarrow{\phi_h} \mathfrak{G}(B_{h(0)})$ belongs to $\mathfrak{G}(B) \cdot R\mathfrak{G}(B_{h(0)})$. It follows that the image of g in $\mathfrak{G}(K)$ belongs to $\mathfrak{G}(B) \cdot R\mathfrak{G}(K)$.

(2) Let $[g] \in \ker(\mathfrak{G}(B)/A^1 \to \mathfrak{G}(K)/A^1)$. Up to shrinking of $X = \operatorname{Spec}(C)$, we can assume that $g \in \mathfrak{G}(C)$. Then there exists then $f \in C$ such that $[g] \in \ker(\mathfrak{G}(C)/A^1 \to \mathfrak{G}(C_f)/A^1)$. As in (1), we apply Panin's theorem [50, th. 2.5] and obtain a diagram (5.1) satisfying the properties (i)–(v). But this time we set $\tilde{g} = \Phi^{-1}(u_*(g)) \in \mathfrak{G}(A)$ and we have $[\tilde{g}] \in \ker(\mathfrak{G}(A)/A^1 \to \mathfrak{G}(A_h)/A^1)$. According to Lemma 5.1, the map $H^1(B[t][x], \mathfrak{G}) \to H^1(B[t]_h[x], \mathfrak{G})$ has trivial kernel so that Lemma 4.3.(1) shows that the map

(5.2)
$$\ker\left(\mathfrak{G}(B[t])/A^1 \to \mathfrak{G}(B[t]_h)/A^1\right) \to \ker\left(\mathfrak{G}(A)/A^1 \to \mathfrak{G}(A_h)/A^1\right)$$

is surjective. Since $\mathfrak{G}(B)/A^1 = \mathfrak{G}(B[t])/A^1$ and $h(1) \in B^{\times}$, we deduce that

$$\operatorname{ker}(\mathfrak{G}(A)/A^1 \to \mathfrak{G}(A_h)/A^1) = 1.$$

We have $[\tilde{g}] = 1 \in \mathfrak{G}(A)/A^1$ and get $[u_*(g)] = 1 \in \mathfrak{G}(A)/A^1$. By applying ϕ_* , the property (ii) yields $[g] = 1 \in \mathfrak{G}(B)/A^1$.

Corollary 5.5. Let k be a field and let G be an affine k-algebraic group. Let A be the local ring at a prime ideal of a polynomial algebra $k[t_1, \ldots, t_d]$. Then the homomorphism

$$G(A)/A^1 \rightarrow G(k(t_1,\ldots,t_d))/A^1$$

is injective.

Proof. If k is infinite, this is the claim of Proposition 4.6. Assume that k is finite. Let G_{red} denote the reduced affine algebraic k-scheme corresponding to G. Since k is perfect, G_{red} is a smooth algebraic k-subgroup of G [45, Prop. 1.26, Cor. 1.39]. Since A is reduced, $G(A) = G_{red}(A)$ and $G(A[u]) = G_{red}(A[u])$, therefore, $G(A)/A^1 =$ $G_{red}(A)/A^1$, and hence we can assume that G is smooth from the start. Let G° be the connected component of the identity $e \in G(k)$. Let $\pi_0(G)$ be the finite étale kscheme of connected components of G Then G° is a smooth geometrically connected algebraic k-subgroup of G, the fiber of the natural map $G \to \pi_0(G)$ at the image of e [45, Prop. 1.31, 1.34]. Since $\pi_0(G)$ is k-finite, we have $\pi_0(G)(A[u]) = \pi_0(G)(A)$, and hence $\pi_0(G)(A)/A^1 = \pi_0(G)(A)$ injects into $\pi_0(G)(K)/A^1 = \pi_0(G)(K)$, where $K = k(t_1, \ldots, t_d)$. Therefore, in order to prove the claim for G, it is enough to prove it for G° . Hence we can assume that G is smooth and connected. Let U be the unipotent radical of G over k, i.e. the largest smooth connected unipotent normal k-subgroup of G. Since k is perfect, the group U is k-split, admits a subnormal series each of those quotients are isomorphic to \mathbb{A}_{k}^{1} [45, 14.63]. Therefore $U(A)/A^{1} = 1$ and $H^{1}(R, U) = 1$ for every k-algebra R. Also, since k is perfect, G/U is a reductive algebraic kgroup [45, Prop. 19.11]. By Lang's theorem, G/U is quasi-split, and therefore either G/U is a k-torus, or it contains a strictly proper parabolic k-subgroup and then satisfies Theorem 5.4 (2). In both cases the map $(G/U)(A)/A^1 \to (G/U)(K)/A^1$ is injective. Now let $g \in G(A)$ be mapped into $A^1G(K) \subseteq G(K)$. By the previous argument, there is $h(u) \in (G/U)(A[u])$ such that h(0) = 1 and h(1) is the image of g in (G/U)(A). Since $H^1(A,U) = H^1(A[u],U) = 1$, there is $g(u) \in G(u)$ such that $g(0) \in U(A)$ and $g(1)g^{-1} \in U(A)$. Since $U(A) \subseteq A^1G(A)$, we conclude that $g \in A^1G(A)$, as required.

6. The case of simply connected semisimple isotropic groups

6.1. Coincidence of equivalence relations. We address the following question.

Question 6.1. Assume that B is regular semilocal and that \mathfrak{G} is semisimple simply connected and strictly isotropic equipped with a strictly proper parabolic B-subgroup. Is the map $K_1^{\mathfrak{G},\mathfrak{P}}(B) \to \mathfrak{G}(B)/R$ an isomorphism? Is the map $\mathfrak{G}(B)/A^1 \to \mathfrak{G}(B)/R$ an isomorphism?

The answer is known to be positive in both cases if B is a field. This is implied by Margaux–Soulé isomorphism [41, Th. 3.10] combined with [23, Th. 7.2].

Theorem 6.2. Assume that B is a semilocal regular domain containing a field k and denote by K its fraction field. Let \mathfrak{G} be a semisimple simply connected B-group having a strictly proper parabolic B-subgroup. Then we have a commutative square of isomorphisms


$$\begin{array}{ccc} G(B)/A^{1} & \xrightarrow{\sim} & G(B)/R \\ & & & & \downarrow \\ & & & \downarrow \\ G(K)/A^{1} & \xrightarrow{\sim} & G(K)/R \end{array}$$

Proof. Let K be the fraction field of B. The bottom horizontal arrow of the square is an isomorphism by the Margaux–Soulé theorem [41, Th. 3.10] combined with [23, Th. 7.2]. On the other hand, the left vertical map is injective by Theorem 5.4 (2). Then the top horizontal arrow is also injective. Since it is surjective by definition, it is an isomorphism. The right vertical arrow is surjective by Theorem 5.4 (1). Hence the vertical arrows are also isomorphisms.

Remark 6.3. The above result does not extend to anisotropic groups. For example, let k be an infinite field and let G be a wound linear algebraic group, i.e. does not contain any subgroups isomorphic to \mathbb{G}_a or \mathbb{G}_m . Then by [25, Corollary 3.8] we have G(k[x]) = G(k) and, consequently, $G(k)/A^1 = G(k)$. This applies in particular to the case of an anisotropic reductive k-group G. On the other hand, the R-equivalence class group of G may be even trivial, e.g. if G is a semisimple anisotropic group of rank ≤ 2 . Indeed, in this case every element of G(k) is R-equivalent to a semisimple regular element, and all maximal tori of G are of rank ≤ 2 and hence rational.

In the same vein, we can establish the following fact.


Corollary 6.4. Let k be a field and let G be a semisimple simply connected k-group strictly of k-rank ≥ 1 . Let A be the localization of $k[x_1, \ldots, x_d]$ at a prime ideal. Then we have a commutative square of isomorphisms



Proof. By [23, th. 5.8] there is an isomorphism $G(k)/A^1 \xrightarrow{\sim} G(k(x_1, \ldots, x_d))/A^1$. Then the claim follows from Theorem 6.2.

Theorem 6.5. Assume that B is a semilocal regular domain containing a field k and that \mathfrak{G} is semisimple simply connected B-group strictly of B-rank ≥ 2 . Then the map $K_1^{\mathfrak{G}}(B) \to \mathfrak{G}(B)/R$ is an isomorphism.

Proof. Let K be the fraction field of B. We consider the commutative diagram



where the bottom isomorphism is [23, th. 7.2]. On the other hand, the left vertical map is injective [63, th. 1.2]. By diagram chase, the top horizontal map is an isomorphism.

6.2. The retract rational case. We now consider the vanishing of Whitehead cosets.

Lemma 6.6. We assume that the base ring B is a semilocal domain. Let \mathfrak{G} be a reductive B-group scheme having a strictly proper B-parabolic subgroup \mathfrak{P} . We consider the following assertions:

- (i) $K_1^{\mathfrak{G},\mathfrak{P}}(F) = 1$ for every *B*-field *F*;
- $(ii) \mathfrak{G}$ satisfies the lifting property;
- (*iii*) (\mathfrak{G}, e) is a retract rational *B*-scheme.

Then the following implications $(i) \Longrightarrow (ii) \Longrightarrow (iii)$ hold.

Proof. $(i) \Longrightarrow (ii)$. Let C be a semilocal B-ring with residue fields F_1, \ldots, F_s . We have to show that the map $\mathfrak{G}(C) \to \prod_{i=1,\ldots,s} \mathfrak{G}(F_i)$ is onto. We are given an element $(g_1, \ldots, g_s) \in \prod_{i=1,\ldots,s} \mathfrak{G}(F_i)$. Our assumption implies that there exists a positive integer d such that

$$g_i = u_{i,1} v_{i,1} u_{i,2} v_{i,2} \dots u_{i,d} v_{i,d}$$

with $u_{i,j} \in R_u(\mathfrak{P})(F_i)$ (resp. $v_{i,j} \in R_u(\mathfrak{P}^-)(F_i)$) for $i = 1, \ldots, s$ and $j = 1, \ldots, d$. Since $R_u(\mathfrak{P})(C) \to \prod_{i=1,\ldots,s} R_u(\mathfrak{P})(F_i)$ is onto (and similarly for $R_u(\mathfrak{P}^-)$), we can lift each $(u_{i,j})_{i=1,\ldots,s}$ in some $u_j \in R_u(\mathfrak{P})(C)$ (resp. $(v_{i,j})_{i=1,\ldots,s}$ in $v_j \in R_u(\mathfrak{P}^-)(C)$). Thus the product $u_1 v_1 u_2 v_2 \ldots u_d v_d$ lifts the g_i 's.

 $(ii) \Longrightarrow (iii)$. This follows from Proposition 2.10.

Proposition 6.7. Assume that B is a semilocal domain and that \mathfrak{G} is semisimple simply connected B-group having a strictly proper parabolic B-subgroup. Let K be the fraction field of B. Then the following assertions are equivalent:

- (i) \mathfrak{G} satisfies the lifting property;
- (ii) $(\mathfrak{G}, 1)$ is a retract rational *B*-scheme;

(iii) \mathfrak{G} is R-trivial on semilocal rings, that is $\mathfrak{G}(C)/R = 1$ for each semilocal B-ring C;

(iv) $\mathfrak{G}(F)/R = 1$ for each B-field F.

Proof. Let \mathfrak{P} be a strictly proper parabolic subgroup scheme of \mathfrak{G} .

 $(i) \Longrightarrow (ii)$. We assume that \mathfrak{G} satisfies the lifting property. Then Proposition 2.10, $(ii) \Longrightarrow (i)$, shows that \mathfrak{G} is retract rational over B.

 $(ii) \Longrightarrow (iii)$. If all residue fields are infinite, this is Proposition 2.14. For the general case, it is enough to show that $\mathfrak{G}(B)/R = 1$. We may assume that $\kappa_1, \ldots, \kappa_b$ are finite fields and that $\kappa_{b+1}, \ldots, \kappa_c$ are infinite. Let $(\mathfrak{U}, 1)$ be an open subset of $(\mathfrak{G}, 1)$ which is a *B*-retract of some open of $(\mathbf{A}_B^N, 0)$. We know that $E_{\mathfrak{P}}(\kappa_i) = \mathfrak{G}(\kappa_i)$ for $i = 1, \ldots, b$ [70, 1.1.2]. We consider the open *B*-subscheme $\mathfrak{V} = \mathfrak{U} E_{\mathfrak{P}}(B)$ of \mathfrak{G} . Since $E_P(\kappa_i)$ is dense in G_{κ_i} for $i = b + 1, \ldots, c$, we have $\mathfrak{V}_{\kappa_i} = \mathfrak{G}_{\kappa_i}$ for $b + 1 = 1, \ldots, c$ Since the map $E_{\mathfrak{P}}(B) \to \prod_{i=1,\ldots,b} E_{\mathfrak{P}}(\kappa_i)$ is onto, we have $\mathfrak{V}(B) = \mathfrak{G}(B)$. Lemma 2.6.(1) shows that $\mathfrak{U}(B)/R = 1$ so that $\mathfrak{V}(B)/R = 1$.

 $(iii) \Longrightarrow (iv)$. Obvious.

 $(iv) \Longrightarrow (i)$. Since (iii) holds in particular for any *B*-field *F*, we have $K_1^{\mathfrak{G},\mathfrak{P}}(F) = 1$ for every *B*-field *F* according to Margaux–Soulé isomorphism [41, Th. 3.10]. Lemma 6.6, $(i) \Longrightarrow (ii)$, implies that *G* satisfies the lifting property for any semilocal *B*-algebra *C*.

This can be refined in the regular case.

Theorem 6.8. Assume that B is a semilocal regular domain containing a field k and that \mathfrak{G} is a semisimple simply connected B-group having a strictly proper parabolic B-subgroup. Let K be the fraction field of B. Then the following assertions are equivalent:

(i) \mathfrak{G} satisfies the lifting property;

(i') \mathfrak{G} satisfies the lifting property for each B-ring C which is a semilocal regular domain and such that B embeds in C;

(ii) $(\mathfrak{G}, 1)$ is a retract rational *B*-scheme;

(iii) \mathfrak{G} is *R*-trivial, that is, $\mathfrak{G}(C)/R = 1$ for each semilocal *B*-ring *C*;

(*iii'*) $\mathfrak{G}(C)/A^1 = 1$ for each *B*-ring *C* which is a semilocal regular domain;

(iv) $\mathfrak{G}(F)/R = 1$ for each B-field F;

(v) \mathfrak{G}_K is a retract rational K-variety.

If, moreover, \mathfrak{G} is strictly of B-rank ≥ 2 , then the above statements are also equivalent to the following:

(iii'') $K_1^{\mathfrak{G}}(C) = 1$ for each B-ring C which is a semilocal regular domain.

Proof. Let \mathfrak{P} be a strictly proper parabolic *B*-subgroup scheme of \mathfrak{G} . We detail only the additional facts from Proposition 6.7 which provides already the equivalences $(i) \iff (ii) \iff (iv)$.

 $(i) \Longrightarrow (i')$. Obvious.

 $(i') \Longrightarrow (ii)$. In the proof of Proposition 2.10, $(ii) \Longrightarrow (i)$, we apply the lifting to a semilocalization of $B[t_1, \ldots, t_n]$ which is a regular semilocal domain which contains B. So the proof of Proposition 6.7, $(i) \iff (iii)$, works so that $(\mathfrak{G}, 1)$ is retract B-rational.

 $(iii) \implies (iii')$. By [41, Th. 3.10] combined with [23, Th. 7.2] we have $\mathfrak{G}(F)/A^1 = \mathfrak{G}(F)/R$ for each *B*-field *F*. Then the claim follows by Theorem 6.2.

 $(iii') \Longrightarrow (iv)$. Obvious.

 $(iv) \Longrightarrow (v)$. The assumption implies that the semisimple simply connected strictly isotropic K-group $G = \mathfrak{G}_K$ satisfies G(E)/R = 1 for all K-fields E. According to [23, cor. 5.10], G is a retract K-rational variety.

 $(v) \Longrightarrow (i')$. Let C be a semilocal regular domain which contains B. It is clear from the proof of the implication $(i) \Longrightarrow (ii)$ of Lemma 6.6 that it is enough to show that $K_1^{\mathfrak{G}}(F) = 1$ for every residue field F of C. Let \hat{C} be the completion of the localization of C at the prime ideal corresponding to F. Then \hat{C} is a regular local ring, and the fraction field \hat{K} of \hat{C} is an extension of K. Since \mathfrak{G}_K is retract rational, we have $\mathfrak{G}(\hat{K})/R = 1$. Then $\mathfrak{G}(\hat{C})/R = 1$ by Theorem 6.2. Since \mathfrak{G} is affine and smooth, the map $\mathfrak{G}(\hat{C}) \to \mathfrak{G}(F)$ is surjective [30, th. I.8]. Hence $\mathfrak{G}(\hat{C})/R \to \mathfrak{G}(F)/R$ is surjective and $\mathfrak{G}(F)/R = 1$. According to [23, Th. 7.2], we have $K_1^{\mathfrak{G}}(F) = \mathfrak{G}(F)/R$.

We assume now that \mathfrak{G} is strictly of *B*-rank ≥ 2 . (*iii*) \Longrightarrow (*iii''*) Follows from Theorem 6.5. (*iii''*) \Longrightarrow (*iii'*). Obvious.

Since a positive answer to the Kneser-Tits problem over fields is known in a bunch of cases, we get the following concrete result.

Corollary 6.9. Assume that B is a connected semilocal ring containing a field k. We assume that \mathfrak{G} is semisimple simply connected isotropic B-group and that \mathfrak{G}_K is absolutely almost K-simple. Then $\mathfrak{G}(B)/A^1 = 1$ in the following cases:

- (1) \mathfrak{G} is quasi-split;
- (2) the components of the anisotropic kernel of \mathfrak{G} are of rank ≤ 2 ;
- (3) $\mathfrak{G} = \operatorname{SL}_m(A)$ where $m \geq 2$ and A is an Azumaya R-algebra of squarefree index;
- (4) \mathfrak{G} is of type B_n , C_n ;
- (5) $\mathfrak{G} = \operatorname{Spin}(q)$ for a regular quadratic form q which is even dimensional (and isotropic);
- (6) $\mathfrak{G}_K = \operatorname{Spin}(A, h)$ where A is an Azumaya R-algebra of degree 2 or 4 equipped with an orthogonal involution of first kind and h is an isotropic regular hermitian form.
- (7) \mathfrak{G} if of type ^{3,6} D_4 or ¹ E_6 ;
- (8) \mathfrak{G} is of type ${}^{2}E_{6}$ with one of the following Tits indices

a)
$$\begin{array}{c} \alpha_2 & \alpha_4 & \alpha_3 & \alpha_1 \\ \bullet & \bullet & \bullet \\ \alpha_5 & \alpha_6 \end{array}$$
 b) $\begin{array}{c} \alpha_2 & \alpha_4 & \alpha_3 & \alpha_1 \\ \bullet & \bullet & \bullet \\ \alpha_5 & \alpha_6 \end{array}$ c) $\begin{array}{c} \alpha_2 & \alpha_4 & \alpha_3 & \alpha_1 \\ \bullet & \bullet & \bullet \\ \alpha_5 & \alpha_6 \end{array}$ c) $\begin{array}{c} \alpha_2 & \alpha_4 & \alpha_3 & \alpha_1 \\ \bullet & \bullet & \bullet \\ \alpha_5 & \alpha_6 \end{array}$ denotes that $\begin{array}{c} \alpha_2 & \alpha_4 & \alpha_3 & \alpha_1 \\ \bullet & \bullet & \bullet \\ \alpha_5 & \alpha_6 \end{array}$ denotes the last case we assume that $6 \in k^{\times}$.

(9) \mathfrak{G} is of type E_7 with one of the following Tits indices

a)
$$\overbrace{\alpha_{7} \ \alpha_{6} \ \alpha_{5} \ \alpha_{4} \ \alpha_{3} \ \alpha_{1}}^{\alpha_{2}}$$
b)
$$\overbrace{\alpha_{7} \ \alpha_{6} \ \alpha_{5} \ \alpha_{4} \ \alpha_{3} \ \alpha_{1}}^{\alpha_{2}}$$
(10)
$$\mathfrak{G} \ is \ of \ type \ E_{8} \ with \ the \ following \ Tits \ indices$$

$$\overbrace{\alpha_{8} \ \alpha_{7} \ \alpha_{6} \ \alpha_{5} \ \alpha_{4} \ \alpha_{3} \ \alpha_{1}}^{\alpha_{2}}$$

$$\overbrace{\alpha_{8} \ \alpha_{7} \ \alpha_{6} \ \alpha_{5} \ \alpha_{4} \ \alpha_{3} \ \alpha_{1}}^{\alpha_{2}}$$

$$\overbrace{\alpha_{8} \ \alpha_{7} \ \alpha_{6} \ \alpha_{5} \ \alpha_{4} \ \alpha_{3} \ \alpha_{1}}^{\alpha_{2}}$$

$$\overbrace{\alpha_{8} \ \alpha_{7} \ \alpha_{6} \ \alpha_{5} \ \alpha_{4} \ \alpha_{3} \ \alpha_{1}}^{\alpha_{2}}$$

If furthermore \mathfrak{G} is strictly of B-rank ≥ 2 , then $K_1^{\mathfrak{G}}(B) = 1$.

30

Proof. We assume firstly that the k-algebra B is a regular domain so that the statement is a case by case application of Theorem 6.8, $(v) \Longrightarrow (iii')$ or $(iv) \Longrightarrow (iii')$. Almost all results of retract rationality over K are quoted in [23, th. 6.1] excepted the following cases.

Third outer E_6 case, i.e. $E_{6,1}^{29}$. This is due to Garibaldi, see [23, Th. 6.2].

Second E_8 case, i.e. $E_{8,2}^{66}$. This is a result by Parimala-Tignol-Weiss [53, §3].

Third E_7 (resp. E_8) case, i.e. $E_{7,1}^{78}$ (resp. $E_{8,2}^{78}$). The *R*-triviality over fields is a result by Alsaody-Chernousov-Pianzola [2, Th. 8.1] and by Thakur [68, Th. 4.2 and Cor. 4.3] independently.

To deduce the general case where B is not necessarily regular, we use Hoobler's trick, see [33, proof of theorem 2] or [35, p. 109]. There exists an henselian pair (C, I) such that C/I = B and $C = \varinjlim C_{\alpha}$ where each C_{α} is a semilocalization of an affine k-space.

We are given a minimal *B*-parabolic subgroup \mathfrak{P} of \mathfrak{G} Denote by G_0 the split Chevalley *k*-form of \mathfrak{G} . Then $(\mathfrak{G}, \mathfrak{P})$ is a form of (G_0, P_0) . Since $\operatorname{Aut}(G_0, P_0)$ is a smooth affine *k*-group [24, lemme 5.1.2], the map

$$H^1(C, \operatorname{Aut}(G_0, P_0)) \to H^1(C/I, \operatorname{Aut}(G_0, P_0))$$

is bijective [66, th. 1]. This implies that there exists a couple $(\widetilde{\mathfrak{G}}, \widetilde{\mathfrak{P}})$ over C such that $(\widetilde{\mathfrak{G}}, \widetilde{\mathfrak{P}}) \times_C B = (\mathfrak{G}, \mathfrak{P})$. Since $\widetilde{\mathfrak{G}}$ is smooth over C, the map $\widetilde{\mathfrak{G}}(C) \to \mathfrak{G}(B)$ is onto and so is $\widetilde{\mathfrak{G}}(C)/A^1 \to \mathfrak{G}(B)/A^1$.

On the other hand, we have $H^1(C, \operatorname{Aut}(G_0)) = \varinjlim H^1(C_\alpha, \operatorname{Aut}(G_0))$ [64, VI_B.10.16]. It follows that there exists α_0 and a couple $(\mathfrak{G}_{\alpha_0}, \mathfrak{P}_{\alpha_0})$ such that $(\mathfrak{G}_{\alpha_0}, \mathfrak{P}_{\alpha_0}) \times_{C_{\alpha_0}} C = (\widetilde{\mathfrak{G}}, \widetilde{\mathfrak{P}})$. We have $\widetilde{\mathfrak{G}}(C) = \varinjlim_{\alpha \geq \alpha_0} \mathfrak{G}_{\alpha_0}(C_\alpha)$. But $\mathfrak{G}_{\alpha_0}(C_\alpha)/A^1 = 1$ by the regular case of the theorem. Since $\varinjlim_{\alpha \geq \alpha_0} \mathfrak{G}_{\alpha_0}(C_\alpha)/A^1 \to \widetilde{\mathfrak{G}}(C)/A^1$ is onto, we conclude that $\widetilde{\mathfrak{G}}(C)/A^1 = 1$.

If furthermore \mathfrak{G} is strictly of B-rank ≥ 2 , then we have similarly $K_1^{\mathfrak{G}_{\alpha_0}}(C_{\alpha}) =$ 1 from the regular case and a composite of surjective maps $\varinjlim_{\alpha \geq \alpha_0} K_1^{\mathfrak{G}_{\alpha_0}}(C_{\alpha}) \longrightarrow K_1^{\mathfrak{G}}(C) \longrightarrow K_1^{\mathfrak{G}}(B)$. Thus $K_1^{\mathfrak{G}}(B) = 1$. \Box

7. Behaviour for henselian pairs

We address the following question with respect to a henselian pair (B, I) [65, 15.11]; this concerns, for example, the case of a nilpotent ideal.

Question 7.1. Let \mathfrak{G} be a reductive *B*-group scheme. Is the map $\mathfrak{G}(B)/R \to \mathfrak{G}(B/I)/R$ an isomorphism?

Note that since \mathfrak{G} is affine and smooth over B, the map $\mathfrak{G}(B) \to \mathfrak{G}(B/I)$ is surjective [30, th. I.8], and hence the map of R-equivalence class groups is surjective.

7.1. The torus case.

Lemma 7.2. Assume that B/I is a normal domain. Let \mathfrak{T} be a B-torus. Then \mathfrak{T} is isotrivial.

Proof. Since B/I is a normal domain, \mathfrak{T} is isotrivial [64, X.5.16], that is, there exists a finite étale cover C_0 of B/I such that $\mathfrak{T}_{C_0} \cong \mathbb{G}^r_{m,C_0}$. Since (B, I) is a henselian couple, C_0 lifts to a finite étale cover C of B [65, Tag 09ZL] and furthermore (C, IC) is an henselian pair (*ibid*, Tag 09XK). According to [10, cor. 3.1.3.(b)], the isomorphism $\mathfrak{T}_{C_0} \cong \mathbb{G}^r_{m,C_0}$ lifts to an isomorphism $\mathfrak{T}_C \cong \mathbb{G}^r_{m,C}$ so that $\mathfrak{T} \times_G C$ is split. Thus \mathfrak{T} is isotrivial.

A first evidence for the question 7.1 is the following fact.

Lemma 7.3. Let \mathfrak{T} be a *B*-torus. Assume that B/I is a regular domain. Then the map $\mathfrak{T}(B)/R \to \mathfrak{T}(B/I)/R$ is an isomorphism.

Proof. The *B*-torus \mathfrak{T} is isotrivial according to Lemma 7.2. Let $1 \to \mathfrak{S} \to \mathfrak{Q} \xrightarrow{\pi} \mathfrak{T} \to 1$ be a flasque resolution. We have a commutative diagram of exact sequences

According to [66, th. 1], the maps $H^1(B,\mathfrak{S}) \to H^1(B/I,\mathfrak{S})$ and $H^1(B,\mathfrak{Q}) \to H^1(B/I,\mathfrak{Q})$ are isomorphisms. By diagram chase we conclude that the map $\mathfrak{T}(B)/\pi(\mathfrak{Q}(B)) \to \mathfrak{T}(B/I)/\pi(\mathfrak{Q}(B/I))$ is an isomorphism.

According to Lemma 2.6.(1) we have $\mathbb{G}_m(B)/R = 1$. Then Lemma 2.4 shows that $R\mathfrak{Q}(B) = \mathfrak{Q}(B)$, hence the inclusion $\pi(\mathfrak{Q}(B)) \subseteq R\mathfrak{T}(B)$. It follows that we deal with a surjection $\mathfrak{T}(B)/\pi(\mathfrak{Q}(B)) \to \mathfrak{T}(B)/R$.

According to Proposition 3.4, we have an isomorphism $\mathfrak{T}(B/I)/\pi(\mathfrak{Q}(B/I)) \xrightarrow{\sim} \mathfrak{T}(B/I)/R$. Then the isomorphism $\mathfrak{T}(B)/\pi(\mathfrak{Q}(B)) \to \mathfrak{T}(B/I)/\pi(\mathfrak{Q}(B/I)) = \mathfrak{T}(B/I)/R$ factors through $\mathfrak{T}(B)/R$, and we conclude that $\mathfrak{T}(B)/\pi(\mathfrak{Q}(B)) = \mathfrak{T}(B)/R$ as well. \Box

7.2. A generalization. Using the case of tori, we obtain the following partial result for R-equivalence of arbitrary reductive groups. We do it by generalizing an argument of Raghunathan [58, §1].

Lemma 7.4. We assume that B/I is a regular domain. Let \mathfrak{G} be a reductive A-group scheme admitting B-subtori $\mathfrak{T}_1, \ldots, \mathfrak{T}_n$ such that $\operatorname{Lie}(\mathfrak{G})(B)$ is generated as B-module by the $\operatorname{Lie}(\mathfrak{T}_i)(B)$'s. Then $\operatorname{ker}(\mathfrak{G}(B) \to \mathfrak{G}(B/I)) \subseteq R\mathfrak{G}(B)$.

Proof. We consider the map of B-schemes

32

 $f:\mathfrak{T}_1\times_B\cdots\times_B\mathfrak{T}_n \quad \to \quad \mathfrak{G}$

 $(t_1,\ldots,t_n) \qquad \mapsto \qquad t_1\ldots t_n.$

For each maximal ideal \mathfrak{m} of B, The differential at $1_{B/\mathfrak{m}}$ is

$$df_{1_k}$$
: Lie $(\mathfrak{T}_1)(B/\mathfrak{m}) \oplus \cdots \oplus$ Lie $(\mathfrak{T}_n)(B/\mathfrak{m}) \longrightarrow$ Lie $(G)(B/\mathfrak{m})$

$$(X_1, \dots, X_n) \qquad \qquad \mapsto \qquad X_1 + \dots + X_n$$

which is onto by construction. It follows that the map f is smooth at $1_{B/\mathfrak{m}}$ for each maximal ideal of \mathfrak{m} . The Jacobian criterion shows that f is smooth in the neighborhood of the unit section of $\mathfrak{T}_1 \times_B \cdots \times_B \mathfrak{T}_n$. The Hensel lemma [30, th. I.8] (see also [10, prop. 3.1.1]) shows that the induced map

$$\ker \left((\mathfrak{T}_1 \times_B \cdots \times_B \mathfrak{T}_n)(B) \to (\mathfrak{T}_1 \times_B \cdots \times_B \mathfrak{T}_n)(B/I) \right) \to \ker \left(\mathfrak{G}(B) \to \mathfrak{G}(B/I) \right)$$

is surjective. The torus case Lemma 7.3 shows that $\ker(\mathfrak{T}_i(B) \to \mathfrak{T}_i(B/I)) \subseteq R\mathfrak{T}_i(B)$ for i = 1, ..., n. Thus $\ker(\mathfrak{G}(B) \to \mathfrak{G}(B/I)) \subseteq R\mathfrak{G}(A)$.

Together with Lemma 3.2.(1), we get the following fact.

Corollary 7.5. Let R be a semilocal ring with infinite residue fields and let \mathfrak{G} be a reductive R-group scheme assumed R-linear. Let (B, J) be a henselian pair where B is an R-algebra such that B/J is a regular domain. Then ker $(\mathfrak{G}(B) \to \mathfrak{G}(B/J)) \subseteq R\mathfrak{G}(B)$.

7.3. The semisimple case. We continue with the henselian pair (A, I). One evidence is the case of the group $\mathrm{SL}_N(\mathcal{A})$ for an Azumaya A-algebra \mathcal{A} of degree invertible in A^{\times} for N >> 0 since Hazrat has proven that the map $\mathrm{SK}_1(\mathcal{A}) \to \mathrm{SL}_1(\mathcal{A}/I)$ is an isomorphism, if A is semilocal [31]. Firstly we make a variation on [26, §3.4].

Lemma 7.6. Let F be a field and let G be a reductive F-group. Let P be a strictly proper parabolic F-subgroup and let P^- be an opposite parabolic subgroup to P. We put $U = \operatorname{rad}_u(P)$ and $U^- = \operatorname{rad}_u(P^-)$. We consider the following commutative diagram

$$0 \longrightarrow \operatorname{Lie}(G) \longrightarrow G(F[\epsilon]) \longrightarrow G(F) \longrightarrow 1$$
$$\bigcup_{E_P(F[\epsilon]) \longrightarrow E_P(F) \longrightarrow 1}$$

and define $V_P = \ker(E_P(F[\epsilon]) \to E_P(F)) \subseteq \operatorname{Lie}(G)$.

(1) The F-subspace V_P is an ideal of Lie(G) which is G(F)-stable. We have $V_P = E_P(F)$. $\text{Lie}(U) + E_P(F)$. $\text{Lie}(U^-)$.

(2) If G is semisimple simply connected, we have $V_P = \text{Lie}(G)$.

Proof. (1) It follows from [64, XXVI.5.1] that $E_P(F[\epsilon])$ (resp. $E_P(F)$) is a normal subgroup of $G(F[\epsilon])$ (resp. G(F)). It implies that V_P is a Lie subalgebra of Lie(G) which is furthermore G(F)-equivariant. Since Lie(U), Lie(U^-) are contained in V_P , it follows that $E_P(F)$. Lie(U) + $E_P(F)$. Lie(U^-) $\subseteq V_P$. Conversely, we are given an element $v \in V_P$. It is of the shape $v = u_1 u_2 \dots u_{2n}$ with $u_{2i+1} \in U(F[\epsilon])$ and $u_{2i} \in U^-(F[\epsilon])$. We have a decomposition $v = v_1(g_2 v_2 g_2^{-1}) \dots (g_{2n} v_{2n} g_{2n}^{-1})$ with $v_{2i+1} \in$ Lie(U), $v_{2i} \in \text{Lie}(U^-)$ and $g_1, \dots, g_{2n} \in E_P(F)$. We have proven that v belongs to $E_P(F)$. Lie(U) + $E_P(F)$. Lie(U^-).

(2) Without loss of generality we can assume that G is almost absolutely F-simple. If F is infinite, we have that $E_P(F)$. Lie(U) = Lie(G) according to [26, lemma 3.3.(3)] so a fortiori $V_P = \text{Lie}(G)$. We can then assume that F is finite so that G is quasi-split. We have then $E_P(F) = G(F)$ according to [70, 1.1.2]. If G is split, the statement is [26, lemma 3.3.(1)]. It remains to deal with the quasi-split non split case, it implies that G is of outer type A, D or E_6 . In particular, all geometrical roots have same length and G is not of type A_1 . Furthermore we can deal with Borel subgroup B of P(according to [54, remark after Theorem 1]). Let T be maximal torus of B, we recall the decomposition $\text{Lie}(G) = \text{Lie}(T) \oplus \text{Lie}(U) \oplus \text{Lie}(U^-)$.

We consider the ideal $V_P \otimes_F F_s$ of $\text{Lie}(G) \otimes_F F_s$. According to [32, prop. 2.6.a], $V_P \otimes_F F_s$ is central or contains $\text{Lie}(T) \otimes_F F_s$. Since V_P is not central, we get that $\text{Lie}(T) \subseteq V_P$. Since $\text{Lie}(U), \text{Lie}(U^-) \subset \text{Lie}(G)$, we conclude that $V_P = \text{Lie}(G)$. \Box

Proposition 7.7. Let R be a semilocal ring and let \mathfrak{G} be a semisimple group scheme over B, such that its simply connected cover morphism $f : \mathfrak{G}^{sc} \to \mathfrak{G}$ is smooth. We assume that \mathfrak{G} has a strictly proper parabolic R-subgroup \mathfrak{P} . Let (B, I) be an henselian couple where B is an R-algebra. Then the map $K_1^{\mathfrak{G},\mathfrak{P}}(B) \to K_1^{\mathfrak{G},\mathfrak{P}}(B/I)$ is an isomorphism.

Proof. Let \mathfrak{P}^- be a parabolic *R*-subgroup of \mathfrak{G} , opposite to \mathfrak{P} . Let $\mathfrak{U} = \operatorname{rad}(\mathfrak{P})$, $\mathfrak{U}^- = \operatorname{rad}(\mathfrak{P}^-)$.

Since \mathfrak{G} is affine smooth, the map $\mathfrak{G}(B) \to \mathfrak{G}(B/I)$ is surjective according to the generalization of Hensel's lemma to henselian couples [30, th. I.8] (see also [10, prop. 3.1.1]), hence $K_1^{\mathfrak{G},\mathfrak{P}}(B) \to K_1^{\mathfrak{G},\mathfrak{P}}(B/I)$ is onto. To show that it is injective, it is enough to prove that ker($\mathfrak{G}(B) \to \mathfrak{G}(B/I)$) $\leq E_{\mathfrak{P}}(B)$, since $E_{\mathfrak{P}}(B)$ surjects onto $E_{\mathfrak{P}}(B/I)$.

Combining the lifting method of [26, lemma 3.5] and Lemma 7.6, there exist $g_1, \ldots, g_{2m} \in E_{\mathfrak{P}}(R)$ such that the product map

 $h: (\mathfrak{U} \times \mathfrak{U}^{-})^{m} \to \mathfrak{G}, \ (u_{1}, \dots, u_{2m}) \mapsto {}^{g_{1}}\!u_{1} \dots {}^{g_{2m}}\!u_{2m}$

is smooth at all $(1, ..., 1)'_{\kappa_i}s$. Then *h* is smooth in the neighborhood of the origin of $(\mathfrak{U} \times \mathfrak{U}^-)^m$. Hensel's lemma yields $\ker(\mathfrak{G}(B) \to \mathfrak{G}(B/I)) \leq E_{\mathfrak{P}}(B)$. \Box

8. Specialization for R-equivalence

8.1. The case of tori. Let A be a henselian local ring with the maximal ideal m and the residue field k. We remind the reader that a torus over a henselian local ring

is isotrivial, this follows of the equivalence of categories between that of A-tori and that of A/m-tori [64, X.4.1] and from the fact that a torus over a field is isotrivial.

Proposition 8.1. Let A be a local ring of residue field k. Let \mathfrak{T} be an A-torus and put $T = \mathfrak{T} \times_A k$. Then

(1) If A is henselian, then the natural map $\mathfrak{T}(A)/R \to \mathfrak{T}(k)/R$ is an isomorphism. In particular we have ker $(\mathfrak{T}(A) \to T(k)) \subseteq R\mathfrak{T}(A)$.

(2) If A is regular and K denotes the fraction field of A, then the natural map $\mathfrak{T}(A)/R \longrightarrow \mathfrak{T}(K)/R$ is an isomorphism.

Proof. (1) Let m be the maximal ideal of A. Since A is henselian, (A, m) is a Henselian pair. Then, since A/m = k is a regular domain, by Lemma 7.3 $\mathfrak{T}(A)/R \to T(k)/R$ is an isomorphism.

(2) We consider a flasque resolution

$$1 \to \mathfrak{S} \to \mathfrak{Q} \xrightarrow{\pi} \mathfrak{T} \to 1.$$

According to Proposition 3.4, we have isomorphisms $\mathfrak{T}(A)/\pi(\mathfrak{Q}(A)) \xrightarrow{\sim} \mathfrak{T}(A)/R \xrightarrow{\sim} H^1(A,\mathfrak{S})$ and $\mathfrak{T}(K)/\pi(\mathfrak{Q}(K)) \xrightarrow{\sim} \mathfrak{T}(K)/R \xrightarrow{\sim} H^1(K,\mathfrak{S})$. Since \mathfrak{S} is flasque, the restriction map $H^1(A,\mathfrak{S}) \to H^1(K,\mathfrak{S})$ is surjective [15, Th. 2.2] and is injective (*ibid*, Th. 4.1). Thus the map $\mathfrak{T}(A)/R \to \mathfrak{T}(K)/R$ is an isomorphism. \Box

Corollary 8.2. We assume that the henselian local ring A is regular noetherian with residue field k and fraction field K. For any A-torus \mathfrak{T} we have two isomorphisms

$$T(k)/R \xleftarrow{\sim} \mathfrak{T}(A)/R \xrightarrow{\sim} \mathfrak{T}(K)/R.$$

8.2. Reduction to the anisotropic case. We come back to the setting of the introduction where A is a henselian local domain of residue field k and fraction field K.

Let \mathfrak{G} be a reductive A-group scheme. Let \mathfrak{P} be a parabolic A-subgroup of \mathfrak{G} and let \mathfrak{L} be a Levi subgroup of \mathfrak{P} . We know that $\mathfrak{L} = Z_{\mathfrak{G}}(\mathfrak{S})$ where \mathfrak{S} is the maximal central A-split subtorus \mathfrak{S} of \mathfrak{L} [64, XXVI]. We put $G = \mathfrak{G} \times_A k$, $P = \mathfrak{P} \times_A k$ and define similarly L and S. According to Corollary 3.8, we have the following commutative diagram where horizontal maps are isomorphisms

By diagram chase, we get the following facts.

Lemma 8.3. (1) If $(\mathfrak{L}/\mathfrak{S})(A)/R \to (L/S)(k)/R$ is injective, then $\mathfrak{G}(A)/R \to G(k)/R$ and $\mathfrak{L}(A)/R \to L(k)/R$ are isomorphisms.

(2) If $(\mathfrak{L}/\mathfrak{S})(A)/R \to (\mathfrak{L}/\mathfrak{S})(K)/R$ is injective (resp. surjective, resp. isomorphism), then $\mathfrak{G}(A)/R \to \mathfrak{G}(K)/R$ is injective (resp. surjective, resp. an isomorphism) and the map $\mathfrak{L}(A)/R \to \mathfrak{L}(K)/R$ is injective (resp. surjective, resp. an isomorphism).

Proof. Since \mathfrak{G} and \mathfrak{L} are smooth A-schemes and A is henselian, the maps $\mathfrak{G}(A)/R \to G(k)/R$ and $\mathfrak{L}(A)/R \to L(k)/R$ are surjective. The rest follows from Corollary 3.8.


It follows that the specialization problem reduces to the case of \mathfrak{L} and even to $\mathfrak{L}/\mathfrak{S}$. In particular, if \mathfrak{P} is minimal, then $\mathfrak{L}/\mathfrak{S}$ is anisotropic.

8.3. The lifting map.

Lemma 8.4. Let A be a henselian local ring with residue field k and let \mathfrak{G} be a reductive A-group. Then ker $(\mathfrak{G}(A) \to G(k)) \subseteq R\mathfrak{G}(A)$.

Proof. If k is infinite, the claim follows from Corollary 7.5. If k is finite, then \mathfrak{G} is quasi-split by [64, XXVI.7.15] combined with the Lang's theorem. Then one has $\mathfrak{G}(A)/R = \mathfrak{G}(k)/R = 1$ by Gauss decomposition [64, XXVI.5.1] combined with the fact that quasi-split tori over A and k are R-trivial.

The above lemma shows that the map $\mathfrak{G}(A) \to \mathfrak{G}(A)/R$ factorizes through G(k), i.e. defines a surjective homomorphism $\phi : G(k) \to \mathfrak{G}(A)/R$. One way to prove that the map $\mathfrak{G}(A)/R \to G(k)/R$ is an isomorphism would be to show that ϕ factorizes through G(k)/R, that is to complete the following diagram



The dotted map is called (when it exists) <u>the lifting map</u>. In what follows we prove the existense of the lifting map in two different cases.

Proposition 8.5. Let A be a henselian local ring with the residue field k, and let \mathfrak{G} be a reductive group over A. Assume that A is equicharacteristic, i.e. A contains a field. Then $\mathfrak{G}(A)/R \to G(k)/R$ is an isomorphism.


Proof. By Lemma 9.4 A is a filtered direct limit of henselian local rings A_i such that the map from A_i to its residue field admits a section. Since \mathfrak{G} is finitely presented over A, and the functor $\mathfrak{G}(-)/R$ commutes with filtered direct limits by Lemma 2.3, we can assume from the start that $A \to k$ admits a section.

We have $\ker(\mathfrak{G}(A) \to G(k)) \subseteq R\mathfrak{G}(A)$ by Lemma 8.4. Since $A \to k$ admits a section, the map $R\mathfrak{G}(A) \to RG(k)$ is surjective. These two statements together imply that $\mathfrak{G}(A)/R \to G(k)/R$ is injective. The surjectivity is obvious.

Theorem 8.6. Let A be a henselian local ring with residue field k, let \mathfrak{G} be a semisimple group scheme over A, such that its simply connected cover morphism $f : \mathfrak{G}^{sc} \to \mathfrak{G}$ is smooth. We assume that \mathfrak{G} has a strictly proper parabolic subgroup \mathfrak{P} .

(1) The map $K_1^{\mathfrak{G},\mathfrak{P}}(A) \to K_1^{G,P}(k)$ is an isomorphism.

(2) If $\mathfrak{G} = \mathfrak{G}^{sc}$, we have a square of isomorphisms



(3) Assume furthermore that A is a domain with fraction field K. There is a natural lifting map $K_1^{G,P}(k) \to K_1^{\mathfrak{G},\mathfrak{P}}(A) \to K_1^{\mathfrak{G},\mathfrak{P}}(K).$

Proof. (1) This is a special case of Proposition 7.7.

(2) If $\mathfrak{G} = \mathfrak{G}^{sc}$, we have the following commutative diagram


where the right vertical isomorphism is [23, th. 7.2]. Since the left vertical map is onto, a diagram chase shows that all maps are isomorphisms.

(3) It is a straightforward consequence.

8.4. The case of DVRs. Assume that A is a henselian DVR and \mathfrak{G} is a reductive group over A. We remind the reader of the existence of a specialization map

$$\varphi: \mathfrak{G}(K)/R \to G(k)/R$$

which is characterized by the property $\varphi([g]) = [\overline{g}]$ for all $g \in G(A)$ [22, th. 0.2]. In other words we have a commutative diagram



This is based on the existence of a specialization map $\mathfrak{X}(A)/R \to \mathfrak{X}(k)/R$ for a projective A-scheme \mathfrak{X} due to Kollár [36] and Madore [39], see also [11, th. 6.1].

Remark 8.7. The quoted reference [22, th. 0.2] requires the assumption that k is not of characteristic 2. This assumption occurs only in the de Concini–Procesi construction of the wonderful compactification of an adjoint semisimple A-group scheme. This is folklore that we can get rid of this assumption by a refinement of [17, th. 3.13]. By descent, the relevant case is that of adjoint Chevalley groups over \mathbb{Z} which is used for example in [61]. Note also that in the field case, there is a construction of the wonderful compactification in [7, §6.1].

Remark 8.8. The existence of the specialization map in the reductive case over a DVR has been established by another method by Colliot-Thèlène, Harbater, Hartmann, Krashen, Parimala, and Suresh which involves simpler compactifications [12, Th. A.10].

Lemma 8.9. Let A be a henselian DVR. For any reductive group \mathfrak{G} over A the map $\mathfrak{G}(A)/R \to \mathfrak{G}(K)/R$ is surjective.

Proof. First case: $G = \mathfrak{G}_k$ is irreducible (that is G is the only parabolic k-subgroup of G). Let S be the maximal central split subtorus of \mathfrak{G}_k . It lifts to a central split subtorus \mathfrak{S} of \mathfrak{G} [64, XI]. Since G/S is anisotropic, we have $(\mathfrak{G}/\mathfrak{S})(A) = (\mathfrak{G}/\mathfrak{S})(K)$ [9]. Hilbert 90 theorem yields $\mathfrak{G}(A)/\mathfrak{S}(A) = \mathfrak{G}(K)/\mathfrak{S}(K)$ hence a decomposition $\mathfrak{G}(K) = \mathfrak{S}(K)\mathfrak{G}(A)$. Since $R\mathfrak{S}(K) = \mathfrak{S}(K)$, we conclude that $\mathfrak{G}(K) = \mathfrak{G}(A)R\mathfrak{G}(K)$.

General case. Let \mathfrak{P} be a minimal parabolic A-subgroup of \mathfrak{G} . Let \mathfrak{P}^- be an opposite parabolic A-subgroup scheme to \mathfrak{P} . Then the Levi subgroup $\mathfrak{L} = \mathfrak{P} \cap \mathfrak{P}^-$ is such that $L = \mathfrak{L}_k$ is irreducible. Let \mathfrak{G} be the maximal central split subtorus of \mathfrak{L} . The first case shows that $(\mathfrak{L}/\mathfrak{S})(A)/R \to (\mathfrak{L}/\mathfrak{S})(K)/R$ is surjective. By Lemma 8.3 this implies the surjectivity of $\mathfrak{G}(A)/R \to \mathfrak{G}(K)/R$. \Box


Proposition 8.10. Let A be a henselian DVR. Let k be the residue field of A and let K be the fraction field of A. Let \mathfrak{G} be a semisimple simply connected A-group scheme having a strictly proper parabolic A-subgroup \mathfrak{P} . Then we have the following commutative diagram of isomorphisms

(8.4)
$$K_{1}^{\mathfrak{G},\mathfrak{P}}(k) \xleftarrow{\sim} K_{1}^{\mathfrak{G},\mathfrak{P}}(A) \xrightarrow{\sim} K_{1}^{\mathfrak{G},\mathfrak{P}}(K)$$
$$\downarrow^{\wr} \qquad \qquad \downarrow^{\wr} \qquad \qquad \downarrow^{\wr}$$
$$\mathfrak{G}(k)/R \xleftarrow{\sim} \mathfrak{G}(A)/R \xrightarrow{\sim} \mathfrak{G}(K)/R$$

Proof. By Theorem 8.6 we have that $\mathfrak{G}(A)/R \to \mathfrak{G}(k)/R$ is an isomorphism. Then it follows from the existence of specialization map (8.3) that $\mathfrak{G}(A)/R \to \mathfrak{G}(K)/R$ is injective. By Lemma 8.9 the map $\mathfrak{G}(A)/R \to \mathfrak{G}(K)/R$ is surjective.

38

Consider the commutative diagram



The bottom horizontal map is an isomorphism as we have used several times [23] and the left vertical map is an isomorphism in view of Proposition 7.7. It follows that $K_1^{\mathfrak{G},\mathfrak{P}}(A) \to \mathfrak{G}(A)/R$ is injective and then an isomorphism. The remaining isomorphisms follow immediately.

Remark 8.11. The surjectivity of the map $K_1^{\mathfrak{G},\mathfrak{P}}(A) \to K_1^{\mathfrak{G},\mathfrak{P}}(K)$ was previously proved in [23, lemme 4.5.1]. Note that it is does not hold for A = k[[t]] and $\mathfrak{G} = \mathrm{GL}_n$ or PGL_n , so seems specific to the semisimple simply connected case.

8.5. Specialization in the equicharacteristic case. Assume that A is a complete regular local ring containing a field k_0 and let K be its fraction field. According to [29, 1.19.6.4], A is k_0 -isomorphic (non-canonically) to a formal series ring $k[[t_1, \ldots, t_d]]$, where k is the residue field of A.

Let \mathfrak{G} be a reductive A-group scheme. There exists a unique reductive k-group G such that $\mathfrak{G} \times_A k[[t_1, \ldots, t_d]] \cong G \times_k k[[t_1, \ldots, t_d]]$ (see the proof of Corollary 8.13 below). Since the fraction field $K = k((t_1, \ldots, t_d))$ of A is a (proper) subfield of the iterated Laurent power series field $k((t_1)) \ldots ((t_d))$, and

$$G(k)/R \to G(k((t_1))\dots((t_d)))/R$$

is an isomorphism [22, cor. 0.3], we can define a specialization map $G(K)/R \rightarrow G(k)/R$ inductively,

$$sp: G(K)/R \to G(k((t_1,\ldots,t_d)))/R \to G(k((t_1,\ldots,t_{d-1})))/R \to \cdots \to G(k)/R$$

However, it is unclear whether this map does not depend of the choice of coordinates t_1, \ldots, t_d . The following theorem solves this problem.

Theorem 8.12. Let k be an arbitrary field. Then for any reductive group G over k and any $d \ge 1$ the natural maps

$$G(k)/R \to G(k[[t_1,\ldots,t_d]])/R \to G(k((t_1,\ldots,t_d)))/R$$

are isomorphisms.

Proof. We set $A = k[[t_1, \ldots, t_d]]$ and $K = k((t_1, \ldots, t_d))$. By Proposition 8.5 we have the isomorphism $G(k)/R \xrightarrow{\sim} G(A)/R$. Theorem 5.4 shows that $G(A)/R \to G(K)/R$ is onto. It remains to prove that the surjective map $G(k)/R \to G(K)/R$ is an isomorphism and we know that it holds in the one dimensional case, i.e. the map $G(k)/R \to G(k((t)))/R$ is an isomorphism [22, cor. 0.3]. Using the embedding

$$K = k((t_1, \dots, t_d)) \hookrightarrow k((t_1)) \dots ((t_d))$$

we get that $G(k)/R \to G(K)/R$ is injective.

Corollary 8.13. Let (A, m) be a complete regular local ring containing a field k_0 . Let k be the residue field of A, and let K be the fraction field of A. Let \mathfrak{G} be a reductive group scheme over A. Then we have two isomorphisms

$$\mathfrak{G}(k)/R \xleftarrow{\sim} \mathfrak{G}(A)/R \xrightarrow{\sim} \mathfrak{G}(K)/R.$$

Proof. According to [29, 1.19.6.4], A is k_0 -isomorphic (non-canonically) to a formal series ring $k[[t_1, \ldots, t_d]]$, where k is the residue field of A. The group \mathfrak{G} is the twisted A-form of a Chevalley reductive group \mathbb{Z} -scheme \mathfrak{G}_0 by a Aut(\mathfrak{G}_0)-torsor \mathfrak{E} . Let $G = \mathfrak{G} \times_A k$ be the restriction of \mathfrak{G} via the residue homomorphism $A \to k$. Since $H^1(\widehat{A}, \operatorname{Aut}(\mathfrak{G}_0)_k) \xrightarrow{\sim} H^1(k, \operatorname{Aut}(\mathfrak{G}_0)_k)$ [64, XXIV.8.1], it follows that \mathfrak{G} is isomorphic to $G \times_k k[[t_1, \ldots, t_d]]$. Then we can apply Theorem 8.12.

Theorem 8.14. Let A be a henselian regular local ring containing a field k_0 . Let k be the residue field of A and let K be the fraction field of A. Let \mathfrak{G} be a reductive group scheme over A. Then we have two isomorphisms

$$\mathfrak{G}(k)/R \xleftarrow{\sim} \mathfrak{G}(A)/R \xrightarrow{\sim} \mathfrak{G}(K)/R$$

In particular, we have a well-defined specialization map $sp: \mathfrak{G}(K)/R \to \mathfrak{G}(k)/R$ and it is an isomorphism.

Proof. By Lemma 9.4 A is a filtered direct limit of henselian regular local rings A_i such that each A_i contains a field and the map from A_i to its residue field admits a section. Since the group scheme \mathfrak{G} and its parabolic subgroups are finitely presented over A, and the functor $\mathfrak{G}(-)/R$ commutes with filtered direct limits, we can assume from the start that $A \to k$ admits a section. Since A is henselian, we have a bijection $H^1(A, \operatorname{Aut}(\mathfrak{G}_0)_k) \xrightarrow{\sim} H^1(k, \operatorname{Aut}(\mathfrak{G}_0)_k)$ [64, XXIV.8.1]. Since $A \to k$ has a section, it follows that \mathfrak{G} is isomorphic to $\mathfrak{G}_k \times_k A$. Clearly, \mathfrak{G} is isotropic if and only if \mathfrak{G}_k is isotropic. Then by Proposition 8.5 $G(A)/R \to G(k)/R$ is an isomorphism and by Theorem 5.4 $\mathfrak{G}(A)/R \to \mathfrak{G}(K)/R$ is surjective.

Let \hat{A} be the completion of A at the maximal ideal and let \hat{K} be its fraction field. Then \hat{A} is a complete regular local ring containing k_0 and k is its residue field. By Corollary 8.13 the maps $G(\hat{A})/R \to G(k)/R$ and $G(\hat{A})/R \to G(\hat{K})/R$ are isomorphisms. Hence $G(A)/R \to G(\hat{A})/R$ is an isomorphism, and consequently $G(A)/R \to G(K)/R$ is injective.

Corollary 8.15. Let B be a regular local ring containing a field k_0 , let L be the fraction field of B and let l be the residue field of B. Let \mathfrak{G} be a reductive group scheme over B. There is a well-defined specialization homomorphism $\mathfrak{G}(L)/R \to \mathfrak{G}(l)/R$.

Proof. Let \hat{B} denote the completion of B with respect to the maximal ideal, and let \hat{L} denote the fraction field of \hat{B} . By Theorem 8.14 the natural maps $\mathfrak{G}(\hat{B})/R \to \mathfrak{G}(l)/R$ and $\mathfrak{G}(\hat{B})/R \to \mathfrak{G}(\hat{L})/R$ are isomorphisms. The specialization map is then induced by these isomorphisms together with the map $\mathfrak{G}(L)/R \to \mathfrak{G}(\hat{L})/R$.

Remark 8.16. Colliot-Thèlène, Harbater, Hartmann, Krashen, Parimala, and Suresh have constructed a specialization homomorphism for arbitrary regular local rings of dimension 2 [12, Prop. A.12].

9. Appendices

9.1. The big Bruhat cell is a principal open subscheme. For split groups and Borel subgroups, this statement goes back to Chevalley, see [8, lemma 4.5].

Lemma 9.1. Let B be a ring and let \mathfrak{G} be a reductive group B-scheme equipped with a pair of opposite parabolic B-subgroups \mathfrak{P}^{\pm} . Then the big cell Ω of \mathfrak{G} attached to \mathfrak{P} and \mathfrak{P}^{-} is a principal open subscheme of \mathfrak{G} . More precisely, there exists $f \in B[\mathfrak{G}]$ such that $\Omega = \mathfrak{G}_{f}$ and f can be chosen $\operatorname{Aut}(\mathfrak{G}, \mathfrak{P}, \mathfrak{P}^{-})$ -invariant.

Proof. Without loss of generality, we can assume that \mathfrak{G} is adjoint. We can assume B noetherian and connected so that $(\mathfrak{G}, \mathfrak{P}, \mathfrak{P}^-)$ is a B-form of $(\mathfrak{G}_0, \mathfrak{P}_0, \mathfrak{P}_0^-)_B$ where \mathfrak{G}_0 is an adjoint Chevalley \mathbb{Z} -group scheme equipped with opposite parabolic \mathbb{Z} -group subschemes $(\mathfrak{P}_0, \mathfrak{P}_0^-)$ related to the Chevalley pinning.

Then $(\mathfrak{G}, \mathfrak{P}, \mathfrak{P}^-)$ is the twist of $(\mathfrak{G}_0, \mathfrak{P}_0, \mathfrak{P}_0^-)_B$ by an $\operatorname{Aut}(\mathfrak{G}_0, \mathfrak{P}_0, \mathfrak{P}_0^-)$ -torsor so that the statement boils down to the split case over \mathbb{Z} . We consider the Levi–subgroup $\mathfrak{L}_0 = \mathfrak{P}_0^+ \cap \mathfrak{P}_0^-$ so that $\operatorname{Aut}(\mathfrak{G}_0, \mathfrak{P}_0, \mathfrak{P}_0^-) = \operatorname{Aut}(\mathfrak{G}_0, \mathfrak{P}_0, \mathfrak{L}_0)$ is the semi-direct product of \mathfrak{L}_0 and a finite constant \mathbb{Z} -group scheme Γ [24, lemme 5.1.2].

According to [9, 3.8.2.(a)] there is a function $f_0 \in \mathbb{Z}[\mathfrak{G}_0]$ such that $\mathbb{Z}[\Omega_0] = \mathbb{Z}[\mathbf{G}_0]_{f_0}$ and satisfying $f_0(1) = 1$. We claim that f_0 is \mathfrak{L}_0 -invariant with respect to the adjoint action. We denote by $\Lambda = \operatorname{Hom}_{\overline{\mathbb{Q}}-gr}(\mathfrak{L}_{\overline{\mathbb{Q}}}, \mathbb{G}_m)$ the lattice of characters and remind the reader of Rosenlicht decomposition [59, th. 3]

$$H^0(\mathfrak{L}_{\overline{\mathbb{Q}}},\mathbb{G}_m)=\overline{\mathbb{Q}}^{\times}\oplus\Lambda$$

which shows that $\Lambda = \{ f \in H^0(\mathfrak{L}_{\overline{\mathbb{Q}}}, \mathbb{G}_m) \mid f(1) = 1 \}$. We observe that the induced action (by the adjoint action) of $\mathfrak{L}_0(\overline{\mathbb{Q}})$ on Λ is trivial. It follows that the map

$$\phi: \mathfrak{L}_0(\overline{\mathbb{Q}}) \to \overline{\mathbb{Q}}[\mathfrak{L}_0]^{\times} \to \Lambda, \ x \mapsto {}^x\!f_0 f_0^{-1}$$

is a group homomorphism. Since $\mathfrak{L}_{0,\overline{\mathbb{Q}}}$ is generated by its maximal tori, we have $\mathfrak{L}_0(\overline{\mathbb{Q}}) = \langle \mathfrak{L}_0(\overline{\mathbb{Q}})^n \rangle$ for all $n \geq 1$. We get that ϕ is zero and this establishes the above claim. Taking the product of Γ -conjugates of f_0 permits to assume that f_0 is $\operatorname{Aut}(\mathfrak{G}_0,\mathfrak{P}_0,\mathfrak{L}_0)$ -invariant. By descent, f_0 gives rise to then to $f \in B[\mathfrak{G}]$ so that $\Omega = \mathfrak{G}_f$.

9.2. Colliot-Thélène and Ojanguren method for functors in pointed sets. In this section we summarize the classic injectivity theorem of Colliot-Thélène and Ojanguren [13, th. 1.1]. Our goal is to make explicit the fact that a certain intermediate step in the proof of this theorem holds under weaker assumptions than the theorem itself. Let k be an infinite field and let $R \mapsto F(R)$ be a covariant functor on the category of k-algebras (commutative, unital) with values in pointed sets. We consider the following properties:

 (\mathbf{P}_1) The functor F commutes with filtered direct limits of k-algebras having flat transition morphisms.

 (\mathbf{P}_2) For each k-field E and for each $n \ge 1$, the map

$$F(E[t_1,\ldots,t_n]) \to F(E(t_1,\ldots,t_n))$$

has trivial kernel;

(P₃) (Patching property) For each finite type flat inclusion $A \hookrightarrow B$ of noetherian integral k-algebras and each non-zero element $f \in A$ such that $A/fA \xrightarrow{\sim} B/fB$, then the map

$$\operatorname{Ker}(F(A) \to F(A_f)) \to \operatorname{Ker}(F(B) \to F(B_f))$$

is onto.

One may consider the following weaker property.

 (\mathbf{P}'_3) (Zariski patching) For each noetherian integral k-algebra A and for each decomposition A = Af + Ag with f non-zero, then the map

$$\operatorname{Ker}(F(A) \to F(A_f)) \to \operatorname{Ker}(F(A_g) \to F(A_{fg}))$$

is onto.

We have $\mathbf{P}_3 \Longrightarrow \mathbf{P}'_3$ by taking $B = A_g$ since we have $B_f = A_{fg}$ and $A/fA \xrightarrow{\sim} B/fB$.

The following theorem was proved by Colliot-Thélène and Ojanguren.

Theorem 9.2. [13, th. 1.1] We assume that F satisfies \mathbf{P}_1 , \mathbf{P}_2 and \mathbf{P}_3 . Let A be a local ring of a smooth L-ring C where L is a k-field. Denote by K the fraction field of A. Then the map $F(A) \to F(K)$ has trivial kernel.

The proof of this theorem relies on the following result.

Proposition 9.3. [13, prop. 1.5] We assume that F satisfies \mathbf{P}_1 , \mathbf{P}_2 and \mathbf{P}'_3 . Let A be the local ring at a prime ideal of a polynomial algebra $L[t_1, \ldots, t_d]$ where L is a k-field. Denote by K the fraction field of A. Then for each integer $n \geq 0$, the map

$$F(A[x_1,\ldots,x_n]) \to F(K(x_1,\ldots,x_n))$$

has trivial kernel.

Proof. The original statement of [13, prop. 1.5] assumes that F satisfies \mathbf{P}_1 , \mathbf{P}_2 and \mathbf{P}_3 , and that A is a maximal localization of $L[t_1, \ldots, t_d]$. The inspection of the proof shows that instead of property \mathbf{P}_3 , only the Zariski patching property \mathbf{P}'_3 was used. Furthermore, since every prime ideal of $L[t_1, \ldots, t_d]$ is an intersection of maximal ideals, and F satisfies \mathbf{P}_1 , the case where A is a localization at a prime ideal follows from the case of maximal localizations [13, p. 101, Première réduction].

42

9.3. Fields of representants for henselian regular local rings. The following fact was brought to our attention by K. Česnavičius.

Lemma 9.4. Let A be a henselian local ring containing a field k_0 . Then A is a filtered direct limit of henselian local rings A_i such that the map from A_i to its residue field admits a section. If A is moreover regular, then the henselian local rings A_i can be chosen regular as well.

Proof. Clearly, we can assume that k_0 is a finite field or \mathbb{Q} without loss of generality. The local ring A is a filtered direct limit of local rings C_i that are localizations of finitely generated k_0 -algebras contained in A. Since A is henselian, we can replace each C_i by its henselization $A_i = (C_i)^h$. Let $k_i = A_i/m_i$ be the residue field of A_i . Then k_i is a finitely generated field extension of k_0 . We claim that $A_i \to k_i$ admits a section. Indeed, since k_0 is perfect, it follows that k_i is separably generated over k_0 , that is, k_i is a finite separable extension of a purely transcendental field extension $L = k_0(t_1, \ldots, t_n)$ of k_0 of finite transcendence degree [72, II, §13, Theorem 31]. Choose arbitrary lifts a_1, \ldots, a_n of t_1, \ldots, t_n to A_i . Then $k_0(a_1, \ldots, a_n) \cong L$ is a subfield of A_i that lifts L. The primitive element theorem states that $k_i = L[b] = L/P(t)$ where P is a separable L-polynomial. Seeing now P as a A-polynomial, the Hensel lemma shows that P(t) has a root $a \in A$ which lifts $b \in L$. We define then a L-map $k_i = L[b] \to A$ by mapping b to a. The composite map $k_i \to A \to k_i = L[b]$ is the identity as desired.

If A is a regular henselian local ring, note that the embedding $k_0 \to A$ is geometrically regular, since k_0 perfect [43, (28.M), (28.N)]. Then by Popescu's theorem [55, 67] A is a filtered direct limit of localizations C_i of smooth k_0 -algebras. Then the henselizations A_i are also regular.

References

- J. Alper, Adequate moduli spaces and geometrically reductive group schemes, Algebr. Geom. 1 (2014), 489–531.
- [2] S. Alsaody, V. Chernousov, A. Pianzola, On the Tits-Weiss Conjecture and the Kneser-Tits Conjecture for E⁷⁸_{7,1} and E⁷⁸_{8,2}, Forum of Mathematics, Sigma 9 (2021), 1–25.
- [3] A. Asok, M. Hoyois, M. Wendt, Affine representability results in A¹-homotopy theory, II: Principal bundles and homogeneous spaces, Geom. Topol. 22 (2018), 1181–1225.
- [4] D.W. Barnes, On Cartan subalgebras of Lie algebras, Math. Z. 101 (1967), 350–355.
- [5] S.M. Bhatwadekar, R.A. Rao, On a question of Quillen, Trans. Amer. Math. Soc. 279 (1983), no. 2, 801–810.
- [6] A. Bouthier, K. Cesnavičius, *Torsors on loop groups and the Hitchin fibration*, to appear in Annales de l'Ecole normale supérieure.
- [7] M. Brion, S. Kumar, Frobenius Splitting Methods in Geometry and Representation Theory, Progress in Mathematics 231, Birkhäuser.
- [8] A. Borel, Properties and linear representations of Chevalley groups. 1970 Seminar on Algebraic Groups and Related Finite Groups, (The Institute for Advanced Study, Princeton, N.J., 1968/69) pp. 1–55, Lecture Notes in Mathematics, Vol. 131 Springer, Berlin.

- [9] F. Bruhat, J. Tits, Groupes algébriques sur un corps local II. Existence d'une donnée radicielle valuée, Publ. Math. Inst. Hautes Études Sci. 60 (1984), 5–184.
- [10] K. Česnavičius, Grothendieck-Serre in the split unramified case, arXiv preprint arXiv:2009.05299 (2020).
- [11] J.-L. Colliot-Thélène, Non rationalité stable sur les corps quelconques, Birational Geometry of Hypersurfaces, Gargnano del Garda, Italy, 2018, A. Hochenegger, M. Lehn, P. Stellari ed., Lecture Notes of the Unione Matematica Italiana, Springer LNM 2019.
- [12] J.-L. Colliot-Thélène, D. Harbater, J. Hartmann, D. Krashen, R. Parimala, V. Suresh Localglobal principles for constant reductive groups over semi-global fields, preprint (2021), arxiv arXiv:2108.12349
- [13] J.-L. Colliot-Thélène, M. Ojanguren, Espaces principaux homogènes localement triviaux, Publ. Math. Inst. Hautes Études Sci. 75 (1992), 97–122.
- [14] J.-L. Colliot-Thélène and J.-J. Sansuc, La R-équivalence sur les tores, Annales Scient. ENS 10 (1977), 175–230.
- [15] J.-L. Colliot-Thélène and J.-J. Sansuc, Principal homogeneous spaces under flasque tori: applications, J. Algebra 106 (1987), 148–205.
- [16] P.M. Cohn, On the structure of GL₂ of a ring, Publ. Math. Inst. Hautes Études Sci. 30 (1966), 365–413.
- [17] C. de Concini, T. A. Springer, Compactification of symmetric varieties, dedicated to the memory of Claude Chevalley, Transform. Groups 4 (1999), 273–300.
- [18] M. Demazure, P. Gabriel, Groupes algébriques, North-Holland (1970).
- [19] R. Fedorov, Affine Grassmannians of group schemes and exotic principal bundles over A¹, Amer. J. Math. 138 (2016), 879–906.
- [20] R. Fedorov, I. Panin, A proof of the Grothendieck-Serre conjecture on principal bundles over regular local rings containing infinite fields, Publ. Math. Inst. Hautes Études Sci. 122 (2015), 169–193.
- [21] P. Gille, Torsors on the affine line, Transform. Groups 7 (2002), 231–245.
- [22] P. Gille, Spécialisation de la R-équivalence pour les groupes réductifs, Trans. Amer. Math. Soc. 35 (2004), 4465–4474.
- [23] P. Gille, Le problème de Kneser-Tits, exposé Bourbaki no 983, Astérisque 326 (2009), 39–81.
- [24] P. Gille, Sur la classification des schémas en groupes semi-simples, "Autour des schémas en groupes, III", Panoramas et Synthèses 47 (2015), 39–110.
- [25] P. Gille, M. Florence, *Residues on Affine Grassmannians*, Journal f
 ür die reine und angewandte Mathematik 776 (2021), 119–150.
- [26] P. Gille, R. Parimala, V. Suresh, Local triviality for G-torsors, Mathematische Annalen 380 (2021), 539-567.
- [27] P. Gille, When is a reductive group scheme linear?, to appear in Michigan Math. Journal.
- [28] A. Grothendieck, J.-A. Dieudonné, Eléments de géométrie algébrique. I, Grundlehren der Mathematischen Wissenschaften 166; Springer-Verlag, Berlin, 1971.
- [29] A. Grothendieck (avec la collaboration de J. Dieudonné), Eléments de Géométrie Algébrique IV, Publ. Math. Inst. Hautes Études Sci. no. 20, 24, 28 and 32 (1964–1967).
- [30] L. Gruson, Une propriété des couples henséliens, Colloque d'Algèbre Commutative (Rennes, 1972), Exp. No. 10, 13 pp; Publ. Sém. Math. Univ. Rennes, 1972.
- [31] R. Hazrat, SK₁ of Azumaya algebras over Hensel pairs, Math. Z. 264 (2010), 295–299.
- [32] G. M. D. Hogeweij, Almost-classical Lie algebras. I, II, Nederl. Akad. Wetensch. Indag. Math. 44 (1982), 441–452, 453–460.
- [33] R. T. Hoobler, The Merkuriev-Suslin theorem for any semi-local ring, Journal of Pure and Applied Algebra 207 (2006), 537–552.

- [34] J.F. Jardine, On the homotopy groups of algebraic groups, J. Algebra 81 (1983), 180–201.
- [35] B. Kahn, K-theory of semi-local rings with finite coefficients and étale cohomology, K-Theory 25 (2002), 99–138.
- [36] J. Kollár, Specialization of zero cycles, Publ. Res. Inst. Math. Sci. 40 (2004), 689–708.
- [37] M. Kontsevich, Yu. Tschinkel, Specialization of birational types, Invent. Math. 217 (2019), no. 2, 415–432.
- [38] M. I. Krusemeyer, Fundamental groups, algebraic K-theory, and a problem of Abhyankar Invent. Math. 19 (1973), 15–47.
- [39] D. Madore, Sur la spécialisation de la R-équivalence, link.
- [40] Y. Manin, Cubic Forms: Algebra, Geometry, Arithmetic, second edition, Elsevier (1986).
- [41] B. Margaux, The structure of the group G(k[t]): variations on a theme of Soulé, Algebra Number Theory **3** (2009), 393–409.
- [42] B. Margaux, Formal torsors under reductive group schemes, Rev. Un. Mat. Argentina 60 (2019), 217–224.
- [43] H. Matsumura, Commutative algebra, Second Edition (Benjamin/Cummings, 1980).
- [44] A. S. Merkurjev, K-theory and algebraic groups, European Congress of Mathematics, Vol. II (Budapest, 1996), 43-72, Progr. Math., 169, Birkhäuser, Basel, 1998.
- [45] J. S. Milne, Algebraic groups, Cambridge Studies in Advanced Mathematics, vol. 170, Cambridge University Press, Cambridge, 2017, The theory of group schemes of finite type over a field.
- [46] F. Morel, A¹-algebraic topology over a field, Lecture Notes in Mathematics, 2052. Springer, Heidelberg, 2012. x+259 pp.
- [47] F. Morel, V. Voevodsky, A¹-homotopy theory of schemes, Publ. Math. Inst. Hautes Études Sci. 90 (1999), 45–143.
- [48] L-F. Moser, Rational triviale Torseure und die Serre-Grothendiecksche Vermutung, Diplomarbeit, 2018.
- [49] J. Nicaise, E. Shinder, The motivic nearby fiber and degeneration of stable rationality, Invent. Math. 217 (2019), no. 2, 377–413.
- [50] I. A. Panin, Nice triples and the Grothendieck-Serre conjecture concerning principal G-bundles over reductive group schemes, Duke Math. J. 168 (2019), 351–375.
- [51] I. A. Panin, Proof of the Grothendieck-Serre conjecture on principal bundles over regular local rings containing a field, Izv. Ross. Akad. Nauk Ser. Mat. 84 (2020), no. 4, 169–186.
- [52] I. Panin, A. Stavrova, N. Vavilov, On Grothendieck-Serre's conjecture concerning principal Gbundles over reductive group schemes: I, Compos. Math. 151 (2015), no. 3, 535–567.
- [53] R. Parimala, J.-P. Tignol, R. M. Weiss, The Kneser-Tits conjecture for groups with Tits-index E⁶⁶_{8.2} over an arbitrary field, Transform. Groups 17 (2012), 209–231.
- [54] V. A. Petrov, A.K. Stavrova, Elementary subgroups in isotropic reductive groups, Algebra i Analiz 20 (2008), 160–188; translation in St. Petersburg Math. J. 20 (2009), 625–644.
- [55] D. Popescu, Letter to the Editor: General Néron desingularization and approximation, Nagoya Math. J. 118 (1990), 45–53.
- [56] D. Quillen, Projective modules over polynomial rings, Invent. Math. 36 (1976), 167–171.
- [57] M. S. Raghunathan, Principal bundles on affine space, C. P. Ramanujam—a tribute, pp. 187–206, Tata Inst. Fund. Res. Studies in Math., 8, Springer, Berlin-New York, 1978.
- [58] M. S. Raghunathan, Principal bundles admitting a rational section, Invent. Math. 116 (1994), 409–423.
- [59] M. Rosenlicht, Toroidal Algebraic Groups, Proc. Amer. Math. Soc. 12 (1961), 984–988.
- [60] D. J. Saltman, Retract rational fields and cyclic Galois extensions, Israel J. Math. 47 (1984), 165–215.

- [61] J. Shalika, R. Takloo-Bighash, Y. Tschinkel, Rational points on compactifications of semi-simple groups, J. Amer. Math. Soc. 20 (2007), 1135–1186.
- [62] A.K. Stavrova, A. Homotopy invariance of non-stable K₁-functors, J. K-Theory 13 (2014), 199–248.
- [63] A.K. Stavrova, \mathbb{A}^1 -invariance of non-stable K_1 -functors in the equicharacteristic case, to appear in Indag. Math.
- [64] Séminaire de Géométrie algébrique de l'I. H. E. S., 1963-1964, schémas en groupes, dirigé par M. Demazure et A. Grothendieck, Lecture Notes in Math. 151-153. Springer (1970).
- [65] Stacks project, https://stacks.math.columbia.edu
- [66] R. Strano, Principal homogeneous spaces over Hensel rings, Proc. Amer. Math. Soc. 87 (1983), 208–212.
- [67] R. G. Swan, Néron-Popescu desingularization, in Algebra and Geometry (Taipei, 1995), Lect. Alg. Geom. 2 (1998), 135–198. Int. Press, Cambridge, MA.
- [68] M. Thakur, Albert algebras and the Tits-Weiss conjecture, preprint (2019), arXiv:1911.04976
- [69] R. W. Thomason, Equivariant resolution, linearization, and Hilbert's fourteenth problem over arbitrary base schemes, Adv. in Math. 65 (1987), no. 1, 16–34.
- [70] J. Tits, Groupes de Whitehead de groupes algébriques simples sur un corps, Séminaire Bourbaki, no. 19 (1976-1977), Talk no. 505, p. 218–236.
- [71] A.E. Tsybyshev, A step toward the non-equicharacteristic Grothendieck-Serre conjecture, Algebra i Analiz **31** (2019), 246-254; translation in St. Petersburg Math. J. **31** (2020), 181–187.
- [72] O. Zariski, P. Samuel, Commutative Algebra I, Graduate Texts in Mathematics, Springer.

UMR 5208 INSTITUT CAMILLE JORDAN - UNIVERSITÉ CLAUDE BERNARD LYON 1 43 BOULE-VARD DU 11 NOVEMBRE 1918 69622 VILLEURBANNE CEDEX - FRANCE Email address: gille@math.univ-lyon1.fr

CHEBYSHEV LABORATORY, ST. PETERSBURG STATE UNIVERSITY, RUSSIA Email address: anastasia.stavrova@gmail.com