

SERRIN-TYPE REGULARITY CRITERIA FOR THE 3D MHD EQUATIONS VIA ONE VELOCITY COMPONENT AND ONE MAGNETIC COMPONENT

HUI CHEN, CHENYIN QIAN, AND TING ZHANG*

ABSTRACT. In this paper, we consider the Cauchy problem to the 3D MHD equations. We show that the Serrin-type conditions imposed on one component of the velocity u_3 and one component of magnetic fields b_3 with

$$u_3 \in L^{p_0,1}(-1, 0; L^{q_0}(B(2))), \quad b_3 \in L^{p_1,1}(-1, 0; L^{q_1}(B(2))),$$

$\frac{2}{p_0} + \frac{3}{q_0} = \frac{2}{p_1} + \frac{3}{q_1} = 1$ and $3 < q_0, q_1 < +\infty$ imply that the suitable weak solution is regular at $(0, 0)$. The proof is based on the new local energy estimates introduced by Chae-Wolf (Arch. Ration. Mech. Anal. 2021) and Wang-Wu-Zhang (arXiv:2005.11906).

Key words: MHD equations; Serrin-type conditions; suitable weak solution; partial regularity; local energy estimates

AMS Subject Classification (2000): 35Q35; 35Q30; 76D03

1. INTRODUCTION

We consider the Cauchy problem to the three-dimensional incompressible magnetohydrodynamics (MHD) equations in $\mathbb{R}^3 \times (0, \infty)$

$$(1.1) \quad \begin{cases} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} - \mu \Delta \mathbf{u} + \nabla \pi = \mathbf{b} \cdot \nabla \mathbf{b}, \\ \partial_t \mathbf{b} + (\mathbf{u} \cdot \nabla) \mathbf{b} - \nu \Delta \mathbf{b} = \mathbf{b} \cdot \nabla \mathbf{u}, \\ \nabla \cdot \mathbf{u} = \nabla \cdot \mathbf{b} = 0, \end{cases}$$

with initial data

$$(1.2) \quad \mathbf{u}|_{t=0} = \mathbf{u}_0, \quad \mathbf{b}|_{t=0} = \mathbf{b}_0.$$

Here $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{b} = (b_1, b_2, b_3)$ and π are nondimensional quantities corresponding to the velocity, magnetic fields and a scalar pressure. $\mu > 0$ is the viscosity coefficient and $\nu > 0$ is the resistivity coefficient. We require $\mu = \nu = 1$ in this paper.

Such system (1.1)–(1.2) describes many phenomena such as the geomagnetic dynamo in geophysics, solar winds and solar flares in astrophysics. G. Duvaut and J. L. Lions [8] constructed a global weak solution and the local strong solution to the initial boundary value problem, and the properties of such solutions have been examined by M. Sermange and R. Temam in [20].

If the magnetic fields $\mathbf{b} = 0$, the system (1.1) degenerates to the incompressible Navier–Stokes equations. A global weak solution to the Navier–Stokes equations was constructed by J. Leray [16]. However, the uniqueness and regularity of such weak solution is still one of the most challenging open problems in the field of mathematical fluid mechanics. One essential

*Corresponding author.

work is usually referred as Serrin–type conditions (see [9, 18, 21, 22] and the references therein.), i.e. if the weak solution \mathbf{u} satisfies

$$(1.3) \quad \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = 1, \quad 3 \leq q \leq \infty,$$

then the weak solution is regular in $(0, T]$. There are several notable results [3, 4, 5, 6, 11, 24] to weaken the above criteria by imposing constraints only on partial components or directional derivatives of velocity field. In particular, D. Chae and J. Wolf [3] made an important progress and obtained the regularity of solution under the condition

$$(1.4) \quad u_3 \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} < 1, \quad 3 < q \leq \infty.$$

W. Wang, D. Wu and Z. Zhang [24] improved to

$$(1.5) \quad u_3 \in L^{p,1}(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = 1, \quad 3 < q < \infty.$$

Throughout this paper, $L^{p,1}$ denotes the Lorentz space with respect to the time variable.

If the magnetic fields $\mathbf{b} \neq 0$, the situation is more complicated due to the coupling effect between the velocity \mathbf{u} and the magnetic fields \mathbf{b} . Some fundamental Serrin–type regularity criteria in term of the velocity only were done in [7, 13, 26, 28]. For instance, if the weak solution (\mathbf{u}, \mathbf{b}) satisfies

$$(1.6) \quad \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = 1, \quad 3 < q \leq \infty,$$

then the solution is regular in $(0, T]$. More regularity criteria can be found in [12, 27] and the references therein.

We now recall the notion of suitable weak solution to the MHD equations (1.1).

Definition 1.1. We say that $(\mathbf{u}, \mathbf{b}, \pi)$ is a suitable weak solution to the MHD equations (1.1) in an open domain $\Omega_T = \Omega \times (-T, 0)$, if

- (1) $\mathbf{u}, \mathbf{b} \in L^\infty(-T, 0; L^2(\Omega)) \cap L^2(-T, 0; H^1(\Omega))$ and $\pi \in L^{\frac{3}{2}}(\Omega_T)$;
- (2) (1.1) is satisfied in the sense of distribution;
- (3) the local energy inequality holds: for any nonnegative test function $\varphi \in C_c^\infty(\Omega_T)$ and $t \in (-T, 0)$,

$$(1.7) \quad \begin{aligned} & \int_{\Omega} (|\mathbf{u}|^2 + |\mathbf{b}|^2) \varphi \, dx + 2 \int_{-T}^t \int_{\Omega} (|\nabla \mathbf{u}|^2 + |\nabla \mathbf{b}|^2) \varphi \, dx \, ds \\ & \leq \int_{-T}^t \int_{\Omega} (|\mathbf{u}|^2 + |\mathbf{b}|^2) \cdot (\partial_s \varphi + \Delta \varphi) + \mathbf{u} \cdot \nabla \varphi (|\mathbf{u}|^2 + |\mathbf{b}|^2 + 2\pi) \, dx \, ds \\ & - 2 \int_{-T}^t \int_{\Omega} (\mathbf{u} \cdot \mathbf{b}) \mathbf{b} \cdot \nabla \varphi \, dx \, ds. \end{aligned}$$

The global existence of suitable weak solution to the MHD equations (1.1) was investigated by C. He and Z. Xin [14]. They also obtained that one-dimensional Hausdorff measure of the possible space-time singular points set for the suitable weak solution is zero.

In this paper, inspired by [3, 24], we focus on the regularity criteria to the MHD equation (1.1) involving the terms u_3 and b_3 only.

Theorem 1.2. *Let $(\mathbf{u}, \mathbf{b}, \pi)$ be a suitable weak solution of the MHD equations (1.1) in $\mathbb{R}^3 \times (-1, 0)$. If \mathbf{u}, \mathbf{b} satisfies*

$$(1.8) \quad u_3 \in L^{p_0,1}(-1, 0; L^{q_0}(B(2)))$$

and

$$(1.9) \quad b_3 \in L^{p_1,1}(-1, 0; L^{q_1}(B(2)))$$

with $\frac{2}{p_0} + \frac{3}{q_0} = \frac{2}{p_1} + \frac{3}{q_1} = 1$, $3 < q_0, q_1 < +\infty$, then the solution (\mathbf{u}, \mathbf{b}) is regular at $(0, 0)$. Here $B(r)$ is the ball in \mathbb{R}^3 with center at origin and radius r .

Remark 1.3. It should be pointed out that the condition $\mu = \nu$ in the MHD equations (1.1) is essential in our estimates. Actually, we adopt the Elsässer variables $\mathbf{Z}^+ = \mathbf{u} + \mathbf{b}$ and $\mathbf{Z}^- = \mathbf{u} - \mathbf{b}$ to obtain the crucial lemma 4.1 that if $u_3 = b_3 = 0$, the solution is regular at $(0, 0)$. However, it is not trivial in the case $\mu \neq \nu$. We will discuss it in our future work.

Remark 1.4. If we replace (1.8) with the subcritical regularity criteria

$$(1.10) \quad u_3 \in L^{p_0}(-1, 0; L^{q_0}(B(2))), \quad \frac{2}{p_0} + \frac{3}{q_0} < 1, \quad 3 < q_0 \leq +\infty,$$

Theorem 1.2 still holds. Actually, we can pick $2 < p_2 < p_0$ and $3 < q_2 < q_0$ with $\frac{2}{p_2} + \frac{3}{q_2} = 1$. Therefore, we can prove it directly by the embedding inequality

$$\|u_3\|_{L^{p_2,1}(-1,0;L^{q_2}(B(2)))} \leq C \|u_3\|_{L^{p_0}(-1,0;L^{q_0}(B(2)))}.$$

Analogously, we can replace (1.9) with the regularity criteria

$$(1.11) \quad b_3 \in L^{p_1}(-1, 0; L^{q_1}(B(2))), \quad \frac{2}{p_1} + \frac{3}{q_1} < 1, \quad 3 < q_1 \leq +\infty.$$

Remark 1.5. By the standard interpolation theory, it is possible to extend the regularity criteria in Theorem 1.2 to weaker one, such as

$$(1.12) \quad u_3 \in L^{p_0,1}(-1, 0; L^{q_0,\infty}(B(2))), \quad b_3 \in L^{p_1,1}(-1, 0; L^{q_1,\infty}(B(2))),$$

with $\frac{2}{p_0} + \frac{3}{q_0} = \frac{2}{p_1} + \frac{3}{q_1} = 1$, $3 < q_0, q_1 < +\infty$. We leave the details to the interested readers.

Thanks to Theorem 1.2 and Remark 1.4, we obtain the following theorem.

Theorem 1.6. *Assume that $\mathbf{u}_0, \mathbf{b}_0 \in H^1(\mathbb{R}^3)$ with $\operatorname{div} \mathbf{u}_0 = \operatorname{div} \mathbf{b}_0 = 0$ and (\mathbf{u}, \mathbf{b}) is the weak solution to the MHD equations (1.1)–(1.2). Then the solution is regular in $\mathbb{R}^3 \times (0, T]$, provided that*

- (1) $u_3 \in L^{p_0,1}(0, T; L^{q_0}(\mathbb{R}^3))$ with $\frac{2}{p_0} + \frac{3}{q_0} = 1$, $3 < q_0 < \infty$, or
 $u_3 \in L^{p_0}(0, T; L^{q_0}(\mathbb{R}^3))$ with $\frac{2}{p_0} + \frac{3}{q_0} < 1$, $3 < q_0 \leq +\infty$;
- (2) $b_3 \in L^{p_1,1}(0, T; L^{q_1}(\mathbb{R}^3))$ with $\frac{2}{p_1} + \frac{3}{q_1} = 1$, $3 < q_1 < \infty$, or
 $b_3 \in L^{p_1}(0, T; L^{q_1}(\mathbb{R}^3))$ with $\frac{2}{p_1} + \frac{3}{q_1} < 1$, $3 < q_1 \leq +\infty$.

Our paper is organized as follows: we recall some notations and preliminary results in Section 2, and establish a key lemma in Section 3; finally, we will complete the proof of Theorem 1.2 in Section 4.

2. NOTATIONS AND PRELIMINARY

In this preparation section, we recall some usual notations and preliminary results.

For two comparable quantities, the inequality $X \lesssim Y$ stands for $X \leq CY$ for some positive constant C . The dependence of the constant C on other parameters or constants are usually clear from the context, and we will often suppress this dependence.

We use the following standard notations in the literature: Given matrix $\mathbf{A} = (A_{ij}) \in \mathbb{R}^{m \times n}$, we denote the norm $|\mathbf{A}| = \left(\sum_{i=1}^m \sum_{j=1}^n A_{ij}^2 \right)^{\frac{1}{2}}$; for two vectors $\mathbf{a} = (a_1, a_2, a_3)$, $\mathbf{b} = (b_1, b_2, b_3)$, $\mathbf{a} \cdot \mathbf{b}$ and $|\mathbf{a}|$ are the usual scalar product and norm in \mathbb{R}^3 respectively, and $\mathbf{a} \otimes \mathbf{b}$ is a matrix with $(\mathbf{a} \otimes \mathbf{b})_{ij} = a_i b_j$, $(\mathbf{a}, \mathbf{b}) = (a_1, a_2, a_3, b_1, b_2, b_3)$; for $x = (x_1, x_2, x_3) \in \mathbb{R}^3$, $x' = (x_1, x_2)$ is the horizontal variable; $B(x_0, R)$ is the ball in \mathbb{R}^3 with center at x_0 and radius R , and $B(R) := B(0, R)$; $B'(x'_0, R) \subset \mathbb{R}^2$ is the ball in the horizontal plane with center at x'_0 and radius R , and $B'(R) := B'(0, R)$; $Q(z_0, R) = B(x_0, R) \times (t_0 - R^2, t_0)$ with $z_0 = (x_0, t_0)$ and $Q(R) := Q(0, R)$; we denote the integral mean value $(f(t))_{B(r)} = \int_{B(r)} f(y, t) dy$.

We also shall use the same notation as that in Chae-Wolf [3]. Set

$$U_n(R) \stackrel{\text{def}}{=} B'(R) \times (-r_n, r_n), \quad Q_n(R) \stackrel{\text{def}}{=} U_n(R) \times (-r_n^2, 0),$$

and

$$A_n(R) \stackrel{\text{def}}{=} Q_n(R) \setminus Q_{n+1}(R),$$

where $r_n = 2^{-n}$, $n \in \mathbb{N}$. We consider Φ_n given by

$$\Phi_n(x, t) \stackrel{\text{def}}{=} \frac{1}{\sqrt{4\pi(-t + r_n^2)}} e^{-\frac{x_3^2}{4(-t + r_n^2)}}, \quad (x, t) \in \mathbb{R}^3 \times (-\infty, 0),$$

which satisfies the fundamental solution of the backward heat equation

$$\partial_t \Phi_n + \partial_3^2 \Phi_n = 0.$$

There exist absolute constants $c_1, c_2 > 0$ such that for $j = 1, \dots, n-1$, it holds

$$(2.1) \quad \begin{aligned} \Phi_n &\leq c_2 r_j^{-1}, & |\partial_3 \Phi_n| &\leq c_2 r_j^{-2}, & \text{in } A_j(R), \\ c_1 r_n^{-1} &\leq \Phi_n \leq c_2 r_n^{-1}, & c_1 r_n^{-2} &\leq |\partial_3 \Phi_n| \leq c_2 r_n^{-2}, & \text{in } Q_n(R). \end{aligned}$$

We denote the energy

$$\begin{aligned} E_n(R) &\stackrel{\text{def}}{=} \sup_{t \in (-r_n^2, 0)} \int_{U_n(R)} |(\mathbf{u}, \mathbf{b})|^2 dx + \int_{-r_n^2}^0 \int_{U_n(R)} |\nabla(\mathbf{u}, \mathbf{b})|^2 dx dt, \\ \mathcal{E} &\stackrel{\text{def}}{=} \sup_{t \in (-1, 0)} \int_{\mathbb{R}^3} |(\mathbf{u}, \mathbf{b})|^2 dx + \int_{-1}^0 \int_{\mathbb{R}^3} |\nabla(\mathbf{u}, \mathbf{b})|^2 dx dt. \end{aligned}$$

The following lemma ensures the energy estimates.

Lemma 2.1. [3, Lemma 3.1] *Let $R \geq \frac{1}{2}$. For $2 \leq p \leq \infty$, $2 \leq q \leq 6$, $\frac{2}{p} + \frac{3}{q} = \frac{3}{2}$, we have*

$$(2.2) \quad \|\mathbf{u}\|_{L^p(-r_n^2, 0; L^q(U_n(R)))}^2 + \|\mathbf{b}\|_{L^p(-r_n^2, 0; L^q(U_n(R)))}^2 \leq CE_n(R).$$

3. A KEY LEMMA

In this section, we apply a similar argument in Cafferalli-Kohn-Nirenberg [1], Chae-Wolf [3], or Wang-Wu-Zhang [24] to prove the following lemma.

Lemma 3.1. *Let $(\mathbf{u}, \mathbf{b}, \pi)$ be a suitable weak solution of (1.1) in $\mathbb{R}^3 \times (-1, 0)$. If \mathbf{u}, \mathbf{b} satisfies*

$$(3.1) \quad u_3 \in L^{p_0,1}(-1, 0; L^{q_0}(B(2))), \quad b_3 \in L^{p_1,1}(-1, 0; L^{q_1}(B(2))),$$

with $\frac{2}{p_0} + \frac{3}{q_0} = \frac{2}{p_1} + \frac{3}{q_1} = 1$, $3 < q_0, q_1 < +\infty$, then

$$r^{-2} \|\mathbf{u}\|_{L^3(Q(r))}^3 + r^{-2} \|\mathbf{b}\|_{L^3(Q(r))}^3 \leq C,$$

for any $0 < r \leq 1$.

For fixed p_0, p_1 , we can pick $\frac{4q_0}{2q_0-5} < p_0^* < p_0$ and $2 < p_1^* < p_1$. Set

$$(3.2) \quad \mathcal{B}_i = r_i^{1-\frac{2}{p_0^*}-\frac{3}{q_0}} \|u_3\|_{L^{p_0^*}(-r_i^2, 0; L^{q_0}(B(2)))} + r_i^{1-\frac{2}{p_1^*}-\frac{3}{q_1}} \|b_3\|_{L^{p_1^*}(-r_i^2, 0; L^{q_1}(B(2)))}.$$

By Lemma A.5, we have

$$(3.3) \quad \sum_{i=0}^{+\infty} \mathcal{B}_i \leq C \left(\|u_3\|_{L^{p_0,1}(-1, 0; L^{q_0}(B(2)))} + \|b_3\|_{L^{p_1,1}(-1, 0; L^{q_1}(B(2)))} \right).$$

Let $\eta(x_3, t) \in C_c^\infty((-1, 1) \times (-1, 0])$ denotes a cut-off function, $0 \leq \eta \leq 1$, and $\eta = 1$ on $(-\frac{1}{2}, \frac{1}{2}) \times (-\frac{1}{4}, 0)$.

In addition, let $\frac{1}{2} \leq \rho < R \leq 1$ be arbitrarily chosen, but $|R - \rho| \leq \frac{1}{2}$. Let $\psi = \psi(x') \in C^\infty(\mathbb{R}^2)$ with $0 \leq \psi \leq 1$ in $B'(R)$ satisfying

$$(3.4) \quad \psi(x') = \psi(|x'|) = \begin{cases} 1 & \text{in } B'(\rho) \\ 0 & \text{in } \mathbb{R}^2 \setminus B'\left(\frac{R+\rho}{2}\right) \end{cases},$$

and

$$|D\psi| \leq \frac{C}{R-\rho}, \quad |D^2\psi| \leq \frac{C}{(R-\rho)^2}.$$

For $j = 0, 1, \dots, n$, denote $\chi_j = \chi_{B'(R)}(x') \cdot \eta(2^j \cdot x_3, 2^{2j} \cdot t)$, where $\chi_{B'(R)}$ is the indicator function of the set $B'(R)$. Let

$$(3.5) \quad \phi_j = \begin{cases} \chi_j - \chi_{j+1}, & \text{if } j = 0, \dots, n-1, \\ \chi_n, & \text{if } j = n. \end{cases}$$

Taking the test function $\varphi = \Phi_n \eta \psi$ in (1.7), we have

$$(3.6) \quad \begin{aligned} & \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \Phi_n \eta \psi \, dx + 2 \int_{-1}^t \int_{U_0(R)} |\nabla(\mathbf{u}, \mathbf{b})|^2 \Phi_n \eta \psi \, dx \, ds \\ & \leq \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \cdot (\partial_s + \Delta)(\Phi_n \eta \psi) \, dx \, ds \\ & \quad + \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \mathbf{u} \cdot \nabla(\Phi_n \eta \psi) - 2(\mathbf{u} \cdot \mathbf{b}) \mathbf{b} \cdot \nabla(\Phi_n \eta \psi) \, dx \, ds \\ & \quad + 2 \int_{-1}^t \int_{U_0(R)} \pi \mathbf{u} \cdot \nabla(\Phi_n \eta \psi) \, dx \, ds. \end{aligned}$$

Next, we shall handle the right side of (3.6) term by term.

3.1. Estimates for nonlinear terms.

Lemma 3.2. *Under the assumptions of Lemma 3.1, we have*

$$(3.7) \quad \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \cdot (\partial_s + \Delta) (\Phi_n \eta \psi) \, dx \, ds \leq C \frac{\mathcal{E}}{(R - \rho)^2}.$$

Proof. By the estimates (2.1) for Φ_n and the fact that $\text{supp } \nabla \eta \subseteq A_0, \text{supp } \phi_j \subseteq Q_j \setminus Q_{j+2}$, we have

$$\begin{aligned} & \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \cdot (\partial_s + \Delta) (\Phi_n \eta \psi) \, dx \, ds \\ &= \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \cdot (\Phi_n \partial_s \eta \psi + 2\partial_3 \Phi_n \partial_3 \eta \psi + \Phi_n \Delta(\eta \psi)) \, dx \, ds \\ &\lesssim \int_{-1}^0 \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \, dx \, ds + (R - \rho)^{-2} \sum_{i=0}^n \int_{-1}^0 \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \Phi_n \phi_i \, dx \, ds \\ &\leq \mathcal{E} + (R - \rho)^{-2} \sum_{i=0}^n r_i^{-1} \int_{-r_i^2}^0 \int_{U_i(R)} |(\mathbf{u}, \mathbf{b})|^2 \, dx \, ds \\ &\lesssim \mathcal{E} + (R - \rho)^{-2} \sum_{i=0}^n r_i \mathcal{E} \\ &\lesssim (R - \rho)^{-2} \mathcal{E}. \end{aligned}$$

□

□

Lemma 3.3. *Under the assumptions of Lemma 3.1, we have*

$$(3.8) \quad \begin{aligned} & \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \mathbf{u} \cdot \nabla (\Phi_n \eta \psi) - 2(\mathbf{u} \cdot \mathbf{b}) \mathbf{b} \cdot \nabla (\Phi_n \eta \psi) \, dx \, ds \\ &\leq C \sum_{i=0}^n \mathcal{B}_i (r_i^{-1} E_i(R)) + C \mathcal{E}^{\frac{3}{2}} + C(R - \rho)^{-1} \mathcal{E}^{\frac{1}{2}} \sum_{i=0}^n r_i^{\frac{1}{2}} (r_i^{-1} E_i(R)), \end{aligned}$$

where \mathcal{B}_i is defined in (3.2).

Proof. We first note that

$$\begin{aligned} & \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \mathbf{u} \cdot \nabla (\Phi_n \eta \psi) - 2(\mathbf{u} \cdot \mathbf{b}) \mathbf{b} \cdot \nabla (\Phi_n \eta \psi) \, dx \, ds \\ &= \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 u_3 \cdot \partial_3 \Phi_n \cdot (\eta \psi) \, dx \, ds \\ &\quad + \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \mathbf{u} \cdot \Phi_n \nabla (\eta \psi) \, dx \, ds \\ &\quad - 2 \int_{-1}^t \int_{U_0(R)} (\mathbf{u} \cdot \mathbf{b}) b_3 \cdot \partial_3 \Phi_n \cdot (\eta \psi) \, dx \, ds \end{aligned}$$

$$\begin{aligned}
& -2 \int_{-1}^t \int_{U_0(R)} (\mathbf{u} \cdot \mathbf{b}) \mathbf{b} \cdot \Phi_n \nabla (\eta \psi) \, dx \, ds \\
(3.9) \quad & \stackrel{\text{def}}{=} I_1 + I_2 + I_3 + I_4.
\end{aligned}$$

By the estimates (2.1) for Φ_n , Hölder inequality and Lemma 2.1, we have

$$\begin{aligned}
I_1 &= \sum_{i=0}^n \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 u_3 \cdot \partial_3 \Phi_n \cdot (\phi_i \psi) \, dx \, ds \\
&\lesssim \sum_{i=0}^n r_i^{-2} \int_{-r_i^2}^0 \|(\mathbf{u}, \mathbf{b})\|_{L^{\frac{2q_0}{q_0-1}}(U_i(R))}^2 \|u_3\|_{L^{q_0}(U_i(R))} \, ds \\
&\lesssim \sum_{i=0}^n r_i^{-\frac{2}{p_0^*} - \frac{3}{q_0}} \|(\mathbf{u}, \mathbf{b})\|_{L^{\frac{4q_0}{3}}(-r_i^2, 0; L^{\frac{2q_0}{q_0-1}}(U_i(R)))}^2 \|u_3\|_{L^{p_0^*}(-r_i^2, 0; L^{q_0}(U_i(R)))} \\
(3.10) \quad &\lesssim \sum_{i=0}^n \mathcal{B}_i(r_i^{-1} E_i(R)).
\end{aligned}$$

For the second term, by (2.1) and Lemma 2.1 again, we have

$$\begin{aligned}
I_2 &= \int_{-1}^t \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^2 \mathbf{u} \cdot \Phi_n \nabla \eta \psi + |(\mathbf{u}, \mathbf{b})|^2 \mathbf{u} \cdot \Phi_n \eta \nabla \psi \, dx \, ds \\
&\lesssim \int_{-1}^0 \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^3 \, dx \, ds + \frac{1}{(R-\rho)} \sum_{i=0}^n r_i^{-1} \int_{-r_i^2}^0 \int_{U_i(R)} |(\mathbf{u}, \mathbf{b})|^3 \, dx \, ds \\
&\lesssim \mathcal{E}^{\frac{3}{2}} + \frac{1}{(R-\rho)} \sum_{i=0}^n r_i^{-1} r_i^{\frac{1}{2}} \|(\mathbf{u}, \mathbf{b})\|_{L^4(-r_i^2, 0; L^3(U_i(R)))}^3 \\
(3.11) \quad &\lesssim \mathcal{E}^{\frac{3}{2}} + \frac{\mathcal{E}^{\frac{1}{2}}}{(R-\rho)} \sum_{i=0}^n r_i^{\frac{1}{2}} (r_i^{-1} E_i(R)).
\end{aligned}$$

As to I_3, I_4 , we have

$$\begin{aligned}
I_3 &\lesssim \sum_{i=0}^n r_i^{-2} \int_{-r_i^2}^0 \|(\mathbf{u}, \mathbf{b})\|_{L^{\frac{2q_1}{q_1-1}}(U_i(R))}^2 \|b_3\|_{L^{q_1}(U_i(R))} \, ds \\
&\lesssim \sum_{i=0}^n r_i^{-\frac{2}{p_1^*} - \frac{3}{q_1}} \|(\mathbf{u}, \mathbf{b})\|_{L^{\frac{4q_1}{3}}(-r_i^2, 0; L^{\frac{2q_1}{q_1-1}}(U_i(R)))}^2 \|b_3\|_{L^{p_1^*}(-r_i^2, 0; L^{q_1}(U_i(R)))} \\
(3.12) \quad &\lesssim \sum_{i=0}^n \mathcal{B}_i(r_i^{-1} E_i(R)),
\end{aligned}$$

which is analogous to (3.10), and

$$I_4 \lesssim \int_{-1}^0 \int_{U_0(R)} |(\mathbf{u}, \mathbf{b})|^3 \, dx \, ds + \frac{1}{(R-\rho)} \sum_{i=0}^n r_i^{-1} \int_{-r_i^2}^0 \int_{U_i(R)} |(\mathbf{u}, \mathbf{b})|^3 \, dx \, ds$$

$$(3.13) \quad \lesssim \mathcal{E}^{\frac{3}{2}} + \frac{\mathcal{E}^{\frac{1}{2}}}{(R-\rho)} \sum_{i=0}^n r_i^{\frac{1}{2}} (r_i^{-1} E_i(R)),$$

which is analogous to (3.11). Combining (3.9) with (3.10), (3.11), (3.12) and (3.13), we obtain (3.8). \square

3.2. Estimates for the pressure. This part is devoted to the estimates regarding the pressure term on the right side of (3.6). We will give a similar argument in [24] here.

Given 3×3 matrix valued function $f = (f_{jk})$, we set

$$\widehat{\mathcal{J}(f)}(\xi) = \sum_{j,k=1,2,3} \frac{\xi_j \xi_k}{|\xi|^2} \mathcal{F} f_{jk},$$

where the Fourier transform is defined by

$$\mathcal{F}g(\xi) = \hat{g}(\xi) = \int_{\mathbb{R}^3} e^{-i(x \cdot \xi)} g(x) dx.$$

Therefore, $\mathcal{J} : L^q(\mathbb{R}^3) \rightarrow L^q(\mathbb{R}^3)$, $1 < q < +\infty$ defines a bounded linear operator with

$$(3.14) \quad \|\mathcal{J}(f)\|_{L^q(\mathbb{R}^3)} \leq C \|f\|_{L^q(\mathbb{R}^3)}.$$

Denote $f = (f_{jk}) = (\mathbf{u} \otimes \mathbf{u} - \mathbf{b} \otimes \mathbf{b}) \cdot \chi_0$,

$$(3.15) \quad \pi_0 = \mathcal{J}(f), \quad \pi_h = \pi - \pi_0.$$

It follows that

$$\Delta \pi_0 = \nabla \cdot \nabla \cdot f = \sum_{j,k=1}^3 \partial_j \partial_k f_{jk} \quad \text{in } \mathbb{R}^3 \times (-1, 0)$$

in the sense of distributions and π_h is harmonic in $Q_1(R)$. Then we have

$$(3.16) \quad \begin{aligned} & \int_{-1}^t \int_{U_0(R)} \pi \mathbf{u} \cdot \nabla (\Phi_n \eta \psi) dx ds \\ &= \int_{-1}^t \int_{U_0(R)} \pi_0 \mathbf{u} \cdot \nabla (\Phi_n \eta \psi) dx ds + \int_{-1}^t \int_{U_0(R)} \pi_h \mathbf{u} \cdot \nabla (\Phi_n \eta \psi) dx ds \\ &= \int_{-1}^t \int_{U_0(R)} \pi_0 u_3 \cdot \partial_3 \Phi_n \eta \psi dx ds + \int_{-1}^t \int_{U_0(R)} \pi_0 \mathbf{u} \cdot \Phi_n \nabla (\eta \psi) dx ds \\ & \quad - \int_{-1}^t \int_{U_0(R)} \nabla \pi_h \cdot \mathbf{u} \cdot (\Phi_n \eta \psi) dx ds \\ & \stackrel{\text{def}}{=} J + K + H. \end{aligned}$$

Lemma 3.4. *Under the assumptions of Lemma 3.1, we have*

$$(3.17) \quad \begin{aligned} & \int_{-1}^t \int_{U_0(R)} \pi \mathbf{u} \cdot \nabla (\Phi_n \eta \psi) dx ds \\ & \leq C \sum_{i=0}^n \mathcal{B}_i (r_i^{-1} E_i(R)) + C \frac{\mathcal{E}^{\frac{1}{2}}}{R-\rho} \sum_{i=0}^n r_i^{\frac{1}{2}} (r_i^{-1} E_i(R)) + \frac{C}{(R-\rho)^3} \mathcal{E}^{\frac{3}{2}}. \end{aligned}$$

Proof. Combining the estimates (3.16), (A.2), (A.7) and (A.12), we have (3.17). \square \square

3.3. The proof of Lemma 3.1. On the basis of the estimates of the nonlinear term and the pressure in subsection 3.1–3.2, we are in position to give the detail proof of Lemma 3.1.

Proof. Gathering (3.6) and the estimates in Lemma 3.2, 3.3 and 3.4, we have

$$\begin{aligned}
r_n^{-1}E_n(\rho) &\leq C \sum_{i=0}^n \mathcal{B}_i (r_i^{-1}E_i(R)) + C \frac{\mathcal{E}^{\frac{1}{2}}}{R-\rho} \sum_{i=0}^n r_i^{\frac{1}{2}} (r_i^{-1}E_i(R)) + C \frac{1 + \mathcal{E}^{\frac{3}{2}}}{(R-\rho)^3} \\
&\leq C \sum_{i=0}^n \mathcal{B}_i (r_i^{-1}E_i(R)) + C \frac{\mathcal{E}^{\frac{3}{4}}}{R-\rho} \sum_{i=0}^n r_i^{-\frac{5}{8}} E_i(R)^{\frac{3}{4}} r_i^{\frac{1}{8}} + C \frac{1 + \mathcal{E}^{\frac{3}{2}}}{(R-\rho)^3} \\
&\leq C \sum_{i=0}^n \mathcal{B}_i (r_i^{-1}E_i(R)) + C \frac{\mathcal{E}^{\frac{3}{4}}}{R-\rho} \left(\sum_{i=0}^n r_i^{-\frac{5}{6}} E_i(R) \right)^{\frac{3}{4}} \left(\sum_{i=0}^n r_i^{\frac{1}{2}} \right)^{\frac{1}{4}} \\
&\quad + C \frac{1 + \mathcal{E}^{\frac{3}{2}}}{(R-\rho)^3} \\
&\leq C_0 \sum_{i=0}^n \left(\mathcal{B}_i + r_i^{\frac{1}{6}} \right) (r_i^{-1}E_i(R)) + C_0 \frac{1 + \mathcal{E}^3}{(R-\rho)^4}.
\end{aligned}$$

In view of (3.3), there exists a sufficient large integer $n_0 \geq 1$ such that

$$(3.18) \quad C_0 \sum_{i=n_0}^{\infty} \left(\mathcal{B}_i + r_i^{\frac{1}{6}} \right) \leq \frac{1}{2}.$$

Then for $n \geq n_0$ we have

$$\begin{aligned}
r_n^{-1}E_n(\rho) &\leq C_0 \sum_{i=n_0}^n \left(\mathcal{B}_i + r_i^{\frac{1}{6}} \right) (r_i^{-1}E_i(R)) \\
&\quad + C_0 \sum_{i=0}^{n_0-1} \left(\mathcal{B}_i + r_i^{\frac{1}{6}} \right) (r_i^{-1}E_i(R)) + C_0 \frac{1 + \mathcal{E}^3}{(R-\rho)^4} \\
(3.19) \quad &\leq C_0 \sum_{i=n_0}^n \left(\mathcal{B}_i + r_i^{\frac{1}{6}} \right) (r_i^{-1}E_i(R)) + \frac{A_0}{(R-\rho)^4},
\end{aligned}$$

where the constant

$$A_0 = C_1 \cdot 2^{n_0} \left(1 + \|u_3\|_{L^{p_0,1}(-1,0;L^{q_0}(B(2)))}^2 + \|b_3\|_{L^{p_1,1}(-1,0;L^{q_1}(B(2)))}^2 + \mathcal{E}^3 \right).$$

As the iteration argument in [10, V. Lemma 3.1], we introduce the sequence $\{\rho_k\}_{k=0}^{+\infty}$ which satisfy

$$\rho_0 = \frac{1}{2}, \quad \rho_{k+1} - \rho_k = \frac{1-\theta}{2} \theta^k,$$

with $\frac{1}{2} < \theta^4 < 1$ and $\lim_{k \rightarrow \infty} \rho_k = 1$. For $n_0 \leq j \leq n$ and $k \geq 1$, we have

$$r_j^{-1}E_j(\rho_k) \leq C_0 \sum_{i=n_0}^j \left(\mathcal{B}_i + r_i^{\frac{1}{6}} \right) (r_i^{-1}E_i(\rho_{k+1})) + A_0 \cdot \frac{16}{(1-\theta)^4} \theta^{-4k}$$

$$(3.20) \quad \leq C_0 \sum_{i=n_0}^n \left(\mathcal{B}_i + r_i^{\frac{1}{6}} \right) \left(r_i^{-1} E_i(\rho_{k+1}) \right) + A_0 \cdot \frac{16}{(1-\theta)^4} \theta^{-4k}.$$

Applying (3.18) and (3.20) to the iteration argument from (3.19), we obtain that for $n \geq n_0$,

$$\begin{aligned} r_n^{-1} E_n(\rho_0) &\leq C_0 \sum_{j=n_0}^n \left(\mathcal{B}_j + r_j^{\frac{1}{6}} \right) \left(r_j^{-1} E_j(\rho_1) \right) + A_0 \cdot \frac{16}{(1-\theta)^4} \\ &\leq \frac{1}{2} C_0 \sum_{j=n_0}^n \left(\mathcal{B}_j + r_j^{\frac{1}{6}} \right) \left(r_j^{-1} E_j(\rho_2) \right) + A_0 \cdot \frac{16}{(1-\theta)^4} \left(1 + \frac{1}{2\theta^4} \right) \\ &\leq \frac{1}{2^{k-1}} C_0 \sum_{j=n_0}^n \left(\mathcal{B}_j + r_j^{\frac{1}{6}} \right) \left(r_j^{-1} E_j(\rho_k) \right) + A_0 \cdot \frac{16}{(1-\theta)^4} \sum_{j=0}^{k-1} \left(\frac{1}{2\theta^4} \right)^j \\ &\leq \frac{1}{2^{k-1}} C_0 \sum_{j=n_0}^n \left(\mathcal{B}_j + r_j^{\frac{1}{6}} \right) \left(r_j^{-1} E_j(1) \right) + A_0 \cdot \frac{16}{(1-\theta)^4} \cdot \frac{2\theta^4}{2\theta^4 - 1}. \end{aligned}$$

Let $k \rightarrow \infty$, we obtain that for $n \geq n_0$,

$$(3.21) \quad r_n^{-1} E_n(\rho_0) \leq A_0 \cdot \frac{16}{(1-\theta)^4} \cdot \frac{2\theta^4}{2\theta^4 - 1}.$$

For $0 < r \leq r_{n_0}$, there exists an integer $j \geq n_0$, such that

$$r_{j+1} < r \leq r_j,$$

which together with (3.21) ensures that

$$r^{-1} \sup_{t \in (-r^2, 0)} \int_{B(r)} |(\mathbf{u}, \mathbf{b})|^2 dx + r^{-1} \int_{Q(r)} |\nabla(\mathbf{u}, \mathbf{b})|^2 dx dt \leq 2 r_j^{-1} E_j(\rho_0) \leq C.$$

For $r_{n_0} < r \leq 1$,

$$r^{-1} \sup_{t \in (-r^2, 0)} \int_{B(r)} |(\mathbf{u}, \mathbf{b})|^2 dx + r^{-1} \int_{Q(r)} |\nabla(\mathbf{u}, \mathbf{b})|^2 dx dt \leq 2^{n_0} \mathcal{E} \leq C.$$

Hence for $0 < r \leq 1$,

$$(3.22) \quad r^{-1} \sup_{t \in (-r^2, 0)} \int_{B(r)} |(\mathbf{u}, \mathbf{b})|^2 dx + r^{-1} \int_{Q(r)} |\nabla(\mathbf{u}, \mathbf{b})|^2 dx dt \leq C.$$

By Sobolev embedding theorem, we have that for $0 < r \leq 1$,

$$(3.23) \quad r^{-2} \|(\mathbf{u}, \mathbf{b})\|_{L^3(Q(r))}^3 \leq C r^{-\frac{3}{2}} \|(\mathbf{u}, \mathbf{b})\|_{L^4(-r^2, 0; L^3(B(r)))}^3 \leq C.$$

The proof is completed. \square \square

4. PROOF OF THEOREM 1.2

Before starting the proof, we give an essential lemma for the 3D MHD equation (1.1).

Lemma 4.1. *Let $\mu = \nu = 1$ in the MHD equations (1.1). Assume that $(\mathbf{u}, \mathbf{b}, \pi)$ is a suitable weak solution in $Q(1)$ with $u_3 = b_3 = 0$. Then the solution is regular in $\overline{Q(\frac{1}{2})}$.*

Proof. Denote \mathcal{S}_{MHD} the singular set of the suitable weak solution, the cylinder $U(x, r) = B'(x', r) \times (x_3 - r, x_3 + r)$ and $U(r) := U(0, r)$. By [14] or [23, Theorem 1.4], we know that the one-dimensional Hausdorff measure of \mathcal{S}_{MHD} is zero.

Assume that the assertion does not hold, i.e. $\mathcal{S}_{\text{MHD}} \cap \overline{Q(\frac{1}{2})} \neq \emptyset$. By a similar argument in [15, Lemma 3.2], there exists $(x_0, t_0) \in \overline{Q(\frac{1}{2})}$ and some ε, ρ with $0 < \varepsilon < \rho < \frac{1}{2}$, such that

$$(x_0, t_0) \in \mathcal{S}_{\text{MHD}} \cap \overline{U(x_0, \rho) \times (t_0 - \rho^2, t_0)} \subseteq U(x_0, \rho - \varepsilon) \times \{t_0\}.$$

Without loss of generality, we may assume $(x_0, t_0) = (0, 0)$. Thus, the solution $(\mathbf{u}, \mathbf{b}, \nabla \pi) \in C^\infty(\overline{U(\rho)} \times [-\rho^2, 0])$ and is singular at $(0, 0)$. Moreover, there exists a constant C , such that

$$(4.1) \quad \sup_{-1 < t < 0} \|(\mathbf{u}, \mathbf{b})\|_{L^2(B(1))}^2 + \|\nabla(\mathbf{u}, \mathbf{b})\|_{L^2(Q(1))}^2 + \|\pi\|_{L^{\frac{3}{2}}(Q(1))} \leq C,$$

and for $(x, t) \in U(\rho) \setminus U(\rho - \varepsilon) \times [-\rho^2, 0]$ or $(x, t) \in U(\rho) \times \{-\rho^2\}$,

$$(4.2) \quad |\mathbf{u}| + |\mathbf{b}| \leq C.$$

We introduce the Elsässer variables $\mathbf{Z}^+ = (Z_1^+, Z_2^+, 0) = \mathbf{u} + \mathbf{b}$ and $\mathbf{Z}^- = (Z_1^-, Z_2^-, 0) = \mathbf{u} - \mathbf{b}$, which satisfy

$$\begin{aligned} \partial_t \mathbf{Z}^+ + \mathbf{Z}^- \cdot \nabla \mathbf{Z}^+ - \Delta \mathbf{Z}^+ + \nabla \pi &= 0, \\ \partial_t \mathbf{Z}^- + \mathbf{Z}^+ \cdot \nabla \mathbf{Z}^- - \Delta \mathbf{Z}^- + \nabla \pi &= 0. \end{aligned}$$

As in [2], we introduce $\overline{\mathbf{Z}^+} = \int_{-\rho}^{\rho} \mathbf{Z}^+ \cdot \beta(x_3) dx_3$, $\overline{\mathbf{Z}^-} = \int_{-\rho}^{\rho} \mathbf{Z}^- \cdot \beta(x_3) dx_3$, where the non-negative cut-off function $\beta(x_3) \in C_c^\infty((-\rho, \rho))$, $\beta = \frac{1}{2\rho - \varepsilon}$ on $(-\rho + \varepsilon, \rho - \varepsilon)$ and $\int_{-\rho}^{\rho} \beta(x_3) dx_3 = 1$. It is easy to find that $\partial_3 \pi = 0$, and

$$\begin{aligned} \partial_t \overline{\mathbf{Z}^+} + \int_{-\rho}^{\rho} \mathbf{Z}^- \cdot \nabla \mathbf{Z}^+ \cdot \beta dx_3 - \Delta \overline{\mathbf{Z}^+} + \nabla \pi &= \int_{-\rho}^{\rho} \mathbf{Z}^+ \cdot \partial_3^2 \beta dx_3, \\ \partial_t \overline{\mathbf{Z}^-} + \int_{-\rho}^{\rho} \mathbf{Z}^+ \cdot \nabla \mathbf{Z}^- \cdot \beta dx_3 - \Delta \overline{\mathbf{Z}^-} + \nabla \pi &= \int_{-\rho}^{\rho} \mathbf{Z}^- \cdot \partial_3^2 \beta dx_3. \end{aligned}$$

Define $\widetilde{\mathbf{Z}^+} = \mathbf{Z}^+ - \overline{\mathbf{Z}^+}$, $\widetilde{\mathbf{Z}^-} = \mathbf{Z}^- - \overline{\mathbf{Z}^-}$, which satisfy

$$(4.3) \quad \partial_t \widetilde{\mathbf{Z}^+} + \mathbf{Z}^- \cdot \nabla \widetilde{\mathbf{Z}^+} + \widetilde{\mathbf{Z}^-} \cdot \nabla \overline{\mathbf{Z}^+} - \int_{-\rho}^{\rho} \widetilde{\mathbf{Z}^-} \cdot \nabla \widetilde{\mathbf{Z}^+} \cdot \beta dx_3 - \Delta \widetilde{\mathbf{Z}^+} = - \int_{-\rho}^{\rho} \mathbf{Z}^+ \cdot \partial_3^2 \beta dx_3$$

$$(4.4) \quad \partial_t \widetilde{\mathbf{Z}^-} + \mathbf{Z}^+ \cdot \nabla \widetilde{\mathbf{Z}^-} + \widetilde{\mathbf{Z}^+} \cdot \nabla \overline{\mathbf{Z}^-} - \int_{-\rho}^{\rho} \widetilde{\mathbf{Z}^+} \cdot \nabla \widetilde{\mathbf{Z}^-} \cdot \beta dx_3 - \Delta \widetilde{\mathbf{Z}^-} = - \int_{-\rho}^{\rho} \mathbf{Z}^- \cdot \partial_3^2 \beta dx_3.$$

Let $\alpha(x_1, x_2) \in C_c^\infty(B'(\rho))$ be a cut-off function with $0 \leq \alpha \leq 1$, $\alpha = 1$ on $B'(\rho - \varepsilon)$. Multiplying (4.3) by $4|\widetilde{\mathbf{Z}^+}|^2 \widetilde{\mathbf{Z}^+} \cdot \varphi^4$ with $\varphi(x) = \alpha(x_1, x_2) \cdot \beta(x_3)$, and integrating the resulting equations over \mathbb{R}^3 , applying (4.1), (4.2), Hölder's inequality and Sobolev embedding theorem, we have

$$\begin{aligned} & \frac{d}{dt} \|\widetilde{\mathbf{Z}^+} \varphi\|_{L^4(U(\rho))}^4 + 4 \int_{\mathbb{R}^3} |\nabla \widetilde{\mathbf{Z}^+}|^2 |\widetilde{\mathbf{Z}^+}|^2 \varphi^4 dx + 2 \int_{\mathbb{R}^3} |\nabla (|\widetilde{\mathbf{Z}^+}|^2 \varphi^2)|^2 dx \\ &= \int_{\mathbb{R}^3} \mathbf{Z}^- \cdot \nabla \alpha^4 \cdot \beta^4 |\widetilde{\mathbf{Z}^+}|^4 - 4 \widetilde{\mathbf{Z}^-} \cdot \nabla \overline{\mathbf{Z}^+} \cdot |\widetilde{\mathbf{Z}^+}|^2 \widetilde{\mathbf{Z}^+} \varphi^4 dx \end{aligned}$$

$$\begin{aligned}
& + 4 \int_{\mathbb{R}^3} \left(\int_{-\rho}^{\rho} \widetilde{\mathbf{Z}}^- \cdot \nabla \widetilde{\mathbf{Z}}^+ \cdot \beta - \mathbf{Z}^+ \cdot \partial_3^2 \beta \, dx_3 \right) \cdot |\widetilde{\mathbf{Z}}^+|^2 \widetilde{\mathbf{Z}}^+ \varphi^4 \, dx + 2 \int_{\mathbb{R}^3} |\widetilde{\mathbf{Z}}^+|^4 |\nabla \varphi^2|^2 \, dx \\
& \lesssim 1 + \left(\|\nabla \widetilde{\mathbf{Z}}^+\|_{L^2(B'(\rho))} + \|\nabla \widetilde{\mathbf{Z}}^+\|_{L^2(U(\rho))} \right) \cdot \|\widetilde{\mathbf{Z}}^-\varphi\|_{L^4(U(\rho))} \|\widetilde{\mathbf{Z}}^+\varphi\|_{L^4(U(\rho))} \\
& \quad \times \|\nabla \left(|\widetilde{\mathbf{Z}}^+|^2 \varphi^2 \right)\|_{L^2(U(\rho))} + \|\widetilde{\mathbf{Z}}^+\varphi\|_{L^4(U(\rho))}^3 \\
& \leq C + C \left(1 + \|\nabla \mathbf{Z}^+\|_{L^2(U(\rho))}^2 \right) \cdot \left(\|\widetilde{\mathbf{Z}}^+\varphi\|_{L^4(U(\rho))}^4 + \|\widetilde{\mathbf{Z}}^-\varphi\|_{L^4(U(\rho))}^4 \right) \\
(4.5) \quad & - \|\nabla \left(|\widetilde{\mathbf{Z}}^+|^2 \varphi^2 \right)\|_{L^2(U(\rho))}^2.
\end{aligned}$$

Analogously, multiplying (4.4) by $4|\widetilde{\mathbf{Z}}^-|^2 \widetilde{\mathbf{Z}}^- \cdot \varphi^4$, and integrating the resulting equations over \mathbb{R}^3 , we have

$$\begin{aligned}
& \frac{d}{dt} \|\widetilde{\mathbf{Z}}^-\varphi\|_{L^4(U(\rho))}^4 + 4 \int_{\mathbb{R}^3} |\nabla \widetilde{\mathbf{Z}}^-|^2 |\widetilde{\mathbf{Z}}^-|^2 \varphi^4 \, dx + 2 \int_{\mathbb{R}^3} |\nabla \left(|\widetilde{\mathbf{Z}}^-|^2 \varphi^2 \right)|^2 \, dx \\
& = \int_{\mathbb{R}^3} \mathbf{Z}^+ \cdot \nabla \alpha^4 \cdot \beta^4 |\widetilde{\mathbf{Z}}^-|^4 - 4 \widetilde{\mathbf{Z}}^+ \cdot \nabla \widetilde{\mathbf{Z}}^- \cdot |\widetilde{\mathbf{Z}}^-|^2 \widetilde{\mathbf{Z}}^- \varphi^4 \, dx \\
& \quad + 4 \int_{\mathbb{R}^3} \left(\int_{-\rho}^{\rho} \widetilde{\mathbf{Z}}^+ \cdot \nabla \widetilde{\mathbf{Z}}^- \cdot \beta - \mathbf{Z}^- \cdot \partial_3^2 \beta \, dx_3 \right) \cdot |\widetilde{\mathbf{Z}}^-|^2 \widetilde{\mathbf{Z}}^- \varphi^4 \, dx + 2 \int_{\mathbb{R}^3} |\widetilde{\mathbf{Z}}^-|^4 |\nabla \varphi^2|^2 \, dx \\
& \leq C + C \left(1 + \|\nabla \mathbf{Z}^-\|_{L^2(U(\rho))}^2 \right) \cdot \left(\|\widetilde{\mathbf{Z}}^+\varphi\|_{L^4(U(\rho))}^4 + \|\widetilde{\mathbf{Z}}^-\varphi\|_{L^4(U(\rho))}^4 \right) \\
(4.6) \quad & - \|\nabla \left(|\widetilde{\mathbf{Z}}^-|^2 \varphi^2 \right)\|_{L^2(U(\rho))}^2.
\end{aligned}$$

Summing up (4.5) and (4.6) and applying Gronwall's inequality, we have that for all $-\rho^2 < t < 0$,

$$(4.7) \quad \|\widetilde{\mathbf{Z}}^+\|_{L^4(U(\rho-\varepsilon))}^4 + \|\widetilde{\mathbf{Z}}^-\|_{L^4(U(\rho-\varepsilon))}^4 \leq C \|\widetilde{\mathbf{Z}}^+\varphi\|_{L^4(U(\rho))}^4 + C \|\widetilde{\mathbf{Z}}^-\varphi\|_{L^4(U(\rho))}^4 \leq C.$$

Accordingly, by (4.2) and (4.7), we have that for $0 < r < \rho - \varepsilon$,

$$\begin{aligned}
& r^{-2} \left(\|\mathbf{u}\|_{L^3(Q(r))}^3 + \|\mathbf{b}\|_{L^3(Q(r))}^3 \right) \\
& \leq C r^{-2} \left(\|\mathbf{Z}^+\|_{L^3(Q(r))}^3 + \|\mathbf{Z}^-\|_{L^3(Q(r))}^3 \right) \\
& \leq C r^{-2} \left(\|\widetilde{\mathbf{Z}}^+\|_{L^3(Q(r))}^3 + \|\widetilde{\mathbf{Z}}^-\|_{L^3(Q(r))}^3 \right) + C r^{-2} \left(\|\overline{\mathbf{Z}}^+\|_{L^3(Q(r))}^3 + \|\overline{\mathbf{Z}}^-\|_{L^3(Q(r))}^3 \right) \\
& \leq C r^{\frac{3}{4}} + C r^{-1} \int_{-r^2}^0 \int_{B'(r)} \left(|\overline{\mathbf{Z}}^+|^3 + |\overline{\mathbf{Z}}^-|^3 \right) \, dx_1 \, dx_2 \, dt \\
& \leq C r^{\frac{3}{4}} + C \sup_{-1 < t < 0} \|(\mathbf{u}, \mathbf{b})\|_{L^2(B(1))}^2 \cdot \left(\int_{-r^2}^0 \int_{-\rho}^{\rho} \int_{B'(r)} |\nabla(\mathbf{u}, \mathbf{b})|^2 \, dx \, dt \right)^{\frac{1}{2}}.
\end{aligned}$$

We can pick a sufficient small $0 < R_0 < 1$, such that for all $0 < r < R_0$,

$$r^{-2} \left(\|\mathbf{u}\|_{L^3(Q(r))}^3 + \|\mathbf{b}\|_{L^3(Q(r))}^3 \right) < \varepsilon_1.$$

By Lemma A.9, we have that the solution is regular at $(0, 0)$, which leads to a contradiction. \square \square

Now, we are in position to prove Theorem 1.2 by standard rigid methods. For $r_k = 2^{-k}$, $k \geq 1$ and $(x, t) \in Q(1)$, we denote the following scaling

$$\begin{aligned}\mathbf{u}_k(x, t) &= r_k \mathbf{u}(r_k x, r_k^2 t), \quad \mathbf{b}_k(x, t) = r_k \mathbf{b}(r_k x, r_k^2 t) \\ \pi_k(x, t) &= r_k^2 \pi(r_k x, r_k^2 t).\end{aligned}$$

By (3.22), we obtain that for $k \geq 1$,

$$(4.8) \quad \sup_{t \in (-1, 0)} \|(\mathbf{u}_k, \mathbf{b}_k)\|_{L^2(B(1))}^2 + \|\nabla(\mathbf{u}_k, \mathbf{b}_k)\|_{L^2(Q(1))}^2 \leq C.$$

Applying Lemma 3.1 and Lemma A.6, we obtain that for $0 < r \leq 1$,

$$r^{-2} \|\pi\|_{L^{\frac{3}{2}}(Q(r))}^{\frac{3}{2}} \leq C,$$

which implies that

$$(4.9) \quad \|\pi_k\|_{L^{\frac{3}{2}}(Q(1))} \leq C.$$

Moreover, $(\mathbf{u}_k, \mathbf{b}_k, \pi_k)$ is a suitable weak solution to the MHD equations (1.1) in $Q(1)$, which converges weakly (by taking subsequences if needed) to some $(\mathbf{v}, \mathbf{B}, \Pi)$,

$$(4.10) \quad \sup_{t \in (-1, 0)} \|(\mathbf{v}, \mathbf{B})\|_{L^2(B(1))}^2 + \|\nabla(\mathbf{v}, \mathbf{B})\|_{L^2(Q(1))}^2 + \|\Pi\|_{L^{\frac{3}{2}}(Q(1))}^{\frac{3}{2}} \leq C.$$

By a similar argument as [17, Theorem 2.2], we can prove that $(\mathbf{v}, \mathbf{B}, \Pi)$ is a suitable weak solution to the MHD equations in $Q(1)$ and $(\mathbf{u}_k, \mathbf{b}_k) \xrightarrow[k \rightarrow \infty]{} (\mathbf{v}, \mathbf{B})$ strongly in $L^3(Q(1))$. By (1.8)–(1.9), we have

$$(4.11) \quad v_3 = B_3 = 0.$$

By Lemma 4.1, the solution (\mathbf{v}, \mathbf{B}) is regular at $(0, 0)$. If we assume that the solution (\mathbf{u}, \mathbf{b}) is singular at $(0, 0)$, $(\mathbf{u}_k, \mathbf{b}_k)$ is also singular at $(0, 0)$. It leads to a contradiction by Lemma A.8.

The proof of Theorem 1.2 is completed. \square

APPENDIX A.

Lemma A.1. [3, Lemma A.2] *Let $0 < r \leq R < +\infty$ and $h : B'(2R) \times (-r, r) \rightarrow \mathbb{R}$ be harmonic. Then for all $0 < \rho \leq \frac{r}{4}$ and $1 \leq \ell \leq q < +\infty$, we get*

$$(A.1) \quad \|h\|_{L^q(B'(R) \times (-\rho, \rho))}^q \leq C \rho r^{2 - \frac{3q}{\ell}} \|h\|_{L^\ell(B'(2R) \times (-r, r))}^q,$$

where C stands for a positive constant depending only on q and ℓ .

We will present the estimates of J , K and H , which is defined in (3.16), in the following several lemmas.

Lemma A.2. *Under the assumptions of Lemma 3.1, we have*

$$(A.2) \quad J \leq C \sum_{i=0}^n \mathcal{B}_i(r_i^{-1} E_i(R)) + C \mathcal{E}^{\frac{3}{2}}.$$

Proof. Let $\pi_{0,j} = \mathcal{J}((\mathbf{u} \otimes \mathbf{u} - \mathbf{b} \otimes \mathbf{b}) \cdot \phi_j)$.

$$\begin{aligned}
J &= \sum_{k=0}^n \int_{-1}^t \int_{U_0(R)} \pi_0 \cdot u_3 \cdot \partial_3 \Phi_n \phi_k \psi \, dx \, ds \\
&= \sum_{k=4}^n \sum_{j=0}^n \int_{-1}^t \int_{U_0(R)} \pi_{0,j} \cdot u_3 \cdot \partial_3 \Phi_n \phi_k \psi \, dx \, ds \\
&\quad + \sum_{k=0}^3 \int_{-1}^t \int_{U_0(R)} \pi_0 \cdot u_3 \cdot \partial_3 \Phi_n \phi_k \psi \, dx \, ds \\
&= \sum_{k=4}^n \sum_{j=0}^{k-4} \int_{-1}^t \int_{U_0(R)} \pi_{0,j} \cdot u_3 \cdot \partial_3 \Phi_n \phi_k \psi \, dx \, ds \\
&\quad + \sum_{k=4}^n \sum_{j=k-3}^n \int_{-1}^t \int_{U_0(R)} \pi_{0,j} \cdot u_3 \cdot \partial_3 \Phi_n \phi_k \psi \, dx \, ds \\
&\quad + \sum_{k=0}^3 \int_{-1}^t \int_{U_0(R)} \pi_0 \cdot u_3 \cdot \partial_3 \Phi_n \phi_k \psi \, dx \, ds \\
(A.3) \quad &\stackrel{\text{def}}{=} J_1 + J_2 + J_3.
\end{aligned}$$

By (2.1), (3.14), Lemma 2.1 and Lemma A.1 with $\rho = r_k, r = r_{j+2}$, we have

$$\begin{aligned}
J_1 &= \sum_{j=0}^{n-4} \sum_{k=j+4}^n \int_{-1}^t \int_{U_0(R)} \pi_{0,j} \cdot u_3 \cdot \partial_3 \Phi_n \phi_k \psi \, dx \, ds \\
&\lesssim \sum_{j=0}^{n-4} \sum_{k=j+4}^n r_k^{-2} \int_{-r_k^2}^0 \|\pi_{0,j}\|_{L^{\frac{q_0}{q_0-1}}(U_k(R))} \|u_3\|_{L^{q_0}(U_k(R))} \, ds \\
&\lesssim \sum_{j=0}^{n-4} \sum_{k=j+4}^n r_k^{-1-\frac{1}{q_0}} r_j^{-1-\frac{1}{2q_0}} \int_{-r_k^2}^0 \|(\mathbf{u}, \mathbf{b})\|_{L^{\frac{4q_0}{2q_0-1}}(U_j(R))}^2 \|u_3\|_{L^{q_0}(U_j(R))} \, ds \\
&\lesssim \sum_{j=0}^{n-4} \sum_{k=j+4}^n r_k^{1-\frac{2}{p_0^*}-\frac{5}{2q_0}} r_j^{-1-\frac{1}{2q_0}} \|(\mathbf{u}, \mathbf{b})\|_{L^{\frac{8q_0}{3}}(-r_j^2, 0; L^{\frac{4q_0}{2q_0-1}}(U_j(R)))}^2 \\
&\quad \times \|u_3\|_{L^{p_0^*}(-r_j^2, 0; L^{q_0}(U_j(R)))} \\
(A.4) \quad &\lesssim \sum_{i=0}^n \mathcal{B}_i(r_i^{-1} E_i(R)).
\end{aligned}$$

By (2.1), (3.14) and Lemma 2.1, we have

$$J_2 = \sum_{k=4}^n \int_{-1}^t \int_{U_0(R)} \mathcal{J}((\mathbf{u} \otimes \mathbf{u} - \mathbf{b} \otimes \mathbf{b}) \cdot \chi_{k-3}) \cdot u_3 \cdot \partial_3 \Phi_n \phi_k \psi \, dx \, ds$$

$$\begin{aligned}
& \lesssim \sum_{k=4}^n r_k^{-2} \int_{-r_k^2}^0 \|\mathcal{L}((\mathbf{u} \otimes \mathbf{u} - \mathbf{b} \otimes \mathbf{b}) \cdot \chi_{k-3})\|_{L^{\frac{q_0}{q_0-1}}(U_k(R))} \|u_3\|_{L^{q_0}(U_k(R))} \, ds \\
& \lesssim \sum_{k=0}^n r_k^{-2} \int_{-r_k^2}^0 \|(\mathbf{u}, \mathbf{b})\|_{L^{\frac{2q_0}{q_0-1}}(U_k(R))}^2 \|u_3\|_{L^{q_0}(U_k(R))} \, ds \\
\text{(A.5)} \quad & \lesssim \sum_{i=0}^n \mathcal{B}_i(r_i^{-1} E_i(R)),
\end{aligned}$$

which is analogous to (3.10). By (2.1) and (3.14), we have

$$\text{(A.6)} \quad J_3 \lesssim \int_{-1}^0 \|\pi_0\|_{L^{\frac{3}{2}}(U_0(R))} \|u_3\|_{L^3(U_0(R))} \, ds \lesssim \int_{-1}^0 \|(\mathbf{u}, \mathbf{b})\|_{L^3(U_0(R))}^3 \, ds \lesssim \mathcal{E}^{\frac{3}{2}}.$$

Summing up all the estimates of (A.3), (A.4), (A.5) and (A.6), we have (A.2). \square \square

Lemma A.3. *Under the assumptions of Lemma 3.1, we have*

$$\text{(A.7)} \quad K \leq \frac{C\mathcal{E}^{\frac{1}{2}}}{R-\rho} \sum_{i=0}^n r_i^{\frac{1}{2}} (r_i^{-1} E_i(R)) + \frac{C}{R-\rho} \mathcal{E}^{\frac{3}{2}}.$$

Proof.

$$\begin{aligned}
K &= \int_{-1}^t \int_{U_0(R)} \pi_0 \mathbf{u} \cdot \Phi_n \eta \nabla \psi \, dx \, ds + \int_{-1}^t \int_{U_0(R)} \pi_0 \mathbf{u} \cdot \Phi_n \nabla \eta \psi \, dx \, ds \\
&\lesssim \sum_{k=0}^n \int_{-1}^t \int_{U_0(R)} \pi_0 \mathbf{u} \cdot \Phi_n \phi_k \nabla \psi \, dx \, ds + \int_{-1}^0 \int_{U_0(R)} |\pi_0| |\mathbf{u}| \, dx \, ds \\
&\lesssim \sum_{k=4}^n \sum_{j=0}^n \int_{-1}^t \int_{U_0(R)} \pi_{0,j} \mathbf{u} \cdot \Phi_n \phi_k \nabla \psi \, dx \, ds \\
&\quad + \sum_{k=0}^3 \int_{-1}^t \int_{U_0(R)} \pi_0 \mathbf{u} \cdot \Phi_n \phi_k \nabla \psi \, dx \, ds + \int_{-1}^0 \int_{U_0(R)} |\pi_0| |\mathbf{u}| \, dx \, ds \\
&\lesssim \sum_{k=4}^n \sum_{j=0}^{k-4} \int_{-1}^t \int_{U_0(R)} \pi_{0,j} \mathbf{u} \cdot \Phi_n \phi_k \nabla \psi \, dx \, ds \\
&\quad + \sum_{k=4}^n \sum_{j=k-3}^n \int_{-1}^t \int_{U_0(R)} \pi_{0,j} \mathbf{u} \cdot \Phi_n \phi_k \nabla \psi \, dx \, ds + \frac{1}{R-\rho} \int_{-1}^0 \int_{U_0(R)} |\pi_0| |\mathbf{u}| \, dx \, ds \\
\text{(A.8)} \quad &\stackrel{\text{def}}{=} K_1 + K_2 + K_3.
\end{aligned}$$

Analogously with (A.4), we have

$$\begin{aligned}
K_1 &= \sum_{j=0}^{n-4} \sum_{k=j+4}^n \int_{-1}^t \int_{U_0(R)} \pi_{0,j} \mathbf{u} \cdot \Phi_n \phi_k \nabla \psi \, dx \, ds \\
&\lesssim \frac{1}{R-\rho} \sum_{j=0}^{n-4} \sum_{k=j+4}^n r_k^{-1} \int_{-r_k^2}^0 \|\pi_{0,j}\|_{L^{\frac{3}{2}}(U_k(R))} \|\mathbf{u}\|_{L^3(U_k(R))} \, ds
\end{aligned}$$

$$\begin{aligned}
&\lesssim \frac{1}{R-\rho} \sum_{j=0}^{n-4} \sum_{k=j+4}^n r_k^{-\frac{1}{3}} r_j^{-\frac{2}{3}} \int_{-r_k^2}^0 \|(\mathbf{u}, \mathbf{b})\|_{L^3(U_j(R))}^3 \, ds \\
&\lesssim \frac{1}{R-\rho} \sum_{j=0}^{n-4} \sum_{k=j+4}^n r_k^{\frac{1}{6}} r_j^{-\frac{2}{3}} \|(\mathbf{u}, \mathbf{b})\|_{L^4(-r_k^2, 0; L^3(U_j(R)))}^3 \\
(A.9) \quad &\lesssim \frac{\mathcal{E}^{\frac{1}{2}}}{R-\rho} \sum_{i=0}^n r_i^{\frac{1}{2}} (r_i^{-1} E_i(R)).
\end{aligned}$$

Analogously with (A.5), we have

$$\begin{aligned}
K_2 &= \sum_{k=4}^n \int_{-1}^t \int_{U_0(R)} \mathcal{J}((\mathbf{u} \otimes \mathbf{u} - \mathbf{b} \otimes \mathbf{b}) \cdot \chi_{k-3}) \mathbf{u} \cdot \Phi_n \phi_k \nabla \psi \, dx \, ds \\
&\lesssim \frac{1}{R-\rho} \sum_{k=0}^n r_k^{-1} \int_{-r_k^2}^0 \|(\mathbf{u}, \mathbf{b})\|_{L^3(U_k(R))}^3 \, ds \\
(A.10) \quad &\lesssim \frac{\mathcal{E}^{\frac{1}{2}}}{R-\rho} \sum_{i=0}^n r_i^{\frac{1}{2}} (r_i^{-1} E_i(R)).
\end{aligned}$$

Analogously with (A.6), we have

$$(A.11) \quad K_3 \lesssim \frac{1}{R-\rho} \int_{-1}^0 \|(\mathbf{u}, \mathbf{b})\|_{L^3(U_0(R))}^3 \, ds \lesssim \frac{1}{R-\rho} \mathcal{E}^{\frac{3}{2}}.$$

Summing up (A.8), (A.9), (A.10) and (A.11), we have (A.7). \square \square

Lemma A.4. *Under the assumptions of Lemma 3.1, we have*

$$(A.12) \quad H \leq \frac{C}{(R-\rho)^3} \mathcal{E}^{\frac{3}{2}}.$$

Proof.

$$\begin{aligned}
H &= - \sum_{k=0}^n \int_{-1}^t \int_{U_0(R)} \nabla \pi_h \cdot \mathbf{u} \cdot (\Phi_n \phi_k \psi) \, dx \, ds \\
&= \sum_{k=0}^3 \int_{-1}^t \int_{U_0(R)} \pi_h \cdot \mathbf{u} \cdot \nabla (\Phi_n \phi_k \psi) \, dx \, ds - \sum_{k=4}^n \int_{-1}^t \int_{U_0(R)} \nabla \pi_h \cdot \mathbf{u} \cdot (\Phi_n \phi_k \psi) \, dx \, ds \\
(A.13) \quad &\stackrel{\text{def}}{=} H_1 + H_2.
\end{aligned}$$

Applying (2.1), Hölder's inequality and the fact that

$$(A.14) \quad \|\pi_h\|_{L^{\frac{3}{2}}(\mathbb{R}^3)} \leq \|\pi\|_{L^{\frac{3}{2}}(\mathbb{R}^3)} + \|\pi_0\|_{L^{\frac{3}{2}}(\mathbb{R}^3)} \lesssim \|(\mathbf{u}, \mathbf{b})\|_{L^3(\mathbb{R}^3)}^2,$$

we have

$$(A.15) \quad H_1 \lesssim \frac{1}{R-\rho} \int_{-1}^t \int_{U_0(R)} |\pi_h| |\mathbf{u}| \, dx \, ds \lesssim \frac{1}{R-\rho} \|\pi_h\|_{L^{\frac{3}{2}}(Q_0(1))} \cdot \|\mathbf{u}\|_{L^3(Q_0(1))} \lesssim \frac{1}{R-\rho} \mathcal{E}^{\frac{3}{2}}.$$

Moreover,

$$\begin{aligned}
H_2 &\lesssim \sum_{k=4}^n r_k^{-1} \int_{Q_k(\frac{R+\rho}{2})} |\nabla \pi_h| \cdot |\mathbf{u}| \, dx \, ds \\
&\lesssim \sum_{k=4}^n r_k^{-1} \|\nabla \pi_h\|_{L^{\frac{3}{2}}(Q_k(\frac{R+\rho}{2}))} \|\mathbf{u}\|_{L^3(Q_k(R))} \\
(A.16) \quad &\lesssim \sum_{k=4}^n r_k^{-\frac{1}{3}} \|\nabla \pi_h\|_{L^{\frac{3}{2}}(-r_k^2, 0; L^\infty(U_k(\frac{R+\rho}{2})))} \|\mathbf{u}\|_{L^3(Q_k(R))}.
\end{aligned}$$

For any $x^* \in U_k(\frac{R+\rho}{2})$, we have $B(x^*, \frac{R-\rho}{4}) \subset U_1(R)$ due to $k \geq 4$ and $|R - \rho| \leq \frac{1}{2}$. Since π_h is harmonic in $U_1(R)$, using the mean value property, we have

$$|\nabla \pi_h|(x^*) \lesssim \frac{1}{|R - \rho|^4} \int_{B(x^*, \frac{R-\rho}{4})} |\pi_h| \, dx \lesssim \frac{1}{(R - \rho)^3} \|\pi_h\|_{L^{\frac{3}{2}}(U_1(R))}.$$

Hence,

$$\begin{aligned}
H_2 &\lesssim \frac{1}{(R - \rho)^3} \sum_{k=4}^n r_k^{-\frac{1}{3}} \|\pi_h\|_{L^{\frac{3}{2}}(-r_k^2, 0; L^{\frac{3}{2}}(U_1(R)))} \|\mathbf{u}\|_{L^3(Q_k(R))} \\
&\lesssim \frac{1}{(R - \rho)^3} \sum_{k=4}^n r_k^{\frac{1}{6}} \|(\mathbf{u}, \mathbf{b})\|_{L^4(-r_k^2, 0; L^3(\mathbb{R}^3))}^3 \\
(A.17) \quad &\lesssim \frac{1}{(R - \rho)^3} \mathcal{E}^{\frac{3}{2}}.
\end{aligned}$$

Summing up (A.13), (A.15) and (A.17), we have (A.12). \square \square

Lemma A.5. [24, Lemma A.3] *For any*

$$1 \leq p_* < p = \frac{2q}{q-3}, \quad 3 < q < +\infty,$$

we have

$$\sum_{k=0}^{+\infty} r_k^{1 - \frac{2}{p_*} - \frac{3}{q}} \left(\int_{-r_k^2}^0 \|f(\cdot, s)\|_{L^q(B(2))}^{p_*} \, ds \right)^{\frac{1}{p_*}} \leq C \|f\|_{L^{p,1}(-1,0; L^q(B(2)))}.$$

For the sake of simplicity, we define

$$(A.18) \quad D(\pi, z_0, r) = r^{-2} \int_{Q(z_0, r)} |\pi|^{\frac{3}{2}} \, dx \, dt, \quad F(\mathbf{u}, \mathbf{b}, z_0, r) = r^{-2} \int_{Q(z_0, r)} |\mathbf{u}|^3 + |\mathbf{b}|^3 \, dx \, dt,$$

$$(A.19) \quad D(\pi, r) \stackrel{\text{def}}{=} D(\pi, (0, 0), r), \quad F(\mathbf{u}, \mathbf{b}, r) \stackrel{\text{def}}{=} C(\mathbf{u}, \mathbf{b}, (0, 0), r).$$

Lemma A.6. *Let $\pi \in L^{\frac{3}{2}}(Q(1))$ solve $\Delta \pi = \nabla \cdot \nabla \cdot (\mathbf{u} \otimes \mathbf{u} - \mathbf{b} \otimes \mathbf{b})$ in the sense of distributions. If there exists a constant K_0 such that for all $0 < r \leq R \leq 1$,*

$$(A.20) \quad F(\mathbf{u}, \mathbf{b}, r) \leq K_0,$$

then for some $\alpha > 0$ and all $0 < r \leq R$,

$$(A.21) \quad D(\pi, r) \leq C \left(\frac{r}{R}\right)^\alpha \cdot D(\pi, R) + CK_0.$$

Proof. We claim that for $0 < 2r < \rho \leq R$,

$$(A.22) \quad D(\pi, r) \leq C_2 \left(\frac{r}{\rho}\right) D(\pi, \rho) + C_2 \left(\frac{\rho}{r}\right)^2 F(\mathbf{u}, \mathbf{b}, \rho).$$

Actually, we write $\pi = \pi_0 + \pi_h$, where $\pi_0 = \mathcal{J}((\mathbf{u} \otimes \mathbf{u} - \mathbf{b} \otimes \mathbf{b}) \cdot \chi_{B(\rho)})$ and

$$(A.23) \quad D(\pi_0, r) \lesssim r^{-2} \|\mathbf{u} \otimes \mathbf{u} - \mathbf{b} \otimes \mathbf{b}\|_{L^{\frac{3}{2}}(Q(\rho))}^{\frac{3}{2}} \lesssim \left(\frac{\rho}{r}\right)^2 F(\mathbf{u}, \mathbf{b}, \rho).$$

π_h is harmonic in $B(\rho)$ and the mean value property of π_h implies that

$$(A.24) \quad D(\pi_h, r) \lesssim r \int_{-r^2}^0 \|\pi_h\|_{L^\infty(B(r))}^{\frac{3}{2}} dt \lesssim \left(\frac{r}{\rho}\right) D(\pi_h, \rho) \lesssim \left(\frac{r}{\rho}\right) (D(\pi, \rho) + D(\pi_0, \rho)).$$

Summing up (A.23) and (A.24), we have (A.22).

Let $\theta \in (0, \frac{1}{2})$. By (A.20) and (A.22), we have that for $0 < r \leq R$,

$$(A.25) \quad D(\pi, \theta r) \leq C_2 \cdot \theta D(\pi, r) + C_2 K_0 \theta^{-2}.$$

We choose θ such that $C_2 \cdot \theta = \frac{1}{2}$. For $\theta R < r \leq R$, we have

$$(A.26) \quad D(\pi, r) \leq \theta^{-2} R^{-2} \int_{Q(R)} |\pi|^{\frac{3}{2}} dx dt.$$

This together with a standard iteration yields (A.21) with $\alpha = -\frac{\ln 2}{\ln \theta} > 0$. \square \square

Lemma A.7. [23, Theorem 1.1] *There exists an absolute constant $\varepsilon_0 > 0$ with the following property. If $(\mathbf{u}, \mathbf{b}, \pi)$ is a suitable weak solution to the MHD equations in $Q(1)$ satisfying that for some $0 < R \leq 1$, $Q(z_0, R) \subseteq Q(1)$ and*

$$(A.27) \quad R^{-2} \int_{Q(z_0, R)} \left(|\mathbf{u}|^3 + |\mathbf{b}|^3 + |\pi|^{\frac{3}{2}} \right) dx dt < \varepsilon_0,$$

then the solution is regular at $z_0 = (x_0, t_0)$.

Lemma A.8 (stability of singularities). *Let $(\mathbf{u}_k, \mathbf{b}_k, \pi_k)$ be a sequence of suitable weak solutions to the MHD equations (1.1) in $Q(1)$ such that $(\mathbf{u}_k, \mathbf{b}_k) \rightarrow (\mathbf{v}, \mathbf{B})$ in $L^3(Q(1))$, $\pi_k \rightarrow \Pi$ in $L^{\frac{3}{2}}(Q(1))$. Assume $(\mathbf{u}_k, \mathbf{b}_k)$ is singular at $z_k = (x_k, t_k)$, $z_k \rightarrow (0, 0)$ as $k \rightarrow \infty$. Then (\mathbf{v}, \mathbf{B}) is singular at $(0, 0)$.*

Proof. The proof is similar with [19, Lemma 2.1]. If (\mathbf{v}, \mathbf{B}) is regular at $(0, 0)$, then there exists $\rho_0 > 0$ and for all $0 < r < \rho_0$,

$$(A.28) \quad r^{-2} \int_{Q(r)} |\mathbf{v}|^3 + |\mathbf{B}|^3 dx dt \leq C r^3.$$

Since $(\mathbf{u}_k, \mathbf{b}_k)$ is singular at z_k , by (A.18) and Lemma A.7, we have that for all $0 < r < \frac{1}{2}$,

$$(A.29) \quad F(\mathbf{u}_k, \mathbf{b}_k, z_k, r) + D(\pi_k, z_k, r) \geq \varepsilon_0.$$

For sufficient large $N = N(r)$ and all $k \geq N$, we have $Q(z_k, r) \in Q(2r)$ and

$$(A.30) \quad F(\mathbf{u}_k, \mathbf{b}_k, 2r) + D(\pi_k, 2r) \geq \frac{\varepsilon_0}{4}.$$

Denote $\tilde{F}(r) = \limsup_{k \rightarrow \infty} F(\mathbf{u}_k, \mathbf{b}_k, r)$ and $\tilde{D}(r) = \limsup_{k \rightarrow \infty} D(\pi_k, r)$. For all $0 < r < 1$, we have

$$(A.31) \quad \tilde{F}(r) + \tilde{D}(r) \geq \frac{\varepsilon_0}{4}.$$

By (A.28), we have that for all $0 < r < \rho < \rho_0$,

$$(A.32) \quad \tilde{F}(r) = F(\mathbf{v}, \mathbf{B}, r) \leq Cr^3 \leq C\rho^3.$$

Applying an analogous argument in Lemma A.6, we have that for all r with $0 < r < \rho$,

$$(A.33) \quad \tilde{D}(r) \leq C \left(\frac{r}{\rho}\right)^\alpha \cdot \tilde{D}(\rho) + C\rho^3.$$

Accordingly, we have for all $0 < r < \rho < \rho_0$,

$$(A.34) \quad C_3 \left(\frac{r}{\rho}\right)^\alpha \cdot \tilde{D}(\rho) + C_3\rho^3 \geq \frac{\varepsilon_0}{4},$$

where the constant C_3 is independent of r and ρ . It leads to a contradiction if we let $r \rightarrow 0$ and then $\rho \rightarrow 0$. The proof is completed. \square

Lemma A.9. [25, Theorem 1.1] *There is an absolute number $\varepsilon_1 > 0$ with the following property. If $(\mathbf{u}, \mathbf{b}, \pi)$ is a suitable weak solution of (1.1) in $Q(1)$ satisfying that for some $0 < R_0 < 1$ and all $0 < r < R_0$,*

$$(A.35) \quad r^{-2} \int_{Q(r)} |\mathbf{u}|^3 dx dt < \varepsilon_1,$$

then the solution (\mathbf{u}, \mathbf{b}) is regular at $(0, 0)$.

ACKNOWLEDGMENTS

Hui Chen was supported by Natural Science Foundation of Zhejiang Province(LQ19A010002). Chenyin Qian was supported by Natural Science Foundation of Zhejiang Province(LY20A010017). Ting Zhang was in part supported by National Natural Science Foundation of China (11771389, 11931010, 11621101).

REFERENCES

- [1] L. Caffarelli, R. Kohn, and L. Nirenberg. Partial regularity of suitable weak solutions of the Navier–Stokes equations. *Comm. Pure Appl. Math.*, 35(6):771–831, 1982.
- [2] C. Cao and E. S. Titi. Global well-posedness of the three-dimensional viscous Primitve equations of large scale ocean and atmosphere dynamics. *Ann. of Math.*, 166(1):245–267, 2007.
- [3] D. Chae and J. Wolf. On the Serrin-type condition on one velocity component for the Navier–Stokes equations. *Arch. Ration. Mech. Anal.*, 2021.
- [4] J.-Y. Chemin and P. Zhang. On the critical one component regularity for 3-D Navier–Stokes system. *Ann. Sci. Éc. Norm. Supér.*, 49(1):131–167, 2016.
- [5] J.-Y. Chemin, P. Zhang, and Z. Zhang. On the critical one component regularity for 3-D Navier–Stokes system: general case. *Arch. Ration. Mech. Anal.*, 224(3):871–905, 2017.
- [6] H. Chen, D. Fang, and T. Zhang. Critical regularity criteria for Navier–Stokes equations in terms of one directional derivative of the velocity. *Math. Methods Appl. Sci.*, 44(6):5123–5132, 2020.
- [7] Q. Chen, C. Miao, and Z. Zhang. On the regularity criterion of weak solution for the 3D viscous magneto-hydrodynamics equations. *Comm. Math. Phys.*, 284(3):919–930, 2008.
- [8] G. Duvaut and J. L. Lions. Inéquations en thermoélasticité et magnétohydrodynamique. *Arch. Ration. Mech. Anal.*, 46(4):241–279, 1972.

- [9] L. Escauriaza, G. A. Seregin, and V. Šverák. Backward uniqueness for parabolic equations. *Arch. Ration. Mech. Anal.*, 169(2):147–157, 2003.
- [10] M. Giaquinta. *Multiple integrals in the calculus of variations and nonlinear elliptic systems*, volume 105 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1983.
- [11] B. Han, Z. Lei, D. Li, and N. Zhao. Sharp one component regularity for Navier–Stokes. *Arch. Ration. Mech. Anal.*, 231(2):939–970, 2019.
- [12] B. Han and N. Zhao. On the critical blow up criterion with one velocity component for 3D incompressible MHD system. *Nonlinear Anal. Real World Appl.*, 51, 2020.
- [13] C. He and Z. Xin. On the regularity of weak solutions to the magnetohydrodynamic equations. *J. Differ. Equ.*, 213(2):235–254, 2005.
- [14] C. He and Z. Xin. Partial regularity of suitable weak solutions to the incompressible magnetohydrodynamic equations. *J. Funct. Anal.*, 227(1):113–152, 2005.
- [15] I. Kukavica, W. Rusin, and M. Ziane. An anisotropic partial regularity criterion for the Navier–Stokes equations. *J. Math. Fluid Mech.*, 19(1):123–133, 2017.
- [16] J. Leray. Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Math.*, 63(1):193–248, 1934.
- [17] F. Lin. A new proof of the Caffarelli-Kohn-Nirenberg theorem. *Comm. Pure Appl. Math.*, 51(3):241–257, 1998.
- [18] G. Prodi. Un teorema di unicità per le equazioni di Navier–Stokes. *Ann. Mat. Pura Appl.*, 48(1):173–182, 1959.
- [19] W. Rusin and V. Šverák. Minimal initial data for potential Navier–Stokes singularities. *J. Funct. Anal.*, 260(3):879–891, 2011.
- [20] M. Sermange and R. Temam. Some mathematical questions related to the MHD equations. *Comm. Pure Appl. Math.*, 36(5):635–664, 1983.
- [21] J. Serrin. On the interior regularity of weak solutions of the Navier–Stokes equations. *Arch. Ration. Mech. Anal.*, 9(1):187–195, 1962.
- [22] S. Takahashi. On interior regularity criteria for weak solutions of the Navier–Stokes equations. *Manuscr. Math.*, 69(3):237–254, 1990.
- [23] V. Vyalov. Partial regularity of solutions to the magnetohydrodynamic equations. *J. Math. Sci.*, 150(1):1771–1786, 2008.
- [24] W. Wang, D. Wu, and Z. Zhang. Scaling invariant Serrin criterion via one velocity component for the Navier–Stokes equations. *eprint arXiv:2005.11906*, 2020.
- [25] W. Wang and Z. Zhang. On the interior regularity criteria for suitable weak solutions of the magnetohydrodynamics equations. *SIAM J. Math. Anal.*, 45(5):2666–2677, 2013.
- [26] Y. Wang. BMO and the regularity criterion for weak solutions to the magnetohydrodynamic equations. *J. Math. Anal. Appl.*, 328(2):1082–1086, 2007.
- [27] Z. Zhang. Remarks on the global regularity criteria for the 3D MHD equations via two components. *Z. Angew. Math. Phys.*, 66(3):977–987, 2015.
- [28] Y. Zhou. Remarks on regularities for the 3D MHD equations. *Discrete Contin. Dyn. Syst.*, 12(5):881–886, 2005.

(H. Chen) SCHOOL OF SCIENCE, ZHEJIANG UNIVERSITY OF SCIENCE AND TECHNOLOGY, HANGZHOU, 310023, PEOPLE’S REPUBLIC OF CHINA

Email address: `chenhui@zust.edu.cn`

(C. Qian) DEPARTMENT OF MATHEMATICS, ZHEJIANG NORMAL UNIVERSITY JINHUA, 321004, CHINA

Email address: `qcyjcsx@163.com`

(T. Zhang) SCHOOL OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU 310027, PEOPLE’S REPUBLIC OF CHINA

Email address: `zhangting79@zju.edu.cn`