

A New Catalog of Asymptotic Giant Branch Stars in Our Galaxy

KYUNG-WON SUH¹

¹*Department of Astronomy and Space Science, Chungbuk National University, Cheongju-City, 28644, Republic of Korea*

(Received June 1, 2021; Revised June 30, 2021; Accepted July 5, 2021)

Submitted to ApJS

ABSTRACT

We present a new catalog of 11,209 O-rich AGB stars and 7172 C-rich AGB stars in our Galaxy identifying more AGB stars in the bulge component and considering more visual carbon stars. For each object, we cross-identify the IRAS, AKARI, MSX, WISE, 2MASS, and AAVSO counterparts. We present the new catalog in two parts: one is based on the IRAS PSC for brighter or more isolated objects; the other one is based on the ALLWISE source catalog for less bright objects or objects in crowded regions. We present various infrared two-color diagrams (2CDs) for the sample stars. We find that the theoretical dust shell models can roughly explain the observations of AGB stars on the various IR 2CDs. We investigate IR properties of SiO and OH maser emission sources in the catalog. For Mira variables in the sample stars, we find that the IR colors get redder for longer pulsation periods. We also study infrared variability of the sample stars using the WISE photometric data in the last 12 yr: the ALLWISE multiepoch data and the Near-Earth Object WISE Reactivation (NEOWISE-R) 2021 data release. We generate light curves using the WISE data at W1 and W2 bands and compute the Lomb-Scargle periodograms for all of the sample stars. From the WISE light curves, we have found useful variation parameters for 3710 objects in the catalog, for which periods were either known or unknown in previous works.

Keywords: Asymptotic giant branch stars (2100); Circumstellar dust (236); Long period variable stars (935); Milky Way Galaxy(1054); Infrared astronomy (786); Radiative transfer (1335)

1. INTRODUCTION

Asymptotic giant branch (AGB) stars are believed to be low-to-intermediate-mass stars ($0.5 - 10 M_{\odot}$) in the last evolutionary phase evolving into post-AGB stars and planetary nebulae (Siess 2006; Höfner & Olofsson 2018). Almost all AGB stars are long-period variables (LPVs) with large amplitude pulsations. As the early-phase AGB stars evolve into the thermally pulsing AGB (TP-AGB) phase, they produce dust grains more effectively and show higher mass-loss rates (e.g., Jiménez-Esteban et al. 2015).

Based on chemistry of the photosphere and/or the circumstellar dust envelope, AGB stars are classified as O-rich AGB (OAGB) or C-rich AGB (CAGB). The spectral energy distributions (SEDs) of OAGB stars show 10 and 18 μm features due to amorphous silicate dust. Low mass-loss rate OAGB (LMOA) stars with thin dust envelopes show the emission features and high mass-loss rate OAGB (HMOA) stars with thick dust envelopes show the absorption features at the same wavelengths (e.g., Suh 1999). To reproduce the

detailed SEDs of LMOA stars, amorphous alumina (Al_2O_3 ; Suh 2016) and Fe-Mg oxides (Th. Posch et al. 2001) dust grains are also necessary. For CAGB stars, SiC and MgS grains as well as featureless amorphous carbon (AMC) dust can reproduce the SEDs (e.g., Suh 2000; Hony et al. 2002).

CAGB stars are generally believed to be the evolutionary successors of OAGB stars. When M-type OAGB stars of intermediate mass range ($1.55 M_{\odot} \leq M < 4 M_{\odot}$: for solar metallicity; Groenewegen et al. 1995) go through carbon dredge-up processes and so the C/O ratio is larger than one, O-rich dust formation ceases and the stars become visual carbon stars in the AGB phase. The visual carbon stars evolve into infrared carbon stars with thick C-rich dust envelopes and high mass-loss rates (e.g., Suh 2000). It is generally thought that S stars (more specifically, intrinsic S stars) are in the intermediate phase between OAGB and CAGB stars (e.g., Suh & Kwon 2011). Though this M-S-C evolutionary sequence could not be right for all AGB stars (e.g., Chan & Kwok 1990), there is much evidence that supports this idea (e.g., Suh 2020).

The Infrared Astronomical Satellite (IRAS), Infrared Space Observatory (ISO), Midcourse Space Experiment (MSX), AKARI, Two-Micron All-Sky Survey (2MASS), and Wide-field Infrared Survey Explorer (WISE) have provided

various IR observational data, which have been useful to identify and study new AGB stars. Using various IR two-color diagrams (2CDs), we can study properties of the central stars and dust envelopes for a large sample of AGB stars (e.g., [Sevenster 2002](#); [Suh & Kwon 2011](#)). Thanks to the optical gravitational lensing experiment (OGLE) projects ([Soszyński et al. 2013a](#)), a larger number of LPVs in the Galactic bulge are identified and studied.

In 2009, WISE ([Wright et al. 2010](#)) started mapping the sky. The ALLWISE multiepoch photometry table obtained in 2009-2010 provided the photometric data at four bands (3.4, 4.6, 12, and 22 μm ; W1, W2, W3, and W4). And the Near-Earth Object WISE Reactivation (NEOWISE-R) mission ([Mainzer et al. 2014](#)) has been providing photometric data at W1 and W2 bands for last seven yr (14 epochs; 2021 data release), two in every year between 2014 and 2020. The ALLWISE multiepoch photometry table and the NEOWISE-R data may allow characterization of the periodic variations of IR emission from AGB stars at the W1 and W2 bands in last 12 yr.

In this paper, we present a new catalog of AGB stars in our Galaxy, identifying more AGB stars in the bulge component and considering more visual carbon stars. Section 2 presents the new catalog in two parts: one is based on the IRAS PSC and the other is based on the ALLWISE source catalog. For each object, we cross-identify the IRAS, AKARI, MSX, WISE, and 2MASS counterparts. Section 3 describes the theoretical radiative transfer models for dust shells around AGB stars. Section 4 presents various infrared 2CDs using the IR photometric data compared with theoretical models. In Section 5, we compare the number distribution of observed IR colors for different classes (or subgroups) of AGB stars to find differences in the IR properties. Section 6 describes spatial distributions of the AGB stars in our Galaxy. Using the new sample stars, we present infrared properties of known pulsating variables in Section 7. In Section 8, we study infrared variability of the sample stars using the WISE photometric data at the W1 and W2 bands in the last 12 yr: the ALLWISE multiepoch data that were acquired between 2009 and 2010 and the NEOWISE-R 2021 data release that was acquired from 2013 until the end of 2020. Section 9 presents the catalog data. Finally, Section 10 summarizes results of the paper.

2. SAMPLE STARS

A catalog of AGB stars for 3003 OAGB, 1168 CAGB, 329 S-type, and 35 silicate carbon stars in our Galaxy was presented by [Suh & Kwon \(2011\)](#) based on the IRAS PSC. [Kwon & Suh \(2012\)](#) presented a revised list of 3373 OAGB stars considering new SiO maser sources. [Kwon & Suh \(2014\)](#) presented a revised sample of 29 silicate carbon stars. [Suh & Hong \(2017\)](#) presented the revised catalog of 3828 OAGB and 1168 CAGB stars (version 2017; <http://web.chungbuk.ac.kr/~kwsuh/agn.htm>).

In this work, we present a new catalog of 11,209 OAGB stars and 7172 CAGB stars in our Galaxy, identifying more

AGB stars in the bulge component and considering more visual carbon stars (see Tables 1 and 2).

Because IRAS has a large beam size and limited sensitivity, it is not possible to find appropriate counterparts for a major portion of newly identified AGB stars, most of which are less bright or smaller objects in crowded regions. Therefore, we present the new revised catalog in two parts: one is based on the IRAS PSC for brighter or more isolated objects; the other is based on the ALLWISE source catalog for less bright objects or objects in crowded regions, for which the IRAS could not observe properly.

Table 1 lists the class name, subgroup name, original reference, and numbers of selected objects for AGB-IRAS objects. Table 2 lists the class name, subgroup name, original reference, and numbers of selected objects for AGB-WISE objects. While IRAS PSC data are available only for the 5908 OAGB-IRAS and 3596 CAGB-IRAS objects, the ALLWISE data are available for most of the AGB-IRAS objects as well as 5301 OAGB-WISE and 3576 CAGB-WISE objects.

Table 1 lists the objects whose original references are based on the IRAS PSC (subgroup names: SH, UR, JB, WU, and AM), all of which are compiled into the AGB-IRAS catalog (see Sections 2.3 and 2.4). Table 3 lists five objects excluded from the list of OAGB Stars (OI-SH) in [Suh & Hong \(2017\)](#).

Tables 1 also lists a major portion of the objects whose original references are not based on the IRAS PSC (subgroup names: ME, ST, OG, and GC), which can be either AGB-IRAS or AGB-WISE objects depending on the existence of the positive IRAS PSC counterpart (see Section 2.2). The objects that can be positively identified with the IRAS PSC sources are compiled into the AGB-IRAS catalog (Table 1). All the other objects with ALLWISE counterparts, which cannot be positively identified with the IRAS PSC sources, are compiled into the AGB-WISE catalog (Table 2).

The sample of 18,381 Galactic AGB stars is composed of 8407 Mira variables (see Tables 1 and 2) according to the American Association of Variable Star Observers (AAVSO) international variable star index (VSX; version 2021 April 19; [Watson et al. 2021](#)). We have also considered the General Catalogue of Variable Stars (GCVS version 5.1; [Samus et al. 2017](#)). In this work, we use the AAVSO which includes all of the LPVs in the GCVS list as well as new lists from other observations.

2.1. Infrared photometric data for the sample stars

IRAS ([Beichman et al. 1988](#)) conducted an all-sky survey and the point source catalog (PSC) provided photometric data at four bands (12, 25, 60, and 100 μm). MSX ([Egan et al. 2003](#)) surveyed the Galactic plane with higher sensitivity and spatial resolution at four MIR bands (8.28, 12.13, 14.65, and 21.34 μm) for 441,879 sources. AKARI ([Murakami et al. 2007](#)) provided PSC data at two bands (9 and 18 μm) and bright-source catalog (BSC) data at four bands (65, 90, 140, and 160 μm). 2MASS ([Cutri et al. 2003](#)) provided fluxes at J (1.24 μm), H (1.66 μm), and K (2.16 μm) bands. The field of view (FOV) pixel sizes of the IRAS, MSX, AKARI PSC,

Table 1. Sample of AGB stars based on the IRAS PSC (AGB-IRAS)

Class	Subgroup	Reference	Number	Selected	Duplicate	Added-Excluded	Remaining ¹
OAGB-IRAS	OI-SH	Suh & Hong (2017)	3828	3828	0	3 ² -5 ³	3826 (1560)
OAGB-IRAS	OI-UR	Urago et al. (2020)	42 ⁴	37 ⁵	0	0	37 (9)
OAGB-IRAS	OI-JB	Jiménez-Esteban et al. (2015)	37	37	32 ⁶	0	5 (0)
OAGB-IRAS	OI-JB	Blommaert et al. (2018)	8	8	7 ⁶	0	1 (0)
OAGB-IRAS	OI-ME	Messineo et al. (2018)	571	166 ⁷	17 ⁶	-22 ⁸	127 (6)
OAGB-IRAS	OI-ST	Stroh et al. (2019)	1427	735 ⁷	59 ⁹ +3 ¹⁰	0	673 (32)
OAGB-IRAS	OI-OG	Soszyński et al. (2013a)	6039 ¹¹	1116 ¹²	59 ⁶	0	1057 (1057)
OAGB-IRAS	OI-WU	Wu et al. (2018)	44	44	25 ⁶	0	19 (19)
OAGB-IRAS	OI-AM ¹³	This work (IRAS PSC)	894	894	731 ¹⁴	0	163 (163)
OAGB-IRAS	OI-all	-	-	-	-	-	5908 (2846)
CAGB-IRAS	CI-SH	Suh & Hong (2017)	1168	1168	0	1 ¹⁵ -3 ² -1 ¹⁶	1165 (340)
CAGB-IRAS	CI-UR	Urago et al. (2020)	42 ⁴	5 ⁵	0	0	5 (0)
CAGB-IRAS	CI-GC	Alksnis et al. (2001)	6891	3379 ¹²	925 ¹⁷ +15 ¹⁸ +20 ¹⁹	-2 ²⁰	2417 (99)
CAGB-IRAS	CI-OG	Soszyński et al. (2013a)	168 ¹¹	9 ¹²	0	0	9 (9)
CAGB-IRAS	CI-all	-	-	-	-	-	3596 (447)
S stars	SI	Suh & Kwon (2011)	329	329	0	1 ²¹	330
silicate carbon stars	SCI	Kwon & Suh (2014)	29	29	0	0	29

¹number in parentheses denotes number of Miras in AAVSO (version 2021-04-19; [Watson et al. 2021](#)). ²three objects in CI-SH are SiO maser sources (OAGB stars) without clear CAGB evidences. ³YSOs or RSGs (see Tables 3). ⁴the number of the original IRAS PSC sources (108) minus duplicate OI-SH sources (66) ⁵color-selected O-AGB or C-AGB stars (see Section 2.3). ⁶in OI-SH. ⁷SiO maser sources with positive IRAS PSC counterparts. ⁸RSGs. ⁹58 objects in OI-SH and one object in OI-JB. ¹⁰IRAS 17105-3746 is in CI-SH (IRAS LRS type C; SiO maser source), IRAS 17001-3651 is an S star (SI), IRAS 15575-5238 is a silicate carbon star (SCI). ¹¹color-selected OAGB or CAGB stars from the OGLE3 sample of Miras in the Galactic bulge (see Section 2.5). One exception is IRAS 18100-2808, which is an OAGB star (IRAS LRS type E) with the CAGB color. ¹²sources with positive IRAS PSC counterparts. ¹³AAVSO Miras with IRAS LRS type E (see Section 2.3). ¹⁴694 in OI-SH, 7 in OI-OG, 6 in OI-ST, 1 in OI-WU, 19 in SI, and 4 in SCI. ¹⁵IRAS 18027-1316 is a typical visual carbon star. ¹⁶IRAS 06176-1036 is a planetary nebula (Red Rectangle). ¹⁷in CI-SH ¹⁸OAGB stars (in OI-SH). ¹⁹two S stars (SI) and 18 silicate carbon stars (SCI). ²⁰duplicate objects. ²¹IRAS 19354+5005 is an intrinsic S star.

Table 2. Sample of AGB stars based on the ALLWISE source catalog (AGB-WISE)

Class	Subgroup	Reference	Number	Selected	Duplicate	Excluded	Remaining ¹
OAGB-WISE	OW-ME	Messineo et al. (2018)	571	196 ²	0	21 ³ +18 ⁴	157 (6)
OAGB-WISE	OW-ST	Stroh et al. (2019)	1427	285 ²	0	54 ⁴	231 (5)
OAGB-WISE	OW-OG	Soszyński et al. (2013a)	6039 ⁵	4923 ⁶	3 ⁷	7 ⁸	4913 (4910 ⁹)
OAGB-WISE	OW-all	-	-	-	-	-	5301 (4921)
CAGB-WISE	CW-GC	Alksnis et al. (2001)	6891	3512 ⁶	0	95 ⁴	3417 (43)
CAGB-WISE	CW-OG	Soszyński et al. (2013a)	168 ⁵	159 ⁶	0	0	159 (157 ¹⁰)
CAGB-WISE	CW-all	-	-	-	-	-	3576 (193)

¹number in parentheses denotes number of Miras in AAVSO. ²SiO maser sources without positive IRAS PSC counterparts. ³RSGs. ⁴no WISE counterparts. ⁵color-selected OAGB or CAGB stars from the OGLE3 sample of Miras in the Galactic bulge (see text). ⁶sources without positive IRAS PSC identification. ⁷in OW-ST. ⁸duplicate objects in cross-matching ALLWISE sources. ⁹For OGLE3 Mira objects, two objects are classified into SRA and one into SR in AAVSO. ¹⁰For OGLE3 Mira objects, one object is classified into SRA and one into RCB: in AAVSO.




Figure 1. Comparison of IRAS and WISE fluxes for AGB-IRAS objects (see Table 1). The number in parentheses denotes the number of the plotted objects with good-quality observed data. See Section 2.2.




Figure 2. A WISE-2MASS 2CD for color-selected AGB-IRAS (OI-UR, CI-UR, OI-OG, and CI-OG) and AGB-WISE (OW-OG and CW-OG) stars (see Tables 1 and 2) compared with theoretical models (see Section 3). The brown line roughly distinguishes between OAGB and CAGB stars. For OAGB models (silicate $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.05, 0.1, 0.5, 1, 3, 7, 15, 30$, and 40 from left to right. For CAGB models (AMC $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.1, 0.5, 1, 2, 3$, and 5 from left to right. For each subgroup, the number of objects is shown. The number in parentheses denotes the number of the plotted objects on the 2CD with good-quality observed colors.




Figure 3. A WISE-2MASS 2CD for color-selected OGLE3 Miras (see Section 2.5) compared with theoretical models (see Section 3). The brown line roughly distinguishes between OAGB and CAGB stars. For OAGB models (silicate $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.05, 0.1, 0.5, 1, 3, 7, 15, 30$, and 40 from left to right. For CAGB models (AMC $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.1, 0.5, 1, 2, 3$, and 5 from left to right. For each subgroup, the number of objects is shown. The number in parentheses denotes the number of the plotted objects on the 2CD with good-quality observed colors. Note that the 321 objects without good-quality observed colors are not plotted and they are not compiled into the new catalog.

Table 3. Five Objects excluded from the list of OAGB stars in Suh & Hong (2017) (OI-SH)

IRAS PSC	Other name	Remark	Reference
05380-0728	HBC 494	YSO	Cieza et al. (2018)
07209-2540	VY CMa	RHG	Humphreys et al. (2021)
16547-4247	-	YSO	Zapata et al. (2019)
18272+0114	OO Ser	YSO	Hodapp et al. (2012)
19312+1950	-	YSO or RSG	Cordiner et al. (2016)

AKARI BSC, and 2MASS images are $0'.75 \times (4'.5-4'.6)$, $18''.3$, $10''$, $30''$, and $2''$, respectively.

WISE (Wright et al. 2010) surveyed the entire sky. The ALLWISE source catalog provided the photometric data at $3.4 \mu\text{m}$ (W1), $4.6 \mu\text{m}$ (W2), $12 \mu\text{m}$ (W3), and $22 \mu\text{m}$ (W4) bands. The FOV pixel sizes are $2''.75$, $2''.75$, $2''.75$, and $5''.5$, and the 5σ photometric sensitivities are 0.068, 0.098, 0.86, and 5.4 mJy for the four WISE bands.

Though IRAS and AKARI data have been very useful for studying AGB stars in our Galaxy (e.g., Suh & Kwon 2011), the number of the cross-identified objects for the new sample

Table 4. IR bands and zero magnitude flux values

Band	$\lambda_{ref} (\mu\text{m})$	ZMF (Jy)	Telescope	Reference
K[2.2]	2.159	666.7	2MASS	Cohen et al. (2003)
W1[3.4]	3.35	306.682	WISE	Jarrett et al. (2011)
W2[4.6]	4.60	170.663	WISE	Jarrett et al. (2011)
MA[8.3]	8.28	58.49	MSX	Egan et al. (2003)
AK[9]	9	56.262	AKARI	Murakami et al. (2007)
IR[12]	12	28.3	IRAS	Beichman et al. (1988)
W3[12] ¹	12 (11.56)	28.3 (29.045)	WISE	Jarrett et al. (2011)
MC[12.1]	12.13	26.51	MSX	Egan et al. (2003)
MD[14.7]	14.65	18.29	MSX	Egan et al. (2003)
AK[18]	18	12.001	AKARI	Murakami et al. (2007)
ME[21.3]	21.34	8.8	MSX	Egan et al. (2003)
W4[22]	22.08	8.284	WISE	Jarrett et al. (2011)
IR[25]	25	6.73	IRAS	Beichman et al. (1988)
IR[60]	60	1.19	IRAS	Beichman et al. (1988)

¹ For W3[12], we use a new reference wavelength and zero magnitude flux for theoretical models (original values are given in parentheses; see section 4.2 in Suh 2020).

stars was very limited because of the relatively large beam sizes and weak sensitivities. On the other hand, the WISE

Table 5. Cross-identified IR sources and AAVSO objects for AGB-IRAS sample stars

Subgroup ¹	IRAS	AKARI	ALLWISE	MSX	AAVSO
OI-SH	3826	3673	3820	2394	2538
OI-UR	37	35	37	28	18
OI-JB	6	6	6	6	5
OI-ME	127	120	127	127	26
OI-ST	673	651	673	673	255
OI-OG	1057	1011	1057	949	1057
OI-AM	163	160	127	102	163
OI-all	5908	5675	5722	4279	4081
CI-SH	1165	1149	1164	753	890
CI-UR	5	5	5	4	1
CI-GC	2417	2393	2413	1848	2117
CI-OG	9	8	9	7	9
CI-all	3596	3555	3591	2612	3017

¹ See Table 1. The numbers of the cross-identified 2MASS sources are the same as the numbers of IRAS PSC sources.

data would be more useful for studying dim objects or objects in crowded regions.

In this paper, we use only good-quality observational data at all wavelength bands for the IRAS, 2MASS, WISE, AKARI, and MSX photometric data ($q = 3$ for IRAS and AKARI; $q = A$ for 2MASS; $q = A$ or B for WISE; $q = 3$ or 4 for MSX).

Table 4 lists the IR bands used in this work. For each band, the reference wavelength (λ_{ref}) and zero magnitude flux (ZMF) value are also shown. The color index is defined by

$$M_{\lambda_1} - M_{\lambda_2} = -2.5 \log_{10} \frac{F_{\lambda_1}/ZMF_{\lambda_1}}{F_{\lambda_2}/ZMF_{\lambda_2}} \quad (1)$$

where ZMF_{λ_i} is the ZMF at given wavelength (λ_i) (see Table 4). We may obtain the color indices from given fluxes from observations (e.g., IRAS, AKARI, and MSX photometric data are in Jy unit) or from theoretical model SEDs using the reference (or effective or isophotal) wavelength and ZMF specified in the the telescope system manual.

2.2. Cross-matches

For all IRAS PSC sources, we have found the AKARI, 2MASS, ALLWISE, MSX, and AAVSO counterparts by using following method. We cross-identify the AKARI PSC or BSC counterpart by finding the nearest source within $60''$ using the position given in the IRAS PSC (version 2.1). Then, we cross-identify the 2MASS, WISE, and MSX counterparts using the position of the available AKARI PSC or BSC counterpart. Only when there is no AKARI counterpart, we have used the position of the IRAS PSC.

Table 6. Cross-identified IR sources and AAVSO objects for AGB-WISE sample stars

Subgroup ¹	ALLWISE	2MASS	AKARI	MSX	AAVSO
OW-ME	157	157	120	157	15
OW-ST	231	231	190	231	29
OW-OG	4913	4913	3159	4171	4913
OW-all	5301	5301	3469	4559	4957
CW-GC	3417	3187	1987	1627	1388
CW-OG	159	159	70	121	159
CW-all	3576	3346	2057	1748	1547

¹ See Table 2.

For the objects whose original references are not based on the IRAS PSC (subgroup names: ME, ST, OG, and GC), we have found the positive IRAS PSC counterparts using the following method. We find the nearest IRAS PSC, AKARI PSC, ALLWISE, 2MASS, MSX, and AAVSO counterparts within $5''$ - $60''$ (depending on the beam size of the telescope) using the position information in the original reference. Because there could be multiple ALLWISE sources for one IRAS PSC source (with a larger beam size), we need to compare the ALLWISE counterpart obtained from the IRAS PSC position (see the previous paragraph) with the counterpart obtained from the position in the original reference. If the ALLWISE (or AKARI or MSX) counterparts obtained from the two positions (from the IRAS PSC and the original reference) are the same, the IRAS PSC source would be a more reliable counterpart. These objects with the positive IRAS PSC counterparts are compiled into the AGB-IRAS catalog (OI-ME, OI-ST, OI-OG, CI-GC, and CI-OG; see Table 1). All other objects with ALLWISE counterparts, which cannot be positively identified with the IRAS PSC sources, are compiled into the AGB-WISE catalog (OW-ME, OW-ST, OW-OG, CW-GC, and CW-OG; see Table 2).

For WISE data, multiple sample objects may have the same cross-matched ALLWISE source. So we have checked all of the duplicate cross-matched objects and selected only one nearest object for the one ALLWISE source (OW-OG and CW-GC in Table 2).

Figure 1 compares the IRAS PSC and WISE fluxes for all of the AGB-IRAS sample stars (see Table 1). The overall comparison is fairly consistent for most objects. Some objects show large deviations (the IRAS flux is much larger than the WISE flux) possibly due to the contamination by the larger beam size of IRAS.

Tables 5 and 6 list the cross-matched counterparts for all AGB-IRAS and AGB-WISE sample stars.

2.3. Catalogs based on IRAS PSC

Based on the catalog of AGB stars presented by Suh & Hong (2017) (3828 OAGB and 1168 CAGB stars), we have revised it using new available literature. We have excluded

five objects from the OAGB list (OI-SH): four young stellar objects (YSOs) or red supergiants (RSGs) and a red hypergiant (RHG) (see Tables 3). We have moved from CI-SH to OI-SH three objects which are SiO maser sources without clear CAGB evidences (see Tables 1).

The IRAS Low Resolution Spectrograph (LRS; $\lambda = 8\text{--}22\ \mu\text{m}$) data have been very useful to identify important dust features of AGB stars. Kwok et al. (1997) used the IRAS LRS data for 11,224 IRAS PSC source to identify new OAGB and CAGB stars. In the IRAS LRS, OAGB stars with silicate dust envelopes are classified into type E ($10\ \mu\text{m}$ in emission) or A ($10\ \mu\text{m}$ in absorption). CAGB stars with SiC grains are classified into type C ($11.3\ \mu\text{m}$ in emission). There are 715 objects that are classified into group C: 713 objects are in the list of CI-SH (see Table 1), one object (IRAS 13136-4426) is an S star (in SI), and the other one (IRAS 22306+5918) is a composite object.

Figure 2 shows a WISE-2MASS 2CD using W3[12]–W4[22] versus K[2.2]–W3[12] for the sample of AGB stars from Suh & Hong (2017) (3828 OAGB stars in OI-SH and 1168 CAGB stars in CI-SH), which can be used to find a rough guide line that distinguishes between OAGB and CAGB stars. The brown line roughly distinguishes between OAGB and CAGB stars.

Urago et al. (2020) presented a list 108 IRAS PSC objects that are believed to be Mira variables because of the regular variability of the light curves at K band. Sixty-six objects are duplicate with Suh & Hong (2017) (64: OI-SH; 2: CI-SH) and 42 objects remain. The 42 objects are classified into OAGB or CAGB using the guide line on the IR 2CD in Figure 2, which resulted in 32 OAGB and 5 CAGB objects. The remaining 5 objects do not have the available colors to plot on the 2CD, but all of them are likely to be OAGB stars because they show O-rich characters in the IRAS LRS or other IR colors. Therefore, there are 37 OAGB (OI-UR) and 5 CAGB (CI-UR) stars in the sample (see Figure 2 and Table 1).

Figure 2 also shows the color-selected AGB-IRAS (OI-OG and CI-OG) and AGB-WISE (OW-OG and CW-OG) stars from Soszyński et al. (2013a) (see Section 2.5).

From all of the Mira variables with known periods in the AAVSO, we identified 10,753 IRAS PSC counterparts. Among them, there are 9 IRAS LRS type A sources (all objects are in OI-SH) and 894 IRAS LRS type-E sources (731 sources are already in OAGB-IRAS). The remaining 163 Miras, which are IRAS LRS type E sources, are likely to be new OAGB stars in our Galaxy because no object shows any evidences for CAGB or S stars. We compile the 163 AAVSO Miras, which are IRAS LRS type-E sources, into the new catalog (OAGB-IRAS: OI-AM).

2.4. New OAGB stars

Jiménez-Esteban et al. (2015) and Blommaert et al. (2018) studied extremely reddened AGB stars in the Galactic bulge and presented lists of OAGB stars. These OAGB stars are compiled into the catalog based on IRAS PSC (OI-JB in OAGB-IRAS; see Table 1).

SiO maser sources are generally believed to be associated with OAGB stars (mostly Mira variables) with a thin silicate dust envelope (e.g., Deguchi et al. 2012; Stroh et al. 2019). But, two CAGB stars (IRAS LRS type C objects in CI-SH) are SiO maser sources: IRAS 01022+6542 (Deguchi et al. 2012) and IRAS 17105-3746 (Stroh et al. 2019).

Messineo et al. (2018) performed the SiO maser survey of late-type stars in the inner Galaxy. Stroh et al. (2019) performed the Bulge Asymmetries and Dynamical Evolution (BAaDE) SiO Maser Survey at 86 GHz with the Atacama Large Millimeter/submillimeter Array (ALMA) and presented a list of 1427 sources with useful information. The OAGB stars (SiO maser sources) from Messineo et al. (2018) and Stroh et al. (2019) are compiled into two separate catalogs based on the IRAS PSC (OI-ME and OI-ST in OAGB-IRAS) and the ALLWISE source catalog (OW-ME and OW-ST in OAGB-WISE) (see Section 2.2).

Wu et al. (2018) performed SiO maser survey toward off-plane OAGB stars around the orbital plane of the Sagittarius stellar stream and detected SiO maser emission from 44 targets, which are IRAS PSC sources.

We have compiled these OAGB objects into the new OAGB catalog (OI-JB, OI-ME, OW-ME, OI-ST, OW-ST, and OI-WU; see Tables 1 and 2).

2.5. LPVs in the Galactic bulge

The OGLE projects detected many LPVs in the bulge of our Galaxy. The fifteenth part of the OGLE-III Catalog of Variable Stars (OIII-CVS) contains 232,406 LPVs detected in the OGLE-II and OGLE-III fields toward the Galactic bulge. The sample consists of 6528 Miras, 33,235 semiregular variables (SRVs) and 192,643 OGLE small amplitude red giants (OSARGs) (Soszyński et al. 2013a).

A giant star is believed to evolve from a OSARG to an SRV and finally to a Mira, limiting the number of its excited modes and increasing its pulsating periods and amplitudes (Soszyński et al. 2013b). Though most of SRVs and a majority of OSARGs could also be in the AGB phase, most Mira variables are surely in the AGB phase and they are usually oscillating in the fundamental mode and occupy a single sequence in the period-luminosity diagram (Soszyński et al. 2013b). In this work, we consider the 6528 OGLE3 Miras in the Galactic bulge for the new catalog of AGB stars.

Using the guide line on the WISE-2MASS 2CD (see Figure 2), we may roughly distinguish between OAGB and CAGB stars for the sample of the 6528 OGLE3 Miras in the Galactic bulge (Soszyński et al. 2013a). Figure 3 shows the 2CD for the sample of 6528 OGLE3 Miras in the Galactic bulge (Soszyński et al. 2013a), but 321 objects without good-quality observed colors (W3[12]–W4[22] and K[2.2]–W3[12]) are not plotted and they are not compiled into the new catalog. The brown line roughly distinguishes between OAGB and CAGB stars. We find that 6038 and 169 objects can be classified into OAGB and CAGB, respectively. One exception is IRAS 18100-2808, which is an OAGB star (IRAS LRS type E) with the CAGB color. Therefore, 6039

and 168 objects can be compiled into the OAGB and CAGB catalogs, respectively.

These OAGB and CAGB stars with the positive IRAS PSC counterparts (see Section 2.2) are compiled into the AGB-IRAS catalog (OI-OG and CI-OG; see Table 1) and others into the AGB-WISE catalog (OW-OG and CW-OG; see Table 2).

Figure 2 shows the WISE-2MASS 2CD for the color-selected sample stars from Soszyński et al. (2013a) in AGB-IRAS (OI-OG, and CI-OG) and AGB-WISE (OW-OG and CW-OG) catalogs. The 2CD also shows the color selected AGB-IRAS objects (OI-UR and CI-UR) from Urago et al. (2020) (see Section 2.3).

2.6. Visual carbon stars

Alksnis et al. (2001) presented general catalog of galactic carbon stars (3rd edition) which consist of 6891 entries, most of which are believed to be visual carbon stars in the AGB phase. Most CAGB stars in previous catalog (CI-SH; Suh & Hong 2017) are infrared carbon stars with relatively thick C-rich dust envelopes.

Objects in Alksnis et al. (2001) are compiled into two separate catalogs based on the IRAS PSC and the ALLWISE source catalog (CI-GC or CW-GC in Tables 1 and 2; see Section 2.2).

Compared with other subgroups, the basic properties (e.g., detailed IR SEDs or pulsations) of the objects from Alksnis et al. (2001) (in CI-GC or CW-GC) are less known, so it is possible that a greater portion of the objects from CI-GC or CW-GC could not be in the AGB phase. Some objects could be in the post-AGB phase or other stages of stellar evolution.

3. THEORETICAL DUST SHELL MODELS

We use the radiative transfer code DUSTY (Ivezić & Elitzur 1997) for a spherically symmetric dust shell around a central star, which is a blackbody. We use the models of Suh (2020) adapted for the new IR bands. In this paper, we briefly describe the theoretical models. See Section 4 in Suh (2020) for details about the models and their limitations.

For all models, we use a continuous power law ($\rho \propto r^{-2}$) dust density distribution and assume that the dust formation temperature (T_c) is 1000 K. For LMOA stars, we also use $T_c=500$ K. The inner radius of the dust shell is set by the T_c and the outer radius of the dust shell is taken to be 10^4 times the inner radius. We use $10 \mu\text{m}$ as the fiducial wavelength of the dust optical depth (τ_{10}). The radii of spherical dust grains are assumed to be $0.1 \mu\text{m}$ uniformly.

For OAGB stars, we use optical constants of warm (SILW) and cold (SILC) silicate dust from Suh (1999). We also use amorphous alumina (Suh 2016) and $\text{Fe}_{0.9}\text{Mg}_{0.1}\text{O}$ (Henning et al. 1995) dust for OAGB stars. For CAGB stars, we use the optical constants of AMC and SiC dust grains from Suh (2000) and Pégourié (1988), respectively. We also use $\text{Mg}_{0.9}\text{Fe}_{0.1}\text{S}$ (Begemann et al. 1994) for CAGB stars. Table 7 summarizes the model parameters.

Figure 4 shows model SEDs for AGB stars ($T_c=1000$ K) for major dust optical depths. For OAGB models, silicate




Figure 4. Theoretical model SEDs for OAGB stars (silicate; $T_c=1000$ K) and CAGB stars (AMC; $T_c=1000$ K) for a number of dust optical depths (see 3). The reference wavelengths for major IR bands are also indicated (see Table 4).

Table 7. Dust shell model parameters

Model	dust	τ_{10}	T_c (K)	T_*^1 (K)
LMOA	O-rich ²	0.001, 0.01, 0.05, 0.1	500	3000
LMOA	SILW	0.001, 0.01, 0.05, 0.1	1000	3000
LMOA	SILW	0.5, 1, 3	1000	2500
CAGB	C-rich ³	0.001, 0.01, 0.1, 0.5	1000	2500
CAGB	C-rich ³	1, 2, 3, 5	1000	2000
HMOA	SILC	7, 15, 30, 40	1000	2000

¹ the black body temperature of the central star ²SILW, SILW + alumina (from Suh 2016), and SILW + $\text{Fe}_{0.9}\text{Mg}_{0.1}\text{O}$ (from Henning et al. 1995). ³AMC, AMC + SiC, and AMC + $\text{Mg}_{0.9}\text{Fe}_{0.1}\text{S}$ (from Begemann et al. 1994).

dust features at 10 and 18 μm are shown for various dust optical depths (τ_{10}). For CAGB models, SiC dust features at 11.3 μm and $\text{Mg}_{0.9}\text{Fe}_{0.1}\text{S}$ dust features at 28 μm are shown for different dust optical depths. The reference wavelengths for major IR bands are also indicated. We may obtain the theoretical model color indices from the model SEDs using the ZMF values at given reference wavelength (see Table 4 and Section 2.1).

4. INFRARED TWO-COLOR DIAGRAMS - COMPARISON BETWEEN THEORY AND OBSERVATIONS

Although the photometric fluxes are less useful than a full SED, the large number of observations at various wavelength bands can be used to form a 2CD, which can be compared with theoretical models. IR 2CDs are useful to statistically distinguish various properties of AGB stars and post-AGB stars (e.g., Suh & Kwon 2011; Suh 2015). Table 4 lists the IR bands used for the IR 2CDs presented in this work. In this work, we use only good-quality observational data at all wavelength bands (see Section 2.1) for plotting IR 2CDs.

Figures 5 - 8 show various IR 2CDs using different combinations of observed IR colors. We compare the observations with the theoretical dust shell models (see Section 3) for AGB stars. We find that the theoretical dust shell model can roughly reproduce the observations of AGB stars on the IR 2CDs using the dust opacity functions of amorphous silicate and amorphous carbon with a mixture of other dust species.

To consider the Galactic extinction processes suggested by Gordon et al. (2009) (for the wavelength range from visual to NIR bands) and Chiar & Tielens (2006) (from NIR to MIR bands), we plot reddening vectors for IR 2CD using NIR data (see Figures 5- 8).

Generally, the stars that have thick dust shells with large dust optical depths are located in the upper-right regions on the IR 2CDs. On all of the IR 2CDs, we also plot the sequences of theoretical dust shell models at increasing dust optical depth for AGB stars (see Section 3).

Because the 10 μm silicate feature changes from emission to absorption when the dust optical depth becomes larger, there is a change in the slope of the theoretical model line for OAGB stars.

We will discuss the meanings of these 2CDs in the following subsections by comparing the observations with the theoretical models.

4.1. IRAS and 2MASS 2CDs

The upper panel of Figure 5 plots AGB stars in an IRAS 2CD using IR[25]–IR[60] versus IR[12]–IR[25]. We find that the basic theoretical model tracks can roughly explain the observed points. This 2CD has been widely used since van der Veen & Habing (1988) (note that the authors did not make zero-magnitude calibrations for their 2CD) divided this 2CD into eight regions of different classes of heavenly bodies. Sevenster (2002) used this 2CD to explain the properties of observed points of AGB stars and post-AGB stars. Using theoretical dust shell models for AGB stars and post-AGB

stars, Suh (2015) presented possible evolutionary tracks from AGB stars to post-AGB stars and to planetary nebulae on this 2CD.

On the IRAS 2CD using IR[25]–IR[60] versus IR[12]–IR[25], CAGB stars are distributed along a curve in the shape of a 'C'. A group of stars in the upper-left region consists of visual carbon stars (most objects in OI-GC; see Table 1) that show excessive flux at 60 μm , due to the remnant of an earlier phase when the stars were OAGB stars (e.g. Chan & Kwok 1990). A group of stars in the lower region, which extends to the right side, consists of infrared carbon stars. The infrared carbon stars on the right side have thick dust envelopes with large dust optical depths.

The lower panel plots an 2MASS-IRAS 2CD using IR[12]–IR[60] versus K[2.2]–IR[12]. The separation between OAGB and CAGB stars is clearer on this 2CD. If we consider Galactic extinction (see the reddening vector on the 2CD), there would be more observed points of OAGB and CAGB stars that would fit the theoretical models well.

4.2. WISE and 2MASS 2CDs

Figure 6 shows 2CDs using WISE and 2MASS colors. The upper panel of Figure 6 shows an WISE 2CDs using W3[12]–W4[22] versus W1[3.4]–W2[4.6]. The lower panel of Figure 6 shows a WISE-2MASS 2CD using W3[12]–W4[22] versus K[2.2]–W3[12].

Generally, the theoretical dust shell models for OAGB and CAGB stars can reproduce the observed points fairly well on these IR 2CDs, but for the W1[3.4]–W2[4.6] color, the theoretical models with small dust optical depths do not reproduce the observations well. On these 2CDs, the theoretical dust shell models for CAGB stars with AMC, SiC, and $\text{Mg}_{0.9}\text{Fe}_{0.1}\text{S}$ dust grains can reproduce a wider range of observed W3[12]–W4[22] colors.

There is a group of CAGB-WISE objects (visual carbon stars in CI-GC and CW-GC) that show different aspects from infrared carbon stars. The objects are in the upper-left regions of the 2CDs that show bluer W1[3.4]–W2[4.6] or K[2.2]–W3[12] colors but redder W3[12]–W4[22] colors, which would be due to detached O-rich dust shells (that are remnants of an earlier phase when the stars were OAGB stars).

4.3. AKARI, MSX, and 2MASS 2CDs

Figure 7 shows an AKARI-2MASS 2CD using AK[9]–AK[18] versus K[2.2]–AK[9]. On this 2CD, the theoretical dust shell models for OAGB stars with thin detached dust shells ($T_c=500$ K) with warm silicate, amorphous alumina, and $\text{Fe}_{0.9}\text{Mg}_{0.1}\text{O}$ dust are more useful to explain the observed points. We find that the theoretical dust shell models for CAGB stars can reproduce only a narrow range of AK[9]–AK[18] colors.

Figure 8 shows MSX-2MASS 2CDs for all IRAS-AGB and WISE-AGB objects. Though good-quality MSX data are available only for a portion of the sample stars, these 2CDs clearly divide between OAGB and CAGB. Compared with other colors, the opacity used for the theoretical models do




Figure 5. IRAS-2MASS 2CDs for all IRAS-AGB stars (see Table 1) in our Galaxy compared with theoretical models (see Section 3). For OAGB models (silicate $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.05, 0.1, 0.5, 1, 3, 7, 15, 30$, and 40 from left to right. For CAGB models (AMC $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.1, 0.5, 1, 2, 3$, and 5 from left to right. For each class, the number of objects is shown. The number in parentheses denotes the number of the plotted objects on the 2CD with good-quality observed colors.




Figure 6. WISE-2MASS 2CDs for all IRAS-AGB and WISE-AGB stars (see Tables 1 and 2) in our Galaxy compared with theoretical models (see Section 3). For OAGB models (silicate $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.05, 0.1, 0.5, 1, 3, 7, 15, 30$, and 40 from left to right. For CAGB models (AMC $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.1, 0.5, 1, 2, 3$, and 5 from left to right. For each class, the number of objects is shown. The number in parentheses denotes the number of the plotted objects on the 2CD with good-quality observed colors.




Figure 7. An AKARI-2MASS 2CD for all IRAS-AGB and WISE-AGB stars (see Tables 1 and 2) in our Galaxy compared with theoretical models (see Section 3). For OAGB models (silicate $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.05, 0.1, 0.5, 1, 3, 7, 15, 30$, and 40 from left to right. For CAGB models (AMC $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.1, 0.5, 1, 2, 3$, and 5 from left to right. For each class, the number of objects is shown. The number in parentheses denotes the number of the plotted objects on the 2CD with good-quality observed colors.

not reproduce the MSX colors using MA[8.3], MD[14.7], and ME[21.3] bands well. Again, the theoretical dust shell models for CAGB stars can reproduce only a narrow range of MA[8.3]–ME[21.3] and MA[8.3]–MD[14.7] colors.

The upper panel of Figure 8 shows a 2CD using $K[2.2]$ –MA[8.3] versus MA[8.3]–ME[21.3]. Lewis et al. (2020) used this 2CD to discuss the line that separates between CAGB and OAGB stars. We also find that the separation between OAGB and CAGB is relatively clear on this 2CD. If we consider Galactic extinction, there would be more observed points of OAGB stars that would fit the OAGB model with thin detached dust shells. The lower panel of Figure 8 shows a 2CD using MA[8.3]–MD[14.7] versus $K[2.2]$ –MA[8.3].

5. NUMBER DISTRIBUTIONS OF IR COLORS

We may compare the number distribution of observed IR colors for different classes (or subgroups) of AGB stars to find differences in the IR properties. We may also compare the number distributions with the theoretical model colors.

5.1. OAGB and CAGB stars

Figure 9 shows number density distributions of observed IR magnitudes at W2[4.6] and W3[12] bands and colors for AGB stars in the AGB-IRAS and AGB-WISE cata-

logs. We choose two IR colors: $W1[3.4]$ – $W2[4.6]$ and $K[2.2]$ – $W3[12]$ (see Figure 6).

Generally, AGB-IRAS objects are brighter at MIR bands and show redder IR colors than AGB-WISE objects. For both IR colors ($W1[3.4]$ – $W2[4.6]$ and $K[2.2]$ – $W3[12]$), CAGB stars are generally bluer than OAGB stars. We also find that AGB-WISE objects are more concentrated toward the bluer colors and the number decreases with the redder colors, whereas numbers for AGB-IRAS objects are more dispersed into redder colors. In general, AGB-IRAS objects look to be more evolved (or more massive) stars with thicker dust envelopes than AGB-WISE objects.

5.2. OH and SiO Maser sources among OAGB stars

A major portion of OAGB stars are SiO, H₂O, OH maser sources (e.g., Kwon & Suh 2012). We use the lists of SiO maser sources compiled by Kwon & Suh (2012) and new SiO maser sources detected from the Galactic bulge (Messineo et al. 2018; Wu et al. 2018; Strohm et al. 2019; see Section 2.5). We have also compiled SiO maser sources detected by the Korean VLBI Network single-dish telescopes (Kim et al. 2010; Cho & Kim 2012; Kim et al. 2013; Yoon et al. 2014; Cho et al. 2017). There are 2527 and 388 SiO maser sources in OAGB-IRAS and OAGB-WISE catalogs, respec-




Figure 8. MSX-2MASS 2CDs for all IRAS-AGB and WISE-AGB stars (see Tables 1 and 2) in our Galaxy compared with theoretical models (see Section 3). For OAGB models (silicate $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.05, 0.1, 0.5, 1, 3, 7, 15, 30$, and 40 from left to right. For CAGB models (AMC $T_c = 1000$ K): $\tau_{10} = 0.001, 0.01, 0.1, 0.5, 1, 2, 3$, and 5 from left to right. For each class, the number of objects is shown. The number in parentheses denotes the number of the plotted objects on the 2CD with good-quality observed colors.




Figure 9. Number density distributions of observed IR magnitudes and colors for OAGB and CAGB stars in AGB-IRAS and AGB-WISE catalogs. For each class, the number of objects is shown. The number in parentheses denotes the number of plotted objects with good-quality observed data. The averaged values and standard deviations of the observed data are also shown.




Figure 10. Number density distributions of observed IR colors for OAGB-IRAS sample stars and OH and SiO maser sources. The group of objects, for which OH (or SiO) maser emission was detected, is denoted by ‘OH (or SiO) maser’ in parentheses. And the group of objects, for which the maser emission was not detected or not observed, is denoted by ‘no OH (or SiO) maser’ in parentheses. The vertical brown lines indicate theoretical OAGB model colors for three dust shell optical depths τ_{10} . For each class, the number of objects is shown. The number in parentheses denotes the number of plotted objects with good-quality observed colors.




Figure 11. Number density distributions of observed IR colors for OAGB stars (OAGB-IRAS and OAGB-WISE) and SiO maser sources. The group of objects, for which SiO maser emission was detected, is denoted by ‘SiO maser’ in parentheses. And the group of objects, for which the maser emission was not detected or not observed, is denoted by ‘no SiO maser’ in parentheses. The vertical brown lines indicate theoretical OAGB model colors for three dust shell optical depths τ_{10} . For each class, the number of objects is shown. The number in parentheses denotes the number of plotted objects with good-quality observed colors.




Figure 12. Number density distributions of observed IR colors for CAGB stars (CAGB-IRAS and CAGB-WISE) and visual carbon stars (CI-GC and CW-GC). The vertical brown lines indicate theoretical CAGB model colors for two dust shell optical depths τ_{10} . For each class, the number of objects is shown. The number in parentheses denotes the number of plotted objects with good-quality observed colors.




Figure 13. Spatial distributions AGB stars (AGB-IRAS and AGB-WISE) in Galactic coordinate.




Figure 14. Number distribution of the Galactic longitude and latitude for OAGB and CAGB stars in AGB-IRAS and AGB-WISE catalogs. The number in parentheses denotes the number of objects.

tively (see Tables 1 and 2). In the OAGB-WISE catalog, all 388 SiO maser sources are in OW-ME and OW-ST.

OH/IR stars are generally considered to be more massive OAGB stars with thicker dust envelopes and higher mass-loss rates (Engels et al. 1983; Blommaert et al. 2018). Chen et al. (2001) presented a list of 1065 OH/IR stars in our Galaxy. The list has been corrected and updated (Suh & Kwon 2011; Kwon & Suh 2012). We also use the catalog of OH/IR stars presented by Engels & Bunzel (2015). There are 2079 OH/IR stars in OAGB-IRAS (1988 in OI-SH, 17 in OI-UR, 3 in OI-JB, 1 in OI-ME, 20 in OI-ST, 18 in OI-OG, 1 in OI-WU, and 31 in OI-AM; see Table 1), but there is only one known OH/IR star (OH 17.434-0.077) in OAGB-WISE (OW-ME) yet.

Figure 10 shows number distributions of IR colors ($K[2.2]-IR[12]$ and $IR[12]-IR[25]$). Most OH maser

sources are in the range of large dust optical depths (or mass-loss rates). However, most SiO maser sources are in the range of moderate dust optical depths (or mass-loss rates) for both colors.

We mark the theoretical model colors on Figures 10 and 11. For OAGB stars, the dust shell (silicate; $T_c=1000$ K) model colors for typical LMOA stars ($\tau_{10}=0.1$ and 1) and HMOA stars ($\tau_{10}=7$) are indicated (see Section 3).

Figure 11 shows number distributions of IR colors ($K[2.2]-W3[12]$ and $W2[4.6]-W3[12]$) for SiO maser sources and undetected (or unobserved) sources among all of the OAGB-IRAS and OAGB-WISE sample stars. Again, most SiO maser sources in OAGB-IRAS are in the range of moderate dust optical depths (or mass-loss rates) for both colors. Because the OAGB-WISE class lacks in objects with thick dust envelopes (or redder IR colors), SiO maser sources




Figure 15. Left panels show period-color relations for AGB-IRAS objects known as Miras. Right panels show period-color relations for 522 OAGB-IRAS objects with periods measured at NIR, MIR, and radio bands. See Section 7. Note that the period-color relation in the upper-right panel is used to find an expected period (EP) in Section 8.4.

look to be in the range of relatively large dust optical depths for the IR colors. Note that SiO maser observations have not been performed yet for a major portion of OAGB-WISE objects, which are OGLE3 Miras in the Galactic bulge (OW-OG; see Table 2).

5.3. Visual carbon stars

Unlike other subgroups of CAGB stars (in CI-SH, CI-UR, CI-OG, and CW-OG; see Tables 1-2), most objects in CI-GC and CW-GC (from Alksnis et al. 2001) are believed to be visual carbon stars in the AGB phase.

To find the differences of the IR colors of visual carbon stars from the colors of other groups that contain a mixture of infrared carbon stars and visual carbon stars, we compare the histograms in Figure 12.

We mark the theoretical model colors on Figure 12. For CAGB stars, the dust shell (AMC; $T_c=1000$ K) model colors for the thin dust shell ($\tau_{10}=0.1$) and thick dust shell ($\tau_{10}=1$) model colors are indicated (see Section 3).

The left panels of Figure 12 show number density distributions of observed IR colors ($IR[12]-IR[25]$, $IR[25]-IR[60]$, $K[2.2]-AK[9]$, and $K[2.2]-MA[8.3]$) for CAGB-IRAS objects. Compared with others, visual carbon stars (in CI-GC) show bluer $IR[12]-IR[25]$, $K[2.2]-AK[9]$, and $K[2.2]-MA[8.3]$ colors and but redder $IR[25]-IR[60]$ colors. This would be because visual carbon stars show exces-

sive flux at $60 \mu\text{m}$ due to the remnant of an earlier phase when they were OAGB stars (see Section 4.1).

The right panels of Figure 12 show number density distributions of observed IR colors ($K[2.2]-W3[12]$ and $W3[12]-W4[22]$) for CAGB-IRAS and CAGB-WISE objects. For $K[2.2]-W3[12]$, non CI-GC objects are in the wide ranges of large dust optical depths, whereas CI-GC objects are in narrow ranges. For CAGB-WISE objects, the difference gets smaller because most of the non CW-GC objects (in CW-OG) are likely to CAGB stars with thin dust shells.

For $W3[12]-W4[22]$, the number density distributions for CAGB-IRAS objects are similar to those for $IR[12]-IR[25]$. For CAGB-WISE objects, some CW-GC objects show redder colors. This would be because visual carbon stars with detached dust shell ($T_c < 500$ K; remnant of an earlier phase when the stars were OAGB stars) may show redder $W3[12]-W4[22]$ (see Section 4.2). Note that the theoretical model colors on Figure 12 are for hot dust shell ($T_c=1000$ K), which are not be applicable to detached dust shell models.

6. SPACIAL DISTRIBUTION OF AGB STARS

Figure 13 shows spacial distributions of AGB stars (AGB-IRAS and AGB-WISE) in Galactic coordinates. Figure 14 shows number distribution of the Galactic longitude and latitude for OAGB and CAGB stars in AGB-IRAS and AGB-WISE catalogs. In the bulge component, there are more




Figure 16. Period-color relations for AGB-IRAS and AGB-WISE objects known as Miras (AAVSO). For each class, the number of objects is shown. The number in parentheses denotes the number of the plotted objects with good-quality observed data. See Section 7.




Figure 17. Number density distributions of magnitudes of the objects, for which the periods are obtained from the WISE light curves. P-W1 (or P-W2) denotes objects whose periods are obtained from the WISE W1 (or W2) light curves. G-W1 (or G-W2) denotes objects whose good-quality periods are obtained from the WISE W1 (or W2) light curves. See Section 8 and Table 8.

OAGB stars than CAGB stars. The lack of OAGB-WISE objects at the Galactic center in the lower-left panel looks to be due to a selection effect of the sample stars.

The histograms for different Galactic latitudes are similar for both OAGB and CAGB stars. All AGB stars are concentrated toward the Galactic disk.

We find that OAGB stars are more concentrated toward the Galactic center and the number decreases with the Galactic longitude, while CAGB stars are distributed more uniformly from the center to large Galactic longitudes. Ishihara et al. (2011) also found that OAGB stars are concentrated toward the Galactic center and that the density decreases with Galactocentric distance, whereas CAGB stars show a relatively uniform distribution within about 8 kpc of Sun.

7. PERIOD-COLOR RELATIONS FOR KNOWN MIRA VARIABLES

It is generally believed that more evolved (or more massive) AGB stars would show the longer pulsation periods, larger pulsation amplitudes, higher mass-loss rates, thicker dust envelopes, and redder IR colors (e.g., De Beck et al. 2010; Suh & Kwon 2013b). Studying IR properties of all types of LPVs in Magellanic clouds, Suh (2020) showed that only Mira variables, among all types of LPVs, show a clear period-color relation (PCR): Miras with longer pulsation periods generally show redder IR colors. This would be because Mira variables are usually oscillating in the fundamental mode and occupy a single sequence in the period-luminosity diagram (Soszyński et al. 2013b; see Section 2.5).

Because most of the known pulsation periods in AAVSO were obtained in optical observations and the longest wavelength band used by OGLE3 observations was the I band (0.8 μm), most Mira variables whose pulsation periods listed in AAVSO are early phase AGB stars with thin dust envelopes.

The left panels of Figure 15 show IR[12]–IR[25] and K[2.2]–IR[12] colors versus pulsation periods for Miras in the AGB-IRAS catalog (see Table 1). For both IR colors, CAGB stars show larger coefficients of determination (R^2), which mean higher strength of the relationship. We find that the Mira variables show fairly strong PCRs.

Jiménez-Esteban et al. (2021) investigated variability properties of the Arecibo sample of OH/IR stars and presented periods for 348 Arecibo sources obtained from observations at NIR bands. All of those sources are in OI-SH except for one object, IRAS 18551+0323, which is a CAGB star (in CI-SH with IRAS LRS type C; this object could be a composite object).

The pulsation periods measured at MIR or radio bands for more evolved or massive AGB stars with thick dust envelopes are available only for a small number of AGB stars. We have compiled pulsation periods of 522 OAGB stars (495 OH/IR stars; 214 AAVSO Miras) measured at NIR, MIR, or radio bands presented by Chen et al. (2001), Kwon & Suh (2010b), Urago et al. (2020), and Jiménez-Esteban et al. (2021). The right panels of Figure 15 show PCRs using IR[12]–IR[25] and K[2.2]–W3[12] colors versus the pulsation periods measured at NIR, MIR, or radio bands for the 522 OAGB stars.

Figure 16 shows PCRs using K[2.2]–W3[12], W1[3.4]–W2[4.6], W2[4.6]–W3[12] and W3[12]–W4[22] colors for Miras in AAVSO. The left and right panels show the PCRs for Miras in AGB-IRAS and AGB-WISE, respectively.

Though there are large scatters, we find that the PCRs for Miras in the sample stars show a noticeable trend: Miras with longer pulsation periods generally show redder IR colors.

8. FINDING IR VARIATIONS OF AGB STARS FROM WISE DATA

Various observational data obtained in the last 50 yr are available for studying variability of AGB stars. There are large amounts of photometric data at visual and NIR bands but the data at MIR bands are available only for a limited number of objects. For AGB stars with thick dust envelopes, the variability can be more properly investigated from the observations at MIR bands (e.g., Engels et al. 1983; Kwon & Suh 2010b).

To study variability of AGB stars at W1[3.4] and W2[4.6] bands during the last 12 yr, we use the ALLWISE multiepoch photometry table obtained in 2009–2010 and the NEOWISE-R data (2021 data release) which give us the photometry data for 14 epochs, two in every year between 2014 and 2020.

We try to find Mira-like variations from the WISE light curves of the sample stars using a simple sinusoidal light curve model with periods longer than 50 days (the shortest period of OGLE3 bulge Miras is 78.31 days). In this work, we use the Lomb-Scargle periodogram which is a commonly used statistical algorithm for detecting and characterizing periodic signals in unevenly spaced observations (e.g., Zechmeister & Kürster 2009; VanderPlas 2018). The Lomb-Scargle Periodograms are computed using the implementations in AstroPy (<https://docs.astropy.org/en/stable/timeseries/lombscargle.html#>). We use the AstroPy computing option of autoperiod using the 'chi2' method, which utilizes the fact that the Lomb-Scargle periodogram at each frequency is equivalent to the least-squares fit of a sinusoid to the data. The advantage of the 'chi2' method is that it allows extensions of the periodogram to multiple Fourier terms.

For each object in the sample, we have generated the light curves using the WISE data and produced the Lomb-Scargle periodograms. But the WISE data for a major part of the sample of AGB stars (mostly bright stars) are either saturated or show scatters too large to provide meaningful variation parameters. Therefore, we need to choose the objects with good-quality variation parameters for which the deviations of the observed points from the derived sinusoidal model light curves are smaller. To find objects with good quality variation parameters at W1[3.4] and W2[4.6] bands (G-W1 and G-W2 objects), we choose the objects with more than 100 observed points and the coefficients of determination (R^2) to fit the sinusoidal model to the observations is larger than 0.6, for which the Lomb-Scargle power is also stronger. Table 8 summarizes the results for the sample stars.

Figure 17 compares the number density distributions of mean magnitudes for all objects whose WISE light curves are

Table 8. Numbers of objects for which good-quality variation parameters are obtained only from the WISE light curves

Subgroup ¹	Number	G-W1 ²	GM-W1 ³		GN-W1 ⁶	G-W2 ²	GM-W2 ³		GN-W2 ⁶
			GM-W1A ⁴	GM-W1B ⁵			GM-W2A ⁴	GM-W2B ⁵	
OI-SH	3826	392 (99)	16 (2)	6	370 (97)	414 (55)	28 (2)	14	372 (53)
OI-UR	37	13 (3)	2 (2)	1	10 (1)	10 (3)	3 (2)	0	7 (1)
OI-JB	5	5 (5)	0	0	5 (5)	4 (1)	0	0	4 (1)
OI-ME	127	7 (3)	0	0	7 (3)	12 (3)	0	0	12 (3)
OI-ST	673	31 (3)	1	0	30 (3)	49 (2)	5	0	44 (2)
OI-OG	1057	76 (12)	37 (7)	39 (5)	0	211 (47)	105 (24)	106 (23)	0
OI-WU	19	0	0	0	0	0	0	0	0
OI-AM	163	0	0	0	0	3 (0)	1	2	0
OI-all	5908	524 (125)	55 (11)	47 (5)	422 (109)	703 (111)	140 (28)	124 (23)	439 (60)
CI-SH	1165	91 (29)	4 (2)	2	85 (27)	50 (6)	1 (1)	2	47 (5)
CI-UR	5	2	0	0	2	0	0	0	0
CI-GC	2417	27 (2)	1	3 (1)	23 (1)	17 (3)	0	1 (1)	16 (2)
CI-OG	9	0	0	0	0	1	1	0	0
CI-all	3596	120 (31)	5 (2)	5 (1)	110 (28)	68 (9)	2 (1)	3 (1)	63 (7)
OW-ME	157	7 (1)	0	0	7 (1)	8 (3)	0	0	8 (3)
OW-ST	231	25 (5)	0	0	25 (5)	19 (3)	0	0	19 (3)
OW-OG	4913	1485 (370)	761 (212)	722 (158)	0	2241 (774)	1100 (417)	1139 (357)	0
OW-all	5301	1517 (376)	761 (212)	722 (158)	34 ⁷ (6)	2268 (780)	1100 (417)	1139 (357)	29 ⁷ (6)
CW-GC	3417	76 (13)	7 (2)	5 (3)	66 (8)	119 (24)	6 (3)	9 (4)	104 (17)
CW-OG	159	63 (20)	30 (8)	31 (12)	0	56 (19)	28 (10)	28 (9)	0
CW-all	3576	139 (33)	37 (10)	36 (15)	66 (8)	175 (43)	34 (13)	37 (13)	104 (17)

¹ See Tables 1 and 2. ² objects with more than 100 observed points for which good-quality ($R^2 > 0.6$) variation parameters (periods and amplitudes) at the WISE W1[3.4] or W2[4.6] band were obtained; the number in parentheses denotes the number of objects with excellent ($R^2 > 0.8$) quality (See Sections 8). ³ G-W1 or G-W2 objects that are known as Mira variables with periods from AAVSO. ⁴ the first peak of the Lomb-Scargle power is the nearest from the known Mira period. ⁵ the second (or up to fourth) peak is the nearest from the known Mira period. ⁶ G-W1 or G-W2 objects with unknown periods or those objects known as non-Mira variables with periods from AAVSO. ⁷ two of them are SRA variables in AAVSO (OGLE3 Miras).

available (P-W1 and P-W2 objects) and those for the objects with good-quality parameters (G-W1 and G-W2). Generally, the objects with good-quality variation parameters are less bright.

Using the WISE data, we have obtained good-quality variation parameters for 3710 objects (G-W1 or G-W2; 885 OAGB-IRAS, 141 CAGB-IRAS, 2468 OAGB-WISE, and 216 CAGB-WISE objects) in the catalog (see Table 8).

There are 2810 objects known as Miras with periods from AAVSO (GM-W1 or GM-W2; 284 OAGB-IRAS, 13 CAGB-IRAS, 2429 OAGB-WISE, and 84 CAGB-WISE objects). For about half of the objects (GM-W1A or GM-W2A objects), the obtained primary periods from the WISE data are similar to the period in AAVSO. For another half (GM-W1B or GM-W2B objects), the obtained primary periods from WISE data are different from the periods in AAVSO (see Section 8.2).

And there are 656 objects with unknown periods (GN-W1 or GN-W2; 441 OAGB-IRAS, 97 CAGB-IRAS, 37 OAGB-WISE, and 81 CAGB-WISE objects) and 244 objects known

as non-Mira variables with periods from AAVSO (GN-W1 or GN-W2; 160 OAGB-IRAS, 31 CAGB-IRAS, 2 OAGB-WISE, and 51 CAGB-WISE objects).

8.1. WISE light curves for known Mira Variables

Kwon & Suh (2010a) and Kwon & Suh (2010b) analyzed L and M band photometric data for 12 OH/IR stars acquired between the 1970s and 2004 and found their periods and amplitudes. Because L [3.4-3.6] and M [4.8-4.9] band wavelengths are near WISE W1[3.4] and W2[4.6] band wavelengths, we have analyzed the combined light curves. But the WISE data for most of these bright AGB stars are either saturated or show too large scatters. We could plot a meaningful combined light curve for only one source (OH 26.5+0.6).

Figure 18 shows the light curve for the OH/IR star OH 26.5+0.6 (in OI-SH) using the data at the L [3.4-3.6] band acquired in 1974-2003 and the WISE W1[3.4] band data acquired in the last 12 yr.

Figure 19 shows the OGLE3 and WISE light curves for IRAS 18007-3012 (in OI-OG). The light curves at I (0.8 μ m) and W2[4.6] bands show similar characteristics, though the




Figure 18. The combined light curve and Lomb-Scargle periodogram for an OH/IR star OH 26.5+0.6 (in OI-SH) using the L [3.4-3.6] band data acquired in 1974-2003 and the WISE W1[3.4] band data acquired in the last 12 yr. In the Lomb-Scargle periodogram, the dashed brown horizontal line indicates the periodogram level corresponding to a maximum peak false alarm probability of 1 %. Refer to [Kwon & Suh \(2010a\)](#) for details of the L band data. See Section 8.1.




Figure 19. Light curves and Lomb-Scargle periodograms for an OAGB-IRAS (OI-OG) object IRAS 18007-3012 (OGLE-BLG-LPV-196468) using the OGLE3 (I band) and WISE ($W2$ band) data. [Soszyński et al. \(2013a\)](#) obtained a period of 188.38 days. See Section 8.1.




Figure 20. WISE light curves and Lomb-Scargle periodograms for six OAGB-IRAS objects known as Miras with periods. In the Lomb-Scargle periodogram, the red X and black cross marks indicate the primary and selected peaks, respectively and the red dashed brown horizontal line indicates the periodogram level corresponding to a maximum peak false alarm probability of 1 %. Upper two panels show OI-UR objects whose periods from [Urago et al. \(2020\)](#) are 268 and 394 days, respectively. Middle two panels show OI-SH and OI-OG objects (see Table 1). For the two objects (OI-OG objects) in lower panels, the second peak of the Lomb-Scargle power is selected for the period (GM-W2B objects). See Section 8.1.




Figure 21. WISE light curves and Lomb-Scargle periodograms for six OAGB-WISE (OW-OG) objects known as Miras with periods. The object name is denoted by the OAGB-WISE identifier (OW-N; see Table 11). In the Lomb-Scargle periodogram, the red X and black cross marks indicate the primary and selected peaks, respectively and the red dashed brown horizontal line indicates the periodogram level corresponding to a maximum peak false alarm probability of 1 %. For three objects in lower panels, the second peak of the Lomb-Scargle power is selected for the period. See Section 8.1.




Figure 22. WISE light curves and Lomb-Scargle periodograms for three CAGB-IRAS objects (objects in upper panels: CI-SH; IRAS 08546+1732: CI-GC) and three CAGB-WISE objects (CW-2323: CW-GC; objects in lower panels: CW-OG) known as Miras with periods. For CAGB-WISE, the object name is denoted by the CAGB-WISE identifier (CW-N; see Table 12). In the Lomb-Scargle periodogram, the red X and black cross marks indicate the primary and selected peaks, respectively and the red dashed brown horizontal line indicates the periodogram level corresponding to a maximum peak false alarm probability of 1 %. Note that the second peak of the Lomb-Scargle power is selected for the periods of the three objects. See Section 8.1.




Figure 23. Properties of variation for OAGB-IRAS objects known as Miras with periods. The left panels show comparisons of the periods from AAVSO and the periods obtained from the WISE light curves. The green vertical (and horizontal) lines indicate the multiples of the interval of WISE observations (6 months). The right panels show the period-amplitude and period-color relations. See Section 8.2.




Figure 24. Properties of variation for OAGB-WISE objects known as Miras with periods. The left panels show comparisons of the periods from AAVSO and the periods obtained from the WISE light curves. The green vertical (and horizontal) lines indicate the multiples of the interval of WISE observations (6 months). The right panels shows the period-amplitude and period-color relations. See Section 8.2.




Figure 25. Properties of variation for CAGB-WISE objects known as Miras with periods. The left panels show comparisons of the periods from AAVSO and the periods obtained from the WISE light curves. The green vertical (and horizontal) lines indicate the multiples of the interval of WISE observations (6 months). The right panels shows the period-amplitude and period-color relations. See Section 8.2.




Figure 26. The period-amplitude relation for all Miras in the Galactic bulge obtained from OGLE3 *I* (0.8 μm) band observations (Soszyński et al. 2013a).

amplitude at the W2[4.6] band is smaller. Unlike the OGLE3 light curve, there are multiple peaks with similar power values in the Lomb-Scargle periodogram obtained from the WISE light curve because the WISE data were taken in a regular interval (every six months).

We have obtained pulsation periods from the WISE light curves for 2810 objects known as Miras with periods from AAVSO (GM-W1 or GM-W2; 284 OAGB-IRAS, 13 CAGB-IRAS, 2429 OAGB-WISE, and 84 CAGB-WISE objects; see Table 8).

Figure 20 shows the light curves for six OAGB-IRAS objects known as Miras. The periods obtained from WISE data are similar to the ones in AAVSO, but there are multiple peaks with similar Lomb-Scargle power values. For the two objects in lower panels (GM-W2B objects), though the period from the primary peak of the Lomb-Scargle power is different from the AAVSO period, the period from the second peak is very similar to the AAVSO period and produce a similar fit to the observations.

Figure 21 shows the light curves for six OAGB-WISE objects known as Miras. The obtained primary period from WISE data is similar to the ones in AAVSO for three objects (GM-W2A objects). For the other three objects (GM-W2B objects), the period from the second peak of Lomb-Scargle power is similar to the AAVSO period.

Figure 22 shows the light curves for for three CAGB-IRAS objects (objects in upper panels: CI-SH; IRAS 08546+1732; CI-GC) and three CAGB-WISE objects (CW-2323; CW-GC; objects in lower panels: CW-OG) known as Miras with periods. For three objects (GM-W2B objects), the period from the second peak of Lomb-Scargle power is similar to the AAVSO period.

For most Mira variables whose primary periods obtained from the WISE light curves are different from AAVSO periods (GM-W1B or GM-W2B objects), we can find a new (second or up to fourth) peak in the Lomb-Scargle power, for

which the new period is similar to the AAVSO period and produces a similar fit (a little worse fit with a smaller R^2 value) to the observations.

Because there are multiple peaks with similar Lomb-Scargle power values, it is not easy to obtain a precise period using only the WISE data (see Figures 20-22). The uncertainties in period and amplitude specified in Figures 18 and 19 are calculated by measuring the smooth imprecision in the selected peak of the Lomb-Scargle power. However, when there are multiple peaks with similar Lomb-Scargle power values, uncertainties expressed in this way cannot be meaningful because the uncertainties of periods can be more affected by the false peaks (VanderPlas 2018).

8.2. Comparison between AAVSO periods and new periods from WISE data for known Miras

If we compare the WISE periods with the AAVSO periods for the objects known as Miras with known periods from AAVSO (GM-W1 or GM-W2 objects; see Table 8), we may check the reliability of the the WISE periods. For AGB-WISE objects, most of AAVSO periods are from OGLE3 *I* (0.8 μm) band observations (Soszyński et al. 2013a).

Figures 23-25 show the relations between AAVSO periods and the periods obtained from the WISE light curves for OAGB-IRAS, OAGB-WISE, and CAGB-WISE objects (known as Miras with known periods from AAVSO), respectively. The figures also show the histograms of the periods obtained from the WISE light curves, period-amplitude relations, and period-color relations. For CAGB-IRAS objects, the numbers of sample stars (GM-W1 or GM-W2; see Table 8) are too small to make proper plots.

The upper-left panels of Figures 23-25 compare the periods from AAVSO (most of them are from OGLE3 *I* band observations) and the periods obtained from the WISE light curves for OAGB-IRAS, OAGB-WISE, and CAGB-WISE objects, respectively. For about a half of the objects (GM-W1A or GM-W2A), the obtained primary periods from the WISE data are similar to the periods in AAVSO. For another half (GM-W1B or GM-W2B), the obtained primary periods from WISE data are different from the periods in AAVSO. The deviations look to occur more severely when the AAVSO or WISE periods are similar to the interval of the WISE observations (6 months).

These deviations could be due to the characteristic of the Lomb-Scargle periodogram with similar multiple peaks, which could be due to the regularity of the WISE observations (6 months). VanderPlas (2018) compared the true period and peak Lomb-Scargle period for 1000 simulated periodic light curves and found that the Lomb-Scargle peak does not always coincide with the true period, and there is noticeable structure among these failures, which is similar to the ones in upper-left panels of Figures 23-25. The noticeable structure is clearer for OAGB-WISE objects because the sample number is much larger than other classes.

But for most GM-W1B or GM-W2B (see Table 8) objects, we can find a new (second or up to fourth) peak in the Lomb-Scargle power values, for which the new period is similar to

the AAVSO period. This would be because the AAVSO periods can be regarded as true periods and the multiple peaks in the Lomb-Scargle power values are very similar (see Figures 20-22). When we select the new peak that is similar to the AAVSO period for the GM-W1B or GM-W2B objects, the periods obtained from the WISE light curves are very similar to the AAVSO periods for most objects.

The lower-left panels of Figures 23-25 show the histograms of the periods obtained from the WISE light curves. GM-W1A or GM-W2A objects show a roughly single peak whereas GM-W1B or GM-W2B objects show multiple peaks. When we select new peaks for GM-W1B or GM-W2B objects, the histogram shows a roughly single peak just like GM-W1A or GM-W2A objects. Again, this effect is clearer for OAGB-WISE objects because the sample number is much larger than other classes.

On the whole, the periods from the WISE data and AAVSO show good correlations for all of the sample stars known as Miras (see upper-left panels of Figures 23-25).

Upper-right panels of Figures 23-25 show the period-amplitude relations. Both the relations for the objects using primary peaks (GM-W1A or GM-W2A) and for the objects using new peaks (GM-W1B or GM-W2B) look similar to the one for Miras in OGLE3 bulge (see Figure 26), which show larger amplitudes. But for some objects, the amplitudes show large deviation from the general trend, especially when obtained periods are similar to the multiples of the interval of WISE observations (6 months). Compared with OAGB-WISE objects, OAGB-IRAS objects show generally larger periods and amplitudes.

Lower-right panels of Figures 23-25 show the PCRs for OAGB-IRAS, OAGB-WISE, and CAGB-WISE objects using the $K[2.2]-W3[12]$ color. The objects show the similar PCRs to those for known Mira variables using AAVSO periods (see Figure 16).

For most objects that are known as Miras with periods in the catalog, we find that the new periods from the WISE W1[3.4] and W2[4.6] light curves are very similar to the periods in AAVSO. They also show very similar period-amplitude relations, though the amplitudes from OGLE3 are generally larger.

Though the reliability of the periods obtained from WISE observations has a weak point due to the regularity of observation (6 months), the new periods obtained from the WISE W1[3.4] and W2[4.6] light curves, whether they are from primary peaks of Lomb-Scargle power or not, could be more reliable than AAVSO periods for some objects depending on the quality of the model fit.

8.3. Candidate objects for new Mira variables

We have obtained new pulsation periods from the WISE light curves for 656 objects with unknown periods and 244 objects known as non-Mira variables with periods from AAVSO (GN-W1 or GN-W2 objects; see Table 8). The 656 objects with unknown periods consist of 441 OAGB-IRAS, 97 CAGB-IRAS, 37 OAGB-WISE, and 81 CAGB-WISE objects. The 244 objects known as non-Mira variables with pe-

riods from AAVSO consist of 160 OAGB-IRAS, 31 CAGB-IRAS, 2 OAGB-WISE, and 51 CAGB-WISE objects. These objects can be candidates for new Mira variables.

Though it is not be easy to obtain precise periods from the WISE light curves because the periodograms show multiple peaks with similar Lomb-Scargle power values, the derived variation parameters would be useful if there are enough observed points that fit the model well (stronger Lomb-Scargle power values or larger coefficients of determination for the model fit).

Figure 27 shows the WISE light curves and periodograms for the six OAGB-IRAS objects (OH/IR stars except for IRAS 16254-48580) with unknown periods, which can be candidate objects for new Miras. IRAS 16254-48580 (in OI-ST), which is a SiO maser source without any known characteristics, shows a Mira-like regular pulsation period (251 or 671 days). Though IRAS 17436-2807 (in OI-SH) was suspected to be a HII region (Walsh et al. 1998), the object is likely to be a typical OH/IR star because it shows a Mira-like regular pulsation period (613 days) obtained from the WISE light curves.

Figure 28 shows the WISE light curves and periodograms for the four CAGB-IRAS (CI-SH) and two CAGB-WISE (CW-GC) objects with unknown periods, which can be candidate objects for new Miras. Note that IRAS 08276-5125 (CI-SH) was suspected to be a SRV with unknown period (in AAVSO).

We have found new periods from the WISE light curves at W2[4.6] for 104 CW-GC objects (GN-W2; see Table 8) from which 33 objects were known to be SRVs with periods in AAVSO.

8.4. Properties of the candidate objects for new Miras

As we discussed in Section 8.2, the obtained primary period from the WISE light curve are different from the true period (AAVSO period) for about a half of the known Mira variables. Likewise, when we use only primary periods, the candidate objects for new Miras do not show the period-amplitude or period-color relations typical for Miras variables (see the upper-left panels of Figures 29-30).

Generally, Mira variables with longer pulsation periods show redder IR colors (see Figures 15-16 and 23-25). Though the PCRs show large scatter, we may roughly estimate the expected periods from the IR colors for the new Mira candidates. We use the relation between IR color ($K[2.2]-W3[12]$) and the period of Miras to estimate the expected period (EP).

We tried to use the PCR from the upper-right panel of Figure 15 for OAGB stars with known periods from radio or IR observations to obtain EP from $K[2.2]-W3[12]$. However, the PCR from the fit line produced too large EPs. After some trials with different slopes and intercepts, we find new PCRs (the blue and red lines in the panel) that produce a better relation between EPs and WISE periods for OAGB-IRAS objects (see the upper-left panel of Figure 29) and CAGB-IRAS objects (see the upper-left panel of Figure 30). Some discrep-




Figure 27. WISE light curves and Lomb-Scargle periodograms for OAGB-IRAS objects with unknown periods (IRAS 16254-4858 in OI-ST, IRAS 17128-3528 in OI-JB, and others in OI-SH). In the Lomb-Scargle periodogram, the red X and black cross marks indicate the primary and selected peaks, respectively and the red dashed brown horizontal line indicates the periodogram level corresponding to a maximum peak false alarm probability of 1 %. See Section 8.3. For four objects, $P(EP)$ (the expected period from the IR color $K[2.2]-W3[12]$; see Section 8.4) is also shown. For two objects, the second peak of the Lomb-Scargle power is selected for the period because it is more similar to EP.




Figure 28. WISE light curves and Lomb-Scargle periodograms for four CAGB-IRAS (CI-SH) and two CAGB-WISE (CW-GC) objects with unknown periods. For CAGB-WISE, the object name is denoted by the CAGB-WISE identifier (CW-N; see Table 12). In the Lomb-Scargle periodogram, the red X and black cross marks indicate the primary and selected peaks, respectively and the red dashed brown horizontal line indicates the periodogram level corresponding to a maximum peak false alarm probability of 1 %. See Section 8.3. For three objects, $P(EP)$ (the expected period from the IR color $K[2.2]-W3[12]$; see Section 8.4) is also shown.




Figure 29. For OAGB-IRAS objects with unknown periods or those objects known as non-Mira variables with periods, the upper-left panel compares the expected periods (EPs) obtained from the IR color ($\text{K}[2.2] - \text{W3}[12]$) with the periods obtained from the WISE light curves. The upper-right panel show the period-amplitude relation. The green vertical lines indicate the multiples of the interval of WISE observations (6 months). The lower panels show the relations between the WISE periods and IR colors ($\text{K}[2.2] - \text{W3}[12]$ and $\text{IR}[12] - \text{IR}[25]$). See Section 8.4.




Figure 30. For CAGB-IRAS objects with unknown periods or those objects known as non-Mira variables with periods, the upper-left panel compares the expected periods (EPs) obtained from the IR color ($\text{K}[2.2] - \text{W3}[12]$) with the periods obtained from the WISE light curves. The upper-right panel show the period-amplitude relation. The green vertical lines indicate the multiples of the interval of WISE observations (6 months). The lower panels show the relations between the WISE periods and IR colors ($\text{K}[2.2] - \text{W3}[12]$ and $\text{IR}[12] - \text{IR}[25]$). See Section 8.4.




Figure 31. WISE light curves and Lomb-Scargle periodograms for six OAGB (in OI-SH) objects with unknown periods. In the Lomb-Scargle periodogram, the red X and black cross marks indicate the primary and selected peaks, respectively and the red dashed brown horizontal line indicates the periodogram level corresponding to a maximum peak false alarm probability of 1 %. $P(EP)$ is the expected period from the IR color $K[2.2]-W3[12]$ (see Section 8.4). For two objects, the second peak of the Lomb-Scargle power is selected for the period because it is more similar to EP.




Figure 32. WISE light curves and Lomb-Scargle periodograms for six CAGB (in CI-SH) objects with unknown periods. In the Lomb-Scargle periodogram, the red X and black cross marks indicate the primary and selected peaks, respectively and the red dashed brown horizontal line indicates the periodogram level corresponding to a maximum peak false alarm probability of 1 %. $P(EP)$ is the expected period from the IR color $K[2.2]-W3[12]$ (see Section 8.4). For three objects, the second peak of the Lomb-Scargle power is selected for the period because it is more similar to EP. Note that the primary peaks at W1 and W2 bands are different for IRAS 19315+1807 (see lower two panels).

ancies were not avoidable because EP cannot be regarded as the true period for all objects.

We use the following PCRs: $K[2.2]-W3[12] = 6.87 * \log_{10}(\text{EP}) - 9.95$ (for OAGB-IRAS) and $K[2.2]-W3[12] = 9.54 * \log_{10}(\text{EP}) - 17.9$ (for CAGB-IRAS); see the upper-right panel of Figure 15. For a major portion of the best quality periodograms of OAGB-IRAS and CAGB-IRAS objects (see Figures 31 and 32 for examples), the WISE periods are similar to EPs when we use these PCRs.

From the GW1-N and GW2-N objects in AGB-IRAS (Table 8), we select the objects with good-quality $K[2.2]-W3[12]$ colors (OAGB-IRAS: 362 from 422 GW1-N objects and 380 from 439 GW2-N objects; CAGB-IRAS: 92 from 110 GW1-N objects and 50 from 63 GW2-N objects).

Figures 29 and 30 show the relations between EPs and the WISE periods, period-amplitude relations, and PCRs for OAGB-IRAS and CAGB-IRAS objects (with unknown periods or known as non-Mira variables with periods from AAVSO; GN-W1 and GN-W2 objects in Table 8), respectively.

The upper-left panels in Figures 29 and 30 compare EPs obtained from the IR colors ($K[2.2]-W3[12]$) and the periods obtained from the WISE light curves. There are objects whose primary WISE period is similar to EP (GN-W1A or GN-W2A) or objects whose new secondary (or up to fourth) period is similar to EP (GN-W1B or GN-W2B). For the GW1-NB or GW2-NB objects, the new WISE periods and EPs show a better correlation when we select the new (secondary or up to fourth) peak of the Lomb-Scargle power that is similar to EP. But the new correlation is still not as good as the one that uses the AAVSO period for the target period (see Figures 23-25) and shows larger deviations. There would be two possible reasons for larger deviations: the target period (EP from $K[2.2]-W3[12]$) cannot be the true period for all objects and some objects could not be actual Mira variables. Though this rough correlation is not as strong as the one between AAVSO and WISE periods, this comparison would be useful to find general characteristics of Mira variables for the candidate objects.

The upper-right panels in Figures 29 and 30 show the period-amplitude relations. And the lower panels in Figures 29 and 30 show PCRs using $K[2.2]-IR[12]$ and $IR[12]-IR[25]$ colors.

When we select the new peak, which is similar to EP, for the GN-W1B or GN-W2B objects, the candidate objects for new Miras show roughly similar period-amplitude relations and PCRs (see Figures 29 and 30) to those for known Mira variables (see Figures 15-16 and 23-25).

Figures 27 and 28 show WISE light curves and periodograms for AGB objects with unknown periods. For related objects with good-quality $K[2.2]-W3[12]$ colors, EP is also shown. We find that the periods obtained from WISE light curves show some deviations from EPs for these objects.

Figure 31 shows WISE light curves and periodograms for six OAGB-IRAS objects (in OI-SH) with unknown periods, which can be candidate objects for new Miras. For these

objects, we find that the periods obtained from WISE light curves are similar to EPs obtained from $K[2.2]-W3[12]$. Note that IRAS 17358-2711 can be identified as OGLE-BLG-ECL-054370 (angular distance: $33''$) with an E type variable (period: 1.08 days) from AAVSO, but they could be different objects.

Figure 32 shows WISE light curves and periodograms for six CAGB-IRAS objects (in CI-SH) with unknown periods, which can be candidate objects for new Miras. Again, we find that the periods obtained from WISE light curves are similar to EPs obtained from $K[2.2]-W3[12]$ for these objects.

9. THE CATALOG DATA

For OAGB-IRAS objects (see Table 1), Table 9 lists the OAGB-IRAS identifier (OI-N), subgroup name, IRAS PSC number, counterparts of AKARI PSC and BSC, best position (right ascension and declination J2000; RA and DEC), source of the best position (AKARI PSC, AKARI BSC, or IRAS PSC; see Section 2.2; BP), ALLWISE counterpart, spectral type (SP), and IRAS LRS type (LRS) from Kwok et al. (1997); variable type (A-Type) and period (A-Period) from AAVSO; and information about SiO and OH maser emission (see Section 5.2). For CAGB-IRAS objects (see Table 1), Table 10 lists the CAGB-IRAS identifier (CI-N) and the same information except for the information about SiO and OH maser emission.

For OAGB-WISE objects (see Table 2), Table 11 lists the OAGB-WISE identifier (OW-N), subgroup name, source number from the original reference (reco/star), ALLWISE source name, ALLWISE source position (W-RA, W-DEC), counterparts of 2MASS, AKARI PSC, MSX, and variable type (A-Type) and period (A-Period) from AAVSO. For CAGB-WISE objects (see Table 2), Table 12 lists the CAGB-WISE identifier (CW-N) and the same information.

For 3710 objects in the catalog (G-W1 or G-W2; see Table 8), the variation parameters obtained from WISE light curves at W1 and W2 bands are listed in Tables 13-16. For each obtained period (P), the amplitude (A) and coefficients of determination (R^2) to fit the sinusoidal model to the observations are also listed.

For OAGB-IRAS objects with variation parameters obtained from WISE light curves, Table 13 lists the OAGB-IRAS identifier (OI-N), variable type (A-Type) and period (A-Period) from AAVSO, expected period from the IR color $K[2.2]-W3[12]$ (EP; see Section 8.4), G-W1(P), G-W1(A), G-W1(R^2), GM-W1(P), GM-W1(A), GM-W1(R^2), GN-W1(P), GN-W1(A), GN-W1(R^2), G-W2(P), G-W2(A), G-W2(R^2), GM-W2(P), GM-W2(A), GM-W2(R^2), GN-W2(P), GN-W2(A), GN-W2(R^2), subgroup name, and IRAS PSC number. For CAGB-IRAS objects with variation parameters obtained from WISE light curves, Table 14 lists the CAGB-IRAS identifier (CI-N) and the same information.

For OAGB-WISE objects with variation parameters obtained from WISE light curves, Table 15 lists the OAGB-WISE identifier (OW-N), variable type (A-Type) and period (A-Period) from AAVSO, G-W1(P), G-W1(A), G-

W1(R^2), GM-W1(P), GM-W1(A), GM-W1(R^2), G-W2(P), G-W2(A), G-W2(R^2), GM-W2(P), GM-W2(A), and GM-W2(R^2). For CAGB-WISE objects with variation parameters obtained from WISE light curves, Table 16 lists the CAGB-WISE identifier (CW-N) and the same information.

Note that G-W1(P) or G-W2(P) listed in Tables 13-16 are primary periods. However, GM-W1(P) and GN-W1(P) (or GM-W2(P) and GN-W2(P)) are selected periods, which can be different from the primary periods (see Sections 8.2 and 8.4).

10. SUMMARY

We have presented a new catalog of 11,209 OAGB stars and 7172 CAGB stars in our Galaxy identifying more AGB stars in the bulge component and considering more visual carbon stars. For each object, we have cross-identified the IRAS, AKARI, MSX, WISE, 2MASS, and AAVSO counterparts.

We have presented the new catalog in two parts: one (AGB-IRAS) is based on the IRAS PSC for brighter or more isolated objects, the other one (AGB-WISE) is based on the ALLWISE source catalog for less bright objects or small objects in crowded regions.

We have performed radiative transfer model calculations for AGB stars using various parameters of central stars and spherically symmetric dust shells.

We have presented various IR 2CDs for the sample stars. We have compared the various sequences of theoretical dust shell models at increasing dust optical depth with the observations of AGB stars on the IR 2CDs. We find that the theoretical dust shell models can roughly explain the observations of AGB stars on the various IR 2CDs.

We have compared number distributions of observed IR magnitudes and colors for AGB stars in the AGB-IRAS and AGB-WISE catalogs. Most AGB-IRAS objects are brighter at MIR bands and they show redder IR colors than AGB-WISE objects. In general, AGB-IRAS objects look to be more evolved (or massive) stars with thicker dust envelopes than AGB-WISE objects.

We have investigated the IR properties of SiO and OH maser emission sources in the OAGB-IRAS and OAGB-WISE catalogs. Almost all known OH maser sources are in the OAGB-IRAS catalog. We have found that most OH maser sources are in the range of large dust optical depths (or mass-loss rates). On the other hand, most SiO maser sources in the OAGB-IRAS catalog are in the range of moderate dust optical depths (or mass-loss rates) for various IR colors. For a major portion of the OAGB stars in the OAGB-WISE catalog, OH or SiO maser observations have not been performed yet.

We have compared the IR properties of visual carbon stars from those of infrared carbon stars. Generally, visual carbon stars show bluer colors than infrared carbon stars because the dust shell optical depths for visual carbon stars are smaller. But some visual carbon stars show redder colors at MIR bands using longer wavelengths, which would be due

to detached circumstellar dust shells that are remnants of an earlier phase when the stars were OAGB stars.

We have investigated number distribution of the Galactic longitude and latitude for AGB stars in the AGB-IRAS and AGB-WISE catalogs. We have found that OAGB stars are more concentrated toward the Galactic center and the number decreases with the Galactic longitude, while CAGB stars are distributed more uniformly from the center to large Galactic longitudes. The histograms for different Galactic latitudes are similar for both OAGB and CAGB stars. All AGB stars are concentrated toward the Galactic disk.

For known Mira variables in the sample stars, we have investigated the period-color relations and found that objects with longer pulsation periods generally show redder colors.

We have investigated infrared variability of the sample stars using the WISE photometric data in the last 12 yr: the ALLWISE multiepoch data that were acquired between 2009 and 2010 and the NEOWISE-R 2021 data release that were acquired from 2013 until the end of 2020.

We have tried to find Mira-like variations from the WISE light curves of the sample stars using a simple sinusoidal light curve model. Using the WISE data at W1 and W2 bands, we have generated the light curves and computed the Lomb-Scargle periodograms for all of the sample stars and found good-quality variation parameters for 3710 objects in the catalog, for which periods were either known or unknown in previous works.

We have obtained pulsation periods from the WISE light curves for 2810 objects, which are known to be Miras with periods from AAVSO. For about a half of the objects, the obtained primary periods from the WISE data are very similar to AAVSO periods. For another half of objects whose primary periods from WISE data are different from AAVSO periods, we can find a new (second or up to fourth) peak in the Lomb-Scargle power values, for which the new period is similar to the AAVSO period. This would be because the AAVSO periods can be regarded as true periods and the multiple peaks in the Lomb-Scargle power values are similar, which would be due to the regularity of the WISE observations (6 months). On the whole, the periods from the WISE data and AAVSO showed very good correlations for all of the sample stars known as Miras.

We have obtained pulsation periods from the WISE light curves for 656 objects with unknown periods and 244 objects known as non-Mira variables with periods from AAVSO. We have found that a major portion of these objects could be candidate objects for new Mira variables because they show similar period-amplitude and period-color relations to those of known Mira variables.

If we perform new photometric observations at L or M band at different pulsation phases of AGB stars, the WISE data would be more useful to find precise periods.

The catalog data are presented in Section 9 (and Tables 9-16). The data will also be accessible through the author's webpage <http://web.chungbuk.ac.kr/~kwsuh/agb.htm>.

I thank the anonymous referee for helpful comments and constructive suggestions. This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT; Ministry of Science and ICT) (No. NRF-2017R1A2B4002328). This research has made use of the VizieR catalogue access tool, CDS, Strasbourg, France. This research has made use of the NASA/ IPAC Infrared Science Archive, which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

Software: For a major part of the computations and figures in this paper, we have used Python codes. We have used the Numpy ([van der Walt et al. 2011](#)), Pandas ([McKinney 2010](#)), AstroPy ([Astropy Collaboration et al. 2013](#)), and Matplotlib ([Hunter 2007](#)) packages.

REFERENCES

- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Alksnis, A., Balklavs, A., Dzervitis, U., et al. 2001, General Catalog of galactic Carbon stars, 3d Ed., *Balt. Astron.*, 10, 1
- Begemann, B., Dorschner, J., Henning, T., Mutschke, H., & Thamm, E. 1994, *ApJ*, 423, L71
- Beichman, C. A., Neugebauer, G., Habing H., Clegg, P. E., & Chester, T. C. 1988, *IRAS Catalogs and Atlases: Explanatory Supplement*, NASA RP-1190 (Washington: NASA)
- Blommaert, J. A. D. L., Groenewegen, M. A. T., Justtanont, K., & Decin, L. 2018, *MNRAS*, 479, 3545
- Cieza, L. A., Ruíz-Rodríguez, D., Perez, S., et al. 2018, *MNRAS*, 474, 4347
- Chan, S. J., & Kwok, S., 1990, *A&A*, 237, 354
- Chen, P. S., Szczerba, R., Kwok, S., & Volk, K. 2001, *A&A*, 368, 1006
- Chiar, J. E., & Tielens, A. G. G. M. 2006, *ApJ*, 637, 774
- Cho, S.-H., & Kim, J. 2012, *AJ*, 144, 129
- Cho, C.-Y., Cho, S.-H., Kim, S., et al. 2017, *ApJS*, 232, 13
- Cohen, M., Wheaton, W. A., & Megeath, S. T. 2003, *AJ*, 126, 1090
- Cordiner, M. A., Boogert, A. C. A., Charnley, S. B., et al. 2016, *ApJ*, 828, 51
- Cutri, R. M., Skrutskie, M. F., Van Dyk, S., et al. 2003, *The IRSA 2MASS All-Sky Point Source Catalog*, NASA/IPAC Infrared Science Archive
- De Beck, E., Decin, L., de Koter, A., et al. 2010, *A&A*, 523, A18
- Deguchi, S., Sakamoto, T., & Hasegawa T., 2012, *PASJ*, 64, 4
- Egan, M. P., Price, S. D., & Kraemer, K. E. 2003, *The Midcourse Space Experiment Point Source Catalog v2.3* (2003 October) Air Force Research Lab. Tech. Rep. AFRL-VS-TR-2003-1589 (Springfield, VA: NTIS)
- Engels, D., Kreysa, E., Schultz, G. V., & Sherwood, W. A. 1983, *A&A*, 124, 123
- Engels, D., & Bunzel, F. 2015, *A&A*, 582, A68
- Gordon, K. D., Cartledge, S., & Clayton, G. C. 2009, *ApJ*, 705, 1320
- Groenewegen, M. A. T., van den Hoek, L. B., & de Jong, T. 1995, *A&A*, 293, 381
- Henning, T., Begemann, B., Mutschke, H., & Dorschner, J. 1995, *A&AS*, 112, 143
- Hodapp, K. W., Chini, R., Watermann, R., & Lemke, R. 2012, *ApJ*, 744, 56
- Höfner, S., & Olofsson, H., 2018, *A&A Rev.*, 26, 1
- Hony, S., Waters, L. B. F. M., & Tielens, A. G. G. M. 2002, *A&A*, 390, 533
- Humphreys, R. M. et al. 2021, *AJ*, 161, 98
- Hunter, J. D. 2007, *CSE*, 9, 90
- Ishihara, D., Kaneda, H., Onaka, T., et al. 2011, *A&A*, 534, A79
- Ivezić, A., & Elitzur, M. 1997, *MNRAS*, 287, 799
- Jarrett, T. H., Cohen, M., Masci, F., et al. 2011, *ApJ*, 735, 112
- Jiménez-Esteban, F. M., & Engels D., 2015, *A&A*, 579, A76
- Jiménez-Esteban, F. M., Engels D., Aguado, D. S., & García-Lario, P. 2021, *MNRAS*, in press
- Kim, J., Cho, S.-H., Oh, C. S., & Byun, D.-Y. 2010, *ApJS*, 188, 209
- Kim, J., Cho, S.-H., & Kim, S. J. 2013, *AJ*, 145, 22
- Kwok, S., Volk, K., & Bidelman, W. P. 1997, *ApJS*, 112, 557
- Kwon, Y.-J., & Suh, K.-W. 2010a, *JKAS*, 43, 123
- Kwon, Y.-J., & Suh, K.-W. 2010b, *Journal of Astronomy and Space Sciences*, 27, 279
- Kwon, Y.-J., & Suh, K.-W. 2012, *JKAS*, 45, 139
- Kwon, Y.-J., & Suh, K.-W. 2014, *JKAS*, 47, 123
- Lewis, M. O., Pihlström, Y. M., Sjouwerman, L. O., & Quiroga-Núñez, L. H. 2020, *ApJ*, 901, 98
- Mainzer, A., Bauer, J., Cutri, R. M., et al. 2014, *ApJ*, 792, 30
- McKinney, W. 2010, in *Proc. 9th Python in Science Conf.*, Data Structures for Statistical Computing in Python, ed. S. van der Walt & J. Millman, 51
- Messineo, M., Habing, H. J., Sjouwerman, L. O., Omont, A., & Menten, K. M. 2018, *A&A*, 619, 35
- Murakami, H., Baba, H., Barthel, P., et al. 2007, *PASJ*, 59, 369
- Pégourié, B. 1988, *A&A*, 194, 335
- Samus, N., Kazarovets, E., Durlevich, O., Kireeva N., & Pastukhova, E., 2017, *ARep*, 61, 80S

- Sevenster, M. N. 2002, *AJ*, 123, 2788
- Siess, L. 2006, *A&A*, 448, 717
- Soszyński, I., Udalski, A., Szymański, M. K., et al. 2013a, *AcA*, 63, 21
- Soszyński, I., Wood, P. R., Udalski, A. 2013b, *ApJ*, 779, 167
- Stroh, M. C., Pihlström, Y. M., Sjouwerman, L. O., et al. 2019, *ApJS*, 244, 25
- Suh, K.-W. 1999, *MNRAS*, 304, 389
- Suh, K.-W. 2000, *MNRAS*, 315, 740
- Suh, K.-W. 2015, *ApJ*, 808, 165
- Suh, K.-W. 2016, *JKAS*, 49, 127
- Suh, K.-W. 2020, *ApJ*, 891, 43
- Suh, K.-W., & Kwon, Y.-J. 2011, *MNRAS*, 417, 3047
- Suh, K.-W., & Kwon, Y.-J. 2013, *JKAS*, 46, 235
- Suh, K.-W., & Hong, J. 2017, *JKAS*, 50, 131
- Th. Posch, F., Kerschbaum, H., Mutschke, J., et al. 2002, *A&A*, 393, L7
- Urago, R., Omodaka, T., Nagayama, T., et al. 2020, *ApJ*, 891, 50
- VanderPlas, J. T. 2018, *ApJS*, 236, 16
- van der Veen, W. E. C. J., & Habing, H. J. 1988, *A&A*, 194, 125
- van der Walt, S., Colbert, S. C., & Varoquaux, G. 2011, *CSE*, 13, 22
- Walsh, A. J., Burton, M. G., Hyland, A. R., & Robinson, G. 1998, *MNRAS*, 301, 640
- Watson, C., Henden, A. A., & Price, A. 2021, *Vizier*, B/vsx/vsx
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868
- Wu, Y. W., Matsunaga, N., Burns, R. A., & Zhang, B. 2017, *MNRAS*, 473, 3325
- Yoon, D.-H., Cho, S.-H., Kim, J., et al. 2014, *ApJS*, 211, 15
- Zapata, L. A., Garay, G., Palau, A., et al. 2019, *ApJ*, 872, 176
- Zechmeister, M., & Kürster, M. 2009, *A&A*, 496, 577

Table 9. OAGB-IRAS objects (16 columns; 5908 rows)⁰

OI-N ¹	Subgroup	IRAS PSC	AKARI PSC	AKARI BSC	RA ²	DEC ²	BP ³	ALLWISE	SP ⁴	LRS ⁵	A-Name ⁶	A-Type ⁷	A-Period ⁸
1	OI-SH	00007+5524	000321.44+554051	000321.04+554052	0.83958	55.68099	akari-p	J000321.45+554052.1	M7e:	E	Y Cas	M	417
2	OI-SH	00017+3949	000420.04+400635		1.08372	40.10991	akari-p	J000420.08+400635.9	M5.5e	F	SV And	M	313
3	OI-SH	00042+4248	000652.74+30502	000652.64+30501	1.71972	43.08394	akari-p	J000652.77+430502.2	M10	E	KU And	M	720
4	OI-SH	00050-2546	000736.22-252940	000736.9-252943	1.90107	-25.49451	akari-p	J000736.26-252939.9	M6e	E	SV Scl	M	411
5	OI-SH	00060-3929	000837.3-391304		2.15578	-39.21801	akari-p	J000837.35-391304.8	M5.5e:	F	V Scl	M	296.1
6	OI-SH	00075+5435	001009.2+545233		2.5386	54.87602	akari-p	J001009.14+545234.3	M6e	F	TT Cas	M	396
7	OI-SH	00127+5437	001524.8+545421		3.85362	54.90609	akari-p	J001524.85+545422.1		E	S Scl	M	367
8	OI-SH	00128-3219	001522.3-320243		3.84295	-32.04537	akari-p	J001522.28-320243.0	M6-e	F	NSV 15060	M	330
9	OI-SH	00138+6544	001636.5+660109	001636.2+660113	4.15224	66.01937	akari-p	J001636.49+660110.5		E		M	
10	OI-SH	00170+6542	001951.2+655929	001951.6+655930	4.9637	65.99148	akari-p	J001951.28+655930.5	(OH)	E		M	
11	OI-SH	00176+6931	002025.5+694756	002025.1+694759	5.10634	69.79903	akari-p	J002025.52+694757.2		F	NSVS 10020257+694759	M	425
12	OI-SH	00193+4033	002147.4+401715	002147.5+401713	5.44763	-40.28772	akari-p	J002147.41+401715.5	M9	E	BE Phe	M:	
13	OI-SH	00205+5530	002314.2+554732	002313.5+554735	5.80928	55.79243	akari-p	J002314.29+554733.1	M7.5e	F	T Cas	M	440
14	OI-SH	00207-6156	002307.7-614017		5.78216	-61.67143	akari-p	J002307.67-614016.9		F	S Tuc	M	242.4
15	OI-SH	00222+6952	002510.1+700851	002508.7+700901	6.29223	70.14756	akari-p	J002510.03+700851.9	M6	F	NQ Cep	M	454
16	OI-SH	00245-0652	002706.4-063616	002706.4-063610	6.77682	-6.60471	akari-p	J002706.42-063617.0	M7	E	UY Cet	SRB	440
17	OI-SH	00309-7956	003240.4-794021		8.16842	-79.67251	akari-p	J003240.38-794021.0			W Hyi	M	280
18	OI-SH	00336+6744	003636.5+680119	003637.2+680125	9.1524	68.02211	akari-p	J003636.61+680119.8		E	V0861 Cas	M	210
19	OI-SH	00340+6251	003659.4+630800	003659.3+630801	9.24777	63.13344	akari-p	J003659.43+630801.9	M6	E	TY Cas	M	645
20	OI-SH	00347+8004	003822.8+802125		9.59516	80.35698	akari-p	J003822.82+802124.9	M5e:	F	Y Cep	M	332.57
21	OI-SH	00381-8018	003939.1-800204		9.91314	-80.03467	akari-p	J003938.99-800204.5		E	X Hyi	M	304
22	OI-SH	00420+7533	004528.0+755022	004526.4+755022	11.36698	75.83947	akari-p	J004528.05+755022.1		E	NSVS 10045283+755024	M	599
23	OI-SH	00428+6854	004600.2+691053	004559.4+691055	11.50094	69.18145	akari-p	J004600.17+691053.8	M8:	E	V0524 Cas	M	490
24	OI-SH	00445+3224	004718.9+324108	004719.1+324108	11.82894	32.68564	akari-p	J004718.91+324108.1	S6/2-e	F	RW And	M	430
25	OI-SH	00453+5317	004810.0+533400		12.04174	53.56691	akari-p	J004809.99+533401.2	M8	F	V0414 Cas	M	195
26	OI-SH	00459+6749	004906.9+680547		12.27877	68.09647	akari-p	J004906.85+680547.3	M8	F	V0865 Cas	M	180
27	OI-SH	00479+4614	005043.3+463029	005043.7+463029	12.68047	46.5083	akari-p	J005043.29+463030.5		E	V0415 And	SRB	460
28	OI-SH	00498+4708	005242.7+472455	005242.5+472502	13.17824	47.41547	akari-p	J005242.78+472456.2	M6Se	E	RV Cas	M	331.68
29	OI-SH	00503+6445	005325.0+650156	005325.2+650201	13.35421	65.03224	akari-p	J005325.18+650155.9	M8/9	E	NSV 15193	M	507
30	OI-SH	00534+6031	005628.3+604710		14.11806	60.78614	akari-p	J005628.33+604709.3		E	V0867 Cas	M	412
5899	OI-AM	20499+4657	205136.6+470909	205136.5+470907	312.90288	47.15266	akari-p	J205136.63+470909.1		E	ZTF J205136.64+470909.3	M	457.3509671
5900	OI-AM	21112+3150	211318.2+320326		318.32608	32.05743	akari-p	J211318.23+320327.2		E	V0472 Cyg	M	297
5901	OI-AM	21444+4752	214619.9+480648	214621.9+480652	326.5831	48.1135	akari-p	J214619.97+480650.0	M7:	E	LP Cyg	M	419
5902	OI-AM	21543+5605	215600.8+561927		329.00362	56.32422	akari-p	J215600.90+561927.5	M9	E	V0720 Cep	M	305
5903	OI-AM	22000+5643	220147.7+565810	220147.9+565811	330.44911	56.96954	akari-p	J220147.74+565810.2	M7	E	YY Cep	M	526.08
5904	OI-AM	22049+4813	220651.6+482755	220652.3+482755	331.71531	48.46548	akari-p	J220651.71+482756.2	M7	E	AP Lac	M	524
5905	OI-AM	22124+7315	221321.8+732958	221321.5+732958	333.34096	73.4995	akari-p	J221321.64+732958.3	M7	E	NSVS J2213225+733019	M	429
5906	OI-AM	22450+5829	224704.2+584513		341.76759	58.75388	akari-p	J224704.07+584514.2	M7-8	E	Mis V1170	M	730
5907	OI-AM	23548-6539	235726.3-652304		359.35966	-65.38461	akari-p	J235726.38-652304.9	M5e	E	R Tuc	M	286.06
5908	OI-AM	23561+6037	235838.2+605342	235837.0+605336	359.65948	60.89513	akari-p	J235838.26+605342.5		E	EU Cas	M	447

⁰Only 14 columns are shown in this example table. In the data file, there are two more columns: OH and SiO maser detection (see Section 5.2). ¹The OAGB-IRAS identifier (see Table 1). ²The best position (right ascension and declination J2000). ³The source of the best position (AKARI PSC, AKARI BSC, or IRAS PSC; see Section 2.2). ⁴The spectral type from Kwok et al. (1997). ⁵The IRAS LRS type from Kwok et al. (1997). ⁶The object name from AAVSO. ⁷The variable type from AAVSO. ⁸The period from AAVSO.

Table 10. CAGB-IRAS objects (14 columns; 3596 rows)

Cl-N ¹	Subgroup	IRAS PSC	AKARI PSC	AKARI BSC	RA ²	DEC ²	Bp ³	ALLWISE	SP ⁴	LRS ⁵	A-Name ⁶	A-Type ⁷	A-Period ⁸
1	Cl-SH	000204+4316	000436+433304		1.15175	43.55133	akari-p	J000436.41+433304.6	C6.4	S	SU And	LC	
2	Cl-SH	00036+6947	0006142+700402	0006131+700409	1.55954	70.06723	akari-p	J000614.21+700402.1	C	C	OR Cep	M	355.32
3	Cl-SH	00050+7357	0007432+741411		1.93001	74.23647	akari-p	J000743.08+741411.5		C	NSVS J0007434+741409	L	476
4	Cl-SH	00084+1851	0010579+183422		2.74137	-18.57287	akari-p	J001057.93+183422.7	M4III	C	AC Cet	LB	
5	Cl-SH	00124+4425	0019540+444234	0019539+444230	4.97506	44.70953	akari-p	J001954.00+444233.8	C4.5J	C	VX And	SRA	375
6	Cl-SH	00210+6221	0023508+623811	0023507+623813	5.96206	62.6364	akari-p	J002350.87+623810.9	(CO)	U			
7	Cl-SH	00247+6922	0027411+693851	0027405+693857	6.92134	69.64762	akari-p	J002741.04+693851.0	(SiC)	C	V0668 Cas	M	650
8	Cl-SH	00248+3518	0027316+353514	0027315+353510	6.88205	35.58736	akari-p	J002731.69+353514.2	C5.4	C	AQ And	SRB	169
9	Cl-SH	00422+5310	0045070+532647	0045069+532652	11.27958	53.44656	akari-p	J004507.07+532647.6	(SiC)	C	V0720 Cas	M	431
10	Cl-SH	00523+6812	0055340+682854		13.89167	68.48174	akari-p	J005533.95+682855.2	C6+-4	C	V0880 Cas	SR	377
11	Cl-SH	00535+5923	0056330+593944	0056329+593951	14.13752	59.66235	akari-p	J005633.03+593943.9	C	C	V0721 Cas	M:	
12	Cl-SH	00596+6135	0102436+615142	0102448+615128	15.68168	61.8618	akari-p	J010243.58+615143.4	C4.3	S	HO Cas	LB	
13	Cl-SH	01022+6542	0105272+655900		16.36364	65.98334	akari-p	J010527.41+655859.7	M7	C	V0888 Cas	M	341
14	Cl-SH	01080+5327	0111035+534339	0111035+534338	17.76461	53.72774	akari-p	J011103.46+534339.9	C4.3e	C	HV Cas	M	527
15	Cl-SH	01105+6241	0113444+625736	0113450+625735	18.4352	62.96007	akari-p	J011344.54+625735.9	C6.3+e	C	NSV 438	M:	280
16	Cl-SH	01132+2530	0116050+254609	0116047+254604	19.02099	25.76919	akari-p	J011605.03+254609.7	C7.2	F	Z Psc	SRB	155.8
17	Cl-SH	01142+6306	0117335+632205	0117337+632211	19.38993	63.36806	akari-p	J011733.56+632205.1	(SiC)	C			
18	Cl-SH	01144+6658	0117515+671352	0117507+671357	19.4647	67.23136	akari-p	J011751.34+671353.2	(HCN)	U			
19	Cl-SH	01156+6237	0118539+625258	0118534+625306	19.7246	62.88294	akari-p	J011853.84+625259.4	C	C	V0829 Cas	M	1060
20	Cl-SH	01215+6430	0124580+644613	0124583+644609	21.24174	64.77053	akari-p	J012458.25+644613.2	C	C	NSVS J0118537+625300	L	279
21	Cl-SH	01246+3248	0126580+323235	0126581+323237	21.74177	-32.54324	akari-p	J012658.05+323236.0	C6.4	C	NSVS 1695145	M:	258
22	Cl-SH	01324+4907	0135288+492242		23.87005	49.37836	akari-p	J013528.78+492242.0	(SiC)	C	R Scl	SRB	370
23	Cl-SH	01377+6503	0136165+651839	0136159+651849	24.06899	65.3109	akari-p	J013616.59+651839.8		C	Dauban V257	VAR	
24	Cl-SH	01348+5543	0138056+555814		24.52342	55.97078	akari-p	J013805.62+555815.2		C	MGAB-V1402	M	300
25	Cl-SH	01411+7104	0145087+711915		26.28656	71.32098	akari-p	J014508.73+711916.1	C	F	PT Cas	M	730
26	Cl-SH	01443+6417	0147555+643255	0147553+643256	26.98151	64.54869	akari-p	J014755.72+643256.1	C	C	NSVS J0145096+711919	L	476
27	Cl-SH	01531+5900	0156381+591533	0156374+591528	29.15886	59.25942	akari-p	J015638.14+591533.9	C5.4e	C	X Cas	M	415
28	Cl-SH	01551+5458	0158294+551258	0158294+551303	29.62284	55.21626	akari-p	J015829.47+551258.8	Ce	F	V0437 Per	M	470
29	Cl-SH	01580+5803	0201281+581814		30.36713	58.30403	akari-p	J020128.16+581814.0	Ce	C	V0666 Cas	M	432.1
30	Cl-SH	02152+2822	0218061+283646	0218062+283645	34.52542	28.61286	akari-p	J021806.05+283645.3	(CO)	C	YY Tri	M	624
-	-	-	-	-	-	-	-	-	-	-	-	-	-
3587	Cl-GC	23573+5642	2359542+565846		359.97602	56.97963	akari-p	J235954.25+565846.6		-	V0533 Cas	SRA	305
3588	Cl-OG	17398+2146	1742510+214729	1742510+214728	265.71265	-21.79161	akari-p	J174250.99+214728.7		-	OGLE-BLG-LPV-020858	M	469.6
3589	Cl-OG	17468+3320	1750084+332118		267.53534	-33.35513	akari-p	J175008.37+332118.3		-	OGLE-BLG-LPV-059872	M	400.4
3590	Cl-OG	17490+3414	1752190+341509		268.07937	-34.25274	akari-p	J175219.08+341510.2		-	OGLE-BLG-LPV-085234	M	555.8
3591	Cl-OG	17514+3354			268.682037	-33.916191	iras	J175443.43+335507.5		-	OGLE-BLG-LPV-114752	M	448
3592	Cl-OG	17552+2814	1758219+281454		269.59127	-28.24834	akari-p	J175821.90+281452.7	(PN)	I	OGLE-BLG-LPV-149402	ZAND+M	416.2
3593	Cl-OG	17570+3056	1800175+305623		270.07295	-30.93992	akari-p	J180017.57+305624.1		-	OGLE-BLG-LPV-169921	M	449.7
3594	Cl-OG	18031+3229	1806270+322914		271.61265	-32.48727	akari-p	J180627.05+322913.8		-	OGLE-BLG-LPV-209822	M	516.9
3595	Cl-OG	18039+3411	1807152+341125		271.81348	-34.19048	akari-p	J180715.23+341125.1		-	OGLE-BLG-LPV-212597	M	378.8
3596	Cl-OG	18204+2133	1823282+213129		275.86784	-21.52491	akari-p	J182328.27+213130.6		-	OGLE-BLG-LPV-231922	M	513.8

¹The CAGB-IRAS identifier (see Table 1). ²The best position (right ascension and declination J2000). ³The source of the best position (AKARI PSC, AKARI BSC, or IRAS PSC; see Section 2.2). ⁴The spectral type from Kwok et al. (1997). ⁵The IRAS LRS type from Kwok et al. (1997). ⁶The object name from AAVSO. ⁷The variable type from AAVSO. ⁸The period from AAVSO.

Table 11. OAGB-WISE objects (12 columns; 5301 rows)

OW-N ¹	Subgroup	reco/star	ALLWISE	W-RA ²	W-DEC ²	2MASS	AKARI PSC	MSX	A-Name	A-Type	A-Period
1	OW-ME	1	J173140.97-320355.7	262.9207368	-32.0654792	17314097-3203559	1731409-320355	G355.7512+00.8670			
2	OW-ME	3	J173707.28-312131.2	264.2803606	-31.3586915	17370728-3121312	1737072-312132	G356.9721+00.2780			
3	OW-ME	4	J173729.36-311716.8	264.3723546	-31.2880029	17372934-3117166	1737292-311717	G357.0739+00.2497			
4	OW-ME	6	J173813.13-293941.4	264.5547352	-29.6615113	17381248-2939385	1738125-293938	G358.5318+00.9902			
5	OW-ME	7	J173817.04-294231.9	264.5710302	-29.7088827	17381707-2942324	1738170-294231	G358.5318+00.9902	GDS J1738177-294232	VAR	
6	OW-ME	9	J173832.49-312042.5	264.6353944	-31.3451427	17383250-3120427	1738324-312043	G357.1459+00.0294			
7	OW-ME	12	J174057.23-294531.4	265.238484	-29.7587303	17405722-2945314	1740571-294532	G358.7654+00.4335			
8	OW-ME	16	J174136.83-292930.9	265.4034951	-29.4919261	17413685-2929309		G359.0679+00.4525			
9	OW-ME	17	J174137.39-293205.7	265.4058107	-29.5349427	17413740-2932057		G359.0325+00.4283			
10	OW-ME	18	J174204.33-295846.1	265.5180671	-29.9794904	17420435-2958463	1742042-295846	G358.7064+00.1100			
11	OW-ME	19	J174206.85-281832.4	265.5285433	-28.3090071	17420685-2818323	1742068-281831	G000.1309+00.9838			
12	OW-ME	20	J174223.29-293935.3	265.5970442	-29.6598205	17422328-2939355	1742232-293936	G359.0141+00.2210			
13	OW-ME	21	J174232.89-294126.0	265.6370525	-29.6905646	17423291-2941251	1742325-294112	G359.0087+00.1783			
14	OW-ME	22	J174232.47-294110.6	265.6352975	-29.6862809	17423247-2941107	1742325-294112	G359.0087+00.1783			
15	OW-ME	24	J174309.81-292403.1	265.7909036	-29.4008852	17430981-2924033	1743097-292403	G359.3228+00.2140			
16	OW-ME	25	J174323.47-285350.1	265.8477983	-28.8972522	17432345-2853503	1743234-285349	G359.7779+00.4361			
17	OW-ME	26	J174325.25-294528.6	265.8552409	-29.7579569	17432525-2945285	1743251-294529	G359.0484+00.0212			
18	OW-ME	27	J174332.70-291539.2	265.8862747	-29.2608935	17433271-2915393	1743326-291539	G359.4858+00.2165			
19	OW-ME	28	J174333.12-295133.1	265.8880322	-29.8592045	17433312-2951331	1743330-295134	G358.9772+00.0984			
20	OW-ME	29	J174334.83-294029.2	265.8951536	-29.6747897	17433479-2940304	1743346-294031	G359.1369+00.0067			
21	OW-ME	30	J174335.12-292447.3	265.896337	-29.4131541	17433512-2924472	1743417-290118	G359.3607+00.1297			
22	OW-ME	32	J174341.73-290118.9	265.9239097	-29.0219439	17434176-2901190		G359.7068+00.3140			
23	OW-ME	33	J174349.53-290319.9	265.9564094	-29.0555301	17434954-2903194		G359.6931+00.2721			
24	OW-ME	34	J174349.98-290121.4	265.9582628	-29.0226238	17434997-2901213		G359.7219+00.2882			
25	OW-ME	35	J174351.10-290028.8	265.9629185	-29.0080118	17435106-2900292		G359.7368+00.2924			
26	OW-ME	36	J174354.00-284128.5	265.975037	-28.6912772	17435399-2841285		G000.0118+00.4492			
27	OW-ME	37	J174355.30-285649.9	265.9804464	-28.9471972	17435531-2856498		G359.7964+00.3112			
28	OW-ME	38	J174357.27-293146.6	265.988648	-29.5296325	17435729-2931465		G359.3035+00.0007			
29	OW-ME	39	J174358.02-293052.2	265.99179	-29.5145069	17435804-2930520		G359.3182+00.0059			
30	OW-ME	40	J174359.29-291951.5	265.997061	-29.3309929	17435926-2919517		G359.4766+00.0982			
5292	OW-OG	232164	J182407.05-212750.1	276.0294118	-21.4639172	18240706-2127500	1824072-212748	G010.7840+03.9197	OGLE-BLG-LPV-232164	M	305.1
5293	OW-OG	232216	J182415.74-214049.6	276.0656208	-21.6804648	18241570-2140498	1824159-214049	G010.6070+04.0495	OGLE-BLG-LPV-232216	M	325.8
5294	OW-OG	232256	J182421.96-212222.4	276.0915081	-21.372902	18242197-2122224	1824221-212222	G010.8916+03.9294	OGLE-BLG-LPV-232256	M	306.8
5295	OW-OG	232307	J182431.29-220041.6	276.1304043	-22.0115722	18243129-2200414	1824312-220041	G010.3404+04.2554	OGLE-BLG-LPV-232307	M	339.4
5296	OW-OG	232329	J182458.95-252439.4	276.2456492	-25.4109592	18245895-2524394		OGLE-BLG-LPV-232329	M	229.2	
5297	OW-OG	232339	J182508.99-253350.7	276.2874842	-25.5641061	18250899-2533505	1825089-253350	OGLE-BLG-LPV-232339	M	197.57	
5298	OW-OG	232340	J182510.62-255542.0	276.2942638	-25.9283545	18251062-2555419	1825106-255541	OGLE-BLG-LPV-232340	M	263.2	
5299	OW-OG	232350	J182547.17-251527.2	276.4465735	-25.2575831	18254717-2515270	1825471-251527	OGLE-BLG-LPV-232350	M	298.8	
5300	OW-OG	232364	J182640.02-251436.3	276.6667693	-25.2434178	18264001-2514363	1826400-251435	OGLE-BLG-LPV-232364	M	199.55	
5301	OW-OG	232397	J183121.84-242406.1	277.8410132	-24.4017158	18312183-2424060	1831218-242406	OGLE-BLG-LPV-232397	M	303.8	

¹The OAGB-WISE identifier (see Table 2). ²The ALLWISE source position (right ascension and declination J2000). See Section 9.

Table 12. CAGB-WISE objects (12 columns; 3576 rows)

CW-N ¹	Subgroup	recno/star	ALLWISE	W-RA ²	W-DEC ²	2MASS	AKARI PSC	MSX	A-Name	A-Type	A-Period
1	CW-GC	1	J000010.87+642554.6	0.0453247	64.4318379	00001089+6425546					
2	CW-GC	3	J000052.97+565807.3	0.2207225	56.9687165	00005295+5658072					
3	CW-GC	6	J000129.81+645507.8	0.3742141	64.9188449	00012982+6455077	0001298+645508	G117.6688+02.5591	ZTF J000129.84+645507.9	SR	253.5952999
4	CW-GC	8	J000203.10+625859.4	0.5129277	62.983188	00020311+6258593	0002032+625858	G117.3577+00.6467	NSV 26200		
5	CW-GC	9	J000207.99+024913.1	0.5333304	-2.8203301	00020802+0249121	0002080-024912				
6	CW-GC	11	J000223.06+560231.9	0.5961116	56.0422064	00022306+5602319					
7	CW-GC	13	J000300.86+303823.4	0.7536194	30.6398393	00030086+3038233			2MASS J00030086+3038233	L	
8	CW-GC	14	J000322.05+084752.4	0.8419097	8.7978933	null			NSV 15007	M:	259
9	CW-GC	15	J000341.96+594413.0	0.9248438	59.736952	00034196+5944130	0003419+594412				
10	CW-GC	19	J000524.99+011003.8	1.3541477	1.1677398	00052499+0110039					
11	CW-GC	21	J000544.48+653057.1	1.4353489	65.515888	00054449+6530569	0005444+653056	G118.2161+03.0626	ZTF J000544.52+653057.2	SR	121.6376906
12	CW-GC	22	J000545.12+592948.0	1.4380045	59.4966744	00054511+5929479	0005452+592948	G117.1563+02.8616	ZTF J000545.13+592948.0	SR	274.403834
13	CW-GC	30	J000911.97+650126.3	2.2999015	65.0239937	00091198+6501262	0009120+650124				
14	CW-GC	31	J001109.46+642925.5	2.7894529	64.4904191	00110946+6429257	0011093+642925	G118.6098+01.9573	NSVS 1612314	LPV	236.7535227
15	CW-GC	32	J001135.45+631830.8	2.897721	63.3085687	00113544+6318308	0011353+631830	G118.4749+00.7823	ZTF J001135.47+631830.7	SR	310.7410887
16	CW-GC	33	J001204.70+613249.6	3.0195937	61.5471352	00120469+6132496	0012046+613249	G118.1186+02.9019	Ms V1359	SR:	
17	CW-GC	36	J001310.30+593635.2	3.2929478	59.6097837	00131045+5936326					
18	CW-GC	37	J001316.63+630200.3	3.3193147	63.0334316	00131664+6302003	0013163+630159	G118.6208+00.4810	ZTF J001316.65+630200.2	SR	255.4633061
19	CW-GC	38	J001322.15+304045.1	3.3423166	-30.6792219	null					
20	CW-GC	41	J001346.40+681730.2	3.4433605	68.2917322	00134645+6817301	0013463+681730		V0966 Cep	SR:	130
21	CW-GC	43	J001431.09+605531.0	3.629571	60.9252917	00143110+6055310	0014310+605530	G118.4594+01.6258	ZTF J001431.10+605531.1	SR	152.8841528
22	CW-GC	44	J001444.57+574812.7	3.6857147	57.8035551	00144456+5748128					
23	CW-GC	46	J001501.62+600821.8	3.7567599	60.1394082	00150169+6008222					
24	CW-GC	47	J001553.66+605328.6	3.9736087	60.8912972	00155366+6053286		G118.4225+02.4038	NSV 107		
25	CW-GC	50	J001718.41+105203.1	4.3267474	10.8675438	00171842+1052032					
26	CW-GC	52	J001751.22+630406.1	4.4634252	63.0683664	00175122+6304060	0017511+630406	G119.1401+00.4449	ZTF J001751.23+630406.2	SR	103.9275228
27	CW-GC	53	J001814.60+594601.9	4.5608417	59.7672053	00181456+5946018					
28	CW-GC	57	J002006.57+615919.6	5.027391	61.9887875	00200656+6159196			ZTF J002006.58+615919.7	SR	209.421136
29	CW-GC	58	J002021.60+011206.7	5.0900124	1.201865	00202160+0112068					
30	CW-GC	60	J002134.74+591448.6	5.3947506	59.246851	00213483+5914483					
3567	CW-OG	224006	J181108.13+261611.8	272.7839129	-26.2699694	18110815+2616119					
3568	CW-OG	224160	J181113.68+251656.4	272.8070377	-25.28234	18111369+2516563		G005.1362+03.5662	OGLE-BLG-LPV-224006	M	222.7
3569	CW-OG	226511	J181316.97+253135.1	273.3207119	-25.5264335	18131697+2531351		G006.0256+03.1142	OGLE-BLG-LPV-224160	M	111.86
3570	CW-OG	226979	J181346.26+253415.6	273.4427696	-25.5710236	18134625+2534156	1813169+253136	G006.0216+03.6336	OGLE-BLG-LPV-226511	RCB:	
3571	CW-OG	227901	J181447.50+282438.8	273.6979355	-28.4107823	18144749+2824389			OGLE-BLG-LPV-226979	M	231.9
3572	CW-OG	228903	J181602.35+240937.0	274.0098284	-24.1603045	18160237+2409370	1814474+282438		OGLE-BLG-LPV-227901	M	359.7
3573	CW-OG	230266	J181744.12+252836.3	274.433864	-25.4767681	18174413+2528361	1816023+240936	G007.5149+03.5473	OGLE-BLG-LPV-228903	M	293.6
3574	CW-OG	230608	J181944.05+222311.2	274.9335824	-22.3864556	18194444+2223039	1817441+252837	G006.5396+04.4973	OGLE-BLG-LPV-230266	M	501.2
3575	CW-OG	231818	J182313.69+252653.8	275.8070795	-25.4482779	18231371+2526540	1819444+222304	G009.4924+03.4501	OGLE-BLG-LPV-230608	M	366.9
3576	CW-OG	231979	J182335.93+262852.2	275.8997362	-26.4811782	18233594+2628521			OGLE-BLG-LPV-231818	M	355.8
									OGLE-BLG-LPV-231979	M	263.1

¹The CAGB-WISE identifier (see Table 2). ²The ALLWISE source position (right ascension and declination J2000). See Section 9.

Table 13. OAGB-IRAS objects with obtained variation parameters from WISE light curves⁰ (24 columns; 885 rows)

OLN ¹	A-Type	A-Period	EP	G-W1(P)	G-W1(A)	G-W1(R ²)	GM-W1(P)	GM-W1(A)	GM-W1(R ²)	GN-W1(P)	GN-W1(A)	GN-W1(R ²)	G-W2(P)	G-W2(A)	G-W2(R ²)
181			880.56	532.78	0.816	0.731				532.78	0.816	0.731			
183	L:		70.48	59.55	0.829	0.677				88.53	0.837	0.615	125.44	0.839	0.700
264			316.24	546.50	0.763	0.726				273.61	0.725	0.629			
295			338.04	1233.38	0.986	0.639				213.70	1.030	0.504			
392			439.42	124.46	1.114	0.937				390.09	1.165	0.902	124.38	1.134	0.940
523	VAR		720.22	1654.50	1.340	0.601				204.20	1.288	0.496			
533			115.72	559.05	0.987	0.628				108.34	0.789	0.335	137.05	1.023	0.643
555	M	383											90.34	1.368	0.787
560			1576.20	207.57	1.315	0.764				1464.18	1.302	0.779			
579			132.85	543.11	1.213	0.808				136.26	1.199	0.771	550.95	1.318	0.627
611			58.82										111.02	0.423	0.698
642			428.80	467.42	1.123	0.766				467.42	1.123	0.766	467.42	1.404	0.660
653			865.06	2001.50	1.214	0.687				199.64	1.193	0.628			
656			341.11										607.19	1.159	0.606
657	VAR		621.10										185.25	0.073	0.624
666			554.24	781.39	1.213	0.775				781.39	1.213	0.775			
688			452.11	570.87	0.896	0.620				570.87	0.896	0.620			
704	VAR		664.36	1420.69	1.050	0.632				208.97	1.084	0.622			
709			359.41	234.67	1.257	0.726				234.67	1.257	0.726			
731				952.69	0.868	0.860							962.30	0.946	0.800
773			125.16	835.88	1.361	0.736									
796			282.57										644.07	0.959	0.623
802			419.00	878.19	1.498	0.710				229.32	1.414	0.565	549.50	0.977	0.648
803			1151.96	683.28	0.883	0.745				683.28	0.883	0.745			
806				821.49	1.427	0.692									
819															
821	VAR		263.99										505.85	0.902	0.608
827	VAR:		1793.77	2290.05	1.148	0.845				2290.05	1.148	0.845	2262.90	1.241	0.750
833			455.15	716.64	0.935	0.611				243.98	0.788	0.615			
835			436.78	546.37	1.035	0.694				546.37	1.035	0.694	540.18	1.265	0.627
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5699	M	301.3											292.65	0.544	0.708
5702	M	331.2											71.90	0.444	0.840
5703	M	278											274.15	0.529	0.757
5711	M	362.9											73.05	0.528	0.754
5716	M	310											306.27	0.229	0.616
5725	M	445.6											444.64	0.877	0.603
5726	M	390.9											124.67	0.467	0.670
5752	M	575											570.73	0.763	0.616
5799	M	444											89.94	0.991	0.782
5861	M	398.8247036											124.54	0.842	0.641

⁰Only 16 columns are shown in this example table. In the data file, there are eight more columns: GM-W2(P), GM-W2(A), GM-W2(R²), GN-W2(P), GN-W2(A), GN-W2(R²), Subgroup, and IRAS PSC. ¹The OAGB-IRAS identifier (see Tables 1 and 9). See Section 9.

Table 14. CAGB-IRAS objects with obtained variation parameters from WISE light curves⁰ (24 columns; 141 rows)

CLN ¹	A-Type	A-Period	EP	G-W1(P)	G-W1(A)	G-W1(R ²)	GM-W1(P)	GM-W1(A)	GM-W1(R ²)	GN-W1(P)	GN-W1(A)	GN-W1(R ²)	G-W2(P)	G-W2(A)	G-W2(R ²)
6				1062.66	1.315	0.935							1068.57	1.730	0.671
18	M	1060		1017.79	1.715	0.940							1001.88	2.180	0.895
34	M			798.05	1.243	0.796		1.715	0.940						
35			1224.47	882.29	1.137	0.780				1.137	0.780				
58			713.21	791.47	1.028	0.606				0.788	0.606				
90	EW	0.5546584	626.50	147.96	0.753	0.702				0.675	0.732				
91			362.81	628.79	0.749	0.716				0.651	0.678				
106	M:		137.15	51.55	0.760	0.698				0.651	0.552				
111			339.67	620.59	0.828	0.764				0.788	0.638		1.014	0.605	
120	SR	89.1313574	563.64	815.10	1.004	0.663				0.659	-0.030				
192			2199.82	898.91	0.855	0.696				0.855	0.696				
210			978.01	839.98	1.115	0.872				1.115	0.872				
230			706.87	485.67	1.007	0.833				1.007	0.833				
233	M	455.3437419		419.91	0.907	0.670		0.907	0.670						
246			397.66	529.24	0.886	0.670				0.819	0.615		531.91	0.912	0.614
251				837.44	1.135	0.895							816.15	1.132	0.603
277	SR	339.9025322	232.80	428.51	0.703	0.704				0.713	0.624		422.86	0.904	0.702
280	ELL—ROT	1.083	696.54	855.04	1.049	0.818				1.049	0.818				
283			257.89	392.57	0.458	0.605				0.495	0.658				
292	M	443.003349		179.95	2.096	0.659	255.25	0.660	0.220						
313			595.25	641.20	0.976	0.883				0.976	0.883		643.35	1.034	0.783
317			290.27	528.43	0.820	0.694				0.779	0.622				
338			325.23	519.87	0.644	0.740				0.597	0.692		524.12	0.550	0.768
348	VAR		379.01	597.37	0.908	0.704				0.914	0.650				
373			352.36	634.81	0.654	0.684				0.666	0.514				
380	EW	0.3634808	765.67	689.44	1.288	0.867				1.288	0.867		686.98	1.183	0.612
396			135.96	87.41	0.907	0.641				1.054	0.638				
406			292.60	577.60	0.668	0.605				0.677	0.585				
407			380.11	104.06	1.378	0.644				0.595	0.594				
-			-	-	-	-	-	-	-	-	-	-	-	-	-
2984	SR	409	147.59	60.08	0.703	0.625				0.425	0.228				
3052	SRB	175.4											177.87	0.775	0.609
3089	SR	858	128.80	175.93	0.641	0.650				0.544	0.506				
3110	SR	214.4446898	110.82	176.23	0.705	0.630				0.630	0.603				
3157			129.65										180.35	3.233	0.684
3203	SR	342	145.61	368.30	1.138	0.617				0.902	0.589				
3267	SR	130.58	86.95	185.12	0.943	0.659				0.588	0.631				
3324	SR	173.7469153	83.53	180.17	1.559	0.677				0.680	0.670				
3410	RCB	37.5	226.32	194.33	0.859	0.778				0.859	0.778		194.92	1.271	0.869
3590	M	555.8											551.19	0.894	0.652

⁰Only 16 columns are shown in this example table. In the data file, there are eight more columns: GM-W2(P), GM-W2(A), GM-W2(R²), GN-W2(P), GN-W2(A), GN-W2(R²), Subgroup, and IRAS PSC. ¹The CAGB-IRAS identifier (see Tables 1 and 10). See Section 9.

Table 15. OAGB-WISE objects with obtained variation parameters from WISE light curves (16 columns; 2468 rows)

OW-N ¹	Subgroup	A-Type	A-Period	G-W1(P)	G-W1(A)	G-W1(R^2)	GM-W1(P)	GM-W1(A)	GM-W1(R^2)	G-W2(P)	G-W2(A)	G-W2(R^2)	GM-W2(P)	GM-W2(A)	GM-W2(R^2)
21	OW-ME		597.79	0.605		0.707				455.23	0.672	0.683			
30	OW-ME									325.59	0.500	0.741			
43	OW-ME	M:								541.79	1.082	0.740			
48	OW-ME														
82	OW-ME		313.27	0.490	0.610					492.84	0.632	0.739			
86	OW-ME		485.32	0.521	0.639					467.50	0.591	0.840			
95	OW-ME		467.50	0.562	0.817					526.87	0.567	0.623			
142	OW-ME		535.74	0.554	0.708					139.83	0.645	0.806			
149	OW-ME		139.94	0.656	0.773					443.63	0.628	0.805			
154	OW-ME		444.66	0.614	0.772					555.94	0.690	0.777			
177	OW-ST		551.13	0.628	0.734					143.78	0.760	0.632			
190	OW-ST		664.94	0.698	0.678					603.87	0.688	0.670			
202	OW-ST		601.97	0.739	0.730										
203	OW-ST		655.68	0.675	0.644					908.00	1.088	0.745			
210	OW-ST		908.00	0.917	0.829					504.86	0.550	0.724			
251	OW-ST		506.20	0.588	0.721					245.42	1.161	0.792			
255	OW-ST		697.99	0.981	0.732					528.60	0.919	0.727			
257	OW-ST		522.81	0.769	0.719										
265	OW-ST		242.93	1.085	0.706					590.05	0.780	0.869			
272	OW-ST		593.72	0.606	0.873					248.58	1.001	0.799			
282	OW-ST		676.37	1.058	0.826										
283	OW-ST		706.43	0.851	0.638					609.51	0.815	0.606			
289	OW-ST														
294	OW-ST		141.24	0.685	0.619										
296	OW-ST		644.39	0.736	0.627					646.58	0.776	0.685			
298	OW-ST		141.65	0.780	0.706					253.85	0.679	0.757			
299	OW-ST		637.94	0.623	0.697					468.83	0.880	0.891			
305	OW-ST		468.83	0.800	0.872					136.68	0.791	0.778			
325	OW-ST		136.49	0.665	0.717										
339	OW-ST		104.57	1.685	0.634										
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5287	OW-OG	M	336.6	341.92	0.513	0.855	341.92	0.513	0.855	342.53	0.362	0.813	342.53	0.362	0.813
5288	OW-OG	M	283.5	278.15	0.498	0.621	278.15	0.498	0.621	278.97	0.394	0.790	278.97	0.394	0.790
5291	OW-OG	M	245.3	236.17	0.192	0.637	236.17	0.192	0.637						
5292	OW-OG	M	305.1	305.29	0.342	0.828	305.29	0.342	0.828	304.80	0.255	0.832	304.80	0.255	0.832
5295	OW-OG	M	339.4	339.49	0.464	0.849	339.49	0.464	0.849	339.49	0.350	0.913	339.49	0.350	0.913
5296	OW-OG	M	229.2	870.31	0.406	0.825	228.82	0.385	0.819	866.36	0.284	0.784	228.82	0.274	0.789
5297	OW-OG	M	197.57	200.46	0.237	0.770	200.46	0.237	0.770						
5298	OW-OG	M	263.2	271.05	0.506	0.719	271.05	0.506	0.719	271.43	0.468	0.722	271.43	0.468	0.722
5299	OW-OG	M	298.8							479.25	0.373	0.808	292.20	0.363	0.781
5300	OW-OG	M	199.55							205.65	0.277	0.693	205.65	0.277	0.693

¹The OAGB-WISE identifier (see Tables 2 and 11). See Section 9.

Table 16. CAGB-WISE objects with obtained variation parameters from WISE light curves (16 columns; 216 rows)

CW-N ¹	Subgroup	A-Type	A-Period	G-W1(P)	G-W1(A)	G-W1(R^2)	GM-W1(P)	GM-W1(A)	GM-W1(R^2)	G-W2(P)	G-W2(A)	G-W2(R^2)	GM-W2(P)	GM-W2(A)	GM-W2(R^2)
7	CW-GC	L		62.76	0.158	0.680									
9	CW-GC	M:	259	245.74	0.611	0.933				245.74	0.345	0.941			
65	CW-GC									151.26	0.117	0.657			
69	CW-GC	SR	335	333.84	0.465	0.828				117.96	0.269	0.832			
94	CW-GC	M	365	121.72	0.588	0.878	365.25	0.890	0.873	122.03	0.436	0.932	367.85	0.393	0.924
176	CW-GC			502.38	0.102	0.673									
197	CW-GC	M	250							259.83	0.335	0.618	259.83	0.335	0.618
252	CW-GC	L:	234	293.99	0.172	0.785									
261	CW-GC			149.61	0.068	0.710									
326	CW-GC														
371	CW-GC	SR	356.4753836							176.59	0.354	0.631			
392	CW-GC	SR	276							177.57	0.371	0.705			
398	CW-GC	SR:								283.23	0.175	0.712			
403	CW-GC									427.38	0.119	0.672			
459	CW-GC	CST													
483	CW-GC	L:		178.72	0.460	0.615									
520	CW-GC	M	374	372.89	0.658	0.778		0.658	0.778	1173.07	0.103	0.623			
523	CW-GC	SR	329	401.80	0.359	0.696				380.26	0.406	0.681	380.26	0.406	0.681
552	CW-GC	VAR								334.13	0.205	0.856			
586	CW-GC	I:		186.36	0.103	0.760				191.38	0.093	0.613			
598	CW-GC	SR	271.08671	64.47	0.282	0.613				186.91	0.163	0.698			
604	CW-GC	SR	421												
608	CW-GC	VAR								71.49	0.164	0.750			
633	CW-GC	VAR		300.98	0.486	0.673				1905.48	0.102	0.757			
648	CW-GC	L		829.16	0.130	0.630				302.40	0.326	0.806			
680	CW-GC	SR	292	95.34	0.275	0.640				818.57	0.102	0.642			
696	CW-GC									112.81	0.246	0.696			
762	CW-GC			566.58	0.133	0.672				1029.98	0.095	0.615			
786	CW-GC														
802	CW-GC	VAR								176.11	0.276	0.741			
-	-	-	-	-	-	-	-	-	-	178.07	0.174	0.608	-	-	-
3557	CW-OG	M	242.5							147.04	0.324	0.718	240.94	0.268	0.651
3558	CW-OG	M	308	113.41	0.313	0.848	301.91	0.319	0.844						
3560	CW-OG	M	229.3	101.42	0.214	0.662	229.92	0.204	0.611	101.42	0.160	0.664	229.64	0.152	0.617
3564	CW-OG	M	145.71	146.70	0.113	0.664	146.70	0.113	0.664						
3568	CW-OG	M	111.86	113.21	0.252	0.772	113.21	0.252	0.772	113.35	0.202	0.784	113.35	0.202	0.784
3569	CW-OG	RCB:		2500.00	0.222	0.775									
3570	CW-OG	M	231.9	229.64	0.360	0.891	229.64	0.360	0.891	229.09	0.287	0.856	229.09	0.287	0.856
3571	CW-OG	M	359.7	73.28	0.429	0.906	363.40	1.396	0.902	362.71	0.975	0.929	362.71	0.975	0.929
3573	CW-OG	M	501.2	300.46	0.510	0.908	460.72	0.498	0.872	300.94	0.485	0.915	459.61	0.480	0.897
3576	CW-OG	M	263.1	257.16	0.315	0.849	257.16	0.315	0.849	611.18	0.198	0.788	258.21	0.204	0.777

¹The CAGB-WISE identifier (see Tables 2 and 12). See Section 9.