EXISTENCE, STABILITY AND REGULARITY OF PERIODIC SOLUTIONS FOR NONLINEAR FOKKER-PLANCK EQUATIONS

ERIC LUÇON AND CHRISTOPHE POQUET

ABSTRACT. We consider a class of nonlinear Fokker-Planck equations describing the dynamics of an infinite population of units within mean-field interaction. Relying on a slow-fast viewpoint and on the theory of approximately invariant manifolds we obtain the existence of a stable periodic solution for the PDE, consisting of probability measures. Moreover we establish the existence of a smooth isochron map in the neighborhood of this periodic solution.

1. Introduction

1.1. **The model.** We are interested in this paper in the existence, stability and regularity of periodic solutions to the following nonlinear PDE on \mathbb{R}^d $(d \ge 1)$:

$$\partial_t u_t = \nabla \cdot \left(\sigma^2 u_t\right) + \nabla \cdot \left(K\left(x - \int_{\mathbb{R}^d} y u_t(\mathrm{d}y)\right) u_t\right) - \delta \nabla \cdot (F(x)u_t). \tag{1.1}$$

Here, $t \geq 0 \mapsto u_t$ is a probability measure-valued process on \mathbb{R}^d , K and σ and diagonal matrices with positive coefficients and $F: \mathbb{R}^d \to \mathbb{R}^d$ is a smooth bounded function with bounded derivatives. Equation (1.1) has a natural probabilistic interpretation: if u_0 is a probability distribution on \mathbb{R}^d , it is well known [30, 37] that u_t is the law of the McKean-Vlasov process X_t where $X_0 \sim u_0$ and

$$dX_t = \delta F(X_t) dt - K (X_t - \mathbb{E}[X_t]) dt + \sqrt{2}\sigma dB_t, \qquad (1.2)$$

The dynamics of the process $(X_t)_{t\geqslant 0}$ is the superposition of a local part $\delta F(X_t) \mathrm{d}t$, where $\delta>0$ is a scaling parameter, a linear interaction term $K\left(X_t-\mathbb{E}[X_t]\right)\mathrm{d}t$, modulated by the intensity matrix K, and an additive noise given by a standard Brownian motion $(B_t)_{t\geqslant 0}$ on \mathbb{R}^d . The difficulty in the analysis of (1.2) lies in its nonlinear character: X_t interacts with its own law, more precisely its own expectation $\mathbb{E}[X_t]$. The long-time dynamics of (1.2) is a longstanding issue in the literature. In particular, the existence of stable equilibria for (1.1) (that is invariant measures for (1.2)) has been studied for various choices of dynamics, interaction and regimes of parameters δ, K, σ , mostly in a context where the corresponding particle dynamics defined in (1.3) below is reversible (see e.g. [39, 7, 11] for further details and references).

The question we address in the present paper concerns the existence of periodic solutions to nonlinear equation such as (1.1). In this case, a major difficulty lies in the fact that the underlying microscopic dynamics is not reversible. From a practical perspective, the emergence of periodicity in such models relates in particular to chemical reactions (Brusselator model [35]), neurosciences [20, 21, 2, 9, 17, 27, 33, 28, 14], and statistical physics (e.g.

Date: January 13, 2022.

²⁰¹⁰ Mathematics Subject Classification. 35K55, 35Q84, 37N25, 60K35, 82C31, 92B20.

Key words and phrases. Mean-field systems, Nonlinear Fokker-Planck equation, McKean-Vlasov process, periodic behavior, Normally hyperbolic manifolds, Isochron map.

spin-flip models [13, 16], see also to [12], where the model considered is in fact not mean-field, but the Ising model with dissipation). An example of particular interest concerns the FitzHugh-Nagumo model [34, 2] (take d=2 and $F(x,y)=\left(x-\frac{x^3}{3}-y,\frac{1}{c}\left(x+a-by\right)\right)$), commonly used as a prototype for excitability in neuronal models [26] or in physics [3]. Roughly speaking, excitability refers to the ability for a neuron to emit spikes (oscillations) in the presence of perturbations (such as noise and/or external input) whereas this neuron would be at rest (steady state) without perturbation. The long-time dynamics of (1.1) in the FizHugh-Nagumo case has been the subject of several previous works (existence of equilibria [31, 33] or periodic solutions [27, 28]) under various asymptotics of the parameters (δ, K, σ) . A crucial feature in this context is the influence of noise and interaction in the emergence and stability of periodic solutions: generically, some balance has to be found in the quantity of noise and interaction that one needs to put in the system in order to observe oscillations (see [26, 27, 28] for further details).

Stability properties and regular isochron map. The purpose of the present paper is to complement the previous results concerning the existence of periodic orbits for (1.1) with accurate stability properties for this periodic solution and with the existence of a sufficiently regular isochron map, properties that are absent in the previous works cited above. We obtain these additional properties by applying a result concerning normally hyperbolic invariant manifolds in Banach spaces proved by Bates, Lu and Zeng [5]. The technical counterpart is that we require assumptions on F and σ that are somehow stricter than the ones used in [35, 27, 33, 28], in the sense that we are considering a dynamics F that is bounded together with all its derivatives (the Brusselator model and FitzHugh Nagumo model considered in the references given above have polynomial growth) as well as nondegenerate noise on all components (while in [33, 28] the noise is only present in one of the two variables).

Large time asymptotics for the mean-field particle system. Standard propagation of chaos results [37] shows that (1.2) is the natural limit of the following mean-field particle system

$$dX_{i,t} = \delta F(X_{i,t}) dt - K \left(X_{i,t} - \frac{1}{N} \sum_{j=1}^{N} X_{j,t} \right) dt + \sqrt{2}\sigma dB_{i,t},$$
 (1.3)

in the sense that one can easily couple (1.3) and (1.2) by choosing the same realization of the noise, so that the resulting error is of order $\frac{1}{\sqrt{N}}$ as $N \to \infty$, at least on any [0,T] with T that can be arbitrarily large but fixed independently from N. At the level of the whole particle system, this boils down to the convergence as $N \to \infty$ of the empirical measure $u_{N,t} = \frac{1}{N} \sum_{i=1}^{N} \delta_{X_{i,t}}$ to u_t , solution to (1.1). Hence, supposing that (1.1) has a periodic solution $(\Gamma_t^{\delta})_{t \geq 0}$, if the empirical measure $u_{N,0}$ is initially close to $\Gamma_{\theta_0}^{\delta}$, $u_{N,t}$ has, for N large, a behavior close to being periodic, since it stays close to $\Gamma_{\theta_0+t}^{\delta}$.

The companion paper [29] of the present work is concerned with the behavior of the empirical measure $u_{N,t}$ on a time scale T that is no longer bounded, but of order N. We show in [29] that $u_{N,Nt}$ is close to $\Gamma^{\delta}_{\theta_0+Nt+\beta^N_t}$, where β^N_t is a random process in \mathbb{R} whose weak limit as $N \to \infty$ has constant drift and diffusion coefficient. This kind of result was already obtained in [15, 8] in the case of the plane rotators model (mean-field noisy interacting oscillators defined on the circle), for which in the scale Nt the empirical measure has a diffusive behavior along the curve of stationary points. Our aim in [29] is to get similar results for models like (1.1) that are defined in \mathbb{R}^d , and are not reversible

(while the plane rotators model is). As we will explain in more details later, the additional stability and regularity results concerning periodic solution to (1.1) obtained in the present paper are crucial for the study of long time behavior of the mean-field particle systems (1.3) made in [29].

1.2. Slow fast viewpoint and application to the FitzHugh-Nagumo model. We give in this paragraph informal intuition on the possibility of emergence of periodic solutions to (1.1). The point of view we adopt here is a slow-fast approach, based on the assumption that the parameter δ in (1.1) is small, as it was already the case in [27, 28]. More precisely, the linear character of the interaction term in (1.1) allows us to decompose the dynamics of (1.1) into its expectation $m_t = \int_{\mathbb{R}^d} x u_t(x)$ and its centered version $p_t(x) = u_t(x - m_t)$: (1.1) is equivalent to the system

$$\begin{cases}
\partial_t p_t = \mathcal{L} p_t - \nabla \cdot (p_t (\delta F_{m_t} - \dot{m}_t)) \\
\dot{m}_t = \delta \int F_{m_t} \, \mathrm{d} p_t
\end{cases},$$
(1.4)

where

$$\mathcal{L}u = \nabla \cdot (\sigma^2 \nabla f) + \nabla \cdot (Kxf), \qquad (1.5)$$

and $F_m(x) = F(x+m)$. Remark that (p_t, m_t) is the weak limit as $N \to \infty$ of the process $\left(\frac{1}{N}\sum_{i=1}^N \delta_{Y_{i,t}}, m_{N,t}\right)$, where

$$m_{N,t} = \frac{1}{N} \sum_{i=1}^{N} X_{i,t}, \text{ and } Y_{i,t} = X_{i,t} - m_{N,t}.$$
 (1.6)

In this set-up, p_t is here the fast variable, while m_t is the slow one. For $\delta = 0$, this system reduces to

$$\begin{cases}
\partial_t p_t^0 = \mathcal{L} p_t^0 \\
m_t^0 = m_0
\end{cases},$$
(1.7)

so $p_t^0 = e^{t\mathcal{L}}p_0$ is the distribution of an Ornstein-Uhlenbeck process, and thus converges exponentially fast to ρ , the density of the Gaussian distribution on \mathbb{R}^d with mean 0 and variance $\sigma^2 K^{-1}$ (see Proposition 1.1 for more details on the contraction properties of \mathcal{L}). So heuristically, taking δ small, at first approximation p_t stays close to ρ while m_t satisfies

$$\dot{m}_t \approx \delta \int_{\mathbb{R}^d} F_{m_t}(x) \rho(x) \, \mathrm{d}x = \delta \int_{\mathbb{R}^d} F(x) \rho(m_t - x) \, \mathrm{d}x = \delta(F * \rho)(m_t). \tag{1.8}$$

For the non-centered PDE (1.1) this approximation means that u_t is close to a Gaussian distribution with variance variance $\sigma^2 K^{-1}$ and mean m_t , where the dynamics of m_t is governed at first order by (1.8). Following this heuristic, we expect a periodic behavior for the system (1.4) if the approximate dynamics of m_t is itself periodic. In this spirit, the main hypothesis we will adopt below is that the solution z_t to

$$\dot{z}_t = \delta \int_{\mathbb{R}^d} F_{z_t}(x) \rho(x) \, \mathrm{d}x = \delta \, \langle F_{z_t}, \rho \rangle \tag{1.9}$$

admits a stable periodic solution $(\alpha_t^{\delta})_{t\in[0,\frac{T_{\alpha}}{\delta}]}$ (more details on this notion of stability, in terms of Floquet exponents, will be given in Section 1.4). In Proposition 1.7 we will show that with these hypotheses, the manifold $\widetilde{\mathcal{M}}^{\delta} = (\rho, \alpha_t^{\delta})_{t\in[0,T_{\alpha}/\delta]}$ is approximately invariant for (1.4).

Let us now describe a situation where the above heuristics is true: in [27, 28] we considered the classical FitzHugh-Nagumo model defined by d=2 and

$$F(x,y) = \left(x - \frac{x^3}{3} - y, \frac{1}{c}(x + a - by)\right). \tag{1.10}$$

A direct calculation shows that in that case, with $K = \operatorname{diag}(k_1, k_2)$ and $\sigma = \operatorname{diag}(\sigma_1, \sigma_2)$,

$$\int_{\mathbb{R}^d} F_{z_1, z_2}(x, y) \rho(x, y) \, \mathrm{d}x \, \mathrm{d}y = \left(\left(1 - \frac{\sigma_1^2}{k_1} \right) z_1 - \frac{z_1^3}{3} - z_2, \frac{1}{c} \left(z_1 + a - b z_2 \right) \right), \tag{1.11}$$

which defines again a FitzHugh-Nagumo model. The additional factor $\frac{\sigma_1^2}{k_1}$ in (1.11) reflects the influence of noise and interaction in the mean-field system (1.2). For an accurate choice of parameters (take e.g. $a=\frac{1}{3},\ b=1$ and c=10), it can be shown that the dynamics of the mean value (1.9) has a unique steady state when $\frac{\sigma_1^2}{k_1}=0$ whereas it admits a stable periodic solution for $\frac{\sigma_1^2}{k_1}$ not too small and not too large, for example $\frac{\sigma_1^2}{k_1}=0.2$. We refer to [27], § 3.4 for a more details of the corresponding bifurcations). The purpose of [27, 28] was to show that the heuristics developed above is true, i.e. the periodicity of (1.9) propagates to (1.4). This emergence of periodic behavior induced by noise and interaction is a signature of excitability: the system (1.1) exhibits a periodic behavior induced by the combined effect of the noise and the interaction, which is not present in the isolated system $\dot{z}_t = F(z_t)$. We refer to [27] for a discussion and references on this phenomenon.

As already said, the point of the present work is to go beyond the existence of oscillations for (1.1), that is to prove regularity for the dynamics around such a limit cycle. Unfortunately the FitzHugh Nagumo model does not satisfy the hypotheses of this present work, since it has polynomial growth at infinity. However it is easy to see that if $\psi : \mathbb{R}_+ \to \mathbb{R}_+$ is a smooth non-increasing function that satisfies $\psi(t) = 1$ for $t \leq 1$ and $\psi(t) = 0$ for $t \geq 2$, then for any $\varepsilon > 0$ the function $x \mapsto F(x)\psi(\varepsilon|x|)$ satisfies our hypotheses, and that $z \mapsto \int_{\mathbb{R}^d} F_z(x)\psi(\varepsilon|x+z|)\rho(x)dx$ converges to $z \mapsto \int_{\mathbb{R}^d} F_z(x)\rho(x)dx$ in $C^1(\mathcal{B}(0,R),\mathbb{R}^d)$ for any ball $\mathcal{B}(0,R)$ centered at 0 with radius R. So, relying on classical results on normally hyperbolic manifolds [18, 19, 40] (a definition of this notion will be provided in Section 1.4), if (1.9) admits a stable limit cycle, then it will also be the case replacing F with $x \mapsto F(x)\psi(\varepsilon|x|)$ for ε small enough.

1.3. Weighted Sobolev norms. We present in this section the Sobolev spaces that we will use in the paper. Let us denote by $|x|_A = (x \cdot Ax)^{1/2}$ the Euclidean norm twisted by some positive matrix A, and, for any $\theta \in \mathbb{R}$, let us define the weight w_θ by

$$w_{\theta}(x) = \exp\left(-\frac{\theta}{2} |x|_{K\sigma^{-2}}^{2}\right). \tag{1.12}$$

We denote as L_{θ}^2 the L^2 -space with weight w_{θ} , that is with norm

$$||h||_{L^{2}_{\theta}} = \left(\int_{\mathbb{R}^{d}} |h(x)|^{2} w_{\theta}(x) dx\right)^{\frac{1}{2}}.$$
 (1.13)

For any $\theta > 0$ we consider the Ornstein-Uhlenbeck operator

$$\mathcal{L}_{\theta}^* f = \nabla \cdot (\sigma^2 \nabla f) - \theta K x \cdot \nabla f. \tag{1.14}$$

It is well know (see for example [1]) that \mathcal{L}_{θ}^* admits the following decomposition: for all $l \in \mathbb{N}^d$,

$$\mathcal{L}_{\theta}^* \psi_l = -\lambda_l \psi_l, \quad \text{with} \quad \lambda_l = \theta \sum_{i=1}^d k_i l_i \quad \text{and} \quad \psi_l(x) := \psi_{l,\theta}(x) = \prod_{i=1}^d h_{l_i} \left(\sqrt{\frac{\theta k_i}{\sigma_i^2}} x_i \right), \tag{1.15}$$

where h_n is the n^{th} renormalized Hermite polynomial:

$$h_n(x) = \frac{(-1)^n}{\sqrt{n!}(2\pi)^{\frac{1}{4}}} e^{\frac{x^2}{2}} \frac{\mathrm{d}^n}{\mathrm{d}x^n} \left\{ e^{-\frac{x^2}{2}} \right\}.$$
 (1.16)

The family $(\psi_{l,\theta})_{l\in\mathbb{N}^d}$ is an orthonormal basis of L^2_{θ} . For f,g with decompositions $f = \sum_{l\in\mathbb{N}^d} f_l \psi_l$ and $g = \sum_{l\in\mathbb{N}^d} g_l \psi_l$, we consider the scalar products

$$\langle f, g \rangle_{H_{\theta}^r} = \langle (a_{\theta} - \mathcal{L}_{\theta}^*)^r f, \bar{g} \rangle_{L_{\theta}^2} = \sum_{l \in \mathbb{N}^d} (a_{\theta} + \lambda_l)^r f_l \bar{g}_l, \tag{1.17}$$

where $a_{\theta} = \theta \operatorname{Tr} K$ and denote by H_{θ}^r the completion of the space of smooth function u satisfying $\|u\|_{H_{\theta}^r} < \infty$. The choice of the constant a_{θ} is made to simplify some technical proofs given in the Appendix A (see the proof of Proposition A.2). Another choice of positive constant would produce an equivalent norm. From Lemma A.1 it is clear that $\|\partial_{x_i} f\|_{H_{\theta}^r} \leq \|f\|_{H_{\theta}^{r+1}}$, and that, if $n \in \mathbb{N}$, the norm $\|f\|_{H_{\theta}^n}$ is in fact equivalent to

$$\sqrt{\sum_{l \in \mathbb{N}^d, \sum_{i=1}^d l_i \leqslant n} \left\| \partial_{x_1}^{l_1} \dots \partial_{x_d}^{l_d} f \right\|_{L_{\theta}^2}^2}.$$
(1.18)

We denote by H_{θ}^{-r} the dual of H_{θ}^{r} . Relying on a "pivot" space structure (for more details, see Appendix A), the product $\langle u, f \rangle_{H_{\theta}^{-r}, H_{\theta}^{r}}$ can be identified with the flat L^{2} product $\langle u, f \rangle$: $L_{-\theta}^{2}$ can be seen as a subset of H_{θ}^{-r} , and for all $f \in H_{\theta}^{r}$ and $u \in L_{-\theta}^{2}$ we have

$$\langle u, f \rangle_{H_{\theta}^{-r}, H_{\theta}^{r}} = \langle u, f \rangle.$$
 (1.19)

This identification allows us to view the operator \mathcal{L}_{θ} defined by

$$\mathcal{L}_{\theta}u = \nabla \cdot (\sigma^2 \nabla f) + \nabla \cdot (\theta K x f), \qquad (1.20)$$

seen as an operator in H_{θ}^{-r} , as the adjoint of \mathcal{L}_{θ}^* , seen as an operator in H_{θ}^r . This is in particular the case for $\mathcal{L} = \mathcal{L}_1$, whose contraction properties will be crucial in the results given in this paper.

Our aim in this paper is to give the existence of a periodic solution for (1.4) viewing p_t as an element of H_{θ}^{-r} . The necessity of considering H_{θ}^{-r} instead of simply taking H_1^{-r} goes back to the companion paper [29], in which we study the long time behavior of the empirical measure $u_{N,t}$ in the same functional spaces. Since this empirical measure involves a sum of Dirac distributions, it can be seen as an element of H_{θ}^{-r} for r > d/2, and we have $\|\delta_x\|_{H_{\theta}^{-r}} \leqslant Cw_{\frac{\theta}{4-\eta}}(x)$ for $\eta > 0$ (see Lemma 2.1 in [29]). Some moment estimates

lead us in [29] to bound terms of the form $\mathbb{E}\left[w_{\frac{m\theta}{4-\eta}}(Y_{i,t})\right]$ with m large and $Y_{i,t}$ defined in (1.6). Since we consider cases where $Y_{i,t}$ has a distribution close to ρ , for this expectation to be bounded we need to consider small values of θ . We need therefore to work in H_{θ}^{-r} for general θ and not only for $\theta = 1$.

It is well known that the semi-group $e^{t\mathcal{L}}$ satisfies, for $\lambda < \min(k_1, \dots, k_d)$ and $u \in H_1^{-r}$ with $\int u = 0$, the contraction property

$$\|e^{t\mathcal{L}}u\|_{H_1^{-r}} \le Ct^{-\frac{\alpha}{2}}e^{-t\lambda}\|u\|_{H_1^{-(r+\alpha)}}.$$
 (1.21)

By obtaining similar estimates (see the following Proposition, which is a particular case of the slightly more general Proposition A.3), we will be able to work in the space H_{θ}^{-r} with any value of θ smaller than 1, but with the constraint of considering values of r larger than a $r_0 > 0$ (independent from θ).

Proposition 1.1. For all $0 < \theta \le 1$ the operator \mathcal{L} is sectorial and generates an analytical semi-group in H_{θ}^{-r} . Moreover we have the following estimates: for any $\alpha \ge 0$, $r \ge 0$ and $\lambda < \min(k_1, \ldots, k_d)$ there exists a constant $C_{\mathcal{L}} > 0$ such that for all $u \in H_{\theta}^{-(r+\alpha)}$,

$$\left\| e^{t\mathcal{L}} u \right\|_{H_{\theta}^{-r}} \leqslant C_{\mathcal{L}} \left(1 + t^{-\alpha/2} e^{-\lambda t} \right) \left\| u \right\|_{H_{\theta}^{-(r+\alpha)}}, \tag{1.22}$$

and for $r \geqslant 1$,

$$||e^{t\mathcal{L}}\nabla u||_{H_{a}^{-r}} \le C_{\mathcal{L}}t^{-\frac{1}{2}}e^{-\lambda t}||u||_{H_{a}^{-r}}.$$
 (1.23)

Moreover for all $r \ge 0$, $0 < \varepsilon \le 1$ and $s \ge 0$,

$$\left\| \left(e^{(t+s)\mathcal{L}} - e^{t\mathcal{L}} \right) u \right\|_{H_{\sigma}^{-r}} \leqslant C_{\mathcal{L}} s^{\varepsilon} t^{-\frac{1}{2} - \varepsilon} e^{-\lambda t} \left\| u \right\|_{H_{\theta}^{-(r+1)}}. \tag{1.24}$$

Finally, there exist $r_0 > 0$ such that for any $0 < \theta \leqslant 1$, for all $r > r_0$, t > 0 and all $u \in H_{\theta}^{-r}$ satisfying $\int u = 0$,

$$\left\| e^{t\mathcal{L}} u \right\|_{H_{\theta}^{-r}} \leqslant C_{\mathcal{L}} e^{-\lambda t} \left\| u \right\|_{H_{\theta}^{-r}}. \tag{1.25}$$

1.4. Main results. With the notation $\mu_t := (p_t, m_t)$ the system (1.4) becomes

$$\begin{cases}
\partial_t p_t = \mathcal{L}p_t + \delta G_1(\mu_t) \\
\dot{m}_t = \delta G_2(\mu_t)
\end{cases},$$
(1.26)

where

$$G(\mu) = G(p, m) = \begin{pmatrix} G_1(p, m) \\ G_2(p, m) \end{pmatrix} = \begin{pmatrix} -\nabla \cdot \left(p \left(F_m - \int F_m p \right) \right) \\ \int F_m p \end{pmatrix}. \tag{1.27}$$

We place ourselves on the space $\mathbf{H}_{\theta}^r := H_{\theta}^r \times \mathbb{R}^d$ endowed with the scalar product

$$\left\langle \left(f,m\right),\left(g,m'\right)\right\rangle _{\mathbf{H}_{\rho}^{r}}:=\left\langle f\,,\,g\right\rangle _{H_{\rho}^{r}}+m\cdot m'.\tag{1.28}$$

We will denote \mathbf{H}_{θ}^{-r} the dual of \mathbf{H}_{θ}^{r} . Clearly $\mathbf{H}_{\theta}^{-r} = H_{\theta}^{-r} \times \mathbb{R}$ and, relying as above on a "pivot" space structure, the product $\langle (\nu, h), (\phi, \psi) \rangle_{\mathbf{H}_{\theta}^{-r}, \mathbf{H}_{\theta}^{r}}$ can be identified with the flat scalar product

$$\langle\!\langle (\nu, h); (\varphi, \psi) \rangle\!\rangle = \langle \nu, \varphi \rangle + h \cdot \psi. \tag{1.29}$$

The following theorem states the existence and uniqueness of mild solutions of (1.26). Its proof, given in Section 2, relies on classical arguments, due to the fact that $G: \mathbf{H}_{\theta}^{-r+1} \to \mathbf{H}_{\theta}^{-r}$ is locally Lispchitz and \mathcal{L} is sectorial (see [36]).

Theorem 1.2. For any initial condition $\mu = (p, m) \in \mathbf{H}_{\theta}^{-r}$ with $\int_{\mathbb{R}^d} p = 1$ there exists a unique maximal mild solution $\mu_t := (p_t, m_t) = T^t(\mu)$ to (1.26) on $[0, t_c]$ for some $t_c > 0$, which satisfies $t \mapsto T^t(\mu) \in \mathcal{C}\left([0, t_c); \mathbf{H}_{\theta}^{-r}\right)$.

Moreover, $\mu \mapsto T^t(\mu)$ is C^2 , and for any R > 0, there exists a $\delta(R) > 0$ such that for all $0 \le \delta \le \delta(R)$ and $\mu_0 = (p_0, m_0)$ satisfying $\|p_0 - \rho\|_{H_{\theta}^{-r}} \le R$ the solution $T^t(\mu_0)$ is well defined for all $t \ge 0$ and there exists a C(R) > 0 such that

$$\sup_{t \ge 0} \|p_t\|_{H_{\theta}^{-r}} \le C(R). \tag{1.30}$$

Remark 1.3. Since we are interested in the existence of a periodic solution made of probability distributions, we will only consider initial condition (p_0, m_0) satisfying $\int p_0 = 1$, and the conservation of mass will induce that $\int_{\mathbb{R}^d} p_t = 1$ for all t. In the same spirit, we will only apply the differential of the semi-group $DT^t(\mu)$ to elements $\nu = (\eta, n) \in \mathbf{H}_{\theta}^{-r}$ that satisfy $\int_{\mathbb{R}^d} \eta = 0$.

As it was previously mentioned, we suppose in the following that the ordinary differential equation (1.9) admits a stable periodic solution $(\alpha_t^{\delta})_{t \in [0, \frac{T_{\alpha}}{\delta}]}$. To state more precisely this hypothesis we rely on Floquet formalism (see for example [38]): let us denote by $\pi_{u+t,u}^{\delta}$ the principal matrix solution associated to the periodic solution α , that it the solution to

$$\partial_t \pi_{u+t,u}^{\delta} = \delta \langle DF_{\alpha_{u+t}^{\delta}}, \rho \rangle \pi_{u+t,u}^{\delta}, \ \pi_{u,u}^{\delta} = I.$$
 (1.31)

The process $\pi_{u+t,u}^{\delta}$ characterizes the linearized dynamics around $(\alpha_t^{\delta})_{t\in[0,\frac{T}{\delta}]}$. We will suppose that this linearized dynamics is a contraction on a supplementary space of the tangent space to $(\alpha_t^{\delta})_{t\in[0,\frac{T}{\delta}]}$. More precisely, we suppose that there exist projections $P_u^{\delta,c}$ and $P_u^{\delta,c}$ for all $u\in[0,\frac{T}{\delta}]$ with $u\mapsto P_u^{\delta,c}$ and $u\mapsto P_u^{\delta,s}$ smooth, that satisfy $P_u^{\delta,s}+P_u^{\delta,c}=I$ $(P_u^{\delta,c}$ being a projection on $\operatorname{vect}(\dot{\alpha}_u^{\delta}))$ and that commute with π^{δ} , i.e.

$$P_u^{\delta,s} + P_u^{\delta,c} = I, \quad P_{u+t}^{\delta,s} \pi_{u+t,u}^{\delta} = \pi_{u+t,u}^{\delta} P_u^{\delta,s}.$$
 (1.32)

We emphasize here the fact that the factor δ in (1.9) is only responsible for a time-scale of the dynamics (the smaller δ , the slower the dynamics, the period being then T_{α}/δ): we have $\alpha_t^{\delta} = \alpha_{\delta t}^1$. Its effect on the projections is only a change of parametrization: $P_u^{\delta,s}$ and $P_u^{\delta,c}$ are defined on $[0,T/\delta)$, and $P_{u/\delta}^{\delta,s} = P_u^{1,s}$, $P_{u/\delta}^{\delta,c} = P_u^{1,c}$ for $u \in [0,T_{\alpha})$.

The stability of the periodic solution $(\alpha_t^{\delta})_{t \in [0, \frac{T_{\alpha}}{\delta}]}$ is expressed by the following hypothesis: there exist positive constants c_{α}, C_{α} and λ_{α} such that

$$\left| \pi_{u+t,u}^{\delta} P_u^{\delta,s} n \right| \leqslant C_{\alpha} e^{-\delta \lambda_{\alpha} t} |n| \quad \text{and} \quad c_{\alpha} |n| \leqslant \left| \pi_{u+t,u}^{\delta} P_u^{\delta,c} n \right| \leqslant C_{\alpha} |n|. \tag{1.33}$$

The factor δ in the rate of contraction is again due to the fact that the factor δ in (1.9) slows the dynamics.

With these hypotheses $(\alpha_t^{\delta})_{t \in [0, \frac{T}{\delta}]}$ is in fact a simple example of Normally Hyperbolic Invariant Manifold (NHIM). We follow here the definition given in [4] for this concept: on a Banach space \mathbf{X} , a smooth compact connected manifold \mathbf{M} is said to be a normally hyperbolic invariant manifold for a continuous semi flow \mathbf{T} (such that $u \mapsto \mathbf{T}^t(\mu)$ is C^1 for all $t \geq 0$) if

- (1) $\mathbf{T}(\mathbf{M}) \subset \mathbf{M}$ for all $t \geq 0$,
- (2) For each $m \in \mathbf{M}$ there exists a decomposition $\mathbf{X} = \mathbf{X}_m^c + \mathbf{X}_m^u + \mathbf{X}_m^s$ of closed subspaces with \mathbf{X}_m^c the tangent space to \mathbf{M} at m,
- (3) For each $m \in \mathbf{M}$ and $t \geq 0$, denoting $m_1 = \mathbf{T}^t(m)$, we have $D\mathbf{T}^t(m)_{|\mathbf{X}_m^\iota|} : \mathbf{X}_m^\iota \to \mathbf{X}_{m_1}^\iota$ for $\iota = c, u, s$, and $D\mathbf{T}^t(m)_{|\mathbf{X}_m^u|}$ is an isomorphism from \mathbf{X}_m^u to $\mathbf{X}_{m_1}^u$.

(4) There exists a $t_0 \ge 0$ and a $\lambda > 0$ such that, for all $t \ge t_0$,

$$\lambda \inf\{|D\mathbf{T}^t(m)[x^u]|: x^u \in \mathbf{X}^u, |x^u| = 1\} > \max\{1, \|D\mathbf{T}^t(m)_{\mathbf{X}_{-\infty}^c}\|\},$$
 (1.34)

$$\lambda \min\{1, \inf |D\mathbf{T}^{t}(m)[x^{c}]| : x^{c} \in \mathbf{X}_{m}^{c}, |x^{c}| = 1\}\} > ||D\mathbf{T}^{t}(m)|_{\mathbf{X}_{m}^{c}}||.$$
(1.35)

(1.34) implies that the semi flow \mathbf{T}^t is expansive at m in the direction \mathbf{X}_m^u at a rate strictly larger than on \mathbf{M} , while (1.35) shows implies that it is contractive at m in the direction \mathbf{X}_m^s at a rate greater than on \mathbf{M} .

This kind of structure is known to be robust under perturbation of the semi-flow: it has been shown in [18, 19] for flows in \mathbb{R}^d , and then generalized in [24] in the case of Riemanian manifolds and in [4, 36] in the infinite dimensional setting. An improvement of these classical results has been obtained in [5] by Bates, Lu and Zeng, who showed that if a system admits a manifold that is approximately invariant and approximately normally hyperbolic (a precise definition of these notions will be given in Section 1.5), then the system possesses an actual normally hyperbolic invariant manifold in a neighborhood.

We will rely on this deep result in our work. Here, the slow fast viewpoint described in Section 1.2 suggests that for δ small the manifold $\widetilde{\mathcal{M}}^{\delta} = \{(\rho, \alpha_t) : t \in [0, T_{\alpha})\}$ is an approximately invariant manifold which is approximately normally hyperbolic (without unstable direction). This statement will be written rigorously in Section 1.5, and proved in Section 3. This idea will allow us to prove for δ small enough the existence of a stable periodic solution to (1.4), as an actual normally hyperbolic invariant manifold in a neighborhood of $\widetilde{\mathcal{M}}^{\delta}$. For a stable periodic solution, conditions (1.34) and (1.35) reduce to the fact that $DT^t(m)$ is bounded from above and below in the direction of the tangent space to the invariant manifold defined by the periodic solution, and is contractive on a stable direction.

Theorem 1.4. There exists $\delta_0 > 0$ such that for r_0 given in Proposition 1.1 and for all $r \geq r_0$, $\delta \in (0, \delta_0)$ and $\theta \in (0, 1]$ the system (1.4) admits a periodic solution $(\Gamma_t^{\delta})_{t \in [0, T_{\delta}]} := (q_t^{\delta}, \gamma_t^{\delta})_{t \in [0, T_{\delta}]}$ in \mathbf{H}_{θ}^{-r} with period $T_{\delta} > 0$. Moreover q_t^{δ} is a probability distribution for all $t \geq 0$, and $t \mapsto \partial_t \Gamma_t^{\delta}$ and $t \mapsto \partial_t^2 \Gamma_t^{\delta}$ are in $C([0, T_{\delta}), \mathbf{H}_{\theta}^{-r})$.

Denoting $\mathcal{M}^{\delta} = \{\Gamma_t^{\delta} : t \in [0, T_{\delta})\}$ and $\Phi_{u+s,u}(\nu) = DT^s(\Gamma_u^{\delta})[\nu]$, there exists families of projection $\Pi_u^{\delta,c}$ and $\Pi_t^{\delta,s}$ that commute with Φ , i.e. that satisfy

$$\Pi_{u+t}^{\delta,\iota} \Phi_{u+t,u} = \Phi_{u+t,u} \Pi_{u}^{\delta,\iota}, \quad \text{for } \iota = c, s.$$
 (1.36)

Moreover $\Pi_t^{\delta,c}$ is a projection on the tangent space to \mathcal{M}^{δ} at Γ_t^{δ} , $\Pi_t^{\delta,c} + \Pi_t^{\delta,s} = I_d$, $t \mapsto \Pi_t^{\delta,c} \in C^1([0,T_{\delta}),\mathcal{B}(\mathbf{H}_{\theta}^{-r}))$, and there exist positive constants $c_{\Phi,\delta}$, $C_{\Phi,\delta}$ and λ_{δ} such that

$$c_{\Phi,\delta} \left\| \Pi_u^{\delta,c}(\nu) \right\|_{\mathbf{H}_o^{-r}} \leqslant \left\| \Phi_{u+t,u} \Pi_u^{\delta,c}(\nu) \right\|_{\mathbf{H}_o^{-r}} \leqslant C_{\Phi,\delta} \left\| \Pi_u^{\delta,c}(\nu) \right\|_{\mathbf{H}_o^{-r}}, \tag{1.37}$$

$$\left\| \Phi_{u+t,u} \Pi_u^{\delta,s}(\nu) \right\|_{\mathbf{H}_{\sigma}^{-r}} \leqslant C_{\Phi,\delta} t^{-\frac{\alpha}{2}} e^{-\lambda_{\delta} t} \left\| \Pi_u^{\delta,s}(\nu) \right\|_{\mathbf{H}_{\sigma}^{-(r+\alpha)}}, \tag{1.38}$$

and

$$\|\Phi_{u+t,u}\nu\|_{\mathbf{H}_{\theta}^{-r}} \leq C_{\Phi,\delta} \left(1 + t^{-\frac{\alpha}{2}} e^{-\lambda_{\delta} t}\right) \|\nu\|_{\mathbf{H}_{\theta}^{-(r+\alpha)}}.$$
 (1.39)

Remark 1.5. The invariant manifold \mathcal{M}^{δ} is located at a distance of order δ from $\widetilde{\mathcal{M}}^{\delta}$ and the period T_{δ} is close to T_{α}/δ (the period of the slow system (1.9)). Moreover $\lambda_{\Phi,\delta}$ is of order δ due to the fact that z_t contracts around α_t with rate $\delta\lambda_{\alpha}$.

In [5] it is in addition proven that the stable manifold of the actual NHIM (in our case \mathcal{M}^{δ} is attractive, the stable manifold is in fact a neighborhood \mathcal{W}^{δ} of \mathcal{M}^{δ}) is foliated by invariant foliations: $\mathcal{W}^{\delta} = \bigcup_{m \in \mathcal{M}^{\delta}} \mathcal{W}^{\delta}_{m}$, where $\nu \in \mathcal{W}^{\delta}_{m}$ if and only if $T^{t}(\nu) - T^{t}(m)$ converges to 0 exponentially fast. This implies the existence of an isochron map Θ^{δ} : $\mathcal{W}^{\delta} \to \mathbb{R}/T_{\delta}\mathbb{Z}$ that satisfies $\Theta^{\delta}(\nu) = t$ if $\nu \in \mathcal{W}^{\delta}_{\Gamma^{\delta}_{t}}$. The deep general result of [5] ensures that Θ^{δ} is Hölder continuous, which is not entirely satisfying in view of the companion paper [29], in which we aim to apply Itô's Lemma to $\Theta^{\delta}(u_{N,t})$. However, the fact that in the present case we simply deal with a stable periodic solution allow us to prove that Θ^{δ} has in our particular case C^{2} regularity, as stated in the following theorem.

Theorem 1.6. For r and δ as in Theorem 1.4, there exists neighborhood $W^{\delta} \in \mathbf{H}_{\theta}^{-r}$ of \mathcal{M}^{δ} and a C^2 mapping $\Theta^{\delta} : W^{\delta} \to \mathbb{R}/T_{\delta}\mathbb{Z}$ that satisfies, for all $\mu \in W^{\delta}$, denoting $\mu_t = T^t \mu$,

$$\Theta^{\delta}(\mu_t) = \Theta^{\delta}(\mu) + t \mod T_{\delta}, \tag{1.40}$$

and there exists a positive constant $C_{\Theta,\delta}$ such that, for all $\mu \in \mathcal{W}^{\delta}$ with $\mu_t = T^t \mu$,

$$\left\| \mu_t - \Gamma_{\Theta^{\delta}(\mu) + t}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_{\Theta, \delta} e^{-\lambda_{\delta} t} \left\| \mu - \Gamma_{\Theta^{\delta}(\mu)}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}}. \tag{1.41}$$

Moreover Θ^{δ} satisfies, for all $\mu \in \mathcal{W}^{\delta}$,

$$\left\| D^2 \Theta^{\delta}(\mu) - D^2 \Theta^{\delta} \left(\Gamma_{\Theta^{\delta}(\mu)}^{\delta} \right) \right\|_{\mathcal{BL}(\mathbf{H}_{a}^{-r})} \leqslant C_{\Theta,\delta} \left\| \mu - \Gamma_{\Theta^{\delta}(\mu)}^{\delta} \right\|_{\mathbf{H}_{a}^{-r}}. \tag{1.42}$$

1.5. An approximately invariant manifold that is approximately normally hyperbolic. In view of the slow fast formalism described in Section 1.2, our aim is to view $\widetilde{\mathcal{M}}^{\delta} = \{(\rho, \alpha_t) : t \in [0, \frac{T}{\delta})\}$ as an approximately invariant and approximately normally hyperbolic manifold, in the sense of [5].

In fact the result of [5] is stated for dynamical systems taking values in a Banach space, while we will consider here solutions (p_t, m_t) to (1.4) elements of \mathbf{H}_{θ}^{-r} that satisfy $\int_{\mathbb{R}^d} p_t = 1$ (since we are interested in probability distributions, recall Remark 1.3), so we will rather consider an affine space. It will not pose any problem, since then $(p_t - \rho, m_t)$ is an element of $\{(v, m) \in \mathbf{H}_{\theta}^{-r} : \int_{\mathbb{R}^d} v = 0\}$ which is a Banach space.

Following the notations of [5] we set

$$\psi(t) := (\rho, \alpha_t^{\delta}), \ t \in \mathbb{R}/T\delta\mathbb{Z}. \tag{1.43}$$

With this notation we have $\widetilde{\mathcal{M}}^{\delta} = \psi(\mathbb{R}/T\delta\mathbb{Z})$. We will consider the projections $\widetilde{\Pi}_u^{\delta,s}$ and $\widetilde{\Pi}_u^{\delta,c}$ defined for $(p,m) \in \mathbf{H}_{\theta}^{-r}$ by

$$\widetilde{\Pi}_{u}^{\delta,s}(p,m) = (p, P_{u}^{\delta,s}m), \qquad \widetilde{\Pi}_{u}^{\delta,c}(p,m) = (0, P_{u}^{\delta,c}m), \tag{1.44}$$

where $P_t^{\delta,s}$ and $P_t^{\delta,c}$ are the projections defined in Section 1.4. The subspaces $\widetilde{\mathbf{X}}_u^{\delta,c} = \widetilde{\Pi}_u^{\delta,c}(\mathbf{H}_{\theta}^{-r})$ and $\widetilde{\mathbf{X}}_u^{\delta,s} = \widetilde{\Pi}_u^{\delta,s}(\mathbf{H}_{\theta}^{-r})$ will correspond to the approximately tangent space and stable space of $\widetilde{\mathcal{M}}^{\delta}$. It is clear that for each $t \in [0, \frac{T}{\delta})$ we have

$$\mathbf{H}_{a}^{-r} = \widetilde{\mathbf{X}}_{t}^{\delta,c} \oplus \widetilde{\mathbf{X}}_{t}^{\delta,s}. \tag{1.45}$$

Consider τ such that

$$e^{-\lambda_{\alpha}\tau} \leqslant \frac{c_{\alpha}}{8C_{\alpha}},$$
 (1.46)

where c_{α} , C_{α} , λ_{α} are given by (1.38). The following proposition states that $\widetilde{\mathcal{M}}^{\delta}$ satisfies the hypotheses given in [5], making it an approximately invariant and approximately normally hyperbolic manifold.

Proposition 1.7. There exists $\delta_0 > 0$ such that for r_0 given in Proposition 1.1 and for all $r \ge r_0$, $\delta \in (0, \delta_0)$ and $\theta \in (0, 1)$, the following assertions are true.

(1) (Definition 2.1. in [5]) There exists a positive constant κ_1 such that for all $u \in \mathbb{R}/\frac{T}{3}\mathbb{Z}$,

$$\left\| T^{\frac{\tau}{\delta}}(\rho, \alpha_u) - (\rho, \alpha_{u + \frac{\tau}{\delta}}) \right\|_{\mathbf{H}_{\rho}^{-r}} \leqslant \kappa_1 \delta. \tag{1.47}$$

(2) (Hypothesis (H2) in [5]) There exist positive constants $\kappa_2, \kappa_3, \kappa_4$ such that for all $s, t \in \mathbb{R}/\frac{T}{\delta}\mathbb{Z}$ such that $|s-u| \leq 1$, $|t-u| \leq 1$, and $\iota = s, c$,

$$\left\| \widetilde{\Pi}_{u}^{\delta,\iota} \right\|_{\mathcal{B}(\mathbf{H}_{o}^{-r})} \leqslant \kappa_{2}, \qquad \left\| \widetilde{\Pi}_{u}^{\delta,\iota} - \widetilde{\Pi}_{s}^{\delta,\iota} \right\|_{\mathcal{B}(\mathbf{H}_{o}^{-r})} \leqslant \kappa_{3} \left\| \psi(t) - \psi(s) \right\|_{\mathbf{H}_{\theta}^{-r}}, \tag{1.48}$$

and

$$\frac{\left\|\psi(t) - \psi(s) - \widetilde{\Pi}_s^{\delta,c}(\psi(t) - \psi(s))\right\|_{\mathbf{H}_{\theta}^{-r}}}{\left\|\psi(t) - \psi(s)\right\|_{\mathbf{H}_{\theta}^{-r}}} \leqslant \kappa_4 \delta. \tag{1.49}$$

(3) (Hypothesis H3 in [5]) There exists a positive constant κ_5 such that for all $u \in \mathbb{R}/\frac{T}{\delta}\mathbb{Z}$,

$$\max \left\{ \left\| \widetilde{\Pi}_{u+\frac{\tau}{\delta}}^{\delta,c} D T^{\frac{\tau}{\delta}}(\rho, \alpha_u)_{|\widetilde{\mathbf{X}}_{u}^{\delta,s}|} \right\|_{\mathcal{B}(\mathbf{H}_{\theta}^{-r})}, \left\| \widetilde{\Pi}_{u+\frac{\tau}{\delta}}^{\delta,s} D T^{\frac{\tau}{\delta}}(\rho, \alpha_u)_{|\widetilde{\mathbf{X}}_{u}^{\delta,c}|} \right\|_{\mathcal{B}(\mathbf{H}_{\theta}^{-r})} \right\} \leqslant \kappa_5 \delta. \quad (1.50)$$

(4) (Hypothesis H3' and C3 in [5]) There exist $a \in (0,1)$ and $\widetilde{\lambda} > 0$ such that for all $u \in \mathbb{R}/\frac{T}{\delta}\mathbb{Z}$,

$$\left\| \left(\widetilde{\Pi}_{u+\frac{\tau}{\delta}}^{\delta,c} D T^{\frac{\tau}{\delta}}(\rho, \alpha_u)_{|\widetilde{\mathbf{X}}_u^{\delta,c}} \right)^{-1} \right\|_{\mathcal{B}(\mathbf{H}_a^{-r})}^{-1} > a, \tag{1.51}$$

and

$$\left\| \widetilde{\Pi}_{u+\frac{\tau}{\delta}}^{\delta,s} D T^{\frac{\tau}{\delta}}(\rho, \alpha_u)_{|\widetilde{\mathbf{X}}_{u}^{\delta,s}|} \right\|_{\mathcal{B}(\mathbf{H}_{\theta}^{-r})} \leqslant \widetilde{\lambda} \min \left(1, \left\| \left(\widetilde{\Pi}_{u+\frac{\tau}{\delta}}^{\delta,c} D T^{\frac{\tau}{\delta}}(\rho, \alpha_u)_{|\widetilde{\mathbf{X}}_{u}^{\delta,c}} \right)^{-1} \right\|_{\mathcal{B}(\mathbf{H}_{\theta}^{-r})}^{-1} \right), \tag{1.52}$$

(5) (Hypothesis H4 in [5]) There exist positive constants κ_6 and κ_7 such that

$$\left\| DT^{\frac{\tau}{\delta}} |_{\mathcal{V}(\widetilde{\mathcal{M}}^{\delta}, 1)} \right\|_{\mathcal{B}(\mathbf{H}_{\theta}^{-r})} \leqslant \kappa_{6}, \quad \left\| D^{2} T^{\frac{\tau}{\delta}} |_{\mathcal{V}(\widetilde{\mathcal{M}}^{\delta}, 1)} \right\|_{\mathcal{BL}\left((\mathbf{H}_{\theta}^{-r})^{2}, \mathbf{H}_{\theta}^{-r}\right)} \leqslant \kappa_{7}. \tag{1.53}$$

where $\mathcal{V}(\widetilde{\mathcal{M}}^{\delta}, R_0)$ denote the R_0 -neighborhood of $\widetilde{\mathcal{M}}^{\delta}$.

(6) (Hypothesis H5 in [5]) For any $\varepsilon > 0$ there exists $\zeta > 0$ such that for all $\mu = (p,m) \in \mathcal{V}(\widetilde{\mathcal{M}}^{\delta},1)$ and $t \in [\frac{\tau}{\delta},\frac{\tau}{\delta}+\zeta]$,

$$\left\| T^{t}(\mu) - T^{\frac{\tau}{\delta}}(\mu) \right\|_{\mathbf{H}_{\theta}^{-r}} \leqslant \varepsilon. \tag{1.54}$$

The first five items of Proposition 1.7 focus on properties of the semi-group $\left(T^{n\frac{\tau}{\delta}}\right)_{n\geqslant 0}$ discretized in time, showing that $\widetilde{\mathcal{M}}^{\delta}$ is an approximately invariant manifold approximately normally hyperbolic for this semi-group, while the last item is an uniform in time bound that implies that this property is also true for the semi-group $\left(T^{t}\right)_{t\geqslant 0}$. More precisely (1)

shows that $\widetilde{\mathcal{M}}^{\delta}$ is approximately invariant for the discrete semi- group, (2) shows that $\widetilde{\mathbf{X}}_{u}^{\delta,c}$ is an approximation of the tangent space to $\widetilde{\mathcal{M}}^{\delta}$ at (ρ, α_{u}) and that ψ does not twist too much, (3) implies that $\widetilde{\mathbf{X}}^{\delta,c}$ and $\widetilde{\mathbf{X}}^{\delta,s}$ are approximately invariant under $\left(DT^{n\frac{\tau}{\delta}}\right)_{n\geqslant 0}$, and (4) implies that $\left(DT^{n\frac{\tau}{\delta}}\right)_{n\geqslant 0}$ contracts more in the direction $\widetilde{\mathbf{X}}^{\delta,s}$ than in the direction $\widetilde{\mathbf{X}}^{\delta,c}$, while it does not contract too much in the direction $\widetilde{\mathbf{X}}^{\delta,c}$. (5) is a technical assumption useful in their proof.

Remark that we do not quote here the hypothesis (H1) of [5] in this Proposition, since it is simply (1.45). Moreover in [5] the authors treat first the inflowing invariant case, and then the overflowing invariant case, while we are here interested in an actual invariant manifold (both inflowing and overflowing), which is why we mix hypotheses (Hi) and (C3), as it is done in Theorem 6.5 of [5].

1.6. Structure of the paper. The proof of Theorem 1.2 concerning the well-posedness of (1.4) is carried out in Section 2. Proposition 1.7 is proven in Section 3. The main result of existence of periodic solutions (Theorem 1.4) is proven in Section 4. The question of regularity of the isochron is addressed in Section 5. The Appendix A gathers technical estimates on the Ornstein-Uhlenbeck operator and some Grönwall type lemmas are listed in Appendix B.

2. Proof of Theorem 1.2

We give in this section the existence, uniqueness and regularity result of Theorem 1.2. We rely here on classical arguments one can find for example in [36] or [23].

Proof of Theorem 1.2. We first remark that $G: \mathbf{H}_{\theta}^{-r} \to \mathbf{H}_{\theta}^{-(r+1)}$ is locally Lispchitz. Indeed, for any $(p,m) \in \mathbf{H}_{\theta}^{-r}$ and any $(\varphi,\psi) \in \mathbf{H}_{\theta}^{r+1}$,

$$\langle \langle G(p,m), (\varphi,\psi) \rangle \rangle = -\langle p(F_m - \int F_m p), \nabla \varphi \rangle + \psi \cdot \int F_m p.$$
 (2.1)

We have $|\int F_m p| \leq ||F_m||_{H^r_{\theta}} ||p||_{\mathbf{H}^{-r}_{\theta}}$, and due to the fact that all derivatives of F are bounded, $||F_m||_{H^r_{\theta}} \leq C_F$ independently from m. Moreover, due to the same reason, we have $||F_m \cdot \nabla \varphi||_{H^r_{\theta}} \leq C_F ||\nabla \varphi||_{H^r_{\theta}}$ independently from m. This means that

$$\left| \left\langle \left\langle G(p,m), (\varphi,\psi) \right\rangle \right\rangle \right| \leqslant C \|p\|_{\mathbf{H}_{\theta}^{-r}} \left(1 + \|p\|_{\mathbf{H}_{\theta}^{-r}} \right) \|\varphi\|_{H_{\theta}^{r+1}} + C \|p\|_{\mathbf{H}_{\theta}^{-r}} |\psi|. \tag{2.2}$$

We deduce $||G(\nu)||_{\mathbf{H}_{\theta}^{-(\mathbf{r}+1)}} \leq C||\nu||_{\mathbf{H}_{\theta}^{-\mathbf{r}}} \left(1 + ||\nu||_{\mathbf{H}_{\theta}^{-\mathbf{r}}}\right)$, and thus that G is locally Lipschitz. Remark that when p is a probability distribution $|\int F_m p| \leq C_F |\int p| \leq C_F$, and in this case G is in fact Globally Lipschitz.

Now, since the operator \mathcal{L} is sectorial in H_{θ}^{-r} , it also the case of the operator $\widetilde{\mathcal{L}}$ in \mathbf{H}_{θ}^{-r} defined by $\widetilde{\mathcal{L}}(p,m) = \mathcal{L}p$, and thus, applying [36, Theorem 47.8], for all initial condition $\mu = (p,m) \in \mathbf{H}_{\theta}^{-r}$ there exists a unique maximal mild solution $\mu_t := (p_t, m_t) = T^t(\mu)$ to (1.26) defined on some time interval $[0, t_c)$ and which satisfies $t \mapsto T^t(\mu) \in \mathcal{C}\left([0, t_c); \mathbf{H}_{\theta}^{-r}\right)$.

Now, for $\mu = (p, m)$ and $\nu = (\eta, n)$, the Frechet differential

$$DG(\mu)[\nu] = \begin{pmatrix} DG_1(\mu)[\nu] \\ DG_2(\mu)[\nu] \end{pmatrix}$$

$$= \begin{pmatrix} -\nabla \cdot \left(\eta \left(F_m - \int F_m p \right) \right) - \nabla \cdot \left(p \left(DF_m[n] - \int F_m \eta - \int DF_m[n] p \right) \right) \\ \int F_m \eta + \int DF_m[n] p \end{pmatrix} (2.3)$$

satisfies, by similar arguments as above (in particular the fact that the derivatives of F_m can be bounded independently from m)

$$||DG(\mu)[\nu]||_{\mathbf{H}_{\theta}^{-(r+1)}} \le C \left(1 + ||\mu||_{\mathbf{H}_{\theta}^{-r}}\right) ||\nu||_{\mathbf{H}_{\theta}^{-r}},$$
 (2.4)

and by [36, Theorem 49.2], $\mu \mapsto T^t(\mu)$ is Frechet differentiable, with derivative $DT^t(\mu)[\nu] = \nu_t := (\eta_t, n_t)$ the unique mild solution to

$$\begin{cases}
\partial_t \eta_t &= \mathcal{L} \eta_t + \delta D G_1(\mu_t) [\nu_t] \\
\dot{n}_t &= \delta D G_2(\mu_t) [\nu_t]
\end{cases}$$
(2.5)

By [36, Theorem 47.5] the solution $\nu_t = (\eta_t, n_t)$ to (2.5) depends continuously on $\mu = (p, m)$, so that the flow $T^t(\mu)$ is C^1 . One can proceed similarly for the second derivative. We have this time, for $\nu_i = (\eta_i, n_i)$, i = 1, 2,

$$D^{2}G_{1}(\mu)[\nu_{1},\nu_{2}] = -\nabla \cdot \left(\eta_{1} \left(DF_{m}[n_{2}] - \int F_{m}\eta_{2} - \int DF_{m}[n_{2}]p\right)\right)$$

$$-\nabla \cdot \left(\eta_{2} \left(DF_{m}[n_{1}] - \int F_{m}\eta_{1} - \int DF_{m}[n_{1}]p\right)\right)$$

$$-\nabla \cdot \left(p\left(D^{2}F_{m}[n_{1},n_{2}] - \int DF_{m}[n_{1}]\eta_{2} - \int DF_{m}[n_{2}]\eta_{1}\right)$$

$$-\int D^{2}F_{m}[n_{1},n_{2}]p\right), \qquad (2.6)$$

and

$$D^{2}G_{2}(\mu)[\nu_{1},\nu_{2}] = \int DF_{m}[n_{1}]\eta_{2} + \int DF_{m}[n_{2}]\eta_{1} + \int D^{2}F_{m}[n_{1},n_{2}]p, \qquad (2.7)$$

so that

$$||D^{2}G(\mu)[\nu_{1},\nu_{2}]||_{\mathbf{H}_{\theta}^{-(r+1)}} \leq C \left(1 + ||\mu||_{\mathbf{H}_{\theta}^{-r}}\right) ||\nu_{1}||_{\mathbf{H}_{\theta}^{-r}} ||\nu_{2}||_{\mathbf{H}_{\theta}^{-r}}, \tag{2.8}$$

and $T^{t}(\mu)$ is C^{2} with $D^{2}T^{t}(\mu)[\nu_{1},\nu_{2}]=\xi_{t}=(\xi_{t}^{1},\xi_{t}^{2})$ where $\xi_{0}=0$ and

$$\partial_t \xi_t = (\mathcal{L}\xi_t^1, 0) + \delta DG(\mu_t)[\xi_t] + \delta D^2 G(\mu_t)[\nu_{1,t}, \nu_{2,t}], \tag{2.9}$$

where $\nu_{i,t} = DT^{t}(\mu_{0})[\nu_{i}]$ for i = 1, 2.

To prove that, for R > 0 and $||p_0 - \rho||_{H_{\theta}^{-r}} \leq R$, this solution is in fact globally defined when δ is taken small enough, remark that it satisfies

$$p_t = e^{t\mathcal{L}} p_0 + \int_0^t e^{(t-s)\mathcal{L}} \nabla \cdot (p_s(\delta F_{m_s} + \dot{m}_s)) \, \mathrm{d}s, \qquad (2.10)$$

and

$$\dot{m}_t = \delta \langle F_{m_t}, p_t \rangle. \tag{2.11}$$

The estimates obtained above imply directly $|\dot{m}_s| \leq \delta C_F ||p_s||_{H_{\theta}^{-r}}$. Using Proposition 1.1 we get:

$$||p_t||_{H_{\theta}^{-r}} \leqslant C_{\mathcal{L}}||p_0||_{H_{\theta}^{-r}} + C_1 \int_0^t \frac{e^{-\lambda(t-s)}}{\sqrt{t-s}} ||p_s(\delta F_{m_s} + \dot{m}_s)||_{H_{\theta}^{-r}} ds$$
(2.12)

$$\leqslant C_2 \left(\|p_0\|_{H_{\theta}^{-r}} + \delta \int_0^t \frac{e^{-\lambda(t-s)}}{\sqrt{t-s}} \|p_s\|_{H_{\theta}^{-r}} \left(1 + \|p_s\|_{H_{\theta}^{-r}} \right) ds \right).$$
 (2.13)

Denote $t_0 = \inf \left\{ t > 0 : \|p_t\|_{H_{\theta}^{-r}} \geqslant 2C_2 \left(R + \|\rho\|_{H_{\theta}^{-r}} \right) \right\}$. By continuity, $t_0 > 0$ and for all $t \in [0, t_0]$,

$$||p_t||_{H_{\theta}^{-r}} \leqslant C_2 \left(R + ||\rho||_{H_{\theta}^{-r}} \right) + \delta \sqrt{\frac{\pi}{\lambda}} 2C_2 \left(R + ||\rho||_{H_{\theta}^{-r}} \right) \left(1 + 2C_2 \left(R + ||\rho||_{H_{\theta}^{-r}} \right) \right). \tag{2.14}$$

For the choice of $\delta > 0$ sufficiently small such that $\delta \sqrt{\frac{\pi}{\lambda}} 2 \left(1 + 2C_2 \left(R + \|\rho\|_{H_{\theta}^{-r}} \right) \right) < 1$, this yields that $t_0 = \infty$, so that (p_t, m_t) is a global solution.

3. Proof of Proposition 1.7

In this section we give the proof of Proposition 1.7 which shows that $\widetilde{\mathcal{M}}^{\delta}$ is an approximately invariant approximately normally hyperbolic manifold. We do not prove the assertions in the order they are given in Proposition 1.7.

Proof of Proposition 1.7.

Proof of (1). Take $p_0 = \rho$ and $m_0 = \alpha_u$. We then have

$$p_t - \rho = \int_0^t e^{(t-s)\mathcal{L}} \nabla \cdot (p_s(\delta F_{m_s} + \dot{m}_s)) \, \mathrm{d}s, \tag{3.1}$$

and

$$\dot{m}_t - \dot{\alpha}_{u+t} = \delta \langle F_{m_t}, p_t \rangle - \delta \langle F_{\alpha_{u+t}}, \rho \rangle. \tag{3.2}$$

As it was already proved in the preceding section, we have $|\dot{m}_s| \leq C_F \delta ||p_s||_{H_{\theta}^{-r}}$, and since Theorem 1.2 with R=1 implies that, choosing δ small enough, $||p_t||_{H_{\theta}^{-r}} \leq C(1)$, we get from Proposition 1.1,

$$||p_{t} - \rho||_{H_{\theta}^{-r}} \leq C_{1} \int_{0}^{t} \frac{e^{-\lambda(t-s)}}{\sqrt{t-s}} ||p_{s}(\delta F_{m_{s}} + \dot{m}_{s})||_{H_{\theta}^{-r}} ds$$

$$\leq C_{1} \delta \int_{0}^{t} \frac{e^{-\lambda(t-s)}}{\sqrt{t-s}} ||p_{s}||_{H_{\theta}^{-r}} \left(1 + ||p_{s}||_{H_{\theta}^{-r}}\right) ds \leq C_{2} \delta. \tag{3.3}$$

Now since

$$\frac{1}{\delta}(\dot{m}_t - \dot{\alpha}_{u+t}) = \langle DF_{\alpha_{u+t}}, \rho \rangle (m_t - \alpha_{u+t}) + \langle F_{m_t} - F_{\alpha_{u+t}} - DF_{\alpha_{u+t}} (m_t - \alpha_{u+t}), \rho \rangle
+ \langle F_{m_t}, p_t - \rho \rangle, \quad (3.4)$$

we have the following mild representation (recall that $m_0 = \alpha_u$):

$$m_t - \alpha_{u+t} = \delta \int_0^t \pi_{u+t,u+s}^{\delta} \left(\langle F_{m_s} - F_{\alpha_{u+s}} - DF_{\alpha_{u+s}} (m_s - \alpha_{u+s}), \rho \rangle + \langle F_{m_s}, p_s - \rho \rangle \right) ds, \quad (3.5)$$

which leads to (recall that the derivatives of F are bounded and that (3.3) is valid for all $t \ge 0$):

$$|m_t - \alpha_{u+t}| \le C_3 \delta \int_0^t |m_s - \alpha_{u+s}|^2 ds + C_3 \delta^2 t.$$
 (3.6)

Consider $t_1 = \inf\{t > 0 : |m_t - \alpha_{u+t}| \ge 2\tau C_4\delta\}$ (recall the definition of τ in (1.46)). By continuity, $t_1 > 0$ and for all $t \le t_1$ we have

$$|m_t - \alpha_{u+t}| \le (4\tau^2 C_3^3 \delta^3 + C_3 \delta^2)t,$$
 (3.7)

which means that $t_1 \geqslant \frac{\tau}{\delta}$ for δ small enough, and implies (1).

Proof of (2). The two first points follow directly form the fact that the projections P_u^c are smooth. For the third point we have

$$\frac{\left\|\psi(t) - \psi(s) - \widetilde{\Pi}_s^{\delta,c}(\psi(t) - \psi(s))\right\|_{\mathbf{H}_{\theta}^{-r}}}{\left\|\psi(t) - \psi(s)\right\|_{\mathbf{H}_{\theta}^{-r}}} = \frac{\left|\alpha_t^{\delta} - \alpha_s^{\delta} - P_s^{\delta,c}(\alpha_t^{\delta} - \alpha_s^{\delta})\right|}{\left|\alpha_t^{\delta} - \alpha_s^{\delta}\right|},\tag{3.8}$$

and since

$$\alpha_t^{\delta} - \alpha_s^{\delta} = \alpha_{\delta t}^1 - \alpha_{\delta s}^1 = \delta(t - s) \frac{d}{du} \alpha_{u|u = \delta s}^1 + O(\delta^2(t - s)), \tag{3.9}$$

and

$$P_s^{\delta,c} \frac{d}{du} \alpha_{u|u=\delta s}^1 = P_{\delta s}^{0,c} \frac{d}{du} \alpha_{u|u=\delta s}^1 = \frac{d}{du} \alpha_{u|u=\delta s}^1, \tag{3.10}$$

the term $\frac{\left\|\psi(t)-\psi(s)-\widetilde{\Pi}_{s}^{\delta,c}(\psi(t)-\psi(s))\right\|_{\mathbf{H}_{\theta}^{-r}}}{\left\|\psi(t)-\psi(s)\right\|_{\mathbf{H}_{\theta}^{-r}}} \text{ is indeed of order } \delta.$

Proof of (5). We choose in the following $R_0 = 1$. For any $(p, m) \in \mathcal{V}(\widetilde{\mathcal{M}}^{\delta}, R_0)$, wich means in particular $||p - \rho||_{H_{\theta}^{-r}} \leq R_0$, we deduce from Theorem 1.2, if δ is small enough, that

$$\sup_{t \ge 0} \|p_t\|_{H_{\theta}^{-r}} \le C(R_0). \tag{3.11}$$

This means in particular, since $m_t = m_0 + \delta \int_0^t \langle F_{m_s}, p_s \rangle ds$, that for $C_4, C_5 > 0$

$$\sup_{t \geqslant 0} |\dot{m}_t| \leqslant \delta C_4, \quad \text{and} \quad \sup_{t \in [0, \frac{\tau}{\Delta}]} |m_t| \leqslant C_5, \tag{3.12}$$

where C_5 depends on τ . Now, using (2.5) we have, with $\mu_s = (p_s, m_s)$,

$$\eta_t = e^{t\mathcal{L}}\eta_0 + \delta \int_0^t e^{(t-s)\mathcal{L}} DG_1(\mu_s)[\eta_s, n_s] \,\mathrm{d}s, \tag{3.13}$$

and

$$n_t = n_0 + \delta \int_0^t DG_2(\mu_s)[\eta_s, n_s] ds.$$
 (3.14)

From (2.4) and Proposition 1.1 (recall that $\int_{\mathbb{R}^d} \eta_0 = 0$, see Remark 1.3), we obtain

$$\|\eta_t\|_{H_{\theta}^{-r}} \leqslant C_{\mathcal{L}} e^{-\lambda t} \|\eta_0\|_{H_{\theta}^{-r}} + C_6 \delta \int_0^t \frac{e^{-\lambda(t-s)}}{\sqrt{t-s}} \left(\|\eta_s\|_{H_{\theta}^{-r}} + |n_s| \right) ds, \tag{3.15}$$

and

$$|n_t| \le |n_0| + C_6 \delta \int_0 \left(\|\eta_s\|_{H_{\theta}^{-r}} + |n_s| \right) ds.$$
 (3.16)

We deduce that, for $\nu_t = DT^t(p, m)[\nu_0] = (\eta_t, n_t)$,

$$\|\nu_t\|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_7 \|\nu_0\|_{\mathbf{H}_{\theta}^{-r}} + C_8 \delta \int_0^t \left(1 + \frac{1}{\sqrt{t-s}}\right) \|\nu_s\|_{\mathbf{H}_{\theta}^{-r}} \, \mathrm{d}s. \tag{3.17}$$

Applying Lemma B.1, we get the desired bound for the $DT^{\frac{\tau}{\delta}}$ with $\kappa_6 = 2C_7 e^{3C_8\tau}$, when δ is small enough.

For the second derivative, recall that $D^2T^t(\mu)[\nu_1,\nu_2] = \xi_t = (\xi_t^1,\xi_t^2)$, where $\xi_0 = 0$ and (recall (2.9))

$$\xi_t^1 = \delta \int_0^t e^{(t-s)\mathcal{L}} \left(DG_1(\mu_S)[\xi_s] + D^2 G_1(\mu_s)[\nu_{1,s}, \nu_{2,s}] \right) ds, \tag{3.18}$$

and

$$\xi_t^2 = \delta \int_0^t \left(DG_2(\mu_S)[\xi_s] + D^2 G_2(\mu_s)[\nu_{1,s}, \nu_{2,s}] \right) ds$$
 (3.19)

where $\mu_t = (p_t, m_t)$, and $\nu_{i,t} = DT^t(\mu_0)[\nu_i]$ for i = 1, 2. This induces for $t \in [0, \frac{\tau}{\delta}]$, recalling (2.4), (2.8) and since $\|\nu_{i,t}\|_{\mathbf{H}_{\theta}^{-r}} \leqslant \kappa_6 \|\nu_{i,0}\|_{\mathbf{H}_{\theta}^{-r}}$,

$$\|\xi_t^1\|_{H_{\theta}^{-r}} \leq \delta C_9 \int_0^t \frac{e^{-\lambda(t-s)}}{\sqrt{t-s}} \left(\|\xi_s\|_{\mathbf{H}_{\theta}^{-r}} + \|\nu_{1,0}\|_{\mathbf{H}_{\theta}^{-r}} \|\nu_{2,0}\|_{\mathbf{H}_{\theta}^{-r}} \right) ds, \tag{3.20}$$

and

$$\left|\xi_{t}^{2}\right| \leq \delta C_{9} \int_{0}^{t} \left(\left\|\xi_{s}\right\|_{\mathbf{H}_{\theta}^{-r}} + \left\|\nu_{1,0}\right\|_{\mathbf{H}_{\theta}^{-r}} \left\|\nu_{2,0}\right\|_{\mathbf{H}_{\theta}^{-r}}\right) ds.$$
 (3.21)

So for $t \leqslant \frac{\tau}{\delta}$,

$$\|\xi_t\|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_{10} \|\nu_{1,0}\|_{\mathbf{H}_{\theta}^{-r}} \|\nu_{2,0}\|_{\mathbf{H}_{\theta}^{-r}} + \delta C_{10} \int_0^t \left(1 + \frac{e^{-\lambda(t-s)}}{\sqrt{t-s}}\right) \|\xi_s\|_{\mathbf{H}_{\theta}^{-r}} \, \mathrm{d}s, \qquad (3.22)$$

and one deduces from Lemma B.1 that $\|\xi_t\|_{\mathbf{H}_{\theta}^{-r}} \leqslant \kappa_7 \|\nu_{1,0}\|_{\mathbf{H}_{\theta}^{-r}} \|\nu_{2,0}\|_{\mathbf{H}_{\theta}^{-r}}$ with $\kappa_7 = 2C_{10}e^{3C_{10}\tau}$ for $t \leqslant \frac{\tau}{\delta}$ and δ small enough, which concludes the proof of (5).

Proof of (3). We are now interested in $DT^{\frac{\tau}{\delta}}(\rho,\alpha_u)(\eta_0,n_0) = (\eta_{\frac{\tau}{\delta}},n_{\frac{\tau}{\delta}}) = \nu_{\frac{\tau}{\delta}}$. From the proof of point (3) we already know that $\sup_{t\in[0,\frac{\tau}{\delta}]}\|\nu_t\|_{\mathbf{H}_{\theta}^{-r}} \leqslant \kappa_6\|\nu_0\|_{\mathbf{H}_{\theta}^{-r}}$, which means, recalling (3.15), that

$$\|\eta_{t}\|_{H_{\theta}^{-r}} \leq C_{\mathcal{L}} e^{-\lambda t} \|\eta_{0}\|_{H_{\theta}^{-r}} + C_{11} \delta \int_{0}^{t} \frac{e^{-\lambda(t-s)}}{\sqrt{t-s}} \left(\|\eta_{0}\|_{H_{\theta}^{-r}} + |n_{0}| \right) ds$$

$$\leq C_{\mathcal{L}} e^{-\lambda t} \|\eta_{0}\|_{H_{\theta}^{-r}} + C_{12} \delta \left(\|\eta_{0}\|_{H_{\theta}^{-r}} + |n_{0}| \right). \tag{3.23}$$

Moreover, since

$$\frac{1}{\delta}\dot{n}_{t} = \langle DF_{\alpha_{u+t}}\left[n_{t}\right], \rho \rangle - \langle DF_{\alpha_{u+t}}\left[n_{t}\right] - DF_{m_{t}}\left[n_{t}\right], \rho \rangle + \langle DF_{m_{t}}\left[n_{t}\right], p_{t} - \rho \rangle + \langle F_{m_{t}}, \eta_{t} \rangle, \quad (3.24)$$

we have the mild representation

$$n_{t} = \pi_{u+t,u}^{\delta} n_{0} + \delta \int_{0}^{t} \pi_{u+t,u+s}^{\delta} \left(-\langle DF_{\alpha_{u+s}} [n_{s}] - DF_{m_{s}} [n_{s}], \rho \rangle + \langle DF_{m_{s}} [n_{s}], p_{s} - \rho \rangle + \langle F_{m_{s}}, \eta_{s} \rangle \right) ds. \quad (3.25)$$

From the proof of point (1), for $t \leq \frac{\tau}{\delta}$, $||p_t - \rho||_{H_{\theta}^{-r}}$ and $|m_t - \alpha_{u+t}|$ are of order δ , and thus we obtain (recall also that $\sup_{t \in [0,\frac{\tau}{\delta}]} |n_t| \leq \kappa_6 ||\nu_0||_{\mathbf{H}_{\theta}^{-r}}$):

$$\left| n_{t} - \pi_{u+t,u}^{\delta} n_{0} \right| \leq C_{13} \delta \int_{0}^{t} \left(\|\eta_{s}\|_{H_{\theta}^{-r}} + \delta |n_{0}| \right) ds$$

$$\leq C_{13} \delta \int_{0}^{t} \left(C_{\mathcal{L}} e^{-\lambda s} \|\eta_{0}\|_{H_{\theta}^{-r}} + C_{12} \delta \left(\|\eta_{0}\|_{H_{\theta}^{-r}} + |n_{0}| \right) + \delta |n_{0}| \right) ds$$

$$\leq C_{14} \delta \left(\|\eta_{0}\|_{H_{\theta}^{-r}} + |n_{0}| \right). \tag{3.26}$$

Suppose now that $(\eta_0, n_0) \in \widetilde{\mathbf{X}}_u^{\delta,s}$, that is $P_u^{\delta,c} n_0 = 0$. Then we have $P_{u+\frac{\tau}{\delta}}^{\delta,c} \pi_{u+\frac{\tau}{\delta},u}^{\delta} n_0 = P_u^{\delta,c} n_0 = 0$, and thus, recalling (3.26) and (3.23),

$$\left| P_{u + \frac{\tau}{\delta}}^{\delta, c} n_{\frac{\tau}{\delta}} \right| = \left| P_{u + \frac{\tau}{\delta}}^{\delta, c} \left(n_{\frac{\tau}{\delta}} - \pi_{u + \frac{\tau}{\delta}, u}^{\delta} n_0 \right) \right| \leqslant C_{15} \delta \left(\| \eta_0 \|_{H_{\theta}^{-r}} + |n_0| \right). \tag{3.27}$$

This shows that

$$\left\| \widetilde{\Pi}_{u + \frac{\tau}{\delta}}^{\delta, c} D T^{\frac{\tau}{\delta}}(\rho, \alpha_u) |_{\widetilde{\mathbf{X}}_u^{\delta, s}} \right\|_{\mathcal{B}(\mathbf{H}_{\rho}^{-r})} \leqslant C_{15} \delta. \tag{3.28}$$

On the other hand, suppose that $(\eta_0, n_0) \in \widetilde{\mathbf{X}}_u^{\delta,c}$, that is $\eta_0 = 0$ and $P_u^{\delta,s} n_0 = 0$. We then have directly $\left\| \eta_{\frac{\tau}{\delta}} \right\|_{H_{\theta}^{-r}} \leqslant C_{12} \delta |n_0|$, and since $P_{u+\frac{\tau}{\delta}}^{\delta,s} \pi_{u+\frac{\tau}{\delta},u}^{\delta} n_0 = P_u^{\delta,s} n_0 = 0$, from (3.26) we deduce

$$\left| P_{u+\frac{\tau}{\delta}}^s n_{\frac{\tau}{\delta}} \right| = \left| P_{u+\frac{\tau}{\delta}}^s \left(n_{\frac{\tau}{\delta}} - \pi_{u+\frac{\tau}{\delta},u}^{\delta} n_0 \right) \right| \leqslant C_{16} \delta^2 \int_0^{\frac{\tau}{\delta}} |n_0| \, \mathrm{d}s \leqslant C_{16} \tau \delta |n_0|. \tag{3.29}$$

This means that

$$\left\| \widetilde{\Pi}_{u + \frac{\tau}{\delta}}^{\delta, s} D T^{\frac{\tau}{\delta}}(\rho, \alpha_u) \right|_{\widetilde{\mathbf{X}}_u^{\delta, c}} \right\|_{\mathcal{B}(\mathbf{H}_{\theta}^{-r})} \leqslant (C_{12} + C_{16}\delta)\delta. \tag{3.30}$$

Proof of (4). On one hand consider $(\eta_0, n_0) \in \widetilde{\mathbf{X}}_u^{\delta, s}$, that is $P_u^{\delta, c} n_0 = 0$. Then, considering δ small enough such that $C_{\mathcal{L}} e^{-\lambda \frac{\tau}{\delta}} \leqslant C_{12} \delta$, by (3.23) we obtain

$$\left\| \eta_{\frac{\tau}{\delta}} \right\|_{H_{\theta}^{-r}} \le 2C_{12}\delta \left(\|\eta_0\|_{H_{\theta}^{-r}} + |n_0| \right).$$
 (3.31)

Moreover, since $P_{u+\frac{\tau}{\delta}}^{\delta,s}\pi_{u+\frac{\tau}{\delta},u}^{\delta}n_0=\pi_{u+\frac{\tau}{\delta},u}^{\delta}n_0$ and $P_u^{\delta,c}n_0=0$ we obtain, by (3.26) and (1.33),

$$\left| P_{u+\frac{\tau}{\delta}}^s n_{\frac{\tau}{\delta}} \right| \leqslant \left| P_{u+\frac{\tau}{\delta}}^s \pi_{u+\frac{\tau}{\delta},u}^{\delta} n_0 \right| + \left| P_{u+\frac{\tau}{\delta}}^s \left(n_{\frac{\tau}{\delta}} - \pi_{u+\frac{\tau}{\delta},u}^{\delta} n_0 \right) \right| \tag{3.32}$$

$$\leq C_{\alpha} e^{-\lambda_{\alpha} \tau} |n_0| + C_{17} \delta \left(\|\eta_0\|_{H_{\theta}^{-r}} + |n_0| \right).$$
 (3.33)

We deduce that for δ small enough

$$\left\| \widetilde{\Pi}_{u + \frac{\tau}{\delta}}^{\delta, s} D T^{\frac{\tau}{\delta}}(\rho, \alpha_u) |_{\widetilde{\mathbf{X}}_u^{\delta, s}} \right\|_{\mathcal{B}(\mathbf{H}_{\rho}^{-r})} \leq 2C_{\alpha} e^{-\lambda_{\alpha} \tau}. \tag{3.34}$$

On the other hand consider $(\eta_0, n_0) \in \widetilde{\mathbf{X}}_u^{\delta,c}$, which means $\eta_0 = 0$ and $P_u^{\delta,s} n_0 = 0$. Then similar arguments as above (recall that this time $\eta_0 = 0$) lead to

$$\left| P_{u+\frac{\tau}{\delta}}^{\delta,c} \left(n_{\frac{\tau}{\delta}} - \pi_{u+\frac{\tau}{\delta},u}^{\delta} n_0 \right) \right| \leqslant C_{18} \delta |n_0|. \tag{3.35}$$

We then obtain, for δ small enough, recalling (1.33),

$$\left| P_{u+\frac{\tau}{\delta}}^{c} n_{\frac{\tau}{\delta}} \right| \geqslant \left(c_{\alpha} - C_{18} \delta \right) |n_{0}| \geqslant \frac{c_{\alpha}}{2} |n_{0}|. \tag{3.36}$$

This means in particular that $\widetilde{\Pi}_{u+\frac{\tau}{\delta}}^{\delta,c}DT^{\frac{\tau}{\delta}}(\rho,\alpha_u)|_{\widetilde{\mathbf{X}}_u^{\delta,c}}$, which is a linear mapping in finite dimensional spaces, is invertible and satisfies

$$\left\| \left(\widetilde{\Pi}_{u+\frac{\tau}{\delta}}^{\delta,c} D T^{\frac{\tau}{\delta}}(\rho, \alpha_u) |_{\widetilde{\mathbf{X}}_u^{\delta,c}} \right)^{-1} \right\|_{\mathcal{B}(\mathbf{H}_{\sigma}^{-r})} \leqslant \frac{2}{c_{\alpha}}. \tag{3.37}$$

We deduce (4) with $a = \frac{c_a}{4}$ and $\tilde{\lambda} = \frac{4C_{\alpha}e^{-\lambda_{\alpha}\tau}}{c_{\alpha}}$, recalling (1.46).

Proof of (6). For any initial condition $\mu = (p_0, m_0) \in \mathcal{V}(\widetilde{\mathcal{M}}^{\delta}, 1)$ recall that Theorem 1.2 implies $\sup_{t \geq 0} \|p_t\|_{H_{\theta}^{-r}} \leq C(1)$. Then for $\frac{\tau}{\delta} \leq t < t', t' - t \leq \zeta$, for some $\zeta \leq 1$ to be chosen later, relying on (3.1), the following is true:

$$||p_{t'} - p_t||_{H_{\theta}^{-r}} \leq ||\left(e^{t'\mathcal{L}} - e^{t\mathcal{L}}\right) p_0||_{H_{\theta}^{-r}} + \int_0^t ||\left(e^{(t'-s)\mathcal{L}} - e^{(t-s)\mathcal{L}}\right) \nabla \cdot (p_s(\delta F_{m_s} + \dot{m}_s))||_{H_{\theta}^{-r}} ds + \int_t^{t'} ||e^{(t'-s)\mathcal{L}} \nabla \cdot (p_s(\delta F_{m_s} + \dot{m}_s))||_{H_{\theta}^{-r}} ds$$
(3.38)

Using Proposition 1.1, the first term above may be bounded as

$$\left\| \left(e^{t'\mathcal{L}} - e^{t\mathcal{L}} \right) p_0 \right\|_{H_{\theta}^{-r}} \leqslant C_{\mathcal{L}} (t' - t)^{\varepsilon} \frac{e^{-\lambda t}}{t^{\frac{1}{2} + \varepsilon}} \left\| p_0 \right\|_{H_{\theta}^{-(r+1)}} \leqslant C_{19} \zeta^{\varepsilon} \delta^{\frac{1}{2} + \varepsilon} \frac{e^{-\lambda \frac{\tau}{\delta}}}{\tau^{\frac{1}{2} + \varepsilon}}. \tag{3.39}$$

Concerning the second term,

$$\int_{0}^{t} \left\| \left(e^{(t'-s)\mathcal{L}} - e^{(t-s)\mathcal{L}} \right) \nabla \cdot \left(p_{s} (\delta F_{m_{s}} + \dot{m}_{s}) \right) \right\|_{H_{\theta}^{-r}} \mathrm{d}s$$

$$\leqslant C_{\mathcal{L}} (t'-t)^{\varepsilon} \int_{0}^{t} \frac{e^{-\lambda(t-s)}}{(t-s)^{\frac{1}{2}+\varepsilon}} \left\| p_{s} (\delta F_{m_{s}} + \dot{m}_{s}) \right\|_{H_{\theta}^{-r}} \mathrm{d}s$$

$$\leqslant C_{20} \delta (t'-t)^{\varepsilon} \int_{0}^{t} \frac{e^{-\lambda(t-s)}}{(t-s)^{\frac{1}{2}+\varepsilon}} \left\| p_{s} \right\|_{H_{\theta}^{-r}} \left(1 + \left\| p_{s} \right\|_{H_{\theta}^{-r}} \right) \mathrm{d}s$$

$$\leqslant C_{21} \delta \zeta^{\varepsilon}. \tag{3.40}$$

Now turning to the third term, relying again on Proposition 1.1,

$$\int_{t}^{t'} \left\| e^{(t'-s)\mathcal{L}} \nabla \cdot (p_{s}(\delta F_{m_{s}} + \dot{m}_{s})) \right\|_{H_{\theta}^{-r}} ds$$

$$\leq C_{22}\delta \int_{t}^{t'} \frac{e^{-\lambda(t'-s)}}{\sqrt{t'-s}} \left\| p_{s} \right\|_{H_{\theta}^{-r}} \left(1 + \left\| p_{s} \right\|_{H_{\theta}^{-r}} \right) ds,$$

$$\leq C_{23}\delta \zeta^{\frac{1}{2}}. \tag{3.41}$$

Gathering (3.39), (3.40), (3.41) into (3.38) yields

$$\|p_{t'} - p_t\|_{H_{\theta}^{-r}} \leqslant \frac{\xi}{2} \tag{3.42}$$

if $\zeta \leq 1$ is chosen sufficiently small. We now turn turn to the control of the mean: since $\dot{m}_t = \delta \int F_{m_t} dp_t$ we have that for $t \leq t' \leq t + \zeta$,

$$m_{t'} - m_t = \delta \int_t^{t'} \langle F_{m_s}, p_s \rangle \, \mathrm{d}s.$$

Since we have the uniform bound $\sup_{s\geq 0} \|p_s\|_{H^{-r}_{\theta}} \leq C(1)$ and since F and its derivatives are bounded, the above quantity is easily bounded by some $C\delta(t'-t)$ which can be made smaller than $\xi/2$, provided ζ is taken small enough.

4. Proof of Theorem 1.4

Proof of Theorem 1.4. From Proposition 1.7 we know that the hypotheses needed in [5] are satisfied for δ small enough, which means that the system (1.4) admits a stable normally hyperbolic manifold \mathcal{M}^{δ} that is at distance δ from $\widetilde{\mathcal{M}}^{\delta}$. Indeed in [5] some constants η, χ, σ need to be small for their result to be true, but in our case these constants are of order δ , so we only need to suppose δ small enough. Moreover \mathcal{M}^{δ} is contructed at a distance δ_0 from $\widetilde{\mathcal{M}}^{\delta}$, with (see [5], Theorem 4.2) δ_0 chosen such that η/ε and ε/δ_0 are bounded for some $\varepsilon > 0$. Since η is in our case of order δ , we can take δ_0 of order δ , and \mathcal{M}^{δ} is indeed at distance δ from $\widetilde{\mathcal{M}}^{\delta}$.

The invariant manifold \mathcal{M}^{δ} is one dimensional, since $\widetilde{\mathcal{M}}^{\delta}$ is, so to prove that it corresponds to a periodic solution it is sufficient to prove that it does not possess any invariant point. But for any $(p_0, m_0) \in \mathcal{M}^{\delta}$ we have, since $\|p - \rho\|_{H_{\theta}^{-r}}$ and $|m_0 - \alpha_u^{\delta}|$ are of order δ for some $u \in [0, \frac{T_{\alpha}}{\delta}]$,

$$\dot{m}_0 = \delta \int F_{m_0} p_0 = \delta \int F_{\alpha_u} \rho + O(\delta^2) = \dot{\alpha}_u + O(\delta^2). \tag{4.1}$$

Since there exists c > 0 such that $|\dot{\alpha}_u^{\delta}/\delta| > c$ independently from u, we have $\dot{m}_0 \neq 0$ for the solutions starting from any point of \mathcal{M}^{δ} , which means that \mathcal{M}^{δ} does not possess any fixed-point, and is thus defined by a periodic solution of positive period T_{δ} , that we denote $\Gamma_t^{\delta} = (q_t^{\delta}, \gamma_t^{\delta})$ for $t \in [0, T_{\delta}]$.

Now, by the Herculean Theorem (see [36], Theorem 47.6), since \mathcal{M}^{δ} is invariant, Γ_t^{δ} is in fact an element of $\mathbf{H}_{\theta}^{-r+2}$ and, by [36] Theorem 48.5, $\partial_t \Gamma_{s+t}^{\delta} = (\partial_t q_{s+t}^{\delta}, \dot{\gamma}_{s+t}^{\delta})$ is in $C([0, T_{\delta}), \mathbf{H}_{\theta}^{-r})$ and is is solution to

$$\begin{cases}
\partial_t \eta_t &= \mathcal{L} \eta_t + \delta D G_1(\Gamma_{s+t}^{\delta})[\nu_t] \\
\dot{n}_t &= \delta D G_2(\Gamma_{s+t}^{\delta})[\nu_t]
\end{cases} ,$$
(4.2)

which means in particular that $\partial_t \Gamma_{s+t}^{\delta} = \Phi_{s+t,s} \partial_t \Gamma_s^{\delta}$. Now $\partial_t \Gamma_{s+t}^{\delta}$ is a periodic solution to (4.2), and the same arguments induce that $\partial_t^2 \Gamma_{s+t}^{\delta}$ is in $C([0, T_{\delta}), \mathbf{H}_{\theta}^{-r})$.

In addition, it is proved in [5] that \mathcal{M}^{δ} is foliated by C^1 invariant foliations: a neighborhood \mathcal{W}^{δ} of \mathcal{M}^{δ} satisfies the decomposition $\mathcal{W}^{\delta} = \bigcup_{s \in [0,T_{\delta})} \mathcal{W}^{\delta}_{s}$, where \mathcal{W}^{δ}_{s} corresponds to the elements of $\mu \in \mathbf{H}^{-r}_{\theta}$ such that $T^{nT_{\delta}}(\mu)$ converges exponentially fast to Γ^{δ}_{s} as n goes to infinity. The projections $\Pi^{\delta,c}_{s}$ and $\Pi^{\delta,s}_{s}$ correspond then respectively to the projections on the tangent space to \mathcal{M}^{δ} and to \mathcal{W}^{δ}_{s} at Γ^{δ}_{s} . The linear operator $\Phi^{\delta}_{s+t,s} = DT^{t}(\Gamma^{\delta}_{s})$ commutes then with these projections, and is bounded from above and below in the direction of the tangent space to \mathcal{M}^{δ} , while it is contractive in the direction of the tangent space to stable foliations.

In addition to the contractive property, the regularization effect of Φ^{δ} given in (1.38) is a consequence of the fact that $\Phi_{t+s,s}\nu = \nu_t$ where $\nu_0 = \nu$ and $\nu_t = (\eta_t, n_t)$ is solution to

$$\begin{cases}
\partial_t \eta_t &= \mathcal{L} \eta_t + \delta D G_1(\Gamma_{s+t}^{\delta})[\nu_t] \\
\dot{n}_t &= \delta D G_2(\Gamma_{s+t}^{\delta})[\nu_t]
\end{cases}$$
(4.3)

The operator $\widetilde{\mathcal{L}}(\eta,n)=(\mathcal{L}\eta,0)$ is sectorial in \mathbf{H}_{θ}^{-r} and thus induces regularization properties for the solutions to (4.3), and thus for Φ^{δ} . More precisely we are in fact exactly in the situation of [23], Theorem 7.2.3 and the following remark. Indeed, for $s\in[0,T_{\delta})$ we can define the operator $U_s^{\delta}=\Phi_{s+T_{\delta},s}^{\delta}$, and we can deduce from above spectral properties for U_t^{δ} . Since Γ^{δ} is a periodic solution U_s^{δ} admits 1 as eigenvalue, with eigenfunction $\partial_s\Gamma_s^{\delta}$ and corresponding projection $\Pi_s^{\delta,c}$, and due to the contractive properties Φ^{δ} the rest of the spectrum of U_s^{δ} is located in a disk centered at 0 with radius $e^{-\lambda_{\delta}T_{\delta}}$. We can then apply Theorem 7.2.3 and the following remark to obtain (1.38) (reducing slightly the value of λ_{δ}).

The C^1 regularity of $s \mapsto \Pi_s^{\delta,c}$ is not a direct consequence the normally hyperbolic results of [5] (they prove that \mathcal{W}_s^{δ} has a Hölder regularity with respect to s), but since we are in the case of a periodic solution we have an explicit formula for $\Pi_s^{\delta,c}$: 1 is an isolated eigenvalue of U_t^{δ} , so for $\mathcal{C}_{\varepsilon}$ the circle centered at 1 with radius $\varepsilon > 0$, with ε small enough, we have

$$\Pi_s^{\delta,c} = \frac{1}{2i\pi} \int_{\mathcal{C}_{\varepsilon}} (\lambda - U_s^{\delta})^{-1} d\lambda.$$
 (4.4)

But applying [23], Theorem 3.4.4., $t \mapsto U_s^{\delta}$ is C^1 , with $\partial_s U_s^{\delta} \zeta = \zeta_{T_{\delta}} = (\zeta_{T_{\delta}}^1, \zeta_{T_{\delta}}^2)$, where $\zeta_0 = \zeta$ and

$$\begin{cases}
\partial_t \zeta_t^1 &= \mathcal{L}\zeta_t^1 + \delta DG_1(\Gamma_{s+t}^{\delta})[\zeta_t] + \delta D^2 G_1(\Gamma_{s+t}^{\delta})[\partial_t \Gamma_{s+t}^{\delta}, \zeta_t] \\
\dot{\zeta}_t^2 &= \delta DG_2(\Gamma_{s+t}^{\delta})[\zeta_t] + \delta D^2 G_2(\Gamma_{s+t}^{\delta})[\partial_t \Gamma_{s+t}^{\delta}, \zeta_t].
\end{cases} , (4.5)$$

and thus $s \mapsto \Pi_s^{\delta,c}$ is also C^1 .

It is not immediate that q_s^{δ} is a probability distribution, since we apply the results of [5] considering solutions $p_t \in H_{\theta}^{-r}$ satisfying $\int_{\mathbb{R}^d} p_t = 1$ but without any hypotheses on non negativeness. However, $\widetilde{\mathcal{M}}^{\delta}$ is in the basin of attraction of \mathcal{M}^{δ} , so any $(q_s^{\delta}, m_s^{\delta}) \in \mathcal{M}^{\delta}$ is the limit in \mathbf{H}_{θ}^{-r} of $(p_t, m_t) = T^t(\rho, \alpha_u^{\delta})$ for some $u \in [0, \frac{T_{\alpha}}{\delta})$. So, since in this case p_t is a probability distribution (recall that it is the probability distribution of $X_t - \mathbb{E}[X_t]$, where X_t satisfies (1.2) with initial distribution ρ), we deduce that $\langle q_s^{\delta}, \varphi \rangle \geqslant 0$ for any smooth function φ with compact support, and thus q_s^{δ} is also a probability distribution. \square

5. Proof of Theorem 1.6

As it was already explained in Section 1.4, the existence of the map Θ^{δ} is a consequence of the foliation property proved in [5]. Θ^{δ} satisfies moreover the relation

$$\Gamma^{\delta}_{\Theta(\mu)} = \lim_{n \to \infty} T^{nT_{\delta}} \mu. \tag{5.1}$$

Our aim in the present section is to prove the C^2 regularity of Θ^{δ} . Following ideas of [22], we will prove uniform in time bounds for the first and second derivatives of the flow T^t , which will induce the regularity of

$$S(\mu) := \lim_{n \to \infty} T^{nT_{\delta}} \mu \tag{5.2}$$

and thus the regularity of Θ^{δ} .

Proof of Theorem 1.6.

Step 1: Let us first show that for some constant $c_1 > 0$

$$\sup_{t \geqslant 0} \sup_{\mu \in \mathcal{V}(\mathcal{M}^{\delta}, \varepsilon)} \|DT^{t}(\mu)\|_{\mathcal{B}(\mathbf{H}_{\theta}^{-r})} \leqslant c_{1}$$
(5.3)

where $\mathcal{V}(\mathcal{M}, \varepsilon) := \left\{ \mu \in \mathbf{H}_{\theta}^{-r}, \operatorname{dist}_{\mathbf{H}_{\theta}^{-r}}(\mu, \mathcal{M}) < \varepsilon \right\}$ is a neighborhood of \mathcal{M} on which the trajectories are attracted to the cycle. For $\mu_0 = (p_0, m_0) \in \mathcal{V}(\mathcal{M}, \varepsilon)$ and $u = \Theta(\mu_0)$, we have, denoting by $\nu_t = (\eta_t, n_t) = DT^t(\mu_0)[\nu_0]$,

$$\nu_t = \Phi_{u+t,u}^{\delta} \nu_0 + \delta \int_0^t \Phi_{u+t,u+s}^{\delta} \left(DG(\mu_s) - DG(\Gamma_{u+s}^{\delta}) \right) \left[\nu_s \right] ds. \tag{5.4}$$

Let us now prove that there exists a constant C_G such that, for $\mu = (p, m)$ and $\Gamma = (q, \gamma)$,

$$||DG(\mu) - DG(\Gamma)||_{\mathcal{B}\left(\mathbf{H}_{\theta}^{-r}, \mathbf{H}_{\theta}^{-(r+1)}\right)} \leqslant C_G ||\mu - \Gamma||_{\mathbf{H}_{\theta}^{-r}}.$$

$$(5.5)$$

We have, for $\nu = (\eta, n)$

$$(DG_1(\mu) - DG_1(\Gamma))[\nu] = -\nabla \cdot (\eta(F_m - F_\gamma)) + \nabla \cdot \left(\eta\left(\int F_m p - \int F_\gamma q\right)\right)$$
 (5.6)

$$-\nabla \cdot (pDF_m[n] - qDF_{\gamma}[n]) \tag{5.7}$$

$$+ \nabla \cdot \left(p \int F_m \eta - q \int F_\gamma \eta \right) \tag{5.8}$$

$$+ \nabla \cdot \left(p \int DF_m[n]p - q \int DF_{\gamma}[n]q \right), \tag{5.9}$$

and

$$(DG_2(\mu) - DG_2(\Gamma))[\nu] = \int (F_m - F_\gamma)\eta + \int DF_m[n]p - \int DF_\gamma[n]q.$$
 (5.10)

For the first term, we obtain

$$\|\nabla \cdot (\eta(F_m - F_\gamma))\|_{H_{\theta}^{-(r+1)}} \leqslant C_1 \|\eta(F_m - F_\gamma)\|_{H_{\theta}^{-r}}, \tag{5.11}$$

and since, for $f \in H_{\theta}^r$,

$$\langle \eta(F_m - F_\gamma), f \rangle \le \|\eta\|_{H_{\theta}^{-r}} \|(F_m - F_\gamma)f\|_{H_{\theta}^r} \le C_2 |m - \gamma| \|\eta\|_{H_{\theta}^{-r}} \|f\|_{H_{\theta}^r},$$
 (5.12)

where we have used the fact that all the derivatives of F are Lipschitz, we get, for some $C_3 > 0$,

$$\|\nabla \cdot (\eta(F_m - F_\gamma))\|_{H_{\theta}^{-(r+1)}} \leqslant C_3 |m - \gamma| \|\eta\|_{H_{\theta}^{-r}}.$$
(5.13)

For the second term, since

$$\left| \int F_m p - \int F_\gamma q \right| \leqslant \left| \int F_m (p - q) \right| + \left| \int (F_m - F_\gamma) q \right| \leqslant C_4 \left(\|p - q\|_{H_{\theta}^{-r}} + |m - \gamma| \right), \tag{5.14}$$

we have

$$\left\| \nabla \cdot \left(\eta \left(\int F_m p - \int F_{\gamma} q \right) \right) \right\|_{H_{\theta}^{-(r+1)}} \leq C_5 \left(\|p - q\|_{H_{\theta}^{-r}} + |m - \gamma| \right). \tag{5.15}$$

The other terms can be tackled in a similar way. Now, since $\mu_0 \in \mathcal{W}_u^{\delta}$, we have for some $C_{\Gamma^{\delta}} > 0$,

$$\left\| \mu_s - \Gamma_{u+s}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_{\Gamma^{\delta}} e^{-\lambda_{\delta} s} \left\| \mu_0 - \Gamma_u^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}}, \tag{5.16}$$

and from the estimates obtained above, we deduce

$$\|\nu_t\|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_6 \|\nu_0\|_{\mathbf{H}_{\theta}^{-r}} + C_6 \delta \int_0^t \left(1 + (t-s)^{-\frac{1}{2}} e^{-\lambda_\delta(t-s)}\right) e^{-\lambda_\delta s} \|\nu_s\|_{\mathbf{H}_{\theta}^{-r}} \, \mathrm{d}s. \tag{5.17}$$

Applying Lemma B.2 for $\phi(u) = u^{-\frac{1}{2}}e^{-\lambda_{\delta}u}$, we obtain from (B.3) that

$$\sup_{t \ge 0} \|\nu_t\|_{\mathbf{H}_{\theta}^{-r}} \le c_1 \|\nu_0\|_{\mathbf{H}_{\theta}^{-r}} \tag{5.18}$$

for some $c_1 > 0$.

Step 2: let us now show that $\left(DT^{nT_{\delta}}\right)_{n \geq 0}$ in the space $C\left(\mathcal{V}(\mathcal{L}^{\delta}, \varepsilon), \mathcal{B}\left(\mathbf{H}_{\theta}^{-r}\right)\right)$ is a Cauchy sequence, which implies that $\mu \mapsto S(\mu)$ is C^{1} (recall (5.2)).

For $n \ge m$ we have

$$\nu_{nT_{\delta}} - \nu_{mT_{\delta}} = \left(\Phi_{u+nT_{\delta},u}^{\delta} - \Phi_{u+mT_{\delta},u}^{\delta}\right)\nu_{0}
+ \delta \int_{mT_{\delta}}^{nT_{\delta}} \Phi_{u+nT_{\delta},u+s}^{\delta} \left(DG(\mu_{s}) - DG(\Gamma_{u+s}^{\delta})\right) \left[\nu_{s}\right] ds
+ \delta \int_{0}^{mT_{\delta}} \left(\Phi_{u+nT_{\delta},u+s}^{\delta} - \Phi_{u+mT_{\delta},u+s}^{\delta}\right) \left(DG(\mu_{s}) - DG(\Gamma_{u+s}^{\delta})\right) \left[\nu_{s}\right] ds.$$
(5.19)

For the first term, we get

$$\|(\Phi_{u+nT_{\delta},u} - \Phi_{u+mT_{\delta},u}) \nu_{0}\|_{\mathbf{H}_{\theta}^{-r}} = \|(\Phi_{u+nT_{\delta},u} - \Phi_{u+mT_{\delta},u}) \Pi_{\delta,u} \nu_{0}\|_{\mathbf{H}_{\theta}^{-r}}$$

$$\leq C_{7} e^{-\lambda_{\delta} m T_{\delta}} \|\nu_{0}\|_{\mathbf{H}_{\theta}^{-r}}.$$
(5.20)

For the second one, using (5.18),

$$\left\| \int_{mT_{\delta}}^{nT_{\delta}} \Phi_{u+nT_{\delta},u+s}^{\delta} \left(DG(\mu_{s}) - DG(\Gamma_{u+s}^{\delta}) \right) \left[\nu_{s} \right] ds \right\|_{\mathbf{H}_{\theta}^{-r}}$$

$$\leq C_{8} \left\| \mu - \Gamma_{u}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} \int_{mT_{\delta}}^{nT_{\delta}} \left(1 + (nT_{\delta} - s)^{-\frac{1}{2}} e^{-\lambda_{\delta}(nT_{\delta} - s)} \right) e^{-\lambda_{\delta} s} ds$$

$$= \frac{C_{8} \left\| \mu - \Gamma_{u}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} e^{-\lambda_{\delta} mT_{\delta}} \left(1 + e^{-\lambda_{\delta}(n-m)T_{\delta}} \left(2\lambda_{\delta} \sqrt{(n-m)T_{\delta}} - 1 \right) \right)$$

$$\leq C_{9} \left\| \mu - \Gamma_{u}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} e^{-\lambda_{\delta} mT_{\delta}}. \tag{5.21}$$

For the last term, remark first that

$$\Phi_{u+nT_{\delta},u+s}^{\delta} - \Phi_{u+mT_{\delta},u+s}^{\delta} = \left(\Phi_{u+nT_{\delta},u+mT}^{\delta} - I_d\right) \prod_{u+mT_{\delta}}^{\delta,s} \Phi_{u+mT_{\delta},u+s}^{\delta}, \tag{5.22}$$

so that, using again (5.18),

$$\left\| \int_{0}^{mT_{\delta}} \left(\Phi_{u+nT_{\delta},u+s}^{\delta} - \Phi_{u+mT_{\delta},u+s}^{\delta} \right) \left(DG(\mu_{s}) - DG(\Gamma_{u+s}) \right) \left[\nu_{s} \right] ds \right\|_{\mathbf{H}_{\theta}^{-r}}$$

$$\leq C_{10} \left\| \mu - \Gamma_{u}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} \int_{0}^{mT_{\delta}} (mT_{\delta} - s)^{-\frac{1}{2}} e^{-\lambda_{\delta}(mT_{\delta} - s)} e^{-\lambda_{\delta} s} ds$$

$$= 2C_{10} \left\| \mu - \Gamma_{u}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} \sqrt{mT_{\delta}} e^{-\lambda_{\delta} mT_{\delta}}. \tag{5.23}$$

Since the constants above are uniform in $\mu \in \mathcal{V}$, we deduce that $(DT^{nT_{\delta}})_{n \geq 0}$ is indeed a Cauchy sequence. Thus S is C^1 with $DS(\mu) = \lim_{n \to \infty} DT^{nT_{\delta}}(\mu)$.

Before moving to the second derivative, let us have a closer look at DS. We have

$$\left\| \Pi_{u+nT_{\delta}}^{\delta,s} \nu_{nT_{\delta}} \right\|_{\mathbf{H}_{\theta}^{-r}} \leq \left\| \Pi_{u+nT_{\delta}}^{\delta,s} \Phi_{u+nT_{\delta},u}^{\delta} \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} + \left\| \int_{0}^{nT_{\delta}} \Pi_{u+nT_{\delta}}^{\delta,s} \Phi_{u+nT_{\delta},u+s}^{\delta} \left(DG(\mu_{s}) - DG(\Gamma_{u+s}^{\delta}) \right) [\nu_{s}] ds \right\|_{\mathbf{H}_{\theta}^{-r}},$$

$$(5.24)$$

and we can bound the right hand side in three steps. Firstly,

$$\left\| \Pi_{u+nT_{\delta}}^{\delta,s} \Phi_{u+nT_{\delta},u}^{\delta} \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_{\Phi,\delta} e^{-\lambda_{\delta} nT_{\delta}} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}}. \tag{5.25}$$

Secondly, since $\sup_{\mu \in \mathcal{V}} \|DG(\mu)\|_{\mathcal{B}\left(\mathbf{H}_{\theta}^{-r}, \mathbf{H}_{\theta}^{-(r+1)}\right)} \leqslant C_G$,

$$\left\| \int_{0}^{\frac{nT_{\delta}}{2}} \Pi_{u+nT_{\delta}}^{\delta} \Phi_{u+nT_{\delta},u+s}^{\delta} \left(DG(\mu_{s}) - DG(\Gamma_{u+s}^{\delta}) \right) \left[\nu_{s} \right] ds \right\|_{\mathbf{H}_{\theta}^{-r}}$$

$$\leq C_{11} \left\| \mu - \Gamma_{u}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} \int_{0}^{\frac{nT_{\delta}}{2}} (nT_{\delta} - s)^{-\frac{1}{2}} e^{-\lambda_{\delta}(nT_{\delta} - s)} ds$$

$$\leq C_{12} \left\| \mu - \Gamma_{u}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} n^{\frac{1}{2}} e^{-\frac{\lambda_{\delta}nT_{\delta}}{2}} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}}. \tag{5.26}$$

Thirdly, by similar arguments as above (replacing mT_{δ} with $n\frac{T_{\delta}}{2}$),

$$\left\| \int_{\frac{nT_{\delta}}{2}}^{nT_{\delta}} \Phi_{u+nT_{\delta},u+s}^{\delta} \left(DG(\mu_{s}) - DG(\Gamma_{u+s}^{\delta}) \right) \left[\nu_{s} \right] ds \right\|_{\mathbf{H}_{\theta}^{-r}}$$

$$\leq C_{13} \left\| \mu - \Gamma_{u}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} e^{-\lambda_{\delta} n \frac{T_{\delta}}{2}} \|\nu_{0}\|_{\mathbf{H}_{\theta}^{-r}}. \quad (5.27)$$

We deduce that $\Pi_{\Theta(\mu)}DS(\mu) = 0$, so that DS has rank 1 and thus there exists a family of linear forms $l_{\mu} \in \mathcal{B}\left(\mathbf{H}_{\theta}^{-r}, \mathbb{R}\right)$ (that depend continuously on μ) such that, for $u = \Theta(\mu)$,

$$DS(\mu)[\nu] = l_{\mu}[\nu]\partial_{u}\Gamma_{u}, \qquad (5.28)$$

and we have proved, for $\nu_t = DT^t(\mu)[\nu_0]$,

$$\|\nu_{nT_{\delta}} - l_{\mu}[\nu_{0}]\partial_{u}\Gamma_{u}\|_{\mathbf{H}_{a}^{-r}} \leqslant C_{13}n^{\frac{1}{2}}e^{-\lambda_{\delta}n^{\frac{T_{\delta}}{2}}}\|\nu_{0}\|_{\mathbf{H}_{a}^{-r}}.$$
(5.29)

With similar computations one can in fact show that

$$\|\nu_t - l_{\mu}[\nu_0] \partial_u \Gamma_{u+t} \|_{\mathbf{H}_{\theta}^{-r}} \leq C_{14} t^{\frac{1}{2}} e^{-\lambda_{\delta} \frac{t}{2}} \|\nu_0\|_{\mathbf{H}_{\theta}^{-r}}.$$
 (5.30)

In the case of $\mu = \Gamma_u^{\delta}$, we deduce in particular that

$$DS(\Gamma_u^{\delta}) = \Pi_u^{\delta,c}. \tag{5.31}$$

In fact, we have proved a more precise estimate: if $\nu_t^2 = DT^t(\mu)[\nu_0]$, $\nu_t^1 = DT^t(\Gamma_u^{\delta})[\nu_0]$ with $u = \Theta(\mu)$, the estimates above lead to

$$\left\| \nu_t^2 - \nu_t^1 - \left(l_{\mu}[\nu_0] - l_{\Gamma_u^{\delta}}[\nu_0] \right) \partial_u \Gamma_{u+t} \right\|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_{15} \left\| \mu - \Gamma_u^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} t^{\frac{1}{2}} e^{-\lambda_{\delta} \frac{t}{2}} \left\| \nu_0 \right\|_{\mathbf{H}_{\theta}^{-r}}. \tag{5.32}$$

Step 3: let us now show that for a constant $c_2 > 0$,

$$\sup_{t \geqslant 0} \sup_{\mu \in \mathcal{V}(\Gamma^{\delta}, \varepsilon)} \|D^2 T^t(\mu)\|_{\mathcal{BL}(\mathbf{H}_{\theta}^{-r})} \leqslant c_2.$$
 (5.33)

From (2.9), we deduce, for $\xi_t = D^2 T^t(\mu)[\nu, w]$, the following mild formulation (recall that $\xi_0 = 0$):

$$\xi_t = \delta \int_0^t \Phi_{u+t,u+s}^{\delta} \left(D^2 G(\mu_s) [\nu_s, w_s] + \left(D G(\mu_s) - D G(\Gamma_{u+s}^{\delta}) \right) \xi_s \right) ds, \tag{5.34}$$

where $\nu_t = DT^t(\mu_0)[\nu]$, $w_t = DT^t(\mu_0)[w]$. With similar arguments as above, we obtain

$$\left\| \int_{0}^{t} \Phi_{u+t,u+s}^{\delta} \left(D^{2} G(\mu_{s}) [\nu_{s}, w_{s}] - D^{2} G(\Gamma_{u+s}^{\delta}) [\nu_{s}, w_{s}] \right) ds \right\|_{\mathbf{H}_{\theta}^{-r}}$$

$$\leq C_{16} \int_{0}^{t} \left(1 + (t-s)^{-\frac{1}{2}} e^{-\lambda_{\delta}(t-s)} \right) \left\| \mu_{s} - \Gamma_{u+s}^{\delta} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| \nu_{s} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| w_{s} \right\|_{\mathbf{H}_{\theta}^{-r}} ds$$

$$\leq C_{16} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| w_{0} \right\|_{\mathbf{H}_{\theta}^{-r}}.$$

$$(5.35)$$

Remark now that

$$\partial_t^2 \Gamma_{u+t}^{\delta} = (\mathcal{L}\partial_t q_{u+t}^{\delta}, 0) + \delta DG(\Gamma_{u+t}^{\delta})[\partial_t \Gamma_{u+t}], \tag{5.36}$$

and

$$\partial_t^3 \Gamma_{u+t}^{\delta} = \left(\mathcal{L} \partial_t^2 q_{u+t}^{\delta}, 0 \right) + \delta DG(\Gamma_{u+t}^{\delta}) \left[\partial_t^2 \Gamma_{u+t}^{\delta} \right] + \delta D^2 G(\Gamma_{u+t}^{\delta}) \left[\partial_t \Gamma_{u+t}^{\delta}, \partial_t \Gamma_{u+t}^{\delta} \right], \tag{5.37}$$

and thus

$$\partial_t^2 \Gamma_{u+t}^{\delta} = \Phi_{u+t,u}^{\delta} \partial_t^2 \Gamma_u^{\delta} + \delta \int_0^t \Phi_{u+t,u+s}^{\delta} D^2 G(\Gamma_{u+s}^{\delta}) [\partial_s \Gamma_{u+s}^{\delta}, \partial_s \Gamma_{u+s}^{\delta}] ds.$$
 (5.38)

So, in particular, since

$$\Pi_u^{\delta,c} \Phi_{u+T_{\delta},u}^{\delta} \partial_u^2 \Gamma_u = \Pi_u^{\delta,c} (\partial_u^2 \Gamma_u), \tag{5.39}$$

we deduce from (5.38) that

$$\Pi_{u}^{\delta,c} \left(\int_{0}^{T_{\delta}} \Phi_{u+T_{\delta},u+s}^{\delta} D^{2} G(\Gamma_{u+s}^{\delta}) [\partial_{s} \Gamma_{u+s}^{\delta}, \partial_{s} \Gamma_{u+s}^{\delta}] ds \right) = 0.$$
 (5.40)

Now, recalling (5.30),

$$\|\nu_t - l_{\mu}[\nu_0] \partial_u \Gamma_{u+t} \|_{\mathbf{H}_a^{-r}} \leqslant C_{14} t^{\frac{1}{2}} e^{-\lambda_{\delta} \frac{t}{2}} \|\nu_0\|_{\mathbf{H}_a^{-r}}, \tag{5.41}$$

$$\|w_t - l_{\mu}[w_0] \partial_u \Gamma_{u+t} \|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_{14} t^{\frac{1}{2}} e^{-\lambda_{\delta} \frac{t}{2}} \|w_0\|_{\mathbf{H}_{\theta}^{-r}},$$
 (5.42)

and we deduce

$$\left\| \int_{0}^{t} \Phi_{u+t,u+s} D^{2} G(\mu_{s}) [\nu_{s}, w_{s}] ds - l_{\mu} [\nu_{0}] l_{\mu} [w_{0}] \int_{0}^{t} \Phi_{u+t,u+s} D^{2} G(\mu_{s}) [\partial_{u} \Gamma_{u+s}, \partial_{u} \Gamma_{u+s}] ds \right\|_{\mathbf{H}_{\theta}^{-r}} \\ \leqslant C_{17} \|\nu_{0}\|_{\mathbf{H}_{\theta}^{-r}} \|w_{0}\|_{\mathbf{H}_{\theta}^{-r}}. \quad (5.43)$$

So, recalling (5.40), and since

$$\left\| \Pi_{u+t}^{\delta,s} \int_{0}^{t} \Phi_{u+t,u+s}^{\delta} D^{2} G(\mu_{s}) [\partial_{u} \Gamma_{u+s}^{\delta}, \partial_{u} \Gamma_{u+s}^{\delta}] \, \mathrm{d}s \right\|_{\mathbf{H}_{\theta}^{-r}}$$

$$\leq C_{18} \int_{0}^{t} (t-s)^{-\frac{1}{2}} e^{-\lambda_{\delta}(t-s)} ds \leq C_{19}, \quad (5.44)$$

we deduce, coming back to (5.34), that

$$\|\xi_{t}\|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_{19} \|\nu_{0}\|_{\mathbf{H}_{\theta}^{-r}} \|w_{0}\|_{\mathbf{H}_{\theta}^{-r}} + \delta \left\| \int_{0}^{t} \Phi_{u+t,u+s}^{\delta} \left(DG(\mu_{s}) - DG(\Gamma_{u+s}^{\delta}) \right) \xi_{s} ds \right\|_{\mathbf{H}_{\theta}^{-r}}.$$
(5.45)

Relying again on (5.16), we deduce that, for some $c_2 > 0$,

$$\|\xi_t\|_{\mathbf{H}_a^{-r}} \leqslant c_2 \|\nu_0\|_{\mathbf{H}_a^{-r}} \|w_0\|_{\mathbf{H}_a^{-r}},$$
 (5.46)

which implies (5.33).

Step 4: let us now prove that $(D^2T^{nT_\delta})_{n\geq 0}$ in the space $C\left(\mathcal{V}(\mathcal{M}^\delta,\varepsilon),\mathcal{BL}\left(\mathbf{H}_{\theta}^{-r}\right)\right)$ is a Cauchy sequence, which implies that $\mu\mapsto S(\mu)$ is C^2 .

We have, for $n \ge m$,

$$\xi_{nT_{\delta}} - \xi_{mT_{\delta}} = \int_{0}^{nT_{\delta}} \Phi_{u+nT_{\delta},u+s}^{\delta} D^{2}G(\Gamma_{u+s}^{\delta})[\nu_{s}, w_{s}] \, \mathrm{d}s \qquad (5.47)$$

$$- \int_{0}^{mT_{\delta}} \Phi_{u+mT_{\delta},u+s}^{\delta} D^{2}G(\Gamma_{u+s}^{\delta})[\nu_{s}, w_{s}] \, \mathrm{d}s$$

$$+ \int_{mT_{\delta}}^{nT_{\delta}} \Phi_{u+nT_{\delta},u+s}^{\delta} \left(D^{2}G(\mu_{s})[\nu_{s}, w_{s}] - D^{2}G(\Gamma_{u+s}^{\delta})[\nu_{s}, w_{s}]\right) \, \mathrm{d}s$$

$$+ \int_{0}^{mT_{\delta}} \left(\Phi_{u+nT_{\delta},u+s}^{\delta} - \Phi_{u+mT_{\delta},u+s}^{\delta}\right)$$

$$\times \left(D^{2}G(\mu_{s})[\nu_{s}, w_{s}] - D^{2}G(\Gamma_{u+s}^{\delta})[\nu_{s}, w_{s}]\right) \, \mathrm{d}s$$

$$+ \int_{mT_{\delta}}^{nT_{\delta}} \Phi_{u+nT_{\delta},u+s}^{\delta} \left(DG(\mu_{s}) - DG(\Gamma_{u+s}^{\delta})\right) \xi_{s} \, \mathrm{d}s$$

$$+ \int_{0}^{nT_{\delta}} \left(\Phi_{u+nT_{\delta},u+s}^{\delta} - \Phi_{u+mT_{\delta},u+s}^{\delta}\right) \left(DG(\mu_{s}) - DG(\Gamma_{u+s}^{\delta})\right) \xi_{s} \, \mathrm{d}s.$$

Let us define

$$R_n^{norm} := \int_0^{nT_\delta} \Pi_{u+nT_\delta}^{\delta,s} \Phi_{u+nT_\delta,u+s}^{\delta} D^2 G(\Gamma_{u+s}^{\delta}) [\partial_s \Gamma_{u+s}^{\delta}, \partial_s \Gamma_{u+s}^{\delta}] \, \mathrm{d}s$$

$$= \sum_{j=0}^{n-1} \left(\Phi_{u+T_\delta,u}^{\delta} \Pi_u^{\delta,s} \right)^j \int_0^{T_\delta} \Phi_{u+T_\delta,u+s}^{\delta} D^2 G(\Gamma_{u+s}^{\delta}) [\partial_s \Gamma_{u+s}^{\delta}, \partial_s \Gamma_{u+s}^{\delta}] \, \mathrm{d}s,$$

$$(5.48)$$

and

$$R_n^{tang}[\nu_0, w_0] := \int_0^{nT_\delta} \Pi_{u+nT_\delta}^{\delta, c} \Phi_{u+nT_\delta, u+s}^{\delta} D^2 G(\Gamma_{u+s}^{\delta})[\nu_s, w_s] \, \mathrm{d}s$$
 (5.49)

$$= \sum_{j=0}^{n-1} \int_0^{T_{\delta}} \Pi_{u+T_{\delta}}^{\delta,c} \Phi_{u+T_{\delta},u+s}^{\delta} D^2 G(\Gamma_{u+s}^{\delta}) [\nu_{jT_{\delta}+s}, w_{jT_{\delta}+s}] \, \mathrm{d}s.$$
 (5.50)

It is clear that

$$||R_n^{norm} - R_m^{norm}||_{\mathbf{H}_a^{-r}} \leqslant Ce^{-\lambda_{\delta} m T_{\delta}}.$$
 (5.51)

Now, for $j \ge 1$, recalling (5.40), (5.41) and (5.42) we have

$$\left\| \int_{0}^{T_{\delta}} \Pi_{u+T_{\delta}}^{\delta,c} \Phi_{u+T_{\delta},u+s}^{\delta} D^{2} G(\Gamma_{u+s}^{\delta}) [\nu_{jT_{\delta}+s}, w_{jT_{\delta}+s}] \, \mathrm{d}s \right\|_{\mathbf{H}_{\theta}^{-r}} \\ \leq C_{20} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| w_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} \int_{0}^{T_{\delta}} \left(1 + (T_{\delta} - s)^{-\frac{1}{2}} e^{-\lambda_{\delta} (T_{\delta} - s)} \right) (jT_{\delta} + s)^{\frac{1}{2}} e^{-\lambda_{\delta} \frac{jT_{\delta} + s}{2}} \, \mathrm{d}s \\ \leq C_{21} \left\| \nu_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| w_{0} \right\|_{\mathbf{H}_{\theta}^{-r}} (j+1)^{\frac{1}{2}} e^{-\lambda_{\delta} j \frac{T_{\delta}}{2}}, \tag{5.52}$$

so that

$$\left\| R_n^{tan}[\nu_0, w_0] - R_m^{tang}[\nu_0, w_0] \right\|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_{22} e^{-\lambda_{\delta} m \frac{T_{\delta}}{4}} \left\| \nu_0 \right\|_{\mathbf{H}_{\theta}^{-r}} \left\| w_0 \right\|_{\mathbf{H}_{\theta}^{-r}}. \tag{5.53}$$

Using similar argument as above, relying on (5.41) and (5.42),

$$\left\| \int_{0}^{nT_{\delta}} \Phi_{u+nT_{\delta},u+s}^{\delta} D^{2} G(\Gamma_{u+s}^{\delta}) [\nu_{s}, w_{s}] ds - R_{n}^{tan} [\nu_{0}, w_{0}] - l_{\mu} [\nu_{0}] l_{\mu} [w_{0}] R_{n}^{norm} ds \right\|_{\mathbf{H}_{\theta}^{-r}}$$

$$\leq C_{23} \|\nu_{0}\|_{\mathbf{H}_{\theta}^{-r}} \|w_{0}\|_{\mathbf{H}_{\theta}^{-r}} \int_{0}^{nT_{\delta}} (nT_{\delta} - s)^{-\frac{1}{2}} e^{-\lambda_{\delta} (nT_{\delta} - s)} s^{\frac{1}{2}} e^{-\lambda_{\delta} \frac{s}{2}} ds$$

$$\leq C_{24} n^{\frac{1}{2}} e^{-\lambda_{\delta} n^{\frac{T_{\delta}}{2}}} \|\nu_{0}\|_{\mathbf{H}_{\theta}^{-r}} \|w_{0}\|_{\mathbf{H}_{\theta}^{-r}}. \tag{5.54}$$

With all these estimates we are able to tackle the first two lines of (5.47):

$$\left\| \int_{0}^{nT_{\delta}} \Phi_{u+nT_{\delta},u+s}^{\delta} D^{2} G(\Gamma_{u+s}^{\delta}) [\nu_{s}, w_{s}] \, \mathrm{d}s - \int_{0}^{mT_{\delta}} \Phi_{u+mT_{\delta},u+s}^{\delta} D^{2} G(\Gamma_{u+s}^{\delta}) [\nu_{s}, w_{s}] \, \mathrm{d}s \right\|_{\mathbf{H}_{\theta}^{-r}} \\ \leqslant C_{25} n^{\frac{1}{2}} e^{-\lambda_{\delta} n^{\frac{T_{\delta}}{4}}} \|\nu_{0}\|_{\mathbf{H}_{\theta}^{-r}} \|w_{0}\|_{\mathbf{H}_{\theta}^{-r}}. \tag{5.55}$$

The other terms can be treated in a straightforward way, with similar estimates as the ones used in Step 2 and Step 3. At the end, one obtains

$$\|\xi_{nT_{\delta}} - \xi_{mT_{\delta}}\|_{\mathbf{H}_{\theta}^{-r}} \le C_{26} m^{\frac{1}{2}} e^{-\lambda_{\delta} m^{\frac{T_{\delta}}{4}}},$$
 (5.56)

with a constant C_{25} uniform in \mathcal{V} . Hence, $\mu \mapsto S(\mu)$ is thus C^2 . Remark that we have in particular

$$D^{2}S(\Gamma_{u})[\nu, w] = \lim_{n \to \infty} \int_{0}^{nT_{\delta}} \Phi_{u+nT_{\delta}, u+s} D^{2}G(\Gamma_{u+s})[\Phi_{u+s, u}\nu, \Phi_{u+s, u}w] \, \mathrm{d}s.$$
 (5.57)

Step 5: from the previous steps, and the fact that $t \mapsto \Gamma_t^{\delta}$ is a C^2 bijection from $\mathbb{R}/T_{\delta}\mathbb{Z}$ to \mathcal{M}^{δ} implies that Θ^{δ} is itself C^2 .

For the last estimate of the Theorem, let us denote $\xi_t^2 = D^2 T^t(\mu)[\nu, w], \ \xi_t^1 = D^2 T^t(\Gamma_u^\delta)[\nu, w], \ \nu_t^2 = D T^t(\mu)[\nu], \ \nu_t^1 = D T^t(\Gamma_u^\delta)[\nu], \ w_t^2 = D T^t(\mu)[w]$ and $w_t^1 = D T^t(\Gamma_u^\delta)[w]$. We then have the decomposition

$$\xi_{t}^{2} - \xi_{t}^{1} = \delta \int_{0}^{t} \Phi_{u+t,u+s}^{\delta} \left(DG(\mu_{s}) - DG \left(\Gamma_{u+s}^{\delta} \right) \right) \xi_{s}^{2} \, \mathrm{d}s$$

$$+ \delta \int_{0}^{t} \Phi_{u+t,u+s}^{\delta} \left(D^{2}G(\mu_{s}) - D^{2}G \left(\Gamma_{u+s}^{\delta} \right) \right) \left[\nu_{s}^{2}, w_{s}^{2} \right] \, \mathrm{d}s$$

$$+ \delta \int_{0}^{t} \Phi_{u+t,u+s}^{\delta} D^{2}G \left(\Gamma_{u+s}^{\delta} \right) \left[\nu_{s}^{2} - \nu_{s}^{1}, w_{s}^{2} \right] \, \mathrm{d}s$$

$$+ \delta \int_{0}^{t} \Phi_{u+t,u+s}^{\delta} D^{2}G \left(\Gamma_{u+s}^{\delta} \right) \left[\nu_{s}^{1}, w_{s}^{2} - w_{s}^{1} \right] \, \mathrm{d}s.$$
(5.58)

Following similar estimates as in the previous steps, relying in particular on (5.32), we obtain

$$\|\xi_t^2 - \xi_t^1\|_{\mathbf{H}_{\theta}^{-r}} \leqslant C_{27} \|\mu - \Gamma_u^{\delta}\|_{\mathbf{H}_{\theta}^{-r}} \|\nu\|_{\mathbf{H}_{\theta}^{-r}} \|w\|_{\mathbf{H}_{\theta}^{-r}}, \tag{5.59}$$

which implies indeed that $\|D^2\Theta^{\delta}(\mu) - D^2\Theta^{\delta}(\Gamma_u^{\delta})\|_{\mathcal{BL}(\mathbf{H}_a^{-r})} \leqslant C_{28} \|\mu - \Gamma_u^{\delta}\|_{\mathbf{H}_a^{-r}}$.

APPENDIX A. ORNSTEIN-UHLENBECK OPERATORS

We first prove the following lemma, which shows the link between the norm $||f||_{H_{\theta}^r}$ and the space derivatives.

Lemma A.1. For all $\theta > 0$, there exists explicit positive constants C_1 , C_2 such that for all $r \ge 0$:

$$C_1 \left(\|u\|_{H_{\theta}^r}^2 + \sum_{i=1}^d \|\partial_{x_i} u\|_{H_{\theta}^r}^2 \right) \leqslant \|u\|_{H_{\theta}^{r+1}}^2 \leqslant C_2 \left(\|u\|_{H_{\theta}^r}^2 + \sum_{i=1}^d \|\partial_{x_i} u\|_{H_{\theta}^r}^2 \right). \tag{A.1}$$

Proof. For u with decomposition $u = \sum_{l \in \mathbb{N}^d} u_l \psi_l$, we have $\partial_{x_i} u = \sum_{l \in \mathbb{N}^d} u_l \partial_{x_i} \psi_l$, and straightforward calculations (recalling (1.15) and using the fact that $h'_n(x) = \sqrt{n}h_{n-1}(x)$) show that

$$\partial_{x_i} \psi_l = \sqrt{l_i} \sqrt{\frac{\theta k_i}{\sigma_i^2}} \psi_{\tilde{l}_i} \mathbf{1}_{l_i \geqslant 1}, \tag{A.2}$$

where we used the notation $\check{l}_i = (l_1, \dots, l_{i-1}, l_i - 1, l_{i+1}, \dots, l_d)$. Then we have the decomposition

$$\partial_{x_i} u = \sqrt{\frac{\theta k_i}{\sigma_i^2}} \sum_{l \in \mathbb{N}^d} \sqrt{l_i} \mathbf{1}_{l_i \geqslant 1} u_l \, \psi_{\tilde{l}_i}, \tag{A.3}$$

so that

$$||u||_{H_{\theta}^{r}}^{2} + \sum_{i=1}^{d} ||\partial_{x_{i}}u||_{H_{\theta}^{r}}^{2} = \sum_{l \in \mathbb{N}^{d}} u_{l}^{2} \left((a_{\theta} + \lambda_{l})^{r} + \sum_{i=1}^{d} l_{i} \mathbf{1}_{l_{i} \geqslant 1} \frac{\theta k_{i}}{\sigma_{i}^{2}} (a_{\theta} + \lambda_{l} - \theta k_{i})^{r} \right). \quad (A.4)$$

Let us prove the upper bound in (A.1): note that for $l_i \ge 1$, we have $\lambda_l \ge \theta k_i$. Hence, since for all $\mu \ge 0$, $r \ge 0$, $\lambda \ge \mu$, $(a_\theta + \lambda - \mu)^r \ge a_\theta^r \frac{(a_\theta + \lambda)^r}{(a_\theta + \mu)^r}$, we deduce that

$$||u||_{H_{\theta}^{r}}^{2} + \sum_{i=1}^{d} ||\partial_{x_{i}}u||_{H_{\theta}^{r}}^{2} \geqslant \sum_{l \in \mathbb{N}^{d}} u_{l}^{2} (a_{\theta} + \lambda_{l})^{r} \left(1 + \sum_{i=1}^{d} l_{i} \mathbf{1}_{l_{i}} \geqslant 1 \frac{\theta k_{i} a_{\theta}^{r}}{\sigma_{i}^{2} (a_{\theta} + \theta k_{i})^{r}} \right),$$

$$\geqslant \sum_{l \in \mathbb{N}^{d}} u_{l}^{2} (a_{\theta} + \lambda_{l})^{r} \left(1 + \frac{a_{\theta}^{r}}{\sigma_{\max}^{2} (a_{\theta} + \theta k_{\max})^{r}} \lambda_{l} \right),$$

so that the upper bound in (A.1) is true for $C_2 := \max\left(\frac{\sigma_{\max}^2(a_{\theta} + \theta k_{\max})^r}{a_{\theta}^r}, a_{\theta}\right)$. Concerning the lower bound in (A.1), we have from (A.4),

$$||u||_{H_{\theta}^r}^2 + \sum_{i=1}^d ||\partial_{x_i} u||_{H_{\theta}^r}^2 \leqslant \sum_{l \in \mathbb{N}^d} u_l^2 (a_{\theta} + \lambda_l)^r \left(1 + \frac{\lambda_l}{\sigma_{\min}^2}\right),$$

so that the upper bound holds for $C_1 := \frac{1}{\min(\sigma_{\min}^2, a_{\theta})}$.

For all $\theta > 0$, the operator $-\mathcal{L}_{\theta}^*$ (recall its definition (1.14) and its decomposition (1.15)) is sectorial in L_{θ}^2 and generates a semigroup $e^{t\mathcal{L}_{\theta}^*}$ satisfying (see e.g. [23]) for all $\alpha \geq 0$, $r \geq 0$, and $\lambda < \theta \min(k_1, \ldots, k_d)$, there exists some C > 0 such that for all $f \in H_{\theta}^r$,

$$\left\| e^{t\mathcal{L}_{\theta}^*} f \right\|_{H_{a}^{r+\alpha}} \leqslant C \left(1 + t^{-\alpha/2} e^{-\lambda t} \right) \|f\|_{H_{\theta}^r}, \tag{A.5}$$

and for all $f \in H_{\theta}^r$ such that $\int f w_{\theta} = 0$,

$$\left\| e^{t\mathcal{L}_{\theta}^*} f \right\|_{H_{\theta}^{r+\alpha}} \leqslant C t^{-\frac{1}{2}} e^{-\lambda t} \|f\|_{H_{\theta}^r}. \tag{A.6}$$

Let $\theta' > 0$. The point of the following result is to state similar contraction results for $\mathcal{L}_{\theta'}^*$ in H_{θ}^r , in the case $\theta' \neq \theta$:

Proposition A.2. For all $0 < \theta \le \theta'$ the following is true: the operator $\mathcal{L}_{\theta'}^*$ generates an analytic semigroup in H_{θ}^r and for all $r \ge 0$, $\alpha \ge 0$ and $\lambda < \theta \min(k_1, \ldots, k_d)$, there exists a constant C > 0 such that for all $f \in H_{\theta}^r$ and t > 0

$$\left\| e^{t\mathcal{L}_{\theta'}^*} f \right\|_{H_{\theta}^{r+\alpha}} \leqslant C \left(1 + t^{-\alpha/2} e^{-\lambda t} \right) \|f\|_{H_{\theta}^r}, \tag{A.7}$$

and for all $r \ge 1$,

$$\left\| \nabla e^{t\mathcal{L}_{\theta'}^*} f \right\|_{H_{\tau}^r} \leqslant C t^{-\frac{1}{2}} e^{-\lambda t} \|f\|_{H_{\theta}^r}. \tag{A.8}$$

Moreover for all $r \ge 0$, $0 < \varepsilon \le 1$ and $s \ge 0$,

$$\left\| \left(e^{(t+s)\mathcal{L}_{\theta'}^*} - e^{t\mathcal{L}_{\theta'}^*} \right) f \right\|_{H_{\theta}^{r+1}} \leqslant C s^{\varepsilon} t^{-\frac{1}{2} - \varepsilon} e^{-\lambda t} \| f \|_{H_{\theta}^r}. \tag{A.9}$$

Finally, there exists $r_0 > 0$ such that for all $r > r_0$, t > 0 and all $f \in H^r_\theta$,

$$\left\| e^{t\mathcal{L}_{\theta'}^*} f - \frac{\int_{\mathbb{R}^d} e^{t\mathcal{L}_{\theta'}^*} f w_{\theta}}{\int_{\mathbb{R}^d} w_{\theta}} \right\|_{H_{\theta}^r} \leqslant e^{-\lambda t} \left\| f - \int_{\mathbb{R}^d} f w_{\theta} \right\|_{H_{\theta}^r}. \tag{A.10}$$

Proof of Proposition A.2. First remark that for all smooth test function u

$$(\mathcal{L}_{\theta}^* - \mathcal{L}_{\theta'}^*) u = (\theta' - \theta) Kx \cdot \nabla u. \tag{A.11}$$

Recalling the decomposition (1.15),since $h'_n(x) = \sqrt{n}h_{n-1}(x)$ and $xh_{n-1}(x) = \sqrt{n}h_n(x) + \sqrt{n-1}h_{n-2}(x)$ (see e.g. [6], p.102), we get, for all $l \in \mathbb{N}$,

$$\left(\mathcal{L}_{\theta}^* - \mathcal{L}_{\theta'}^*\right) \psi_{\theta,l} = \left(\theta' - \theta\right) \sum_{i=1}^d k_i \sqrt{\frac{\theta k_i}{\sigma_i^2}} \sqrt{l_i} x_i \psi_{\theta,l_{\downarrow_i}} \tag{A.12}$$

$$= (\theta' - \theta) \sum_{i=1}^{d} k_i \left(l_i \psi_{\theta,l} + \sqrt{l_i (l_i - 1)} \psi_{\theta,l_{\downarrow \downarrow_i}} \right), \tag{A.13}$$

where we used the following notations for the shifts w.r.t. the ith coordinate

$$l_{\downarrow i} = (l_1, \dots, l_{i-1}, l_i - 1, l_{i+1}, \dots, l_d), \tag{A.14}$$

$$l_{\uparrow_i} = (l_1, \dots, l_{i-1}, l_i + 1, l_{i+1}, \dots, l_d)$$
(A.15)

and the convention $\psi_l = 0$ if $l_i < 0$ for some $i \in \{1, \dots, d\}$. In particular we have, recalling that $\lambda_{\theta,l} = \theta \sum_{i=1}^d k_i l_i$,

$$-\mathcal{L}_{\theta'}^* \psi_{\theta,l} = \frac{\theta' \lambda_{\theta,l}}{\theta} \psi_{\theta,l} + (\theta' - \theta) \sum_{i=1}^d k_i \sqrt{l_i (l_i - 1)} \psi_{\theta,l_{\parallel i}}, \tag{A.16}$$

So, we deduce that for $f = \sum_{l} f_{l} \psi_{\theta,l}$, with $f_{l} \in \mathbb{C}$ for all l,

$$\left\| \left(\mathcal{L}_{\theta}^* - \frac{\theta}{\theta'} \mathcal{L}^* \right) f \right\|_{H_{\theta}^r}^2 = \left(1 - \frac{\theta}{\theta'} \right)^2 \sum_{l} (a_{\theta} + \lambda_{\theta, l})^r \left| \sum_{i=1}^d \theta k_i \sqrt{(l_i + 1)(l_i + 2)} f_{l_{\uparrow i}} \right|^2.$$

Setting

$$I(r,f) := \sum_{l} (a_{\theta} + \lambda_{\theta,l})^r \left(\sum_{i=1}^d \theta k_i \sqrt{(l_i + 1)(l_i + 2)} f_{l_{\Pi_i}} \right)^2.$$
 (A.17)

Using Jensen's inequality, we obtain (recalling that $a_{\theta} = \theta \operatorname{Tr}(K)$),

$$I(r,f) \leqslant \sum_{l_{1},\dots,l_{d}=0}^{+\infty} (a_{\theta} + \lambda_{\theta,l})^{r} \left(\sum_{i=1}^{d} \theta k_{i}(l_{i}+1) \right) \left(\sum_{i=1}^{d} \theta k_{i}(l_{i}+2) \left| f_{l_{\Pi_{i}}} \right|^{2} \right)$$

$$= \sum_{l_{1},\dots,l_{d}=0}^{+\infty} (a_{\theta} + \lambda_{\theta,l})^{r+1} \left(\sum_{i=1}^{d} \theta k_{i}(l_{i}+2) \left| f_{l_{\Pi_{i}}} \right|^{2} \right)$$

$$= \sum_{l_{1},\dots,l_{d}=2}^{+\infty} (a_{\theta} + \lambda_{\theta,l-2})^{r+1} \left(\sum_{i=1}^{d} \theta k_{i}l_{i} \left| f_{l_{1}-2,\dots,l_{i},l_{i+1}-2,l_{d}-2} \right|^{2} \right)$$

$$= \sum_{i=1}^{d} \sum_{l_{1},\dots,l_{d}=2}^{+\infty} (a_{\theta} + \lambda_{\theta,l-2})^{r+1} \theta k_{i}l_{i} \left| f_{l_{1}-2,\dots,l_{i},l_{i+1}-2,l_{d}-2} \right|^{2}$$

$$= \sum_{i=1}^{d} \sum_{l_{i}=2}^{+\infty} \sum_{\substack{l_{p}=0\\p\neq i}}^{+\infty} \left(a_{\theta} + \lambda_{\theta,l_{\square_{i}}} \right)^{r+1} \theta k_{i}l_{i} \left| f_{l} \right|^{2}.$$

Now we use that $\lambda_{\theta,l_{\perp i}} \leq \lambda_{\theta,l}$ for any l and i, so that

$$I(r,f) \leqslant \sum_{i=1}^{d} \sum_{\substack{l_i=2 \ p \neq i}}^{+\infty} \sum_{l_p=0}^{+\infty} (a_{\theta} + \lambda_{\theta,l})^{r+1} \theta k_i l_i |f_l|^2,$$
(A.18)

This sum is anyway smaller than

$$I(r,f) \leqslant \sum_{i=1}^{d} \sum_{\substack{l_i=0 \ p \neq i}}^{+\infty} \sum_{\substack{l_p=0 \ p \neq i}}^{+\infty} (a_{\theta} + \lambda_{\theta,l})^{r+1} \theta k_i l_i |f_l|^2 = \sum_{l_1=0}^{+\infty} \dots \sum_{l_d=0}^{+\infty} (a_{\theta} + \lambda_{\theta,l})^{r+1} \lambda_{\theta,l} |f_l|^2,$$

$$\leqslant \sum_{l_1=0}^{+\infty} \dots \sum_{\substack{l_d=0}}^{+\infty} (a_{\theta} + \lambda_{\theta,l})^{r+2} |f_l|^2 = \|(a_{\theta} - \mathcal{L}_{\theta}^*)f\|_{H_{\theta}^r}^2. \tag{A.19}$$

Coming back to (A.17), we obtain

$$\left\| \left(\mathcal{L}_{\theta}^* - \frac{\theta}{\theta'} \mathcal{L}_{\theta'}^* \right) f \right\|_{H_0^r} \leqslant \left(1 - \frac{\theta}{\theta'} \right) \left\| (a_{\theta} - \mathcal{L}_{\theta}^*) f \right\|_{H_{\theta}^r}, \tag{A.20}$$

which implies in particular that

$$\left\| \left(a_{\theta} - \frac{\theta}{\theta'} \mathcal{L}_{\theta'}^* \right) f \right\|_{H_{\theta}^r} \leqslant \left(2 - \frac{\theta}{\theta'} \right) \left\| \left(a_{\theta} - \mathcal{L}_{\theta}^* \right) f \right\|_{H_{\theta}^r} = \left(2 - \frac{\theta}{\theta'} \right) \left\| f \right\|_{H_{\theta}^{r+2}}. \tag{A.21}$$

Let us now look for a lower bound. Using $(a+b)^2 \geqslant \frac{\varepsilon}{1+\varepsilon}a^2 - \varepsilon b^2$ $(\varepsilon > 0)$, we get

$$\left\| \left(a_{\theta} - \frac{\theta}{\theta'} \mathcal{L}_{\theta'}^* \right) f \right\|_{H_{\theta}^r}^2 = \sum_{l} (a_{\theta} + \lambda_{\theta, l})^r \left((a_{\theta} + \lambda_{\theta, l}) f_l + \left(1 - \frac{\theta}{\theta'} \right) \sum_{i=1}^d \theta k_i \sqrt{(l_i + 1)(l_i + 2)} f_{l_{\uparrow i}} \right)^2$$

$$\geqslant \frac{\varepsilon}{1 + \varepsilon} \sum_{l} (a_{\theta} + \lambda_{\theta, l})^{r+2} f_l^2 - \varepsilon \left(1 - \frac{\theta}{\theta'} \right)^2 I(r, f). \tag{A.22}$$

Hence, recalling (A.19), we obtain

$$\left\| \left(a_{\theta} - \frac{\theta}{\theta'} \mathcal{L}_{\theta'}^* \right) f \right\|_{H_{\theta}^r}^2 \geqslant \varepsilon \left(\frac{1}{1 + \varepsilon} - \left(1 - \frac{\theta}{\theta'} \right)^2 \right) \left\| (a_{\theta} - \mathcal{L}_{\theta}^*) f \right\|_{H_{\theta}^r}^2 \tag{A.23}$$

So for $\varepsilon > 0$ small enough (depending only on θ, θ'), there exists a constant $c_{\theta,\theta'} > 0$ such that we have $\|\left(a_{\theta} - \frac{\theta}{\theta'}\mathcal{L}_{\theta'}^*\right)f\|_{H_{\theta}^r} \geqslant c_{\theta,\theta'}\|(a_{\theta} - \mathcal{L}_{\theta}^*)f\|_{H_{\theta}^r}$. This means, together with (A.21), that

$$c_{\theta,\theta'} \|f\|_{H^{r+2}_{\theta}} \leqslant \left\| \left(a_{\theta} - \frac{\theta}{\theta'} \mathcal{L}^*_{\theta'} \right) f \right\|_{H^r_{\theta}} \leqslant \left(2 - \frac{\theta}{\theta'} \right) \|f\|_{H^{r+2}_{\theta}}. \tag{A.24}$$

In particular 0 is in the resolvent set of $a_{\theta} - \frac{\theta}{\theta'} \mathcal{L}_{\theta'}^*$, and $a_{\theta} - \frac{\theta}{\theta'} \mathcal{L}_{\theta'}^*$ has a compact resolvent, since it is the case for $a_{\theta} - \mathcal{L}_{\theta}^*$. So $\mathcal{L}_{\theta'}^*$ has a discrete spectrum, composed of a sequence of eigenvalues with modulus going to infinity. But any eigenfunction ψ of $\mathcal{L}_{\theta'}^*$ in H_{θ}^r is also an eigenfunction of $\mathcal{L}_{\theta'}^*$ in $H_{\theta'}^r$, and thus the eigenvalues of $\mathcal{L}_{\theta'}^*$ in H_{θ}^r are the $\lambda_{l,\theta'}$'s, with

associated eigenfunctions the $\psi_{l,\theta'}$'s. So in particular $\mathcal{L}_{\theta'}^*$ is sectorial, and thus generates an analytic semigroup $e^{t\mathcal{L}_{\theta'}^*}$ in H_{θ}^r .

Let us now prove that the interpolation spaces induced by $\mathcal{L}_{\theta'}^*$ and \mathcal{L}_{θ}^* in H_{θ}^r are equivalent. Since the operator $\left(\frac{\theta}{\theta'}\mathcal{L}_{\theta'}^* - \mathcal{L}_{\theta}^*\right)\left(a_{\theta} - \frac{\theta}{\theta'}\mathcal{L}_{\theta'}^*\right)^{-1}$ (and thus $(1+(\theta\mathcal{L}^*-\mathcal{L}_{\theta}^*)(a_{\theta}-\theta\mathcal{L}^*)^{-1})^{\alpha}$ for $\alpha \geq 0$) is bounded in H_{θ}^r , we obtain

$$||f||_{H_{\theta}^{r+\alpha}} = \left\| \left(a_{\theta} - \mathcal{L}_{\theta}^{*} \right)^{\alpha/2} f \right\|_{H_{\theta}^{r}}$$

$$= \left\| \left(1 + \left(\frac{\theta}{\theta'} \mathcal{L}_{\theta'}^{*} - \mathcal{L}_{\theta}^{*} \right) \left(a_{\theta} - \frac{\theta}{\theta'} \mathcal{L}_{\theta'}^{*} \right)^{-1} \right)^{\alpha/2} \left(a_{\theta} - \frac{\theta}{\theta'} \mathcal{L}_{\theta'}^{*} \right)^{\alpha/2} f \right\|_{H_{\theta}^{r}}$$

$$\leq C \left\| \left(a_{\theta} - \frac{\theta}{\theta'} \mathcal{L}_{\theta'}^{*} \right)^{\alpha/2} f \right\|_{H_{\theta}^{r}}.$$
(A.25)

The inverse bound $\left\|\left(a_{\theta} - \frac{\theta}{\theta'}\mathcal{L}_{\theta'}^*\right)^{\alpha/2} f\right\|_{H_{\theta}^r} \leqslant C\|f\|_{H_{\theta}^{r+\alpha}}$ follows from similar arguments.

We are now in condition to prove (A.5), applying [23], Th. 1.4.3. Indeed, $\mathcal{L}_{\theta'}^*$ has a real spectrum smaller than $-\theta' \min(k_1, \dots, k_d)$ on the the subspace of H_{θ}^r generated by the eigenfunctions $\psi_{l,\theta'}$ with $l \neq 0$, so applying this Theorem we get, denoting $\mathcal{P}_{\theta'} f = f - \frac{\int_{\mathbb{R}^d} f w_{\theta'}}{\left(\int_{\mathbb{R}^d} w_{\theta'}\right)^2}$ the projection on this subspace (which is an element of $\mathcal{B}(H_{\theta}^r)$ for $0 < \theta \leqslant \theta'$),

$$\left\| e^{t\mathcal{L}_{\theta'}^*} \mathcal{P}_{\theta'} f \right\|_{H_{\theta}^{r+\alpha}} \leqslant C_1 \left\| \left(a_{\theta} - \frac{\theta}{\theta'} \mathcal{L}_{\theta'}^* \right)^{\alpha/2} e^{t\mathcal{L}_{\theta'}^*} \mathcal{P}_{\theta'} f \right\|_{H_{\theta}^r} \leqslant C_2 t^{-\alpha/2} e^{-\lambda t} \left\| f \right\|_{H_{\theta}^r}. \tag{A.26}$$

This implies (A.5), since $e^{t\mathcal{L}_{\theta'}^*} (1 - \mathcal{P}_{\theta'}) f = (1 - \mathcal{P}_{\theta'}) f$.

The proof of (A.6) relies on the classical identity, valid for $f \in L^2_{\theta'}$.

$$e^{t\mathcal{L}_{\theta'}^*} f = \mathbb{E}\left[f\left(e^{-t\theta'K}x + \sqrt{1 - e^{-2t\theta'K}}G_{\theta'}\right)\right],\tag{A.27}$$

where $G_{\theta'}$ a gaussian variable on \mathbb{R}^d with mean 0 and variance $(\theta'K)^{-1}\sigma^2$. This implies directly, for $f \in H^r_{\theta}$ with $r \geqslant 1$,

$$\nabla e^{t\mathcal{L}_N^*} f = e^{-t\theta' K} e^{t\mathcal{L}_N^*} \nabla f, \tag{A.28}$$

and thus, denoting $\underline{K} = \min\{k_1, \dots, k_d\}$,

$$\begin{split} \left\| \nabla e^{t\mathcal{L}_N^*} f \right\|_{H_{\theta}^r} &\leqslant e^{-\theta' \underline{K}t} \left\| e^{t\mathcal{L}_N^*} \nabla f \right\|_{H_{\theta}^r} \leqslant C \left(1 + t^{-\frac{1}{2}} e^{-\lambda t} \right) e^{-\theta' \underline{K}t} \left\| \nabla f \right\|_{H_{\theta}^{r-1}} \\ &\leqslant C' t^{-\frac{1}{2}} e^{-\lambda t} \left\| f \right\|_{H_{x}^{r}}. \quad (A.29) \end{split}$$

For the proof of the third assertion, since [23], Th. 1.4.3. implies that for $0 < \varepsilon \le 1$,

$$\left\| \left(e^{s\mathcal{L}_{\theta'}^*} - 1 \right) f \right\|_{H_{\theta}^r} \leqslant C_{\varepsilon} s^{\varepsilon} \left\| f \right\|_{H_{\theta}^{r+2\varepsilon}}, \tag{A.30}$$

we obtain, since $\left(e^{s\mathcal{L}_{\theta'}^*}-1\right)\mathcal{P}_{\theta'}f=\left(e^{s\mathcal{L}_{\theta'}^*}-1\right)f$ and $\mathcal{P}_{\theta'}$ commutes with $e^{t\mathcal{L}_{\theta'}^*}$,

$$\left\| \left(e^{(t+s)\mathcal{L}_{\theta'}^*} - e^{t\mathcal{L}_{\theta'}^*} \right) f \right\|_{H_{\theta}^{r+1}} = \left\| \left(e^{s\mathcal{L}_{\theta'}^*} - 1 \right) e^{t\mathcal{L}_{\theta'}^*} \mathcal{P}_{\theta'} f \right\|_{H_{\theta}^{r+1}} \leqslant C_{\varepsilon} s^{\varepsilon} \left\| e^{t\mathcal{L}_{\theta'}^*} \mathcal{P}_{\theta'} f \right\|_{H_{\theta}^{r+1+2\varepsilon}}$$

$$\leqslant C s^{\varepsilon} t^{-\frac{1}{2} - \varepsilon} e^{-\lambda t} \left\| f \right\|_{H_{\theta}^{r}}. \quad (A.31)$$

The last assertion is not a direct consequence of the estimates obtained above, since the hypothesis $\int_{\mathbb{R}^d} f w_{\theta} = 0$ is not well adapted to the eigenfunctions $\psi_{l,\theta'}$ of $\mathcal{L}_{\theta'}^*$. In particular, having $\int_{\mathbb{R}^d} f w_{\theta} = 0$ does not imply $\int_{\mathbb{R}^d} e^{t\mathcal{L}_{\theta'}^*} f w_{\theta} = 0$, while it is the case when $\theta = \theta'$. We will only be able to obtain this estimates for r large enough, via direct calculations. Remark first that

$$\frac{1}{2} \frac{d}{dt} \left\| e^{t\mathcal{L}_{\theta'}^*} f - \frac{\int_{\mathbb{R}^d} e^{t\mathcal{L}_{\theta'}^*} f w_{\theta}}{\int_{\mathbb{R}^d} w_{\theta}} \right\|_{H_{\theta}^r}^2 \\
= \left\langle \mathcal{L}_{\theta'}^* e^{t\mathcal{L}_{\theta'}^*} f - \frac{\int_{\mathbb{R}^d} \mathcal{L}_{\theta'}^* e^{t\mathcal{L}_{\theta'}^*} f w_{\theta}}{\int_{\mathbb{R}^d} w_{\theta}}, e^{t\mathcal{L}_{\theta'}^*} f - \frac{\int_{\mathbb{R}^d} e^{t\mathcal{L}_{\theta'}^*} f w_{\theta}}{\int_{\mathbb{R}^d} w_{\theta}} \right\rangle_{H_{\theta}^r}. \quad (A.32)$$

Recalling that $\mathcal{L}_{\theta'}^* a = 0$ and remarking that $\left\langle a, e^{t\mathcal{L}_{\theta'}^*} f - \frac{\int_{\mathbb{R}^d} e^{t\mathcal{L}_{\theta'}^*} f w_{\theta}}{\int_{\mathbb{R}^d} w_{\theta}} \right\rangle_{H_{\theta}^r} = 0$ for any constant a, we get

$$\frac{1}{2} \frac{d}{dt} \left\| e^{t\mathcal{L}_{\theta'}^*} f - \frac{\int_{\mathbb{R}^d} e^{t\mathcal{L}_{\theta'}^*} f w_{\theta}}{\int_{\mathbb{R}^d} w_{\theta}} \right\|_{H_{\theta}^T}^2 \\
= \left\langle \mathcal{L}_{\theta'}^* \left(e^{t\mathcal{L}_{\theta'}^*} f - \frac{\int_{\mathbb{R}^d} e^{t\mathcal{L}_{\theta'}^*} f w_{\theta}}{\int_{\mathbb{R}^d} w_{\theta}} \right), e^{t\mathcal{L}_{\theta'}^*} f - \frac{\int_{\mathbb{R}^d} e^{t\mathcal{L}_{\theta'}^*} f w_{\theta}}{\int_{\mathbb{R}^d} w_{\theta}} \right\rangle_{H_0^T}, \quad (A.33)$$

so the proof of the last assertion reduces to the study of $\langle \mathcal{L}_{\theta'}^* f, f \rangle_{H_{\theta}^r}$ with $\int_{\mathbb{R}^d} f w_{\theta} = 0$. Now for f satisfying $\int_{\mathbb{R}^d} f w_{\theta} = 0$, with decomposition $f = \sum_{l \neq 0} f_l \psi_{l,\theta}$, we get

$$-\frac{\theta}{\theta'} \langle \mathcal{L}_{\theta'}^* f, f \rangle_{H_{\theta}^r} = \sum_{l \neq 0} (a_{\theta} + \lambda_{\theta, l})^r \lambda_{\theta, l} |f_l|^2 + \left(1 - \frac{\theta}{\theta'}\right) \left(\sum_{l \neq 0} (a_{\theta} + \lambda_{\theta, l})^r \sum_{i=1}^d \theta k_i \sqrt{(l_i + 1)(l_i + 2)} f_l \bar{f}_{l_{\parallel i}}\right). \tag{A.34}$$

Now remark that for the second term, using Cauchy Schwarz inequality, we get

$$\left| \sum_{l \neq 0} (a_{\theta} + \lambda_{\theta,l})^r \sum_{i=1}^d \theta k_i \sqrt{(l_i + 1)(l_i + 2)} f_l \bar{f}_{l \uparrow_i} \right|$$

$$= \left| \sum_{l \neq 0} \left\{ (a_{\theta} + \lambda_{\theta,l})^{r/2} \sqrt{\lambda_{\theta,l}} f_l \right\} \left\{ \frac{(a_{\theta} + \lambda_{\theta,l})^{r/2}}{\sqrt{\lambda_{\theta,l}}} \sum_{i=1}^d \theta k_i \sqrt{(l_i + 1)(l_i + 2)} \bar{f}_{l \uparrow_i} \right\} \right|$$

$$\leqslant \left| \sum_{l \neq 0} (a_{\theta} + \lambda_{\theta,l})^r \lambda_{\theta,l} |f_l|^2 \right|^{\frac{1}{2}} \left| \sum_{l \neq 0} \frac{(a_{\theta} + \lambda_{\theta,l})^r}{\lambda_{\theta,l}} \left(\sum_{i=1}^d \theta k_i \sqrt{(l_i + 1)(l_i + 2)} \bar{f}_{l \uparrow_i} \right)^2 \right|^{\frac{1}{2}}.$$

Using Jensen's inequality

$$\left| \sum_{l \neq 0} \frac{(a_{\theta} + \lambda_{\theta,l})^r}{\lambda_{\theta,l}} \left(\sum_{i=1}^d \theta k_i \sqrt{(l_i + 1)(l_i + 2)} \bar{f}_{l_{\uparrow i}} \right)^2 \right|$$

$$\leqslant \sum_{l \neq 0} \frac{(a_{\theta} + \lambda_{\theta,l})^r}{\lambda_{\theta,l}} \left(\sum_{i=1}^d \theta k_i (l_i + 2) \right) \sum_{i=1}^d \theta k_i (l_i + 2) \left| f_{l_{\uparrow i}} \right|^2,$$

$$= \sum_{l \neq 0} \frac{(a_{\theta} + \lambda_{\theta,l})^r}{\lambda_{\theta,l}} \left(2a_{\theta} + \lambda_{\theta,l} \right) \sum_{i=1}^d \theta k_i (l_i + 2) \left| f_{l_{\uparrow i}} \right|^2,$$

Denoting by $\mathcal{N}_i := \uparrow \uparrow_i (\mathbb{N}^d \setminus \{0\}) = \{l \in \mathbb{N}^d, l_i \geqslant 2, \sum_{j=1}^d l_j \geqslant 3\}$, we obtain

$$\left| \sum_{l \neq 0} \frac{(a_{\theta} + \lambda_{\theta,l})^r}{\lambda_{\theta,l}} \left(\sum_{i=1}^d \theta k_i \sqrt{(l_i + 1)(l_i + 2)} \bar{f}_{l_{\Pi_i}} \right)^2 \right|$$

$$\leq 2 \sum_{i=1}^d \sum_{l \in \mathcal{N}_i} \frac{\left(a_{\theta} + \lambda_{\theta,l_{\coprod_i}} \right)^r}{\lambda_{\theta,l_{\coprod_i}}} \left(2a_{\theta} + \lambda_{\theta,l_{\coprod_i}} \right) \theta k_i l_i |f_l|^2$$

$$= \sum_{i=1}^d \sum_{l \in \mathcal{N}_i} \frac{\left(a_{\theta} + \lambda_{\theta,l} - 2\theta k_i \right)^r}{\lambda_{\theta,l} - 2\theta k_i} \left(2a_{\theta} + \lambda_{\theta,l} - 2\theta k_i \right) \theta k_i l_i |f_l|^2$$

$$= \sum_{i=1}^d \sum_{l \in \mathcal{N}_i} b_{\theta,l,i} \left(a_{\theta} + \lambda_{\theta,l} \right)^r \theta k_i l_i |f_l|^2,$$

where

$$b_{\theta,l,i} := \frac{(a_{\theta} + \lambda_{\theta,l} - 2\theta k_i)^r (2a_{\theta} + \lambda_{\theta,l} - 2\theta k_i)}{(a_{\theta} + \lambda_{\theta,l})^r (\lambda_{\theta,l} - 2\theta k_i)} = \left(1 - \frac{2\theta k_i}{a_{\theta} + \lambda_{\theta,l}}\right)^r \left(1 + \frac{2a_{\theta}}{\lambda_{\theta,l} - 2\theta k_i}\right). \tag{A.35}$$

Now, for $l \in \mathcal{N}_i$, we have

$$a_{\theta} + \lambda_{\theta,l} \leqslant d\theta k_{max} + \lambda_{\theta,l} \leqslant (d+1) \frac{k_{max}}{k_{min}} \lambda_{\theta,l},$$
 (A.36)

and

$$\lambda_{\theta,l} - 2\theta k_i \geqslant \theta \left(\sum_{j=1}^{d} l_j - 2\right) k_{min} \geqslant \frac{k_{min}}{3k_{max}} \lambda_{\theta,l},$$
 (A.37)

so that

$$b_{\theta,l,i} \leqslant \left(1 - \frac{2k_{min}^2}{(d+1)k_{max}} \cdot \frac{1}{\sum_{j=1}^d k_j l_j}\right)^r \left(1 + \frac{6d k_{max}^2}{k_{min}} \frac{1}{\sum_{j=1}^d k_j l_j}\right). \tag{A.38}$$

Now, observe that for $c_1, c_2 > 0$, $x \mapsto \left(1 - \frac{c_1}{x}\right)^r \left(1 + \frac{c_2}{x}\right)$ is strictly increasing with limit 1 as $x \to \infty$, provided that $r > \frac{c_2}{c_1}$. Hence, taking r large enough in (A.38) (r depending only on K and d, not on l, i nor θ) we have $|b_{\theta,l,i}| \leq 1$, which means that the second term of the right-hand side of (A.34) is bounded as follows:

$$\left| \sum_{l \neq 0} (a_{\theta} + \lambda_{\theta,l})^r \sum_{i=1}^d \theta k_i \sqrt{(l_i + 1)(l_i + 2)} f_l \bar{f}_{l_{\Pi_i}} \right| \leqslant \sum_{l \neq 0} (a_{\theta} + \lambda_{\theta,l})^r \lambda_{\theta,l} |f_l|^2. \tag{A.39}$$

We deduce from (A.34) and this estimate that

$$\left| -\frac{\theta}{\theta'} \left\langle \mathcal{P}_{\theta}^{\perp} \mathcal{L}_{\theta'}^{*} \mathcal{P}_{\theta}^{\perp} f, \bar{f} \right\rangle_{H_{\theta}^{r}} - \sum_{l \neq 0} (a_{\theta} + \lambda_{\theta, l})^{r} \lambda_{\theta, l} |f_{l}|^{2} \right| \leq \left(1 - \frac{\theta}{\theta'} \right) \sum_{l \neq 0} (a_{\theta} + \lambda_{\theta, l})^{r} \lambda_{\theta, l} |f_{l}|^{2}, \tag{A.40}$$

which means that

$$-\operatorname{Re}\left\langle \mathcal{L}_{\theta'}^{*}f, \bar{f}\right\rangle_{H_{\theta}^{r}} \geqslant \sum_{l\neq 0} (a_{\theta} + \lambda_{\theta, l})^{r} \lambda_{\theta, l} |f_{l}|^{2} \geqslant \theta k_{min} ||f||_{H_{\theta}^{r}}^{2}. \tag{A.41}$$

As already stated in Section 1.3, we rely in this paper on a "pivot" space structure (see [10], pp.81-82): observe first that for $u \in L^2_{-\theta}$, $v \in L^2_{\theta} \mapsto \int_{\mathbb{R}^d} uv \mathrm{d}x$ defines a continuous linear form on L^2_{θ} . Respectively, for $u \in (L^2_{\theta})'$, the mapping $\psi \mapsto Tu(\psi) := \langle u\,,\,\psi w_{-\theta}\rangle$ defines a continuous linear form on L^2 (that is the usual L^2 space without weight, i.e. $w \equiv 1$ in (1.13)). By Riesz Theorem, there exists $v \in L^2$, such that $Tu(\psi) = \int v\psi = \int \widetilde{v}\widetilde{\psi}, \ \psi \in L^2$, for $\widetilde{v} := vw_{\theta/2} \in L^2_{-\theta}, \ \widetilde{\psi} = \psi w_{-\theta/2} \in L^2_{\theta}$. This observation permits the identification of $(L^2_{\theta})'$ with $L^2_{-\theta}$ (and hence, $\langle \cdot, \cdot \rangle_{(L^2_{\theta})' \times L^2_{\theta}}$ with $\langle \cdot, \cdot \rangle_{L^2} = \langle \cdot, \cdot \rangle$). Now, since $H^r_{\theta} \to L^2_{\theta}$ is dense, we have a dense injection $(L^2_{\theta})' \to H^{-r}_{\theta}$. With the identification $(L^2_{\theta})' \approx L^2_{-\theta}$, we obtain, for all $u \in L^2_{-\theta} \subset H^{-r}_{\theta}$ and all $f \in H^r_{\theta}$,

$$\langle u, f \rangle_{H_{\theta}^{-r} \times H_{\theta}^{r}} = \langle u, f \rangle.$$

Remark in particular that if $u \in L^2_{-\theta}$, then for all $f \in H^{r+1}_{\theta}$ we have

$$|\langle \partial_{x_i} u, f \rangle| = |-\langle u, \partial_{x_i} f \rangle| \leqslant C ||u||_{H_a^{-r}} ||f||_{H_a^{r+1}}, \tag{A.42}$$

so that if $u \in H_{\theta}^{-r}$ then $\nabla u \in H_{\theta}^{-(r+1)}$ with

$$\|\nabla u\|_{H_{a}^{-(r+1)}} \leqslant C\|u\|_{H_{a}^{-r}}.\tag{A.43}$$

With this structure, since L^2_{θ} is reflexive the closure of $\mathcal{L}_{\theta'}$, seen as an operator on $(L^2_{\theta})'$, is the adjoint of $\mathcal{L}^*_{\theta'}$ ([25], Th. 5.29) and is thus sectorial and defines an analytical semi-group $e^{t\mathcal{L}_{\theta'}}$ in H^{-r}_{θ} . In the same way, since H^r_{θ} is reflexive, the adjoint of $e^{t\mathcal{L}^*_{\theta'}}$ seen as an operator on H^r_{θ} is $e^{t\mathcal{L}_{\theta'}}$ seen as an operator on H^{-r}_{θ} ([32], Cor. 10.6).

From Proposition A.2 and the structure described above we deduce directly the following estimates for the semi-group induced by $\mathcal{L}_{\theta'}$ (recall (1.20)) in H_{θ}^{-r} and t > 0.

Proposition A.3. For all $0 < \theta \le \theta'$ the operator $\mathcal{L}_{\theta'}$ is sectorial and generates an analytical semi-group in H_{θ}^{-r} . Moreover we have the following estimates: for any $r \ge 0$, $\alpha \ge 0$ and $\lambda < \theta \min(k_1, \ldots, k_d)$ there exists a constant C > 0 such that for all $u \in H_{\theta}^{-(r+\alpha)}$,

$$\|e^{t\mathcal{L}_{\theta'}}u\|_{H_{\theta}^{-r}} \le C\left(1 + t^{-\alpha/2}e^{-\lambda t}\right)\|u\|_{H_{\theta}^{-(r+\alpha)}},$$
 (A.44)

and for all $r \ge 1$,

$$\|e^{t\mathcal{L}_{\theta'}}\nabla u\|_{H_{\theta}^{-r}} \le Ct^{-\frac{1}{2}}e^{-\lambda t}\|u\|_{H_{\theta}^{-r}}.$$
 (A.45)

Moreover for all $r \ge 0$, $0 < \varepsilon \le 1$ and $s \ge 0$,

$$\left\| \left(e^{(t+s)\mathcal{L}_{\theta'}} - e^{t\mathcal{L}_{\theta'}} \right) u \right\|_{H_{\theta}^{-r}} \leqslant C s^{\varepsilon} t^{-\frac{1}{2} - \varepsilon} e^{-\lambda t} \left\| u \right\|_{H_{\theta}^{-(r+1)}}. \tag{A.46}$$

Finally, there exist $r_0 > 0$, C > 0 such that for any $0 < \theta \leqslant \theta'$, for all $r > r_0$, t > 0 and all $u \in H_{\theta}^{-r}$ satisfying $\int u = 0$,

$$\|e^{t\mathcal{L}_{\theta'}}u\|_{H_{\theta}^{-r}} \leqslant Ce^{-\lambda t} \|u\|_{H_{\theta}^{-r}}.$$
 (A.47)

Proof of Proposition A.3. The spectral structure of $\mathcal{L}_{\theta'}$ follows directly from the one of $\mathcal{L}_{\theta'}^*$. To prove the first estimate of the proposition it is now sufficient to remark that for all $f \in H^r_{\theta}, u \in L^2_{-\theta}$,

$$\left| \left\langle e^{t\mathcal{L}_{\theta'}} u, f \right\rangle \right| = \left| \left\langle u, e^{t\mathcal{L}_{\theta'}^*} f \right\rangle \right| \leqslant C \left(1 + t^{-\alpha/2} e^{-\lambda t} \right) \|f\|_{H_{\theta}^r} \|u\|_{H_{\theta}^{-(r+\alpha)}}.$$

For the second point,

$$\left| \left\langle e^{t\mathcal{L}_{\theta'}} \nabla u, f \right\rangle \right| = \left| \left\langle u, \nabla e^{t\mathcal{L}_{\theta'}^*} f \right\rangle \right| \leqslant C t^{-\alpha/2} e^{-\lambda t} \left\| f \right\|_{H_{\theta}^r} \left\| u \right\|_{H_{\theta}^{-(r+\alpha)}}.$$

The third point follows from similar estimates. For the last point, remark that if $\langle u, 1 \rangle = 0$,

$$\left| \langle e^{t\mathcal{L}_{\theta'}} u, f \rangle \right| = \left| \left\langle u, e^{t\mathcal{L}_{\theta'}^*} f \right\rangle \right| = \left| \left\langle u, e^{t\mathcal{L}_{\theta'}^*} f - \frac{\int_{\mathbb{R}^d} e^{t\mathcal{L}_{\theta'}^*} f w_{\theta}}{\int_{\mathbb{R}^d} w_{\theta}} \right\rangle \right|$$

$$\leq C t^{-\alpha/2} e^{-\lambda t} \left\| f - \int f w_{\theta} \right\|_{H_{\theta}^{-}} \|u\|_{H_{\theta}^{-(r+\alpha)}}, \quad (A.48)$$

and $||f - \int f w_{\theta}||_{H_{\rho}^{r}} \leq 2||f||_{H_{\theta}^{r}}$.

Appendix B. Grönwall Lemma

Lemma B.1. Let $t \mapsto y_t$ be a nonnegative and continuous function on [0,T) satisfying, for all $t \in [0,T)$ and some positive constants c_0 and c_1 ,

$$y_t \leqslant c_0 + c_1 \int_0^t \left(1 + \frac{1}{\sqrt{t-s}} \right) y_s \, \mathrm{d}s.$$
 (B.1)

Then for all $t \in [0,T)$, $y_t \leq 2c_0e^{\alpha t}$ with $\alpha = 2c_1 + 4c_1^2\left(\Gamma\left(\frac{1}{2}\right)\right)^2$, where Γ is the usual special function $\Gamma(r) = \int_0^\infty x^{r-1}e^{-x}dx$.

For the proof of this Lemma, see [20], Lemma 5.2.

Lemma B.2. Let $a, b, \lambda > 0$ and ϕ a nonnegative measurable function on $[0, +\infty)$ such that ϕ is integrable on $[0, +\infty)$. Suppose that $t \ge 0 \mapsto u_t$ is a nonnegative function satisfying

$$u_t \leqslant a + b \int_0^t (1 + \phi(t - s)) e^{-\lambda s} u_s ds.$$
 (B.2)

Then, there exists some constant $C(b, \phi) > 0$ such that

$$\sup_{t \geqslant 0} u_t \leqslant 2a \exp\left(\frac{C(b,\phi)}{\lambda}\right). \tag{B.3}$$

Proof of Lemma B.2. Define $A = A(b, \phi) \ge 0$ such that $\int_0^{+\infty} \phi(u) \mathbf{1}_{\{\phi(u) \ge A\}} du \le \frac{1}{2b}$. Then, for all $v \le t$

$$\begin{aligned} u_{v} &\leqslant a + b \int_{0}^{v} e^{-\lambda s} u_{s} \mathrm{d}s + b \int_{0}^{v} \phi(v - s) \mathbf{1}_{\{\phi(v - s) \geqslant A\}} e^{-\lambda s} u_{s} \mathrm{d}s \\ &+ b \int_{0}^{v} \phi(v - s) \mathbf{1}_{\{\phi(v - s) \leqslant A\}} e^{-\lambda s} u_{s} \mathrm{d}s, \\ &\leqslant a + b \int_{0}^{v} e^{-\lambda s} u_{s} \mathrm{d}s + b \sup_{s \leqslant v} u_{s} \int_{0}^{v} \phi(v - s) \mathbf{1}_{\{\phi(v - s) \geqslant A\}} \mathrm{d}s + bA \int_{0}^{v} e^{-\lambda s} u_{s} \mathrm{d}s, \\ &\leqslant a + b (1 + A) \int_{0}^{v} e^{-\lambda s} u_{s} \mathrm{d}s + \frac{1}{2} u_{v}^{*} \leqslant a + b (1 + A) \int_{0}^{t} e^{-\lambda s} u_{s}^{*} \mathrm{d}s + \frac{1}{2} u_{t}^{*}, \end{aligned}$$

where we have defined $u^*(s) := \sup_{r \leq s} u_r$. Since the last inequality is true for all $v \leq t$, we get

$$u_t^* \le 2a + 2b(1+A) \int_0^t e^{-\lambda s} u_s^* ds.$$

The usual Grönwall lemma applied to $t \mapsto u_t^*$ gives the conclusion, for $C(b, \phi) = 2b(1 + A(b, \phi))$.

ACKNOWLEDGEMENTS

Both authors benefited from the support of the ANR-19-CE40-0023 (PERISTOCH), C. Poquet from the ANR-17-CE40-0030 (Entropy, Flows, Inequalities), E. Luçon from the ANR-19-CE40-002 (ChaMaNe).

References

- [1] D. Bakry, I. Gentil, M. Ledoux Analysis and Geometry of Markov Diffusion Operators. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 348. Springer, 2014.
- [2] J. Baladron, D. Fasoli, O. Faugeras, and J. Touboul. Mean-field description and propagation of chaos in networks of Hodgkin-Huxley and FitzHugh-Nagumo neurons. The Journal of Mathematical Neuroscience, 2(1):10, 2012.
- [3] S. Barland, O. Piro, M. Giudici, J. R. Tredicce, and S. Balle. Experimental evidence of van der polfitzhugh-nagumo dynamics in semiconductor optical amplifiers. *Phys. Rev. E*, 68:036209, Sep 2003.
- [4] P.W. Bates, K. Lu, and C. Zeng. Existence and persistence of invariant manifolds for semiflows in Banach space. *American Mathematical Soc.*, 1998, Volume 645.
- [5] P.W. Bates, K. Lu, and C. Zeng. Approximately invariant manifolds and global dynamics of spike states. *Inventiones mathematicae*, volume 174, pages 355–422, 2008.
- [6] R. Beals and R. Wong. Special Functions and Orthogonal Polynomials. Cambridge University Press, 2016.
- [7] S. Benachour, B. Roynette, and P. Vallois. Nonlinear self-stabilizing processes. II. Convergence to invariant probability. *Stochastic Process. Appl.*, 75(2):203–224, 1998.
- [8] L. Bertini, G. Giacomin and C. Poquet. Synchronization and random long time dynamics for mean-field plane rotators. *Probability Theory and Related Fields*, volume 160(3-4), 593–653, 2014.
- [9] M. Bossy, O. Faugeras, and D. Talay. Clarification and complement to "Mean-field description and propagation of chaos in networks of Hodgkin-Huxley and FitzHugh-Nagumo neurons". J. Math. Neurosci., 5:Art. 19, 23, 2015.
- [10] H. Brezis. Analyse fonctionnelle. Collection Mathématiques Appliquées pour la Maîtrise. [Collection of Applied Mathematics for the Master's Degree]. Masson, Paris, 1983. Théorie et applications. [Theory and applications].
- [11] P. Cattiaux, A. Guillin, and F. Malrieu. Probabilistic approach for granular media equations in the non-uniformly convex case. *Probability Theory and Related Fields*, 140(1):19–40, Jan 2008.
- [12] R. Cerf, P. Dai Pra, M. Formentin, and D. Tovazzi. Rhythmic behavior of an Ising Model with dissipation at low temperature. ALEA, Lat. Am. J. Probab. Math. Stat. 18, 439-467 (2021)
- [13] F. Collet, P. Dai Pra and M. Formentin. Collective periodicity in mean-field models of cooperative behavior. *Nonlinear Differ. Equ. Appl.* DEA22(5), 1461–1482, 2015.
- [14] Q. Cormier, E. Tanré, R. Veltz. Hopf bifurcation in a Mean-Field model of spiking neurons. arXiv preprint, arXiv:2008.11116.
- [15] R. Dahms. Long time behavior of a spherical mean field model. PhD Thesis, Technische Universität Berlin, Fakultät II Mathematik und Naturwissenschaften, 2002.
- [16] P. Dai Pra, M. Formentin and G. Pelino Oscillatory behavior in a model of non-Markovian mean field interacting spins. *Journal of Statistical Physics*, 179 (3), 690–712, 2020.
- [17] S. Ditlevsen and E. Löcherbach. Multi-class oscillating systems of interacting neurons. Stochastic Processes and their Applications, 127(6):1840–1869, 2017.
- [18] N. Fenichel. Persistence and smoothness of invariant manifolds for flows. *Indiana Univ. Math. J.*, 21:193–226, 1972.
- [19] N. Fenichel. Geometric singular perturbation theory for ordinary differential equations. Journal of Differential Equations, 31:53–98, 1979.
- [20] G. Giacomin, K. Pakdaman, X. Pellegrin, Xavier and C. Poquet, Transitions in active rotator systems: invariant hyperbolic manifold approach. SIAM Journal on Mathematical Analysis, volume 44, pages 4165–4194, 2012.
- [21] G. Giacomin and C. Poquet. Noise, interaction, nonlinear dynamics and the origin of rhythmic behaviors. *Brazilian Journal of Probability and Statistics*, 29(2):460–493, 2015.
- [22] J. Guckenheimer. Isochrons and phaseless sets. *Journal of Mathematical Biology*, 1(3):259–273, Sept. 1975.
- [23] D. Henry. Geometric theory of semilinear parabolic equations. Springer, 1981.
- [24] M. W. Hirsch, C. C. Pugh, and M. Shub. *Invariant manifolds*. Lecture Notes in Mathematics, Vol. 583. Springer-Verlag, Berlin, 1977.
- [25] T. Kato. Perturbation theory for linear operators. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.

- [26] B. Lindner, J. Garcia-Ojalvo, A. Neiman, and L. Schimansky-Geier. Effects of noise in excitable systems. *Physics Reports*, 392(6):321 – 424, 2004.
- [27] E. Lucon and C. Poquet. Emergence of oscillatory behaviors for excitable systems with noise and mean-field interaction, a slow-fast dynamics approach. Communications in Mathematical Physics, vol. 373, no 3, p. 907–969, 2020.
- [28] E. Luçon and C. Poquet. Periodicity induced by noise and interaction in the kinetic mean-field FitzHugh-Nagumo model. *Annals of Applied Probability*, Vol. 31, No. 2, 561–593 (2021).
- [29] E. Luçon and C. Poquet. Periodicity and longtime diffusion for mean field systems in \mathbb{R}^d . arXiv e-prints, 2021.
- [30] H. P. McKean. A Class of Markov Processes Associated with Nonlinear Parabolic Equations. Proc. Natl. Acad. Sci. USA, 56(6), 1907–1911, 1966.
- [31] S. Mischler, C. Quiñinao, and J. Touboul. On a Kinetic Fitzhugh-Nagumo Model of Neuronal Network. Communications in Mathematical Physics, 342(3):1001–1042, 2016.
- [32] A. Pazy. Semigroups of linear operators and applications to partial differential equations, volume 44 of Applied Mathematical Sciences. Springer-Verlag, New York, 1983.
- [33] C. Quiñinao and J. D. Touboul. Clamping and Synchronization in the strongly coupled FitzHugh-Nagumo model. SIAM J. Appl. Dyn. Syst., 19(2), 788–827, 2020.
- [34] C. Rocsoreanu, A. Georgescu, and N. Giurgiţeanu. The FitzHugh-Nagumo model, volume 10 of Mathematical Modelling: Theory and Applications. Kluwer Academic Publishers, Dordrecht, 2000. Bifurcation and dynamics.
- [35] M. Scheutzow. Periodic behavior of the stochastic Brusselator in the mean-field limit. Probability theory and related fields, 72(3):425–462, 1986.
- [36] G.R. Sell and Y. You. Dynamics of evolutionary equations. Springer Science & Business Media, 2013.
- [37] Sznitman, A.-S. Topics in propagation of chaos. *Ecole d'été de probabilités de Saint-Flour XIX-1989*, Springer, 165–251, 1991.
- [38] G. Teschl. Ordinary differential equations and dynamical systems, volume 140 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2012.
- [39] J. Tugaut. Convergence to the equilibria for self-stabilizing processes in double-well landscape. The Annals of Probability, 41(3A):1427 – 1460, 2013.
- [40] S. Wiggins. Normally hyperbolic invariant manifolds in dynamical systems. Springer Science & Business Media, volume 105, 2013.

Université de Paris, Laboratoire MAP5 (UMR CNRS 8145), 75270 Paris, France & FP2M, CNRS FR 2036, eric.lucon@u-paris.fr.

UNIVERSITÉ CLAUDE BERNARD LYON 1, CNRS UMR 5208, INSTITUT CAMILLE JORDAN, F-69622 VILLEURBANNE, FRANCE, POQUET@MATH.UNIV-LYON1.FR