

# Normal and stable approximation to subgraph counts in superpositions of Bernoulli random graphs

Mindaugas Bloznelis<sup>1</sup>, and Joonas Karjalainen<sup>2</sup>, and Lasse Leskelä<sup>2</sup>

<sup>1</sup>Institute of Informatics, Vilnius University  
Naugarduko 24, LT-03225 Vilnius, Lithuania

<sup>2</sup>Department of Mathematics and Systems Analysis  
School of Science, Aalto University  
Otakaari 1, FI-02150 Espoo, Finland

## Abstract

The clustering property of complex networks indicates the abundance of small dense subgraphs in otherwise sparse networks. For a community-affiliation network defined by a superposition of Bernoulli random graphs, which has a nonvanishing global clustering coefficient and a power-law degree distribution, we establish normal and  $\alpha$ -stable approximations to the number of small cliques, cycles and more general 2-connected subgraphs.

## 1 Introduction and results

Mathematical modeling of complex networks aims at explaining and reproducing characteristic properties of large real-world networks, e.g., power-law degree distributions and clustering. By clustering property we mean the tendency of nodes to cluster together by forming relatively small groups with a high density of ties within a group. Locally, in the vicinity of a vertex, clustering can be measured by the local clustering coefficient, the probability that two randomly selected neighbors of the vertex are adjacent. Globally, the fraction of wedges (paths of length 2) that induce triangles defines the global clustering coefficient, which represents the probability that endpoints of a randomly selected wedge are adjacent. Non-vanishing clustering coefficients indicate the abundance of triangles and other small dense subgraphs in otherwise sparse networks. The problem of determining the asymptotic distribution of dense subgraph counts in sparse complex networks is of considerable interest, but there are very few results obtained so far.

In the present study we establish normal and  $\alpha$ -stable approximations of the numbers of  $k$ -cliques,  $k$ -cycles and more general 2-connected subgraphs in a community-affiliation network model defined by a superposition of Bernoulli random graphs [3, 15, 16].

To the best of our knowledge this is the first systematic study of an  $\alpha$ -stable approximation to subgraph counts in a theoretical model of sparse affiliation networks. We note that in the network model considered, the clustering property and the power-law degree distribution, two basic properties of complex networks, are essential for an  $\alpha$ -stable limit to emerge.

## 1.1 Network model

Let  $(X, Q)$  be a random vector with values in  $\{0, 1, 2, \dots\} \times [0, 1]$  and let

$$\{G(x, p) : x \in \{1, 2, \dots\}, p \in [0, 1]\}$$

be a family of Bernoulli random graphs independent of  $(X, Q)$ . We set  $[x] = \{1, 2, \dots, x\}$  to be the vertex set of  $G(x, p)$ . Recall that in  $G(x, p)$  every pair of vertices  $\{i, j\} \subset [x]$  is declared adjacent independently at random with probability  $p$ . For notational convenience we introduce the empty graph  $G_\emptyset$  having no vertices and set  $G(0, p) = G_\emptyset$  for any  $p \in [0, 1]$ . We define the mixture of Bernoulli random graphs  $G(X, Q)$  in a natural way.

Let  $(X_1, Q_1), (X_2, Q_2), \dots$  be a sequence of independent copies of  $(X, Q)$ . Given  $X_1, \dots, X_m$ , let  $\mathcal{V}_{n,i} = \mathcal{V}_{n,i}(X_i)$ ,  $1 \leq i \leq m$ , be independent random subsets of  $[n]$  defined as follows. For  $X_i \leq n$  we select  $\mathcal{V}_{n,i}$  uniformly at random from the class of subsets of  $[n]$  of size  $X_i$ . For  $X_i > n$  we set  $\mathcal{V}_{n,i} = [n]$ . We denote  $\tilde{X}_i = |\mathcal{V}_{n,i}| = X_i \wedge n$ . Let  $G_{n,i}$ ,  $1 \leq i \leq m$ , be independent random graphs with vertex sets  $\mathcal{V}_{n,i}$  defined as follows. We obtain  $G_{n,i}$  by a one-to-one mapping of vertices of  $G(\tilde{X}_i, Q_i)$  to the elements of  $\mathcal{V}_{n,i}$  and by retaining the adjacency relations of  $G(\tilde{X}_i, Q_i)$ . We denote by  $\mathcal{E}_{n,i}$  the edge set of  $G_{n,i}$ . Finally, let  $G_{[n,m]} = (V, \mathcal{E})$  be the random graph with the vertex set  $V = [n]$  and edge set  $\mathcal{E} = \mathcal{E}_{n,1} \cup \dots \cup \mathcal{E}_{n,m}$ . Therefore  $G_{[n,m]}$  is the superposition of  $G_{n,1}, \dots, G_{n,m}$ . We call the contributing random graphs  $G_{n,i}$  layers or communities.

The random graph  $G_{[n,m]}$  represents a null model of a community-affiliation network [15, 16], which has attracted considerable attention in the literature. In the particular case where  $Q \equiv 1$  the random graph  $G_{[n,m]}$  goes back to the ‘passive’ model of random intersection graph [5]. In the parameter regime  $m = \Theta(n)$  as  $m, n \rightarrow +\infty$  the random graph  $G_{[n,m]}$  admits a power-law degree distribution with tunable power-law exponent, nonvanishing global clustering coefficient and tunable clustering spectrum [3]. Moreover, it admits a limiting bidegree distribution with (stochastically dependent) power-law marginals shown in [4]. The present paper continues the study of the random graph  $G_{[n,m]}$  and focuses on the asymptotic distributions of (dense) subgraph counts.

## 1.2 Results

Let  $F = (\mathcal{V}_F, \mathcal{E}_F)$  be a graph with vertex set  $\mathcal{V}_F$  and edge set  $\mathcal{E}_F$ . We denote  $v_F = |\mathcal{V}_F|$  and  $e_F = |\mathcal{E}_F|$ . We assume in what follows that  $F$  is 2-connected. That is,  $F$  is connected and, moreover, it stays connected even if we remove any one of its vertices. We call  $F$  balanced if  $e_F/v_F = \max\{e_H/v_H : H \subset F \text{ with } e_H \geq 1\}$ . For example, the cycle  $\mathcal{C}_k$  and clique  $\mathcal{K}_k$  (where  $k$  stands for the number of vertices) are 2-connected and balanced. Let  $N_F$  be the number of copies of  $F$  in  $G(X, Q)$ . Denote  $\sigma_F^2 = \mathbf{Var}N_F$ , the variance of  $N_F$ . We write  $\sigma_F^2 < \infty$  if the variance is finite and  $\sigma_F^2 = \infty$  otherwise. We use the shorthand notation  $N_F^* := \mathbf{E}(N_F|X, Q) = a_F \binom{X}{v_F} Q^{e_F}$ , where  $a_F$  stands for the number of distinct copies of  $F$  in the complete graph on  $v_F$  vertices. We have, for example, that  $N_{\mathcal{C}_k}^* = (X)_k Q^k / (2k)$  and  $N_{\mathcal{K}_k}^* = (X)_k Q^{\binom{k}{2}} / k!$ . Here and below  $(x)_k = x(x-1)\dots(x-k+1)$  denotes the falling factorial.

Let  $\mathcal{N}_F$  be the number of copies of  $F$  in  $G_{[n,m]}$ . Our first result establishes the asymptotic normality of  $\mathcal{N}_F$ .

**Theorem 1.** *Let  $\nu > 0$ . Let  $n, m \rightarrow +\infty$  and assume that  $m/n \rightarrow \nu$ . Let  $F$  be a 2-connected graph with  $v_F \geq 3$  vertices. Assume that  $\mathbf{E}X < \infty$  and  $0 < \sigma_F^2 < \infty$ . Assume, in addition, that*

$$\mathbf{E} \left( X^{1+s} \left(1 - \frac{1}{2e_F}\right) Q^s \right) < \infty, \quad \text{for each } s = 1, 2, \dots, v_F - 1. \quad (1)$$

*Then  $(\mathcal{N}_F - \mathbf{E}\mathcal{N}_F) / (\sigma_F \sqrt{m})$  converges in distribution to the standard normal distribution.*

**Remark 1.** For a balanced graph  $F$ , the finite variance condition  $\sigma_F^2 < \infty$  is equivalent to the second moment condition  $\mathbf{E}(N_F^*)^2 < \infty$ . In particular, we have

$$\sigma_F^2 < \infty \Leftrightarrow \mathbf{E}(X^{2v_F} Q^{2e_F}) < \infty. \quad (2)$$

**Remark 2.** In the special case, where  $F$  is a clique on  $k \geq 3$  vertices ( $F = \mathcal{K}_k$ ) condition (1) can be replaced by the following one

$$\mathbf{E}\left(X^{r - \frac{\hat{r}}{k(k-1)}} Q^{\hat{r}}\right) < \infty, \quad \text{for each } r = 2, \dots, k. \quad (3)$$

where we denote  $\hat{r} := \binom{r-1}{2} + 1$ .

The proof of Remarks 1 and 2 is presented in Section 2. Let us briefly explain the result and conditions of Theorem 1. Let  $N_{F,i}$  be the number of copies of  $F$  in  $G(X_i, Q_i)$  and define  $S_F = N_{F,1} + \dots + N_{F,m}$ . The moment condition  $\mathbf{E}X < \infty$  and the assumption  $m \approx \nu n$  control the amount of overlap between the different layers  $G_{n,i}$  and ensures that (the layer sizes)  $\tilde{X}_i = X_i$ ,  $1 \leq i \leq m$  with high probability. The principal contribution to the subgraph count  $\mathcal{N}_F$  comes from the subgraph counts  $N_{F,i}$  of individual layers (recall that  $F$  is 2-connected). Therefore we have  $\mathcal{N}_F \approx S_F$ . To make this approximation rigorous we introduce conditions (1) and (3) aimed at controlling the number of overlaps of different copies of  $F$  in  $G_{[n,m]}$ . (The combinatorial origin of (1), (3) is explained in Lemmas 1–4). Finally, the asymptotic normality of  $\mathcal{N}_F$  follows from the asymptotic normality of  $S_F$ . The latter is guaranteed by the second moment condition  $\sigma_F^2 < \infty$ .

In the case where  $F$  is balanced and the random variable  $N_F^*$  has an infinite second moment, we can obtain an  $\alpha$ -stable limiting distribution for the subgraph count  $\mathcal{N}_F$ . In Theorem 2 below we assume that for some  $a > 0$  and  $0 < \alpha < 2$ , we have

$$\mathbf{P}\{N_F^* > t\} = (a + o(1))t^{-\alpha} \quad \text{as } t \rightarrow +\infty. \quad (4)$$

Let  $N_{F,i}^* = \mathbf{E}(N_F | X_i, Q_i)$ ,  $1 \leq i \leq m$ , be iid copies of  $N_F^*$  and put  $S_F^* = N_{F,1}^* + \dots + N_{F,m}^*$ . It is well known (Theorem 2 in § 35 of [6]) that the distribution of  $m^{-1/\alpha}(S_F^* - B_m)$  converges to a stable distribution, say  $G_{\alpha,a}$ , which is defined by  $a$  and  $\alpha$ . Here  $B_m = m\mathbf{E}N_F^* = \mathbf{E}N_F$  for  $1 < \alpha < 2$  and  $B_m \equiv 0$  for  $0 < \alpha < 1$ . For  $\alpha = 1$  we have  $B_m = c_{\alpha,a}^* \ln m$ , where the constant  $c_{\alpha,a}^* > 0$  depends on  $a$  and  $\alpha$ .

Our second result establishes an  $\alpha$ -stable approximation to the distribution of  $\mathcal{N}_F$ .

**Theorem 2.** Let  $\nu > 0$ . Let  $n, m \rightarrow +\infty$  and assume that  $m/n \rightarrow \nu$ . Let  $F$  be a balanced and 2-connected graph with  $v_F \geq 3$  vertices. Let  $a > 0$  and  $0 < \alpha < 2$ . Assume that  $\mathbf{E}X < \infty$  and that (4) holds. Assume, in addition, that

$$\mathbf{E}\left(X^{1+s\left(1-\frac{1}{\alpha e_F}\right)} Q^s\right) < \infty \quad \text{for each } s = 1, \dots, v_F - 1. \quad (5)$$

Then  $(\mathcal{N}_F - B_m)/m^{1/\alpha}$  converges in distribution to  $G_{\alpha,a}$ .

**Remark 3.** In the special case, where  $F$  is a clique on  $k \geq 3$  vertices ( $F = \mathcal{K}_k$ ) condition (5) can be replaced by the following one

$$\mathbf{E}\left(X^{r - \hat{r} \frac{2}{\alpha k(k-1)}} Q^{\hat{r}}\right) < \infty \quad \text{for each } r = 2, \dots, k. \quad (6)$$

where  $\hat{r} = \binom{r-1}{2} + 1$ .

The result of Theorem 2 is obtained using the same approximation  $\mathcal{N}_F \approx S_F$  as above. In addition, we use the observation that condition (4) implies  $S_F \approx S_F^*$ . To make this approximation rigorous we apply exponential large deviation bounds [10] to individual subgraph counts  $N_{F,i}$  conditionally given  $(X_i, Q_i)$ , see Lemma 5. The  $\alpha$ -stable limit of  $S_F^*$  is now guaranteed by condition (4) and Theorem 2 in § 35 of [6].

We briefly comment on technical conditions (1), (3), (5) and (6). The mixed moments defined there appear in our upper bounds on the expected number of overlaps of different copies of  $F$  in  $G_{[n,m]}$ , see Lemmas 1, 4 and inequality (11) in the proof below. More precisely, we use these moments to upper bound the quantity  $h_F$  of (11). Alternatively, one can impose conditions on the rate of decay of  $h_F$  directly. We note that for particular graphs  $F$  the moment conditions (1), (3), (5), (6) can be relaxed.

Let us examine Theorems 1 and 2 in the special case where the marginals  $X, Q$  of  $(X, Q)$  are independent and  $\mathbf{P}\{Q > 0\} > 0$ . We first consider Theorem 1. The finite variance condition  $\sigma_F^2 < \infty$  of Theorem 1 reduces to the moment condition  $\mathbf{E}X^{2v_F} < \infty$ . Indeed, by the simple inequality  $N_F \leq (X)_{v_F}$ , we have that  $\mathbf{E}X^{2v_F} < \infty \Rightarrow \mathbf{E}N_F^2 < \infty \Rightarrow \sigma_F^2 < \infty$ . On the other hand, by the variance identity  $\mathbf{Var}N_F = \mathbf{Var}N_* + \mathbf{E}(\mathbf{Var}(N_F|X, Q))$ , we have that  $\sigma_F^2 < \infty \Rightarrow \mathbf{E}(N_*)^2 < \infty$ , where the latter inequality (for independent  $X$  and  $Q$ ) implies  $\mathbf{E}X^{2v_F} < \infty$ . Moreover, the moment condition  $\mathbf{E}X^{2v_F} < \infty$  implies (1). Therefore Theorem 1 establishes the asymptotic normality under the minimal second moment condition  $\sigma_F^2 < \infty$ .

We now turn to Theorem 2. For independent  $X$  and  $Q$  condition (4) of Theorem 2 is equivalent to the condition

$$\mathbf{P}\{X > t\} = (b + o(1))t^{-\gamma} \quad \text{as } t \rightarrow +\infty, \quad (7)$$

where  $\gamma = \alpha v_F$  and where  $b$  solves the equation  $a = b(a_F/v_F!)^{\gamma/v_F} \mathbf{E}Q^{\gamma e_F/v_F}$ . Note that  $\mathbf{E}X < \infty$  implies  $\gamma > 1$ . Furthermore, the inequality  $v_F \leq e_F$  (which holds for any 2-connected  $F$  with  $v_F \geq 3$ ) combined with  $\gamma > 1$  implies  $\alpha e_F > 1$ . Observe that for  $\alpha e_F > 1$  condition (5) reads as  $\mathbf{E}X^{1+(v_F-1)(1-\frac{1}{\alpha e_F})} < \infty$ . In view of (7) the latter expectation is finite whenever

$$1 + (v_F - 1) \left(1 - \frac{1}{\alpha e_F}\right) < \gamma. \quad (8)$$

We have arrived to the following corollary.

**Corollary 1.** *Let  $\nu > 0$ . Let  $n, m \rightarrow +\infty$  and assume that  $m/n \rightarrow \nu$ . Let  $F$  be a 2-connected graph with  $v_F \geq 3$  vertices. Assume that  $X$  and  $Q$  are independent and  $\mathbf{P}\{Q > 0\} > 0$ .*

(i) *If  $\mathbf{E}X^{2v_F} < \infty$  then  $(\mathcal{N}_F - \mathbf{E}\mathcal{N}_F)/(\sigma_F \sqrt{m})$  converges in distribution to the standard normal distribution.*

(ii) *Let  $b > 0$  and  $1 < \gamma < 2v_F$ . Assume that (7) holds. Assume, in addition, that  $F$  is balanced and (8) holds, where  $\alpha = \gamma/v_F$ . Then  $(\mathcal{N}_F - B_m)/m^{1/\alpha}$  converges in distribution to  $G_{\alpha,a}$ . Here  $B_m$  and  $G_{\alpha,a}$  are the same as in Theorem 2, with  $a = b(a_F/v_F!)^{\gamma/v_F} \mathbf{E}Q^{\gamma e_F/v_F}$ .*

It is relevant to mention that the moment condition  $\mathbf{E}X < \infty$  together with the assumption  $m \approx \nu n$  imply the existence of an asymptotic degree distribution of  $G_{[n,m]}$  as  $n, m \rightarrow +\infty$ . An asymptotic power-law degree distribution is obtained if we choose an appropriate distribution for the layer type  $(X, Q)$ . Furthermore, under an additional moment condition  $\mathbf{E}X^3 Q^2 < \infty$  the random graph  $G_{[n,m]}$  has a non-vanishing global clustering coefficient, see [3]. Therefore, Theorems 1 and 2 establish the limit distributions of subgraph counts in a highly clustered complex network.

Finally, we discuss an important question about the relation between the community size  $X$  and strength  $Q$ . In Theorems 1 and 2, no assumption has been made about the stochastic dependence between the marginals  $X$  and  $Q$  of the bivariate random vector  $(X, Q)$  defining the random graph  $G_{[n,m]}$ . To simplify the model we can assume that  $X$  and  $Q$  are independent, see Corollary 1 above. However, for network modelling purposes, various types of dependence between  $X$  and  $Q$  are of interest. For example, a negative correlation between  $X$  and  $Q$  would emphasize small strong communities and large weak communities, a pattern likely to occur in real networks with overlapping communities. Assuming that  $Q$  is proportional to a negative power of  $X$ , for example,  $Q = \min\{1, bX^{-\beta}\}$  for some  $\beta \geq 0$  and  $b > 0$  (cf. [15], [16]), one obtains a mathematically tractable network model admitting tunable power-law degree and bidegree distributions and rich clustering spectrum [3, 4].

*Related work.* Asymptotic distributions of subgraph counts in Bernoulli random graphs is a well established area of research, see, e.g., [9], [14] and references therein. For a recent

development we refer to [8], [12], [13]. A significant difference between the sparse Bernoulli random graphs and complex networks is that the former ones have none or very few copies of a triangle or a larger clique, while the latter ones often have abundant numbers of those. The abundance of dense subgraphs in otherwise sparse complex networks is related to the clustering property. The global and local clustering coefficients are expressed in terms of counts of triangles and wedges. Therefore, a rigorous asymptotic analysis of clustering coefficients in large random networks reduces to that of the triangle counts and wedge counts. In particular, the bivariate asymptotic normality for triangle and wedge counts in a related sparse random intersection graph was shown in [2], and related  $\alpha$ -stable limits were established in [1]. Another line of research pursued in [7, 11] addresses the concentration of subgraph counts in  $G_{[n,m]}$ .

The rest of the paper is organized as follows: In Section 2 we formulate and prove Theorems 1, 2 and Remarks 1, 2, 3. We mention that combinatorial Lemmas 2, 3 and inequality (16), see below, may be of independent interest.

## 2 Proofs

### 2.1 Notation

Before the proof we introduce some notation. We denote for short  $\mathbb{X} = (X_1, \dots, X_m)$  and  $\mathbb{Q} = (Q_1, \dots, Q_m)$ . By  $\mathbf{E}^*(\cdot) = \mathbf{E}(\cdot | \mathbb{X}, \mathbb{Q})$  and  $\mathbf{P}^*(\cdot) = \mathbf{P}(\cdot | \mathbb{X}, \mathbb{Q})$  we denote the conditional expectation and probability given  $(\mathbb{X}, \mathbb{Q})$ . Recall that  $a_F$  stands for the number of copies of  $F$  in the complete graph on  $v_F$  vertices  $\mathcal{K}_{v_F}$ . For example  $a_{\mathcal{K}_k} = 1$  and  $a_{\mathcal{C}_k} = (k-1)!/2$ . Given  $F$ , for any positive sequences  $\{a_n\}$  and  $\{b_n\}$  we denote  $a_n \succ b_n$  (respectively  $a_n \prec b_n$ ) whenever for sufficiently large  $n$  we have  $c_1 \leq a_n/b_n \leq c_2$  (respectively  $a_n \leq c_2 b_n$ ), where constants  $0 < c_1 < c_2$  may only depend on  $v_F$ .

Recall that  $N_F$  and  $N_{F,i}$  denote the numbers of copies of  $F$  in  $G(X, Q)$  and  $G(X_i, Q_i)$ , respectively. Furthermore,  $N_F^* = \mathbf{E}(N_F | X, Q)$ ,  $N_{F,i}^* = \mathbf{E}(N_{F,i} | X_i, Q_i)$ , and

$$S_F = N_{F,1} + \dots + N_{F,m}, \quad S_F^* = N_{F,1}^* + \dots + N_{F,m}^*.$$

Note that  $N_{F,i}^* = \mathbf{E}^*(N_{F,i})$  and  $S_F^* = \mathbf{E}^*(S_F)$ . Finally, let  $\tilde{N}_i$  be the number of copies of  $F$  in  $G_{n,i}$  and let  $\tilde{S} = \tilde{N}_1 + \dots + \tilde{N}_m$ .

We can identify the indices  $1 \leq i \leq m$  with colors, and assign (the edges of) each  $G_{n,i}$  the color  $i$ . The colored graph is denoted by  $G_{n,i}^*$ . The union of colored graphs  $G_{n,1}^* \cup \dots \cup G_{n,m}^*$  defines a multigraph, denoted by  $G_{[n,m]}^*$ , that admits parallel edges of different colors. Furthermore each edge  $u \sim v$  of  $G_{[n,m]}^*$  is assigned the set of colors that correspond to parallel edges of  $G_{[n,i]}^*$  connecting  $u$  and  $v$ .

A subgraph  $H \subset G_{[n,m]}^*$  is called monochromatic if it is a subgraph of some  $G_{n,i}$  and none of edges of  $H$  are assigned more than one color. Otherwise  $H$  is called polychromatic.  $\mathcal{N}_{F,M}$  and  $\mathcal{N}_{F,P}$  stand for the numbers of monochromatic and polychromatic copies of  $F$  in  $G_{[n,m]}^*$ . A subgraph  $H^* \subset G_{[n,m]}^*$  is called monochromatic if it is a subgraph of some  $G_{n,i}^*$ . It is called polychromatic if it contains edges of different colors. Given  $H^* \subset G_{[n,m]}^*$ , let  $H_0 \subset G_{[n,m]}$  be the graph obtained from  $H^*$  by merging parallel edges. We call  $H_0$  the projection of  $H^*$ . Let  $\mathcal{N}_{F,P}^*$  be the number of polychromatic copies of  $F$  in  $G_{[n,m]}^*$ . Note that there can be several monochromatic and/or polychromatic copies of  $F$  in  $G_{[n,m]}^*$  sharing the same projection  $F_0$ . We fix a copy  $F_0 \subset G_{[n,m]}$  of  $F$  and denote by  $h_F$  the expected number of polychromatic subgraphs of  $G_{[n,m]}^*$  whose projection is  $F_0$ . Clearly, the number  $h_F$  does not depend on the location of  $F_0$ .

### 2.2 Proofs

We start with an outline of the proof. We approximate  $\mathcal{N}_F \approx \tilde{S}_F$  and  $\tilde{S}_F \approx S_F$ . In the case where  $\mathbf{E}N_F^2 < \infty$  we deduce the normal approximation to the sum  $S_F$  (of iid random

variables) by the standard central limit theorem. In the case where  $N_F$  has an infinite variance we further approximate  $S_F \approx S_F^*$  and deduce the  $\alpha$ -stable approximation by the generalized central limit theorem (see Theorem 2 in § 35 of [6]).

*Approximation*  $\mathcal{N}_F \approx \tilde{S}_F$ . The approximation follows from the simple observation that

$$\mathcal{N}_F = \mathcal{N}_{F,M} + \mathcal{N}_{F,P}, \quad \mathcal{N}_{F,M} \leq \tilde{S}_F \leq \mathcal{N}_{F,M} + \mathcal{N}_{F,P}^*, \quad \mathcal{N}_{F,P} \leq \mathcal{N}_{F,P}^*. \quad (9)$$

We only comment on the second inequality. To see why it holds true, let us inspect every copy of  $F$  in  $G_{[n,m]}^*$  that belongs to two or more layers  $G_{n,i}$ . Let  $F_0 \subset G_{[n,m]}$  be such a copy. Clearly, the number of polychromatic subgraphs  $F^*$  in  $G_{[n,m]}^*$ , whose projection  $F_0^*$  is  $F_0$ , is larger than the number of monochromatic ones. Hence  $\tilde{S}_F \leq \mathcal{N}_{F,M} + \mathcal{N}_{F,P}^*$ . From (9) we conclude that

$$|\tilde{S}_F - \mathcal{N}_F| \leq \mathcal{N}_{F,P}^*. \quad (10)$$

In order to assess the accuracy of the approximation  $\mathcal{N}_F \approx \tilde{S}_F$ , we evaluate the expected value of  $\mathcal{N}_{F,P}^*$ . Let  $\mathcal{K}_{[n]}$  be a clique on the vertex set  $V = [n]$ . We couple  $G_{[n,m]} \subset \mathcal{K}_{[n]}$  and fix a subgraph  $F_0 = (\mathcal{V}_0, \mathcal{E}_0) \subset \mathcal{K}_{[n]}$  with vertex set  $\{1, \dots, v_F\} \subset V$ , which is a copy of  $F$ . We have, by symmetry,

$$\mathbf{E}\mathcal{N}_{F,P}^* = \binom{n}{v_F} a_F h_F. \quad (11)$$

Each  $F^* \subset G_{[n,m]}^*$  whose projection is  $F_0$  is defined by the partition of the edge set  $\mathcal{E}_0$  into non-empty color classes, say,  $B_1 \cup \dots \cup B_r = \mathcal{E}_0$ , and the vector of distinct colors  $(i_1, \dots, i_r) \subset [m]^r$  such that all the edges in  $B_j$  are of the color  $i_j$  (edges of  $B_j$  belong to  $G_{n,i_j}^*$ ). Denote by  $\tilde{B} = (B_1, \dots, B_r)$  and  $\tilde{i} = (i_1, \dots, i_r)$  the partition and its coloring. The polychromatic subgraph  $F^*$  defined by the pair  $(\tilde{B}, \tilde{i})$  is denoted  $F(\tilde{B}, \tilde{i})$ . The probability that such a subgraph is present in  $G_{[n,m]}^*$  is

$$h(\tilde{B}, \tilde{i}) := \mathbf{P}\{F(\tilde{B}, \tilde{i}) \in G_{[n,m]}^*\} = \prod_{j=1}^r \frac{1}{\binom{n}{v_j}} \mathbf{E}\left(\binom{\tilde{X}_{i_j}}{v_j} Q_{i_j}^{b_j}\right). \quad (12)$$

Here  $b_j := |B_j|$ , and  $v_j$  is the number of distinct vertices incident to edges from  $B_j$ . We have

$$h_F = \mathbf{E}\left(\sum_{(\tilde{B}, \tilde{i})} \mathbb{I}_{\{F(\tilde{B}, \tilde{i}) \in G_{[n,m]}^*\}}\right) = \sum_{(\tilde{B}, \tilde{i})} h(\tilde{B}, \tilde{i}). \quad (13)$$

Here the sum runs over all possible polychromatic  $F^*$  whose projection  $F_0^*$  is  $F_0$ . We upper bound  $h_F$  in Lemmas 1 and 4 below.

*Approximation*  $\tilde{S}_F \approx S_F$ . For  $1 \leq i \leq m$  we couple  $G(\tilde{X}_i, Q_i) \subset G(X_i, Q_i)$  and  $\tilde{N}_i \leq N_i$  so that  $G(\tilde{X}_i, Q_i) \neq G(X_i, Q_i)$  and  $\tilde{N}_i \neq N_i$  whenever  $X_i > n$ . For  $m = O(n)$ , the event  $\mathcal{A}_n := \{\max_{1 \leq i \leq m} X_i > n\}$  has probability

$$\mathbf{P}\{\mathcal{A}_n\} \leq \sum_{i=1}^m \mathbf{P}\{X_i > n\} \leq \frac{m}{n} \mathbf{E}(X_1 \mathbb{I}_{\{X_1 > n\}}) = o(1), \quad (14)$$

hence  $\mathbf{P}\{\tilde{S}_F \neq S_F\} = o(1)$ . In (14) we used the fact that  $\mathbf{E}X_1 < \infty \Rightarrow \mathbf{E}(X_1 \mathbb{I}_{\{X_1 > n\}}) = o(1)$ .

*Proof of Theorem 1 and Remark 2.* By Lemma 1 (respectively, Lemma 4), we have  $h_F = o(n^{0.5-v_F})$ . Invoking this bound in (11) we obtain  $\mathcal{N}_{F,P}^* = o_P(\sqrt{m})$ . Next, from (10) we obtain that  $(\mathcal{N}_F - S_F) = o_P(\sqrt{m})$ . Then an application of (14) shows  $(\mathcal{N}_F - S_F) = o_P(\sqrt{m})$ . Finally, we apply the classical central limit theorem to the sum of iid random variables  $S_F$  to get the asymptotic normality of  $(\mathcal{N}_F - \mathbf{E}\mathcal{N}_F)/(\sigma_F \sqrt{m})$ .  $\square$

*Proof of Theorem 2 and Remark 3.* By Lemma 1 (respectively, Lemma 4), we have  $h_F = o(n^{\frac{1}{\alpha}-v_F})$ . Proceeding as in the proof of Theorem 1 above, we obtain  $(\mathcal{N}_F - S_F) = o_P(m^{1/\alpha})$ . Furthermore, by Lemma 5, the iid random variables  $N_{F,1}, N_{F,2}, \dots$  obey the power law (29) and therefore  $(S_F - B_m)/m^{1/\alpha}$  converges in distribution to  $G_{\alpha,a}$  (see Theorem 2 in § 35 of [6]). Hence  $m^{-1/\alpha}(\mathcal{N}_F - B_m)$  converges in distribution to  $G_{\alpha,a}$ .  $\square$

*Proof of Remark 1.* We have  $\sigma_F^2 = \mathbf{Var}N_F = \mathbf{Var}N_F^* + \mathbf{E}(\Delta_F^*)^2$ , where  $\Delta_F^* := N_F - N_F^*$ . Therefore  $\sigma_F^2 < \infty \Rightarrow \mathbf{Var}N_F^* < \infty \Rightarrow \mathbf{E}(N_F^*)^2 < \infty$ . To prove that  $\mathbf{E}(N_F^*)^2 < \infty \Rightarrow \sigma_F^2 < \infty$ , it suffices to show that  $\mathbf{E}(\Delta_F^*)^2 < \infty$ . By Lemma 3.5 of [9], we have  $\mathbf{E}^*(\Delta_F^*)^2 \prec (N_F^*)^2/\Phi_F(X, Q)$ , where  $\Phi_F(X, Q) = \min_{H \subset F} X^{v_H} Q^{e_H}$ . Furthermore, from the inequality (27), see below, which holds for balanced  $F$ , we obtain

$$\mathbf{E}^*(\Delta_F^*)^2 \prec \frac{(N_F^*)^2}{\min\{(N_F^*)^{2/v_F}, N_F^*\}} = \max\{(N_F^*)^{2-2/v_F}, N_F^*\} \leq \max\{1, (N_F^*)^2\}.$$

Hence  $\mathbf{E}(N_F^*)^2 < \infty$  implies  $\mathbf{E}(\Delta_F^*)^2 = \mathbf{E}(\mathbf{E}^*(\Delta_F^*)^2) < \infty$ .  $\square$

### 2.3 Auxiliary lemmas

In Lemmas 1 and 4 we upper bound the moments  $h_F$  for 2-connected  $F$  and for  $F = \mathcal{K}_k$ , respectively. Clearly, the result of Lemma 1 applies to  $F = \mathcal{K}_k$  as well, but the bound of Lemma 4 is tighter for large  $k$ .

**Lemma 1.** *Let  $F$  be a 2-connected graph with  $v_F \geq 3$  vertices. Let  $n, m \rightarrow +\infty$ . Assume that  $m = O(n)$ .*

(i) *Assume that (1) holds. Then  $h_F = o(n^{0.5-v_F})$ .*

(ii) *Assume that  $0 < \alpha < 2$ , and that (5) holds. Then  $h_F = o(n^{\frac{1}{\alpha}-v_F})$ .*

In the proof we use the simple fact that for any  $s, t, \tau > 0$ , the moment condition  $\mathbf{E}(X^s Q^t) < \infty$  implies

$$\mathbf{E}((\min\{X, n\})^{s+\tau} Q^t) = o(n^\tau). \quad (15)$$

Denote  $\tilde{X} := \min\{X, n\}$ . To see why (15) holds, choose  $0 < \delta < \tau/(s+\tau)$  and split the expectation

$$\mathbf{E}(\tilde{X}^{s+\tau} Q^t) = \mathbf{E}(\tilde{X}^{s+\tau} Q^t \mathbb{I}_{\{X < n^\delta\}}) + \mathbf{E}(\tilde{X}^{s+\tau} Q^t \mathbb{I}_{\{X \geq n^\delta\}}) =: I_1 + I_2.$$

Inequalities  $\tilde{X} \leq n$  and  $\mathbf{E}(X^s Q^t) < \infty$  imply  $I_2 \leq n^\tau \mathbf{E}(X^s Q^t \mathbb{I}_{\{X \geq n^\delta\}}) = n^\tau \cdot o(1)$ . Inequality  $\tilde{X} \leq X$  implies  $I_1 \leq n^{\delta(s+\tau)} = o(n^\tau)$ .

*Proof of Lemma 1.* The proofs of statements (i) and (ii) are identical. Therefore we only prove statement (i).

We start with establishing an auxiliary inequality (16) below, which may be interesting in itself. Let  $r \geq 2$ . Given a partition  $\tilde{B} = (B_1, \dots, B_r)$  of the edge set  $\mathcal{E}_0$  of the graph  $F_0 = (\mathcal{V}_0, \mathcal{E}_0)$ , and given  $i \in [r]$ , let  $V_i$  be the set of vertices incident to the edges from  $B_i$ . Let  $\rho_i$  be the number of (connected) components of the graph  $Z_i = (V_i, B_i)$  and put  $v_i = |V_i|$ . We claim that

$$v_1 + \dots + v_r \geq v_F + \rho_1 + \dots + \rho_r. \quad (16)$$

To establish the claim we consider the list  $H_1, H_2, \dots, H_t$  of components of  $Z_1, \dots, Z_r$  arranged in an arbitrary order. Here  $t := \rho_1 + \dots + \rho_r$ . Therefore, each graph  $H_i$  is a component of some  $Z_j$  and their union  $H_1 \cup \dots \cup H_t = Z_1 \cup \dots \cup Z_r = F_0$ . Let us consider the sequence of graphs  $\bar{H}_j := H_1 \cup \dots \cup H_j$ , for  $j = 1, \dots, t-1$ . Let  $\bar{\rho}_j$  and  $\bar{v}_j$  denote the number of components and the number of vertices of  $\bar{H}_j$ . Let  $v'_j$  denote the number of vertices of  $H_j$ . We use the observation that

$$\bar{v}_j \leq \bar{v}_{j-1} + v'_j + \bar{\rho}_j - \bar{\rho}_{j-1} - 1 \quad \text{for } j = 2, \dots, t-1. \quad (17)$$

Indeed,  $\bar{\rho}_{j-1} = \bar{\rho}_j$  means that the vertex set of (the connected graph)  $H_j$  intersects with exactly one component of  $\bar{H}_{j-1}$ . Consequently,  $H_j$  and  $\bar{H}_{j-1}$  have at least one common vertex and therefore (17) holds. Similarly,  $\bar{\rho}_{j-1} - \bar{\rho}_j = y > 0$  means that the vertex set of  $H_j$  intersects with exactly  $y + 1$  different components of  $\bar{H}_{j-1}$ . Consequently,  $H_j$  and  $\bar{H}_{j-1}$  have at least  $y + 1$  common vertices and (17) holds again. The remaining case  $\bar{\rho}_{j-1} - \bar{\rho}_j = -1$  is realized by the configuration where the vertex sets of  $H_j$  and  $\bar{H}_{j-1}$  have no common elements. In this case (17) follows from the identity  $\bar{v}_j = \bar{v}_{j-1} + v'_j$ .

By summing up the inequalities (17), we obtain (using  $\bar{\rho}_1 = 1$ ) that

$$\bar{v}_{t-1} \leq v'_1 + \cdots + v'_{t-1} + \bar{\rho}_{t-1} - t + 1.$$

Note that given  $\bar{H}_{t-1}$  with  $\bar{\rho}_{t-1}$  components, the vertex set of  $H_t$  must intersect with each component in two or more points, in order to make the union  $\bar{H}_{t-1} \cup H_t = F_0$  2-connected. Consequently, we have

$$\bar{v}_t \leq \bar{v}_{t-1} + v'_t - 2\bar{\rho}_{t-1}.$$

Finally, we obtain

$$v_F = \bar{v}_t \leq v'_1 + \cdots + v'_t - \bar{\rho}_{t-1} - t + 1.$$

Now the claim follows from the identity  $v'_1 + \cdots + v'_t = v_1 + \cdots + v_r$  and the inequality  $\bar{\rho}_{t-1} \geq 1$ .

Let us prove statement (i). Given  $(\tilde{B}, \tilde{i})$ , we obtain from (12) and (16) (recall the notation  $b_j = |B_j|$ ) that

$$h(\tilde{B}, \tilde{i}) \leq \frac{1}{n^{v_1 + \cdots + v_r}} \prod_{j=1}^r \mathbf{E} \left( \tilde{X}^{v_j} Q^{b_j} \right) \leq \frac{1}{n^{v_F + \rho_1 + \cdots + \rho_r}} \prod_{j=1}^r \mathbf{E} \left( \tilde{X}^{v_j} Q^{b_j} \right).$$

Given  $\tilde{B} = (B_1, \dots, B_r)$ , we estimate the sum over all possible colorings (there are  $(m)_r$  of them)

$$\begin{aligned} \sum_{\tilde{i}} h(\tilde{B}, \tilde{i}) &\prec \frac{(m)_r}{n^{v_F + \rho_1 + \cdots + \rho_r}} \prod_{j=1}^r \mathbf{E} \left( \tilde{X}^{v_j} Q^{b_j} \right) \asymp n^{-v_F} \prod_{j=1}^r \frac{\mathbf{E} \left( \tilde{X}^{v_j} Q^{b_j} \right)}{n^{\rho_j - 1}} \\ &= n^{0.5 - v_F} \prod_{j=1}^r \frac{\mathbf{E} \left( \tilde{X}^{v_j} Q^{b_j} \right)}{n^{\rho_j - 1 + (b_j / (2e_F))}} = o(n^{0.5 - v_F}). \end{aligned}$$

In the second last identity we used  $b_1 + \cdots + b_r = e_F$ , while the last bound follows by the chain of inequalities

$$\begin{aligned} n^{1 - \rho_j} \mathbf{E} \left( \tilde{X}^{v_j} Q^{b_j} \right) &\leq \mathbf{E} \left( \tilde{X}^{v_j + 1 - \rho_j} Q^{b_j} \right) \leq \mathbf{E} \left( \tilde{X}^{v_j + 1 - \rho_j} Q^{v_j - \rho_j} \right) \\ &= o \left( n^{(v_j - \rho_j) / (2e_F)} \right) = o \left( n^{b_j / (2e_F)} \right). \end{aligned}$$

Here in the first step we used  $\tilde{X} \leq n$ ; in the second step we used  $Q \leq 1$  and  $b_j \geq v_j - \rho_j$  (the latter inequality is based on the observation that any graph with  $v_j$  vertices and  $\rho_j$  components has at least  $v_j - \rho_j$  edges); the third step follows by (15) from the moment condition (1) applied to  $s = v_j - \rho_j$ ; the last step follows from the inequality  $b_j \geq v_j - \rho_j$ .

Finally, we conclude that

$$h_F = \sum_{\tilde{B}} \sum_{\tilde{i}} h(\tilde{B}, \tilde{i}) = o(n^{0.5 - v_F}), \quad (18)$$

because the number of partitions  $\tilde{B}$  of the edge set of a given graph  $F$  is always finite.  $\square$

Before showing an upper bound for  $h_F$ ,  $F = \mathcal{K}_k$ , we introduce some notation. Given an integer  $b \geq 1$ , let  $b^*$  be the minimal number of vertices that a graph with  $b$  edges may have. Let  $H_b$  be such a graph. It has a simple structure described below. Let  $k_b \geq 2$  be the largest integer satisfying  $b \geq \binom{k_b}{2}$ . Then

$$b = \binom{k_b}{2} + \Delta_b$$

for some integer  $0 \leq \Delta_b \leq k_b - 1$ . For  $\Delta_b = 0$  we have  $b^* = k_b$  and  $H_b = \mathcal{K}_{k_b}$  (clique on  $b^* = k_b$  vertices). For  $\Delta_b > 0$ , graph  $H_b$  is a union of  $\mathcal{K}_{k_b}$  and a star  $\mathcal{K}_{1, \Delta_b}$ , such that all the vertices of the star except for the central vertex belong to the vertex set of  $\mathcal{K}_{k_b}$ . In this case  $b^* = k_b + 1$ . In other words, one obtains  $H_b$  from  $\mathcal{K}_{k_b+1}$  by deleting  $k_b - \Delta_b$  edges sharing a common endpoint. The next two lemmas establish useful properties of the function  $b \rightarrow b^*$ .

**Lemma 2.** *For integers  $s \geq t \geq 1$  we have*

$$s^* + t^* \geq (s + t - 1)^* + 2. \quad (19)$$

*Proof.* In the proof we consider graphs  $H_s$  and  $H_t$  that have disjoint vertex sets so that the union  $H_s \cup H_t$  has  $s^* + t^*$  vertices.

Note that for  $t = 1$  both sides of (19) are equal. In order to show (19) for  $s \geq t \geq 2$  we consider the chain of neighboring pairs

$$(s, t) \rightarrow (s + 1, t - 1) \rightarrow \cdots \rightarrow (s + t - 1, 1). \quad (20)$$

In a step  $(x, y) \rightarrow (x + 1, y - 1)$  we remove an edge from  $H_y$  and add it to  $H_x$ . A simple analysis of the step  $(H_x, H_y) \rightarrow (H_{x+1}, H_{y-1})$  shows that

$$(x + 1)^* + (y - 1)^* = x^* + y^* + 1 \quad \text{whenever} \quad \Delta_x = 0, \Delta_y \neq 1, \quad (21)$$

$$(x + 1)^* + (y - 1)^* = x^* + y^* - 1 \quad \text{whenever} \quad \Delta_x \neq 0, \Delta_y = 1, \quad (22)$$

$$(x + 1)^* + (y - 1)^* = x^* + y^* \quad \text{in the remaining cases.} \quad (23)$$

We call a step  $(x, y) \rightarrow (x + 1, y - 1)$  positive (respectively negative or neutral) if (22) (respectively (21) or (23)) holds. Therefore, as we move in (20) from left to right, every positive (negative) step decreases (increases) the total number of vertices in the union  $H_x \cup H_y$ .

Let us now traverse (20) from right to left. We observe that the first non-neutral step encountered is positive (if we encounter a non-neutral step at all). Furthermore, after a negative step the first non-neutral step encountered is positive. Note that it may happen that the last encountered non-neutral step is negative. Therefore, the total number of positive steps is at least as large as the number of negative ones. This proves (19).  $\square$

**Lemma 3.** *Let  $k \geq 3$  and  $r \geq 2$ . Let  $B_1 \cup \cdots \cup B_r$  be a partition of the edge set of the clique  $\mathcal{K}_k$ . Denote  $b_i = |B_i|$ ,  $1 \leq i \leq r$ , and  $\varkappa = \binom{k}{2}$ . We have*

$$b_1^* + \cdots + b_r^* \geq (\varkappa - (r - 1))^* + 2(r - 1) \geq k + r. \quad (24)$$

*Proof.* The first inequality of (24) follows from (19) and the identity  $b_1 + \cdots + b_r = \varkappa$ . The second inequality is simple. Indeed, for  $r \geq k$  the inequality follows from  $2(r - 1) \geq k + r - 2$  and  $(\varkappa - (r - 1))^* \geq 2$ . For  $r \leq k - 1$  we have  $\varkappa - (r - 1) \geq \binom{k-1}{2} + 1$  and therefore  $(\varkappa - (r - 1))^* \geq k$ .  $\square$

Now we are ready to bound  $h_F$  for  $F = \mathcal{K}_k$ .

**Lemma 4.** *Let  $k \geq 3$ ,  $0 < \alpha \leq 2$ , and  $A > 0$ . Let  $n, m \rightarrow +\infty$ . Assume that  $m \leq An$ . Let  $F = \mathcal{K}_k$ . Then (6) implies the bound  $h_F = o\left(n^{\frac{1}{\alpha} - k}\right)$ . Note that for  $\alpha = 2$  condition (6) is the same as (3).*

*Proof.* For  $F = \mathcal{K}_k$  we have  $e_F = \binom{k}{2}$ . We observe that (6) implies

$$\mathbf{E}\left(X^{b^*-b/(\alpha e_F)}Q^b\right) < \infty \quad \text{for each} \quad 1 \leq b < \binom{k}{2}. \quad (25)$$

Note that  $\hat{s} = \binom{s-1}{2} + 1$  is the smallest integer  $t$  such that  $t^* = s$ . In particular, for any  $b$  with  $b^* = s$  we have  $b \geq \hat{s}$ . Therefore, given  $2 \leq s \leq k$ , the moment condition  $\mathbf{E}\left(X^{s-\hat{s}/(\alpha e_F)}Q^{\hat{s}}\right) < \infty$  implies  $\mathbf{E}\left(X^{s-b/(\alpha e_F)}Q^b\right) < \infty$  for any  $b$  satisfying  $b^* = s$ . In this way (6) yields (25)

Let us bound  $h_{\mathcal{K}_k}$  from above. Given a partition  $\tilde{B} = (B_1, \dots, B_r)$  of the edge set  $\mathcal{E}_0$  of  $\mathcal{K}_k = ([k], \mathcal{E}_0)$ , let  $v_j$  be the number of vertices incident to the edges from  $B_j$  and let  $b_j = |B_j|$ . For any vector  $\tilde{i} = (i_1, \dots, i_r)$  of distinct colors we have

$$h(\tilde{B}, \tilde{i}) \leq \prod_{j=1}^r \frac{\mathbf{E}(\tilde{X}^{v_j} Q^{b_j})}{n^{v_j}} \leq \prod_{j=1}^r \frac{\mathbf{E}(\tilde{X}^{b_j^*} Q^{b_j})}{n^{b_j^*}} \leq \frac{1}{n^{k+r}} \prod_{j=1}^r \mathbf{E}(\tilde{X}^{b_j^*} Q^{b_j}).$$

Here the first inequality follows from  $(\tilde{X})_t/(n)_t \leq \tilde{X}^t/n^t$ , since  $\tilde{X} \leq n$ . The second inequality follows from the obvious inequality  $b_j^* \leq v_j$  and the fact that  $\tilde{X} \leq n$ . The last inequality follows from the inequality  $b_1^* + \dots + b_r^* \geq k+r$  of Lemma 3.

For each  $r$ -partition  $\tilde{B}$  as above we bound the sum over all possible colorings  $\tilde{i}$  (there are  $(m)_r$  of them)

$$\sum_{\tilde{i}} h(\tilde{B}, \tilde{i}) \leq \frac{(m)_r}{n^{k+r}} \prod_{j=1}^r \mathbf{E}(\tilde{X}^{b_j^*} Q^{b_j}) \leq \frac{A^r}{n^k} \prod_{j=1}^r \mathbf{E}(\tilde{X}^{b_j^*} Q^{b_j}) = o\left(n^{\frac{1}{\alpha}-k}\right). \quad (26)$$

In the very last step we used the bounds (with  $e_F = b_1 + \dots + b_r = \binom{k}{2}$ )

$$\mathbf{E}\left(\tilde{X}^{b_j^*} Q^{b_j}\right) = o\left(n^{\frac{b_j}{\alpha e_F}}\right)$$

that follow from the moment conditions  $\mathbf{E}\left(X^{b_j^* - \frac{b_j}{\alpha e_F}} Q^{b_j}\right) < \infty$ , see (25), via (15). Finally, proceeding as in (18) above, we obtain the desired bound  $h_F = o\left(n^{\frac{1}{\alpha}-k}\right)$  from (26).  $\square$

## 2.4 Power-law tails

Recall that given a graph  $F = (\mathcal{V}_F, \mathcal{E}_F)$ , we denote by  $v_F = |\mathcal{V}_F|$  the number of vertices and by  $e_F = |\mathcal{E}_F|$  the number of edges. Let  $\Psi_F = \Psi_F(n, p) = n^{v_F} p^{e_F}$ , and define

$$\Phi_F = \Phi_F(n, p) = \min_{H \subset F, e_H \geq 1} \Psi_H, \quad m_F = \max_{H \subset F, e_H \geq 1} (e_H/v_H).$$

Here the minimum/maximum is taken over all subgraphs  $H \subset F$  with  $e_H \geq 1$ . Recall that  $F$  is called balanced if  $m_F = e_F/v_F$ . For a balanced  $F$  we have for any  $H \subset F$  with  $e_H \geq 1$  that

$$\Psi_H = (np^{e_H/v_H})^{v_H} \geq (np^{e_F/v_F})^{v_H} = \Psi_F^{v_H/v_F}.$$

Hence

$$\Phi_F \geq \min\{\Psi_F^{2/v_F}, \Psi_F\}. \quad (27)$$

**Lemma 5.** *Let  $k \geq 3$  be an integer. Let  $a > 0$  and  $0 < \alpha < 2$ . Assume that  $F$  is balanced and connected. Assume that*

$$\mathbf{P}\{N_F^* > t\} = (a + o(1))t^{-\alpha} \quad \text{as} \quad t \rightarrow +\infty. \quad (28)$$

Then

$$\mathbf{P}\{N_F > t\} = (a + o(1))t^{-\alpha} \quad \text{as} \quad t \rightarrow +\infty. \quad (29)$$

We remark that for  $0 < \alpha < 2$ , the tail asymptotics (29) implies that  $N_F$  belongs to the domain of attraction of an  $\alpha$ -stable distribution. Indeed, the left tail of  $N_F$  vanishes since  $\mathbf{P}\{N_F \geq 0\} = 1$ . Therefore, the conditions of Theorem 2 in § 35 Chapter 7 of [6] are satisfied.

*Proof.* We denote the conditional expectation and probability given  $(X, Q)$  by  $\mathbf{E}^*$  and  $\mathbf{P}^*$ . Furthermore, we denote  $k = v_F$  and  $\Delta_F^* = N_F - N_F^*$ . In the proof we often use the fact (see Lemma 3.5 of [9]) that

$$\mathbf{E}^*(\Delta_F^*)^2 \asymp \frac{(N_F^*)^2}{\Phi_F(X, Q)}(1 - Q). \quad (30)$$

We also use the simple relation  $N_F^* \asymp a_F \Psi_F(X, Q)$ .

To prove (29) we show that the contribution of  $\Delta_F^*$  to the sum  $N_F = N_F^* + \Delta_F^*$  is negligible compared to  $N_F^*$  and, therefore, the tail asymptotic (29) is determined by (28). For this purpose we apply exponential large deviation bounds for subgraph counts in Bernoulli random graphs [9, 10].

Given large  $t > 0$  and small  $\varepsilon > 0$ , introduce event  $\mathcal{H} = \{-\varepsilon N_F^* \leq \Delta_F^* \leq \varepsilon t\}$  and split

$$\begin{aligned} \mathbf{P}\{N_F > t\} &= \mathbf{P}\{N_F > t, \mathcal{H}\} + \mathbf{P}\{N_F > t, \Delta_F^* < -\varepsilon N_F^*\} + \mathbf{P}\{N_F > t, \Delta_F^* > \varepsilon t\} \\ &=: P_1 + P_2 + P_3. \end{aligned} \quad (31)$$

We first consider  $P_1$ . Replacing  $\Delta_F^*$  by its extreme values (on  $\mathcal{H}$ ) yields the inequalities

$$\mathbf{P}\{(1 - \varepsilon)N_F^* > t, \mathcal{H}\} \leq P_1 \leq \mathbf{P}\{N_F^* > t(1 - \varepsilon), \mathcal{H}\}. \quad (32)$$

We note that the right side of (32) is at most  $\mathbf{P}\{N_F^* > t(1 - \varepsilon)\}$  and the left side is at least

$$\mathbf{P}\{(1 - \varepsilon)N_F^* > t\} - P_2' - P_3',$$

where

$$P_2' := \mathbf{P}\{(1 - \varepsilon)N_F^* > t, \Delta_F^* < -\varepsilon N_F^*\}, \quad P_3' := \mathbf{P}\{(1 - \varepsilon)N_F^* > t, \Delta_F^* > \varepsilon t\}.$$

Hence, we have

$$\mathbf{P}\{(1 - \varepsilon)N_F^* > t\} - P_2' - P_3' \leq P_1 \leq \mathbf{P}\{N_F^* > t(1 - \varepsilon)\}. \quad (33)$$

Invoking the simple inequalities  $P_2 \leq P_2'$  and  $P_3 \leq P_3'$ , we obtain from (31), (33) that

$$\mathbf{P}\{(1 - \varepsilon)N_F^* > t\} - P_2' \leq \mathbf{P}\{N_F > t\} \leq \mathbf{P}\{N_F^* > t(1 - \varepsilon)\} + P_2' + P_3'. \quad (34)$$

We show below that for any  $0 < \varepsilon < 1$

$$P_2' = o(t^{-\alpha}) \quad \text{and} \quad P_3 = o(t^{-\alpha}) \quad \text{as} \quad t \rightarrow +\infty. \quad (35)$$

Note that (28) and (34) together with (35) imply (29). It remains to show (35).

*Proof of  $P_2' = o(t^{-\alpha})$ .* Given  $(X, Q)$  with  $0 < Q < 1$  (cases 0 and 1 are trivial), we apply Janson's inequality (Theorem 2.14 of [9]) to  $p_\varepsilon^* := \mathbf{P}^*\{\Delta_F^* < -\varepsilon N_F^*\}$ . In what follows, we assume that the random graph  $G(X, Q)$  and complete graph  $\mathcal{K}_X$  are both defined on the same vertex set of size  $X$  and that  $X \geq 1$ . Let

$$\bar{\delta} := \mathbf{E}^*(N_F^2) - \delta, \quad \delta := \sum_{F' \subset \mathcal{K}_X} \sum_{\substack{F'' \subset \mathcal{K}_X \\ \mathcal{E}_{F'} \cap \mathcal{E}_{F''} = \emptyset}} \mathbf{E}^*(\mathbb{I}_{F'} \mathbb{I}_{F''}).$$

Here the sum runs over ordered pairs  $(F', F'')$  of subgraphs of  $\mathcal{K}_X$  such that  $F'$  and  $F''$  are copies of  $F$  and their edge sets  $\mathcal{E}_{F'}$  and  $\mathcal{E}_{F''}$  are disjoint. Furthermore,  $\mathbb{I}_{F'}$  stands for the indicator of the event that  $F'$  is present in  $G(X, Q)$ . Janson's inequality implies

$$\mathbf{P}^*\{\Delta_F^* < -\eta N_F^*\} \leq e^{-(\eta N_F^*)^2 / \bar{\delta}}, \quad \forall \eta \in (0, 1). \quad (36)$$

Next we bound  $\bar{\delta}$  from above. The (variance) identity  $\mathbf{E}^*(N_F^2) - (N_F^*)^2 = \mathbf{E}^*(\Delta_F^*)^2$  implies

$$\bar{\delta} = \mathbf{E}^*(\Delta_F^*)^2 + (N_F^*)^2 - \delta. \quad (37)$$

Furthermore, using the observation that  $V_{F'} \cap V_{F''} = \emptyset$  implies  $\mathcal{E}_{F'} \cap \mathcal{E}_{F''} = \emptyset$ , and that the latter relation implies  $\mathbf{E}^*(\mathbb{I}_{F'} \mathbb{I}_{F''}) = (\mathbf{E}^* \mathbb{I}_{F'}) (\mathbf{E}^* \mathbb{I}_{F''}) = Q^{2e_F}$  we bound  $\delta$  from below:

$$\delta \geq \sum_{F' \subset \mathcal{K}_X} \sum_{\substack{F'' \subset \mathcal{K}_X \\ V_{F'} \cap V_{F''} = \emptyset}} \mathbf{E}^*(\mathbb{I}_{F'} \mathbb{I}_{F''}) = a_F^2 \binom{X}{k} \binom{X-k}{k} Q^{2e_F} = \frac{(X-k)_k}{(X)_k} (N_F^*)^2.$$

Then we lower bound the fraction

$$\frac{(X-k)_k}{(X)_k} \geq \left(1 - \frac{k}{X-k}\right)^k \geq 1 - \frac{k^2}{X-k}, \quad \text{for } X \geq 2k,$$

and obtain that  $\delta \geq (N_F^*)^2 (1 - k^2(X-k)^{-1})$ . Invoking this bound in (37) we obtain

$$\bar{\delta} \leq \mathbf{E}^*(\Delta_F^*)^2 + (N_F^*)^2 k^2 (X-k)^{-1}.$$

Hence the ratio in the exponent of (36) satisfies

$$\frac{(N_F^*)^2}{\bar{\delta}} \geq \frac{(N_F^*)^2}{2 \max\{\mathbf{E}^*(\Delta_F^*)^2, (N_F^*)^2 k^2 (X-k)^{-1}\}} = \frac{1}{2} \min \left\{ \frac{(N_F^*)^2}{\mathbf{E}^*(\Delta_F^*)^2}, \frac{X-k}{k^2} \right\}. \quad (38)$$

We will show below that there exists  $c_k > 0$  (independent of  $t$ ) such that  $N_F^* > t$  implies

$$\frac{(N_F^*)^2}{\mathbf{E}^*(\Delta_F^*)^2} > c_k t^{2/k}. \quad (39)$$

We also note that  $N_F^* > t$  implies  $X > (t/a_F)^{1/k}$  (we use  $a_F \binom{X}{k} \geq a_F \binom{X}{k} Q^{e_F} = N_F^*$ ). Therefore, on the event  $N_F^* > t$  the right side of (38) is at least

$$\frac{1}{2} \min \left\{ c_k t^{2/k}, \frac{(t/a_F)^{1/k} - k}{k^2} \right\} \quad (40)$$

and this quantity scales as  $t^{1/k}$  as  $t \rightarrow +\infty$ . Finally, from (36), (38), (40) we obtain that on the event  $N_F^* > t$ ,

$$p_\varepsilon^* \leq e^{-\varepsilon^2 \Theta(t^{1/k})} = o(t^{-\alpha}) \quad \text{as } t \rightarrow +\infty.$$

We conclude that  $P'_2 = o(t^{-\alpha})$ . It remains to show (39). We observe that inequalities  $N_F^* \leq a_F \Psi_F(X, Q)$  and  $N_F^* > t$  imply  $\Psi_F(X, Q) > t/a_F > 1$ , where the last inequality holds for  $t > a_F$ . Then (27) implies  $\Phi_F(X, Q) \geq (\Psi_F(X, Q))^{2/k}$  and (30) implies

$$\frac{(N_F^*)^2}{\mathbf{E}^*(\Delta_F^*)^2} \asymp \frac{\Phi_F(X, Q)}{1-Q} \geq \Phi_F(X, Q) \geq \Psi_F^{2/k}(X, Q) \geq (t/a_F)^{2/k}.$$

*Proof of  $P_3 = o(t^{-\alpha})$ .* In the proof we apply exponential inequalities for upper tails of subgraph counts in Bernoulli random graphs [10]. For reader's convenience, we state the result of [10] we will use. Let  $\Delta_F$  be the maximum degree of  $F$ . Let

$$M_F(n, p) = \begin{cases} 1 & \text{if } p < n^{-1/m_F}, \\ \min_{H \subset F} (\Psi_H(n, p))^{1/\alpha_H^*} & \text{if } n^{-1/m_F} \leq p \leq n^{-1/\Delta_F}, \\ n^2 p^{\Delta_F} & \text{if } p \geq n^{-1/\Delta_F}. \end{cases}$$

Here  $\alpha_H^*$  is the fractional independence number of a graph  $H$ , see [10]. We do not define the fractional independence number here as we only use the upper bound  $\alpha_H^* \leq v_H - 1$  that

holds for any  $H$  with  $e_H > 0$ , see formula (A.1) in [10]. Let  $\xi_F$  be the number of copies of  $F$  in  $G(n, p)$ . By Theorems 1.2 and 1.5 of [10], for any  $\eta > 0$  there exists  $c_{\eta, F} > 0$  such that uniformly in  $p$  and  $n \geq k$  (recall that  $k = v_F$  is the number of vertices of  $F$ ) we have

$$\mathbf{P}\{\xi_F \geq (1 + \eta)\mathbf{E}\xi_F\} \leq e^{-c_{\eta, F} M_F(n, p)}. \quad (41)$$

We will apply (41) to the number  $N_F$  of copies of  $F$  in  $G(X, Q)$  conditionally given  $X, Q$ , see (46) below.

We write, for short,  $s = \varepsilon t$  and estimate  $P_3 \leq \mathbf{P}\{\Delta_F^* > s\}$ . Let  $\eta > 0$ . We split

$$\mathbf{P}\{\Delta_F^* > s\} = \mathbf{P}\{\Delta_F^* > \eta N_F^*, \Delta_F^* > s\} + \mathbf{P}\{\Delta_F^* \leq \eta N_F^*, \Delta_F^* > s\} =: P_{31} + P_{32}$$

and estimate the probabilities  $P_{31}$  and  $P_{32}$  separately. The second probability

$$P_{32} \leq \mathbf{P}\{N_F^* > s/\eta\} = \eta^\alpha (a + o(1))s^{-\alpha} \quad (42)$$

can be made negligibly small by choosing  $\eta$  arbitrarily small.

Now we upper bound the remaining probability  $P_{31}$ . Introduce the events

$$\mathcal{A}_1 = \{Q \leq X^{-1/m_F}\}, \quad \mathcal{A}_{21} = \{X^{-1/m_F} < Q < X^{-1/\Delta_F}\}, \quad \mathcal{A}_{22} = \{Q \geq X^{-1/\Delta_F}\},$$

and put  $\mathcal{A}_2 = \mathcal{A}_{21} \cup \mathcal{A}_{22}$  (note that  $\Delta_F \geq 2m_F = 2e_F/v_F$ ). We split

$$P_{31} = \tilde{P}_1 + \tilde{P}_2, \quad \tilde{P}_i := \mathbf{P}\{\Delta_F^* > \eta N_F^*, \Delta_F^* > s, \mathcal{A}_i\} \quad (43)$$

and estimate  $\tilde{P}_1$  and  $\tilde{P}_2$  separately. We firstly consider  $\tilde{P}_1$ . The inequality  $Q \leq X^{-1/m_F}$  implies  $\Psi_F(X, Q) \leq 1$ . Consequently, (27) implies  $\Phi_F(X, Q) \geq \Psi_F(X, Q)$ . The latter inequality together with (30) imply  $\mathbf{E}^*(\Delta_F^*)^2 \leq c_k \Psi_F(X, Q) \leq c_k$  for some  $c_k > 0$ . Hence, on the event  $\mathcal{A}_1$  we have  $\mathbf{E}^*(\Delta_F^*)^2 \leq c_k$ . Finally, by Markov's inequality,

$$\tilde{P}_1 \leq \mathbf{P}\{\Delta_F^* > s, \mathcal{A}_1\} = \mathbf{E}(\mathbb{I}_{\mathcal{A}_1} \mathbf{E}^* \mathbb{I}_{\{\Delta_F^* > s\}}) \leq \mathbf{E}(\mathbb{I}_{\mathcal{A}_1} \mathbf{E}^*(\Delta_F^*)^2 s^{-2}) \leq c_k s^{-2}. \quad (44)$$

We secondly consider  $\tilde{P}_2$ . The inequality  $X^{-1/m_F} < Q$  implies  $\Psi_F(X, Q) > 1$ . For balanced  $F$  this yields  $\Psi_H(X, Q) > 1$  for every  $H \subset F$  with  $e_H > 0$ . Then, by using  $\alpha_H^* \leq v_H - 1$  we obtain

$$\min_{H \subset F: e_H > 0} (\Psi_H(X, Q))^{1/\alpha_H^*} \geq \min_{H \subset F: e_H > 0} (\Psi_H(X, Q))^{1/v_H} = (\Psi_F(X, Q))^{1/v_F}.$$

In the last step we used the fact that  $F$  is balanced once again. Hence, on the event  $\mathcal{A}_{21}$  we have (recall that  $v_F = k$ )

$$M_F(X, Q) \geq (\Psi_F(X, Q))^{1/k}. \quad (45)$$

We observe that (45) holds on the event  $\mathcal{A}_{22}$  as well. Indeed, the inequality  $Q \geq X^{-1/\Delta_F}$  yields  $M_F(X, Q) \geq X^2 Q^{\Delta_F} \geq X$ . Now the inequality  $X^{v_F} \geq \Psi_F(X, Q)$  implies (45).

From (41) and (45) we obtain the exponential bound

$$\mathbf{P}^*\{\Delta_F^* > \eta N_F^*\} \leq e^{-c_{\eta, F} M_F(X, Q)} \leq e^{-c_{\eta, F} (\Psi_F(X, Q))^{1/k}}. \quad (46)$$

Let us bound  $\tilde{P}_2$  from above. We fix a (large) number  $B > 0$  and introduce the events

$$\mathcal{B}_1 = \{\Psi_F(X_1, Q_1) > B \ln^k s\}, \quad \mathcal{B}_2 = \{\Psi_F(X_1, Q_1) \leq B \ln^k s\}.$$

We then split

$$\tilde{P}_2 = \tilde{P}_{21} + \tilde{P}_{22}, \quad \tilde{P}_{2i} = \mathbf{P}\{\Delta_F^* > \eta N_F^*, \Delta_F^* > s, \mathcal{A}_2, \mathcal{B}_i\},$$

and bound  $\tilde{P}_{21}$  from above, by using (46),

$$\begin{aligned} \tilde{P}_{21} &\leq \mathbf{P}\{\Delta_F^* > \eta N_F^*, \mathcal{A}_2, \mathcal{B}_1\} = \mathbf{E}(\mathbb{I}_{\mathcal{B}_1} \mathbb{I}_{\mathcal{A}_2} \mathbf{P}^*\{\Delta_F^* > \eta N_F^*\}) \\ &\leq \mathbf{E}\left(\mathbb{I}_{\mathcal{B}_1} e^{-c_{\eta, F} (\Psi_H(X_1, Q_1))^{1/k}}\right) \leq e^{-c_{\eta, F} B^{1/k} \ln s}. \end{aligned} \quad (47)$$

It remains to upper bound  $\tilde{P}_{22}$ . The inequality  $\Psi_F(X, Q) > 1$ , which holds on the event  $\mathcal{A}_2$ , implies (see (27))  $\Phi_F(X, Q) \geq (\Psi_F(X, Q))^{2/k}$ . Furthermore, (30) implies

$$\mathbf{E}^*(\Delta_F^*)^2 \leq c_F (\Psi_F(X, Q))^{2-(2/k)} (1 - Q),$$

where  $c_F > 0$  depends only on  $F$ . Note that on the event  $\mathcal{B}_2$  the right side is upper bounded by  $c_F (B \ln^k s)^{2-(2/k)}$ . Hence, by Markov's inequality,

$$\mathbf{P}^*(\Delta_F^* > s) \leq s^{-2} \mathbf{E}^*(\Delta_F^*)^2 \leq c_F B^{2-(2/k)} s^{-2} \ln^{k-2} s.$$

Finally, we obtain

$$\tilde{P}_{22} \leq \mathbf{P}\{\Delta_F^* > s, \mathcal{A}_2, \mathcal{B}_2\} = \mathbf{E}(\mathbb{1}_{\mathcal{A}_2} \mathbb{1}_{\mathcal{B}_2} \mathbf{P}^*\{\Delta_F^* > s\}) \leq c_F B^{2-(2/k)} s^{-2} \ln^{2k-2} s. \quad (48)$$

We complete the proof by showing that for any  $0 < \varepsilon < 1$  the probability  $P_3$ , which depends on  $\varepsilon$ , satisfies  $P_3 = o(t^{-\alpha})$  as  $t \rightarrow +\infty$ . Recall that  $s = \varepsilon t$ . We have for any  $\eta > 0$

$$\begin{aligned} \limsup_{t \rightarrow +\infty} t^\alpha P_3 &\leq \limsup_{t \rightarrow +\infty} t^\alpha \mathbf{P}\{\Delta_F^* > \varepsilon t\} = \varepsilon^{-\alpha} \limsup_{s \rightarrow +\infty} s^\alpha \mathbf{P}\{\Delta_F^* > s\} \\ &\leq \varepsilon^{-\alpha} \limsup_{s \rightarrow +\infty} s^\alpha (\tilde{P}_1 + \tilde{P}_{21} + \tilde{P}_{22} + P_{32}) \leq (\eta/\varepsilon)^\alpha a. \end{aligned} \quad (49)$$

Hence  $\limsup_{t \rightarrow +\infty} t^\alpha P_3 = 0$ . The last inequality of (49) follows from (42), (44), (47), and (48). Indeed, given  $\eta > 0$ , we choose  $B = B(\eta)$  (in (47), (48)) large enough so that  $c_{\eta, F} B^{1/k} > 2$ . Then  $\tilde{P}_{21} \leq s^{-2}$  and  $\limsup_s s^\alpha \tilde{P}_{21} = 0$ . We also mention the obvious relations  $\limsup_s s^\alpha \tilde{P}_1 = 0$  and  $\limsup_s s^\alpha \tilde{P}_{22} = 0$ .  $\square$

## References

- [1] Bloznelis, M. and Kurauskas, V. (2016): Clustering coefficient of random intersection graphs with infinite degree variance. *Internet Mathematics* doi:10.24166/im.02.2017.
- [2] Bloznelis, M. and Jaworski, J. (2018): The asymptotic normality of the global clustering coefficient in sparse random intersection graphs. *Algorithms and models for the web graph*, 16–29, Lecture Notes in Comput. Sci., 10836, Springer.
- [3] Bloznelis, M. and Leskelä, L. (2019): Clustering and percolation on superpositions of Bernoulli random graphs, [arXiv:1912.13404](https://arxiv.org/abs/1912.13404).
- [4] Bloznelis, M., Karjalainen, J., Leskelä, L. (2021+): Assortativity and bidegree distributions on Bernoulli random graph superpositions. *Probability in the Engineering and Informational Sciences*, doi:10.1017/S0269964821000310.
- [5] Godehardt, E., Jaworski, J. (2001): Two models of random intersection graphs and their applications. *Electronic Notes in Discrete Mathematics* 10, 129–132.
- [6] Gnedenko, B. V. and Kolmogorov, A. N. (1954): *Limit distributions for sums of independent random variables*. Addison-Wesley, Cambridge.
- [7] Gröhn, T., Karjalainen, J., Leskelä, L.: Clique and cycle frequencies in a sparse random graph model with overlapping communities, [arXiv:1911.12827](https://arxiv.org/abs/1911.12827).
- [8] Hladký, J., Pelekis, Ch., Šileikis, M. (2021): A limit theorem for small cliques in inhomogeneous random graphs. *J. Graph Theory* 97, 578–599.
- [9] Janson, S., Łuczak, T., Ruciński, A. (2000): *Random Graphs*, Wiley, New York.
- [10] Janson, S., Oleszkiewicz, K., Ruciński, A. (2004): Upper tails for subgraph counts in random graphs. *Isr. J. Math.* 142, 61–92.
- [11] Karjalainen, J., van Leeuwen, J. S. H., Leskelä, L. (2018): Parameter estimators of random intersection graphs with thinned communities. In A. Bonato, P. Prałat, A. Raigorodskii (Eds.): *Algorithms and models for the web graph - 15th International workshop, WAW 2018, Lecture Notes in Comput. Sci. 10836*, Springer (2018), 44–58.

- [12] Privault, N. and Serafin, G. (2020): Normal approximation for sums of weighted U-statistics: Application to Kolmogorov bounds in random subgraph counting. *Bernoulli* 26, 587–615.
- [13] Rollin, A. (2021+): Kolmogorov bounds for the normal approximation of the number of triangles in the Erdős-Rényi random graph, *Probability in the Engineering and Informational Sciences*, doi:10.1017/S0269964821000061.
- [14] Ruciński, A. (1988): When are small subgraphs of a random graph normally distributed? *Probability Theory and Related Fields* 78, 1–10.
- [15] Yang, J. and Leskovec, J. (2012): Community-affiliation graph model for overlapping network community detection. In *2012 IEEE 12th International Conference on Data Mining*, pages 1170–1175. IEEE.
- [16] Yang, J. and Leskovec, J. (2014): Structure and overlaps of ground-truth communities in networks. *ACM Trans. Intell. Syst. Technol.* 5(2), 1–35 .