

## CLOSED FORMULAE FOR CERTAIN FERMAT-PELL EQUATIONS

FERNANDO SZECHTMAN

ABSTRACT. Given positive integers  $j, k$ , with  $j \geq 2$ , we show that there are positive integers  $d, e$  such that  $\sqrt{d}$  has continued fraction expansion  $\sqrt{d} = [e, \overline{k, \dots, k, 2e}]$ , with period  $j$ , if and only if  $k$  is even or  $3 \nmid j$ , in which case we give closed formulae to find all such  $d, e$  as well as the smallest solution in positive integers to the Fermat-Pell equation  $X^2 - dY^2 = (-1)^j$ .

## 1. INTRODUCTION

Given a positive integer  $d$  that is not a perfect square, its square root  $\sqrt{d}$  has an infinite periodic continued fraction expansion

$$\sqrt{d} = [e, \overline{a_1, \dots, a_{j-1}, 2e}],$$

where  $e = [\sqrt{d}]$  is the integral part of  $\sqrt{d}$ ,  $j \geq 1$  is the period of this expansion, and

$$a_1 = a_{j-1}, a_2 = a_{j-2}, \dots$$

are positive integers [S]. Set

$$(1) \quad \begin{pmatrix} y & x \\ z & w \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 2e \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & a_{j-1} \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ 1 & a_1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & e \end{pmatrix}.$$

Then  $(x, y) \in \mathbb{N} \times \mathbb{N}$  is a solution to the Fermat-Pell equation

$$(2) \quad X^2 - dY^2 = (-1)^j.$$

Moreover, if  $j$  is even, then

$$(3) \quad X^2 - dY^2 = -1$$

has no integral solutions, and  $(x_1, y_1) \in \mathbb{Z} \times \mathbb{Z}$  is a solution to

$$(4) \quad X^2 - dY^2 = 1$$

if and only if

$$(5) \quad x_1 + y_1\sqrt{d} = \pm(x + y\sqrt{d})^k, \quad k \in \mathbb{Z}.$$

Furthermore, if  $j$  is odd, then  $(x_1, y_1) \in \mathbb{Z} \times \mathbb{Z}$  is a solution to (3) (resp. (4)) if and only if (5) holds with  $k$  odd (resp. even) [S]. Thus

$$(6) \quad x + y\sqrt{d}$$

is the smallest real number such that  $x, y$  are both positive integers and  $(x, y)$  is a solution to (2). For this reason, we will refer to  $(x, y)$  as the *smallest solution* to (2).

For certain values of  $d$ , a closed formula for (6) is available. For instance, if  $d = e^2 + 1$ , then  $(x, y) = (e, 1)$  is the smallest solution to (3),  $\sqrt{d} = [e, \overline{2e}]$ , and  $j = 1$ . Also, if  $d = e^2 + 2e$ , then

---

2010 *Mathematics Subject Classification.* 11Y65, 11A55, 11D09, 11Y50, 11B37.

*Key words and phrases.* Fermat-Pell equation, continued fraction.

This research was partially supported by an NSERC grant.

$(x, y) = (e + 1, 1)$  is the smallest solution to (4),  $\sqrt{d} = [e, \overline{1, 2e}]$ , and  $j = 2$ . More generally, if  $d = e^2 + \frac{2e}{k}$ , where  $k \in \mathbb{N}$  and  $k|2e$ , then

$$(7) \quad (x, y) = (ke + 1, k)$$

is the smallest solution to (4),  $\sqrt{d} = [e, \overline{k, 2e}]$ , and  $j = 2$ , provided  $k \neq 2e$ . The following table illustrates a few instances, with  $j = 2$ , of the latter phenomenon.

$e$	$k$	$d$	$x$	$y$
1	1	3	2	1
2	1	8	3	1
2	2	6	5	2
3	1	15	4	1
3	2	12	7	2
3	3	11	10	3
4	1	24	5	1
4	2	20	9	2
4	4	18	17	4

We seek a closed formula, such as (7), rather than algorithm, such as (1), for the smallest solution to (2), available for a suitable family of values for  $d$ . The preceding examples suggest that we consider the case

$$(8) \quad \sqrt{d} = [e, \overline{k, \dots, k, 2e}], \text{ with period } j \geq 2.$$

Note that  $j \geq 2$  implies  $k \neq 2e$ . However, given  $j, k \in \mathbb{N}$ , with  $j \geq 3$ , there may not exist any  $d, e \in \mathbb{N}$  such that (8) holds (unlike the case  $j = 2$ ). Indeed, let  $d \in \mathbb{N}$ , not a perfect square, with

$$\sqrt{d} = [a_0, a_1, a_2, \dots], \text{ with period } j \geq 1,$$

and set

$$(9) \quad \begin{pmatrix} q_{-2} & p_{-2} \\ q_{-1} & p_{-1} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

as well as

$$(10) \quad \begin{pmatrix} q_{n-1} & p_{n-1} \\ q_n & p_n \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & a_n \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ 1 & a_0 \end{pmatrix}, \quad n \geq 0.$$

Then

$$q_n = a_n q_{n-1} + q_{n-2}, \quad p_n = a_n p_{n-1} + p_{n-2}, \quad n \geq 0,$$

and, according to (1),

$$(x, y) = (q_{j-1}, p_{j-1})$$

is the smallest solution to (2). Suppose, if possible, that

$$\sqrt{d} = [e, \overline{1, 1, 2e}]$$

for some  $d, e \in \mathbb{N}$ . By above,

$n$	$q_n$	$p_n$
-2	1	0
-1	0	1
0	1	$e$
1	1	$e + 1$
2	2	$2e + 1$

and  $(x, y) = (2e + 1, 2)$  is a solution to (2). Thus  $(2e + 1)^2 - 4d = -1$ , which implies

$$d = \frac{4e^2 + 4e + 2}{4},$$

against the fact that  $d \in \mathbb{N}$ . The above discussion leads to the following

**Goal.** Given  $j, k \in \mathbb{N}$ , with  $j \geq 2$ , we wish to find necessary and sufficient conditions for the existence of  $d, e \in \mathbb{N}$  such that (8) holds. Moreover, when these conditions are satisfied, we aim to find all possible values for  $d, e$ , as well as a closed formula for the smallest solution  $(x, y)$  to the Fermat-Pell equation (2).

For instance, when  $k \in \mathbb{N}$ , we look for all  $e, d \in \mathbb{N}$  so that

$$(11) \quad \sqrt{d} = [e, \overline{k, k, 2e}], \text{ with period } j = 3,$$

as well as the smallest solution to (2). Assuming (11) holds, the algorithm (1) produces

$n$	$q_n$	$p_n$
-2	1	0
-1	0	1
0	1	$e$
1	$k$	$ke + 1$
2	$k^2 + 1$	$(k^2 + 1)2e + k$

where  $q_n, p_n$  are defined in (9) and (10), so

$$(12) \quad (x, y) = ((k^2 + 1)e + k, k^2 + 1)$$

is the smallest solution to (2). Thus

$$((k^2 + 1)e + k)^2 - d(k^2 + 1)^2 = -1,$$

which implies

$$(13) \quad d = \frac{((k^2 + 1)e + k)^2 + 1}{(k^2 + 1)^2} = e^2 + \frac{2(k^2 + 1)ke + k^2 + 1}{(k^2 + 1)^2} = e^2 + \frac{2ke + 1}{k^2 + 1}.$$

Now

$$2ke \equiv -1 \pmod{k^2 + 1}$$

is solvable if and only if  $k$  is even, in which case there is a unique solution  $e \equiv \frac{k}{2} \pmod{k^2 + 1}$ , whence

$$(14) \quad e = \frac{k}{2} + \ell(k^2 + 1), \quad \ell \geq 1 \quad (\text{to ensure period } j = 3).$$

Suppose, conversely, that  $x, y, d, e$  are as indicated in (12)-(14). Then, by construction, we have  $x/y = [e, k, k]$ . Since  $x^2 - dy^2 = -1$ , it follows from [S, Exercises 7.7.15, 7.7.17] that  $\sqrt{d} = [e, \overline{k, k, 2e}]$ , with period  $j = 3$  as  $2e \neq k$ . (There is a typing error at the beginning of [S, Exercise 7.7.17]: It should say  $X^2 - dY^2 = \pm 1$  instead of  $X^2 - dY^2 = 1$ .)

The following table lists a few examples.

$e$	$k$	$d$	$x$	$y$
6	2	41	32	5
11	2	130	57	5
16	2	269	82	5
19	4	370	327	17
36	4	1313	616	17
40	6	1613	1486	37
69	8	4778	4493	65

## 2. THE GENERAL CASE

Given  $k \in \mathbb{N}$  and  $j = m + 1 \geq 2$ , suppose that (8) holds, and let  $(x, y)$  be the smallest solution to (2). Set

$$(15) \quad f_{-2} = 1, f_{-1} = 0, f_n = kf_{n-1} + f_{n-2}, \quad n \geq 0.$$

Then the algorithm (1) yields

$$(16) \quad (x, y) = (f_m + ef_{m-1}, f_m).$$

We digress here to record a few properties of the sequence (15). We readily see by induction that

$$(17) \quad \gcd(f_n, f_{n-1}) = 1, \quad n \geq -1$$

as well as

$$\begin{pmatrix} 0 & 1 \\ 1 & k \end{pmatrix}^n = \begin{pmatrix} f_{n-2} & f_{n-1} \\ f_{n-1} & f_n \end{pmatrix}, \quad n \geq 1,$$

which implies

$$(18) \quad f_{n-1}^2 + (-1)^n = f_n f_{n-2}, \quad n \geq 1 \text{ (Cassini identity)}.$$

Arguing by induction, and making use of (15) and (18), we easily derive

$$(19) \quad (-1)^n f_{n-2} f_{n-1} \equiv k \pmod{f_n}, \quad n \geq 2.$$

The following table displays  $f_n$  for all  $-2 \leq n \leq 11$ .

$n$	$f_n$
-2	1
-1	0
0	1
1	$k$
2	$k^2 + 1$
3	$k^3 + 2k$
4	$k^4 + 3k^2 + 1$
5	$k^5 + 4k^3 + 3k$
6	$k^6 + 5k^4 + 6k^2 + 1$
7	$k^7 + 6k^5 + 10k^3 + 4k$
8	$k^8 + 7k^6 + 15k^4 + 10k^2 + 1$
9	$k^9 + 8k^7 + 21k^5 + 20k^3 + 5k$
10	$k^{10} + 9k^8 + 28k^6 + 35k^4 + 15k^2 + 1$
11	$k^{11} + 10k^9 + 36k^7 + 56k^5 + 35k^3 + 6k$

As the table suggests, it is easily shown by induction that

$$f_n = a_n k^n + a_{n-2} k^{n-2} + a_{n-4} k^{n-4} + \cdots, \quad n \geq 0,$$

where

$$a_{n-2i} = \binom{1}{0} + \binom{i}{1} + \binom{i+1}{2} + \cdots + \binom{n-i-1}{n-2i} = \binom{1}{0} + \sum_{1 \leq s \leq n-2i} \binom{s+(i-1)}{s}.$$

We now resume our prior discussion. Since (16) is a solution to (2), we have

$$d = \frac{(f_m e + f_{m-1})^2 + (-1)^m}{f_m^2} = e^2 + \frac{2f_m f_{m-1} e + f_{m-1}^2 + (-1)^m}{f_m^2}.$$

By virtue of (18), the above may be rewritten as follows:

$$(20) \quad d = e^2 + \frac{2f_{m-1}e + f_{m-2}}{f_m}.$$

The congruence equation

$$(21) \quad 2f_{m-1}e \equiv -f_{m-2} \pmod{f_m}$$

is solvable if and only if

$$\gcd(f_m, 2f_{m-1}) \mid f_{m-2}.$$

By (17), the above translates into

$$\gcd(f_m, 2) \mid f_{m-2},$$

which is true except only when  $f_m$  is even and  $f_{m-2}$  is odd. Suppose first  $k$  is even. Then  $f_m, f_{m-2}$  are both even if  $m$  is odd (no constant term) and both odd if  $m$  is even (constant term 1). Suppose next  $k$  is odd. Then (15) yields the following parity for  $f_n$ :

$$\begin{array}{ccccccc} f_0 & f_1 & f_2 & f_3 & f_4 & f_5 & \dots \\ \text{odd} & \text{odd} & \text{even} & \text{odd} & \text{odd} & \text{even} & \dots \end{array}$$

We deduce that

$$f_m \text{ is even and } f_{m-2} \text{ is odd} \Leftrightarrow k \text{ is odd and } m \equiv 2 \pmod{3}.$$

Suppose henceforth that  $k$  is even or  $m \not\equiv 2 \pmod{3}$ . Since  $\gcd(f_m, f_{m+1}) = 1$ , we may multiply (21) by  $f_{m+1}$  and obtain the equivalent equation

$$2f_{m-1}f_{m+1}e \equiv -f_{m-2}f_{m+1} \pmod{f_m}.$$

By (18), this translates into

$$2(f_m^2 + (-1)^{m+1})e \equiv -f_{m-2}f_{m+1} \pmod{f_m},$$

that is,

$$2e \equiv (-1)^m f_{m-2}f_{m+1} \equiv (-1)^m f_{m-2}(kf_m + f_{m-1}) \equiv (-1)^m f_{m-2}f_{m-1} \pmod{f_m}.$$

In light of (19), this is equivalent to

$$(22) \quad 2e \equiv k \pmod{f_m}.$$

Three cases arise.

CASE 1.  $k$  is even and  $f_m$  is odd. Then all solutions to (22) are of the form

$$(23) \quad e = \frac{k}{2} + \ell f_m, \quad \ell \geq 1 \quad (\text{to ensure period } j).$$

By (20) and (23), we see that

$$d = e^2 + \frac{(k + 2\ell f_m)f_{m-1} + f_{m-2}}{f_m} = e^2 + \frac{2\ell f_m f_{m-1} + kf_{m-1} + f_{m-2}}{f_m} = e^2 + \frac{2\ell f_m f_{m-1} + f_m}{f_m},$$

so

$$(24) \quad d = e^2 + 2\ell f_{m-1} + 1, \quad \ell \geq 1.$$

CASE 2.  $k$  is even and  $f_m$  is even. Then all solutions to (22) are of the form

$$(25) \quad e = \frac{k}{2} + \ell \frac{f_m}{2}, \quad \ell \geq 1,$$

in which case the above calculations yield

$$(26) \quad d = e^2 + \ell f_{m-1} + 1, \quad \ell \geq 1.$$

CASE 3.  $k$  is odd. In this case  $m \not\equiv 2 \pmod 3$ , so  $f_m$  is also odd. Then  $k + f_m$  is even, and all solutions to (22) are of the form

$$(27) \quad e = \frac{k + f_m}{2} + \ell f_m, \quad \ell \geq 0,$$

in which case the above calculations yield

$$(28) \quad d = e^2 + (2\ell + 1)f_{m-1} + 1, \quad \ell \geq 0.$$

Suppose, conversely, that  $x, y$  are as in (16) and that  $d, e$  are as indicated in (23)-(28). Then, by construction, we have  $x/y = [e, k, \dots, k]$ , where  $k$  is repeated  $m$  times. Since  $x^2 - dy^2 = -1$ , it follows from [S, Exercises 7.7.15, 7.7.17] that  $\sqrt{d} = [e, \overline{k, \dots, k, 2e}]$ , with period  $j$  as  $2e \neq k$ .

We have proven the following

**Theorem.** *Let  $j, k \in \mathbb{N}$ , with  $j = m + 1 \geq 2$ , and let  $f_{-2}, f_{-1}, f_0, \dots$  be defined as in (15). Then there exist  $d, e \in \mathbb{N}$  such that*

$$\sqrt{d} = [e, \overline{k, \dots, k, 2e}], \text{ with period } j \geq 2,$$

*if and only if  $k$  is even or  $m \not\equiv 2 \pmod 3$ , in which case all such  $d, e$  are given by (23) and (24) if  $k$  is even and  $f_m$  is odd, by (25) and (26) if  $k$  is even and  $f_m$  is even, and by (27) and (28) if  $k$  is odd; moreover, the smallest solution to (2) is as indicated in (16).*

The following table illustrates a few instances of the above result.

$m$	$k$	$f_{m-1}$	$f_m$	$\ell$	$e$	$d$	$f_m e + f_{m-1}$
3	1	2	3	0	2	7	8
3	1	2	3	1	5	32	17
3	1	2	3	2	8	75	26
3	1	2	3	3	11	136	35
3	2	5	12	4	25	646	305
3	3	10	33	0	18	335	604
3	4	17	72	1	38	1462	2753
3	5	26	135	0	70	4927	9476
4	1	3	5	0	3	13	18
4	1	3	5	1	8	74	43
4	2	12	29	1	30	925	882
4	3	33	109	0	56	3170	6137
5	2	29	70	2	71	5100	4999
5	2	29	70	4	141	19998	9899
5	2	29	70	6	211	44696	14799
5	4	305	1292	2	1294	1675047	1672153
6	1	8	13	0	7	58	99
6	1	8	13	1	20	425	268
6	2	70	169	1	170	29041	28800
7	1	13	21	0	11	135	244
7	2	169	408	2	409	167620	167041
9	1	34	55	0	28	819	1574
10	1	55	89	0	45	2081	4060
12	1	144	233	0	117	13834	27405
13	1	233	377	0	189	35955	71486

## REFERENCES

- [S] H. M. Stark, *An Introduction to Number Theory*, Markham, Chicago, 1970.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF REGINA, CANADA  
*Email address:* `fernando.szechtman@gmail.com`