

# Families of extensions of the Kantorovich-Rubinstein and Lipschitz norms

Dávid Terjék  
Alfréd Rényi Institute of Mathematics  
Budapest, Hungary  
dterjek@renyi.hu

October 25, 2021

## Abstract

We propose a family of extensions of the Kantorovich-Rubinstein norm from the space of zero-charge countably additive measures on a compact metric space to the space of all countably additive measures, and a family of extensions of the Lipschitz norm from the quotient space of Lipschitz functions on a compact metric space to the space of all Lipschitz functions. These families are parameterized by  $p, q \in [1, \infty]$ , and if  $p, q$  are Hölder conjugates, then the dual of the resulting  $p$ -Kantorovich space is isometrically isomorphic to the resulting  $q$ -Lipschitz space.

## 1 Introduction

Given a compact metric space  $(X, d)$ , the vector space  $\mathcal{M}(X, 0)$  of countably additive measures  $\mu$  on the Borel  $\sigma$ -algebra of  $X$  such that  $\mu(X) = 0$  can be normed by the Kantorovich-Rubinstein norm

$$\|\mu\|_{KR} = \inf_{\pi \in \mathcal{M}(X \times X), \pi(X \times \cdot) - \pi(\cdot \times X) = \mu} \left\{ \int d(x, y) d\pi(x, y) \right\},$$

the theory of which was developed in [Kantorovich and Rubinstein, 1957] and [Kantorovich and Rubinstein, 1958]. See [Kantorovich and Akilov, 1982, Section VIII.4] or [Cobzaş et al., 2019, Section 8.4] for summaries. The topological dual space can be identified with the quotient space of Lipschitz functions with respect to constant functions, or equivalently the space  $\text{Lip}(X, x_0)$  of Lipschitz functions vanishing at an arbitrary base point  $x_0 \in X$ , equipped with the Lipschitz norm

$$\|f\|_L = \sup_{x, y \in X, x \neq y} \left\{ \frac{|f(x) - f(y)|}{d(x, y)} \right\}.$$

We propose to extend the Kantorovich-Rubinstein norm to the space  $\mathcal{M}(X)$  of all countably additive measures on  $X$  as

$$\|\mu\|_{pK} = \inf_{\xi \in \mathcal{M}(X,0)} \left\{ (\|\xi\|_{KR}^p + \|\mu - \xi\|_{TV}^p)^{\frac{1}{p}} \right\} \quad (1)$$

with  $p \in [1, \infty]$  and  $\|\cdot\|_{TV}$  being the total variation norm, and to extend the Lipschitz norm to the space  $\text{Lip}(X)$  of all Lipschitz functions on  $X$  as

$$\|f\|_{qL} = (\|f\|_L^q + \|f\|_\infty^q)^{\frac{1}{q}} \quad (2)$$

with  $q \in [1, \infty]$  and  $\|\cdot\|_\infty$  being the sup norm. The limiting cases  $p, q = \infty$  are interpreted as usual. In the following, we will prove some properties of the resulting  $p$ -Kantorovich  $(\mathcal{M}(X), \|\cdot\|_{pK})$  and  $q$ -Lipschitz  $(\text{Lip}(X), \|\cdot\|_{qL})$  spaces, including that if  $p, q$  are Hölder conjugates, i.e.  $\frac{1}{p} + \frac{1}{q} = 1$ , then the topological dual of the  $p$ -Kantorovich space can be identified with the  $q$ -Lipschitz space.

The theory of the  $p = 1$  and  $q = \infty$  case was developed by L. G. Hanin in a series of papers [Hanin, 1992, Hanin, 1994, Hanin, 1997, Hanin, 1999]. See [Cobzaş et al., 2019, Section 8.5] for a summary. The case  $p = \infty$  and  $q = 1$  was proposed in dual form in [Chitescu et al., 2014]. Equivalence to the primal form was shown in [Terjék, 2021]. For  $\mu, \nu$  probability measures,  $\|\mu - \nu\|_{\infty K}$  is also known as the Fortet-Mourier distance.

## 2 $p$ -Kantorovich and $q$ -Lipschitz norms

The following propositions show that (1) and (2) define families of equivalent norms, with the spaces  $(\mathcal{M}(X), \|\cdot\|_{pK})$  and  $(\text{Lip}(X), \|\cdot\|_{qL})$  being incomplete and complete, respectively, while the pointwise product turns  $q$ -Lipschitz spaces into Gelfand algebras.

**Proposition 1.** *Given Hölder conjugates  $p, q \in [1, \infty]$ , for any  $\mu \in \mathcal{M}(X)$  and  $f \in \text{Lip}(X)$ , one has*

$$\left| \int f d\mu \right| \leq \|\mu\|_{pK} \|f\|_{qL}. \quad (3)$$

*Proof.* For any  $f \in \text{Lip}(X)$ ,  $\mu \in \mathcal{M}(X)$  and  $\xi \in \mathcal{M}(X, 0)$ , one has

$$\begin{aligned} \left| \int f d\mu \right| &\leq \left| \int f d\xi \right| + \left| \int f d(\mu - \xi) \right| \leq \|f\|_L \|\xi\|_{KR} + \|f\|_\infty \|\mu - \xi\|_{TV} \\ &\leq (\|f\|_L^q + \|f\|_\infty^q)^{\frac{1}{q}} (\|\xi\|_{KR}^p + \|\mu - \xi\|_{TV}^p)^{\frac{1}{p}} = \|f\|_{qL} (\|\xi\|_{KR}^p + \|\mu - \xi\|_{TV}^p)^{\frac{1}{p}} \end{aligned}$$

where the Hölder inequality for  $\mathbb{R}^2$  was used. Since this holds for any  $\xi$ , the proposition follows from the definition of  $\|\cdot\|_{pK}$ .  $\square$

**Proposition 2.** *Given  $p, q \in [1, \infty]$ ,  $\|\cdot\|_{pK}$  is a norm on  $\mathcal{M}(X)$  and  $\|\cdot\|_{qL}$  is a norm on  $\text{Lip}(X)$ .*

*Proof.* Both functionals are clearly seminorms. By Proposition 1 they are separating, so that they are norms as well.  $\square$

**Proposition 3.** *Given  $p_1, p_2 \in [1, \infty]$ , the norms  $\|\cdot\|_{p_1K}$  and  $\|\cdot\|_{p_2K}$  on  $\mathcal{M}(X)$  are equivalent. Given  $q_1, q_2 \in [1, \infty]$ , the norms  $\|\cdot\|_{q_1L}$  and  $\|\cdot\|_{q_2L}$  on  $\text{Lip}(X)$  are equivalent.*

*Proof.* Suppose that  $p_1 \leq p_2$  and  $q_1 \leq q_2$ . One has

$$\begin{aligned} 2^{-\frac{1}{p_1}} (\|\xi\|_{KR}^{p_1} + \|\mu - \xi\|_{TV}^{p_1})^{\frac{1}{p_1}} &\leq \max\{\|\xi\|_{KR}, \|\mu - \xi\|_{TV}\} \leq (\|\xi\|_{KR}^{p_2} + \|\mu - \xi\|_{TV}^{p_2})^{\frac{1}{p_2}} \\ &\leq 2^{\frac{1}{p_2}} \max\{\|\xi\|_{KR}, \|\mu - \xi\|_{TV}\} \leq 2^{\frac{1}{p_2}} (\|\xi\|_{KR}^{p_1} + \|\mu - \xi\|_{TV}^{p_1})^{\frac{1}{p_1}} \end{aligned}$$

for any  $\mu \in \mathcal{M}(X)$  and  $\xi \in \mathcal{M}(X, 0)$ . Similarly, one has

$$\begin{aligned} 2^{-\frac{1}{q_1}} (\|f\|_L^{q_1} + \|f\|_\infty^{q_1})^{\frac{1}{q_1}} &\leq \max\{\|f\|_L, \|f\|_\infty\} \leq (\|f\|_L^{q_2} + \|f\|_\infty^{q_2})^{\frac{1}{q_2}} \\ &\leq 2^{\frac{1}{q_2}} \max\{\|f\|_L, \|f\|_\infty\} \leq 2^{\frac{1}{q_2}} (\|f\|_L^{q_1} + \|f\|_\infty^{q_1})^{\frac{1}{q_1}} \end{aligned}$$

for any  $f \in \text{Lip}(X)$ .  $\square$

**Proposition 4.** *Given  $p, q \in [1, \infty]$ , the normed space  $(\mathcal{M}(X), \|\cdot\|_{pK})$  is not complete if  $X$  is an infinite set, while the normed space  $(\text{Lip}(X), \|\cdot\|_{qL})$  is complete for any  $X$ , i.e. a Banach space.*

*Proof.* The following is an adaptation of the proof of [Cobzaş et al., 2019, Theorem 8.4.7]. If  $X$  is infinite, it has an accumulation point  $x$ . Let  $(x_n)$  be a sequence in  $X \setminus \{x\}$  converging to  $x$ . Since  $\|\delta_x - \delta_{x_n}\|_{pK} \leq d(x, x_n)$ , one has  $\lim_{n \rightarrow \infty} \|\delta_x - \delta_{x_n}\|_{pK} = 0$ . On the other hand,  $\|\delta_x - \delta_{x_n}\|_{TV} = 2$ , so that  $\delta_{x_n}$  does not converge to  $\delta_x$  with respect to the topology induced by  $\|\cdot\|_{TV}$ , meaning that  $\|\cdot\|_{TV}$  and  $\|\cdot\|_{pK}$  are not equivalent norms. If  $(\mathcal{M}(X), \|\cdot\|_{pK})$  was complete, the identity operator from  $(\mathcal{M}(X), \|\cdot\|_{pK})$  to  $(\mathcal{M}(X), \|\cdot\|_{TV})$  would be an isomorphism by the Banach isomorphism theorem, which would imply the equivalence of the norms  $\|\cdot\|_{TV}$  and  $\|\cdot\|_{pK}$ .

The following is an adaptation of the proof of [Cobzaş et al., 2019, Theorem 8.1.3]. Suppose that  $(f_n)$  is a Cauchy sequence with respect to  $\|\cdot\|_{qL}$ . Then it is also Cauchy with respect to  $\|\cdot\|_\infty$ , hence converges uniformly to some bounded function  $f : X \rightarrow \mathbb{R}$ . Since it is also Cauchy with respect to  $\|\cdot\|_L$ , by [Cobzaş et al., 2019, Lemma 8.1.4] it also converges to the same  $f$  with respect to  $\|\cdot\|_L$ , and  $f$  is Lipschitz. Consequently,  $(f_n)$  converges to  $f$  with respect to both  $\|\cdot\|_\infty$  and  $\|\cdot\|_L$ , hence with respect to  $\|\cdot\|_{qL}$  as well, so that the space  $(\text{Lip}(X), \|\cdot\|_{qL})$  is complete.  $\square$

**Corollary 5.** *Given  $q \in [1, \infty]$ , defining the product of  $f, g \in \text{Lip}(X)$  as  $fg(x) = f(x)g(x)$  turns  $(\text{Lip}(X), \|\cdot\|_{qL})$  into a complete normed algebra whose product is continuous, i.e. a Gelfand algebra.*

*Proof.* The result is known for  $q = 1$  and  $q = \infty$ , see [Weaver, 2018, Section 7.1]. The result follows by the equivalence of  $q$ -Lipschitz norms by Proposition 3 and the completeness of  $(\text{Lip}(X), \|\cdot\|_{qL})$  by Proposition 4.  $\square$

### 3 Duality of $p$ -Kantorovich and $q$ -Lipschitz spaces

The following proposition is needed to prove duality.

**Proposition 6.** *Given  $p \in [1, \infty]$ , the set of all measures with finite support is dense in  $(\mathcal{M}(X), \|\cdot\|_{pK})$ .*

*Proof.* By [Hanin, 1999, Lemma 2], the set of all measures with finite support is dense in  $(\mathcal{M}(X), \|\cdot\|_{1K})$ . Since the norms  $\|\cdot\|_{1K}$  and  $\|\cdot\|_{pK}$  are equivalent for any  $p \in [1, \infty]$  by Proposition 3, they generate the same topology, implying the proposition.  $\square$

We are going to apply techniques of convex analysis to show that the dual of the  $p$ -Kantorovich space can be identified with the  $q$ -Lipschitz space if  $\frac{1}{p} + \frac{1}{q} = 1$ .

**Theorem 7.** *Given Hölder conjugates  $p, q \in [1, \infty]$ , for any  $f \in Lip(X)$  the functional  $u_f : (\mathcal{M}(X), \|\cdot\|_{pK}) \rightarrow \mathbb{R}$  defined by  $u_f(\mu) = \int f d\mu$  is linear and continuous with  $\|u_f\|_{pK}^* = \|f\|_{qL}$ . Moreover, every continuous linear functional  $v$  on  $(\mathcal{M}(X), \|\cdot\|_{pK})$  is of the form  $v(\mu) = u_f(\mu)$  for a uniquely determined function  $f \in Lip(X)$  with  $\|v\|_{pK}^* = \|f\|_{qL}$ . Consequently, the mapping  $(f \rightarrow u_f)$  is an isometric isomorphism of  $(Lip(X), \|\cdot\|_{qL})$  onto the topological dual  $(\mathcal{M}(X), \|\cdot\|_{pK})^*$ , i.e.*

$$(Lip(X), \|\cdot\|_{qL}) \cong (\mathcal{M}(X), \|\cdot\|_{pK})^*. \quad (4)$$

*Proof.* It follows from (3) that  $u_f$  is a bounded and linear functional on  $(\mathcal{M}(X), \|\cdot\|_{pK})$ .

Consider the duality of  $(\mathcal{M}(X), \|\cdot\|_{1K})$  and  $(Lip(X), \|\cdot\|_{\infty L})$  given by [Hanin, 1999, Theorem 1]. For  $p \in [1, \infty]$ , let the indicators  $\iota_p : (\mathcal{M}(X), \|\cdot\|_{1K}) \rightarrow \overline{\mathbb{R}}$  be defined as

$$\iota_p(\mu) = \begin{cases} 0 & \text{if } \|\mu\|_{pK} \leq 1 \\ \infty & \text{otherwise} \end{cases}.$$

Their convex conjugates  $\iota_p^* : (Lip(X), \|\cdot\|_{\infty L}) \rightarrow \overline{\mathbb{R}}$  are defined as

$$\iota_p^*(f) = \sup_{\mu \in \mathcal{M}(X)} \left\{ \int f d\mu - \iota_p(\mu) \right\},$$

so that  $\iota_p^*(f) = \sup_{\|\mu\|_{pK} \leq 1} \left\{ \int f d\mu \right\} = \|u_f\|_{pK}^*$  is exactly the dual norm of  $u_f$ . We claim that  $\iota_p^*(f) = \|f\|_{qL}$ , which would prove that the linear map  $(f \rightarrow u_f)$  is an isometry.

Let  $H : (\mathcal{M}(X, 0), \|\cdot\|_{KR}) \times (\mathcal{M}(X), \|\cdot\|_{1K}) \rightarrow \mathbb{R}^2$  be defined as

$$H(\xi_1, \xi_2) = (\|\xi_1\|_{KR}, \|\xi_2\|_{TV}).$$

Let  $G : \mathbb{R}^2 \rightarrow \overline{\mathbb{R}}$  be defined as

$$G(x) = \begin{cases} 0 & \text{if } \|x\|_p \leq 1, \\ \infty & \text{otherwise,} \end{cases}$$

i.e.  $G$  is the indicator of the unit ball of the  $l^p$  norm on  $\mathbb{R}^2$ , so that its conjugate is  $G^*(y) = \|y\|_q$ , i.e. the  $l^q$  norm. With the usual ordering on  $\mathbb{R}^2$ ,  $H$  is clearly convex while  $g$  is proper, convex and increasing. Then the mapping  $\varphi = g \circ H$  is convex. We are going to invoke [Zalinescu, 2002, Theorem 2.8.10]. In the notation of the theorem,  $D = Y_0 = \mathbb{R}^2$  in our case, so that condition (vi) of the theorem clearly holds. This implies that the conjugate  $\varphi^* : (\text{Lip}(X, x_0), \|\cdot\|_L) \times (\text{Lip}(X), \|\cdot\|_{\infty L}) \rightarrow \overline{\mathbb{R}}$  is

$$\varphi^*(f_1, f_2) = \min_{y \in \mathbb{R}_+^2} \{(y_1 \|\cdot\|_{KR} + y_2 \|\cdot\|_{TV})^*(f_1, f_2) + \|y\|_q\}.$$

By [Zalinescu, 2002, Theorem 2.3.1(v)], [Zalinescu, 2002, Theorem 2.3.1(viii)] and the well known conjugate relations

$$(\xi_1 \rightarrow \|\xi_1\|_{KR})^* = \left( f_1 \rightarrow \begin{cases} 0 & \text{if } \|f_1\|_L \leq 1, \\ \infty & \text{otherwise} \end{cases} \right)$$

and

$$(\xi_2 \rightarrow \|\xi_2\|_{TV})^* = \left( f_2 \rightarrow \begin{cases} 0 & \text{if } \|f_2\|_{\infty} \leq 1, \\ \infty & \text{otherwise} \end{cases} \right),$$

the conjugate of the mapping

$$(x_1, x_2) \rightarrow y_1 \|\xi_1\|_{KR} + y_2 \|\xi_2\|_{TV}$$

is the mapping

$$(f_1, f_2) \rightarrow \begin{cases} 0 & \text{if } \|f_1\|_L \leq y_1 \text{ and } \|f_2\|_{\infty} \leq y_2, \\ \infty & \text{otherwise,} \end{cases}$$

so that one has

$$\varphi^*(f_1, f_2) = \min_{y \in \mathbb{R}_+^2, \|f_1\|_L \leq y_1, \|f_2\|_{\infty} \leq y_2} \{\|y\|_q\} = (\|f_1\|_L^q + \|f_2\|_{\infty}^q)^{\frac{1}{q}}.$$

Consider the linear map  $A \in L((\mathcal{M}(X, 0), \|\cdot\|_{KR}) \times (\mathcal{M}(X), \|\cdot\|_{1K}), (\mathcal{M}(X), \|\cdot\|_{1K}))$  defined as

$$A(\xi_1, \xi_2) = \xi_1 + \xi_2,$$

which is clearly bounded, hence continuous.

Its adjoint  $A^* \in L((\text{Lip}(X), \|\cdot\|_{\infty L}), (\text{Lip}(X, x_0), \|\cdot\|_L) \times (\text{Lip}(X), \|\cdot\|_{\infty L}))$  is given by

$$A^*f = (f - f(x_0), f).$$

By [Zalinescu, 2002, Theorem 2.3.1(ix)], the conjugate of

$$A\varphi(\mu) = \inf_{(\xi_1, \xi_2) \in (\mathcal{M}(X, 0), \|\cdot\|_{KR}) \times (\mathcal{M}(X), \|\cdot\|_{1K}), \mu = A(\xi_1, \xi_2)} \{\varphi(\xi_1, \xi_2)\} = \iota_p(\mu)$$

is the mapping

$$\varphi^* \circ A^*(f) = (\|f - f(x_0)\|_L^q + \|f\|_{\infty}^q)^{\frac{1}{q}} = \|f\|_{qL},$$

proving the claim, so that  $(f \rightarrow u_f)$  is an isometry.

To see that  $(f \rightarrow u_f)$  is onto  $\text{Lip}(X)$ , take any  $v \in (\mathcal{M}(X), \|\cdot\|_{pK})^*$  and set  $f(x) = v(\delta_x)$  for all  $x \in X$ . One has  $|f(x)| \leq \|v\|_{pK}^* \|\delta_x\|_{pK} = \|v\|_{pK}^*$  and  $|f(x) - f(y)| \leq \|v\|_{pK}^* \|\delta_x - \delta_y\|_{pK} \leq \|v\|_{pK}^* d(x, y)$  for any  $x, y \in X$ , so that  $f \in \text{Lip}(X)$ . One has  $u_f(\delta_x) = v(\delta_x)$ , so that by linearity,  $u_f(\mu) = v(\mu)$  for any  $\mu \in \mathcal{M}(X)$  with finite support. Since such measures are dense in  $(\mathcal{M}(X), \|\cdot\|_{pK})$  by Proposition 6, one has  $u_f = v$ , completing the proof.  $\square$

We get the following dual representation of  $\|\cdot\|_{pK}$  as a corollary.

**Corollary 8.** *Given  $\mu \in \mathcal{M}(X)$ , one has the dual representation*

$$\|\mu\|_{pK} = \sup_{f \in \text{Lip}(X), \|f\|_{qL} \leq 1} \left\{ \int f d\mu \right\}, \quad (5)$$

and there exists  $f_* \in \text{Lip}(X)$  such that  $\|f_*\|_{qL} = 1$  and  $\int f_* d\mu = \|\mu\|_{pK}$ .

*Proof.* This follows from Proposition 7 and the Hahn-Banach theorem.  $\square$

## 4 Optimality conditions

The following proposition shows that the infimum in (1) is always attained.

**Proposition 9.** *For any  $\mu \in \mathcal{M}(X)$ , there exists  $\xi_* \in \mathcal{M}(X, 0)$  such that  $\|\mu\|_{pK} = (\|\xi_*\|_{KR}^p + \|\mu - \xi_*\|_{TV}^p)^{\frac{1}{p}}$ .*

*Proof.* The following is an adaptation of the proof of [Hanin, 1999, Proposition 6]. Let  $\xi_n$  be a sequence in  $\mathcal{M}(X, 0)$  such that  $\|\mu\|_{pK} = \lim_{n \rightarrow \infty} (\|\xi_n\|_{KR}^p + \|\mu - \xi_n\|_{TV}^p)^{\frac{1}{p}}$ . Without loss of generality one can assume that  $\|\xi_n\|_{TV} \leq R$  for all  $n$  for some constant  $R > 0$ . By the weak-\* compactness of the  $\|\cdot\|_{TV}$ -ball of radius  $R$  by [Cobzaş et al., 2019, Theorem 8.4.25], up to a subsequence,  $\xi_n$  weak-\* converges to some  $\xi_* \in \mathcal{M}(X, 0)$ . Let  $a = \liminf_{n \rightarrow \infty} \|\xi_n\|_{KR}$  and  $b = \liminf_{n \rightarrow \infty} \|\mu - \xi_n\|_{TV}$ , so that  $\|\mu\|_{pK} \geq (a^p + b^p)^{\frac{1}{p}}$ . For every  $n$  and any  $f \in \text{Lip}(X)$  with  $\|f\|_L \leq 1$  one has  $|\int f d\xi_n| \leq \|\xi_n\|_{KR}$ , so that  $|\int f d\xi_*| \leq a$ , hence  $\|\xi_*\|_{KR} \leq a$ . Similarly, for every  $n$  and any  $f \in \text{Lip}(X)$  with  $\|f\|_\infty \leq 1$  one has  $|\int f d(\mu - \xi_n)| \leq \|\mu - \xi_n\|_{TV}$ , so that  $|\int f d(\mu - \xi_*)| \leq b$ , hence  $\|\mu - \xi_*\|_{TV} \leq b$ . Thus  $(\|\xi_*\|_{KR}^p + \|\mu - \xi_*\|_{TV}^p)^{\frac{1}{p}} \leq (a^p + b^p)^{\frac{1}{p}} \leq \|\mu\|_{pK}$ , implying the proposition.  $\square$

It is well known that for any  $\xi \in \mathcal{M}(X, 0)$  there exists  $\pi_* \in \mathcal{M}(X \times X)$  such that  $\pi_*(X \times \cdot) - \pi_*(\cdot \times X) = \xi$  and  $\|\xi\|_{KR} = \int d(x, y) d\pi_*(x, y)$  (see e.g. [Kantorovich and Akilov, 1982, Section VIII.4]). By this and Proposition 9, there exists an optimal pair  $\xi_* \in \mathcal{M}(X, 0)$ ,  $\pi_* \in \mathcal{M}(X \times X)$  such that  $\pi_*(X \times \cdot) - \pi_*(\cdot \times X) = \xi_*$  and

$$\|\mu\|_{pK} = \left( \left( \int d(x, y) d\pi_*(x, y) \right)^p + \|\xi_*\|_{TV}^p \right)^{\frac{1}{p}},$$

while by Proposition 8 there exists an optimal  $f_* \in Lip(X)$  such that  $\|\mu\|_{pK} = \int f_* d\mu$  and  $\|f\|_{qL} = 1$ . The following proposition characterizes such optimal variables, generalizing [Hanin, 1999, Proposition 7].

**Proposition 10.** *Given  $\mu \in \mathcal{M}(X)$ , the measures  $\xi_* \in \mathcal{M}(X, 0)$ ,  $\pi_* \in \mathcal{M}(X \times X)$  with  $\pi_*(X \times \cdot) - \pi_*(\cdot \times X) = \xi_*$  are optimal if and only if there exists a  $f_* \in Lip(X)$  such that the conditions*

$$(i) \quad \|f_*\|_{qL} = 1,$$

$$(ii) \quad \|f_*\|_L \|\xi_*\|_{KR} + \|f_*\|_\infty \|\mu - \xi_*\|_{TV} = (\|\xi_*\|_{KR}^p + \|\mu - \xi_*\|_{TV}^p)^{\frac{1}{p}},$$

$$(iii) \quad f_*(x) - f_*(y) = \|f_*\|_L d(x, y) \text{ if } (x, y) \in \text{supp}(\pi_*) \text{ and}$$

$$(iv) \quad f_*(x) = \pm \|f_*\|_\infty \text{ if } x \in \text{supp}((\mu - \xi_*)_\pm)$$

are satisfied. In this case,  $f_*$  is optimal, i.e.  $\int f_* d\mu = \|\mu\|_{pK}$ .

*Proof.* Let  $f_* \in Lip(X)$  be a function satisfying the above conditions. Then one has

$$\begin{aligned} \|\mu\|_{pK} &\geq \int f_* d\mu = \int f_* d\xi_* + \int f_* d(\mu - \xi_*) \\ &= \int f_*(x) - f_*(y) d\pi_*(x, y) + \|f_*\|_\infty \|\mu - \xi_*\|_{TV} \\ &= \|f_*\|_L \int d(x, y) d\pi_*(x, y) + \|f_*\|_\infty \|\mu - \xi_*\|_{TV} \\ &\geq \|f_*\|_L \|\xi_*\|_{KR} + \|f_*\|_\infty \|\mu - \xi_*\|_{TV} \\ &= (\|\xi_*\|_{KR}^p + \|\mu - \xi_*\|_{TV}^p)^{\frac{1}{p}} \geq \|\mu\|_{pK}, \end{aligned}$$

so that the conditions are sufficient.

Now let  $\xi_*$ ,  $\pi_*$  and  $f_*$  be optimal. Clearly (i) is satisfied. On the other hand, one has

$$\begin{aligned} \|\mu\|_{pK} &= \int f_* d\mu = \int f_* d\xi_* + \int f_* d(\mu - \xi_*) \\ &\leq \int f_*(x) - f_*(y) d\pi_*(x, y) + \|f_*\|_\infty \|\mu - \xi_*\|_{TV} \\ &\leq \|f_*\|_L \int d(x, y) d\pi_*(x, y) + \|f_*\|_\infty \|\mu - \xi_*\|_{TV} \\ &= \|f_*\|_L \|\xi_*\|_{KR} + \|f_*\|_\infty \|\mu - \xi_*\|_{TV} \\ &\leq (\|f\|_L^q + \|f\|_\infty^q)^{\frac{1}{q}} (\|\xi_*\|_{KR}^p + \|\mu - \xi_*\|_{TV}^p)^{\frac{1}{p}} = \|\mu\|_{pK}, \end{aligned}$$

implying (ii), (iii) and (iv). This shows that the conditions are necessary as well.  $\square$

## Acknowledgements

The author is supported by the Hungarian National Excellence Grant 2018-1.2.1-NKP-00008 and by the Hungarian Ministry of Innovation and Technology NRDI Office within the framework of the Artificial Intelligence National Laboratory Program.

## References

- [Chitescu et al., 2014] Chitescu, I., Ioana, L., Miculescu, R., and Nita, L. (2014). Monge–Kantorovich norms on spaces of vector measures. *Results in Mathematics*, 70:349–371.
- [Cobzaş et al., 2019] Cobzaş, Ş., Miculescu, R., and Nicolae, A. (2019). *Lipschitz Functions*. Lecture Notes in Mathematics. Springer International Publishing.
- [Hanin, 1992] Hanin, L. G. (1992). Kantorovich-Rubinstein norm and its application in the theory of Lipschitz spaces. *Proceedings of the American Mathematical Society*, 115(2):345–352.
- [Hanin, 1994] Hanin, L. G. (1994). On isometric isomorphism between the second dual to the “small” Lipschitz space and the “big” Lipschitz space. In *Nonselfadjoint Operators and Related Topics*, pages 316–324. Springer.
- [Hanin, 1997] Hanin, L. G. (1997). Duality for general Lipschitz classes and applications. *Proceedings of the London Mathematical Society*, 75(1):134–156.
- [Hanin, 1999] Hanin, L. G. (1999). An extension of the Kantorovich norm. *Contemporary Mathematics*, 226:113–130.
- [Kantorovich and Akilov, 1982] Kantorovich, L. V. and Akilov, G. P. (1982). *Functional Analysis*. Pergamon Press.
- [Kantorovich and Rubinstein, 1957] Kantorovich, L. V. and Rubinstein, G. S. (1957). On a functional space and certain extremum problems. *Dokl. Akad. Nauk SSSR*, 115:6:1058–1061.
- [Kantorovich and Rubinstein, 1958] Kantorovich, L. V. and Rubinstein, G. S. (1958). On a space of completely additive functions. *Vestnik Leningrad. Univ*, 13:52–59.
- [Terjék, 2021] Terjék, D. (2021). A primal representation of the Monge-Kantorovich norm.
- [Weaver, 2018] Weaver, N. (2018). *Lipschitz Algebras*. WORLD SCIENTIFIC, 2nd edition.
- [Zalinescu, 2002] Zalinescu, C. (2002). *Convex Analysis in General Vector Spaces*. World Scientific.