On Einstein hypersurfaces of $I \times_f \mathbb{Q}^n(c)$

V. BORGES AND A. DA SILVA

ABSTRACT. In this paper, we investigate Einstein hypersurfaces of the warped product $I \times_f \mathbb{Q}^n(c)$, where $\mathbb{Q}^n(c)$ is a space form of curvature c. We prove that M has at most three distinct principal curvatures and that it is locally a multiply warped product with at most two fibers. We also show that exactly one or two principal curvatures on an open set imply constant sectional curvature on that set. For exactly three distinct principal curvatures this is no longer true, and we classify such hypersurfaces provided it does not have constant sectional curvature and a certain principal curvature vanishes identically.

1. INTRODUCTION AND MAIN RESULTS

The investigation of Einstein hypersurfaces of space forms $\mathbb{Q}^{n+1}(c)$ goes back to the mid 30's [6, 14] and a complete local classification was obtained in 1968, by Ryan [13]. His proof goes through showing that these hypersurfaces have at most two distinct principal curvatures at each point. Using the integral manifolds of the distributions corresponding to the principal curvatures, he shows that either M has constant sectional curvature, which is not smaller than c, or c > 0 and the Einstein metric splits off as certain Riemannian products between spheres.

A complete classification is also known when the ambient space is a complex or quaternionic space form of non zero curvature. In this direction, the first result came out in 1982, when Cecil and Ryan [4] showed that there is no real Einstein hypersurface in the complex projective space \mathbb{CP}^n . Three years later, Montiel [11] showed the same result in the complex hyperbolic space \mathbb{CH}^n . Martinéz and Pérez [10], in turn, showed in 1985 that certain geodesic hyperspheres are the only real Einstein hypersurfaces of the quaternionic projective space \mathbb{QP}^n . The quaternionic hyperbolic space \mathbb{QH}^n , on the other hand, admits no real Einstein hypersurface, as proved in 1996 by Ortega and Peréz [12].

Einstein hypersurfaces of the cylinders $\mathbb{R} \times \mathbb{S}^n$ and $\mathbb{R} \times \mathbb{H}^n$ have also recently been classified [9]. Here the classical theory of isoparametric hypersurfaces of space forms, developed by Cartan [3], plays an important role. Another important point in this classification is the fact that the orthogonal projection of ∂_t on the Einstein hypersurface is a principal direction, where ∂_t is the field spanning the factor \mathbb{R} .

²⁰²⁰ Mathematics Subject Classification. Primary 53C42, 53B25, 53C25; Secondary 53C15.

Key words and phrases. Isometric Immersions, Einstein Manifolds, Warped Product, Multiply Warped Product, Frobenius Theorem.

Their main result asserts that Einstein hypersurfaces in these ambient have constant sectional curvature, which allowed the use of a previous classification.

In this paper we investigate Einstein manifolds M^n , satisfying $Ric = \rho g$, which are hypersurfaces of the warped product $I \times_f \mathbb{Q}^n(c)$. As three dimensional Einstein manifolds have constant sectional curvature, our main results are concerned with the case where $n \ge 4$. Here, $f: I \to \mathbb{R}$ is a positive smooth function, $\mathbb{Q}^n(c)$ is space form of constant curvature $c \in \{-1, 0, 1\}$ and I is an interval. This encloses a large variety of ambient spaces. For instance, space forms and cylinders can be obtained in this way by choosing suitable warping functions. However, we notice that our arguments do not apply to the case where the ambient $I \times_f \mathbb{Q}^n(c)$ has constant sectional curvature.

In order to state the main results of this paper, let us still denote by ∂_t the direction spanning I. Consider the orthogonal decomposition $\partial_t = T + \theta \mathcal{N}$ on points of M, where \mathcal{N} is the unit normal field of M. Both T and θ play an important role in the investigation of hypersurfaces of $I \times_f \mathbb{Q}^n(c)$. For instance, when T vanishes identically, M is isometric to $\{t\} \times \mathbb{Q}^n(c)$ with the induced metric (see [5, 8]), which has constant sectional curvature. In this case we say that M is a *slice*. Our main theorem is the following one.

Theorem 1. Let (M^n, g) , $n \ge 4$, be an Einstein hypersurface of $I \times_f \mathbb{Q}^n(c)$ and suppose that $I \times_f \mathbb{Q}^n(c)$ does not have constant sectional curvature. Let $U \subset M$ be an open set where T does not vanish. Then

- (1) M has at least 2 and at most 3 distinct principal curvatures on U;
- (2) T is a principal direction of M on U with multiplicity 1;
- (3) if M has exactly 3 distinct principal curvatures on U, say λ_n , λ_1 and λ_2 , then their multiplicities 1, p_1 and p_2 , respectively, are constant, and U is locally isometric to $J \times N_1^{p_1} \times N_2^{p_2}$, with metric

$$g = ds^2 + \varphi_1(s)^2 g_1 + \varphi_2(s)^2 g_2,$$

where $J \subset \mathbb{R}$ is an interval, $(N_i^{p_i}, g_i)$ is a space form and $\varphi_i : J \to \mathbb{R}$ is a positive smooth function, $i \in \{1, 2\}$.

A first consequence of item (1) is that if M is totally umbilical, then T vanishes. Consequently,

Corollary 1. Let (M^n, g) , $n \ge 4$, be an Einstein hypersurface of $I \times_f \mathbb{Q}^n(c)$ and suppose that $I \times_f \mathbb{Q}^n(c)$ does not have constant sectional curvature. If M is totally umbilical, then it is a slice.

In order to prove Theorem 1, we show that the distributions defined by the principal curvatures are involutive and manage to show that the metric of M splits as a multiply warped product (see Lemma 1 and Proposition 4). We remark that T being a principal direction is essential in our proof. It is exactly the case if

the ambient space does not have constant sectional curvature (see Proposition 2). Our tools are the fundamental equations describing a hypersurface of $I \times_f \mathbb{Q}^n(c)$, recently obtained in [8] ([5], for f constant), and Frobenius theorem.

When M has only two distinct principal curvatures in $U \subset M$, we are able to show that U is locally conformally flat. Consequently we have the following.

Theorem 2. Let (M^n, g) , $n \ge 4$, be an Einstein hypersurface of $I \times_f \mathbb{Q}^n(c)$ and suppose that $I \times_f \mathbb{Q}^n(c)$ does not have constant sectional curvature. Let $U \subset M$ be an open set where T does not vanish and where M has exactly two principal curvatures. Then, M has constant sectional curvature on U.

When M has three distinct principal curvatures on U, Theorem 1 says that U is locally isometric to a multiply warped product with one dimensional base and two fibers of constant sectional curvatures, say, k_1 and k_2 , and warping functions φ_1 and φ_2 . The curvature of such spaces can be expressed in terms of k_1 , k_2 , φ_1 , φ_2 and their derivatives. Such expressions can be found, for example, in [1]. When the principal curvature corresponding to T, denoted by λ_n , vanishes, these formulas for the curvature of M together with Gauss equation yield the following result.

Theorem 3. Let (M^n, g) , $n \ge 4$, be an Einstein hypersurface of $I \times_f \mathbb{Q}^n(c)$ and suppose that $I \times_f \mathbb{Q}^n(c)$ does not have constant sectional curvature. Let $U \subset M$ be an open set where T does not vanish and where λ_n vanishes. Then, M does not have constant sectional curvature in U if and only if the following are true

(1) $c = 1, p_1 \ge 2, p_2 \ge 2, n \ge 5, \theta \equiv 0$ and f is a solution of

$$\left(\frac{df}{dt}\right)^2 + \frac{\rho}{n-1}f^2 = \frac{n-3}{n-2};$$

(2) there are positive constants k_1 and k_2 , so that U is locally isometric to

(1.1)
$$I \times_{\varphi_1} \mathbb{S}^{p_1}\left(\frac{1}{\sqrt{k_1}}\right) \times_{\varphi_2} \mathbb{S}^{p_2}\left(\frac{1}{\sqrt{k_2}}\right),$$

where φ_i and the principal curvature λ_i , whose multiplicity is p_i , are given by

$$\varphi_i = \sqrt{\frac{(p_i - 1)k_i}{n - 3}} f, \quad \lambda_i = (-1)^i \sqrt{\frac{p_j - 1}{p_i - 1}} \frac{1}{f},$$

where $i \in \{1, 2\}$ and j = 3 - i.

We remark that three distinct principal curvatures do not occur neither in cylinders [9] nor in space forms [13]. This is in contrast with example (1.1).

The paper is organized as follows. In Section 2 we state the structure equations for the existence of an immersion of M^n in $I \times_f \mathbb{Q}^n(c)$. Our first result, which could be useful elsewhere, concerns hypersurfaces for which T is a principal direction. In Section 3 we investigate extrinsic and intrinsic properties of M. We prove that Mhas at most three distinct principal curvatures and that it is locally isometric to a multiply warped product with at most two fibers (see [1] for definitions). In Section 4, we prove the main theorems. We also rewrite the proof of the main result of [9]. See Theorem 5.

2. Preliminaries

The goal of this section is to fix notation and to set the main tools used along this paper.

2.1. Notations. Consider the simply connected, complete, *n*-dimensional space form $\mathbb{Q}^n(c)$ having constant curvature $c \in \{-1, 0, 1\}$ and a smooth real function $f: I \subseteq \mathbb{R} \to \mathbb{R}^+$. The warped product $I \times_f \mathbb{Q}^n(c)$ is the product manifold $I \times \mathbb{Q}^n(c)$ endowed with metric $\langle , \rangle = dt^2 + f(t)^2 g$, where g denotes the metric of $\mathbb{Q}^n(c)$.

Let M^n be an oriented hypersurface of $I \times_f \mathbb{Q}^n(c)$ with Levi-Civita connection ∇ . We set our convention for the curvature operator \mathcal{R} as

$$\mathcal{R}(X,Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]} Z.$$

Let \mathcal{N} be a normal unit vector field to M in $I \times_f \mathbb{Q}^n(c)$ and let A be the shape operator associated with \mathcal{N} . The mean curvature of M^n is defined by

$$H = \frac{1}{n}Tr(A).$$

If ∂_t denotes the field tangent to I in $I \times \mathbb{Q}^n(c)$, then it can be decomposed at points of M as

$$\partial_t = T + \theta \mathcal{N},$$

where T is tangent to M and $\theta = \langle \partial_t, \mathcal{N} \rangle$. The function θ is called *angle function*.

2.2. Structure equations. In this subsection we state the structure equations to obtain a Riemannian manifold (M^n, g) as a hypersurface of $I \times_f \mathbb{Q}^n(c)$. This result was first obtained for f constant in [5], and then for general f in [8].

Let (M^n, \langle, \rangle) be a Riemannian manifold with Levi-Civita connection ∇ and (0, 4)-curvature tensor \mathcal{R} . We say that M satisfies the *structure conditions* if there are: a smooth (1, 1)-tensor $A : \mathfrak{X}(M) \to \mathfrak{X}(M)$, a constant $c \in \{-1, 0, 1\}$, an interval $I \subset \mathbb{R}$ and smooth functions $f : I \to (0, +\infty), \theta : M \to \mathbb{R}$ and $\pi : M \to I$, with $T = \nabla \pi$, satisfying

- (A) A is self adjoint with respect to \langle, \rangle ;
- (B) $|T|^2 + \theta^2 = 1;$
- (C) $\nabla_X T = \frac{f'}{f} (X \langle X, T \rangle T) + \theta A X, \ \forall X \in \mathfrak{X}(M);$
- (D) $X(\theta) = -\langle AT, X \rangle \frac{f'}{f} \theta \langle X, T \rangle, \ \forall X \in \mathfrak{X}(M);$
- (E) For any $X, Y \in \mathfrak{X}(M)$,

$$(\nabla_X A)Y - (\nabla_Y A)X = \theta b \big(\langle T, X \rangle Y - \langle T, Y \rangle X \big);$$

(F) For any $X, Y, Z, W \in TM$,

$$\mathcal{R}(X,Y,Z,W) = a(\langle X,Z \rangle \langle Y,W \rangle - \langle Y,Z \rangle \langle X,W \rangle) + b(\langle X,Z \rangle \langle Y,T \rangle \langle W,T \rangle - \langle Y,Z \rangle \langle X,T \rangle \langle W,T \rangle - \langle X,W \rangle \langle Y,T \rangle \langle Z,T \rangle + \langle Y,W \rangle \langle X,T \rangle \langle Z,T \rangle) + \langle AY,Z \rangle \langle AX,W \rangle - \langle AY,W \rangle \langle AX,Z \rangle,$$

where

(2.1)
$$a = \frac{(f')^2 - c}{f^2}, \qquad b = \frac{f''}{f} - \frac{(f')^2}{f^2} + \frac{c}{f^2}.$$

We observe that b vanishes identically if and only if $I \times_f \mathbb{Q}^n(c)$ has constant sectional curvature (see Proposition 2 of [8]).

The result below is the main tool in investigating hypersurfaces in the warped product $I \times_f \mathbb{Q}^n(c)$. Its proof can be found in [8].

Theorem 4 ([8]). Let (M^n, \langle, \rangle) be a Riemannian manifold satisfying the structure conditions. Then, for each point $p \in M$, there exists a neighborhood U of p in M, an isometric immersion $\chi : U \to I \times_f \mathbb{Q}^n(c)$ and a normal vector field \mathcal{N} along χ such that

- (1) \mathcal{N} has length one in $I \times_f \mathbb{Q}^n(c)$;
- (2) $\pi_1 \circ \chi = \pi$, where $\pi_1 : I \times_f \mathbb{Q}^n(c) \to I$ is the projection in the first component;
- (3) The shape operator associated with \mathcal{N} is A;
- (4) (E) is the Codazzi equation and (F) is the Gauss equation;
- (5) Along χ one has $\partial_t = T + \theta \mathcal{N}$.

Conversely, if such immersion exists, then M satisfies the structure equations.

By tracing (F) we obtain as a consequence of Theorem 4 the following.

Corollary 2. The Ricci tensor Ric of M in $I \times_f \mathbb{Q}^n(c)$ is given by

$$Ric(X,Y) = -((n-1)a + |T|^2b)\langle X,Y\rangle - (n-2)b\langle X,T\rangle\langle Y,T\rangle$$
$$+ nH\langle AX,Y\rangle - \langle AX,AY\rangle.$$

2.3. Hypersurfaces of $I \times_f \mathbb{Q}^n(c)$ for which T is a principal direction. In this subsection we investigate hypersurfaces of the warped product $I \times_f \mathbb{Q}^n(c)$ which have $T \neq 0$ as principal direction on an open set.

Proposition 1. Let M^n be a hypersurface of $I \times_f \mathbb{Q}^n(c)$ and $U \subset M$ an open set. Assume that T is an eigenvector of $A : \mathfrak{X}(M) \to \mathfrak{X}(M)$, the shape operator of M, satisfying $AT = \lambda_n T$ on U. The following is true on U:

- (1) The integral curves of $\frac{T}{|T|}$ are geodesics;
- (2) λ_n is smooth and $\nabla \lambda_n \stackrel{|}{=} \left(\nabla_{\frac{T}{|T|}} A \right) \left(\frac{T}{|T|} \right) = \left\langle \frac{T}{|T|}, \nabla \lambda_n \right\rangle \frac{T}{|T|};$

(3)
$$\nabla \theta = -\left(\lambda_n + \frac{f'}{f}\theta\right)T.$$

In particular, $|T|^2$, θ and λ_n are constant on each connected component of regular levels of π contained in U.

Proof. Using (B) and (C) we have

$$\nabla_T T = \frac{f'}{f} (T - |T|^2 T) + \theta A T = \theta \left(\lambda_n + \frac{f'}{f} \theta \right) T,$$

what implies that the integral curves of $\frac{T}{|T|}$ are geodesics, proving (1). In order to prove (2) we start noting that since λ_n can be written as

$$\lambda_n = \frac{\langle AT, T \rangle}{|T|^2},$$

it is smooth. To prove the first equality of (2) we start differentiating the identity above with respect to $X \in \mathfrak{X}(M)$ to obtain

$$\begin{split} |T|^4 \langle \nabla \lambda_n, X \rangle &= (\langle \nabla_X (AT), T \rangle + \langle AT, \nabla_X T \rangle) |T|^2 - 2 \langle \nabla_X T, T \rangle \langle AT, T \rangle \\ &= (\langle \nabla_X (AT), T \rangle - \lambda_n \langle T, \nabla_X T \rangle) |T|^2, \end{split}$$

which gives

$$|T|^2 \langle \nabla \lambda_n, X \rangle = \langle \nabla_X (AT), T \rangle - \langle AT, \nabla_X T \rangle.$$

On the other hand, it follows from (E) that

$$\begin{split} \langle \nabla_X (AT), T \rangle &= \langle (\nabla_X A)T + A(\nabla_X T), T \rangle \\ &= \langle (\nabla_T A)X, T \rangle + \theta b \left\langle \langle T, X \rangle T - |T|^2 X, T \right\rangle + \left\langle \nabla_X T, AT \right\rangle \\ &= \left\langle (\nabla_T A)T, X \right\rangle + \left\langle \nabla_X T, AT \right\rangle, \end{split}$$

where in the last line we have used the identity $\langle (\nabla_Z A)X, Y \rangle = \langle (\nabla_Z A)Y, X \rangle$, $\forall X,Y,Z\in\mathfrak{X}(M). \text{ Consequently, } |T|^2\left<\nabla\lambda_n,X\right>=\left<(\nabla_TA)T,X\right>, \text{ as we claimed.}$

In order to finish the proof of (2) notice that since the integral curves of $\frac{T}{|T|}$ are geodesics, we have

$$\left(\nabla_{\frac{T}{|T|}}A\right)\left(\frac{T}{|T|}\right) = \nabla_{\frac{T}{|T|}}\left(A\left(\frac{T}{|T|}\right)\right) = \nabla_{\frac{T}{|T|}}\left(\lambda_n \frac{T}{|T|}\right) = \frac{\langle T, \nabla \lambda_n \rangle}{|T|^2}T,$$

which is the second equality of (2).

Now we prove (3). Using identity (D) we have

$$\langle \nabla \theta, X \rangle = X(\theta) = -\left\langle AT + \frac{f'}{f}\theta T, X \right\rangle = \left\langle -\left(\lambda_n + \frac{f'}{f}\theta\right)T, X \right\rangle,$$

for any $X \in \mathfrak{X}(M)$, which implies (3).

3. EINSTEIN HYPERSURFACES OF $I \times_f \mathbb{Q}^n(c)$

In this section we investigate Einstein hypersurfaces of $I \times_f \mathbb{Q}^n(c)$ assuming $b \neq 0$. This is equivalent to saying that $I \times_f \mathbb{Q}^n(c)$ does not have constant sectional curvature (see Proposition 2 of [8]). A complete local classification of Einstein hypersurfaces in spaces of constant sectional curvature can be found in [13]. Recall that as three dimensional Einstein manifolds have constant sectional curvature, we will assume from now on that $n \geq 4$.

3.1. Extrinsic structure of M. In this subsection we investigate properties of the principal curvatures of the immersion on an open set $U \subset M$ where T and b do not vanish. We start with the following proposition which, in particular, allows us to use Proposition 1.

Proposition 2. Let M^n , $n \ge 4$, be an Einstein hypersurface of $I \times_f \mathbb{Q}^n(c)$. If $T \ne 0$ and $b \ne 0$ at $x_0 \in M^n$, then T is an eigenvector of the shape operator A at x_0 . Furthermore, A has at least 2 and most 3 distinct principal curvatures at x_0 , say, λ_1, λ_2 and λ_n , which satisfy

(3.1)
$$\lambda_i^2 - nH\lambda_i + \rho + (n-1)a + |T|^2 b = 0, \ i \in \{1, 2\},$$

(3.2) $\lambda_n^2 - nH\lambda_n + \rho + (n-1)(a+|T|^2b) = 0.$

In particular, the multiplicity of λ_n as an eigenvalue of A at x_0 is 1.

Proof. Let $\{e_i\}_{i=1}^n$ be an orthonormal basis of principal directions at $x_0 \in M$, with $Ae_i = \lambda_i e_i$. If we write $T = \sum_{i=1}^n t_i e_i$, Corollary 2 gives the equality

$$R_{ij} = -(n-1)a\delta_{ij} - b\left(|T|^2\delta_{ij} + (n-2)t_it_j\right) + nH\lambda_i\delta_{ij} - \lambda_i\lambda_j\delta_{ij}$$

Since M^n is an Einstein hypersurface satisfying $Ric = \rho g$, we must have

(3.3)
$$\left[\lambda_i\lambda_j - nH\lambda_i + \rho + (n-1)a + |T|^2b\right]\delta_{ij} = -b(n-2)t_it_j.$$

Hence, we conclude that $t_i t_j = 0$ for all $i \neq j$. As $T \neq 0$ at x_0 , there is only one coefficient $t_k \neq 0$. Without loss of generality, assume that k = n. Thus, $AT = \lambda_n T$ at x_0 . Then, it follows from (3.3) that (3.1) and (3.2) hold. Now, since all λ_i , $i \in \{1, \ldots, n-1\}$, satisfy the second degree equation (3.1), there are at most two of them, say, λ_1 and λ_2 . Since n > 3, $b \neq 0$ and $T \neq 0$, (3.1) and (3.2) imply that $\lambda_1 \neq \lambda_n$ and $\lambda_2 \neq \lambda_n$, which finishes the proof.

As T is a principal direction of M, Proposition 1 says that λ_n and $|T|^2$ are constant on the connected regular levels of π . The next proposition asserts that the same is true for λ_1 and λ_2 . This is a consequence of (3.1) and (3.2).

Proposition 3. Let M^n be an Einstein hypersurface of $I \times_f \mathbb{Q}^n(c)$. Assume that $U \subset M$ is an open set where $T \neq 0$ and $b \neq 0$, and Σ_t is a connected level of π inside U. Then the principal curvatures of $M \subset I \times_f \mathbb{Q}^n(c)$ are constant on Σ_t . If there is, in addition, $x_0 \in \Sigma_t$ so that $\lambda_n(x_0) = 0$, then:

(1) If $b(x_0) < 0$, then $p_1 = n - 1$, $p_2 = 0$ on Σ_t .

(2) If $b(x_0) > 0$, then p_1 and p_2 are constant on Σ_t , $p_1 \ge 2$, $p_2 \ge 2$ and n > 4. Here, $p_i = p_i(x)$ denotes the multiplicity of $\lambda_i(x)$, $i \in \{1, 2\}$.

Proof. We start by observing that the functions a and b are constant on Σ_t and since T is an eigenvector of A, it follows from Proposition 1 that $|T|^2$, θ and λ_n are constant on Σ_t .

Suppose that $\lambda_n(x_0) \neq 0$ for some $x_0 \in \Sigma_t$. Then we have $\lambda_n(x) = \lambda_n(x_0) \neq 0$, $\forall x \in \Sigma_t$. If $X \in \mathfrak{X}(\Sigma_t)$, then (3.2) gives $X(H)\lambda_n = 0$ and then, H is constant on Σ_t which, together with (3.1), implies that the remaining principal curvatures of A are also constant on Σ_t .

Now assume that $\lambda_n(x_0) = 0$ for some $x_0 \in \Sigma_t$. As λ_n is constant on Σ_t , we have $\lambda_n(x) = 0$, $\forall x \in \Sigma_t$. Consequently, for each $x \in \Sigma_t$ we have

$$(3.4) nH = p_1\lambda_1 + p_2\lambda_2,$$

with $p_1, p_2 \in \{0, \ldots, n-1\}$ and $p_1 + p_2 = n-1$. Using (3.2) we also have $\rho = -(n-1)(a + |T|^2b)$, which by (3.1) implies that λ_1 and λ_2 are solution of the equation for y

(3.5)
$$y^2 - nHy - (n-2)|T|^2b = 0.$$

Observe that the equation above implies that $(\lambda_1 - \lambda_2)(\lambda_1 + \lambda_2 - nH) = 0.$

In what follows we proceed as in Theorem 3.1 of [13] (pages 374 and 375). We consider two situations according to the sign of $b(x_0)$.

Assume that $b(x_0) < 0$. In this case b(x) < 0, $\forall x \in \Sigma_t$. Assume that $\lambda_1 \neq \lambda_2$ at some point $x \in \Sigma_t$, that is $p_1 \ge 1$ and $p_2 \ge 1$. Since b(x) < 0, λ_1 and λ_2 have the same sign. As a consequence of identity (3.4) and equation (3.5), we have $\lambda_1 + \lambda_2 = nH = p_1\lambda_1 + p_2\lambda_2$, and then $(p_1 - 1)\lambda_1 + (p_2 - 1)\lambda_2 = 0$, which implies $p_1 = 1$ and $p_2 = 1$. In this case $n - 1 = p_1 + p_2 = 2$, what is a contradiction. Hence, the multiplicity of λ_1 on Σ_t is n - 1. Using equation (3.5) we obtain $\lambda_1^2 = -|T|^2 b$. This shows that λ_1 is constant on Σ_t .

Assume that $b(x_0) > 0$. In this case b(x) > 0, $\forall x \in \Sigma_t$. Assume that the multiplicity of λ_1 is n-1 at some point $x \in \Sigma_t$. Proceeding as before we conclude $0 \leq \lambda_1^2 = -|T|^2 b < 0$, what is a contradiction. Then $\lambda_1 \neq \lambda_2$ for all $x \in \Sigma_t$. In this case equation (3.5) implies that

(3.6)
$$\begin{aligned} \lambda_1 \lambda_2 &= -(n-2)|T|^2 b, \\ (p_1 - 1)\lambda_1 + (p_2 - 1)\lambda_2 &= 0 \end{aligned}$$

and Proposition 2.2 of [13] allows us to conclude that the multiplicities of λ_1 and λ_2 are constant on Σ_t , since they are the eigenvalues of the restriction of A to Σ_t . Assume that $p_1 = 1$. Then $p_2 = n - 2$ and, using (3.6), we get $\lambda_2 = 0$ and $(n-2)|T|^2b = 0$, which is a contradiction. The same argument applies if $p_2 = 1$.

8

Consequently $p_1 \ge 2$ and $p_2 \ge 2$, what gives $n \ge 5$. Solving (3.6) we get

$$\lambda_i^2 = \frac{(p_j - 1)(n - 2)}{p_i - 1} |T|^2 b_i$$

with $i \in \{1, 2\}$ and j = 3 - i. This shows that λ_i is constant on Σ_t .

3.2. Local Intrinsic Structure of M^n . In this subsection we prove that an Einstein hypersurface of $I \times_f \mathbb{Q}^n(c)$ is locally isometric to a multiply warped product whose number of fibers is either 1 or 2, according to M having either 2 or 3 principal curvatures, respectively.

Assume that $U \subset M^n$ is an open set for which T and b do not vanish, and that A has three distinct eigenvalues λ_n , λ_1 and λ_2 .

In what follows we consider the distributions D_i , $i \in \{1, 2, n\}$, which assign to each $x \in U$ the vector spaces

$$D_{i}(x) = \{ v \in T_{x}M; (A(x) - \lambda_{i}(x)I)v = 0 \}.$$

Note that the distributions defined above are mutually orthogonal. Note also that D_n is involutive, since it is smooth and has constant rank 1. The involutivity of D_i , $i \in \{1, 2\}$, is proved below.

Lemma 1. Assume that in the open set U the shape operator A has three distinct eigenvalues λ_n , λ_1 and λ_2 and that T and b do not vanish. The distributions D_i , $i \in \{1, 2, n\}$, are smooth, involutive and have rank p_1 , p_2 and 1, respectively. Consequently, if $x_0 \in U$ and J, $N_1^{p_1}$ and $N_2^{p_2}$ are the integral manifolds of D_n , D_1 and D_2 through x_0 , respectively, then, shrinking U, if necessary, it is diffeomorphic to $J \times N_1^{p_1} \times N_2^{p_2}$.

Proof. First, recall that it follows from Proposition 1 and Proposition 2 that λ_n is smooth and has multiplicity 1. Arguing in a similar way to [13, Proposition 2.2], it follows that λ_1 and λ_2 have constant multiplicities on U, say, p_1 and p_2 , respectively, and are smooth functions in this set.

Since for each $x \in U$ the distributions D_1 , D_2 and D_n are the eigenspaces of A associated to λ_1 , λ_2 and λ_n , respectively, they have constant ranks p_1 , p_2 and 1, respectively, in U. These distributions are smooth, once λ_1 , λ_2 and λ_n are. To see this, let $x \in U$ and consider differentiable vector fields E_1, \ldots, E_n around x, linearly independent at x, so that $E_1(x) \in D_n(x)$, $E_2(x), \ldots, E_{p_1+1}(x) \in D_1(x)$ and $E_{p_1+2}(x), \ldots, E_n(x) \in D_2(x)$. Define the smooth vector fields

$$F_1 = (A - \lambda_1 I) \circ (A - \lambda_2 I) E_1,$$

$$F_i = (A - \lambda_2 I) \circ (A - \lambda_n I) E_i, \ i \in \{2, \dots, p_1 + 1\},$$

$$F_i = (A - \lambda_1 I) \circ (A - \lambda_n I) E_i, \ i \in \{p_1 + 2, \dots, n\}.$$

We claim that $\{F_1\}, \{F_2, \ldots, F_{p_1+1}\}$ and $\{F_{p_1+2}, \ldots, F_n\}$ generate D_n, D_1 and D_2 , respectively, around x, from what follows that the distributions are smooth. The

claim follows from $(A - \lambda_i I) \circ (A - \lambda_j I)(TM) \subset D_k$, where i, j, k are distinct indices in $\{1, 2, n\}$. These inclusions, in turn, follow from

(3.7)
$$(A - \lambda_1 I) \circ (A - \lambda_2 I) \circ (A - \lambda_n I) = 0,$$

which reflects the fact that λ_n , λ_1 and λ_2 are the only eigenvalues of A.

Now, to see that D_i , $i \in \{1, 2\}$, is involutive, consider $X, Y \in D_i$. It follows from (E) that $(\nabla_X A)Y = (\nabla_Y A)X$. Consequently,

$$\begin{split} (A - \lambda_i I)[X, Y] =& A(\nabla_X Y) - A(\nabla_Y X) - \lambda_i \nabla_X Y + \lambda_i \nabla_Y X \\ = & \nabla_X (AY) - (\nabla_X A) Y - \nabla_Y (AX) + (\nabla_Y A) X \\ & -\lambda_i \nabla_X Y + \lambda_i \nabla_Y X \\ =& \lambda_i \nabla_X Y + X(\lambda_i) Y - \lambda_i \nabla_Y X - Y(\lambda_i) X \\ & -\lambda_i \nabla_X Y + \lambda_i \nabla_Y X \\ =& 0, \end{split}$$

where we have used Proposition 3 to assure that $X(\lambda_i) = Y(\lambda_i) = 0$. This implies that $[X, Y] \in D_i$, and then D_i is involutive.

The lemma follows from a well known application of Frobenius Theorem. See [7] for further details. $\hfill \Box$

Lemma 1 gives a topological decomposition of M on U. Now we seek for a compatible decomposition for the metric g on U. The first step in this task is given in the following lemma.

Lemma 2. Let $X_i \in D_i$, $i \in \{1, 2\}$, and $T \in D_n$. Then,

(1)
$$\nabla_T X_i = [T, X_i] + \left(\frac{f'}{f} + \theta \lambda_i\right) X_i;$$

(2) $\nabla_{X_j} X_i \in D_i, \text{ with } i, j \in \{1, 2\} \text{ and } i \neq j$

Proof. Item (1) follows immediately from (C). To prove item (2), observe that from (E), $(\nabla_{X_i} A)X_j = (\nabla_{X_j} A)X_i$, and from Proposition 3, $\nabla_{X_2}(\lambda_1 X_1) = \lambda_1 \nabla_{X_2} X_1$ and $\nabla_{X_1}(\lambda_2 X_2) = \lambda_2 \nabla_{X_1} X_2$. Using these identities we have

$$\begin{aligned} (A - \lambda_2 I)(\nabla_{X_1} X_2) = &\nabla_{X_1} (AX_2) - (\nabla_{X_1} A)X_2 - \lambda_2 \nabla_{X_1} X_2 \\ = &\nabla_{X_1} (\lambda_2 X_2) - (\nabla_{X_1} A)X_2 - \lambda_2 \nabla_{X_1} X_2 \\ = &\nabla_{X_2} (\lambda_1 X_1) - (\nabla_{X_2} A)X_1 - \lambda_1 \nabla_{X_2} X_1 \\ = &\nabla_{X_2} (AX_1) - (\nabla_{X_2} A)X_1 - \lambda_1 \nabla_{X_2} X_1 \\ = &(A - \lambda_1 I)(\nabla_{X_2} X_1). \end{aligned}$$

Observe that from (3.7), we get $(A - \lambda_i I)(TM) \subset D_j \oplus D_k$, i, j, k all distinct in $\{1, 2, n\}$. This implies that $(A - \lambda_1 I)(\nabla_{X_2} X_1) = (A - \lambda_2 I)(\nabla_{X_1} X_2) \in D_n$. But

since,

$$\begin{split} \langle (A - \lambda_1 I) \nabla_{X_2} X_1, T \rangle &= \langle \nabla_{X_2} X_1, AT \rangle - \lambda_1 \langle \nabla_{X_2} X_1, T \rangle \\ &= (\lambda_n - \lambda_1) \langle \nabla_{X_2} X_1, T \rangle \\ &= -(\lambda_n - \lambda_1) \langle X_1, \nabla_{X_2} T \rangle \\ &= -(\lambda_n - \lambda_1) \left(\frac{f'}{f} \langle X_1, X_2 \rangle + \theta \langle X_1, AX_2 \rangle \right) \\ &= -(\lambda_n - \lambda_1) \theta \lambda_2 \langle X_1, X_2 \rangle \\ &= 0, \end{split}$$

we have $(A - \lambda_1 I)(\nabla_{X_2} X_1) = (A - \lambda_2 I)(\nabla_{X_1} X_2) = 0$, and then $\nabla_{X_2} X_1 \in D_1$ and $\nabla_{X_1} X_2 \in D_2$, finishing the proof.

If J is an integral curve of T with arc length parameter s, one has $T = |T|\partial_s$ and, from Propositions 1 and 3, λ_1 , λ_2 , |T| and θ are functions only of s in U. On the other hand, the restrictions of f and f' to M obey the relation $\frac{df}{ds} = |T|f'$. With this in mind, the decomposition of g in U is given in the next result.

Proposition 4. There are Riemannian metrics g_1 and g_2 on N_1 and N_2 , the manifolds of Lemma 1, respectively, so that the open set U with the Riemannian metric induced by M is locally isometric to $J \times N_1^{p_1} \times N_2^{p_2}$ with the metric

(3.8)
$$g = ds^2 + \varphi_1(s)^2 g_1 + \varphi_2(s)^2 g_2,$$

where

(3.9)
$$\frac{1}{\varphi_i} \frac{d\varphi_i}{ds} = \frac{1}{|T|^2 f} \frac{df}{ds} + \frac{\theta \lambda_i}{|T|}, \ i \in \{1, 2\}.$$

The manifold $J \times N_1^{p_1} \times N_2^{p_2}$ regarded with the metric (3.8) is called multiply warped product and it is denoted by $J \times_{\varphi_1} N_1 \times_{\varphi_2} N_2$.

Proof. Let $x_0 \in U$ and consider a coordinate system $(s, x_1, \ldots, x_{p_1}, y_1, \ldots, y_{p_2})$ around x_0 for which $x_0 = (0, \ldots, 0)$, where $s \in J$ is the arc length parameter, $(x_1, \ldots, x_{p_1}) \in N_1^{p_1}$ and $(y_1, \ldots, y_{p_2}) \in N_2^{p_2}$. Consequently, $\partial_s \in D_n, \partial_{x_1}, \ldots, \partial_{x_{p_1}} \in D_1$ and $\partial_{y_1}, \ldots, \partial_{y_{p_2}} \in D_2$. Let g_{ij} be the representation of the metric of M in these coordinates. If $i, j \in \{1, \ldots, p_1\}$ and $k, l \in \{1, \ldots, p_2\}$, then item (2) of Lemma 2 gives $\nabla_{\partial_{x_i}} \partial_{y_k} = \nabla_{\partial_{y_k}} \partial_{x_i} \in D_1 \cap D_2 = \{0\}$, and then

$$\begin{aligned} \partial_{y_k}(g_{ij}) &= \left\langle \nabla_{\partial_{y_k}} \partial_{x_i}, \partial_{x_j} \right\rangle + \left\langle \partial_{x_i}, \nabla_{\partial_{y_k}} \partial_{x_j} \right\rangle = 0, \\ \partial_{x_i}(g_{kl}) &= \left\langle \nabla_{\partial_{x_i}} \partial_{y_k}, \partial_{y_l} \right\rangle + \left\langle \partial_{y_k}, \nabla_{\partial_{x_i}} \partial_{y_l} \right\rangle = 0 \end{aligned}$$

Thus, g_{ij} restricted to $N_1^{p_1}$ does not depend on the coordinates (y_1, \ldots, y_{p_2}) , and g_{kl} restricted to $N_2^{p_2}$ does not depend on the coordinates (x_1, \ldots, x_{p_1}) . On the

other hand, using item (1) of Lemma 2 and $[\partial_{x_i}, \partial_s] = 0$ we obtain

$$\partial_s(g_{ij}) = \left\langle \nabla_{\partial_s} \partial_{x_i}, \partial_{x_j} \right\rangle + \left\langle \partial_{x_i}, \nabla_{\partial_s} \partial_{x_j} \right\rangle$$
$$= \frac{2}{|T|} \left(\frac{f'}{f} + \theta \lambda_1 \right) g_{ij},$$

which gives

$$\partial_s(\ln|g_{ij}|) = 2\left(\frac{1}{|T|^2 f}\frac{df}{ds} + \frac{\theta\lambda_1}{|T|}\right).$$

Analogously,

$$\partial_s(\ln|g_{kl}|) = 2\left(\frac{1}{|T|^2 f}\frac{df}{ds} + \frac{\theta\lambda_2}{|T|}\right).$$

Note that $\partial_s(\ln |g_{ij}|)$ and $\partial_s(\ln |g_{kl}|)$ depend only on s. Integrating them from 0 to $s \in J$ and using the expressions above we get

(3.10)
$$g_{ij}(s, x_1, \dots, x_{p_1}, y_1, \dots, y_{p_2}) = \varphi_1(s)^2 g_{ij}(0, x_1, \dots, x_{p_1}, 0, \dots, 0), \\ g_{kl}(s, x_1, \dots, x_{p_1}, y_1, \dots, y_{p_2}) = \varphi_2(s)^2 g_{kl}(0, 0, \dots, 0, y_1, \dots, y_{p_2}),$$

where φ_i is given by (3.9). Now, consider the Riemannian metrics

(3.11)
$$(g_1)_{ij}(x_1, \dots, x_{p_1}) = g_{ij}(0, x_1, \dots, x_{p_1}, 0, \dots, 0) (g_2)_{kl}(y_1, \dots, y_{p_2}) = g_{kl}(0, 0, \dots, 0, y_1, \dots, y_{p_2}),$$

on $N_1^{p_1}$ and $N_2^{p_2}$, respectively. Using (3.10), (3.11), $g(\partial_s, \partial_s) = 1$, $g(\partial_s, \partial_{y_k}) = 0$, $g(\partial_s, \partial_{x_i}) = 0$ and $g(\partial_{x_i}, \partial_{y_k}) = 0$, we obtain (3.8), finishing the proof. \Box

If $U \subset M^n$ is an open set where A has exactly two distinct eigenvalues at each $x \in U$, say, λ_n and λ , applying similar arguments one has

Proposition 5. There is a Riemannian metric g_N on N^{n-1} and so that the open set U with the Riemannian metric induced by M is locally isometric to $J \times N^{n-1}$ with the metric

$$g = ds^2 + \varphi(s)^2 g_N,$$

where

$$\frac{1}{\varphi}\frac{d\varphi}{ds} = \frac{1}{|T|^2f}\frac{df}{ds} + \frac{\theta\lambda}{|T|}$$

In the next section we apply Proposition 4 and Proposition 5 to prove Theorem 1, Theorem 2 and Theorem 3.

EINSTEIN HYPERSURFACES

4. Applications

In this section we apply the results obtained in the last section. An immediate consequence is the following one.

Proof of Theorem 1. Follows essentially from Proposition 2 and Proposition 4. \Box

Next, we prove Theorem 2, Theorem 3 and use our arguments to prove the main theorem of [9].

Proof of Theorem 2. By Proposition 2 and Proposition 5 it follows that U can be taken isometric to $J \times_{\varphi} N^{n-1}$. Now consider orthogonal vector fields $X, Y \in \mathfrak{X}(N)$. Using the formulas for the sectional curvature of a multiply warped product (which may be found for example in [1], formulas (3), where their curvature tensor differs from ours by a sign) one gets

$$\left(sec_N(X,Y) - \left(\frac{d\varphi}{ds}\right)^2\right)\frac{1}{\varphi^2} = sec_M(X,Y) = \lambda^2 - a,$$

where in the right hand side we have used the Gauss formula or, more precisely, equation (F). Rewriting the equation above we get

$$\left(\frac{d\varphi}{ds}\right)^2 + (\lambda^2 - a)\varphi^2 = sec_N(X, Y).$$

Now we observe that from the equation above, there is a constant k such that $sec_N(X,Y) = k$, for any orthonormal vector fields $X, Y \in \mathfrak{X}(N)$. As a consequence, the warped product $J \times_{\varphi} N^{n-1}$ is locally conformally flat [1, 2, 15], and since it is isometric to U, which is Einstein, M has constant sectional curvature on U. \Box

When M has three distinct principal curvatures on U, Proposition 4 says that it is a multiply warped product with one dimensional base and two fibers. The curvature of such spaces can be expressed in terms of the warping functions (i.e., φ_1 and φ_2) and the curvatures of the fibers. Such expressions can be found, for example, in [1]. This and Gauss formula yield the following result.

Proposition 6. Let (M^n, g) , $n \ge 4$, be an Einstein hypersurface of $I \times_f \mathbb{Q}^n(c)$ and suppose that T and b do not vanish on an open set $U \subset M$. If M has exactly three distinct principal curvatures on U, say, λ_1 , λ_2 and λ_n , with multiplicities p_1 , p_2 and 1, respectively, then U is locally isometric to $J \times_{\varphi_1} N_1^{p_1} \times_{\varphi_2} N_2^{p_2}$, where $(N_i^{p_i}, g_{N_i})$ is a manifold of constant sectional curvature provided $p_i > 1$. Furthermore, the following equations are satisfied

(4.1)
$$\frac{d^2\varphi_i}{ds^2} = (a+b|T|^2 - \lambda_i\lambda_n)\varphi_i$$

(4.2)
$$\left(\frac{d\varphi_i}{ds}\right)^2 + (\lambda_i^2 - a)\varphi_i^2 = k_i, \text{ if } p_i > 1$$

(4.3)
$$\frac{d\varphi_1}{ds}\frac{d\varphi_2}{ds} = (a - \lambda_1\lambda_2)\varphi_1\varphi_2$$

where $i \in \{1, 2\}$.

Proof. The idea is to use the curvature formulas of a multiply warped product $J \times_{\varphi_1} N_1^{p_1} \times_{\varphi_2} N_2^{p_2}$ (which can be found in [2]) together with Gauss equation (equation (F)). The existence of the constant k_i follows in a similar fashion to the proof of Theorem 2.

Equations (4.1)-(4.3) seem to play an important role in the investigation of Einstein hypersurfaces of $I \times_f \mathbb{Q}^n(c)$ with 3 distinct principal curvatures. One application is in the proof of Theorem 3, which deals with the case where λ_n vanishes.

Proof of Theorem 3. Since M does not have constant sectional curvature, it follows from Theorem 1 and Theorem 2 that M has three distinct principal curvatures. Now, Proposition 3 and the assumption $\lambda_n \equiv 0$ assures that $p_1 \geq 2$ and $p_2 \geq 2$, and equations (3.1) and (3.2) imply that

(4.4)
$$a + |T|^{2}b = -\frac{\rho}{n-1},$$
$$\lambda_{1}\lambda_{2} = -(n-2)|T|^{2}b < 0,$$
$$(p_{1}-1)\lambda_{1} + (p_{2}-1)\lambda_{2} = 0.$$

Since $\lambda_1 \lambda_2 < 0$, we will assume that $\lambda_1 < 0 < \lambda_2$. Manipulating (4.4) one gets,

$$a = -\frac{\rho}{n-1} + \frac{1}{n-2}\lambda_1\lambda_2$$

and then

$$\lambda_i^2 - a = \frac{(n-3)p_j}{(n-2)(p_j-1)}\lambda_i^2 + \frac{\rho}{n-1}.$$

Fix $i \in \{1, 2\}$ and let j = 3 - i. Inserting the relations above into (4.1), (4.2) and (4.3) we obtain

(4.5)
$$\frac{d^2\varphi_i}{ds^2} = -\frac{\rho}{n-1}\varphi_i,$$

(4.6)
$$\left(\frac{d\varphi_i}{ds}\right)^2 + \left(\frac{(n-3)p_j}{(n-2)(p_j-1)}\lambda_i^2 + \frac{\rho}{n-1}\right)\varphi_i^2 = k_i,$$

(4.7)
$$\frac{d\varphi_1}{ds}\frac{d\varphi_2}{ds} = -\left(\frac{\rho}{n-1} + \frac{n-3}{n-2}\lambda_1\lambda_2\right)\varphi_1\varphi_2.$$

Using (4.5) and (4.6), we obtain

$$\frac{(n-3)p_j}{(n-2)(p_j-1)}\frac{d}{ds}\left(\lambda_i^2\varphi_i^2\right) = \frac{d}{ds}\left(k_i - \left(\frac{d\varphi_i}{ds}\right)^2 - \frac{\rho}{n-1}\varphi_i^2\right) = 0.$$

Consider positive constants A_1 and A_2 satisfying

(4.8)
$$\varphi_1 = -\frac{A_1}{\lambda_1} \quad \text{and} \quad \varphi_2 = \frac{A_2}{\lambda_2}.$$

In view of the last equation of (4.4) we get

(4.9)
$$\varphi_2 = \frac{(p_2 - 1)A_2}{(p_1 - 1)A_1}\varphi_1$$

Using this equality in (3.9) one has

$$\frac{1}{|T|^2 f} \frac{df}{ds} + \frac{\theta \lambda_1}{|T|} = \frac{1}{\varphi_1} \frac{d\varphi_1}{ds} = \frac{1}{\varphi_2} \frac{d\varphi_2}{ds} = \frac{1}{|T|^2 f} \frac{df}{ds} + \frac{\theta \lambda_2}{|T|}$$

and since $\lambda_1 \neq \lambda_2$, one obtains $\theta \equiv 0$. Therefore, $|T|^2 \equiv 1$, s = t and then

(4.10)
$$\frac{1}{f}\frac{df}{dt} = \frac{1}{\varphi_1}\frac{d\varphi_1}{dt} = \frac{1}{\varphi_2}\frac{d\varphi_2}{dt}$$

proving the existence of a constant $B_i > 0$ so that $\varphi_i = B_i f$. On the other hand, using the last equation of (4.4), (4.8) and (4.9) in (4.7) we get

(4.11)
$$B_i^2\left(\left(\frac{df}{dt}\right)^2 + \frac{\rho}{n-1}f^2\right) = \left(\frac{d\varphi_i}{dt}\right)^2 + \frac{\rho}{n-1}\varphi_i^2 = \frac{(p_i - 1)(n-3)A_i^2}{(p_j - 1)(n-2)},$$

and using the last equation of (4.6) and (4.8) we obtain

(4.12)
$$\left(\frac{d\varphi_i}{dt}\right)^2 + \frac{\rho}{n-1}\varphi_i^2 = k_i - \frac{(n-3)p_j A_i^2}{(n-2)(p_j-1)}.$$

From (4.11) and (4.12) we conclude that

(4.13)
$$\frac{(p_1-1)A_1}{B_1} = \frac{(p_2-1)A_2}{B_2}, \quad k_i = \frac{(n-3)A_i^2}{p_j-1} > 0.$$

On the other hand, by (2.1), (4.3) and (4.10) we have

$$\frac{(f')^2 - c}{f^2} = \frac{1}{\varphi_1 \varphi_2} \frac{d\varphi_2}{dt} \frac{d\varphi_2}{dt} + \lambda_1 \lambda_2 = \frac{(f')^2}{f^2} + \lambda_1 \lambda_2$$

and from (4.8) and $\varphi_i = B_i f$,

$$-\frac{c}{f^2} = \lambda_1 \lambda_2 = -\frac{A_1 A_2}{B_1 B_2 f^2} < 0,$$

from where it follows that c = 1 and $A_1A_2 = B_1B_2$. This together with (4.13) implies that the constants A_i and B_i are given by

$$A_i = \sqrt{\frac{p_j - 1}{p_i - 1}} B_i, \qquad B_i = \sqrt{\frac{(p_i - 1)k_i}{n - 3}}.$$

Using these expressions in $\varphi_i = B_i f$, (4.8) and (4.11) we prove the necessity. The proof that (1.1) is immersed in $I \times_f \mathbb{S}^n$ is a simple computation, where one has to check all conditions of Theorem 4.

The last result of this paper concerns cylinders, that is, the case where f constant.

Theorem 5 ([9]). Let (M^n, g) , $n \ge 4$, be an Einstein manifold locally immersed in $I \times_f \mathbb{Q}^n(c)$ and suppose that T does not vanish on an open set $U \subset M$. If f is constant and $c \ne 0$, then M has constant sectional curvature on U.

Proof. Assume by contradiction that M does not have constant sectional curvature on U and that f is constant. Without loss of generality, assume that $f \equiv 1$.

Theorem 1 and Theorem 2 assure that M has three distinct principal curvatures on a open set $U \subset M$. On the other hand, if λ_n vanishes, U is locally isometric to (1.1), by Theorem 3. But since f is constant, it follows from (4.5) that its Einstein constant is 0, what is a contradiction. Thus, λ_n does not vanish. Consequently, it follows from item (3) of Proposition 1 that θ is not constant.

On the other hand, from $f \equiv 1$, Proposition 4 and Proposition 6 we have

$$\frac{d\varphi_i}{ds} = \frac{\theta\lambda_i}{|T|}\varphi_i, \qquad \frac{d\varphi_1}{ds}\frac{d\varphi_2}{ds} = -(c+\lambda_1\lambda_2)\varphi_1\varphi_2,$$

and then $\lambda_1\lambda_2 = -c|T|^2$. Since we already have $\lambda_1\lambda_2 = \rho - (n-1)c + c|T|^2$, which follows from (3.1), we get $2c|T|^2 = (n-1)c - \rho$, which implies that θ is constant, and this is a contradiction. Thus, M has constant sectional curvature. \Box

References

- Brozos-Vázquez, M.; García-Río, E.; Vázquez-Lorenzo, R.: Warped product metrics and locally conformally flat structures. *Matemática Contemporânea*. 28, 91-110, (2005).
- [2] Brozos-Vázquez, M.; García-Río, E.; Vázquez-Lorenzo, R.: Some remarks on locally conformally flat static space-times. *Journal of mathematical physics*. 46(2), 022501, (2005).
- [3] Cartan, É. Familles de surfaces isoparamétriques dans les espaces à courbure constante. Annali di Matematica Pura ed Applicata. 17(1), 177-191, (1938).
- [4] Cecil, T. E.; Ryan, P. J.: Focal sets and real hypersurfaces in complex projective space. Transactions of the American Mathematical Society. 269(2), 481-499, (1982).
- [5] Daniel, B.: Isometric immersions into Sⁿ×ℝ and Hⁿ×ℝ and applications to minimal surfaces. Transactions of the American Mathematical Society. 361, no. 12, 6255-6282, (2009).
- [6] Fialkow, A.: Hypersurfaces of a space of constant curvature. Annals of Mathematics. 39(4), 762–85, (1938).
- [7] Kobayashi, S.; Nomizu, K.: Foundations of differential geometry Vol. 1, No. 2. New York. London, (1963).
- [8] Lawn, M. and Ortega, M.: A fundamental theorem for hypersurfaces in semi-Riemannian warped products. *Journal of Geometry and Physics.* 90, 55–70, (2015).
- [9] Leandro Neto, B.; Pina, R.; dos Santos, J.P.: Einstein Hypersurfaces of Sⁿ × ℝ and Hⁿ × ℝ. Bull Braz Math Soc. New Series 52, 537–546 (2021).
- [10] Martínez, A.; Pérez, J. D.: Real hypersurfaces in quaternionic projective space. Annali di matematica pura ed applicata. 145(1), 355-384, (1986).
- [11] Montiel, S.: Real hypersurfaces of a complex hyperbolic space. Journal of the Mathematical Society of Japan. 37(3), 515-535, (1985).
- [12] Ortega, M.; de Dios Pérez, J.: On the Ricci tensor of a real hypersurface of quaternionic hyperbolic space. *Manuscripta Mathematica*, **93(1)**, 49-57, (1997).

- [13] Ryan, P. J.: Homogeneity and some curvature conditions for hypersurfaces. Tohoku Mathematical Journal. 21(3). 363-388, (1969).
- [14] Thomas, T. Y. On closed spaces of constant mean curvature. American Journal of Mathematics. 58.4, 702-704, (1936).
- [15] Yau, S. T. Remarks on conformal transformations. Journal of Differential Geometry. 8(3), 369-381, (1973).

(V. Borges) FACULDADE DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO PARÁ, 66075-110 BE-LÉM, PARÁ, BRAZIL.

 $Email \ address: \verb+valterborges@ufpa.br+$

(A. da Silva) FACULDADE DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO PARÁ, 66075-110 BE-LÉM, PARÁ, BRAZIL.

Email address: adamsilva@ufpa.br