KILLING MAGNETIC CURVES IN NON-FLAT LORENTZIAN-HEISENBERG SPACES

MURAT ALTUNBAŞ

ABSTRACT. We obtain some explicit formulas for Killing magnetic curves in non-flat Lorentzian-Heisenberg spaces.

2020 Mathematics Subject Classification: 53C25, 53B30

1. INTRODUCTION

Let (M, g) be a three-dimensional (semi-)Riemannian manifold. The magnetic curves γ on M are generalizations of geodesics which satisfy the following differential equation:

(1.1)
$$\nabla_{\gamma'}\gamma' = \varphi(\gamma'),$$

where ∇ is the Levi-Civita connection of g and φ is a skew-symmetric (1, 1)-tensor field. The tensor field φ is known as the Lorentz force and Equation (1.1) is said to be the Lorentz equation. Magnetic curves were investigated by several authors in Riemannian and semi-Riemannian manifolds (see [10], [11], [12], [13]).

Moreover, when the magnetic fields (which will be explained later) correspond to a Killing vector, the curves γ which fulfill Equation (1.1) are said to be Killing magnetic curves. Studying Killing magnetic curves is an actual topic of research in pure mathematics and theoretical physics. In [14], Romaniuc and Munteanu considered Killing magnetic curves in three-dimensional Euclidean space. In [15], the same authors studied these curves in three-dimensional Minkowski space. In [5], Erjavec gave some characterizations about Killing magnetic curves in $SL(2,\mathbb{R})$. In [6] and [7], Killing magnetic curves were investigated in Sol space and almost cosymplectic Sol space, respectively. In [9], Munteanu and Nistor classified Killing magnetic curves in $S^2 \times \mathbb{R}$. In [3], Calvaruso, Munteanu and Perrone obtained a complete classification for the Killing magnetic curves in three-dimensional almost paracontact manifolds. In [2], Bejan and Romaniuc proved that equipped with a Killing vector field V, any arc length parameterized spacelike or timelike curve, normal to V, is a magnetic trajectory associated to V in a Walker manifold. And finally, in [4], Derkaoui and Hathout occured explicit formulas for Killing magnetic curves in Heisenberg group.

In this paper, we determine the Killing magnetic curves in the three-dimensional Lorentzian-Heisenberg space. It is known that Lorentzian-Heisenberg space can be equipped with three non-isometric metrics. We will consider two of them which are non-flat.

MURAT ALTUNBAŞ

2. Preliminaries

Let (M, g) be a three-dimensional semi-Riemannian manifold. The magnetic curves on (M, g) are trajectories of charged particles, moving on M under the action of a magnetic field F. A magnetic field on M is a closed 2-form F on M, to which one can associate a skew-symmetric (1, 1)-tensor field φ on M, uniquely determined by

$$F(X,Y) = g(\varphi(X),Y),$$

for all $X, Y \in \chi(M)$. Here, the tensor field φ is called the Lorentz force.

The magnetic trajectories of F are regular curves γ in M which satisfy the Lorentz equation

(2.1) $\nabla_{\mathbf{t}} \mathbf{t} = \varphi(\mathbf{t}),$

where $\mathbf{t} = \gamma'$ is the speed vector of γ .

Furthermore, to have a positively oriented orthonormal frame field $\{e_1, e_2, e_3\}$ and represent the vectors X and Y as $X = x_1e_1 + x_2e_2 + x_3e_3$ and $Y = y_1e_1 + y_2e_2 + y_3e_3$, the vector product of two vector fields $X = (x_1, x_2, x_3)$ and $Y = (y_1, y_2, y_3)$ is given by

$$X \wedge Y = (x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_2y_1 - x_1y_2).$$

The mixed product of the vector fields $X, Y, Z \in \chi(M)$ is then defined by

$$g(X \wedge Y, Z) = dv_g(X, Y, Z),$$

where dv_g denotes a volume on M.

A vector field V is called a Killing vector field if it satisfy the Killing equation

$$g(\nabla_X V, Y) + g(\nabla_Y V, X) = 0$$

for all $X, Y \in \chi(M)$, where ∇ is the Levi-Civita connection of the metric g.

Let $F_V = i_V dvg$ be the Killing magnetic force corresponding to the Killing magnetic vector field V on M, where i denotes the inner product. The Lorentz force of F_V is described as

$$\varphi(X) = V \wedge X$$

for all $X \in \chi(M)$. Therefore, Equation (2.1) can be rewritten as

(2.2)
$$\nabla_{\mathbf{t}} \mathbf{t} = V \wedge \mathbf{t}$$

and solutions of above equation are called Killing magnetic curves corresponding to the Killing vector fields V.

For shortness, we will call these curves as V-magnetic curves in this paper.

3. Metrics on Lorentzian-Heisenberg spaces

Each left-invariant Lorentzian metric on the 3-dimensional Heisenberg group H_3 is isometric to one of the following metrics:

$$g_{1} = -\frac{1}{\lambda^{2}}dx^{2} + dy^{2} + (xdy + dz)^{2},$$

$$g_{2} = \frac{1}{\lambda^{2}}dx^{2} + dy^{2} - (xdy + dz)^{2}, \ \lambda > 0,$$

$$g_{3} = dx^{2} + (xdy + dz)^{2} - ((1 - x)dy - dz)^{2}.$$

Furthermore, the Lorentzian metrics g_1, g_2, g_3 are non-isometric and the Lorentzian metric g_3 is flat [1]. We will deal with the metrics g_1 and g_2 (i.e. non-flat cases).

Remark 1. According to the coordinate change $u = \lambda^{-1}x$, v = y, w = z + 2xy, we rewrite the metrics as

(3.1)
$$g_1 = -du^2 + dv^2 + \lambda^2 (vdu - udv)^2, g_2 = du^2 + dv^2 - \lambda^2 (vdu - udv)^2, \ \lambda > 0.$$

4. The metric g_1

An orthonormal basis on (H_3, g_1) is given by

(4.1)
$$e_1 = \frac{\partial}{\partial z}, \ e_2 = \frac{\partial}{\partial y} - x \frac{\partial}{\partial z}, \ e_3 = \lambda \frac{\partial}{\partial x},$$

where the vector e_3 is timelike. The non-zero components of the Levi-Civita connection ∇ of the metric g_1 are given by

(4.2)
$$\nabla_{e_1} e_2 = \nabla_{e_2} e_1 = \frac{\lambda}{2} e_3,$$
$$\nabla_{e_1} e_3 = \nabla_{e_3} e_1 = \frac{\lambda}{2} e_2,$$
$$\nabla_{e_2} e_3 = -\nabla_{e_3} e_2 = \frac{\lambda}{2} e_1.$$

The Lie algebra of Killing vector fields of (H_3, g_1) admits as basis

(4.3)
$$V_1 = \frac{\partial}{\partial z}, V_2 = \frac{\partial}{\partial y}, V_3 = \lambda \frac{\partial}{\partial x} - \lambda y \frac{\partial}{\partial z},$$
$$V_4 = \lambda^2 y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y} - \frac{1}{2} (x^2 + \lambda^2 y^2) \frac{\partial}{\partial z}.$$

Using (4.1), we rewrite equations (4.3) as follows:

$$V_1 = e_1, V_2 = xe_1 + e_2, V_3 = -\lambda ye_1 + e_3,$$

$$V_4 = \frac{1}{2}(x^2 - \lambda^2 y^2)e_1 + xe_2 + \lambda ye_3.$$

If $\gamma : I \to (H_3, g_1), \ \gamma(t) = (x(t), y(t), z(t))$ is a regular curve, then its speed vector is described as

$$\mathbf{t} = \gamma'(t) = (x'(t), y'(t), z'(t))$$

and

(4.4)
$$\mathbf{t} = \gamma'(t) = (z' + xy')e_1 + y'e_2 + \frac{x'}{\lambda}e_3.$$

From equations (4.2), we have

(4.5)
$$\nabla_{\mathbf{t}}\mathbf{t} = (z' + xy')'e_1 + (y'' + x'(z' + xy'))e_2 + (\frac{x''}{\lambda} + \lambda y'(z' + xy'))e_3.$$

In the following subsections, we obtain some formulas for V_i -magnetic curves (i = 1, ..., 4) in (H_3, g_1) . To solve the differential equations, we need help of *Wolfram Mathematica*.

4.1. V_1 -magnetic curves. Using $(4.3)_1$ and (4.4), we occur

(4.6)
$$V_1 \wedge \mathbf{t} = -\frac{x'}{\lambda} e_2 - y' e_3.$$

The equation $\nabla_{\mathbf{t}} \mathbf{t} = V_1 \wedge \mathbf{t}$ gives us the following system:

(4.7)
$$S_1: \begin{cases} y'' + x'((z' + xy') + \frac{1}{\lambda}) = 0, \\ x'' + y'(\lambda^2(z' + xy') + \lambda) = 0, \\ (z' + xy')' = 0. \end{cases}$$

By integrating $(S_1)_3$ and putting it in $(S_1)_{1,2}$, we obtain

$$S_1: \begin{cases} y'' + x'(c + \frac{1}{\lambda}) = 0, \\ x'' + y'\lambda(\lambda c + 1) = 0, \\ z' + xy' = c \text{ (constant)} \end{cases}$$

Solution of the system $(S_1)_{1,2}$ is

(4.8)
$$\begin{cases} x(t) = -\frac{\lambda}{\lambda c+1} [k_1 \cosh((\lambda c+1)t) + k_2 \sinh((\lambda c+1)t)] + k_3, \\ y(t) = \frac{1}{\lambda c+1} [k_1 \sinh((\lambda c+1)t) + k_2 \cosh((\lambda c+1)t)] + k_4, \end{cases}$$

where k_i , i = 1, ..., 4 are constants. Setting equations (4.8) in $(S_1)_3$ and by integration, we get

$$z(t) = (c + \frac{\lambda}{2(\lambda c + 1)}(k_1^2 - k_2^2))t + \frac{\lambda}{(\lambda c + 1)^2} [\frac{(k_1^2 + k_2^2)}{4}\sinh(2(\lambda c + 1)t) \\ + \frac{k_1k_2}{2}\cosh(2(\lambda c + 1)t)] + \frac{1}{\lambda c + 1}[k_1k_3\sinh((\lambda c + 1)t) \\ + k_2k_3\cosh((\lambda c + 1)t)] + k_5,$$

where k_5 is a constant.

If $c = -\frac{1}{\lambda}$, the system S_1 reduces to

$$S_1: \begin{cases} y'' = 0, \\ x'' = 0, \\ z' + xy' = -\frac{1}{\lambda} \end{cases}$$

Its general solution

$$S_1: \begin{cases} x(t) = k_1 t + k_2, \\ y(t) = k_3 t + k_4, \\ z(t) = -\frac{k_1 k_3}{2} t^2 - (\frac{1}{\lambda} + k_2 k_3) t + k_5, \end{cases}$$

where k_i , i = 1, ..., 5 are constants. Therefore, we state the following theorem.

Theorem 1. All V_1 -magnetic curves of (H_3, g_1) satisfy the following equations: (i) If $c = -\frac{1}{\lambda}$, then

$$\gamma(t) = \begin{pmatrix} x(t) = k_1 t + k_2, \\ y(t) = k_3 t + k_4, \\ z(t) = -\frac{k_1 k_3}{2} t^2 - (\frac{1}{\lambda} + k_2 k_3) t + k_5 \end{pmatrix}.$$

$$\begin{aligned} \text{(ii) If } c \neq -\frac{1}{\lambda}, \text{ then} \\ \gamma(t) &= \begin{pmatrix} x(t) = -\frac{\lambda}{\lambda c+1} [k_1 \cosh((\lambda c+1)t) + k_2 \sinh((\lambda c+1)t)] + k_3, \\ y(t) = \frac{1}{\lambda c+1} [k_1 \sinh((\lambda c+1)t) + k_2 \cosh((\lambda c+1)t)] + k_4, \\ z(t) &= (c + \frac{\lambda}{2(\lambda c+1)} (k_1^2 - k_2^2))t + \frac{\lambda}{(\lambda c+1)^2} [\frac{(k_1^2 + k_2^2)}{4} \sinh((\lambda c+1)t)] \\ &+ \frac{k_1 k_2}{2} \cosh(2(\lambda c+1)t)] + \frac{1}{(\lambda c+1)} [k_1 k_3 \sinh((\lambda c+1)t)] \\ &+ k_2 k_3 \cosh((\lambda c+1)t)] + k_5 \end{pmatrix}, \end{aligned}$$

where k_i , i = 1, ..., 5 are constants.

4.2. V_2 -magnetic curves. According to $(4.3)_2$ and (4.4), we have

(4.9)
$$V_2 \wedge \mathbf{t} = \frac{x'}{\lambda} e_1 - \frac{xx'}{\lambda} e_2 - (xy' - (z' + xy'))e_3.$$

From the equation $\nabla_{\mathbf{t}} \mathbf{t} = V_2 \wedge \mathbf{t}$, we get

(4.10)
$$S_2: \begin{cases} y'' + x'(z' + xy') = -\frac{xx'}{\lambda}, \\ (\lambda y' - 1)(z' + xy') = -xy' - \frac{x''}{\lambda}, \\ (z' + xy')' = \frac{x'}{\lambda}. \end{cases}$$

By integrating $(S_2)_3$, we deduce

$$z' + xy' = \frac{x}{\lambda} + c,$$

where c is a constant. Putting the last equation in $(S_2)_{1,2}$ we get

$$\bar{S}_2: \begin{cases} y'' = -x'(\frac{2x}{\lambda} + c),\\ (\lambda y' - 1)(\frac{x}{\lambda} + c) = -xy' - \frac{x''}{\lambda}, \end{cases}$$

and

(4.11)
$$\bar{S}_2: \begin{cases} y' = -\frac{x^2}{\lambda} - xc, \\ x'' - 2x^3 - x - \lambda c(3x^2 + c\lambda x + 1) = 0 \end{cases}$$

Without loss of generality, we can suppose that c = 0. In this case, the equation $x'' - 2x^3 - x = 0$ involves Jacobi elliptic functions as solutions. So, we can express the following proposition.

Proposition 1. The Killing magnetic curves in (H_3, g_1) corresponding to the Killing vector field $V_2 = xe_1 + e_2$ are solutions of the system of differential equations (4.10).

4.3. V_3 -magnetic curves. From $(4.3)_3$ and (4.4), we have

(4.12)
$$V_3 \wedge \mathbf{t} = -y'e_1 + (yx' + (z' + xy'))e_2 + \lambda yy'e_3.$$

Using the equation $\nabla_{\mathbf{t}} \mathbf{t} = V_3 \wedge \mathbf{t}$, we obtain

(4.13)
$$S_3: \begin{cases} y'' + x'(z' + xy') = yx' + (z' + xy'), \\ \frac{x''}{\lambda} + \lambda y'(z' + xy') = \lambda yy', \\ (z' + xy')' = -y'. \end{cases}$$

By integrating $(S_3)_3$, we occur

$$z' + xy' = -y + c,$$

where c is a constant. Putting the last equation in $(S_3)_{1,2}$, we get

(4.14)
$$\bar{S}_3: \begin{cases} x' = \lambda^2 y^2 - \lambda^2 yc, \\ y'' - 2x'y + y + c(x'-1) = 0. \end{cases}$$

MURAT ALTUNBAŞ

Without loss of generality, we can assume that c = 0. In this case, when we try to solve the system \bar{S}_3 , *i.e.*, the equation $y'' - 2\lambda^2 y^3 + y = 0$, we encounter Jacobi elliptic functions. Therefore, we write the following proposition.

Proposition 2. The Killing magnetic curves in (H_3, g_1) corresponding to the Killing vector field $V_3 = -\lambda y e_1 + e_3$ are solutions of the system of differential equations (4.13).

4.4. V_4 -magnetic curves. From (4.3)₃ and (4.4), we write

(4.15)
$$V_4 \wedge \mathbf{t} = (\frac{xx'}{\lambda} - \lambda yy')e_1 - (\frac{1}{2}(x^2 - \lambda^2 y^2)\frac{x'}{\lambda} - \lambda y(z' + xy'))e_2 - (\frac{1}{2}(x^2 - \lambda^2 y^2)y' - x(z' + xy'))e_3.$$

From the equation $\nabla_{\mathbf{t}} \mathbf{t} = V_4 \wedge \mathbf{t}$, we get

(4.16)
$$S_4: \begin{cases} y'' + x'(z' + xy') = -\frac{1}{2}(x^2 - \lambda^2 y^2)\frac{x'}{\lambda} + \lambda y(z' + xy'), \\ \frac{x''}{\lambda} + \lambda y'(z' + xy') = -\frac{1}{2}(x^2 - \lambda^2 y^2)y' + x(z' + xy'), \\ (z' + xy')' = (\frac{xx'}{\lambda} - \lambda yy'). \end{cases}$$

By integrating $(S_4)_3$, we obtain

(4.17)
$$z' + xy' = \frac{x^2}{2\lambda} - \frac{\lambda y^2}{2} + c$$

where c is a constant. Putting the last equation in $(S_4)_{1,2}$, we get

(4.18)
$$\bar{S}_4: \begin{cases} y'' + \frac{x'}{\lambda}(x^2 - \lambda^2 y^2) = \frac{1}{2}y(x^2 - \lambda^2 y^2) + c(\lambda y - x'), \\ x'' + \lambda y'(x^2 - \lambda^2 y^2) = \frac{1}{2}x(x^2 - \lambda^2 y^2) + c(\lambda x - \lambda^2 y') \end{cases}$$

It seems very difficult to solve the system \bar{S}_4 in general case. For a particular case $x = \lambda y$, we deduce

(4.19)
$$\bar{S}_4: \begin{cases} x'' + c\lambda x' - c\lambda x = 0, \\ y'' + c\lambda y' - c\lambda y = 0. \end{cases}$$

By solving the second equation of the above system, we get

$$y(t) = k_1 e^{-\frac{t}{2}(c\lambda + \sqrt{c\lambda(4+c\lambda)})} + k_2 e^{\frac{t}{2}(-c\lambda + \sqrt{c\lambda(4+c\lambda)})},$$

where k_1 and k_2 are constants. From (4.17), we obtain

$$z(t) = ct - \frac{\lambda y^2}{2}$$

= $ct - \frac{\lambda}{2} (k_1 e^{-\frac{t}{2}(c\lambda + \sqrt{c\lambda(4+c\lambda)})} + k_2 e^{\frac{t}{2}(-c\lambda + \sqrt{c\lambda(4+c\lambda)})})^2$

Therefore, the solution of the system \bar{S}_4 is given by

$$(4.20) \qquad \bar{S}_4: \begin{cases} x(t) = \lambda \left(k_1 e^{-\frac{t}{2}(c\lambda + \sqrt{c\lambda(4+c\lambda)})} + k_2 e^{\frac{t}{2}(-c\lambda + \sqrt{c\lambda(4+c\lambda)})} \right), \\ y(t) = k_1 e^{-\frac{t}{2}(c\lambda + \sqrt{c\lambda(4+c\lambda)})} + k_2 e^{\frac{t}{2}(-c\lambda + \sqrt{c\lambda(4+c\lambda)})}, \\ z(t) = ct - \frac{\lambda}{2} (k_1 e^{-\frac{t}{2}(c\lambda + \sqrt{c\lambda(4+c\lambda)})} + k_2 e^{\frac{t}{2}(-c\lambda + \sqrt{c\lambda(4+c\lambda)})})^2. \end{cases}$$

Hence, we write the following proposition.

6

Proposition 3. The Killing magnetic curves in (H_3, g_1) corresponding to the Killing vector field $V_4 = \frac{1}{2}(x^2 - \lambda^2 y^2)e_1 + xe_2 + \lambda ye_3$ are solutions of the system of differential equations (4.16). Moreover, the space curves given by parametric equations (4.20) are V_4 -magnetic curves in (H_3, g_1) .

In the last section, we follow the steps explained in the strategy mentioned in this section for the metric g_2 .

5. The metric g_2

We have an orthonormal basis on (H_3, g_2)

(5.1)
$$e_1 = \frac{\partial}{\partial y} - x \frac{\partial}{\partial z}, \ e_2 = \lambda \frac{\partial}{\partial x}, \ e_3 = \frac{\partial}{\partial z},$$

where the vector e_3 is timelike. The non-zero components of the Levi-Civita connection ∇ of the metric g_2 are given by

(5.2)
$$\nabla_{e_1}e_2 = -\nabla_{e_2}e_1 = \frac{\lambda}{2}e_3,$$
$$\nabla_{e_1}e_3 = \nabla_{e_3}e_1 = \frac{\lambda}{2}e_2,$$
$$\nabla_{e_2}e_3 = \nabla_{e_3}e_2 = -\frac{\lambda}{2}e_1.$$

The Lie algebra of Killing vector fields of (H_3, g_2) admits as basis

(5.3)
$$V_1 = \frac{\partial}{\partial z}, V_2 = \frac{\partial}{\partial y}, V_3 = \lambda \frac{\partial}{\partial x} - \lambda y \frac{\partial}{\partial z},$$
$$V_4 = -\lambda^2 y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y} + \frac{1}{2} (-x^2 + \lambda^2 y^2) \frac{\partial}{\partial z}$$

Using (5.1), we rewrite Equations (5.3) as follows:

$$V_1 = e_3, V_2 = e_1 + xe_3, V_3 = e_2 - \lambda ye_3,$$

$$V_4 = xe_1 - \lambda ye_2 + \frac{1}{2}(x^2 + \lambda^2 y^2)e_3.$$

If $\gamma : I \to (H_3, g_2), \ \gamma(t) = (x(t), y(t), z(t))$ is a regular curve, then its speed vector is described as

$$\mathbf{t} = \gamma'(t) = (x'(t), y'(t), z'(t))$$

and

(5.4)
$$\mathbf{t} = \gamma'(t) = y'e_1 + \frac{x'}{\lambda}e_2 + (z' + xy')e_3.$$

From equations (5.2), we have

(5.5)
$$\nabla_{\mathbf{t}} \mathbf{t} = (y'' - x'(z' + xy'))e_1 + (\frac{x''}{\lambda} + \lambda y'(z' + xy'))e_2 + (z' + xy')'e_3.$$

5.1. V_1 -magnetic curves. We have

(5.6)
$$V_1 \wedge \mathbf{t} = -\frac{x'}{\lambda}e_1 + y'e_2$$

From the equation $\nabla_{\mathbf{t}} \mathbf{t} = V_1 \wedge \mathbf{t}$, we get

(5.7)
$$S_1: \begin{cases} y'' - x'((z'+xy') - \frac{1}{\lambda}) = 0, \\ x'' + y'(\lambda^2(z'+xy') - \lambda) = 0, \\ (z'+xy')' = 0. \end{cases}$$

By integrating $(S_1)_3$ and putting it in $(S_1)_{1,2}$, we obtain

$$S_1: \begin{cases} y'' - x'(c - \frac{1}{\lambda}) = 0, \\ x'' + y'\lambda(\lambda c - 1) = 0, \\ z' + xy' = c \text{ (constant)} \end{cases}$$

Solution of the system $(S_1)_{1,2}$ is

(5.8)
$$S_1: \begin{cases} x(t) = \frac{\lambda}{\lambda c - 1} [k_1 \cos((\lambda c - 1)t) + k_2 \sin((\lambda c - 1)t)] + k_3, \\ y(t) = \frac{1}{\lambda c - 1} [k_1 \sin((\lambda c - 1)t) - k_2 \cos((\lambda c - 1)t)] + k_4, \end{cases}$$

where k_i , i = 1, ..., 4 are constants. Setting equations (5.7) in $(S_1)_3$ and by integration, we get

$$z(t) = (c - \frac{\lambda}{2(\lambda c - 1)}(k_1^2 - k_2^2))t - \frac{\lambda}{(\lambda c - 1)^2} [\frac{(k_1^2 - k_2^2)}{4}\sin(2(\lambda c - 1)t) - \frac{k_1k_2}{2}\cos(2(\lambda c - 1)t)] + \frac{1}{\lambda c - 1}(k_1k_3\sin((\lambda c - 1)t)) - k_2k_3\cos((\lambda c - 1)t)) + k_5,$$

where k_5 is a constant. If $c = \frac{1}{\lambda}$, the system S_1 reduces to

$$S_1: \begin{cases} y'' = 0, \\ x'' = 0, \\ z' + xy' = \frac{1}{\lambda}. \end{cases}$$

Its general solution

$$S_1: \begin{cases} x(t) = k_1 t + k_2, \\ y(t) = k_3 t + k_4, \\ z(t) = -\frac{k_1 k_3}{2} t^2 + (\frac{1}{\lambda} - k_2 k_3) t + k_5, \end{cases}$$

where k_i , i = 1, ..., 5 are constants. So, we have proved the theorem below.

Theorem 2. All V_1 -magnetic curves of (H_3, g_2) satisfy the following equations: (i) If $c = \frac{1}{\lambda}$, then

$$\gamma(t) = \begin{pmatrix} x(t) = k_1 t + k_2, \\ y(t) = k_3 t + k_4, \\ z(t) = -\frac{k_1 k_3}{2} t^2 + (\frac{1}{\lambda} - k_2 k_3) t + k_5 \end{pmatrix}$$
(ii) If $c \neq \frac{1}{\lambda}$, then

$$\gamma(t) = \begin{pmatrix} x(t) = \frac{\lambda}{\lambda c - 1} [k_1 \cos((\lambda c - 1)t) + k_2 \sin((\lambda c - 1)t)] + k_3, \\ y(t) = \frac{1}{\lambda c - 1} [k_1 \sin((\lambda c - 1)t) - k_2 \cos((\lambda c - 1)t)] + k_4, \\ z(t) = (c - \frac{\lambda}{2(\lambda c - 1)} (k_1^2 - k_2^2))t - \frac{\lambda}{(\lambda c - 1)^2} [\frac{(k_1^2 - k_2^2)}{4} \sin(2(\lambda c - 1)t)] \\ -\frac{k_1 k_2}{2} \cos(2(\lambda c - 1)t)] + \frac{1}{\lambda c - 1} (k_1 k_3 \sin((\lambda c - 1)t)) \\ -k_2 k_3 \cos((\lambda c - 1)t)) + k_5 \end{pmatrix}$$

,

where k_i , i = 1, ..., 5 are constants.

Remark 2. These curves were considered by Lee in [8] according to corresponding metric g_2 in (3.1).

5.2. V_2 -magnetic curves. Direct computations give

(5.9)
$$V_2 \wedge \mathbf{t} = -\frac{xx'}{\lambda}e_1 + (xy' - (z' + xy'))e_2 - \frac{x'}{\lambda}e_3.$$

The equation $\nabla_{\mathbf{t}} \mathbf{t} = V_2 \wedge \mathbf{t}$ concludes

(5.10)
$$S_2: \begin{cases} y'' - x'(z' + xy') = -\frac{xx'}{\lambda}, \\ (\lambda y' + 1)(z' + xy') = xy' - \frac{x''}{\lambda}, \\ (z' + xy')' = -\frac{x'}{\lambda}. \end{cases}$$

By integrating $(S_2)_3$, we obtain

$$z' + xy' = -\frac{x}{\lambda} + c$$

where c is a constant. Putting the last equation in $(S_2)_{1,2}$, we get

(5.11)
$$\bar{S}_2: \begin{cases} y' = -\frac{x^2}{\lambda} + xc, \\ x'' + 2x^3 - x - \lambda c(3x^2 - c\lambda x - 1) = 0. \end{cases}$$

Without loss of generality, we can assume that c = 0. This system \bar{S}_2 , *i.e.*, the equation $x'' + 2x^3 - x = 0$ involves Jacobi elliptic functions. So, we write the following proposition.

Proposition 4. The Killing magnetic curves in (H_3, g_2) corresponding to the Killing vector field $V_2 = e_1 + xe_3$ are solutions of the system of differential equations (5.10).

5.3. V_3 -magnetic curves. We have

(5.12)
$$V_3 \wedge \mathbf{t} = (yx' + (z' + xy'))e_1 - \lambda yy'e_2 + y'e_3.$$

From the equation $\nabla_{\mathbf{t}} \mathbf{t} = V_3 \wedge \mathbf{t}$, we get

(5.13)
$$S_3: \begin{cases} y'' - x'(z' + xy') = yx' + (z' + xy') \\ \frac{x''}{\lambda} + \lambda y'(z' + xy') = -\lambda yy', \\ (z' + xy')' = y'. \end{cases}$$

By integrating $(S_3)_3$, we deduce

$$z' + xy' = y + c,$$

where c is a constant. Putting the last equation in $(S_3)_{1,2}$, we have

(5.14)
$$\bar{S}_3: \begin{cases} x' = -\lambda^2 y^2 - \lambda^2 y c, \\ y'' - 2x'y - y - c(x'+1) = 0 \end{cases}$$

Without loss of generality, we can suppose that c = 0. Then, the system \bar{S}_3 , *i.e.*, the equation $y'' + 2\lambda^2 y^3 - y = 0$ has solutions which include Jacobi elliptic functions. Thus, we give the proposition below.

Proposition 5. The Killing magnetic curves in (H_3, g_2) corresponding to the Killing vector field $V_3 = e_2 - \lambda y e_3$ are solutions of the system of differential equations (5.13).

5.4. V_4 -magnetic curves. We write (5.15)

$$V_4 \wedge \mathbf{t} = (-\frac{1}{2}(x^2 + \lambda^2 y^2)\frac{x'}{\lambda} - \lambda y(z' + xy'))e_1 + ((x^2 + \lambda^2 y^2)\frac{y'}{2} - x(z' + xy'))e_2 - (\frac{xx'}{\lambda} + \lambda yy')e_3.$$

The equation $\nabla_{\mathbf{t}} \mathbf{t} = V_4 \wedge \mathbf{t}$ gives us

(5.16)
$$S_4: \begin{cases} y'' - x'(z'+xy') = -\frac{1}{2}(x^2+\lambda^2y^2)\frac{x'}{\lambda} - \lambda y(z'+xy'), \\ \frac{x''}{\lambda} + \lambda y'(z'+xy') = \frac{1}{2}(x^2+\lambda^2y^2)y' - x(z'+xy'), \\ (z'+xy')' = -\frac{xx'}{\lambda} - \lambda yy'. \end{cases}$$

By integrating $(S_4)_3$, we obtain

(5.17)
$$z' + xy' = -\frac{x^2}{2\lambda} - \frac{\lambda y^2}{2} + c$$

where c is a constant. Putting the last equation in $(S_4)_{1,2}$, we get

(5.18)
$$\bar{S}_4: \begin{cases} y'' + \frac{x'}{\lambda}(x^2 + \lambda^2 y^2) = \frac{1}{2}y(x^2 + \lambda^2 y^2) + c(-\lambda y + x'), \\ x'' - \lambda y'(x^2 + \lambda^2 y^2) = \frac{1}{2}x(x^2 + \lambda^2 y^2) - c(\lambda x + \lambda^2 y'). \end{cases}$$

It seems a true challenge to solve the system \bar{S}_4 in general case. However, we can find a special solution by considering $c = \lambda = 1$. In this case,

$$x(t) = \cos \frac{\sqrt{2}}{2}t, \ y(t) = \sin \frac{\sqrt{2}}{2}t$$

will be a solution for the system \bar{S}_4 . Using these relations in (5.17), we get

$$z(t) = \frac{2 - \sqrt{2}}{4}t - \frac{1}{4}\sin\sqrt{2}t + k_1$$

where k_1 is a constant. So, we find a solution as follows:

(5.19)
$$\begin{cases} x(t) = \cos \frac{\sqrt{2}}{2}t, \\ y(t) = \sin \frac{\sqrt{2}}{2}t, \\ z(t) = \frac{2-\sqrt{2}}{4}t - \frac{1}{4}\sin\sqrt{2}t + k_1. \end{cases}$$

Therefore, we can state the last propositon of the paper.

Proposition 6. The space curves given by parametric equations

$$\gamma(t) = \begin{pmatrix} x(t) = \cos\frac{\sqrt{2}}{2}t, \\ y(t) = \sin\frac{\sqrt{2}}{2}t, \\ z(t) = \frac{2-\sqrt{2}}{4}t - \frac{1}{4}\sin\sqrt{2}t + k_1 \end{pmatrix}$$

are V_4 -magnetic curves in (H_3, g_2) , where k_1 is an arbitrary constants.

Acknowledgement 1. The author would like to thank Professor Zlatko Erjavec and Professor Jun-ichi Inoguchi for their valuable suggestions.

References

- W. Batat and S. Rahmani, Isometries, Geodesics and Jacobi Fields of Lorentzian Heisenberg Group, Mediterranean Journal of Mathematics, 8 (2011), 411–430.
- [2] C. Bejan and S.L. Druta Romaniuc, Walker manifolds and Killing magnetic curves, Differential Geometry and Its Applications, 35 (2014), 106-116.
- [3] G. Calvaruso, M. I. Munteanu, A. Perrone, Killing magnetic curves in three-dimensional almost paracontact manifolds, Journal of Mathematical Analysis and Applications, 426 (1) (2015), 423-439.

10

- [4] K. Derkaoui and F.Hathout, Explicit formulas for Killing magnetic curves in Heisenberg group, International Journal of Geometric Methods in Modern Physics, (2021), https://doi.org/10.1142/S0219887821501358.
- [5] Z. Erjavec, On Killing magnetic curves in SL(2, ℝ) geometry, Reports on Mathematical Physics, 84 (3), (2019), 333-350.
- [6] Z. Erjavec and J. Inoguchi, Killing magnetic curves in Sol space, Mathematical Physics, Analysis and Geometry, 21 (15) (2018).
- [7] Z. Erjavec and J. Inoguchi, *Killing magnetic curves in almost cosymplectic Sol space*. Results in Mathematics. 75, 113, (2020).
- J-E. Lee, Slant curves and contact magnetic curves in Sasakian Lorentzian 3-manifolds. Symmetry 11, no. 6: 784, (2019), https://doi.org/10.3390/sym11060784.
- [9] M. I. Munteanu and A. Nistor, The classification of Killing magnetic curves in S² × ℝ, Journal of Geometry and Physics, 62 (2012), 170-182.
- [10] Z. Ozdemir, I. Gok, Y. Yayli and F. N. Ekmekci, Notes on magnetic curves in 3D semi-Riemannian manifolds, Turkish Journal of Mathematics, 39 (3) (2015), 412-426.
- [11] C. Ozgur, On magnetic curves in 3-dimensional Heisenberg group, Proceedings of the Institute of Mathematics and Mechanics (PIMM), National Academy of Sciences of Azerbaijan, 43 (2), (2017), 278-286.
- [12] S.L. Druta-Romaniuc, J. Inoguchi, M. I. Munteanu and A. I. Nistor, *Magnetic curves in cosymplectic manifolds*, Reports on Mathematical Physics, 78 (2016) 33-48.
- [13] S. L. Druta-Romaniuc, J. Inoguchi, M. I. Munteanu and A. I. Nistor, Magnetic curves in Sasakian manifolds, Journal of Nonlinear Mathematical Physics, 22 (3) (2015), 428–447.
- [14] S. L. Druta-Romaniuc and M.I. Munteanu, Magnetic curves corresponding to Killing magnetic fields in E³, Journal of Mathematical Physics, 52, (2011), 113506.
- [15] S. L. Druta-Romaniuc and M.I. Munteanu, Killing magnetic curves in a Minkowski 3-space, Nonlinear Analysis: Real World Applications, 14 (2013), 383-396.

(M. Altunbas) Department of Mathematics, Faculty of Sciences and Arts, Erzincan University, 25240 Erzurum, Turkey

Email address, M. Altunbas: maltunbas@erzincan.edu.tr