

ON COMPLEMENTED COPIES OF THE SPACE c_0 IN SPACES $C_p(X, E)$

CHRISTIAN BARGETZ, JERZY KĄKOL, AND DAMIAN SOBOTA

ABSTRACT. We study the question for which Tychonoff spaces X and locally convex spaces E the space $C_p(X, E)$ of continuous E -valued functions on X contains a complemented copy of the space $(c_0)_p = \{x \in \mathbb{R}^\omega : x(n) \rightarrow 0\}$, both endowed with the pointwise topology. We provide a positive answer for a vast class of spaces, extending classical theorems of Cembranos, Freniche, and Domański and Drewnowski, proved for the case of Banach and Fréchet spaces $C_k(X, E)$. For given infinite Tychonoff spaces X and Y , using the results of Henriksen and Woods and analysing relations between the space $C_p(X, C_p(Y))$ and the space $SC_p(X \times Y)$ of separately continuous real-valued functions on the product $X \times Y$, we show that $C_p(X, C_p(Y))$ contains a complemented copy of $(c_0)_p$ if and only if any of the spaces $C_p(X)$ and $C_p(Y)$ contains such a subspace. This approach applies also to show that, contrary to the case of the compact-open topology, for every infinite compact spaces X and Y the space $C_p(X \times Y)$ cannot be mapped onto the space $C_p(X, C_p(Y))$ by a continuous linear operator.

1. INTRODUCTION

For a topological space X and a locally convex space E , by $C_p(X, E)$ and $C_k(X, E)$ we denote the space of all continuous E -valued functions on X endowed with the pointwise topology (i.e. the topology inherited from the product E^X) and the compact-open topology, respectively. Similarly, if X is a compact space and E is a Banach space, by $C(X, E)$ we mean the Banach space of all continuous E -valued functions on X endowed with the sup-norm topology. If $E = \mathbb{R}$, then we simply write $C(X)$, $C_p(X)$ and $C_k(X)$ instead of $C(X, \mathbb{R})$, $C_p(X, \mathbb{R})$ and $C_k(X, \mathbb{R})$, respectively.

Recall the following result due to Cembranos ([10]) and Freniche ([15, Corollary 2.5]).

Theorem 1.1 (Cembranos, Freniche). *For an infinite compact space X and an infinite-dimensional Banach space E , the Banach space $C(X, E)$ contains a complemented copy of the Banach space c_0 .*

Theorem 1.1 was extended by Domański and Drewnowski ([11, Theorem 1, p. 260]) to Fréchet spaces of the form $C_k(X, E)$.

Theorem 1.2 (Domański–Drewnowski). *Let X be a Tychonoff space which contains an infinite compact subset and E a Fréchet space which is not Montel. Then the Fréchet space $C_k(X, E)$ contains a complemented copy of the space c_0 .*

Both of the theorems have important consequences. E.g., since for each Tychonoff space X and locally compact space Y , the spaces $C_k(X \times Y)$ and $C_k(X, C_k(Y))$ are linearly homeomorphic (in short, *isomorphic*), see [29, Corollary 2.5.7], Theorem 1.1 yields that for infinite compact spaces X and Y the Banach space $C(X \times Y)$ contains a complemented copy of c_0 . Similarly, if X is a Tychonoff space containing an infinite compact subset and Y is a locally compact and σ -compact Tychonoff space, then $C_k(Y)$ is an infinite-dimensional Fréchet space

which is not a Montel space ([18, Theorem 11.8.7]), and hence Theorem 1.2 implies that $C_k(X \times Y) \approx C_k(X, C_k(Y))$ contains a complemented copy of c_0 .

Let the space $(c_0)_p = \{x \in \mathbb{R}^\omega : x(n) \rightarrow 0\}$ be endowed with the pointwise topology inherited from \mathbb{R}^ω . The present paper deals with the following “delicate” question, motivated by Theorems 1.1 and 1.2:

(*) *When does a given space $C_p(X, E)$ contain a complemented copy of the space $(c_0)_p$?*

Theorems 1.1 and 1.2 were derived by their authors from the celebrated Josefson–Nissenzweig theorem or one of its many variants. Recall that the theorem states that every infinite-dimensional Banach space E admits a weakly* null sequence $(e'_n)_{n \in \omega}$ in the dual space E' of E such that $\|e'_n\| = 1$ for every $n \in \omega$. Following Banach and Gabrielyan [3] we say that a locally convex space E has the *Josefson–Nissenzweig Property* (JNP, in short) if the identity map $(E', \sigma(E', E)) \rightarrow (E', \beta^*(E', E))$ on the topological dual E' of E is not sequentially continuous. The Josefson–Nissenzweig theorem implies that every infinite-dimensional Banach space has the JNP, see Section 4 for more details concerning the property.

Following the argument of Domański and Drewnowski, we show in Theorem 6.2 that for a Tychonoff space X containing an infinite compact subset and a barrelled locally convex space E with the Josefson–Nissenzweig property, the space $C_p(X, E)$ contains a complemented copy of $(c_0)_p$. This not only generalises Theorem 1.1 (see Corollary 6.5) and provides a C_p -variant of Theorem 1.2 (Corollary 6.4), but also has several applications. For example, we get that if X is an infinite Tychonoff space and Y is an infinite compact space, then the space $C_p(X, C(Y))$ contains a complemented copy of \mathbb{R}^ω or a complemented copy of $(c_0)_p$ (Corollary 6.6; see also Corollary 6.7 for a $C_k(X, C_k(Y))$ -version). We immediately get from this that the space $C_p(X, C(Y))$ is isomorphic to $C_p(X, C(Y)) \times \mathbb{R}$ (Corollary 6.9), which extends the number of known cases of locally convex spaces E for which $E \approx E \times \mathbb{R}$, see the remark following Corollary 6.9 and concerning a problem of Arkhangel'ski.

We also investigate the case of spaces of the form $C_p(X, E_w)$, where E_w means a locally convex space E endowed with its weak topology $w = \sigma(E, E')$. Using again a variant of the Josefson–Nissenzweig theorem due to Domański and Drewnowski (see Lemma 7.1), we claim in Theorem 7.2 that for every Tychonoff space X containing an infinite compact subset and locally convex space E containing a copy of the Banach space ℓ_1 , the space $C_p(X, E_w)$ contains a complemented copy of $(c_0)_p$. As a corollary we get that if X is a Tychonoff space which contains an infinite compact subset and Y is an infinite Tychonoff space containing a non-scattered compact subset, then also $C_p(X, C_k(Y)_w)$ contains a complemented copy of $(c_0)_p$, see Corollary 7.3. This applies to show that, e.g., there is no continuous linear surjection from $C_p(\beta\omega, C_p(\beta\omega))$ onto $C_p(\beta\omega, C(\beta\omega)_w)$, as well as that for every infinite compact space X the spaces $C_p(X)$ and $C(X)_w$ are not isomorphic (which is a special case of Krupski–Marciszewski's [26, Corollary 3.2], cf. Remark 3.10).

In the case of spaces of the form $C_p(X, C_p(Y))$ question (*) is fully answered. To achieve this, we first show that for every Tychonoff spaces X and Y the space $C_p(X, C_p(Y))$ is isomorphic to the space $SC_p(X \times Y)$ of separately continuous real-valued functions on the product $X \times Y$ endowed with the pointwise topology (Theorem 3.1), which, on the other hand, by Henriksen and Woods [16, Proposition 2.1], is equal to the space $C_p(X \times Y, \sigma)$, where σ denotes the weak topology on the product $X \times Y$ induced by all separately continuous real-valued functions. The results of [16] allow us then to obtain Theorem 4.2 stating that for infinite X and Y the space $C_p(X \times Y, \sigma)$ has the JNP if and only if $C_p(X)$ has the JNP or $C_p(Y)$ has

the JNP. Finally, using the characterisation of those spaces $C_p(Z)$ which have the JNP, due to Banach, Kąkol and Śliwa (Theorem 5.1), we get in Theorem 5.2 that $C_p(X, C_p(Y))$ contains a complemented copy of $(c_0)_p$ if and only if any of the spaces $C_p(X)$ and $C_p(Y)$ contains such a copy.

As a by-product to the above arguments we obtain that the corresponding C_p -variant of the above mentioned [29, Corollary 2.5.7] (concerning isomorphisms between spaces $C_k(X \times Y)$ and $C_k(X, C_k(Y))$) fails, that is, in Corollary 3.8 we prove that for every infinite compact space X and every infinite pseudocompact space Y the space $C_p(X \times Y)$ cannot be linearly mapped onto the space $C_p(X, C_p(Y))$, which, in particular, implies that these function spaces cannot be isomorphic.

The plan of the paper is as follows. The next section contains most of the basic definitions and symbols which we frequently use in the paper. In Section 3 we investigate relations between spaces of the form $C_p(X \times Y)$ and $C_p(X, C_p(Y))$. Section 4 is devoted to the Josefson–Nissenzweig property of $C_p(X, C_p(Y))$ -spaces. In Sections 5, 6, and 7 we study the question of when the space $(c_0)_p$ is complemented in spaces of the form $C_p(X, C_p(Y))$, $C_p(X, E)$, and $C_p(X, E_w)$, respectively.

2. PRELIMINARIES

The cardinality of a set X is denoted by $|X|$. By ω we denote the first infinite cardinal number, i.e., the cardinality of the space of natural numbers \mathbb{N} . Usually we identify ω with \mathbb{N} , so ω is an infinite countable discrete topological space, and thus topological notions like the Čech–Stone compactification $\beta\omega$ of ω make sense. We denote the remainder $\beta\omega \setminus \omega$ of this compactification by ω^* .

We assume that all topological spaces we consider are **Tychonoff**, that is, completely regular and Hausdorff. If (X, τ) is a topological space (with the topology τ) and $Y \subseteq X$, then by $\tau \upharpoonright Y$ we mean the restriction of τ on Y , that is, $\tau \upharpoonright Y = \{U \cap Y : U \in \tau\}$.

We also assume that Fréchet spaces are locally convex. If E is a locally convex space, by E' we denote its topological dual. For a Tychonoff space X we write $L_p(X) = C_p(X)'$. Recall that each $\varphi \in L_p(X)$ may be uniquely represented as a finite linear combination of point-measures on X , i.e. $\varphi = \sum_{x \in X} \alpha_x^\varphi \delta_x$, where $\alpha_x^\varphi \in \mathbb{R}$ and only for finitely many $x \in X$ we have $\alpha_x^\varphi \neq 0$. If $A \subseteq X$, then we set $\varphi \upharpoonright A = \sum_{x \in A} \alpha_x^\varphi \delta_x$. We put $\text{supp}(\varphi) = \{x \in X : \alpha_x^\varphi \neq 0\}$, and define the norm $\|\varphi\|$ of φ by the formula

$$\|\varphi\| = \sum_{x \in \text{supp}(\varphi)} |\alpha_x^\varphi|.$$

For a locally convex space E , E_w denotes the pair $(E, \sigma(E, E'))$, i.e. the space E endowed with its weak topology. It is easy to see that for every Tychonoff space X we have $C_p(X) = (C_p(X))_w$.

Note that if \mathcal{P} is a family of semi-norms on a locally convex space E inducing the topology of E , then the pointwise topology of $C_p(X, E)$ is induced by the family \mathcal{Q} of semi-norms given for every $f \in C_p(X, E)$ by the formula:

$$q_{x_1, \dots, x_n; p}(f) = \max_{i=1, \dots, n} p(f(x_i)),$$

where $p \in \mathcal{P}$ and $x_1, \dots, x_k \in X$; see [35, Section I.2]. In particular, if $E = C_p(Y)$ for Tychonoff Y , then the topology on E is induced by the semi-norms:

$$p_{y_1, \dots, y_m}(f) = \max_{i=1, \dots, m} |f(y_i)|,$$

with $y_1, \dots, y_m \in Y$, whence the topology on $C_p(X, E)$ is given by the semi-norms of the form:

$$q_{x_1, \dots, x_n; y_1, \dots, y_m}(f) = \max_{i=1, \dots, n} \max_{j=1, \dots, m} |f(x_i)(y_j)|,$$

where $x_1, \dots, x_n \in X$ and $y_1, \dots, y_m \in Y$.

If E and F are both locally convex, we write $E \approx F$ if E and F are *isomorphic* (i.e. linearly homeomorphic). We also say that E *contains a complemented copy of F* (or that F *is complemented in E*) if there are closed linear subspaces E_1 and E_2 of E such that E is a direct algebraic sum of E_1 and E_2 (i.e. $E = E_1 \oplus E_2$ as a vector space and $E_1 \cap E_2 = \{0\}$), E_1 is isomorphic to F , and the natural projection from E onto E_1 is continuous.

By S we denote the non-trivial convergent sequence, i.e. the space $S = \{0\} \cup \{1/n : n \in \omega, n > 0\}$ the usual topology inherited from the real line. By c_0 we mean the usual Banach space of all real-valued sequences convergent to 0, i.e. $c_0 = \{x \in \mathbb{R}^\omega : x(n) \rightarrow 0\}$ endowed with the supremum norm. The symbol $(c_0)_p$ denotes the space c_0 with the pointwise topology inherited from the product \mathbb{R}^ω . Note that $(c_0)_p \approx C_p(S)$ (see [38, p.397]).

All other necessary and possibly non-standard notions will be defined where we need them.

3. $C_p(X \times Y)$ VERSUS $C_p(X, C_p(Y))$

It is well known that if X is a Tychonoff space and Y is a locally compact Hausdorff space, the spaces $C_k(X \times Y)$ and $C_k(X, C_k(Y))$ are isomorphic (see [29, Corollary 2.5] or [12, Theorems 3.4.8–3.4.10]). Motivated by this result, we will analyse the relation between the spaces $C_p(X \times Y)$ and $C_p(X, C_p(Y))$ for general Tychonoff spaces X and Y .

We start with the following standard definition. Let X and Y be Tychonoff spaces. Recall that a function $f: X \times Y \rightarrow \mathbb{R}$ is *separately continuous* if the functions $Y \ni y \mapsto f(x_0, y)$ and $X \ni x \mapsto f(x, y_0)$ are continuous for every $x_0 \in X$ and $y_0 \in Y$. Of course, if a function is continuous, then it is separately continuous, but the converse may not hold. A standard example of a separately continuous but not continuous function is the function $s: [0, 1]^2 \rightarrow \mathbb{R}$ defined as follows:

$$s(x, y) = \begin{cases} \frac{2xy}{x^4 + y^4}, & \text{if } (x, y) \neq (0, 0) \\ 0, & \text{if } (x, y) = (0, 0). \end{cases}$$

s is not continuous at $(0, 0)$ as well as it is not bounded, since $s(1/n, 1/n) = n^2$ for every $n > 0$. Given two infinite compact spaces X and Y , using the function s and appropriate mappings from X and Y into $[0, 1]$, one can easily construct a separately continuous function on the product $X \times Y$ which is not continuous. For more information on separately continuous functions we refer the reader to [30, 33, 34, 16, 5, 14], and the references therein.

Fix two Tychonoff spaces X and Y . By $SC(X \times Y)$ we denote the space of all real-valued separately continuous functions on $X \times Y$ and by $SC_p(X \times Y)$ we mean $SC(X \times Y)$ endowed with the pointwise topology. Note that $C(X \times Y)$ is a linear subspace of $SC(X \times Y)$. Since $C_p(X \times Y)$ is dense in the product $\mathbb{R}^{X \times Y}$, it is also dense in $SC_p(X \times Y)$.

We consider the mapping $A: SC_p(X \times Y) \rightarrow C_p(X, C_p(Y))$ defined as follows: for every $f \in SC_p(X \times Y)$, $A(f)$ is the only mapping in $C_p(X, C_p(Y))$ such that $A(f)(x)(y) = f(x, y)$

for every $x \in X$ and $y \in Y$, that is,

$$SC_p(X \times Y) \ni f \xrightarrow{A} [x \mapsto [y \mapsto f(x, y)]] \in C_p(X, C_p(Y)).$$

Theorem 3.1. *The mapping A is a well-defined isomorphism between the spaces $SC_p(X \times Y)$ and $C_p(X, C_p(Y))$.*

Proof. We first need to show that A is well-defined, that is, it takes values in the space $C_p(X, C_p(Y))$. Fix $f \in SC_p(X \times Y)$ and $x \in X$. Since the function $f(x, \cdot)$ is continuous on Y , we have $A(f)(x) \in C_p(Y)$, so we only need to show that $A(f)$ is continuous at x . Let $y_1, \dots, y_k \in Y$ and $\varepsilon > 0$ be given. Since $f(\cdot, y_i)$ is continuous for every $i = 1, \dots, k$, there is an open neighborhood U of x in X such that

$$|f(x, y_i) - f(x', y_i)| < \varepsilon$$

for every $x' \in U$ and $i = 1, \dots, k$, or, in other words,

$$|A(f)(x)(y_i) - A(f)(x')(y_i)| < \varepsilon,$$

which implies that

$$A(f)[U] \subseteq \{g \in C_p(Y) : |A(f)(x)(y_i) - g(y_i)| < \varepsilon, \forall i = 1, \dots, k\}.$$

This shows that $A(f): X \rightarrow C_p(Y)$ is continuous at x and hence that A is well-defined.

It is clear that A is a linear mapping. A is injective, since $A(f) = A(g)$ implies that $f(x, y) = A(f)(x)(y) = A(g)(x)(y) = g(x, y)$ for all $(x, y) \in X \times Y$, i.e. $f = g$. Let now $g \in C_p(X, C_p(Y))$ and put $f(x, y) = g(x)(y)$ for every $(x, y) \in X \times Y$. It is immediate that $f(x, \cdot)$ is continuous on Y for every $x \in X$. Fix $y \in Y$; we need to show that $f(\cdot, y)$ is continuous on X . Let $x \in X$ and $\varepsilon > 0$. Since g is continuous at x , there is an open neighborhood U of x in X such that

$$g[U] \subseteq \{h \in C_p(Y) : |g(x)(y) - h(y)| < \varepsilon\}.$$

In particular, for every $x' \in U$ we have:

$$|f(x, y) - f(x', y)| = |g(x)(y) - g(x')(y)| < \varepsilon,$$

which proves that $f(\cdot, y)$ is continuous at x . It follows that A is surjective, and hence that it is an algebraic isomorphism of the spaces $SC_p(X \times Y)$ and $C_p(X, C_p(Y))$. It remains to be proved that A is a topological isomorphism.

Recall that the topology on $C_p(X, C_p(Y))$ is generated by the semi-norms

$$q_{x_1, \dots, x_n; y_1, \dots, y_m}(g) = \max_{\substack{i=1, \dots, n \\ j=1, \dots, m}} |g(x_i)(y_j)|,$$

where $x_1, \dots, x_n \in X$ and $y_1, \dots, y_m \in Y$, and hence for every $f \in SC_p(X \times Y)$ we have

$$\begin{aligned} q_{x_1, \dots, x_n; y_1, \dots, y_m}(A(f)) &= \max_{\substack{i=1, \dots, n \\ j=1, \dots, m}} |(A(f))(x_i)(y_j)| = \max_{\substack{i=1, \dots, n \\ j=1, \dots, m}} |f(x_i, y_j)| \\ &= p_{(x_1, y_1), \dots, (x_n, y_m)}(f), \end{aligned}$$

which shows that A is continuous. Now, since

$$p_{(x_1, y_1), \dots, (x_n, y_n)}(f) \leq q_{x_1, \dots, x_n; y_1, \dots, y_n}(A(f))$$

for every $f \in SC_p(X \times Y)$, $n \in \omega$, $x_1, \dots, x_n \in X$ and $y_1, \dots, y_n \in Y$, we get that A^{-1} is also continuous. It follows that A is a linear homeomorphism. \square

Since $C_p(X \times Y)$ is dense in $SC_p(X \times Y)$, we immediately get the following

Corollary 3.2. *For every Tychonoff spaces X and Y , the space $C_p(X \times Y)$ embeds as a dense linear subspace of the space $C_p(X, C_p(Y))$.*

If X and Y are infinite compact, then it follows from our discussion at the beginning of this section that $C_p(X \times Y) \neq SC_p(X \times Y)$ and hence that $A[C_p(X \times Y)] \neq C_p(X, C_p(Y))$. We will now show that it is not accidental, as $C_p(X \times Y)$ in the case of infinite compact spaces X and Y cannot be linearly mapped onto $C_p(X, C_p(Y))$. However, in order to prove this, we need to introduce another topology on the product $X \times Y$.

Let X and Y be Tychonoff spaces. By σ let us denote the weak topology on $X \times Y$ generated by the family of all separately continuous functions. This topology was investigated by Henriksen and Woods [16], who proved among others that:

- $(X \times Y, \sigma)$ is a Tychonoff space ([16, Theorem 4.8]);
- $(X \times Y, \sigma)$ is not pseudocompact provided that both X and Y are infinite ([16, Corollary 5.9.b]);
- $C(X \times Y, \sigma) = SC(X \times Y)$, i.e. every continuous function on $(X \times Y, \sigma)$ is separately continuous on $X \times Y$ and *vice versa* ([16, Proposition 2.1]).

By the last listed result and Theorem 3.1 we immediately get the following theorem, which allows us to treat spaces of the form $C_p(X, C_p(Y))$ the same way as spaces of the form $C_p(X \times Y)$.

Theorem 3.3. *Let X and Y be Tychonoff spaces. Then, the space $C_p(X \times Y, \sigma)$ is isomorphic to the space $C_p(X, C_p(Y))$.*

Using the above result, we will now prove that $C_p(X, C_p(Y)) \approx C_p(Y, C_p(X))$ for every Tychonoff spaces X and Y .

Lemma 3.4. *For every Tychonoff spaces X and Y the spaces $(X \times Y, \sigma)$ and $(Y \times X, \sigma)$ are homeomorphic.*

Proof. Let $\varphi: X \times Y \rightarrow Y \times X$ be given by the formula $\varphi(x, y) = (y, x)$ for every $(x, y) \in X \times Y$. Clearly, φ is a bijection. We first show that φ is σ - σ -continuous. Note that for every $f \in \mathbb{R}^{Y \times X}$ we have:

$$(*) \quad f \in SC(Y \times X) \text{ if and only if } f \circ \varphi \in SC(X \times Y).$$

Fix $(x_0, y_0) \in X \times Y$ and let $f_1, \dots, f_n \in SC(Y \times X)$ and $\varepsilon > 0$. The set

$$V(\varphi(x_0, y_0); f_1, \dots, f_n; \varepsilon) = \{(v, u) \in Y \times X: |f_i(\varphi(x_0, y_0)) - f_i(v, u)| < \varepsilon, i = 1, \dots, n\},$$

is an open basic neighborhood of $\varphi(x_0, y_0)$ in $(Y \times X, \sigma)$. By (*), the set

$$U((x_0, y_0); f_1 \circ \varphi, \dots, f_n \circ \varphi; \varepsilon) = \{(u, v) \in X \times Y: |(f_i \circ \varphi)(x_0, y_0) - (f_i \circ \varphi)(u, v)| < \varepsilon, i = 1, \dots, n\}$$

is an open basic neighborhood of (x_0, y_0) in $(X \times Y, \sigma)$, and it holds:

$$\varphi[U((x_0, y_0); f_1 \circ \varphi, \dots, f_n \circ \varphi; \varepsilon)] = V(\varphi(x_0, y_0); f_1, \dots, f_n; \varepsilon),$$

which proves that φ is continuous.

Using the fact that for every $g \in \mathbb{R}^{X \times Y}$ we have

$$g \in SC(X \times Y) \text{ if and only if } g \circ \varphi^{-1} \in SC(Y \times X),$$

we prove similarly that φ^{-1} is σ - σ -continuous. \square

Corollary 3.5. *For every Tychonoff spaces X and Y the spaces $C_p(X \times Y, \sigma)$ and $C_p(Y \times X, \sigma)$ are isomorphic.*

By Theorem 3.3, we immediately obtain the following

Theorem 3.6. *For every Tychonoff spaces X and Y the spaces $C_p(X, C_p(Y))$ and $C_p(Y, C_p(X))$ are isomorphic.*

Recall that every continuous linear mapping between two topological vector spaces is uniformly continuous ([37, Ex. 134.iv]). Also, if W and Z are such Tychonoff spaces that W is pseudocompact and there exists a uniformly continuous surjection from $C_p(W)$ onto $C_p(Z)$, then Z is pseudocompact ([37, Ex. 193]). Since the space $(X \times Y, \sigma)$ is never pseudocompact for infinite X and Y , Theorem 3.3 implies the following result.

Theorem 3.7. *Let X and Y be infinite pseudocompact spaces such that the product $X \times Y$ is pseudocompact. Then, the space $C_p(X \times Y)$ cannot be mapped onto the space $C_p(X, C_p(Y))$ by a continuous linear map. In particular, $C_p(X \times Y)$ is not isomorphic to $C_p(X, C_p(Y))$.*

The following corollary follows from the fact that the product of a compact space and a pseudocompact space is pseudocompact.

Corollary 3.8. *If X is an infinite compact space and Y an infinite pseudocompact space, then the space $C_p(X \times Y)$ cannot be mapped onto the space $C_p(X, C_p(Y))$ by a continuous linear map.*

Corollary 3.9. *If X and Y are infinite compact spaces, then $C_p(X \times Y)$ is not isomorphic to $C_p(X, C_p(Y))$.*

We cannot drop the compactness assumption in Corollary 3.9. Indeed, if X and Y are infinite and Y is discrete, then the canonical mapping $A: C_p(X \times Y) \rightarrow C_p(X, C_p(Y))$ defined above is easily seen to be an isomorphism. Also, [16, Corollary 6.15] states that if X is a (locally) separable Tychonoff space and Y is a P-space, then $C(X \times Y) = SC(X \times Y)$ and hence σ coincides with the product topology on $X \times Y$ —it follows by Theorem 3.3 that the spaces $C_p(X \times Y)$ and $C_p(X, C_p(Y))$ are isomorphic. In particular, if X is a metric compact space or $X = \beta\omega$ and Y a P-space, then $C_p(X \times Y) \approx C_p(X, C_p(Y))$.

Remark 3.10. Krupski and Marciszewski [26, Corollary 3.2] proved that for all infinite compact spaces X and Y the locally convex spaces $C_p(X)$ and $C(Y)_w$ are never isomorphic. It follows that $C_p(X \times Y)$ and $C(X \times Y)_w$ are not isomorphic. Since $C(X \times Y)_w$ is isomorphic to $C(X, C(Y))_w$, we get that $C_p(X \times Y)$ is not isomorphic to $C(X, C(Y))_w$. Thus, Corollary 3.9 may be thought of as a C_p -counterpart of the result of Krupski and Marciszewski. This observation leads to the following relevant question, some issues of which we will study in Section 7.

Problem 3.11. *Let X and Y be infinite compact spaces. Are the spaces $C_p(X \times Y)$ and $C_p(X, C(Y)_w)$ isomorphic?*

The next two examples may serve as an illustration to Corollary 3.9.

Example 3.12. Let $X = \beta\omega \setminus \omega$. In [16, Example 5.7] it was proved that $|SC(X \times X)| > |C(X \times X)|$, hence we get that $C_p(X, C_p(X))$ has cardinality strictly greater than $C_p(X \times X)$ and thus the spaces cannot be isomorphic.

Example 3.13. Let $X = [0, 1]^\omega$ be endowed with the product topology. From Corollary 3.9 it follows that $C_p(X \times X)$ is not isomorphic to $C_p(X, C_p(X))$. However, [37, Ex. 300] implies that $C_p(X \times X)$ is isomorphic to $C_p(X)$, and hence $C_p(X \times X)$ is isomorphic to a complemented subspace of $C_p(X, C_p(X))$ (by [35, Theorem I.4.4]). A similar argument may be provided for any uncountable metrizable totally-disconnected compact space X , e.g. for the Cantor space (see [37, Ex. 297]).

Problem 3.14. *For which infinite compact spaces X , is the space $C_p(X \times X)$ isomorphic to a complemented subspace of the space $C_p(X, C_p(X))$? In particular, does there exist an infinite compact space X such that $C_p(X)$ is not isomorphic to $C_p(X \times X)$ and $C_p(X \times X)$ is isomorphic to a complemented subspace of $C_p(X, C_p(X))$?*

In Corollary 5.5 we describe a class of infinite compact spaces X such that $C_p(X \times X)$ is not complemented in $C_p(X, C_p(X))$. Note that if X is an infinite compact space such that $C_p(X \times X)$ is complemented in $C_p(X, C_p(X))$, then the latter may be linearly mapped onto the former—a phenomenon which by Corollary 3.8 cannot occur in the reverse direction.

We finish the section with the following observation.

Proposition 3.15. *Let E be a normed space and X a Tychonoff space. Then, the original pointwise topology of $C_p(X, E)$ coincides with the weak topology of $C_p(X, E)$ if and only if E is finite-dimensional space.*

Proof. If E is finite-dimensional, say E is isomorphic to \mathbb{R}^n for some $n \in \omega$, then we have:

$$C_p(X, E) \approx C_p(X, \mathbb{R}^n) \approx C_p(X)^n.$$

Hence, the pointwise topology and weak topology coincide in $C_p(X, E)$. Conversely, if both of the topologies coincide, then E is finite-dimensional, since E is isomorphic to a complemented subspace of $C_p(X, E)$ by [35, Theorem I.4.4]. \square

Corollary 3.16. *If X is a Tychonoff space and Y is an infinite compact space, then the spaces $C_p(X \times Y)$ and $C_p(X, C(Y))$ are not isomorphic.*

4. THE JOSEFSON–NISSENZWEIG PROPERTY

Following Banach and Gabrielyan [3], we say that a locally convex space E has *the Josefson–Nissenzweig property*, or *the JNP* in short, if the identity map

$$(E', \sigma(E', E)) \rightarrow (E', \beta^*(E', E))$$

is not sequentially continuous, i.e. if the dual space E' contains a weak* null sequence which does not converge for the topology $\beta^*(E', E)$. The topology $\beta^*(E', E)$ can be described as the topology of uniform convergence on so-called *barrel-bounded sets*, i.e. on sets which have the property that they are absorbed by all barrels—see [3, p. 3] for more details. Note that the JNP is an isomorphic property, since the transpose of an isomorphism is also an isomorphism for the above topologies—see e.g. Corollary 8.6.6 in [18, p. 161]. By the famous Josefson–Nissenzweig Theorem, see e.g. [19, 32, 8], every infinite-dimensional Banach space has the JNP. On the other hand, no Montel space enjoys the JNP. Indeed: First, since Montel spaces are barrelled, the topologies $\beta^*(E', E)$ and $\beta(E', E)$ coincide. Second, the strong dual of a Montel space is again Montel and hence on bounded sets the topologies $\sigma(E', E)$ and $\beta(E', E)$ coincide. In [6], Bonet, Lindström and Valdivia showed that for Fréchet spaces this is already a characterisation of the JNP, i.e. a Fréchet space has the JNP if and only if it is not Montel; see also [27] for a similar characterisation of non-Schwarz Fréchet spaces.

In the case of $C_p(X)$ -spaces, Banach and Gabrielyan proved the following characterisation of the Josefson–Nissenzweig property.

Theorem 4.1 ([3, Theorem 3.4]). *For every Tychonoff space X , the space $C_p(X)$ has the JNP if and only if there exists a sequence of functionals $(\varphi_n)_{n \in \omega}$ in the dual space $L_p(X)$ of $C_p(X)$ such that $\varphi_n(f) \rightarrow 0$ for every $f \in C(X)$ and $\|\varphi_n\| = 1$ for every $n \in \omega$.*

If a sequence $(\varphi_n)_{n \in \omega}$ of functionals on a given space $C_p(X)$ has the properties as in the above theorem, then it will be called a *JN-sequence* on X . The existence of JN-sequences on arbitrary spaces was intensively studied in [4], [21] and [22], where several classes of Tychonoff spaces were recognized to (not) admit such sequences (e.g. it appears that for every infinite compact spaces X and Y their product admits a JN-sequence, see [21, Theorem 11.3]). Note that if X is pseudocompact and $C_p(X)$ has the JNP, then every JN-sequence on X satisfies already the conclusion of the Josefson–Nissenzweig theorem for the Banach space $C(X)$ (by the virtue of the Riesz representation of continuous functionals on $C(X)$).

The following theorem provides a simple criterion for a space $C_p(X \times Y, \sigma)$ to have the JNP.

Theorem 4.2. *Let X and Y be infinite Tychonoff spaces. Then, the space $C_p(X \times Y, \sigma)$ has the JNP if and only if the space $C_p(X)$ has the JNP or the space $C_p(Y)$ has the JNP.*

Proof. Assume first that $C_p(X \times Y, \sigma)$ has the JNP and let $(\varphi_n)_{n \in \omega}$ be a JN-sequence on $(X \times Y, \sigma)$. Put:

$$\mathcal{S} = \bigcup_{n \in \omega} \text{supp}(\varphi_n).$$

By [21, Lemma 4.11] (cf. also [22, Proposition 4.1]), every continuous function $f \in C(X \times Y, \sigma)$ is bounded on \mathcal{S} , that is, $f[\mathcal{S}]$ is a bounded subset of \mathbb{R} . By [16, Corollary 5.9.a], \mathcal{S} must be contained in finitely many vertical sections or finitely many horizontal sections of the product $X \times Y$.

Let us assume that the former happens, i.e. there are $x_1, \dots, x_m \in X$ such that $\mathcal{S} \subseteq \bigcup_{i=1}^m \{x_i\} \times Y$. Since $\|\varphi_n\| = 1$ for every $n \in \omega$, there are $i_0 \in \{1, \dots, m\}$ and a subsequence $(\varphi_{n_k})_{k \in \omega}$ such that $\|\varphi_{n_k} \upharpoonright \{x_{i_0}\} \times Y\| \geq 1/m$ for every $k \in \omega$. We claim that the sequence $(\psi_k)_{k \in \omega}$ defined for every $k \in \omega$ by the formula

$$\psi_k = (\varphi_{n_k} \upharpoonright \{x_{i_0}\} \times Y) / \|\varphi_{n_k} \upharpoonright \{x_{i_0}\} \times Y\|$$

is a JN-sequence on $\{x_{i_0}\} \times Y$. Obviously, $\|\psi_k\| = 1$ for every $k \in \omega$. Let $f \in C_p(\{x_{i_0}\} \times Y)$. Since X is Tychonoff, there is a function $g \in C_p(X)$ such that $g(x_{i_0}) = 1$ and $g(x_i) = 0$ for every $i \in \{1, \dots, m\} \setminus \{i_0\}$. For every $(x, y) \in X \times Y$ we put:

$$F(x, y) = g(x) \cdot f(x_{i_0}, y).$$

It follows that $F \in SC(X \times Y) = C(X \times Y, \sigma)$, $F(x_{i_0}, \cdot) \equiv f(x_{i_0}, \cdot)$, and $F(x_i, \cdot) \equiv 0$ for every $i \in \{1, \dots, m\} \setminus \{i_0\}$. Since $(\varphi_{n_k})_{k \in \omega}$ is also a JN-sequence on $(X \times Y, \sigma)$ and $\mathcal{S} \subseteq \bigcup_{i=1}^m \{x_i\} \times Y$, we get that

$$|\psi_k(f)| \leq |\varphi_{n_k}(F)| \cdot m \rightarrow 0$$

as $k \rightarrow \infty$. This proves that $(\psi_k)_{k \in \omega}$ is a JN-sequence on $\{x_{i_0}\} \times Y$ —it follows easily that $C_p(Y)$ has the JNP, too.

If \mathcal{S} is contained in finitely many horizontal sections of the product $X \times Y$, then we proceed similarly and prove that $C_p(X)$ has the JNP.

Let us assume now that, conversely, $C_p(X)$ has the JNP, that is, there is a JN-sequence (φ_n) on X . For each $n \in \omega$ write:

$$\varphi_n = \sum_{i=1}^{k_n} \alpha_i^n \delta_{x_i^n}.$$

Fix $y \in Y$ and define the functional ψ_n on $C_p(X \times Y, \sigma)$ by the similar formula:

$$\psi_n = \sum_{i=1}^{k_n} \alpha_i^n \delta_{(x_i^n, y)}.$$

We claim that $(\psi_n)_{n \in \omega}$ is a JN-sequence on $(X \times Y, \sigma)$. Of course, $\|\psi_n\| = 1$ for every $n \in \omega$. Let $f \in C_p(X \times Y, \sigma) = SC_p(X \times Y)$. Then, the function $F: X \rightarrow \mathbb{R}$ defined for every $x \in X$ by $F(x) = f(x, y)$ is continuous on X and thus $\lim_{n \rightarrow \infty} \varphi_n(F) = 0$. Since for every $n \in \omega$ we have $\psi_n(f) = \varphi_n(F)$, we get that $\lim_{n \rightarrow \infty} \psi_n(f) = 0$, too. It follows that the space $C_p(X \times Y, \sigma)$ has the JNP.

We prove similarly that if $C_p(Y)$ has the JNP, then $C_p(X \times Y, \sigma)$ has the JNP. \square

The first part of the proof suggest the following corollary.

Corollary 4.3. *Let X and Y be infinite Tychonoff spaces. Let $(\varphi_n)_{n \in \omega}$ be a sequence in $L_p(X \times Y, \sigma)$ such that $\|\varphi_n\| = 1$ for every $n \in \omega$. Set $\mathcal{S} = \bigcup_{n \in \omega} \text{supp}(\varphi_n)$. If \mathcal{S} is contained in neither finitely many vertical sections nor finitely many horizontal sections, then there exists $f \in SC(X \times Y)$ such that $\limsup_{n \rightarrow \infty} |\varphi_n(f)| > 0$.*

We will now study the Josefson–Nissenzweig property of the spaces $C_p(X, E)$ for X Tychonoff and E locally convex. Note that by [13, Proposition 19] the topological dual space of $C_p(X, E)$ is algebraically isomorphic to the space $L_p(X) \otimes E'$, so each functional $\varphi \in C_p(X, E)'$ may be represented as a finite sum of tensor products

$$\sum_{i,j} \alpha_{i,j} (\delta_{x_i} \otimes e'_j),$$

where $x_i \in X$ and $e'_j \in E'$. Hence, if $E = C_p(Y)$ for some Tychonoff space Y , then each $\varphi \in C_p(X, E)'$ may be written as a finite sum

$$\sum_{i,j} \alpha_{i,j} (\delta_{x_i} \otimes \delta_{y_j}).$$

Let us define the norm of φ (in $L_p(X) \otimes L_p(Y)$) in the following natural way:

$$\|\varphi\| = \sum_{i,j} |\alpha_{i,j}|.$$

Let X and Y be Tychonoff spaces and assume that $C_p(X \times Y, \sigma)$ has the JNP, that is, there is a sequence $(\varphi_n)_{n \in \omega}$ in $L_p(X \times Y, \sigma)$ such that $\varphi_n(f) \rightarrow 0$ for every $f \in C_p(X \times Y, \sigma)$ and $\|\varphi_n\| = 1$ for every $n \in \omega$. For each $n \in \omega$ write:

$$\varphi_n = \sum_{i=1}^{k_n} \alpha_i^n \delta_{(x_i^n, y_i^n)}.$$

By Theorem 3.1, the operator $A: C_p(X \times Y, \sigma) \rightarrow C_p(X, C_p(Y))$ is an isomorphism, so the sequence $(\psi_n)_{n \in \omega}$ given for every $n \in \omega$ by the formula:

$$\psi_n = A'(\varphi_i) = \sum_{i=1}^{k_n} \alpha_i^n ((\delta_{x_i^n} \otimes \delta_{y_i^n}) \circ A),$$

where A' is the transpose of A , is weak* convergent to 0 (see [25, Theorem 21.5]). Obviously, $\|\psi_n\| = 1$ for every $n \in \omega$.

Let us now assume that, conversely, the space $C_p(X, C_p(Y))$ admits a weak* null sequence $(\psi_n)_{n \in \omega}$ in $L_p(X) \otimes L_p(Y)$ such that $\|\psi_n\| = 1$ for every $n \in \omega$. By a very similar argument as above (this time applied to the transpose of A^{-1}), one may obtain a JN-sequence on the space $(X \times Y, \sigma)$. This way we obtain the following

Proposition 4.4. *For every Tychonoff spaces X and Y the space $C_p(X \times Y, \sigma)$ has the JNP if and only if there exists a sequence $(\psi_n)_{n \in \omega}$ in the dual $L_p(X) \otimes L_p(Y)$ which is weakly* convergent to 0 and such that $\|\psi_n\| = 1$ for every $n \in \omega$.*

Since the Josefson–Nissenzweig property for locally convex spaces is preserved under topological isomorphisms, Proposition 4.4 and Theorem 4.1 yield the following characterisation of the property for $C_p(X, C_p(Y))$ -spaces.

Proposition 4.5. *For every Tychonoff space X and Y the space $C_p(X, C_p(Y))$ has the JNP if and only if there exists a sequence $(\psi_n)_{n \in \omega}$ in $L_p(X) \otimes L_p(Y)$ such that $\|\psi_n\| = 1$ for every $n \in \omega$ and $\psi_n(f) \rightarrow 0$ for every $f \in C_p(X, C_p(Y))$.*

The following corollary is an immediate consequence of Theorem 4.2.

Corollary 4.6. *For every infinite Tychonoff spaces X and Y , the space $C_p(X, C_p(Y))$ has the JNP if and only if $C_p(X)$ has the JNP or $C_p(Y)$ has the JNP.*

For general locally convex spaces E , we still have one of the implications observed in the above corollary.

Proposition 4.7. *Let X be a Tychonoff space and let E be a locally convex space. If $C_p(X)$ has the JNP or E has the JNP, then the same is true for the space $C_p(X, E)$.*

Proof. Assume that E has the Josefson–Nissenzweig property. This allows us to pick a weak* null sequence $(\varphi_n)_{n \in \omega}$ in E' and a barrel-bounded set $B \subset E$ on which it does not converge to zero uniformly. Let $f \in C_p(X)$ and $x \in X$ be such that $f(x) = 1$. For each $n \in \omega$ set $\psi_n = \delta_x \otimes \varphi_n$; of course, $\psi_n \in L_p(X) \otimes E'$. Note that for every $g \in C_p(X, E)$ we have

$$\langle \psi_n, g \rangle = \langle \varphi_n, g(x) \rangle \rightarrow 0$$

as $n \rightarrow \infty$, so $(\psi_n)_{n \in \omega}$ is a weak* null sequence in $(C_p(X, E))'$.

We now exhibit a barrel-bounded subset of $C_p(X, E)$ on which $(\psi_n)_{n \in \omega}$ does not converge uniformly. The tensor product $C_p(X) \otimes E$ embeds (as a vector space) into the space $C_p(X, E)$ by the formula

$$C_p(X) \otimes E \ni \sum_i f_i \otimes e_i \mapsto [x \mapsto \sum_i f_i(x) e_i] \in C_p(X, E).$$

Thus, the subspace of $C_p(X) \otimes E$ given by the formula

$$f \otimes E = \{f \otimes e: e \in E\}$$

may be thought of as a subspace of $C_p(X, E)$. Moreover, by [35, Proposition I.4.3.(a)], $f \otimes E$ is complemented (hence closed) in $C_p(X, E)$ —a continuous linear projection is given by the mapping $g \mapsto f \otimes g(x)$ since $f(x) = 1$ by choice of f and x . Note that $f \otimes E$ is isomorphic to E . For every barrel $H \subset C_p(X, E)$ the intersection $H \cap f \otimes E$ is also a barrel in $f \otimes E$. By the assumption, $H \cap f \otimes E$ absorbs $f \otimes B$. This shows that $f \otimes B$ is absorbed by every barrel in $C_p(X, E)$, which means that it is barrel-bounded. Moreover, since $f(x) = 1$, we have:

$$\sup_{g \in f \otimes B} |\langle \psi_n, g \rangle| = \sup_{e \in B} |\langle \delta_x \otimes \varphi_n, f \otimes e \rangle| = \sup_{e \in B} |\langle \varphi_n, f(x)e \rangle| = \sup_{e \in B} |\langle \varphi_n, e \rangle| \not\rightarrow 0$$

for $n \rightarrow \infty$, which shows that $(\psi_n)_{n \in \omega}$ is a sequence witnessing the JNP of $C_p(X, E)$.

The proof for the case when $C_p(X)$ has the JNP is analogous. \square

We do not know whether the converse to Proposition 4.7 holds.

Problem 4.8. *Let X be a Tychonoff space and E a locally convex space. Assume that the space $C_p(X, E)$ has the JNP. Does it follow that $C_p(X)$ has the JNP or E has the JNP?*

Having in mind Theorem 4.1 and Proposition 4.5, we also ask the following.

Problem 4.9. *Can the Josefson–Nissenzweig property of the spaces $C_p(X, E)$ for X Tychonoff and E an arbitrary locally convex space be characterised in terms of some special “JN-sequences” in $L_p(X) \otimes E'$, like it is done in the case of $E = C_p(Y)$ for Y Tychonoff?*

5. COMPLEMENTED COPIES OF $(c_0)_p$ IN $C_p(X, C_p(Y))$

The following theorem of Banach, Kałol and Śliwa characterizes in terms of the complemented copies of the space $(c_0)_p$ those infinite Tychonoff spaces X for which the space $C_p(X)$ has the JNP.

Theorem 5.1 ([4, Theorem 1]). *Let X be an infinite Tychonoff space. Then, the following conditions are equivalent:*

- (1) $C_p(X)$ has the JNP;
- (2) $C_p(X)$ contains a complemented copy of $(c_0)_p$;
- (3) $C_p(X)$ admits a continuous linear surjection onto $(c_0)_p$.

This theorem, together with Theorems 3.3 and 4.2 and the fact that the JNP is preserved under isomorphisms, implies the following criterion for a given space $C_p(X, C_p(Y))$ to contain a complemented copy of $(c_0)_p$. Of course, by Theorem 3.3, the following result will remain true if we exchange each occurrence of $C_p(X, C_p(Y))$ by $C_p(X \times Y, \sigma)$.

Theorem 5.2. *Let X and Y be infinite Tychonoff spaces. Then, the following conditions are equivalent:*

- (1) $C_p(X, C_p(Y))$ has the JNP;
- (2) $C_p(X, C_p(Y))$ contains a complemented copy of $(c_0)_p$;
- (3) $C_p(X, C_p(Y))$ admits a continuous linear surjection onto $(c_0)_p$;
- (4) $C_p(X)$ contains a complemented copy of $(c_0)_p$ or $C_p(Y)$ contains a complemented copy of $(c_0)_p$;
- (5) $C_p(X)$ admits a continuous linear surjection onto $(c_0)_p$ or $C_p(Y)$ admits a continuous linear surjection onto $(c_0)_p$;
- (6) $C_p(X)$ has the JNP or $C_p(Y)$ has the JNP.

Corollary 5.3. *For any infinite Tychonoff space X , the space $C_p(X)$ contains a complemented copy of $(c_0)_p$ if and only if $C_p(X, C_p(X))$ contains a complemented copy of $(c_0)_p$.*

By [21, Corollary 11.5], for any infinite compact space X the space $C_p(X \times X)$ contains a complemented copy of $(c_0)_p$. Since $C_p(\beta\omega)$ does not have any complemented copies of $(c_0)_p$, Theorem 5.2 yields the following example.

Example 5.4. The space $C_p(\beta\omega, C_p(\beta\omega))$ does not have any complemented copy of $(c_0)_p$, even though the space $C_p(\beta\omega \times \beta\omega)$ does have such a copy.

As a corollary to Theorem 5.2 we obtain also the following result providing a large class of counterexamples to Problem 3.14.

Corollary 5.5. *If X is such an infinite compact space that $C_p(X)$ does not have the JNP, then $C_p(X, C_p(X))$ does not contain any complemented copy of $C_p(X \times X)$.*

Proof. Assume that $C_p(X, C_p(X))$ contains a complemented copy of $C_p(X \times X)$. Since X is compact, by [21, Corollary 11.5], $C_p(X \times X)$ (and hence $C_p(X, C_p(X))$) contains a complemented copy of $(c_0)_p$. By Corollary 5.3, $C_p(X)$ also contains a complemented copy of $(c_0)_p$. By Theorem 1, $C_p(X)$ has the JNP, a contradiction. \square

Example 5.6. $C_p(\beta\omega \times \beta\omega)$ is not complemented in $C_p(\beta\omega, C_p(\beta\omega))$.

If X is a Tychonoff space and κ a cardinal number, then we assume that $C_p(X)^\kappa$ is equipped with the standard product topology, that is, the topology given by basic open sets of the form $\prod_{\xi < \kappa} U_\xi$, where each U_ξ is open in $C_p(X)$ and for all but finitely many $\xi < \kappa$ we have $U_\xi = C_p(X)$. Using Theorem 5.1, it is easy to see that for every infinite Tychonoff space X the space $C_p(X)^2 \approx C_p(X \sqcup X)$, where $X \sqcup X$ denotes the topological disjoint union of two copies of X , contains a complemented copy of $(c_0)_p$ if and only if $C_p(X)$ contains such a copy. The following proposition generalizes this result to any infinite power of $C_p(X)$.

Proposition 5.7. *Let X be an infinite Tychonoff space and κ an infinite cardinal number. Then, the space $C_p(X)^\kappa$ contains a complemented copy of $(c_0)_p$ if and only if $C_p(X)$ contains a complemented copy of $(c_0)_p$.*

Proof. If $C_p(X)$ contains a complemented copy of the space $(c_0)_p$, then trivially $C_p(X)^\kappa$ does it, too, since $C_p(X)$ is complemented in $C_p(X)^\kappa$.

Conversely, assume that $C_p(X)$ does not contain any complemented copies of $(c_0)_p$. Note that for any infinite discrete space Y the space $C_p(Y)$ does not have the JNP, so if we equip the set $\kappa = \{\xi: \xi < \kappa\}$ with the discrete topology, then the space $C_p(\kappa)$ does not have the JNP. By Theorem 5.1, it follows that $C_p(\kappa)$ does not contain complemented copies of $(c_0)_p$. Theorem 5.2 yields that the space $C_p(X, C_p(\kappa))$ does not contain such a copy as well. Since $C_p(\kappa) \approx \mathbb{R}^\kappa$, we get the following equalities:

$$C_p(X)^\kappa = C_p(X, \mathbb{R}^\kappa) \approx C_p(X, \mathbb{R}^\kappa) \approx C_p(X, C_p(\kappa)),$$

which implies that $C_p(X)^\kappa$ does not contain any complemented copies of the space $(c_0)_p$. \square

6. COMPLEMENTED COPIES OF $(c_0)_p$ IN $C_p(X, E)$

We start with the following simple observation.

Proposition 6.1. *Let E be a locally convex space and X a Tychonoff space containing an infinite compact subset. Then $C_p(X, E)$ contains a closed copy of the space $(c_0)_p$.*

Proof. It is known that for such a space X , the space $C_p(X)$ contains a closed copy of the space $(c_0)_p$ (see the proof of [23, Theorem 5] for $\mathbb{K} = \mathbb{R}$). The thesis follows now from the fact that $C_p(X)$ is complemented in $C_p(X, E)$ (see [35, Theorem I.4.4]). \square

Let us now prove the main result of this section. The proof follows closely the arguments presented in the proof of Theorem 1 in Domański and Drewnowski [11].

Theorem 6.2. *Let E be a barrelled locally convex space with the Josefson–Nissenzweig property and let X be a Tychonoff space containing an infinite compact set. Then $C_p(X, E)$ contains a complemented copy of $(c_0)_p$.*

Proof. Since E has the JNP, there is a weak* null sequence $(x_n^*)_{n \in \omega}$ in E' which is not convergent for the $\beta^*(E', E)$ -topology. Taking into account that E is barrelled this means that the sequence is not strongly convergent and hence we may choose a bounded sequence $(x_n)_{n \in \omega}$ in E with

$$\langle x_n^*, x_n \rangle = 1.$$

Let $K \subset X$ be an infinite compact set. Using the Tychonoff property, we may choose a sequence of continuous functions $\varphi_n: X \rightarrow [0, 1]$ with disjoint support and $\varphi_n(t_n) = 1$ for some $t_n \in K$.

We set

$$J: (c_0)_p \rightarrow C_p(X, E), \quad (a_n)_{n \in \omega} \mapsto \sum_{n \in \omega}^J a_n \varphi_n(t) x_n$$

and

$$P: C_p(X, E) \rightarrow (c_0)_p, \quad f \mapsto^P (\langle x_n^*, f(t_n) \rangle)_{n \in \omega}.$$

The mapping J is well-defined, since the sequence $(a_n x_n)_{n \in \omega}$ is bounded and the functions φ_n have disjoint support—this implies that the series converges uniformly on X , and hence it defines a continuous function from X into E . Note that the condition of the supports of φ_n implies that this function only takes values of the form $a_n \varphi_n(t) x_n$. Let us see why P is well-defined, too. Since f is continuous the set $f(K) \subset E$ is compact. Since on equicontinuous subsets of E' the weak* topology coincides with the topology of uniform convergence on precompact subsets of E (see e.g. [17, Proposition 3.9.8]) and the sequence $(x_n^*)_{n \in \omega}$, being a weak* null sequence, is equicontinuous, we may conclude that $\langle x_n^*, e \rangle \rightarrow 0$ uniformly for $e \in f(K)$.

We now check that these mappings are continuous. Let p be a continuous semi-norm on E and $C > 0$ such that $p(x_n) \leq C$ for all n (such C exists, since $(x_n)_{n \in \omega}$ is a bounded sequence). For each $t \in X$, since the functions φ_n have disjoint supports, there is at most one m_t with $\varphi_{m_t}(t) \neq 0$. We pick this m_t if it exists and set $m_t = 1$ otherwise (of course, any other m_t works as well in this case), and obtain that for every $(a_n)_{n \in \omega} \in (c_0)_p$ and $t \in X$ we have:

$$p\left(J((a_n)_{n \in \omega})(t)\right) = p\left(\sum_{n \in \omega} a_n \varphi_n(t) x_n\right) \leq C |a_{m_t}| \leq C \max_{n \leq N} |a_n|$$

for all $N \geq m_t$. It follows that J is continuous. Now, let $f \in C_p(X, E)$ be given and $N \in \omega$ be arbitrary. Observe that since the functionals x_n^* are continuous there is a continuous semi-norm p on E and $\tilde{C} > 0$ with

$$\max_{n \leq N} |(P(f))_n| = \max_{n \leq N} |\langle x_n^*, f(t_n) \rangle| \leq \tilde{C} \max_{n \leq N} p(f(t_n)),$$

which implies the continuity of P .

Since

$$\begin{aligned} P(J((a_n)_{n \in \omega})) &= \left(\left\langle x_n^*, \sum_{m=1}^{\infty} a_m \varphi_m(t_n) x_m \right\rangle \right)_{n \in \omega} \\ &= (a_n \langle x_n^*, x_n \rangle)_{n \in \omega} = (a_n)_{n \in \omega}, \end{aligned}$$

we have that J is an isomorphic embedding and P is a surjection. Since $JPJP = JP$, the mapping JP is a projection onto the image of J . \square

Remark 6.3. Note the following comment concerning Theorem 6.2.

(1) If X is infinite discrete and E is a Banach space, then $C_p(X, E) \approx E^X$, which does not contain any closed copies of $(c_0)_p$, as the latter is not complete.

(2) The existence of an infinite compact subspace in X is not necessary for Theorem 6.2. Indeed, if X is a pseudocompact space with all compact subsets being finite and K is an infinite compact space, then:

$$C_p(X, C(K)) \approx C_k(X, C_k(K)) \approx C_k(X \times K),$$

which, by [22, Proposition 2.10], contains a complemented copy of $(c_0)_p$.

(3) If $C_p(Y)$ has the JNP for an infinite Tychonoff space Y , then $C_p(Y)$ is not barrelled (by the Buchwalter–Schmets theorem [9] and [21, Lemma 4.11]), so it does not satisfy the assumptions of Theorem 6.2, even though the space $C_p(X, C_p(Y))$ still has the JNP for every space X (by Theorem 5.2).

As a corollary we immediately obtain the following C_p -variant of Domański–Drewnowski's Theorem 1.2.

Corollary 6.4. *Let X be a Tychonoff space containing an infinite compact set and E a Fréchet space which is not a Montel space. Then $C_p(X, E)$ contains a complemented copy of $(c_0)_p$.*

Proof. Since by [3, Theorem 2.3] a Fréchet space has the JNP if and only if it is not a Montel space, the claim is a special case of Theorem 6.2. \square

Using the Closed Graph Theorem one can easily derive a special case of Theorem 1.2 for Fréchet spaces $C_k(X, E)$. Thus, if X is compact and E infinite-dimensional Banach, we obtain Cembranos' and Freniche's Theorem 1.1.

Corollary 6.5. *Let X be a Tychonoff space containing an infinite compact subspace and let E be a Fréchet space which is not Montel. If $C_k(X, E)$ is an infinite-dimensional Fréchet space, then $C_k(X, E)$ contains a complemented copy of the Banach space c_0 .*

Proof. Let τ_p and τ_k denote the pointwise topology of the space $C_p(X, E)$ and the compact-open topology of $C_k(X, E)$, respectively. By Theorem 6.2, $C_p(X, E)$ contains a (closed) complemented subspace G which is isomorphic to $(c_0)_p$, i.e. $(G, \tau_p \upharpoonright G) \approx (c_0)_p$. Let ξ be the Banach topology on G such that $\tau_p \upharpoonright G \subseteq \xi$ and $(G, \xi) \approx c_0$. The identity map $I: (G, \tau_p \upharpoonright G) \rightarrow (G, \tau_k \upharpoonright G)$ has closed graph, so also $I: (G, \xi) \rightarrow (G, \tau_k \upharpoonright G)$ has closed graph. The Closed Graph Theorem ([7, Theorem 4.1.10]) assures that both I and I^{-1} are continuous, so $(G, \xi) = (G, \tau_k \upharpoonright G)$, which completes the proof. \square

Theorem 6.2 yields also the following dichotomy.

Corollary 6.6. *Let X and Y be infinite Tychonoff spaces. Assume that Y is compact. Then $C_p(X, C(Y))$ contains a complemented copy of \mathbb{R}^ω or a complemented copy of $(c_0)_p$.*

Proof. We need to consider the following three cases:

Case 1. *X is not pseudocompact.* Then $C_p(X, C(Y))$ contains a complemented copy of \mathbb{R}^ω (since $C_p(X)$ contains such a copy and is complemented in $C_p(X, C(Y))$).

Case 2. *X is pseudocompact and X contains an infinite compact subset.* We apply Theorem 6.2 to get a complemented copy of $(c_0)_p$.

Case 3. *X is pseudocompact and every compact set in X is finite.* Since $X \times Y$ is pseudocompact, we apply [22, Proposition 2.10] to deduce that $C_p(X, C(Y)) \approx C_k(X \times Y)$ contains a complemented copy of $(c_0)_p$. \square

Recall that a Tychonoff space X is σ -compact if X is covered by an increasing sequence of compact subsets. We provide the following extension of [22, Proposition 2.10].

Corollary 6.7. *Let X and Y be infinite locally compact σ -compact spaces. Then $C_k(X \times Y)$ contains a complemented copy of \mathbb{R}^ω or a complemented copy of the Banach space c_0 .*

Proof. By [29, Corollary 2.5.7], the spaces $C_k(X \times Y)$ and $C_k(X, C_k(Y))$ are isomorphic. Assume first that Y is non-discrete. Since $C_k(Y)$ is not Montel (see [18, Theorem 11.8.7]) and it is a Fréchet space [7, 10.1.25], we apply Corollary 6.5 to obtain a complemented copy of the Banach space c_0 (note that the space $C_k(X, C_k(Y))$ is Fréchet by [35, Theorem IV.1.3.(b)]). On the other hand, if Y is discrete, then Y is countable, and hence the space $\mathbb{R}^\omega = C_k(Y)$ is complemented in $C_k(X, C_k(Y)) \approx C_k(X \times Y)$. \square

Remark 6.8. If X is a locally compact and σ -compact space without infinite compact subsets, then X must necessarily be discrete and countable, and hence homeomorphic to ω . It follows that for such X and any locally convex space E , the space $C_p(X, E) = C_k(X, E)$ contains a complemented copy of \mathbb{R}^ω (cf. Case 1 in the proof of Corollary 6.6).

Since $(c_0)_p \approx C_p(S)$ is linearly homeomorphic to $C_p(S) \times \mathbb{R}$, we obtain the following application of Corollary 6.5.

Corollary 6.9. *If X and Y are as in Corollary 6.6, then the space $C_p(X, C(Y))$ is isomorphic to the space $C_p(X, C(Y)) \times \mathbb{R}$.*

The above result may be thought of as a (partial) positive answer to a problem of Arkhangel'ski for spaces of the form $C_p(X, E)$. Recall that Arkhangel'ski (see [1]) asked if it is true that for every infinite (compact) space K the space $C_p(K)$ is linearly homeomorphic to $C_p(K) \times \mathbb{R}$. For a wide class of spaces the answer to this question is affirmative, e.g. if the space X contains a non-trivial convergent sequence, or X is not pseudocompact (see [1, Section 4]), yet, in general, the answer is negative even for compact spaces K —namely, Marciszewski [28] showed that there exists an infinite compact space K such that $C_p(K)$ cannot be mapped onto $C_p(K) \times \mathbb{R}$ by a continuous linear surjection.

We finish this section with the following natural question motivated by Theorems 5.1 and 5.2.

Problem 6.10. *Let X be a Tychonoff space (containing an infinite compact subspace) and E a locally convex space. Assume that $C_p(X, E)$ has the JNP. Does it follow that $C_p(X, E)$ contains a complemented copy of $(c_0)_p$?*

7. COMPLEMENTED COPIES OF $(c_0)_p$ IN $C_p(X, E_w)$

In order to provide a “weak” version of Theorem 6.2, we need the following auxiliary result proved by Domański and Drewnowski ([11, Lemma 1]). The lemma plays a similar role in

proving Theorem 7.2 as [6, Theorem 2] does in the proof of Theorem 6.2. As usual, by ℓ_1 we mean the standard Banach space of all summable real-valued sequences.

Lemma 7.1 ([11, Lemma 1]). *Let E be a locally convex space containing a copy of the Banach space ℓ_1 . Then there exists an equicontinuous sequence $(u_n)_n$ in E' which converges uniformly on each weakly compact set in E_w but on some bounded set B in E one has $\sup_{x \in B} |u_n(x)| = 1$ for each $n \in \omega$.*

Following the main idea of the proof of Theorem 6.2, Lemma 7.1 can be used to obtain the next result.

Theorem 7.2. *Let X be a Tychonoff space which contains an infinite compact subset. Let E be a locally convex space which contains an isomorphic copy of the Banach space ℓ_1 . Then $C_p(X, E_w)$ contains a complemented copy of $(c_0)_p$.*

Kąkol and Moll [24, Theorem 3] showed that for a Tychonoff space Y the space $C_k(Y)$ does not contain a copy of ℓ_1 if and only if every compact subset of Y is scattered. Hence, we have the following

Corollary 7.3. *Let X be a Tychonoff space which contains an infinite compact subset and let Y be an infinite Tychonoff space containing a non-scattered compact subset. Then $C_p(X, C_k(Y)_w)$ contains a complemented copy of $(c_0)_p$.*

Recall that a Banach space E is a *Grothendieck space* if every weakly* convergent sequence in the dual space E' is weakly convergent. If X is a compact space such that $C(X)$ is a Grothendieck space (e.g. if $X = \beta\omega$ or $X = \omega^*$), then X is non-scattered (as otherwise it would contain a non-trivial convergent sequence and hence $C(X)$ would not be Grothendieck) and $C_p(X)$ does not contain a complemented copy of $(c_0)_p$ (see [21, Section 6]). It follows that both the spaces $C_p(X, C_k(X)_w) = C_p(X, C(X)_w)$ and $C_p(X, C(X))$ contain a complemented copy of $(c_0)_p$ (Corollary 7.3 and Theorem 6.2, respectively), even though the space $C_p(X, C_p(X))$ does not contain such a copy (Corollary 5.3).

The next corollary implies that there is no continuous linear surjection from the space $C_p(\beta\omega, C_p(\beta\omega))$ onto $C_p(\beta\omega, C(\beta\omega)_w)$ (cf. Problem 3.11).

Corollary 7.4. *Let X be a non-scattered compact space such that $C_p(X)$ does not contain a complemented copy of $(c_0)_p$. Then there is no continuous linear surjection from $C_p(X, C_p(X))$ onto $C_p(X, C(X)_w)$.*

Proof. For the sake of contradiction assume that there exists a continuous linear surjection from $C_p(X, C_p(X))$ onto $C_p(X, C(X)_w)$. Since X is non-scattered, by Corollary 7.3, $C_p(X, C(X)_w)$ contains a complemented copy of $(c_0)_p$, hence there exists a continuous linear surjection from $C_p(X, C(X)_w)$ onto $(c_0)_p$. It follows that $C_p(X, C_p(X))$ can be mapped onto $(c_0)_p$ by a continuous linear map. We apply Theorem 5.2 to derive that $C_p(X)$ contains a complemented copy of $(c_0)_p$, a contradiction. \square

The last corollary yields a variant of Krupski–Marciszewski’s result ([26, Corollary 3.2]), already mentioned in Section 3.

Corollary 7.5. *Let X be an infinite compact space. Then the spaces $C_p(X)$ and $C(X)_w$ are not isomorphic.*

Proof. Assume that an isomorphism $T: C_p(X) \rightarrow C(X)_w$ exists. Consider two cases:

Case 1. X is not scattered. Define a linear map

$$S: C_p(X, C_p(X)) \rightarrow C_p(X, C(X)_w)$$

by the formula $S(f) = T \circ f$ for each continuous $f: X \rightarrow C_p(X)$. By [29, Theorem 2.2.4] the map S is continuous. Note that S is a surjection. Indeed, choose arbitrary continuous $h: X \rightarrow C(X)_w$ and define the continuous function $f(x) = T^{-1}(h(x))$ for each $x \in X$ —clearly, $S(f)(x) = T(f(x)) = h(x)$ for every $x \in X$.

By Corollary 7.4, the space $C_p(X)$ contains a complemented copy of $(c_0)_p$. Since T is an isomorphism, also the space $C(X)_w$ contains a complemented copy F of $(c_0)_p$. As both the weak topology τ_w of $C(X)$ and the normed topology τ_∞ of $C(X)$ have the same bounded sets, we have $\tau_w \upharpoonright F = \tau_\infty \upharpoonright F$, a contradiction (since $C(X)_w$ does not contain an infinite-dimensional normed subspace).

Case 2. X is scattered. By [2, Theorem III.1.2] the space $C_p(X)$ is Fréchet–Urysohn. Hence, the space $C(X)_w$ is also Fréchet–Urysohn. By [20, Lemma 14.6] the weak and the Banach topologies of $C(X)$ coincide which implies that X is finite, a contradiction. \square

Acknowledgements. The first and the third named authors are supported by the Austrian Science Fund (FWF): I 4570-N. The second named author is supported by the GAČR project 20-22230L RVO: 67985840.

REFERENCES

- [1] A. V. Arkhangel'ski, C_p -theory, in: *Recent Progress in General Topology*, North-Holland, 1992, 1–56.
- [2] A. V. Arkhangel'ski, *Topological Function Spaces*, Kluwer, Dordrecht, 1992.
- [3] T. Banach, S. Gabrielyan, *The Josefson–Nissenzweig property for locally convex spaces*, preprint (arXiv:2003.06764), 2020.
- [4] T. Banach, J. Kałol, W. Śliwa, *Josefson–Nissenzweig property for C_p -spaces*, *RACSAM* **113** (2019), 3015–3030.
- [5] A. Bareche, A. Bouziad, *Some results on separate and joint continuity*, *Topol. Appl.* **157** (2010), no. 1, 327–335.
- [6] J. Bonnet, M. Lindström, M. Valdivia, *Two theorems of Josefson–Nissenzweig type for Fréchet spaces*, *Proc. Amer. Math. Soc.* **117** (1993), 363–364.
- [7] J. Bonnet, P. Pérez Carreras, *Barrelled Locally Convex Spaces*, North-Holland Mathematics Studies **131**, North-Holland, Amsterdam, 1987.
- [8] J. Bourgain, J. Diestel, *Limited operators and strict cosingularity*, *Math. Nachr.* **119** (1984), 55–58.
- [9] H. Buchwalter, J. Schmets, *Sur quelques propriétés de l'espace $C_s(T)$* , *J. Math. Pures et Appl.* **52** (1973), 337–352.
- [10] P. Cembranos, *$C(K, E)$ contains a complemented copy of c_0* , *Proc. Amer. Math. Soc.* **91** (1984), no. 4, 556–558.
- [11] P. Domański, L. Drewnowski, *Fréchet spaces of continuous vector-valued functions. Complementability in dual Fréchet spaces and injectivity*, *Studia Math.* **102** (1992), 257–267.
- [12] R. Engelking, *General Topology*, Heldermann Verlag, Berlin, 1989.
- [13] J. C. Ferrando, J. Kałol, *Distinguished metrizable spaces E and injective tensor products $C_p(x) \otimes_\varepsilon E$* , preprint, 2021.
- [14] O. Filipchuk, V. Maslyuchenko, O. Myronyk, *Joint continuity of separately continuous mappings with values in completely regular spaces*, *Tatra Mountains Math. Publ.* **68** (2017), no. 1, 47–58.
- [15] F.J. Freniche, *Barrelledness of the space of vector valued and simple functions*, *Math. Ann.* **267** (1984), 479–486.
- [16] M. Henriksen, R.G. Woods, *Separate versus joint continuity: A tale of four topologies*, *Topol. Appl.* **97** (1999), 175–205.
- [17] J. Horváth, *Topological Vector Spaces and Distributions*, vol. 1, Addison-Wesley Publishing Co., Reading, Mass.–London–Don Mills, Ont., 1966.
- [18] H. Jarchow, *Locally Convex Spaces*, B.G. Teubner, Stuttgart, 1981.

- [19] B. Josefson, *Weak sequential convergence in the dual of a Banach space does not imply norm convergence*, Arkiv Mat. **13** (1975), 79–89.
- [20] J. Kąkol, W. Kubiś, M. Lopez-Pellicer, *Descriptive Topology in Selected Topics of Functional Analysis*, Developments in Mathematics, Springer, New York, 2011.
- [21] J. Kąkol, D. Sobota, L. Zdomskyy, *The Josefson–Nissenzweig theorem, Grothendieck property, and finitely supported measures on compact spaces*, preprint (arXiv:2009.07552), 2020.
- [22] J. Kąkol, W. Marciszewski, D. Sobota, L. Zdomskyy, *On complemented copies of the space c_0 in spaces $C_p(X \times Y)$* , Israel J. Math., accepted, 2021.
- [23] J. Kąkol, W. Śliwa, *On metrizable subspaces and quotients of non-Archimedean spaces $C_p(X, \mathbf{K})$* , RACSAM **114** (2020), 125, 1–23.
- [24] J. Kąkol, S. Moll-Lopez, *A note on the weak topology of spaces $C_k(X)$ of continuous functions*, RACSAM (2021), 115–125.
- [25] J.L. Kelley, I. Namioka, *Linear Topological Spaces*, Springer, Berlin–Heidelberg, 1963.
- [26] M. Krupski, W. Marciszewski, *On the weak and pointwise topologies in function spaces. II*, J. Math. Anal. Appl. **452** (2017), no. 1, 646–658.
- [27] M. Lindström, T. Schlumprecht, *A Josefson–Nissenzweig theorem for Fréchet spaces*, Bull. London Math. Soc. **25** (1993), 55–58.
- [28] W. Marciszewski, *A function space $C_p(X)$ not linearly homeomorphic to $C_p(X) \times \mathbb{R}$* , Fund. Math. **153** (1997), 125–140.
- [29] R.A. McCoy, I. Ntantu, *Topological Properties of Spaces of Continuous Functions*, Lecture Notes in Math. **1315**, Springer, Berlin–Heidelberg, 1988.
- [30] I. Namioka, *Separate continuity and joint continuity*, Pacific J. Math. **51** (1974), no. 2, 515–531.
- [31] L. Narici, E. Beckenstein, *Topological Vector Spaces*, Marcel Dekker, 1985.
- [32] A. Nissenzweig, *w^* sequential convergence*, Israel J. Math. **22** (1975), 266–272.
- [33] Z. Piotrowski, *Separate and joint continuity*, Real Anal. Exch. **11** (1985), no. 2, 293–322.
- [34] Z. Piotrowski, *Separate and joint continuity, II*, Real Anal. Exch. **15** (1989), no. 1, 248–258.
- [35] J. Schmets, *Spaces of Vector-Valued Continuous Functions*, Lecture Notes in Math. **1003**, Springer, Berlin–Heidelberg, 1983.
- [36] Z. Semadeni, *Banach Spaces of Continuous Functions*, Monografie Matematyczne **55**, PWN—Polish Scientific Publishers, Warszawa, 1971.
- [37] V. V. Tkachuk, *A C_p -Theory Problem Book. Functional Equivalencies*, vol. 4, Springer, Zurich, 2016.
- [38] J. van Mill, *The Infinite-Dimensional Topology of Function Spaces*, North-Holland Mathematical Library **64**, North-Holland, Amsterdam, 2001.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF INNSBRUCK, INNSBRUCK, AUSTRIA.

Email address: christian.bargetz@uibk.ac.at

FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, A. MICKIEWICZ UNIVESITY, POZNAŃ, POLAND,
AND INSTITUTE OF MATHEMATICS, CZECH ACADEMY OF SCIENCES, PRAGUE, CZECH REPUBLIC.

Email address: kakol@amu.edu.pl

KURT GÖDEL RESEARCH CENTER, DEPARTMENT OF MATHEMATICS, VIENNA UNIVERSITY, VIENNA, AUSTRIA.

Email address: damian.sobota@univie.ac.at