

# A LIOUVILLE THEOREM FOR THE NEUMANN PROBLEM OF MONGE-AMPÈRE EQUATIONS

**Huaiyu Jian**

Department of Mathematical Sciences, Tsinghua University  
Beijing 100084, China

**Xushan Tu**

Department of Mathematical Sciences, Tsinghua University  
Beijing 100084, China

**ABSTRACT.** In this paper, we study the Neumann problem of Monge-Ampère equations in Semi-space. For two dimensional case, we prove that its viscosity convex solutions must be a quadratic polynomial. When the space dimension  $n \geq 3$ , we show that the conclusion still holds if either the boundary value is zero or the viscosity convex solutions restricted on some  $n - 2$  dimensional subspace is bounded from above by a quadratic function.

**AMS Mathematics Subject Classification:** 35J60, 35B50.

**Running head:** Liouville theorem for Monge-Ampère equation

# A Liouville theorem for the Neumann problem of Monge-Ampère equations

Huaiyu Jian Xushan Tu

## 1. INTRODUCTION

The Bernstein theorem for Monge-Ampère equation  $\det D^2u(x) = 1$  in  $\mathbb{R}^n$  concludes that its solution must be a quadratic polynomial, which was proved in [13, 7, 20, 21, 8]. This type of theorems was extended to the Hessian equation[1], special Lagrangian equation[24], affine maximal hypersurface equation[22] and more general Monge-Ampère equation  $\det D^2u(x) = f(x)$  [5, 6, 9, 17], and it is called Liouville theorem if additional assumption on the solution is necessary, just as the Liouville theorem for harmonic functions. The Liouville theorem for Monge-Ampère equations in semi-space  $\mathbb{R}_+^n$  with Dirichlet boundary value on  $\partial\mathbb{R}_+^n$  was studied in [16, 19].

In this paper, we study the Liouville theorem for the Neumann problem

$$(1.1) \quad \begin{cases} \det D^2u(x) = 1 & \text{in } \mathbb{R}_+^n, \\ D_nu(x', 0) = ax_1 & \text{on } \partial\mathbb{R}_+^n. \end{cases}$$

Here and below,  $\mathbb{R}_+^n := \{(x', x_n) \in \mathbb{R}^n : x_n > 0\}$ ,  $D_nu = D_{e_n}u = \frac{\partial u}{\partial x_n}$ ,  $a \in \mathbb{R}$  is a constant, and  $e_n$  is the standard unit vector on  $x_n$ -axis.

When  $n \geq 3$ , we need to assume that  $u$  is quadratic growth in the auxiliary subspace at the infinite, i.e.,

$$(1.2) \quad \lim_{z \in \mathbb{R}^{n-2}, z \rightarrow \infty} \frac{u(0, z, 0)}{|z|^2} < \infty.$$

We use  $\Theta$  to denote the family of functions satisfying (1.2) .

We will use the standard concept for (Alesandrov) generalized solutions and viscosity solutions to the first equation of (1.1). See the book [11] for example. Caffarelli[2] indicated that the generalized solution and viscosity solution to the first equation of (1.1) are equivalent, which is proved in detail in [11]. This fact will be used again and again. To describe the Neumann boundary value condition in a weak sense, we choose the viscosity solution.

We say that  $u$  is a viscosity subsolution ( or supersolution) of problem (1.1) if  $u \in C(\overline{\mathbb{R}_+^n})$  is a convex function, and for any  $x_0 \in \overline{\mathbb{R}_+^n}$  and any convex function  $\varphi \in C^2(\overline{\mathbb{R}_+^n})$ , the condition that  $u - \varphi$  has a local maximum ( or minimum) at  $x_0$ , implies

$$\begin{cases} \det D^2\varphi(x_0) \geq (\text{or } \leq) 1 & \text{if } x_0 \in \mathbb{R}_+^n, \\ D_n\varphi(x_0) \geq (\text{or } \leq) \phi(x_0) & \text{if } x_0 \in \partial\mathbb{R}_+^n. \end{cases}$$

If  $u \in C(\overline{\mathbb{R}_+^n})$  is both viscosity subsolution and viscosity supersolution, we call it a viscosity solution.

The main result of this paper is the following

**Theorem 1.1.** *Let  $u$  be a viscosity solution to problem (1.1). If either  $n = 2$ , or when  $n \geq 3$ ,  $u \in \Theta$  or  $a = 0$ , then  $u$  is a quadratic polynomial.*

In a future paper [14], we will use Theorem 1.1 and perturbation method to prove the global Schauder estimate for the viscosity solution to the Neumann problem of Monge-Ampère equation  $\det D^2u(x) = f(x)$  in a bounded convex domain. Such estimate was obtained in [18, 23] for Dirichlet boundary value problem.

Observe that the condition (1.2) does not depend on the base point 0, this is because

$$2u(a, b + z'', 0) \leq u(0, 2z'', 0) + u(2a, 2b, 0)$$

by the convexity, and (1.2) is invariant after subtracting any linear functions. Therefore, under a suitable linear transformation, our theorem actually applies to the following problem:

$$(1.3) \quad \begin{cases} \det D^2u(x) = c & \text{in } \mathbb{R}_+^n, \\ D_\alpha u(x', 0) = \beta \cdot x' + b & \text{on } \partial\mathbb{R}_+^n, \end{cases}$$

where  $c > 0$  and  $b \in R$  are constants, while  $\alpha$  and  $\beta$  are constant vectors such that  $\alpha \cdot e_n > 0$  and  $\beta \cdot e_n = 0$ . In this situation, the assumption (1.2) changes to

$$(1.4) \quad \lim_{z' \in H'', z' \rightarrow \infty} \frac{u(z', 0)}{|z'|^2} < \infty$$

for some  $n - 2$  dimensional subspace  $H'' \subset R^n$  which is not parallel to  $\beta$ . And Theorem 1.1 is turned to

**Theorem 1.2.** *Let  $u \in C(\overline{\mathbb{R}_+^n})$  be a continuous, convex viscosity solution to problem (1.3). Suppose that either  $n = 2$ , or when  $n \geq 3$ ,  $u$  satisfies (1.4) or  $\beta = 0$ , then  $u$  is a quadratic polynomial.*

Our start point is to consider the level set at boundary point. First, we need to prove that  $\nabla u(0)$  is well defined for a viscosity solution. Then the main task is to study the geometry of the height section

$$(1.5) \quad S_h(0) := \{x \in \overline{\mathbb{R}_+^n} \mid u(x) < u(0) + \nabla u(0) \cdot x + h\}$$

by blowing down  $u$  at the infinite point of auxiliary subplane and then blowing up at origin. As a result, we will finally prove that  $D_n u(x)$  is a linear function. The case  $n = 2$  or  $a = 0$  is much more easy, which will be proved in Remark 2.16 and Remark 8.2 respectively.

This paper is arranged as follows. In section 2, we will show the basic properties for the viscosity solutions of problem (1.1) and list a few well-known Lemmas to be used. In section 3, we will prove the strict convexity of solution to (1.1) away from the boundary and then introduce the linearized problem for  $D_n u(x)$  to obtain a boundary estimate of the normal derivatives of the solutions. In section 4, we will study a general global strict convexity lemma and provide an example showing that the growth assumption for the strict convexity is optimal. In section 5, we will prove the height section of solution to (1.1) has a good shape. In section 6, we will use the result for case  $a = 0$  to prove a stability result for small  $a$ . In section 7, we will prove the global uniformly convexity of solution to (1.1). In the last section, we will improve the famous  $C^{1,\alpha}$  estimate techniques[2, 4], so as to prove that the blow down-up limit has certain  $C^2$  regularity on a cone with vertex 0, which implies that  $u$  is a quadratic polynomial.

## 2. PRELIMINARIES

We start at notations and definitions, then list some basic results. They will be used throughout this paper.

A point in  $\mathbb{R}^n$  is written as

$$x = (x_1, \dots, x_{n-1}, x_n) = (x', x_n)$$

and as  $x = (x_1, x'', x_n)$  if  $n \geq 3$ .

Let  $\mathbb{R}_+^n = \{x \in \mathbb{R}^n \mid x_n > 0\}$  be the upper half-space. Then  $\overline{\mathbb{R}_+^n} = \{x \in \mathbb{R}^n \mid x_n \geq 0\}$ . We will take the convention

$$R^{n-1} = \partial\mathbb{R}_+^n = \{x \in \mathbb{R}^n \mid x_n = 0\}, \quad R^{n-2} = \{x \in \mathbb{R}^n \mid x_1 = x_n = 0\}$$

from time to time.

Denote by  $\mathcal{I}$  (or  $\mathcal{I}'$ ,  $\mathcal{I}''$ ) the identity matrices of size  $n$  (or  $n-1$ ,  $n-2$ ),  $B_r(x)$  (or  $B'_r(x')$ ,  $B''_r(x'')$ ) the ball of radius  $r$  centered at  $x$  (or  $x'$ ,  $x''$ ) in  $\mathbb{R}^n$  (or  $\mathbb{R}^{n-1}$ ,  $\mathbb{R}^{n-2}$  respectively),  $B_r^+(x) = \{y \in B_r(x) : y_n > 0\}$  and  $\mathcal{P}x = x'$  is the projection mapping onto the  $\mathbb{R}^{n-1}$ .

The oscillation function of a function  $W$  over a domain  $E$  is denoted by

$$Osc_E W = \sup_{y, z \in E} |W(y) - W(z)|.$$

Suppose that  $u$  satisfies the assumption of Theorem 1.1. Since  $u$  is continuous up to boundary, we have

$$(2.1) \quad S_1(0) \supset B_{\kappa_0}^+(0)$$

for some  $\kappa_0 > 0$ , where we have used notation (1.5) (Also see (2.4) below).

When  $n \geq 3$ , the quadratic growth condition (1.2) will be replaced by

$$(2.2) \quad u(0, x'', 0) < \frac{K}{2}(|x''|^2 + 1), \forall x'' \in \mathbb{R}^{n-2}$$

for a constant  $K > 0$ . We will use  $\sigma_i : [0, \infty) \rightarrow [0, \infty)$ , ( $i = 0, 1, 2$ ), to denote some universal strictly increasing  $C^0$  function with  $\sigma_i(0) = 0$  that depends only on  $(n, a, \kappa_0, K)$ .

$\lambda$  and  $\Lambda$  will denote known positive constants. We always use the convention that  $c$  or  $C$  denotes various positive constants depending only on  $n, a, \kappa_0, K, \lambda$  and  $\Lambda$ , which are often called as *universal constants*.

If not specified, the  $\det D^2u$  will be understood as the general Monge-Ampère measure of convex functions[10, 11], which yields the (Alexandrov) generalized solution to the equation  $\det D^2u = f$ , and it is equivalent to the viscosity solution in the domain where  $f$  is continuous. If  $u$  is a viscosity subsolution (or supersolution) of the equation  $\det D^2u = f$ , we say that  $u$  is a viscosity solution to the inequalities  $\det D^2u \geq$  (or  $\leq$ )  $f$ , or  $u$  solves these inequalities in viscosity senses.

The sub-differential of  $u$  at point  $x_0$  is defined by

$$\partial u(x_0) := \{p \mid L := u(x_0) + p \cdot (x - x_0) \text{ is a support plane of } u \text{ at } x_0\}.$$

We define  $\nabla u(x_0)$  as a fixed element of  $\partial u(x_0)$  for interior point. When  $n = 2$  or when  $n \geq 3$  and the assumption (1.2) is satisfied, we will show the viscosity solution to (1.1),  $u$ , is strictly convex in  $\mathbb{R}_+^n$ . Then the classical interior estimate [2, 4] means that  $u \in C_{loc}^\infty(\mathbb{R}_+^n)$  and  $\nabla u$  is uniquely determined in  $\mathbb{R}_+^n$ .

Take a point  $y = (y', 0) \in \partial \mathbb{R}_+^n$ . If  $u$  is convex, viscosity solution to problem(1.1), by the definition we have

$$(2.3) \quad \sup_{a \in \partial u(y)} \{a \cdot e_n\} \leq ay_1,$$

and by Lemma 3.2 in Section 3, one may define

$$D_n u(y) = \overline{\lim}_{t \rightarrow 0^+} D_n u(y + te_n) := \lim_{t \rightarrow 0^+} \sup \{p \cdot e_n : p \in \partial u(y + te_n)\}.$$

And in Corollary 3.3, we will prove that there exists  $p_y \in \partial u(y)$  such that

$$p \cdot e_n = ay_1 = D_n u(y).$$

Hence, we define  $\nabla u(y)$  as a fixed  $p_y$  for any boundary point  $y \in \partial \mathbb{R}_+^n$ .

In this way, the section of  $u$  centered at  $y \in \overline{\mathbb{R}_+^n}$  with height  $h > 0$  can be defined by

$$(2.4) \quad S_h^u(y) := \{x \in \overline{\mathbb{R}_+^n} \mid u(x) < u(y) + \nabla u(y) \cdot (x - y) + h\},$$

and its Neumann boundary is defined by

$$(2.5) \quad G_h^u(y) := S_h^u(y) \cap \partial \mathbb{R}_+^n.$$

Sometimes, the notations  $S_h^u(0)$ ,  $G_h^u(0)$ ,  $B_r(0)$  and  $B_r^+(0)$  will be shorten as  $S_h$ ,  $G_h$ ,  $B_r$  and  $B_r^+$  respectively.

When we are going to prove Theorem 1.1, we may assume

$$(2.6) \quad a \geq 0, \quad u(0) = 0, \quad \nabla u(0) = 0 \quad \text{and} \quad u \geq 0 \quad \text{in} \quad \overline{\mathbb{R}_+^n}$$

by considering the function  $u(x, -x_n)$  and subtracting the linear function  $u(0) + \nabla u(0) \cdot x$ .

**Definition 2.1** (Good Shape). *We say that a section  $S_h^u(0)$  is of good shape if it satisfies the following two property:*

$S_h^u(0)$  has finite density at 0, i.e.,

$$c \leq \frac{\text{Vol}(S_h^u(0))}{h^{\frac{n}{2}}} \leq C,$$

and

$$c\mathcal{P}S_h^u(0) \subset \mathcal{P}G_h^u(0) \cap (-\mathcal{P}G_h^u(0)).$$

When blowing up at infinity, we will need to modify the class  $\Theta$  to  $\Theta(\delta)$ .

**Definition 2.2** ( $\Theta(\delta)$  Class). *Suppose that  $n \geq 3$ ,  $b \in \mathbb{R}$  and  $u$  satisfies*

$$D_n u(x', 0) = bx_1, \quad \forall x' \in \mathbb{R}^{n-1}$$

and

$$u(0, z'', 0) \leq z''^T \mathcal{Q} z'' + \delta, \quad \forall z'' \in \mathbb{R}^{n-2}$$

for some  $(n-2) \times (n-2)$  symmetric matrix  $\mathcal{Q}$  and  $\delta > 0$ . We say  $u \in \Theta(\delta)$  if the pair  $(b, \mathcal{Q})$  satisfies

$$0 \leq b \leq c \quad \text{and} \quad b^2 \det \mathcal{Q} \leq C.$$

**Definition 2.3.** *We say that a closed set  $E \in \mathbb{R}^n$  is balance about  $x$  up to a constant  $\kappa$ , if*

$$t(x - E) \subset E - x \quad \text{for} \quad t \leq c\kappa.$$

When  $\kappa$  is universal, we say that  $E \in \mathbb{R}^n$  is balance about  $x$ .

The following lemmas will be used in our article.

Use  $\text{Vol}_k(E)$  to denote the  $k$ -dimensional Lebesgue measure and denote  $\text{Vol}(E) = \text{Vol}_n(E)$ .

**Lemma 2.4.** *Given any convex set  $E \in \mathbb{R}^n$ . Suppose the line  $l(t) := x + te_n$  intersects  $E$  at the points  $p, q$ . Then*

$$(2.7) \quad |p_n - q_n| \cdot \text{Vol}_{n-1}(\mathcal{P}E) \leq C \text{Vol}(E).$$

*In addition, if  $\mathcal{P}E$  is balance about  $\mathcal{P}q$  up to a constant  $\kappa$ , then we have the reverse inequality*

$$(2.8) \quad |p_n - q_n| \cdot \text{Vol}_{n-1}(\mathcal{P}E) \geq c\kappa \text{Vol}(E).$$

*Proof.* Consider the non-negative concave function defined on  $\mathcal{P}E$

$$L(x') = \sup\{(y - z) \cdot e_n \mid y, z \in E, \text{ such that } \mathcal{P}y = \mathcal{P}z = x'\}.$$

Then

$$\text{Vol}(E) \sim \|L\|_{L^\infty} \text{Vol}_{n-1}(\mathcal{P}E),$$

which yields (2.7). Now, suppose that  $\mathcal{P}E$  is balance about  $\mathcal{P}q$  up to the constant  $\kappa$ . By concavity  $L(\mathcal{P}q) \geq c\kappa \|L\|_{L^\infty}$ , and (2.8) follows.  $\square$

**Lemma 2.5.** *Let  $u \in C(\overline{B_1^+(0)})$  be a convex function. Suppose that*

$$0 \leq u \leq \sigma_1(|x|) \text{ in } B_c^+(0)$$

*and*

$$\overline{\lim}_{t \rightarrow 0^+} D_n u(x', t) \geq -\sigma_0(|x'|), \forall (x', 0) \in \overline{B_c^+(0)}.$$

*Then*

$$(2.9) \quad |\nabla u(y)| \leq C \left( \inf_{|y| \leq t \leq c} \frac{2\sigma_1(1+t)}{t} + \sigma_0(|y'|) \right) \text{ in } B_c^+(0).$$

*Here we do not require  $\sigma_i(0) = 0$  ( $i = 0, 1$ ).*

*Proof.* Given  $y \in B_c^+(0)$ ,  $c \leq \frac{1}{2}$ , let  $p \in \partial u(y)$ . The assumption and the convexity implies that

$$p_n \geq -\sigma_0(|y'|).$$

Suppose that  $|p'| > 0$ . Note that for every constant  $a \in [0, 1]$ , the point  $z = y + a|y|(\frac{p'}{|p'|}, 1)$  is in  $B_1^+(0)$ . Hence, by the convexity we have

$$a|y|(p_n + |p'|) = p \cdot (z - y) \leq u(z) - u(y) \leq 2\sigma_1((1+a)|y|).$$

Letting  $t = a|y|$ , we obtain the desired (2.9). We may ignore the estimation for  $y \in \partial\Omega$ , because  $\nabla u(y)$  in this paper is determined by taking the internal limit. (See the above definition for  $D_n u(y)$  and Corollary 3.3).  $\square$

Similar to Lemma 2.5, we can prove

**Lemma 2.6.** Write  $x \in \mathbb{R}^n$  as  $x = (x^1, \dots, x^k)$ ,  $x^j \in \mathbb{R}^{a_j}$ ,  $\sum_{j=1}^k a_j = n$ . Let  $u \in C(\bar{B}_r(0))$  be a convex function, satisfying

$$u(x) \leq u(0) + \sum_{i=1}^k \sigma_i(|x^i|).$$

Then

$$|\nabla_{x_j} u(0)| \leq C \inf_{0 \leq t \leq cr} \frac{2\sigma_j(Ct)}{t}.$$

**Corollary 2.7.** Let  $\epsilon > 0$  be small constant,  $u \in C(\overline{B_1^+(0)})$  and  $v \in C^2(\overline{B_1^+(0)})$  be two convex functions. Suppose that

$$\|u - v\|_{L^\infty} \leq \epsilon$$

and

$$\overline{\lim}_{t \rightarrow 0^+} D_n u(x', t) \geq D_n v(x', 0) - \epsilon^{\frac{1}{2}}, \forall (x, 0) \in \overline{B_1^+(0)}.$$

Then

$$(2.10) \quad |\nabla u(y) - \nabla v(y)| \leq C_v \epsilon^{\frac{1}{2}} \text{ in } B_c^+(0),$$

where  $C_v$  is a constant which may depend on  $\|v\|_{C^2}$ .

*Proof.* Given point  $y \in B_c^+(0)$ . Consider the function

$$U^y(x) = u(x) + 2\epsilon - v(y) - \nabla v(y) \cdot (x - y).$$

Then by the assumption we have

$$0 \leq U^y(x) \leq 2\epsilon + v(x) + 2\epsilon - v(y) - \nabla v(y) \cdot (x - y) \leq C_v |x - y|^2 + 4\epsilon,$$

and

$$\overline{\lim}_{t \rightarrow 0^+} D_n U^y(x', t) \geq D_n v(x', 0) - \epsilon^{\frac{1}{2}} - D_n v(y', 0) \geq -C_v |x' - y'| - \epsilon^{\frac{1}{2}}.$$

Regarding  $y$  as origin and applying Lemmas 2.5 and 2.6 to  $U^y(x)$ , we get the desired results.  $\square$

The following is well-known John's Lemma. See [10, 11, 12] for example.

**Lemma 2.8** (John's Lemma). *If  $\Omega$  is a bounded convex set with non-empty interior in  $\mathbb{R}^n$  and  $E$  is the ellipsoid of smallest volume containing  $\Omega$ , then after an affine transformation  $T$ ,*

$$B_{c(n)} \subset T(\Omega) \subset B_{C(n)} := T(E).$$

**Lemma 2.9.** *If  $\Omega$  is a bounded convex set in  $\mathbb{R}^n$  and  $u, v \in C(\overline{\Omega})$  are convex functions in  $\Omega$  satisfying*

$$\begin{cases} \det D^2v \leq \det D^2u & \text{in } \Omega \\ u \leq v & \text{on } \partial\Omega \end{cases},$$

*then  $u \leq v$  on  $\overline{\Omega}$ .*

*Proof.* See Theorem 2.10 in [10]. □

**Lemma 2.10** ( Comparison principle for mixed problem). *Supposed that  $\Omega$  is a bounded convex set in  $\mathbb{R}^n$ ,  $G_1 \subset \partial\Omega$  and  $G_2 = \partial\Omega \setminus G_1$  are piecewise  $C^1$  smooth, and  $\beta$  is inward vector field on  $\partial\Omega_2$ . If  $u, v$  are continuous, convex function defined on  $\overline{\Omega}$  such that either  $u$  or  $v$  is  $C^2$  smooth, satisfying*

$$(2.11) \quad \begin{cases} \det D^2v < \det D^2u < \infty & \text{in } \Omega \\ u < v & \text{on } G_1 \\ D_\beta v < D_\beta u & \text{on } G_2 \end{cases}$$

*in the viscosity sense, then  $u < v$  on  $\overline{\Omega}$ .*

*Proof.* On the contrary to the conclusion, we suppose that  $w = u - v$  archives positive maximum value  $m$  at a point  $x_0 \in \overline{\Omega}$ . According to the assumption and the first two inequalities in (2.11) we can take  $x_0 \in \partial\Omega_2$  such that  $u(x_0) - v(x_0) = m$ . This means that  $u - [v(x_0) + \nabla v(x_0) \cdot (x - x_0) + m]$  archives the maximum at  $x_0$  when  $v$  is  $C^2$  smooth, or  $v - [u(x_0) + \nabla u(x_0) \cdot (x - x_0) - m]$  archives the minimum at  $x_0$  when  $u$  is  $C^2$  smooth, which contradicts the third inequality in (2.6) by the definition of viscosity solution. □

**Lemma 2.11.** *If  $\Omega$  is a bounded convex domain in  $\mathbb{R}^n$  and  $u$  is a convex function satisfying*

$$\det D^2u > \lambda \text{ in } \Omega,$$

*then*

$$(2.12) \quad \text{Vol}(\Omega) \leq C(\lambda) \|u\|_{L^\infty}^{\frac{n}{2}}.$$

*In particular,*

$$(2.13) \quad \text{Vol}(S_h) \leq Ch^{\frac{n}{2}}.$$

*Proof.* (2.13) was proved in Corollary 3.2.5 in [11], and (2.12) can be proved as its proof. □

**Lemma 2.12.** *Suppose that  $\Omega$  is a convex domain in  $\mathbb{R}^n$  such that  $B_c(0) \subset \Omega \subset B_C(0)$  and  $u \in C(\bar{\Omega})$  is a convex function. If*

$$\lambda \leq \det D^2u \leq \Lambda \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

then

$$u(x) \geq -C \operatorname{dist}(x, \partial\Omega)^{\frac{1}{n}}.$$

*Proof.* This is the Aleksandrove's maximum principle. See Theorem 1.4.2 in [11].  $\square$

**Lemma 2.13.** *Suppose that  $B_c(0) \subset \Omega \subset B_C(0)$  and  $u$  is a convex function defined on convex domain  $\bar{\Omega}$  such that*

$$\lambda \leq \det D^2u = f(x) \leq \Lambda \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega.$$

If  $\|f\|_{C^{k,\alpha}(\Omega)} \leq C$  for some  $k \geq 0$  and  $\alpha \in (0, 1)$ , then

$$\|u\|_{C^{k+2,\alpha}(\frac{1}{2}\Omega)} \leq C.$$

*Proof.* When  $k = 0$ , this is the famous interior  $C^{2,\alpha}$  estimate in [3]. Also see [15]. Then the case of  $k \geq 1$  follows from the standard regularity result for linear elliptic equations.  $\square$

**Lemma 2.14.** *Let  $u \in C(B_1(0))$  be a convex solution to*

$$0 < \lambda \leq \det D^2u \leq \Lambda.$$

For  $x \in B_1(0)$  and  $p \in \partial u(x)$ , define  $\Sigma := \{y \in B_1(0) \mid u(y) = u(x) + p \cdot (y - x)\}$ . Then, either  $\Sigma = \{x\}$  or  $\Sigma$  has no extremal point inside  $B_1(0)$ .

*Proof.* See [2].  $\square$

The following result is already well-known. See [13, 7, 20, 21, 8] or the books [10, 11].

**Theorem 2.15.** *Let  $u \in C(\mathbb{R}^n)$  be a continuous, convex solution. If  $\det D^2u = 1$  in  $\mathbb{R}^n$ , then  $u$  is a quadratic polynomial.*

**Remark 2.16.** *The case  $a = 0$  of Theorem 1.1 can be obtained directly from Theorem 2.15. In fact, Lemma 3.2 in next section allow us to reflect  $u$ , the solution to problem (1.1), with respect to  $\partial\mathbb{R}_+^n$ . Then we get a convex function  $\bar{u}$ . Corollary 3.3 also means that  $D_n\bar{u}(\partial\mathbb{R}_+^n) \subset \partial\mathbb{R}_+^n$ . Thus  $\bar{u}$  solves the equation*

$$\det D^2\bar{u} = 1, \text{ in } \mathbb{R}^n.$$

Hence  $\bar{u}$  is a quadratic polynomial by Theorem 2.15, so is  $u$ .

### 3. INTERIOR STRICT CONVEXITY AND NORMAL DERIVATIVE ESTIMATE

In this section, we first prove that a viscosity solution to problem (1.1) is interior strict convexity, then give its normal derivative estimates.

**Lemma 3.1.** *Suppose  $u$  is a viscosity solution to problem (1.1). Then the graph of  $u$  can't contains any ray, and  $u$  is strictly convex away from the boundary  $\partial\mathbb{R}_+^n$ .*

*Proof.* If the graph of  $u$  contains a ray  $l$  and take  $y \in l' = \mathcal{P}l$ , the function

$$u^y(x) = u(x) - u(y) - \nabla u(y) \cdot (y - x)$$

is bounded in the tubular domain  $E := (l' - y + B_1(y)) \cap \overline{\mathbb{R}_+^n}$  and

$$\det D^2 u^y = 1 \text{ in } E,$$

which contradicts Lemma 2.11 since  $E$  has infinite volume.

Suppose  $y \in \mathbb{R}_+^n$ . Let

$$(3.1) \quad h_y^u := \sup\{h \mid S_t^u(y) \subset \mathbb{R}_+^n, \forall t < h\}.$$

Since  $u$  is strictly convex in the interior of sections[2], it is enough for the strict convexity of  $u$  away from the boundary  $\partial\mathbb{R}_+^n$  to show  $h_y^u > 0$  for  $y \in \mathbb{R}_+^n$ .

Otherwise, if  $h_y = 0$ , the convex set

$$\Gamma := \{x \in \mathbb{R}_+^n \mid u(x) = u(y) + \nabla u(y) \cdot (x - y)\}$$

should contain more than one point. Lemma 2.14 implies that the extremal point of  $\Gamma$  is not in interior. There are two cases: either  $\Gamma$  contains a line  $l$  parallel to  $\mathbb{R}^{n-1}$ , or  $\Gamma$  contains a ray  $l$  starting from point  $z = (z', 0)$  on  $\partial\mathbb{R}_+^n$ . Both cases contradicts the previous fact that the graph of  $u$  can't contains any ray.  $\square$

**Lemma 3.2.** *Suppose  $u$  is a viscosity solution to problem (1.1) and  $y \in \partial\mathbb{R}_+^n$ . Then  $D_n u(y) := \overline{\lim}_{t \rightarrow 0^+} D_n u(y', t)$  is well-defined. Moreover,*

$$(3.2) \quad D_n u(y) = ay_1$$

and

$$(3.3) \quad u(y + te_n) \geq u(y) + tay_1, \quad \forall t > 0.$$

*Proof.* Without loss of generality we may assume  $y = 0$ . By Lemma 3.1 and the convexity of  $u$ ,  $D_n u(te_n)$  are well-definend and continuous for  $t > 0$ . Hence,

$$D_n u(0) := \overline{\lim}_{t \rightarrow 0^+} D_n u(te_n) = \lim_{t \rightarrow 0^+} D_n u(te_n)$$

is well-defined. Notice that  $ay_1|_{y=0} = 0$ . Were if (3.2) false, then

$$\lim_{t \rightarrow 0^+} D_n u(te_n) < -2\epsilon$$

for some  $\epsilon > 0$ . Consider the tubular domain

$$\Gamma = \{(x', 0) + st_0 e_n \mid x' \in B'_r(0), s \in (0, 1)\}$$

and choose the function

$$v(x) = u(0) + \frac{2 \operatorname{Osc}_{B_r^+(0)} u}{r^2} |x'|^2 - \epsilon x_n,$$

where  $t_0$  and  $r = r(\epsilon, t_0, u)$  are positive and so small that

$$D_n u(t_0 e_n) < -2\epsilon t_0, \quad 2 \operatorname{Osc}_{B_r^+(t_0 e_n)} u \leq \epsilon t_0 \text{ and } |a|r < \epsilon,$$

which implies

$$\sup_{y \in B_r(t_0 e_n) \cup \partial B_r^+(0)} (u(y) - v(y)) < 0.$$

Notice that  $v$  is linear along the  $e_n$  direction and  $u$  is convex, we find that  $v$  is strictly greater than  $u$  on the boundary  $\partial\Gamma_1 := \overline{\partial\Gamma} \setminus \overline{\mathbb{R}^{n-1}}$ . Then,  $v(0) = u(0)$  means that  $v + c_0$  will touch  $u$  by above at some point  $z \in \overline{\Gamma} \setminus \partial\Gamma_1$  for a proper  $c_0 \in \mathbb{R}$ . Moreover,  $\det D^2 v = 0 < 1 = \det D^2 u$  implies  $z \notin \Gamma$ . Therefore,  $z \in \partial\Gamma \cap \mathbb{R}^{n-1}$ . However,

$$D_n v(z) = -\epsilon < -|a|r \leq az_1 = D_n u(z),$$

which contradicts the definition of viscosity supersolution. Thus, (3.2) is proved and (3.3) follows by a direct integral.  $\square$

**Corollary 3.3.** *If  $u$  is a viscosity solution to problem (1.1) and  $y \in \partial\mathbb{R}_+^n$ , then there exists an element  $p \in \partial u(y)$  such that  $p \cdot e_n = D_n u(y)$ .*

*Proof.* Without loss of generality we may assume  $y = 0$ . Let  $p_t = \nabla u(te_n)$  and  $l_t(x) := u(te_n) + p_t \cdot (x - te_n)$  be the support function of  $u$  at  $te_n$ . Lemmas 3.2 and 2.5 implies that  $p_t$  is uniformly bounded as  $t \rightarrow 0^+$ . Then, as  $t \rightarrow 0^+$ ,  $\{l_t\}$  is pre-compact and contains a subsequence that converges to some support function  $l(x) := u(0) + p \cdot x$  of  $u$  at 0. Hence  $p \in \partial u(0)$  and  $D_n u(te_n) = p_t \cdot e_n$  monotonically decreases to  $p \cdot e_n$ . Furthermore, by (2.3) and (3.2) we see that  $p \cdot e_n = 0 = D_n u(0)$ .  $\square$

**Lemma 3.4.** *Suppose  $u$  is a viscosity solution to problem (1.1). Then  $D_n u \in C(\overline{\mathbb{R}_+^n})$  and  $v(x) := D_n u(x) - D_n u(x', 0)$  is a viscosity solution to following linearized problem:*

$$(3.4) \quad \begin{cases} U^{ij} D_{ij} v = 0 & \text{in } \mathbb{R}_+^n \\ v \geq 0 & \text{on } \overline{\mathbb{R}_+^n} \\ v = 0 & \text{on } \partial\mathbb{R}_+^n \end{cases},$$

where  $U^{ij}$  is the cofactor of the Hessian  $D^2 u$ .

*Proof.* First, Lemmas 3.1 and 2.13 means that  $u \in C_{loc}^\infty(\mathbb{R}_+^n)$ . Then it follows from the proof of Corollary 3.3 that the family of continuous function  $f_t(x') = D_n u(x', t)$  point-wisely decreases to function  $f_0(x') = D_n u(x', 0) = ax_1$  monotonically as  $t \rightarrow 0^+$ . Finally, the finite cover Lemma implies that this convergence is uniformly. Therefore  $D_n u$  is continuous up to boundary.

The linearized equation  $U^{ij} D_{ij} v = 0$  comes directly from the equation  $\det D^2 u = 1$ . The other two in (3.4) follows from the convexity of  $u$  and the continuity of  $D_n u$  at the boundary.  $\square$

Next, we introduce two functions,  $w^{y,+}(x)$  and  $w^{y,-}(x)$ , which will be used as the upper/below barriers of  $v$  at  $y$  to the linearized problem (3.4).

Given  $y \in G_{\frac{1}{2}}^u(0)$  and consider the functions

$$(3.5) \quad u^y(x) = u(x) - u(y) - \nabla u(y) \cdot (y - x)$$

and

$$(3.6) \quad w^y(x) = u^y(x) - \frac{n}{2} x_n D_n u^y.$$

Define

$$(3.7) \quad \begin{cases} w^{y,+}(x) = C_1[w^y(x) + C_2 x_n] \\ w^{y,-}(x) = c_1[x_n - c_2 w^y(x)] \end{cases}.$$

Since  $\det D^2 u = 1$ , we have  $U^{ij} D_{ij} D_n u = 0$  in  $\mathbb{R}_+^n$ . Hence

$$(3.8) \quad U^{ij} D_{ij} w^y = U^{ij} D_{ij} w^0 = n \det D^2 u - \frac{n}{2} (U^{in} D_{in} u + U^{ni} D_{ni} u) = 0 \text{ in } \mathbb{R}_+^n.$$

**Lemma 3.5.** *Suppose that  $u$  is a viscosity solution to problem (1.1) and (2.6) is satisfied. If*

$$B_c^+(0) \subset S_1(0) \subset B_C^+(0)$$

and

$$(3.9) \quad \|D_n u\|_{L^\infty(S_1(0))} \leq C,$$

then,  $w^{0,+}(x)$  defined in (3.7) is the upper barrier of  $v$  at 0 to the linearized problem (3.4) in  $S_1(0)$ . (See (3.11) below). This implies

$$(3.10) \quad D_n u(te_n) \leq Ct \quad \text{and} \quad u(te_n) \leq Ct^2$$

for all  $t \geq 0$  such that  $te_n \in S_1(0)$ .

*Proof.* It follows from (3.9) that

$$v(x) = D_n u(x) - D_n u(x', 0) \leq C, \quad \forall x \in S_1(0)$$

and

$$u(x) - \frac{n}{2}x_n D_n u \geq 1 - Cx_n, \quad \forall x \in \partial S_1(0) \setminus G_1(0).$$

In (3.7) we choose  $C_1$  and  $C_2$  such that they are larger than the  $C$ . Then,  $w^{0,+}(x) \geq C_1$  on  $\partial S_1(0) \setminus G_1(0)$  by (3.6). This, together with (3.5)-(3.8), we see that

$$(3.11) \quad \begin{cases} U^{ij} D_{ij} w^{0,+} = 0 & \text{in } S_1(0) \\ w^{0,+} \geq v & \text{on } \partial S_1(0) \setminus G_1(0) . \\ w^{0,+} \geq 0 = v & \text{on } G_1(0) \end{cases}$$

Hence, the comparison principle implies that  $v(x) \leq w^{0,+}(x)$  in  $S_1(0)$ , which, together with the convexity, yields the desired (3.10).  $\square$

**Lemma 3.6.** *Under the assumption of Lemma 3.5, there exists a small  $\bar{\sigma} > 0$  depending on  $\sigma_0$  such that if*

$$(3.12) \quad v(x) \geq \sigma_0(x_n) - \bar{\sigma} \text{ in } S_1(0),$$

then  $w^{0,-}(x)$  defined in (3.7) is the lower barrier of  $v$  at 0 to the linearized problem (3.4) in  $S_1(0)$  (see (3.14) below). This implies that

$$(3.13) \quad v(te_n) \geq ct \quad \text{and} \quad u(te_n) \geq ct^2$$

for all  $t \geq 0$  such that  $te_n \in S_1(0)$ .

*Proof.* Denote  $K = Cn \|D_n u\|_{L^\infty(S_1(0))}$ . Assume that  $\bar{\sigma} > 0$  is small and satisfies

$$\sigma_0^{-1}(2\bar{\sigma}) \leq \frac{1}{2K}.$$

Recall that  $\sigma_0$  denotes a strictly increasing and continuous function from  $[0, +\infty)$  to  $[0, +\infty)$  with  $\sigma_0(0) = 0$ . Let  $c_0 = \sigma_0^{-1}(2\bar{\sigma})$ . By (3.12) and the convexity we have

$$v(x) \geq \max\{\sigma_0(x_n) - \bar{\sigma}, 0\} \geq \bar{\sigma} \chi_{\{x_n \geq c_0\}},$$

where  $\chi_E$  denotes the characteristic function of the set  $E$ , i.e.,  $\chi_E(x) = 1$  if  $x \in E$ , otherwise  $\chi_E(x) = 0$ . Take  $c_1 = \frac{\bar{\sigma}}{2c}$  and  $c_2 = 2c_0$ . We have

$$w^0(x) \geq 1 - Kx_n, \quad \text{on } \partial S_1(0) \setminus G_1(0).$$

Hence on  $\partial S_1(0) \setminus G_1(0)$ , we see that

$$\begin{aligned} w^{0,-}(x) &\leq c_1[x_n + c_2(Kx_n - 1)] \\ &\leq c_1[(2x_n - c_2)\chi_{\{x_n \leq c_0\}} + 2x_n\chi_{\{x_n \geq c_0\}}] \\ &\leq 2c_1 C \chi_{\{x_n \geq c_0\}} \\ &= \bar{\sigma} \chi_{\{x_n \geq c_0\}} \leq v(x). \end{aligned}$$

In conclusion, we have

$$(3.14) \quad \begin{cases} U^{ij}D_{ij}w^{0,-} = 0 & \text{in } S_1(0) \\ w^{0,-} \leq v & \text{on } \partial S_1(0) \setminus G_1(0) . \\ w^{0,-} \leq 0 = v & \text{on } G_1(0) \end{cases}$$

The comparison principle implies that  $w^{0,-}(x) \leq v(x)$  in  $S_1(0)$ . In particular,

$$v(te_n) \geq w^{0,-}(te_n) \geq c_1(t - Cc_2t) \geq \frac{c_1}{2}t$$

when  $c_2 = 2\sigma_0^{-1}(2\bar{\sigma}) > 0$  is small, which is equivalent to  $\bar{\sigma} > 0$  is small enough. This proves (3.13).  $\square$

**Corollary 3.7.** *Under the assumption of Lemma 3.5, there exists a small  $\rho > 0$  such that if*

$$(3.15) \quad \sup_{B'_\rho(0)} \|u(y', 0)\| \leq \sigma_0(\rho)\rho,$$

then for any  $y' \in B'_\rho(0)$  and  $y = (y, 0)$ , the function  $w^{y,+}(x)$  defined in (3.7) is the upper barrier of  $v$  at  $y$  to the linearized problem (3.4). This also implies that  $u^y(y + te_n) \leq Ct^2$  for small  $t > 0$  if  $u(y) = 0$  and  $\nabla u(y) = 0$ .

*Proof.* If  $\rho$  is small, the assumption implies that

$$\sup_{B'_\rho(0)} \|\nabla u(y', 0)\| \leq \min\{\sigma_0(\rho), C\rho\} \ll \frac{1}{2}.$$

Then, the section

$$S_{\frac{1}{2}}(y) \subset S_{1-C|\nabla u(y)|}(y) \subset S_1(0).$$

Regarding  $y$  as 0 in Lemma 3.5, we see that this corollary is a direct consequence of Lemma 3.5.  $\square$

Similarly, we have

**Corollary 3.8.** *Under the assumption of Lemma 3.6, there exists a small  $\rho > 0$  such that if (3.15) holds, then for any  $y' \in B'_\rho(0)$  and  $y = (y, 0)$ , the function  $w^{y,-}(x)$  defined in (3.7) is the lower barrier of  $v$  at  $y$  to the linearized problem (3.4). This also implies  $u^y(y + te_n) \geq ct^2$  for small  $t > 0$  if  $u(y) = 0$  and  $\nabla u(y) = 0$ .*

#### 4. STRICT CONVEXITY LEMMA AND EXAMPLES

Recalling that if  $u \in C(\overline{B_1^+(0)})$  is a convex function and satisfies

$$(4.1) \quad \lambda \leq \det D^2 u \leq \Lambda \text{ in } B_1^+(0),$$

we can naturally *define*

$$D_n u(x', 0) = \overline{\lim}_{t \rightarrow 0^+} D_n u(x', t), \quad \forall x' \in B_1'(0).$$

See Lemma 3.2 for the proof. In this section, we first study the strict convexity for such functions. Then we provide an example showing that the growth assumption for the strict convexity is optimal.

**Lemma 4.1.** *Given positive constants  $H, M_1$  and  $M_2$ . Suppose that  $u \in C(\overline{B_1^+(0)})$  is a convex function satisfying (4.1) and*

$$D_n u(x', 0) \geq -M_2 \text{ in } B_1'(0).$$

*There exists  $\sigma = \sigma(H, M_1, M_2, \lambda, \Lambda, n) > 0$  such that if  $u$  satisfies*

$$0 \leq u(x_1, x'', 0) \leq \sigma + M_1 |x''|^2 \text{ in } B_1'(0),$$

*then*

$$\sup_{B_1^+(0)} u \geq H.$$

*Proof.* Assume by way of contradiction that there exists  $\sigma_k \rightarrow 0^+$  and convex functions  $u_k$  defined on  $B_1^+(0) \subset \mathbb{R}^n$ , satisfying

$$0 < \lambda \leq \det D^2 u_k \leq \Lambda \text{ in } B_1^+(0), \quad D_n u_k(x', 0) \geq -M_2 \text{ in } B_1'(0),$$

$$0 \leq u_k(x_1, x'', 0) \leq \sigma_k + M_1 |x''|^2 \text{ in } B_1'(0),$$

and

$$\sup_{B_1^+(0)} u_k \leq H.$$

Applying Lemma 2.5 we see that  $u_k$  is locally Lipschitz with

$$\|u_k\|_{Lip(\overline{B_{4c}^+(0)})} \leq C(H + M_2).$$

Hence,

$$u_k(x) \leq \sigma_k + M_1 |x''|^2 + C(H + M_2) |x_n| \text{ in } B_{4c}^+(0).$$

Let  $h_k = \sigma_k^{\frac{1}{2}}$  and  $\mathcal{D}_k = \text{diag}(1, h_k^{\frac{1}{2}} \mathcal{I}'', h_k)$ . Consider the functions

$$v_k(x) = \frac{u_k(\mathcal{D}_k x)}{h_k}, \quad x \in \Omega_k := \mathcal{D}_k^{-1} B_{4c}^+(0).$$

Then  $0 < \lambda \leq \det D^2 v_k \leq \Lambda$  in  $\Omega_k$ ,

$$v_k(x) \leq \sigma_k^{\frac{1}{2}} + M_1 |x''|^2 + C(H + M_2) |x_n| \text{ in } \mathcal{D}_k^{-1} B_1^+(0)$$

and

$$D_n v_k(x', 0) \geq -M_2 \text{ on } \mathcal{D}_k^{-1} B_1'(0).$$

Similar to Lemma 2.5, for any given  $R > 0$ , we can prove that

$$\|v_k\|_{Lip(B_R(0) \cap \Omega_k)} \leq C(H + M_1 + M_2) R^2.$$

Therefore,  $v_k$  is locally uniformly Lipschitz, and there is a subsequence that locally uniformly converges to a convex function  $v$  defined on  $\Omega := [-4c, 4c] \times \mathbb{R}^{n-1}$ . Moreover, we have

$$(4.2) \quad 0 < \lambda \leq \det D^2 v \leq \Lambda,$$

and

$$(4.3) \quad v(x) \leq M_1 |x''|^2 + C(H + M_2) |x_n| \text{ in } B_{4c}^+(0), \quad D_n v(x', 0) \geq -M_2 \text{ on } B_R'(0).$$

Furthermore, (4.3) implies that

$$(4.4) \quad a(x_1) = \lim_{t \rightarrow 0^+} \frac{v(x_1, 0, t)}{t}$$

is well-defined, and

$$-M_2 \leq a(x_1) \leq C(H + M_2).$$

Again, we can consider the blow-up of  $v$  along axis  $e_1$  by taking

$$\bar{h}_k = 2^k, \quad \bar{\mathcal{D}}_k := \text{diag}(1, \bar{h}_k^{\frac{1}{2}} \mathcal{I}'', \bar{h}_k), \quad \bar{v}_k(x) = \frac{v(\bar{\mathcal{D}}_k x)}{\bar{h}_k}, \quad x \in \bar{\mathcal{D}}_k^{-1} \Omega.$$

The limit  $\bar{v}$  still satisfies (4.2), (4.3). And (4.4) gives

$$\bar{v}(x_1, 0, x_n) = a(x_1) x_n.$$

Obviously,  $\nabla \bar{v}(e_n) \cdot e_n = a(x_1)$ . The non-negative function

$$w(x) = \bar{v}(x) - \bar{v}(e_n) - \nabla \bar{v}(e_n) \cdot x,$$

is zero on the line  $te_n$ . This is,  $w$  is bounded on a convex domain with infinite volume. However,  $0 < \lambda \leq \det D^2 w \leq \Lambda$ , which contradicts with Lemma 2.11. □

**Lemma 4.2.** *Given positive constants  $H$  and  $M_1$ . Suppose  $u \in C(\overline{B_1(0)})$  is a convex function satisfying*

$$\lambda \leq \det D^2 u \leq \Lambda \text{ in } B_1(0).$$

*There exists  $\sigma = \sigma(H, M_1, M_2, \lambda, \Lambda, n) > 0$  such that if  $u$  satisfies*

$$0 \leq u(x_1, x'', 0) \leq \sigma + M_1 |x''|^2 \text{ on } B_1'(0),$$

then

$$\sup_{B_1(0)} u \geq H.$$

*Proof.* Assume by way of contradiction that  $\sup_{B_1(0)} u < H$ . Then, the assumption of Lemma 4.2 immediately implies that  $u \in Lip(B_c(0))$  and  $|\nabla' u(0)| \leq C(\sigma + \sigma^{\frac{1}{2}})$  (See Lemma 2.6). After subtracting the supporting function  $u(0) + \nabla u(0) \cdot x$  at 0,  $u$  satisfies the assumption of Lemma 4.1 with  $M_2 = \sigma + \sigma^{\frac{1}{2}}$ . Applying Lemma 4.1 we obtain a contradiction, completing the proof.  $\square$

We point that the constant  $\sigma$  in the growth assumption of Lemma 4.1 and 4.2 can not be allowed to be zero. Otherwise, the constant  $H$  may tend to  $\infty$ , yielding a contradiction. Moreover, the following example shows that the growth exponent 2 is optimal.

**Example 4.3.** Suppose  $n \geq 3$ , constants  $1 < a, b < \infty$  and  $\delta > 0$  is small such that

$$\frac{1}{a} = \frac{1}{2} + \frac{1}{n-2}\delta, \quad 1 - \delta = \frac{1}{b}.$$

Let

$$E_1 := \{x' \in \mathbb{R}^{n-1} : |x''|^a \geq |x_1|^b\}, \quad E_2 := \{x' \in \mathbb{R}^{n-1} : |x_1|^b \geq |x''|^a\}.$$

Consider the function

$$(4.5) \quad W_{a,b}(x') = \begin{cases} |x''|^a + |x''|^{a-\frac{2a}{b}}|x_1|^2, & x \in E_1 \\ \frac{2b+a-ab}{b}|x_1|^b + \frac{ab-a}{b}|x_1|^{b-\frac{2b}{a}}|x''|^2, & x \in E_2 \end{cases}$$

and

$$(4.6) \quad W(x) = (1 + x_n^2)W_{a,b}(x').$$

We will show that there is a  $\rho = \rho(\delta) > 0$  such that  $W$  is a convex function in  $\mathbb{R}^{n-1} \times (-\rho, \rho)$  satisfying

$$(4.7) \quad c(\delta) \leq \det D^2W \leq C(\delta), \quad D_n W(x', 0) = 0 \text{ in } \mathbb{R}^{n-1} \times (-\rho, \rho).$$

*Proof.* Notice that  $W \in C^1(\mathbb{R}^n) \cap C^2(\mathbb{R}^n \setminus \{x' = 0\})$ ,  $\nabla W(\{x' = 0\}) \subset \{x' = 0\}$  and the  $n$ -dim Hausdroff measure of the set  $\nabla W(\{x' = 0\})$  is zero. The second equation in (4.7) is obvious. Hence, it is enough to show that  $W$  is convex and satisfies the first inequality in (4.7) in the sense of (Alexandrov) generalized solution. For this purpose, we need to calculate  $\det D^2W$  at any point  $x' \neq 0$ .

It is easy to see that

$$D^2W = \begin{bmatrix} (1 + x_n^2)D'^2W_{a,b} & 2x_n D'W_{a,b} \\ 2x_n D'W_{a,b} & 2W_{a,b} \end{bmatrix}.$$

So the convexity of  $W$  and  $\det D^2W$  is reduced to the study of the matrix

$$\mathcal{A} = \begin{bmatrix} (1+x_n^2)D'^2W_{a,b} - 2W_{a,b}^{-2}x_n^2D'W_{a,b} \otimes D'W_{a,b} & 0 \\ 0 & 2W_{a,b} \end{bmatrix}.$$

First, we assume  $x' \in E_1$ . Then

$$|x''|^a \leq W_{a,b} \leq 2|x''|^a, \quad \forall x' \in E_1.$$

Set the matrix functions

$$\mathcal{K}'_1(x'') = \text{diag}\{|x''|^{a-\frac{2a}{b}}, |x''|^{a-2}\mathcal{I}''\}, \quad \mathcal{K}_1(x'') = \text{diag}\{\mathcal{K}'_1, |x''|^a\}.$$

Then by the assumption on  $a$  and  $b$  we have

$$(4.8) \quad \det \mathcal{K}_1 = |x''|^{2a(\frac{n}{2}-\frac{1}{b}-\frac{n-2}{a})} \equiv 1.$$

Observing that  $C(a, b)$  can be written as  $C(\delta)$ , we have

$$0 \leq 2W_{a,b}^{-2}D'W_{a,b} \otimes D'W_{a,b} \leq C(\delta)\mathcal{K}'_1, \quad D'^2W_{a,b} \leq C(\delta)\mathcal{K}'_1, \quad \forall x' \in E_1.$$

Hence, in  $E_1$  we have

$$(4.9) \quad C(\delta)\mathcal{K}_1 \geq \mathcal{A} \geq \begin{bmatrix} (1+x_n^2)D'^2W_{a,b} - x_n^2C(\delta)\mathcal{K}'_1 & 0 \\ 0 & C(\delta)|x''|^a \end{bmatrix}.$$

Since  $|x_n| \leq \rho$  will be small, by (4.8)-(4.9) we can conclude the convexity of  $W$  and the first inequality in (4.7) after we prove

$$(4.10) \quad D'^2W_{a,b} \geq \epsilon(\delta)\mathcal{K}'_1, \quad \forall x' \in E_1.$$

Denote

$$\mathcal{B}_1 = \sqrt{\mathcal{K}'_1}, \quad X = |x''|^{-1}x'', \quad y = |x_1||x''|^{-\frac{a}{b}}$$

and

$$m = a[\mathcal{I}'' + (a-2)X \otimes X] + (a - \frac{2a}{b})[\mathcal{I}'' + (a - \frac{2a}{b} - 2)X \otimes X]y.$$

By a direct calculation we have

$$(4.11) \quad \mathcal{B}_1^{-1}D'^2W_{a,b}\mathcal{B}_1^{-1} = \begin{bmatrix} 2 & 2(a - \frac{2a}{b})yX \\ 2(a - \frac{2a}{b})yX^T & m \end{bmatrix}.$$

Observe that

$$(4.12) \quad m - 2(a - \frac{2a}{b})^2|y|^2X \otimes X = f_1\mathcal{I}'' - f_2X \otimes X$$

where,  $f_1 = 2a - \frac{2a}{b}$  and

$$f_2 = 2(a - \frac{2a}{b})^2y^2 - (a - \frac{2a}{b})(a - \frac{2a}{b} - 2)y^2 - a(a-2).$$

Notice that

$$\begin{aligned} 2\left(a - \frac{2a}{b}\right)^2 - \left(a - \frac{2a}{b}\right)\left(a - \frac{2a}{b} - 2\right) &= \left(a - \frac{2a}{b}\right)\left(a - \frac{2a}{b} + 2\right) \\ &= a\left(a - \frac{2a}{b}\right)\left(\frac{1}{n-2} - 1\right)\delta \\ &\geq 0. \end{aligned}$$

Obviously,  $|y| \leq 1$  in  $E_1$ . Recall  $\delta$  is small,  $a \geq \frac{2a}{b} - a \geq \frac{7}{4} > \frac{3}{2}$ , thus we get

$$f_1 - f_2 \geq \left(a - \frac{1}{2}\right)^2 - \left(\frac{2a}{b} - a - \frac{1}{2}\right)^2 \geq c\left[a - \left(\frac{2a}{b} - a\right)\right] \geq c\delta a.$$

Since

$$\mathcal{I}'' - X \otimes X \geq 0,$$

we obtain

$$m - 2\left(a - \frac{2a}{b}\right)^2 |y|^2 X \otimes X \geq ca\delta \mathcal{I}''.$$

Replacing this in (4.12) and using (4.10)-(4.11), we finally obtain (4.10).

Second, we assume  $x \in E_2$ . Then we have

$$c(\delta)|x_1|^b \leq W_{a,b} \leq C(\delta)|x_1|^b.$$

Denote

$$\mathcal{K}'_2 = \text{diag}\{|x_1|^{b-2}, |x_1|^{b-\frac{2b}{a}} \mathcal{I}''\} \quad \text{and} \quad \mathcal{K}_2 = \text{diag}\{\mathcal{K}'_2, |x_1|^b\}.$$

Then

$$\det \mathcal{K}_2 = |x_1|^{2b\left(\frac{n}{2} - \frac{1}{b} - \frac{n-2}{a}\right)} \equiv 1.$$

Similar to the first case in  $E_1$ , it is enough to show that

$$(4.13) \quad D'^2 W_{a,b} \geq \epsilon(\delta) \mathcal{K}'_2 \quad \text{in } E_2.$$

Denote

$$\mathcal{B}_2 = \sqrt{\mathcal{K}'_2}, \quad Y = |x''|^{-1} x'', \quad y = |x_1|^{-\frac{b}{a}} |x''|$$

and

$$d = (2 - a\delta)b(b-1) + a\delta\left(b - \frac{2b}{a}\right)\left(b - \frac{2b}{a} - 1\right)y^2.$$

By a direct calculation we obtain

$$(4.14) \quad \mathcal{B}_2^{-1} D'^2 W_{a,b} \mathcal{B}_2^{-1} = \begin{bmatrix} d & 2a\delta\left(b - \frac{2b}{a}\right)yY \\ 2a\delta\left(b - \frac{2b}{a}\right)yY & 2a\delta \mathcal{I}'' \end{bmatrix}.$$

Write

$$d - 2a\delta\left(b - \frac{2b}{a}\right)^2 y^2 |Y|^2 = g_1 - g_2 y^2,$$

where

$$\begin{aligned} g_2 &= a\delta\left(b - \frac{2b}{a}\right)\left(b - \frac{2b}{a} - 1\right) - 2a\delta\left(b - \frac{2b}{a}\right)^2 \\ &= \delta b^2(2-a)\left(\frac{2}{a} + \frac{1}{b} - 1\right) \\ &\geq 0 \end{aligned}$$

Obviously,  $|y| \leq 1$  in  $E_1$ . If  $\delta > 0$  is small, we have

$$\begin{aligned} d - 2a\delta\left(b - \frac{2b}{a}\right)^2 y^2 |Y|^2 &\geq b^2\delta\left[(2-a\delta) + a\frac{2\delta}{n-2}\left(1 - \frac{2}{a} - \frac{1}{b}\right) - 2a\left(\frac{2\delta}{n-2}\right)^2\right] \\ &\geq cb^2\delta, \end{aligned}$$

which, together with (4.14), implies the desired (4.13).  $\square$

**Remark 4.4.** *Suppose that  $R > 0$ ,  $W$  is given by (4.5)-(4.6), and consider the Dirichlet problem*

$$(4.15) \quad \begin{cases} \det D^2 W_R^+ = c(\delta) & \text{in } B'_R(0) \times [-\rho, \rho] \\ W_R^+ = W & \text{on } \partial(B'_R(0) \times [-\rho, \rho]). \end{cases}$$

*It is known that there exists a convex solution  $W_R^+$ , which is still symmetric about  $x_n = 0$  since so is  $W$ . This is to say*

$$D_n W_R^+(x', 0) = 0.$$

*Comparison principle (Lemma 2.9) implies*

$$W_R^+ \geq W \geq 0$$

*if  $c(\delta) > 0$  is small. The convexity means*

$$W_R^+(x', t) \leq \max\{W_R^+(x', \rho), W_R^+(x', -\rho)\} \leq 4W(x', 0), \quad x \in B'_R(0), t \in [-\rho, \rho].$$

*and it follows that*

$$W_R^+(te_n) = 0.$$

*This mean  $W_R^+$  has the same asymptotic behavior around 0 as  $W$ .*

## 5. GOOD SHAPE LEMMA

In this section, we assume that  $u$  is a viscosity solution to problem (1.1), satisfying (2.6) and the hypothesis of Theorem 1.1. We are going to prove that its height section  $S_h := S_h^u(0)$  (defined by (2.4)) is of good shape for large  $h$ , then we will introduce the normalization family  $(u_h, S_1^h)$  of  $(u, S_h)$ .

At first, we mention here that the good shape property is invariant under any linear transform of the form  $\mathcal{A} = \text{diag}\{\mathcal{A}', a_n\}$  with  $c \leq a_n$ ,  $\det \mathcal{A}' \leq C$ .

Let  $p_h$  be the mass center of  $S_h$ , and consider the *sliding transformation*

$$\mathcal{A}_h^c x = x - \sum_{i=1}^{n-1} \frac{p_h \cdot e_i}{p_h \cdot e_n} x_n e_i.$$

Take  $y_h \in \overline{\mathcal{P}(\mathcal{A}_h^c S_h)}$  such that

$$|y_h \cdot e_1| = \sup\{x \cdot e_1 \mid x \in \mathcal{P}(\mathcal{A}_h^c S_h)\}$$

and consider another sliding transformation

$$\mathcal{A}_h x = x - \sum_{i=2}^{n-1} \frac{y_h \cdot e_i}{y_h \cdot e_1} x_1 e_i.$$

Apply John's lemma (Lemma 2.8) to  $\{x'' \in \mathbb{R}^{n-2} \mid (x_1, x'', 0) \in \mathcal{A}_h \mathcal{P}(\mathcal{A}_h^c S_h)\}$ , we can find a new coordinates  $\{e_i\}_{i=1}^n$  which keeps  $e_1$  and  $e_n$  (or only  $e_n$  for  $\mathcal{A}_h^c$ ) invariant such that

$$\mathcal{D}_h B_1(q_h) \subset \mathcal{A}_h \mathcal{A}_h^c S_h \subset \mathcal{D}_h B_{C(n)}(q_h) \subset \mathcal{D}_h B_{2C(n)}^+(0),$$

where  $q_h = \mathcal{A}_h \mathcal{A}_h^c p_h$ ,  $\mathcal{D}_h := \text{diag}(d_1(h), \dots, d_n(h))$  is the diagonal matrix satisfying

$$\prod_{i=1}^n d_i(h) = \det \mathcal{D}_h = \text{Vol}(S_h).$$

It follows from Lemma 2.11 that  $\text{Vol}(S_h) \leq Ch^{\frac{n}{2}}$ . Denote

$$\tau_h = \frac{(\det \mathcal{D}_h)^{\frac{1}{n}}}{h^{\frac{1}{2}}}, \quad \bar{\mathcal{D}}_h = \tau_h^{-1} \mathcal{D}_h := \text{diag}(\bar{d}_1(h), \dots, \bar{d}_n(h)).$$

We first consider the rescaling  $(u_h, S_1^h)$  of  $(u, S_h)$  given by

$$(5.1) \quad u_h(x) = \frac{u(\mathcal{A}_h^{-1} \bar{\mathcal{D}}_h x)}{h}, \quad x \in S_1^h := \bar{\mathcal{D}}_h^{-1} \mathcal{A}_h S_h,$$

then we have

$$(5.2) \quad \mathcal{A}_h^* B_{\tau_h}(x_h) \subset S_1^h \subset \mathcal{A}_h^* B_{C(n)\tau_h}(x_h) \subset \{x_n \leq 2\tau_n C(n)\}$$

for some  $x_h$ , where

$$\mathcal{A}_h^* = \bar{\mathcal{D}}_h^{-1} \mathcal{A}_h \mathcal{A}_h^c \mathcal{A}_h^{-1} \bar{\mathcal{D}}_h$$

is still a sliding transformation of the form  $\mathcal{A}_h^* x = x - \sum_{i=1}^{n-1} \beta_i^h x_n e_i$ .

We use the notation  $G_1^h := S_1^h \cap \partial \mathbb{R}_+^n$  similar to (2.5). And we will simplify  $S_t^h, G_t^h$  by  $S_t, G_t$  when the dependence on  $h$  and  $u_h$  is self-evident from the context. In Section 7, we will use the corresponding notations  $S_t^R, G_t^R$  for the  $(u_R, S_1^R)$  as we just defined  $S_1^h, G_1^h$  for  $(u, S_1)$ .

Now, the Neumann problem (1.1) becomes

$$(5.3) \quad \begin{cases} \det D^2 u_h = 1 & \text{in } S_1^h \\ D_n u_h = a_h x_1 & \text{on } G_1^h \\ u_h = 1 & \text{on } \partial S_1^h \setminus G_1^h \end{cases}$$

where

$$(5.4) \quad a_h = \frac{\bar{d}_1(h) \bar{d}_n(h)}{h} a.$$

Recall that if  $n \geq 3$ ,  $u \in \Theta$  and (2.2) is satisfied. Let

$$\delta_h = \frac{K}{h}, \quad \bar{Q}_h = \frac{C \bar{D}_h''^T I'' \bar{D}_h''}{h}.$$

We have

$$(5.5) \quad u_h(0, z, 0) \leq z^T \bar{Q}_h z + \delta_h, \quad \forall z \in \mathbb{R}^{n-2}$$

and

$$(5.6) \quad a_h^2 \det \bar{Q}_h = C^n \tau_h^{-2n} \frac{(\det \mathcal{D}_h)^2}{h^n} a^2 \leq C.$$

If  $h \geq 2K$ , (2.2) also implies  $d_i(h) \geq ch^{\frac{1}{2}}$  for  $2 \leq i \leq n-1$ . More precisely,

$$\prod_{i=2}^{n-1} \bar{d}_i(h) \geq C \tau_h^{-\frac{n-2}{2}} \text{Vol}_{n-2}(S_h(0) \cap \mathbb{R}^{n-2}) \geq Ch^{\frac{n-2}{2}}.$$

Therefore  $\det \bar{Q}_h \geq c$  and  $0 \leq a_h \leq C$ . To conclude, we have obtained that

$$(5.7) \quad 0 \leq a_h \leq C, \quad u_h \in \Theta(\delta_h), \quad \forall h \geq 2K.$$

The Good Shape Lemma is stated as follows.

**Theorem 5.1** (Good Shape). *For large  $h$ , the normalization  $(u_h, S_1^h)$  given by (5.1) satisfies*

$$(5.8) \quad \mathcal{P}S_1^h(0) \subset C\mathcal{P}G_1^h(0) \text{ and } S_1^h(0) \subset B_{C(n)\tau_h}(0),$$

$$(5.9) \quad c \leq \text{Vol}(S_1^h(0)) \leq C \text{ and } c \leq \tau_h \leq C,$$

$$(5.10) \quad \mathcal{P}G_1^h(0) \subset -C\mathcal{P}G_1^h(0).$$

We are going to prove the theorem by the coming three lemmas.

**Lemma 5.2.** *The normalization  $(u_h, S_1^h)$  given by (5.1) satisfies (5.8) for large  $h$ . More precisely,*

$$\mathcal{P}S_1^h(0) \subset (1 + C|a_h|)\mathcal{P}G_1^h(0) \subset C\mathcal{P}G_1^h(0).$$

*Proof.* If there exists a unit direction  $e, e \perp e_n$  such that

$$\frac{\sup\{s \mid se \in \mathcal{P}S_1^h(0)\}}{\sup\{s \mid te \in \mathcal{P}G_1^h(0), \forall t \in (0, s)\}} \geq M > 1.$$

We assume that the maximum values are attained at points  $y$  and  $z$  respectively:

$$y = y_e e + y_n e_n \in \overline{\partial S_1^h(0)}, \quad z = z_e e \in \overline{\partial G_1^h(0)}.$$

It is clear that

$$(5.11) \quad u(z) = u(y) = 1, \quad y_e \geq M z_e, \quad 0 < y_n \leq C,$$

where we have used (5.2) for the last inequality.

Since the transforms  $\mathcal{A}_h^*$  leaves  $\mathbb{R}^{n-1}$  invariant, we have  $|z| \leq C$ . Let

$$l_{y,0} = \{sy, s \in (0, 1)\},$$

$$l_z^{e_n} = \{z + te_n, t \in [0, \infty)\}.$$

There exists a point  $p = sy = z + ty_n e_n \in l_{y,0} \cap l_z^{e_n}$  with  $s, t \in (0, \frac{1}{M}]$ . By convexity, (2.6) and (5.11) we have

$$u(p) \leq su(y) + (1-s)u(0) \leq \frac{1}{M}.$$

However, Lemma 3.2, (5.11), the convexity again, and the fact  $|z| \leq C$  implies that

$$u(p) \geq u(z) + ty_n D_n u(z) \geq 1 + ta_n y_n z \cdot e_1 \geq 1 - Ct|a_h|y_n \geq 1 - \frac{C|a_h|}{M}.$$

These two inequalities mean that  $M \leq 1 + C|a_h| \leq C$  by (5.7), which implies  $S_1^h(0) \subset B_{C(n)\tau_h}(0)$ . Combining this and (5.2) we have

$$S_1^h \subset \{x_n \leq C\tau_h\} \cap \{CG_1^h \times [0, \infty)\} \subset B_{C\tau_h}(0).$$

□

**Lemma 5.3.** *The normalization  $(u_h, S_1^h)$  given by (5.1) satisfies (5.9) for large  $h$ .*

*Proof.* Since  $\det \mathcal{D}_h = \text{Vol}(S_h) \leq Ch^{\frac{n}{2}}$  by Lemma 2.11, we see that  $\tau_h \leq C$  by its definition. Notice that

$$\tau_h^n \sim \text{Vol}(S_1^h(0)) = \tau_h^n \mathcal{A}_h \sim \tau_h^n.$$

It is enough for (5.9) to prove that  $\tau_h \geq c$ .

Consider the function  $v$  defined by

$$v(x) = \tau_h^{-1} \left( \frac{|x|^2}{8C_n^2} + \frac{x_n}{2C_n} \right).$$

If  $\tau_h > 0$  is small, by (5.3), (5.7) and (5.8) we have

$$v \leq \tau_h^{-1} \left( \frac{\tau_h^2}{8} + \frac{\tau_h}{2} \right) < 1 \leq u_h \text{ on } \partial S_1^h \setminus G_1^h,$$

$$\det D^2 v \geq c \tau_h^{-n} > 1 = \det D^2 u_h \text{ in } S_1^h,$$

and

$$D_n v \geq \frac{1}{2C_n \tau_h} > C \geq D_n u_h \text{ on } G_1^h.$$

By comparison principle (Lemma 2.10)  $v < u_h$  in  $S_1^h$ , which contradicts  $v(0) = 0 = u_h(0)$ . Therefore  $\tau_h \geq c$ . □

**Lemma 5.4.** *The normalization  $(u_h, S_1^h)$  given by (5.1) satisfies (5.10) for large  $h$ .*

*Proof.* Lemmas 5.2 and 5.3 mean that  $\tau_h$  is between two positive universal constants, and

$$(5.12) \quad \mathcal{P}S_1^h \subset C\mathcal{P}G_1^h \text{ and } B_c(x_h) \subset S_1^h \subset B_C(0)$$

for some  $x_h \in R_+^n$ . Therefore, we only need to show  $\mathcal{P}G_1^h$  is balance about 0, this is

$$(5.13) \quad \mathcal{P}G_1^h \supset B'_c(0).$$

Let  $M \geq 4n^{4n}C^{4n}$  is large universal,  $\theta = M^{-\frac{1}{n+1}}$ ,  $m = M^{-1}$  and  $\delta = \frac{1}{M^3}$  is small enough. On the contrary to (5.13), there would exist a  $p = (p_1, p'') \in \partial\mathcal{P}G_1^h$  such that

$$(5.14) \quad |p| = \text{dist}(0, p) = \text{dist}(0, \partial\mathcal{P}G_1^h) \leq \delta.$$

Without loss of generality, we may assume that  $p_1 \geq 0$ . There are two cases:

Case 1:  $|a_h| \geq CM^{-1}$ . Then we have

$$\max\{d_2(h), \dots, d_{n-1}(h)\} \leq CMh^{\frac{1}{2}},$$

which means that

$$B''_{cM^{-1}}(0) \subset \mathcal{P}S_1^h(0) \subset B'_C(0).$$

By convexity and (5.14), we get

$$\sup\{x_1 | x \in G_1^h\} \leq CM\delta \leq \frac{C}{M}.$$

Therefore, (5.7) and the Neumann condition in (5.3) implies

$$|D_n u_h(x', 0)| \leq \frac{C}{M}, \text{ on } x_1 \geq -\frac{1}{M}.$$

Case 2:  $|a_h| \leq CM^{-1}$ . Then

$$|D_n u_h(x', 0)| \leq \frac{C}{M}, \text{ on } G_1^h.$$

Next, we consider a new coordinate with zero as based point, axis  $e_1 = p/|p|$  and  $e_i (i \geq 2)$  unchanged. For simplicity, we still use the notation  $(e_1, e_2, \dots, e_n)$ .

In both cases, consider the domain  $E = \{x \in S_1^h | x_1 > -\frac{1}{M}\}$ . Then  $\partial E = E_1 \cup E_2 \cup E_3$ , where

$$\begin{aligned} E_1 &= \overline{E} \cap \{x_1 = -\frac{1}{M}\}, \\ E_2 &= \overline{E} \cap G_1^h \cap \{x_1 > -\frac{1}{M}\}, \\ E_3 &= \overline{E} \cap (\partial S_1^h \setminus G_1^h). \end{aligned}$$

For the function

$$v(x) = \left[ Q(Mx_1) + \theta \sum_{i=2}^n Q\left(\frac{x_i}{C}\right) \right]$$

where  $Q(t) = \frac{1}{4nC^2}(\frac{t^2}{2} + 2t)$ , we claim that

$$(5.15) \quad \begin{cases} \det D^2 u_h < \det D^2 v & \text{in } E \\ D_{-x} u_h < D_{-x} v & \text{on } E_1 \\ D_n u_h < D_n v & \text{on } E_2 \\ v < u_h & \text{on } E_3 \end{cases}.$$

By comparison principle (Lemma 2.10) we have  $v < u_h$  in  $E$ . This would contradict  $v(0) = u_h(0)$  and hence (5.13) is proved.

Finally, let us verify (5.15). First, by convexity and (2.6) we see that

$$D_{-x} u_h = -Du_h \cdot x \leq u_h(0) - u_h(x) \leq 0$$

holds in viscosity sense. Applying this, the results of Case 1 and 2, and the largeness of  $M$ , we have:

$$\text{in } E, \det D^2 v \geq CM^2 \theta^{n-1} > 1 = \det D^2 u_h;$$

$$\text{on } E_1, D_{-x} v \geq c - \theta \sum_{i=2}^{n-1} C > 0 \geq D_{-x} u_h;$$

$$\text{on } E_2, D_n v \geq c\theta > \frac{c}{M} \geq D_n u_h;$$

$$\text{on } E_3, v \leq \frac{3}{4} < 1 = u_h. \quad \square$$

In this way, we have proved Theorem 5.1. Since  $c \leq \tau_h \leq C$ , we can still take

$$\mathcal{D}_h = \text{diag}\{d_1(h), \dots, d_n(h)\}$$

in (5.1) instead of  $\bar{\mathcal{D}}_h$ . This yields the definition of normalization family.

**Definition 5.5.** *The normalization family  $(u_h, S_1^h)$  of  $(u, S_h)$  is defined as*

$$(5.16) \quad u_h(x) = \frac{u(\mathcal{D}_h \mathcal{A}_h^{-1} x)}{h}, \quad x \in S_1^h := \mathcal{A}_h \mathcal{D}_h^{-1} S_h.$$

By (2.1), (5.3)-(5.10), we have the following properties for large  $h$ :

$$(5.17) \quad B_c^+(0) \subset S_1^h \subset B_C(0),$$

$$(5.18) \quad \begin{cases} \det D^2 u_h = 1 & \text{in } S_1^h \\ D_n u_h = a_h x_1 & \text{on } G_1^h \\ u_h = 1 & \text{on } \partial S_1^h \setminus G_1^h \end{cases}$$

where

$$(5.19) \quad a_h = \frac{d_1(h)d_n(h)}{h}a,$$

$$(5.20) \quad u_h(0, z, 0) \leq z^T \mathcal{Q}_h z + \delta_h$$

where

$$(5.21) \quad \delta_h = \frac{K}{h}, \quad \bar{\mathcal{Q}} = \frac{C\mathcal{D}_h''^T I'' \mathcal{D}_h''}{h},$$

and

$$(5.22) \quad 0 \leq a_h \leq C \text{ and } a_h^2 \det \mathcal{Q}_h \leq C.$$

Moreover, Theorem 5.1 still holds for the normalization family  $(u_h, S_1^h)$ .

## 6. STATIONARY THEOREM

In this section, we study the normalization family  $(u_h, S_1^h)$  given by (5.16) in the case  $a_h$  is small. For simplify we omit  $h$  and write it by  $(u, S_1)$ . This leads us to study the standard model

$$(6.1) \quad \begin{cases} \det D^2 u = 1 & \text{in } S_1 \\ D_n u = a x_1 & \text{on } G_1 \\ u = 1 & \text{on } \partial S_1 \setminus G_1 \end{cases}$$

and

$$(6.2) \quad u(0) = 0, \quad \nabla u(0) = 0, \quad u \geq 0 \text{ in } S_1, \quad B_c^+(0) \subset S_1 \subset B_C(0).$$

For every  $a \in R$ , set

$$(6.3) \quad Q_a(x) := \frac{1}{2} \left[ \sqrt{1 + a^2(x_1^2 + x_n^2)} + 2ax_1x_n + \sum_{i=2}^{n-1} x_i^2 \right].$$

Obviously,  $\det D^2 Q_a(x) \equiv 1$  and  $D_n Q_a(x', 0) = ax_1$ .

We will prove the following

**Theorem 6.1.** *Suppose that  $u \in \overline{(S_1)}$  is a viscosity solution to problem (6.1) and (6.2) is satisfied. There exists universal constants  $b_0 > 0$ ,  $c_0 > 0$  such that if*

$$(6.4) \quad \|u - Q_a\|_{L^\infty(S_1^{Q_a}(0))} + \|D_n u - D_n Q_a\|_{L^\infty(S_1^{Q_a}(0))} \leq b_0,$$

then  $u \in C^{2,\alpha}(\overline{B_{c_0}^+(0)})$  for some  $\alpha > 0$  and

$$\|u\|_{C^{2,\alpha}(\overline{B_{c_0}^+(0)})} \leq C.$$

With this theorem, we will also show

**Theorem 6.2.** *Suppose that  $u \in \overline{(S_1)}$  is a viscosity solution to problem (6.1) and (6.2) is satisfied. There exists universal constants  $\delta_0 > 0$ ,  $c_0 > 0$  such if  $a \in (-\delta_0, \delta_0)$ , then  $u \in C^{2,\alpha}(\overline{B_{c_0}^+(0)})$  for some  $\alpha > 0$  and*

$$(6.5) \quad \|u\|_{C^{2,\alpha}(\overline{B_{c_0}^+(0)})} \leq C.$$

Denote

$$\mathcal{F}_a^*(x) := \{Q \mid Q(x) = \sum_{i,j=1}^n a^{ij} x_i x_j \text{ is a quadratic polynomial,}$$

and  $Q$  solves the first and second equations of problem (6.1)\}

and

$$\mathcal{F}_a(x) := \{l(x') + Q(x) \mid Q \in \mathcal{F}_a^*, l(x') \text{ is a linear function, } l(0) = 0\}.$$

Theorem 6.1 relies on the following Lemma.

**Lemma 6.3.** *Let  $u \in C(\overline{S_1(0)})$  be a viscosity solution to problem (6.1), and satisfy (6.2). For any small constant  $\mu > 0$ , there exists a small  $\epsilon_0 = \epsilon_0(\mu) > 0$  such that if*

$$(6.6) \quad \|u - P\|_{L^\infty(S_1)} \leq \epsilon, \quad \|D_n u - D_n Q\|_{L^\infty(S_1)} \leq C\epsilon$$

for some  $\epsilon \in (0, \epsilon_0)$  and  $P(x) = l(x') + Q(x) \in \mathcal{F}_a$ , then

$$(6.7) \quad |Dl| \leq C\epsilon^{\frac{1}{2}},$$

and

$$(6.8) \quad |u - P^0| \leq C\epsilon\mu^{\frac{3}{2}}, \quad |D_n u - D_n Q^0| \leq C\epsilon\mu \text{ in } S_\mu^{Q^0}$$

for some  $P^0 = l^0 + Q^0 \in \mathcal{F}_a$ . Moreover,

$$(6.9) \quad |D^2 Q - D^2 Q^0| \leq 2\epsilon \text{ in } S_\mu^{Q^0}.$$

*Proof.* By virtue of the fact  $l(0) = 0$ , (6.6) gives  $|Dl| \leq C\epsilon$ . One can choose  $x = (x', 0) \in G_1$  such that

$$-l(x') = \frac{1}{n} |Dl| |x'| \text{ and } |x'| \sim \epsilon^{\frac{1}{2}}.$$

It follows from (6.2) and (6.6) that

$$\begin{aligned} \frac{1}{n}|Dl||x'| &= -l(x') \leq u(x', 0) - l(x') \\ &\leq |u(x, 0) - l(x') - Q(x', 0)| + |Q(x', 0)| \\ &\leq \epsilon + C|x'|^2 \leq C|x'|^2. \end{aligned}$$

Therefore  $|Dl| \leq C\epsilon^{\frac{1}{2}}$  and (6.7) holds.

In following discussion, we will replace  $u$  by  $u - l(x')$  and need only to consider the case  $P = Q$  in (6.6). (6.2) and (6.6) implies that  $c\mathcal{I} \leq D^2Q \leq C\mathcal{I}$ . For simplicity, we further assume  $Q = Q_a$  (see (6.3)). This is because we can always find a new coordinate with the aid of the sliding transform  $\mathcal{B}x := x - 2D''Q^{-1}D_{1i}Qx_1e_i$ , under which  $Q$  may be transformed to  $Q_a$ .

Let

$$(6.10) \quad u_0 := u - P = u - Q, \quad v_\epsilon := \frac{u_0}{\epsilon}, \quad \epsilon > 0.$$

In fact, we may assume  $\|u - Q\|_{L^\infty(S_1)} > 0$  and take  $\epsilon = \|u - Q\|_{L^\infty(S_1)}$ . We will show the system  $(v_\epsilon, D_nv_\epsilon)$  could be well approximated by  $(w, D_nw)$ , a solution of some linearized problem. More precisely, let  $F = S_{\frac{1}{2}}^Q(0)$  and

$$Lw := \sqrt{1 + a^2}(w_{11} + w_{nn}) + aw_{1n} + \Delta''w,$$

where  $\Delta'' := \sum_{i=2}^{n-1} \partial_{x_i}^2$ . We will show that as if  $\epsilon \rightarrow 0^+$ , the corresponding  $(v_\epsilon, D_nv_\epsilon)$  will have a subsequence converging uniformly to  $(w, D_nw)$ , a solution of the problem:

$$(6.11) \quad Lw = 0 \text{ in } F, \quad w(0) = 0, \quad D_nw = 0 \text{ on } \{x_n = 0\}.$$

**Step 1.** We first consider the convergence on compact subset  $E = \{x_n \geq C\epsilon^{\frac{1}{2}}\} \cap F$ , where  $\epsilon > 0$  is sufficiently small. For any point  $y$  such that  $B_{C\rho}(y) \subset E$  for some  $\rho \gg c\epsilon^{\frac{1}{2}}$ . Then  $B_{c\rho}(y) \subset S_{\rho^2}^Q(y) \subset B_{C\rho}(y)$  by the fact  $Q = Q_a$ . It follows from (6.1) that

$$(6.12) \quad 0 = \det D^2u - \det D^2Q = Tr(AD^2(u - Q)) = \epsilon Tr(AD^2v_\epsilon),$$

where

$$A = [A_{ij}]_{n \times n} := \int_0^1 cof((1-t)D^2Q + tD^2u)dt,$$

and  $cofM$  denotes the cofactor matrix of  $M$ .

Since (6.6) gives us  $|u - Q| \leq \epsilon$  in  $S_1$ , we see that

$$B_{2c\rho}(y) \subset S_{\rho^2/2}^u(y) \subset B_{C\rho}(y).$$

Hence, the classical interior estimate (Lemma 2.13 and its proof in [3]) means that

$$\|u\|_{C^k(B_{c\rho}(y))} \leq C\rho^{2-k} \text{ and } c\mathcal{I} \leq D^2u \leq C\mathcal{I} \text{ in } B_{c\rho}(y).$$

Therefore, the operator  $\bar{L}f := A_{ij}f_{ij}$  is uniformly elliptic with smooth coefficients. By (6.12), we obtain

$$(6.13) \quad \|v_\varepsilon\|_{C^k(B_{c\rho}(y))} \leq C\rho^{-k},$$

which means

$$|D^2u - D^2Q| \leq C\varepsilon\rho^{-2} \text{ and } |A - D^2Q| \leq C\varepsilon\rho^{-2} \text{ in } B_{c\rho}(y).$$

Let  $\varepsilon \rightarrow 0^+$ , we have  $(v_\varepsilon, D_nv_\varepsilon)$  converges to  $(w, D_nw)$  in  $B_{c\rho}(y)$ . By (6.10), we see that

$$w(0) = 0, \quad Lw = 0 \text{ in } B_{c\rho}(y).$$

**Step 2.** We give the uniform control of  $(v_\varepsilon, D_nv_\varepsilon)$  near  $\{x_n = 0\}$ , so as to show

$$(6.14) \quad |D_nv_\varepsilon(x)| \leq C|x_n| \text{ in } S_{\frac{1}{4}}(0)$$

and

$$(6.15) \quad \text{Osc}_{B_r(x)} v_\varepsilon \leq C \max\{r^{\frac{2}{3}}, \varepsilon\} \text{ for } x \in G_{\frac{1}{4}}.$$

Given  $y \in G_{\frac{1}{2}}(0)$ . It follows from (6.6) and Corollary 2.7 that

$$|\nabla u - \nabla P| \leq C\varepsilon^{\frac{1}{2}} \text{ in } S_{\frac{1}{2}}(y)$$

if  $\varepsilon$  is small enough. Hence, the function  $u^y(x)$  defined in (3.5) still satisfies

$$u^y(x) \geq c \text{ on } \partial S_{\frac{1}{4}}(y).$$

As in Lemma 3.4,  $V := D_nv_\varepsilon$  satisfies

$$(6.16) \quad \begin{cases} U^{ij}D_{ij}(\pm V) = 0 & \text{in } S_{\frac{1}{4}}(0) \\ |V| \leq C & \text{on } \partial S_{\frac{1}{4}}(0) \setminus G_{\frac{1}{4}}(0) . \\ \pm V = 0 & \text{on } G_{\frac{1}{4}}(0) \end{cases}$$

Notice that the second in (6.16) is equivalent to  $V \leq C$  or  $-V \leq C$ , which comes from (6.6). Similar to Corollary 3.7, we see that the function  $w^{y,+}(x) = C_1[w^y(x) + C_2x_n]$  defined in (3.7) is a supersolution for  $V$  and  $-V$  in (6.16) respectively.

Hence we get

$$|D_nv_\varepsilon(x)| = |V(x)| \leq w^{(x',0),+}(x', x_n) \leq Cx_n, \text{ in } S_{\frac{1}{4}}(0).$$

This proves (6.14).

To prove (6.15), we observe that (6.14) implies

$$(6.17) \quad |v_\varepsilon(x', 0) - v_\varepsilon(x', x_n)| \leq Cx_n^2 \text{ in } S_{\frac{1}{4}}(0).$$

It is enough to estimate  $|v_\varepsilon(p) - v_\varepsilon(q)|$  for  $p = (p', 0), q = (q', 0)$  in  $G_{\frac{1}{2}}(0)$ . Suppose that

$$|p' - q'| = r, \quad \rho = C \max\{r^{\frac{1}{3}}, \varepsilon^{\frac{1}{2}}\}, \quad A = p + \rho e_n, \quad B = q + \rho e_n.$$

As  $\rho \geq C\epsilon^{\frac{1}{2}}$ , (6.13) implies

$$|v_\epsilon(A) - v_\epsilon(B)| \leq \frac{r}{\rho},$$

which, together with (6.17), gives

$$|v_\epsilon(p) - v_\epsilon(q)| \leq C\rho^2 + \frac{r}{\rho} \leq C \max\{r^{\frac{2}{3}}, \epsilon\}.$$

In summary, we have proved (6.14) and (6.15). By this and the result of Step 1 we conclude that as  $\epsilon \rightarrow 0^+$ ,  $(v_\epsilon, D_n v_\epsilon)$  has a subsequence converging to a solution  $w$  of problem (6.11) in  $S_{\frac{1}{4}}(0)$  uniformly.

**Step 3.** The classical regular results then shows the function  $w \in C_{loc}^3$ . Since  $w(0) = 0$ ,  $D_n w(0) = 0$  and  $D' D_n w(0) = 0$  by (6.11), there exists a linear function

$$l^0(x) = \sum_{i=1}^{n-1} a_i x_i$$

and a quadratic function

$$R(x) = \sum_{1 \leq i, j \leq n-1} a_{ij} x_i x_j + b_n x_n^2$$

such that

$$|w(x) - l^0(x) - R(x)| \leq C|x|^3 \text{ and } |D_n w(x) - 2b_n x_n| \leq C|x|^2.$$

Here,  $a_i, a_{ij}, b_n$  are bounded by a universal constant, and

$$\sqrt{1 + a^2}(a_{11} + b_n) + \sum_{i=2}^{n-1} a_{ii} = 0$$

by (6.11). As  $\epsilon \rightarrow 0^+$ , the uniform convergence means

$$|v_\epsilon(x) - l^0(x) - R(x)| \leq \sigma + C|x|^3 \text{ and } |D_n v_\epsilon(x) - D_n R| \leq \sigma + C|x|^2,$$

where  $\sigma = \sigma(\epsilon) \rightarrow 0^+$ . If we let  $\mu \geq \sigma^{\frac{1}{3}} + \epsilon^{\frac{1}{3}}$ , then

$$(6.18) \quad |u - [Q + \epsilon(l^0 + R)]| \leq C\epsilon\mu^{\frac{3}{2}} \text{ in } S_\mu^Q,$$

and

$$(6.19) \quad |D_n u - (D_n Q + \epsilon D_n R)| \leq C\epsilon\mu \text{ in } S_\mu^Q.$$

**Step 4.** Let

$$f(t) := \det [D^2 Q + \epsilon D^2 R + t\mathcal{I}].$$

Notice that  $|f(0) - 1| \leq C\epsilon^2$ ,  $f'(t) \sim C$ . The equation  $f(t) = 1$  is solvable. The solution  $t_0$  satisfies

$$(6.20) \quad |t_0| \leq C\epsilon^2.$$

Let

$$(6.21) \quad Q^0(x) = Q(x) + \epsilon R(x) + \frac{t_0}{2}|x|^2, \quad P^0 = \epsilon l^0 + Q^0.$$

Then

$$\det D^2 P^0 = 1, \quad D_n P^0 = D_n Q = ax_1 \text{ on } \{x_n = 0\},$$

which means that  $P^0 \in \mathcal{F}_a$ .

Finally, we combine (6.18), (6.19), (6.20) and (6.21) to obtain

$$|u(x) - P^0(x)| \leq C\epsilon^2\mu + C\epsilon\mu^{\frac{3}{2}} \leq C\epsilon\mu^{\frac{3}{2}} \text{ in } S_\mu^{Q^0},$$

and

$$|D_n u(x) - D_n P^0(x)| \leq C\epsilon^2\mu^{\frac{1}{2}} + C\epsilon\mu \leq C\epsilon\mu \text{ in } S_\mu^{Q^0}.$$

Noticing that  $S_{\mu/2}^{Q^0} \subset S_\mu^{Q^0}$  and replacing  $\mu$  with  $\mu/2$ , we have proved the desired (6.8), while (6.9) follows directly from (6.21).  $\square$

**Proof of Theorem 6.1:** Let  $C, \mu$ , and  $\epsilon_0(\mu)$  are the same as in Lemma 6.3. Choose  $0 < \alpha < 1$  and  $0 < \mu \leq (4C)^{\frac{2}{\alpha-1}}$ . Let

$$C_\infty = \max\{\prod_{j=0}^\infty (1 + 4C^2\mu^{\frac{j\alpha}{2}}), e^{4C}\}.$$

Observing that

$$\log C + \sum_{j=0}^k \log(1 + 4CC_\infty^2\epsilon_0\mu^{\frac{k\alpha}{2}-2}) \leq 2C + \epsilon_0 C(\mu)C_\infty^3 \leq \log C_\infty,$$

we can assume that  $\epsilon_0$  is sufficiently small such that the sequence

$$C_0 = C, \quad C_k = (1 + 4CC_\infty^2\epsilon_0\mu^{\frac{k\alpha}{2}-2})C_{k-1}, \quad k = 1, 2, 3, \dots$$

is bounded by  $C_\infty$ .

We will prove by mathematical induction that there exists a sequence of constants  $a_k$ , transformations

$$\mathcal{M}_k = \text{diag}\{\mathcal{M}'_k, \mathcal{M}_{k,nn}\}, \quad \det \mathcal{M}_k = 1$$

(where  $\mathcal{M}'_k$  is a  $n-1$  square matrix), and linear functions  $l_k = \sum_{i=1}^{n-1} a_i^k x_i$  such that at height  $h_k := \mu^k$ , the normalization solution  $(u_k, \Omega_k)$  of  $(u, \Omega)$  given by

$$(6.22) \quad u_k(x) = \frac{u(\mathcal{T}_k x)}{h_k}, \quad x \in \Omega_k := \mathcal{T}_k^{-1} S_{h_k}(0), \quad \mathcal{T}_k = h_k^{\frac{1}{2}} \mathcal{M}_k$$

satisfying problem (6.1) and

$$(6.23) \quad \|u_k - l_k - Q_{a_k}\|_{L^\infty(S_1^{P_k})} + \|D_n u_k - D_n Q_{a_k}\|_{L^\infty(S_1^{P_k})} \leq \epsilon_k := \epsilon_0 \mu^{\frac{k\alpha}{2}},$$

for  $P_k = l_k + Q_k \in \mathcal{F}_a$ , where

$$Q_k(x) := \frac{1}{2} \left[ \sqrt{1 + a_k^2(x_1^2 + x_2^2)} + 2a_k x_1 x_2 + \sum_{i=2}^{n-1} x_i^2 \right].$$

Moreover,

$$(6.24) \quad \begin{cases} |b_{k-1}| \leq C(C_\infty)\epsilon_{k-1}^{\frac{1}{2}}, & |a_k| \leq C_k, & |\mathcal{M}_k\mathcal{M}_{k-1}^{-1} - \mathcal{I}| \leq C_k\epsilon_{k-1}. \\ C_k^{-1}B_1(0) \cap \Omega_k \subset S_1^{u_k}(0) \subset C_k B_1(0). \end{cases}$$

The case  $k = 0$  is the assumption of Theorem 6.1. If the case of  $k$  holds we will prove it for the case  $k + 1$ . We check for  $\epsilon_k$  whether  $u_k$  satisfies the conditions in Lemma 6.3. By induction hypothesis (6.23), we have

$$|u_k - P_k| \leq \epsilon_k \text{ in } \Omega_k \cap S_1^{P_k}.$$

Apply lemma 6.3 to  $u_k$ , we obtain the quadratic function  $\bar{P}_{k+1} = \bar{l}_{k+1} + \bar{Q}_{k+1} \in \mathcal{F}_a$  satisfies

$$(6.25) \quad \begin{cases} \text{dist}_\mu(u_k, \bar{P}_{k+1}, 0) \leq C\epsilon_k\mu^{\frac{3}{2}} \leq \epsilon_{k+1}\mu, \\ |b_k| \leq C\epsilon_k^{\frac{1}{2}} \text{ and } |D^2Q_k - D^2\bar{Q}_{k+1}| \leq C\epsilon_k. \end{cases}$$

Then there exists  $a_k$  and positive matrix  $\mathcal{B}_{k+1} = \text{diag}\{\mathcal{B}'_k, \mathcal{B}_{k,nn}\}$  satisfies

$$\mathcal{B}_{k+1}^T D^2Q_k \mathcal{B}_{k+1} = D^2Q_{k+1}$$

and

$$|a_{k+1} - a_k| \leq CC_k\epsilon_k \text{ and } |\mathcal{B}_{k+1} - \mathcal{I}| \leq CC_k\epsilon_k.$$

In particular,  $|a_{k+1}| \leq C_{k+1}$ . Take  $\mathcal{M}_{k+1} = \mathcal{M}_k\mathcal{B}_{k+1}$ ,  $\mathcal{T}_{k+1} = \mu^{\frac{1}{2}}\mathcal{T}_k\mathcal{B}_{k+1}$  and let

$$u_{k+1}(x) = \frac{u(\mathcal{T}_{k+1}x)}{h_{k+1}}, \quad S_1^{u_{k+1}}(0) = \mathcal{T}_{k+1}^{-1}S_{h_{k+1}}(0), \quad h_{k+1} = \mu^{(k+1)\alpha}.$$

Then we have

$$(1 - CC_\infty^2\mu^{-1}\epsilon_k)C_k^{-1}B_1(0) \cap \Omega_{k+1} \subset S_1^{u_{k+1}}(0) \subset (1 + CC_\infty^2\mu^{-1}\epsilon_k)C_k B_1(0),$$

which is

$$C_{k+1}^{-1}B_1(0) \cap \Omega_{k+1} \subset S_1^{u_{k+1}}(0) \subset C_{k+1}B_1(0).$$

Using the above facts and (6.25), one may verify by direct calculations that (6.23)-(6.24) holds for  $k + 1$ .

It follows that  $|a_k| \leq C_k$ ,  $|\mathcal{M}_k| \leq C_k$  converge geometrically to  $a_\infty$ ,  $\mathcal{M}_\infty$  and

$$|\mathcal{M}_\infty\mathcal{M}_k^{-1} - \mathcal{I}| \leq C_k\epsilon_k \leq C_\infty h_k^{\frac{\alpha}{2}}.$$

We replace each  $\mathcal{M}_k$  by  $\mathcal{M}_\infty$  and  $Q_k$  by  $Q_\infty$ , the inductive hypothesis still holds with a large universal constant. This is

$$|u(x) - h_k^{\frac{1}{2}}l_k(\mathcal{M}_\infty^{-1}x) - Q_\infty(\mathcal{M}_\infty^{-1}x)| \leq C_1C_\infty^2h_k^{1+\frac{\alpha}{2}} \text{ in } B_{ch_k^{\frac{1}{2}}}(0).$$

Recalling that  $|Dl_k| \leq C\epsilon_{k-1}^{\frac{1}{2}}$ , we obtain

$$|u(x) - Q_\infty(\mathcal{M}_\infty^{-1}x)| \leq C_1C_\infty^2h_k^{1+\frac{\alpha}{2}} \text{ in } B_{ch_k^{\frac{1}{2}}}(0),$$

which means

$$(6.26) \quad |u(x) - Q_\infty(\mathcal{M}_\infty^{-1}x)| \leq C_1 C_\infty^2 |x|^{2+\alpha}.$$

The above induction argument and Lemma 6.3 could be applied to any point  $y \in G_{c_0}(0)$ . As a result, we may obtain that

$$|u(y+x) - Q^y(x)| \leq C|x|^{2+\alpha}$$

for a quadratic polynomial depending on the based point  $y$ . In fact, consider  $p = y + \theta e_n$  with  $\theta > 0$  is small enough. The height section  $S_{c_1\theta^2}^{Q^y}(p)$  is contained in  $S_1$  with

$$B_{c\theta^2}(p) \subset S_{c_1\theta^2}^{Q^y}(p) \subset B_{C\theta^2}(p).$$

By (6.26) we have

$$\begin{cases} \det D^2u(x) = 1 & \text{in } S_{c_1\theta^2}^{Q^y}(p) \\ |u - Q^y(x)| \leq C|\theta|^{2+\alpha} & \text{on } \partial S_{c_1\theta^2}^{Q^y}(p) \end{cases}.$$

By interior  $C^{2,\alpha}$  theory and its perturbation method (See [3] for the details),  $\|u\|_{C^{2,\alpha}(B_{c\theta^2}(p))} \leq C$ . Since the constant  $C$  is independent of  $y$  and  $\theta$ ,  $u \in C^{2,\alpha}(\overline{B_{c_0}^+(0)})$  for a small  $c_0 > 0$ .

**Proof of Theorem 6.2:** Choose  $\delta_0 = b_0^{\delta n}$  and  $c_0 = cb_0$ , where  $\alpha, \mu$ , and  $b_0$  are the same as in the proof of Theorem 6.1. Take  $a \in (-\delta_0, \delta_0)$ . We are going to construct a quadratic approximation of  $u$  near 0 as (6.4), and applying Theorem 6.1 to complete the proof of Theorem 6.2.

Lemma 5.2 implies

$$\mathcal{P}S_1(0) \subset (1 + C\delta_0)G_1(0).$$

For  $(x', 0) \in \mathcal{P}S_1(0)$ , let  $g(x')$  be the maximum/minimum value such that  $(x', g(x')) \in S_1(0)$ . We consider the convex set

$$E = \{(x', x_n) | (x', 0) \in \mathcal{P}S_1(0), 0 \leq x_n \leq g(x')\},$$

and

$$S = \{(x', \pm x_n) | (x', x_n) \in E\}.$$

Then

$$\partial S_1 \setminus G_1 \subset E_{C\delta_0} = \{x \in \mathbb{R}^n | \text{dist}(x, \partial E) \leq C\delta_0\}.$$

By the standard theory (see [10, 11] for example) we see that there is a convex function  $v$  symmetric about the hyperplane  $\{x_n = 0\}$  solving

$$\begin{cases} \det D^2v = 1 & \text{in } S \\ v = 1 & \text{on } \partial S \end{cases}.$$

Moreover, by (6.2) and Lemma 2.13,  $v$  is smooth in the interior and Lemma 2.12 tells us

$$v(x) \geq 1 - C \operatorname{dist}(x, \partial S)^{\frac{1}{n}}.$$

Thus,  $v$  is also a solution to the following Neumann problem

$$(6.27) \quad \begin{cases} \det D^2 v = 1 & \text{in } S_1 \\ 1 - b_0^6 \leq v \leq 1 & \text{on } \partial S_1 \setminus G_1. \\ D_n v = 0 & \text{on } G_1 \end{cases}$$

Since  $|a| < \delta_0$  and  $b_0 > 0$  is small, then  $v^\pm := v \pm 2b_0^6(2C - x_n)$  must be supersolution/subsolution of  $u$  to (6.1), which implies

$$(6.28) \quad \|v - u\|_{L^\infty(S_1)} \leq Cb_0^6$$

by the comparison principle (Lemma 2.10). Let  $w = u - v$ , by Corollary 2.7 we have

$$(6.29) \quad |\nabla w| \leq Cb_0^3 \text{ in } B_c^+(0).$$

By (6.27) we have  $D_{in}v = 0$  for  $i \neq n$  in  $G_1$ . Let

$$\mathcal{M} = D_{ij}v(0), \quad R(x) = x^T \mathcal{M} x, \quad \mathcal{M} = \mathcal{B}^2,$$

where  $\mathcal{B} = (b_{ij})_{n \times n}$  may be chosen such that it is bounded symmetric matrix with  $b_{in} = 0$  for  $i \neq n$  because of the same property of  $\mathcal{M}$ . Consider the diagonal matrices  $\mathcal{B}_1$  defined by

$$\mathcal{B}_1^2 = \left[ \frac{1}{2} \operatorname{diag} \{ \sqrt{1+a^2}, I'', \sqrt{1+a^2} \} \right]^{-1}$$

and the function

$$\tilde{u}(x) = \frac{u(b_0(\mathcal{B}\mathcal{B}^{-1})x)}{b_0^2}, \quad Q(x) = \frac{Q_a(b_0(\mathcal{B}\mathcal{B}_1)^{-1}x)}{b_0^2}, \quad x \in B_C(0),$$

where  $Q_a$  is the same as in (6.3). Letting  $E_0 = B_{Cb_0}(0)$  and noticing  $v \in C^3(B_c(0))$ , we see that

$$\begin{aligned} \|\tilde{u} - Q\|_{L^\infty(E_0)} &\leq \frac{1}{b_0^2} \|u - R\|_{L^\infty(E_0)} + C|a| \\ &\leq \frac{1}{b_0^2} \|u - v\|_{L^\infty(E_0)} + Cb_0 + C|a| \\ &\leq Cb_0 \end{aligned}$$

and

$$\begin{aligned} \|D_n \tilde{u} - D_n Q\|_{L^\infty(E_0)} &\leq \frac{1}{b_0} \|D_n u - x_n D_{nn} v\|_{L^\infty(E_0)} + C|a| \\ &\leq \frac{1}{b_0} \|D_n u - D_n v\|_{L^\infty(E_0)} + Cb_0 + C|a| \\ &\leq Cb_0. \end{aligned}$$

By Theorem 6.1, we have  $\tilde{u} \in C^{2,\alpha}(\overline{B_c^+(0)})$ , therefore  $u \in C^{2,\alpha}(\overline{B_{c_0}^+(0)})$  for  $c_0 = cb_0$ .

## 7. UNIVERSAL STRICT CONVEXITY

Throughout this section, we assume that  $u$  is a viscosity solution to problem (1.1), satisfying (1.2) (or (2.2)), (2.1) and (2.6). By (2.1) and Theorem 5.1 we may assume

$$(7.1) \quad B_c^+(0) \subset S_1(0) \subset B_C(0).$$

We will study the geometry of the section  $S_R(0)$  for large  $R$ . In particular, we will show the uniformly strict convexity for the normalization family  $(u_R, S_1^R)$  defined by (5.16).

Recall the notations from the beginning in Section 5 to (5.1). When we fix a  $R$ , we will use the notation  $\mathcal{A}_R$ ,  $d_i(R)$  and  $\mathcal{D}_R$  instead of  $\mathcal{A}_h$ ,  $d_i(h)$  and  $\mathcal{D}_h$ . We define the section  $S_t^R(0)$  of  $(u_R, S_1^R)$  as we defined  $S_1^h(0)$  of  $(u, S_h)$  by (5.1) in Section 5. Set

$$d_i^R(t) = \frac{d_i(tR)}{d_i(R)}, \quad \mathcal{D}_t^R = \mathcal{D}_{tR}(\mathcal{D}_R)^{-1} \text{ and } \mathcal{A}_t^R = (D_R)^{-1} \mathcal{A}_{tR}(\mathcal{A}_R)^{-1} D_R,$$

The normalization of  $(u_R, S_1^R)$  at height  $t$  is essentially the same as  $(u_{tR}, S_1^{tR})$ :

$$u_{tR}(x) := \frac{u_R((\mathcal{A}_t^R)^{-1} \mathcal{D}_t^R x)}{t}, \quad x \in S_1^{tR} := (\mathcal{D}_t^R)^{-1} \mathcal{A}_t^R S_1^R.$$

Here is the main result of this section.

**Theorem 7.1.** *Given large constants  $R_0 > 0$  and  $C_0 > 1$ . There exists a small constant  $\eta > 0$  such that for any  $R \geq R_0$ ,*

$$(7.2) \quad C_0 \eta S_R(0) \subset S_{\eta R}(0) \subset (1 - \eta) S_R(0).$$

Moreover, there exist  $\bar{\delta} = \delta_{R_0}$  and  $\gamma \in (0, 1)$  such that

$$(7.3) \quad c|x|^{\frac{1+\gamma}{\gamma}} - C\delta_R/\bar{\delta} \leq u_R(x) \leq C|x|^{1+\gamma} + C\delta_R/\bar{\delta}, \quad \forall x \in S_1^R,$$

where  $\delta_R$  denotes a constant depending on  $R$  and tends to 0 as  $R \rightarrow \infty$ .

**Claim 7.2.** *(7.2) is a scaling invariant property. Using iterative techniques, one can see that (7.2) and (7.3) are equivalent. More precisely, regardless of the term  $C\delta_R/\bar{\delta}$ , the left side of (7.2) and the right side of (7.3) are equivalent to*

$$u_R(x) \leq \sigma_1(|x|)|x| \text{ for all large } R,$$

and the right side of (7.2) and the left side of (7.3) are equivalent to

$$u_R(x) \geq \sigma_0(|x|) \text{ for all large } R,$$

where  $\sigma_0, \sigma_1$  are some universal strictly increasing  $C^0$  function with  $\sigma_i(0) = 0$  ( $i = 0, 1$ ).

The right side of (7.2) implies  $S_{\eta^k}^{\eta^{-k}R} \subset B_{(1-\eta)^k C}(0)$ . Thus, if we need to obtain the global type estimate of  $u_R$  in  $S_1^R$ , we only need to prove the corresponding local estimate of  $u_{\eta^{-k}R}$  in  $S_{\eta^k}^{\eta^{-k}R}$  for some universal large  $k$ . Especially, we can obtain that  $D_n u_R$  is uniformly bounded in  $S_1^R(0)$ .

By Lemma 2.5 and the Neumann boundary value condition, the right side of (7.3) provides a universal  $C^{1,\gamma}$  module for  $u_R$  at 0.

To prove Theorem 7.1, in addition to the notations  $\delta_0, c_0$  in Theorem 6.2 and  $a_R$  in (5.19) we need the following lemma.

**Lemma 7.3.** *If  $R \geq R_0$  is large, then  $a_R \geq c\delta_0 a$  and*

$$(7.4) \quad cR^{\frac{1}{2}}\mathcal{I}'' \leq D_R'' \leq CR^{\frac{1}{2}}\mathcal{I}''.$$

*Proof.* Assume by way of contradiction that  $a_R \leq c\delta_0 a \leq \delta_0$ . Then Theorem 6.2, together with (2.6) and (7.1), implies that the normalization solution  $u_R$  is  $C^{2,\alpha}$  near 0. This is

$$B_c^+(0) \subset t^{-\frac{1}{2}}S_t^R(0) \subset B_C^+(0), \quad \forall t \in (0, cc_0^2).$$

We recall (7.1) and  $S_1(0) = \mathcal{D}_R^{-1}\mathcal{A}_R S_{\frac{1}{R}}^R(0)$ . If  $R \geq Cc_0^{-2}$  is large, we can take  $t = \frac{1}{R}$  to obtain

$$c\mathcal{I} \leq \mathcal{D}_R^{-1}\mathcal{A}_R R^{\frac{1}{2}} \leq C\mathcal{I}.$$

This gives  $d_i(R) \leq CR^{\frac{1}{2}}$  for  $2 \leq i \leq n-1$ . Hence,

$$a_R = \frac{d_1(R)d_n(R)}{R}a \geq Ca,$$

which is a contradiction.

Thus  $a_R \geq c\delta_0 a$ , which implies  $\prod_{i=2}^{n-2} d_i(R) \leq C_1\delta_0^{-1}R^{\frac{n-2}{2}}$ . Noticing that  $d_i(R) \geq cR^{\frac{1}{2}}$  for  $2 \leq i \leq n-1$  (see (2.2)), we see that (7.4) holds for  $C = (C_1\delta_0^{-1})^{\frac{1}{n-2}}$ .  $\square$

**Remark 7.4.** *By Lemma 7.3 we could improve (2.2) to*

$$c|z''|^2 - \bar{\delta}^{-1} \leq u(0, z'', 0) \leq C|z''|^2,$$

which implies

$$(7.5) \quad c|z''|^2 - \delta_R/\bar{\delta} \leq u_R(0, z'', 0) \leq C|z''|^2 \text{ for } R \geq \bar{\delta}.$$

At this moment, the above also holds when  $a = 0$ .

**Proof of Theorem 7.1:** Due to Claim 7.2, it is sufficient to prove (7.3), which will be completed by four steps. According to Theorem 5.1, we only need to prove (7.3) in normal direction and on tangent plane.

**Step 1.** Let  $E_R = S_1^R \setminus S_{\delta_R/\bar{\delta}}^R$ , we first prove that

$$(7.6) \quad u_R(x) \geq \sigma_0(\max\{|x_1|, |x_n|\}) - C\delta_R \text{ on } E_R$$

for some universal module  $\sigma_0$ .

Assume  $q = (q', 0) = (q_1, q'', 0) \in \mathcal{P}E_R$  with  $q_1 \neq 0$ , let

$$\mathcal{B}x = x - \sum_{i=2}^{n-1} \frac{q_i}{q_1} x_1 e_i$$

and consider the positive function

$$v(x) = \frac{u(\frac{q}{2} + \frac{q_1 \mathcal{B}^{-1}x}{8C})}{q_1^2}, \quad x \in B_1^+(0).$$

By the convexity, (7.5) and the assumption on  $u$ , we have

$$0 \leq v(x) \leq \frac{C}{q_1^2}$$

and

$$v(x_1, x'', 0) \leq \frac{u(q)}{q_1^2} + C(|x''|^2 + \frac{\delta_R}{q_1^2}) \leq C|x''|^2 + \frac{u(q) + C\delta_R}{q_1^2}.$$

The Neumann boundary value condition guarantees

$$D_n v(x_1, x'', 0) \geq -C.$$

Regarding  $\frac{u(q) + C\delta_R}{q_1^2}$  as small constant and  $q_1^{-1}$  as universal constant, we use Lemma 4.1 to obtain

$$\frac{u(q) + C\delta_R}{q_1^2} \geq \sigma\left(\frac{C}{q_1^2}\right).$$

Letting  $\sigma_0(t) = t^2\sigma(C/t^2)$ , we get

$$(7.7) \quad u(q', 0) \geq \hat{\sigma}_0(q_1) - C\delta_R, \quad \forall (q', 0) \in \mathcal{P}E_R.$$

Next, assume  $q = q_n e_n$  and consider the positive function

$$v(x) = \frac{u(\frac{3q}{4} + \frac{q_n x}{8C})}{q_n^2}, \quad x \in B_1(0).$$

Similarly, we have

$$0 \leq v(x) \leq \frac{C}{q_n^2} \text{ and } v(0, x'', x_n) \leq C|x''|^2 + \frac{u(q) + C\delta_R}{q_n^2}.$$

Using Lemma 4.2, we get

$$(7.8) \quad u(q_n e_n) \geq \sigma_0(q_n) - C\delta_R$$

for  $\sigma_0(t) = t^2\sigma(C/t^2)$ .

Since  $S_t$  is of good shape for  $t \geq \delta_R/\bar{\delta}$ , combining (7.7) and (7.8), we have proved the desired (7.6).

**Step 2.** We first choose a large  $R_0$  such that  $\bar{\delta}R_0^{-1}$  is small enough, and define  $\delta_R = (\bar{\delta}R^{-1})$ . Then  $\delta_R \leq \delta R_0$  for  $R \geq R_0$ .

For simplicity, we only consider the point  $x \in E := S_1^R(0) \setminus S_{\delta_R}^R(0)$  and assume  $t \geq \delta_R$  in the rest of this Step. By iteration, (7.6) means that there exists universal  $\gamma \in (0, 1)$  such that

$$u_R(x) \geq c \max\{|x_1|, |x_n|\}^{\frac{1+\gamma}{\gamma}},$$

which is

$$\max\{d_1^R(t), d_n^R(t)\} \leq Ct^{\frac{\gamma}{1+\gamma}}.$$

Recalling (7.4) and  $\Pi_{i=1}^n d_i^R(t) = t^{\frac{n}{2}}$ , we have

$$\min\{d_1^R(t), d_n^R(t)\} \geq ct^{\frac{1}{1+\gamma}}.$$

This gives (7.3) in the normal direction.

**Step 3.** We now have

$$(7.9) \quad ct^{\frac{1}{1+\gamma}} \leq d_1^R(t) \leq Ct^{\frac{\gamma}{1+\gamma}},$$

whose right side gives

$$u_R(x', 0) \geq c|x_1|^{\frac{1+\gamma}{\gamma}}.$$

Noticing (7.5) and that  $u_R$  is locally Lipschitz, we get

$$(7.10) \quad u_R(x', 0) \geq \max\{c|x''|^2 - \frac{C\delta_R}{\delta}|x_1|, 0\}.$$

Let  $E_1 = \{x' \mathcal{P}(E) : |x_1| \leq \frac{c^2}{4}|x''|^2\}$  and  $E_2 = \mathcal{P}(E) \setminus E_1$ . The above two inequalities implies that

$$\begin{aligned} u_R(x', 0) &\geq \max\{c|x''|^2 - C|x_1|, 0\}\chi_{E_1} + c|x_1|^{\frac{1+\gamma}{\gamma}}\chi_{E_2} \\ &\geq \frac{c}{2}|x''|^2\chi_{E_1} + c|x_1|^{\frac{1+\gamma}{\gamma}}\chi_{E_2} \\ &\geq c|x'|^{\frac{2(1+\gamma)}{\gamma}}, \end{aligned}$$

Which proves the right side of (7.3) on the tangent plane.

**Step 4.** Given constant  $M$ , we claim that there exist universal  $\varepsilon$ , such that

$$\sup\{s \mid se_1 \in G_t^R(0)\} \geq M\varepsilon$$

for some  $t \in [\varepsilon, \frac{1}{2}]$ . Then This claim, (7.5) and (5.8) implies that

$$\varepsilon^{-1}G_\varepsilon^R(0) \supset t^{-1}G_t^R(0) \supset cMB_1(0) \supset cM\mathcal{P}(S_1^R(0)),$$

which proves the left side of (7.3) on the tangent plane.

On the contrary to this claim, by the balance property of  $G_t^R(0)$  we get

$$u_R(x_1, 0) \geq M^{-1}|x_1| \text{ for } x_1 \geq CM\varepsilon.$$

For simplicity, we may require the point  $x \in S_1^R(0) \setminus (S_{CM\varepsilon}^R(0) \cap \{|x_1| \leq 2CM\varepsilon\})$  and assume constant  $t \geq CM\varepsilon$  in the rest of this Step. Then by the convexity and (7.5) we have

$$u_R(x_1, x'', 0) \geq 2u_R(2x_1, 0, 0) - u_R(0, -x'', 0) \geq 2M^{-1}|x_1| - C|x''|^2.$$

This together with (7.10) implies that

$$G_t^R(0) \subset \{c(|x''|^2 - t) \leq |x_1| \leq CM(|x''|^2 + t)\}.$$

Since  $0 \in G_t^R(0)$ , the above relation then implies that

$$G_t^R(0) \subset \{|x_1| \leq CM^4t\}.$$

Thus,  $d_1^R(t) \leq CM^4t$ . Taking  $t = \varepsilon$  is small enough, this contradicts (7.9). In this way, we completes the proof of Theorem 7.1.

## 8. LIOUVILLE THEOREM

In this section, we will modify the techniques developed from the classical  $C^{1,\alpha}$  estimate [4] to prove Theorem 1.1. We always suppose that  $u$  is a viscosity solution to problem (1.1) and the hypothesis of Theorem 1.1 is satisfied. As we have said, we may assume (2.6). Recall the normalization family  $(u_R, S_1^R)$  defined by (5.16), which satisfies (5.17)-(5.22).

Consider the function

$$(8.1) \quad v_R(x) = D_n u_R(x) - a_R x_1,$$

and let

$$(8.2) \quad \begin{aligned} m_R &= \inf \left\{ \frac{v_R(x)}{x_n} \mid x \in S_1^R(0) \right\}, \\ M_R &= \sup \left\{ \frac{v_R(x)}{x_n} \mid x \in S_1^R(0) \right\}, \\ \omega_R &= \frac{M_R}{m_R} - 1. \end{aligned}$$

Our goal is to prove that

$$(8.3) \quad c \leq m_R \leq M_R \leq C$$

and  $\omega_R$  is strictly increasing in  $R$ . For this purpose, we need the auxiliary function like those from (3.5) to (3.7). Given  $y \in G_{\frac{1}{2}}^R(0)$ , let

$$(8.4) \quad u^{R,y}(x) = u_R(x) - u_R(y) - \nabla u_R(y) \cdot (y - x)$$

and define the functions

$$(8.5) \quad \begin{aligned} w^{R,y}(x) &= [u^{R,y}(x) - \frac{n}{2}x_n D_n u^{R,y}(x)], \\ w^{R,y,+}(x) &= C_1[w^{R,y}(x) + C_2 x_n], \\ w^{R,y,-}(x) &= c_1[x_n - c_2 w^{R,y}(x)]. \end{aligned}$$

In order to show that  $w^{R,y,\pm}$  are supersolution/subsolution to the following linearized problem (8.17) for  $v_R$ , we need the following two lemmas.

**Lemma 8.1.** *Suppose that  $u$  is a viscosity solution to the first equation of problem (1.1) and (2.6) is satisfied. Assume that*

$$(8.6) \quad B_c^+(0) \subset S_1^u(0) \subset B_C^+(0) \text{ and } \|u\|_{Lip(B_C(0))} \leq C.$$

Define  $h_x^u = \sup\{h \mid S_t^u(x) \subset B_1^+(0), \forall t < h\}$ . There exists small universal constants  $\delta > 0, \rho > 0$  such that if

$$(8.7) \quad c|x|^{\frac{1+\gamma}{\gamma}} - \delta \leq u(x) \leq C|x|^{1+\gamma} + \delta, \quad \forall x \in S_1^u(0),$$

then

$$h_x^u \geq \sigma_0(x_n) \text{ in } B_\rho^+(0).$$

*Proof.* If the result were false, we could find a constant  $\tau > 0$ , a sequence of convex viscosity solutions  $u_k$  satisfying (8.6)-(8.7) and points  $p_k \in B_\rho(0)$  such that

$$p_k \cdot e_n \geq \tau, \quad \text{and } h_{p_k}^{u_k} \rightarrow 0.$$

Letting  $k \rightarrow \infty$ , we get a limit  $v$  which still satisfies our assumption and  $p_k$  converges to a point  $p$  such that  $h_p^v = 0$ . Let  $\Sigma := S_0^v(p)$ . Then  $\Sigma$  is not a single point set. (8.7) implies that  $\Sigma \subset B_{\delta_1}(0)$  for some constant  $\delta_1 > 0$  depending on  $\delta, \rho$ . Hence we can find an interior extremal point

$$q \in \{y \mid y \in \Sigma, y_n = \sup_{x \in \Sigma} x_n\},$$

which contradicts Lemma 2.14. □

**Lemma 8.2.** *For large  $R \geq R_0$ , one has*

$$(8.8) \quad \sigma_0(x_n) \leq v_R(x) \leq C \text{ in } S_1^R(0).$$

*Proof.* As we have discussed in Claim 7.2, we need only to prove (8.8) for  $y \in S_{\sigma_0(|c_0|)}^R(0) \subset B_{c_0}(0)$  for a small universal  $c_0$ . According to Lemma 2.5, the Neumann boundary value condition and (7.3) provides a  $C^1$  module for  $u_R$  at 0. This is,

$$(8.9) \quad |\nabla u(x)| \leq C|x|^\gamma, \text{ for } x \text{ near } 0,$$

which implies the right side of (8.8).

As in Lemma 8.1 we denote

$$(8.10) \quad h_y^R = \sup\{h \mid S_t^R(y) \subset \mathbb{R}_+^n, \forall t < h\}.$$

We will find a universal  $C^0$  module  $\sigma_0$  such that the following three inequalities hold in  $S_1^R(0)$ :

$$(8.11) \quad h_x^R \geq \sigma_0(x_n),$$

$$(8.12) \quad \sigma_0(x_n)\mathcal{I} \leq D^2 u_R(x) \leq \frac{1}{\sigma_0(x_n)}\mathcal{I},$$

and

$$(8.13) \quad v_R(x) \geq \sigma_0(x_n).$$

Again, it is sufficient to prove (8.10)-(8.12) for  $x$  around 0. Suppose  $c_0 > 0$  is small and  $y \in S_{c_0}^R(0)$ , then (7.3) and (8.9) means

$$|\nabla u_R(y)| + u_R(y) \leq c_1 := c_0 + Cc_0^\gamma$$

and  $c_1$  is still small. Thus, the section  $S_{h_y^R}^R(y) \cap B_C(0)$  is contained in  $S_{c_1}^R(0) \subset S_{\frac{1}{2}}^R(0)$  and will touch  $\partial\mathbb{R}_+^n$  at some point  $z = (z', 0) \in G_{c_1}^R(0)$ .

Let  $l_{zy}$  be the line segment connecting  $z$  and  $y$ . Extend  $l_{zy}$  such that it intersects  $\partial B_c^+(0)$  at point  $q$ . Set

$$w(x) = u_R(x) - u_R(y) - \nabla u_R(y) \cdot (x - y).$$

Then

$$(8.14) \quad w(x) \leq h_y^R, \quad x \in l_{yz}.$$

Furthermore, it follows from Theorem 7.1 and Lemma 8.1 that

$$(8.15) \quad h_y^R \geq \sigma_0(y_n)$$

for  $\sigma_0(t) := ct^{\frac{2}{n}}\sigma(ct^{-\frac{2}{n}})$ .

The classical estimate result (see Lemma 2.11) gives  $\text{Vol}(S_{h_y^R}^R(y)) \sim (h_y^R)^{\frac{n}{2}}$ . Notice that  $S_{h_y^R}^R \subset B_C^+(0)$ , thus  $S_{h_y^R}^R \supset B_{c(h_y^R)^{\frac{1}{2}}}^+(0)$  and the classical interior estimate (See Lemma 2.13 and its proof) then gives

$$(8.16) \quad \hat{\sigma}_0(y_n)\mathcal{I} \leq D^2u_R(y) \leq \frac{1}{\hat{\sigma}_0(y_n)}\mathcal{I}$$

for  $\hat{\sigma}_0(y_n) = \min\{[\sigma_0(y_n)]^{n-2}, \sigma_0(y_n)\}$ .

Now, up to a bounded linear transform which maps in  $S_{\sigma_0}^R(0)$  to  $S_1^{\sigma_0 R}$ , (8.15)-(8.16) and the statement in Claim 7.2 implies that (8.11) and (8.12) in  $S_1^R(0)$ .

Finally, let us prove the (8.13). Similarly as Lemma 3.4, the non-negative function  $v_R(x)$  in (8.1) solves the linearized problem:

$$(8.17) \quad \begin{cases} U_R^{ij}D_{ij}v_R = 0 & \text{in } S_1^R(0) \\ 0 \leq v_R \leq C & \text{in } S_1^R(0) \\ v_R = 0 & \text{on } G_1^R(0) \end{cases} .$$

Notice that we also have  $v_R(ce_n) \geq c$  by the convexity. Due to (8.12), the first equation in (8.17) is uniformly elliptic with norm of the elliptic coefficients depending on  $\sigma_0(y_n)$  in the domain  $E(y_n) = S_1^R(0) \cap \{x_n \geq \frac{y_n}{2}\}$ . We apply the classical Harnack inequality to obtain

$$(8.18) \quad v_R(y', y_n) \geq C(\|U_R^{ij}\|_{E(y_n)}, \|U_R^{ij,-1}\|_{E(y_n)}, \frac{y_n}{2})v_R(ce_n) \geq \sigma_0(y_n)$$

in  $S_{\frac{1}{2}} \cap \{x_n \geq \frac{y_n}{2}\}$  for some new  $\sigma_0$ . Thus, the proof is completed.  $\square$

**Proof of Theorem 1.1:** It follows from Lemma 8.2 that  $v_R(x)$  satisfies the following problem:

$$(8.19) \quad \begin{cases} U_R^{ij}D_{ij}v_R = 0 & \text{in } S_1^R(0) \\ \sigma_0(x_n) \leq v_R \leq C & \text{on } \partial S_1^R(0) \setminus G_1^R(0) \\ v_R = 0 & \text{on } G_1^R(0) \end{cases} .$$

Then as Corollary 3.7 and 3.8 we see that the function  $w^{R,y,\pm}(x)$  in (8.5) are super-solution/subsolution of problem (8.19) in  $S_c^R(0)$ , and  $cx_n \leq v_R(x) \leq Cx_n$  in  $S_c^R(0)$ . Using the definition (8.2), we get

$$c \leq m_R \leq M_R \leq C,$$

which is the desired (8.3).

Next, we are going to prove

$$(8.20) \quad \omega_{\rho R} \leq (1 - \theta)\omega_R$$

for some universal  $\rho > 0$  and  $\theta \in (0, 1)$ .

Let

$$\begin{aligned} m_{R,t} &= \inf\left\{\frac{v_R(x)}{x_n} \mid x \in S_t^R(0)\right\}, \\ M_{R,t} &= \sup\left\{\frac{v_R(x)}{x_n} \mid x \in S_t^R(0)\right\}, \\ \omega_{R,t} &= \frac{M_{R,t}}{m_{R,t}} - 1. \end{aligned}$$

It is clear that for all  $t \in (0, 1)$ ,

$$m_{R,1} \leq m_{R,t} \leq M_{R,t} \leq M_{R,1}$$

and

$$\omega_{R,t} = \omega_{Rt}.$$

To prove (8.20), it is sufficient to prove

$$(8.21) \quad \omega_{R,\rho} \leq (1 - \theta)\omega_{R,1}$$

for some universal  $\rho > 0$  and  $\theta \in (0, 1)$ . Assume  $\omega_{R,1} > 0$  and consider the two non-negative functions:

$$(8.22) \quad \begin{aligned} H_1(x) &= \frac{v_R(x) - m_{R,1}x_n}{M_{R,1} - m_{R,1}}, \\ H_2(x) &= \frac{M_{R,1}x_n - v_R(x)}{M_{R,1} - m_{R,1}}. \end{aligned}$$

It follows from (8.19) that each  $H_i$  ( $i = 1, 2$ ) solves

$$(8.23) \quad \begin{cases} U_R^{ij} D_{ij} H_i = 0 & \text{in } S_1^R(0) \\ 0 \leq H_i \leq 1 & \text{in } S_1^R(0) \\ H_i = 0 & \text{on } G_1^R(0) \end{cases}.$$

While  $H_1 + H_2 = x_n$ , we may assume  $H_1(ce_n) \geq \frac{\epsilon}{2}$  without loss of generality.

Since (8.12) means that the first equation in (8.23) is universal elliptic away from  $G_1^R(0)$ , as (8.18) we have

$$(8.24) \quad H_1(x) \geq \sigma_0(x_n)H_1(ce_n) \text{ in } B_{c/2}^+(0).$$

Again, for any  $y = (y', 0) \in G_{c_0}^R(0)$  with  $c_0 > 0$  small enough, we could find a universal constant  $\bar{c} > 0$  such that  $B_{c(\bar{c})/2}^+(0) \subset S_{\bar{c}}^R(y) \subset B_{c/2}^+(0)$ .

With some universal constant  $C_1, C_2, c_1, c_2$  depending on  $\bar{c}$  and  $\sigma_0(x_n)$ , the function  $w^{R,y,\pm}(x)$  in (8.5) is still supersolution/subsolution of  $H_1$  to the following problem:

$$(8.25) \quad \begin{cases} U_R^{ij} D_{ij} H_1 = 0 & \text{in } S_{\bar{c}}^R(y) \\ \sigma_0(x_n) \leq H_1 \leq 1 & \text{on } \partial S_{\bar{c}}^R(y) \setminus G_{\bar{c}}^R(y) . \\ H_1 = 0 & \text{on } G_{\bar{c}}^R(y) \end{cases}$$

In conclusion, we have

$$(8.26) \quad H_1(y + te_n) \geq w^{R,y,-}(y + te_n) \geq \frac{c_1}{2}t \text{ for } t \leq t_0.$$

On the other hand, it follows from (7.2) (or (7.3)) that

$$(8.27) \quad S_{\rho}^R(0) \subset G_{c_0}^R(0) \times [0, t_0) \text{ for } \rho = c \min\{c_0^{\frac{(1+\gamma)^2}{\gamma}}, t_0\}.$$

Combining (8.26) and (8.27), we have

$$\frac{v_R(x) - m_{R,1}x_n}{M_{R,1} - m_{R,1}} \geq \theta x_n \text{ in } S_{\rho}^R(0)$$

for  $\theta = \frac{cc_1}{2}$ . That is

$$m_{R,\rho} \geq m_{R,1} + \theta(M_{R,1} - m_{R,1}) \text{ in } S_{\rho}^R(0),$$

which yields the desired (8.21). In this way, (8.20) has been proved.

By (8.3) and (8.20) we see that

$$\omega_R = \omega_{\rho^k \rho^{-k} R} \leq (1 - \theta)^k \omega_{R \rho^{-k}} \leq C(1 - \theta)^k.$$

Letting  $k \rightarrow 0$ , we get  $\omega_R = 0$  for any  $R$ . Hence,  $v_R(x) = D_n u_R - a_R x_1$  is a linear function in  $S_R(0)$ . Then letting  $R \rightarrow \infty$ , we get the limit function  $\hat{v}(x) := D_n u(x) - a x_1 = c x_n$  for some  $c > 0$ .

Lemma 3.1 means that  $u$  is strict convex and smooth away from  $\partial \mathbb{R}_+^n$ . Thus

$$u(x) = f(x') + a x_1 x_n + c x_n^2$$

for some smooth function  $f$  in  $\mathbb{R}^{n-1}$ . Let

$$g(x') = f(x') - \frac{a^2}{4c} x_1^2.$$

Since  $D^2 u(x', 1) > 0$ , we have  $D^2 g(x') > 0$ . Therefore,  $g$  is a convex, smooth function solving

$$\det D^2 g = 1 \text{ in } \mathbb{R}^{n-1}.$$

The classical results (Theorem 2.15) shows that  $g$  is quadratic polynomial of  $x'$ , which implies that  $u$  is a quadratic polynomial of  $x$ . The proof of Theorem 1.1 is completed.

**Remark 8.3.** *In the case  $n = 2$ , there are two other proofs for Theorem 1.1, which depends on the strict convex Lemma 4.1 and 4.2. With the two lemmas, one can show the solution is  $C^1$  up to boundary and is of super-linearity growth at infinite. This fact, together with partial Legendre transform and the Liouville theorem for harmonic functions, will gives us the first proof. As for the second proof, we need to use Wang's function[25] to construct supersolution/subsolution and so prove that the  $u$  is  $C^{1,1}$  up to boundary, then by blowing up at infinity to obtain  $D^2u = c\mathcal{I}$ .*

As the end of this paper, we give the two proofs in more details.

**Proof 1.** We write a point in  $\mathbb{R}^2$  as  $(x, y)$  and use the classical partial Legendre transform in the  $x$ -variable. For any  $p \in R$ , define

$$u^*(p, y) := \sup_{x \in R} \{px - u(x, y)\}.$$

The uniformly strict convexity of  $u$  at boundary and the smoothness of  $u$  at interior also implies the superlinearity at infinite, which guarantees the uniqueness and existence of  $x = X(p, y)$  such that the above supremum is attained.

By a calculation the two dimensional case of problem (1.1),

$$\det D^2u = 1 \text{ in } \mathbb{R}_+^2, \quad D_2u(x, 0) = ax \text{ on } R$$

is turned to the problem for  $u^*$ :

$$\begin{cases} \Delta u^* = 1 & \text{in } \mathbb{R}_+^2 \\ D_{v^*}u^* = 0 & \text{on } \mathbb{R} \end{cases},$$

where  $v^* = e_2 + ae_1$ . Differentiating  $u^*$  with respect to  $v^*$  and letting  $V = D_{v^*}u^*$  we obtain

$$\begin{cases} \Delta V = 0 & \text{in } \mathbb{R}_+^2 \\ V = 0 & \text{on } \mathbb{R} \end{cases}.$$

Note that

$$V = D_{v^*}u^*(p, y) = aX(p, y) - u_2(X, y) = u_2(X, 0) - u_2(X, y) \leq 0.$$

We can use the Liouville theorem for harmonic functions to see that the only solutions to above problem are

$$V(x, y) = D_{v^*}u^*(x, y) \equiv A$$

for some constant  $A$ . Then

$$u^*(p, y) = g(p - ay) + Ay$$

for some function  $g : [0, \infty) \rightarrow R$ . Recalling  $\Delta u^* = 1, D_{v^*}u^* = 0$ , we have

$$g(0) = 0 \text{ and } (1 + a^2)g'' = 1,$$

which implies that  $u^*$  is a quadratic polynomial, and so is the  $u$ .

**Proof 2.** As we have said, when  $n = 2$ , one could use strict convexity Lemma 4.1 and 4.2 to show that the solution  $u$  to problem (1.1) is uniformly  $C^{1,\alpha}$  up to boundary[4].

We consider Wang's singular function with bounded positive Monge-Ampère measure for large  $a \gg 2$ :

$$W_{a,b}(p, q) = \begin{cases} p^a + \frac{a^2-1}{a(a-2)}p^{a-\frac{2a}{b}}q^2, & |p|^a \geq |q|^b \\ \frac{4a-5}{2(a-2)}q^b + \frac{1}{2a}q^{b-\frac{2b}{a}}p^2, & |q|^b \geq |p|^a \end{cases} .$$

Let

$$W(x_1, x_2) = CaW_{a,b}(\tau x_1, x_2/\tau),$$

where  $\tau > 0$  is small and satisfies  $a\tau^a \leq c$ . One can use this function to construct a supersolution ( $W^+$ )/subsolution ( $W^-$ ) at boundary point:

$$\begin{cases} w^+(x) = u(x) - W(x_1, x_2) + Mx_2 \\ w^-(x) = \epsilon_1(W(x_1, x_2) - u(x) + \epsilon_2x_2) \end{cases} .$$

This will give the uniform estimate  $c \leq D_{nn}u \leq C$  on the boundary. Therefore  $u$  is  $C^{1,1}$  and then in  $C^{2,\alpha}$ . After blowing up at infinite, this together with the superlinearity will imply  $D^2u = c\mathcal{I}$  and the proof is completed.

## REFERENCES

- [1] Bao J.G., Chen J.Y., Guan B. and Ji M., Liouville property and regularity of a Hessian quotient equation, *American J. Math.* **125**(2003), 301-316.
- [2] Caffarelli L.A., A localization property of viscosity solution to the Monge-Ampère equation and their strict convexity, *Ann. of Math.* **131** (1990), 129-134.
- [3] Caffarelli L. A., Interior  $W^{2,p}$  estimates for solutions of Monge-Ampère equation, *Ann. of Math.* (2) **131** (1990), 135-150.
- [4] Caffarelli L.A., Some regularity properties of solutions of Monge Ampère equation, *Comm. pure Appl. Math.* **44** (1991), 965-969.
- [5] Caffarelli L.A. and Li Y.Y., A Liouville theorem for solutions of the Monge-Ampère equation with periodic data, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **21** (2004), 97-120.
- [6] Caffarelli L.A. and Li Y.Y., An extension to a theorem of Jörgens, Calabi, and Pogorelov, *Comm. Pure Appl. Math.* **56** (2003), 549-583.
- [7] Calabi E. , Improper affine hyperspheres of convex type and a generalization of a theorem by K. Jörgens, *Mich. Math. J.* **5** (1958), 105–126.
- [8] Cheng, S.Y., Yau, S.T., On the regularity of the Monge-Ampère equation  $\det \frac{\partial^2 u}{\partial x_i \partial x_j} = F(x, u)$ , *Comm. Pure Appl. Math.* **30** (1977), 41-68.
- [9] Chou K.S. and Wang X.-J., Entire solutions to Monge Ampère equations, *Comm. Pure Appl. Math.* **49** (1996), 529-539.
- [10] Figalli A., The Monge-Ampère equation and its Applications, European Math Soc Publ House, CH-8092 Zurich, Switzerland, 2017.
- [11] Gutiérrez, C.E., The Monge-Ampère equation, Birkhäuser, Boston, 2001.
- [12] John F., Extremum problems with inequalities as subsidiary conditions, Studies and Essays Presented to R. Courant on his 60th Birthday, p.187-204, Interscience Publishers, New York, 1948.
- [13] Jörgens K., Über die Lösungen der Differentialgleichung  $rt - s^2 = 1$ , *Math. Ann.* **127** (1954), 130-134.
- [14] Jian H.Y and Tu X.S., Regularity for Monge-Ampère Equations with Neumann Data, Preprint, May 2021.
- [15] Jian H.Y. and Wang X.-J., Continuity estimates for the Monge-Ampère equation, *SIAM J. Math. Anal.* **39** (2007), 608-626.
- [16] Jian H.Y. and Wang X.-J., A Bernstein theorem for the Monge-Ampère equation and regularity of affine spheres, *J. Diff. Geom.* **93** (2013), 431-469,
- [17] Jian H.Y. and Wang X.-J., Existence of entire solutions to the Monge-Ampère equation, *American J. Math.* (4) **136** (2014), 1093-1106.
- [18] Savin O., pointwise  $C^{2,\alpha}$  estimates at the boundary for the Monge-Ampère equation, *J. Amer. Math. Soc.* (1) **26**, 63-99 (2013).
- [19] Savin O., A localization theorem and boundary regularity for a class of degenerate Monge-Ampère equations, *J. Differ. Equ.* (2) **256** (2014), 327-388.
- [20] Pogorelov A.V., The Minkowski multidimensional problem, J. Wiley, New York, 1978.
- [21] Pogorelov A.V., An analog of Minkowski's problem for infinite complete convex hypersurfaces, *Soviet Math. Dokl.* **21** (1980), 138-140.
- [22] Trudinger, N.S., Wang, X.-J., The Bernstein problem for and affine maximal hypersurfaces, *Invent. Mat.* **140** (2) (2000) , 399-422.

- [23] Trudinger, N.S., Wang, X.-J., Boundary regularity for the Monge-Ampere and affine maximal surface equations, *Ann. Math.* (2) 167 (2008), 993-1028.
- [24] Yuan Y., A Bernstein problem for special Lagrangian equations, *Invent. Mat.* **150** (2002), 117-125.
- [25] Wang X.-J., Some counterexamples to the regularity of Monge-Ampère equations, *Proc. Amer. Math. Soc.* **123** (3) (1995), 841-845.

HUAIYU JIAN: DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, CHINA.

XUSHAN TU: DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, CHINA.

*Email address:* `hjian@tsinghua.edu.cn`; `txs17@mails.tsinghua.edu.cn`.