

On an affinity principle by Krasnoselskii

P. Amster and J. Epstein

Departamento de Matemática
Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires
& IMAS-CONICET
Ciudad Universitaria - Pabellón I, 1428, Buenos Aires, Argentina

Abstract

An abstract formulation of a duality principle established by Krasnoselskii is presented. Under appropriate conditions, it shall be shown that, if the solutions of a nonlinear functional equation can be obtained by finding fixed points of certain operators in possibly different Banach spaces, then these operators share some topological properties.

1 Introduction

Let us consider the T -periodic problem for the system

$$x'(t) = f(t, x(t)) \quad (1)$$

where $f : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is smooth and T -periodic in its first coordinate. As is well known, there are several approaches to search solutions as fixed points of certain mappings, specifically:

1. **In the phase space.** Define the Poincaré map $P : \text{dom}(P) \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by $P(x_0) := \Phi(T, x_0)$. Here, Φ denotes the associated flow, that is, $\Phi(t, x_0) := x(t)$, where x is the unique solution of (1) satisfying the initial condition $x(0) = x_0$. Due to the periodicity of f , if x_0 is a fixed point of P then the corresponding solution $x(t)$ is globally defined and T -periodic. Conversely, if $x(t)$ is a T -periodic solution of (1), then $x_0 := x(0) \in \text{dom}(P)$ and $P(x_0) = x_0$.

2. **In a functional space.** Fix an appropriate Banach space X , e.g.

$$X = C[0, T] := C([0, T], \mathbb{R}^n)$$

or

$$X = C_T := \{x \in C(\mathbb{R}, \mathbb{R}^n) : x(t+T) \equiv x(t)\}$$

and define an operator $K : X \rightarrow X$ in such a way that the fixed points of K can be identified with the set of T -periodic solutions of (1). A typical choice of such operators is $K_1 : C[0, T] \rightarrow C[0, T]$ given by

$$K_1(x)(t) := x(T) + \int_0^t f(s, x(s)) ds. \quad (2)$$

In both situations, solutions can be found by the use of different fixed points theorems or, more generally, employing degree theory. To this end, one may try, in the first case, to find an open bounded set U such that $\overline{U} \subset \text{dom}(P)$ and

$$\deg_B(I - P, U, 0) \neq 0,$$

where \deg_B denotes the Brouwer degree and, in the second case, an open bounded set $U_X \subset X$ such that

$$\deg_{LS}(I - K, U_X, 0) \neq 0,$$

where \deg_{LS} is now the (infinite dimensional) Leray-Schauder degree.

At this point, it is worthy recalling the following paragraph of [1]:

Every approach to the specific problem in question has its advantages and drawbacks, and therefore the picture we obtain when investigating a problem becomes more complete if we manage to find internal connections between different approaches and use the positive aspects of each of them.

This is the goal of the so-called theory of affinity, whose origin can be traced back to the works of M. A. Krasnoselskii in the early sixties (see [1, 3, 4] and the references therein). In the previous context, a remarkable result is a relatedness principle, also called *duality principle* (see [5]) which, roughly speaking, says that, despite their different nature, the mapping $I - P$ and the operator $I - K$ have the same degree. In more rigorous terms, it is required that the sets U and U_X have *common core*, which means:

1. P and K have no fixed points on ∂U and ∂U_X respectively.
2. If $x_0 \in U$ is a fixed point of P , then the corresponding solution $x(t)$ satisfies $\alpha(x) \in U_X$, where α maps the T -periodic functions with their related elements of X . The definition of α depends on the choice of X ; for example, if $X = C_T$ then $\alpha(x) := x$ and if $X = C[0, T]$ then $\alpha(x) := x|_{[0, T]}$.
3. If $x \in U_X$ is a fixed point of K , then $x_0 := x(0) \in U$.

In other words, when $X = C[0, T]$, the last two properties say that if a T -periodic solution has initial value in U then its restriction to $[0, T]$ belongs to U_X and vice versa.

In this context, one of the early results by Krasnoselskii reads

$$\deg_B(I - P, U, 0) = \deg_{LS}(I - K_1, U_1, 0),$$

where K_1 is the operator defined by (2) and the open bounded sets $U \subset \mathbb{R}^n$ and $U_1 \subset C[0, T]$ have common core. This equivalence can be extended to other operators, both in $C[0, T]$ and C_T or a variety of spaces, according to the equation under study. The proof relies on the homotopy invariance of the degree; however, this is not straightforward because K_1 and P are defined in different spaces. In order to avoid this difficulty, the key idea consists in noticing that, if one identifies \mathbb{R}^n with the subset of constant functions of $C[0, T]$, then P can be identified with a compact operator K_2 which is, in turn, homotopic to K_1 . Then, the proof follows from the fact that, since $\text{Im}(K_2) \subset \mathbb{R}^n$, the Leray-Schauder degree of $I - K_2$ coincides with the Brouwer degree of $(I - K_2)|_{\mathbb{R}^n}$. Regarded from an abstract point of view, this procedure can be understood as a conjugacy between operators and, as we shall see, this allows to extend the relatedness properties to other situations.

The main goal of this paper is to give a general version of Krasnoselskii relatedness principle in abstract Banach spaces in such a way that, when applied to the specific cases treated in the literature, the original results are retrieved. We refer to [3] for a very complete exposition, including various boundary conditions. The scope of the applications is not limited to ODEs: we shall present simple examples that include a nonlocal boundary problem for an elliptic PDE and a system of delay differential equations.


The paper is organized as follows. In the next section, we shall present the abstract setting of the Krasnoselskii results. Section 3 is devoted to the particular problem $L(x) = N(x)$ which includes a wide range of semilinear equations and is frequently treated in the context of the coincidence degree theory [2]. Finally, some applications are given in Section 4. In particular, a relatedness principle for a delay equation shall not be deduced from the problem $L(x) = N(x)$ but, instead, from the abstract formulation given in section 2. Finally, in section 5 we introduce further comments and pose some questions that shall be treated in subsequent works.

Throughout the paper we shall use the generic notation ‘deg’ to refer to the topological degree and reserve \deg_B and \deg_{LS} only for those cases in which we want to emphasize the fact that we are working in a finite or an infinite dimensional space, respectively. Also, since we shall be dealing with different spaces, in the second section we shall use, for the sake of clarity, the notation id_X to refer to the identity operator in the space X .

2 The abstract result

In this section, we shall prove an abstract theorem that compares the degree of certain operators in a general context. In the first place, let us establish an useful lemma:

Lemma 2.1. (*Conjugacy lemma*) *Let X and \tilde{X} be Banach spaces, let S be a subset of X and let $U \subset \tilde{X}$ be bounded and open such that $S \cap \partial U = \emptyset$. Consider the diagram*



where:

- π and i are linear and continuous, with $\pi \circ i = \text{id}_X$,
- K is compact such that $\text{fp}(K) = S$, where fp stands for the set of fixed points of a mapping,
- $\tilde{K} = i \circ K \circ \pi$
- $V_r = \pi^{-1}(U) \cap B(0, r)$.

Then $\text{fp}(\tilde{K}) = i(S)$. Moreover, there exists $r_0 > 0$ such that if $r > r_0$ then $\partial V_r \cap i(S) = \emptyset$ and

$$\deg(\text{id}_X - K, U, 0) = \deg(\text{id}_{\tilde{X}} - \tilde{K}, V_r, 0).$$

Proof. It is readily seen that $i \circ \pi : \tilde{X} \rightarrow \tilde{X}$ is a projector; thus, the subspace $E := i(X) \subset \tilde{X}$ is closed and $(i \circ \pi)|_E = \text{id}_E$. Let $\tilde{x} \in \text{fp}(\tilde{K})$ and $x := \pi(\tilde{x})$, then

$$\tilde{x} = (i \circ K \circ \pi)(\tilde{x}).$$

Because $\tilde{x} \in E$, it follows that $\tilde{x} = i(\pi(\tilde{x})) = i(x)$ and hence

$$x = (\pi \circ i)(x) = \pi(\tilde{x}) = (K \circ \pi)(\tilde{x}) = K(x).$$

Thus, $x \in S$ and, consequently, $\tilde{x} \in i(S)$. Conversely, assume that $\tilde{x} = i(x)$ for some x such that $x = K(x)$, then $\pi(\tilde{x}) = x$ and

$$\tilde{x} = i(x) = i(K(x)) = i(K(\pi(\tilde{x}))) = \tilde{K}(\tilde{x}).$$

Since U is bounded, we may fix r_0 such that $i(U) \subset B(r_0, 0)$ and take $r > r_0$. It is clear that $\partial V_r \cap i(S) = \emptyset$, $V_r \cap i(S) = i(S \cap U)$ and $V_r \cap E = i(U)$. Moreover, using the properties of the Leray-Schauder degree, we conclude:

$$\begin{aligned} \deg(\text{id}_{\tilde{X}} - \tilde{K}, V_r, 0) &= \deg((\text{id}_{\tilde{X}} - \tilde{K})|_E, V_r \cap E, 0) \\ &= \deg(i \circ (\text{id}_X - K) \circ \pi, i(U), 0) \\ &= \deg(\text{id}_X - K, U, 0). \end{aligned}$$

□

As a corollary, we obtain the following

Proposition 2.2. *In the situation of the previous lemma, let $\tilde{U} \subset \tilde{X}$ be open and bounded such that $\partial\tilde{U} \cap i(S) = \emptyset$ and $\tilde{U} \cap i(S) = i(S \cap U)$. Then*

$$\deg(\text{id}_{\tilde{X}} - \tilde{K}, \tilde{U}, 0) = \deg(\text{id}_X - K, U, 0).$$

Proof. Let V_r be as in the previous lemma, then for $r \gg 0$ we know that

$$\deg(\text{id}_{\tilde{X}} - \tilde{K}, V_r, 0) = \deg(\text{id}_X - K, U, 0).$$

Because the fixed points of \tilde{K} in \tilde{U} coincide with the fixed points of K in V_r , the proof follows from the excision property of the degree. \square

In what follows, we shall consider diagrams \mathcal{B} of the form

$$\begin{array}{ccc} & S \subset X_0 & \\ \alpha_1 \swarrow & & \searrow \alpha_2 \\ \alpha_1(S) \subset X_1 & \xrightleftharpoons[\mu]{\pi} & X_2 \supset \alpha_2(S) \end{array}$$

where:

- X_0, X_1, X_2 are Banach spaces.
- α_1, α_2 are linear and continuous, μ is compact, π is linear, continuous and onto.
- Restricted to S , $\alpha_1(S)$ and $\alpha_2(S)$, the diagram is commutative, namely:

$$\pi \circ \alpha_1(x) = \alpha_2(x), \quad \mu \circ \alpha_2(x) = \alpha_1(x)$$

for every $x \in S$.

In the applications, S shall be the set of solutions of a differential equation under certain boundary condition, X_0 a space of functions satisfying the boundary condition and X_1, X_2 specific spaces in which solutions can be found as fixed points of accurate operators. For example, in the periodic case we may take $X_0 = C_T$, X_1 as the space of continuous functions over one period and X_2 as the phase space.

Definition 2.3. *Let $U_i \subset X_i$ be open and bounded and consider the sets $S_i := \alpha_i(S) \cap U_i$ with $i = 1, 2$. We shall say that U_1 and U_2 have **common core** with respect to the diagram \mathcal{B} if*


1. $\alpha_i(S) \cap \partial U_i = \emptyset$
2. There exists $S_0 \subset S$ such that $\alpha_i(S_0) = S_i$.

Remark 2.4. In particular, the last condition ensures that

$$(\mu \circ \pi)|_{S_1} = \text{id}_{S_1}, \quad (\pi \circ \mu)|_{S_2} = \text{id}_{S_2}.$$

In the previous setting, we may also consider fixed point operators $K_i : X_i \rightarrow X_i$, that is, such that $\text{fp}(K_i) = \alpha_i(S)$. The resulting diagram \mathcal{B} shall be called a **Krasnoselskii diagram** if:

1. U_i have common core with respect to \mathcal{B} .
2. K_i are fixed point operators.
3. π has a continuous right inverse $i \in L(X_2, X_1)$, that is: $\pi \circ i = \text{id}_{X_2}$.



As a special case, taking

$$K_1 := \mu \circ \pi, \quad K_2 := \pi \circ \mu$$

it is readily verified that $\alpha_i(S) \subset \text{fp}(K_i)$. Indeed, if $x = \alpha_1(\tilde{x})$, then

$$(\mu \circ \pi)(x) = (\mu \circ \pi \circ \alpha_1)(\tilde{x}) = (\mu \circ \alpha_2)(\tilde{x}) = \alpha_1(\tilde{x}) = x.$$

The inclusion $\alpha_2(S) \subset \text{fp}(K_2)$ is analogous.

This allows to give a direct proof of the following result, which can be regarded as a particular case of [4, Thm 26.1] for the composition of arbitrary operators in Banach spaces.

Theorem 2.5. In the previous situation, assume that \mathcal{B} is a Krasnoselskii diagram for $K_1 = \mu \circ \pi$ and $K_2 = \pi \circ \mu$. Then

$$\deg(\text{id}_{X_1} - K_1, U_1, 0) = \deg(\text{id}_{X_2} - K_2, U_2, 0)$$

Proof. Define $\tilde{K} := i \circ K_2 \circ \pi$, then, by Lemma 2.1, there exists r_0 such that, for $r > r_0$ and $V_r = \pi^{-1}(U_2) \cap B(0, r)$,

$$\deg(\text{id}_{X_1} - \tilde{K}, V_r, 0) = \deg(\text{id}_{X_2} - K_2, U_2, 0).$$

Fix $r > r_0$ such that $i(U_2)$ and U_1 are contained in $B(0, r)$ and consider the homotopy $H_\lambda := \lambda K_1 + (1 - \lambda)\tilde{K}$.

Claim: The homotopy is admissible, that is: $\text{fp}(H_\lambda) \cap \partial V_r = \emptyset$ for all $\lambda \in [0, 1]$. Indeed, let $\lambda \in [0, 1]$ and $x \in \overline{V_r}$ such that

$$x = \lambda K_1(x) + (1 - \lambda) \tilde{K}(x). \quad (3)$$

Because $\pi(V_r) = U_2$, it follows that $\pi(x) \in \overline{U_2}$. Next, apply π on both sides of (3) to obtain

$$\begin{aligned} \pi(x) &= \lambda (\pi \circ K_1)(x) + (1 - \lambda) (\pi \circ \tilde{K})(x) \\ &= \lambda (\pi \circ \mu \circ \pi)(x) + (1 - \lambda) (\pi \circ i \circ \pi \circ \mu \circ \pi)(x) \\ &= \lambda K_2(\pi(x)) + (1 - \lambda) K_2(\pi(x)) \\ &= K_2(\pi(x)). \end{aligned}$$

Thus, $\pi(x) \in \text{fp}(K_2|_{\overline{U_2}}) = S_2 \subset U_2$, whence $x \in \pi^{-1}(U_2)$. Moreover, it is seen from (3) that

$$\|x\| \leq \lambda \|K_1(x)\| + (1 - \lambda) \|\tilde{K}(x)\|.$$

Since U_1 and U_2 have common core, $\pi(x) = S_2 = \alpha_2(S_0)$, then

$$K_1(x) = \mu \circ \pi(x) \in \alpha_1(S_0) = S_1 \subset U_1 \subset B(0, r).$$

On the other hand, because $\pi(x) \in \text{fp}(K_2)$ we obtain:

$$\tilde{K}(x) = (i \circ K_2 \circ \pi)(x) = i(\pi(x)) \in i(U_2) \subset B(0, r).$$

We conclude that $\|x\| < r$ and the claim follows.

Thus, using the homotopy invariance of the degree, we deduce:

$$\deg(\text{id}_{X_1} - \tilde{K}, V_r, 0) = \deg(\text{id}_{X_1} - K_1, V_r, 0).$$

Finally, from the fact that $V_r \cap \alpha_1(S) = U_1 \cap \alpha_1(S)$ and the excision property it follows that

$$\deg(\text{id}_{X_1} - K_1, V_r, 0) = \deg(\text{id}_{X_1} - K_1, U_1, 0)$$


□

3 The problem $L(x) = N(x)$

Many semilinear boundary value problems for differential equations can be expressed in the form

$$L(x) = N(x) \quad (4)$$

as described by the following diagram:



Here,

- $L : D \rightarrow Y$ is a linear operator, where $D := \text{dom}(L) \subset X$ is dense and L is onto the Banach space Y .
- $\ker(L)$ is closed and there exists a continuous projector $\pi_{\ker(L)} : X \rightarrow X$ onto $\ker(L)$. It follows that $X = \ker(L) \oplus Z$, where $Z = \ker(\pi_{\ker(L)})$ is closed.
- $V : Y \rightarrow Z$ is linear and compact such that $L \circ V = \text{id}_Y$ and $\text{Im}(V) = Z$.
- N is continuous.

Remark 3.1. In particular, due to the Hahn-Banach Theorem, a continuous projector $\pi_{\ker(L)}$ always exists when $\ker(L)$ is finite dimensional. In some situations, it may occur that V is not compact, although all the results are still valid under the weaker assumption that $V \circ N$ is compact. This is analogous to what is called, in the context of the coincidence degree theory, the L -compactness of the operator N .

Proposition 3.2. In the previous setting, let $K_0 := \pi_{\ker(L)} + V \circ N$. Then the solutions of (4) are the fixed points of K_0 . In other words, (4) is equivalent to the equation

$$K_0(x) = x. \quad (5)$$

Proof. Let $x \in \text{fp}(K_0)$, then $x = \pi_{\ker(L)}(x) + (V \circ N)(x)$ and, applying L , we deduce that $L(x) = N(x)$. Conversely, let x be a solution of (4) and set $z := x - \pi_{\ker(L)}(x) \in Z$. It follows that $L(z) = L(x) = N(x)$; thus, $z = (V \circ N)(x)$, that is, $x = \pi_{\ker(L)}(x) + (V \circ N)(x)$. \square

Finally, we shall need a condition regarding the uniqueness of the solutions. This is the role of the following assumption, that can be interpreted as an abstract Gronwall-like inequality. Specifically, we shall say that the previous diagram satisfies the condition (J) if, for all $\eta \geq 0$,

$$x - y = \eta V(N(x) - N(y)) \implies x = y. \quad (J)$$

Special solutions

In this section, we shall analyze the problem of finding solutions of (4) that satisfy a certain linear condition, e. g. periodic solutions of an ODE or a DDE, Dirichlet or Neumann boundary conditions for ODEs or PDEs, etc.

Let us consider the problem

$$L(x) = N(x) \quad (6)$$

$$\delta(x) = 0 \quad (7)$$

where $\delta : X \rightarrow X$ is a bounded linear operator such that $\text{Im}(\delta) \subset \ker(L)$. As an example, we may consider the Dirichlet problem

$$x''(t) = f(x(t), x'(t)) \quad (8)$$

$$x(0) = x(1) = 0, \quad (9)$$

with $X = C^1[0, 1]$. Observe that $\ker(L) = \{b + ta : a, b \in \mathbb{R}^n\} \cong \mathbb{R}^{2n}$. Thus, if we define $\pi_{\ker(L)}(x) := x(0) + tx'(0) \cong (x(0), x'(0))$ and $\delta(x) = x(0) + tx(1)$, then (8)-(9) is written in the form (6)-(7).

Proposition 3.3. *In the previous situation, define $\pi := \pi_{\ker(L)} + \delta$ and $K := \pi + V \circ N$, then (6)-(7) is equivalent to the fixed point problem $K(x) = x$.*

Proof. Let $x \in \text{fp}(K)$, then $x = \pi(x) + (V \circ N)(x)$. Because $\text{Im}(\pi) \subset \ker(L)$, applying L we obtain $L(x) = N(x)$. On the other hand, if we apply $\pi_{\ker(L)}$ instead, we deduce that $\pi_{\ker(L)}(x) = \pi_{\ker(L)}(x) + \delta(x)$ which, in turn, implies $\delta(x) = 0$. Conversely, if x solves (6)-(7), then $x = \pi_{\ker(L)}(x) + (V \circ N)(x)$. Since $\delta(x) = 0$, this obviously implies $x = \pi_{\ker(L)}(x) + \delta(x) + (V \circ N)(x) = K(x)$. \square

Lemma 3.4. *Consider as before a diagram associated to (6)-(7) satisfying (J) and assume there exists $\mu : \ker(L) \rightarrow X$ continuous such that*

1. $\mu(\ker(L)) = \{x \in X : x \text{ is a solution of (6)}\}$.
2. $\pi_{\ker(L)} \circ \mu = \text{id}_{\ker(L)}$.

Let \mathcal{B} be the diagram given by

$$\begin{array}{ccc}
 & S \subset \ker(\delta) & \\
 \alpha_1 \swarrow & & \searrow \alpha_2 \\
 \alpha_1(S) \subset X & \xrightleftharpoons[\mu]{\pi} & \ker(L) \supset \alpha_2(S)
 \end{array}$$

where:

- $S = \{x \in X : x \text{ is a solution of (6)-(7)}\}$.
- $\pi = \delta + \pi_{\ker(L)}$
- α_1 denotes the inclusion of $\ker(\delta)$ in X
- $\alpha_2 = \pi_{\ker(L)}|_{\ker(\delta)}$.


Then \mathcal{B} commutes when restricted to S , $\alpha_1(S)$ and $\alpha_2(S)$.

Proof. Let $x \in S$, then

$$(\pi \circ \alpha_1)(x) = \pi(x) = \delta(x) + \pi_{\ker(L)}(x) = \pi_{\ker(L)}(x) = \alpha_2(x).$$

On the other hand, calling $k := \pi_{\ker(L)}(x)$, it is seen that $(\mu \circ \alpha_2)(x) = \mu(k)$. It follows from the definition $L(\mu(k)) = N(\mu(k))$ and $\pi_{\ker(L)}(\mu(k)) = k = \pi_{\ker(L)}(x)$. Because $x \in S$, it follows that $x - \mu(k) = V(N(x) - N(\mu(k)))$ and we deduce from (J) that $x = \mu(k) = (\mu \circ \alpha_2)(x)$. \square

Lemma 3.5. *In the situation of the previous Lemma, assume that the mapping $\pi = \delta + \pi_{\ker(L)}$ is onto. Furthermore, assume that $U_1 \subset X$ and $U_2 \subset \ker(L)$ are bounded open sets with common core with respect to \mathcal{B} . Consider the diagram \mathcal{K} given by*



with S , π , α_1 and α_2 as before and

- i is a right inverse of π
- $K = \pi + V \circ N$.
- $K_2 = \pi \circ \mu$.

Then \mathcal{K} is a Krasnoselskii diagram.

Proof. From the previous lemma, we only have to verify that K and K_2 are fixed point operators. This is clear for K , from Proposition 3.3 and the fact that α_1 is simply the inclusion. Since we already know that $\alpha_2(S) \subset \text{fp}(K_2)$, it only remains to prove that $\text{fp}(K_2) \subset \alpha_2(S)$. To this end, let us consider $k \in \text{fp}(K_2)$, that is, $k = (\pi \circ \mu)(k)$. Let $x = \mu(k)$, then x is a solution of (6) and satisfies $\alpha_2(x) = \pi_{\ker(L)}(x) = k$. Moreover,

$$k = \pi(\mu(k)) = (\pi_{\ker(L)} + \delta)(\mu(k)) = k + \delta(\mu(k)) = k + \delta(x).$$

We deduce that $\delta(x) = 0$, whence $x \in S$ and $k = \alpha_2(x) \in \alpha_2(S)$. \square

Lemma 3.6. *In the situation of the previous lemma, consider the diagram \mathcal{K}_1 obtained from \mathcal{K} by replacing the operator K by $K_1 := \mu \circ \pi$. Then \mathcal{K}_1 is a Krasnoselskii diagram.*

Proof. Let $x \in \text{fp}(K_1)$, that is $x = (\mu \circ \pi)(x) = \mu(\pi_{\ker(L)}(x) + \delta(x))$. Applying $\pi_{\ker(L)}$ at both sides, we obtain $\pi_{\ker(L)}(x) = \pi_{\ker(L)}(x) + \delta(x)$ and consequently $\delta(x) = 0$. Moreover, since $x = \mu(\pi(x))$, we deduce that x is a solution of (6) and hence $x \in S = \alpha_1(S)$. Thus, the proof is complete, because we already know that $\alpha_1(S) \subset \text{fp}(K_1)$. \square

We are now in condition of establishing the main result of this section.

Theorem 3.7. *In the situation of the previous lemmas, assume that μ is compact. Then*

$$\deg(I - K, U_1, 0) = \deg(I - K_2, U_2, 0).$$

Proof. From the previous lemma and Theorem 2.5, we know that

$$\deg(I - K_1, U_1, 0) = \deg(I - K_2, U_2, 0).$$

Thus, it suffices to verify that

$$\deg(I - K_1, U_1, 0) = \deg(I - K, U_1, 0).$$

To this end, let us define $H_\lambda := \lambda K + (1 - \lambda)K_1$ over the set $\overline{U_1}$ and suppose that $x \in \text{fp}(H_\lambda)$, that is

$$x = \lambda [\pi(x) + V \circ N(x)] + (1 - \lambda) (\mu \circ \pi)(x). \quad (10)$$

On the one hand, applying $\pi_{\ker(L)}$ we obtain:

$$\pi_{\ker(L)}(x) = \lambda \pi(x) + (1 - \lambda) \pi(x) = \pi(x) \implies \delta(x) = 0. \quad (11)$$

On the other hand, applying L instead, we obtain:

$$L(x) = \lambda N(x) + (1 - \lambda)(L \circ \mu \circ \pi)(x) \quad (12)$$

Let $y = (\mu \circ \pi)(x)$, then it follows from the definition of μ that $L(y) = N(y)$ and $\pi_{\ker(L)}(y) = \pi(x)$; moreover, because $\delta(x) = 0$, it follows that $\pi_{\ker(L)}(y) =$

$\pi_{\ker(L)}(x)$. Thus, writing (12) as $L(x) = \lambda N(x) + (1 - \lambda)L(y)$, we deduce the equality

$$L(x) - L(y) = \lambda(N(x) - N(y)), \quad (13)$$

that is

$$x - y = \lambda V(N(x) - N(y)) \quad (14)$$

and, from (J), we conclude that $x = y$ and $L(x) = N(x)$. This implies that $x \in S = \alpha_1(S)$ and hence $x \notin \partial U_1$. Thus, the proof follows from the homotopy invariance of the Leray-Schauder degree. \square

4 Applications


4.1 Periodic solutions of ODEs

Consider the problem

$$x'(t) = f(t, x(t)) \quad (15)$$

$$x(0) = x(T) \quad (16)$$

where $f : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous and T -periodic in t . For simplicity, we shall also assume that f is globally Lipschitz in x with constant l . A diagram for this equation is the following



where:

- $L : \text{dom}(L) \subset C[0, T] \rightarrow C[0, T]$ is given by $Lx = x'$ and $\text{dom}(L) = C^1[0, T]$.
- $N : C[0, T] \rightarrow C[0, T]$ is the Nemitskii operator associated to f , namely $N(x)(t) := f(t, x(t))$.
- $\ker(L) = \{x(t) \equiv c : c \in \mathbb{R}^n\} \cong \mathbb{R}^n$.
- $Z = \{x \in C[0, T] \mid x(0) = 0\}$.
- $V(x)(t) := \int_0^t x(s) ds$.
- $\pi_{\ker(L)}(x) \equiv x(0)$.

- $\delta(x) := x(T) - x(0)$

Here, we may consider the flow Φ associated to the system, that is $\Phi(t, x_0) = x(t)$, where x is the unique solution of (15) such that $x(0) = x_0$, and define $\mu(x_0)(t) := \Phi(t, x_0)$. Since $\ker(L) \cong \mathbb{R}^n$, we may assume $\mu : \ker(L) \rightarrow C[0, T]$. From the Arzelà-Ascoli Theorem, it is clear that μ is compact.

Lemma 4.1. *Condition (J) is satisfied.*

Proof. Let $x, y \in C[0, T]$ satisfy $(x - y)(t) = \eta \int_0^t (f(s, x(s)) - f(s, y(s))) ds$ for some $\eta \geq 0$, then

$$|x(t) - y(t)| = \eta \left| \int_0^t (f(s, x(s)) - f(s, y(s))) ds \right| \quad (17)$$

$$\leq \eta \int_0^t |f(s, x(s)) - f(s, y(s))| ds \quad (18)$$

$$\leq \eta l \int_0^t |x(s) - y(s)| ds. \quad (19)$$

From Gronwall's lemma, we deduce that $|x(t) - y(t)| = 0$ for all t . \square

Lemma 4.2. $\pi = \pi_{\ker(L)} + \delta$ is onto.

Proof. This is trivial from the definition, because $\pi(x) = x(T)$. \square

Thus, we may consider a diagram \mathcal{B} as in Lemmas 3.4 and 3.5 and assume that $U_1 \subset C[0, T]$ and $U_2 \subset \ker(L)$ have common core with respect to \mathcal{B} . Here, it is seen that

$$K(x)(t) = \pi(x) + (V \circ N)(x)(t) = x(T) + \int_0^t f(r, x(r)) dr$$

and

$$K_2(x_0) = (\pi \circ \mu)(x_0) = \Phi(T, x_0)$$

coincides, via the isomorphism $\ker(L) \cong \mathbb{R}^n$, with the Poincaré map. Then, by Theorem 3.7, the original result by Krasnoselskii mentioned in the introduction is retrieved:

$$\deg_{LS}(I - K, U_1, 0) = \deg_B(I - P, U_2, 0).$$

Moreover, observe that

$$K_1(x)(t) = (\mu \circ \pi)(x)(t) = \Phi(t, x(T)),$$

which yields another one of the equivalences presented in [3]:

$$\deg_{LS}(I - K_1, U_1, 0) = \deg_B(I - P, U_2, 0).$$

Other fixed point operators

In the previous example, it is observed that the operator $\pi(x) = x(T)$ is a projector onto $\ker(L)$. With this in mind, we may extend the definition of K to a more general family of fixed point operators:

Lemma 4.3. *In the previous situation, let $\gamma : C[0, T] \rightarrow C[0, T]$ be a projector onto $\ker(L)$ and define the (compact) operator*

$$K_\gamma := \gamma + [(\pi - \gamma) + \text{id}_X] \circ V \circ N$$

Then K_γ is a fixed point operator respect to \mathcal{B} and

$$\deg_{LS}(I - K_\gamma, U_1, 0) = \deg_B(I - P, U_2, 0).$$

Proof. Let $x \in \text{fp}(K_\gamma)$, then

$$x = \gamma(x) + [(\pi - \gamma) \circ V \circ N](x) + (V \circ N)(x) \quad (20)$$

and applying L we obtain $L(x) = N(x)$. On the other hand, applying γ we obtain

$$\gamma(x) = \gamma(x) + ((\pi - \gamma) \circ V \circ N)(x) + (\gamma \circ V \circ N)(x)$$

and hence

$$(\pi \circ V \circ N)(x) = 0.$$

Finally, applying $\pi_{\ker(L)}$, from the previous equality and employing the fact that $\pi_{\ker(L)} \circ V \equiv 0$ we obtain

$$\pi_{\ker(L)}(x) = \gamma(x) - (\gamma \circ V \circ N)(x).$$

In the same way, because π is a projector we also obtain from (20) that

$$\pi(x) = \gamma(x) - (\gamma \circ V \circ N)(x)$$

and consequently $\delta(x) = \pi(x) - \pi_{\ker(L)}(x) = 0$, that is, $x \in \alpha_1(S)$.

Conversely, let $x \in \alpha_1(S)$ and write $C[0, T] = \ker(\gamma) \oplus \text{Im}(\gamma) = \ker(\gamma) \oplus \ker(L)$. The operator $V_\gamma := V - \gamma \circ V$ is a right inverse of L with range in $Z_\gamma := \ker(\gamma)$, then the equation $L(x) = N(x)$ is equivalent to $x = \gamma(x) + (V_\gamma \circ N)(x)$, that is

$$x = \gamma(x) + (V \circ N)(x) - (\gamma \circ V \circ N)(x). \quad (21)$$

Moreover, we know from Proposition 3.3 that

$$x = \pi(x) + (V \circ N)(x)$$

and applying π we deduce that $(\pi \circ V \circ N)(x) = 0$. Combined with (21), this implies

$$x = \gamma(x) + [(\pi - \gamma) \circ V \circ N](x) + (V \circ N)(x)$$

and hence $x \in \text{fp}(K_\gamma)$.

Next, consider the homotopy $H_\lambda = \lambda K + (1 - \lambda)K_\gamma$ with $\lambda \in [0, 1]$. It is clear that $H_\lambda = K_{\tilde{\gamma}}$ for $\tilde{\gamma} = \lambda\pi + (1 - \lambda)\gamma$, which is also a projector. Thus,

$\text{fp}(H_\lambda) = \alpha_1(S)$, which does not intersect ∂U_1 ; hence the degree is well defined along the homotopy and

$$\deg_{LS}(I - K_\gamma, U_1, 0) = \deg_{LS}(I - K, U_1, 0) = \deg_B(I - P, U_2, 0).$$

□

A different operator can be obtained by simply noticing that, if $L(x) = N(x)$, then x satisfies the boundary condition if and only if $\overline{N(x)} = 0$, where $\overline{\varphi}$ denotes the average of a continuous function φ , namely $\overline{\varphi} := \frac{1}{T} \int_0^T \varphi(t) dt$. This motivates the definition of

$$K_3(x) := \overline{x} + T\overline{N(x)} + V[N(x) - \overline{N(x)}] - \overline{V[N(x) - \overline{N(x)}]},$$

which is clearly compact and, as one easily verifies, it is a fixed point operator.

Corollary 4.3.1. *In the previous situation,*

$$\deg_{LS}(I - K_3, U_1, 0) = \deg_B(I - P, U_2, 0).$$

Proof. Let $K_4(x) := \overline{x} + T\overline{N(x)} + V(N(x)) - \overline{V(N(x))}$, then $K_4 = K_\gamma$ for the projector $\gamma(x) := \overline{x}$. Then K_4 is a fixed point operator and

$$\deg_{LS}(I - K_4, U_1, 0) = \deg_B(I - P, U_2, 0).$$

Next, consider the homotopy

$$H_\lambda(x) = \overline{x} + T\overline{N(x)} + V\left(N(x) - \lambda\overline{N(x)}\right) - \overline{V\left(N(x) - \lambda\overline{N(x)}\right)},$$

which is also a fixed point operator for all $\lambda \in [0, 1]$. As before, we deduce that H_λ is admissible and $H_0 = K_4$, whence

$$\deg_{LS}(I - K_3, U_1, 0) = \deg_{LS}(I - K_4, U_1, 0) = \deg_B(I - P, U_2, 0).$$

□

The latter operator is closely related to the standard Lyapunov-Schmidt decomposition for the problem $x'(t) = f(t, x(t))$ in C_T , which can be expressed as

$$\overline{N_T x} = 0, \quad x - \overline{x} = \tilde{V}(N_T x - \overline{N_T x})$$

where $N_T : C_T \rightarrow C_T$ is given as before by $N_T(x)(t) := f(t, x(t))$ and the linear operator \tilde{V} is defined for $\varphi \in C_T$ such that $\overline{\varphi} = 0$ as the unique $x \in C_T^1$ such that $x' = \varphi$ and $\overline{x} = 0$. Thus, the problem is equivalently written as

$$x = \overline{x} + T\overline{N_T(x)} + \tilde{V}(N_T x - \overline{N_T x}) := K_5(x).$$

In order to prove the relatedness principle for this operator, it proves convenient to notice that

$$K_5(x) = K_3(x|_{[0, T]})$$

extended T -periodically to \mathbb{R} . Moreover, we may define

$$\tilde{S} := \{x \in C_T : x \text{ is a solution of (6)}\} = \text{fp}(K_5)$$

and recall that S is the set of solutions of (6)-(7) in $X = C[0, T]$.

Proposition 4.4. *Let $\tilde{U} \subset C_T$ and $U \subset C[0, T]$ be open bounded sets such that*

- $\partial\tilde{U} \cap \tilde{S} = \emptyset$ and $\partial U \cap S = \emptyset$.
- For $x \in C_T$,
$$x \in \tilde{U} \cap \tilde{S} \iff x|_{[0, T]} \in U \cap S.$$

Then

$$\deg_{LS}(I - K_3, U, 0) = \deg_{LS}(I - K_5, \tilde{U}, 0).$$

Proof. Let $X_0 = \ker(\delta) = \{x \in C[0, T] : x(0) = x(T)\}$, then $\text{Im}(K_3) \subset X_0$. From the properties of the degree, we know that

$$\deg_{LS}(I - K_3, U, 0) = \deg_{LS}((I - K_3)|_{X_0}, U \cap X_0, 0).$$

Next, set $\tilde{\pi} : C_T \rightarrow X_0$ as the restriction and $\tilde{i} : X_0 \rightarrow C_T$ as the T -periodic extension, then the result is a direct consequence of Proposition 2.2 and the fact that $K_5 = \tilde{i} \circ K_3 \circ \tilde{\pi}$. \square

Let us consider now the fixed point operator defined as follows. For arbitrary $\eta \neq 0$, let $K^\eta : C_T \rightarrow C_T$ be given by $K^\eta(x) = y$, where x is the unique T -periodic solution of the problem

$$y'(t) + \eta y(t) = f(t, x(t)) + \eta x(t).$$

In order to analyze the affinity with respect to the preceding operators, let us consider firstly the case $\eta > 0$ and take again the linear homotopy

$$H_\lambda(x) := \lambda K^\eta(x) + (1 - \lambda)K_5(x).$$

Suppose $x = H_\lambda(x)$, then taking average we obtain

$$\bar{x} = \lambda \left[\bar{x} + \frac{\overline{N_T x}}{\eta} \right] + (1 - \lambda)[\bar{x} + T \overline{N_T x}]$$

that is

$$0 = (1 - \lambda)T \overline{N_T(x)} + \lambda \frac{\overline{N_T(x)}}{\eta},$$

whence $\overline{N_T x} = 0$. Setting $y = K^\eta(x)$, it follows that

$$x' = N_T(x) + \lambda \eta (x - y)$$

and

$$y' = N_T(x) + \eta (x - y).$$

Thus,

$$(x - y)' = (\lambda - 1)(x - y).$$

When $\lambda < 1$, the fact $x - y \in C_T$ implies $x = y$ and hence $x' = N_T(x)$; on the other hand, the case $\lambda = 1$ corresponds to a fixed point of K^η , so we also

deduce that $x' = N_T(x)$. In consequence, x is a solution and K^η is homotopic to K_5 , provided that there are no solutions on $\partial\tilde{U}$.

Now assume, instead, that $\eta < 0$, then we may define

$$\hat{K}_5(x) := \bar{x} - T\overline{N_T(x)} + \tilde{K}(N_T(x) - \overline{N_Tx})$$

and it follows exactly as before that K^η is homotopic to \hat{K}_5 . Furthermore, if we define \hat{K}_3 by changing in K_3 the sign of the term $T\overline{N_T(x)}$ and set U and \tilde{U} as before, then

$$\deg_{LS}(I - \hat{K}_5, \tilde{U}, 0) = \deg_{LS}(I - \hat{K}_3, U, 0).$$

Next we may take, in the original diagram, $\delta(x) := x(0) - x(T)$. Then the mapping

$$\pi(x) = 2x(0) - x(T)$$

is, as before, a projector and taking $\gamma(x) = \bar{x}$ we obtain the corresponding operators

$$K_\gamma(x) = \bar{x} + [-V(N(x))(T) - \overline{V(N(x))} + V(N(x))] = \hat{K}_3(x)$$

and

$$K_2(x_0) = 2x_0 - \Phi(T, x_0).$$

Thus,

$$\deg_{LS}(I - \hat{K}_3, U, 0) = \deg_B(I - K_2, U_2, 0).$$

Moreover, observe that

$$I - K_2(x_0) = \Phi(T, x_0) - x_0 = -(I - P)(x_0),$$

whence

$$\deg_B(I - K_2, U_2, 0) = (-1)^n \deg_B(I - P, U_2, 0).$$

Summarizing, we have established:

Proposition 4.5. *Let U and \tilde{U} be as before, then*

$$\deg_{LS}(I - K^\eta, \tilde{U}, 0) = \text{sgn}(\eta)^n \deg_{LS}(I - K_3, U, 0).$$


4.2 Second order equation with Dirichlet conditions

Let us consider the problem

$$x'' = f(t, x(t)) \tag{22}$$

$$x(0) = x(1) = 0, \tag{23}$$

where $f : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous and Lipschitz in x with constant l . A diagram for this equation reads



where:

- $X = C^1[0, 1]$ and $Lx := x''$ is defined in $\text{dom}(L) = C^2[0, T]$.
- $N : X \rightarrow X$ is the Nemitskii operator $N(x)(t) := f(t, x(t))$.
- $\ker(L) = \{x(t) = ta + b : a \in \mathbb{R}^n, b \in \mathbb{R}^n\} \cong \mathbb{R}^n \times \mathbb{R}^n$.
- $Z = \{x \in C^1[0, 1] : x(0) = x'(0) = 0\}$.
- $Vx(t) = \int_0^t \int_0^r x(s) ds dr$.
- $\pi_{\ker(L)}(x) = tx'(0) + x(0) \cong (x'(0), x(0))$.
- $\delta(x) = tx(1) + x(0) \cong (x(1), x(0))$.

In this setting, we may define $\mu(ta + b)$ as the unique solution of (22) with initial condition $x(0) = b$, $x'(0) = a$. It is clear that μ is well defined and compact.

The following lemma is a direct consequence of Gronwall's inequality:

Lemma 4.6. *Assume that $u \in C[0, 1]$ satisfies*

$$0 \leq u(t) \leq c^2 V(u)(t) \quad \forall t \in [0, 1]. \quad (24)$$

Then $u \equiv 0$.

Proof. It suffices to observe that $V(u)(t) = \int_0^t (t-s)u(s)ds$ and apply the standard Gronwall lemma. \square

As a consequence, we deduce that the preceding diagram satisfies (J). Indeed, if for some $\eta \geq 0$ one has

$$x(t) - y(t) = \eta \int_0^t \int_0^r [f(s, x(s)) - f(s, y(s))] ds dr,$$

then

$$|x(t) - y(t)| \leq \eta l \int_0^t \int_0^r |x(s) - y(s)| ds dr$$

and Lemma 4.6 applies.

Lemma 4.7. $\pi = \pi_{\ker(L)} + \delta$ is onto.

Proof. By definition, $\pi(x) = t[x(1) + x'(0)] + 2x(0)$. Given $y(t) = ta + b$, then $x(t) = t^2(a - \frac{b}{2}) + \frac{b}{2}$ satisfies $\pi(x) = y$. \square

In the previous setting, consider the diagram \mathcal{B} associated to (22)-(23) and assume that the open bounded sets $U_1 \subset C[0, T]$ and $U_2 \subset \ker(L)$ have common core with respect to \mathcal{B} . In the situation of Theorem 3.7, we have that

$$K(x)(t) = t[x(1) + x'(0)] + 2x(0) + \int_0^t \int_0^r f(s, x(s)) ds dr,$$

$$K_2(ta + b) = (\pi \circ \mu)(ta + b)$$

and it follows that

$$\deg_{LS}(I - K, U_1, 0) = \deg_B(I - K_2, U_2, 0).$$

Since we are dealing with a Dirichlet problem, it is worthy comparing the latter operator with the shooting operator $\mathcal{S} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by $\mathcal{S}(a) := x(1)$, where $x(t)$ is the unique solution of the equation such that $x(0) = 0$ and $x'(0) = a$. For convenience, we may write $\mathcal{S}(a) = \Phi^1(1, 0, a)$, where $\Phi(t, b, a) = (\Phi^1, \Phi^2)$ denotes the associated flow. Moreover, we may define the (open) mapping $\varphi : \ker(L) \rightarrow \mathbb{R}^n$ given by $\varphi(ta + b) := a$.

Proposition 4.8. *In the previous situation,*

$$\deg_B(\mathcal{S}, \varphi(U_2), 0) = \deg_B(I - K_2, U_2, 0).$$

Proof. Consider the subspace $W := \{ta : a \in \mathbb{R}^n\} \subset \ker(L)$ and define the function $g : \ker(L) \rightarrow W$ given by $g(ta + b) = t(a - \mathcal{S}(a))$. From the properties of the Brouwer degree,

$$\deg_B(I - g, U_2, 0) = \deg_B((I - g)|_W, U_2 \cap W, 0).$$

Because $(I - g)(ta) = t\mathcal{S}(a)$ and $\varphi|_W$ is an isomorphism, the latter degree is clearly equal to $\deg_B(\mathcal{S}, \varphi(U_2), 0)$. On the other hand, observe that

$$(I - K_2)(ta + b) = -(t\Phi^1(1, b, a) + b)$$

and hence

$$\deg_B(I - K_2, U_2, 0) = (-1)^{2n} \deg_B(F, U_2, 0) = \deg_B(F, U_2, 0),$$

where $F(ta + b) := t\Phi^1(1, b, a) + b$. Thus, it suffices to verify that F and $I - g$ are homotopic. To this end, suppose that

$$\lambda F(ta + b) + (1 - \lambda)(t\mathcal{S}(a) + b) = 0,$$

then

$$\lambda\Phi^1(1, a, b) + (1 - \lambda)\mathcal{S}(a) = 0 \quad \text{and} \quad b = 0.$$

Next, recall that $\Phi^1(1, a, 0) = \mathcal{S}(a)$; hence, we conclude that $\mathcal{S}(a) = 0$ and it follows from the assumptions that $ta \notin \partial U_2$. \square

Several other examples of fixed point operators for this problem shall be defined in section 5.

4.3 An elliptic problem with nonlocal conditions

Let us consider the elliptic system

$$\Delta u(x) = f(x, u(x)) \quad x \in \Omega \quad (25)$$

with the nonlocal condition

$$u|_{\partial\Omega} = \text{constant}, \quad \int_{\partial\Omega} \frac{\partial u}{\partial \nu} dS = 0. \quad (26)$$

Here, $\Omega \subset \mathbb{R}^d$ is a smooth bounded domain, $f : \overline{\Omega} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is smooth and we set

$$X := \{u \in C^{1,\alpha}(\overline{\Omega}) : u|_{\partial\Omega} = \text{constant}\}, \\ Y = C^\alpha(\overline{\Omega})$$

for some $\alpha \in (0, 1)$;

$L : D \rightarrow Y$ given by $Lu := \Delta u$, with $D := X \cap C^{2,\alpha}(\overline{\Omega})$;

$$N(u)(x) := f(x, u(x))$$

and $V : C^\alpha(\overline{\Omega}) \rightarrow D$ defined by $V\varphi := u$, the unique solution of the problem

$$\Delta u = \varphi, \quad u|_{\partial\Omega} = 0.$$

It is clear that $\ker(L)$ is the set of constant functions, which shall be identified with \mathbb{R}^n , and we may consider

$$\pi_{\ker(L)}(u) := u|_{\partial\Omega}$$

and

$$\delta(u) := \int_{\partial\Omega} \frac{\partial u}{\partial \nu} dS.$$

In order to define a function μ as before, we shall assume the following monotonicity assumption

$$\langle f(x, v) - f(x, u), v - u \rangle \geq -\kappa |u - v|^2 \quad (27)$$

where $\kappa < \lambda_1^{(D)}$, the first eigenvalue of $-\Delta$ for the Dirichlet conditions. Then $\mu : \mathbb{R}^n \rightarrow D$ is defined by $\mu(c) := u_c$, the unique solution of (25) such that $u|_{\partial\Omega} = c$. The fact that μ is well defined is known, and can be regarded as one of those “uniqueness implies existence” results. Indeed, if u and v are solutions of (25) such that $u = v$ on $\partial\Omega$, then

$$\langle \Delta(u - v), u - v \rangle_x = \langle f(\cdot, u) - f(\cdot, v), u - v \rangle_x \geq -\kappa |u(x) - v(x)|^2;$$

thus,

$$\int_{\Omega} |\nabla(u - v)(x)|^2 dx \leq \kappa \|u - v\|_{L^2}^2$$

and from the Poincaré inequality it follows that $u \equiv v$. Moreover, writing

$$\Delta u(x) = f(x, u(x)) - f(x, c) + f(x, c)$$

and multiplying by $u(x) - c$ it is seen that

$$\|\nabla(u - c)\|_{L^2}^2 \leq \kappa \|u - c\|_{L^2}^2 + k \|u - c\|_{L^2}^2$$

for some constant k . This yields *a priori* bounds for the solutions and the existence of u_c follows from a standard truncation argument. Observe, furthermore, that the same computation shows that condition (J) is satisfied: if $u - v = \eta V(N(u) - N(v))$ with $\eta \geq 0$, then $u = v = 0$ on $\partial\Omega$ and

$$\Delta(u - v)(x) = \eta(f(x, u(x)) - f(x, v(x))).$$

As before, it is deduced that $u \equiv v$.

In this setting, one has

$$\pi(u) = \pi_{\ker(L)}(u) + \delta(u) = u|_{\partial\Omega} + \int_{\partial\Omega} \frac{\partial u}{\partial \nu} dS,$$

which is clearly onto, and

$$K(u) = \pi(u) + (V \circ N)(u) = v,$$

where v is the unique solution of (25) such that

$$v|_{\partial\Omega} = u|_{\partial\Omega} + \int_{\partial\Omega} \frac{\partial u}{\partial \nu} dS.$$

It is readily seen that K is a fixed point operator. Moreover, we may also compute $K_1 = \pi \circ \mu : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $K_2 = \mu \circ \pi : X \rightarrow X$ explicitly as

$$K_1(c) = c + \int_{\partial\Omega} \frac{\partial u_c}{\partial \nu} dS,$$

$$K_2(u) = u_{\pi(u)},$$

that is, $K_2(u)$ is the unique solution of (25) whose value over the boundary of Ω is equal to $u|_{\partial\Omega} + \int_{\partial\Omega} \frac{\partial u}{\partial \nu} dS$. From the previous results, we deduce:

Proposition 4.9. *Assume that $U_1 \subset \mathbb{R}^n$ and $U_2 \subset X$ are open bounded sets with common core, then*

$$\deg_{LS}(I - K_2, U_2, 0) = \deg_{LS}(I - K, U_2, 0) = \deg_B(I - K_1, U_1, 0).$$

As in the preceding examples, several other fixed point operators can be defined; in particular, the one that arises on the Lyapunov-Schmidt reduction. In the first place, notice that π is a projector, so a fixed point operator K_γ can be defined in the same way as before. Thus, taking $\gamma(u) = \bar{u}$ yields the operator

$$\bar{u} + |\Omega| \overline{Nu} + V(N(u)) - \overline{V(N(u))}$$

which, in turn, is homotopic to

$$K_3(u) = \bar{u} + |\Omega| \overline{Nu} + V(N(u) - \overline{Nu}) - \overline{V(N(u) - \overline{Nu})}.$$

Finally, observe again that the latter operator is conjugated to the one defined by the same formula, after replacing D by

$$\text{dom}(L) = C^{2,\alpha}(\bar{\Omega}) \cap \delta^{-1}(0),$$

which corresponds to the Lyapunov-Schmidt decomposition and, as in section 4.1, may be called K_5 . In this case, if one considers the operators $K^\eta : u \mapsto v$, defined by

$$\Delta v(x) + \eta v(x) = f(x, u(x)) + \eta u(x)$$

where η is not an eigenvalue of $-\Delta$ for the conditions (26), then K^η is linearly homotopic to K_5 when $\eta > 0$. Interestingly, the interaction of η with the spectrum of $-\Delta$ is not relevant in this case: an intuitive explanation of this fact shall be given in section 5.

4.4 Delay differential equations

Let us consider the problem of finding periodic solutions to the delay equation

$$x'(t) = f(t, x(t), x(t - \tau)) \quad t \geq 0 \quad (28)$$

$$x_0 = x_T \quad (29)$$

where $f : \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous and Lipschitz in the last two variables and T -periodic in t , with $T \geq \tau$. Here, as usual, we employ the notation $x_t \in C[-\tau, 0]$ for the function defined by $x_t(s) := x(s + t)$; thus, a solution of (28) satisfying (29) is extended to a periodic solution of (28). In this case, we may consider the diagram \mathcal{B}_τ as follows

$$\begin{array}{ccc} & S \subset C_T & \\ \alpha_1 \swarrow & & \searrow \alpha_2 \\ \alpha_1(S) \subset C[0, T] & \xrightleftharpoons[\mu]{\pi} & C[-\tau, 0] \supset \alpha_2(S) \end{array}$$

where:

- $C_T := \{x \in C(\mathbb{R}, \mathbb{R}^n) : x(t + T) = x(t)\}$.
- $S \subset C_T$ is the set of T -periodic solutions of (28).
- α_1 denotes the restriction to the interval $[0, T]$.
- α_2 denotes the restriction to the interval $[-\tau, 0]$, that is $\alpha_2(x) := x_0$.
- $\pi(x) := x_T$

- $\tilde{\mu}(y)$ is the solution of (28) with initial condition $x_0 = y$, defined in $[-\tau, T]$ and $\mu(y) := \tilde{\mu}(y)|_{[0, T]}$.


It is easy to verify that μ is compact and π is linear and continuous. Furthermore, observe that it has a right inverse $i : C[-\tau, 0] \rightarrow C[0, T]$ given by

$$i(y)(t) = \mathbb{I}_{[T-\tau, T]}(t)y(t-T) + \mathbb{I}_{[0, T-\tau)}(t)y(-\tau)$$

where \mathbb{I} stands for the indicator function.

We claim that the diagram, restricted to S , $\alpha_1(S)$ and $\alpha_2(S)$, is commutative. Indeed, for $x \in S$ it is seen that $(\pi \circ \alpha_1)(x) = x_T$. Because $x \in C_T$, it follows that $x_T = x_0$ and hence $(\pi \circ \alpha_1)(x) = \alpha_2(x)$. On the other hand, $(\mu \circ \alpha_2)(x) = \mu(x_0)$, which defines the unique solution of (28) with initial condition x_0 , restricted to $[0, T]$. In other words, $(\mu \circ \alpha_2)(x) = x|_{[0, T]} = \alpha_1(x)$.

Next, consider the following diagram \mathcal{K}_τ given by



where:

- i is defined as before.
- $K_1 = \mu \circ \pi$ and $K_2 = \mu \circ \pi$.
- U_i have common core with respect to \mathcal{B}_τ .

Lemma 4.10. \mathcal{K}_τ is a Krasnoselskii diagram.

Proof. We already know that the linear continuous function i is a right inverse of π ; thus, it only remains to verify that K_1 and K_2 are fixed point operators. Let $x \in S$, then

$$K_1(\alpha_1(x)) = \mu(\pi \circ \alpha_1(x)) = \mu(x_T) = \mu(x_0) = x|_{[0, T]} = \alpha_1(x).$$

Conversely, let $\hat{x} \in \text{fp}(K_1)$, that is, $\hat{x} = (\mu \circ \pi)(\hat{x}) = \mu(\hat{x}_T)$. Evaluation at $t = 0$ yields

$$\hat{x}(0) = \mu(\hat{x}_T)(0) = \hat{x}_T(0) = \hat{x}(T).$$

Let us consider now the function $x : [-\tau, T]$ defined by $x|_{[-\tau, 0]} = \hat{x}_T$ and $x|_{(0, T]} = \hat{x}$. Since $\hat{x}(0) = \hat{x}(T)$, it is deduced that x is continuous. Moreover, over $[0, T]$ we have that $x = \hat{x} = \mu(\hat{x}_T)$, which is the restriction of the unique solution y of (28) satisfying $y_0 = \hat{x}_T = x_0$. We conclude that $x = y$ and, because $x_0 = x_T$, it follows that x can be extended periodically to a solution of (28), that is, $x \in S$ and $\alpha_1(x) = \hat{x}$.

Concerning K_2 , we proceed in an analogous way. Let $x \in S$, then

$$K_2(\alpha_2(x)) = (\pi \circ \mu)(x_T) = \pi(\mu(x_0)) = \pi(x|_{[0, T]}) = x_T = x_0 = \alpha_2(x).$$

Conversely, let $y \in \text{fp}(K_2)$, that is $y = (\pi \circ \mu)(y)$. Let \tilde{y} be the unique solution of (28) such that $\tilde{y}_0 = y$, then $y = \pi(\mu(y)) = \tilde{y}_T$. This implies that \tilde{y} is extended periodically to an element $\tilde{y} \in S$ and $y = \alpha_2(\tilde{y})$. \square

As in section 4.1, an affinity principle is obtained between K_1 and K_2 . Note that K_2 corresponds again to the Poincaré operator P although, in contrast with the non-delayed case, now P is defined over the infinite-dimensional space $C[-\tau, 0]$.

Corollary 4.10.1.

$$\deg_{LS}(I - K_1, U_1, 0) = \deg_{LS}(I - P, U_2, 0)$$

Next, let us analyze the operator $K : C[0, T] \rightarrow C[0, T]$ given by

$$K(x)(t) = x(T) + \int_0^t f(u, x(u), x(r(u)))du,$$

$$\text{where } r(u) =: \begin{cases} u - \tau + T & \text{if } 0 \leq u < \tau \\ u - \tau & \text{if } \tau \leq u \leq T. \end{cases}$$

Notice that, despite the fact that r is not continuous, the operator K is well defined and compact.

Lemma 4.11. *K is a fixed point operator for \mathcal{B}_τ*

Proof. Let $x \in S$, then $x'(t) = f(t, x(t), x(t - \tau))$ for all $t \in \mathbb{R}$, that is

$$x(t) = x(0) + \int_0^t f(u, x(u), x(u - \tau))du$$

and, because $x \in C_T$, it follows that $x|_{[0, T]} = \alpha_1(x)$ is a fixed point of K .

Conversely, if $\hat{x} \in \text{fp}(K)$, then clearly $\hat{x}(T) = \hat{x}(0)$. Moreover, if

$$x(t) := \mathbb{I}_{[-\tau, 0]}(t)\hat{x}_T(t) + \mathbb{I}_{[0, T]}(t)\hat{x}(t),$$

then $x_0 = \hat{x}_T = x_T$ and $x(r(u)) = x(u - \tau)$ for $u \in [0, T]$. Because $\hat{x} = K(\hat{x})$, for $t \in [0, T]$ it follows that

$$x(t) = \hat{x}(t) = \hat{x}(T) + \int_0^t f(u, \hat{x}(u), \hat{x}(r(u)))du = x(T) + \int_0^t f(u, x(u), x(u - \tau))du.$$

Thus, x can be extended to a T -periodic solution of (28) and hence $\hat{x} \in \alpha_1(S)$. \square

Theorem 4.12.

$$\deg_{LS}(I - K, U_1, 0) = \deg_{LS}(I - K_2, U_2, 0)$$

Proof. From the previous corollary, it suffices to verify that $\deg_{LS}(I - K, U_1, 0) = \deg_{LS}(I - K_1, U_1, 0)$. To this end, consider again the linear homotopy $H_\lambda = \lambda K + (1 - \lambda)K_1$ with $\lambda \in [0, 1]$. Suppose that $\hat{x} \in \overline{U}_1$ is a fixed point of H_λ , then

$$\hat{x}(t) = \lambda \left(\hat{x}(T) + \int_0^t f(u, \hat{x}(u), \hat{x}(r(u))) du \right) + (1 - \lambda)(\mu \circ \pi)(\hat{x})(t).$$

Evaluation at $t = 0$ yields $\hat{x}(0) = \hat{x}(T)$. As in the previous theorem, let $x := \hat{x}_T \mathbb{I}_{[-\tau, 0)} + \hat{x} \mathbb{I}_{[0, T]} \in C[-\tau, T]$, then

$$x(t) = \lambda \left(x(T) + \int_0^t f(u, x(u), x(u - \tau)) du \right) + (1 - \lambda)\tilde{\mu}(x_T)(t) \quad (30)$$

for all $t \in [0, T]$ and $x_0 = x_T$. Now consider $y = \tilde{\mu}(x_T)$, then

$$y'(t) = f(t, y(t), y(t - \tau)), \quad \forall t \in [0, T] \quad (31)$$

$$y_0 = x_0 \quad (32)$$

Moreover, from (30), we obtain

$$x'(t) = \lambda f(t, x(t), x(t - \tau)) + (1 - \lambda)f(t, y(t), y(t - \tau)). \quad (33)$$

Because y is a solution of the latter equation and $y_0 = x_0$, it follows from the uniqueness that $y = x$.

Remark 4.13. As before, using Gronwall's lemma it follows, more generally, that condition (J) is fulfilled.

As a conclusion, we deduce that x , extended periodically to \mathbb{R} , belongs to S and $\alpha_1(x) = \hat{x}$. Moreover, since $\alpha_1(S) \cap \partial(U_1) = \emptyset$ we conclude that $\hat{x} \in U_1$; thus, H_λ has no fixed points on ∂U_1 and the result follows. \square

4.4.1 Other fixed point operators in $C[0, T]$ and C_T

For notational convenience, let us define

$$Vx(t) = \int_0^t x(s) ds$$

$$N_r(x)(t) := f(t, x(t), x(r(t)))$$

with r as before and observe that, although $N_r(x)$ is not a continuous function, the operator $K_6 : C[0, T] \rightarrow C[0, T]$ given by

$$K_6(x)(t) := \bar{x} + T\overline{N_r(x)} + V(N_r(x))(t) - \overline{V(N_r(x))}$$

is well defined and compact. In the same way, we may define the operators $K_7 : C[0, T] \rightarrow C[0, T]$ and $K_8 : C_T \rightarrow C_T$ given by

$$K_7(x)(t) := \bar{x} + T\overline{N_r(x)} + V(N_r(x) - \overline{N_r(x)})(t) - \overline{V(N_r(x) - \overline{N_r(x)})}$$

$$K_8(x)(t) := \bar{x} + T\overline{N(x)} + V(N(x) - \overline{N(x)})(t) - \overline{V(N(x) - \overline{N(x)})}.$$

The operators are analogous, respectively, to K_4 , K_3 and K_5 in section 4.1. The proof of the following proposition is left for the reader.

Proposition 4.14. *K_6, K_7 and K_8 are fixed point operators and*

$$\deg_{LS}(I - K_6, U, 0) = \deg_{LS}(I - K_7, U, 0) = \deg_{LS}(I - K_8, \tilde{U}, 0)$$

provided that the open bounded sets $U \subset C[0, T]$ and $\tilde{U} \subset C_T$ have common core.

5 Final comments and open questions

1. In section 4.1, an alternative proof of the case $\eta < 0$ in Proposition 4.5 may be performed as follows. In the first place, observe that an analogue of Corollary 4.3.1 is obtained for \hat{K}_3 if we take, in the original diagram,

$$\hat{\pi}_{\ker(L)}(x) = x(T), \quad \hat{\delta}(x) = x(0) - x(T),$$

$$\hat{V}(x)(t) = - \int_t^T x(s)ds, \quad \hat{\mu}(x_T)(t) = \hat{\Phi}(t, x_T),$$

where $\hat{\Phi}(t, A)$ is defined as the unique solution of the equation such that $\hat{\Phi}(T, A) = A$. Then $\hat{\pi}(x) = x(0)$ and

$$\hat{K}(x)(t) = x(0) - \int_t^T f(s, x(s))ds,$$

while

$$\hat{K}_1(x)(t) = \hat{\Phi}(t, x(0)).$$

Thus, the abstract result implies that, if the common core assumption is satisfied, then the degree of $I - \hat{K}_1$ coincides with the degree of $I - \hat{K}_2$, where

$$\hat{K}_2(A) = (\hat{\mu} \circ \hat{\pi}) = \hat{\Phi}(0, A).$$

In other words, \hat{K}_2 can be identified with the inverse \hat{P} of the Poincaré map; thus, the degree computation follows from the next lemma:

Lemma 5.1. *Let $\Omega \subset \mathbb{R}^n$ be open and bounded such that P has no fixed points on $\partial\Omega$, then*

$$\deg_B(I - \hat{P}, P(\Omega), 0) = (-1)^n \deg_B(I - P, \Omega, 0).$$

Proof. It suffices to observe that P and \hat{P} have the same fixed points and, due to the properties of the Brouwer degree, we may assume that 0 is a regular value of $I - P$. Because the degree over small balls coincides with the degree of its linearization, we only need to prove that

$$\deg_B(I - P_L, B_r(0), 0) = (-1)^n \deg_B(I - \hat{P}_L, P_L(B_r(0)), 0),$$

where P_L is the Poincaré map associated to a linear system $x'(t) = B(t)x(t)$, where B is a T -periodic matrix such that P_L has no nontrivial fixed points. Let $M(t)$ be the fundamental matrix of the system $x' = Bx$ such that $M(0) = I$, then $P_L(x_0) = M(T)x_0$, so the degree of $I - P_L$ is simply computed by

$$\deg_B(I - P_L, B_r(0), 0) = \operatorname{sgn}(\det(I - M(T))).$$

On the other hand,

$$\begin{aligned} \deg_B(I - \hat{P}_L, P_L(B_r(0)), 0) &= \operatorname{sgn}(\det(I - M(T)^{-1})) \\ &= (-1)^n \operatorname{sgn} \det(M(T)^{-1}) \operatorname{sgn}(\det(I - M(T))). \end{aligned}$$

Thus, the result follows from the fact that $M(t)$ is invertible for all t and, by continuity,

$$\operatorname{sgn}(\det(M(T))) = \operatorname{sgn}(\det(M(0))) = 1.$$

□

Also, it is worth noticing that, in the context of the coincidence degree theory, it is well known that if the assumption

$$x' \neq \lambda N_T(x), \quad x \in \partial \tilde{U}, \lambda \in (0, 1] \quad (34)$$

holds, then

$$\deg_{LS}(I - K_5, \tilde{U}, 0) = \deg_B(\phi, \tilde{U} \cap \mathbb{R}^n, 0)$$

and

$$\deg_{LS}(I - \hat{K}_5, \tilde{U}, 0) = \deg_B(-\phi, \tilde{U} \cap \mathbb{R}^n, 0)$$

where $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is defined by

$$\phi(x) := -TN_T(x).$$

In other words, when (34) is satisfied, taking $U, \tilde{U} \subset C_T$ as in the previous proposition it is seen in a direct way that

$$\deg_{LS}(I - K_5, \tilde{U}, 0) = (-1)^n \deg_{LS}(I - \hat{K}_5, U, 0).$$

2. Concerning the Dirichlet problem in section 4.2, several comments can be made. To begin, observe that the explicit form of the operator K_2 is

$$K_2(ta + b) = (\pi \circ \mu)(ta + b) = t[x(1) + x'(0)] + 2x(0) = t[x(1) + a] + 2b,$$

where $x = \mu(ta + b) = \Phi^1(\cdot, b, a)$. In particular, if $K_2(ta + b) = ta + b$, then $b = 0$: in other words, the fixed points of K_2 belong to the subspace

$W = \{ta : a \in \mathbb{R}^n\} \subset \ker(L)$. Notice, furthermore, that $K_2(W) \subset W$. This yields a direct computation of degree as follows. According to the standard properties of the Brouwer degree, may assume that 0 is a regular value of $I - K_2$, then

$$\deg_B(I - K_2, U_2, 0) = \sum_{ta \in S_2} \operatorname{sgn}(\det(I - DK_2(ta)))$$

where $S_2 := \{ta \in U_2 : K_2(ta) = ta\}$. A simple computation shows that $DK_2(ta)$ can be identified with the $\mathbb{R}^{2n \times 2n}$ matrix given by

$$\begin{pmatrix} D_a \Phi^1(1, 0, a) + I_n & D_b \Phi^1(1, 0, a) \\ 0 & 2I_n \end{pmatrix}$$

where $I_n \in \mathbb{R}^{n \times n}$ denotes the identity matrix. This implies that

$$\operatorname{sgn}(\det(I - DK_2(ta))) = \operatorname{sgn}(\det(D_a \Phi^1(1, 0, a))).$$

Moreover, taking into account the isomorphism $\varphi|_W : ta \mapsto a$ it is clear that the zeros of \mathcal{S} in $\varphi(\bar{U}_2)$ coincide with the elements of $\varphi(S_2)$. Because $D\mathcal{S}(a) = D_a \Phi^1(1, 0, a)$, we conclude that

$$\deg(\mathcal{S}, \varphi(U_2), 0) = \sum_{a \in \varphi(S_2)} \operatorname{sgn}(\det(D_a \Phi^1(1, 0, a)))$$

and so the result is deduced.

3. Still regarding section 4.2, we may define a variety of fixed point operators as follows. Firstly, inspired by Proposition 2.2, we may define an operator \tilde{K} as a conjugate of the function g defined in the proof of Proposition 4.8. With this in mind, let us define $\tilde{i} : W \rightarrow C^1[0, 1]$ given by $\tilde{i}(ta) := \mu(ta) = \Phi^1(\cdot, 0, a)$ and $\tilde{\pi} : C^1[0, 1] \rightarrow W$ given by $\tilde{\pi}(x) := tx'(0)$. Then $\tilde{\pi} \circ \tilde{i} = \operatorname{id}_W$ and, according to the definition $g(ta) := t(a - \mathcal{S}(a))$, we may compute $\tilde{K} = \tilde{i} \circ g \circ \tilde{\pi}$ to obtain:

$$\tilde{K}(x) = \Phi^1(\cdot, 0, x'(0) - \mathcal{S}(x'(0))).$$

It is interesting to observe that \tilde{K} and $K := \pi + V \circ N$ satisfy the affinity principle, although they are not linearly homotopic. The same happens with the operator $K_1 = \mu \circ \pi$, which reads

$$K_1(x) = \mu[t(x(1) + x'(0)) + 2x(0)].$$

Indeed, from the equality $x = \lambda K_1(x) + (1 - \lambda)\tilde{K}(x)$ one obtains $x(0) = 2\lambda x(0)$, but the conclusion $x(0) = 0$ does not necessarily hold when $\lambda = \frac{1}{2}$.

More generally, a family of fixed point operators related with all the preceding ones can be obtained in the following way. Consider the mapping $\delta_\theta := \theta \circ \delta$, where $\theta : \ker(L) \rightarrow \ker(L)$ is an arbitrary orientation-preserving isomorphism. It is well known that θ and the identity belong to the same connected component of the linear group $GL(\ker(L))$ and, thus, we may define $\pi_\lambda := \pi_{\ker(L)} + h(\lambda) \circ \delta$, where $h : [0, 1] \rightarrow GL(\ker(L))$ is continuous with $h(0) = I$ and $h(1) = \theta$. Observe that π_λ may not be necessarily

onto but, however, the corresponding K_λ is a fixed point operator: if $x = K_\lambda(x)$, then $L(x) = N(x)$; moreover, applying $\pi_{\ker(L)}(x)$ it follows that $h(\lambda)(\delta(x)) = 0$ and, consequently, $\delta(x) = 0$. If one takes, instead, an element $\theta \in GL(\ker(L))$ that does not preserve orientation, then it belongs to the same component of $\hat{\delta}(x) = -tx(1) + x(0)$ and it follows that $\deg(I - K_2(\theta), U_2, 0) = (-1)^n \deg(\mathcal{S}, \varphi(U_2), 0)$.

It is clear that one may also replace the projector $\pi_{\ker(L)}$ by another one, although the application of the conjugacy lemma relies on the existence of μ . On the other hand, a family of fixed point operators of different nature is given by $K^\eta : x \mapsto y$, where y is the unique solution of the problem

$$y''(t) + \eta y(t) = f(t, x(t)) + \eta x(t), \quad y(0) = y(1) = 0$$

and η is not an eigenvalue of $-L$ for the Dirichlet condition. Here, the degree of $I - K^\eta$ will depend on the position of η with respect to the spectrum of $-L$; this study shall be the subject of a forthcoming paper.

4. Analogously, an operator K^η was also defined in section 4.3, assuming that $-\eta$ does not belong to the spectrum of $-\Delta$ under the nonlocal conditions (26). As mentioned, in this case the degree of $I - K^\eta$ depends only on the sign of η : more precisely, following the ideas of section 4.1 it can be proven that the degree of $I - K^\eta$ coincides with the degree of $I - K_5$, multiplied by $\text{sgn}(\eta)^n$. For $\eta > 0$, we already observed that K^η is homotopic to K_5 ; when $\eta < 0$, an intuitive explanation reads as follows. For simplicity, assume that $d = 1$ (ODE case), which corresponds, when $\Omega = (0, 2\pi)$, to the second order 2π -periodic problem. With this in mind, consider the linear autonomous problem

$$u''(t) = Au(t)$$

with $A \in \mathbb{R}^{n \times n}$ and suppose there are no nontrivial T -periodic solutions. If one defines as before the mapping $u \mapsto v$ by solving the equation

$$v'' + \eta v = Au + \eta u,$$

then $(I - K^\eta)(u) = u - v$ is computed in terms of the Fourier expansion of u given by

$$u = a_0 + \sum_{n \in \mathbb{N}} [\cos(nt)a_n + \sin(nt)b_n]$$

as

$$(I - K^\eta)(u) = \left(I - \frac{A}{\eta}\right) a_0 + \sum_{n \in \mathbb{N}} \left(\frac{\eta I - A}{\eta - n^2}\right) [\cos(nt)a_n + \sin(nt)b_n].$$

This implies that the subspaces $S_n := \text{span}\{\cos(nt), \sin(nt)\}$ are invariant and, furthermore, the associated matrix of $(I - K^\eta)|_{S_n}$ is

$$M_n = \begin{pmatrix} \frac{\eta I - A}{\eta - n^2} & 0 \\ 0 & \frac{\eta I - A}{\eta - n^2} \end{pmatrix}$$

which satisfies $\text{sgn}(\det(M_n)) > 0$. In other words, the degree computation depends only on the sign of $\det\left(I - \frac{A}{\eta}\right)$.

Acknowledgement

The authors would like to thank Prof. Rafael Ortega for the reading of a first version of the manuscript and his fruitful comments. This work was partially supported by projects UBACyT 20020190100039BA and PIP 11220130100006CO CONICET.

References

- [1] N. Bobylev and M. A. Krasnoselskii, *Principles of affinity in nonlinear problems*. Journal of Mathematical Sciences 83 No. 4 (1997), 485–521.
- [2] R. Gaines and J. Mawhin, *Coincidence degree and nonlinear differential equations*. Springer, 1977.
- [3] M. A. Krasnoselskii, *The operator of translation along the trajectories of differential equations*, American Mathematical Society, Providence RI, 1968.
- [4] M. A. Krasnoselskii and P. P. Zabreiko, *Geometrical methods of nonlinear analysis*. Springer-Verlag, Berlin, 1984.
- [5] J. Mawhin, *Topological degree methods in nonlinear boundary value problems*. CMBS Conf. Math. No. 40, Amer. Math. Soc., Providence, 1979.