

Structure Calculations Without Effective Interactions

J. A. Secrest¹, J. M. Conroy² and H. G. Miller²

¹*Department of Physics and Astronomy, Georgia Southern University,
Armstrong Campus, Savannah, Georgia, USA and*

²*Department of Physics, State University of New
York at Fredonia, Fredonia, New York, USA*

(Dated: November 12, 2021)

Abstract

Good approximate eigenstates of a Hamiltonian operator which possesses a point as well as a continuous spectrum have been obtained using the Lanczos algorithm. Iterating with the bare Hamiltonian operator yields spurious solutions which can easily be identified. The rms radius of the ground state eigenvector, for example, is calculated using the bare operator.

The Hamiltonian operators in nuclear and atomic physics are usually only bounded from below and possess a point as well as continuous spectrum. Unlike in atomic physics where in a properly chosen basis, the Hamiltonian matrix is diagonally dominant this is definitely not the case in nuclear physics. The generally accepted procedure for determining the lower-lying eigenstates in this case is to first map the Hamiltonian operator onto a finite dimensional subspace space spanned by a number of \mathcal{L}^2 basis states, *e.g.* to construct an effective Hamiltonian operator[1–4] which must then be diagonalized. This procedure is fraught with difficulties as initially the effective Hamiltonian operator is non-linear, non-hermetian and upon diagonalization guarantees at best only good approximate eigenvalues of the lowest lying eigenstates. The eigenstates of the effective Hamiltonian are the projection of the exact eigenstates onto the subspace. Hence to calculate any other observable the corresponding operator must also be mapped on the same subspace.

Consider an operator, \hat{H} , which possesses a continuum (or continua) as well as a point spectrum. The subspace spanned by its bound state eigenfunctions, \mathcal{H}_B , is by itself certainly not complete. As the composition of this space is generally not known beforehand, a set of basis states which is complete and spans a space, \mathcal{F} , is chosen to construct a matrix representation of the operator, \hat{H} , to be diagonalized. Mathematically this corresponds to projecting the operator \hat{H} onto the space \mathcal{F} . Clearly the eigenpairs obtained from diagonalizing the projected operator, \hat{H}_P , need not all be the same as those of the operator \hat{H} . However, because the set of basis states is complete, any state contained in \mathcal{H}_B can be expanded in terms of this set of basis states. Hence \mathcal{H}_B may also be regarded as a subspace of \mathcal{F} and the complete diagonalization of \hat{H}_P will yield not only the exact eigenstates of \hat{H} but additional spurious eigensolutions. Note these spurious eigenfunctions are eigenfunctions of \hat{H}_P but not of \hat{H} . Furthermore in this case the Rayleigh-Ritz bounds discussed in the paper by Krauthauser and Hill[5] apply now to the eigenstates of \hat{H}_P .

It is interesting to note that the same problem occurs in the Lanczos algorithm[6, 7] when it is applied to operators which possess a bound state spectrum as well as a continuum[8]. This is not surprising as the Lanczos algorithm can also be considered as an application of the Rayleigh-Ritz method[9]. In this case an orthonormalized set of Krylov basis vectors is used to construct iteratively a matrix representation of the operator which is then diagonalized. Again spurious states can occur for precisely the same reasons given above. In this case we have proposed identifying the exact bound states in the following manner[8]. After each

iteration, for each of the converging eigenpairs $(e_{l\lambda}, |e_{l\lambda}\rangle)$, $\Delta_{l\lambda} = |e_{l\lambda}^2 - \langle e_{l\lambda} | \hat{H}^2 | e_{l\lambda} \rangle|$ (where l is the iteration number) is calculated and a determination is made as to whether Δ is converging toward zero or not. For the exact bound states of \hat{H} , Δ must be identically zero while the other eigenstates of the projected operator should converge to some non-zero positive value. This method has been successfully implemented to identify spurious states in non-relativistic[8] as well as relativistic[10] eigenvalue problems. A similar procedure can be implemented in any Rayleigh-Ritz application. One simply must check to see whether the eigensolutions from the diagonalization of \hat{H}_P are also eigensolutions of \hat{H}^2 .

To illustrate this a simple example is presented. Consider the following Hamiltonian operator

$$\hat{H} = \hat{H}_0 + \hat{V} \quad (1)$$

where H_0 is the harmonic oscillator Hamiltonian with $\hbar\omega = 1$ and

$$V = V_0 \exp -ar^2 \quad (2)$$

with $V_0 = -11$ and $a = 1$. Rather than simply attempt to obtain a coordinate state representation of the eigenstate[8], approximate eigenstates have been back expanded in a basis as is the case in more realistic many body calculations. For a single particle the Hamiltonian matrix in an s wave oscillator basis

$$\phi(x) = \frac{1}{\sqrt{2^v \sqrt{\pi} v!}} H_v(x) e^{-x^2/2} \quad (3)$$

where $H_v(x)$ are the Hermite polynomials. an n -dimensional representation of \hat{H} can easily be constructed.

Choosing 30 basis states the Lanczos algorithm is used to iteratively obtain approximate eigenstates of \hat{H} . Approximate energy eigenvalues of the ground and excited states were determined (see Fig 1). After each iteration Δ^2 (see Fig. 2) is calculated. It is clear that the ground state and excited state converge to relatively small values while the unphysical spurious state converges to a relatively large non-zero value. After 11 iterations the approximate values of energy eigenvalues have converged to $e_{0,8} = 12.397$ and $e_{2,11} = -3.579$ with a convergence factor of $\leq 10^{-3}$.

Having ascertained the ground state eigenfunction the rms radius can easily be calculated (see Fig3). Note that the convergence is much slower than that of the eigen-




FIG. 1: The iterative convergence behavior of the energy ground, excited, and spurious state eigenvalues obtained from the Lanczos algorithm.




FIG. 2: The iterative convergence behavior of Δ^2 of the ground states, excited, and spurious state. Note that the ground and excited state converge to a relatively small value while the spurious state converges to a relatively large non-zero value.

values. This is undoubtedly due to the fact that the convergence rate of the approximate ground state eigenfunction is considerably slower than that of the approximate ground state eigenvalue. After 30 iterations the RMS radius is 0.322.

For a bare operator which contains a bound as well as a continuum spectrum the Lanczos algorithm has been used to obtain its approximate eigenstates. Unphysical spurious eigenstates which may appear can easily be identified. Since the approximate eigenfunctions are




FIG. 3: The convergence behavior of the RMS radius

obtained the expectation value of other observables can easily be calculated without recourse to the use of effective operators.



- [1] C. Bloch and J. Horowitz, Nucl. Phys. **8**, 91 (1958).
- [2] T. H. Shucan and H. Weidenmueller, Ann. Physics **76**, 483 (1973).
- [3] P. J. Ellis and E. Osnes, Rev. Mod. Phys. **49**, 777 (1977).
- [4] L. Corraggio, A. Covello, A. Gargano, N. Itaco, and T. Kuo, Progress in Particle and Nuclear Physics **62**, 135 (2009).
- [5] C. Krauthauser and R. Hill, Can. J. Phys. **80**, 181 (2002).
- [6] C. Lanczos, J. Res. Nat. Bur. Stand. **45**, 255 (1950).
- [7] W. R. Davis, ed., *Cornelius Lanczos Collected Published Papers with Commentaries*, vol. V (North Carolina State University, Raleigh, NC, 1998).
- [8] R. C. Andrew and H. G. Miller, Phys. Lett. A **318**, 487 (2003).
- [9] B. N. Parlett, *The Symmetric Eigenvalue Problem* (Prentice Hall, Englewood Cliffs, N. J., 1980).
- [10] R. C. Andrew, H. G. Miller, and G. D. Yen, Chem. Phys. Lett. **475**, 298 (2009).