

# STOCHASTIC NAVIER-STOKES EQUATIONS FOR TURBULENT FLOWS IN CRITICAL SPACES

ANTONIO AGRESTI AND MARK VERAAR

ABSTRACT. In this paper we study the stochastic Navier-Stokes equations on the  $d$ -dimensional torus with gradient noise, which arises in the study of turbulent flows. Under very weak smoothness assumptions on the data we prove local well-posedness in the critical case  $\mathbb{B}_{q,p}^{d/q-1}$  for  $q \in [2, 2d)$  and  $p$  large enough. Moreover, we obtain new regularization results for solutions, and new blow-up criteria which can be seen as a stochastic version of the Serrin blow-up criteria. The latter is used to prove global well-posedness with high probability for small initial data in critical spaces in any dimensions  $d \geq 2$ . Moreover, for  $d = 2$  we obtain new global well-posedness results and regularization phenomena, which unify and extend several earlier results.

## CONTENTS

1. Introduction	2
1.1. Scaling and criticality	3
1.2. Overview of the main results	4
1.3. Notation	6
2. Statement of the main results	7
2.1. The Helmholtz projection and function spaces	7
2.2. Main assumptions and definitions	8
2.3. Local existence and regularization	10
2.4. Stochastic Serrin's blow-up criteria	12
2.5. Global existence results	13
3. Maximal $L^p$ -regularity for the turbulent Stokes couple	14
3.1. Assumptions and main results	15
3.2. Proof of Theorem 3.2	17
3.3. Proof of Lemma 3.6	19
4. Proofs of the main results	24
4.1. Local well-posedness and the proof of Theorem 2.4	24
4.2. Higher order regularity and the proof of Theorem 2.7	29
4.3. Blow-up criteria and Theorems 2.9	32
4.4. Global well-posedness, and Theorems 2.11 and 2.12	33

*Date:* December 30, 2021.

2010 *Mathematics Subject Classification.* Primary: 35Q30, Secondary: 60H15, 35B65, 76M35.

*Key words and phrases.* Stochastic Navier-Stokes equations, Kraichnan's turbulence, turbulent flows, critical spaces, Serrin blow-up criteria, stochastic maximal regularity, stochastic evolution equations.

The first author has been partially supported by the Nachwuchsring – Network for the promotion of young scientists – at TU Kaiserslautern. The second author is supported by the VIDI subsidy 639.032.427 of the Netherlands Organisation for Scientific Research (NWO).

Appendix A. Global well-posedness for $d = 2$ with $L^\infty$ -noise	37
A.1. Assumptions and main result	37
A.2. Maximal $L^2$ -regularity for the turbulent Stokes system	39
A.3. Proof of Theorem A.3	41
References	44

## 1. INTRODUCTION

In this paper we provide a comprehensive treatment of (analytically and probabilistically) *strong* solutions to stochastic Navier-Stokes equations for turbulent incompressible flows on the  $d$ -dimensional torus  $\mathbb{T}^d$ :

$$(1.1) \quad \begin{cases} du = [\nu \Delta u - (u \cdot \nabla)u - \nabla p] dt + \sum_{n \geq 1} [(b_n \cdot \nabla)u - \nabla \tilde{p}_n + g_n(u)] dw_t^n, \\ \operatorname{div} u = 0, \\ u(\cdot, 0) = u_0. \end{cases}$$

Here  $u := (u^k)_{k=1}^d : [0, \infty) \times \Omega \times \mathbb{T}^d \rightarrow \mathbb{R}^d$  denotes the unknown velocity field,  $p, \tilde{p}_n : [0, \infty) \times \Omega \times \mathbb{T}^d \rightarrow \mathbb{R}$  the unknown pressures,  $(w_t^n : t \geq 0)_{n \geq 1}$  a given sequence of independent standard Brownian motions and

$$(b_n \cdot \nabla)u := \left( \sum_{j=1}^d b_n^j \partial_j u^k \right)_{k=1}^d, \quad (u \cdot \nabla)u := \left( \sum_{j=1}^d u_j \partial_j u^k \right)_{k=1}^d.$$

Actually, we will consider (1.1) in a slightly more general form (see (2.1) below). For instance we replace  $\nu$  by variable viscosity in divergence form, and we allow an additional deterministic forcing term.

The relation between Navier-Stokes equations and hydrodynamic turbulence is one of the most challenging problems in fluid mechanics. It is believed that the onset of the turbulence is related to the randomness of background movement. The mathematical study of stochastic perturbations of Navier-Stokes equations began with the work of Bensoussan and Temam in [BT73] and it was later extended in several directions, see e.g. [BP00, BCF91, BCF92, Cho78, CP01, FG95, IF79, WS00, BF17] and the references therein. In the mathematical literature, usually, the random perturbation of the Navier-Stokes equations is postulated and does not have a clear physical motivation. In [MR01, MR04], Mikulevicius and Rozovskii found a new approach to derive the stochastic Navier-Stokes equations. Indeed, there it is assumed that the dynamics of the fluid particles is given by the *stochastic flow*

$$(1.2) \quad \begin{cases} d\mathcal{X}(t, x) = u(t, \mathcal{X}(t, x))dt + b(t, \mathcal{X}(t, x)) \circ dW, \\ \mathcal{X}(0, x) = x, \quad x \in \mathbb{T}^d, \end{cases}$$

where  $W$  and  $\circ$  denote white-noise and Stratonovich integration, respectively. Using Newton's second law the stochastic Euler and Navier-Stokes equations for  $u$  are derived. Roughly speaking, (1.2) says that the velocity field splits into a sum of slow oscillating part  $u dt$  and a fast oscillating part  $b \circ dW$ . As noticed in [MR04], such splitting can be traced back to the work of Reynolds in 1880. In more recent works, closely related models were proposed by Kraichnan [Kra68, Kra94] in the

study of turbulence transportation of passive scalars and then developed further by other authors, see e.g. [GK96, GV00]. The corresponding theory of turbulence transportation is called Kraichnan's theory, and in this theory, one typically has  $b = ((b_n^j)_{j=1}^d)_{n \geq 1}$  and for all  $n \geq 1$ ,

$$(1.3) \quad (b_n^j)_{j=1}^d \in H^{\frac{d}{2} + \eta, 2}(\mathbb{T}^d; \mathbb{R}^d) \cap \{\operatorname{div} \xi = 0 \text{ in } \mathcal{D}'\} \hookrightarrow C^\eta(\mathbb{T}^d; \mathbb{R}^d),$$

where  $\eta > 0$  is small and  $H$  denotes Bessel-potential spaces, cf. [MR05, Section 1]. We refer to [BF20, Fla08, MR01] for additional motivations and to [MR04] for a derivation of (1.1) from (1.2) via Newton's law. With this application in mind, we focus on the case where  $(b_n^j)_{n \geq 1}$  belongs to a certain Sobolev space that embeds into a Hölder space  $C^\eta(\mathbb{T}^d; \mathbb{R}^d)$  where  $\eta > 0$  is small. In light of this, the vorticity formulation of (1.1) (i.e. for the unknown  $\operatorname{curl} u$ ) is in general not available. The reader interested in the Navier-Stokes equations in vorticity form may consult [FL21, Fla15, Fla11, HLN21] and the references therein.

The problem (1.1) will be considered in the Itô formulation. In the literature the Stratonovich formulation of (1.1) is also used frequently, i.e. in the former case one replaces  $\sum_{n \geq 1} (b_n \cdot \nabla) u \, dw_t^n$  by  $\sum_{n \geq 1} (b_n \cdot \nabla) u \circ dw_t^n$  and similar for the  $g$ -term. Recall that, at least formally,

$$(1.4) \quad \sum_{n \geq 1} (b_n \cdot \nabla) u \circ dw_t^n = \operatorname{div}(a_b \cdot \nabla u) \, dt + \sum_{n \geq 1} (b_n \cdot \nabla) u \, dw_t^n,$$

where  $a_b := (\frac{1}{2} \sum_{n \geq 1} b_n^j b_n^i)_{i,j=1}^d$  provided  $b_n$  is  $(t, \omega)$ -independent and  $\operatorname{div} b_n = 0$ , see e.g. the computations at the beginning of Section 3.1 in [FGL21]. The additional term  $\operatorname{div}(a_b \cdot \nabla u) \, dt$  is referred as the Itô correction. By (1.3), the divergence free condition on  $b_n$  is physically justified. Since in (2.1) we will also consider variable viscosity in divergence form, our results also cover the case of noise in Stratonovich form provided one takes into account the relative Itô's correction.

It is not possible to discuss all relevant papers on stochastic Navier-Stokes equations here, and we mostly restrict ourselves to the ones where gradient noise is included. There are several important papers concerned with martingale solutions to (1.1), see e.g. [MR99, MR05, BM13, FG95] and the references therein. In these papers global existence is the main point, and uniqueness is open unless  $d = 2$ . See also [HLN19] for a rough path theory approach.

**1.1. Scaling and criticality.** Before we discuss some of our main results in more detail, we will discuss scaling and criticality in the case (1.1) would be considered on  $\mathbb{R}^d$  instead of  $\mathbb{T}^d$  as the scaling is easier in the  $\mathbb{R}^d$ -case. This will also motivate the function spaces and the growth condition on  $g$  used in the paper. In the deterministic setting (e.g.  $b \equiv 0$  and  $g \equiv 0$ ), it is well-known that (local smooth) solutions to (1.1) are (roughly) invariant under the map  $u \mapsto u_\lambda$  where  $\lambda > 0$  and

$$(1.5) \quad u_\lambda(t, x) = \lambda^{1/2} u(\lambda t, \lambda^{1/2} x), \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d.$$

In the PDE literature (see e.g. [Can04, LR16, PW18, Tri13]), Banach spaces of functions (locally) invariant under the induced map  $u_0 \mapsto \lambda^{1/2} u_0(\lambda^{1/2} \cdot)$  are called *critical* for (1.1). Examples of such spaces are the Besov spaces  $B_{q,p}^{d/q-1}(\mathbb{R}^d; \mathbb{R}^d)$  and the Lebesgue space  $L^d(\mathbb{R}^d; \mathbb{R}^d)$ . Indeed, the corresponding homogeneous spaces satisfy

$$\|u_{0,\lambda}\|_{\dot{B}_{q,p}^{d/q-1}(\mathbb{R}^d; \mathbb{R}^d)} \widetilde{\sim} \|u_0\|_{\dot{B}_{q,p}^{d/q-1}(\mathbb{R}^d; \mathbb{R}^d)}, \quad \|u_{0,\lambda}\|_{L^d(\mathbb{R}^d; \mathbb{R}^d)} \widetilde{\sim} \|u_0\|_{L^d(\mathbb{R}^d; \mathbb{R}^d)},$$

where the implicit constants do not depend on  $\lambda > 0$ .

In the stochastic case a similar behavior appears. More precisely, one can check that if  $u$  is a (local smooth) solution to (1.1) on  $\mathbb{R}^d$ , then  $u_\lambda$  is a (local smooth) solution to (1.1) on  $\mathbb{R}^d$ , where  $(w^n)_{n \geq 1}$  is replaced by the *scaled* noise  $(\beta_{t,\lambda}^n)_{n \geq 1}$  where  $\beta_{t,\lambda}^n := \lambda^{-1/2} w_{\lambda t}^n$  for  $t \geq 0$  and  $n \geq 1$ . Moreover, the scaling (1.5) also suggests that the nonlinearity  $g_n(u)$  has to grow quadratically in  $u$  in order to keep the same scaling. Indeed, let  $g_n(u) = G_n(u, u)$  where  $G_n : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  is bilinear. In such a case, roughly speaking one has, for all  $\lambda > 0$ ,  $x \in \mathbb{R}^d$  and  $n \geq 1$ ,

$$\begin{aligned} \int_0^{t/\lambda} g_n(u_\lambda(s, x)) d\beta_{s,\lambda}^n &= \int_0^{t/\lambda} \lambda G_n(u_\lambda(\lambda s, \lambda^{1/2} x), u_\lambda(\lambda s, \lambda^{1/2} x)) d\beta_{s,\lambda}^n \\ &= \lambda^{1/2} \int_0^t G_n(u_\lambda(s, \lambda^{1/2} x), u_\lambda(s, \lambda^{1/2} x)) dw_s^n, \end{aligned}$$

which agrees with the scaling of the deterministic nonlinearities

$$\int_0^{t/\lambda} (u_\lambda(s, x) \cdot \nabla) u_\lambda(s, x) ds = \lambda^{1/2} \int_0^t (u(s, \lambda^{1/2} x) \cdot \nabla) u(s, \lambda^{1/2} x) ds.$$

A similar scaling argument also holds for the remaining deterministic and stochastic integrals. In particular, the  $b$ -term has the correct scaling under the map (1.5).

The above discussion shows that the stochastic perturbations of gradient type and/or of quadratic type in  $u$  respect the scaling of the deterministic Navier-Stokes equations. In the present paper we will consider both situations. Furthermore, for our results to hold, we do not need  $g_n$  to be bilinear. A quadratic growth condition will be sufficient.

Finally, we mention that our methods can be also extended to the case where  $g_n$  grows more than quadratically. However, the above scaling argument fails, and the corresponding critical spaces change accordingly to the growth of  $g_n$  (see [AV21a, Subsection 5.2] for related results). Since we are not aware of physical motivations for such behaviour of  $g_n$ , we leave the details to the interested reader.

**1.2. Overview of the main results.** The aim of this paper is to provide a first systematic treatment of the stochastic Navier-Stokes equations (1.1) arising in study of turbulent flows in the *strong* setting (more precisely  $L^q(H^{s,q})$ -setting). Below we briefly list our main results.

- (a) Local well-posedness in *critical* spaces (Theorem 2.4);
- (b) Regularization with *optimal* gain in the Sobolev scale (Theorems 2.4 and 2.7);
- (c) Stochastic *Serrin's* blow-up criteria (Theorem 2.9);
- (d) Global well-posedness for small data in *critical* spaces (Theorem 2.11);
- (e) Global well-posed in case  $d = 2$  with *rough* data (Theorem 2.12);

(a): In Theorem 2.4 (see also Theorem 4.1) we show existence of *strong unique* solutions to (1.1) for initial values  $u_0$  belonging to the critical Besov space  $B_{q,p}^{d/q-1}(\mathbb{T}^d; \mathbb{R}^d)$ , where  $q$  varies in a certain range (depending on the smoothness of  $b_n$ ) and  $p$  can be chosen as large as one wants. The embedding  $L^d(\mathbb{T}^d; \mathbb{R}^d) \hookrightarrow B_{d,p}^0(\mathbb{T}^d; \mathbb{R}^d)$  for  $p \geq d$  shows that our results also hold for the critical space  $L^d(\mathbb{T}^d; \mathbb{R}^d)$ . Let us remark that in all cases, we have the restriction  $q < 2d$  and therefore we allow critical spaces with smoothness up to  $-\frac{1}{2}$ . We are not aware of the optimality of this threshold. Note that this behavior differs from the deterministic case where  $q$  can be chosen to be large. Next we compare the result of Theorem 2.4 to the existing

literature. In [MR04] the authors showed existence of strong solutions for divergence free vector field for initial values from the (non critical) space  $W^{1,q}(\mathbb{T}^d; \mathbb{R}^d)$  under a restrictive assumption on the regularity of  $b$ . In particular, the smoothness assumption there does not satisfy (1.3) if  $\eta$  is small. Existence of strong solutions to (1.1) in an  $L^p(L^q)$ -setting has been also considered in [DZ20, Wan20]. However, in the latter references, the Kraichnan term  $(b_n \cdot \nabla)u$  is not included.

(b): One question which will be answered in this paper is how the regularity of  $b_n$  affects the regularity of  $u$ . The term  $(b_n \cdot \nabla)u$  suggests that  $u$  can gain at most one order of smoothness compared to  $b_n$ . In Theorem 2.7 we prove an *optimal* gain of regularity in the Sobolev scale, i.e. for all  $\eta > 0$  and  $\xi \in (2, \infty)$  we have

$$(b_n^j)_{n \geq 1} \in H^{\eta, \xi}(\mathbb{T}^d; \ell^2) \implies u \in L_{\text{loc}}^r(0, \sigma; H^{\eta+1, \xi}(\mathbb{T}^d; \mathbb{R}^d)) \text{ a.s.}$$

where  $r \in (2, \infty)$  is arbitrary and  $\sigma$  denotes the explosion time of  $u$ . In the Hölder scale, by Theorem 2.7 and Remark 2.2, we prove a *sub-optimal* gain of regularity, i.e. for all  $\eta > 0$  and  $\varepsilon > 0$  and  $\theta \in (0, 1/2)$  we have

$$(b_n^j)_{n \geq 1} \in C^\eta(\mathbb{T}^d; \ell^2) \implies u(t) \in C_{\text{loc}}^{\theta, 1+\eta-\varepsilon}((0, \sigma) \times \mathbb{T}^d; \mathbb{R}^d) \text{ a.s.}$$

The above results are *new* even in the important case  $d = 2$  where  $\sigma = \infty$  a.s. Indeed, for  $(b_n^j)_{n \geq 1} \in C^\eta(\mathbb{T}^d; \ell^2)$  with  $\eta > 0$  small, the regularization cannot be proved with classical bootstrap methods, see the text below Theorem 2.7. However, in the case  $b_n$  is regular enough, (some) gain of regularity can be proved by standard methods (see e.g. [Mik09]) or via the previously mentioned vorticity formulation of (1.1). But for rough  $(b_n^j)_{n \geq 1}$  these approaches are not applicable.

(c): To the best of our knowledge, blow-up criteria for stochastic Navier-Stokes equations have not been studied so far in the literature. In Theorem 2.9 we provide an extension to the classical Serrin's criteria (see e.g. [LR16, Theorem 11.2]).

(d): In Theorem 2.11 we prove global existence in probability for solutions to (1.1) under smallness assumption on  $u_0$  and on the *sublinear* part of  $g_n$ . In deterministic framework this result is well-known and for  $H^{1/2}$ -data (here  $d = 3$ ) is due to Fujita and Kato in the seminal paper [FK64] and later extended to  $B_{q,p}^{d/q-1}$ -data and even to  $BMO^{-1}$ -data by Koch-Tataru [KT01]. In the stochastic framework, under smallness assumptions on the noise, some results in this direction can be found in [Kim10, KXZ21, DZ20]. Note that in [KXZ21, DZ20] the transport noise term  $(b_n \cdot \nabla)u$  is not allowed, and in [Kim10] a small gradient noise term can be used, where an additional smallness condition on the data is used, but in non critical spaces. In Theorem 2.11 we are able to give an extension of the previous results, where we also allow  $b$  to be *not* small and  $g_n$  of quadratic type. As in (a), the smallness condition on  $u_0$  is given in a  $B_{q,p}^{d/q-1}$ -norm with smoothness up to  $-\frac{1}{2}$ .

(e): In Theorem 2.12 we prove *global* existence for (1.1) for initial data which is much *rougher* than data from the classical energy space  $L^2(\mathbb{T}^2; \mathbb{R}^2)$ . More precisely, we prove that there exists  $\zeta \in (2, 4)$  (depending on the smoothness of  $b_n$ ) such that (1.1) is globally well-posed for data in  $B_{\zeta,p}^{2/\zeta-1}(\mathbb{T}^2; \mathbb{R}^2)$  where  $p$  is large. Since  $L^2(\mathbb{T}^2; \mathbb{R}^2) \hookrightarrow B_{\zeta,p}^{2/\zeta-1}(\mathbb{T}^2; \mathbb{R}^2)$  for  $p \geq 2$ , this result extends the well-known global existence result for  $L^2$ -data in two dimensions. For the reader's convenience, we also provide a self-contained proof of the latter result in Appendix A which also improves and unifies several existing results, see Remark A.4. In the deterministic case similar results have been shown in [GP02] and also for rough forcing terms in [BF09].

To prove our main results we apply the full power of the local well-posedness theory which we recently developed in [AV21a] together with the blow-up criteria and new bootstrapping techniques of our paper [AV21b]. In these works we completely revised the theory of abstract stochastic evolution equations for semi- and quasi-linear SPDEs, and is the stochastic analogue of [PSW18] of Prüss, Simonett and Wilke. One of the main ingredients in our theory is the so-called stochastic maximal  $L^p$ -regularity for the linear part, which we prove in a  $H^{s,q}$ -setting. In case of the Navier-Stokes equations this linear part is the turbulent Stokes system, for which we will prove stochastic maximal  $L^p$ -regularity in an  $H^{s,q}(\mathbb{T}^d; \mathbb{R}^d)$ -setting in Section 3, which can be seen as one of the main result of our work as well.

**1.3. Notation.** Here we collect some notation which will be used throughout the paper. Further notation will be introduced where needed. As usual, we write  $A \lesssim_P B$  (resp.  $A \gtrsim_P B$ ) whenever there is a constant  $C$  only depending on the parameter  $P$  such that  $A \leq CB$  (resp.  $A \geq CB$ ). We write  $C_P$  or  $C(P)$  if  $C$  depends only on  $P$ . Finally,  $B_Y(y, \eta) := \{y' \in Y : d_Y(y, y') < \eta\}$  if  $(Y, d_Y)$  is a metric space.

For  $a, \kappa \in \mathbb{R}$ , the weights will be denoted by  $w_\kappa^a = |t - a|^\kappa$  and  $w_\kappa := w_\kappa^0$ . For a Banach space  $X$ ,  $a, b \in \mathbb{R}$  and an interval  $(a, b) \subseteq \mathbb{R}_+$ ,  $L^p(a, b, w_\kappa^a; X)$  denotes the set of all strongly measurable maps  $f : (a, b) \rightarrow X$  such that

$$\|f\|_{L^p(a, b, w_\kappa^a; X)} := \left( \int_a^b \|f(t)\|_X^p w_\kappa^a(t) dt \right)^{1/p} < \infty.$$

Moreover,  $W^{1,p}(a, b, w_\kappa^a; X) \subseteq L^p(a, b, w_\kappa^a; X)$  denotes the set of all  $f$  such that  $f' \in L^p(a, b, w_\kappa^a; X)$  and we set

$$\|f\|_{W^{1,p}(a, b, w_\kappa^a; X)} := \|f\|_{L^p(a, b, w_\kappa^a; X)} + \|f'\|_{L^p(a, b, w_\kappa^a; X)}.$$

Let  $[\cdot, \cdot]_\theta$  and  $(\cdot, \cdot)_{\theta, p}$  be the complex and real interpolation functor, see e.g. [BL76, Tri95]. For each  $\theta \in (0, 1)$ , we set

$$H^{\theta, p}(a, b, w_\kappa^a; X) := [L^p(a, b, w_\kappa^a; X), W^{1,p}(a, b, w_\kappa^a; X)]_\theta.$$

The above spaces can also be equivalently introduced by using Fourier methods on  $\mathbb{R}$  and restrictions, see e.g. [ALV21, Definition 3.1] and the references therein.

For  $\mathcal{A} \in \{L^p, H^{\theta, p}\}$ , we denote by  $\mathcal{A}_{\text{loc}}^\theta(a, b; X)$  (resp.  $\mathcal{A}_{\text{loc}}^\theta([a, b]; X)$ ) the set of all strongly measurable maps  $f : (a, b) \rightarrow X$  such that  $f \in \mathcal{A}^\theta(c, d; X)$  for all  $a < c < d < b$  (resp.  $f \in \mathcal{A}^\theta(a, d; X)$  for all  $a < d < b$ ).

The  $d$ -dimensional torus is denoted by  $\mathbb{T}^d$ . For  $m \geq 1$  and  $s, r \in (0, 1)$ ,  $C_{\text{loc}}^{s, r}((a, b) \times \mathbb{T}^d; \mathbb{R}^m)$  denotes the space of all maps  $v : (a, b) \times \mathbb{T}^d \rightarrow \mathbb{R}^m$  such that for all  $a < a' < b' < b$  we have

$$|v(t, x) - v(t', x')| \lesssim_{a', b'} |t - t'|^s + |x - x'|^r, \quad t, t' \in [a', b'], x, x' \in \mathbb{T}^d.$$

Finally, we collect the main probabilistic notation. Through the paper, we fix a filtered probability space  $(\Omega, \mathcal{A}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$  and we let  $\mathbb{E}[\cdot] := \int_\Omega \cdot d\mathbb{P}$ . A mapping  $\tau : \Omega \rightarrow [0, \infty)$  is said to be a stopping time if  $\{\tau \leq t\} \in \mathcal{F}_t$  for all  $t \geq 0$ . For a stopping time  $\tau$  we let

$$\llbracket 0, \tau \rrbracket := [0, \tau] \times \Omega := \{(t, \omega) : 0 \leq t \leq \tau(\omega)\}.$$

Similar definitions, hold for  $\llbracket 0, \tau \rrbracket$ ,  $[0, \tau) \times \Omega$  etc.  $\mathcal{P}$  denotes the progressive  $\sigma$ -algebra on the above mentioned filtered probability space.

## 2. STATEMENT OF THE MAIN RESULTS

In this section we state our main results on the following stochastic Navier-Stokes equations on the  $d$ -dimensional torus  $\mathbb{T}^d$

$$(2.1) \quad \begin{cases} du = [\operatorname{div}(a \cdot \nabla u) - \operatorname{div}(u \otimes u) + f_0(\cdot, u) + \operatorname{div}(f(\cdot, u)) - \nabla p + \partial_h \tilde{p}] dt \\ \quad + \sum_{n \geq 1} [(b_n \cdot \nabla)u + g_n(\cdot, u) - \nabla \tilde{p}_n] dw_t^n, \\ \operatorname{div} u = 0, \\ u(\cdot, 0) = u_0. \end{cases}$$

Here  $u = (u^k)_{k=1}^d : [0, \infty) \times \Omega \times \mathbb{T}^d \rightarrow \mathbb{R}^d$  is the unknown velocity field,  $p, \tilde{p}_n : [0, \infty) \times \Omega \times \mathbb{T}^d \rightarrow \mathbb{R}$  the unknown pressures,  $(w^n)_{n \geq 1}$  is a sequence of standard independent Brownian motions on the previously mentioned filtered probability space  $(\Omega, \mathcal{A}, (\mathcal{F}_t)_{t \geq 0}, \mathbf{P})$ ,  $u \otimes u := (u^i u^j)_{i,j=1}^d$ , and

$$(2.2) \quad \begin{aligned} \operatorname{div}(a \cdot \nabla u) &= \left( \sum_{i,j=1}^d \partial_i (a^{i,j} \partial_j u^k) \right)_{k=1}^d, & (b_n \cdot \nabla)u &= \left( \sum_{j=1}^d b_n^j \partial_j u^k \right)_{k=1}^d, \\ \operatorname{div}(f(\cdot, u)) &= \left( \sum_{j=1}^d \partial_j (f_j^k(\cdot, u)) \right)_{k=1}^d, & \partial_h \tilde{p} &= \left( \sum_{n \geq 1} \sum_{j=1}^d \partial_j \tilde{p}_n h_n^{j,k} \right)_{k=1}^d \end{aligned}$$

where  $a^{ij}, b_n^j, h_n^{j,k}$  are given functions. As explained near (1.3) the  $b_n^j$  are used to model turbulence. The  $a^{ij}$  model a variable viscosity term and may also take into account Itô's corrections in case of stochastic Navier-Stokes equations in the Stratonovich form, see (1.4) and the discussion below it. Finally,  $\partial_h \tilde{p}$  models the effect of the turbulent part of the pressure  $\tilde{p}_n$  (see [MR04, Example 1] and [MR05]). The additional forcing terms  $f_0(u)$  and  $\operatorname{div}(f(\cdot, u))$  can be used to model the effect of gravity or the Coriolis force.

This section is organized as follows. In Subsections 2.1 and 2.2 we fix our notation and describe the spaces in which we consider (2.1). In Subsection 2.3 we present local existence and regularization phenomena for solutions to (2.1) with  $u_0$  in critical spaces (see Theorem 2.4), in Subsection 2.4 we state a stochastic version of the Serrin's criteria and in Subsection 2.5 we state global our existence result for small initial data if  $d \geq 3$ , and for general rough initial data if  $d = 2$ . The proofs will be given Section 4.

Let us emphasise that even in the case  $a^{ij} = \delta_{i,j}$ ,  $f_0 = 0$ ,  $f = 0$ , and  $h_n^{i,j} = 0$ , and with  $b \neq 0$  or  $g \neq 0$ , our well-posedness and regularity results are new:

- the wide class of initial data we can treat has not been considered before;
- new higher order regularity results are proved.

Moreover, the global well-posedness results for  $d = 2$  are also new even with the above simplifications.

**2.1. The Helmholtz projection and function spaces.** For any  $s \in \mathbb{R}$ ,  $p \in [1, \infty]$ ,  $q \in (1, \infty)$  and integers  $d, m \geq 1$ , let  $L^q(\mathbb{T}^d; \mathbb{R}^m)$ ,  $H^{s,q}(\mathbb{T}^d; \mathbb{R}^m)$  and  $B_{q,p}^s(\mathbb{T}^d; \mathbb{R}^m)$  denote the Lebesgue, Bessel-potential and Besov spaces on  $\mathbb{T}^d$  with values in  $\mathbb{R}^d$ , respectively. We refer to [ST87, Chapter 3] for details on periodic function spaces.

We denote by  $\mathbb{P}$  the *Helmholtz projection* which, for  $f = (f^n)_{n=1}^d \in C^\infty(\mathbb{T}^d; \mathbb{R}^d)$  and  $n \in \{1, \dots, d\}$ , is given by  $\mathbb{P}f = ((\mathbb{P}f)^n)_{n=1}^d$  where

$$(2.3) \quad (\widehat{\mathbb{P}f})^n(k) := \widehat{f}^n(k) - \sum_{j=1}^d \frac{k_j k_n}{|k|^2} \widehat{f}^j(k), \quad k \in \mathbb{Z}^d \setminus \{0\}, \quad (\widehat{\mathbb{P}f})^n(0) := \widehat{f}^n(0).$$

Here,  $\widehat{f}(k)$  denotes the  $k$ -th Fourier coefficient of  $f$ . Note that  $\operatorname{div}(\mathbb{P}f) = 0$  for all  $f \in C^\infty(\mathbb{T}^d; \mathbb{R}^d)$ , and by duality  $\mathbb{P} : \mathcal{D}'(\mathbb{T}^d; \mathbb{R}^d) \rightarrow \mathcal{D}'(\mathbb{T}^d; \mathbb{R}^d)$  where  $\mathcal{D}'(\mathbb{T}^d; \mathbb{R}^d) = (C^\infty(\mathbb{T}^d; \mathbb{R}^d))'$ . Moreover, by standard Fourier multiplier theorems (see [HNVW16, Theorem 5.7.11]),  $\mathbb{P}$  restricts uniquely to a bounded linear operator

$$(2.4) \quad \mathbb{P} : H^{s,q}(\mathbb{T}^d; \mathbb{R}^d) \rightarrow H^{s,q}(\mathbb{T}^d; \mathbb{R}^d).$$

For  $s \in \mathbb{R}$ ,  $q \in (1, \infty)$  and  $\mathcal{Y} \in \{L^q, H^{s,q}, B_{q,p}^s\}$ , we set

$$(2.5) \quad \mathbb{Y}(\mathbb{T}^d) := \left\{ f \in \mathcal{Y}(\mathbb{T}^d; \mathbb{R}^d) : \operatorname{div} f = 0 \text{ in } \mathcal{D}'(\mathbb{T}^d) \right\}, \quad \|f\|_{\mathbb{Y}(\mathbb{T}^d)} := \|f\|_{\mathcal{Y}(\mathbb{T}^d; \mathbb{R}^d)}.$$

Note that  $\mathbb{Y}(\mathbb{T}^d) = \mathbb{P}(\mathcal{Y}(\mathbb{T}^d; \mathbb{R}^d))$  for  $\mathcal{Y} \in \{L^q, H^{s,q}, B_{q,p}^s\}$  and  $s, q$  as above. By standard interpolation theory (see [ST87, Chapter 3.6] and [Tri95, Theorem 1.2.4])

$$(2.6) \quad \begin{aligned} \mathbb{H}^{s,q}(\mathbb{T}^d) &= [\mathbb{H}^{s_0,q}(\mathbb{T}^d), \mathbb{H}^{s_1,q}(\mathbb{T}^d)]_\theta, \\ \mathbb{B}_{q,p}^s(\mathbb{T}^d) &= (\mathbb{H}^{s_0,q}(\mathbb{T}^d), \mathbb{H}^{s_1,q}(\mathbb{T}^d))_{\theta,p}, \end{aligned}$$

for all  $s_0, s_1 \in \mathbb{R}$ ,  $\theta \in (0, 1)$ ,  $s := (1 - \theta)s_0 + \theta s_1$  and  $p, q \in (1, \infty)$ .

**2.2. Main assumptions and definitions.** Here we collect the main assumptions and definitions. Further hypotheses will be given where needed. Below  $\mathcal{P}$  and  $\mathcal{B}$  denotes the progressive and Borel  $\sigma$ -algebra, respectively.

**Assumption 2.1.** *Let  $d \geq 2$ . We say that Assumption 2.1( $p, \kappa, q, \delta$ ) holds if  $\delta \in (-1, 0]$ , and one of the following two is satisfied:*

- $q \in [2, \infty)$ ,  $p \in (2, \infty)$  and  $\kappa \in [0, \frac{p}{2} - 1)$ ;
- $q = p = 2$  and  $\kappa = 0$ ,

and the following conditions hold:

- (1) For all  $i, j \in \{1, \dots, d\}$  and  $n \geq 1$ , the mappings  $a^{i,j}, \phi_n^j, h_n^{i,j} : \mathbb{R}_+ \times \Omega \times \mathbb{T}^d \rightarrow \mathbb{R}$  are  $\mathcal{P} \otimes \mathcal{B}(\mathbb{T}^d)$ -measurable;
- (2) There exist an  $M > 0$  and  $\eta > \max\{d/\xi, -\delta\}$ , with  $\xi \in [2, \infty)$  such that a.s. for all  $t \in \mathbb{R}_+$  and  $i, j \in \{1, \dots, d\}$ ,

$$\|a^{i,j}(t, \cdot)\|_{H^{\eta, \xi}(\mathbb{T}^d)} + \|(b_n^j(t, \cdot))_{n \geq 1}\|_{H^{\eta, \xi}(\mathbb{T}^d; \ell^2)} + \|(h_n^{i,j}(t, \cdot))_{n \geq 1}\|_{H^{\eta, \xi}(\mathbb{T}^d; \ell^2)} \leq M.$$

- (3) There exists  $\nu > 0$  such that a.s. for all  $t \in \mathbb{R}_+$ ,  $\Upsilon \in \mathbb{R}^d$  and  $x \in \mathbb{T}^d$ ,

$$\sum_{i,j=1}^d \left[ a^{i,j}(t, x) - \frac{1}{2} \left( \sum_{n \geq 1} b_n^i(t, x) b_n^j(t, x) \right) \right] \Upsilon_i \Upsilon_j \geq \nu |\Upsilon|^2.$$

- (4) For all  $j \in \{0, \dots, d\}$  and  $n \geq 1$ , the maps  $f_j, g_n : \mathbb{R}_+ \times \Omega \times \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  are  $\mathcal{P} \otimes \mathcal{B}(\mathbb{T}^d \times \mathbb{R}^d)$ -measurable;
- (5) For all  $j \in \{0, \dots, d\}$ ,  $f_j(\cdot, 0) \in L^\infty(\mathbb{R}_+ \times \Omega \times \mathbb{T}^d; \mathbb{R}^d)$ ,  $(g_n)_{n \geq 1}(\cdot, 0) \in L^\infty(\mathbb{R}_+ \times \Omega \times \mathbb{T}^d; \ell^2)$ , and a.s. for all  $t \in \mathbb{R}_+$ ,  $x \in \mathbb{T}^d$  and  $y, y' \in \mathbb{R}^d$ ,

$$|f_j(t, x, y) - f_j(t, x, y')| + \|(g_n(t, x, y) - g_n(t, x, y'))_{n \geq 1}\|_{\ell^2} \lesssim (1 + |y| + |y'|)|y - y'|.$$



The parameter  $p$  will be used for  $L^p$ -integrability in time with weight  $t^\kappa dt$ . The parameter  $q$  will be used for integrability in space with Sobolev smoothness  $1 + \delta$ . In the main results below we will always use  $\delta \in [-\frac{1}{2}, 0]$ . However, in Section 4 we will also consider  $\delta \in (-1, 0]$ . By Sobolev embeddings and Assumption 2.1(2) we have

$$(2.7) \quad \|a^{i,j}\|_{C^{\eta-\frac{d}{\xi}}(\mathbb{T}^d)} + \|(b_n^j)_{n \geq 1}\|_{C^{\eta-\frac{d}{\xi}}(\mathbb{T}^d; \ell^2)} \lesssim_{\eta, \xi} M.$$

*Remark 2.2.*

- If Assumption 2.1 holds for  $\delta = 0$ , then it also holds for some  $\delta < 0$ .
- The regularity assumption on  $a^{i,j}, b^j$  in (2) also fit naturally in the case of stochastic Navier-Stokes equations in Stratonovich form, see (1.4) and the text below it. Indeed, if  $b^j \in H^{\eta, \xi}(\mathbb{T}^d; \ell^2)$  uniformly in  $(t, \omega)$ , then by (2.7) and [AV21c, Proposition 4.1(1)] the Itô correction satisfies  $a_b^{i,j} = \frac{1}{2} \sum_{n \geq 1} b_n^i b_n^j \in H^{\eta, \xi}(\mathbb{T}^d)$  uniformly in  $(t, \omega)$ .
- Assumption 2.1 also covers the case of  $b^j$  with  $\gamma$ -Hölder smoothness which is the natural one as explained in the introduction. Indeed,  $C^\gamma(\mathbb{T}^d; \ell^2) \hookrightarrow H^{\eta, \xi}(\mathbb{T}^d; \ell^2)$  for all  $\gamma > \eta$  and  $\xi \in (1, \infty)$ . Note that typically  $\gamma \in (0, 1)$  is small in applications. A similar remark holds for  $a^{i,j}$ .
- The quadratic growth of  $f_j$  and  $g_n$  shows that we can allow nonlinearities which are as strong as the convection term  $\operatorname{div}(u \otimes u)$  (see Subsection 1.1).

To introduce the notion of solutions to (2.1) we rewrite it as a stochastic evolution equation for the unknown velocity field  $u$ . To this end, to have a divergence free stochastic part we need

$$\nabla \tilde{p}_n = (I - \mathbb{P})[(b_n \cdot \nabla)u + g_n(\cdot, u)]$$

where  $I$  is the identity operator. Therefore, letting  $h_n^{j,k} := (h_n^{j,k})_{j=1}^d$  and setting

$$(2.8) \quad \tilde{f}(\cdot, u) := \left( \sum_{n \geq 1} \left[ (I - \mathbb{P})[(b_n \cdot \nabla)u + g_n(\cdot, u)] \right] \cdot h_n^{j,k} \right)_{k=1}^d,$$

we obtain  $\tilde{f}(\cdot, u) = \partial_h \tilde{p}$ . Applying the Helmholtz projection to the first equation in (2.1), at least formally, the system (2.1) is equivalent to

$$(2.9) \quad \begin{cases} du = \mathbb{P}[\operatorname{div}(a \cdot \nabla u) - \operatorname{div}(u \otimes u) + f_0(\cdot, u) + \operatorname{div}(f(\cdot, u)) + \tilde{f}(\cdot, u)] dt \\ \quad + \sum_{n \geq 1} \mathbb{P}[(b_n \cdot \nabla)u + g_n(\cdot, u)] dw_t^n, \\ u(\cdot, 0) = u_0. \end{cases}$$

We are in position to define solutions to (2.1). We recall that the sequence of independent Brownian motions  $(w^n)_{n \geq 1}$  induces uniquely an  $\ell^2$ -cylindrical Brownian motion  $W_{\ell^2}$  (see e.g. [AV21a, Example 2.12]).

**Definition 2.3.** *Let Assumption 2.1( $p, \kappa, q, \delta$ ) be satisfied.*

- (1) *Let  $\sigma$  be a stopping time and  $u : [0, \sigma) \times \Omega \rightarrow \mathbb{H}^{1+\delta, q}(\mathbb{T}^d)$  be a stochastic process.  $(u, \sigma)$  is called a local  $(p, \kappa, q, \delta)$ -solution to (2.1) if there exists a sequence of stopping times  $(\sigma_j)_{j \geq 1}$  such that the following hold.*

- $\sigma_j \leq \sigma$  a.s. for all  $j \geq 1$  and  $\lim_{j \rightarrow \infty} \sigma_j = \sigma$  a.s.;
- For all  $j \geq 1$ , the process  $\mathbf{1}_{[0, \sigma_j] \times \Omega} u$  is progressively measurable;

(2.10)  $\bullet$  a.s. for all  $j \geq 1$ , we have  $u \in L^p(0, \sigma_j, w_\kappa; \mathbb{H}^{1+\delta, q}(\mathbb{T}^d))$  and

$$-\operatorname{div}(u \otimes u) + f_0(\cdot, u) + \operatorname{div}(f(\cdot, u)) + \tilde{f}(\cdot, u) \in L^p(0, \sigma_j, w_\kappa; H^{-1+\delta, q}(\mathbb{T}^d; \mathbb{R}^d)),$$

$$(g_n(\cdot, u))_{n \geq 1} \in L^p(0, \sigma_j, w_\kappa; H^{\delta, q}(\mathbb{T}^d; \ell^2(\mathbb{R}^d))).$$

(2.11)  $\bullet$  a.s. for all  $j \geq 1$  the following equality holds for all  $t \in [0, \sigma_j]$ :

$$u(t) - u_0 = \int_0^t \mathbb{P}[\operatorname{div}(a \cdot \nabla u) - \operatorname{div}(u \otimes u) + f_0(\cdot, u) + \operatorname{div}(f(\cdot, u)) + \tilde{f}(\cdot, u)] ds$$

$$+ \int_0^t (\mathbf{1}_{[0, \sigma_j]} \mathbb{P}[(b_n \cdot \nabla)u + g_n(\cdot, u)])_{n \geq 1} dW_{\ell^2}(s).$$

(2)  $(u, \sigma)$  is called a  $(p, \kappa, q, \delta)$ -solution to (2.1) if for any other local  $(p, \kappa, q, \delta)$ -solution  $(v, \tau)$  to (2.1) we have  $\tau \leq \sigma$  a.s. and  $u = v$  a.e. on  $[0, \sigma] \times \Omega$ .

Note that  $(p, q, \delta, \kappa)$ -solutions are *unique* by definition. By Assumption 2.1 and [AV21c, Proposition 4.1] a.s. for all  $j \geq 1$ ,

$$\operatorname{div}(a \cdot \nabla u) \in L^p(0, \sigma_j; H^{-1+\delta, q}(\mathbb{T}^d; \mathbb{R}^d)),$$

$$((b_n \cdot \nabla)u)_{n \geq 1} \in L^p(0, \sigma_j; H^{\delta, q}(\mathbb{T}^d; \ell^2(\mathbb{R}^d))).$$

Thus the deterministic integrals in (2.11) are well-defined as Bochner integrals with values in  $H^{-1+\delta, q}(\mathbb{T}^d; \mathbb{R}^d)$ . The stochastic integral is well-defined as an  $H^{\delta, q}$ -valued stochastic integral (see [NVW15, Theorem 4.7] and use the identity (4.7) below). By Assumption 2.1, Lemma 4.2 and [AV21a, Lemma 4.10 and 4.12], (2.10) holds provided

$$u \in \bigcap_{\theta \in [0, 1/2)} H^{\theta, p}(0, \sigma_j, w_\kappa; \mathbb{H}^{1+s-2\theta, q}(\mathbb{T}^d)) \quad \text{a.s. for all } j \geq 1.$$

**2.3. Local existence and regularization.** The following is our main local well-posedness result. It is formulated for the space of critical initial data. The general case is covered by Theorem 4.1, where  $\delta \in (-1, 1/2)$  is included as well. Global well-posedness for small initial data will be discussed afterwards in Theorem 2.11, and the special case  $d = 2$  is considered in Theorem 2.12.

**Theorem 2.4** (Local well-posedness). *Let Assumption 2.1( $p, \kappa, q, \delta$ ) be satisfied and suppose that one of the following conditions holds:*

- $\bullet$   $\delta \in [-\frac{1}{2}, 0]$ ,  $\frac{d}{2+\delta} < q < \frac{d}{1+\delta}$ ,  $\frac{2}{p} + \frac{d}{q} \leq 2 + \delta$ , and  $\kappa = \kappa_c = -1 + \frac{p}{2}(2 + \delta - \frac{d}{q})$ ;
- $\bullet$   $\delta = \kappa = 0$  and  $p = q = d = 2$ .

Then for all  $u_0 \in L^0_{\mathcal{F}_0}(\Omega; \mathbb{B}_{q, p}^{\frac{d}{q}-1}(\mathbb{T}^d))$ , (2.1) has a unique  $(p, \kappa_c, q, \delta)$ -solution  $(u, \sigma)$  such that a.s.  $\sigma > 0$  and

$$(2.12) \quad u \in L^p_{\text{loc}}([0, \sigma], w_{\kappa_c}; \mathbb{H}^{1+\delta, q}(\mathbb{T}^d)) \cap C([0, \sigma]; \mathbb{B}_{q, p}^{\frac{d}{q}-1}(\mathbb{T}^d)).$$

Moreover,  $(u, \sigma)$  instantaneously regularizes in time and space: a.s.

$$(2.13) \quad u \in H^{\theta, r}_{\text{loc}}(0, \sigma; \mathbb{H}^{1-2\theta, \zeta}(\mathbb{T}^d)) \quad \text{for all } \theta \in [0, 1/2), r, \zeta \in (2, \infty),$$

$$(2.14) \quad u \in C^{\theta_1, \theta_2}_{\text{loc}}((0, \sigma) \times \mathbb{T}^d; \mathbb{R}^d) \quad \text{for all } \theta_1 \in [0, 1/2), \theta_2 \in (0, 1).$$

In case  $p > 2$ , one can check that by the conditions on  $p$  and  $q$  one has  $\kappa_c \in [0, \frac{p}{2} - 1)$ . The assertions (2.13)-(2.14) show an instantaneous regularization for solutions to (2.1). By Sobolev embedding one can check that (2.13) implies (2.14).

In particular,  $u$  becomes *almost*  $C^1$  in space for rough initial data  $u_0$ , rough coefficient  $a^{ij}$ ,  $h^{ij}$ , and rough nonlinearities  $f_0, f$  and  $g$ . If  $\eta > \max\{d/\xi, 1/2\}$  in Assumption 2.1, then we can set  $\delta = -1/2$ , and therefore taking  $q \uparrow 2d$  it suffices to know that for some  $p > 2$  and  $\varepsilon > 0$ ,

$$u_0 \in L^0_{\mathcal{F}_0}(\Omega; \mathbb{B}^{\varepsilon - \frac{1}{2}}_{2d,p}(\mathbb{T}^d)).$$

It would be interesting to know whether this is optimal.

Setting  $q = d$  it follows from  $L^d(\mathbb{T}^d; \mathbb{R}^d) \hookrightarrow B^0_{d,p}(\mathbb{T}^d; \mathbb{R}^d)$  for  $p \in [d, \infty)$  that Theorem 2.4 (with  $\delta < 0$ ) also applies to initial data from the scaling invariant space  $L^0_{\mathcal{F}_0}(\Omega; \mathbb{L}^d(\mathbb{T}^d))$ . By Remark 2.2, it is enough to know that Assumption 2.1( $p, \kappa, d, 0$ ) holds where  $\kappa = -1 + \frac{p}{2}(1 + \delta)$  and  $p \geq d$ .

*Remark 2.5.* If  $p > 2$ , then it follows from the proof of Theorem 2.4 that the following weighted regularity holds up to  $t = 0$ :

$$u \in H^{\theta,p}_{\text{loc}}([0, \sigma]; w_{\kappa_c}; \mathbb{H}^{1+\delta,q}(\mathbb{T}^d)) \quad \text{for all } \theta \in [0, 1/2).$$

Next we consider *higher order regularity*. We are particular interested in studying how the regularity of  $b_n$  in the transport term  $(b_n \cdot \nabla)u$  effects the regularity of  $u$ . By regularization one could expect that  $u$  gains one order of smoothness compared to  $b_n$ . The following result shows that this is indeed the case. Of course in order to prove such a result we also need further smoothness of the nonlinearities  $f_0, f, g$ . We emphasize that we do not impose further conditions on the initial data.

The next assumption roughly says that  $f_0, f$  and  $(g_n)$  are  $C^{[\eta+1]}$  in the  $y$ -variable, where  $\eta > 0$  is some fixed number.

**Assumption 2.6.** *Let  $\eta > 0$ ,  $f_0, f$  and  $g_n$  be as in Assumption 2.1. We assume that  $f_0, f$  and  $g_n$  are  $x$ -independent,  $C^{[\eta+1]}$  in  $y$  and, a.s. for all  $N \geq 1$  there is a  $C_N > 0$  such that*

$$\sum_{j=1}^{[\eta+1]} |\partial_y^j f_0(t, y)| + |\partial_y^j f(t, y)| + \|(\partial_y^j g_n(t, y))_{n \geq 1}\|_{\ell^2} \leq C_N, \quad |v| \leq N, \quad t \geq 0.$$

**Theorem 2.7** (Higher order regularity). *Let the assumptions of Theorem 2.4 be satisfied, where  $(\eta, \xi)$  are such that Assumption 2.1(2) holds, i.e.  $\eta > \max\{\xi/d, -\delta\}$ ,  $\xi \in [2, \infty)$ , and there exists an  $M$  such that*

$$\|a^{i,j}(t, \cdot)\|_{H^{\eta,\xi}(\mathbb{T}^d)} + \|(b_n^j(t, \cdot))_{n \geq 1}\|_{H^{\eta,\xi}(\mathbb{T}^d; \ell^2)} + \|(h_n^{i,j}(t, \cdot))_{n \geq 1}\|_{H^{\eta,\xi}(\mathbb{T}^d; \ell^2)} \leq M.$$

*Furthermore, suppose that Assumption 2.6 holds. Let  $(u, \sigma)$  be the  $(p, \kappa_c, q, \delta)$ -solution to (2.1) provided by Theorem 2.4. Then a.s.*

$$(2.15) \quad u \in H^{\theta,r}_{\text{loc}}(0, \sigma; \mathbb{H}^{1+\eta-2\theta,\xi}(\mathbb{T}^d)) \quad \text{for all } \theta \in [0, 1/2), \quad r \in (2, \infty),$$

$$(2.16) \quad u \in C^{\theta_1, \theta_2 + \eta - \frac{d}{\xi}}_{\text{loc}}((0, \sigma) \times \mathbb{T}^d; \mathbb{R}^d) \quad \text{for all } \theta_1 \in [0, 1/2), \quad \theta_2 \in (0, 1).$$

From the above theorem one can see how the regularity of order  $\eta$  of the coefficients appears in (2.15) and (2.16). In particular, (2.15) with  $\theta = 0$  shows that the regularity of  $u$  is one order higher than the regularity of  $(a, b, h)$ .

*Remark 2.8.* If  $u_0 \in L^0_{\mathcal{F}_0}(\Omega; \mathbb{B}_{\xi, r}^{1+\eta-\frac{2}{r}}(\mathbb{T}^d))$  for some fixed  $r \in (2, \infty)$ , then one can check from the proofs that the regularity result (2.15) (for the fixed  $r$ ) holds locally on  $[0, \sigma)$  instead of  $(0, \sigma)$ . However, this will not be used in the sequel.

**2.4. Stochastic Serrin's blow-up criteria.** In this section we state some blow-up criteria for (2.1) which can be seen as a stochastic analogue of the *Serrin's* criteria for Navier-Stokes equations (see e.g. [LR16, Theorem 11.2]).

**Theorem 2.9** (Stochastic Serrin's criteria). *Let the conditions of Theorem 2.4 be satisfied and let  $(u, \sigma)$  be the  $(p, \kappa_c, q, \delta)$ -solution to (2.1). Suppose that Assumption 2.1( $p_0, 0, q_0, \delta_0$ ) holds in one of the following cases:*

- $\delta_0 \in [-\frac{1}{2}, 0]$ ,  $\frac{d}{2+\delta_0} < q_0 < \frac{d}{1+\delta_0}$  and  $\frac{2}{p_0} + \frac{d}{q_0} \leq 2 + \delta_0$ ;
- $\delta_0 = 0$  and  $p_0 = q_0 = d = 2$ .

Then for each  $T, \varepsilon \in (0, \infty)$ :

$$(2.17) \quad \mathbb{P}\left(\varepsilon < \sigma < T, \|u\|_{L^{p_0}(\varepsilon, \sigma; H^{\gamma_0, q_0}(\mathbb{T}^d; \mathbb{R}^d))} < \infty\right) = 0, \quad \text{where } \gamma_0 = \frac{2}{p_0} + \frac{d}{q_0} - 1.$$

In particular, for all  $T, \varepsilon \in (0, \infty)$  the following hold:

- (1)  $\mathbb{P}\left(\varepsilon < \sigma < T, \sup_{t \in [\varepsilon, \sigma]} \|u(t)\|_{B_{q_1, \infty}^{d/q_0-1}(\mathbb{T}^d; \mathbb{R}^d)} < \infty\right) = 0$  if  $q_1 > q_0$ ;
- (2)  $\mathbb{P}\left(\varepsilon < \sigma < T, \|u\|_{L^{p_0}(\varepsilon, \sigma; L^{q_0}(\mathbb{T}^d; \mathbb{R}^d))} < \infty\right) = 0$  if  $p_0$  satisfies  $\frac{2}{p_0} + \frac{d}{q_0} = 1$ .

Item (2) is the stochastic version of the well-known Serrin's criteria. It is a special case of (2.17) where we can even allow negative smoothness. In particular, one can take  $p_0 = p$ ,  $q_0 = q$  and  $\delta_0 = 0$ , but the above formulation is much more flexible as it allows us to transfer information on  $\sigma$  between different settings. An interesting choice is  $p_0 = q_0 = 2$  and  $\delta_0 = 0$  in case  $d = 2$ , and will be used in Theorem 2.12 for  $d = 2$ . Note that by (2.13)-(2.14), the paths of  $u$  are regular enough on  $(0, \sigma)$  to consider any of the above norms.

*Remark 2.10.*

- (1) Choosing  $q_0, p_0$  large enough, one has  $\frac{2}{p_0} + \frac{d}{q_0} < 1$  (thus  $\gamma_0 < 0$ ) and (2.17) provides blow-up criteria in spaces of *negative* smoothness. The same holds for (1) if  $q_0 > d$ . To see how far one can go note that for  $\delta_0 = -\frac{1}{2}$  and  $q_0 \rightarrow 2d$ ,  $p_0 \rightarrow \infty$  one would obtain  $\frac{d}{q_0} - 1 \rightarrow -\frac{1}{2}$  in (1) and  $\gamma_0 \rightarrow -\frac{1}{2}$  in (2.17).
- (2) The *space-time Sobolev index* (keeping in mind the parabolic scaling) of the space  $L^{p_0}(0, t; H^{\gamma_0, q_0})$  appearing in (2.17) and (2) is given by  $-\frac{2}{p_0} + \gamma_0 - \frac{d}{q_0} = -1$ . This space-time Sobolev index is equivalent to the Sobolev index of the space of initial data  $\mathbb{B}_{q, p}^{d/q-1}(\mathbb{T}^d)$ , which means that these blow-up criteria are optimal. In contrary, the space-time Sobolev index of  $L^\infty(0, t; B_{q_1, p_0}^{d/q_0-1})$  in (1) is given by  $-1 + d(\frac{1}{q_0} - \frac{1}{q_1}) > -1$  which is sub-optimal. We do not know if (1) holds in the borderline case  $q_1 = q_0$ .

In the deterministic setting, blow-up criteria in the critical space  $L^3$  (here  $d = 3$  for simplicity) are known to be valid even with quantitative growth assumption in the  $L^\infty((\varepsilon, \sigma); L^3)$ -norm, see [Tao19, Theorem 1.4]. It would be interesting to extend Theorem 2.9 into this direction by exploiting the structure of the equation (2.1).

**2.5. Global existence results.** Global well-posedness for the deterministic Navier-Stokes equations is known for small initial data. The next result is its stochastic analogue.

**Theorem 2.11** (Global well-posedness for small and rough initial data). *Let Assumption 2.1( $p, \kappa, q, \delta$ ) be satisfied and suppose that one of the following conditions holds:*

- $\delta \in [-\frac{1}{2}, 0]$ ,  $\frac{d}{2+\delta} < q < \frac{d}{1+\delta}$ ,  $\frac{2}{p} + \frac{d}{q} \leq 2 + \delta$ , and  $\kappa = \kappa_c = -1 + \frac{p}{2}(2 + \delta - \frac{d}{q})$ ;
- $\delta = \kappa = 0$  and  $p = q = d = 2$ .

Assume that there are  $M_1, M_2 > 0$  such that a.s. for all  $t \geq 0$  and  $y \in \mathbb{R}^d$ ,

$$(2.18) \quad \sum_{j=0}^d |f_j(t, y)| + \|(g_n(t, y))_{n \geq 1}\|_{\ell^2} \leq M_1(1 + |y|) + M_2|y|^2.$$

Let  $u_0 \in L^p_{\mathcal{F}_0}(\Omega; \mathbb{B}_{q,p}^{\frac{d}{q}-1}(\mathbb{T}^d))$ . Let  $(u, \sigma)$  be the  $(p, \kappa_c, q, \delta)$ -solution to (2.1) provided by Theorem 2.4. Let  $0 < \varepsilon < T < \infty$ . Then there exists  $C_{\varepsilon, T} > 0$  independent of  $u_0, u$  such that

$$\mathbb{E} \|u_0\|_{\mathbb{B}_{q,p}^{\frac{d}{q}-1}(\mathbb{T}^d; \mathbb{R}^d)}^p + M_1^p \leq C_{\varepsilon, T} \implies \mathbb{P}(\sigma \geq T) > 1 - \varepsilon.$$

In other words, the previous result shows that if  $u_0$  and  $M_1$  are close to 0, then  $u$  exists up to  $T$  with probability  $> 1 - \varepsilon$ . In Corollary 4.6 we also provide an estimate for  $u$  on  $\{\sigma \geq T\}$  in the  $H^{\theta, p}(H^{1+\delta-2\theta, q})$ -norm. Recall that  $\mathbb{L}^d(\mathbb{T}^d; \mathbb{R}^d) \hookrightarrow \mathbb{B}_{d,p}^0(\mathbb{T}^d)$  for all  $p \geq d$ . Thus Theorem 2.11 also proves global existence with high probability under a smallness assumption on  $u_0$  in the scaling invariant Lebesgue space  $L^d(\mathbb{T}^d; \mathbb{R}^d)$ .

Next we turn our attention to the two-dimensional setting in which one can avoid smallness of the initial data. A variation of a classical result for  $\mathbb{L}^2$ -initial data can be found in Theorem A.3 in the appendix, where the results are partly based on [AV21a]. The main new contributions in the result below are:

- We obtain global existence for initial data from  $\mathbb{B}_{q,p}^{2/q-1}(\mathbb{T}^2)$  for suitable  $p \in [2, \infty)$  and  $q \in [2, 4)$ . By Sobolev embedding the latter space contains  $\mathbb{L}^2(\mathbb{T}^2)$ . By the restriction  $q < 4$  below, we see that we can allow smoothness of negative order  $-\frac{1}{2} + \varepsilon$  for any  $\varepsilon > 0$ .
- We prove arbitrary spatial regularity, and time regularity of order  $C^{\frac{1}{2}-\varepsilon}$  for any  $\varepsilon > 0$ , where the initial data is still assumed to be irregular.

**Theorem 2.12** (Global well-posedness in two dimensions with rough initial data). *Let  $d = 2$ . Suppose that Assumption 2.1( $p, \kappa, q, \delta$ ) and suppose that one of the following conditions holds:*

- $\delta \in [-\frac{1}{2}, 0]$ ,  $\frac{2}{2+\delta} < q < \frac{2}{1+\delta}$ ,  $\frac{2}{p} + \frac{2}{q} \leq 2 + \delta$ , and  $\kappa = \kappa_c = -1 + \frac{p}{2}(2 + \delta - \frac{2}{q})$ ;
- $\delta = \kappa = 0$  and  $p = q = 2$ .

Assume that  $f$  and  $g$  additionally satisfy the following linear growth assumption: there exists a  $C > 0$  such that a.s. for all  $t \geq 0$ ,  $x \in \mathbb{T}^2$  and  $y \in \mathbb{R}^2$ ,

$$\sum_{j \in \{0,1,2\}} \sum_{k \in \{1,2\}} |f_j^k(t, x, y)| + \|(g_n(t, x, y))_{n \geq 1}\|_{\ell^2} \leq C(1 + |y|).$$

Then for all  $u_0 \in L^0_{\mathcal{F}_0}(\Omega; \mathbb{B}_{q,p}^{\frac{2}{q}-1}(\mathbb{T}^2))$  there exists a unique global  $(p, q, \delta, \kappa_c)$ -solution  $u$  of (2.1) with

$$u \in L^p_{\text{loc}}([0, \infty), w_{\kappa_c}; \mathbb{H}^{1+\delta, q}(\mathbb{T}^2)) \cap C([0, \infty); \mathbb{B}_{q,p}^{\frac{2}{q}-1}(\mathbb{T}^2)).$$

Moreover,

$$\begin{aligned} u &\in H^{\theta, r}_{\text{loc}}(0, \infty; \mathbb{H}^{1-2\theta, \zeta}(\mathbb{T}^2)) \quad \text{for all } \theta \in [0, 1/2), \quad r, \zeta \in (2, \infty), \\ u &\in C^{\theta_1, \theta_2}_{\text{loc}}((0, \infty) \times \mathbb{T}^2; \mathbb{R}^2) \quad \text{for all } \theta_1 \in [0, 1/2), \quad \theta_2 \in (0, 1). \end{aligned}$$

Furthermore, if the conditions of Theorem 2.7 hold for  $d = 2$ , then  $u$  satisfies (2.15) and (2.16) with  $\sigma = \infty$ .

In particular, note that if  $a^{ij}$ ,  $b^j$ ,  $h^{ij}$ ,  $f_j$ , and  $g$  satisfy the conditions of Theorem 2.7 for each  $\eta > 0$  and  $\xi \in [2, \infty)$ , then a.s. the paths of  $u$  have regularity up to any order in space:

$$u \in C^{\theta, \infty}_{\text{loc}}((0, \infty) \times \mathbb{T}^2; \mathbb{R}^2), \quad \text{for all } \theta \in (0, 1/2).$$

The above regularity results of solutions to (2.1) appears to be new. It is not possible to prove this by classical bootstrapping arguments based on  $L^2$ -theory since the nonlinearity  $\text{div}(u \otimes u)$  is critical in this setting. Indeed, after one proves existence of a *global* solution  $u$  for which the paths satisfy a.s.  $u \in L^2(0, T; H^{1,2}) \cap C([0, T]; L^2)$ , one is tempted to prove further regularity by analyzing the term  $\text{div}(u \otimes u)$ . Note that the following embeddings are sharp in dimension two

$$L^2(0, T; H^{1,2}) \cap C([0, T]; L^2) \hookrightarrow L^4(0, T; H^{1/2,2}) \hookrightarrow L^4(0, T; L^4).$$

Therefore, we can only conclude  $\text{div}(u \otimes u) \in L^2(0, T; H^{-1,2})$  which is not good enough to bootstrap regularity. In order to prove the desired smoothness we will apply our recent bootstrapping method from [AV21b, Sections 6.2 and 6.3] which is based on arguments involving weights in time.

We conclude by remarking that in general, one can not improve the starting point  $u \in L^2(0, T; H^{1,2}) \cap C([0, T]; L^2)$  a.s. of the above argument by using energy methods. Indeed, as remarked in Section 1, our regularity assumptions on  $a, b$ , and  $h$  do *not* allow us to reformulate (2.1) in terms of the *vorticity*, i.e. for the unknown  $\text{curl } u$ . Thus besides the usual energy estimate in the space  $L^2(0, T; H^1) \cap C([0, T]; L^2)$ , no other estimates are available for solutions to (2.1) in general.

### 3. MAXIMAL $L^p$ -REGULARITY FOR THE TURBULENT STOKES COUPLE

Before we turn to the main results on the stochastic Navier-Stokes equations with gradient noise (2.1), we analyse the linear part of the problem, which is central in our approach. In order to be as flexible as possible in later applications we obtain  $L^p((s, T), w_{\kappa_c}^s, H^{1+\delta, q})$ -regularity estimates for any  $p, q \geq 2$ ,  $\kappa_c \in [0, \frac{p}{2} - 1)$  and  $\delta \in \mathbb{R}$ . We consider the following *turbulent Stokes system* on  $\mathbb{T}^d$ :

$$(3.1) \quad \begin{cases} du + \mathbb{P}Au \, dt = f \, dt + \sum_{n \geq 1} (\mathbb{P}\mathcal{B}_n u + g_n) dw_t^n, \\ u(s) = 0, \end{cases}$$

Here  $s \in [0, \infty)$ ,  $(w^n)_{n \geq 1}$  is a sequence of independent standard Brownian motion on  $(\Omega, \mathcal{A}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ ,  $\mathbb{P}$  denotes the Helmholtz projection, and for  $v = (v^k)_{k=1}^d$ ,

$$(3.2) \quad \begin{aligned} \mathcal{A}(t)v &= \left( -\operatorname{div}(a(t, \cdot) \cdot \nabla v^k) - \sum_{n \geq 1} \left[ (I - \mathbb{P})((b_n(t) \cdot \nabla)v) \right] \cdot h_n^k \right)_{k=1}^d, \\ \mathcal{B}_n(t)v &= \left( (b_n(t) \cdot \nabla)v \right)_{k=1}^d, \end{aligned}$$

where  $a, b$  and  $h$  are given.

**3.1. Assumptions and main results.** Suppose that the following conditions hold:

**Assumption 3.1.** *Let  $d \geq 2$ ,  $\delta \in \mathbb{R}$  and  $0 \leq s < T < \infty$ . Suppose that one of the following holds*

- $q \in [2, \infty)$ ,  $p \in (2, \infty)$  and  $\kappa \in [0, \frac{p}{2} - 1]$ ;
- $q = p = 2$  and  $\kappa = 0$ ,

and that the following conditions hold:

- (1) For all  $i, j \in \{1, \dots, d\}$  and  $n \geq 1$ , the mappings  $a^{i,j}, b_n^j, h_n^{i,j} : \mathbb{R}_+ \times \Omega \times \mathbb{T}^d \rightarrow \mathbb{R}$  are  $\mathcal{P} \otimes \mathcal{B}(\mathbb{T}^d)$ -measurable;
- (2) Let  $\xi \in [q', \infty)$  and  $\eta > \frac{d}{q}$  be such that one of the following hold:
  - $\delta \leq 0$  and  $\eta > -\delta$ ;
  - $\delta > 0$ ,  $\eta \geq \delta$ ,  $\xi \leq q$  and  $\eta - \frac{d}{\xi} \geq \delta - \frac{d}{q}$ .

There exists an  $M > 0$  such that a.s. for all  $t \in \mathbb{R}_+$  and  $i, j \in \{1, \dots, d\}$ ,

$$\begin{aligned} &\|a^{i,j}(t, \cdot)\|_{H^{\eta, \xi}(\mathbb{T}^d)} + \|(b_n^j(t, \cdot))_{n \geq 1}\|_{H^{\eta, \xi}(\mathbb{T}^d; \ell^2)} \\ &\quad + \|(h_n^{i,j}(t, \cdot))_{n \geq 1}\|_{H^{\eta, \xi}(\mathbb{T}^d; \ell^2)} \leq M. \end{aligned}$$

- (3) There exists  $\nu > 0$  such that a.s. for all  $t \in \mathbb{R}_+$ ,  $\Upsilon \in \mathbb{R}^d$  and  $x \in \mathbb{T}^d$ ,

$$\sum_{i,j=1}^d \left[ a^{i,j}(t, x) - \frac{1}{2} \left( \sum_{n \geq 1} b_n^i(t, x) b_n^j(t, x) \right) \right] \Upsilon_i \Upsilon_j \geq \nu |\Upsilon|^2.$$

By Sobolev embeddings and Assumption 3.1(2), for all  $i, j \in \{1, \dots, d\}$ ,

$$(3.3) \quad a^{i,j} \in C^\alpha(\mathbb{T}^d), \quad \text{and} \quad (b_n^j)_{n \geq 1}, (h_n^{i,j})_{n \geq 1} \in C^\alpha(\mathbb{T}^d; \ell^2),$$

where  $\alpha := \eta - \frac{d}{\xi} > 0$ . Note that Assumption 3.1 coincides with part of Assumption 2.1( $p, \kappa, q, \delta$ ) in the case  $\delta \in (-1, 0]$ , and of course the nonlinear parts of Assumption 2.1(4)-(5) do not play a role here. The case  $\delta > 0$  will be used to prove the high-order regularity result of Theorem 2.7.

Under the previous assumptions, we consider the *turbulent Stokes couple* as:

$$(3.4) \quad \begin{aligned} (A_{\mathbb{S}}, B_{\mathbb{S}}) : \mathbb{R}_+ \times \Omega &\rightarrow \mathcal{L}(\mathbb{H}^{1+\delta, q}(\mathbb{T}^d), \mathbb{H}^{-1+\delta, q}(\mathbb{T}^d) \times \mathbb{H}^{\delta, q}(\mathbb{T}^d; \ell^2)) \\ (A_{\mathbb{S}}, B_{\mathbb{S}})u &:= (\mathbb{P}\mathcal{A}(\cdot)u, (\mathbb{P}\mathcal{B}_n(\cdot)u)_{n \geq 1}), \quad u \in \mathbb{H}^{1+\delta, q}, \end{aligned}$$

where  $\mathbb{H}^{s, q}$  is as in (2.5) with  $\mathcal{Y} = H^{s, q}$ . Note that by Assumption 3.1,

$$(3.5) \quad \begin{aligned} &\|A_{\mathbb{S}}(t)v\|_{\mathbb{H}^{-1+\delta, q}(\mathbb{T}^d)} + \|B_{\mathbb{S}}(t)v\|_{\mathbb{H}^{\delta, q}(\mathbb{T}^d; \ell^2)} \\ &\stackrel{(2.4)}{\lesssim} \max_{i,j,k} \left( \|a^{i,j}(t, \cdot) \partial_j v^k\|_{H^{\delta, q}(\mathbb{T}^d)} + \|(b_n^j(t, \cdot) \partial_j v^k)_{n \geq 1}\|_{H^{\delta, q}(\mathbb{T}^d; \ell^2)} \right) \\ &\quad + \left\| \left( [(I - \mathbb{P})[(b_n \cdot \nabla)v]]^k h_n^{j,k} \right)_{n \geq 1} \right\|_{H^{\delta, q}(\mathbb{T}^d; \ell^2)} \lesssim_{\delta, q} M \|v\|_{\mathbb{H}^{1+\delta, q}(\mathbb{T}^d)}, \end{aligned}$$

where we used [AV21c, Corollary 4.2 and Proposition 4.1(3)] if  $\delta > 0$  and  $\delta \leq 0$ , respectively. By Assumption 3.1(1), the mapping  $(A_S, B_S)$  defined in (3.4) is strongly progressively measurable.

For any stopping time  $\tau : \Omega \rightarrow [s, T]$ ,  $f \in L^0_{\mathcal{F}}(\Omega; L^1(s, \tau; \mathbb{H}^{-1+\delta, q}(\mathbb{T}^d)))$ ,  $g \in L^0_{\mathcal{F}}(\Omega; L^2(s, \tau; \mathbb{H}^{-1+\delta, q}(\mathbb{T}^d; \ell^2)))$  we say that  $u \in L^0_{\mathcal{F}}(\Omega; L^2(s, \tau; \mathbb{H}^{1+\delta, q}))$  is a *strong solution to (3.1) on  $[s, \tau] \times \Omega$*  if a.s. for all  $t \in [s, \tau]$

$$u(t) + \int_s^t A_S(r)u(r)dr = \int_s^t f(r)dr + \int_s^t (B_S(r)u(r) + g(r))dW_{\ell^2}(r),$$

where  $W_{\ell^2}$  is the  $\ell^2$ -cylindrical Brownian motion associated with  $(w^n)_{n \geq 1}$ , see [AV21a, Example 2.12]. Note that the stochastic integral is well-defined in  $\mathbb{H}^{\delta, q}(\mathbb{T}^d)$  due to (3.5), [Ond04] or [NVW15, Theorem 4.7] and  $q \geq 2$ .

The aim of this section is to prove the following.

**Theorem 3.2** (Stochastic maximal  $L^p$ -regularity for the turbulent Stokes system). *Suppose Assumption 3.1 holds. Then for each  $s \in [0, T]$ , and each progressively measurable  $f \in L^p(\Omega \times (s, T), w_{\kappa}^s; \mathbb{H}^{-1+\delta, q})$ , and  $g \in L^p(\Omega \times (s, T), w_{\kappa}^s; \mathbb{H}^{\delta, q}(\ell^2))$ , there exists a unique strong solution  $u$  to (3.1) on  $[s, T]$ . Moreover, letting*

$$J_{p, q, \kappa}(f, g) := \|f\|_{L^p(\Omega \times (s, T), w_{\kappa}^s; \mathbb{H}^{-1+\delta, q})} + \|g\|_{L^p(\Omega \times (s, T), w_{\kappa}^s; \mathbb{H}^{\delta, q}(\ell^2))},$$

the following results hold, where the constants  $C_i$  do not depend on  $(s, f, g)$ :

(1) in case  $p \in (2, \infty)$ ,  $q \in [2, \infty)$ ,  $\theta \in [0, \frac{1}{2})$ ,

$$\begin{aligned} \|u\|_{L^p(\Omega; L^p(s, T, w_{\kappa}^s; \mathbb{H}^{1+\delta, q}))} &\leq C_1 J_{p, q, \kappa}(f, g), \\ \|u\|_{L^p(\Omega; H^{\theta, p}(s, T, w_{\kappa}^s; \mathbb{H}^{1-2\theta+\delta, q}))} &\leq C_2(\theta) J_{p, q, \kappa}(f, g), \quad \theta \in [0, 1/2), \\ \|u\|_{L^p(\Omega; C([s, T]; \mathbb{B}_{q, p}^{1+\delta-2(1+\kappa)/p}))} &\leq C_3 J_{p, q, \kappa}(f, g), \\ \|u\|_{L^p(\Omega; C([s+\varepsilon, T]; \mathbb{B}_{q, p}^{1+\delta-2/p}))} &\leq C_4(\varepsilon) J_{p, q, \kappa}(f, g), \quad \varepsilon \in (s, T). \end{aligned}$$

(2) in case  $p = q = 2$  (and thus  $\kappa = 0$ ),

$$\begin{aligned} \|u\|_{L^2(\Omega; L^2(s, T; \mathbb{H}^{1+\delta, 2}))} &\leq C_5 J_{2, 2, 0}(f, g), \\ \|u\|_{L^2(\Omega; C([s, T]; \mathbb{H}^{\delta, 2}))} &\leq C_6 J_{2, 2, 0}(f, g). \end{aligned}$$

The above result can be seen as an extension of [Mik02], where  $p = q$  was considered and where the  $\mathbb{R}^d$ -case was considered. The proof of Theorem 3.2 uses similar ideas as in [AV21c, Section 5] in which we combined perturbation and localization arguments to obtain stochastic maximal regularity for parabolic problems on  $\mathbb{T}^d$ . In the present case, additional difficulties arise due the nonlocal nature of the Helmholtz projection  $\mathbb{P}$ . The  $\mathbb{R}^d$ -case can also be covered with the same method if one imposes the following condition:

*Remark 3.3* (The  $\mathbb{R}^d$ -case). Theorem 3.2 also holds for (3.1) with  $\mathbb{T}^d$  replaced by  $\mathbb{R}^d$  if the coefficients become constants as  $|x| \rightarrow \infty$ : There exists progressively measurable maps  $\widehat{a}^{i,j} : [s, T] \times \Omega \rightarrow \mathbb{R}$ ,  $\widehat{b}_n^j : [s, T] \times \Omega \rightarrow \mathbb{R}$  such that

$$\lim_{|x| \rightarrow \infty} \sup_{\omega \in \Omega} \sup_{r \in [s, T]} |a^{i,j}(r, \omega, x) - \widehat{a}^{i,j}(r, \omega)| + \|(b_n^j(r, \omega, x) - \widehat{b}_n^j(r, \omega))_{n \geq 1}\|_{\ell^2} = 0,$$

which is needed in our localization argument (see Steps 3 and 4 of the proof of Lemma 3.6). In case  $a^{i,j} = \delta^{i,j}$  and  $(b_n^j)_{n \geq 1} \in H^{\eta, \xi}(\mathbb{R}^d; \ell^2)$  for some  $\eta > 0$ ,  $\xi \in [q', \infty)$  such that  $\eta > \frac{d}{\xi}$ , then the above condition is satisfied with  $\widehat{b}_n^j \equiv 0$ . Thus the case of Kraichnan's turbulence on  $\mathbb{R}^d$  is covered, see (1.3).



Our methods can also be extended to the non-divergence case setting. For this one needs to modify the assumptions on  $a^{i,j}$  in Assumption 3.1(2). A detailed statement for related non-divergence systems can be found in [AV21c, Section 5].

Theorem 3.2 will be proved in Subsection 3.2 below. The core of the proof is the a priori estimate on small time intervals of Lemma 3.6 which will be proven in Subsection 3.3.

**3.2. Proof of Theorem 3.2.** In the rest of this section, for  $s \in \mathbb{R}$  and  $q \in (1, \infty)$ , we write  $L^q$ ,  $H^{s,q}$  and  $B_{q,p}^s$  instead of  $L^q(\mathbb{T}^d; \mathbb{R}^d)$ ,  $H^{s,q}(\mathbb{T}^d; \mathbb{R}^d)$  and  $B_{q,p}^s(\mathbb{T}^d; \mathbb{R}^d)$ .

Recall that  $\mathbb{P}$  denotes the Helmholtz projection, see (2.3). For future convenience, we recall another construction of  $\mathbb{P}$ . Let  $f \in \mathcal{D}'(\mathbb{T}^d; \mathbb{R}^d)$ . Consider the following elliptic problem on  $\mathbb{T}^d$ :

$$(3.6) \quad \begin{cases} \Delta \psi = \operatorname{div} f, \\ \langle 1, \psi \rangle = 0, \end{cases}$$

where  $\langle \cdot, \cdot \rangle$  denotes the duality of  $\mathcal{D}'(\mathbb{T}^d)$  and  $\mathcal{D}'(\mathbb{T}^d)$ . If we set  $\mathbb{Q}f := \psi$ , then the Helmholtz projection can be equivalently defined by

$$(3.7) \quad \mathbb{P}f := f - \nabla \mathbb{Q}f.$$

Indeed, by standard Fourier methods, one can check that (2.3) and (3.7) coincide. Moreover  $\nabla \mathbb{Q}$  is a projection (i.e.  $\nabla \mathbb{Q} = (\nabla \mathbb{Q})^2$ ) and  $\mathbb{Q}$  restricts to a map

$$(3.8) \quad \mathbb{Q} : H^{s,q} \rightarrow H^{s+1,q}.$$

Below we collect some lemmata which will be useful in the proof of the main results. The following concerns an operator appearing several times in the proofs.

**Lemma 3.4.** *Let  $s \in \mathbb{R}$ ,  $q \in (1, \infty)$  and  $\phi \in C^\infty(\mathbb{T}^d)$ . For any  $f \in H^{s,q}(\mathbb{T}^d; \mathbb{R}^d)$ , we set  $\mathbb{J}_\phi f := \psi$ , where  $\psi \in \mathcal{D}'(\mathbb{T}^d; \mathbb{R}^d)$  is the unique solution to the following elliptic problem on  $\mathbb{T}^d$ :*

$$(3.9) \quad \begin{cases} \Delta \psi = \nabla \phi \cdot f - \langle \nabla \phi, f \rangle, \\ \langle 1, \psi \rangle = 0. \end{cases}$$

Then  $\|\mathbb{J}_\phi f\|_{H^{s+2,q}} \lesssim_{s,q} \|f\|_{H^{s,q}}$ .

*Proof.* Let us denote by  $\Delta_N^{-1}$  the operator defined by  $(\widehat{\Delta_N^{-1} f})(0) = 0$  and  $(\widehat{\Delta_N^{-1} f})(k) = \frac{1}{|k|^2} \widehat{f}(k)$  for  $\mathbb{Z}^d \ni k \neq 0$ . By standard Fourier techniques, one can show that for each  $s \in \mathbb{R}$  and  $q \in (1, \infty)$ ,

$$(3.10) \quad \Delta_N^{-1} : \{f \in H^{s,q} : \widehat{f}(0) = 0\} \rightarrow \{f \in H^{s+2,q} : \widehat{f}(0) = 0\}.$$

The claim follows by noticing that  $\mathbb{J}_\phi f = \Delta_N^{-1}(\nabla \phi \cdot f - \langle \nabla \phi, f \rangle)$ .  $\square$

The following simple extension result is taken from [AV21c, Lemma 5.8 and Remark 5.6].

**Lemma 3.5** (Extension operator). *Let  $\alpha \in (0, N]$  and  $\zeta \in (1, \infty)$  be such that  $\alpha > \frac{d}{\zeta}$ . Let  $X$  be a Hilbert space. Then, for any  $y \in \mathbb{T}^d$  and any  $r \in (0, \frac{1}{8})$  there exists an extension operator*

$$E_{y,r}^{\mathbb{T}^d} : H^{\alpha,\zeta}(B_{\mathbb{T}^d}(y, r); X) \rightarrow H^{\alpha,\zeta}(\mathbb{T}^d; X)$$

which satisfies the following properties:

- (1)  $E_{y,r}^{\mathbb{T}^d} f|_{B_{\mathbb{T}^d}(y,r)} = f$ ,  $E_{y,r}^{\mathbb{T}^d} c \equiv c$  for any  $f \in H^{\alpha,\zeta}(B_{\mathbb{T}^d}(y,r); X)$  and  $c \in X$ ;
- (2)  $\|E_{y,r}^{\mathbb{T}^d}\|_{\mathcal{L}(H^{\alpha,\zeta}(B_{\mathbb{T}^d}(y,r); X), H^{\alpha,\zeta}(\mathbb{T}^d; X))} \leq C_r$  for some  $C_r$  independent of  $y$ ;
- (3)  $\|E_{y,r}^{\mathbb{T}^d}\|_{\mathcal{L}(C(\overline{B_{\mathbb{T}^d}(y,r)}); X), C(\mathbb{T}^d; X)} \leq C$  for some  $C$  independent of  $y, r$ .

The above lemma also holds if  $X$  is a Banach space with the UMD property. For details on the latter we refer to [HNVW16, Chapter 4].

The key step in the proof of Theorem 3.2 is the following a priori estimate for solutions to (3.1) on small time intervals.

**Lemma 3.6.** *Let Assumption 3.1 be satisfied. Then there exist  $T^*, C > 0$  depending only on the parameters  $q, p, \kappa, d, \delta, \nu, \eta, \xi, M$  such that for any  $t \in [s, T)$ ,  $\ell \in \{0, \kappa\}$ , any stopping time  $\tau : \Omega \rightarrow [t, t + T^*) \wedge T$ ,*

$$(3.11) \quad f \in L^p_{\mathcal{F}}(\llbracket t, \tau \rrbracket, w_{\ell}^t; \mathbb{H}^{-1+\delta, q}), \quad g \in L^p_{\mathcal{F}}(\llbracket t, \tau \rrbracket, w_{\ell}^t; \mathbb{H}^{\delta, q}(\ell^2))$$

and any strong solution  $u \in L^p_{\mathcal{F}}(\llbracket t, \tau \rrbracket, w_{\ell}^t; \mathbb{H}^{1+\delta, q})$  to (3.1) on  $\llbracket t, \tau \rrbracket$  one has

$$\|u\|_{L^p(\llbracket t, \tau \rrbracket, w_{\ell}^t; \mathbb{H}^{1+\delta, q})} \leq C \|f\|_{L^p(\llbracket t, \tau \rrbracket, w_{\ell}^t; \mathbb{H}^{-1+\delta, q})} + C \|g\|_{L^p(\llbracket t, \tau \rrbracket, w_{\ell}^t; \mathbb{H}^{\delta, q}(\ell^2))}.$$

We first show how Lemma 3.6 implies Theorem 3.2.

*Proof of Theorem 3.2.* We refer to [AV21c, Definition 2.3] for the stochastic maximal  $L^p$ -regularity notation  $\mathcal{SMR}_{p,\kappa}(s, T)$  and  $\mathcal{SMR}_{p,\kappa}^{\bullet}(s, T)$ . Here we are using  $X_j = \mathbb{H}^{-1+2j+\delta, q}(\mathbb{T}^d)$ . Also note that  $X_{\kappa,p}^{\text{Tr}} := (X_0, X_1)_{1-\frac{1+\kappa}{p}, p} = \mathbb{B}_{q,p}^{1+\delta-2\frac{1+\kappa}{p}}$  and  $X_{\theta} := [X_0, X_1]_{\theta} = \mathbb{H}^{1+\delta-2\theta, q}$  for  $\theta \in (0, 1)$ . Since  $\Delta$  and  $\mathbb{P}$  commute on  $\mathbb{H}^{1+\delta, q}$  (see (2.3)), the Stokes operator  $\mathbb{S} := -\Delta : \mathbb{H}^{s+2, q} \subseteq \mathbb{H}^{s, q} \rightarrow \mathbb{H}^{s, q}$ , is well-defined, and by the periodic version of [HNVW17, Theorem 10.2.25]  $1 + \mathbb{S}$  has a bounded  $H^{\infty}$ -calculus of angle zero. Therefore, by [AV20, Theorem 7.16] we have

$$(3.12) \quad \mathbb{S} \in \mathcal{SMR}_{p,\kappa}^{\bullet}(s, T), \quad \text{for all } 0 \leq s < T < \infty$$

with constant only depending on  $q, \kappa, p, d$  and  $T$ .

The assertions of Theorem 3.2 follow if we can prove that  $(A_{\mathbb{S}}, B_{\mathbb{S}}) \in \mathcal{SMR}_{p,\ell}^{\bullet}(t, t + T^*)$  (see [AV21b, Proposition 3.9]). By [AV21c, Proposition 3.1], it is enough to show that  $(A_{\mathbb{S}}, B_{\mathbb{S}}) \in \mathcal{SMR}_{p,\ell}(t, t + T^*)$  for all  $\ell \in \{0, \kappa\}$ ,  $t \in [0, T - T^*)$  where  $T^*$  is as in Lemma 3.6. To prove the latter, we use the method of continuity (see [AV21b, Proposition 3.13]).

For any  $\lambda \in [0, 1]$  and  $v \in \mathbb{H}^{1+\delta, q}$ , we set

$$\begin{aligned} A_{\mathbb{S}, \lambda} v &:= (1 - \lambda) \mathbb{S} v + \lambda A_{\mathbb{S}} v \\ &= -\mathbb{P}(\text{div}(a_{\lambda}(\cdot) \cdot \nabla v)) - \left( \sum_{n \geq 1} \left[ (I - \mathbb{P})((b_n(t) \cdot \nabla) v) \right] \cdot \lambda h_n^{\cdot k} \right)_{k=1}^d, \\ B_{\mathbb{S}, \lambda} v &:= \lambda B_{\mathbb{S}} v = \left( \mathbb{P}(\lambda (b_n \cdot \nabla) v) \right)_{n \geq 1} \end{aligned}$$

where  $a_{\lambda}^{i,j} = \lambda \delta^{i,j} + (1 - \lambda) a^{i,j}$  for all  $i, j \in \{1, \dots, d\}$ .

Note that  $(A_{\mathbb{S}, 0}, B_{\mathbb{S}, 0}) = (\mathbb{S}, 0)$ , and  $(A_{\mathbb{S}, \lambda}, B_{\mathbb{S}, \lambda})$  satisfies the Assumption 3.1 uniformly w.r.t.  $\lambda \in [0, 1]$ . Thus Lemma 3.6 ensures that the a priori estimate (3.6) holds for strong solutions to (3.1) on  $\llbracket t, \tau \rrbracket$  with  $C$  independent of  $\lambda \in [0, 1]$ . The claim of this step follows from the method of continuity in [AV21b, Proposition 3.13] and (3.12).  $\square$

**3.3. Proof of Lemma 3.6.** We will next prove the key estimate. In principle we use the perturbation method of our recent work [AV21c, Lemma 5.4], but due to the non-local behavior of the Helmholtz projection, there are several complications in the proof.

*Proof of Lemma 3.6 in case  $h = 0$ .* The proof will be divided into several steps. Moreover, for exposition convenience, we set  $s = 0$  and in Steps 1-6 we only consider the case  $\delta \leq 0$  and in Step 7 we discuss the modifications needed for the case  $\delta > 0$ .

We will use the method where we freeze the coefficients and use comparison on a small ball. For any  $y \in \mathbb{T}^d$ ,  $r \in (0, \frac{1}{8})$ ,  $B(y, r) := B_{\mathbb{T}^d}(y, r)$ , and for  $v \in \mathbb{H}^{2-\delta, q}$ , we set

$$(3.13) \quad \begin{aligned} \mathcal{A}_y(t)v &:= -\operatorname{div}(a(t, y) \cdot \nabla v), & \mathcal{B}_y(t) &:= ((b_n(t, y) \cdot \nabla)v)_{n \geq 1}, \\ \mathcal{A}_{y,r}^E(t) &:= -\operatorname{div}(a_{y,r}^E(t, \cdot) \cdot \nabla v), & \mathcal{B}_{y,r,\ell}^E(t) &:= ((b_{n,y,r}^E(t, \cdot) \cdot \nabla)v)_{n \geq 1}, \end{aligned}$$

where

$$a_{y,r}^E := \left( \mathbb{E}_{y,r}^{\mathbb{T}^d}(a^{i,j}(t, \cdot)) \right)_{i,j=1}^d, \quad b_{n,y,r}^E := \left( \mathbb{E}_{y,r}^{\mathbb{T}^d}(b_n^j(t, \cdot)) \right)_{j=1}^d, \quad n \geq 1,$$

and  $\mathbb{E}_{y,r}^{\mathbb{T}^d}$  is the extension operator of Lemma 3.5. The operators  $\mathcal{A}_y$  and  $\mathcal{B}_y$  have a ‘‘frozen coefficient at  $y \in \mathbb{T}^d$ ’’ and  $\mathcal{A}_{y,r}^E$ ,  $\mathcal{B}_{y,r,\ell}^E$  are the operators whose coefficients are the extensions of  $a^{i,j}|_{B(y,r)}$ ,  $b_n^j|_{B(y,r)}$ .

Let  $t \in [s, T)$ ,  $\ell \in \{0, \kappa\}$  and let  $\tau : \Omega \rightarrow [t, t^*]$  be a stopping time with  $t^* := T \wedge (t + T^*)$  where  $T^* > 0$  will be chosen in Step 6. We use same notation as in [AV21a, Subsection 3.3] in case  $(\mathbb{P}\mathcal{A}, \mathbb{P}\mathcal{B}) := (\mathbb{P}\mathcal{A}, (\mathbb{P}\mathcal{B}_n)_{n \geq 1}) \in \mathcal{SMR}_{p,\ell}(t, T)$  with  $X_j = \mathbb{H}^{-1+2j+\delta, q}$  where  $j \in \{0, 1\}$ . The solution operator  $\mathcal{R}_{t,(\mathbb{P}\mathcal{A}, \mathbb{P}\mathcal{B})}$  associated to the couple  $(\mathbb{P}\mathcal{A}, \mathbb{P}\mathcal{B})$  (see (3.2)) maps

$$L_{\mathcal{F}}^p([t, T], w_\ell^t; \mathbb{H}^{-1+\delta, q}) \times L_{\mathcal{F}}^p([t, T], w_\ell^t; \mathbb{H}^{\delta, q}(\ell^2)) \rightarrow L_{\mathcal{F}}^p([t, T], w_\ell^t; \mathbb{H}^{1+\delta, q}),$$

where we set  $\mathcal{R}_{t,(\mathbb{P}\mathcal{A}, \mathbb{P}\mathcal{B})}(f, g) := u$  with  $u$  is the unique strong to (3.1) on  $[t, \tau]$ , and  $f$  and  $g$  are as in (3.11). Moreover, we set  $K_{(\mathbb{P}\mathcal{A}, \mathbb{P}\mathcal{B})}^{p,\ell}(t, T) = \|\mathcal{R}_{t,(\mathbb{P}\mathcal{A}, \mathbb{P}\mathcal{B})}(\cdot, \cdot)\|$ . For  $f$  and  $g$  as in (3.11), and for  $\tau$  as above, we set

$$(3.14) \quad N_{f,g}(t, \tau) := \|f\|_{L^p([t, \tau], w_\ell^t; \mathbb{H}^{-1+\delta, q})} + \|g\|_{L^p([t, \tau], w_\ell^t; \mathbb{H}^{\delta, q}(\ell^2))}.$$

To abbreviate the dependencies let  $\mathcal{S} = \{q, p, \kappa, d, \delta, \nu, \eta, \xi, M\}$ , and whenever we wish to emphasise the dependencies we write  $C(\mathcal{S})$  instead of  $C$ .

*Step 1:* There exists  $C_1(\mathcal{S}) > 0$  such that for each  $y \in \mathbb{T}^d$ , one has  $(\mathbb{P}\mathcal{A}_y, \mathbb{P}\mathcal{B}_y) \in \mathcal{SMR}_{p,\ell}(t, T)$  and  $K_{(\mathbb{P}\mathcal{A}_y, \mathbb{P}\mathcal{B}_y)}^{p,\ell}(t, T) \leq C_1$ .

Due to (3.6)-(3.7), one can check that

$$\mathbb{P}\mathcal{A}_y v = \mathcal{A}_y \mathbb{P}v, \quad \mathbb{P}\mathcal{B}_y v = \mathcal{B}_y \mathbb{P}v, \quad \text{a.e. on } [t, T] \text{ for all } v \in H^{1+\delta, q}.$$

Therefore the result is immediate from [AV21c, Lemma 5.4].

*Step 2:* There exists  $\chi(\mathcal{S}) > 0$  for which the following holds:

If  $y \in \mathbb{T}^d$  and  $r \in (0, \frac{1}{8})$  are such that a.s. for all  $t \in (0, T)$ ,  $i, j \in \{1, \dots, d\}$ ,

$$(3.15) \quad \|a^{i,j}(t, \cdot) - a^{i,j}(t, y)\|_{L^\infty(B(y,r))} + \|(b_n^j(t, \cdot) - b_n^j(t, y))_{n \geq 1}\|_{L^\infty(B(y,r); \ell^2)} \leq \chi,$$

then  $(\mathbb{P}\mathcal{A}_{y,r}^E, \mathbb{P}\mathcal{B}_{y,r}^E) \in \mathcal{SMR}_{p,\ell}(t, T)$  and  $K_{(\mathbb{P}\mathcal{A}_{y,r}^E, \mathbb{P}\mathcal{B}_{y,r}^E)}^{p,\ell}(t, T) \leq C_2(\mathcal{S})$ .

To prove this we apply the perturbation result of [AV21c, Theorem 3.2]. To this end, we write

$$(3.16) \quad \mathbb{P}\mathcal{A}_{y,r}^E = \mathbb{P}\mathcal{A}_y + \mathbb{P}(\mathcal{A}_{y,r}^E - \mathcal{A}_y), \quad \mathbb{P}\mathcal{B}_{y,r}^E = \mathbb{P}\mathcal{B}_y + \mathbb{P}(\mathcal{B}_{y,r}^E - \mathcal{B}_y).$$

By (2.4), for each  $v \in \mathbb{H}^{1+\delta,q}$ ,

$$\|\mathbb{P}(\mathcal{A}_{y,r}^E - \mathcal{A}_y)v\|_{\mathbb{H}^{-1+\delta,q}} \lesssim \sum_{i,j=1}^d \left\| (a^{i,j}(t,y) - \mathbb{E}_{y,r}^{\mathbb{T}^d}(a^{i,j}(t,\cdot)))\partial_j v \right\|_{H^{\delta,q}}.$$

Next we estimate each term separately:

$$\begin{aligned} & \left\| (a^{i,j}(t,y) - \mathbb{E}_{y,r}^{\mathbb{T}^d}(a^{i,j}(t,\cdot)))\partial_j v \right\|_{H^{\delta,q}} \\ & \stackrel{(i)}{=} \left\| \mathbb{E}_{y,r}^{\mathbb{T}^d}(a^{i,j}(t,y) - a^{i,j}(t,\cdot))\partial_j v \right\|_{H^{\delta,q}} \\ & \stackrel{(ii)}{\lesssim} \left( \left\| \mathbb{E}_{y,r}^{\mathbb{T}^d}(a^{i,j}(t,y) - a^{i,j}(t,\cdot)) \right\|_{L^\infty} \|\partial_j v\|_{H^{\delta,q}} \right. \\ & \quad \left. + \left\| \mathbb{E}_{y,r}^{\mathbb{T}^d}(a^{i,j}(t,y) - a^{i,j}(t,\cdot)) \right\|_{H^{\eta,\xi}} \|\partial_j v\|_{H^{\delta-\varepsilon,q}} \right) \\ & \stackrel{(iii)}{\leq} \chi \|v\|_{H^{1+\delta,q}} + C_r M \|v\|_{H^{1+\delta-\varepsilon,q}} \\ & \stackrel{(iv)}{\leq} 2\chi \|v\|_{H^{1+\delta,q}} + C_{r,M,\chi} \|v\|_{H^{-1+\delta,q}}, \end{aligned}$$

where in (i) we used Lemma 3.5(1), in (ii) [AV21c, Proposition 4.1(3)] for some  $\varepsilon > 0$  and in (iii) Lemma 3.5(2)-(3) and (3.15). Finally, in (iv) we used a standard interpolation inequality.

For the  $\mathcal{B}$ -term, recall that  $\xi \geq q'$  and  $\eta > -\delta$ . Thus employing similar arguments, by Assumption 2.1(2) and [AV21c, Proposition 4.1(3)] with  $H = \ell^2$ , one obtains

$$\|\mathbb{P}(\mathcal{B}_{y,r}^E - \mathcal{B}_y)v\|_{\mathbb{H}^{\delta,q}(\ell^2)} \leq 2\chi \|v\|_{H^{1+\delta,q}} + C_{M,r,\chi} \|v\|_{H^{-1+\delta,q}}.$$

The claim of Step 2 follows from Step 1, [AV21c, Theorem 3.2] and the above estimates.

*Step 3:* Let  $\chi$  be as in step 2. There exist an integer  $\Lambda := \Lambda(\mathcal{S}) \geq 1$ ,  $(y_\lambda)_{\lambda=1}^\Lambda \subseteq \mathbb{T}^d$ ,  $(r_\lambda)_{\lambda=1}^\Lambda \subseteq (0, \frac{1}{8})$  depending only on  $\mathcal{S}$  such that  $\mathbb{T}^d \subseteq \cup_{\lambda=1}^\Lambda B_\lambda$ , where  $B_\lambda := B_{\mathbb{T}^d}(y_\lambda, r_\lambda)$ , and a.s. for all  $t \in (0, T)$ ,  $i, j \in \{1, \dots, d\}$ ,

$$\|a^{i,j}(t, y_\lambda) - a^{i,j}(t, \cdot)\|_{L^\infty(B_\lambda)} + \|(b_n^j(t, y_\lambda) - b_n^j(t, \cdot))_{n \geq 1}\|_{L^\infty(B_\lambda; \ell^2)} \leq \chi.$$

In particular, for all  $\lambda \in \{1, \dots, \Lambda\}$ ,

$$(\mathbb{P}\mathcal{A}_\lambda^E, \mathbb{P}\mathcal{B}_\lambda^E) := (\mathbb{P}\mathcal{A}_{x_\lambda, r_\lambda}^E, \mathbb{P}\mathcal{B}_{x_\lambda, r_\lambda}^E) \in \mathcal{SMR}_{p,\ell}(t, T)$$

with  $K_{(\mathbb{P}\mathcal{A}_\lambda^E, \mathbb{P}\mathcal{B}_\lambda^E)}^{p,\ell}(t, T) \leq C_3(\mathcal{S})$ .

The last claim follows from the first one and Step 2. Let  $\alpha := \eta - \frac{d}{\xi} > 0$  due Assumption 3.1(1). By Sobolev embeddings,  $H^{\eta,\xi}(H) \hookrightarrow C^\alpha(H)$  where  $H \in \{\mathbb{R}; \ell^2\}$ . We denote by  $\tilde{C}$  the embedding constant. Thus for any  $y \in \mathbb{T}^d$ ,

$$\begin{aligned} & \|a^{i,j}(t, y) - a^{i,j}(t, \cdot)\|_{L^\infty(B(y,r))} + \|b^j(t, y) - b^j(t, \cdot)\|_{L^\infty(B(y,r); \ell^2)} \\ & \leq ([a^{i,j}(t, \cdot)]_{C^\alpha(B(y,r))} + [b^j(t, \cdot)]_{C^\alpha(B(y,r); \ell^2)}) r^\alpha \leq \tilde{C} M r^\alpha \leq \chi, \end{aligned}$$

where the last inequality follows by choosing  $r := \min\{(\frac{\chi}{CM})^{1/\alpha}, \frac{1}{8}\}$ . Since  $\chi = \chi(\mathcal{S})$ , it follows that  $r = r(\mathcal{S})$ . To conclude, it remains to note that  $\mathbb{T}^d$  can be covered by finitely many balls of the form  $B(y, r)$ .

*Step 4:* Let  $(\pi_\lambda)_{\lambda=1}^\Lambda$  be a smooth partition of the unity subordinate to the covering  $(B_\lambda)_{\lambda=1}^\Lambda$  (see Step 3). Let  $\mathcal{R}_\lambda := \mathcal{R}_{t, (\mathbb{P}\mathcal{A}_\lambda^E, \mathbb{P}\mathcal{B}_\lambda^E)}$  be the solution operator associated to

$(\mathbb{P}\mathcal{A}_\lambda^E, \mathbb{P}\mathcal{B}_\lambda^E) \in \mathcal{SMR}_{p,\ell}(t, T)$ . Recall that  $f, g$  are as in (3.11),  $\tau$  is a stopping time with values in  $[t, (t + T^*) \wedge T]$  and  $u \in L^p_{\mathcal{D}}(\llbracket t, \tau \rrbracket, w_\ell^t; \mathbb{H}^{1+\delta, q})$  is a strong solution to (3.1) on  $\llbracket t, \tau \rrbracket$ . Then for any  $\lambda \in \{1, \dots, \Lambda\}$  the following holds

$$\mathbb{P}(\pi_\lambda u) = \mathcal{R}_\lambda(0, F_\lambda u, G_\lambda u) + \mathcal{R}_\lambda(0, \mathbb{P}f_\lambda, \mathbb{P}g_\lambda), \quad \text{a.e. on } \llbracket t, \tau \rrbracket,$$

where  $\mathbb{J}_\lambda := \mathbb{J}_{\pi_\lambda}$  (see Lemma 3.4),  $g_\lambda := (g_{\lambda, n})_{n \geq 1} := (g_n \pi_\lambda)_{n \geq 1}$ ,  $f_\lambda := \pi_\lambda f$  and

$$\begin{aligned} F_\lambda u &:= -\mathbb{P}\mathcal{A}_\lambda^E \nabla \mathbb{J}_\lambda u + \mathbb{P}[\mathcal{A}, \pi_\lambda]u - \mathbb{P}[(\nabla \pi_\lambda) \mathbb{Q}(\mathcal{A}u)], \\ (3.17) \quad G_{\lambda, n} u &:= \mathbb{P}\mathcal{B}_{\lambda, n}^E \nabla \mathbb{J}_\lambda u + \mathbb{P}[\pi_\lambda, \mathcal{B}_n]u + \mathbb{P}[(\nabla \pi_\lambda) \mathbb{Q}(\mathcal{B}_n u)], \\ G_\lambda u &:= (G_{\lambda, n} u)_{n \geq 1}, \end{aligned}$$

with  $\mathbb{Q}$  as in (3.7), and  $[\cdot, \cdot]$  denotes the commutator.

To begin, set  $u_\lambda := \pi_\lambda u$ . Note that  $\mathbb{P}(\mathcal{A}u) = \mathcal{A}u - \nabla[\mathbb{Q}(\mathcal{A}u)]$  and  $\mathbb{P}(\mathcal{B}_n u) = \mathcal{B}_n u - \nabla[\mathbb{Q}(\mathcal{B}_n u)]$ . Using these identities and multiplying (3.1) by  $\pi_\lambda$ , one obtains

$$\begin{aligned} (3.18) \quad du_\lambda + \mathcal{A}u_\lambda dr &= \left( [\mathcal{A}, \pi_\lambda]u + \nabla(\pi_\lambda \mathbb{Q}(\mathcal{A}u)) - (\nabla \pi_\lambda) \mathbb{Q}(\mathcal{A}u) + f_\lambda \right) dr \\ &+ \sum_{n \geq 1} \left( \mathcal{B}_n u_\lambda + [\pi_\lambda, \mathcal{B}_n]u - \nabla(\pi_\lambda \mathbb{Q}_n) + (\nabla \pi_\lambda) \mathbb{Q}(\mathcal{B}_n u) + g_{\lambda, n} \right) dw_r^n, \end{aligned}$$

on  $\mathbb{T}^d$ . Since  $\text{supp}(u_\lambda) \subseteq B_\lambda$ , for each  $\lambda \in \{1, \dots, \Lambda\}$  and  $n \geq 1$  one has (see (3.13))

$$\mathcal{A}u_\lambda = \mathcal{A}_{x_\lambda, r_\lambda}^E u_\lambda = \mathcal{A}_\lambda^E u_\lambda, \quad \mathcal{B}_n u_\lambda = \mathcal{B}_{n, x_\lambda, r_\lambda}^E u_\lambda = \mathcal{B}_{\lambda, n}^E u_\lambda.$$

Moreover, note that the Helmholtz decomposition gives

$$u_\lambda = \pi_\lambda u = \nabla \mathbb{Q}(\pi_\lambda u) + \mathbb{P}(\pi_\lambda u) \stackrel{(i)}{=} \nabla \mathbb{J}_\lambda u + \mathbb{P}u_\lambda.$$

Here in (i) we used  $\text{div } u = 0$  in  $\mathcal{D}'(\mathbb{T}^d)$ , (3.9) and

$$\text{div } u_\lambda = \text{div } u_\lambda - \langle 1, \text{div } u_\lambda \rangle = \nabla \pi_\lambda \cdot u - \langle 1, \nabla \pi_\lambda \cdot u \rangle.$$

Thus  $u_\lambda = \nabla \mathbb{J}_\lambda u + v_\lambda$  with  $v_\lambda := \mathbb{P}u_\lambda$ . Applying the operator  $\mathbb{P}$  to (3.18), we get

$$\begin{aligned} dv_\lambda + \mathbb{P}\mathcal{A}_\lambda^E v_\lambda dr &= \left( -\mathbb{P}\mathcal{A}_\lambda^E \nabla \mathbb{J}_\lambda u + \mathbb{P}[\mathcal{A}, \pi_\lambda]u - \mathbb{P}[(\nabla \pi_\lambda) \mathbb{Q}(\mathcal{A}u)] + \mathbb{P}f_\lambda \right) dr \\ &+ \sum_{n \geq 1} \left( \mathbb{P}\mathcal{B}_{\lambda, n}^E v_\lambda + \mathbb{P}\mathcal{B}_{\lambda, n}^E \nabla \mathbb{J}_\lambda u + \mathbb{P}[\pi_\lambda, \mathcal{B}_n]u + \mathbb{P}[(\nabla \pi_\lambda) \mathbb{Q}(\mathcal{B}_n u)] + \mathbb{P}g_{\lambda, n} \right) dw_r^n, \\ &= \left( F_\lambda u + \mathbb{P}f_\lambda \right) dr + \sum_{n \geq 1} \left( \mathbb{P}\mathcal{B}_{\lambda, n}^E v_\lambda + G_{\lambda, n} u + \mathbb{P}g_{\lambda, n} \right) dw_r^n \end{aligned}$$

with initial value  $v_\lambda(t) = 0$ . Recall that by Step 3, one has  $(\mathbb{P}\mathcal{A}_\lambda^E, (\mathbb{P}\mathcal{B}_{\lambda, n}^E)_{n \geq 1}) \in \mathcal{SMR}_{p,\ell}(t, T)$ . Thus the claim follows from [AV21a, Proposition 3.12] and the previous displayed formula.

*Step 5:* There exists  $\varepsilon(\mathcal{S}) \in (0, 1)$ ,  $C_5(\mathcal{S}) > 0$  such that for each  $\lambda \in \{1, \dots, \Lambda\}$ ,  $v \in \mathbb{H}^{1+\delta, q}$  and  $t \in [0, T)$ ,

$$\|F_\lambda v\|_{\mathbb{H}^{-1+\delta, q}} + \|(G_{\lambda, n} v)_{n \geq 1}\|_{\mathbb{H}^{\delta, q}(\ell^2)} \leq C_5 \|v\|_{\mathbb{H}^{1+\delta-\varepsilon, q}}, \quad \text{a.e. on } \llbracket t, T \rrbracket.$$

First consider  $F_\lambda$ . Recall that  $\Lambda = \Lambda(\mathcal{S}) < \infty$  by Step 3. Thus it is enough to prove suitable estimates for  $F_\lambda$  where  $\lambda \in \{1, \dots, \Lambda\}$  is fixed. Let us write  $F_\lambda := F_{\lambda, \mathbb{J}} + F_{\lambda, \mathcal{A}} + F_{\lambda, \mathbb{Q}}$ , where

$$F_{\lambda, \mathbb{J}} v := -\mathbb{P}(\mathcal{A} \nabla \mathbb{J}_\lambda v), \quad F_{\lambda, \mathcal{A}} v := \mathbb{P}[\mathcal{A}, \pi_\lambda]v, \quad F_{\lambda, \mathbb{Q}} v := -\mathbb{P}[(\nabla \pi_\lambda) \mathbb{Q}(\mathcal{A}v)].$$

Then by the pointwise multiplication result of [AV21c, Proposition 4.1(3)] and  $\eta > -\delta$  and  $\xi \in [q', \infty)$  (see Assumption 3.1(2)),

$$\begin{aligned} \|\mathbb{F}_{\lambda, \mathbb{J}} v\|_{\mathbb{H}^{-1+\delta, q}} &\lesssim_{\delta, q} \max_{i, j} \|a^{i, j}(t, \cdot) \partial_j \nabla \mathbb{J}_\lambda v\|_{H^{\delta, q}} \\ &\lesssim_{\delta, q} M \max_j \|\partial_j \nabla \mathbb{J}_\lambda v\|_{H^{\delta, q}} \lesssim_{\delta, q} M \|v\|_{H^{\delta, q}}, \end{aligned}$$

where in the last estimate we applied Lemma 3.4. To estimate  $\mathbb{F}_{\lambda, \mathcal{A}}$ , note that

$$[\pi_\lambda, \mathcal{A}]v = \sum_{i, j=1}^d \left[ \partial_i (a^{i, j} (\partial_j \pi_\lambda) v) + (\partial_j v) a^{i, j} (\partial_i \pi_\lambda) \right], \quad \text{in } \mathcal{D}'(\mathbb{T}^d).$$

By [AV21c, Proposition 4.1(3)] we obtain

$$\|\mathbb{P}[\partial_i (a^{i, j} (\partial_j \pi_\lambda) v)]\|_{\mathbb{H}^{-1+\delta, q}} \lesssim_{\delta, q} \|a^{i, j} (\partial_j \pi_\lambda) v\|_{H^{\delta, q}} \lesssim_{\delta, q} M \|v\|_{H^{\delta, q}} \leq CC_{a, b} \|v\|_{H^{\delta, q}},$$

Letting  $\varepsilon \in (0, 1)$  be such that  $\eta \geq -\delta + \varepsilon$ , in a similar way we obtain

$$\begin{aligned} (3.19) \quad \|\mathbb{P}(\partial_j v a^{i, j} \partial_i \pi_\lambda)\|_{\mathbb{H}^{-1+\delta, q}} &\lesssim_{\delta, q} \|a^{i, j} \partial_j v\|_{H^{-1+\delta, q}} \\ &\lesssim_{\delta, q} \|a^{i, j} \partial_j v\|_{H^{\delta-\varepsilon, q}} \\ &\lesssim_{\delta, q} M \|\partial_j v\|_{H^{\delta-\varepsilon, q}} \lesssim_{\delta, q} M \|v\|_{H^{1+\delta-\varepsilon, q}}, \end{aligned}$$

from which we obtain the required bound for  $\mathbb{F}_{\lambda, \mathcal{A}}$ . By (3.8),  $\mathbb{F}_{\lambda, \mathbb{Q}}$  can be estimated as follows:

$$\|\mathbb{F}_{\lambda, \mathbb{Q}} v\|_{\mathbb{H}^{-1+\delta, q}} \lesssim \|\mathbb{Q}(\mathcal{A}v)\|_{H^{-1+\delta, q}} \lesssim \|a^{i, j} \partial_j v\|_{H^{-1+\delta, q}} \lesssim \|v\|_{H^{1+\delta-\varepsilon, q}},$$

where the last estimate can be proved similarly as in (3.19). Combining the previous estimates one obtains the claim for  $\mathbb{F}_\lambda$ .

By (3.17) we have  $\mathbb{G}_{\lambda, n} := \mathbb{G}_{\lambda, \mathbb{Q}, n} + \mathbb{G}_{\lambda, \mathbb{J}, n} + \mathbb{G}_{\lambda, \mathcal{B}, n}$  where

$$\mathbb{G}_{\lambda, \mathbb{Q}, n} v := \mathbb{P}[(\nabla \pi_\lambda) \mathbb{Q}(\mathcal{B}_n v)], \quad \mathbb{G}_{\lambda, \mathbb{J}, n} v := \mathbb{P} \mathcal{B}_n \nabla \mathbb{J}_\lambda v, \quad \mathbb{G}_{\lambda, \mathcal{B}, n} v := \mathbb{P}[(\partial_j \pi_\lambda) b_n^j v]$$

where  $v \in \mathbb{H}^{1+\delta, q}$  and we used that  $[\pi_\lambda, \mathcal{B}_n]v = (\partial_j \pi_\lambda) b_n^j v$ . Now the estimate for  $\mathbb{G}_\lambda$  can be proved in a similar way as we did for  $\mathbb{F}_\lambda$ .

*Step 6: Conclusion.* Let  $(u, f, g, \tau, t)$  be as at the beginning of the proof. By Step 3 we know that  $(\mathbb{P} \mathcal{A}_\lambda^E, \mathbb{P} \mathcal{B}_\lambda^E) \in \mathcal{SMR}_{p, \kappa}(t, T)$  for all  $\lambda \in \{1, \dots, \Lambda\}$ . Combining the latter with Steps 4-5, one can see that there exist  $\varepsilon(\mathcal{S}) \in (0, 1)$ ,  $C_6(\mathcal{S}) > 0$  such that for all  $\lambda \in \{1, \dots, \Lambda\}$ ,

$$(3.20) \quad \begin{aligned} \|\mathbb{P}(\pi_\lambda u)\|_{L^p(\llbracket t, \tau \rrbracket, w_\ell^t; \mathbb{H}^{1+\delta, q})} &\leq C_6 N_{\mathbb{F}_\lambda u, \mathbb{G}_\lambda u}(t, \tau) + C_6 N_{f, g}(t, \tau) \\ &\leq C_5 C_6 \|u\|_{L^p(\llbracket t, \tau \rrbracket, w_\ell^t; H^{1+\delta-\varepsilon, q})} + C_6 N_{f, g}(t, \tau) \end{aligned}$$

where  $N_{f, g}(t, \tau)$  is as in (3.14). As in Step 5, since  $\operatorname{div} u = 0$  in  $\mathcal{D}'(\mathbb{T}^d)$  a.e. on  $\llbracket t, \tau \rrbracket$ , one has  $\pi_\lambda u = \mathbb{P}(\pi_\lambda u) + \nabla \mathbb{J}_\lambda u$ . Since  $(\pi_\lambda)_{\lambda=1}^\Lambda$  is a partition of unity, we can write  $u = \sum_{\lambda=1}^\Lambda u_\lambda$ , and hence the previous considerations yield,

$$(3.21) \quad \begin{aligned} &\|u\|_{L^p(\llbracket t, \tau \rrbracket, w_\ell^t; H^{1+\delta, q})} \\ &\leq \sum_{\lambda=1}^\Lambda \left( \|\mathbb{P}(\pi_\lambda u)\|_{L^p(\llbracket t, \tau \rrbracket, w_\ell^t; H^{1+\delta, q})} + \|\nabla \mathbb{J}_\lambda u\|_{L^p(\llbracket t, \tau \rrbracket, w_\ell^t; H^{1+\delta, q})} \right) \\ &\stackrel{(i)}{\leq} \Lambda C_6 N_{f, g}(t, \tau) + \Lambda C_5 C_6 \|u\|_{L^p(\llbracket t, \tau \rrbracket, w_\ell^t; H^{1+\delta-\varepsilon, q})} + \Lambda C_7 \|u\|_{L^p(\llbracket t, \tau \rrbracket, w_\ell^t; H^{\delta, q})} \\ &\stackrel{(ii)}{\leq} \tilde{C}_6 N_{f, g}(t, \tau) + \frac{1}{4} \|u\|_{L^p(\llbracket t, \tau \rrbracket, w_\ell^t; H^{1+\delta, q})} + \tilde{C}_6 \|u\|_{L^p(\llbracket t, \tau \rrbracket, w_\ell^t; H^{-1+\delta, q})}, \end{aligned}$$

where in (i) we used (3.20) and Lemma 3.4 and in (ii) a standard interpolation inequality. Since  $u$  is a strong solution to (3.1) on  $[[t, \tau]]$ , by (3.5) and [AV21a, Lemma 3.14] there exists  $(c_s(\mathcal{S}))_{s>0}$  that  $\lim_{s \downarrow 0} c_s = 0$  and

$$(3.22) \quad \|u\|_{L^p([t, \tau], w_\ell^t; H^{-1+\delta, q})} \leq c_{T^*} \|u\|_{L^p([t, \tau], w_\ell^t; H^{1+\delta, q})} + c_{T^*} N_{f, g}(t, \tau)$$

where we have used that  $\tau - t \leq T^*$ .

Combining (3.21)-(3.22), one obtains

$$\|u\|_{L^p([t, \tau], w_\ell^t; H^{1+\delta, q})} \leq \tilde{C}_6 N_{f, g}(t, \tau) + \left(\frac{1}{4} + c_{T^*} \tilde{C}_6\right) \|u\|_{L^p([t, \tau], w_\ell^t; H^{1+\delta, q})},$$

By choosing  $T^*(\mathcal{S}) > 0$  so that  $c_{T^*} \leq 1/(4\tilde{C}_6)$ , the previous formula and (2.5) imply the claimed a priori estimate of Lemma 3.6.

*Step 7: The case  $\delta > 0$ .* To prove Lemma 3.6 for  $\delta > 0$ , one can argue as in Steps 1-6. The only changes appear in Steps 2 and 5, where instead of [AV21c, Proposition 4.1(3)] we use [AV21c, Proposition 4.1(1)].  $\square$

*Proof of Lemma 3.6 for general  $h$ .* Let us divide the proof of this step into three cases. In each case we use the previously obtained estimate in the case  $h = 0$ , and view the non-zero  $h$  as a lower order perturbation. Below, for all  $v \in \mathbb{H}^{1+\delta, q}$ , we set

$$\tilde{f}_b(t)v := \left( \sum_{n \geq 1} \left[ (I - \mathbb{P})((b_n(t) \cdot \nabla)v) \right] \cdot h_n^{j, k} \right)_{k=1}^d.$$

*Case  $\delta \leq 0$ .* Pick  $\varepsilon \in (0, 1)$  such that  $\eta > -\delta + \varepsilon$ . Let  $M$  be as in Assumption 3.1(2). Note that for each  $v \in \mathbb{H}^{1+\delta, q}(\mathbb{T}^d)$ ,

$$\begin{aligned} \|\mathbb{P}[\tilde{f}_b v]\|_{\mathbb{H}^{-1+\delta, q}} &\lesssim \max_k \left\| \sum_{n \geq 1} \left[ (I - \mathbb{P})[(b_n \cdot \nabla)v] \right] \cdot h_n^{j, k} \right\|_{\mathbb{H}^{\delta-\varepsilon, q}} \\ &\stackrel{(i)}{\lesssim} \max_{j, k} \| (h_n^{j, k})_{n \geq 1} \|_{H^{\eta, \xi}(\ell^2)} \| (b_n^j \partial_j v)_{n \geq 1} \|_{H^{\delta-\varepsilon, q}} \\ &\stackrel{(ii)}{\lesssim} M \max_j \| (b_n^j)_{n \geq 1} \|_{H^{\eta, \xi}(\ell^2)} \| \partial_j v \|_{H^{\delta-\varepsilon, q}} \lesssim M^2 \|v\|_{\mathbb{H}^{1+\delta-\varepsilon, q}}, \end{aligned}$$

where the implicit constants depend only on  $s, q, d$ . Moreover, in (i) – (ii) we used that  $\eta > -\delta$ ,  $\xi \geq q'$  and the pointwise multiplication result of [AV21c, Proposition 4.1(3)] and the text below it.

By Young's inequality and standard interpolation result, for all  $\gamma > 0$  we have

$$\|v\|_{\mathbb{H}^{1+\delta-\varepsilon, q}} \leq \gamma \|v\|_{\mathbb{H}^{1+\delta, q}} + C_\gamma \|v\|_{\mathbb{H}^{-1+\delta, q}} \stackrel{(2.5)}{=} \gamma \|v\|_{\mathbb{H}^{1+\delta, q}} + C_\gamma \|v\|_{\mathbb{H}^{-1+\delta, q}}.$$

Thus the claim of this step in case  $\delta \leq 0$  follows by combining the above estimate with the estimate in the case  $h = 0$ . Indeed, since  $u$  is a solution to (3.1), setting

$$N_{f, g} := \|f\|_{L^p((t, \tau), w_\ell^t; \mathbb{H}^{-1+\delta, q})} + \|g\|_{L^p((t, \tau), w_\ell^t; \mathbb{H}^{\delta, q}(\ell^2))},$$

the estimate for  $h = 0$  and the above estimate for  $\mathbb{P}[\tilde{f}_b u]$  imply

$$\begin{aligned} \|u\|_{L^p((t, \tau), w_\ell^t; \mathbb{H}^{1+\delta, q})} &\leq CN_{f, g} + C \|\mathbb{P}[\tilde{f}_b u]\|_{L^p((t, \tau), w_\ell^t; \mathbb{H}^{-1+\delta, q})} \\ &\leq CN_{f, g} + C\gamma \|u\|_{L^p((t, \tau), w_\ell^t; \mathbb{H}^{1+\delta, q})} + C'_\gamma \|u\|_{L^p((t, \tau), w_\ell^t; \mathbb{H}^{-1+\delta, q})}. \end{aligned}$$

Choosing  $\gamma = 1/(2C)$ , we get

$$\|u\|_{L^p((t, \tau), w_\ell^t; \mathbb{H}^{1+\delta, q})} \leq 2CN_{f, g} + 2C'_\gamma \|u\|_{L^p((t, \tau), w_\ell^t; \mathbb{H}^{-1+\delta, q})}.$$

Reasoning as in Step 6 of Lemma 3.6 one can also adsorb the remaining lower order term by using  $\tau - t \leq T^*$  and choosing  $T^*$  sufficiently small.

Case  $\delta \in (0, 1]$ . In this case, for all  $v \in \mathbb{H}^{1+\delta, q}$ ,

$$\begin{aligned} \|\mathbb{P}[\tilde{f}_b v]\|_{\mathbb{H}^{-1+\delta, q}} &\lesssim \|\tilde{f}_b v\|_{L^q} \\ &\lesssim \max_{i,j} \left( \|(h_n^{i,j})_{n \geq 1}\|_{L^\infty(\ell^2)} \|(b_n^j)_{n \geq 1}\|_{L^\infty(\ell^2)} \right) \|\nabla v\|_{L^q} \stackrel{(3.3)}{\lesssim} \|v\|_{H^{1, q}}. \end{aligned}$$

Since  $\delta > 0$ ,  $\|v\|_{H^{1, q}}$  is lower order compared to  $\|v\|_{\mathbb{H}^{1+\delta, q}}$ . Thus the claim follows from standard interpolation estimates and Young's inequality as in the case  $\delta \leq 0$ .

Case  $\delta > 1$ . The proof goes as in the case  $\delta \leq 0$ , where one should replace [AV21c, Proposition 4.1(3)] by [AV21c, Corollary 4.2].  $\square$

#### 4. PROOFS OF THE MAIN RESULTS

**4.1. Local well-posedness and the proof of Theorem 2.4.** Theorem 2.4 will be derived from a more general local well-posedness which will be proved using the methods of [AV21a, AV21b]. For  $\delta \in (-1, 0]$  and  $q \in [2, \infty)$  we define

$$\begin{aligned} (4.1) \quad X_0 &= \mathbb{H}^{-1+\delta, q}, \quad X_1 = \mathbb{H}^{1+\delta, q} \quad A(\cdot)v = A_{\mathbb{S}}v, \quad B(\cdot)v = B_{\mathbb{S}}v, \\ F(\cdot, v) &= \mathbb{P}[f_0(\cdot, v) + \operatorname{div}(f(\cdot, v)) + f_{g,h}(\cdot, v)] - \mathbb{P}(\operatorname{div}(v \otimes v)), \\ G(\cdot, v) &= (\mathbb{P}[g_n(\cdot, v)])_{n \geq 1}, \end{aligned}$$

Here  $(A_{\mathbb{S}}, B_{\mathbb{S}})$  is as in (3.2) and (3.4), and

$$f_{g,h}(\cdot, v) = \left( \sum_{n \geq 1} [(I - \mathbb{P})[g_n(\cdot, v)]] \cdot h_n^{\cdot, k} \right)_{k=1}^d.$$

Note that the linear part of  $\tilde{f}$  (see (2.8)) is in the operator  $A$ , and the nonlinear part in  $f_{g,h}$ . In this way (2.9) can be formulated as the semilinear stochastic evolution equation (see Definition 2.3 for the definition of  $(p, \kappa, q, \delta)$ -solutions):

$$(4.2) \quad \begin{cases} du + Au \, dt = F(\cdot, u)dt + (Bu + G(u))dW_{\ell^2}, \\ u(0) = u_0. \end{cases}$$

Through this section we set

$$\begin{aligned} X_{\kappa, p}^{\operatorname{Tr}} &:= (X_0, X_1)_{1 - \frac{1+\kappa}{p}, p} = \mathbb{B}_{q, p}^{1+\delta - 2\frac{1+\kappa}{p}}, \quad \text{and} \\ X_\theta &:= [X_0, X_1]_\theta = \mathbb{H}^{1+\delta - 2\theta, q}, \quad \text{for } \theta \in (0, 1), \end{aligned}$$

where we used (2.6). The following is our main result on local existence and regularization. Below we set  $1/0 := \infty$ .

**Theorem 4.1** (Local well-posedness). *Let Assumption 2.1( $p, \kappa, q, \delta$ ) hold with*

$$(4.3) \quad \delta \in (-1, 0], \quad \frac{d}{2+\delta} < q < \frac{d}{-\delta} \quad \text{and} \quad 2\frac{1+\kappa}{p} + \frac{d}{q} \leq 2 + \delta.$$

*Then for every  $u_0 \in L^0_{\mathcal{F}_0}(\Omega; \mathbb{B}_{q, p}^{1+\delta - 2\frac{1+\kappa}{p}}(\mathbb{T}^d))$ , (4.2) has a  $(p, \kappa, q, \delta)$ -solution  $(u, \sigma)$  such that  $\sigma > 0$  a.s. and*

$$(4.4) \quad u \in L^p_{\operatorname{loc}}([0, \sigma), w_\kappa; \mathbb{H}^{1+\delta, q}(\mathbb{T}^d)) \cap C([0, \sigma); \mathbb{B}_{q, p}^{1+\delta - 2\frac{1+\kappa}{p}}(\mathbb{T}^d)), \quad \text{a.s.}$$



and the trace space  $\mathbb{B}_{q,p}^{1+\delta-2\frac{1+\kappa}{p}}(\mathbb{T}^d)$  is critical for (4.2) if and only if the second part of (4.3) holds with equality. Moreover,  $(u, \sigma)$  instantaneously regularizes in time and space in the following sense: a.s.

$$(4.5) \quad u \in H_{\text{loc}}^{\theta,r}(0, \sigma; \mathbb{H}^{1-2\theta,\zeta}(\mathbb{T}^d)) \quad \text{for all } \theta \in [0, 1/2), r, \zeta \in (2, \infty),$$

$$(4.6) \quad u \in C_{\text{loc}}^{\theta_1, \theta_2}((0, \sigma) \times \mathbb{T}^d; \mathbb{R}^d) \quad \text{for all } \theta_1 \in [0, 1/2), \theta_2 \in (0, 1).$$

By Sobolev embedding it is straightforward to see that (4.5) implies (4.6). The definition of criticality which we use is taken from [AV21a, Section 4.1] and some details can be found in the proof of Theorem 4.1.

In order to prove the theorem, we first prove a lemma for the nonlinearities. For the noise part we use the language of  $\gamma$ -radonifying operators for which we refer to [HNVW17, Chapter 9]. The following identification (as sets and isomorphically) will be used several times below:

$$(4.7) \quad \gamma(\ell^2, H^{s,q}(\mathbb{T}^d)) = H^{s,q}(\mathbb{T}^d; \ell^2)$$

where  $s \in \mathbb{R}$  and  $q \in (1, \infty)$ . The identity (4.7) follows from [HNVW17, Proposition 9.3.1] and the fact that  $(I - \Delta)^{s/2}$  induces uniquely an isomorphism between  $H^{s,q}(\mathbb{T}^d; \ell^2)$  and  $L^q(\mathbb{T}^d; \ell^2)$ .

By (2.4) and the ideal property of  $\gamma$ -spaces [HNVW17, Theorem 9.1.10], we also note that  $\mathbb{P}$  extends to a bounded mapping from  $\gamma(\ell^2, H^{s,q}(\mathbb{T}^d; \mathbb{R}^d))$  into  $\gamma(\ell^2, \mathbb{H}^{s,q}(\mathbb{T}^d))$ .

To prove the theorem we will first show that the nonlinearities  $F$  and  $G$  have the right mapping properties:

**Lemma 4.2.** *Let Assumption 2.1( $p, \kappa, q, \delta$ ) be satisfied and suppose that (4.3) holds. Set  $\beta = \frac{1}{2}(1 - \frac{\delta}{2} + \frac{d}{2q})$ . Then there is a constant  $C$  such that for all  $v, v' \in \mathbb{H}^{\theta,q}$ ,*

$$\begin{aligned} \|F(\cdot, v) - F(\cdot, v')\|_{X_0} &\leq C(1 + \|v\|_{X_\beta} + \|v'\|_{X_\beta})\|v - v'\|_{X_\beta} \\ \|F(\cdot, v)\|_{X_0} &\leq C(1 + \|v\|_{X_\beta}^2) \\ \|G(\cdot, v) - G(\cdot, v')\|_{\gamma(\ell^2, X_{1/2})} &\leq C(1 + \|v\|_{X_\beta} + \|v'\|_{X_\beta})\|v - v'\|_{X_\beta}. \\ \|G(\cdot, v)\|_{\gamma(\ell^2, X_{1/2})} &\leq C(1 + \|v\|_{X_\beta}^2). \end{aligned}$$

Note that  $\beta \in (0, 1)$  by the conditions on the parameters in (4.3).

*Proof.* Note that  $X_\beta = \mathbb{H}^{\theta,q}$  with  $\theta = \frac{d}{2q} + \frac{\delta}{2}$ . First we check the estimate for  $F$ . Let  $F = F_1 + F_2 + F_3$  where

$$(4.8) \quad \begin{aligned} F_1(v) &= \mathbb{P}\text{div}(v \otimes v) \\ F_2(v) &= \mathbb{P}[f_0(\cdot, v) + \text{div}(f(\cdot, v))] \\ F_3(v) &= \mathbb{P}[f_{g,h}(\cdot, v)] \end{aligned}$$

for all  $v \in \mathbb{H}^{1+\delta,q}$ . First we estimate  $F_1$ . For all  $v, v' \in X_1$ ,

$$(4.9) \quad \begin{aligned} \|F_1(v) - F_1(v')\|_{\mathbb{H}^{-1+\delta,q}} &\lesssim \|(v \otimes v) - (v' \otimes v')\|_{H^{\delta,q}} \\ &\stackrel{(i)}{\lesssim} \|(v \otimes v) - (v' \otimes v')\|_{L^\lambda} \\ &\lesssim \|v \otimes (v - v')\|_{L^\lambda} + \|(v - v') \otimes v\|_{L^\lambda} \\ &\leq (\|v\|_{L^{2\lambda}} + \|v'\|_{L^{2\lambda}})\|v - v'\|_{L^{2\lambda}} \\ &\stackrel{(ii)}{\lesssim} (\|v\|_{\mathbb{H}^{\theta,q}} + \|v'\|_{\mathbb{H}^{\theta,q}})\|v - v'\|_{\mathbb{H}^{\theta,q}}, \end{aligned}$$

where in (i) and (ii) we have used the Sobolev embedding with  $-\frac{d}{\lambda} = \delta - \frac{d}{q}$  and  $\theta - \frac{d}{q} = -\frac{d}{2\lambda}$ . Therefore, we find  $\theta = \frac{d}{2q} + \frac{\delta}{2}$ . To ensure that both Sobolev embeddings hold note that  $\lambda \in (q/2, q]$  and  $\theta > 0$  by (4.3). The estimate for  $\|F_1(v)\|_{X_0}$  follows since  $F_1(0) = 0$ .

We prove a similar estimate for  $F_2$ . Indeed, by Assumption 2.1,

$$\begin{aligned}
(4.10) \quad \|F_2(\cdot, v) - F_2(\cdot, v')\|_{\mathbb{H}^{-1+\delta, q}} &\lesssim \sum_{j=0}^d \|f_j(\cdot, v) - f_j(\cdot, v')\|_{L^\lambda} \\
&\lesssim \|(1 + |v| + |v'|)|v - v'|\|_{L^\lambda} \\
&\lesssim (1 + \|v\|_{L^{2\lambda}} + \|v'\|_{L^{2\lambda}})\|v - v'\|_{L^{2\lambda}} \\
&\lesssim (1 + \|v\|_{\mathbb{H}^{\theta, q}} + \|v'\|_{\mathbb{H}^{\theta, q}})\|v - v'\|_{\mathbb{H}^{\theta, q}},
\end{aligned}$$

where  $\lambda$  and  $\theta$  are as before. The estimate for  $\|F_2(\cdot, v)\|_{X_0}$  follows as well by using the boundedness of  $f_j(\cdot, 0)$ .

Next we consider  $F_3$ . Since  $h \in L^\infty(\mathbb{T}^d; \ell^2)$  (see Assumption 2.1)

$$\begin{aligned}
(4.11) \quad \|F_3(\cdot, v) - F_3(\cdot, v')\|_{\mathbb{H}^{-1+\delta, q}} &\lesssim \|f_{g, h}(\cdot, v) - f_{g, h}(\cdot, v')\|_{L^\lambda} \\
&\lesssim \|h\|_{L^\infty(\mathbb{T}^d; \ell^2)} \|(1 + |v| + |v'|)|v - v'|\|_{L^\lambda} \\
&\lesssim (1 + \|v\|_{\mathbb{H}^{\theta, q}} + \|v'\|_{\mathbb{H}^{\theta, q}})\|v - v'\|_{\mathbb{H}^{\theta, q}}
\end{aligned}$$

The estimate for  $\|F_3(\cdot, v)\|_{X_0}$  follows as well by using the boundedness of  $f_{g, h}(\cdot, 0)$ .

Combining the estimates (4.9), (4.10) and (4.11), we obtain the required estimate for  $F$  by the definition of  $\beta$ .

It remains to estimate  $G$ . Recall that  $X_{1/2} = \mathbb{H}^{\delta, q}$  (see (2.6)) and let  $\beta, \theta$  be as above. Then, Assumption 2.1 and (4.7) yield that for all  $v, v' \in \mathbb{H}^{1+\delta, q}$ ,

$$\begin{aligned}
(4.12) \quad \|G(\cdot, v) - G(\cdot, v')\|_{\gamma(\ell^2, \mathbb{H}^{\delta, q})} &\lesssim \|g(\cdot, v) - g(\cdot, v')\|_{\gamma(\ell^2, L^\lambda)} \\
&\lesssim (1 + \|v\|_{L^{2\lambda}} + \|v'\|_{L^{2\lambda}})\|v - v'\|_{L^{2\lambda}} \\
&\lesssim (1 + \|v\|_{X_\beta} + \|v'\|_{\mathbb{H}^{\theta, q}})\|v - v'\|_{\mathbb{H}^{\theta, q}}.
\end{aligned}$$

□

The proof of Theorem 4.1 is split into three parts. In Part (A) we prove local well-posedness. In Parts (B) and (C) we prove (4.5) in the case  $[p > 2]$  and  $[q = p = d = 2, \delta = 0]$ , respectively. The proof of (4.6) follows as a simple consequence as already mentioned before.

*Proof of Theorem 4.1 Part (A) – Local well-posedness.* The proof is divided in several steps.

*Step 1: the conditions (HF) and (HG) of [AV21a, Section 4.1] hold with (4.1), and the trace space  $X_{\kappa, p}^{\text{Tr}} = \mathbb{B}_{q, p}^{1+\delta-2\frac{1+\kappa}{p}}$  is critical for (4.2) if and only if*

$$(4.13) \quad 2\frac{1+\kappa}{p} + \frac{d}{q} = 2 + \delta.$$

To prove this we use Lemma 4.2 and consider two cases. Recall  $\beta = \frac{1}{2}(1 - \frac{\delta}{2} + \frac{d}{2q})$ .

(1) If  $1 - \frac{1+\kappa}{p} \geq \beta$ , one can check that (4.13) does not hold by using  $q > d/(2 + \delta)$ .

We prove the desired mapping properties of  $F$  and  $G$  and non-criticality. Using that  $X_{1 - \frac{1+\kappa}{p} + \varepsilon} \hookrightarrow X_\beta$  for each  $\varepsilon > 0$ , we get that Lemma 4.2 holds with  $\beta$  replaced by  $1 - \frac{1+\kappa}{p} + \varepsilon$ . Letting  $\rho_j = 1, \beta_j = \varphi_j = 1 - \frac{1+\kappa}{p} + \varepsilon$  for  $j \in \{1, 2\}$ ,

where  $\varepsilon > 0$  is such that  $\varepsilon < \frac{1+\kappa}{2p}$ , one can check  $\rho_1\left(\varphi_1 - 1 + \frac{1+\kappa}{p}\right) + \beta_1 < 1$ , which means that  $X_{\kappa,p}^{\text{Tr}}$  is not critical.

(2) If  $1 - \frac{1+\kappa}{p} < \beta$ , then we set  $\rho_j = 1, \beta_j = \varphi_j = \beta$  and by Lemma 4.2 the condition in [AV21a, (4.2)] becomes

$$(4.14) \quad \frac{1+\kappa}{p} \leq \frac{\rho_j + 1}{\rho_j}(1 - \beta) = 1 - \frac{d}{2q} + \frac{\delta}{2}.$$

Note that (4.14) is equivalent to the second part of (4.3). Finally, the corresponding trace space  $\mathbb{B}_{q,p}^{-1+\delta-2\frac{1+\kappa}{p}}$  is critical for (2.1) if and only if the equality in (4.14) holds.

*Step 2: Application of [AV21a, Theorem 4.8].* By Step 1 and Theorem 3.2 the conditions of the latter are satisfied. Therefore, there is a  $(p, \kappa, q, \delta)$ -solution  $(u, \sigma)$  and (4.4) holds. The assertions on criticality of the trace space also follows from Step 1.  $\square$

Next we prove Part (B) of Theorem 4.1, i.e. we show (4.5)-(4.6) in the case  $p > 2$ . Here we exploit the results in [AV21b, Section 6] in the following way.

- Bootstrap regularity in time via [AV21b, Proposition 6.8] (Step 1a below) and [AV21b, Corollary 6.5] (Step 1b below).
- Bootstrap high-order integrability in space. Here we apply [AV21b, Theorem 6.3] finitely many times in the  $(\mathbb{H}^{-1,q_j}, \mathbb{H}^{1,q_j}, r, \alpha)$ -setting with  $(q_j)_{j \geq 1}$  such that  $q_{j+1} - q_j \geq c > 0$  (see Step 2).
- Bootstrap regularity in space by applying [AV21b, Theorem 6.3] on the shifted scales  $Y_i = \mathbb{H}^{-1+\delta+2i}$  and  $\hat{Y}_i = \mathbb{H}^{-1+2i}$  (see Step 3).

*Proof of Theorem 4.1 Part (B) – Proof of (4.5) in the case  $p > 2$ .* Let  $(u, \sigma)$  be the  $(p, \kappa, q, \delta)$ -solution to (4.2) provided by Part (A). In the proof below we use Theorem 3.2 without further reference to obtain the required stochastic maximal regularity in different settings.

*Step 1: For all  $r \in (2, \infty)$ ,*

$$(4.15) \quad u \in \bigcap_{\theta \in (0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1+\delta-2\theta, q}), \text{ a.s.}$$

To prove this regularization effect in time, in the case  $\kappa = 0$  we divide the argument into two sub-steps. In the latter case, we will first use [AV21b, Proposition 6.8] to create a weighted setting with a slight increase in integrability. After that we will apply [AV21b, Corollary 6.5] to extend the integrability to arbitrary order. In the case  $\kappa > 0$ , Step 1a below can be skipped.

*Step 1a: If  $\kappa = 0$ , then (4.15) holds for some  $r > p$ .* Let  $\alpha > 0$  be such that

$$Y_i = X_i = \mathbb{H}^{-1+2i-\delta, q}, \quad i \in \{0, 1\}, \quad \frac{1}{p} = \frac{1+\alpha}{r}, \quad \text{and} \quad \frac{1}{r} = \beta_1 - 1 + \frac{1}{p},$$

where  $\beta_1 \in (0, 1)$  is as in Part (A) of the proof. Then  $\alpha < \frac{r}{2} - 1$ . Now by Part (A) of the proof we can apply [AV21b, Proposition 6.8] to obtain (4.15).

*Step 1b: (4.15) holds for all  $r \in (2, \infty)$ .* Let  $r > p, \alpha > 0$  be as in Step 1. Let  $\hat{r} \in [r, \infty)$  be arbitrary and let  $\hat{\alpha} \in [0, \frac{r}{2} - 1)$  be such that  $\frac{1+\hat{\alpha}}{\hat{r}} < \frac{1+\alpha}{r}$ . Step 1 of the above proof Part (A) and the above Step 1a show that [AV21b, Corollary 6.5] applies with  $Y_i = \mathbb{H}^{-1+\delta+2i, q}(\mathbb{T}^d)$ ,  $\kappa, r, \alpha$  and  $\hat{r}, \hat{\alpha}$  as above.

*Step 2: For all  $r \in (2, \infty)$  and  $\zeta \in (2, \frac{d}{-\delta})$ ,*

$$(4.16) \quad u \in \bigcap_{\theta \in [0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1+\delta-2\theta, \zeta}), \text{ a.s.}$$

From Step 2, we know that (4.16) holds for all  $r \in (2, \infty)$  and  $\zeta = q$ . Fix  $\alpha > 0$  and choose  $r > p$  such that  $2\frac{1+\alpha}{r} + \frac{d}{q} < 2 + \delta$ . By Step 1 of Part (A) we know that for all  $\zeta \in [q, d/(-\delta))$ , (HF), (HG), and non-criticality holds in the  $(\mathbb{H}^{-1+\delta, \zeta}, \mathbb{H}^{1+\delta, \zeta}, r, \alpha)$ -setting. To prove the claim, it suffices to show the existence of  $\varepsilon > 0$  depending only on  $q, \delta$  such that for any  $\zeta \in [q, d/(-\delta))$ ,

$$(4.17) \quad u \in \bigcap_{\theta \in [0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1+\delta-2\theta, \zeta}) \text{ a.s.} \implies u \in \bigcap_{\theta \in [0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1+\delta-2\theta, \zeta+\varepsilon}) \text{ a.s.}$$

To prove (4.17), we apply [AV21b, Theorem 6.3]. Due to the previous choice of  $r, \alpha$ , Step 1 of Part (A), and the left-hand side of (4.17), [AV21b, Theorem 6.3] is applicable with

$$Y_i = \mathbb{H}^{1+\delta-2i, \zeta}, \quad \widehat{Y}_i = \mathbb{H}^{1+\delta-2i, \zeta+\varepsilon}, \quad \widehat{r} = r, \quad \widehat{\alpha} = \alpha$$

where  $\varepsilon > 0$  is chosen so that

$$Y_r^{\text{Tr}} = \mathbb{B}_{\zeta, r}^{1+\delta-\frac{2}{r}} \hookrightarrow \mathbb{B}_{\zeta+\varepsilon, r}^{1+\delta-2\frac{1+\alpha}{r}} = \widehat{Y}_{\widehat{\alpha}, \widehat{r}}^{\text{Tr}}.$$

By Sobolev embeddings, the previous display holds provided  $\frac{1}{\zeta+\varepsilon} - \frac{1}{\zeta} \leq 2\frac{\alpha}{dr}$ . To satisfy the latter, it is enough to choose  $\varepsilon := 2\frac{\alpha}{dr}$ .

*Step 3: For all  $\zeta, r \in (2, \infty)$ ,*

$$(4.18) \quad u \in \bigcap_{\theta \in [0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1-2\theta, \zeta}), \text{ a.s.}$$

If  $\delta = 0$ , then (4.18) follows from (4.16). In case  $\delta < 0$  first fix  $\zeta \in (d, d/(-\delta))$ . Let  $r \geq p$  be so large that  $r > \frac{2}{1+\delta}$  and

$$(4.19) \quad \frac{2}{r} + \frac{d}{\zeta} < 2 + \delta.$$

This is possible since  $\zeta > d$  and  $\delta > -1$ . Set  $\widehat{\alpha} := r(\frac{1}{r} - \frac{\delta}{2}) - 1 \in [0, \frac{r}{2} - 1)$ . Then by (4.19) and Step 1 of Part (A), (HF), (HG), and non-criticality hold in the  $(\mathbb{H}^{-1, q}, \mathbb{H}^{1, q}, r, \widehat{\alpha})$ -setting. Now [AV21b, Theorem 6.3] with

$$X_i = Y_i = \mathbb{H}^{-1+\delta+2i, q}, \quad \widehat{Y}_i = \mathbb{H}^{-1+2i, q}, \quad \widehat{r} = r, \quad \alpha = 0, \quad \widehat{\alpha} \text{ as above.}$$

gives (4.18) with the above  $\zeta$  and  $r$ . To check the conditions (1)-(3) in [AV21b, Theorem 6.3], note that (1) and (2) follow from (4.16), Step 1 of Part (A) and the above choice of  $Y_i, \widehat{Y}_i, r, \widehat{r}, \alpha, \widehat{\alpha}$ . To check (3) note that  $\frac{1+\widehat{\alpha}}{r} = \frac{1}{r} - \frac{\delta}{2}$ . Thus

$$Y_r^{\text{Tr}} = \mathbb{B}_{q, r}^{1+\delta-\frac{2}{r}} = \mathbb{B}_{q, r}^{1-2\frac{1+\widehat{\alpha}}{r}} = \widehat{Y}_{\widehat{\alpha}, \widehat{r}}^{\text{Tr}}.$$

The second part of condition (3) holds due to [AV21b, Lemma 6.2(4)] since  $\frac{1+\widehat{\alpha}}{r} = \frac{1}{r} - \frac{\delta}{2}$ ,  $\widehat{Y}_{1+\frac{\delta}{2}} = \mathbb{H}^{1+\delta, q} = Y_1$  and  $\widehat{Y}_0 = \mathbb{H}^{-1, q} = Y_{1+\frac{\delta}{2}}$ .

In order to obtain (4.18) for all  $\zeta < \infty$ , we can again apply (4.17) but this time with  $\delta = 0$ .  $\square$

Next we prove Part (C) of Theorem 4.1, i.e. we show (4.5) in the case  $q = p = 2$  and  $\kappa = 0$ . Here we follow the arguments in [AV21b, Section 7] employing the results in [AV21b, Section 6], see [AV21b, Roadmap 7.4] for a strategy summary.

*Proof of Theorem 4.1 Part (C) – Proof of (4.5) in the case  $q = p = d = 2$ ,  $\delta = 0$ .* Let  $(u, \sigma)$  be the  $(2, 0, 2, 0)$ -solution to (4.2) provided by Part (A). It is enough to prove that for all  $\varepsilon \in (-\frac{1}{2}, 0)$  for all  $r \in (2, \infty)$ ,

$$(4.20) \quad u \in \bigcap_{\theta \in [0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1+\varepsilon-2\theta}) \quad \text{a.s.}$$

where, to shorthand the notation, we wrote  $\mathbb{H}^{1+\varepsilon-2\theta}$  instead of  $\mathbb{H}^{1+\varepsilon-2\theta, 2}$ . Indeed, afterwards we can apply Steps 2 and 3 of Part (B) with  $\delta = \varepsilon$  and  $q = 2$ .

*Step 1: For each  $\varepsilon \in (0, \frac{1}{2})$ , (4.20) holds with  $r = 4$ .* Let  $\alpha > 0$  be such that

$$(4.21) \quad Y_i = \mathbb{H}^{-1+2i-\varepsilon}, \quad X_i = \mathbb{H}^{-1+2i}, \quad p = 2, \quad r = 4, \quad \frac{1}{2} = \frac{1+\alpha}{4} + \frac{\varepsilon}{2}.$$

Since  $\varepsilon \in (0, \frac{1}{2})$ , (4.21) yields  $\alpha \in (0, 1) = (0, \frac{r}{2} - 1)$ . To conclude Step 1, it is enough to check the assumptions of [AV21b, Proposition 6.8]. However, these follow from Step 1 of Part (A), Theorem 3.2, and the fact that  $\varphi_j = \beta_j = \frac{1}{2}(1 - \frac{d}{2q}) = \frac{3}{4}$  for  $j \in \{1, 2\}$ .

*Step 2: For each  $\varepsilon \in (0, \frac{1}{2})$ , (4.20) holds with arbitrary  $r > 2$ .* Let  $r = 4$  and  $\alpha \in (0, 1)$  be as in Step 1. Let  $\hat{r} \in [4, \infty)$  be arbitrary and let  $\hat{\alpha} \geq 0$  such that

$$Y_i = \mathbb{H}^{-1-\varepsilon+2i}, \quad X_i = \mathbb{H}^{-1+2i}, \quad \hat{r} \in (4, \infty), \quad \frac{1+\hat{\alpha}}{\hat{r}} < \frac{1+\alpha}{r}.$$

Then  $\hat{\alpha} \in [0, \frac{\hat{r}}{2} - 1)$ . It remains to apply [AV21b, Corollary 6.5]. Note that the assumptions follow from Step 1 of Part (A), and

$$Y_r^{\text{Tr}} = \mathbb{B}_{2,r}^{1-\varepsilon-\frac{2}{r}} \hookrightarrow \mathbb{L}^2 = X_{0,2}^{\text{Tr}},$$

where we used  $1 - \varepsilon > \frac{2}{r}$ . □

*Proof of Theorem 2.4.* It suffices to set  $\kappa = \kappa_c$  in Theorem 4.1. One can check that this is admissible under the conditions on  $p, q, \delta$  in (4.3). □

In Theorem 2.4 we omitted  $\delta \in (-1, -\frac{1}{2})$ . The reason for this is that it does not enlarge the class of critical spaces with smoothness  $> -\frac{1}{2}$  for the initial data for which we can treat (2.1). Indeed, in case  $\delta \in (-1, -\frac{1}{2})$ , then the restrictions  $q < \frac{d}{-\delta}$  and  $2\frac{1+\kappa}{p} + \frac{d}{q} \leq 2 + \delta$  (see (4.3)) implies that the smoothness of the space of the initial data satisfies  $1 + \delta - 2\frac{1+\kappa}{p} > \frac{d}{q} - 1 > -1 - \delta > -\frac{1}{2}$ .

**4.2. Higher order regularity and the proof of Theorem 2.7.** For the proof of Theorem 2.7 we need the following lemma, which is the analogue of Lemma 4.2 but with  $\delta$  replaced by a higher order smoothness  $s > 0$ .

**Lemma 4.3.** *Suppose that Assumption 2.1( $p, \kappa, q, \delta$ ) and Assumption 2.6 holds. Let  $X_j = \mathbb{H}^{-1+2j+s, q}$  with  $s \in (0, \eta]$ . Assume that*

$$(4.22) \quad q \in (d, \infty), \quad p \in (2, \infty), \quad \kappa \in \left[0, \frac{p}{2} - 1\right) \quad \text{satisfy} \quad 1 + s - \frac{2(1+\kappa)}{p} - \frac{d}{q} > 0.$$

Then for all  $n \geq 1$  there exists a constant  $C_n$  such that for all  $v, v' \in X_1$  satisfying  $\|v\|_{X_{\kappa,p}^{\text{Tr}}}, \|v'\|_{X_{\kappa,p}^{\text{Tr}}} \leq n$ ,

$$\begin{aligned} \|F(\cdot, v)\|_{X_0} &\leq C_n(1 + \|v\|_{X_{\kappa,p}^{\text{Tr}}}), \\ \|F(\cdot, v) - F(\cdot, v')\|_{X_0} &\leq C_n\|v - v'\|_{X_{\kappa,p}^{\text{Tr}}}, \\ \|G(\cdot, v)\|_{\gamma(\ell^2, X_{1/2})} &\leq C_n(1 + \|v\|_{X_{\kappa,p}^{\text{Tr}}}), \\ \|G(\cdot, v) - G(\cdot, v')\|_{\gamma(\ell^2, X_{1/2})} &\leq C_n\|v - v'\|_{X_{\kappa,p}^{\text{Tr}}}, \end{aligned}$$

Note that this lemma will also imply non-criticality of  $F$  and  $G$  since using the notation of [AV21a, Section 4.1] we can take  $F = F_{\text{Tr}}$  and  $G = G_{\text{Tr}}$  and hence the critical parts  $F_c$  and  $G_c$  vanish.

*Proof.* Note that  $X_\beta = \mathbb{H}^{s+2\beta,q}$ ,  $X_{\kappa,p}^{\text{Tr}} = \mathbb{B}_{q,p}^{1+s-2(1+\kappa)/p}$ . Due to the last condition in (4.22) and Sobolev embedding we have

$$(4.23) \quad X_{\kappa,p}^{\text{Tr}} \hookrightarrow C^\varepsilon \cap H^{s,q}, \quad \text{for some } \varepsilon > 0.$$

We prove the claimed estimate for  $F$ . The estimate for  $G$  follows similarly. Fix  $n \geq 1$  and  $v, v' \in X_1$  satisfying  $\|v\|_{X_{\kappa,p}^{\text{Tr}}}, \|v'\|_{X_{\kappa,p}^{\text{Tr}}} \leq n$ . As in (4.8) we write  $F = F_1 + F_2 + F_3$ . Note that

$$\begin{aligned} \|F_1(v) - F_1(v')\|_{\mathbb{H}^{-1+s,q}} &\lesssim_{s,q} \|(v - v') \otimes v\|_{H^{s,q}} + \|v' \otimes (v - v')\|_{H^{s,q}} \\ &\stackrel{(i)}{\lesssim}_{s,q} \|v - v'\|_{L^\infty} \|v\|_{H^{s,q}} + \|v\|_{H^{s,q}} \|v - v'\|_{L^\infty} \\ &\stackrel{(4.23)}{\lesssim}_n \|v - v'\|_{X_{\kappa,p}^{\text{Tr}}} \end{aligned}$$

where in (i) we used [AV21c, Proposition 4.1(1)] and in (ii) Sobolev embedding. The estimate for  $F_1(v)$  follows as well since  $F_1(0) = 0$ .

Next we prove the estimates for  $F_2$ . We only consider the  $\text{div}(f(\cdot, v))$  part, since the other term is similar. Let us write

$$\begin{aligned} f(\cdot, v) - f(\cdot, v') &= \Phi(v, v')(v - v'), \\ \text{where } \Phi(v, v') &:= \int_0^1 \partial_y f(\cdot, (1-t)v + tv') dt. \end{aligned}$$

By repeating the argument for  $F_1$ , it is enough to show that

$$\|\Phi(v, v')\|_{H^{s,q}} \leq C_n.$$

The latter follows from Assumption 2.6, [BMS10, Theorem 1] or [Tay11, Proposition 10.2, Chapter 13] and (4.23). The estimate for  $F_2(\cdot, v)$  can be proved more directly. The estimates for  $F_3$  are proved similarly.  $\square$

After this preparation we will now be able to prove the higher order regularity by an iteration argument based on [AV21b, Theorem 6.3].

*Proof of Theorem 2.7.* As usual we only prove (2.15), since (2.16) follows from (2.15) and Sobolev embeddings.

Let  $\eta, \xi$  be as in Assumption 2.1. Let  $\eta_0 \geq 0$  and  $\xi_0 \in (d, \infty)$  be given by

$$\xi_0 = \begin{cases} d+1, & \text{if } \xi \vee q \leq d, \\ \xi \vee q & \text{otherwise,} \end{cases} \quad \text{and} \quad \eta - \frac{d}{\xi} = \eta_0 - \frac{d}{\xi_0}.$$

Note that  $\eta_0 \in (0, \eta]$  since  $\xi_0 \geq \xi$  and  $\eta > \frac{d}{\xi}$ . We split the proof into two steps. In the following steps, without further mentioning it, we apply Theorem 3.2 several times in order to check stochastic maximal regularity of  $(A, B) = (A_{\mathbb{S}}, B_{\mathbb{S}})$ . This is needed in order to check the conditions of our bootstrapping result [AV21b, Theorem 6.3]. In case  $\xi_0 = \xi$  we also have  $\eta_0 = \eta$  and Step 2 below is not needed.

*Step 1: For all  $r \in (2, \infty)$ ,*

$$u \in \bigcap_{\theta \in [0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1+\eta_0, \xi_0}) \quad \text{a.s.}$$

To show this fix  $N \in \mathbb{N}$  so large that  $\eta_0 < N$ . Set  $s_k = k\frac{\eta_0}{N}$  for  $k \in \{0, \dots, N\}$ . By induction and (2.12) in Theorem 2.4, it is enough to prove that for each  $k \in \{1, \dots, N\}$ ,

(4.24)

$$u \in \bigcap_{\theta \in [0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1+s_{k-1}-2\theta, \xi_0}) \text{ a.s.} \Rightarrow u \in \bigcap_{\theta \in [0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1+s_k-2\theta, \xi_0}) \text{ a.s.}$$

To prove (4.24) we apply [AV21b, Theorem 6.3] with  $X_j = \mathbb{H}^{-1+2j+\delta, q}$ ,  $Y_j = \mathbb{H}^{-1+s_{k-1}, \xi_0}$ ,  $\widehat{Y}_j = \mathbb{H}^{-1+s_k, \xi_0}$  for  $j \in \{0, 1\}$  and  $\alpha = 0$ ,  $r = \widehat{r}$ ,  $\widehat{\alpha}$  to be chosen below. Note that  $\widehat{Y}_j \hookrightarrow Y_j \hookrightarrow X_j$  for all  $j \in \{0, 1\}$ . Since  $\xi_0 > d$  and  $\eta_0 < N$ , one can find  $r \geq p$  such that  $1 - \frac{d}{\xi_0} > \frac{2}{r}$  and  $\frac{\eta_0}{N} < 1 - \frac{2}{r}$ . Set  $\widehat{\alpha} = \frac{r}{2} \frac{\eta_0}{N} \in [0, \frac{r}{2} - 1)$ . It remains to check the conditions (1)–(3) of [AV21b, Theorem 6.3]. Conditions (1) and (2) are satisfied by the induction hypothesis and Lemma 4.3 which is applicable since

$$1 + s_k - \frac{d}{\xi_0} - 2\frac{1+\widehat{\alpha}}{r} = 1 + s_{k-1} - \frac{d}{\xi_0} - \frac{2}{r} > 0, \quad \text{for all } k \in \{1, \dots, N\},$$

where we used  $s_{k-1} \geq 0$  and  $1 - \frac{d}{\xi_0} > \frac{2}{r}$ . To check condition (3) note that [AV21b, (6.1)] follows from [AV21b, Lemma 6.2(4)],  $\widehat{Y}_{1-\frac{\eta_0}{2N}} = Y_1$ ,  $Y_{\frac{\eta_0}{2N}} = \widehat{Y}_0$ , and the choice of the parameters. For (3) it remains to observe that

$$Y_r^{\text{Tr}} = \mathbb{B}_{\xi, r}^{1+s_{k-1}-\frac{2}{r}} = \mathbb{B}_{\xi, r}^{1+s_k-2\frac{1+\widehat{\alpha}}{r}} = \widehat{Y}_{\widehat{\alpha}, r}^{\text{Tr}}.$$

Therefore, we can conclude (4.24).

*Step 2: Proof of (2.15).* Fix  $L \in \mathbb{N}$  such that  $\eta - \eta_0 < L$ . Let  $\eta_k = \eta_0 + k\frac{\eta - \eta_0}{L}$  where  $k \in \{1, \dots, L\}$ . Define  $\xi_0 \geq \xi_1 \geq \dots \geq \xi_L = \xi$  by

$$(4.25) \quad \eta_k - \frac{d}{\xi_k} = \eta - \frac{d}{\xi},$$

By induction and Step 1 it is enough to prove that for each  $k \in \{1, \dots, L\}$ ,

(4.26)

$$u \in \bigcap_{\theta \in [0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1+\eta_{k-1}-2\theta, \xi_{k-1}}) \text{ a.s.} \Rightarrow u \in \bigcap_{\theta \in [0, 1/2)} H_{\text{loc}}^{\theta, r}(0, \sigma; \mathbb{H}^{1+\eta_k-2\theta, \xi_k}) \text{ a.s.}$$

Choose  $r$  so large that  $\frac{\eta - \eta_0}{L} < 1 - \frac{2}{r}$  (this is possible since  $\eta - \eta_0 < L$ ). Set  $\widehat{\alpha} := \frac{\eta - \eta_0}{L} \frac{r}{2} \in [0, \frac{r}{2} - 1)$ . As in the previous step, (4.26) follows from [AV21b, Theorem 6.3] applied with  $X_j = \mathbb{H}^{-1+2j+\delta, q}$ ,  $Y_j = \mathbb{H}^{-1+2j+\eta_{k-1}, \xi_{k-1}}$ ,  $\widehat{Y}_j = \mathbb{H}^{-1+2j+\eta_k, \xi_k}$  and  $r = \widehat{r}$ ,  $\alpha = \widehat{\alpha}$  as above. Note that by (4.25) and Sobolev embeddings,  $\widehat{Y}_j \hookrightarrow Y_j \hookrightarrow X_j$  for  $j \in \{0, 1\}$ . It remains to check the conditions (1)–(3) of [AV21b, Theorem 6.3].

Conditions (1)-(2) follows from Lemma 4.3, which is applicable since

$$1 + \eta_k - 2\frac{1 + \widehat{\alpha}}{r} - \frac{d}{\xi_k} \geq \eta_k - \frac{d}{\xi_k} \stackrel{(4.25)}{>} 0,$$

where we used  $\frac{1 + \widehat{\alpha}}{r} < \frac{1}{2}$ . To check condition (3) note that [AV21b, (6.1)] follows from [AV21b, Lemma 6.2(1)], and the fact that

$$Y_r^{\text{Tr}} = \mathbb{B}_{\xi_{k-1}, r}^{1 + \eta_{k-1} - \frac{2}{r}} \hookrightarrow \mathbb{B}_{\xi_k, r}^{1 + \eta_{k-1} - \frac{2}{r}} \hookrightarrow \mathbb{B}_{\xi_k, r}^{1 + \eta_k - 2\frac{1 + \widehat{\alpha}}{r}} = \widehat{Y}_{\widehat{\alpha}, r}^{\text{Tr}},$$

where we used the definition of  $\widehat{\alpha}$  and the fact that  $\xi_{k-1} \geq \xi_k$ .  $\square$

**4.3. Blow-up criteria and Theorems 2.9.** The proof of Theorem 2.9 is based on the blow-up criteria given in [AV21b, Section 4] and the extrapolation result in [AV21b, Lemma 6.10]. We begin by specializing the [AV21b, Lemma 6.10] to the present situation. Let  $(u, \sigma)$  be the  $(p, \kappa_c, q, \delta)$ -solution (2.1) provided by Theorem 2.4. Consider the following stochastic Navier-Stokes equations starting at  $\varepsilon > 0$ :

$$(4.27) \quad \begin{cases} dv + Av dt = F(\cdot, v)dt + (Bv + G(v))dW_{\ell^2}, \\ v(\varepsilon) = \mathbf{1}_{\sigma > \varepsilon} u(\varepsilon). \end{cases}$$

Definition 2.3 extends naturally to the problem (4.27). By either (2.13) or (2.14),

$$(4.28) \quad \mathbf{1}_{\sigma > \varepsilon} u(\varepsilon) \in L_{\mathcal{F}_\varepsilon}^0(\Omega; \mathbb{B}_{q_0, p_0}^{\frac{d}{q_0} - 1}(\mathbb{T}^d)) \text{ a.s. for all } p_0, q_0 \in (2, \infty).$$

Thus Theorem 4.1 gives the existence and uniqueness of a  $(p_0, 0, q_0, \delta_0)$ -solution  $(v, \tau)$  under conditions on the parameters.

**Lemma 4.4** (Extrapolating life-span). *Let Assumption 2.1( $p, \kappa_c, q, \delta$ ) hold with*

$$-\frac{1}{2} \leq \delta \leq 0, \quad \frac{d}{2 + \delta} < q < \frac{d}{1 + \delta}, \quad \frac{2}{p} + \frac{d}{q} \leq 2 + \delta, \quad \kappa = \kappa_c = -1 + \frac{p}{2}(2 + \delta - \frac{d}{q}).$$

*Let Assumption 2.1( $p_0, \kappa_0, q_0, \delta_0$ ) hold with*

$$\delta_0 \in (-1, 0], \quad \frac{d}{2 + \delta_0} < q_0 < \frac{d}{-\delta_0} \quad \text{and} \quad 2\frac{1 + \kappa_0}{p_0} + \frac{d}{q_0} \leq 2 + \delta_0.$$

*Let  $(u, \sigma)$  be the  $(p, \kappa_c, q, \delta)$ -solution to (2.1). Fix  $\varepsilon > 0$  and let  $(v, \tau)$  be the  $(p_0, \kappa_0, q_0, \delta_0)$ -solution to (4.27). Then*

$$\tau = \sigma \text{ a.s. on } \{\sigma > \varepsilon\} \quad \text{and} \quad u = v \text{ a.e. on } \llbracket \varepsilon, \sigma \rrbracket.$$

*Proof.* The claim follows by applying [AV21b, Lemma 6.10] with  $X_j = \mathbb{H}^{-1+2j+\delta, q}(\mathbb{T}^d)$ ,  $Y_j = \mathbb{H}^{-1+2j+\delta_0, q_0}(\mathbb{T}^d)$ ,  $\widehat{Y}_j = \mathbb{H}^{-1+2j, \widehat{q}_0}(\mathbb{T}^d)$ ,  $r = p_0$ ,  $\widehat{r} = \widehat{p}_0$ ,  $\alpha = \kappa_0$  and  $\widehat{\alpha} = 0$  where  $\widehat{p}_0, \widehat{q}_0 > 2$  are large enough. To check this, note that the required regularity assumptions are satisfied thanks to (2.13) applied to  $(u, \sigma)$  and  $(v, \tau)$ . The required stochastic maximal regularity also holds in the  $(Y_0, Y_1, p_0, \kappa_0)$ -setting by Theorem 3.2 and the conditions on the parameters. Moreover, by Lemma 4.2 the conditions (HF)-(HG) of [AV21b, Section 4.1] hold in the  $(X_0, X_1, p, \kappa_c)$ -setting,  $(Y_0, Y_1, p_0, \kappa_0)$ -setting, and the  $(\widehat{Y}_0, \widehat{Y}_1, \widehat{p}_0, 0)$ -setting if  $\widehat{p}_0, \widehat{q}_0$  are large enough. The required embeddings  $\widehat{Y}_{\widehat{p}_0}^{\text{Tr}} \hookrightarrow X_{\kappa_c, p}^{\text{Tr}}$  and  $\widehat{Y}_1 \hookrightarrow X_{1 - \frac{\kappa}{p}}$  also hold if  $\widehat{p}_0, \widehat{q}_0$  are chosen large enough.  $\square$

Next we prove Theorem 2.9 by applying our blow-up criteria from [AV21b].



*Proof of Theorem 2.9.* By Definition 2.3, (2.1) is understood as a stochastic evolution equation [AV21b, (4.1)] with the choice (4.1),  $X_j = \mathbb{H}^{-1+2j+\delta, q}$ ,  $\kappa_c = -1 + \frac{p}{2}(2 + \delta - \frac{d}{q})$ , and  $H = \ell^2$ . As in the proof of Lemma 4.4, Lemma 4.2 implies that (HF)-(HG) in [AV21b, Section 4.1] hold in the  $(X_0, X_1, p, \kappa_c)$ -setting with  $\varphi_j = \beta_j = \frac{1}{2}(1 - \frac{\delta}{2} + \frac{d}{2q})$ ,  $\rho_j = 1$  for  $j \in \{1, 2\}$ . The same holds with  $(p, \kappa_c, q, \delta)$  replaced by  $(p_0, \kappa_{0,c}, q_0, \delta_0)$  where  $\kappa_{0,c} := -1 + \frac{p_0}{2}(2 + \delta_0 - \frac{d}{q_0})$  for the translated problem (4.28). By the condition  $q_0 < \frac{d}{1+\delta_0}$ , one can check that  $\kappa_{0,c} < \frac{p_0}{2} - 1$ .

Finally note that Theorem 3.2 can be applied in these two different settings, which will be used in Step 1 below when we apply the blow-up criteria. Below  $\varepsilon, T \in (0, \infty)$  are fixed.

*Step 1: Proof of (2.17).* [AV21b, Theorem 4.11] applied to the  $(p, \kappa_{c,0}, q_0, \delta_0)$ -solution  $(v, \tau)$  to (4.27) gives

$$\mathbb{P}\left(\varepsilon < \tau < T, \|v\|_{L^{p_0}(\varepsilon, \tau; \mathbb{H}^{\gamma_0, q_0}(\mathbb{T}^d; \mathbb{R}^d))} < \infty\right) = 0,$$

where we used  $[\mathbb{H}^{-1+\delta_0, q_0}, \mathbb{H}^{1+\delta_0, q_0}]_{1-\frac{\kappa_{0,c}}{p_0}} = \mathbb{H}^{\gamma_0, q_0}$  and that  $\operatorname{div} v = 0$  on  $[\varepsilon, \sigma]$ . Therefore, the required result follows from Lemma 4.4.

*Step 2: Proof of (1)-(2).*

(1): Let  $q_1 > q_0$ . Since  $B_{q_1, \infty}^{\frac{d}{q_0}-1}(\mathbb{T}^d; \mathbb{R}^d) \hookrightarrow B_{q_2, \infty}^{\frac{d}{q_0}-1}(\mathbb{T}^d; \mathbb{R}^d)$  for  $q_2 < q_1$ , replacing  $q_1$  by a smaller number if necessary, we may assume  $q_1 \in (q_0, \frac{d}{1+\delta})$ . Pick  $p_1 \in (2, \infty)$  so large that

$$\frac{2}{p_1} + \frac{d}{q_1} - 1 < \frac{d}{q_0} - 1.$$

Note that  $B_{q_1, \infty}^{d/q_0-1} \hookrightarrow H^{\gamma_1, q_1}$  where  $\gamma_1 = \frac{2}{p_1} + \frac{d}{q_1} - 1$  and therefore

$$\mathbb{P}\left(\varepsilon < \sigma < T, \sup_{t \in [\varepsilon, \sigma]} \|u(t)\|_{B_{q_1, \infty}^{\frac{d}{q_0}-1}} < \infty\right) \leq \mathbb{P}\left(\varepsilon < \sigma < T, \|u\|_{L^{p_1}(\varepsilon, \sigma; H^{\gamma_1, q_1})} < \infty\right) = 0,$$

where in the last equality we apply (2.17) with  $(p_0, q_0, \gamma_0)$  replaced by  $(p_1, q_1, \gamma_1)$ .

(2): Follows from (2.17).  $\square$

**4.4. Global well-posedness, and Theorems 2.11 and 2.12.** Below we use the notation (4.1). Also recall from Lemma 4.2 that  $\beta = \frac{1}{2}(1 + \frac{\delta}{2} - \frac{d}{2q})$  and  $X_\beta = \mathbb{H}^{\frac{d}{2q} + \frac{\delta}{2}, q}$ .

We first deduce Theorem 2.12 from the blow-up criterium (2.17) in Theorem 2.9 and the energy estimates provided by Theorem A.3.

*Proof of Theorem 2.12.* Let  $(u, \sigma)$  be the solution provided by Theorem 2.4. By combining Lemma 4.4 with  $p_0 = q_0 = 2$ ,  $\delta_0 = 0$  and Theorem A.3(2) from the appendix, we obtain that for all  $\varepsilon > 0$ ,

$$(4.29) \quad u \in L^2(\varepsilon, \sigma; H^{1,2}(\mathbb{T}^2; \mathbb{R}^2)) \quad \text{a.s. on } \{\sigma > \varepsilon\}.$$

Thus for all  $0 < \varepsilon < T < \infty$ ,

$$(4.30) \quad \mathbb{P}(\varepsilon < \sigma < T) \stackrel{(4.29)}{=} \mathbb{P}\left(\varepsilon < \sigma < T, \|u\|_{L^2(\varepsilon, \sigma; H^{1,2}(\mathbb{T}^2; \mathbb{R}^2))} < \infty\right) = 0$$

where the last equality follows from (2.17) of Theorem 2.9 with  $p_0 = q_0 = 2$  and  $\gamma_0 = 1$ . Since  $\sigma > 0$  a.s. by Theorem 2.4, by (4.30) we obtain  $\mathbb{P}(\sigma < \infty) = 0$ .

The remaining assertions follow from Theorems 2.4 and 2.7.  $\square$

To prove Theorem 2.11 we need a suitable  $L^p$ -space which allows us to bound the nonlinearities in (2.1). For each  $t \in (0, \infty)$  it is defined as

$$(4.31) \quad \mathcal{X}(t) := L^{2p}(0, t, w_{\kappa_c}; X_\beta), \quad \text{where } \kappa_c = -1 + \frac{p}{2} \left( 2 + \delta - \frac{d}{q} \right).$$

By Lemma 4.2 the above space coincide with the abstract one introduced in [AV21a, Subsection 4.3], where we also showed

$$\bigcap_{\theta \in [0, 1/2)} H^{\theta, p}(0, t; w_{\kappa_c}; X_{1-\theta}) \subseteq \mathcal{X}(t).$$

In particular, the solution  $(u, \sigma)$  provided by Theorem 2.4 satisfies a.s. for all  $t \in (0, \sigma)$ ,  $u \in \mathcal{X}(t)$ .

Arguing as in Lemma 4.2 and using (2.18) one can check that we can find a constant  $C_0$  such that for all  $v \in X_\beta$

$$N(v) := \|F(\cdot, v)\|_{X_0} + \|G(\cdot, v)\|_{\gamma(\ell^2, X_{1/2})} \leq C_0 M_1 (1 + \|v\|_{X_\beta}) + C_0 \|v\|_{X_\beta}^2$$

Therefore, by Hölder's inequality there exists a constant  $C_1$  (also depending on  $T, p, \kappa_c$ ) such that for all  $t \in (0, T)$  and  $v \in \mathcal{X}(t)$  a.s.

$$(4.32) \quad \|N(v)\|_{L^p(0, t, w_{\kappa_c})}^p \leq C_1 M_1^p (1 + \|v\|_{\mathcal{X}(t)}^p) + C_1 \|v\|_{\mathcal{X}(t)}^{2p}.$$

This estimate motivates once more the definition of  $\mathcal{X}(t)$  and plays a crucial role in the analysis below.

The next result is a special case of [AV21b, Lemma 5.3], and provides an a priori estimate for the linear part of the equation based on maximal  $L^p$ -regularity.

**Lemma 4.5.** *Let Assumption 2.1( $p, \kappa_c, q, \delta$ ) be satisfied. Fix  $T \in (0, \infty)$ . Let  $A$  and  $B$  be as in (4.1). Then there exists  $K > 0$  depending only on  $p, q, \delta$  and the constant  $C_2$  in Theorem 3.2 such that the following holds:*

*For any stopping time  $\tau : \Omega \rightarrow [0, T]$ , and any*

$$v_0 \in L^0_{\mathcal{F}_0}(\Omega; X_{\kappa_c, p}^{\text{Tr}}), f \in L^p_{\mathcal{F}}(\llbracket 0, \tau \rrbracket, w_{\kappa_c}; X_0), g \in L^p_{\mathcal{F}}(\llbracket 0, \tau \rrbracket, w_{\kappa_c}; \gamma(\ell^2, X_{1/2})),$$

*and any  $(p, \kappa_c, q, \delta)$ -solution  $v \in L^p_{\mathcal{F}}(\llbracket 0, \tau \rrbracket, w_{\kappa_c}; X_1)$  to*

$$\begin{cases} dv + Av dt = f dt + (Bv + g) dW_{\ell^2}, \\ v(0) = v_0, \end{cases}$$

*on  $\llbracket 0, \tau \rrbracket$  satisfies the following estimate*

$$\|v\|_{L^p(\Omega; \mathcal{X}(\tau))}^p \leq K^p (\|v_0\|_{L^p(\Omega; X_{\kappa_c, p}^{\text{Tr}})}^p + \|f\|_{L^p(\llbracket 0, \tau \rrbracket, w_{\kappa_c}; X_0)}^p + \|g\|_{L^p(\llbracket 0, \tau \rrbracket, w_{\kappa_c}; \gamma(\ell^2, X_{1/2}))}^p).$$

Next we turn to the proof of Theorem 2.11.

*Proof of Theorem 2.11.* Throughout the proof we fix  $T, \varepsilon \in (0, \infty)$ . To obtain theorem we claim that it is sufficient to prove the existence of  $C_{\varepsilon, T}, r_{\varepsilon, T} > 0$  independent of  $u$  and  $u_0$  such that

$$(4.33) \quad \mathbb{E} \|u_0\|_{B_{1,p}^{\frac{d}{q}-1}(\mathbb{T}^d; \mathbb{R}^d)}^p + M_1^p \leq C_{\varepsilon, T} \implies \mathbb{P}(\mathcal{O}) > 1 - \varepsilon.$$

where  $\mathcal{O} = \{\|u\|_{\mathcal{X}(\sigma \wedge T)} \leq r_{\varepsilon, T}\}$ . In Step 1 we will show that the theorem follows from (4.33) by applying the Serrin criterium (2.17) of Theorem 2.9.

*Step 1:* (4.33) implies the claim in Theorem 2.11. By (4.32) on  $\mathcal{O}$  we find

$$\|N(u)\|_{L^p(0, T, w_{\kappa_c})}^p \leq C_1 r_{\varepsilon, T} + C_1 =: C_2.$$

Define the stopping time  $\tau$  by  $\tau = \inf\{t \in [0, \sigma) : \|N(u)\|_{L^p(0,t,w_{\kappa_c})}^p \geq C_2 + 1\}$ , where we set  $\inf \emptyset = \sigma$ . Then  $\tau = \sigma$  on  $\mathcal{O}$ .

By Theorem 3.2,  $(A, B)$  has stochastic maximal  $L^p$ -regularity. Since  $(u, \sigma)$  is a  $(p, \kappa_c, q, \delta)$ -solution to (2.1), as in [AV21a, Proposition 3.12(2)] it follows that a.s. on  $[0, \tau)$

$$du + Audt = \mathbf{1}_{[0,\tau)} F(\cdot, u)dt + (Bu + \mathbf{1}_{[0,\tau)} G(\cdot, u))dW_{\ell^2}$$

and  $u(0) = u_0$ . Now Theorem 3.2 with  $\theta = \frac{\kappa_c}{p}$  gives  $u \in L^p(\Omega; H^{\frac{\kappa_c}{p}, p}(0, \tau, w_{\kappa_c}; X_{1-\frac{\kappa_c}{p}}))$ .

Using  $\tau = \sigma$  on  $\mathcal{O}$ , by Sobolev embedding [AV21a, Proposition 2.7] we obtain

(4.34)

$$u \in H^{\frac{\kappa_c}{p}, p}(0, \sigma, w_{\kappa_c}; X_{1-\frac{\kappa_c}{p}}) \hookrightarrow L^p(0, \sigma; X_{1-\frac{\kappa_c}{p}}) = L^p(0, \sigma; \mathbb{H}^{\gamma, q}(\mathbb{T}^d)) \text{ a.s. on } \mathcal{O},$$

where  $\gamma = 1 + \delta - 2\frac{\kappa_c}{p} = \frac{2}{p} + \frac{d}{q} - 1$ . It follows that

$$\begin{aligned} \mathbb{P}(\{\sigma < T\} \cap \mathcal{O}) &\stackrel{(i)}{=} \lim_{s \downarrow 0} \mathbb{P}\left(\left\{s < \sigma < T, \|u\|_{L^p(s, \sigma; H^{\gamma, q})} < \infty\right\} \cap \mathcal{O}\right) \\ &\leq \limsup_{s \downarrow 0} \mathbb{P}\left(s < \sigma < T, \|u\|_{L^p(s, \sigma; H^{\gamma, q})} < \infty\right) \stackrel{(ii)}{=} 0. \end{aligned}$$

Here in (i) we used  $\sigma > 0$  a.s. (see Theorem 2.4) and (4.34). In (ii) we used (2.17) of Theorem 2.9 with  $p_0 = p$ ,  $q_0 = q$  and  $\gamma_0 = \gamma$ . Therefore,  $\sigma \geq T$  on  $\mathcal{O}$  and this gives the result of the theorem, and thus completes Step 1.

Next we turn to the proof of (4.33). Before we continue it is important to recall that from the discussion below (4.31) a.s. for all  $t \in (0, \sigma)$ ,  $u \in \mathcal{X}^c(t)$ .

*Step 2: Let  $K$  be as in Lemma 4.5. Let  $M_1$  and  $C_1$  be as in (4.32). Suppose that  $K^p C_1 M_1^p \leq \frac{1}{2}$ . Then there exists  $R > 0$  depending only on  $K$  and  $C_1$  such that for any  $N \geq 1$  and any stopping time  $\mu$  satisfying  $0 \leq \mu \leq \sigma$  and  $\|u\|_{\mathcal{X}(\mu)} \leq N$  a.s.,*

$$\mathbb{E}[\psi_R(\|u\|_{\mathcal{X}(\mu)}^p)] \leq \mathbb{E}\|u_0\|_{B_{q,p}^{d/q-1}}^p + M_1^p, \quad \text{with } \psi_R(x) = \frac{x}{R} - x^2.$$

In order to prove the assertion in Step 2 we use a localization argument. Set

$$\sigma_n := \inf\{t \in [0, \sigma) : \|u\|_{L^p(0,t,w_{\kappa_c}; X_1)} + \|u\|_{\mathcal{X}(t)} \geq n\}, \quad n \geq 1,$$

where we set  $\inf \emptyset = \sigma$ , and let  $\mu_n := \mu \wedge \sigma_n$ . Then  $\lim_{n \rightarrow \infty} \sigma_n = \sigma$  a.s.,  $\lim_{n \rightarrow \infty} \mu_n = \mu$ , and for all  $n \geq 1$ ,

$$\|u\|_{\mathcal{X}(\sigma_n)} + \|u\|_{L^p(0, \sigma_n, w_{\kappa_c}; X_1)} \leq n.$$

As in Step 1 it follows from Lemma 4.5 and (4.32) that

$$\begin{aligned} \mathbb{E}\|u\|_{\mathcal{X}(\mu_n)}^p &\leq K^p (\mathbb{E}\|u_0\|_{B_{q,p}^{d/q-1}}^p + \mathbb{E}\|F(\cdot, u)\|_{L^p([0, \mu_n], w_c; X_0)}^p \\ &\quad + \mathbb{E}\|G(\cdot, u)\|_{L^p([0, \mu_n], w_c; X_{1/2})}^p) \\ &\leq K^p \|u_0\|_{B_{q,p}^{d/q-1}}^p + K^p C_1 M_1^p (1 + \mathbb{E}\|u\|_{\mathcal{X}(\mu_n)}^p) + K^p C_1 \mathbb{E}\|u\|_{\mathcal{X}(\mu_n)}^{2p} \end{aligned}$$

Since  $K^p C_1 M_1^p \leq \frac{1}{2}$  we obtain

$$\mathbb{E}\|u\|_{\mathcal{X}(\mu_n)}^p \leq 2K^p \|u_0\|_{B_{q,p}^{d/q-1}}^p + 2C_1 M_1^p + 2K^p C_1 \mathbb{E}\|u\|_{\mathcal{X}(\mu_n)}^{2p}.$$

Letting  $n \rightarrow \infty$ , the desired estimate follows after division by  $R = 2K^p \max\{1, C_1\}$ .

*Step 3:* (4.33) holds. Let  $\psi_R$  be as in Step 2. It is easy to check that  $\psi_R$  has a unique maximum on  $\mathbb{R}_+$  given by  $\frac{1}{4R^2}$  which is attained at  $x = \frac{1}{2R}$ . Set  $r_{\varepsilon,T} = (2R)^{-1/p}$  and thus  $\mathcal{O} = \{\|u\|_{\mathcal{X}(\sigma)} \leq (2R)^{-1/p}\}$  and let

$$(4.35) \quad \mu := \inf\{t \in [0, \sigma) : \|u\|_{\mathcal{X}(t)} \geq (2R)^{-1/p}\},$$

where  $\inf \emptyset := \sigma$ . We prove (4.33) with  $C_{\varepsilon,T} = \frac{\varepsilon}{4R^2}$ . To derive a contradiction suppose that

$$(4.36) \quad \mathbb{E}\|u_0\|_{B_{q,p}^{d/q-1}}^p + M_1^p \leq \frac{\varepsilon}{4R^2}, \quad \text{and} \quad \mathbb{P}(\mathcal{O}) \leq 1 - \varepsilon.$$

From the definition of  $\mathcal{O}$  and (4.35), we find that  $\mu < \sigma$  a.s. on  $\Omega \setminus \mathcal{O}$ . Moreover,

$$(4.37) \quad \psi_R(\|u\|_{\mathcal{X}(\mu)}^p) = \frac{1}{4R^2} \quad \text{a.s. on } \Omega \setminus \mathcal{O}, \quad \text{and} \quad \psi_R(\|u\|_{\mathcal{X}(\mu)}^p) \geq 0, \quad \text{a.s. on } \mathcal{O}.$$

Therefore,

$$\begin{aligned} \mathbb{E}\left[\psi_R(\|u\|_{\mathcal{X}(\mu)}^p)\right] &= \mathbb{E}\left[\psi_R(\|u\|_{\mathcal{X}(\mu)}^p)\mathbf{1}_{\mathcal{O}}\right] + \mathbb{E}\left[\psi_R(\|u\|_{\mathcal{X}(\mu)}^p)\mathbf{1}_{\Omega \setminus \mathcal{O}}\right] \\ &\stackrel{(4.37)}{\geq} \mathbb{P}(\Omega \setminus \mathcal{O}) \frac{1}{4R^2} \stackrel{(4.36)}{>} \frac{\varepsilon}{4R^2} \stackrel{(4.36)}{\geq} \mathbb{E}\|u_0\|_{B_{q,p}^{d/q-1}}^p + M_1^p. \end{aligned}$$

The latter contradicts Step 2. Thus  $\mathbb{P}(\mathcal{O}) > 1 - \varepsilon$  as desired.  $\square$

Additionally, we can prove an  $L^p$ -estimate for the solution.

**Corollary 4.6.** *Suppose that the conditions of Theorem 2.11 hold. Let  $C_{\varepsilon,T}$  be as in Theorem 2.11. If*

$$\mathbb{E}\|u_0\|_{B_{q,p}^{d/q-1}(\mathbb{T}^d; \mathbb{R}^d)}^p + M_1^p \leq C_{\varepsilon,T},$$

*then there exists a stopping time  $\tau \leq \sigma$  such that  $\mathbb{P}(\tau \geq T) > 1 - \varepsilon$  and*

$$\mathbb{E}\|u\|_{H^{\theta,p}(0,\tau,w_{\kappa_c}; \mathbb{H}^{1+\delta-2\theta,q})}^p \lesssim_{\theta} \mathbb{E}\|u_0\|_{B_{q,p}^{d/q-1}(\mathbb{T}^d; \mathbb{R}^d)}^p + M_1^p \quad \text{for all } \theta \in [0, 1/2),$$

*where in case  $p = q = d = 2$ , the LHS should be replaced by  $\mathbb{E}\|u\|_{C([0,\tau]; \mathbb{L}^2(\mathbb{T}^d))}^p$ .*

*Proof.* We use the same notation as in the proof of Theorem 2.11. Let  $\tau$  be stopping time defined by the following variation of (4.35):

$$\tau := \inf\{t \in [0, \sigma) : \|u\|_{\mathcal{X}(t)} \geq (2R)^{-1/p} + 1\},$$

where  $\inf \emptyset := \sigma$ . Then from Step 3 of the previous proof one can check that  $\tau = \sigma$  on  $\mathcal{O}$ , and thus

$$\mathbb{P}(\tau \geq T) \geq \mathbb{P}(\{\tau \geq T\} \cap \mathcal{O}) = \mathbb{P}(\{\sigma \geq T\} \cap \mathcal{O}) = \mathbb{P}(\sigma \geq T) > 1 - \varepsilon.$$

Also note that  $\|u\|_{\mathcal{X}(\tau)}^p \leq (2R)^{-1} + 1$ . Therefore, by (4.32), we get

$$\begin{aligned} \mathbb{E}\|F(\cdot, u)\|_{L^p(0,\tau,w_{\kappa_c}; X_0)}^p + \mathbb{E}\|G(\cdot, u)\|_{L^p(0,\tau,w_{\kappa_c}; \gamma(\ell^2, X_{1/2}))}^p \\ \leq C_1 M_1^p (1 + L) + C_1 L^2 < \infty \end{aligned}$$

Thus the result follows from Theorem 3.2.  $\square$

APPENDIX A. GLOBAL WELL-POSEDNESS FOR  $d = 2$  WITH  $L^\infty$ -NOISE

The aim of this appendix is to prove global well-posedness for the stochastic Navier-Stokes equations (2.1) in two dimensions with  $L^2$ -data and  $L^\infty$ -noise of gradient type. Although such results are essentially known, our approach based on maximal  $L^2$ -regularity techniques seems new and provides an improvement of several of the known results. We refer to Remark A.4 below for a comparison with other approaches.

The results presented in this appendix are used to prove Theorems 2.12 and will be presented more or less independently of the proofs given in the paper.

Here we consider the following stochastic Navier-Stokes equations on a domain  $\mathcal{O} \subseteq \mathbb{R}^d$

$$(A.1) \quad \begin{cases} du = [\operatorname{div}(a \cdot \nabla u) - \nabla p - \operatorname{div}(u \otimes u) + f_0(\cdot, u) + \operatorname{div}(f(\cdot, u)) + \partial_h \tilde{p}] dt \\ \quad + \sum_{n \geq 1} [(b_n \cdot \nabla) u - \nabla \tilde{p}_n + g_n(\cdot, u)] dw_t^n, \\ u = 0, \quad \text{on } \partial \mathcal{O}, \\ \operatorname{div} u = 0, \\ u(\cdot, 0) = u_0, \end{cases}$$

where  $u = (u^k)_{k=1}^d : [0, \infty) \times \Omega \times \mathcal{O} \rightarrow \mathbb{R}^d$  is unknown velocity field,  $p, \tilde{p}_n : [0, \infty) \times \Omega \times \mathcal{O} \rightarrow \mathbb{R}$  are the unknown pressures,  $(w^n)_{n \geq 1}$  is a sequence of standard independent Brownian motions on a filtered probability space  $(\Omega, \mathcal{A}, (\mathcal{F}_t)_{t \geq 0}, \mathbf{P})$ ,  $(u \otimes u)_{i,j} = u^i u^j$  and  $\operatorname{div}(a \cdot \nabla u)$ ,  $(b_n \cdot \nabla) u$ ,  $\operatorname{div}(f(\cdot, u))$ ,  $\partial_h \tilde{p}$  are as in (2.2). Here  $\mathcal{O}$  can be either an open set in  $\mathbb{R}^d$  (not necessarily bounded) or a compact manifold without boundaries (e.g. the torus  $\mathbb{T}^d$ ). In the latter case the boundary conditions in (A.1) have to be omitted.

If  $\mathcal{O} \subseteq \mathbb{R}^d$ , then under suitable regularity assumptions  $\mathcal{O}$  other boundary conditions can be considered, e.g. perfect-slip or Navier boundary conditions (see Remark A.8 below). For brevity, this will not be pursued here. Let us remark that the no-slip condition allows us to avoid any further regularity assumptions on  $\mathcal{O}$ .

**A.1. Assumptions and main result.** We begin by listing the assumption needed in this appendix.

**Assumption A.1.** *Let  $d \geq 2$  be an integer.*

- (1) *For all  $n \geq 1$  and  $i, j \in \{1, \dots, d\}$ , the maps  $a^{i,j}, \phi_n^j, h_n^{j,k} : \mathbb{R}_+ \times \Omega \times \mathcal{O} \rightarrow \mathbb{R}$  are  $\mathcal{P} \otimes \mathcal{B}(\mathcal{O} \times \mathbb{R}^d)$ -measurable and there exist  $M, \nu > 0$  such that a.s. for all  $t \in \mathbb{R}_+$ ,  $x \in \mathcal{O}$  and  $\Upsilon \in \mathbb{R}^d$ ,*

$$|a^{i,j}(t, x)| + \sum_{n \geq 1} \left( |\phi_n^j(t, x)| + |h_n^{i,j}(t, x)| \right) \leq M,$$

$$\sum_{i,j=1}^d \left[ a^{i,j}(t, x) - \frac{1}{2} \left( \sum_{n \geq 1} b_n^i(t, x) b_n^j(t, x) \right) \right] \Upsilon_i \Upsilon_j \geq \nu |\Upsilon|^2.$$

- (2) *For all  $j \in \{0, \dots, d\}$  and  $n \geq 1$ ,  $f_j, g_n : \mathbb{R}_+ \times \Omega \times \mathcal{O} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  are  $\mathcal{P} \otimes \mathcal{B}(\mathcal{O} \times \mathbb{R}^d)$ -measurable. Moreover,  $f_j(\cdot, 0) \in L_{\text{loc}}^2([0, \infty); L^2(\Omega \times \mathcal{O}))$ ,  $(g_n(\cdot, 0))_{n \geq 1} \in L_{\text{loc}}^2([0, \infty); L^2(\Omega \times \mathcal{O}; \ell^2))$  and, a.s. for all  $t \in \mathbb{R}_+$ ,  $x \in \mathbb{T}^2$ ,  $y, y' \in \mathbb{R}^2$ ,*

$$|f_j(t, x, y) - f_j(t, x, y')| + \|(g_n(t, x, y) - g_n(t, x, y'))_{n \geq 1}\|_{\ell^2} \lesssim (c_{\mathcal{O}} + |y| + |y'|)|y - y'|,$$

where  $c_{\mathcal{O}} = 0$  if  $\mathcal{O}$  is unbounded and  $c_{\mathcal{O}} = 1$  otherwise.

To formulate our main result we introduce the needed function spaces. Set

$$H_0^1(\mathcal{O}) := \overline{C_0^\infty(\mathcal{O})}^{H^1(\mathcal{O})}, \quad \text{and} \quad H_0^1(\mathcal{O}; \mathbb{R}^d) := (H_0^1(\mathcal{O}))^d.$$

Next we introduce the Helmholtz projection. For each  $F \in L^2(\mathcal{O}; \mathbb{R}^d)$ , let  $\psi_F$  be the unique distribution such that  $\nabla \psi_F \in L^2(\mathcal{O}; \mathbb{R}^d)$  and

$$(A.2) \quad \int_{\mathcal{O}} \nabla \psi_F \cdot \nabla \varphi \, dx = \int_{\mathcal{O}} F \cdot \nabla \varphi \, dx, \quad \text{for all } \varphi \in H^1(\mathcal{O}).$$

The existence of a unique solution  $\psi_F$  to the above follows from the Riesz representation theorem. Note that (A.2) is the weak formulation of

$$\begin{cases} \Delta \psi_F = \operatorname{div} F, & \text{on } \mathcal{O}, \\ \partial_\nu \psi_F = 0, & \text{on } \partial \mathcal{O}. \end{cases}$$

The Helmholtz projection is given by  $\mathbb{P}F := F - \nabla \psi_F$ . Using (A.2), one sees that  $\mathbb{P}$  is an orthonormal projection on  $L^2(\mathcal{O}; \mathbb{R}^d)$ . Finally, we let

$$\mathbb{L}^2(\mathcal{O}) := \mathbb{P}(L^2(\mathcal{O}; \mathbb{R}^d)), \quad \mathbb{H}_0^1(\mathcal{O}) := H_0^1(\mathcal{O}; \mathbb{R}^d) \cap \mathbb{L}^2(\mathcal{O}), \quad \mathbb{H}^{-1}(\mathcal{O}) := (\mathbb{H}_0^1(\mathcal{O}))^*.$$

Next we introduce  $L^2$ -solutions to (2.1) which, roughly speaking, are *strong* in the probabilistic sense and *weak* in the analytic sense. As in Subsection (2.2), we view (A.1) as a problem for the unknown  $u$ . To this end, for  $h_n^{i,j} = (h_n^{i,j})_{j=1}^d$ , let us set

$$\tilde{f}(u) := (\tilde{f}^j(u))_{j=1}^d := \left( \sum_{n \geq 1} \left[ (I - \mathbb{P})[(b_n \cdot \nabla)u + g_n(\cdot, u)] \right] \cdot h_n^{i,j} \right)_{j=1}^d.$$

**Definition A.2.** *Let Assumption A.1 be satisfied.*

(1) *Let  $\sigma$  be a stopping time and let  $u : [0, \sigma) \times \Omega \rightarrow \mathbb{H}_0^1(\mathcal{O})$  be a stochastic process.*

*We say that  $(u, \sigma)$  is an  $L^2$ -local solution to (A.1) if there exists a sequence of stopping times  $(\sigma_\ell)_{\ell \geq 1}$  for which the following hold:*

- $\sigma_\ell \leq \sigma$  a.s. for all  $\ell \geq 1$  and  $\lim_{\ell \rightarrow \infty} \sigma_\ell = \sigma$  a.s.;
- For all  $\ell \geq 1$ , the process  $\mathbf{1}_{[0, \sigma_\ell] \times \Omega} u$  is progressively measurable;
- $u \in L^2(0, \sigma_\ell; \mathbb{H}_0^1(\mathcal{O})) \cap C([0, \sigma_\ell]; L^2(\mathcal{O}))$  a.s. and for all  $i, j \in \{1, \dots, d\}$

$$(A.3) \quad \begin{aligned} f_0^i(u) + f_j^i(u) + \tilde{f}^i(u) - u^i u^j &\in L^2(0, \sigma_\ell; L^2(\mathcal{O}; \mathbb{R}^d)), \quad \text{a.s.} \\ ((g_n^j(\cdot, u))_{n \geq 1}) &\in L^2(0, \sigma_\ell; L^2(\mathcal{O}; \ell^2)), \quad \text{a.s.} \end{aligned}$$

- a.s. for all  $n \geq 1$ ,  $t \in [0, \sigma_\ell]$  and  $\psi \in \mathbb{H}_0^1(\mathcal{O})$  the following identity holds

$$(A.4) \quad \begin{aligned} &\int_{\mathcal{O}} (u^k(t) - u_0^k) \psi_k \, dx + \int_0^t \int_{\mathcal{O}} a^{i,j} \partial_j u^k \partial_i \psi_k \, dx ds \\ &= \int_0^t \int_{\mathcal{O}} (u^j u^k + f_j^k(u)) \partial_j \psi_k \, dx ds + \int_0^t \int_{\mathcal{O}} (f_0^k(u) + \tilde{f}^k(u)) \psi_k \, dx ds \\ &+ \sum_{n \geq 1} \int_0^t \mathbf{1}_{[0, \sigma_n]} \int_{\mathcal{O}} (b_n^j \partial_j u^k + g_n^k(\cdot, u)) \psi_k \, dx dw_s^n, \end{aligned}$$

where we use the Einstein summation convention for the sums over  $i, j, k \in \{0, 1, \dots, d\}$ .

(2) *An  $L^2$ -local solution  $(u, \sigma)$  to (A.1) is called an  $L^2$ -maximal solution if for any other local solution  $(v, \tau)$  to (A.1) one has  $\tau \leq \sigma$  a.s. and  $v = u$  on  $[0, \tau) \times \Omega$ .*

Note that  $L^2$ -maximal solutions are *unique*. Moreover, (A.3) ensures that all the integrals and series appearing in (A.4) are well-defined and convergent.

Next we present the main result of this appendix.

**Theorem A.3** (Global solutions in 2D). *Assume that  $d = 2$  and that  $\mathcal{O} \subseteq \mathbb{R}^2$  is an open set, or a compact  $d$ -dimensional manifold without boundary (e.g. the torus  $\mathbb{T}^d$ ). Let Assumption A.1 be satisfied. Then for each  $u_0 \in L^0_{\mathcal{F}_0}(\Omega; \mathbb{L}^2(\mathcal{O}))$  there exists an  $L^2$ -maximal solution  $(u, \sigma)$  to (A.1) such that  $\sigma > 0$  a.s. and*

$$(A.5) \quad u \in C([0, \sigma]; \mathbb{L}^2(\mathcal{O})) \cap L^2_{\text{loc}}([0, \sigma]; \mathbb{H}_0^1(\mathcal{O})) \quad \text{a.s.}$$

Moreover, if there exist  $\Xi \in L^2_{\text{loc}}([0, \infty); L^2(\Omega \times \mathcal{O}))$  and  $C > 0$  such that a.s. for all  $j \in \{0, 1, 2\}$ ,  $t \in \mathbb{R}_+$ ,  $x \in \mathcal{O}$  and  $y \in \mathbb{R}$ ,

$$(A.6) \quad |f_j(t, x, y)| + \|(g_n(t, x, y))_{n \geq 1}\|_{\ell^2} \leq \Xi(t, x) + C|y|,$$

then the following hold:

- (1) the  $L^2$ -maximal solution  $(u, \sigma)$  to (A.1) is global in time, i.e.  $\sigma = \infty$  a.s.;
- (2) if  $u_0 \in L^2(\Omega; \mathbb{L}^2(\mathcal{O}))$ , then for each  $T \in (0, \infty)$  there exists  $C_T > 0$  independent of  $u, u_0$  such that

$$\mathbb{E} \left[ \sup_{t \in [0, T]} \|u(t)\|_{L^2(\mathcal{O})}^2 \right] + \mathbb{E} \int_0^T \|\nabla u(t)\|_{L^2(\mathcal{O})}^2 dt \leq C_T(1 + \mathbb{E}\|u_0\|_{L^2(\mathcal{O})}^2).$$

The proof of Theorem A.3 will be given in Subsection A.3 below. Note that under further regularity conditions on the coefficients  $a^{ij}$ ,  $b^j$  and  $h^{ij}$ , the above result can be improved and this is the content of Theorem 2.12 combined with Theorems 2.4 and 2.7. In the following remark we give a comparison with the existing results.

*Remark A.4.* The existence of  $L^2$ -maximal solutions to (A.1) with nonlinearities of quadratic growth (see Assumption A.1(2)) seems new. (2) should be compared with [MR05, Theorem 2.2] if  $\mathcal{O} = \mathbb{R}^d$  and [BM13, Section 6 and Corollary 7.7] if  $\mathcal{O}$  is an unbounded domain. Some earlier related results appeared in [BCF91, BCF92] using some smoothness of  $b$ . Unlike in the above references we do not need any regularity assumption on  $b$  besides boundedness and measurability. The regularity assumption on  $b$  are needed in [MR05, BM13] to construct martingale solutions to (A.1) for general  $d \geq 2$ . The case  $d = 2$  is only considered afterwards.

In case  $f$  and  $g$  are globally Lipschitz, Theorem A.3 also follows from the monotone operator approach (see e.g. [LR15, Chapter 4]). The extensions to the non-Lipschitz setting in [LR15, Chapter 5] are not applicable since gradient noise is not considered there.

Below, we write  $L^2$ ,  $\mathbb{H}_0^1$ ,  $L^2(\ell^2)$  instead of  $L^2(\mathcal{O})$ ,  $\mathbb{H}_0^1(\mathcal{O})$ ,  $L^2(\mathcal{O}; \ell^2)$  etc. for brevity.

**A.2. Maximal  $L^2$ -regularity for the turbulent Stokes system.** In this subsection we consider the linear part of (A.1) and prove stochastic maximal  $L^2$ -regularity estimates which will be used for Theorem A.3. We consider the *turbulent Stokes system* on  $\mathcal{O} \subseteq \mathbb{R}^d$ :

$$(A.7) \quad \begin{cases} du + A_{\mathbb{S}}u dt = f dt + \sum_{n \geq 1} (B_{\mathbb{S}, n}u + g_n) dw_t^n, \\ u(s) = 0. \end{cases}$$

Here  $s \in [0, \infty)$ ,  $(A_{\mathbb{S}}, B_{\mathbb{S}}) = (A_{\mathbb{S}}, (B_{\mathbb{S},n})_{n \geq 1}) : \mathbb{R}_+ \times \Omega \rightarrow \mathcal{L}(\mathbb{H}_0^1, \mathbb{H}^{-1} \times \mathbb{L}^2(\ell^2))$  is the *turbulent Stokes couple*, which for all  $v \in \mathbb{H}_0^1$  and  $\psi \in \mathbb{H}_0^1$ , is given by

$$\begin{aligned} \langle \psi, A_{\mathbb{S}}v \rangle &:= \sum_{i,j=1}^d \int_{\mathcal{O}} a^{i,j} \partial_j u^k \partial_i \psi_k dx - \sum_{n \geq 1} \sum_{j=1}^d \int_{\mathcal{O}} \left[ (I - \mathbb{P})[(b_n \cdot \nabla)u] \cdot h_n^{i,j} \psi_j dx, \right. \\ B_{\mathbb{S}}(t)v &:= \left( \mathbb{P}[(b_n \cdot \nabla)v] \right)_{n \geq 1}, \quad h_n^{i,j} := (h_n^{i,j})_{i=1}^d, \end{aligned}$$

where  $\langle \cdot, \cdot \rangle$  is the pairing in the duality  $(\mathbb{H}_0^1)^* = \mathbb{H}^{-1}$  and  $\mathbb{L}^2(\ell^2) := \mathbb{P}(L^2(\mathcal{O}; \ell^2(\mathbb{N}; \mathbb{R}^d)))$ . Note that if Assumption A.1(1) holds,

$$\|A_{\mathbb{S}}v\|_{\mathbb{H}^{-1}} + \|(B_{\mathbb{S},n}v)_{n \geq 1}\|_{\mathbb{L}^2(\ell^2)} \lesssim M \|v\|_{\mathbb{H}_0^1}.$$

Recall that  $\mathcal{P}$  denotes the progressive  $\sigma$ -algebra. For a stopping time  $\tau$ , we say that  $u \in L^2_{\mathcal{P}}(\llbracket 0, \tau \rrbracket; \mathbb{H}_0^1)$  is a *strong solution* to (A.7) on  $\llbracket 0, \tau \rrbracket$  if a.s. for all  $t \in [s, \tau]$

$$u(t) + \int_s^t A_{\mathbb{S}}(r)u(r) dr = \int_s^t f(r) dr + \sum_{n \geq 1} \int_s^t (B_{\mathbb{S},n}(r)u(r) + g_n(r)) dw_r^n.$$

Next we show that the turbulent Stokes couple  $(A_{\mathbb{S}}, B_{\mathbb{S}})$  has stochastic maximal  $L^2$ -regularity on  $[s, T]$ .

**Proposition A.5** (Maximal  $L^2$ -regularity). *Let  $d \geq 2$ . Let Assumption A.1(1) be satisfied. Assume that  $\mathcal{O} \subseteq \mathbb{R}^d$  is an open set, or a compact  $d$ -dimensional manifold without boundary (e.g. the torus  $\mathbb{T}^d$ ). Let  $0 \leq s < T < \infty$ . Then for all*

$$f \in L^2_{\mathcal{P}}(\llbracket s, T \rrbracket; \mathbb{H}^{-1}), \quad \text{and} \quad g \in L^2_{\mathcal{P}}(\llbracket s, T \rrbracket; \mathbb{L}^2(\ell^2)),$$

*there exists a unique strong solution  $u \in L^2_{\mathcal{P}}(\llbracket s, T \rrbracket; \mathbb{H}_0^1) \cap L^2(\Omega; C(\llbracket s, T \rrbracket; \mathbb{L}^2))$ . Moreover, for any stopping time  $\tau : \Omega \rightarrow [s, T]$  and any strong solution  $u$  to (A.7) on  $\llbracket s, \tau \rrbracket$  one has*

$$\|u\|_{L^2(\llbracket s, \tau \rrbracket; \mathbb{H}_0^1)} + \|u\|_{L^2(\Omega; C(\llbracket s, \tau \rrbracket; \mathbb{L}^2))} \lesssim \|f\|_{L^2(\llbracket s, \tau \rrbracket; \mathbb{H}^{-1})} + \|g\|_{L^2(\llbracket s, \tau \rrbracket; \mathbb{L}^2(\ell^2))}$$

*where the implicit constant is independent of  $(f, g)$ .*

*Proof.* This follows from the monotone operator approach to SPDEs (see [LR15, Theorem 4.2.4]). Indeed, it suffices to check the coercivity assumption: a.s. for all  $t \in \mathbb{R}_+$  and  $v \in \mathbb{H}_0^1$ ,

$$\begin{aligned} & - \langle v, A_{\mathbb{S}}v \rangle + \frac{1}{2} \sum_{n \geq 1} \|B_{\mathbb{S},n}v\|_{L^2}^2 \\ & \stackrel{(i)}{\leq} - \sum_{i,j=1}^d \int_{\mathcal{O}} a^{i,j} \partial_i v \cdot \partial_j v dx + \frac{1}{2} \sum_{n \geq 1} \|(b_n \cdot \nabla)v\|_{L^2}^2 + C \|h(t)\|_{L^\infty(\ell^2)} \|\nabla v\|_{L^2} \|v\|_{L^2} \\ & \stackrel{(ii)}{\leq} -\nu \|\nabla v\|_{L^2}^2 + \frac{\nu}{2} \|\nabla v\|_{L^2}^2 + C_\nu M \|v\|_{L^2}^2 = -\frac{\nu}{2} \|\nabla v\|_{L^2}^2 + C_\nu M \|v\|_{L^2}^2 \end{aligned}$$

where in (i) we used  $\|\mathbb{P}\|_{\mathcal{L}(L^2)} = 1$  and in (ii) we used Assumption A.1(1).  $\square$

We conclude by pointing out another proof of the above result.

*Remark A.6.* Instead of the monotone operator approach one can also prove Proposition A.5 via the method of continuity for stochastic maximal regularity (see [AV21b, Proposition 3.13]). Indeed, one can reason as in the proof of Theorem 3.2 and the required a priori estimate can be obtained from Itô's formula for  $u \mapsto \|u\|_{L^2}^2$ .



**A.3. Proof of Theorem A.3.** To estimate the nonlinearities in (A.1) we need the following consequence of Sobolev embedding. Here  $[\cdot, \cdot]_\theta$  denotes the complex interpolation functor, see e.g. [BL76].

**Lemma A.7** (Sobolev embeddings). *Let  $\theta \in (0, 1)$  and  $q \in [2, \infty)$  be such that  $\theta - \frac{d}{2} \geq -\frac{d}{q}$ . Then  $[\mathbb{H}^{-1}, \mathbb{H}_0^1]_{\frac{1+\theta}{2}} \hookrightarrow L^q$ .*

*Proof.* From general theory on coercive forms (see [Are04, Section 5.5.2]) we obtain  $[\mathbb{H}^{-1}, \mathbb{H}_0^1]_{1/2} = \mathbb{L}^2$ . Therefore, by reiteration for complex interpolation (see [BL76, Theorem 4.6.1])

$$[\mathbb{H}^{-1}, \mathbb{H}_0^1]_{(1+\theta)/2} = [\mathbb{L}^2, \mathbb{H}_0^1]_\theta \hookrightarrow [L^2, H_0^1]_\theta.$$

Using the boundedness of the zero-extension operator  $E_0 : L^2(\mathcal{O}) \rightarrow L^2(\mathbb{R}^d)$  and  $E_0 : H_0^1(\mathcal{O}) \rightarrow H^1(\mathbb{R}^d)$ , and Sobolev embedding (see [BL76, Theorem 6.4.5]),

$$\|f\|_{L^q} \leq \|E_0 f\|_{L^q(\mathbb{R}^d)} \lesssim \|E_0 f\|_{[L^2(\mathbb{R}^d); H^1(\mathbb{R}^d)]_\theta} \leq \|f\|_{[L^2; H_0^1]_\theta} \leq \|f\|_{[\mathbb{L}^2; \mathbb{H}_0^1]_\theta}.$$

□

The result of Lemma A.7 holds under more general boundary conditions if there exists an extension operator (e.g.  $\mathcal{O}$  is a Lipschitz domain,  $\mathcal{O} = \mathbb{T}^d$ , or  $\mathcal{O} = \mathbb{R}_+^d$ ).

*Proof of Theorem A.3.* The proof is divided into several steps.

*Step 1: There exists an  $L^2$ -maximal solutions  $(u, \sigma)$  to (A.1) such that  $\sigma > 0$  a.s. and that (A.5) holds.* By Definition A.2 the claim of this step follows from [AV21a, Theorem 4.9] with the choice  $p = 2$ ,  $\kappa = 0$ ,  $X_0 = \mathbb{H}^{-1}$ ,  $X_1 = \mathbb{H}_0^1$ ,  $H = \ell^2$  and for  $v \in \mathbb{H}_0^1$ ,  $\psi \in \mathbb{H}_0^1$ ,

$$(A.8) \quad \begin{aligned} Av &= A_{\mathbb{S}}v, & Bv &= B_{\mathbb{S}}v, & G(v) &= (\mathbb{P}[g_n(\cdot, v)])_{n \geq 1}, \\ \langle \psi, F(\cdot, v) \rangle &= \int_{\mathcal{O}} (u^j u^i + f_j^i(u)) \partial_j \psi_i \, dx ds + \int_{\mathcal{O}} (f_0^i(u) + \tilde{f}^i(u)) \psi_i \, dx ds, \end{aligned}$$

where  $A_{\mathbb{S}}$  and  $B_{\mathbb{S}}$  are as below (A.7). Indeed, it remains to check the assumptions of [AV21a, Theorem 4.9]. The stochastic maximal  $L^2$ -regularity follows from Proposition A.5. Next we check hypothesis [AV21a, (HF)-(HG)]. To this end, note that by Assumption A.1(2) we have, for all  $v, v' \in \mathbb{H}_0^1$ ,

$$(A.9) \quad \begin{aligned} &\|F(\cdot, v) - F(\cdot, v')\|_{\mathbb{H}^{-1}} + \|G(\cdot, v) - G(\cdot, v')\|_{\mathbb{L}^2(\ell^2)} \\ &\lesssim (c_{\mathcal{O}} + \|v\|_{L^4} + \|v'\|_{L^4}) \|v - v'\|_{L^4} \\ &\lesssim (c_{\mathcal{O}} + \|v\|_{[\mathbb{H}^{-1}, \mathbb{H}_0^1]_{\frac{3}{4}}} + \|v'\|_{[\mathbb{H}^{-1}, \mathbb{H}_0^1]_{\frac{3}{4}}}) \|v - v'\|_{[\mathbb{H}^{-1}, \mathbb{H}_0^1]_{\frac{3}{4}}} \end{aligned}$$

where in the last step we used Lemma A.7 with  $d = 2$ . Thus [AV21a, (HF)-(HG)] are satisfied with  $m_G = m_F = 1$ ,  $\rho_j = 1$ ,  $\beta_j = \varphi_j = \frac{3}{4}$  where  $j \in \{1, 2\}$ . Hence the claim of this step follows from [AV21a, Theorem 4.9].

It remains to prove (1)-(2) in Theorem A.3 in case (A.6) holds. To this end, it is enough to show that if  $(u, \sigma)$  is an  $L^2$ -maximal solution to (A.1) and  $u_0 \in L^2(\Omega; L^2)$ , then for each  $T \in (0, \infty)$

$$(A.10) \quad \mathbb{E} \left[ \sup_{t \in [0, T \wedge \sigma)} \|u(t)\|_{L^2}^2 \right] + \mathbb{E} \int_0^{T \wedge \sigma} \|\nabla u(t)\|_{L^2}^2 \, dt \lesssim_T 1 + \mathbb{E} \|u_0\|_{L^2}^2.$$

Indeed, if (A.10) is true, then in Step 2 we prove that (1) holds and therefore (2) follows from (1) and (A.10).

*Step 2:* If (A.10) holds, then (1) is true. Recall that (2.1) is understood as a stochastic evolution equations on  $\mathbb{H}^{-1}$  with the choice (A.8). Thus by localization we may assume  $u_0 \in L^\infty(\Omega; \mathbb{L}^2)$  (see [AV21b, Proposition 4.13]). Hence, for all  $T$  and  $\varepsilon > 0$ ,

$$\mathbb{P}(\varepsilon < \sigma < T) \stackrel{(A.10)}{=} \mathbb{P}\left(\varepsilon < \sigma < T, \sup_{t \in [0, \sigma]} \|u(t)\|_{\mathbb{L}^2} + \|u\|_{L^2(0, \sigma; \mathbb{H}_0^1)} < \infty\right) = 0$$

where in the last equality we used [AV21b, Theorem 4.10(3)]. Since  $T, \varepsilon > 0$  are arbitrary and  $\sigma > 0$  a.s. by Step 1, the previous yields  $\sigma = \infty$  as desired.

To prove (A.10), we need a localization argument. Fix  $T \in (0, \infty)$ . For each  $j \geq 1$ , let  $\sigma_j$  be the stopping time given by

$$(A.11) \quad \sigma_j := \inf \{t \in [0, \sigma) : \|\nabla u\|_{L^2(0, t; L^2)} + \|u(t)\|_{L^2} \geq j\} \wedge T,$$

where  $\inf \emptyset := \sigma$ . By Step 1, (A.5) holds and therefore  $\lim_{j \rightarrow \infty} \sigma_j = \sigma$  a.s. Thus to obtain (A.10), by Gronwall and Fatou's lemmas, it is enough to show the existence of a constant  $C > 0$  independent of  $j, u, u_0$  such that

$$(A.12) \quad \mathbb{E}y(t) \leq C(1 + \mathbb{E}\|u_0\|_{L^2}^2) + C \int_0^t \mathbb{E}y(s) ds, \quad t \in [0, T],$$

where

$$(A.13) \quad y(t) = \sup_{r \in [0, t \wedge \sigma_j]} \|u(t)\|_{L^2}^2 + \int_0^{t \wedge \sigma_j} \|\nabla u(s)\|_{L^2}^2 ds.$$

In the remaining steps  $j \geq 1$  is fixed and we set  $\varrho := \sigma_j$ .

*Step 3:* We apply Itô's formula to obtain the identity (A.15) below. Let us begin by collecting some useful facts. Note that  $\|\nabla u\|_{L^2(0, \varrho; L^2)} + \sup_{t \in [0, \varrho]} \|u(t)\|_{L^2} \leq j$  a.s. by (A.11) (recall  $\varrho = \sigma_j$ ), and by Lemma A.7 with  $d = 2$  and standard interpolation inequalities,

$$L^2(0, t; \mathbb{H}_0^1) \cap L^\infty(0, t; \mathbb{L}^2) \hookrightarrow L^4(0, t; [\mathbb{L}^2, \mathbb{H}_0^1]_{1/2}) \hookrightarrow L^4(0, t; L^4)$$

for each  $t > 0$ . Combining the above observations with (A.9) we have

$$(A.14) \quad \mathbf{1}_{[0, \varrho]} F(\cdot, u) \in L^2((0, T) \times \Omega; \mathbb{H}^{-1}), \quad \mathbf{1}_{[0, \varrho]} G(\cdot, u) \in L^2((0, T) \times \Omega; \mathbb{L}^2(\ell^2)),$$

where  $F$  and  $G$  are as in (A.8). By (A.14), Itô's formula [LR15, Theorem 4.2.5], and Assumption A.1(1) we have, a.s. for all  $t \in [0, T]$ ,

$$\begin{aligned} & \|u_\varrho(t)\|_{L^2}^2 - \|u_0\|_{L^2}^2 + 2\nu \int_0^t \mathbf{1}_{[0, \varrho]} \|\nabla u\|_{L^2}^2 ds \\ & \leq \int_0^t \mathbf{1}_{[0, \varrho]} \left[ 2|(f_0(u), u)_{L^2}| + 2 \sum_{k=1}^2 |(f_k(u), \partial_k u)_{L^2}| + \|g(u)\|_{L^2(\ell^2)}^2 \right] ds \\ (A.15) \quad & + \sum_{n \geq 1} \int_0^t \mathbf{1}_{[0, \varrho]} ((b_n \cdot \nabla)u, \mathbb{P}g_n(u))_{L^2} ds \\ & + 2 \sum_{n \geq 1} \int_0^t \mathbf{1}_{[0, \varrho]} (g_n(u), u)_{L^2} dw_s^n + 2 \sum_{n \geq 1} \int_0^t \mathbf{1}_{[0, \varrho]} ((b_n \cdot \nabla)u, u)_{L^2} dw_s^n \\ & =: I_t + II_t + III_t + IV_t \end{aligned}$$

where  $u_\varrho := u(\cdot \wedge \varrho)$ ,  $(f, g)_{L^2} = \sum_{k=1}^2 \int_{\mathcal{O}} f^k g^k dx$  and we used the cancellation

$$(A.16) \quad \sum_{i,j=1}^2 \int_{\mathcal{O}} u^i u^j \partial_j u^i dx = 0, \quad \text{for all } u \in \mathbb{H}_0^1.$$

The above follows from  $\operatorname{div}(u) = 0$  and by integrations by parts and using that  $u \cdot \nu = 0$  on  $\partial\mathcal{O}$ , where  $\nu$  denotes the outer normal vector on  $\partial\mathcal{O}$ .

*Step 4: Proof of (A.12) with  $C$  independent of  $u, u_0, j$ .* Recall that  $\varrho := \sigma_j$ . Due to (A.6), one can readily check that there exists  $C > 0$  independent of  $j, u, u_0$  such that for all  $t \in [0, T]$ ,

$$(A.17) \quad \mathbb{E} \left[ \sup_{s \in [0, t]} |I_s| \right] \leq \frac{\nu}{2} \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|\nabla u(s)\|_{L^2}^2 ds + C \left( 1 + \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|u(s)\|_{L^2}^2 ds \right)$$

By Assumption A.1(1), we have

$$(A.18) \quad \begin{aligned} \mathbb{E} \left[ \sup_{s \in [0, t]} |III_s| \right] &\leq \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|(b_n(s) \cdot \nabla)u(s)\|_{L^2(\ell^2)} \|u(s)\|_{L^2} ds \\ &\leq M \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|\nabla u(s)\|_{L^2} \|u(s)\|_{L^2} ds \\ &\leq \frac{\nu}{2} \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|\nabla u(s)\|_{L^2}^2 ds + C_{M, \nu} \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|u(s)\|_{L^2}^2 ds. \end{aligned}$$

Therefore by taking expectations in (A.15), using  $\mathbb{E}[III_t] = \mathbb{E}[IV_t] = 0$ , and (A.17)-(A.18), one obtains

$$(A.19) \quad \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|\nabla u(s)\|_{L^2}^2 ds \leq C \left( 1 + t + \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|u(s)\|_{L^2}^2 ds \right)$$

where  $C$  is independent of  $u, u_0, j$ .

Next we take  $\mathbb{E}[\sup_{s \in [0, t]} |\cdot|]$  in (A.15). Due to (A.17)-(A.18), it remains to estimate  $III$  and  $IV$ . By the Burkholder-Davis-Gundy inequality, for all  $t \in [0, T]$ ,

$$\begin{aligned} \mathbb{E} \left[ \sup_{s \in [0, t]} |III_s| \right] &\lesssim \mathbb{E} \left[ \int_0^t \mathbf{1}_{[0, \varrho]} \|(b_n(s) \cdot \nabla)u(s)\|_{L^2(\ell^2)}^2 \|u(s)\|_{L^2}^2 ds \right]^{1/2} \\ &\leq M \mathbb{E} \left[ \left( \sup_{s \in [0, \varrho \wedge t]} \|u(s)\|_{L^2}^2 \right)^{1/2} \left( \int_0^t \mathbf{1}_{[0, \varrho]} \|\nabla u(s)\|_{L^2}^2 ds \right)^{1/2} \right] \\ &\leq \frac{1}{4} \mathbb{E} y(t) + C_M \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|\nabla u(s)\|_{L^2}^2 ds \\ &\stackrel{(A.19)}{\leq} \frac{1}{4} \mathbb{E} y(t) + C'_M \left( 1 + \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|u(s)\|_{L^2}^2 ds \right). \end{aligned}$$

where  $y$  is as in (A.13). A similar argument applies to  $IV$ . Indeed, by (A.6), for all  $t \in [0, T]$ ,

$$\begin{aligned} \mathbb{E} \left[ \sup_{s \in [0, t]} |IV_s| \right] &\leq \mathbb{E} \left[ \left( \sup_{s \in [0, \varrho \wedge t]} \|u(s)\|_{L^2}^2 \right)^{1/2} \left( \int_0^t \mathbf{1}_{[0, \varrho]} \|g(s, u(s))\|_{L^2(\ell^2)}^2 ds \right)^{1/2} \right] \\ &\leq \frac{1}{4} \mathbb{E} y(t) + C \left( 1 + \mathbb{E} \int_0^t \mathbf{1}_{[0, \varrho]} \|u(s)\|_{L^2}^2 ds \right) \end{aligned}$$

Taking  $\mathbb{E}[\sup_{s \in [0, t]} |\cdot|]$  in (A.15) and collecting the previous estimates as well as using that  $u_\varrho = u$  on  $[0, \varrho]$  and  $\varrho = \sigma_j$  we get (A.12).  $\square$

*Remark A.8.* As already mentioned at the beginning of this appendix several other boundary conditions can be used in (A.1). We mention two possible choices in case  $a^{i,j} = \delta^{i,j}$  (the Kronecker's delta): *perfect slip* boundary conditions

$$u \cdot \nu = 0, \quad P_{\partial\mathcal{O}}[(\nabla u - (\nabla u)^T)\nu] = 0, \quad \text{on } \partial\mathcal{O},$$

or *Navier* boundary conditions

$$u \cdot \nu = 0, \quad P_{\partial\mathcal{O}}[(\nabla u + (\nabla u)^T)\nu] + \alpha u = 0, \quad \text{on } \partial\mathcal{O},$$

where  $\alpha \geq 0$ ,  $\nu$  is the outer normal vector field on  $\partial\mathcal{O}$  and  $P_{\partial\mathcal{O}} := I - \nu \otimes \nu$  is the orthogonal projection on the tangent plane of  $\partial\mathcal{O}$ . In case  $\mathcal{O}$  is a bounded  $C^3$ -domain, the weak formulation of the Stokes operators subject to one of the previous boundary conditions has been given in Subsection 2.3 and 3.2 of [PW18], respectively. Using the methods in the latter one can check that in case  $a^{i,j} = \delta^{i,j}$  and  $\mathcal{O}$  is a  $C^3$ -bounded domain, then the results of Theorem A.3 extends to the case of perfect-slip or Navier boundary conditions. Indeed, (A.16) still holds since  $u \cdot \nu = 0$  on  $\partial\mathcal{O}$  in both cases and one can replace the use of Lemma A.7 by standard Sobolev embeddings due to the regularity of  $\mathcal{O}$ .

*Acknowledgements.* The authors thank Max Sauerbrey for careful reading and helpful comments.

#### REFERENCES

- [ALV21] A. Agresti, N. Lindemulder, and M. Veraar. On the trace embedding and its applications to evolution equations. *arXiv preprint arXiv:2104.05063*, 2021.
- [AV20] A. Agresti and M.C. Veraar. Stability properties of stochastic maximal  $L^p$ -regularity. *J. Math. Anal. Appl.*, 482(2):123553, 35, 2020.
- [AV21a] A. Agresti and M.C. Veraar. Nonlinear parabolic stochastic evolution equations in critical spaces Part I. Stochastic maximal regularity and local existence. Accepted for publication in *Nonlinearity*, arXiv preprint arXiv:2001.00512, 2021.
- [AV21b] A. Agresti and M.C. Veraar. Nonlinear parabolic stochastic evolution equations in critical spaces Part II. Blow criteria and instantaneous regularization. arXiv preprint arXiv:2012.04448, 2021.
- [AV21c] A. Agresti and M.C. Veraar. Stochastic maximal  $L^p(L^q)$ -regularity for second order systems with periodic boundary conditions. arXiv preprint arXiv:2106.01274, 2021.
- [Are04] W. Arendt. Semigroups and evolution equations: functional calculus, regularity and kernel estimates. In *Evolutionary equations. Vol. I*, Handb. Differ. Equ., pages 1–85. North-Holland, Amsterdam, 2004.
- [BT73] A. Bensoussan and R. Temam. Équations stochastiques du type Navier-Stokes. *J. Functional Analysis*, 13:195–222, 1973.
- [BL76] J. Bergh and J. Löfström. *Interpolation spaces. An introduction*. Springer-Verlag, Berlin, 1976. Grundlehren der Mathematischen Wissenschaften, No. 223.
- [BF20] L. A. Bianchi and F. Flandoli. Stochastic Navier-Stokes equations and related models. *Milan J. Math.*, 88(1):225–246, 2020.
- [BMS10] G. Bourdaud, M. Moussai, and W. Sickel. Composition operators on Lizorkin-Triebel spaces. *J. Funct. Anal.*, 259(5):1098–1128, 2010.
- [BCF91] Z. Brzeźniak, M. Capiński, and F. Flandoli. Stochastic partial differential equations and turbulence. *Math. Models Methods Appl. Sci.*, 1(1):41–59, 1991.
- [BCF92] Z. Brzeźniak, M. Capiński, and F. Flandoli. Stochastic Navier-Stokes equations with multiplicative noise. *Stochastic Anal. Appl.*, 10(5):523–532, 1992.
- [BF09] Z. Brzeźniak and B. Ferrario. 2D Navier-Stokes equation in Besov spaces of negative order. *Nonlinear Anal.*, 70(11):3902–3916, 2009.
- [BF17] Z. Brzeźniak and B. Ferrario. A note on stochastic Navier-Stokes equations with not regular multiplicative noise. *Stoch. Partial Differ. Equ. Anal. Comput.*, 5, 2017.

- [BM13] Z. Brzeźniak and E. Motyl. Existence of a martingale solution of the stochastic Navier-Stokes equations in unbounded 2D and 3D domains. *J. Differential Equations*, 254(4):1627–1685, 2013.
- [BP00] Z. Brzeźniak and S. Peszat. Strong local and global solutions for stochastic Navier-Stokes equations. In *Infinite dimensional stochastic analysis (Amsterdam, 1999)*, volume 52 of *Verh. Afd. Natuurkd. 1. Reeks. K. Ned. Akad. Wet.*, pages 85–98. R. Neth. Acad. Arts Sci., Amsterdam, 2000.
- [Can04] M. Cannone. Harmonic analysis tools for solving the incompressible Navier-Stokes equations. In *Handbook of mathematical fluid dynamics. Vol. III*, pages 161–244. North-Holland, Amsterdam, 2004.
- [CP01] M. Capiński and S. Peszat. On the existence of a solution to stochastic Navier-Stokes equations. *Nonlinear Anal.*, 44, 2001.
- [Cho78] P. L. Chow. Stochastic partial differential equations in turbulence related problems. In *Probabilistic analysis and related topics, Vol. 1*, pages 1–43. 1978.
- [DZ20] L. Du and T. Zhang. Local and global strong solutions to the stochastic incompressible Navier-Stokes equations in critical Besov space. *J. Math. Anal. Appl.*, 481, 2020.
- [Fla08] F. Flandoli. An introduction to 3D stochastic fluid dynamics. In *SPDE in hydrodynamic: recent progress and prospects*, pages 51–150. Springer, 2008.
- [Fla11] F. Flandoli. *Random perturbation of PDEs and fluid dynamic models*, volume 2015 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 2011. Lectures from the 40th Probability Summer School held in Saint-Flour, 2010, École d’Été de Probabilités de Saint-Flour.
- [Fla15] F. Flandoli. A stochastic view over the open problem of well-posedness for the 3D Navier-Stokes equations. In *Stochastic analysis: a series of lectures*, volume 68 of *Progr. Probab.*, pages 221–246. Birkhäuser/Springer, Basel, 2015.
- [FGL21] F. Flandoli, L. Galeati, and D. Luo. Eddy heat exchange at the boundary under white noise turbulence. *arXiv preprint arXiv:2103.08098*, 2021.
- [FG95] F. Flandoli and D. Gatarek. Martingale and stationary solutions for stochastic Navier-Stokes equations. *Probab. Theory Related Fields*, 102, 1995.
- [FL21] F. Flandoli and D. Luo. High mode transport noise improves vorticity blow-up control in 3D Navier-Stokes equations. *Probab. Theory Related Fields*, 180, 2021.
- [FK64] H. Fujita and T. Kato. On the Navier-Stokes initial value problem. I. *Arch. Rational Mech. Anal.*, 16:269–315, 1964.
- [GP02] I. Gallagher and F. Planchon. On global infinite energy solutions to the Navier-Stokes equations in two dimensions. *Arch. Ration. Mech. Anal.*, 161(4):307–337, 2002.
- [GK96] K. Gawędzki and A. Kupiainen. Universality in turbulence: an exactly solvable model. In *Low-dimensional models in statistical physics and quantum field theory*, volume 469 of *Lecture Notes in Phys.*, pages 71–105. Springer, Berlin, 1996.
- [GV00] K. Gawędzki and M. Vergassola. Phase transition in the passive scalar advection. *Phys. D*, 138(1-2):63–90, 2000.
- [HLN19] M. Hofmanová, J.-M. Leahy, and T. Nilssen. On the Navier-Stokes equation perturbed by rough transport noise. *J. Evol. Equ.*, 19, 2019.
- [HLN21] M. Hofmanová, J.-M. Leahy, and T. Nilssen. On a rough perturbation of the Navier-Stokes system and its vorticity formulation. *Ann. Appl. Probab.*, 31(2):736–777, 2021.
- [HNVW16] T.P. Hytönen, J.M.A.M. van Neerven, M.C. Veraar, and L. Weis. *Analysis in Banach spaces. Vol. I. Martingales and Littlewood-Paley theory*, volume 63 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer, 2016.
- [HNVW17] T.P. Hytönen, J.M.A.M. van Neerven, M.C. Veraar, and L. Weis. *Analysis in Banach spaces. Vol. II. Probabilistic Methods and Operator Theory.*, volume 67 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer, 2017.
- [IF79] A. Inoue and T. Funaki. On a new derivation of the Navier-Stokes equation. *Comm. Math. Phys.*, 65(1):83–90, 1979.
- [Kim10] J. U. Kim. Strong solutions of the stochastic Navier-Stokes equations in  $\mathbb{R}^3$ . *Indiana Univ. Math. J.*, 59, 2010.
- [KT01] H. Koch and D. Tataru. Well-posedness for the Navier-Stokes equations. *Adv. Math.*, 157(1):22–35, 2001.
- [Kra68] R. H. Kraichnan. Small-scale structure of a scalar field convected by turbulence. *The Physics of Fluids*, 11(5):945–953, 1968.

- [Kra94] R. H. Kraichnan. Anomalous scaling of a randomly advected passive scalar. *Physical review letters*, 72(7):1016, 1994.
- [KXZ21] I. Kukavica, F. Xu, and M. Ziane. Global existence for the stochastic Navier–Stokes equations with small  $L^p$  data. *Stochastics and Partial Differential Equations: Analysis and Computations*, pages 1–30, 2021.
- [LR16] P. G. Lemarié-Rieusset. *The Navier-Stokes problem in the 21st century*. CRC Press, Boca Raton, FL, 2016.
- [LR15] W. Liu and M. Röckner. *Stochastic partial differential equations: an introduction*. Universitext. Springer, Cham, 2015.
- [Mik02] R. Mikulevicius. On the Cauchy problem for stochastic Stokes equations. *SIAM J. Math. Anal.*, 34, 2002.
- [Mik09] R. Mikulevicius. On strong  $H_2^1$ -solutions of stochastic Navier-Stokes equation in a bounded domain. *SIAM J. Math. Anal.*, 41(3):1206–1230, 2009.
- [MR01] R. Mikulevicius and B. Rozovskii. On equations of stochastic fluid mechanics. In *Stochastics in finite and infinite dimensions*, Trends Math., pages 285–302. Birkhäuser Boston, Boston, MA, 2001.
- [MR04] R. Mikulevicius and B. Rozovskii. Stochastic Navier-Stokes equations for turbulent flows. *SIAM J. Math. Anal.*, 35(5):1250–1310, 2004.
- [MR05] R. Mikulevicius and B. Rozovskii. Global  $L_2$ -solutions of stochastic Navier-Stokes equations. *Ann. Probab.*, 33(1):137–176, 2005.
- [MR99] R. Mikulevicius and B. L. Rozovskii. Martingale problems for stochastic PDE’s. In *Stochastic partial differential equations: six perspectives*, volume 64 of *Math. Surveys Monogr.*, pages 243–325. Amer. Math. Soc., Providence, RI, 1999.
- [NVW15] J.M.A.M. van Neerven, M.C. Veraar, and L.W. Weis. Stochastic integration in Banach spaces—a survey. In *Stochastic analysis: a series of lectures*, volume 68 of *Progr. Probab.*, pages 297–332. Birkhäuser/Springer, Basel, 2015.
- [Ond04] M. Ondreját. Uniqueness for stochastic evolution equations in Banach spaces. *Dissertationes Math. (Rozprawy Mat.)*, 426:63, 2004.
- [PSW18] J. Prüss, G. Simonett, and M. Wilke. Critical spaces for quasilinear parabolic evolution equations and applications. *J. Differential Equations*, 264(3):2028–2074, 2018.
- [PW18] J. Prüss and M. Wilke. On critical spaces for the Navier-Stokes equations. *J. Math. Fluid Mech.*, 20(2):733–755, 2018.
- [ST87] H.-J. Schmeisser and H. Triebel. *Topics in Fourier analysis and function spaces*. A Wiley-Interscience Publication. John Wiley & Sons, Ltd., Chichester, 1987.
- [Tao19] T. Tao. Quantitative bounds for critically bounded solutions to the Navier-Stokes equations. *arXiv preprint arXiv:1908.04958*, 2019.
- [Tay11] M. E. Taylor. *Partial differential equations III. Nonlinear equations*, volume 117 of *Applied Mathematical Sciences*. Springer, New York, second edition, 2011.
- [Tri95] H. Triebel. *Interpolation theory, function spaces, differential operators*. Johann Ambrosius Barth, Heidelberg, second edition, 1995.
- [Tri13] H. Triebel. *Local function spaces, heat and Navier-Stokes equations*, volume 20 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2013.
- [Wan20] W. Wang. Global existence and analyticity of mild solutions for the stochastic Navier-Stokes-Coriolis equations in Besov spaces. *Nonlinear Anal. Real World Appl.*, 52:103048, 15, 2020.
- [WS00] E. Weinan and Y. G. Sinai. Recent results on mathematical and statistical hydrodynamics. *Russian Mathematical Surveys*, 55(4):635, 2000.

DEPARTMENT OF MATHEMATICS, TECHNISCHE UNIVERSITÄT KAISERSLAUTERN, PAUL-EHRlich-STRASSE 31, 67663 KAISERSLAUTERN, GERMANY  
 Email address: antonio.agresti92@gmail.com

DELFT INSTITUTE OF APPLIED MATHEMATICS, DELFT UNIVERSITY OF TECHNOLOGY, P.O. BOX 5031, 2600 GA DELFT, THE NETHERLANDS.  
 Email address: M.C.Veraar@tudelft.nl