

UNIQUENESS AND NON-UNIQUENESS RESULTS FOR DYADIC MHD MODELS

MIMI DAI AND SUSAN FRIEDLANDER

ABSTRACT. We construct non-unique Leray-Hopf solutions for some dyadic models for magnetohydrodynamics when the intermittency dimension δ is less than 1. In contrast, uniqueness of Leray-Hopf solution is established in the case of $\delta \geq 1$. Analogous results on uniqueness and non-uniqueness of Leray-Hopf solution are also obtained for dyadic models of MHD with fractional diffusion.

KEY WORDS: magnetohydrodynamics; intermittency; dyadic model; uniqueness and non-uniqueness.

CLASSIFICATION CODE: 35Q35, 76D03, 76W05.

1. INTRODUCTION

1.1. Magnetohydrodynamics. In geophysics and astrophysics, incompressible magnetohydrodynamics (MHD) governed by the equations

$$u_t + (u \cdot \nabla)u - (B \cdot \nabla)B + \nabla P = \nu \Delta u + f, \quad (1.1a)$$

$$B_t + (u \cdot \nabla)B - (B \cdot \nabla)u = \mu \Delta B, \quad (1.1b)$$

$$\nabla \cdot u = 0, \quad \nabla \cdot B = 0, \quad (1.1c)$$

is a fundamental model in the investigation of electrically conducting fluids. In this system the vector fields u and B represent the fluid velocity and magnetic field, respectively; the scalar function P denotes the pressure; the parameters ν and μ denote respectively the viscosity and the magnetic resistivity; and f stands for an external force acting on the fluid. When $B = 0$, system (1.1a)-(1.1c) becomes the Navier-Stokes equation (NSE) (1.4) which will be discussed later.

It is evident that the MHD system inherits challenges from the NSE, but also exhibits its own complexity which is mainly caused by the nonlinear interactions between the fluid velocity field and the magnetic field. The unsolved problems for the NSE usually also hang in the air for the MHD system. In particular, it is not clear whether either the NSE or MHD has a classical solution for all the time, given arbitrary initial data. The concept of Leray-Hopf solution for the NSE was introduced by Leray [21] and Hopf [17]. A Leray-Hopf solution is a weak solution in the standard distributional sense, which satisfies the basic energy inequality. Such a concept is naturally adapted to other partial differential equations. Since the pioneering work of Leray, the well-posedness problem for the Leray-Hopf solutions to the NSE in three dimensional (3D) space is still open. In particular, the uniqueness of a Leray-Hopf solution to the 3D NSE remains unsolved. This is the case for the 3D MHD (1.1a)-(1.1c) as well. Nonetheless, wild weak solutions have been constructed for the ideal MHD, i.e. $\nu = \mu = 0$ and $f = 0$ in (1.1a)-(1.1c) by Beekie, Buckmaster and Vicol [3]. The weak solutions constructed in [3] have finite total

energy, but do not conserve the magnetic helicity which is an invariant quantity for smooth solutions.

The main objective of this paper is to investigate the problem of uniqueness of Leray-Hopf solutions for the diffusive dyadic models of the MHD system (1.1a)-(1.1c). The following dyadic model for the MHD system was proposed in [10],

$$\begin{aligned} \frac{d}{dt}a_j + \nu\lambda_j^2a_j + \kappa_1(\lambda_j^\theta a_j a_{j+1} - \lambda_{j-1}^\theta a_{j-1}^2) \\ - \kappa_2(\lambda_j^\theta b_j b_{j+1} - \lambda_{j-1}^\theta b_{j-1}^2) = f_j, \end{aligned} \quad (1.2a)$$

$$\frac{d}{dt}b_j + \mu\lambda_j^2b_j + \kappa_2(\lambda_j^\theta a_j b_{j+1} - \lambda_j^\theta b_j a_{j+1}) = 0 \quad (1.2b)$$

for $j \geq 0$, $\lambda_j = \lambda^j$ with a fixed constant $\lambda > 1$, and $a_{-1} = b_{-1} = 0$. The variables (a_j, b_j) are quantities related to the energy in the sense that $\frac{1}{2}a_j^2$ and $\frac{1}{2}b_j^2$ are the kinetic energy and magnetic energy in the j -th shell, respectively. The parameter θ is defined as $\theta = \frac{5-\delta}{2}$, where $\delta \in [0, 3]$ is the intermittency dimension for the 3D turbulent vector field (cf. [5]). Naturally, $\theta \in [1, \frac{5}{2}]$. Notice that smaller δ corresponds to larger θ , and hence stronger nonlinearity. The parameters κ_1 and κ_2 placed in front of the nonlinear terms represent the energy transfer direction and strength among shells. Similar dyadic models have been presented by physicists for the MHD system, for instance, see [16, 23].

Denote the total energy by

$$E(t) = \frac{1}{2} \sum_{j=0}^{\infty} (a_j^2 + b_j^2)$$

and the flux through the j -th shell by

$$\Pi_j = \lambda_j^\theta (\kappa_1 a_j^2 - \kappa_2 b_j^2) a_{j+1}, \quad j \geq 0.$$

The energy balance for the j -th shell of the system (1.2a)-(1.2b) is

$$\frac{d}{dt}(a_j^2 + b_j^2) = -\nu\lambda_j^2a_j^2 - \mu\lambda_j^2b_j^2 + \Pi_{j-1} - \Pi_j + f_j a_j.$$

Thus system (1.2a)-(1.2b) obeys the formal energy law

$$\frac{d}{dt}E(t) = -\nu \sum_{j=0}^{\infty} \lambda_j^2 a_j^2 - \mu \sum_{j=0}^{\infty} \lambda_j^2 b_j^2 + \sum_{j=0}^{\infty} f_j a_j.$$

It is clear to see that the energy is invariant for (1.2a)-(1.2b) if $\nu = \mu = 0$ and $f_j = 0$ for $j \geq 0$. We will consider the four particular cases of the general model (1.2a)-(1.2b) with $\kappa_1 = \pm 1$ and $\kappa_2 = \pm 1$.

We will provide a definition of Leray-Hopf solutions for dyadic models in analogy with the Leray-Hopf solution for the original MHD equations (1.1a)-(1.1c). The main goal is to: (i) establish global in time existence of Leray-Hopf solutions for the dyadic models; (ii) show the uniqueness of Leray-Hopf solution when $\theta \leq 2$; (iii) construct non-unique Leray-Hopf solutions in the case of $\theta > 2$. Philosophically, the process of constructing non-uniqueness resembles the convex integration method in the sense that it takes advantage of the forcing term in the construction. Technically it is much simpler than convex integration since no iteration or approximation is involved.

1.2. Main results for dyadic MHD models. In this part, we lay out the results regarding Leray-Hopf solutions for the dyadic MHD model (1.2a)-(1.2b). First, for any initial data with finite total energy, we show the existence of global Leray-Hopf solutions.

Theorem 1.1. *Let $\theta > 0$, $a^0 = \{a_j^0\}_{j \geq 0} \in l^2$ and $b^0 = \{b_j^0\}_{j \geq 0} \in l^2$. For any $T > 0$, assume*

$$\sum_{j=0}^{\infty} \lambda_j^{-2} \int_0^T f_j^2(t) dt < \infty,$$

i.e. $f \in L^2(0, T; H^{-1})$. Then there exists a Leray-Hopf solution to system (1.2a)-(1.2b) accompanied with the initial data (a^0, b^0) on $[0, T]$.

The next result concerns the weak-strong type of uniqueness.

Theorem 1.2. *Let $\theta > 0$. Let $(a(t), b(t))$ and $(u(t), v(t))$ be Leray-Hopf solutions to (1.2a)-(1.2b) with the same initial data $(a^0, b^0) \in l^2 \times l^2$. Assume in addition that there is a number J such that*

$$|a_j(t)| \leq C_0 \lambda_j^{2-\theta}, \quad |b_j(t)| \leq C_0 \lambda_j^{2-\theta} \quad \text{for } j \geq J, \quad t \in [0, T] \quad (1.3)$$

with a constant C_0 depending on λ and θ . Then

$$a_j \equiv u_j, \quad b_j \equiv v_j, \quad \text{on } [0, T] \quad \text{for all } j \geq 0.$$

As a consequence of Theorem 1.2, the uniqueness of the Leray-Hopf solution in the case of $\theta \leq 2$ follows immediately.

Theorem 1.3. *Let $0 < \theta \leq 2$. Let $a^0 = \{a_j^0\}_{j \geq 0} \in l^2$, $b^0 = \{b_j^0\}_{j \geq 0} \in l^2$ and $f \in L^2(0, T; H^{-1})$. Then the Leray-Hopf solution to (1.2a)-(1.2b) is unique.*

When $\theta > 2$, we adapt the construction strategy of [14] for the dyadic NSE and show that the dyadic MHD models have more than one Leray-Hopf solutions. Specifically, we prove:

Theorem 1.4. *Let $\theta > 2$. Let $a^0 = 0$ and $b^0 = 0$, i.e. $a_j^0 = b_j^0 = 0$ for all $j \geq 0$. There exists $T > 0$ and functions $\{f_j(t)\}$ satisfying*

$$\sum_{j=0}^{\infty} \lambda_j^{-2} \int_0^T f_j^2(t) dt < \infty,$$

such that system (1.2a)-(1.2b) with initial data (a^0, b^0) has at least two Leray-Hopf solutions $(a(t), b(t))$, one of which has non-vanishing $a(t)$ and $b(t)$ on $[0, T]$.

Remark 1.5. The solutions constructed in Theorem 1.4 satisfy the energy identity.

Remark 1.6. We see that the threshold value of θ that separates the uniqueness and non-uniqueness results is $\theta = 2$. Notice that since $\theta = \frac{5-\delta}{2}$, $\theta = 2$ corresponds to the intermittency dimension $\delta = 1$. In fact, there is evidence that $\delta = 1$ is critical for 3D turbulent flows, see [9].

1.3. Weak solutions for dyadic NSE. The incompressible Navier-Stokes equation

$$\begin{aligned} u_t + (u \cdot \nabla)u + \nabla P &= \nu \Delta u + f, \\ \nabla \cdot u &= 0, \end{aligned} \tag{1.4}$$

is a central topic in the study of fluids. In (1.4), the vector field u , scalar function P , parameter ν and vector valued function f denote the fluid velocity, pressure, viscosity and external forcing, respectively. Although there has been much progress in the past century concerning fundamental properties of the NSE, many significant questions remain open. Partly for this reason various so called dyadic models have been proposed. One such model for oceanographic turbulence was presented by Desnyanskiy and Novikov [12] in 1974 and later with motivation from harmonic analysis by Katz and Pavlović [19]. This model takes the form

$$\frac{d}{dt}a_j + \nu \lambda_j^2 a_j + \lambda_j^\theta a_j a_{j+1} - \lambda_{j-1}^\theta a_{j-1}^2 = f_j, \tag{1.5}$$

for $j \geq 1$ and $a_0 = 0$. A crucial property of this particular model is the persistence of positivity, namely that with nonnegative forcing a solution starting from positive initial data remains positive for all time. This attribute of the system (1.5) was essential for the proof of many interesting results, for example see [1, 2, 4, 6, 7, 8]. However, as was recently observed by Filonov and Khodunov [14] the energy cascade in a turbulent fluid is a random process with no physical reason for the conservation of positivity. Hence there is an intrinsic desirability for techniques that do not depend on positivity. In [14] the authors introduced a novel approach that does not depend on positivity. They proved for (1.5) existence and uniqueness of Leray-Hopf solution with $\theta \leq 2$ and showed that there exist more than one Leray-Hopf solution with $\theta > 2$. Specifically, they proved the following theorem:

Theorem 1.7. [14] *Let $\theta > 2$ and $a^0 = 0$. There exists $T > 0$ and functions $f_j(t)$ satisfying $f = \{f_j\}_{j \geq 0} \in L^2(0, T; H^{-1})$ such that the dyadic NSE model (1.5) with initial data a^0 has at least two Leray-Hopf solutions.*

Returning to the dyadic MHD model (1.2a)-(1.2b) the delicate interactions between the velocity and the magnetic fields preclude the possibility of making sign choice of the parameters that ensure the persistence of positivity. The techniques that we use to prove the results stated in Theorems 1.2 - 1.4 are motivated by the approach used for the NSE in [14] which does not depend on positivity. We observe that in this context the complexity of the nonlinear coupling of the two fields is actually a benefit that gives us additional freedom in constructing a scheme used to prove non-uniqueness.

1.4. Dyadic models with fractional Laplacian scaling. We note that the dyadic MHD equations (1.2a)-(1.2b) can also be rescaled to

$$\begin{aligned} \frac{d}{dt}a_j + \nu \lambda_j^{2\alpha} a_j + \kappa_1 (\lambda_j a_j a_{j+1} - \lambda_{j-1} a_{j-1}^2) \\ - \kappa_2 (\lambda_j b_j b_{j+1} - \lambda_{j-1} b_{j-1}^2) &= f_j, \\ \frac{d}{dt}b_j + \mu \lambda_j^{2\alpha} b_j + \kappa_2 (\lambda_j a_j b_{j+1} - \lambda_j b_j a_{j+1}) &= 0, \end{aligned} \tag{1.6}$$

for $j \geq 1$, $a_0 = b_0 = 0$ and $\alpha = \frac{1}{\theta}$.

The analogous dyadic model for the fractional MHD with diffusion terms $(-\Delta)^\alpha u$ and $(-\Delta)^\beta B$ is

$$\begin{aligned} \frac{d}{dt} a_j + \nu \lambda_j^{2\alpha} a_j + \kappa_1 (\lambda_j a_j a_{j+1} - \lambda_{j-1} a_{j-1}^2) \\ - \kappa_2 (\lambda_j b_j b_{j+1} - \lambda_{j-1} b_{j-1}^2) = f_j, \\ \frac{d}{dt} b_j + \mu \lambda_j^{2\beta} b_j + \kappa_2 (\lambda_j a_j b_{j+1} - \lambda_j b_j a_{j+1}) = 0, \end{aligned} \quad (1.7)$$

with $j \geq 1$, $a_0 = b_0 = 0$, and $\alpha > 0, \beta > 0$. Obviously (1.6) is a special case of (1.7) with $\alpha = \beta$. With slight modifications of the proof for Theorem 1.1, we can prove that:

Theorem 1.8. *Let $\alpha > 0$ and $\beta > 0$. Let $a^0 = \{a_j^0\}_{j \geq 0} \in l^2$ and $b^0 = \{b_j^0\}_{j \geq 0} \in l^2$. Assume $f \in L^2(0, T; H^{-\alpha})$ for any $T > 0$. Then there exists a Leray-Hopf solution to system (1.7) accompanied with the initial data (a^0, b^0) on $[0, T]$.*

In analogy with Theorem 1.2, we can prove the following weak-strong type of uniqueness for a Leray-Hopf solution to (1.7).

Theorem 1.9. *Let $\alpha > 0$ and $\beta > 0$. Let $(a(t), b(t))$ and $(u(t), v(t))$ be Leray-Hopf solutions to (1.7) with the same initial data $(a^0, b^0) \in l^2 \times l^2$. Assume in addition that there is a number J such that*

$$|a_j(t)| \leq C_0 \left(\lambda_j^{2\alpha-1} + \lambda_j^{2\beta-1} \right), \quad |b_j(t)| \leq C_0 \lambda_j^{\alpha+\beta-1} \quad (1.8)$$

for all $j \geq J$ and $t \in [0, T]$, with a constant C_0 depending on λ and θ . Then

$$a_j \equiv u_j, \quad b_j \equiv v_j, \quad \text{on } [0, T] \text{ for all } j \geq 0.$$

The following uniqueness of a Leray-Hopf solution to (1.7) with $\alpha \geq \frac{1}{2}$ and $\beta \geq \frac{1}{2}$ is an immediate consequence of Theorem 1.9.

Theorem 1.10. *Let $\alpha \geq \frac{1}{2}$ and $\beta \geq \frac{1}{2}$. Let $a^0 = \{a_j^0\}_{j \geq 0} \in l^2$, $b^0 = \{b_j^0\}_{j \geq 0} \in l^2$ and $f \in L^2(0, T; H^{-\alpha})$. Then the Leray-Hopf solution to (1.7) is unique.*

We also construct non-unique Leray-Hopf solutions to (1.7) for appropriate values of α and β . Namely, we will show:

Theorem 1.11. *Let $0 < \alpha \leq \beta < \frac{1}{2}$ and $3\beta - \alpha < 1$. Let $a^0 = 0$ and $b^0 = 0$. There exists $T > 0$ and functions $\{f_j(t)\}$ satisfying*

$$\sum_{j=0}^{\infty} \lambda_j^{-2\alpha} \int_0^T f_j^2(t) dt < \infty,$$

such that system (1.7) with initial data (a^0, b^0) has at least two Leray-Hopf solutions.

When $0 < \alpha = \beta < \frac{1}{2}$, we automatically have the same result for system (1.6).

Remark 1.12. In view of Theorem 1.10 and Theorem 1.11, the value of $\frac{1}{2}$ for α and β is a sharp threshold to separate uniqueness from non-uniqueness result for system (1.6). On the other hand, for (1.7) the additional conditions of $\alpha \leq \beta$ and $3\beta - \alpha < 1$ leave some gap where neither uniqueness nor non-uniqueness is known to hold.

1.5. **Organisation of the paper.** We provide an outline of the rest of the paper.

- Section 2 introduces notations and definitions of solutions for dyadic systems.
- Section 3 is devoted to a proof of Theorem 1.1 on the existence of Leray-Hopf solutions.
- Section 4 addresses the weak-strong uniqueness and uniqueness of Leray-Hopf solution for system (1.2a)-(1.2b) with $\theta \leq 2$.
- In Section 5 we construct non-unique Leray-Hopf solutions for (1.2a)-(1.2b) with $\theta > 2$.
- Section 6 outlines constructions to establish firstly conditions for uniqueness and secondly conditions for non-uniqueness of Leray-Hopf solutions to the dyadic model for MHD with fractional diffusions.

2. NOTATIONS AND NOTION OF SOLUTIONS

2.1. **Notations.** The space l^2 is endowed with the standard scalar product and norm,

$$(u, v) := \sum_{n=1}^{\infty} u_n v_n, \quad |u| := \sqrt{(u, u)}.$$

It is regarded as the energy space in this paper. We use H^s to represent the space of sequences equipped with the scalar product

$$(u, v)_s := \sum_{n=1}^{\infty} \lambda_n^{2s} u_n v_n$$

and norm

$$\|u\|_s := \sqrt{(u, u)_s}.$$

Notice that $H^0 = l^2$.

2.2. **Notion of solutions.** In the following, we introduce the concept of solutions for dyadic systems.

Definition 2.1. A pair of l^2 -valued functions $(a(t), b(t))$ defined on $[0, \infty)$ is said to be a weak solution of (1.2a)-(1.2b) if a_j and b_j satisfy (1.2a)-(1.2b) and $a_j, b_j \in C^1([0, \infty))$ for all $j \geq 0$.

Definition 2.2. A Leray-Hopf solution $(a(t), b(t))$ of (1.2a)-(1.2b) on $[0, T)$ is a weak solution satisfying

$$a_j, b_j \in L^\infty([0, T]; l^2) \cap L^2([0, T]; H^1), \quad \forall j \geq 0,$$

and

$$\begin{aligned} & \|a(t)\|_{l^2}^2 + \|b(t)\|_{l^2}^2 + 2\nu \int_0^t \|a(\tau)\|_{H^1}^2 d\tau + 2\mu \int_0^t \|b(\tau)\|_{H^1}^2 d\tau \\ & \leq \|a(0)\|_{l^2}^2 + \|b(0)\|_{l^2}^2 \end{aligned}$$

for all $0 \leq t < T$.

Definition 2.3. A Leray-Hopf solution $(a(t), b(t))$ of (1.6) on $[0, T)$ is a weak solution satisfying

$$a_j, b_j \in L^\infty([0, T]; l^2) \cap L^2([0, T]; H^\alpha), \quad \forall j \geq 0,$$

and

$$\begin{aligned} & \|a(t)\|_{l^2}^2 + \|b(t)\|_{l^2}^2 + 2\nu \int_0^t \|a(\tau)\|_{H^\alpha}^2 d\tau + 2\mu \int_0^t \|b(\tau)\|_{H^\alpha}^2 d\tau \\ & \leq \|a(0)\|_{l^2}^2 + \|b(0)\|_{l^2}^2 \end{aligned}$$

for all $0 \leq t < T$.

Weak solution and Leray-Hopf solution of other dyadic systems in the paper can be defined analogously.

To reduce the number of parameters, we take $\nu = \mu = 1$ in the rest of the paper since they do not affect the estimates or constructions.

3. EXISTENCE OF LERAY-HOPF SOLUTIONS

In this section, we apply the Galerkin approximating approach to show the existence of Leray-Hopf solutions to (1.2a)-(1.2b). Since the value of κ_1 and κ_2 does not play a role in the proof, without loss of generality, we set $\kappa_1 = -\kappa_2 = 1$. Fix any integer $N \geq 1$. Denote the sequences

$$a^N(t) = \{a_j^N(t)\}_{j \geq 0}, \quad b^N(t) = \{b_j^N(t)\}_{j \geq 0}, \quad \text{with } a_j^N = b_j^N \equiv 0, \quad \forall j \geq N+1.$$

That is,

$$\begin{aligned} a^N(t) &= (a_0^N(t), a_1^N(t), a_2^N(t), \dots, a_N^N(t), 0, 0, 0, \dots), \\ b^N(t) &= (b_0^N(t), b_1^N(t), b_2^N(t), \dots, b_N^N(t), 0, 0, 0, \dots). \end{aligned}$$

Consider the truncated system for $(a^N(t), b^N(t))$,

$$\begin{aligned} \frac{d}{dt} a_j^N &= -\lambda_j^2 a_j^N - \lambda_j^\theta a_j^N a_{j+1}^N + \lambda_{j-1}^\theta (a_{j-1}^N)^2 - \lambda_j^\theta b_j^N b_{j+1}^N \\ &\quad + \lambda_{j-1}^\theta (b_{j-1}^N)^2 + f_j, \quad 0 \leq j \leq N \\ \frac{d}{dt} b_j^N &= -\lambda_j^2 b_j^N + \lambda_j^\theta a_j^N b_{j+1}^N - \lambda_j^\theta b_j^N a_{j+1}^N, \quad 0 \leq j \leq N \\ a_j^N(0) &= a_j^0, \quad b_j^N(0) = b_j^0, \quad 0 \leq j \leq N. \end{aligned} \tag{3.1}$$

By convention, $a_{-1}^N = b_{-1}^N = 0$.

In the following, we proceed with the standard Galerkin approximating framework: (i) for any $N \geq 1$, there is a solution $(a^N(t), b^N(t))$ to (3.1) with $a^N(t)$ and $b^N(t)$ in the space $L^\infty(0, T; l^2) \cap L^2(0, T; H^1)$ and satisfying the corresponding energy inequality; (ii) we pass the sequence $\{(a^N(t), b^N(t))\}_{N \geq 1}$ (or a subsequence of it) to a limit $(a(t), b(t))$; (iii) the limit $(a(t), b(t))$ is shown to be a Leray-Hopf solution of (1.2a)-(1.2b).

The integral form of (3.1) is

$$\begin{aligned} a_j^N(t) &= a_j^0 + \int_0^t \left(-\lambda_j^2 a_j^N(\tau) - \lambda_j^\theta a_j^N(\tau) a_{j+1}^N(\tau) + \lambda_{j-1}^\theta (a_{j-1}^N(\tau))^2 \right. \\ &\quad \left. - \lambda_j^\theta b_j^N(\tau) b_{j+1}^N(\tau) + \lambda_{j-1}^\theta (b_{j-1}^N(\tau))^2 + f_j(\tau) \right) d\tau, \\ b_j^N(t) &= b_j^0 + \int_0^t \left(-\lambda_j^2 b_j^N(\tau) + \lambda_j^\theta a_j^N(\tau) b_{j+1}^N(\tau) - \lambda_j^\theta b_j^N(\tau) a_{j+1}^N(\tau) \right) d\tau, \end{aligned} \quad (3.2)$$

for $0 \leq j \leq N$. Denote

$$\begin{aligned} F^N(a^N, b^N, t) &= (F_0^N(a^N, b^N, t), F_1^N(a^N, b^N, t), \dots, F_N^N(a^N, b^N, t)), \\ G^N(a^N, b^N) &= (G_0^N(a^N, b^N), G_1^N(a^N, b^N), \dots, G_N^N(a^N, b^N)), \end{aligned}$$

with

$$\begin{aligned} F_j^N(a^N, b^N, t) &= -\lambda_j^2 a_j^N(t) - \lambda_j^\theta a_j^N(t) a_{j+1}^N(t) + \lambda_{j-1}^\theta (a_{j-1}^N(t))^2 \\ &\quad - \lambda_j^\theta b_j^N(t) b_{j+1}^N(t) + \lambda_{j-1}^\theta (b_{j-1}^N(t))^2 + f_j(t), \\ G_j^N(a^N, b^N) &= -\lambda_j^2 b_j^N(t) + \lambda_j^\theta a_j^N(t) b_{j+1}^N(t) - \lambda_j^\theta b_j^N(t) a_{j+1}^N(t), \end{aligned}$$

for $0 \leq j \leq N$. Denote $a^{0,N} = (a_0^0, a_1^0, \dots, a_N^0)$, $b^{0,N} = (b_0^0, b_1^0, \dots, b_N^0)$ and $f^N = (f_0, f_1, \dots, f_N)$. Thus, system (3.2) can be written as

$$\begin{aligned} a^N(t) &= a^{0,N} + \int_0^t F^N(a^N(\tau), b^N(\tau), \tau) d\tau, \\ b^N(t) &= b^{0,N} + \int_0^t G^N(a^N(\tau), b^N(\tau)) d\tau. \end{aligned} \quad (3.3)$$

Denote the map

$$M_N(a^N, b^N)(t) = \begin{pmatrix} a^{0,N} + \int_0^t F^N(a^N(\tau), b^N(\tau), \tau) d\tau \\ b^{0,N} + \int_0^t G^N(a^N(\tau), b^N(\tau)) d\tau \end{pmatrix}.$$

Notice that there exists a constant C_N depending on N such that

$$|F^N(a^N, b^N, t)| \leq C_N (|a^N| + |a^N|^2 + |b^N|^2) + |f^N|, \quad (3.4)$$

$$|G^N(a^N, b^N, t)| \leq C_N (|b^N| + |a^N|^2 + |b^N|^2), \quad (3.5)$$

and moreover

$$\begin{aligned} & \left| F^N(a^N, b^N, t) - F^N(\tilde{a}^N, \tilde{b}^N, t) \right| \\ & \leq C_N \left(1 + |a^N| + |\tilde{a}^N| + |b^N| + |\tilde{b}^N| \right) \left(|a^N - \tilde{a}^N| + |b^N - \tilde{b}^N| \right), \end{aligned} \quad (3.6)$$

$$\begin{aligned} & \left| G^N(a^N, b^N) - G^N(\tilde{a}^N, \tilde{b}^N) \right| \\ & \leq C_N \left(1 + |a^N| + |b^N| + |\tilde{a}^N| + |\tilde{b}^N| \right) \left(|a^N - \tilde{a}^N| + |b^N - \tilde{b}^N| \right). \end{aligned} \quad (3.7)$$

Choose

$$R_N = 2|a^{0,N}| + 2|b^{0,N}| + 2 \int_0^T |f^N(t)| dt, \quad (3.8)$$

and

$$t_{N,1} = \frac{1}{2C_N(2R_N + 1)}. \quad (3.9)$$

Consider the map $M_N(a^N, b^N)$ on the following closed subset of the space of continuous functions $C([0, t_{N,1}]; \mathbb{R}^N)$

$$B_N = \{(u, v) \in C([0, t_{N,1}]; \mathbb{R}^N) \times C([0, t_{N,1}]; \mathbb{R}^N) : \|u\|_C \leq R_N, \|v\|_C \leq R_N.\}$$

We claim that M_N is a contraction mapping on B_N . Indeed, for any $(a^N, b^N) \in B_N$, it follows from (3.4), (3.8) and (3.9) that

$$\begin{aligned} & \left| a^{0,N} + \int_0^t F^N(a^N(\tau), b^N(\tau), \tau) d\tau \right| \\ & \leq |a^{0,N}| + \int_0^t |F^N(a^N(\tau), b^N(\tau), \tau)| d\tau \\ & \leq |a^{0,N}| + tC_N(R_N + 2R_N^2) + \int_0^t |f^N(t)| dt \\ & \leq \frac{1}{2}R_N + \frac{1}{2C_N(2R_N + 1)}C_N(R_N + 2R_N^2) \\ & = R_N; \end{aligned}$$

and similarly, by (3.5), (3.8) and (3.9)

$$\begin{aligned} & \left| b^{0,N} + \int_0^t G^N(a^N(\tau), b^N(\tau)) d\tau \right| \\ & \leq |b^{0,N}| + \int_0^t |G^N(a^N(\tau), b^N(\tau))| d\tau \\ & \leq |b^{0,N}| + tC_N(R_N + 2R_N^2) \\ & \leq \frac{1}{2}R_N + \frac{1}{2C_N(2R_N + 1)}C_N(R_N + 2R_N^2) \\ & = R_N. \end{aligned}$$

Thus, M_N maps B_N to itself. On the other hand, the property of contraction follows from (3.6), (3.7) and the choice of time $t_{N,1}$ in (3.9). Therefore, system (3.3) has a solution $(a^N(t), b^N(t))$ on $[0, t_{N,1}]$, and so does system (3.1). Next we show that the solution satisfies the energy inequality. Multiplying the first equation of (3.1) by a_j^N and the second one by b_j^N , taking the sum for $0 \leq j \leq N$ and integrating over $[0, t]$, we obtain

$$\begin{aligned} & \sum_{j=0}^N (a_j^N(t)^2 + b_j^N(t)^2) + 2 \sum_{j=0}^N \int_0^t \lambda_j^2 (a_j^N(\tau)^2 + b_j^N(\tau)^2) d\tau \\ & = \sum_{j=0}^N \left((a_j^{0,N})^2 + (b_j^{0,N})^2 \right) + 2 \sum_{j=0}^N \int_0^t f_j(\tau) a_j^N(\tau) d\tau. \end{aligned} \tag{3.10}$$

Applying the Cauchy-Schwarz inequality, we have

$$2 \sum_{j=0}^N \int_0^t f_j(\tau) a_j^N(\tau) d\tau \leq \sum_{j=0}^N \int_0^t \lambda_j^{-2} f_j^2(\tau) d\tau + \sum_{j=0}^N \int_0^t \lambda_j^2 a_j^N(\tau)^2 d\tau.$$

Hence, it follows from (3.10)

$$\begin{aligned} & \sum_{j=0}^N (a_j^N(t)^2 + b_j^N(t)^2) + \sum_{j=0}^N \int_0^t \lambda_j^2 (a_j^N(\tau)^2 + b_j^N(\tau)^2) d\tau \\ & \leq \sum_{j=0}^N \left((a_j^{0,N})^2 + (b_j^{0,N})^2 \right) + \sum_{j=0}^N \int_0^t \lambda_j^{-2} f_j^2(\tau) d\tau. \end{aligned} \quad (3.11)$$

We can iterate the process above to construct the solution on time intervals $[t_{N,1}, t_{N,2}]$, $[t_{N,2}, t_{N,3}]$, ..., $[t_{N,k}, t_{N,k+1}]$, ..., and finally reach the time T . Indeed, we observe from the energy inequality (3.11) that

$$|a^N(t_k)| + |b^N(t_k)| \leq |a^{0,N}| + |b^{0,N}| + k \left(\sum_{j=0}^N \int_0^t \lambda_j^{-2} f_j^2(\tau) d\tau \right)^{\frac{1}{2}}.$$

Hence, according to (3.9) and (3.8), at the $k+1$ -th iteration, we have

$$\begin{aligned} & t_{N,k+1} - t_{N,k} \\ & \geq \left[C_N \left(8|a^{0,N}| + 8|b^{0,N}| + 4 \int_0^T |f^N(t)| dt + 2 + 4k \left(\sum_{j=0}^N \int_0^t \lambda_j^{-2} f_j^2(\tau) d\tau \right)^{\frac{1}{2}} \right) \right]^{-1} \\ & \gtrsim \frac{1}{k}. \end{aligned}$$

Therefore, the sum $\sum_k (t_{N,k+1} - t_{N,k})$ diverges and will reach T after a certain number of iterations. In conclusion, we obtain a solution $(a^N(t), b^N(t))$ of (3.1) on the interval $[0, T]$, which satisfies the energy inequality (3.11) for all $t \in [0, T]$.

The next step is to extract a limit from the sequence $\{(a^N(t), b^N(t))\}_{N \geq 1}$. In view of the integral equations in (3.2) and the energy inequality (3.11), we know $a_j^N, b_j^N \in C^1[0, T]$ for any $0 \leq j \leq N$ and $a^N, b^N \in L^\infty(0, T; l^2) \cap L^2(0, T; H^1)$ for any $N \geq 1$. As a consequence, there exists a subsequence $\{(a^{N_k}(t), b^{N_k}(t))\}_{k \geq 1}$ which converges to $(a(t), b(t))$ in $C[0, T]$ such that (by employing a diagonal process)

$$a_j^{N_k} \rightarrow a_j, \quad b_j^{N_k} \rightarrow b_j, \quad \text{in } C(0, T) \text{ as } k \rightarrow \infty, \quad \forall j \geq 0.$$

The last step is to show that the limit $(a(t), b(t))$ is a Leray-Hopf solution of (1.2a)-(1.2b). Replacing N by N_k in (3.2) and taking the limit $k \rightarrow \infty$, we see that $(a(t), b(t))$ satisfies the integral system

$$\begin{aligned} a_j(t) &= a_j^0 + \int_0^t \left(-\lambda_j^2 a_j(\tau) - \lambda_j^\theta a_j(\tau) a_{j+1}(\tau) + \lambda_{j-1}^\theta (a_{j-1}(\tau))^2 \right. \\ & \quad \left. - \lambda_j^\theta b_j(\tau) b_{j+1}(\tau) + \lambda_{j-1}^\theta (b_{j-1}(\tau))^2 + f_j(\tau) \right) d\tau, \\ b_j(t) &= b_j^0 + \int_0^t \left(-\lambda_j^2 b_j(\tau) + \lambda_j^\theta a_j(\tau) b_{j+1}(\tau) - \lambda_j^\theta b_j(\tau) a_{j+1}(\tau) \right) d\tau, \end{aligned}$$

for all $j \geq 0$. Hence, $(a(t), b(t))$ satisfies system (1.2a)-(1.2b). Moreover, $a_j, b_j \in C^1[0, T]$ for all $j \geq 0$. In addition, taking the limit in the energy inequality (3.11) yields

$$a, b \in L^\infty(0, T; l^2) \cap L^2(0, T; H^1).$$

Notice that $a^{N_k} \in L^\infty(0, T; l^2) \cap L^2(0, T; H^1)$ for all k and N_k . Thus, the sequence $\{a_j^{N_k}\}_{k \geq 1}$ converges weakly in $L^2(0, T)$ for any fixed $j \geq 0$, and the limit coincides with a_j . Consequently, we have

$$\sum_{j=0}^{\infty} \int_0^t f_j(\tau) a_j^{N_k}(\tau) d\tau \rightarrow \sum_{j=0}^{\infty} \int_0^t f_j(\tau) a_j(\tau) d\tau, \quad \text{as } k \rightarrow \infty, \quad \forall t \in [0, T]. \quad (3.12)$$

Passing the limit in (3.10) and applying (3.12), it leads to the energy inequality satisfied by the limit $(a(t), b(t))$

$$\begin{aligned} & \sum_{j=0}^{\infty} (a_j(t)^2 + b_j(t)^2) + 2 \sum_{j=0}^{\infty} \int_0^t \lambda_j^2 (a_j(\tau)^2 + b_j(\tau)^2) d\tau \\ & \leq \sum_{j=0}^{\infty} \left((a_j^0)^2 + (b_j^0)^2 \right) + 2 \sum_{j=0}^{\infty} \int_0^t f_j(\tau) a_j(\tau) d\tau. \end{aligned}$$

It completes the proof of Theorem 1.1.

4. WEAK-STRONG UNIQUENESS

In order to show the weak-strong uniqueness, a standard argument involving Grönwall's inequality will be applied to the difference $(a(t) - u(t), b(t) - v(t))$ of the two solutions $(a(t), b(t))$ and $(u(t), v(t))$.

Proof of Theorem 1.2: As in the previous section, we set $\kappa_1 = -\kappa_2 = 1$. We start with the energy balance through the j -th shell

$$\begin{aligned} & \frac{d}{dt} \left((a_j - u_j)^2 + (b_j - v_j)^2 \right) \\ & = \left((2a_j a'_j + 2b_j b'_j) + (2u_j u'_j + 2v_j v'_j) \right) \\ & \quad - \left((2a_j u'_j + 2a'_j u_j) + (2b_j v'_j + 2b'_j v_j) \right) \end{aligned} \quad (4.1)$$

and continue to estimate the four groups on the right hand side. In view of equations (1.2a)-(1.2b) satisfied by (a_j, b_j) and (u_j, v_j) respectively, we have

$$a_j a'_j + b_j b'_j = -\lambda_j^2 (a_j^2 + b_j^2) - \lambda_j^\theta (a_j^2 + b_j^2) a_{j+1} + \lambda_{j-1}^\theta (a_{j-1}^2 + b_{j-1}^2) a_j + f_j a_j, \quad (4.2)$$

$$u_j u'_j + v_j v'_j = -\lambda_j^2 (u_j^2 + v_j^2) - \lambda_j^\theta (u_j^2 + v_j^2) u_{j+1} + \lambda_{j-1}^\theta (u_{j-1}^2 + v_{j-1}^2) u_j + f_j u_j, \quad (4.3)$$

$$\begin{aligned} (a_j u_j)' & = -2\lambda_j^2 a_j u_j - \lambda_j^\theta a_j u_j a_{j+1} - \lambda_j^\theta a_j u_j u_{j+1} \\ & \quad - \lambda_j^\theta b_j u_j b_{j+1} - \lambda_j^\theta a_j v_j v_{j+1} + \lambda_{j-1}^\theta a_{j-1}^2 u_j + \lambda_{j-1}^\theta b_{j-1}^2 u_j \\ & \quad + \lambda_{j-1}^\theta u_{j-1}^2 a_j + \lambda_{j-1}^\theta v_{j-1}^2 a_j + f_j (a_j + u_j), \end{aligned} \quad (4.4)$$

$$(b_j v_j)' = -2\lambda_j^2 b_j v_j + \lambda_j^\theta a_j v_j b_{j+1} + \lambda_j^\theta b_j u_j v_{j+1} - \lambda_j^\theta b_j v_j a_{j+1} - \lambda_j^\theta b_j v_j u_{j+1}. \quad (4.5)$$

Combining (4.1)-(4.5) and grouping the terms appropriately gives

$$\begin{aligned}
& \frac{d}{dt} ((a_j - u_j)^2 + (b_j - v_j)^2) + 2\lambda_j^2(a_j - u_j)^2 + 2\lambda_j^2(b_j - v_j)^2 \\
&= -\lambda_j^\theta(a_j^2 + b_j^2)a_{j+1} + \lambda_{j-1}^\theta(a_{j-1}^2 + b_{j-1}^2)a_j \\
&\quad - \lambda_j^\theta(u_j^2 + v_j^2)u_{j+1} + \lambda_{j-1}^\theta(u_{j-1}^2 + v_{j-1}^2)u_j \\
&\quad + (-\lambda_j^\theta a_j u_j a_{j+1} - \lambda_j^\theta a_j u_j u_{j+1} + \lambda_{j-1}^\theta a_{j-1}^2 u_j + \lambda_{j-1}^\theta u_{j-1}^2 a_j) \\
&\quad + (-\lambda_j^\theta b_j u_j b_{j+1} - \lambda_j^\theta a_j v_j v_{j+1} + \lambda_j^\theta a_j v_j b_{j+1} + \lambda_j^\theta b_j u_j v_{j+1}) \\
&\quad + (\lambda_{j-1}^\theta b_{j-1}^2 u_j + \lambda_{j-1}^\theta v_{j-1}^2 a_j - \lambda_j^\theta b_j v_j a_{j+1} - \lambda_j^\theta b_j v_j u_{j+1}).
\end{aligned} \tag{4.6}$$

We further rearrange the terms in the last three parentheses of (4.6) to create terms in differences, for instance, $a_j - u_j$ and $b_j - v_j$. Shifting the sub-index j to $j+1$ in the last two terms of

$$(-\lambda_j^\theta a_j u_j a_{j+1} - \lambda_j^\theta a_j u_j u_{j+1} + \lambda_{j-1}^\theta a_{j-1}^2 u_j + \lambda_{j-1}^\theta u_{j-1}^2 a_j),$$

we have

$$\begin{aligned}
& -\lambda_j^\theta a_j u_j a_{j+1} - \lambda_j^\theta a_j u_j u_{j+1} + \lambda_{j-1}^\theta a_{j-1}^2 u_j + \lambda_{j-1}^\theta u_{j-1}^2 a_j \\
&= -\lambda_j^\theta a_j u_j a_{j+1} - \lambda_j^\theta a_j u_j u_{j+1} + \lambda_j^\theta a_j^2 u_{j+1} + \lambda_j^\theta u_j^2 a_{j+1} \\
&= \lambda_j^\theta (a_j u_{j+1} (a_j - u_j) - u_j a_{j+1} (a_j - u_j)) \\
&= \lambda_j^\theta (a_j - u_j) (a_j u_{j+1} - a_j a_{j+1} + a_j a_{j+1} - u_j a_{j+1}) \\
&= -\lambda_j^\theta a_j (a_j - u_j) (a_{j+1} - u_{j+1}) + \lambda_j^\theta (a_j - u_j)^2 a_{j+1}.
\end{aligned} \tag{4.7}$$

Similarly, with a shift of sub-index in the first two terms of

$$(\lambda_{j-1}^\theta b_{j-1}^2 u_j + \lambda_{j-1}^\theta v_{j-1}^2 a_j - \lambda_j^\theta b_j v_j a_{j+1} - \lambda_j^\theta b_j v_j u_{j+1}),$$

we obtain

$$\begin{aligned}
& \lambda_j^\theta b_j^2 u_{j+1} + \lambda_j^\theta v_j^2 a_{j+1} - \lambda_j^\theta b_j v_j a_{j+1} - \lambda_j^\theta b_j v_j u_{j+1} \\
&= \lambda_j^\theta b_j u_{j+1} (b_j - v_j) - \lambda_j^\theta v_j a_{j+1} (b_j - v_j) \\
&= \lambda_j^\theta (b_j - v_j) (b_j u_{j+1} - b_j a_{j+1} + b_j a_{j+1} - v_j a_{j+1}) \\
&= -\lambda_j^\theta b_j (b_j - v_j) (a_{j+1} - u_{j+1}) + \lambda_j^\theta (b_j - v_j)^2 a_{j+1}.
\end{aligned} \tag{4.8}$$

We rearrange the terms of

$$(-\lambda_j^\theta b_j u_j b_{j+1} - \lambda_j^\theta a_j v_j v_{j+1} + \lambda_j^\theta a_j v_j b_{j+1} + \lambda_j^\theta b_j u_j v_{j+1})$$

as

$$\begin{aligned}
& -\lambda_j^\theta b_j u_j b_{j+1} - \lambda_j^\theta a_j v_j v_{j+1} + \lambda_j^\theta a_j v_j b_{j+1} + \lambda_j^\theta b_j u_j v_{j+1} \\
&= \lambda_j^\theta a_j v_j (b_{j+1} - v_{j+1}) - \lambda_j^\theta b_j u_j (b_{j+1} - v_{j+1}) \\
&= \lambda_j^\theta (a_j v_j - a_j b_j + a_j b_j - b_j u_j) (b_{j+1} - v_{j+1}) \\
&= -\lambda_j^\theta a_j (b_j - v_j) (b_{j+1} - v_{j+1}) + \lambda_j^\theta b_j (a_j - u_j) (b_{j+1} - v_{j+1}).
\end{aligned} \tag{4.9}$$

Since $(a(t), b(t))$ and $(u(t), v(t))$ are Leray-Hopf solutions, we have that the following two series with telescope sums vanish,

$$\begin{aligned} \sum_{j=0}^{\infty} \int_0^t (-\lambda_j^\theta (a_j^2 + b_j^2) a_{j+1} + \lambda_{j-1}^\theta (a_{j-1}^2 + b_{j-1}^2) a_j) d\tau &= 0, \\ \sum_{j=0}^{\infty} \int_0^t (-\lambda_j^\theta (u_j^2 + v_j^2) u_{j+1} + \lambda_{j-1}^\theta (u_{j-1}^2 + v_{j-1}^2) u_j) d\tau &= 0. \end{aligned} \quad (4.10)$$

Integrating (4.6) over $[0, t]$, taking the sum for $j \geq 0$, using the fact $a_{-1} = b_{-1} = u_{-1} = v_{-1} = 0$, shifting the sub-index in the terms with sub-index $j - 1$, and applying (4.7)-(4.10), we deduce

$$\begin{aligned} & \sum_{j=0}^{\infty} ((a_j(t) - u_j(t))^2 + (b_j(t) - v_j(t))^2) \\ & + 2 \sum_{j=0}^{\infty} \lambda_j^2 \int_0^t (a_j(\tau) - u_j(\tau))^2 + (b_j(\tau) - v_j(\tau))^2 d\tau \\ & = - \sum_{j=0}^{\infty} \int_0^t \lambda_j^\theta a_j(\tau) (a_j(\tau) - u_j(\tau)) (a_{j+1}(\tau) - u_{j+1}(\tau)) d\tau \\ & \quad - \sum_{j=0}^{\infty} \int_0^t \lambda_j^\theta b_j(\tau) (b_j(\tau) - v_j(\tau)) (a_{j+1}(\tau) - u_{j+1}(\tau)) d\tau \\ & \quad + \sum_{j=0}^{\infty} \int_0^t \lambda_j^\theta (a_j(\tau) - u_j(\tau))^2 a_{j+1}(\tau) d\tau \\ & \quad + \sum_{j=0}^{\infty} \int_0^t \lambda_j^\theta (b_j(\tau) - v_j(\tau))^2 a_{j+1}(\tau) d\tau \\ & \quad - \sum_{j=0}^{\infty} \int_0^t \lambda_j^\theta a_j(\tau) (b_j(\tau) - v_j(\tau)) (b_{j+1}(\tau) - v_{j+1}(\tau)) d\tau \\ & \quad + \sum_{j=0}^{\infty} \int_0^t \lambda_j^\theta b_j(\tau) (a_j(\tau) - u_j(\tau)) (b_{j+1}(\tau) - v_{j+1}(\tau)) d\tau. \end{aligned} \quad (4.11)$$

We claim that the series on the right hand side of (4.11) are well-defined. Indeed, since $(a(t), b(t))$ and $(u(t), v(t))$ are Leray-Hopf solutions, it is clear that

$$\sum_{j=0}^{\infty} \lambda_j^2 \int_0^t a_j^2(\tau) + b_j^2(\tau) d\tau < \infty, \quad \sum_{j=0}^{\infty} \lambda_j^2 \int_0^t u_j^2(\tau) + v_j^2(\tau) d\tau < \infty.$$

As a consequence, applying the assumption (1.3), we infer

$$\begin{aligned}
& \sum_{j=J}^{\infty} \int_0^t |\lambda_j^\theta a_j(\tau) (a_j(\tau) - u_j(\tau)) (a_{j+1}(\tau) - u_{j+1}(\tau))| d\tau \\
& \leq C \sum_{j=J}^{\infty} \int_0^t |\lambda_j^2 (a_j(\tau) - u_j(\tau)) (a_{j+1}(\tau) - u_{j+1}(\tau))| d\tau \\
& \leq 4C \sum_{j=J}^{\infty} \lambda_j^2 \int_0^t a_j^2(\tau) + b_j^2(\tau) + u_j^2(\tau) + v_j^2(\tau) d\tau \\
& < \infty
\end{aligned}$$

for a constant C . Other series can be shown to converge analogously. Next, we estimate these series starting from the J -th shell. We only need to show details for one of them, for instance, thanks to the assumption (1.3)

$$\begin{aligned}
& \sum_{j=J}^{\infty} \int_0^t |\lambda_j^\theta b_j(\tau) (b_j(\tau) - v_j(\tau)) (a_{j+1}(\tau) - u_{j+1}(\tau))| d\tau \\
& \leq C_0 \sum_{j=J}^{\infty} \int_0^t |\lambda_j^2 (b_j(\tau) - v_j(\tau)) (a_{j+1}(\tau) - u_{j+1}(\tau))| d\tau \\
& \leq \frac{C_0}{2} \sum_{j=J}^{\infty} \int_0^t \lambda_j^2 (b_j(\tau) - v_j(\tau))^2 d\tau + \frac{C_0}{2\lambda^2} \sum_{j=J}^{\infty} \int_0^t \lambda_{j+1}^2 (a_{j+1}(\tau) - u_{j+1}(\tau))^2 d\tau.
\end{aligned}$$

Similarly, the other series have the estimates

$$\begin{aligned}
& \sum_{j=J}^{\infty} \int_0^t |\lambda_j^\theta a_j(\tau) (a_j(\tau) - u_j(\tau)) (a_{j+1}(\tau) - u_{j+1}(\tau))| d\tau \\
& \leq \frac{C_0}{2} \sum_{j=J}^{\infty} \int_0^t \lambda_j^2 (a_j(\tau) - u_j(\tau))^2 d\tau + \frac{C_0}{2\lambda^2} \sum_{j=J}^{\infty} \int_0^t \lambda_{j+1}^2 (a_{j+1}(\tau) - u_{j+1}(\tau))^2 d\tau, \\
& \sum_{j=J}^{\infty} \int_0^t |\lambda_j^\theta a_j(\tau) (b_j(\tau) - v_j(\tau)) (b_{j+1}(\tau) - v_{j+1}(\tau))| d\tau \\
& \leq \frac{C_0}{2} \sum_{j=J}^{\infty} \int_0^t \lambda_j^2 (b_j(\tau) - v_j(\tau))^2 d\tau + \frac{C_0}{2\lambda^2} \sum_{j=J}^{\infty} \int_0^t \lambda_{j+1}^2 (b_{j+1}(\tau) - v_{j+1}(\tau))^2 d\tau, \\
& \sum_{j=J}^{\infty} \int_0^t |\lambda_j^\theta b_j(\tau) (a_j(\tau) - u_j(\tau)) (b_{j+1}(\tau) - v_{j+1}(\tau))| d\tau \\
& \leq \frac{C_0}{2} \sum_{j=J}^{\infty} \int_0^t \lambda_j^2 (a_j(\tau) - u_j(\tau))^2 d\tau + \frac{C_0}{2\lambda^2} \sum_{j=J}^{\infty} \int_0^t \lambda_{j+1}^2 (b_{j+1}(\tau) - v_{j+1}(\tau))^2 d\tau, \\
& \sum_{j=J}^{\infty} \int_0^t |\lambda_j^\theta (a_j(\tau) - u_j(\tau))^2 a_{j+1}(\tau)| d\tau \leq C_0 \lambda^{2-\theta} \sum_{j=J}^{\infty} \int_0^t \lambda_j^2 (a_j(\tau) - u_j(\tau))^2 d\tau, \\
& \sum_{j=J}^{\infty} \int_0^t |\lambda_j^\theta (b_j(\tau) - v_j(\tau))^2 a_{j+1}(\tau)| d\tau \leq C_0 \lambda^{2-\theta} \sum_{j=J}^{\infty} \int_0^t \lambda_j^2 (b_j(\tau) - v_j(\tau))^2 d\tau.
\end{aligned}$$

Combining the estimates above and (4.11), we obtain

$$\begin{aligned}
& \sum_{j=0}^{\infty} ((a_j(t) - u_j(t))^2 + (b_j(t) - v_j(t))^2) \\
& + 2 \sum_{j=0}^{\infty} \lambda_j^2 \int_0^t (a_j(\tau) - u_j(\tau))^2 + (b_j(\tau) - v_j(\tau))^2 d\tau \\
& \leq C_1 \sum_{j=0}^J \int_0^t (a_j(\tau) - u_j(\tau))^2 + (b_j(\tau) - v_j(\tau))^2 d\tau \\
& + C_0 (1 + \lambda^{2-\theta} + \lambda^{-2}) \sum_{j=J}^{\infty} \lambda_j^2 \int_0^t (a_j(\tau) - u_j(\tau))^2 + (b_j(\tau) - v_j(\tau))^2 d\tau
\end{aligned} \tag{4.12}$$

where the constant C_1 is given by

$$C_1 = 32\lambda_J^\theta \sup_{0 \leq j \leq J+1} (\|a_j\|_C + \|b_j\|_C) \leq 32\lambda_J^\theta (\|a^0\|_{l^2} + \|b^0\|_{l^2}).$$

We take C_0 such that $C_0 (1 + \lambda^{2-\theta} + \lambda^{-2}) \leq 2$. Hence, it follows from (4.12) that

$$\begin{aligned}
& \sum_{j=0}^{\infty} ((a_j(t) - u_j(t))^2 + (b_j(t) - v_j(t))^2) \\
& \leq C_1 \sum_{j=0}^{\infty} \int_0^t (a_j(\tau) - u_j(\tau))^2 + (b_j(\tau) - v_j(\tau))^2 d\tau.
\end{aligned}$$

Therefore, Grönwall's inequality implies that

$$a_j \equiv u_j, \quad b_j \equiv v_j, \quad \forall j \geq 0.$$

□

Proof of Theorem 1.3: Since $0 < \theta \leq 2$, for any Leray-Hopf solution $(a(t), b(t))$, there exists $J > 0$ such that

$$|a_j(t)| \leq C_0 \lambda_j^{2-\theta}, \quad |b_j(t)| \leq C_0 \lambda_j^{2-\theta}, \quad \forall j \geq J.$$

That is, assumption (1.3) is satisfied and hence uniqueness follows.

□

5. NON-UNIQUENESS OF LERAY-HOPF SOLUTIONS FOR $\theta > 2$

We prove Theorem 1.4 in this section. We adapt the construction scheme for the dyadic NSE in [14] in order to construct a solution $(a(t), b(t))$ of (1.2a)-(1.2b) with zero initial data such that both $a(t)$ and $b(t)$ are non-vanishing. We first present the proof for the special case $\kappa_1 = -\kappa_2 = 1$ and then point out modifications to prove other cases when changing the signs of κ_1 and κ_2 .

Fix $T = \frac{1}{\lambda^2 - 1}$. Define

$$t_j = \lambda_j^{-2} T, \quad j \geq 0.$$

We note

$$\begin{aligned}
t_{j-1} - t_j &= \lambda_j^{-2}, \quad j \geq 1, \\
(0, T) &= \cup_{j=1}^{\infty} [t_j, t_{j-1}).
\end{aligned}$$

For $p, q \in C_0^\infty(0, 1)$ and constant $\rho > \lambda^\theta$, we construct a_j and b_j as follows,

$$a_j(t) = \begin{cases} 0, & t < t_{j+1}, \\ \lambda_{j+1}^{2-\theta} p(\lambda_{j+1}^2(t - t_{j+1})), & t_{j+1} < t < t_j, \\ -\lambda_j^{2-\theta} q(\lambda_j^2(t - t_j)), & t_j < t < t_{j-1}, \\ 0, & t > t_{j-1}. \end{cases} \quad (5.1)$$

$$b_j(t) = \begin{cases} 0, & t < t_{j+1}, \\ \rho^{-j-1} h_1(\lambda_{j+1}^2(t - t_{j+1})), & t_{j+1} < t < t_j, \\ \rho^{-j} h_2(\lambda_j^2(t - t_j)), & t_j < t < t_{j-1}, \\ \rho^{-j+1} h_3(\lambda_{j-1}^2(t - t_{j-1})), & t_{j-1} < t < t_{j-2}, \\ \rho^{-j+1} h_3(1)e^{-\lambda_j^2(t - t_{j-2})}, & t > t_{j-2}, \end{cases} \quad (5.2)$$

such that h_1, h_2 and h_3 satisfy the ODE system on $[0, 1]$

$$\frac{d}{dt} h_1 + (\lambda^{-2} - \lambda^{-\theta} q) h_1 - \lambda^{-\theta} p h_2 = 0, \quad (5.3a)$$

$$\frac{d}{dt} h_2 + h_2 + q h_3 = 0, \quad (5.3b)$$

$$\frac{d}{dt} h_3 + \lambda^2 h_3 = 0, \quad (5.3c)$$

$$h_1(0) = 0, \quad h_2(0) = c_0, \quad h_3(0) = d_0. \quad (5.3d)$$

In addition, we assume

$$h_1(1) = \rho c_0, \quad h_2(1) = \rho d_0. \quad (5.4)$$

With (a_j, b_j) constructed in (5.1)-(5.2), we define the forcing by

$$f_j = \frac{d}{dt} a_j + \lambda_j^2 a_j + \lambda_j^\theta a_j a_{j+1} + \lambda_j^\theta b_j b_{j+1} - \lambda_{j-1}^\theta a_{j-1}^2 - \lambda_{j-1}^\theta b_{j-1}^2 \quad (5.5)$$

for all $j \geq 0$.

Lemma 5.1. *Let a_j and b_j be constructed as in (5.1)-(5.2). Then, the following properties hold:*

- (i) $a_j \in C_0^\infty(0, T)$ for all $j \geq 0$;
- (ii) b_j are piecewise smooth and $b_j \in H^1(0, T)$ for all $j \geq 0$;
- (iii)

$$a_j(0) = b_j(0) = 0, \quad \forall j \geq 0;$$

(iv)

$$a_j(t) = O(\lambda_j^{2-\theta}), \quad a'_j(t) = O(\lambda_j^{4-\theta}), \quad b_j(t) = O(\rho^{-j}), \quad j \rightarrow \infty.$$

Proof: Since $p, q \in C_0^\infty(0, T)$, we only need to verify the values of the functions at t_{j+1}, t_j, t_{j-1} and t_{j-2} . The functions b_j are piecewise smooth and continuous at these times; hence, $b_j \in H^1(0, T)$. It is obvious to see (iii) and (iv) from (5.1)-(5.2). \square

Lemma 5.2. *The functions a_j and b_j defined in (5.1)-(5.2) satisfy*

$$\frac{d}{dt} b_j + \lambda_j^2 b_j - \lambda_j^\theta a_j b_{j+1} + \lambda_j^\theta b_j a_{j+1} = 0$$

for all $j \geq 0$.

Proof: Denote

$$A_j(t) = \frac{d}{dt}b_j(t) + \lambda_j^2 b_j(t) - \lambda_j^\theta a_j(t)b_{j+1}(t) + \lambda_j^\theta b_j(t)a_{j+1}(t).$$

For $t < t_{j+1}$, we see $a_j(t) = b_j(t) = 0$ from (5.1) and (5.2), and hence $A_j(t) = 0$.

For $t_{j+1} < t < t_j$, we denote $\tau = \lambda_{j+1}^2(t - t_{j+1}) \in (0, 1)$. It follows from (5.1)-(5.2) that

$$\begin{aligned} A_j(t) &= \rho^{-j-1}\lambda_{j+1}^2 \frac{d}{d\tau}h_1(\tau) + \rho^{-j-1}\lambda_{j+1}^2 h_1(\tau) \\ &\quad - \rho^{-j-1}\lambda_{j+1}^2 \lambda^{-\theta} p(\tau)h_2(\tau) - \rho^{-j-1}\lambda_{j+1}^2 \lambda^{-\theta} q(\tau)h_1(\tau) \\ &= \rho^{-j-1}\lambda_{j+1}^2 \left(\frac{d}{d\tau}h_1(\tau) + \lambda^{-2}h_1(\tau) - \lambda^{-\theta}q(\tau)h_1(\tau) - \lambda^{-\theta}p(\tau)h_2(\tau) \right) \\ &= 0 \end{aligned}$$

thanks to (5.3a).

For $t_j < t < t_{j-1}$, we denote $\tau = \lambda_j^2(t - t_j) \in (0, 1)$. We note $a_{j+1}(t) = 0$ by (5.1). Moreover, we have

$$\begin{aligned} A_j(t) &= \rho^{-j}\lambda_j^2 \frac{d}{d\tau}h_2(\tau) + \rho^{-j}\lambda_j^2 h_2(\tau) + \rho^{-j}\lambda_j^2 q(\tau)h_3(\tau) \\ &= \rho^{-j}\lambda_j^2 \left(\frac{d}{d\tau}h_2(\tau) + h_2(\tau) + q(\tau)h_3(\tau) \right) \\ &= 0 \end{aligned}$$

where we applied (5.3b).

For $t_{j-1} < t < t_{j-2}$, we denote $\tau = \lambda_{j-1}^2(t - t_{j-1}) \in (0, 1)$. On this interval, we have $a_j(t) = a_{j+1}(t) = 0$, and by (5.2) and (5.3c)

$$A_j(t) = \rho^{-j+1}\lambda_{j-1}^2 \frac{d}{d\tau}h_3(\tau) + \rho^{-j+1}\lambda_{j-1}^2 h_3(\tau) = 0.$$

For $t > t_{j-2}$, we note $a_j(t) = a_{j+1}(t) = 0$, and

$$A_j(t) = -\rho^{-j+1}\lambda_j^2 h_3(1)e^{-\lambda_j^2(t-t_{j-2})} + \rho^{-j+1}\lambda_j^2 h_3(1)e^{-\lambda_j^2(t-t_{j-2})} = 0.$$

□

Lemma 5.3. *The forcing $f = \{f_j(t)\}_{j \geq 0}$ constructed in (5.5) satisfies*

$$\sum_{j=0}^{\infty} \lambda_j^{-2} \int_0^T f_j^2(t) dt < \infty.$$

Proof: It follows from (5.1)-(5.2), (5.5) and straightforward computations that

$$f_j(t) = \begin{cases} 0, & t < t_{j+1}, \\ O(\lambda_j^{4-\theta}), & t_{j+1} < t < t_{j-2}, \\ O(\lambda_j^{-\theta}), & t > t_{j-2}. \end{cases}$$

Hence,

$$\begin{aligned} \lambda_j^{-2} \int_0^T f_j^2(t) dt &= \lambda_j^{-2} \int_{t_{j+1}}^{t_{j-2}} O(\lambda_j^{8-2\theta}) dt + \lambda_j^{-2} \int_{t_{j-2}}^T O(\lambda_j^{-2\theta}) dt \\ &= O(\lambda_j^{4-2\theta}) + O(\lambda_j^{-2-2\theta}). \end{aligned}$$

Since $4 - 2\theta < 0$ for $\theta > 2$, it is clear that

$$\sum_{j=0}^{\infty} \lambda_j^{-2} \int_0^T f_j^2(t) dt \leq \sum_{j=0}^{\infty} (O(\lambda_j^{4-2\theta}) + O(\lambda_j^{-2-2\theta})) < \infty.$$

□

Lemma 5.4. *There exist functions $p, q \in C_0^\infty(0, 1)$ and constants c_0 and d_0 with $c_0^2 + d_0^2 \neq 0$ such that there exists a unique solution $h = (h_1, h_2, h_3)$ of system (5.3a)-(5.3d) satisfying (5.4) and $h \in C^\infty([0, 1]; \mathbb{R}^3)$.*

Proof: It is obvious from (5.3c) and the initial data $h_3(0) = d_0$ that

$$h_3(t) = d_0 e^{-\lambda^2 t}.$$

It then follows from (5.3b) and $h_2(0) = c_0$ that

$$h_2(t) = c_0 e^{-t} - \int_0^t e^{s-t} q(s) h_3(s) ds = c_0 e^{-t} - d_0 e^{-t} \int_0^t e^{(1-\lambda^2)s} q(s) ds.$$

Since $h_2(1) = \rho d_0$, we have the constraint

$$c_0 - d_0 \int_0^1 e^{(1-\lambda^2)s} q(s) ds = \rho d_0. \quad (5.6)$$

In the end, we solve (5.3a) with $h_1(0) = 0$ as

$$h_1(t) = \int_0^t e^{-\int_s^t (\lambda^{-2} - \lambda^{-\theta} q(\tau)) d\tau} \lambda^{-\theta} p(s) h_2(s) ds.$$

The assumption $h_1(1) = \rho c_0$ gives another constraint,

$$\int_0^1 e^{-\int_s^1 (\lambda^{-2} - \lambda^{-\theta} q(\tau)) d\tau} \lambda^{-\theta} p(s) h_2(s) ds = \rho c_0. \quad (5.7)$$

We note that in the case of constant p and q , equations (5.6)-(5.7) have a unique solution (c_0, d_0) . Thus, by a continuity argument, we know that there exist functions $p, q \in C_0^\infty(0, 1)$ such that (5.6) and (5.7) are satisfied for some constants c_0, d_0 with $c_0^2 + d_0^2 \neq 0$. Since $p, q \in C_0^\infty(0, 1)$, it is clear that $h_1, h_2, h_3 \in C^\infty(0, 1)$. □

Proof of Theorem 1.4: Let $a = (a_j)_{j \geq 0}$ and $b = (b_j)_{j \geq 0}$ be constructed as in (5.1)-(5.2). According to Lemma 5.2, we have shown (a, b) satisfies the model (1.2a)-(1.2b) with $\kappa_1 = -\kappa_2 = 1$ and with forcing f_j defined in (5.5). It is shown in Lemma 5.1 that $b \in H^1(0, T)$. We are left to show that $a \in l^2 \cap H^1$ and (a, b) satisfies the energy estimate.

Since $\theta > 2$,

$$a_j(t) = \begin{cases} 0, & t < t_{j+1}, \\ O(\lambda_j^{2-\theta}), & t_{j+1} < t < t_{j-1}, \\ 0, & t > t_{j-1}, \end{cases}$$

which implies

$$\sup_{t \in [0, T]} \sum_{j \geq 0} a_j^2(t) < \infty.$$

Hence, we have $a \in l^2$.

Notice

$$t_{j-1} - t_j = \lambda_j^{-2}, \quad t_j - t_{j+1} = \lambda_{j+1}^{-2}, \quad T - t_{j-1} = \frac{1}{\lambda^2 - 1} \left(1 - \frac{1}{\lambda_{j-1}^2} \right) < \frac{1}{\lambda^2 - 1}.$$

As a consequence, we have

$$\int_0^T a_j^2(t) dt = \int_{t_{j+1}}^{t_{j-1}} a_j^2(t) dt = \int_{t_{j+1}}^{t_{j-1}} O(\lambda_j^{4-2\theta}) dt = O(\lambda_j^{2-2\theta}).$$

Hence,

$$\sum_{j=0}^{\infty} \int_0^T \lambda_j^2 a_j^2(t) dt = \sum_{j=0}^{\infty} O(\lambda_j^{4-2\theta}) < \infty$$

provided $\theta > 2$. That is, $a \in H^1(0, T)$.

Next, we show that (a, b) satisfies the energy identity. Since $(a(t), b(t))$ is a solution of (1.2a)-(1.2b), it follows

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} (a_j^2(t) + b_j^2(t)) &= -\lambda_j^2 a_j^2 - \lambda_j^2 b_j^2 - \lambda_j^\theta a_j^2 a_{j+1} + \lambda_{j-1}^\theta a_{j-1}^2 a_j \\ &\quad - \lambda_j^\theta b_j^2 a_{j+1} + \lambda_{j-1}^\theta b_{j-1}^2 a_j + f_j a_j; \end{aligned}$$

and hence

$$\begin{aligned} &\frac{1}{2} (a_j^2(t) + b_j^2(t)) - \frac{1}{2} (a_j^2(0) + b_j^2(0)) \\ &= - \int_0^t \lambda_j^2 a_j^2 dt - \int_0^t \lambda_j^2 b_j^2 dt - \int_0^t \lambda_j^\theta a_j^2 a_{j+1} dt + \int_0^t \lambda_{j-1}^\theta a_{j-1}^2 a_j dt \\ &\quad - \int_0^t \lambda_j^\theta b_j^2 a_{j+1} dt + \int_0^t \lambda_{j-1}^\theta b_{j-1}^2 a_j dt + \int_0^t f_j a_j dt. \end{aligned} \quad (5.8)$$

Again, notice

$$a_j(t) = \begin{cases} 0, & t < t_{j+1}, \\ O(\lambda_j^{2-\theta}), & t_{j+1} < t < t_{j-1}, \\ 0, & t > t_{j-1}, \end{cases}$$

and

$$a_{j+1}(t) = \begin{cases} 0, & t < t_{j+2}, \\ O(\lambda_{j+1}^{2-\theta}), & t_{j+2} < t < t_j, \\ 0, & t > t_j. \end{cases}$$

Thus, we have

$$\int_0^t \lambda_j^\theta |a_j^2 a_{j+1}| dt = \int_{t_{j+2}}^{t_{j-1}} \lambda_j^\theta O(\lambda_j^{6-3\theta}) dt = O(\lambda_j^{\theta-2+6-3\theta}) < \infty$$

since $\theta > 2$. Obviously, we also have

$$\int_0^t \lambda_{j-1}^\theta |a_{j-1}^2 a_j| dt < \infty.$$

On the other hand, we note

$$b_j(t) = \begin{cases} 0, & t < t_{j+1}, \\ O(\rho^{-j-1}), & t_{j+1} < t < t_j, \\ O(\rho^{-j}), & t_j < t < t_{j-1}, \\ O(\rho^{-j+1}), & t > t_{j-1}. \end{cases}$$

Thus, we have

$$\begin{aligned} \int_0^t \lambda_j^\theta |b_j^2 a_{j+1}| dt &= \int_{t_{j+1}}^{t_j} \lambda_j^\theta O(\rho^{-2j-2} \lambda_{j+1}^{2-\theta}) dt + \int_{t_j}^{t_{j-1}} \lambda_j^\theta O(\rho^{-2j} \lambda_{j+1}^{-\theta}) dt \\ &\quad + \int_{t_{j-1}}^T \lambda_j^\theta O(\rho^{-2j+2} \lambda_{j+1}^{-\theta}) dt \\ &= O(\rho^{-2j-2}) + O(\rho^{-2j} \lambda_j^{-2}) + O(\rho^{-2j+2}) \\ &< \infty. \end{aligned}$$

Similarly,

$$\int_0^t \lambda_{j-1}^\theta |b_{j-1}^2 a_j| dt < \infty.$$

Therefore, we can take the sum of (5.8) over $j \geq 0$ and obtain

$$\begin{aligned} &\frac{1}{2} \sum_{j=0}^{\infty} (a_j^2(t) + b_j^2(t)) - \frac{1}{2} \sum_{j=0}^{\infty} (a_j^2(0) + b_j^2(0)) \\ &= - \sum_{j=0}^{\infty} \int_0^t \lambda_j^2 a_j^2 dt - \sum_{j=0}^{\infty} \int_0^t \lambda_j^2 b_j^2 dt + \sum_{j=0}^{\infty} \int_0^t f_j a_j dt. \end{aligned}$$

Thus, we conclude (a, b) is a Leray-Hopf solution of (1.2a)-(1.2b) with zero initial data; however, $a \neq 0$ and $b \neq 0$. Non-uniqueness then follows. Indeed, for such forcing $f(t)$ as in (5.5), considering $b(t) \equiv 0$ in (1.2a)-(1.2b), the forced dyadic model of the NSE has a solution $\tilde{a}(t)$. Hence, $(\tilde{a}(t), 0)$ is a trivial solution for the dyadic MHD model (1.2a)-(1.2b). \square

Remark 5.5. When κ_1 and κ_2 take different signs, we can choose the same constructions for $(a_j(t), b_j(t))$ as in (5.1) and (5.2). The difference comes in the ODE system (5.3a)-(5.3d) for the profile functions h_1, h_2 and h_3 . For instance, if $\kappa_1 = \kappa_2 = 1$ the functions h_1, h_2 and h_3 satisfy the following system

$$\frac{d}{dt} h_1 + (\lambda^{-2} + \lambda^{-\theta} q) h_1 + \lambda^{-\theta} p h_2 = 0, \quad (5.9a)$$

$$\frac{d}{dt} h_2 + h_2 - q h_3 = 0, \quad (5.9b)$$

$$\frac{d}{dt} h_3 + \lambda^2 h_3 = 0, \quad (5.9c)$$

$$h_1(0) = 0, \quad h_2(0) = c_0, \quad h_3(0) = d_0, \quad (5.9d)$$

accompanied with

$$h_1(1) = \rho c_0, \quad h_2(1) = \rho d_0. \quad (5.10)$$

We note that the structure of system (5.9a)-(5.9d) remains similar to that of system (5.3a)-(5.3d). Thus in analogy with Lemma 5.4, it is not hard to show the existence

of a solution (h_1, h_2, h_3) to system (5.9a)-(5.9d) satisfying (5.10). The rest analysis of Section 5 also holds for system (1.2a)-(1.2b) with $\kappa_1 = \kappa_2 = 1$.

6. UNIQUENESS AND NON-UNIQUENESS RESULTS FOR THE DYADIC MHD MODEL WITH FRACTIONAL LAPLACIANS

6.1. Uniqueness. The weak-strong uniqueness stated in Theorem 1.9 under assumption (1.8) can be proved by following the steps described in Section 4. We briefly present the main steps and emphasize why assumption (1.8) is required for the uniqueness.

Let $(a(t), b(t))$ and $(u(t), v(t))$ be two Leray-Hopf solutions of (1.7) with $(a(t), b(t))$ satisfying (1.8). The difference of the two solutions satisfies the energy estimate

$$\begin{aligned}
& \sum_{j=0}^{\infty} ((a_j(t) - u_j(t))^2 + (b_j(t) - v_j(t))^2) \\
& + 2 \sum_{j=0}^{\infty} \int_0^t \lambda_j^{2\alpha} (a_j(\tau) - u_j(\tau))^2 + \lambda_j^{2\beta} (b_j(\tau) - v_j(\tau))^2 d\tau \\
= & - \sum_{j=0}^{\infty} \int_0^t \lambda_j a_j(\tau) (a_j(\tau) - u_j(\tau)) (a_{j+1}(\tau) - u_{j+1}(\tau)) d\tau \\
& - \sum_{j=0}^{\infty} \int_0^t \lambda_j b_j(\tau) (b_j(\tau) - v_j(\tau)) (a_{j+1}(\tau) - u_{j+1}(\tau)) d\tau \\
& + \sum_{j=0}^{\infty} \int_0^t \lambda_j (a_j(\tau) - u_j(\tau))^2 a_{j+1}(\tau) d\tau + \sum_{j=0}^{\infty} \int_0^t \lambda_j (b_j(\tau) - v_j(\tau))^2 a_{j+1}(\tau) d\tau \\
& - \sum_{j=0}^{\infty} \int_0^t \lambda_j a_j(\tau) (b_j(\tau) - v_j(\tau)) (b_{j+1}(\tau) - v_{j+1}(\tau)) d\tau \\
& + \sum_{j=0}^{\infty} \int_0^t \lambda_j b_j(\tau) (a_j(\tau) - u_j(\tau)) (b_{j+1}(\tau) - v_{j+1}(\tau)) d\tau \\
\equiv & I_1 + I_2 + I_3 + I_4 + I_5 + I_6.
\end{aligned} \tag{6.1}$$

With $(a(t), b(t))$ and $(u(t), v(t))$ being Leray-Hopf solutions of (1.7), it holds

$$\sum_{j=0}^{\infty} \int_0^t (\lambda_j^{2\alpha} a_j^2(\tau) + \lambda_j^{2\beta} b_j^2(\tau)) d\tau < \infty, \quad \sum_{j=0}^{\infty} \int_0^t (\lambda_j^{2\alpha} u_j^2(\tau) + \lambda_j^{2\beta} v_j^2(\tau)) d\tau < \infty.$$

Thus combining assumption (1.8) we know that all of the series on the right hand side of (6.1) are well-defined. Moreover, these series can be estimated in the following way by using (1.8). For example, we estimate I_2 thanks to the condition

$|b_j| \leq C_0 \lambda_j^{\alpha+\beta-1}$ of (1.8),

$$\begin{aligned} & \sum_{j=J}^{\infty} \int_0^t \lambda_j |b_j(\tau)(b_j(\tau) - v_j(\tau))(a_{j+1}(\tau) - u_{j+1}(\tau))| d\tau \\ & \leq C_0 \sum_{j=J}^{\infty} \int_0^t \left| \lambda_j^{\alpha+\beta} (b_j(\tau) - v_j(\tau))(a_{j+1}(\tau) - u_{j+1}(\tau)) \right| d\tau \\ & \leq \frac{C_0}{2} \sum_{j=J}^{\infty} \int_0^t \lambda_j^{2\beta} (b_j(\tau) - v_j(\tau))^2 d\tau + \frac{C_0}{2\lambda^{2\alpha}} \sum_{j=J}^{\infty} \int_0^t \lambda_{j+1}^{2\alpha} (a_{j+1}(\tau) - u_{j+1}(\tau))^2 d\tau. \end{aligned}$$

The term I_6 can be handled similarly. The condition $|a_j| \leq C_0 \lambda_j^{2\alpha-1}$ is posed to estimate I_1 and I_3 , and $|a_j| \leq C_0 \lambda_j^{2\beta-1}$ is for I_4 and I_5 . With the estimates, it follows from (6.1) that

$$\begin{aligned} & \sum_{j=0}^{\infty} ((a_j(t) - u_j(t))^2 + (b_j(t) - v_j(t))^2) \\ & + 2 \sum_{j=0}^{\infty} \lambda_j^2 \int_0^t (a_j(\tau) - u_j(\tau))^2 + (b_j(\tau) - v_j(\tau))^2 d\tau \\ & \leq C_1 \sum_{j=0}^J \int_0^t (a_j(\tau) - u_j(\tau))^2 + (b_j(\tau) - v_j(\tau))^2 d\tau \tag{6.2} \\ & + C_0 (1 + \lambda^{2\alpha-1} + \lambda^{-2\alpha}) \sum_{j=J}^{\infty} \int_0^t \lambda_j^{2\alpha} (a_j(\tau) - u_j(\tau))^2 d\tau \\ & + C_0 (1 + \lambda^{2\beta-1} + \lambda^{-2\beta}) \sum_{j=J}^{\infty} \int_0^t \lambda_j^{2\alpha} (b_j(\tau) - v_j(\tau))^2 d\tau \end{aligned}$$

with a constant $C_1 > 0$. The constant C_0 can be chosen small enough such that $C_0 (1 + \lambda^{2\alpha-1} + \lambda^{-2\alpha}) \leq 2$ and $C_0 (1 + \lambda^{2\beta-1} + \lambda^{-2\beta}) \leq 2$. Consequently, we have from (6.2) that

$$\begin{aligned} & \sum_{j=0}^{\infty} ((a_j(t) - u_j(t))^2 + (b_j(t) - v_j(t))^2) \\ & \leq C_1 \sum_{j=0}^{\infty} \int_0^t (a_j(\tau) - u_j(\tau))^2 + (b_j(\tau) - v_j(\tau))^2 d\tau. \end{aligned}$$

Grönwall's inequality immediately implies that $a_j \equiv u_j$ and $b_j \equiv v_j$ for all $j \geq 0$.

6.2. Non-uniqueness. The construction scheme to prove Theorem 1.11 is similar to that presented in Section 5. The main effort is to determine the scaling in constructing a_j and b_j . To be complete, we specify the constructions as follows. For $T = \frac{1}{\lambda^{2\beta-1}}$, we take the partition

$$t_j = \lambda_j^{-2\beta} T, \quad j \geq 0,$$

with

$$t_{j-1} - t_j = \lambda_j^{-2\beta}, \quad j \geq 1; \quad (0, T) = \cup_{j=1}^{\infty} [t_j, t_{j-1}).$$

For $p, q \in C_0^\infty(0, 1)$ and constant $\rho > \lambda$, we choose a_j and b_j as

$$a_j(t) = \begin{cases} 0, & t < t_{j+1}, \\ \lambda_{j+1}^{2\beta-1} p \left(\lambda_{j+1}^{2\beta} (t - t_{j+1}) \right), & t_{j+1} < t < t_j, \\ -\lambda_j^{2\beta-1} q \left(\lambda_j^{2\beta} (t - t_j) \right), & t_j < t < t_{j-1}, \\ 0, & t > t_{j-1}. \end{cases} \quad (6.3)$$

$$b_j(t) = \begin{cases} 0, & t < t_{j+1}, \\ \rho^{-j-1} h_1 \left(\lambda_{j+1}^{2\beta} (t - t_{j+1}) \right), & t_{j+1} < t < t_j, \\ \rho^{-j} h_2 \left(\lambda_j^{2\beta} (t - t_j) \right), & t_j < t < t_{j-1}, \\ \rho^{-j+1} h_3 \left(\lambda_{j-1}^{2\beta} (t - t_{j-1}) \right), & t_{j-1} < t < t_{j-2}, \\ \rho^{-j+1} h_3(1) e^{-\lambda_j^{2\beta} (t - t_{j-2})}, & t > t_{j-2}, \end{cases} \quad (6.4)$$

where h_1, h_2 and h_3 are functions satisfying the following ODE system on $[0, 1]$

$$\frac{d}{dt} h_1 + (\lambda^{-2\beta} - \lambda^{-1} q) h_1 - \lambda^{-1} p h_2 = 0, \quad (6.5a)$$

$$\frac{d}{dt} h_2 + h_2 + q h_3 = 0, \quad (6.5b)$$

$$\frac{d}{dt} h_3 + \lambda^{2\beta} h_3 = 0, \quad (6.5c)$$

$$h_1(0) = 0, \quad h_2(0) = c_0, \quad h_3(0) = d_0, \quad (6.5d)$$

$$h_1(1) = \rho c_0, \quad h_2(1) = \rho d_0. \quad (6.5e)$$

We take the forcing f_j as

$$f_j = \frac{d}{dt} a_j + \lambda_j^{2\alpha} a_j + \lambda_j a_j a_{j+1} + \lambda_j b_j b_{j+1} - \lambda_{j-1} a_{j-1}^2 - \lambda_{j-1} b_{j-1}^2 \quad (6.6)$$

for all $j \geq 0$.

For f_j defined by (6.6), we can show that $(a(t), b(t))$ with components constructed as in (6.3)-(6.4) is a Leray-Hopf solution of system (1.7) with non-vanishing $b(t)$. We state the main ingredients to prove Theorem 1.11 in the following Lemmas, the proof of which are omitted.

Lemma 6.1. *Let $\rho > \lambda > 1$ and $0 < \alpha \leq \beta < \frac{1}{2}$. The following properties hold for a_j and b_j as constructed in (6.3)-(6.4):*

- (i) $a_j \in C_0^\infty(0, T)$ and $a_j \in H^\alpha(0, T)$ for all $j \geq 0$;
- (ii) b_j are piecewise smooth and $b_j \in H^\beta(0, T)$ for all $j \geq 0$;
- (iii)

$$a_j(0) = b_j(0) = 0, \quad \forall j \geq 0;$$

(iv)

$$a_j(t) = O(\lambda_j^{2\beta-1}), \quad a'_j(t) = O(\lambda_j^{4\beta-1}), \quad b_j(t) = O(\rho^{-j}), \quad j \rightarrow \infty.$$

Lemma 6.2. *The functions a_j and b_j defined in (6.3)-(6.4) satisfy system (1.7) with forcing f_j defined by (6.6).*

Lemma 6.3. *Let $\rho > \lambda > 1$. Assume $0 < \alpha \leq \beta < \frac{1}{2}$ and $3\beta - \alpha < 1$. The forcing f_j defined by (6.6) satisfies*

$$\sum_{j=0}^{\infty} \lambda_j^{-2\alpha} \int_0^T f_j^2(t) dt < \infty.$$

Lemma 6.4. *Let $\rho > \lambda > 1$ and $0 < \alpha \leq \beta < \frac{1}{2}$. There exist functions $p, q \in C_0^\infty(0, 1)$ and constants c_0, d_0 satisfying $c_0^2 + d_0^2 \neq 0$ such that there exists a unique solution $h = (h_1, h_2, h_3) \in C^\infty([0, 1]; \mathbb{R}^3)$ of system (6.5a)-(6.5e).*

Lemma 6.5. *Let $\rho > \lambda > 1$ and $0 < \alpha \leq \beta < \frac{1}{2}$. Then $(a(t), b(t))$ satisfies the energy identity*

$$\begin{aligned} & \frac{1}{2} \sum_{j=0}^{\infty} (a_j^2(t) + b_j^2(t)) - \frac{1}{2} \sum_{j=0}^{\infty} (a_j^2(0) + b_j^2(0)) \\ &= - \sum_{j=0}^{\infty} \int_0^t \lambda_j^{2\alpha} a_j^2 dt - \sum_{j=0}^{\infty} \int_0^t \lambda_j^{2\beta} b_j^2 dt + \sum_{j=0}^{\infty} \int_0^t f_j a_j dt \end{aligned}$$

with f_j defined by (6.6).

ACKNOWLEDGEMENT

M. Dai is partially supported by the NSF grants DMS-1815069 and DMS-2009422. S. Friedlander is partially supported by the NSF grant DMS-1613135. S. Friedlander is grateful to IAS for its hospitality in 2020-2021. M. Dai is also grateful to IAS for its hospitality in 2021-2022.

REFERENCES

- [1] D. Barbato, F. Flandoli, and F. Morandin. *Energy dissipation and self-similar solutions for an unforced inviscid dyadic model*. Trans. Amer. Math. Soc., 363 (4): 1925–1946, 2011.
- [2] D. Barbato, F. Morandin, and M. Romito. *Smooth solutions for the dyadic model*. Nonlinearity, 24 (11): 3083–3097, 2011.
- [3] R. Beekie, T. Buckmaster, and V. Vicol. *Weak solutions of ideal MHD which do not conserve magnetic helicity*. Ann. of PDE, 6 (1), <https://doi.org/10.1007/s40818-020-0076-1>, 2020.
- [4] A. Cheskidov. *Blow-up in finite time for the dyadic model of the Navier-Stokes equations*. Trans. Amer. Math. Soc., 360 (10): 5101–5120, 2008.
- [5] A. Cheskidov and M. Dai. *Kolmogorov’s dissipation number and the number of degrees of freedom for the 3D Navier-Stokes equations*. Proceedings of the Royal Society of Edinburgh, Section A, Vol. 149, Issue 2: 429–446, 2019.
- [6] A. Cheskidov and S. Friedlander. *The vanishing viscosity limit for a dyadic model*. Physica D, 238:783–787, 2009.
- [7] A. Cheskidov, S. Friedlander, and N. Pavlović. *Inviscid dyadic model of turbulence: the fixed point and Onsager’s conjecture*. J. Math. Phys., 48 (6): 065503, 16, 2007.
- [8] A. Cheskidov, S. Friedlander, and N. Pavlović. *An inviscid dyadic model of turbulence: the global attractor*. Discrete Contin. Dyn. Syst., 26 (3): 781–794, 2010.
- [9] A. Cheskidov and R. Shvydkoy. *A unified approach to regularity problems for the 3D Navier-Stokes and Euler equations: the use of Kolmogorov’s dissipation range*. J. Math. Fluid Mech., Vol.16, Issue 2: 263–273, 2014.
- [10] M. Dai. *Blow-up of a dyadic model with intermittency dependence for the Hall MHD*. Physica D: Nonlinear Phenomena, Vol. 428: 133066, 2021.
- [11] M. Dai and S. Friedlander. *Dyadic models for ideal MHD*. arXiv:2104.09440. To appear in the Journal of Mathematical Fluid Mechanics.

- [12] V. N. Desnyanskiy and E. A. Novikov. *Evolution of turbulence spectra toward a similarity regime*. Izv. Akad. Nauk SSSR, Fiz. Atmos. Okeana, 10: 127–136, 1974.
- [13] N. Filonov. *Uniqueness of the Leray-Hopf solution for a dyadic model*. Transactions of the American Mathematical Society, Vol. 369 (12): 8663–8684, 2017.
- [14] N. Filonov and P. Khodunov. *Non-uniqueness of Leray-Hopf solutions for a dyadic model*. St. Petersburg Math. J., Vol. 32: 371–387, 2021.
- [15] S. Friedlander and N. Pavlović. *Blowup in a three-dimensional vector model for the Euler equations*. Comm. Pure Appl. Math., 57 (6): 705–725, 2004.
- [16] C. Gloaguen, J. Léorat, A. Pouquet and R. Grappin. *A scalar model for MHD turbulence*. Physica D.: Nonlinear Phenomena, 17(2):154–182, 1985.
- [17] E. Hopf. *Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen*. Math. Nachr., 4:213–231, 1951.
- [18] I. Jeong and D. Li. *A blow-up result for dyadic models of the Euler equations*. Communications in Mathematical Physics, 337:1027–1034, 2015.
- [19] N. Katz and N. Pavlović. *Finite time blow-up for a dyadic model of the Euler equations*. Trans. Amer. Math. Soc., 357 (2): 695–708, 2005.
- [20] A. Kiselev and A. Zlatoš. *On discrete models of the Euler equation*. Int. Math. Res. Not., 38: 2315–2339, 2005.
- [21] J. Leray. *Sur le mouvement d’un liquide visqueux emplissant l’espace*. Acta Math., 63(1):193–248, 1934.
- [22] L. Onsager. *Statistical hydrodynamics*. Nuovo Cimento (9), 6(Supplemento, 2(Convegno Internazionale di Meccanica Statistica)):279–287, 1949.
- [23] F. Plunian, R. Stepanov and P. Frick. *Shell models of magnetohydrodynamic turbulence*. Physics Reports, vol. 523, 2013.

DEPARTMENT OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, IL 60607, USA

Email address: mdai@uic.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089, USA

Email address: susanfri@usc.edu