The energy conservation and regularity for the Navier-Stokes equations

*Wenke Tan¹ and Zhaoyang Yin^{2,3}

¹Key Laboratory of Computing and Stochastic Mathematics (Ministry of Education), School of Mathematics and Statistics, Hunan Normal University, Changsha, Hunan 410081, P. R. China

²Department of Mathematics, Sun Yat-sen University, Guangzhou, 510275, China ³Faculty of Information Technology, Macau University of Science and Technology, Macau, China

Abstract

In this paper, we consider the energy conservation and regularity of the weak solution u to the Navier-Stokes equations in the endpoint case. We first construct a divergence-free field u(t, x) which satisfies $\lim_{t\to T} \sqrt{T-t} ||u(t)||_{BMO} < \infty$ and $\lim_{t\to T} \sqrt{T-t} ||u(t)||_{L^{\infty}} = \infty$ to demonstrate that the Type II singularity is admissible in the endpoint case $u \in L^{2,\infty}(BMO)$. Secondly, we prove that if a suitable weak solution u(t, x) satisfying $||u||_{L^{2,\infty}([0,T];BMO(\Omega))} < \infty$ for arbitrary $\Omega \subseteq \mathbb{R}^3$ then the local energy equality is valid on $[0,T] \times \Omega$. As a corollary, we also prove $||u||_{L^{2,\infty}([0,T];BMO(\mathbb{R}^3))} < \infty$ implies the global energy equality on [0,T]. Thirdly, we show that as the solution u approaches a finite blowup time T, the norm $||u(t)||_{BMO}$ must blow up at a rate faster than $\frac{c}{\sqrt{T-t}}$ with some absolute constant c > 0. Furthermore, we prove that if $||u_3||_{L^{2,\infty}([0,T];BMO(\mathbb{R}^3))} = M < \infty$ then there exists a small constant c_M depended on M such that if $||u_h||_{L^{2,\infty}([0,T];BMO(\mathbb{R}^3))} \leq c_M$ then u is regular on $(0,T] \times \mathbb{R}^3$.

Key words: Navier-Stokes equations, Suitable weak solutions, Energy equality, Regularity.

1 Introduction

We consider the incompressible Navier-Stokes in the domain $\Omega \subseteq \mathbb{R}^3$

(1.1)
$$\begin{cases} \partial_t u - \Delta u + u \cdot \nabla u + \nabla P = 0, \\ \nabla \cdot u = 0, \\ u(x,t)|_{\partial\Omega} = 0, \\ u(0,x) = u_0(x) \end{cases}$$

where the unknowns u, P denote the velocity vector field, pressure respectively.

It is well-known that if u_0 is smooth enough, then problems (1.1) have a unique solution on [0, T) for some T > 0; see, for example, [1, 9, 12, 13, 35] and the references therein. The concepts of weak solutions of (1.1) and their regularity were already introduced in the fundamental paper of Leray [1]. Pioneering works of Leray [1] and Hopf [11] showed the global existence of a weak solution called

^{*}Email Address: tanwenkeybfq@163.com (W. Tan), mcsyzy@mail.sysu.edu.cn (Z. Yin).

Leray-Hopf solution. In three dimensions, however, the question of regularity and uniqueness of weak solutions is an outstanding open problem in mathematical fluid mechanics.

On the one hand, in Leray's paper [1], he proved that if [0, T) is the maximal existence interval of a smooth solution, then for p > 3, there exists $c_p > 0$ such that

$$||u(t,\cdot)||_{L^p(\mathbb{R}^3)} \ge \frac{c_p}{|T-t|^{\frac{p-3}{2p}}}.$$

This means that as the solution u approaches a finite blowup time T, the norm $||u(t)||_{L^p}$ with 3 < p must blow up at a rate $\frac{c_p}{|T-t|^{\frac{p-3}{2p}}}$ with some absolute constant $c_p > 0$. In general, if u satisfies

$$||u(t,\cdot)||_{L^{p}(\mathbb{R}^{3})} \leq \frac{C}{|T-t|^{\frac{p-3}{2p}}}$$

the regularity of the solution at t = T remains unknown. The well-known Ladyzhenskaya-Prodi-Serrin criteria [21, 26, 29] showed that if $u \in L^r([0,T], L^s(\mathbb{R}^3))$ for $\frac{2}{r} + \frac{3}{s} \leq 1, s > 3$ then u is regular on [0, T]. For the endpoint case p = 3, Escauriaza, Serengin and Sverak [6] proved that the $L^{\infty}L^3$ solutions are smooth. This result was improved by Tao [33] showed that as the solution u approaches a finite blowup time T, the critical norm $||u(t)||_{L^3}$ must blow up at a rate $(\log \log \log \log \frac{1}{T-t})^c$ with some absolute constant c > 0. The other endpoint case $p = \infty$ was improved by Kozono and Taniuchi [14] proved that $u \in L^2([0,T]; BMO(\mathbb{R}^3))$ implies the regularity of the solution u to (1.1). Since the condition $||u(t)||_{L^p(\mathbb{R}^3)} \leq \frac{c_p}{|T-t|^{\frac{p-3}{2p}}}$ only requires $u \in L^{q,\infty}(L^p)$ for $\frac{3}{p} + \frac{2}{q} = 1, p \geq 3$, it is natural to generalize the classical Ladyzhenskaya-Prodi-Serrin type criterion into Lorentz spaces. In [15], Kim and Kozono proved the local boundedness of a weak solution u under the assumption that $||u||_{L^{r,\infty}([0,T];L^{s,\infty}(\mathbb{R}^3))}$ is sufficiently small for some (r,s) with $\frac{2}{r} + \frac{3}{s} = 1$ and $3 \leq s < \infty$. The limiting case of the regularity criteria derived by Kim and Kozono was proved by He and Wang [10] i.e. any weak solution u to the Navier-Stokes equations is regular under the assumption that $||u||_{L^{2,\infty}([0,T];L^{\infty}(\mathbb{R}^3))}$ is sufficiently small. This results were improved by Wang and Zhang [36] which showed that $||u_3||_{L^{r,\infty}([0,T];L^{s,\infty}(\mathbb{R}^3))} \leq M$ and $||u_h||_{L^{r,\infty}([0,T];L^{s,\infty}(\mathbb{R}^3))} \leq c_M$ with $\frac{2}{r} + \frac{3}{s} = 1$ and $3 < s \leq \infty$ imply the regularity of the suitable weak solution u to Navier-Stokes equations, where $u_h = (u_1, u_2)$, $u = (u_h, u_3)$ and c_M is a small constant depending on M. Collecting the results of [14, 10, 36], one should notice that the regularity in the endpoint case $u \in L^{2,\infty}BMO$ is left open. It is worth pointing out that although the spaces $L^{2,\infty}L^{\infty}$ and $L^{2,\infty}BMO$ are on the same scale under the scaling (1.6), but we can construct a divergence free field u(t,x) which satisfies $\lim_{t\to T} \sqrt{T-t} ||u(t)||_{BMO} < \infty$ and $\lim_{t\to T} \sqrt{T-t} ||u(t)||_{L^{\infty}} = \infty$ (see Proposition 1.1). It is surprising to some extent because the

conditions $\lim_{t\to T} \sqrt{T-t} ||u(t)||_{BMO} < \infty$ and $\lim_{t\to T} \sqrt{T-t} ||u(t)||_{L^{\infty}} = \infty$ implies that some Type II blow-up are admissible in the invariant space $L^{2,\infty}BMO$. Inspiring by the above observation, it is of interest to investigate the blow-up rate for $||u(t)||_{BMO}$ as the solution u approaches a finite blow-up time T. This is one of our goals in this paper

On the other hand, it is well-known that the Leray-Hopf weak solutions are weak continuous in L^2 space, but the strong continuity of u in L^2 space is still an open problem in the mathematic theory of Navier-Stokes equations. A sufficient condition for strong continuity of u in L^2 space is the energy equality, as should be expected from the physical point of view. The question of energy equality has of course also been extensively studied for Leray-Hopf solutions of the 3-dimensional Navier-Stokes equations. Lions [23] and Ladyzhenskaya [20] proved independently that such solutions satisfy

the (global) energy equality under the additional assumption $u \in L^4L^4$. Shinbrot in [30] proved the energy equality under the extrapolated version of the Lions-Ladyzhenskaya condition, namely $\frac{2}{r} + \frac{2}{s} \leq 1$ for $s \geq 4$. The endpoint case $u \in L^2 L^\infty$ was generalized to $u \in L^2(BMO)$ by Kozono and Taniuchi [14]. Kukavica [16] proved sufficiency of the weaker but dimensionally equivalent criterion $P \in L^2 L^2$. Cheskidov, Friedlander and Shvydkoy [5] proved energy equality for $u \in L^3 D(A^{\frac{5}{12}})$ on a bounded domain; an extension to exterior domains was proved in [8] by Farwig and Taniuchi. (Here A denotes the Stokes operator.) Seregin and Šverák [27] proved energy equality (regularity, in fact) for suitable weak solutions whose associated pressure is bounded from below in some sense; this paper makes use of the low-dimensionality of the singular set for suitable weak solutions that is guaranteed by the celebrated Caffarelli-Kohn-Nirenberg Theorem [3]. Leslie and Shvydkoy [17]; Shvydkoy [31] proved energy equality under assumptions on the size and (or structure of the singularity set) in addition to the integrability of the solution. Noticing the gap between the energy conservation criteria $\frac{1}{q} + \frac{1}{p} = \frac{1}{2}$ and the regularity criteria $\frac{2}{q} + \frac{3}{p} = 1$, it is natural to consider the energy conservation at blow-up time. Recently, Leslie and Shvydkoy [18] proved that any solution to the 3-dimensional Navier-Stokes Equations in \mathbb{R}^3 which is Type I (i.e. $||u(t)||_{L^{\infty}} \leq \frac{C}{\sqrt{T-t}}$) in time must satisfy the energy equality at the first blowup time T. This result was improved by Cheskidov and Luo [24] by showing $||u(t)||_{L^{2,\infty}(B^0_{\infty,\infty})} < \infty$ implies energy equality. Based the example constructed in Proposition 1.1, we thus consider the energy conservation at some potential Type II blow-up time T satisfying $\limsup_{t\to T^-} \sqrt{T-t} ||u(t)||_{BMO} < \infty \text{ and } \limsup_{t\to T^-} \sqrt{T-t} ||u(t)||_{L^{\infty}} = \infty.$

The tool that we use to address the question of energy conservation is the energy measure which was first introduced in [32]. The potential failure of energy equality for a solution u of (1.1) can be quantified using a so-called energy measure \mathcal{E} , which was defined to be the weak-* limit of the measures $|u(t)|^2 dx$ as t approaches the first possible blowup time. Precisely, assume [0, T) is the maximal existence interval of a smooth solution to (1.1). It is clear that $|u(t)|^2 dx$ is a bounded sequence of Radon measures, so that when $t_k \to T^-$ there exists a subsequence $|u(t_k)|^2 dx$ which converges weak-* to some Radon measure \mathcal{E} which is called the energy measure at time T.

We introduce the following two quantities, the lower local dimension $d(x, \mathcal{E})$ of \mathcal{E} at $x \in \Omega$, and the concentration dimension D of \mathcal{E} in Ω , defined respectively by

(1.2)
$$d(x,\mathcal{E}) = \liminf_{r \to 0^+} \frac{\ln \mathcal{E}(B_r(x))}{\ln r},$$

(1.3)
$$D = \inf\{\dim_{\mathcal{H}}(S) : S \subset \Omega \text{ compact, and } \mathcal{E}(S) > 0\}$$

with the convention that D = 3 if the collection over which the infimum is taken is empty. The local dimension is a standard geometric measure theoretic quantity, see [25], while the concentration dimension which assigns a numerical value to the concentration of the energy measure, namely the smallest Hausdorff dimension of a set of positive \mathcal{E} -measure, was first introduced in [32], together with the energy measure itself.

It is worth pointing out that although the definitions of $d(x, \mathcal{E})$ and D are local, the results in [18] require the solution u must be defined on whole space \mathbb{R}^3 . Hence, the methods which are dependent on a expanding iterative arguments developed in [18] can not deal with the energy measure on the domains with boundary. The methods developed in [24] have similar limitations. In order to consider energy measure and the energy measure and the local energy equality. Our methods are applicable to other systems for divergence free fields, in particular, to the three-dimensional MHD equations [34].

1.1 Definitions and Notations

Before presenting the main results, we give some notations and definitions. We first recall the definitions of weak and suitable weak solutions of (1.1).

Definition 1.1. Let T > 0. The function u is called a Leray-Hopf weak solution of (1.1) in $[0, T] \times \Omega$ if

 $1.u \in L^{\infty}(0,T; L^{2}(\Omega)) \cap L^{2}(0,T; H^{1}(\Omega);$

2. There exists a distribution P such that (u, P) satisfies (1.1) in the sense of distributions; 3.u satisfies the energy inequality: for a.e. $t_0 \in [0, t]$ including $t_0 = 0$,

(1.4)
$$\int_{\Omega} |u(t,x)|^2 dx + 2 \int_0^t \int_{\Omega} |\nabla u|^2 dx dt \le \int_{\Omega} |u(t_0,x)|^2 dx.$$

Furthermore, the pair (u, P) is called a suitable weak solution to (1.1) if $u \in L^3([0, T]; L^3(\Omega)), P \in L^{\frac{3}{2}}([0, T] \times \Omega)$ and the following local energy inequality holds

(1.5)
$$\int_{\Omega} |u(t,x)|^2 \phi dx + 2 \int_0^t \int_{\Omega} |\nabla u|^2 \phi dx dt \le \int_0^t \int_{\Omega} |u|^2 (\partial_t \phi + \Delta \phi) + (|u|^2 + 2P)u \cdot \nabla \phi dx dt$$

for every nonnegative $\phi \in C_0^{\infty}((0,T] \times \Omega)$.

We say a point $z_0 = (t_0, x_0) \in [0, T] \times \Omega$ is a regular point of solution u to (1.1) if there exists a non-empty neighborhood $\mathcal{O}_{z_0} \subset [0, T] \times \Omega$ of z_0 such that $u \in L^{\infty}(\mathcal{O}_{z_0})$. The complement of the set of regular points will be called the singular set.

Let (u, P) be a solution of (1.1). Introduce the scaling

(1.6)
$$u_{\lambda}(t,x) = \lambda u(\lambda^2 t, \lambda x); \quad P_{\lambda}(t,x) = \lambda^2 P(\lambda^2 t, \lambda x),$$

for arbitrary $\lambda > 0$. Then the family (u, P) is also a solution of (1.1). Setting

$$B_r(x_0) = \{ x \in \mathbb{R}^3 : |x - x_0| < r \}, B_r = B_r(0), \ B = B_1, Q_r(z_0) = B_r(x_0) \times (t_0 - r^2, t_0), \ Q_r = Q_r(0), \ Q = Q_1.$$

We introduce the following invariant quantities, which are invariant under the natural scaling (1.6):

$$\begin{aligned} A(u,r,z) &= \sup_{t-r^2 \le s \le t} \frac{1}{r} \int_{B_r(x) \times \{s\}} |u|^2 dx; \ B(u,r,z) = \frac{1}{r} \int \int_{Q_r(z)} |\nabla u|^2 dx dt, \\ C(u,r,z) &= \frac{1}{r^2} \int \int_{Q_r(z)} |v|^3 dx dt; \ D(P,r,z) = \frac{1}{r^2} \int \int_{Q_r(z)} |P|^{\frac{3}{2}} dx dt. \end{aligned}$$

For simplicity, we introduce the notations

$$A(u,r) = A(u,r,0); \ B(u,r) = B(u,r,0); \ C(u,r) = C(u,r,0); \ D(P,r) = D(P,r,0)$$

Let $\Omega \subset \mathbb{R}^n$ and $\Omega_T = [0,T] \times \Omega$. we use $L^q L^p(\Omega_T)$ to denote the space of measurable functions with the following norm

$$||f||_{L^{q}L^{p}(\Omega_{T})} = \begin{cases} (\int_{0}^{T} (\int_{\Omega} |f(t,x)|^{p} dx)^{\frac{q}{p}} dt)^{\frac{1}{q}}, \ 1 \leq q < \infty, \\ ess \sup_{t \in [0,T]} ||f(t,\cdot)||_{L^{p}(\Omega)}, \ q = \infty. \end{cases}$$

The Lorentz space $L^{r,s}(\Omega)$ is the space of measurable functions with the following norm:

$$||f||_{L^{r,s}(\Omega)} = \begin{cases} (\int_0^\infty \sigma^{s-1} |\{x \in \Omega : |f(x)| > \sigma\}|^{\frac{s}{r}} d\sigma)^{\frac{1}{s}}, \ 1 \le s < \infty, \\ \sup_{\sigma > 0} \sigma |\{x \in \Omega : |f(x)| > \sigma\}|^{\frac{1}{r}}, \ s = \infty. \end{cases}$$

We say that a local integrable function f is in $BMO(\mathbb{R}^3)$ if it satisfies

$$\sup_{R>0, x_0 \in \mathbb{R}^3} \frac{1}{|B_R(x_0)|} \int_{B_R(x_0)} |f(x) - [f]_{B_R(x_0)} |dx < \infty.$$

A local integrable function f is in $BMO_{loc}(\mathbb{R}^3)$ if $f \in BMO(\Omega')$ for any $\Omega' \subset \subset \mathbb{R}^3$. Moreover, f(t, x) is in $L^{2,\infty}([0,T]; BMO)$ if $||f(t, \cdot)||_{BMO} \in L^{2,\infty}([0,T])$.

Throughout this paper, C is a positive constant and C(A1, A2, ...) is a positive constant depending on A1, A2, ...

1.2 Main results

Before the statements of our main results, we show some examples to explain that some potential type II singularity are admissible in the space $L^{2,\infty}([0,T]; BMO(\mathbb{R}^3))$.

Proposition 1.1. Let $u(x,t) = \mathbb{P}\left\{\frac{1}{\sqrt{T-t}} | \frac{x}{\sqrt{T-t}} | \sqrt{T-t} (-\ln | \frac{x}{\sqrt{T-t}} |)^s \phi(\frac{x}{\sqrt{T-t}}) \right\}, \ 0 < s < \frac{2}{3}.$ Here \mathbb{P} is the Leray projection and ϕ is a radial cut-off function which equals 1 in $B_{\frac{1}{2}}$ and vanishes outside of $B_{\frac{3}{4}}$. Then, it holds

$$||u(t)||_{L^{2}}^{2} \sim \sqrt{T-t}, \ ||\nabla u(t)||_{L^{2}}^{2} \sim \frac{1}{\sqrt{T-t}},$$
$$||u(t)||_{L^{\infty}} \geq \frac{1}{(T-t)^{1+\frac{s}{2}}}, \ ||u(t)||_{BMO} \leq \frac{1}{\sqrt{T-t}}.$$

In particular, it is holding that

$$\lim_{t \to T} \sqrt{T-t} ||u(t)||_{(BMO(\mathbb{R}^3))} < \infty \text{ and } \lim_{t \to T} \sqrt{T-t} ||u(t)||_{L^{\infty}} = \infty.$$

Noticing that $||u(t)||_{BMO} \leq \frac{C}{\sqrt{T-t}}$ implies $u \in L^{2,\infty}(BMO)$, we choose naturally the space $L^{2,\infty}(BMO)$ as the workspace. We now state our main results.

Theorem 1.1. Let (u, P) be a suitable weak solution to (1.1) on $[0, T] \times \Omega$ and $||u||_{L^{2,\infty}([0,T];BMO(\Omega))} = M < \infty$. Then it is holding that

(1.7)

$$\sup_{t-r^{2} \leq s < t} \frac{1}{r} \int_{B_{r}(x)} |u|^{2} dx + \frac{1}{r} \int \int_{Q_{r}(z)} |\nabla u|^{2} dx dt \\
+ \frac{1}{r^{2}} \int \int_{Q_{r}(z)} |u|^{3} dx dt + \frac{1}{r^{2}} \int \int_{Q_{r}(z)} |P|^{\frac{3}{2}} dx dt \\
\leq C(\rho, M),$$

where $Q_{2r}(z) \subset Q_{\rho}(z) \subset (0,T] \times \Omega$. In particular, fix $z \in (0,T] \times \Omega$, there exists a constant r_0 depending on $d_z = dist(z, \partial([0,T] \times \Omega))$ such that if $r < r_0$ then it is holding that

(1.8)

$$\sup_{t-r^{2} \leq s < t} \frac{1}{r} \int_{B_{r}(x)} |u|^{2} dx + \frac{1}{r} \int \int_{Q_{r}(z)} |\nabla u|^{2} dx dt \\
+ \frac{1}{r^{2}} \int \int_{Q_{r}(z)} |u|^{3} dx dt + \frac{1}{r^{2}} \int \int_{Q_{r}(z)} |P|^{\frac{3}{2}} dx dt \\
\leq C(M).$$

Moreover, we have

(1.9) $d(x,\mathcal{E}) \ge 1; \quad D \ge 1.$

As an application of Theorem 1.1, we first show that

Theorem 1.2. Assume $\Omega \subseteq \mathbb{R}^3$. Let (u, P) be a solution to the Navier-Stokes equations on $[0, T) \times \Omega$ which is regular on the time interval [0, T). If u experiences singularity in time at t = T and $||u||_{L^{2,\infty}([0,T];BMO(\Omega))} = M < \infty$ (or $\limsup_{t \to T^-} \sqrt{T-t} ||u(t)||_{BMO(\Omega)} < \infty$), then it is holding

$$\int_{\Omega} |u(t)|^2 \phi dx + 2 \int_0^t \int_{\Omega} |\nabla u|^2 \phi dx dt$$
$$= \int_0^t \int_{\Omega} |u|^2 (\partial_t \phi + \Delta \phi) dx dt + \int_0^t \int_{\Omega} (|u^2| + 2P) u \cdot \nabla \phi dx dt \text{ for all } t \in [0, T]$$

where $\phi \in C_0^{\infty}((0,T] \times \Omega)$.

The second application of Theorem 1.1 is

Theorem 1.3. Assume (u, P) be a smooth solution of the Navier-Stokes equations (1.1) on $[0, T) \times \mathbb{R}^3$ and $u \in L^{2,\infty}([0,T); BMO(\mathbb{R}^3))$ (or $\limsup_{t\to T^-} \sqrt{T-t} ||u(t)||_{BMO(\mathbb{R}^3)} < \infty$) then the energy equality must hold on interval [0,T]

(1.10)
$$\int_{\mathbb{R}^3} |u(t,\cdot)|^2 dx + 2 \int_0^t \int_{\mathbb{R}^3} |\nabla u|^2 dx dt = \int_{\mathbb{R}^3} |u_0|^2 dx \text{ for } t \in [0,T].$$

Thirdly, we show a necessary condition for a solution u of Navier-Stokes equations (1.1) developing a singularity at time T

Theorem 1.4. Let (u, P) be a solution to the Navier-Stokes equations (1.1) on $[0, T) \times \mathbb{R}^3$ which is regular on the time interval [0, T). If u satisfies $||u(t)||_{L^{2,\infty}([0,T];BMO(\mathbb{R}^3))} \leq c$ for suitable small constant c then u can be continued to the regular solution beyond t = T. Especially, if u satisfies

$$\limsup_{t \to T^-} \sqrt{T - t} ||u(t)||_{BMO(\mathbb{R}^3)} \le c,$$

then u can be continued to the regular solution beyond t = T.

Remark 1.1. Theorem 1.4 means that as the solution u of (1.1) approaches a finite blow-up time T, the norm $||u(t)||_{BMO(\mathbb{R}^3)}$ must blow up at a rate $\frac{c}{\sqrt{T-t}}$.

Furthermore, we can allow the vertical part of the velocity u to be large.

Theorem 1.5. Let (u, P) be a suitable weak solution to (1.1) on $[-1, 0] \times \mathbb{R}^3$. Assume that u satisfies

(1.11)
$$||u_3||_{L^{2,\infty}([-1,0];BMO(\mathbb{R}^3))} = M < \infty$$

Then there exists a suitable small constant ε_1 depending on M such that if $||u_h||_{L^{2,\infty}([-1,0];BMO(\mathbb{R}^3))} \leq \varepsilon_1$ then u is regular in $[-1,0] \times \mathbb{R}^3$.

By using Theorem 1.3, we obtain some corollaries.

Corollary 1.1. Assume (u, P) be a Leray-Hopf weak solution to the Navier-Stokes equations (1.1) on $[0,T] \times \mathbb{R}^3$. If u experiences a finite number of singularities at $\{t_i\}, 1 \leq i \leq N$ in [0,T] and u satisfies $\limsup_{t \to t_i} \sqrt{t_i - t} ||u(t)||_{BMO(\mathbb{R}^3)} < \infty$, then u must satisfy energy equality on [0,T].

Proof. Without loss of generality, we assume $t_0 \in (0, T)$ and u experiences singularity at t_0 and T. It is clear that u is regular in $[0, t_0)$ and (t_0, T) . By Theorem 1.3, we have that the energy equality must hold on $[0, t_0]$ and $[t_0 + \epsilon, T]$ for $\epsilon \in (0, \frac{1}{2}(T - t_0))$

(1.12)
$$\int_{\mathbb{R}^3} |u(s)|^2 dx + \int_0^s \int_{\mathbb{R}^3} |\nabla u|^2 dx dt = \int_{\mathbb{R}^3} |u_0|^2 dx \text{ for } s \in [0, t_0],$$

(1.13)
$$\int_{\mathbb{R}^3} |u(\tau)|^2 dx + \int_{t_0+\epsilon}^{\tau} \int_{\mathbb{R}^3} |\nabla u|^2 dx dt = \int_{\mathbb{R}^3} |u(t_0+\epsilon)|^2 dx \text{ for } \tau \in [t_0+\epsilon,T].$$

To show the energy equality on [0, T], it is sufficient to prove

(1.14)
$$\lim_{\epsilon \to 0^+} ||u(t_0 + \epsilon)||^2_{L^2(\mathbb{R}^3)} = ||u(t_0)||^2_{L^2(\mathbb{R}^3)}$$

On the one hand, by the weak lower semicontinuity of the L^2 norm of u, we know that

(1.15)
$$||u(t_0)||^2_{L^2(\mathbb{R}^3)} \le \liminf_{\epsilon \to 0^+} ||u(t_0 + \epsilon)||^2_{L^2(\mathbb{R}^3)}.$$

On the other hand, it is well-known that the Leray-Hopf weak solutions on \mathbb{R}^3 satisfy the following strong energy inequality

(1.16)
$$||u(t)||_{L^{2}(\mathbb{R}^{3})}^{2} + \int_{s}^{t} \int_{\mathbb{R}^{3}} |\nabla u(\tau, x)|^{2} dx d\tau \leq ||u(s)||_{L^{2}(\mathbb{R}^{3})}^{2} \text{ for a.e. } s \in [0, t].$$

Noticing that (1.12) implies $\lim_{s \to t_0^-} ||u(s)||_{L^2(\mathbb{R}^3)}^2 = ||u(t_0)||_{L^2(\mathbb{R}^3)}^2$. This means by (1.16)

(1.17)
$$||u(t)||_{L^{2}(\mathbb{R}^{3})}^{2} + \int_{t_{0}}^{t} \int_{\mathbb{R}^{3}} |\nabla u(\tau, x)|^{2} dx d\tau \leq ||u(t_{0})||_{L^{2}(\mathbb{R}^{3})}^{2} \text{ for } t \geq t_{0}.$$

This implies that

(1.18)
$$||u(t_0)||^2_{L^2(\mathbb{R}^3)} \ge \limsup_{\epsilon \to 0^+} ||u(t_0 + \epsilon)||^2_{L^2(\mathbb{R}^3)}$$

According to (1.18) and (1.15), we conclude that (1.14) is holding.

If u is a suitable weak solution, by using Theorem 1.1 and a similar argument as [18], we can obtain a stronger conclusion.

Corollary 1.2. Assume (u, P) be a suitable weak solution to the Navier-Stokes equations (1.1) on $[0,T] \times \mathbb{R}^3$ and $u \in L^{2,\infty}([0,T], BMO(\mathbb{R}^3))$, then u must satisfy the energy equality on [0,T].

Proof. From Theorem 1.1, we obtain that $u \in L^{\infty}((0,T]; \mathcal{M}^{2,1})$, where $\mathcal{M}^{2,1}$ is the Morrey space with integrability 2 and rate 1. Since the Morrey space $\mathcal{M}^{2,1}$ is invariant under shifts $f \to f(\cdot - x_0)$ and the rescaling $f(x) \to \lambda f(\lambda x)$, we have by Cannone's Theorem [4] that $u \in L^{\infty}B_{\infty,\infty}^{-1}$. Consequently, interpolation with the enstrophy space $L^2H^1 = L^2B_{2,2}^1$ puts the solution u into the Onsager-critical class $L^3((0,T]B_{3,3}^1)$, from which we conclude that

$$\int_{\mathbb{R}^3} |u(t,\cdot)|^2 dx + 2 \int_0^t \int_{\mathbb{R}^3} |\nabla u|^2 dx dt = \int_{\mathbb{R}^3} |u(t_0,\cdot)|^2 dx \text{ for a.e. } t_0 \in (0,t].$$

By using the fact $\lim_{t_0\to 0} ||u(t_0)||_{L^2} = ||u_0||_{L^2}$, we deduce the energy conservation

$$\int_{\mathbb{R}^3} |u(t,\cdot)|^2 dx + 2 \int_0^t \int_{\mathbb{R}^3} |\nabla u|^2 dx dt = \int_{\mathbb{R}^3} |u_0|^2 dx \text{ for } t \in [0,T].$$

Remark 1.2. Proposition 1.1 and Theorem 1.3 imply that though the solution u to (1.1) experiences some potential type II singularity at first blow up time T, the energy equality may still holds on [0, T]. Hence, Theorem 1.1-1.3 and Corollary 1.1-1.2 improve considerably the recent results in [18, 14]. Theorem 1.4-1.5 show a precise blow up rate for the BMO norm of the solution u to (1.1) and improve considerably the results in [36, 10, 15].

Remark 1.3. In recent work of Cheskidov-Luo [24] showed that if $u \in L^{2,\infty}([0,T]; B^0_{\infty,\infty}(\mathbb{R}^3))$, then the weak solution u of (1.1) equations satisfies energy equality. Where, we illustrate that $BMO \not\subset B^0_{\infty,\infty}$. For example, let $h(x) = \ln |x|$. Obviously, h is in $BMO(\mathbb{R}^3)$. We shall show that h is not in $B^0_{\infty,\infty}(\mathbb{R}^3)$. Indeed,

$$\|\ln |x|\|_{B^0_{\infty,\infty}} = \sup_{j \ge -1} \|\Delta_j \ln |x|\|_{L^\infty},$$

where

$$\Delta_j \ln |x| = \mathcal{F}^{-1} \left(\varphi(2^{-j}\xi) \widehat{\ln} |\xi| \right) = 2^{jd} \int_{\mathbb{R}^d} \check{\varphi} \left(2^j (x-y) \right) \ln |y| dy.$$

Let $2^j y = z$, one has

$$\begin{aligned} \Delta_j \ln |x| &= \int_{\mathbb{R}^d} \check{\varphi} \left(2^j x - z \right) \ln \left(2^{-j} z \right) dz \\ &= \int_{\mathbb{R}^d} \check{\varphi} \left(2^j x - z \right) \left(\ln 2^{-j} + \ln z \right) dz \\ &= \int_{\mathbb{R}^d} \check{\varphi} \left(2^j x - z \right) \ln 2^{-j} dz + \int_{\mathbb{R}^d} \check{\varphi} \left(2^j x - z \right) \ln z dz. \end{aligned}$$

Thus we have

$$\Delta_{j}\ln(0) = \int_{\mathbb{R}^{d}} \check{\varphi}(-z) \ln 2^{-j} dz + \int_{\mathbb{R}^{d}} \check{\varphi}(-z) \ln z dz$$
$$= -j \ln 2 \int_{\mathbb{R}^{d}} \check{\varphi}(-z) dz + \int_{\mathbb{R}^{d}} \check{\varphi}(-z) \ln z dz.$$

Due to $\check{\varphi} \in \mathcal{S}(\mathbb{R}^d)$, one has that

$$\int_{\mathbb{R}^d} \check{\varphi}\left(-z\right) \ln z dz < \infty.$$

We now choose $\int_{\mathbb{R}^d} \check{\varphi}(-z) \, dz \neq 0$, then

$$\lim_{j \to \infty} |j \ln 2 \int_{\mathbb{R}^d} \check{\varphi}(-z) \, dz| = \infty.$$

Clearly,

$$\left\|\Delta_j \ln |x|\right\|_{L^{\infty}} \sim j$$

and thus

$$\|\ln |x|\|_{B^0_{\infty,\infty}} = \sup_{j \ge -1} \|\Delta_j \ln |x|\|_{L^\infty} = \infty.$$

It turns out that our result on energy conservation is of independent interest compare to the result of Cheskidov and Luo.

Our paper is organized as follows: in Section 2, we establish some uniform bounds on invariant quantities A(r), B(r), C(r) and D(r); in Section 3, we prove Theorem 1.1-1.5 and Proposition 1.1.

2 The uniform bounds on invariant quantities

Our goals in this section are to show some uniform bounds on invariant quantities under the assumption $u \in L^{2,\infty}([0,T]; BMO(\Omega))$.

Lemma 2.1. Let (u, P) be a suitable weak solution of (1.1) in $[0, T] \times \Omega$ satisfying

$$||u||_{L^{2,\infty}([0,T];BMO(\Omega))} = M < \infty$$

Then it is holding

$$C(u,r,z) \leq C \frac{r}{\rho} C(u,\rho,z) + C(\frac{\rho}{r})^{\frac{3}{2}} M^{\frac{3}{2}} A^{\frac{3}{4}}(u,\rho,z)$$

for $0 < r < \rho$ and $Q_{\rho}(z) \subset [0,T] \times \Omega$.

Proof. At almost every time we estimate

(2.1)
$$\int_{B_r(x)} |u|^3 dx \le \int_{B_r(x)} |u - u_\rho|^3 dx + C|B_r||u_\rho(x)|^3 = I_1 + I_2$$

where $u_{\rho} = \frac{1}{|B_{\rho}|} \int_{B_{\rho}(x)} u dy$. For I_1 , we have

(2.2)
$$I_{1} = \int_{B_{r}(x)} |u - u_{\rho}|^{\frac{3}{2}} |u - u_{\rho}|^{\frac{3}{2}} dx$$
$$\leq (\int_{B_{\rho}(x)} |u - u_{\rho}|^{2})^{\frac{3}{4}} (\frac{1}{|B_{\rho}|} \int_{B_{\rho}(x)} |u - u_{\rho}|^{6})^{\frac{1}{4}} |B_{\rho}|^{\frac{1}{4}}$$
$$\leq \rho^{\frac{3}{2}} A(\rho, z)^{\frac{3}{4}} ||u(t)||^{\frac{3}{2}}_{BMO(\mathbb{R}^{3})}.$$

For I_2 , we have

(2.3)
$$I_{2} = |B_{r}(x)| \frac{1}{|B_{\rho}(x)|} \int_{B_{\rho}(x)} u dy|^{3}$$
$$\leq C(\frac{r}{\rho})^{3} \int_{B_{\rho}(x)} |u|^{3} dx.$$

Summing up the estimates for I_1 and I_2 and integrating with respect to time from $t - r^2$ to t, we obtain

$$(2.4) \quad C(u,r,z) = \frac{1}{r^2} \int \int_{Q_r(z)} |u|^3 dx dt \le C \frac{r}{\rho} C(u,\rho,z) + (\frac{\rho}{r})^{\frac{3}{2}} r^{\frac{-1}{2}} \int_{t-r^2}^t ||u(s)||_{BMO}^{\frac{3}{2}} ds A^{\frac{3}{4}}(u,\rho,z).$$

Using the assumption $||u||_{L^{2,\infty}([0,T];BMO(\mathbb{R}^3))} = M$, we deduce

$$(2.5) r^{-\frac{1}{2}} \int_{t-r^{2}}^{t} ||u||_{BMO(\mathbb{R}^{3})}^{\frac{3}{2}} ds \\ = \frac{3}{2} r^{-\frac{1}{2}} \int_{0}^{\infty} \sigma^{\frac{1}{2}} |\{s \in [t-r^{2},t] : ||u(s,\cdot)||_{BMO(\mathbb{R}^{3})} > \sigma\}| d\sigma \\ = \frac{3}{2} r^{-\frac{1}{2}} \{\int_{0}^{R} \sigma^{\frac{1}{2}} |\{s \in [t-r^{2},t] : ||u(s,\cdot)||_{BMO(\mathbb{R}^{3})} > \sigma\}| d\sigma \\ + \int_{R}^{\infty} \sigma^{\frac{1}{2}} |\{s \in [t-r^{2},t] : ||u(s,\cdot)||_{BMO(\mathbb{R}^{3})} > \sigma\}| d\sigma \} \\ \leq r^{-\frac{1}{2}} (R^{\frac{3}{2}}r^{2} + 3R^{-\frac{1}{2}}M^{2}) \\ = 2(3^{\frac{3}{4}})M^{\frac{3}{2}} \end{aligned}$$

where we choose $R = \sqrt{3}r^{-1}M$. Substituting (2.5) into (2.4), we complete the proof of this lemma.

By using Lemma 2.1, we deduce some uniform bounds on invariant quantities:

Lemma 2.2. Let (u, P) be a suitable weak solution of (1.1) in $[0, T] \times \Omega$ satisfying

$$||u||_{L^{2,\infty}([0,T];BMO(\Omega))} = M < \infty.$$

Then it is holding

(2.6)

$$A(u,r,z) + B(u,r,z) + C^{\frac{7}{6}}(u,r,z) + D^{\frac{8}{7}}(P,r,z) \le C(d_z,M) \text{ for } Q_{2r}(z) \subset Q_{d_z}(z) \subset (0,T] \times \Omega$$

where $d_z = dist(z, \partial((0, T] \times \Omega)))$. In particular, fix $z \in (0, T] \times \Omega$, there exists a constant $r_0 > 0$ depending on d_z , such that if $r < r_0$ and $Q_r(z) \subset (0, T] \times \mathbb{R}^3$ then it is holding

(2.7)
$$A(u,r,z) + B(u,r,z) + C^{\frac{7}{6}}(u,r,z) + D^{\frac{8}{7}}(P,r,z) \le C(M).$$

Proof. Without loss of generality, we set z = 0 and $\rho \leq d_z$. Let $\phi(t, x) = \chi(t, x)\psi(t, x)$ where χ is cutoff function which equals 1 in $Q_{\frac{1}{2}\rho}$ and vanishes outside of $Q_{\frac{3}{4}\rho}$. Then let $\psi = (4\pi(r^2 - t))^{-\frac{3}{2}}e^{-\frac{|x|^2}{4(r^2 - t)}}$. Direct computations show that $\phi \geq 0$ and

$$\begin{aligned} \partial_t \phi + \triangle \phi = 0 \ in \ Q_{\frac{1}{2}\rho}, \\ |\partial_t \phi + \triangle \phi| \leq C\rho^{-5} \ in \ Q_{\rho}, \\ C^{-1}r^{-3} \leq \phi \leq Cr^{-3}; \ |\nabla \phi| \leq Cr^{-4} \ in \ Q_r, \\ \phi \leq C\rho^{-3}; \ |\nabla \phi| \leq C\rho^{-4} \ in \ Q_{\rho} - Q_{\frac{3}{2}\rho}. \end{aligned}$$

Using ϕ as a test function in the local energy inequality (1.5), we obtain

(2.8)
$$A(u,r) + B(u,r) \leq C(\frac{r}{\rho})^2 A(u,\rho) + C(\frac{\rho}{r})^2 C(u,\rho) + C(\frac{\rho}{r})^2 C^{\frac{1}{3}}(u,\rho) D^{\frac{2}{3}}(P,\rho)$$
$$\leq C(\frac{r}{\rho})^2 A(u,\rho) + C(\frac{\rho}{r})^2 C(u,\rho) + C(\frac{\rho}{r})^2 D(P,\rho).$$

For $C(u, \rho)$, we have by using Lemma 2.1

(2.9)
$$C(u,r) \le C\frac{r}{\rho}C(u,\rho) + C(\frac{\rho}{r})^{\frac{3}{2}}M^{\frac{3}{2}}A^{\frac{3}{4}}(u,\rho).$$

It remains to show some bounds on D(u,r). Let $\eta(x)$ be a cut-off function which equals 1 in $B_{\frac{3\rho}{4}}$ and vanishes outside of B_{ρ} . Let P_1 satisfy $-\Delta P_1 = \partial_{x_i}\partial_{x_j}(u_iu_j\eta)$ and $P_2 = P - P_1$. Then, it is clear that $\Delta P_2 = 0$ in $B_{\frac{3\rho}{4}}$. by using the Calderón-Zygmund inequality, we have

$$\int_{B_{\rho}} |P_1|^{\frac{3}{2}} dx \le C(\int_{B_{\rho}} |u|^3 dx).$$

By the properties of harmonic functions, we infer that for $r \leq \frac{\rho}{2}$,

$$\int_{B_r} |P_2|^{\frac{3}{2}} dx \le Cr^3 \sup_{x \in B_r} |P_2(x)|^{\frac{3}{2}} \le C(\frac{r}{\rho})^3 \int_{B_\rho} |P_2|^{\frac{3}{2}} dx.$$

It then follows that for $0 < r \leq \frac{\rho}{2}$

$$\begin{split} &\int_{B_r} |P|^{\frac{3}{2}} dx \\ \leq & C(\int_{B_{\rho}} |u|^3 dx) + C(\frac{r}{\rho})^3 \int_{B_{\rho}} |P - P_1|^{\frac{3}{2}} dx \\ \leq & C(\int_{B_{\rho}} |u|^3 dx) + C(\frac{r}{\rho})^3 \int_{B_{\rho}} |P|^{\frac{3}{2}} dx. \end{split}$$

Integrating with respect to t from $-r^2$ to 0, we obtain, using Hölder inequality,

$$\int_{Q_r} |P|^{\frac{3}{2}} dx dt \le C \int_{Q_{\rho}} |u^3| dx dt + C(\frac{r}{\rho})^3 \int_{Q_{\rho}} |P|^{\frac{3}{2}} dx dt.$$

This implies

(2.10)
$$D(P,r) \le C\frac{r}{\rho}D(P,\rho) + C(\frac{\rho}{r})^2C(u,\rho).$$

According to (2.10), we have by using Young's inequality

(2.11)
$$D(P,r)^{\frac{8}{7}} \leq C(\frac{r}{\rho})^{\frac{8}{7}} D^{\frac{8}{7}}(P,\rho) + C(\frac{\rho}{r})^{\frac{16}{7}} C(u,\rho)^{\frac{8}{7}} \leq C(\frac{r}{\rho})^{\frac{8}{7}} D^{\frac{8}{7}}(P,\rho) + (\frac{r}{\rho})^{\frac{7}{6}} C(u,\rho)^{\frac{7}{6}} + C(\frac{\rho}{r})^{168}$$

Similar computations show by using (2.9)

(2.12)
$$C^{\frac{7}{6}}(u,r) \le C(\frac{r}{\rho})^{\frac{7}{6}}C^{\frac{7}{6}}(u,\rho) + (\frac{r}{\rho})^{2}A(u,\rho) + C(\frac{\rho}{r})^{21}M^{14}$$

By using Young's inequality, we deduce from (2.8)

$$(2.13) A(u,r) + B(u,r) \le C(\frac{r}{\rho})^2 A(u,\rho) + (\frac{r}{\rho})^{\frac{7}{6}} C^{\frac{7}{6}}(u,\rho) + (\frac{r}{\rho})^{\frac{8}{7}} D^{\frac{8}{7}}(P,\rho) + C((\frac{\rho}{r})^{21} + (\frac{\rho}{r})^{24}).$$

Summing up the estimates (2.11), (2.12) and (2.13), we obtain by defining G(r) = A(u,r) + C(u,r) $B(u,r) + C^{\frac{7}{6}}(u,r) + D^{\frac{8}{7}}(P,r)$

(2.14)
$$G(r) \le C(\frac{r}{\rho})^{\frac{8}{7}}G(\rho) + (C+M^{14})(\frac{\rho}{r})^{168}$$

where we have used the fact $\frac{r}{\rho} < 1$. Fix $\theta = \min\{\frac{1}{2}, \frac{1}{C^7}\}$ and set $r = \theta^k \rho$ for $k \in \mathbb{N}$, (2.14) yields

(2.15)
$$G(\theta^k \rho) \le \theta G(\theta^{k-1} \rho) + C(1 + M^{14}) \theta^{-168}.$$

By a standard iterative argument, we deduce that

(2.16)
$$G(r) \le C \frac{r}{\rho} G(\rho) + C(1 + M^{14}) \text{ for } r \le \frac{\rho}{2}.$$

This means (2.6).

In particular, by choosing r_0 satisfying $C \frac{r_0}{d_z} G(d_z) \leq M$, we obtain (2.7).

We conclude this section by giving a uniform bound on C(u, r, z) + D(P, r, z) when $||u||_{L^{2,\infty}(BMO(\mathbb{R}^3))}$ is suitable small.

Lemma 2.3. Let (u, P) be a suitable weak solution of (1.1) in $[0, T] \times \mathbb{R}^3$. Fix $\varepsilon > 0$, there exists constant c and r^* such that if

$$||u||_{L^{2,\infty}([0,T];BMO(\mathbb{R}^3))} \le c,$$

then it is holding

(2.17)
$$C(u,r,z) + D(P,r,z) \le \varepsilon, \text{ for } 0 < r \le r^*,$$

where r^* is depending on d_z .

Proof. Without loss of generality, we assume $c \leq 1$. Firstly, from Lemma 2.2, we have that there exists $r_0 > 0$ depending on d_z such that if $\rho \leq r_0$, then it is holding

(2.18)
$$C(u, \rho, z) + D(P, \rho, z) + A(u, \rho, z) + B(u, \rho, z) \le C.$$

Then we deduce by choosing $\rho \leq r_0$ in (2.9) and using (2.18)

$$C(u, r, z) \leq \frac{r}{\rho} C(u, \rho, z) + C(\frac{\rho}{r})^{\frac{3}{2}} c^{\frac{3}{2}} A^{\frac{3}{4}}(u, \rho, z) \leq \frac{r}{\rho} C(u, \rho, z) + C c^{\frac{3}{2}} (\frac{\rho}{r})^{\frac{3}{2}}$$

for $r \leq \rho$. We now take $r = 2^{-k} r_0$ and obtain by a standard iterative argument that

(2.19)

$$C(u, 2^{-k}r_0, z) \leq \frac{1}{2}C(u, 2^{-k+1}r_0, z) + C(2c)^{\frac{3}{2}} \leq 2^{-k}C(u, r_0, z) + \sum_{j=0}^{k} 2^{-j}C(2c)^{\frac{3}{2}} \leq 2^{-k}C(u, r_0, z) + C(2c)^{\frac{3}{2}}.$$

Similarly, by using (2.10) and (2.19), we get

$$(2.20) \qquad D(u, 2^{-k_0 k} r_0, z) \\ \leq \frac{1}{2} D(u, 2^{(-k_0(k-1))} r_0, z) + C 2^{2k_0} (2^{-k_0(k-1)} C(u, r_0, z) + C (2c)^{\frac{3}{2}}) \\ \leq 2^{-k} D(u, 2^{-k_0} r_0, z) + C 2^{2k_0} (2^{-k_0(k-1)} C(u, r_0, z) + C (2c)^{\frac{3}{2}})$$

where we choose k_0 satisfying $C2^{-k_0} \leq \frac{1}{2}$.

Summing up (2.19) and (2.20), we obtain by using (2.18) that

$$C(u, 2^{-kk_0}r_0, z) + D(P, 2^{-kk_0}r_0, z)$$

$$\leq (2^{-k} + 2^{-kk_0})(C(u, r_0, z) + D(P, 2^{-k_0}r_0, z)) + C2^{2k_0}(2^{-kk_0+k_0}C(u, r_0, z) + (2c)^{\frac{3}{2}})$$

$$\leq C(2^{-k} + 2^{-kk_0}) + C2^{2k_0}(2^{-kk_0+k_0}C + (2c)^{\frac{3}{2}}).$$

We now choose K satisfying $C(2^{-K} + 2^{-Kk_0} + 2^{-Kk_0+3k_0}) \leq \frac{\varepsilon}{2}$ and $r^* = 2^{-Kk_0}r_0$, then we take $c = \frac{1}{2}(\frac{\varepsilon}{2^{2k_0+1}C})^{\frac{2}{3}}$ and conclude the Lemma 2.3.

3 The proof of main theorems

We first recall two well-known results about the suitable weak solutions to (1.1).

Proposition 3.1. [22] Let (u^k, P^k) be a sequence of weak solutions of (1.1) in Q_1 such that

$$||u^{k}||_{L^{\infty}([-1,0];L^{2}(B))} + ||P^{k}||_{L^{\frac{3}{2}}(Q)} + ||\nabla u^{k}||_{L^{2}(Q)} \le E$$

and (u^k, P^k) satisfies the local energy inequality (1.5). Suppose that (u, P) is the weak limit of (u^k, P^k) ; then (u, P) is a suitable weak solution of (1.1) on Q.

Proposition 3.2. [3, 22, 19] Let (u, P) be a suitable weak solution of (1.1) in $Q_r(z)$. There exists $\varepsilon_0 > 0$ such that if

$$\frac{1}{r^2} \int_{Q_r(z)} (|u|^3 + |P|^{\frac{3}{2}}) dx dt \le \varepsilon_0,$$

then u is regular in $Q_{\frac{r}{2}}(z)$.

We now begin to prove Theorem 1.1.

Proof. The estimates (1.8)-(1.9) are direct conclusions of lemma 2.2.

For any $z = (x, T) \in \Omega \times \{T\}$, set $d_x = dist(x, \partial \Omega)$. From (1.8) and the weak lower semicontinuity of the L^2 norm of u, we get

$$\mathcal{E}(B_r(x)) \leq \lim_{t \to T^-} \int_{B_r(x)} |u(t)|^2 dy \leq C(d_x, M) r \text{ for } r \leq \frac{d_x}{2}.$$

By the definitions of local dimension $d(x, \mathcal{E})$, it is clear that $d(x, \mathcal{E}) \ge 1$.

We now prove $D \geq 1$. For any compact $S \subset \Omega$ with $\dim_{\mathcal{H}}(S) = s^* < 1$, by the definition of Hausdorff dimension, we have $H^{s_1}(S) = 0$ for $s^* < s_1 < 1$. Then fixing $0 < \delta < \operatorname{dist}(S, \partial\Omega)$, there exists a collection of open balls $\{B_{r_i}(x_i)\}_{i=1}^{\infty}$ satisfying $r_i \leq \delta$ such that $S \subset \bigcup_{i=1}^{\infty} B_{r_i}(x_i) \subset \Omega$ and $\sum_{i=1}^{\infty} r_i^{s_1} \leq 1$. From this facts, we deduce

$$\mathcal{E}(S) \le \sum_{i=1}^{\infty} \mathcal{E}(B_{r_i}(x_i)) \le C(M) \sum_{i=1}^{\infty} r_i \le C(M) \delta^{1-s_1} \sum_{i=1}^{\infty} r_i^{s_1} \to 0 \text{ as } \delta \to 0.$$

This means $D \geq 1$.

The proof of Theorem 1.2

Proof. Assume $supp\phi(T, x) \equiv \overline{\Omega}_0 \subset \Omega$ and $d = dist(supp\phi, \partial(\Omega \times [0, T]))$. From the proof of Lemma 2.2, it is clear that

(3.1)
$$\sup_{T-r^2 \le t < T, x \in \bar{\Omega}_0} \frac{1}{r} \int_{B_r(x)} |u(t)|^2 dy \le C(M, d) \text{ for } 0 < r \le \frac{d}{2}.$$

By the weak lower semicontinuity of the L^2 norm of u, we see that to prove Theorem 1.2 it is enough to prove

(3.2)
$$\lim_{t \to T^{-}} \int_{\bar{\Omega}_{0}} |u(t)|^{2} dx = \int_{\bar{\Omega}_{0}} |u(T)|^{2} dx.$$

	н

Let Σ be the set of all singular points of u at time T. By the well-known Caffarelli-Kohn-Nirenberg Theorem [3], we have

 $H^1(\Sigma) = 0.$

This fact implies that for each $\epsilon > 0$, there exists a countable family of sets of the form

$$B_i^{\epsilon} = B_{r_i^{\epsilon}}(x_i^{\epsilon}) \times \{t = T\}$$

such that

(3.3)
$$r_i^{\epsilon} \le \frac{d}{2}, \ \Sigma \cap (\bar{\Omega}_0 \times \{t = T\}) \subset \cup B_i^{\epsilon}, \ \sum r_i^{\epsilon} < \epsilon.$$

Fix $\varepsilon > 0$ and let

$$\epsilon = \frac{\varepsilon}{8C(||u_0||_{L^2}, M, d)}.$$

Then, by (3.1) and (3.3), we have

(3.4)

$$\begin{aligned} |\int_{\cup B_{i}^{\epsilon}} |u(t)|^{2} dx - \int_{\cup B_{i}^{\epsilon}} |u(T)|^{2} dx| \\
\leq \sum_{i} \int_{B_{i}^{\epsilon}} |u(t)|^{2} dx + \sum_{i} \int_{B_{i}^{\epsilon}} |u(T)|^{2} dx \\
\leq 2C(||u_{0}||_{L^{2}}, M, d) \sum_{i} r_{i}^{\epsilon} \\
< 2C(||u_{0}||_{L^{2}}, M, d)\epsilon \\
\leq \frac{\varepsilon}{4}
\end{aligned}$$

for all $t \in [T - (\frac{d}{2})^2, T]$, where we have used the fact $\int_{\cup B_i^{\epsilon}} |u(T)|^2 dx \leq \liminf_{t \to T^-} \int_{\cup B_i^{\epsilon}} |u(t)|^2 dx$ due to the weak lower semicontinuity of the L^2 norm of u.

 Set

$$\omega^{\epsilon} \equiv (\bar{\Omega}_0 \times \{t = T\}) - \cup B_i^{\epsilon}.$$

For each $z \in \omega^{\epsilon}$, there exists a non-empty neighborhood \mathcal{O}_z such that u is Hölder continuous on $\mathcal{O}_z \cap ([0,T] \times \overline{\Omega}_0)$. Since ω^{ϵ} is compact, there exists a non-empty neighborhood $\mathcal{O}_{\omega}^{\epsilon}$ of the set ω^{ϵ} such that u is continuous in $\overline{\mathcal{O}}_{\omega}^{\epsilon} \cap ([0,T] \times \overline{\Omega}_0)$. Hence,

(3.5)
$$|\int_{\omega^{\epsilon}} |u(t)|^2 dx - \int_{\omega^{\epsilon}} |u(T)|^2 dx| < \frac{\varepsilon}{2}$$

for |T - t| small enough. Combining (3.4) and (3.5), we obtain

$$(3.6) \qquad \lim_{t \to T^{-}} |\int_{\Omega_{0}} |u(t)|^{2} dx - \int_{\Omega_{0}} |u(T)|^{2} dx| \\\leq \limsup_{t \to T^{-}} |\int_{\omega^{\epsilon}} |u(t)|^{2} dx - \int_{\omega^{\epsilon}} |u(T)|^{2} dx| + \limsup_{t \to T^{-}} |\int_{\cup B_{i}^{\epsilon}} |u(t)|^{2} dx - \int_{\cup B_{i}^{\epsilon}} |u(T)|^{2} dx| \\<\varepsilon.$$

This implies (3.2) by taking $\varepsilon \to 0$.

We now begin to prove Theorem 1.3

Proof. Since $u \in L^{2,\infty}([0,T), BMO(\mathbb{R}^3))$, we deduce that from Theorem 1.2

$$(3.7) \qquad \int_{\mathbb{R}^3} |u(t)|^2 \phi dx + 2 \int_0^t \int_{\mathbb{R}^3} |\nabla u|^2 \phi dx dt$$
$$= \int_{\mathbb{R}^3} |u_0|^2 \phi dx + \int_0^t \int_{\mathbb{R}^3} |u|^2 (\partial_t \phi + \Delta \phi) dx dt + \int_0^t \int_{\mathbb{R}^3} (|u^2| + 2P) u \cdot \nabla \phi dx dt \text{ for all } t \in [0, T]$$

where $\phi \in C_0^{\infty}([0,T] \times \mathbb{R}^3)$. Let $\phi(t,x) = \eta(x)$ be a cut-off function which equals 1 in B_R and vanishes outside of B_{2R} for R > 0. Then we obtain that

(3.8)
$$\int_{\mathbb{R}^3} |u(t)|^2 \eta dx + 2 \int_0^t \int_{\mathbb{R}^3} |\nabla u|^2 \eta dx dt$$
$$= \int_{\mathbb{R}^3} |u_0|^2 \eta dx + \int_0^t \int_{\mathbb{R}^3} |u|^2 \Delta \eta dx dt + \int_0^t \int_{\mathbb{R}^3} (|u^2| + 2P) u \cdot \nabla \eta dx dt \text{ for all } t \in [0, T]$$

Noticing that

(3.9)
$$|\int_0^t \int_{\mathbb{R}^3} |u|^2 \Delta \eta dx dt | \leq C|T| \sup_{t \in [0,T)} ||u(t)||_{L^2}^2 \frac{1}{R^2},$$

$$(3.10) \qquad |\int_0^t \int_{\mathbb{R}^3} (|u^2| + 2P)u \cdot \nabla \eta dx dt| \le C(||u||_{L^3([0,T];L^3(\mathbb{R}^3))}^3 + ||P||_{L^{\frac{3}{2}}([0,T];L^{\frac{3}{2}}(\mathbb{R}^3))}^{\frac{3}{2}}) \frac{1}{R}$$

Taking $R \to \infty$, we deduce that from (3.8)

(3.11)
$$\int_{\mathbb{R}^3} |u(t)|^2 dx + 2 \int_0^t \int_{\mathbb{R}^3} |\nabla u|^2 dx dt = \int_{\mathbb{R}^3} |u_0|^2 dx.$$

We now prove Theorem 1.4:

Proof. Fix $z \in (0,T] \times \mathbb{R}^3$. From Lemma 2.3, there exists c > 0 and $0 < r^* < d_z$ such that if $||u||_{L^{2,\infty}([0,T];BMO(\mathbb{R}^3))} \leq c$ then it holds

(3.12)
$$C(u,r,z) + D(P,r,z) < \varepsilon_0 \text{ for } 0 < r < r^*.$$

By Proposition 3.2, u is regular on $[0,T] \times \mathbb{R}^3$. This complete the proof of Theorem 1.4.

We now turn to the proof of Theorem 1.5, this proof is based on the uniform bounds on invariant quantities established in Lemma 2.2 and a standard blow-up argument.

Proof. If the conclusion of the theorem is false, then there exists a constant M and a sequence (u^k, P^k) whose elements are suitable weak solutions of (1.1) in $[-1, 0] \times \mathbb{R}^3$ and satisfy

$$||u_3^k||_{L^{2,\infty}([-1,0];BMO(\mathbb{R}^3))} \le M, \ ||u_h^k||_{L^{2,\infty}([-1,0];BMO(\mathbb{R}^3))} \le \frac{1}{k}.$$

Furthermore, we can assume u^k is singular at (0,0). According to Lemma 2.2, there exists a sequence of $\{r_k\}$ satisfying

$$C(u^k, r) + D(P^k, r) \le C(M) \text{ for } 0 < r \le r_k$$

Set $v^k = r_k u(r_k^2 t, r_k x)$ and $\Pi^k = r_k^2 P(r_k^2 t, r_k x)$, then it is clear that (v^k, Π^k) are suitable weak solutions to (1.1) on Q and satisfy

$$||v_3^k||_{L^{2,\infty}([-1,0];BMO(B))} \le M, ||v_h^k||_{L^{2,\infty}([-1,0];BMO(B))} \le \frac{1}{k};$$

$$C(v^k, 1) + D(\Pi^k, 1) \le C(M).$$

By the local energy inequality (1.5), we obtain

$$||\partial_t v^k||_{L^{\frac{3}{2}}}([-1,0]; H^{-2}(B_{\frac{3}{4}})) + ||v^k||_{L^{\infty}([-1,0]; L^2(B_{\frac{3}{4}}))} + ||\nabla v^k||_{L^2([-1,0]; L^2(B_{\frac{3}{4}}))} \le C(M).$$

According to the well-known Aubin-Lions lemma and Proposition 3.1, by passing to a subsequence if necessary, we may deduce that there exists a suitable weak solution (v, Π) to (1.1) such that

$$v^{k} \to v \text{ in } L^{3}(Q_{\frac{3}{4}}), v^{k} \rightharpoonup v \in L^{\infty}([-\frac{9}{16}, 0]; L^{2}(B_{\frac{3}{4}})),$$
$$\Pi^{k} \to \Pi \text{ in } L^{\frac{3}{2}}(Q_{\frac{3}{4}}), \nabla v^{k} \to \nabla v \text{ in } L^{2}(Q_{\frac{3}{4}})$$

as $k \to \infty$. In particular, we have by Fatou's lemma

$$||v_h||_{L^{2,\infty}([-1,0];BMO(B_{\frac{3}{4}}))} \le \liminf_{k \to \infty} ||v_h^k||_{L^{2,\infty}([-1,0];BMO(B_{\frac{3}{4}}))} = 0.$$

Therefore, we get $v_h = b(t)$ with $b(t) \in L^{\infty}[-\frac{9}{16}, 0]$. Then by $\nabla \cdot v = 0$, we obtain $\partial_3 v_3 = 0$. Thus, we deduce that (v, Π) satisfies

(3.13)
$$\begin{cases} \partial_t v - \Delta v + \nabla \Pi = -(b_1(t), b_2(t), 0) \cdot \nabla(0, 0, v_3) \text{ in } Q_{\frac{3}{4}}, \\ \nabla \cdot v = 0. \end{cases}$$

By the classical result concerning linear Stokes equations [20] and the fact $b(t) \in L^{\infty}[-\frac{9}{16}, 0]$, we have that $|v| \leq C(M)$ in $Q_{\frac{1}{2}}$. However, (0,0) is a singular point of v^k , hence, it is holding by using Proposition 3.2, (2.10) and the fact $v^k \to v$ in $L^3(Q_{\frac{3}{4}})$

(3.14)
$$\varepsilon_0 < \liminf_{k \to \infty} \frac{1}{r^2} \int_{Q_r} (|v^k|^3 + |\Pi^k|^{\frac{3}{2}}) dx dt$$
$$\leq \liminf_{k \to \infty} (C(v, r) + \frac{r}{\rho} D(\Pi^k, \rho) + \frac{\rho^2}{r^2} C(v^k, \rho))$$
$$= C(v, r) + \frac{\rho^2}{r^2} C(v, \rho) + \liminf_{k \to \infty} \frac{r}{\rho} D(\Pi^k, \rho)$$

for $0 < 2r < \rho < \frac{1}{2}$. From Lemma 2.2 and the fact $|v| \leq C(M)$ in $Q_{\frac{1}{2}}$, we get $D(\Pi^k, \rho) \leq C(M)$ and

$$\varepsilon_0 < C(M)(r^3 + \frac{\rho^5}{r^2} + \frac{r}{\rho}).$$

By taking $\rho = \sqrt{r}$, we infer

$$\varepsilon_0 < C(M)r^{\frac{1}{2}}.$$

which, for sufficiently small r, is a contradiction.

Finally, we prove Proposition 1.1.

Proof. Firstly, a direct computation shows that

(3.15)
$$||u(t)||_{L^2}^2 \sim \sqrt{T-t}, \quad ||\nabla u(t)||_{L^2}^2 \sim \frac{1}{\sqrt{T-t}}.$$

Secondly, we observe that $||u(\lambda x)||_{L^{\infty}} = ||u(x)||_{L^{\infty}}$ and $||u(\lambda x)||_{BMO} = ||u(x)||_{BMO}$. It is clear that

$$||u(t)||_{L^{\infty}} = \frac{1}{\sqrt{T-t}} ||\mathbb{P}(|x|^{\sqrt{T-t}}(-\ln|x|)^{s}\phi(x))||_{L^{\infty}},$$

$$||u(t)||_{BMO} = \frac{1}{\sqrt{T-t}} ||\mathbb{P}(|x|^{\sqrt{T-t}}(-\ln|x|)^{s}\phi(x))||_{BMO}.$$

Let $v = \frac{1}{\sqrt{T-t}} |x|^{\sqrt{T-t}} (-\ln |x|)^s \phi(x)$, it is clear that $||u||_{L^{\infty}} = ||\mathbb{P}v||_{L^{\infty}}$ and $||u||_{BMO} = ||\mathbb{P}v||_{BMO}$. We first show $||\sqrt{T-t}\nabla v||_{L^3(\mathbb{R}^3)}$ is bounded. Direct computations show that

$$\begin{split} &\int_{\mathbb{R}^3} |\sqrt{T - t} \nabla v|^3 dx \\ \leq & C(\int_{\mathbb{R}^3} (T - t)^{\frac{3}{2}} |x|^{3\sqrt{T - t} - 3} (-\ln|x|)^{3s} \phi^3(x) dx + \int_{\mathbb{R}^3} |x|^{3\sqrt{T - t} - 3} (-\ln|x|)^{3s - 3} \phi^3(x) dx \\ &+ \int_{\mathbb{R}^3} |x|^{3\sqrt{T - t}} (-\ln|x|)^{3s} |\nabla \phi|^3 dx) \\ \leq & C \frac{4\pi}{3} ((T - t)^{\frac{3}{2}} \int_0^{\frac{3}{4}} r^{3\sqrt{T - t} - 1} (-\ln r)^{3s} dr + \int_0^{\frac{3}{4}} r^{3\sqrt{T - t} - 1} (-\ln r)^{3s - 3} dr + \int_0^{\frac{3}{4}} r^{3\sqrt{T - t} + 2} (-\ln r)^{3s} dr) \\ \leq & C (I_1 + I_2 + I_3). \end{split}$$

For I_1 , it holds by setting $\eta = -3\sqrt{T-t}\ln r$ that

(3.16)

$$I_{1} = (T-t)^{\frac{3}{2}} \int_{0}^{\frac{3}{4}} r^{3\sqrt{T-t}-1} (-\ln r)^{3s} dr$$

$$= (T-t)^{\frac{3}{2}} \int_{3\sqrt{T-t}\ln\frac{4}{3}}^{\infty} e^{-\eta} (\frac{\eta}{3\sqrt{T-t}})^{3s} \frac{d\eta}{3\sqrt{T-t}}$$

$$\leq 3^{-3s-1} (T-t)^{\frac{3-3s-1}{2}} \int_{0}^{\infty} e^{-\eta} \eta^{3s} d\eta$$

$$= 3^{-3s-1} (T-t)^{\frac{3-3s-1}{2}} \Gamma(3s+1).$$

For I_2 , we have that

(3.17)
$$I_{2} = \int_{0}^{\frac{3}{4}} r^{3\sqrt{T-t}-1} (-\ln r)^{3s-3} dr$$
$$\leq \int_{0}^{\frac{3}{4}} r^{-1} (-\ln r)^{3s-3} dr$$
$$\leq \frac{1}{2-3s} (\ln 4 - \ln 3)^{3s-2}.$$

It is obviously that

(3.18)
$$I_3 = \int_0^{\frac{3}{4}} r^{3\sqrt{T-t}+2} (-\ln r)^{3s} dr \le C.$$

Combining with the estimates (3.16)-(3.18), we obtain

(3.19)
$$\int_{\mathbb{R}^3} |\sqrt{T-t}\nabla v|^3 dx \le C$$

for $0 < s < \frac{2}{3}$. Using the above estimate and the boundedness of \mathbb{P} on L^3 , we get

$$||\sqrt{T-t}\nabla u(t)||_{L^3} = ||\sqrt{T-t}\mathbb{P}\nabla v||_{L^3} \le C_1||\sqrt{T-t}\nabla v||_{$$

We thus obtain by using the fact $\dot{W}^{1,3}(\mathbb{R}^3) \hookrightarrow BMO(\mathbb{R}^3)$ that

(3.20)
$$||u(t)||_{BMO(\mathbb{R}^3)} \le \frac{C}{\sqrt{T-t}}.$$

We now consider the estimate of $||u(t)||_{L^{\infty}}$. By the definition of Leray projection, we see that

(3.21)
$$u_i = (\mathbb{P}v)_i = v_i - \mathbb{R}_i \mathbb{R}_j v_j$$

where \mathbb{R}_i is the Riesz transform. On the one hand, we deduce by choosing $|x| = e^{-\frac{s}{\sqrt{T-t}}}$

(3.22)
$$||v(t)||_{L^{\infty}} \sim \frac{1}{(T-t)^{\frac{1+s}{2}}}$$

On the other hand, by the definition of Riesz transform, we have

$$(3.23) \qquad \mathbb{R}_{i}\mathbb{R}_{j}v_{j}(0) = \frac{1}{\sqrt{T-t}} \int_{\mathbb{R}^{3}} \frac{y_{i}y_{j}}{|y|^{5}} |y|^{\sqrt{T-t}} (-\ln|y|)^{s} \phi_{j}(|y|) dy \\ = \frac{1}{\sqrt{T-t}} \int_{S^{2}} \omega_{i}\omega_{j}d\omega \int_{0}^{1} r^{\sqrt{T-t}-1} (-\ln r)^{s} \phi_{j}(r) dr \\ = \frac{1}{(T-t)^{\frac{1}{2}}} \int_{S^{2}} |\omega_{i}|^{2} d\omega \int_{0}^{1} r^{\sqrt{T-t}-1} (-\ln r)^{s} \phi_{j}(r) dr \\ \ge \frac{1}{(T-t)^{\frac{1}{2}}} \int_{S^{2}} |\omega_{i}|^{2} d\omega \int_{0}^{\frac{1}{2}} r^{\sqrt{T-t}-1} (-\ln r)^{s} dr \\ = \frac{1}{(T-t)^{1+\frac{s}{2}}} \frac{4\pi}{9} \int_{\sqrt{T-t}\ln 2}^{\infty} e^{-\tau} \tau^{s} d\tau \\ \ge \frac{2\pi}{9} \frac{1}{(T-t)^{1+\frac{s}{2}}} \Gamma(1+s).$$

Collecting (3.21)-(3.23) implies

(3.24)
$$||u(t)||_{L^{\infty}} \ge \frac{1}{(T-t)^{1+\frac{s}{2}}}.$$

Combining with (3.15), (3.20), (3.24) completes the proof of Proposition 1.1.

Acknowledgments Tan was supported by the Construct Program of the Key Discipline in Hunan Province and the National Natural Science Foundation of China (No. 11871209). Yin was partially supported by the National Natural Science Foundation of China (No. 11671407), the Macao Science and Technology Development Fund (No. 0091/2018/A3), and Guangdong Province of China Special Support Program (No. 8-2015), and the key project of the Natural Science Foundation of Guangdong province (No. 2016A030311004).

References

- [1] J. Leray, Sur le mouvement d'un liquide visqueux emplissant l'espace, Acta. Math. 63 (1934) 183-248.
- [2] T. Buckmaster and V. Vicol, Nonuniqueness of weak solutions to the Navier-Stokes equation, Ann.Math. 189 (2019) 101-144.
- [3] L. Caffarelli, R. Kohn, and L. Nirenberg, Partial regularity of suitable weak solutions of the Navier-Stokes equations, *Comm.Pure Appl.Math.* 35 (1982) 771-831.
- [4] M. Cannone, Ondelettes, paraproduits et Navier-Stokes. DiderotEditeur, Paris (1995) With apreface by Yves Meyer
- [5] A. Cheskidov, S. Friedlander and R. Shvydkoy. On the energy equality for weak solutions of the 3D Navier-Stokes equations, Advances in mathematical fluid mechanics Springer, Berlin, (2010) 171-175.

- [6] L. Escauriaza, G. Seregin, V. Šverák, $L^{3,\infty}$ solutions to the Navier-Stokes equations and backward uniqueness, *Russian Math. Surveys* 58 (2003) 186-212.
- [7] E.B. Fabes, B.F. Jones, and N.M. Riviere, The initial value problem for the Navier-Stokes equations with data in L^p, Arch. Rational Mech. Anal. 45 (1972) 222-240.
- [8] R. Farwig and Y. Taniuchi, On the energy equality of Navier-Stokes equations in general unbounded domains, *Arch.Math.* (Basel) 95(5) (2010) 447-456.
- [9] H. Fujita and T. Kato, On the nonstationary Navier-Stokes system, Rend. Sem. Mat. Univ. Padova 32 (1962) 243-260.
- [10] C. He and Y.Wang, Limiting case for the regularity criterion of the Navier-Stokes equations and the magnetohydrodynamic equations, *Sci. China Math.* 53 (2010) 1767-1774.
- [11] E. Hopf, Uber die Aufangswertaufgabe f
 ür die hydrodynamischen Grundgleichungen, Math. Nachr. 4 (1950-51) 213-231.
- [12] S. Kaniel and M. Shinbrot, Smoothness of weaksolutions of the Navier-Stokes equations, Arch. Rational Mech. Anal. 24 (1967) 302-324.
- [13] A.A. Kiselev and O.A. Ladyzhenskaya, On the existence and uniqueness of the solution of the nonstationary problem for a viscous incompressible fluid, *Izv. Akad. Nauk SSSR Ser. Mat.* 21 (1957) 655-680.
- [14] H. Kozono and Y. Taniuchi, Biliniear estimates in BMO and the Navier-Stokes equations, Math.z. 235 (2000) 173-194.
- [15] H. Kim and H. Kozono, Interior regularity criteria in weak spaces for the Navier-Stokes equa- tions, *Manuscripta Math.* bf 115 (2004) 85-100.
- [16] I. Kukavica, Role of the pressure for validity of the energy equality for solutions of the Navier-Stokes equation, J. Dynam. Differential Equations 18(2) (2006) 461-482.
- [17] T.M. Leslie and R. Shvydkoy, Conditions implying energy equality for weak solutions of the Navier-Stokes equations. SIAM J. Math. Anal. 50(1) (2018) 870-890.
- [18] T.M. Leslie and R. Shvydkoy, The energy measure for the Euler and Navier-Stokes equations, Arch. Rational Mech. Anal. 230 (2018) 459-492.
- [19] O. A. Ladyzhěnskaya and G. A. Seregin, On partial regularity of suitable weak solutions to the three-dimensional Navier-Stokes equations, J. Math. Fluid Mech. 1 (1999) 356-387.
- [20] O.A.Ladyzhĕnskaja, V.A.Solonnikov and N.N.Ural'ceva, Linear and quasilinear equations of parabolic type. Translated from the Russianby S.Smith.T ranslations of Mathematical Monographs, AmericanMathematicalSociety, Providence, R.I. 23 1968.
- [21] O. A. Ladyzhěnskaya, Uniqueness and smoothness of generalized solutions of Navier-Stokes equations, Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 5 (1967) 169-185.
- [22] F. H. Lin, A new proof of the Caffarelli-Kohn-Nirenberg theorem, Comm. Pure Appl. Math. 51 (1998) 241-257.
- [23] J.L. Lions, Sur la régularité et l'unicité des solutions turbulentes des équations de Navier-Stokes, Rend.Sem.Mat.Univ. Padova. 30 (1960) 16-23.

- [24] A. Cheslidov, X. Luo, Energy equality for the Navier-Stokes equations in weak-in-time Onsager spaces, *Nonlinearity* 33 (2020) 1388-1403.
- [25] P. Mattila, Geometry of sets and measures in Euclidean spaces: Fractals and rectifiability, Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge 44 (1995).
- [26] G. Prodi, Un teorema di unicitá per le equazioni di Navier-Stokes, Ann. Mat. Pure Appl. (4)48 (1959) 173-182.
- [27] G.A. Seregin and V. Šverák, Navier-Stokes equations with lower bounds on the pressure. Arch. Ration. Mech. Anal. 163(1) (2002) 65-86.
- [28] J. Serrin, The initial value problem for the Navier-Stokes equations, Nonlinear Problems (Proceedings of a Symposium, Madison, Wis.), R. T. Langer, ed., University of Wisconsin, Madison (1963) 69-98.
- [29] J. Serrin, On the interior regularity of weak solutions of the Navier-Stokes equations, Arch. Rat. Mech. Anal. 9 (1962) 187-195.
- [30] M. Shinbrot, The energy equation for the Navier-Stokes system, SIAMJ. Math. Anal.5 (1974) 948-954.
- [31] R. Shvydkoy. A geometric condition implying an energy equality for solutions of the 3D Navier-Stokes equation, J. Dynam. Differential Equations 21(1) (2009) 117-125.
- [32] R. Shvydkoy, A study of energy concentration and drain in incompressible fluids, *Nonlinearity* 26(2) (2013) 425-436.
- [33] T. Tao, Quantitative bounds for critically bounded solutions to the Navier-Stokes equations, arXiv:1908.04958v2
- [34] W. Tan and F. Wu, Energy conservation and regularity for the 3D magneto-hydrodynamics equations, *preprint*
- [35] W. von Wahl, The Equations of Navier-Stokes and Abstract Parabolic Equations, Aspects of Mathematics No. E8, *Vieweg, Braunschweig* 1985.
- [36] W. Wang, Z. Zhang, On the interior regularity criteria and the number of singular points to the Navier-Stokes equations, J. Anal. Math. 132 (2014) 139-170.