Local estimates for conformal *Q***-curvature equations**

Tianling Jin * and Hui Yang

July 9, 2021

Abstract

We derive local estimates of positive solutions to the conformal Q-curvature equation

$$(-\Delta)^m u = K(x) u^{\frac{n+2m}{n-2m}} \quad \text{in } \Omega \setminus \Lambda$$

near their singular set Λ , where $\Omega \subset \mathbb{R}^n$ is an open set, K(x) is a positive continuous function on Ω , Λ is a closed subset of \mathbb{R}^n , $2 \leq m < n/2$ and m is an integer. Under certain flatness conditions at critical points of K on Λ , we prove that $u(x) \leq C[\operatorname{dist}(x,\Lambda)]^{-(n-2m)/2}$ when the upper Minkowski dimension of Λ is less than (n-2m)/2.

Keywords: conformal Q-curvature equations, local estimates, singular set.

Mathematics Subject Classification (2010): 35J91; 35B40; 45M05

1 Introduction and main results

In this paper, we study the higher order conformal Q-curvature equation

$$(-\Delta)^m u = K(x) u^{\frac{n+2m}{n-2m}}, \quad u > 0 \quad \text{in } \Omega \setminus \Lambda$$
 (1.1)

with a singular set Λ , where $\Omega \subset \mathbb{R}^n$ is an open set, K(x) is a positive continuous function on Ω , Λ is a closed subset of \mathbb{R}^n , $1 \leq m < n/2$ and m is an integer. Throughout the paper, K(x) is assumed to be bounded between two positive constants in Ω .

When m = 1, Eq. (1.1) is the conformal scalar curvature equation which reads as

$$-\Delta u = K(x)u^{\frac{n+2}{n-2}}, \quad u > 0 \quad \text{in } \Omega \backslash \Lambda.$$
 (1.2)

This equation appears in the problem of finding a metric conformal to the flat metric δ_{ij} on \mathbb{R}^n such that K(x) is the scalar curvature of the new metric $u^{4/(n-2)}\delta_{ij}$. The classical works of Schoen and Yau [47–49] on the Yamabe problem and conformally flat manifolds have indicated the

^{*}T. Jin was partially supported by Hong Kong RGC grant GRF 16302217.

importance of studying the equation (1.2) with a singular set. In particular, an interesting question is to understand how u(x) tends to infinity when x approaches the singular set. When $K(x) \equiv 1$ and $\Lambda = \{0\}$ is an isolated singularity in Ω , Caffarelli, Gidas and Spruck in the pioneering paper [6] proved that every singular solution u of (1.2) is asymptotically radially symmetric near 0, and further proved that u is asymptotic to a radial singular solution of the same equation on $\mathbb{R}^n \setminus \{0\}$. Later, Korevaar, Mazzeo, Pacard and Schoen in [32] studied refined asymptotics and expanded such a singular solution u to the first order. Han, Li and Li [24] recently established the expansions up to arbitrary orders for such a singular solution u. Subsequent to [6], other second-order Yamabe type equations with isolated singularities related to (1.2) have also been studied; see, for example, [9, 24, 25, 33, 34, 40, 52] and the references therein. When $K(x) \equiv 1$ and Λ is a general singular set with Newtonian capacity zero, Chen and Lin [10] proved that any solution u of (1.2) satisfies the following a priori estimate

$$u(x) \le C[\operatorname{dist}(x,\Lambda)]^{-\frac{n-2}{2}},\tag{1.3}$$

and showed that u is asymptotically symmetric near Λ based on this estimate. In a series of masterful papers [11–13,39], Chen and Lin studied the equation (1.2) in the case when K(x) is a non-constant positive function. They first proved under some flatness conditions on K(x) that every C^2 solution u of (1.2) satisfies the a priori estimate (1.3) via the method of moving planes, and then applied the Pohozaev identity to describe the precise asymptotic behavior of solutions when the singularity set Λ is isolated. By using the method of moving spheres, Zhang in [54] simplified and improved the argument of Chen-Lin [11] to derive the local estimate (1.3) for the equation (1.2) under some flatness assumptions on K(x). In [51], Taliaferro and Zhang further gave conditions on K(x) to characterize the precise behavior of solutions of (1.2) near an isolated singularity.

When m=2, Eq. (1.1) is a fourth-order equation with critical Sobolev exponent. One of the motivations to study this equation arises from the problem of finding a metric which is conformal to the flat metric on \mathbb{R}^n such that K(x) is the fourth-order Q curvature of the new metric $u^{4/(n-4)}\delta_{ij}$ (see [5,45]). For $m\geq 3$, the higher-order equation (1.1) also arises in the study of similar problem in conformal geometry (see, e.g., [22]). We refer to Gursky-Malchiodi [23] and Hang-Yang [26] for the recent progress of the fourth-order Q curvature problem on compact Riemannian manifolds. In [8], Chang, Hang and Yang have shown that the Hausdorff dimension of singular set $\dim_H(\Lambda) < (n-4)/2$ is a necessary condition for the existence of a complete conformal metric whose scalar curvature and fourth-order Q curvature both have a positive lower bound. For higher values $m\geq 3$ and a smooth k-dimensional submanifold Λ , González-Mazzeo-Sire [21] showed that $\Gamma(\frac{n}{4}-\frac{k}{2}+\frac{m}{2})/\Gamma(\frac{n}{4}-\frac{k}{2}-\frac{m}{2})>0$ is necessary to have a complete conformal metric under some conditions, which holds in particular when k<(n-2m)/2.

The problem of characterizing asymptotic behavior of solutions of (1.1) for $m \geq 2$ is significantly more challenging due to the lack of maximum principle. Recently, this problem has also attracted much attention. When $K(x) \equiv 1$ and $\Lambda = \{0\}$ is an isolated singularity in Ω , Jin and Xiong in [30] proved sharp blow up rates and the asymptotic radial symmetry of singular solutions of (1.1) near the singularity 0 under the sign assumptions

$$(-\Delta)^s u \ge 0 \quad \text{in } \Omega \setminus \{0\} \quad \text{for all } s = 1, \dots, m - 1. \tag{1.4}$$

Based on the a priori estimates of Jin and Xiong, in the case of $K(x) \equiv 1$, Andrade-do Ó [1] and

Ratzkin [46] further proved for m=2 that singular solutions of (1.1) satisfying (1.4) are asymptotic to positive singular solutions of $(-\Delta)^2 u = u^{(n+4)/(n-4)}$ on $\mathbb{R}^n \setminus \{0\}$. All these singular solutions on $\mathbb{R}^n \setminus \{0\}$ have been classified by Frank and König in [20] using ODE analysis according to the radial symmetry result of Lin [38]. When $K(x) \equiv 1$ and Λ is a general singular set with the upper Minkowski dimension being less than (n-2m)/2, Du and Yang in [16] established the following a priori estimate

$$u(x) \le C[\operatorname{dist}(x,\Lambda)]^{-\frac{n-2m}{2}} \tag{1.5}$$

for any solution u of (1.1) satisfying (1.4) in $\Omega \setminus \Lambda$, and proved further that u is asymptotically symmetric near Λ . When m=2 and $\Omega=\mathbb{R}^n$, the estimate (1.5) was obtained in Chang, Han and Yang [9] under the assumption that the metric $u^{4/(n-4)}\delta_{ij}$ has nonnegative scalar curvature.

Our interest in this paper is the situation where the Q-curvature K(x) is a non-constant positive function. Under some conditions on the order of flatness at critical points of K on Λ , we will establish the local estimate (1.5) for any solution of (1.1) near its singular set Λ when the upper Minkowski dimension of Λ is less than (n-2m)/2.

We first recall the definition of the Minkowski dimension (see, e.g., [31,41]). Suppose $E \subset \mathbb{R}^n$ is a compact set, the λ -dimensional Minkowski r-content of E is defined by

$$\mathcal{M}_r^{\lambda}(E) = \inf \left\{ lr^{\lambda} \mid E \subset \bigcup_{k=1}^l B(x_k, r), \ x_k \in E \right\},$$

and the upper and lower Minkowski dimensions are defined, respectively, as

$$\overline{\dim}_M(E) = \inf \Big\{ \lambda \ge 0 \ \big| \ \limsup_{r \to 0} \mathcal{M}_r^{\lambda}(E) = 0 \Big\},$$

$$\underline{\dim}_{M}(E) = \inf \Big\{ \lambda \geq 0 \; \big| \; \liminf_{r \to 0} \mathcal{M}_{r}^{\lambda}(E) = 0 \Big\}.$$

If $\overline{\dim}_M(E) = \underline{\dim}_M(E)$, then the common value, denoted by $\dim_M(E)$, is the Minkowski dimension of E. Recall also that for a compact set $E \subset \mathbb{R}^n$, we have the relation $\dim_H(E) \leq \underline{\dim}_M(E) \leq \overline{\dim}_M(E)$, where $\dim_H(E)$ is the Hausdorff dimension of E. We also introduce a notation $\mathcal{C}^{\alpha}(\Omega)$ with $\alpha > 0$ which will be used later.

Definition 1.1.

- 1. If α is a positive integer, then $C^{\alpha}(\Omega)$ is the usual space $C^{\alpha}(\Omega)$.
- 2. If $\alpha > [\alpha]$, then $\mathcal{C}^{\alpha}(\Omega)$ is the set of all functions $f \in C^{[\alpha]}(\Omega)$ satisfying

$$|\nabla^{[\alpha]}f(x) - \nabla^{[\alpha]}f(y)| \le c(|x-y|)|x-y|^{\alpha-[\alpha]}, \quad x,y \in \Omega,$$

where $c(\cdot)$ is a nonnegative continuous function with c(0) = 0.

We will use $B_r(x)$ to denote the open ball of radius r in \mathbb{R}^n with center x and write $B_r(0)$ as B_r for short. From now on, without loss of generality, we take the domain $\Omega = B_2$.

Our assumption on the Q-curvature K(x) is as follows:

1. For
$$n = 2m + 1$$
, $K \in \mathcal{C}^{\frac{1}{2}}(B_2)$.

- 2. For n = 2m + 2, $K \in C^1(B_2)$.
- 3. For $n \ge 2m + 3$, $K \in C^1(B_2)$ and one of the following is satisfied:
 - (K1) If $x \in \Lambda$ is a critical point of K, then there exists a neighborhood N of x such that

$$|c_1|y - x|^{\alpha - 1} \le |\nabla K(y)| \le c_2|y - x|^{\alpha - 1}$$
 (1.6)

and for any $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon)$ such that for $|z - y| < \delta |y - x|$ we have

$$|\nabla K(z) - \nabla K(y)| < \varepsilon |y - x|^{\alpha - 1},\tag{1.7}$$

where $\alpha > 1$, $y, z \in N$, c_1 and c_2 are two positive constants.

(K2) $K \in \mathcal{C}^{\alpha}(B_2)$ with $\alpha = \frac{n-2m}{2}$. In addition, for $n \geq 2m+4$, if $x \in \Lambda$ is a critical point of K, then there exists a neighborhood N of x such that

$$|\nabla^{i}K(y)| \le c(|y-x|)|\nabla K(y)|^{\frac{\alpha-i}{\alpha-1}}, \quad 2 \le i \le [\alpha], \quad y \in N, \tag{1.8}$$

where $c(\cdot)$ is a nonnegative continuous function satisfying c(0) = 0.

Then our main result in this paper is

Theorem 1.2. Suppose that $1 \le m < n/2$ and m is an integer. Let $\Lambda \subset B_{1/2}$ be a compact set with the upper Minkowski dimension $\overline{\dim}_M(\Lambda)$ (not necessarily an integer), $\overline{\dim}_M(\Lambda) < \frac{n-2m}{2}$, or $\Lambda \subset B_{1/2}$ be a smooth k-dimensional closed manifold with $k \le \frac{n-2m}{2}$. Let K satisfy the above assumption and let $u \in C^{2m}(B_2 \setminus \Lambda)$ be a solution of

$$(-\Delta)^m u = K(x) u^{\frac{n+2m}{n-2m}}, \quad u > 0 \quad \text{in } B_2 \backslash \Lambda.$$
 (1.9)

Suppose

$$(-\Delta)^s u \ge 0 \quad \text{in } B_2 \backslash \Lambda, \quad s = 1, \dots, m - 1. \tag{1.10}$$

Then there exists a constant C > 0 such that

$$u(x) \le C[\operatorname{dist}(x,\Lambda)]^{-\frac{n-2m}{2}} \tag{1.11}$$

for all $x \in B_1 \setminus \Lambda$, where $dist(x, \Lambda)$ is the distance between x and Λ .

Remark 1.3. For $n \ge 2m+4$, if we only assume $K \in C^1(B_2)$, then Theorem 1.2 does not hold. Indeed, when $n \ge 2m+4$ and $\Lambda = \{0\}$, Du and Yang in [17] have shown the existence of $K \in C^1(B_2)$ such that Eq. (1.9) has a C^{2m} solution u(x) which satisfies (1.10) but does not satisfy (1.11), and can even be constructed to be arbitrarily large near its singularity 0. Such large singular solution of (1.9) extends a similar result of Taliaferro [50] for m=1. On the other hand, Theorem 1.2 also indicates that Theorem 1.3 of [17] is not true in dimension n=2m+1 or n=2m+2 for $m\ge 2$.

Remark 1.4. When m=2, a solution u of (1.9) defines a conformal metric $g_{ij}=u^{\frac{4}{n-4}}\delta_{ij}$ which has Q-curvature $\frac{2}{n-4}K(x)$. Assuming that the scalar curvature of g_{ij} is positive, then one can obtain the sign condition $-\Delta u>0$. We also mention that under the assumption of positive scalar curvature, Gursky and Malchiodi in [23] studied the positivity of the Paneitz operator and its Green's function.

Remark 1.5. The assumption (K2) also covers the situation of $K(x) \equiv 1$. In this case, Mazzeo and Pacard in [42] have constructed singular solutions of (1.9) for m = 1 when Λ is a smooth submanifold of dimension $k \leq (n-2)/2$, and Hyder-Sire [27] recently proved for m = 2 that Eq. (1.9) has singular solutions which satisfy (1.10) when Λ is a smooth submanifold of dimension k < (n-4)/2.

As mentioned earlier, Eq. (1.9) is more challenging for $m \geq 2$ to study since the maximum principle is lacking. Notice also that the sign conditions (1.10) may change when performing the Kelvin transformation. Thus, even under the assumption (1.10), it is still difficult to apply the method of moving planes or moving spheres directly to the local differential equation (1.9). Inspired by the work of Jin-Li-Xiong [29,30], we will rewrite the differential equation (1.9) into the local integral equation (1.12) below and derive local estimates of singular solutions to this integral equation. Similar idea has also been used in [16] to study the case when K(x) is identically a positive constant and the singular set Λ is not isolated.

Suppose the dimension $n \ge 1$, $0 < \sigma < \frac{n}{2}$ is a real number, and Σ is a closed set in \mathbb{R}^n . We consider the local integral equation

$$u(x) = \int_{B_2} \frac{K(y)u(y)^{\frac{n+2\sigma}{n-2\sigma}}}{|x-y|^{n-2\sigma}} dy + h(x), \quad u(x) > 0, \quad x \in B_2 \backslash \Sigma,$$
 (1.12)

where $u \in L^{\frac{n+2\sigma}{n-2\sigma}}(B_2) \cap C(B_2 \setminus \Sigma)$ and $h \in C^1(B_2)$ is a positive function. Under the assumptions in Theorem 1.2, by a similar argument as in [16] we can show $u \in L^{\frac{n+2\sigma}{n-2\sigma}}(B_2)$ and can rewrite the equation (1.9) locally into the integral equation (1.12) after some scaling (see Theorem 2.3 in the next section).

Next we state the corresponding result for singular solutions of the integral equation (1.12). Denote \mathcal{L}^n the *n*-dimensional Lebesgue measure on \mathbb{R}^n . We assume that the positive *Q*-curvature function K(x) satisfies the one of the following:

(K1) $K \in C^1(B_2)$. If $x \in \Sigma$ is a critical point of K, then there exists a neighborhood N of x such that

$$c_1|y-x|^{\alpha-1} \le |\nabla K(y)| \le c_2|y-x|^{\alpha-1}$$
 (1.13)

and for any $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon)$ such that for $|z - y| < \delta |y - x|$ we have

$$|\nabla K(z) - \nabla K(y)| < \varepsilon |y - x|^{\alpha - 1}, \tag{1.14}$$

where $\alpha > 1$, $y, z \in N$, c_1 and c_2 are two positive constants.

(K2) $K \in \mathcal{C}^{\alpha}(B_2)$ with $\alpha = \frac{n-2\sigma}{2}$. In addition, for $n \geq 2\sigma + 4$, if $x \in \Sigma$ is a critical point of K, then there exists a neighborhood N of x such that

$$|\nabla^{i}K(y)| \le c(|y-x|)|\nabla K(y)|^{\frac{\alpha-i}{\alpha-1}}, \quad 2 \le i \le [\alpha], \quad y \in N, \tag{1.15}$$

where $c(\cdot)$ is a nonnegative continuous function satisfying c(0) = 0.

Theorem 1.6. Suppose $n \geq 2$, $1 \leq \sigma < n/2$, and Σ is a closed set in \mathbb{R}^n with $\mathcal{L}^n(\Sigma) = 0$. Suppose that K(x) satisfies the assumption (K1) or (K2) above. Let $h \in C^1(B_2)$ be a positive

function and $u \in L^{\frac{n+2\sigma}{n-2\sigma}}(B_2) \cap C(B_2 \setminus \Sigma)$ be a positive solution of (1.12). Then there exists a constant C > 0 such that

$$u(x) \le C[\operatorname{dist}(x,\Sigma)]^{-\frac{n-2\sigma}{2}} \tag{1.16}$$

for all $x \in B_1 \backslash \Sigma$, where $dist(x, \Sigma)$ is the distance between x and Σ .

Remark 1.7. For the integral equation (1.12), we only assume that the singular set Σ has n-dimensional Lebesgue measure zero. This is a weaker condition than singular set of Newtonian capacity zero, which is used in [10, 11, 54] to study the second-order scalar curvature equation (1.2). In particular, if one obtains a solution $u \in C^{2m}(B_2 \setminus \Lambda) \cap L^{\frac{n+2m}{n-2m}}(B_2)$ to (1.9) satisfying (1.10) by whatever method with $\mathcal{L}^n(\Lambda) = 0$, then u is a distributional solution in B_2 and thus Theorem 1.6 can give the local estimate (1.11) of u near Λ even though the Newtonian capacity of Λ might be greater than 0. For example, Pacard [44] have constructed solutions with such high dimensional singular set to (1.2) when $K(x) \equiv 1$ in dimension 4 and in dimension 6.

Remark 1.8. In Theorem 1.6, we assume that the order of the integral equation (1.12) satisfies $1 \le \sigma < n/2$ which is sufficient to apply to higher order Q-curvature equation (1.1). When $0 < \sigma < 1$, equation (1.12) is closely related to the fractional Nirenberg problem (see [29]). For the fractional case $0 < \sigma < 1$, our proof of Theorem 1.6 encounters a difficulty due to the more singular properties of the integral kernel $G(0,\lambda;\xi,z)$ defined in (3.9). More specifically, the negativity of Φ_{λ} in (4.37) is very important when using the moving sphere method, but in the case of $0 < \sigma < 1$, the negative part of the integral in (4.37) cannot control the positive parts according to the estimates of $G(0,\lambda;\xi,z)$ in Lemma 3.1. We plan to deal with this difficulty in future work. We also mention that fractional order critical equations with singularities in the case $K(x) \equiv 1$ have been studied in [2, 3, 7, 15, 18, 28] and so on.

We will prove Theorem 1.6 in the spirit of the works of Chen, Lin, Taliaferro and Zhang [11,39,51,54] by using the moving sphere method of an integral form introduced by Li [35], which is inspired by Li-Zhu [37] and Li-Zhang [36]. One difference is that the authors of [11,39,51,54] dealt directly with the second-order equation (1.2), while we work with the integral equation (1.12) by exploring its various specific features. Thus we need some analysis techniques to overcome the difficulties caused by the absence of a maximum principle. Another difference is the non-locality of the integral equation (1.12).

The rest of this paper is organized as follows. In Section 2, we show the integral representation for singular positive solutions to the differential equation (1.9). In Section 3, we provide some preliminary estimates for the integral kernel involved in the moving sphere method of integral form. In Section 4, we give a detailed proof of Theorem 1.6 under the assumption (K1). In Section 5, we present the proof of Theorem 1.6 under the assumption (K2). Finally, Theorem 1.2 follows from Theorem 1.6 and Theorem 2.3 via the covering and rescaling arguments.

2 An integral representation for singular solutions

In this section, we show that every singular positive solution of the differential equation (1.9) satisfies the integral equation (1.12) in some local sense under suitable assumptions. We first prove that under the assumptions of Theorem 1.2 ((1.10) is not needed here), $u \in L^{\frac{n+2m}{n-2m}}_{loc}(B_2)$ and u is a distributional solution in the entire ball B_2 .

Proposition 2.1. Suppose that $1 \leq m < n/2$ and m is an integer. Let $\Lambda \subset B_{1/2}$ be a compact set with the upper Minkowski dimension $\overline{\dim}_M(\Lambda)$ (not necessarily an integer), $\overline{\dim}_M(\Lambda) < \frac{n-2m}{2}$, or $\Lambda \subset B_{1/2}$ be a smooth k-dimensional closed manifold with $k \leq \frac{n-2m}{2}$. Let K be a positive continuous function on \overline{B}_2 and let $u \in C^{2m}(\overline{B}_2 \backslash \Lambda)$ be a positive solution of (1.9). Then $u \in L^{\frac{n+2m}{n-2m}}(B_2)$ and u is a distributional solution in the entire ball B_2 , i.e., we have

$$\int_{B_2} u(-\Delta)^m \varphi dx = \int_{B_2} K u^{\frac{n+2m}{n-2m}} \varphi dx \tag{2.1}$$

for every $\varphi \in C_c^{\infty}(B_2)$.

Proof. The proof is very similar to that of [16, Proposition 2.1], so we omit the details. See also some related arguments in [4,53].

Suppose n > 2m. Let $G_m(x, y)$ be the Green function of $(-\Delta)^m$ on B_2 under the Navier boundary condition:

$$\begin{cases} (-\Delta)^m G_m(x,\cdot) = \delta_x & \text{in } B_2, \\ G_m(x,\cdot) = -\Delta G_m(x,\cdot) = \dots = (-\Delta)^{m-1} G_m(x,\cdot) = 0 & \text{on } \partial B_2, \end{cases}$$
 (2.2)

where δ_x is the Dirac measure to the point $x \in B_2$. Then, for any $u \in C^{2m}(B_2) \cap C^{2m-2}(\overline{B}_2)$ we have

$$u(x) = \int_{B_2} G_m(x, y) (-\Delta)^m u(y) dy + \sum_{i=1}^m \int_{\partial B_2} H_i(x, y) (-\Delta)^{i-1} u(y) dS_y, \qquad (2.3)$$

where

$$H_i(x,y) = -\frac{\partial}{\partial \nu_y} (-\Delta_y)^{m-i} G_m(x,y) \quad \text{for } x \in B_2, y \in \partial B_2.$$

In particular, taking $u \equiv 1$ in (2.3) yields

$$\int_{\partial B_2} H_1(x, y) dS_y = 1 \quad \text{for all } x \in B_2.$$
 (2.4)

Moreover, a direct computation gives

$$G_m(x,y) = c_{n,m}|x-y|^{2m-n} + A_m(x,y),$$
(2.5)

 $c_{n,m}=rac{\Gamma(rac{n}{2}-m)}{2^{2m}\pi^{n/2}\Gamma(m)},$ $A_m(x,y)$ is smooth in $B_2\times B_2$, and

$$H_i(x,y) \ge 0, \quad i = 1, \dots, m.$$
 (2.6)

Proposition 2.2. Under the assumptions of Proposition 2.1, we have

$$u(x) = \int_{B_2} G_m(x, y) K(y) u(y)^{\frac{n+2m}{n-2m}} dy + \sum_{i=1}^m \int_{\partial B_2} H_i(x, y) (-\Delta)^{i-1} u(y) dS_y$$
 (2.7)

for all $x \in B_2 \backslash \Lambda$.

Proof. For any $x \in B_2 \setminus \Lambda$, we define

$$v(x) = \int_{B_2} G_m(x, y) K(y) u(y)^{\frac{n+2m}{n-2m}} dy + \sum_{i=1}^m \int_{\partial B_2} H_i(x, y) (-\Delta)^{i-1} u(y) dS_y.$$

Because $u(y)^{\frac{n+2m}{n-2m}} \in L^1(B_2)$ and the Riesz potential $|x|^{2m-n}$ is weak type $\left(1, \frac{n}{n-2m}\right)$, we obtain $v \in L^{\frac{n}{n-2m}}(B_2) \cap L^1(B_2)$. Define w = u - v. By Proposition 2.1 we have that w satisfies

$$(-\Delta)^m w = 0 \quad \text{in } B_2$$

in the distributional sense, i.e., for any $\varphi \in C_c^{\infty}(B_2)$,

$$\int_{B_2} w(-\Delta)^m \varphi dx = 0.$$

It follows from the regularity for polyharmonic functions (see, e.g., Mitrea [43]) that w is smooth and satisfies $(-\Delta)^m w = 0$ in B_2 . Note that $w = -\Delta w = \cdots = (-\Delta)^{m-1} w = 0$ on ∂B_2 , hence $w \equiv 0$. This implies that u = v in $B_2 \setminus \Lambda$.

Now we show that u satisfies the integral equation (1.12) in some local sense under the additional assumption (1.10).

Theorem 2.3. Suppose that $1 \le m < n/2$ and m is an integer. Let $\Lambda \subset B_{1/2}$ be a compact set with the upper Minkowski dimension $\overline{\dim}_M(\Lambda)$ (not necessarily an integer), $\overline{\dim}_M(\Lambda) < \frac{n-2m}{2}$, or $\Lambda \subset B_{1/2}$ be a smooth k-dimensional closed manifold with $k \le \frac{n-2m}{2}$. Let K be a positive continuous function on \overline{B}_2 and let $u \in C^{2m}(B_2 \setminus \Lambda)$ be a positive solution of

$$(-\Delta)^m u = K(x) u^{\frac{n+2m}{n-2m}} \quad \text{in } B_2 \backslash \Lambda. \tag{2.8}$$

Suppose

$$(-\Delta)^s u \ge 0 \quad \text{in } B_2 \setminus \Lambda, \quad s = 1, \dots, m - 1. \tag{2.9}$$

Then there exists $\tau > 0$ (independent of $x \in \Lambda$) such that for any $x_0 \in \Lambda$ we have

$$u(x) = c_{n,m} \int_{B_{\tau}(x_0)} \frac{K(y)u(y)^{\frac{n+2m}{n-2m}}}{|x-y|^{n-2m}} dy + h_1(x) \quad \text{for } x \in B_{\tau}(x_0) \backslash \Lambda,$$
 (2.10)

where $h_1(x)$ is a positive smooth function in $B_{\tau}(x_0)$.

Proof. We may assume, without loss of generality, that $u \in C^{2m}(\overline{B}_2 \backslash \Lambda)$ and u > 0 in $\overline{B}_2 \backslash \Lambda$. Otherwise, we just consider the equation in a smaller ball.

From the assumptions on the singular set Λ we obtain $\operatorname{Cap}(\Lambda) = 0$, where $\operatorname{Cap}(\Lambda)$ is the Newtonian capacity of Λ (see, e.g., [19]). Since u > 0 and $-\Delta u \geq 0$ in $B_2 \setminus \Lambda$, the maximum principle (see, e.g., [10, Lemma 2.1]) leads to

$$u(x) \ge c_0 := \inf_{\partial B_2} u > 0 \quad \text{ for all } x \in \overline{B}_2 \backslash \Lambda.$$

By Proposition 2.1 we have $u^{\frac{n+2m}{n-2m}} \in L^1(B_2)$. Thus, there exists $0 < \tau < \frac{1}{4}$ independent of $z \in B_1$ such that

$$\int_{B_{\tau}(z)} |A_m(x,y)| K(y) u(y)^{\frac{n+2m}{n-2m}} dy < \frac{c_0}{2} \quad \text{ for all } x \in B_{\tau}(z) \subset B_{3/2},$$

where $A_m(x,y)$ is defined in (2.5). For every $x_0 \in \Lambda$, using Proposition 2.2 we can write

$$u(x) = c_{n,m} \int_{B_{\tau}(x_0)} \frac{K(y)u(y)^{\frac{n+2m}{n-2m}}}{|x-y|^{n-2m}} dy + h_1(x) \quad \text{for } x \in B_{\tau}(x_0) \setminus \Lambda,$$

where

$$h_{1}(x) = \int_{B_{\tau}(x_{0})} A_{m}(x, y) K(y) u(y)^{\frac{n+2m}{n-2m}} dy + c_{n,m} \int_{B_{2} \setminus B_{\tau}(x_{0})} G_{m}(x, y) K(y) u(y)^{\frac{n+2m}{n-2m}} dy$$

$$+ \sum_{i=1}^{m} \int_{\partial B_{2}} H_{i}(x, y) (-\Delta)^{i-1} u(y) dS_{y}$$

$$\geq -\frac{c_{0}}{2} + \int_{\partial B_{2}} H_{1}(x, y) u(y) dS_{y}$$

$$\geq -\frac{c_{0}}{2} + \inf_{\partial B_{2}} u = \frac{c_{0}}{2} > 0 \quad \text{for } x \in B_{\tau}(x_{0}),$$

where we have used the sign conditions (2.9) in the first inequality and (2.4) in the second inequality. It is easy to check that h_1 is smooth in $B_{\tau}(x_0)$ and satisfies $(-\Delta)^m h_1 = 0$ in $B_{\tau}(x_0)$. The proof of Theorem 2.3 is completed.

Proof of Theorem 1.2. It follows from Theorem 1.6 and Theorem 2.3 via the covering and rescaling arguments. \Box

3 Preliminary results

For $x \in \mathbb{R}^n$, $\lambda > 0$ and a function u, we denote

$$\xi^{x,\lambda} = x + \frac{\lambda^2(\xi - x)}{|\xi - x|^2} \quad \text{for } x \neq \xi \in \mathbb{R}^n \quad \text{and} \quad \Omega^{x,\lambda} = \{\xi^{x,\lambda} : \xi \in \Omega\}.$$
 (3.1)

Let u be a positive function, its Kelvin transformation is defined as

$$u^{x,\lambda}(\xi) = \left(\frac{\lambda}{|\xi - x|}\right)^{n - 2\sigma} u(\xi^{x,\lambda}). \tag{3.2}$$

Note that $(\xi^{x,\lambda})^{x,\lambda} = \xi$ and $(u^{x,\lambda})^{x,\lambda} \equiv u$. If x = 0, we use the notations $\xi^{\lambda} = \xi^{0,\lambda}$ and $u^{\lambda} = u^{0,\lambda}$, i.e.,

$$\xi^{\lambda} = \frac{\lambda^2 \xi}{|\xi|^2} \quad \text{and} \quad u^{\lambda}(\xi) = \left(\frac{\lambda}{|\xi|}\right)^{n-2\sigma} u\left(\frac{\lambda^2 \xi}{|\xi|^2}\right).$$
 (3.3)

Suppose $0 < \sigma < n/2$, $u \in L^{\frac{n+2\sigma}{n-2\sigma}}(B_2) \cap C(B_2 \backslash \Sigma)$ is a positive solution of (1.12) and $h \in C^1(B_2)$ is a positive function. If we extend both u and K to be identically 0 outside B_2 , then

$$u(x) = \int_{\mathbb{R}^n} \frac{K(y)u(y)^{\frac{n+2\sigma}{n-2\sigma}}}{|x-y|^{n-2\sigma}} dy + h(x) \quad \text{for } x \in B_2 \backslash \Sigma.$$
 (3.4)

Making a change of variables, we also have the following two identities (see, e.g., [35]),

$$\left(\frac{\lambda}{|\xi - x|}\right)^{n - 2\sigma} \int_{|z - x| \ge \lambda} \frac{K(z)u(z)^{\frac{n + 2\sigma}{n - 2\sigma}}}{|\xi^{x, \lambda} - z|^{n - 2\sigma}} dz = \int_{|z - x| \le \lambda} \frac{K(z^{x, \lambda})u^{x, \lambda}(z)^{\frac{n + 2\sigma}{n - 2\sigma}}}{|\xi - z|^{n - 2\sigma}} dz \tag{3.5}$$

and

$$\left(\frac{\lambda}{|\xi-x|}\right)^{n-2\sigma} \int_{|z-x|<\lambda} \frac{K(z)u(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|\xi^{x,\lambda}-z|^{n-2\sigma}} dz = \int_{|z-x|>\lambda} \frac{K(z^{x,\lambda})u^{x,\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|\xi-z|^{n-2\sigma}} dz. \tag{3.6}$$

Thus, we obtain

$$u^{x,\lambda}(\xi) = \int_{\mathbb{R}^n} \frac{K(z^{x,\lambda})u^{x,\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|\xi - z|^{n-2\sigma}} dz + h^{x,\lambda}(\xi) \quad \text{for } \xi \in (B_2 \backslash \Sigma)^{x,\lambda}.$$
 (3.7)

Hence, for any $x \in B_1$ and $\lambda < 1$, we have for any $\xi \in B_2 \setminus (\Sigma \cup \Sigma^{x,\lambda} \cup B_{\lambda}(x))$ that

$$u(\xi) - u^{x,\lambda}(\xi) = \int_{|z-x| \ge \lambda} G(x,\lambda;\xi,z) \left[K(z)u(z)^{\frac{n+2\sigma}{n-2\sigma}} - K(z^{x,\lambda})u^{x,\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right] dz + h^{x,\lambda}(\xi) - h(\xi),$$
(3.8)

where

$$G(x,\lambda;\xi,z) := \frac{1}{|\xi - z|^{n-2\sigma}} - \left(\frac{\lambda}{|\xi - x|}\right)^{n-2\sigma} \frac{1}{|\xi^{x,\lambda} - z|^{n-2\sigma}}, \quad |\xi - x|, |z - x| > \lambda > 0. \quad (3.9)$$

It is elementary to check that

$$G(x,\lambda;\xi,z) > 0$$
 for all $|\xi - x|, |z - x| > \lambda > 0$. (3.10)

We also give the following estimates for the integral kernel $G(0, \lambda; \xi, z)$. Their proofs are elementary, and we include them here for completeness.

Lemma 3.1. Assume $0 < \sigma < n/2$ and $\lambda > 0$. Then for $\xi, z \in \mathbb{R}^n \setminus \overline{B_{\lambda}}$, we have

(1) There exists a positive constant $C_1 = C_1(n, \sigma)$ such that for $|\xi - z| < \frac{1}{3}(|\xi| - \lambda)$,

$$G(0,\lambda;\xi,z) \ge C_1 \frac{1}{|\xi-z|^{n-2\sigma}}.$$
(3.11)

(2) There exist two positive constants $C_2 = C_2(n, \sigma)$ and $C_3 = C_3(n, \sigma)$ such that for $|\xi - z| \ge \frac{1}{3}(|\xi| - \lambda)$ and $|\xi| \le 10\lambda$,

$$C_2 \frac{(|\xi| - \lambda)(|z|^2 - \lambda^2)}{\lambda |\xi - z|^{n - 2\sigma + 2}} \le G(0, \lambda; \xi, z) \le C_3 \frac{(|\xi| - \lambda)(|z|^2 - \lambda^2)}{\lambda |\xi - z|^{n - 2\sigma + 2}}.$$
(3.12)

Moreover, if we only assume $\lambda < |\xi| \le 10\lambda$ and $|z| > \lambda$, then we have the second inequality of (3.12).

(3)
$$G(0, \lambda; \xi, z) = G(0, \lambda; z, \xi)$$
.

Remark 3.2. By the symmetry of $G(0, \lambda; \xi, z)$, we can reverse the roles of ξ and z in (1) and (2) of Lemma 3.1 and the corresponding conclusions still hold.

Proof. (1) Note that for $|\xi - z| \le \frac{1}{3}(|\xi| - \lambda)$ and $|\xi| > \lambda$, we have $|\xi^{\lambda} - z| \ge |\xi^{\lambda} - \xi| - |\xi - z| \ge 2|\xi - z|$. Thus

$$G(0, \lambda; \xi, z) \ge (1 - 2^{2\sigma - n})|\xi - z|^{2\sigma - n}$$
.

(2) The kernel $G(0, \lambda; \xi, z)$ can be rewritten as

$$G(0,\lambda;\xi,z) = |\xi - z|^{2\sigma - n} - \left| \frac{\lambda}{|\xi|} \xi - \frac{|\xi|}{\lambda} z \right|^{2\sigma - n}.$$
 (3.13)

By a direct calculation, we have the following formula

$$\left| \frac{\lambda}{|\xi|} \xi - \frac{|\xi|}{\lambda} z \right|^2 - |\xi - z|^2 = \frac{(|\xi|^2 - \lambda^2)(|z|^2 - \lambda^2)}{\lambda^2}, \quad \xi \neq 0.$$
 (3.14)

This obviously implies that the kernel $G(0, \lambda; \xi, z)$ is positive. Let $f(t) = t^{-(n-2\sigma)/2}$, then by the mean value theorem and (3.14) we get

$$G(0,\lambda;\xi,z) = f\left(|\xi-z|^2\right) - f\left(\left|\frac{\lambda}{|\xi|}\xi - \frac{|\xi|}{\lambda}z\right|^2\right)$$

$$= \frac{n-2\sigma}{2} \frac{(|\xi|^2 - \lambda^2)(|z|^2 - \lambda^2)}{\lambda^2 \left(\theta|\xi-z|^2 + (1-\theta)\left|\frac{\lambda}{|\xi|}\xi - \frac{|\xi|}{\lambda}z\right|^2\right)^{\frac{n-2\sigma+2}{2}}}$$

for some $\theta \in (0,1)$. When $|\xi - z| \ge \frac{1}{3}(|\xi| - \lambda)$ and $\lambda < |\xi| \le 10\lambda$ we have

$$\left|\frac{\lambda}{|\xi|}\xi - \frac{|\xi|}{\lambda}z\right| \leq \left|\frac{\lambda}{|\xi|}\xi - \frac{|\xi|}{\lambda}\xi\right| + \left|\frac{|\xi|}{\lambda}\xi - \frac{|\xi|}{\lambda}z\right| \leq 43|\xi - z|$$

and thus

$$G(0,\lambda;\xi,z) \ge C_2 \frac{(|\xi|-\lambda)(|z|^2-\lambda^2)}{\lambda|\xi-z|^{n-2\sigma+2}}$$

with some constant $C_2 = C_2(n, \sigma)$. On the other hand, if we only assume $\lambda < |\xi| \le 10\lambda$ and $|z| > \lambda$, then by (3.14) we obtain $|\xi - z| \le \left| \frac{\lambda}{|\xi|} \xi - \frac{|\xi|}{\lambda} z \right|$ and so

$$G(0,\lambda;\xi,z) \le C_1 \frac{(|\xi|-\lambda)(|z|^2-\lambda^2)}{\lambda|\xi-z|^{n-2\sigma+2}}$$

for some constant $C_1 = C_1(n, \sigma)$.

(3) Note that

$$\frac{|z|}{\lambda} \frac{|\xi|}{\lambda} |\xi^{\lambda} - z^{\lambda}| = \left| \frac{|z|}{|\xi|} \xi - \frac{|\xi|}{|z|} z \right| = |\xi - z| \quad \text{for } \xi, z \neq 0, \tag{3.15}$$

where we have used the basic identity $\left|\frac{|z|}{|\xi|}\xi - \frac{|\xi|}{|z|}z\right|^2 = |\xi - z|^2$. This implies that

$$\frac{|\xi|}{\lambda}|\xi^{\lambda} - z| = \frac{|z|}{\lambda}|z^{\lambda} - \xi| \quad \text{for } \xi, z \neq 0, \tag{3.16}$$

from which we obtain $G(0, \lambda; \xi, z) = G(0, \lambda; z, \xi)$. Lemma 3.1 is proved.

Lemma 3.3. Assume $0 < \sigma < n/2$. Let

$$U(y) = \left(\frac{1}{1 + |y - Re|^2}\right)^{\frac{n-2\sigma}{2}},$$

where $e \neq 0$ and R|e| > 10. Then there exists a positive constant $C = C(n, \sigma, R|e|)$ such that for $\lambda_0 = R|e| - 2$, we have

$$U(y) - U^{\lambda_0}(y) \ge C(|y|^2 - \lambda_0^2)|y|^{-(n-2\sigma+2)}$$
 for all $|y| \ge \lambda_0$

and

$$\frac{\partial (U - U^{\lambda_0})}{\partial u} > C > 0 \quad on \, \partial B_{\lambda_0},$$

where ν denotes the unit outer normal vector of ∂B_{λ_0} .

For $\lambda_1 = R|e| + 2$, we have

$$U(y) - U^{\lambda_1}(y) < 0$$
 for all $|y| > \lambda_1$.

Proof. Using the mean value theorem and the formula (3.14), we get that for any $\lambda > 0$,

$$U(y) - U^{\lambda}(y) = \left(\frac{1}{1 + |y - Re|^2}\right)^{\frac{n - 2\sigma}{2}} - \left(\frac{1}{\frac{|y|^2}{\lambda^2} + \frac{|y|^2}{\lambda^2}|y^{\lambda} - Re|^2}\right)^{\frac{n - 2\sigma}{2}}$$

$$= \frac{n - 2\sigma}{2} \cdot \frac{(|y|^2 - \lambda^2)(R^2|e|^2 - \lambda^2 + 1)}{\lambda^2 n^{(n - 2\sigma + 2)/2}},$$
(3.17)

where η is some number between $1+|y-Re|^2$ and $\frac{|y|^2}{\lambda^2}+\frac{|y|^2}{\lambda^2}|y^\lambda-Re|^2$. When $\lambda=\lambda_0$, it is easy to see that $R^2|e|^2-\lambda^2+1>0$ and $\eta\leq c|y|^2$ for some constant c=c(R)>0. Hence there exists $C=C(n,\sigma,R)>0$ such that

$$U(y) - U^{\lambda_0}(y) \ge C(|y|^2 - \lambda_0^2)|y|^{-(n-2\sigma+2)}$$
 for any $|y| \ge \lambda_0$.

This also implies that

$$\frac{\partial (U - U^{\lambda_0})}{\partial \nu} > C > 0 \quad \text{on } \partial B_{\lambda_0}$$

for another constant $C=C(n,\sigma,R)>0$, where ν means the unit outer normal vector of ∂B_{λ_0} . When $\lambda=\lambda_1$, we have $R^2|e|^2-\lambda^2+1<0$ and by (3.17),

$$U(y) - U^{\lambda_1}(y) < 0 \qquad \text{for any } |y| > \lambda_1.$$

Lemma 3.3 is proved.

4 Local estimates under the assumption (K1)

In this section, by using the method of moving spheres introduced by Li-Zhu [35, 37], we shall prove Theorem 1.6 under the assumption (K1) in the spirit of the works of Chen-Lin [11, 39] and Zhang [54]. Unlike [11,39,54] dealing directly with second-order differential equations, we study the problem in a framework of integral equations. In particular, analysis techniques for integral equations are needed to overcome the lack of maximum principle.

Proof of Theorem 1.6 under the assumption (K1). Suppose by contradiction that there exists a sequence $\{x_j\}_{j=1}^{\infty} \subset B_1 \setminus \Sigma$ such that

$$d_j := \operatorname{dist}(x_j, \Sigma) \to 0 \quad \text{ as } j \to \infty,$$

but

$$d_j^{\frac{n-2\sigma}{2}}u(x_j)\to\infty$$
 as $j\to\infty$.

Without loss of generality we may assume that $0 \in \Sigma$ and $x_j \to 0$ as $j \to \infty$. Since the following proof is very long, we first explain the idea and the sketch of the proof.

In Step 1, by using the blow up analysis we show that x_j can be chosen as the local maximum points of u.

In Step 2, we show that $\nabla K(0)=0$ under the assumption of $K\in C^1(B_2)$. If not, then we assume without loss of generality that $\nabla K(0)=e=(1,0,\ldots,0)$. Let w_j be the scaled and shifted function in (4.17) for some sufficiently large R>0 and let w_j^λ be the Kelvin transformation of w_j in (4.19). Define $\varphi_\lambda(y)=w_j(y)-w_j^\lambda(y)$ and $\Phi_\lambda(y)$ as in (4.37) with $\lambda\in[R-2,R+2]$, then $\varphi_\lambda+\Phi_\lambda$ satisfies the integral inequality (4.39). More importantly, Φ_λ is negative and satisfies the estimates in Lemma 4.3 under the current assumption that $\nabla K(0)=(1,0,\ldots,0)$. By Lemmas 4.1 and 4.3, for $\lambda=\lambda_0=R-2$ we have

$$\varphi_{\lambda} + \Phi_{\lambda} \ge 0 \quad \text{in } \Pi_{j} \backslash \overline{B}_{\lambda},$$

$$\tag{4.1}$$

and for $\lambda_1 = R + 2$ we have

$$\varphi_{\lambda_1}(y^*) + \Phi_{\lambda_1}(y^*) < 0 \quad \text{ for some } y^* \in B_{2\lambda_1} \backslash \overline{B}_{\lambda_1}.$$
 (4.2)

Thus we can start moving the sphere continuously for $\varphi_{\lambda} + \Phi_{\lambda}$ from $\lambda = \lambda_0$ as long as (4.1) holds, and the sphere must stop at some $\bar{\lambda} \in [\lambda_0, \lambda_1)$. Furthermore, with the help of the negativity of Φ_{λ} in Lemma 4.3 and of the lower bounds of the remainder term J_{λ} in Lemma 4.4, the moving sphere procedure may continue beyond $\bar{\lambda}$ where we reach a contradiction.

In Step 3, we complete the proof of Theorem 1.6 under the assumption of (K1). By Step 2 we know that the origin 0 is a critical point of K. Without loss of generality we may assume that $\lim_{j\to\infty} |\nabla K(x_j)|^{-1} \nabla K(x_j) = (1,0,\ldots,0)$. We will follow the notation in Step 2. As in Step 2, the main idea is still to use the moving sphere method for $\varphi_{\lambda} + \Phi_{\lambda}$, but under the current assumption of K, the function Φ_{λ} may not be negative. However, we can rewrite Φ_{λ} as $\Phi_{1,\lambda} + \Phi_{2,\lambda}$ where $\Phi_{1,\lambda}$ is negative and $\Phi_{2,\lambda}$ has a good estimate (see Lemmas 4.5 and 4.6). Based on the estimates of $\Phi_{2,\lambda}$, we consider two cases for $\alpha > 1$ separately.

(1) $\alpha < 2\sigma$ or $\alpha \ge (n-2\sigma)/2$. In this case, by choosing a sufficiently small $\varepsilon > 0$ we construct the following function

$$H_{\lambda}(y) = -\varepsilon M_j^{-1}(\lambda^{2\sigma-n} - |y|^{2\sigma-n}) + \Phi_{\lambda}(y), \quad y \in \Pi_j \setminus \overline{B}_{\lambda}$$

which can be negative. Using the lower estimates of the remainder term J_{λ} in Lemma 4.4, the method of moving spheres can be applied to $\varphi_{\lambda} + H_{\lambda}$ to reach a contradiction.

(2) $2\sigma \leq \alpha < (n-2\sigma)/2$. In this case, $\Phi_{2,\lambda}$ on $\Pi_j \backslash B_{2l_j}$ is too large to construct H_{λ} as in the first case, where $l_j := u(x_j)^{\frac{2}{n-2\sigma}}|x_j| \to \infty$. To deal with this difficulty, a different idea is needed. Note that Φ_{λ} is negative on $B_{2l_j} \backslash \overline{B}_{\lambda}$ and could be non-negative on $\Pi_j \backslash B_{2l_j}$, thus when moving the sphere, the trouble is

$$\int_{\mathcal{O}_{\lambda}} G(0,\lambda;y,z) b_{\lambda}(z) \varphi_{\lambda}(z) dz, \tag{4.3}$$

where $\mathcal{O}_{\lambda} = \{y \in \Pi_{j} \backslash B_{2l_{j}} : w_{j}(y) < w_{j}^{\lambda}(y)\}$. To control this integral, we add a function T_{λ} defined in (4.89) to both sides of the integral inequality (4.39). Let $H_{\lambda} := \Phi_{1,\lambda} + T_{\lambda}$, then $\varphi_{\lambda} + H_{\lambda}$ satisfies the integral inequality (4.95). Now when the method of moving spheres is applied to $\varphi_{\lambda} + H_{\lambda}$ (starting from $\lambda_{0} = R - 2$), the troublesome integral (4.3) can be controlled by

$$M_j^{-\frac{2\alpha}{n-2\sigma}} \int_{\Omega_j \setminus B_{2l_j}} G(0,\lambda;y,z) |z|^{\alpha-2\sigma-n} dz$$

with the help of Lemmas 4.5 and 4.6. This will lead to a contradiction.

Now we return to the detailed proof of Theorem 1.6 under the assumption (K1).

Step 1. We show that x_j can be chosen as the local maximum points of u. Moreover, the functions $u(x_j)^{-1}u(x_j+u(x_j)^{-\frac{2}{n-2\sigma}}y)$ converge in $C^2_{loc}(\mathbb{R}^n)$, after passing a subsequence, to a positive function $U_0 \in C^2(\mathbb{R}^n)$ where U_0 satisfies

$$\begin{cases} U_0(y) = \int_{\mathbb{R}^n} \frac{K(0)U_0(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz & \text{for } y \in \mathbb{R}^n, \\ \max_{\mathbb{R}^n} U_0 = U_0(0) = 1. \end{cases}$$
(4.4)

Define

$$s_j(x) := \left(\frac{d_j}{2} - |x - x_j|\right)^{\frac{n-2\sigma}{2}} u(x), \quad |x - x_j| \le \frac{d_j}{2}.$$

Since u is positive and continuous in $\overline{B}_{d_j/2}(x_j)$, we can find a point $\bar{x}_j \in \overline{B}_{d_j/2}(x_j)$ such that

$$s_j(\bar{x}_j) = \max_{|x-x_j| \le \frac{d_j}{2}} s_j(x) > 0.$$

Let $2\mu_j := \frac{d_j}{2} - |\bar{x}_j - x_j|$. Then

$$0 < 2\mu_j \leq \frac{d_j}{2} \quad \text{ and } \quad \frac{d_j}{2} - |x - x_j| \geq \mu_j \quad \forall \ |x - \bar{x}_j| \leq \mu_j.$$

By the definition of s_i , we have

$$(2\mu_j)^{\frac{n-2\sigma}{2}}u(\bar{x}_j) = s_j(\bar{x}_j) \ge s_j(x) \ge \mu_j^{\frac{n-2\sigma}{2}}u(x) \quad \forall |x - \bar{x}_j| \le \mu_j. \tag{4.5}$$

Hence

$$2^{\frac{n-2\sigma}{2}}u(\bar{x}_j) \ge u(x) \qquad \forall |x - \bar{x}_j| \le \mu_j. \tag{4.6}$$

We also have

$$(2\mu_j)^{\frac{n-2\sigma}{2}}u(\bar{x}_j) = s_j(\bar{x}_j) \ge s_j(x_j) = \left(\frac{d_j}{2}\right)^{\frac{n-2\sigma}{2}}u(x_j) \to \infty \quad \text{as } j \to \infty. \tag{4.7}$$

Now, we consider

$$v_{j}(y) = \frac{1}{u(\bar{x}_{j})} u\left(\bar{x}_{j} + \frac{y}{u(\bar{x}_{j})^{\frac{2}{n-2\sigma}}}\right), \ f_{j}(y) = \frac{1}{u(\bar{x}_{j})} h\left(\bar{x}_{j} + \frac{y}{u(\bar{x}_{j})^{\frac{2}{n-2\sigma}}}\right) \text{ in } \Theta_{j},$$

where

$$\Theta_j = \left\{ y \in \mathbb{R}^n : \bar{x}_j + \frac{y}{u(\bar{x}_j)^{\frac{2}{n-2\sigma}}} \in B_2 \backslash \Sigma \right\}.$$

We extend v_j to be identically 0 outside Θ_j and K to be identically 0 outside B_2 . Then v_j satisfies $v_j(0) = 1$ and

$$v_{j}(y) = \int_{\mathbb{R}^{n}} \frac{K(\bar{x}_{j} + u(\bar{x}_{j})^{-\frac{2}{n-2\sigma}} z) v_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y - z|^{n-2\sigma}} dz + f_{j}(y) \quad \text{for } y \in \Theta_{j}.$$
 (4.8)

Moreover, it follows from (4.6) and (4.7) that

$$v_j(y) \le 2^{\frac{n-2\sigma}{2}} \quad \text{in } B_{\bar{R}_j},\tag{4.9}$$

where

$$\bar{R}_j := \mu_j u(\bar{x}_j)^{\frac{2}{n-2\sigma}} \to \infty \quad \text{as } j \to \infty.$$

Clearly $u(\bar{x}_j) \to \infty$ as $j \to \infty$, and $\|f_j\|_{C^1(B_{\bar{R}_j})} \to 0$ as $j \to \infty$ since

$$||f_{j}||_{C^{1}(B_{\bar{R}_{j}})} \leq \frac{1}{u(\bar{x}_{j})} ||h||_{L^{\infty}(B_{\mu_{j}}(\bar{x}_{j}))} + \frac{1}{u(\bar{x}_{j})^{\frac{n-2\sigma+2}{n-2\sigma}}} ||\nabla h||_{L^{\infty}(B_{\mu_{j}}(\bar{x}_{j}))}$$

$$\leq \frac{1}{u(\bar{x}_{j})} ||h||_{L^{\infty}(B_{3/2})} + \frac{1}{u(\bar{x}_{j})^{\frac{n-2\sigma+2}{n-2\sigma}}} ||\nabla h||_{L^{\infty}(B_{3/2})} \to 0 \text{ as } j \to \infty.$$

$$(4.10)$$

Claim 1: There exists a subsequence of v_j , still denoted by v_j , such that v_j in $C^2_{loc}(\mathbb{R}^n)$ converges to a positive function $v \in C^2(\mathbb{R}^n)$ where v satisfies

$$v(y) = \int_{\mathbb{R}^n} \frac{K(0)v(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz \quad \text{for } y \in \mathbb{R}^n.$$
 (4.11)

Since for any R>0 we have $v_j(y)\leq 2^{\frac{n-2\sigma}{2}}$ in B_R for all large j, by regularity results in [29, Section 2.1] there exists $v\geq 0$ such that, up to a subsequence,

$$v_j \to v \quad \text{in } C^2_{loc}(\mathbb{R}^n).$$

Clearly v(0) = 1. To show that v satisfies the integral equation (4.11), we will use some arguments of [29, Proposition 2.9]. Write (4.8) as

$$v_{j}(y) = \int_{B_{r}} \frac{K(\bar{x}_{j} + u(\bar{x}_{j})^{-\frac{2}{n-2\sigma}}z)v_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y - z|^{n-2\sigma}} dz + F_{j}(r, y) \quad \text{for } y \in \Theta_{j},$$
(4.12)

where

$$F_{j}(r,y) = \int_{B^{c}} \frac{K(\bar{x}_{j} + u(\bar{x}_{j})^{-\frac{2}{n-2\sigma}}z)v_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y - z|^{n-2\sigma}} dz + f_{j}(y).$$

Then, for $y \in B_{r/2}$ we have

$$F_{j}(r,y) = \int_{B_{r}^{c}} \frac{K(\bar{x}_{j} + u(\bar{x}_{j})^{-\frac{2}{n-2\sigma}}z)v_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|z|^{n-2\sigma}} \frac{|z|^{n-2\sigma}}{|y-z|^{n-2\sigma}}dz + f_{j}(y)$$

$$\leq C \int_{B_{r}^{c}} \frac{K(\bar{x}_{j} + u(\bar{x}_{j})^{-\frac{2}{n-2\sigma}}z)v_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|z|^{n-2\sigma}}dz + ||f_{j}||_{L^{\infty}(B_{r/2})}$$

$$\leq Cv_{j}(0) + ||f_{j}||_{L^{\infty}(B_{r/2})}$$

for all large j. Similarly, for $y \in B_{r/2}$,

$$|\nabla_y F_j(r,y)| \le C(r)v_j(0) + \|\nabla f_j\|_{L^{\infty}(B_{r/2})}.$$

These together with (4.9) and (4.10) imply that $||F_j(r,\cdot)||_{C^1(B_{r/2})} \leq C(r)$ for all j large. Thus, after passing to a subsequence, $F_j(r,\cdot) \to F(r,\cdot)$ in $C^{1/2}(B_{r/2})$. Hence, letting $j \to \infty$ in (4.12) we obtain

$$F(r,y) = v(y) - \int_{B_r} \frac{K(0)v(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz \quad \text{for } y \in B_{r/2}.$$
 (4.13)

Furthermore, $F(r, y) \ge 0$ and it is non-increasing in r. For r >> |y|,

$$\frac{r^{n-2\sigma}}{(r+|y|)^{n-2\sigma}} (F_j(r,0) - f_j(0)) \leq F_j(r,y) - f_j(y)
= \int_{B_r^c} \frac{K(\bar{x}_j + u(\bar{x}_j)^{-\frac{2}{n-2\sigma}} z) v_j(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y|^{n-2\sigma}} \frac{|y|^{n-2\sigma}}{|y-z|^{n-2\sigma}} dz
\leq \frac{r^{n-2\sigma}}{(r-|y|)^{n-2\sigma}} (F_j(r,0) - f_j(0)).$$

Let j tend to ∞ , we get

$$\frac{r^{n-2\sigma}}{(r+|y|)^{n-2\sigma}}F(r,0) \le F(r,y) \le \frac{r^{n-2\sigma}}{(r-|y|)^{n-2\sigma}}F(r,0),$$

which leads to $\lim_{r\to\infty} F(r,y) = \lim_{r\to\infty} F(r,0) =: C_0 \ge 0$. Sending $r\to +\infty$ in (4.13) and using Lebesgue's monotone convergence theorem, we obtain

$$v(y) = \int_{\mathbb{R}^n} \frac{K(0)v(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz + C_0 \quad \text{for } y \in \mathbb{R}^n.$$

If $C_0 > 0$, then $v(y) \ge C_0 > 0$ for any $y \in \mathbb{R}^n$ and hence

$$1 = v(0) \ge \int_{\mathbb{R}^n} \frac{K(0)C_0^{\frac{n+2\sigma}{n-2\sigma}}}{|z|^{n-2\sigma}} dz = +\infty.$$

This is impossible. Therefore, $C_0 = 0$ and Claim 1 is established.

Since v(0) = 1, by the classification results in [14] or [35], v must be of the form

$$v(y) = \left(\frac{a_0}{1 + a_0^2 |y - y_0|^2}\right)^{\frac{n - 2\sigma}{2}} \tag{4.14}$$

for some $y_0 \in \mathbb{R}^n$ and some $a_0 \ge 1$. Obviously, v has an absolute maximum at y_0 . It implies that $v_j(y)$ must have a local maximum at a point y_j near y_0 when j is large. Replacing \bar{x}_j by

$$\tilde{x}_j := \bar{x}_j + u(\bar{x}_j)^{-\frac{2}{n-2\sigma}} y_j,$$

then $\{\tilde{x}_i\}$ are local maximum points of u for large j. Moreover, by (4.7) we have for large j that

$$u(\tilde{x}_j) = v_j(y_j)u(\bar{x}_j) \ge \frac{a_0^{(n-2\sigma)/2}}{2}u(\bar{x}_j) \ge \frac{a_0^{(n-2\sigma)/2}}{2}u(x_j)$$
(4.15)

and $\tilde{x}_j \in B_{\mu_j/2}(\bar{x}_j) \subset B_{d_j/2}(x_j)$ which implies $\frac{1}{2}d_j \leq \operatorname{dist}(\tilde{x}_j, \Sigma) \leq \frac{3}{2}d_j$ and $B_{\mu_j/2}(\tilde{x}_j) \subset B_{\mu_j}(\bar{x}_j)$. Consequently,

$$\operatorname{dist}(\tilde{x}_j, \Sigma)^{\frac{n-2\sigma}{2}} u(\tilde{x}_j) \to \infty \text{ and } \tilde{x}_j \to 0 \quad \text{as } j \to \infty.$$

Furthermore, by (4.6), (4.15) and (4.7) we have

$$u(\tilde{x}_j) \ge \frac{1}{2} \left(\frac{a_0}{2}\right)^{\frac{n-2\sigma}{2}} u(x) \quad \forall |x - \tilde{x}_j| \le \frac{\mu_j}{2}$$

and

$$R_j := \frac{1}{2} \mu_j u(\tilde{x}_j)^{\frac{2}{n-2\sigma}} \to \infty \quad \text{as } j \to \infty.$$

Using the proof of Claim 1 we know that the functions $u(\tilde{x}_j)^{-1}u(\tilde{x}_j+u(\tilde{x}_j)^{-\frac{2}{n-2\sigma}}y)$ converge in $C^2_{loc}(\mathbb{R}^n)$, after passing a subsequence, to a positive function $U_0 \in C^2(\mathbb{R}^n)$ which satisfies (4.4). It follows from the classification results in [14] or [35] that, modulo a positive constant,

$$U_0(y) = \left(\frac{1}{1+|y|^2}\right)^{\frac{n-2\sigma}{2}}. (4.16)$$

Thus the conclusion of Step 1 is proved. From now on, we consider x_i as \tilde{x}_i .

Step 2. We show $\nabla K(0) = 0$ assuming $K \in C^1(B_2)$ and $n > 2\sigma$.

Suppose $\nabla K(0) \neq 0$. We may assume without loss of generality that

$$\nabla K(0) = e = (1, 0, \dots, 0).$$

Let

$$w_{j}(y) = M_{j}^{-1}u\left(x_{j} + M_{j}^{-\frac{2}{n-2\sigma}}(y - Re)\right), \ h_{j}(y) = M_{j}^{-1}h\left(x_{j} + M_{j}^{-\frac{2}{n-2\sigma}}(y - Re)\right) \text{ in } \Omega_{j},$$

$$(4.17)$$

where $M_j = u(x_j)$, R > 10 is a large positive constant to be determined later, and

$$\Omega_j = \left\{ y \in \mathbb{R}^n : x_j + M_j^{-\frac{2}{n-2\sigma}} (y - Re) \in B_2 \backslash \Sigma \right\}. \tag{4.18}$$

By Step 1, w_j converges in C^2 norm to the bubble $U_1 = U_0(\cdot - Re)$ on every compact subset of \mathbb{R}^n . We also extend w_j to be identically 0 outside Ω_j and K to be identically 0 outside B_2 . For $\lambda > 0$, let

$$w_j^{\lambda}(y) = \left(\frac{\lambda}{|y|}\right)^{n-2\sigma} w_j \left(\frac{\lambda^2 y}{|y|^2}\right), \quad h_j^{\lambda}(y) = \left(\frac{\lambda}{|y|}\right)^{n-2\sigma} h_j \left(\frac{\lambda^2 y}{|y|^2}\right) \tag{4.19}$$

and let

$$K_j(y) = K\left(x_j + M_j^{-\frac{2}{n-2\sigma}}(y - Re)\right).$$
 (4.20)

Then

$$w_{j}(y) = \int_{\mathbb{R}^{n}} \frac{K_{j}(z)w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz + h_{j}(y) \quad \text{for } y \in \Omega_{j}.$$
 (4.21)

By (3.7), w_i^{λ} satisfies

$$w_j^{\lambda}(y) = \int_{\mathbb{R}^n} \frac{K_j(z^{\lambda}) w_j^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz + h_j^{\lambda}(y) \quad \text{for } y \in \Omega_j \backslash \overline{B}_{\lambda}, \tag{4.22}$$

where $z^{\lambda} = \frac{\lambda^2 z}{|z|^2}$ is the inversion of z with respect to ∂B_{λ} .

The following lemma gives the estimates on the difference between w_j and w_j^{λ} with $\lambda = R-2$ or R+2.

Lemma 4.1. Let $\lambda_0 = R - 2$ and $\lambda_1 = R + 2$. Then there exist $\varepsilon_0 = \varepsilon_0(n, \sigma, \min_{B_2 \setminus \Sigma} u, K, R) > 0$ and $j_0 = j_0(n, \sigma, \min_{B_2 \setminus \Sigma} u, K, R) > 1$ such that for all $j \geq j_0$,

$$w_j(y) - w_j^{\lambda_0}(y) \ge \varepsilon_0(|y| - \lambda_0)|y|^{2\sigma - 1 - n} + \varepsilon_0 M_j^{-1}(\lambda_0^{2\sigma - n} - |y|^{2\sigma - n}), \quad y \in \Omega_j \setminus \overline{B}_{\lambda_0}.$$
 (4.23)

Moreover, there exists $y^* \in B_{2\lambda_1} \setminus \overline{B}_{\lambda_1}$ such that for all $j \geq j_0$,

$$w_j(y^*) - w_j^{\lambda_1}(y^*) \le -\varepsilon_0. \tag{4.24}$$

Proof. Since w_j converges in C^2 norm to $U_1 = U_0(\cdot - Re)$ on any compact subset of \mathbb{R}^n , by Lemma 3.3 there exists $\varepsilon_1 = \varepsilon_1(n, \sigma, R) > 0$ such that for any fixed $R_1 \gg R$,

$$w_j(y) - w_j^{\lambda_0}(y) \ge \varepsilon_1(|y| - \lambda_0)|y|^{2\sigma - 1 - n}, \quad \lambda_0 < |y| \le R_1$$
 (4.25)

for sufficiently large j. Clearly, Lemma 3.3 also implies that (4.24) is true.

Next we show that (4.23) holds for $y \in \Omega_j \backslash B_{R_1}$. Firstly, it is easy to see that there exists a small $\varepsilon_2 = \varepsilon_2(n, \sigma, R) > 0$ such that for any $R_1 \gg R$,

$$|U_1(y) - |y|^{2\sigma - n}| \le \frac{\varepsilon_2}{2}|y|^{2\sigma - n}, \quad |y| \ge R_1$$
 (4.26)

and

$$U_1^{\lambda_0}(y) \le (1 - 3\varepsilon_2)|y|^{2\sigma - n}, \quad |y| \ge R_1.$$
 (4.27)

Consequently, we obtain that for large j,

$$w_i^{\lambda_0}(y) \le (1 - 2\varepsilon_2)|y|^{2\sigma - n}, \quad |y| \ge R_1.$$
 (4.28)

On the other hand, because w_j converges to U_1 in $C^2(B_{R_1})$, for any $y \in \Omega_j \backslash B_{R_1}$ we have that when j is sufficiently large,

$$\left(1 - \frac{\varepsilon_2}{8}\right) w_j(y) \ge \left(1 - \frac{\varepsilon_2}{8}\right) \int_{B_{R_1}} \frac{K_j(z) w_j(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y - z|^{n-2\sigma}} dz$$

$$\ge \left(1 - \frac{\varepsilon_2}{4}\right) \int_{B_{R_1}} \frac{K(0) w_j(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y - z|^{n-2\sigma}} dz$$

$$\ge \left(1 - \frac{\varepsilon_2}{2}\right) \int_{B_{R_1}} \frac{K(0) U_1(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y - z|^{n-2\sigma}} dz$$

$$\ge \left(1 - \frac{\varepsilon_2}{2}\right) U_1(y) - \int_{\{|z| \ge R_1\}} \frac{K(0) U_1(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y - z|^{n-2\sigma}} dz.$$
(4.29)

A simple computation yields that for $y \in \Omega_j \backslash B_{R_1}$

$$\int_{\{|z| \ge R_1\}} \frac{U_1(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz \le \left(\int_{\{|z| \ge R_1, |y-z| \ge \frac{|y|}{2}\}} + \int_{\{|z| \ge R_1, |y-z| \le \frac{|y|}{2}\}} \right) \frac{U_1(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz \\
\le \frac{C}{|y|^{n-2\sigma}} \int_{\{|z| \ge R_1\}} \frac{dz}{(1+|z|^2)^{(n+2\sigma)/2}} + \frac{C}{|y|^n} \\
\le \frac{C}{|y|^{n-2\sigma}} \cdot \frac{1}{R_1^{2\sigma}}$$

for some positive constant $C = C(n, \sigma, R)$. This, together with (4.26) and (4.29), gives

$$w_{j}(y) \geq \frac{\varepsilon_{2}}{8}w_{j}(y) + (1 - \varepsilon_{2})|y|^{2\sigma - n} + \frac{\varepsilon_{2}}{4}|y|^{2\sigma - n} - \frac{C}{|y|^{n - 2\sigma}} \cdot \frac{1}{R_{1}^{2\sigma}}$$

$$\geq \frac{\varepsilon_{2}}{8}w_{j}(y) + (1 - \varepsilon_{2})|y|^{2\sigma - n} \quad \text{for } y \in \Omega_{j} \backslash B_{R_{1}}$$

$$(4.30)$$

by choosing $R_1 \gg R$ large enough and then fixing it. Moreover, from the equation (1.12) we have $u(x) \geq 4^{2\sigma-n} \int_{B_2} K(y) u(y)^{\frac{n+2\sigma}{n-2\sigma}} dy =: A_0 > 0$ for all $x \in B_2 \backslash \Sigma$, so by the definition of w_j we obtain

$$w_j(y) \ge \varepsilon_3 M_j^{-1} (\lambda_0^{2\sigma - n} - |y|^{2\sigma - n}) \quad \text{for } y \in \Omega_j \backslash B_{R_1}.$$
 (4.31)

with some constant $\varepsilon_3 = \varepsilon_3(n, \sigma, A_0, R)$. It follows from (4.28), (4.30) and (4.31) that there exists a small $\varepsilon_0 > 0$ such that for large j,

$$w_j(y) - w_j^{\lambda_0}(y) \ge \varepsilon_0 |y|^{2\sigma - n} + \varepsilon_0 M_j^{-1} (\lambda_0^{2\sigma - n} - |y|^{2\sigma - n}) \quad \text{ for } y \in \Omega_j \backslash B_{R_1}.$$

This together with (4.25) implies that (4.23) holds by choosing ε_0 sufficiently small. Lemma 4.1 is established.

Next we will use the moving sphere method for $\lambda \in [R-2,R+2]$ to derive a contradiction. Denote

$$\Pi_j := \left\{ y \in \mathbb{R}^n : x_j + M_j^{-\frac{2}{n-2\sigma}} (y - Re) \in B_1 \backslash \Sigma \right\} \subset \Omega_j$$
(4.32)

and

$$\Sigma_j := \left\{ y \in \mathbb{R}^n : x_j + M_j^{-\frac{2}{n-2\sigma}} (y - Re) \in \Sigma \right\}. \tag{4.33}$$

Note that we have $\mathcal{L}^n(\Sigma_j) = 0$ due to $\mathcal{L}^n(\Sigma) = 0$. It follows from the same arguments as in Lemma 3.1 of [30] that for all large j, there holds

$$h_j^{\lambda}(y) \le h_j(y) \quad \forall y \in \Pi_j \backslash \overline{B}_{\lambda}, \ \lambda \in [R-2, R+2].$$
 (4.34)

Let $\varphi_{\lambda}(y) = w_j(y) - w_j^{\lambda}(y)$, where we omit j in the notation for brevity. By (3.8), (4.34) and $K_j \equiv 0$ on $\Omega_j^c \setminus \Sigma_j$, we have for $\lambda \in [R-2, R+2]$ and $y \in \Pi_j \setminus \overline{B}_{\lambda}$ that

$$\varphi_{\lambda}(y) \geq \int_{B_{\lambda}^{c}} G(0,\lambda;y,z) \left(K_{j}(z) w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - K_{j}(z^{\lambda}) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz$$

$$= \int_{B_{\lambda}^{c}} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz$$

$$+ \int_{B_{\lambda}^{c}} G(0,\lambda;y,z) \left(K_{j}(z) - K_{j}(z^{\lambda}) \right) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} dz$$

$$= \int_{\Pi_{j} \backslash B_{\lambda}} G(0,\lambda;y,z) b_{\lambda}(z) \varphi_{\lambda}(z) dz + J_{\lambda}(y) - \Phi_{\lambda}(y), \tag{4.35}$$

where

$$b_{\lambda}(y) = K_{j}(y) \frac{w_{j}(y)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(y)^{\frac{n+2\sigma}{n-2\sigma}}}{w_{j}(y) - w_{j}^{\lambda}(y)},$$
(4.36)

$$\Phi_{\lambda}(y) = \int_{\Omega_{j} \backslash B_{\lambda}} G(0, \lambda; y, z) \left(K_{j}(z^{\lambda}) - K_{j}(z) \right) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} dz$$
(4.37)

and

$$J_{\lambda}(y) = \int_{\Omega_{j}\backslash\Pi_{j}} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz$$

$$- \int_{\Omega_{j}^{c}} G(0,\lambda;y,z) K_{j}(z^{\lambda}) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} dz.$$

$$(4.38)$$

Note that $b_{\lambda}(y)$ is always non-negative. Thus, φ_{λ} satisfies the integral inequality

$$\varphi_{\lambda}(y) + \Phi_{\lambda}(y) \ge \int_{\Pi_{j} \backslash B_{\lambda}} G(0, \lambda; y, z) b_{\lambda}(z) \varphi_{\lambda}(z) dz + J_{\lambda}(y), \quad y \in \Pi_{j} \backslash \overline{B}_{\lambda}, \tag{4.39}$$

where $\lambda \in [R-2, R+2]$.

Now we show some estimates for Φ_{λ} on $\Pi_{j} \backslash \overline{B}_{\lambda}$ in order to apply the moving sphere method to the function $\varphi_{\lambda} + \Phi_{\lambda}$. We first need to establish the estimates on $K_{j}(z^{\lambda}) - K_{j}(z)$ and on $w_{j}^{\lambda}(z)$ for $z \in \Omega_{j} \backslash B_{\lambda}$. For this purpose we define a special domain in $\Omega_{j} \backslash B_{\lambda}$. Let

$$D_{\lambda} = \{z \in \mathbb{R}^n : \lambda < |z| < 2\lambda, z_1 > 2|z'| \text{ where } z' = (z_2, \dots, z_n)\}.$$

Claim 2: There exists a constant C > 0 independent of λ such that for all large j,

$$K_j(z^{\lambda}) - K_j(z) \le -CM_j^{-\frac{2}{n-2\sigma}}(|z| - \lambda), \quad z \in D_{\lambda}, \tag{4.40}$$

$$|K_j(z^{\lambda}) - K_j(z)| \le CM_j^{-\frac{2}{n-2\sigma}}(|z| - \lambda), \quad z \in \Omega_j \backslash B_{\lambda}.$$
(4.41)

Proof. The first estimate follows from the mean value theorem and $\lim_{x\to 0} \nabla K(x) = \nabla K(0) = e$. The second estimate follows from the mean value theorem.

Lemma 4.2. There exist constants $C_1, C_2 > 0$ independent of λ such that for all large j,

$$C_1|z|^{2\sigma-n} \le w_j^{\lambda}(z) \le C_2|z|^{2\sigma-n} \quad \text{for } z \in \Omega_j \setminus (B_{\lambda} \cup D_{\lambda})$$
 (4.42)

and

$$C_1\left(\frac{\lambda}{|z|}\right)^{n-2\sigma} \left(\frac{1}{1+(z_1-\lambda)^2+|z'|^2}\right)^{\frac{n-2\sigma}{2}} \le w_j^{\lambda}(z) \le 2 \quad \text{for } z \in D_{\lambda}. \tag{4.43}$$

Proof. Since the functions $w_j(y)$ converge in $C^2_{loc}(\mathbb{R}^n)$ to $U_1(y) = (1 + |y - Re|^2)^{-\frac{n-2\sigma}{2}}$, for $z \in D_\lambda$ and for large j we have

$$\frac{1}{2} \left(\frac{\lambda}{|z|} \right)^{n-2\sigma} \left(\frac{1}{1 + \left| \frac{\lambda^2 z_1}{|z|^2} - R \right|^2 + \left| \frac{\lambda^2 z'}{|z|^2} \right|^2} \right)^{\frac{n-2\sigma}{2}} \le \left(\frac{\lambda}{|z|} \right)^{n-2\sigma} w_j \left(\frac{\lambda^2 z}{|z|^2} \right) = w_j^{\lambda}(z) \le 2.$$
(4.44)

Note that $|\lambda - R| \le 2$ and for $|z| > \lambda$,

$$\left| \frac{\lambda^2 z_1}{|z|^2} - \lambda \right|^2 = \frac{\lambda^2}{|z|^2} \left(\frac{\lambda^2}{|z|^2} z_1^2 - 2z_1 \lambda + \frac{|z|^2}{\lambda^2} \lambda^2 \right)$$

$$\leq (z_1 - \lambda)^2 + (|z|^2 - \lambda^2) \left(1 - \frac{z_1^2}{|z|^2} \right)$$

$$\leq (z_1 - \lambda)^2 + |z'|^2,$$

from which we obtain

$$1 + \left| \frac{\lambda^2 z_1}{|z|^2} - R \right|^2 + \left| \frac{\lambda^2 z'}{|z|^2} \right|^2 \le 10(1 + (z_1 - \lambda)^2 + |z'|^2).$$

This together with (4.44) gives the proof of (4.43).

Next we establish (4.42). Firstly one can verify the following elementary estimate:

$$|z^{\lambda} - Re|^2 \ge C\lambda^2, \quad z \in \Omega_i \setminus (B_{\lambda} \cup D_{\lambda})$$
 (4.45)

for some constant C>0 independent of λ . Because $w_j(y)$ converges in $C^2_{loc}(\mathbb{R}^n)$ to $U_1(y)=(1+|y-Re|^2)^{-\frac{n-2\sigma}{2}}$, we have for large j that

$$\frac{1}{2} \left(\frac{\lambda}{|z|} \right)^{n-2\sigma} \left(\frac{1}{1 + |z^{\lambda} - Re|^{2}} \right)^{\frac{n-2\sigma}{2}} \le w_{j}^{\lambda}(z) \le 2 \left(\frac{\lambda}{|z|} \right)^{n-2\sigma} \left(\frac{1}{1 + |z^{\lambda} - Re|^{2}} \right)^{\frac{n-2\sigma}{2}}, \tag{4.46}$$

where $z \in \Omega_i \backslash B_\lambda$. By (4.45) we get

$$w_j^{\lambda}(z) \le 2\left(\frac{\lambda}{|z|}\right)^{n-2\sigma} \left(\frac{1}{1+C\lambda^2}\right)^{\frac{n-2\sigma}{2}} \le C_2|z|^{2\sigma-n}, \quad z \in \Omega_j \setminus (B_{\lambda} \cup D_{\lambda})$$

for another constant $C_2>0$ independent of λ . On the other hand, using $|z^{\lambda}-Re|\leq 3\lambda$ for $|z|\geq \lambda$ we obtain

$$w_j^{\lambda}(z) \ge \frac{1}{2} \left(\frac{\lambda}{|z|} \right)^{n-2\sigma} \left(\frac{1}{1+9\lambda^2} \right)^{\frac{n-2\sigma}{2}} \ge C_1 |z|^{2\sigma-n}, \quad z \in \Omega_j \setminus (B_{\lambda} \cup D_{\lambda})$$

for some constant $C_1 > 0$ independent of λ . Lemma 4.2 is proved.

Lemma 4.2 tells us that w_j^{λ} is bigger on D_{λ} and is smaller on $\Omega_j \setminus (B_{\lambda} \cup D_{\lambda})$, while $K_j(z^{\lambda}) - K_j(z)$ is negative on D_{λ} by Claim 2. These combined with Lemma 3.1 will lead to a crucial fact that Φ_{λ} is non-positive. Indeed, the positive parts of the integral in (4.37) can be controlled by the negative one. More precisely, we have

Lemma 4.3. There exists a constant C > 0 independent of λ such that for all large j,

$$\Phi_{\lambda}(y) \leq \begin{cases} -CM_{j}^{-\frac{2}{n-2\sigma}}(|y|-\lambda)\lambda^{-n}\log\lambda & \text{for } y \in B_{4\lambda}\backslash \overline{B}_{\lambda}, \\ -CM_{j}^{-\frac{2}{n-2\sigma}}|y|^{2\sigma-n}\lambda^{1-2\sigma}\log\lambda & \text{for } y \in \Pi_{j}\backslash \overline{B}_{4\lambda} \end{cases}$$

and

$$|\Phi_{\lambda}(y)| \leq \begin{cases} CM_{j}^{-\frac{2}{n-2\sigma}}(|y|-\lambda)\lambda^{2\sigma} & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ CM_{j}^{-\frac{2}{n-2\sigma}}|y|^{2\sigma-n}\lambda^{n+1} & \text{for } y \in \Pi_{j} \backslash \overline{B}_{4\lambda}. \end{cases}$$

Proof. Let

$$Q_{\lambda}(z) = \left(K_{j}(z^{\lambda}) - K_{j}(z)\right) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \quad \text{for } z \in \Omega_{j} \backslash B_{\lambda}.$$
(4.47)

Then Φ_{λ} can be written as

$$\Phi_{\lambda}(y) = \int_{\Omega_{j} \setminus B_{\lambda}} G(0, \lambda; y, z) Q_{\lambda}(z) dz.$$

By Claim 2 and Lemma 4.2 we have

$$Q_{\lambda}(z) \le -CM_{j}^{-\frac{2}{n-2\sigma}}(|z|-\lambda) \left(\frac{1}{1+(z_{1}-\lambda)^{2}+|z'|^{2}}\right)^{\frac{n+2\sigma}{2}} \quad \text{for } z \in D_{\lambda}$$
 (4.48)

and

$$|Q_{\lambda}(z)| \le \begin{cases} CM_{j}^{-\frac{2}{n-2\sigma}}(|z|-\lambda) & \text{for } z \in D_{\lambda}, \\ CM_{j}^{-\frac{2}{n-2\sigma}}(|z|-\lambda)|z|^{-2\sigma-n} & \text{for } z \in \Omega_{j} \setminus (B_{\lambda} \cup D_{\lambda}). \end{cases}$$
(4.49)

These lead to

$$\Phi_{\lambda}(y) \leq -CM_{j}^{-\frac{2}{n-2\sigma}} \int_{D_{\lambda}} G(0,\lambda;y,z) (|z|-\lambda) \left(\frac{1}{1+(z_{1}-\lambda)^{2}+|z'|^{2}}\right)^{\frac{n+2\sigma}{2}} dz \\
+ CM_{j}^{-\frac{2}{n-2\sigma}} \int_{\Omega_{j} \setminus (B_{\lambda} \cup D_{\lambda})} G(0,\lambda;y,z) (|z|-\lambda) |z|^{-2\sigma-n} dz.$$
(4.50)

We consider the following two cases separately.

Case 1: $\lambda < |y| \le 4\lambda$. When $z \in D_{\lambda}$ and $|y-z| < \frac{1}{3}(|y|-\lambda)$, by Lemma 3.1 we have

$$G(0,\lambda;y,z) \ge C|y-z|^{2\sigma-n} \ge C \frac{(|y|-\lambda)^2}{(|y|-\lambda)^{n-2\sigma+2}} \ge C \frac{(|y|-\lambda)(|z|-\lambda)}{\lambda^{n-2\sigma+2}}.$$

When $z \in D_{\lambda}$ and $|y - z| \ge \frac{1}{3}(|y| - \lambda)$, by Lemma 3.1 we also have

$$G(0,\lambda;y,z) \ge C \frac{(|y|-\lambda)(|z|^2-\lambda^2)}{\lambda|y-z|^{n-2\sigma+2}} \ge C \frac{(|y|-\lambda)(|z|-\lambda)}{\lambda^{n-2\sigma+2}}.$$

Thus, in either case we have

$$G(0,\lambda;y,z) \ge C \frac{(|y|-\lambda)(|z|-\lambda)}{\lambda^{n-2\sigma+2}}.$$
(4.51)

To estimate $G(0, \lambda; y, z)$ for $z \in \Omega_j \backslash B_{\lambda}$, we define three sets

$$\mathcal{A}_1 = \{ z \in \Omega_j \backslash B_\lambda : |z - y| < (|y| - \lambda)/3 \},$$

$$\mathcal{A}_2 = \{ z \in \Omega_j \backslash B_\lambda : |z - y| \ge (|y| - \lambda)/3 \text{ and } |z| \le 8\lambda \},$$

$$\mathcal{A}_3 = \{ z \in \Omega_j \backslash B_\lambda : |z| \ge 8\lambda \}.$$

Then, from Lemma 3.1 we have

$$G(0,\lambda;y,z) \leq \begin{cases} C|y-z|^{2\sigma-n} & \text{if } z \in \mathcal{A}_{1}, \\ C\frac{(|y|-\lambda)(|z|^{2}-\lambda^{2})}{\lambda|y-z|^{n-2\sigma+2}} \leq C\frac{(|y|-\lambda)(|z|-\lambda)}{|y-z|^{n-2\sigma+2}} & \text{if } z \in \mathcal{A}_{2}, \\ C\frac{(|y|-\lambda)(|z|^{2}-\lambda^{2})}{\lambda|y-z|^{n-2\sigma+2}} \leq C\frac{|y|-\lambda}{\lambda}|z|^{2\sigma-n} & \text{if } z \in \mathcal{A}_{3}. \end{cases}$$
(4.52)

Hence, combining (4.50), (4.51) and (4.52) we get

$$\Phi_{\lambda}(y) \leq -CM_{j}^{-\frac{2}{n-2\sigma}} \int_{D_{\lambda}} \frac{(|y|-\lambda)(|z|-\lambda)^{2}}{\lambda^{n-2\sigma+2}} \left(\frac{1}{1+(z_{1}-\lambda)^{2}+|z'|^{2}}\right)^{\frac{n+2\sigma}{2}} dz
+ CM_{j}^{-\frac{2}{n-2\sigma}} \int_{\mathcal{A}_{1}\backslash D_{\lambda}} |y-z|^{2\sigma-n} (|z|-\lambda)|z|^{-2\sigma-n} dz
+ CM_{j}^{-\frac{2}{n-2\sigma}} \int_{\mathcal{A}_{2}\backslash D_{\lambda}} \frac{(|y|-\lambda)(|z|-\lambda)^{2}}{|y-z|^{n-2\sigma+2}} |z|^{-2\sigma-n} dz
+ CM_{j}^{-\frac{2}{n-2\sigma}} \int_{\mathcal{A}_{3}} \frac{|y|-\lambda}{\lambda} |z|^{2\sigma-n} (|z|-\lambda)|z|^{-2\sigma-n} dz
=: I_{1} + I_{2} + I_{3} + I_{4}.$$
(4.53)

For I_1 we have

$$I_{1} \leq -CM_{j}^{-\frac{2}{n-2\sigma}} \frac{(|y|-\lambda)}{\lambda^{n-2\sigma+2}} \int_{D_{\lambda}} \frac{(|z|-\lambda)^{2}}{(1+(z_{1}-\lambda)^{2}+|z'|^{2})^{\frac{n+2\sigma}{2}}} dz$$

$$\leq -CM_{j}^{-\frac{2}{n-2\sigma}} \cdot \frac{(|y|-\lambda)}{\lambda^{n-2\sigma+2}} \cdot \frac{1}{\lambda^{2\sigma-2}} \int_{D_{\lambda}} \frac{(|z|-\lambda)^{2}}{(1+(z_{1}-\lambda)^{2}+|z'|^{2})^{\frac{n+2\sigma}{2}}} dz$$

$$\leq -CM_{j}^{-\frac{2}{n-2\sigma}} (|y|-\lambda) \frac{\log \lambda}{\lambda^{n}},$$
(4.54)

where we used the fact $\sigma \ge 1$ in the second inequality. Since $|z| - \lambda \le 4(|y| - \lambda)/3$ when $z \in \mathcal{A}_1$, for I_2 we have

$$I_{2} \leq CM_{j}^{-\frac{2}{n-2\sigma}} \int_{\mathcal{A}_{1}\backslash D_{\lambda}} |y-z|^{2\sigma-n} (|z|-\lambda)|z|^{-2\sigma-n} dz$$

$$\leq CM_{j}^{-\frac{2}{n-2\sigma}} (|y|-\lambda)\lambda^{-2\sigma-n} \int_{|z-y|\leq \lambda} |y-z|^{2\sigma-n} dz$$

$$\leq CM_{j}^{-\frac{2}{n-2\sigma}} (|y|-\lambda)\frac{1}{\lambda^{n}}.$$

$$(4.55)$$

Since $|z| - \lambda \le 4|z - y|$ when $z \in A_2$, for I_3 we have

$$I_{3} \leq CM_{j}^{-\frac{2}{n-2\sigma}}(|y|-\lambda) \int_{\mathcal{A}_{2}\backslash D_{\lambda}} \frac{1}{|y-z|^{n-2\sigma}} |z|^{-2\sigma-n} dz$$

$$\leq CM_{j}^{-\frac{2}{n-2\sigma}}(|y|-\lambda) \frac{1}{\sqrt{n}}.$$

$$(4.56)$$

For I_4 we have

$$I_4 \le CM_j^{-\frac{2}{n-2\sigma}} (|y| - \lambda) \frac{1}{\lambda} \int_{A_2} |z|^{1-2n} dz \le CM_j^{-\frac{2}{n-2\sigma}} (|y| - \lambda) \frac{1}{\lambda^n}. \tag{4.57}$$

Substituting these estimates into (4.53) and taking R sufficiently large, we obtain

$$\Phi_{\lambda}(y) \leq -CM_{j}^{-\frac{2}{n-2\sigma}}(|y|-\lambda)\frac{\log\lambda}{\lambda^{n}} \quad \text{ for } \lambda < |y| \leq 4\lambda.$$

Next we estimate $|\Phi_{\lambda}(y)|$. It is clear that we only need to give the estimation of

$$\widetilde{I}_1 := \int_{D_{\lambda}} G(0, \lambda; y, z) |Q_{\lambda}(z)| dz.$$

By (4.49) and (4.52) we get

$$\begin{split} \widetilde{I}_1 &\leq CM_j^{-\frac{2}{n-2\sigma}} \int_{D_{\lambda}} G(0,\lambda;y,z) (|z|-\lambda) dz \\ &\leq CM_j^{-\frac{2}{n-2\sigma}} \left(\int_{\mathcal{A}_1} |y-z|^{2\sigma-n} (|z|-\lambda) dz + \int_{\mathcal{A}_2} \frac{(|y|-\lambda)(|z|-\lambda)^2}{|y-z|^{n-2\sigma+2}} dz \right) \\ &\leq CM_j^{-\frac{2}{n-2\sigma}} (|y|-\lambda) \int_{|z-y| \leq 12\lambda} |y-z|^{2\sigma-n} dz \\ &\leq CM_j^{-\frac{2}{n-2\sigma}} (|y|-\lambda) \lambda^{2\sigma}, \end{split}$$

where we used the facts that $|z| - \lambda \le 4(|y| - \lambda)/3$ for $z \in \mathcal{A}_1$ and $|z| - \lambda \le 4|z - y|$ for $z \in \mathcal{A}_2$ in the third inequality. This, together with (4.55), (4.56) and (4.57), yields

$$|\Phi_{\lambda}(y)| \le CM_i^{-\frac{2}{n-2\sigma}}(|y|-\lambda)\lambda^{2\sigma} \quad \text{for } \lambda < |y| \le 4\lambda.$$

Case 2: $|y| > 4\lambda$. When $z \in D_{\lambda}$, we have $|y - z| > (|z| - \lambda)/3$. By the symmetry of $G(0, \lambda; y, z)$ and (2) of Lemma 3.1, we obtain

$$G(0,\lambda;y,z) = G(0,\lambda;z,y) \ge C \frac{(|z|-\lambda)(|y|^2-\lambda^2)}{\lambda|y-z|^{n-2\sigma+2}} \ge C \frac{|z|-\lambda}{\lambda}|y|^{2\sigma-n} \quad \text{for } z \in D_{\lambda}.$$

$$(4.58)$$

Moreover, we also have

$$G(0, \lambda; y, z) \le C|y - z|^{2\sigma - n} \quad \text{for } z \in \Omega_j \backslash B_{\lambda}.$$
 (4.59)

Therefore, by (4.50),

$$\Phi_{\lambda}(y) \leq -CM_{j}^{-\frac{2}{n-2\sigma}} \int_{D_{\lambda}} \frac{(|z|-\lambda)^{2}}{\lambda} |y|^{2\sigma-n} \left(\frac{1}{1+(z_{1}-\lambda)^{2}+|z'|^{2}}\right)^{\frac{n+2\sigma}{2}} dz
+CM_{j}^{-\frac{2}{n-2\sigma}} \int_{\Omega_{j}\setminus (B_{\lambda}\cup D_{\lambda})} |y-z|^{2\sigma-n} (|z|-\lambda)|z|^{-2\sigma-n} dz
=: II_{1} + II_{2}.$$
(4.60)

For II_1 we have

$$II_{1} \leq -CM_{j}^{-\frac{2}{n-2\sigma}}|y|^{2\sigma-n}\lambda^{-1} \int_{D_{\lambda}} \frac{(|z|-\lambda)^{2}}{(1+(z_{1}-\lambda)^{2}+|z'|^{2})^{\frac{n+2\sigma}{2}}} dz
\leq -CM_{j}^{-\frac{2}{n-2\sigma}}|y|^{2\sigma-n}\lambda^{1-2\sigma} \int_{D_{\lambda}} \frac{(|z|-\lambda)^{2}}{(1+(z_{1}-\lambda)^{2}+|z'|^{2})^{\frac{n+2}{2}}} dz
\leq -CM_{j}^{-\frac{2}{n-2\sigma}}|y|^{2\sigma-n} \frac{\log \lambda}{\lambda^{2\sigma-1}},$$

where we used the fact $\sigma \geq 1$ in the second inequality. For II_2 we have

$$\begin{split} II_2 &\leq CM_j^{-\frac{2}{n-2\sigma}} \int_{\Omega_j \setminus (B_\lambda \cup D_\lambda)} |y-z|^{2\sigma-n} |z|^{1-2\sigma-n} dz \\ &= CM_j^{-\frac{2}{n-2\sigma}} \sum_{k=1}^4 \int_{\mathcal{S}_k \setminus D_\lambda} |y-z|^{2\sigma-n} |z|^{1-2\sigma-n} dz, \end{split}$$

where

$$\mathcal{S}_1 = \{z \in \Omega_j \backslash B_\lambda : |z| < |y|/2\},$$

$$\mathcal{S}_2 = \{z \in \Omega_j \backslash B_\lambda : |z| > 2|y|\},$$

$$\mathcal{S}_3 = \{z \in \Omega_j \backslash B_\lambda : |z - y| \le |y|/2\},$$

$$\mathcal{S}_4 = \{z \in \Omega_j \backslash B_\lambda : |z - y| \ge |y|/2 \text{ and } |y|/2 \le |z| \le 2|y|\}.$$

Direct computations give

$$\int_{\mathcal{S}_1} |y-z|^{2\sigma-n} |z|^{1-2\sigma-n} dz \leq C |y|^{2\sigma-n} \int_{\mathcal{S}_1} |z|^{1-2\sigma-n} dz \leq C \frac{|y|^{2\sigma-n}}{\lambda^{2\sigma-1}} \quad \text{if } \sigma > \frac{1}{2},$$

$$\int_{\mathcal{S}_2} |y-z|^{2\sigma-n} |z|^{1-2\sigma-n} dz \leq C \int_{\mathcal{S}_2} |z|^{2\sigma-n} |z|^{1-2\sigma-n} dz \leq C |y|^{1-n} \leq C \frac{|y|^{2\sigma-n}}{\lambda^{2\sigma-1}} \quad \text{if } \sigma \geq \frac{1}{2},$$

$$\int_{\mathcal{S}_3} |y-z|^{2\sigma-n} |z|^{1-2\sigma-n} dz \leq C |y|^{1-2\sigma-n} \int_{\mathcal{S}_3} |y-z|^{2\sigma-n} dz \leq C |y|^{1-n} \leq C \frac{|y|^{2\sigma-n}}{\lambda^{2\sigma-1}} \quad \text{if } \sigma \geq \frac{1}{2},$$
 and

$$\int_{\mathcal{S}_4} |y-z|^{2\sigma-n} |z|^{1-2\sigma-n} dz \leq C|y|^{1-2\sigma-n} \int_{\mathcal{S}_4} |y-z|^{2\sigma-n} dz \leq C|y|^{1-n} \leq C \frac{|y|^{2\sigma-n}}{\lambda^{2\sigma-1}} \ \text{if} \ \sigma \geq \frac{1}{2}.$$

These estimates imply that

$$II_2 \le CM_j^{-\frac{2}{n-2\sigma}} |y|^{2\sigma-n} \frac{1}{\lambda^{2\sigma-1}}.$$
 (4.61)

Hence, by taking R sufficiently large we have

$$\Phi_{\lambda}(y) \le -CM_j^{-\frac{2}{n-2\sigma}} |y|^{2\sigma - n} \frac{\log \lambda}{\lambda^{2\sigma - 1}} \quad \text{for } |y| > 4\lambda.$$

To bound $|\Phi_{\lambda}(y)|$, we only need to give the estimation of

$$\widetilde{II}_1 := \int_{D_{\lambda}} G(0, \lambda; y, z) |Q_{\lambda}(z)| dz.$$

By (4.49) and (4.59) we get

$$\begin{split} \widetilde{II}_{1} &\leq CM_{j}^{-\frac{2}{n-2\sigma}} \int_{D_{\lambda}} |y-z|^{2\sigma-n} (|z|-\lambda) dz \\ &\leq CM_{j}^{-\frac{2}{n-2\sigma}} |y|^{2\sigma-n} \int_{D_{\lambda}} (|z|-\lambda) dz \leq CM_{j}^{-\frac{2}{n-2\sigma}} |y|^{2\sigma-n} \lambda^{n+1}, \end{split}$$

where we have used $|y-z| \ge |y|/2$ for $z \in D_{\lambda}$. Combining this with (4.61) we obtain

$$|\Phi_{\lambda}(y)| \le CM_j^{-\frac{2}{n-2\sigma}} |y|^{2\sigma-n} \lambda^{n+1} \quad \text{for } |y| > 4\lambda.$$

The proof of Lemma 4.3 is completed.

In the proof of Lemma 4.3, R is chosen so large that Φ_{λ} is negative for any $\lambda \in [R-2, R+2]$. Recall $\lambda_0 = R-2$ and $\lambda_1 = R+2$. By Lemmas 4.1 and 4.3 we know that for large j,

$$\varphi_{\lambda_0}(y) + \Phi_{\lambda_0}(y) \ge 0, \quad y \in \Pi_j \setminus \overline{B}_{\lambda_0}.$$

We define

$$\bar{\lambda} := \sup\{\mu \ge \lambda_0 \mid \varphi_{\lambda}(y) + \Phi_{\lambda}(y) \ge 0, \ \forall \ y \in \Pi_j \backslash \overline{B}_{\lambda}, \ \forall \ \lambda_0 \le \lambda \le \mu\}.$$

Then, $\bar{\lambda}$ is well defined and $\bar{\lambda} \geq \lambda_0$ for all large j. Furthermore, from Lemmas 4.1 and 4.3 we have that $\bar{\lambda} < \lambda_1$ for all large j. Using the continuity we obtain

$$\varphi_{\bar{\lambda}}(y) + \Phi_{\bar{\lambda}}(y) \ge 0, \quad y \in \Pi_i \backslash \overline{B}_{\bar{\lambda}}.$$

This together with Lemma 4.3 implies that

$$\varphi_{\bar{\lambda}}(y) \ge 0, \qquad y \in \Pi_j \backslash \overline{B}_{\bar{\lambda}}.$$
(4.62)

It follows from (4.39) that for $\bar{\lambda} \leq \lambda \leq \lambda_1$ and $y \in \Pi_j \backslash \overline{B}_{\lambda}$,

$$\varphi_{\lambda}(y) + \Phi_{\lambda}(y) \ge \int_{\Pi_{j} \backslash B_{\lambda}} G(0, \lambda; y, z) b_{\lambda}(z) \varphi_{\lambda}(z) dz + J_{\lambda}(y), \tag{4.63}$$

where J_{λ} is given by

$$J_{\lambda}(y) = \int_{\Omega_{j}\backslash\Pi_{j}} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz$$
$$- \int_{\Omega_{j}^{c}} G(0,\lambda;y,z) K_{j}(z^{\lambda}) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} dz.$$

Lemma 4.4. There exists a constant C > 0 depending on R such that for any $\lambda_0 \le \lambda \le \lambda_1$ and for all large j,

$$J_{\lambda}(y) \ge \begin{cases} C(|y| - \lambda)u(x_{j})^{-1}, & \text{if } \lambda \le |y| \le \lambda_{1} + 1, \\ Cu(x_{j})^{-1}, & \text{if } |y| > \lambda_{1} + 1, \ y \in \Pi_{j}. \end{cases}$$
(4.64)

Proof. For any $z \in \mathbb{R}^n \setminus (\Pi_j \cup \Sigma_j)$ and $\lambda_0 \leq \lambda \leq \lambda_1$, we have $|z| \geq \frac{1}{2}u(x_j)^{\frac{2}{n-2\sigma}}$ for large j and hence

$$w_j^{\lambda}(z) \le \left(\frac{\lambda}{|z|}\right)^{n-2\sigma} \max_{B_{\lambda_1}} w_j \le Cu(x_j)^{-2},$$

where C > 0 depends on R. On the other hand, by the equation (1.12) we have

$$u(x) \ge 4^{2\sigma - n} \int_{B_2} K(y) u(y)^{\frac{n+2\sigma}{n-2\sigma}} dy =: A_0 > 0 \quad \text{ for all } x \in B_2 \backslash \Sigma, \tag{4.65}$$

and by the definition of w_i , we obtain

$$w_j(y) \ge \frac{A_0}{u(x_j)}$$
 for $y \in \Omega_j \backslash \Pi_j$. (4.66)

Therefore, for large j we have

$$w_j(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_j^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \ge \frac{1}{2} w_j(z)^{\frac{n+2\sigma}{n-2\sigma}} \quad \text{in } \Omega_j \backslash \Pi_j.$$

Since $G(0, \lambda; y, z) = 0$ for $|y| = \lambda$ and

$$y \cdot \nabla_y G(0, \lambda; y, z) \Big|_{|y| = \lambda} = (n - 2\sigma)|y - z|^{2\sigma - n - 2} (|z|^2 - |y|^2) > 0$$

for $|z| \ge \lambda_1 + 2$, and by the positivity and smoothness of G we obtain

$$\frac{\delta_1}{|y-z|^{n-2\sigma}}(|y|-\lambda) \le G(0,\lambda;y,z) \le \frac{\delta_2}{|y-z|^{n-2\sigma}}(|y|-\lambda) \tag{4.67}$$

for $\lambda_0 \le \lambda \le |y| \le \lambda_1 + 1$ and $\lambda_1 + 2 \le |z| \le M < \infty$, where $0 < \delta_1 \le \delta_2 < \infty$. Moreover, if M is large, then

$$0 < c \leq y \cdot \nabla_y(|y-z|^{n-2\sigma}G(0,\lambda;y,z)) \leq C < \infty$$

for all $|z| \ge M$ and $\lambda_0 \le \lambda \le |y| \le \lambda_1 + 1$. Hence, (4.67) also holds for $\lambda_0 \le \lambda \le |y| \le \lambda_1 + 1$ and $|z| \ge M$. Moreover, by the definition of $G(0,\lambda;y,z)$, we can verify that for $|y| \ge \lambda_1 + 1$ and $|z| \ge \lambda_1 + 2$,

$$\frac{\delta_3}{|y-z|^{n-2\sigma}} \le G(0,\lambda;y,z) \le \frac{1}{|y-z|^{n-2\sigma}}$$
(4.68)

for some $\delta_3 \in (0,1)$.

Denote $\rho_j := u(x_j)^{\frac{2}{n-2\sigma}}$ and $A_1 := \min_{B_2} K > 0$. Then, for $\lambda \le |y| \le \lambda_1 + 1$ and for large j, we have

$$J_{\lambda}(y) \geq \frac{A_1}{2} \left(\frac{A_0}{u(x_j)}\right)^{\frac{n+2\sigma}{n-2\sigma}} \int_{\Omega_j \backslash \Pi_j} \frac{\delta_1}{|y-z|^{n-2\sigma}} (|y|-\lambda) dz$$

$$-C \int_{\Omega_j^c} \frac{\delta_2}{|y-z|^{n-2\sigma}} (|y|-\lambda) \left(\frac{\lambda}{|z|}\right)^{n+2\sigma} dz$$

$$\geq C(|y|-\lambda) u(x_j)^{-\frac{n+2\sigma}{n-2\sigma}} \int_{\left\{\frac{5}{4}\rho_j \leq |z| \leq \frac{7}{4}\rho_j\right\} \backslash \Sigma_j} \frac{1}{|y-z|^{n-2\sigma}} dz$$

$$-C(|y|-\lambda) \int_{\left\{|z| \geq \frac{7}{4}\rho_j\right\} \cup \Sigma_j} \frac{1}{|y-z|^{n-2\sigma}} \left(\frac{1}{|z|}\right)^{n+2\sigma} dz$$

$$\geq C(|y|-\lambda) u(x_j)^{-1} - C(|y|-\lambda) u(x_j)^{-\frac{2n}{n-2\sigma}}$$

$$\geq C(|y|-\lambda) u(x_j)^{-1},$$

where we have used $\mathcal{L}^n(\Sigma_j) = 0$ and $u(x_j) \to \infty$ as $j \to \infty$. Similarly, for $|y| \ge \lambda_1 + 1$ and $y \in \Pi_j$, we have

$$J_{\lambda}(y) \ge Cu(x_j)^{-1} - Cu(x_j)^{-\frac{2n}{n-2\sigma}} \ge Cu(x_j)^{-1}.$$

Lemma 4.4 is established.

By (4.62), (4.63) and (4.64), there exists a small $\varepsilon_1 \in (0, \lambda_1 - \bar{\lambda})$ (which depends on j) such that

$$\varphi_{\bar{\lambda}}(y) + \Phi_{\bar{\lambda}}(y) \ge J_{\bar{\lambda}}(y) \ge Cu(x_j)^{-1} \ge \frac{\varepsilon_1}{|y|^{n-2\sigma}} \quad \forall |y| \ge \lambda_1 + 1, y \in \Pi_j.$$

By the above estimate and the explicit formulas for $w_j^{\lambda}(y)$ and $\Phi_{\lambda}(y)$, there exists $\varepsilon_2 \in (0, \varepsilon_1)$ such that for any $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon_2$,

$$\varphi_{\lambda}(y) + \Phi_{\lambda}(y) = \left[\varphi_{\bar{\lambda}}(y) + \Phi_{\bar{\lambda}}(y)\right] + \left[\Phi_{\lambda}(y) - \Phi_{\bar{\lambda}}(y)\right] + \left[w_{j}^{\bar{\lambda}}(y) - w_{j}^{\lambda}(y)\right]$$

$$\geq \frac{\varepsilon_{1}}{2|y|^{n-2\sigma}} \quad \forall |y| \geq \lambda_{1} + 1, \ y \in \Pi_{j}.$$

$$(4.69)$$

This, together with Lemma 4.3, also implies that for any $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon_2$,

$$w_j(y) - w_j^{\lambda}(y) \ge \frac{\varepsilon_1}{2|y|^{n-2\sigma}} \quad \forall |y| \ge \lambda_1 + 1, \ y \in \Pi_j.$$
 (4.70)

For $\varepsilon \in (0, \varepsilon_2)$ which we choose below, by (4.63), (4.64) and (4.70) we have, for $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$

and for $\lambda \leq |y| \leq \lambda_1 + 1$,

$$\begin{split} \varphi_{\lambda}(y) + \Phi_{\lambda}(y) &\geq \int_{\lambda \leq |z| \leq \lambda_{1} + 1} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz \\ &+ \int_{\lambda_{1} + 2 \leq |z| \leq \lambda_{1} + 3} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz \\ &\geq -C \int_{\lambda \leq |z| \leq \lambda + \varepsilon} G(0,\lambda;y,z) (|z| - \lambda) dz \\ &+ \int_{\lambda + \varepsilon \leq |z| \leq \lambda_{1} + 1} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}^{\bar{\lambda}}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz \\ &+ A_{1} \int_{\lambda_{1} + 2 \leq |z| \leq \lambda_{1} + 3} G(0,\lambda;y,z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz, \end{split}$$

where $A_1 = \min_{B_2} K > 0$, and we also used

$$|w_j(z)^{\frac{n+2\sigma}{n-2\sigma}} - (w_j)_{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}| \le C(|z| - \lambda)$$

in the second inequality. By (4.70) there exists $\hat{\delta} > 0$ (which depends on j) such that

$$w_j(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_j^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \ge \hat{\delta}$$
 for $\lambda_1 + 2 \le |z| \le \lambda_1 + 3$.

Since $\|w_j\|_{C^1(B_{\lambda_1+2})} \leq C$ and $\|K_j\|_{L^\infty(B_{\lambda_1+2})} \leq C$ for some constant C independent of j, there exists some constant C > 0 independent of both ε and j such that for any $\bar{\lambda} \leq \bar{\lambda} \leq \bar{\lambda} + \varepsilon$,

$$K_{j}(z)|w_{j}^{\bar{\lambda}}(z)|^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)|^{\frac{n+2\sigma}{n-2\sigma}}| \leq C(\lambda - \bar{\lambda}) \leq C\varepsilon \quad \forall \lambda \leq |z| \leq \lambda_{1} + 1.$$

For any $\lambda \leq |y| \leq \lambda_1 + 1$, one can estimate the integrals of the kernel G (or, see [30]) as follows:

$$\int_{\lambda+\varepsilon \le |z| \le \lambda_1 + 1} G(0, \lambda; y, z) dz \le \left| \int_{\lambda+\varepsilon \le |z| \le \lambda_1 + 1} \left(\frac{1}{|y-z|^{n-2\sigma}} - \frac{1}{|y^{\lambda} - z|^{n-2\sigma}} \right) dz \right| + \int_{\lambda+\varepsilon \le |z| \le \lambda_1 + 1} \left| \left(\frac{\lambda}{|y|} \right)^{n-2\sigma} - 1 \right| \frac{1}{|y^{\lambda} - z|^{n-2\sigma}} dz \\ \le C(\varepsilon^{2\sigma - 1} + |\ln \varepsilon| + 1)(|y| - \lambda)$$

and

$$\int_{\lambda \le |z| \le \lambda + \varepsilon} G(0, \lambda; y, z) (|z| - \lambda) dz \le \left| \int_{\lambda \le |z| \le \lambda + \varepsilon} \left(\frac{|z| - \lambda}{|y - z|^{n - 2\sigma}} - \frac{|z| - \lambda}{|y^{\lambda} - z|^{n - 2\sigma}} \right) dz \right|$$

$$+ \varepsilon \int_{\lambda \le |z| \le \lambda + \varepsilon} \left| \left(\frac{\lambda}{|y|} \right)^{n - 2\sigma} - 1 \right| \frac{1}{|y^{\lambda} - z|^{n - 2\sigma}} dz$$

$$\le C(|y| - \lambda) \varepsilon^{2\sigma/n} + C\varepsilon(|y| - \lambda)$$

$$\le C(|y| - \lambda) \varepsilon^{2\sigma/n}.$$

Therefore, by (4.67) we have for $\lambda < |y| \le \lambda_1 + 1$ that

$$\varphi_{\lambda}(y) + \Phi_{\lambda}(y) \ge -C\varepsilon^{2\sigma/n}(|y| - \lambda) + \delta_{1}\hat{\delta}(|y| - \lambda) \int_{\lambda_{1} + 2 \le |z| \le \lambda_{1} + 3} \frac{1}{|y - z|} dz$$
$$\ge \left(\delta_{1}\hat{\delta}c - C\varepsilon^{2\sigma/n}\right)(|y| - \lambda) \ge 0$$

if ε is sufficiently small. This and (4.69) contradict the definition of $\bar{\lambda}$. Hence the proof of Step 2 is finished.

Step 3. In this step, we will complete the proof of Theorem 1.6 under the assumption of (K1). By Step 2 we have $\lim_{j\to\infty} |\nabla K(x_j)| = |\nabla K(0)| = 0$. From the assumption (K1) we know that there exists a neighborhood N of 0 such that (1.6) and (1.7) hold. Here we can suppose $N=B_2$, since otherwise we just consider the integral equation on N and then make a rescaling argument. That is, we suppose K satisfies

$$c_1|y|^{\alpha-1} \le |\nabla K(y)| \le c_2|y|^{\alpha-1} \quad \text{for } y \in B_2,$$
 (4.71)

and for any $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon)$ such that for $|z - y| < \delta|y|$ we have

$$|\nabla K(z) - \nabla K(y)| < \varepsilon |y|^{\alpha - 1}, \tag{4.72}$$

where $\alpha > 1$, $y, z \in B_2$, c_1 and c_2 are two positive constants.

Without loss of generality, we may assume

$$\lim_{j \to \infty} \frac{\nabla K(x_j)}{|\nabla K(x_j)|} = e = (1, 0, \dots, 0). \tag{4.73}$$

As in Step 2, let w_j and h_j be defined as in (4.17) and let Ω_j be defined as in (4.18). Then w_j still satisfies (4.21) on Ω_j where K_j is defined as in (4.20). Moreover, on every compact subset of \mathbb{R}^n , w_j converges in C^2 norm to $U_1 = U_0(\cdot - Re)$ by Step 1. We also extend w_j to be identically 0 outside Ω_j and K to be identically 0 outside B_2 .

Define w_j^{λ} and h_j^{λ} as in (4.19), then w_j^{λ} satisfies (4.22). It is clear that Lemma 4.1 still holds. Let $\varphi_{\lambda}(y) = w_j(y) - w_j^{\lambda}(y)$ with $\lambda \in [R-2,R+2]$, then φ_{λ} satisfies the following integral inequality for large j,

$$\varphi_{\lambda}(y) + \Phi_{\lambda}(y) \ge \int_{\Pi_{\beta} \backslash B_{\lambda}} G(0, \lambda; y, z) b_{\lambda}(z) \varphi_{\lambda}(z) dz + J_{\lambda}(y), \quad y \in \Pi_{j} \backslash \overline{B}_{\lambda}, \tag{4.74}$$

where Π_i , b_λ , Φ_λ and J_λ are respectively given as in (4.32), (4.36), (4.37) and (4.38).

In order to apply the moving sphere method, as in Step 2, we need to establish some estimates for Φ_{λ} on $\Pi_{j} \backslash \overline{B}_{\lambda}$. Define a special domain D_{λ} in $\Omega_{j} \backslash B_{\lambda}$ as follows,

$$D_{\lambda} = \{ z \in \mathbb{R}^n : \lambda < |z| < 2\lambda, z_1 > 2|z'| \text{ where } z' = (z_2, \dots, z_n) \}.$$

Define $l_j := M_j^{\frac{2}{n-2\sigma}} |x_j|$ where $M_j = u(x_j)$, then $l_j \to \infty$ and so $l_j \gg R$ when j is large. Under our current assumption (4.73), we have

Claim 3: There exists a constant C > 0 independent of λ such that for all large j,

$$K_j(z^{\lambda}) - K_j(z) \le -CM_j^{-\frac{2}{n-2\sigma}} |x_j|^{\alpha-1} (|z| - \lambda), \quad z \in D_{\lambda}.$$
 (4.75)

$$|K_j(z^{\lambda}) - K_j(z)| \le CM_j^{-\frac{2}{n-2\sigma}} |x_j|^{\alpha-1} (|z| - \lambda), \quad z \in B_{2l_j} \setminus \overline{B}_{\lambda}. \tag{4.76}$$

$$|K_j(z^{\lambda}) - K_j(z)| \le CM_j^{-\frac{2\alpha}{n-2\sigma}} |z|^{\alpha}, \quad z \in \Omega_j \backslash B_{2l_j}. \tag{4.77}$$

Proof. (1) For $z \in D_{\lambda}$, by the mean value theorem and the assumptions (4.71)-(4.72) on K we obtain that for j large,

$$K_{j}(z^{\lambda}) - K_{j}(z) = M_{j}^{-\frac{2}{n-2\sigma}} \nabla K \left(x_{j} + M_{j}^{-\frac{2}{n-2\sigma}} \left((1-\theta)z^{\lambda} + \theta z - Re \right) \right) \cdot (z^{\lambda} - z)$$

$$\leq M_{j}^{-\frac{2}{n-2\sigma}} \nabla K(x_{j}) \cdot (z^{\lambda} - z) + \varepsilon M_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} |z^{\lambda} - z|$$

$$\leq -CM_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} (|z| - \lambda)$$

by taking ε sufficiently small, where $\theta \in (0,1)$ and C > 0 is a constant.

(2) For $z \in B_{2l_j} \setminus \overline{B}_{\lambda}$, by the mean value theorem and the assumption (4.71) on K we obtain that

$$|K_j(z^{\lambda}) - K_j(z)| \le M_j^{-\frac{2}{n-2\sigma}} \left| \nabla K \left(x_j + M_j^{-\frac{2}{n-2\sigma}} \left((1-\theta)z^{\lambda} + \theta z - Re \right) \right) \right| |z^{\lambda} - z|$$

$$\le C M_j^{-\frac{2}{n-2\sigma}} |x_j|^{\alpha-1} (|z| - \lambda)$$

for some constant C > 0, where $\theta \in (0, 1)$.

(3) For $z \in \Omega_j \backslash B_{2l_j}$, by the mean value theorem and the assumption (4.71) on K we obtain that

$$|K_{j}(z^{\lambda}) - K_{j}(z)| \leq M_{j}^{-\frac{2}{n-2\sigma}} \left| \nabla K \left(x_{j} + M_{j}^{-\frac{2}{n-2\sigma}} \left((1-\theta)z^{\lambda} + \theta z - Re \right) \right) \right| |z^{\lambda} - z|$$

$$\leq C M_{j}^{-\frac{2}{n-2\sigma}} \left| x_{j} + M_{j}^{-\frac{2}{n-2\sigma}} \left((1-\theta)z^{\lambda} + \theta z - Re \right) \right|^{\alpha-1} |z^{\lambda} - z|$$

$$\leq C M_{j}^{-\frac{2\alpha}{n-2\sigma}} |z|^{\alpha}$$

for some constant C > 0, where $\theta \in (0, 1)$. Claim 3 is proved.

It is also clear that Lemma 4.2 still holds. If we let

$$Q_{\lambda}(z) = \left(K_{j}(z^{\lambda}) - K_{j}(z) \right) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \quad \text{for } z \in \Omega_{j} \backslash B_{\lambda},$$

then by Claim 3 and Lemma 4.2 we have

$$Q_{\lambda}(z) \le -CM_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} (|z| - \lambda) \left(\frac{1}{1 + (z_{1} - \lambda)^{2} + |z'|^{2}} \right)^{\frac{n+2\sigma}{2}} \quad \text{for } z \in D_{\lambda} \quad (4.78)$$

and

$$|Q_{\lambda}(z)| \leq \begin{cases} CM_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} (|z|-\lambda) & \text{for } z \in D_{\lambda}, \\ CM_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} (|z|-\lambda) |z|^{-2\sigma-n} & \text{for } z \in B_{2l_{j}} \setminus (B_{\lambda} \cup D_{\lambda}), \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} |z|^{\alpha-2\sigma-n} & \text{for } z \in \Omega_{j} \setminus B_{2l_{j}}. \end{cases}$$

$$(4.79)$$

We split $\Phi_{\lambda}(y)$ into two parts:

$$\Phi_{\lambda}(y) = \int_{B_{2l_{j}} \backslash B_{\lambda}} G(0, \lambda; y, z) Q_{\lambda}(z) dz + \int_{\Omega_{j} \backslash B_{2l_{j}}} G(0, \lambda; y, z) Q_{\lambda}(z) dz$$

$$=: \Phi_{1,\lambda}(y) + \Phi_{2,\lambda}(y). \tag{4.80}$$

By the estimates of $Q_{\lambda}(z)$ on $B_{2l_j}\backslash B_{\lambda}$, one can see easily that the estimates of $\Phi_{1,\lambda}$ are very similar to those of Φ_{λ} in Lemma 4.3 of Step 2. Hence we have the following:

Lemma 4.5. There exists a constant C > 0 independent of λ such that for all large j,

$$\Phi_{1,\lambda}(y) \le \begin{cases} -CM_j^{-\frac{2}{n-2\sigma}} |x_j|^{\alpha-1} (|y|-\lambda)\lambda^{-n} \log \lambda & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ -CM_j^{-\frac{2}{n-2\sigma}} |x_j|^{\alpha-1} |y|^{2\sigma-n} \lambda^{1-2\sigma} \log \lambda & \text{for } y \in \Pi_j \backslash \overline{B}_{4\lambda} \end{cases}$$

and

$$|\Phi_{1,\lambda}(y)| \leq \begin{cases} CM_j^{-\frac{2}{n-2\sigma}} |x_j|^{\alpha-1} (|y|-\lambda) \lambda^{2\sigma} & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ CM_j^{-\frac{2}{n-2\sigma}} |x_j|^{\alpha-1} |y|^{2\sigma-n} \lambda^{n+1} & \text{for } y \in \Pi_j \backslash \overline{B}_{4\lambda}. \end{cases}$$

Proof. We only need to replace $M_j^{-\frac{2}{n-2\sigma}}$ and $\Omega_j \backslash B_\lambda$ in Lemma 4.3 by $M_j^{-\frac{2}{n-2\sigma}} |x_j|^{\alpha-1}$ and $B_{2l_j} \backslash B_\lambda$, respectively.

The estimates of $\Phi_{2,\lambda}$ depend on α . For the sake of application, let's define

$$g_{\lambda}(y) = M_j^{-\frac{2\alpha}{n-2\sigma}} \int_{\Omega_j \setminus B_{2l_j}} G(0,\lambda;y,z) |z|^{\alpha-2\sigma-n} dz \quad \text{for } \lambda < |y| \le 3L_j,$$

where $L_j:=M_j^{\frac{2}{n-2\sigma}}$. Then $\Omega_j\subset B_{3L_j}$ for large j and $|\Phi_{2,\lambda}(y)|\leq Cg_{\lambda}(y)$ for some constant C>0 depending only on K,n,σ .

Lemma 4.6. For $\lambda < |y| \le 4\lambda$, we have

$$g_{\lambda}(y) \leq \begin{cases} CM_{j}^{-\frac{2n}{n-2\sigma}} (|y| - \lambda)\lambda^{-1} & \text{if } \alpha > n, \\ CM_{j}^{-\frac{2n}{n-2\sigma}} \log M_{j} (|y| - \lambda)\lambda^{-1} & \text{if } \alpha = n, \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} l_{j}^{\alpha - n} (|y| - \lambda)\lambda^{-1} & \text{if } 1 < \alpha < n. \end{cases}$$

$$(4.81)$$

For $4\lambda \leq |y| \leq l_j$, we have

$$g_{\lambda}(y) \leq \begin{cases} CM_{j}^{-\frac{2n}{n-2\sigma}} & \text{if } \alpha > n, \\ CM_{j}^{-\frac{2n}{n-2\sigma}} \log M_{j} & \text{if } \alpha = n, \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} l_{j}^{\alpha-n} & \text{if } 1 < \alpha < n. \end{cases}$$

$$(4.82)$$

For $l_i \leq |y| \leq 3L_i$, we have

$$g_{\lambda}(y) \leq \begin{cases} CM_{j}^{-\frac{2n}{n-2\sigma}} & \text{if } \alpha > n, \\ CM_{j}^{-\frac{2n}{n-2\sigma}} \log M_{j} & \text{if } \alpha = n, \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} |y|^{\alpha-n} & \text{if } 2\sigma < \alpha < n, \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} |y|^{2\sigma-n} \log |y| & \text{if } \alpha = 2\sigma, \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} l_{j}^{\alpha-2\sigma} |y|^{2\sigma-n} & \text{if } 1 < \alpha < 2\sigma. \end{cases}$$

$$(4.83)$$

Proof. (1) $\lambda < |y| \le 4\lambda$. By Lemma 3.1 we have

$$G(0,\lambda;y,z) \leq C \frac{(|y|-\lambda)(|z|^2-\lambda^2)}{\lambda|y-z|^{n-2\sigma+2}} \leq C \frac{|y|-\lambda}{\lambda}|z|^{2\sigma-n}.$$

Hence

$$\begin{split} g_{\lambda}(y) &\leq CM_{j}^{-\frac{2\alpha}{n-2\sigma}}(|y|-\lambda)\lambda^{-1}\int_{\Omega_{j}\backslash B_{2l_{j}}}|z|^{\alpha-2n}dz\\ &\leq CM_{j}^{-\frac{2\alpha}{n-2\sigma}}(|y|-\lambda)\lambda^{-1}\int_{2l_{j}\leq|z|\leq3L_{j}}|z|^{\alpha-2n}dz\\ &\leq \begin{cases} CM_{j}^{-\frac{2n}{n-2\sigma}}(|y|-\lambda)\lambda^{-1} & \text{if }\alpha>n,\\ CM_{j}^{-\frac{2n}{n-2\sigma}}\log M_{j}(|y|-\lambda)\lambda^{-1} & \text{if }\alpha=n,\\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}}l_{j}^{\alpha-n}(|y|-\lambda)\lambda^{-1} & \text{if }1<\alpha$$

 $(2) \ 4\lambda \leq |y| \leq l_j. \text{ Since } |z| \geq 2l_j \text{ and thus } |y| \leq |z|/2, \text{ we have } G(0,\lambda;y,z) \leq C|y-z|^{2\sigma-n} \leq C|z|^{2\sigma-n}. \text{ Hence}$

$$g_{\lambda}(y) \leq CM_{j}^{-\frac{2\alpha}{n-2\sigma}} \int_{\Omega_{j} \backslash B_{2l_{j}}} |z|^{\alpha-2n} dz$$

$$\leq \begin{cases} CM_{j}^{-\frac{2n}{n-2\sigma}} & \text{if } \alpha > n, \\ CM_{j}^{-\frac{2n}{n-2\sigma}} \log M_{j} & \text{if } \alpha = n, \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} l_{j}^{\alpha-n} & \text{if } 1 < \alpha < n. \end{cases}$$

(3) $l_j \leq |y| \leq 3L_j$. We separate the region $\Omega_j \backslash B_{2l_j}$ into four parts:

$$\mathcal{E}_1 = \{ z \in \Omega_j \backslash B_{2l_j} : |z| < |y|/2 \},$$

$$\mathcal{E}_2 = \{ z \in \Omega_j \backslash B_{2l_j} : |z| > 2|y| \},$$

$$\mathcal{E}_3=\{z\in\Omega_j\backslash B_{2l_j}:|z-y|\leq |y|/2\},$$

$$\mathcal{E}_4=\{z\in\Omega_j\backslash B_{2l_j}:|z-y|\geq |y|/2\text{ and }|y|/2\leq |z|\leq 2|y|\}.$$

Note that some of these sets may be empty. Then

$$\begin{split} g_{\lambda}(y) &\leq CM_{j}^{-\frac{2\alpha}{n-2\sigma}} \int_{\Omega_{j} \backslash B_{2l_{j}}} |y-z|^{2\sigma-n} |z|^{\alpha-2\sigma-n} dz \\ &\leq CM_{j}^{-\frac{2\alpha}{n-2\sigma}} \left(|y|^{2\sigma-n} \int_{\mathcal{E}_{1}} |z|^{\alpha-2\sigma-n} dz + \int_{\mathcal{E}_{2}} |z|^{2\sigma-n} |z|^{\alpha-2\sigma-n} dz \\ &+ |y|^{\alpha-2\sigma-n} \int_{\mathcal{E}_{3}} |y-z|^{2\sigma-n} dz + |y|^{2\sigma-n} \int_{\mathcal{E}_{4}} |z|^{\alpha-2\sigma-n} dz \right) \\ &\leq CM_{j}^{-\frac{2\alpha}{n-2\sigma}} \left(|y|^{2\sigma-n} \int_{2l_{j}}^{|y|/2} r^{\alpha-2\sigma-1} dr + \int_{2|y|}^{3L_{j}} r^{\alpha-n-1} dr \right) \\ &+ |y|^{\alpha-n} + |y|^{2\sigma-n} \int_{|y|/2}^{2|y|} r^{\alpha-2\sigma-1} dr \right) \\ &\leq \begin{cases} CM_{j}^{-\frac{2n}{n-2\sigma}} & \text{if } \alpha > n, \\ CM_{j}^{-\frac{2n}{n-2\sigma}} \log M_{j} & \text{if } \alpha = n, \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} |y|^{\alpha-n} & \text{if } 2\sigma < \alpha < n, \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} |y|^{2\sigma-n} \log |y| & \text{if } \alpha = 2\sigma, \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} l_{j}^{\alpha-2\sigma} l_{j}^{\alpha-2\sigma} |y|^{2\sigma-n} & \text{if } 1 < \alpha < 2\sigma. \end{cases} \end{split}$$

The proof of Lemma 4.6 is finished.

Next we consider two cases for $\alpha > 1$ separately.

Case 1: $\alpha < 2\sigma$ or $\alpha \ge (n-2\sigma)/2$. We construct a function H_{λ} which is non-positive on $\Pi_j \backslash \overline{B}_{\lambda}$. Let

$$H_{\lambda}(y) = -\varepsilon M_{j}^{-1}(\lambda^{2\sigma - n} - |y|^{2\sigma - n}) + \Phi_{\lambda}(y), \quad y \in \Pi_{j} \setminus \overline{B}_{\lambda}$$
 (4.84)

for some small $\varepsilon \in (0, \varepsilon_0/4)$ where ε_0 is defined in Lemma 4.1. It is clear that Lemma 4.4 still holds, and by using Lemma 4.4 we can choose a small $\varepsilon > 0$ such that for any $\lambda \in [\lambda_0, \lambda_1]$,

$$J_{\lambda}(y) - \varepsilon M_j^{-1}(\lambda^{2\sigma - n} - |y|^{2\sigma - n}) \ge \frac{1}{2} J_{\lambda}(y), \quad y \in \Pi_j \backslash \overline{B}_{\lambda}. \tag{4.85}$$

Here we recall that $\lambda_0 = R - 2$ and $\lambda_1 = R + 2$. Then by Lemmas 4.5 and 4.6 we have for $\lambda \in [\lambda_0, \lambda_1]$ and for large j,

$$H_{\lambda}(y) < 0, \quad y \in \Pi_j \backslash \overline{B}_{\lambda}.$$
 (4.86)

Moreover, it follows from Lemmas 4.1, 4.5 and 4.6 that for large j,

$$\varphi_{\lambda_0}(y) + H_{\lambda_0}(y) \ge 0, \quad y \in \Pi_j \backslash \overline{B}_{\lambda_0}.$$

By (4.74) and (4.85), $\varphi_{\lambda} + H_{\lambda}$ satisfies

$$\varphi_{\lambda}(y) + H_{\lambda}(y) \ge \int_{\Pi_{j} \backslash B_{\lambda}} G(0, \lambda; y, z) b_{\lambda}(z) \varphi_{\lambda}(z) dz + \frac{1}{2} J_{\lambda}(y) \quad \text{for } y \in \Pi_{j} \backslash \overline{B}_{\lambda}, \quad (4.87)$$

where $\lambda \in [\lambda_0, \lambda_1]$. We define

$$\bar{\lambda} := \sup\{\mu \ge \lambda_0 \mid \varphi_{\lambda}(y) + H_{\lambda}(y) \ge 0, \ \forall \ y \in \Pi_j \setminus \overline{B}_{\lambda}, \ \forall \ \lambda_0 \le \lambda \le \mu\}.$$

Then, $\bar{\lambda}$ is well defined and $\bar{\lambda} \geq \lambda_0$ for all large j. Furthermore, by (4.86) and Lemma 4.1 we have $\bar{\lambda} < \lambda_1$ for all large j. Then, as in Step 2, applying the moving sphere method one can derive a contradiction to the definition of $\bar{\lambda}$. The proof of Theorem 1.6 in Case 1 is finished.

Case 2: $2\sigma \le \alpha < (n-2\sigma)/2$. By (4.79) we can choose a constant $\hat{C} > 0$ such that

$$\hat{C}M_j^{-\frac{2\alpha}{n-2\sigma}}|z|^{\alpha-2\sigma-n} - Q_{\lambda}(z) \ge M_j^{-\frac{2\alpha}{n-2\sigma}}|z|^{\alpha-2\sigma-n} \quad \text{for } z \in \Omega_j \backslash B_{2l_j}. \tag{4.88}$$

Let

$$T_{\lambda}(y) = \hat{C}M_{j}^{-\frac{2\alpha}{n-2\sigma}} \int_{\Omega_{j} \setminus B_{2l_{j}}} G(0,\lambda;y,z)|z|^{\alpha-2\sigma-n} dz \quad \text{for } \lambda < |y| \le 3L_{j}.$$
 (4.89)

As a consequence of Lemma 4.6, we have the following estimates for T_{λ} .

Lemma 4.7. Suppose $2\sigma \le \alpha < (n-2\sigma)/2$. For $\lambda < |y| \le 4\lambda$, we have

$$T_{\lambda}(y) \le CM_j^{-\frac{2}{n-2\sigma}} |x_j|^{\alpha-1} (|y| - \lambda) \lambda^{-n}.$$
 (4.90)

For $4\lambda \leq |y| \leq 2l_j$, we have

$$T_{\lambda}(y) \le CM_j^{-\frac{2}{n-2\sigma}} |x_j|^{\alpha-1} |y|^{2\sigma-n} \lambda^{1-2\sigma}.$$
 (4.91)

For $2l_j \leq |y| \leq 3L_j$, we have

$$T_{\lambda}(y) \le \begin{cases} CM_{j}^{-\frac{2\alpha}{n-2\sigma}} |y|^{\alpha-n} & \text{if } 2\sigma < \alpha < (n-2\sigma)/2, \\ CM_{j}^{-\frac{4\sigma}{n-2\sigma}} |y|^{2\sigma-n} \log |y| & \text{if } \alpha = 2\sigma. \end{cases}$$
(4.92)

Proof. When $\lambda < |y| \le 4\lambda$, by (4.81) we obtain

$$T_{\lambda}(y) \le CM_{j}^{-\frac{2\alpha}{n-2\sigma}} l_{j}^{\alpha-1} \cdot l_{j}^{1-n}(|y|-\lambda)\lambda^{-1} \le CM_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1}(|y|-\lambda)\lambda^{-n},$$

where we used $l_j=M_j^{\frac{2}{n-2\sigma}}|x_j|$ and $l_j\to\infty$. When $4\lambda\leq |y|\leq 3L_j$, by (4.82) and (4.83) we have

$$T_{\lambda}(y) \leq \begin{cases} CM_{j}^{-\frac{2\alpha}{n-2\sigma}} |y|^{\alpha-n} & \text{if } 2\sigma < \alpha < (n-2\sigma)/2, \\ CM_{j}^{-\frac{4\sigma}{n-2\sigma}} |y|^{2\sigma-n} \log |y| & \text{if } \alpha = 2\sigma. \end{cases}$$

In particular, for $4\lambda \leq |y| \leq 2l_j$ we have

$$\begin{split} T_{\lambda}(y) &\leq C M_{j}^{-\frac{2\alpha}{n-2\sigma}} |y|^{\alpha-n} \log |y| \\ &\leq C M_{j}^{-\frac{2\alpha}{n-2\sigma}} |y|^{\alpha-2\sigma} |y|^{2\sigma-n} \log |y| \\ &\leq C M_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} |y|^{2\sigma-n} l_{j}^{1-2\sigma} \log l_{j} \\ &\leq C M_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} |y|^{2\sigma-n} \lambda^{1-2\sigma}, \end{split}$$

where we used $\sigma > \frac{1}{2}$ and $l_j \to \infty$. Lemma 4.7 is proved.

Define

$$H_{\lambda}(y) := \Phi_{1,\lambda}(y) + T_{\lambda}(y) \quad \text{for } y \in \Pi_{j} \setminus \overline{B}_{\lambda}.$$

Then from Lemmas 4.5 and 4.7 we obtain

$$H_{\lambda}(y) \leq \begin{cases} -CM_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} (|y|-\lambda)\lambda^{-n} \log \lambda & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ -CM_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} |y|^{2\sigma-n} \lambda^{1-2\sigma} \log \lambda & \text{for } y \in B_{2l_{j}} \backslash B_{4\lambda}, \\ CM_{j}^{-\frac{2\alpha}{n-2\sigma}} |y|^{\alpha-n} & \text{for } y \in \Pi_{j} \backslash B_{2l_{j}}, \ 2\sigma < \alpha < (n-2\sigma)/2, \\ CM_{j}^{-\frac{4\sigma}{n-2\sigma}} |y|^{2\sigma-n} \log |y| & \text{for } y \in \Pi_{j} \backslash B_{2l_{j}}, \ \alpha = 2\sigma \end{cases}$$

$$(4.93)$$

and

$$|H_{\lambda}(y)| \leq \begin{cases} CM_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} (|y|-\lambda) \lambda^{2\sigma} & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ CM_{j}^{-\frac{2}{n-2\sigma}} |x_{j}|^{\alpha-1} |y|^{2\sigma-n} \lambda^{n+1} & \text{for } y \in B_{2l_{j}} \backslash B_{4\lambda}, \\ \theta_{j} |y|^{2\sigma-n} & \text{for } y \in \Pi_{j} \backslash B_{2l_{j}}, \end{cases}$$
(4.94)

where $\{\theta_j\}$ is a positive sequence satisfying $\theta_j \to 0$ as $j \to \infty$. Remark that in order to obtain (4.93), we could take R larger and then fix it. Furthermore, by (4.74) and (4.88) we see that φ_{λ} satisfies

$$\varphi_{\lambda}(y) + H_{\lambda}(y) \ge \int_{\Pi_{j} \backslash B_{\lambda}} G(0, \lambda; y, z) b_{\lambda}(z) \varphi_{\lambda}(z) dz
+ M_{j}^{-\frac{2\alpha}{n-2\sigma}} \int_{\Omega_{j} \backslash B_{2l_{j}}} G(0, \lambda; y, z) |z|^{\alpha-2\sigma-n} dz + J_{\lambda}(y), \quad y \in \Pi_{j} \backslash \overline{B}_{\lambda},$$
(4.95)

where $\lambda \in [\lambda_0, \lambda_1]$ with $\lambda_0 = R - 2$ and $\lambda_1 = R + 2$. Moreover, it follows from Lemma 4.1 and (4.94) that for large j,

$$\varphi_{\lambda_0}(y) + H_{\lambda_0}(y) \ge 0, \quad y \in \Pi_j \backslash \overline{B}_{\lambda_0}.$$

We define

$$\bar{\lambda} := \sup \{ \mu \ge \lambda_0 \mid \varphi_{\lambda}(y) + H_{\lambda}(y) \ge 0, \ \forall \ y \in \Pi_j \backslash \overline{B}_{\lambda}, \ \forall \ \lambda_0 \le \lambda \le \mu \}.$$

Then, $\bar{\lambda}$ is well defined and $\bar{\lambda} \geq \lambda_0$ for all large j. Furthermore, by (4.93) and Lemma 4.1 we have $\bar{\lambda} < \lambda_1$ for all large j. Using the continuity we obtain

$$\varphi_{\bar{\lambda}}(y) + H_{\bar{\lambda}}(y) \ge 0, \quad y \in \Pi_j \backslash \overline{B}_{\bar{\lambda}}.$$

This together with Lemma 4.5 and (4.93) yields

$$\varphi_{\bar{\lambda}}(y) \ge -H_{\bar{\lambda}}(y) \ge \begin{cases} 0, & y \in B_{2l_j} \setminus (\overline{B}_{\bar{\lambda}} \cup \Sigma_j), \\ -T_{\bar{\lambda}}(y), & y \in \Pi_j \setminus B_{2l_j}, \end{cases}$$
(4.96)

where Σ_j is the singular set of w_j .

Let

$$\mathcal{O}_{\lambda} = \{ y \in \Pi_j \backslash B_{2l_j} : w_j(y) < w_j^{\lambda}(y) \}.$$

By (4.95) and (4.96),

$$\begin{split} \varphi_{\bar{\lambda}}(y) + H_{\bar{\lambda}}(y) &\geq \int_{\Pi_{j} \backslash B_{2l_{j}}} G(0, \bar{\lambda}; y, z) b_{\bar{\lambda}}(z) \varphi_{\bar{\lambda}}(z) dz \\ &+ M_{j}^{-\frac{2\alpha}{n-2\sigma}} \int_{\Omega_{j} \backslash B_{2l_{j}}} G(0, \bar{\lambda}; y, z) |z|^{\alpha-2\sigma-n} dz + J_{\bar{\lambda}}(y) \\ &\geq - \int_{\mathcal{O}_{\bar{\lambda}}} G(0, \bar{\lambda}; y, z) b_{\bar{\lambda}}(z) T_{\bar{\lambda}}(z) dz + \int_{(\Pi_{j} \backslash B_{2l_{j}}) \backslash \mathcal{O}_{\bar{\lambda}}} G(0, \bar{\lambda}; y, z) b_{\bar{\lambda}}(z) \varphi_{\bar{\lambda}}(z) dz \\ &+ M_{j}^{-\frac{2\alpha}{n-2\sigma}} \int_{\Omega_{j} \backslash B_{2l_{j}}} G(0, \bar{\lambda}; y, z) |z|^{\alpha-2\sigma-n} dz + J_{\bar{\lambda}}(y), \quad y \in \Pi_{j} \backslash \overline{B}_{\bar{\lambda}}. \end{split}$$

Note that on $\mathcal{O}_{\bar{\lambda}}$, by Lemma 4.2 we have

$$0 \le b_{\bar{\lambda}}(z) = K_j(z) \frac{w_j(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_j^{\bar{\lambda}}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{w_j(z) - w_j^{\bar{\lambda}}(z)} \le Cw_j^{\bar{\lambda}}(z)^{\frac{4\sigma}{n-2\sigma}} \le C|z|^{-4\sigma}.$$

Thus, Lemma 4.7 gives

$$\begin{split} &-\int_{\mathcal{O}_{\bar{\lambda}}}G(0,\bar{\lambda};y,z)b_{\bar{\lambda}}(z)T_{\bar{\lambda}}(z)dz+M_{j}^{-\frac{2\alpha}{n-2\sigma}}\int_{\Omega_{j}\backslash B_{2l_{j}}}G(0,\bar{\lambda};y,z)|z|^{\alpha-2\sigma-n}dz\\ &\geq M_{j}^{-\frac{2\alpha}{n-2\sigma}}\int_{\mathcal{O}_{\bar{\lambda}}}G(0,\bar{\lambda};y,z)\left(|z|^{\alpha-2\sigma-n}-C|z|^{\alpha-4\sigma-n}\log|z|\right)dz\geq 0 \end{split}$$

when j is large. Therefore

$$\varphi_{\bar{\lambda}}(y) + H_{\bar{\lambda}}(y) \ge J_{\bar{\lambda}}(y), \quad y \in \Pi_i \backslash \overline{B}_{\bar{\lambda}}.$$

Lemma 4.4 is clearly still true. It follows from Lemma 4.4 that there exists a small $\varepsilon_1 \in (0, \lambda_1 - \lambda)$ (which depends on j) such that

$$\varphi_{\bar{\lambda}}(y) + H_{\bar{\lambda}}(y) \ge J_{\bar{\lambda}}(y) \ge Cu(x_j)^{-1} \ge \frac{\varepsilon_1}{|y|^{n-2\sigma}} \quad \forall |y| \ge \lambda_1 + 1, y \in \Pi_j.$$

By the above estimate and the explicit formulas for $w_j^{\lambda}(y)$ and $H_{\lambda}(y)$, there exists $\varepsilon_2 \in (0, \varepsilon_1)$ such that for any $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon_2$,

$$\varphi_{\lambda}(y) + H_{\lambda}(y) = \left[\varphi_{\bar{\lambda}}(y) + H_{\bar{\lambda}}(y)\right] + \left[H_{\lambda}(y) - H_{\bar{\lambda}}(y)\right] + \left[w_{j}^{\bar{\lambda}}(y) - w_{j}^{\lambda}(y)\right]$$

$$\geq \frac{\varepsilon_{1}}{2|y|^{n-2\sigma}} \quad \forall |y| \geq \lambda_{1} + 1, \ y \in \Pi_{j}.$$

$$(4.97)$$

This, together with Lemma 4.5 and (4.93), implies that for any $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon_2$,

$$w_{j}(y) - w_{j}^{\lambda}(y) \ge -H_{\lambda}(y) + \frac{\varepsilon_{1}}{2|y|^{n-2\sigma}} \ge \begin{cases} \frac{\varepsilon_{1}}{2|y|^{n-2\sigma}}, & y \in B_{2l_{j}} \backslash \Sigma_{j}, \ |y| \ge \lambda_{1} + 1, \\ -T_{\lambda}(y), & y \in \Pi_{j} \backslash B_{2l_{j}}. \end{cases}$$
(4.98)

For $\varepsilon \in (0, \varepsilon_2)$ which we choose below, by (4.95), (4.98) and (4.64), we have, for $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$ and for $\lambda \leq |y| \leq \lambda_1 + 1$,

$$\begin{split} \varphi_{\lambda}(y) + H_{\lambda}(y) &\geq \int_{\lambda \leq |z| \leq \lambda_{1} + 1} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz \\ &+ \int_{\lambda_{1} + 2 \leq |z| \leq \lambda_{1} + 3} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz \\ &- \int_{\mathcal{O}_{\lambda}} G(0,\lambda;y,z) b_{\lambda}(z) T_{\lambda}(z) dz + \int_{(\Pi_{j} \backslash B_{2l_{j}}) \backslash \mathcal{O}_{\lambda}} G(0,\lambda;y,z) b_{\lambda}(z) \varphi_{\lambda}(z) dz \\ &+ M_{j}^{-\frac{2\alpha}{n-2\sigma}} \int_{\Omega_{j} \backslash B_{2l_{j}}} G(0,\lambda;y,z) |z|^{\alpha-2\sigma-n} dz. \end{split}$$

On the set \mathcal{O}_{λ} , Lemma 4.2 leads to

$$0 \le b_{\lambda}(z) \le C w_j^{\lambda}(z)^{\frac{4\sigma}{n-2\sigma}} \le C|z|^{-4\sigma},$$

thus, using Lemma 4.7 yields that for j large,

$$-\int_{\mathcal{O}_{\lambda}} G(0,\lambda;y,z) b_{\lambda}(z) T_{\lambda}(z) dz + M_{j}^{-\frac{2\alpha}{n-2\sigma}} \int_{\Omega_{j} \setminus B_{2l_{j}}} G(0,\lambda;y,z) |z|^{\alpha-2\sigma-n} dz$$

$$\geq M_{j}^{-\frac{2\alpha}{n-2\sigma}} \int_{\mathcal{O}_{\lambda}} G(0,\lambda;y,z) \left(|z|^{\alpha-2\sigma-n} - C|z|^{\alpha-4\sigma-n} \log|z| \right) dz \geq 0.$$

Hence, for $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$ and for $\lambda \leq |y| \leq \lambda_1 + 1$, we have

$$\begin{split} \varphi_{\lambda}(y) + H_{\lambda}(y) &\geq \int_{\lambda \leq |z| \leq \lambda_{1}+1} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}\right) dz \\ &+ \int_{\lambda_{1}+2 \leq |z| \leq \lambda_{1}+3} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}\right) dz \\ &\geq -C \int_{\lambda \leq |z| \leq \lambda+\varepsilon} G(0,\lambda;y,z) (|z|-\lambda) dz \\ &+ \int_{\lambda+\varepsilon \leq |z| \leq \lambda_{1}+1} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}^{\bar{\lambda}}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}\right) dz \\ &+ A_{1} \int_{\lambda_{1}+2 \leq |z| \leq \lambda_{1}+3} G(0,\lambda;y,z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}\right) dz, \end{split}$$

where $A_1 = \min_{B_2} K > 0$, and we have used (4.96) and

$$|w_j(z)^{\frac{n+2\sigma}{n-2\sigma}} - (w_j)_{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}| \le C(|z| - \lambda)$$

in the second inequality. By (4.98) there exists $\hat{\delta} > 0$ (which depends on j) such that

$$w_j(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_j^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \ge \hat{\delta}$$
 for $\lambda_1 + 2 \le |z| \le \lambda_1 + 3$.

Since $\|w_j\|_{C^1(B_{\lambda_1+2})} \leq C$ and $\|K_j\|_{L^\infty(B_{\lambda_1+2})} \leq C$ for some constant C independent of j, there exists some constant C>0 independent of both ε and j such that for any $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$,

$$K_{j}(z)|w_{j}^{\bar{\lambda}}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}| \leq C(\lambda - \bar{\lambda}) \leq C\varepsilon \quad \forall \lambda \leq |z| \leq \lambda_{1} + 1.$$

For any $\lambda \leq |y| \leq \lambda_1 + 1$, we can estimate the kernel G as in Step 2 to obtain

$$\int_{\lambda+\varepsilon < |z| < \lambda_1 + 1} G(0, \lambda; y, z) dz \le C(\varepsilon^{2\sigma - 1} + |\ln \varepsilon| + 1)(|y| - \lambda)$$

and

$$\int_{\lambda < |z| < \lambda + \varepsilon} G(0,\lambda;y,z) (|z| - \lambda) dz \le C(|y| - \lambda) \varepsilon^{2\sigma/n}.$$

Thus, using (4.67) we have for $\lambda < |y| \le \lambda_1 + 1$ that

$$\varphi_{\lambda}(y) + H_{\lambda}(y) \ge -C\varepsilon^{2\sigma/n}(|y| - \lambda) + \delta_{1}\hat{\delta}(|y| - \lambda) \int_{\lambda_{1} + 2 \le |z| \le \lambda_{1} + 3} \frac{1}{|y - z|} dz$$
$$\ge \left(\delta_{1}\hat{\delta}c - C\varepsilon^{2\sigma/n}\right)(|y| - \lambda) \ge 0$$

if ε is sufficiently small. This and (4.97) contradict the definition of $\bar{\lambda}$, and so the proof in Case 2 is finished. The proof of Theorem 1.6 under the assumption (K1) is completed.

5 Local estimates under the assumption (K2)

In this section, by using the method of moving spheres introduced by Li-Zhu [35, 37], we shall prove Theorem 1.6 under the assumption (K2) in the spirit of the works of Chen-Lin [11] and Taliaferro-Zhang [51]. Similar to Section 4, we have to set up a framework to fit the integral equation rather than dealing directly with differential equations as in [11, 51]. In addition to developing integral techniques to overcome the lack of maximum principle, the non-locality of integral equation will bring a new difficulty under the present assumption (K2).

Proof of Theorem 1.6 under the assumption (K2). Suppose by contradiction that there exists a sequence $\{x_j\}_{j=1}^{\infty} \subset B_1 \setminus \Sigma$ such that

$$d_j := \operatorname{dist}(x_j, \Sigma) \to 0 \quad \text{ as } j \to \infty,$$

but

$$d_j^{\frac{n-2\sigma}{2}}u(x_j)\to\infty$$
 as $j\to\infty$.

Without loss of generality we may assume that $0 \in \Sigma$ and $x_j \to 0$ as $j \to \infty$.

Step 1. We show that x_j can be chosen as the local maximum points of u. Moreover, the functions $u(x_j)^{-1}u(x_j+u(x_j)^{-\frac{2}{n-2\sigma}}y)$ converge in $C^2_{loc}(\mathbb{R}^n)$, after passing a subsequence, to a positive function $U_0 \in C^2(\mathbb{R}^n)$ where U_0 satisfies

$$\begin{cases} U_0(y) = \int_{\mathbb{R}^n} \frac{K(0)U_0(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz & \text{for } y \in \mathbb{R}^n, \\ \max_{\mathbb{R}^n} U_0 = U_0(0) = 1. \end{cases}$$
 (5.1)

Step 2. For $n > 2\sigma$, we have $\nabla K(0) = 0$ under the assumption of $K \in C^1(B_2)$.

The proofs of these two steps are the same as those of Step 1 and Step 2 in Section 4, we omit their proofs. Since the following proof is long, we first explain the idea and the sketch of the proof. Let w_j be the scaled function in (5.2) (here we don't need to shift it) and let w_j^{λ} be the Kelvin transformation of w_j in (5.4). In order to take full advantage of the property in the assumption (K2) that $c(\cdot)$ is sufficiently small near 0, we will restrict the integral equation (1.12) to a small ball. Thus, for some small $\tau \in (0, \frac{1}{4})$ we denote

$$\Pi_j := \left\{ y \in \mathbb{R}^n : x_j + u(x_j)^{-\frac{2}{n-2\sigma}} y \in B_\tau \backslash \Sigma \right\}.$$

Note that the set Π_j depends on τ . Let $\varphi_\lambda = w_j - w_j^\lambda$ and let Φ_λ be as in (5.15) where $\lambda \in [1/2, 2]$. Then $\varphi_\lambda + \Phi_\lambda$ satisfies the integral inequality (5.17). Moreover, by Lemma 5.2 the remainder term J_λ has a lower bound which is independent of τ . After performing the above restriction to (1.12), we need extra efforts to obtain estimates for Φ_λ . Based on the assumption (K2) about the function K, we will complete the proof in three situations. Denote $M_j = u(x_j)$.

(1) $2\sigma < n < 2\sigma + 2$. By the assumption (K2) we know that $K \in \mathcal{C}^{\alpha}(B_2)$ with $\alpha = \frac{n-2\sigma}{2} < 1$. Furthermore, Φ_{λ} satisfies the estimates in Lemma 5.3. By choosing sufficiently small $\varepsilon_1 > 0$ and $\tau > 0$ we can construct

$$H_{\lambda}(y) = -\varepsilon_1 M_j^{-1}(\lambda^{2\sigma - n} - |y|^{2\sigma - n}) + \Phi_{\lambda}(y), \quad y \in \Pi_j \setminus \overline{B}_{\lambda}$$

such that $H_{\lambda} < 0$ for all $\lambda \in [1/2, 2]$. With the help of Lemma 5.2, the method of moving spheres can be applied to $\varphi_{\lambda} + H_{\lambda}$ to reach a contradiction.

- (2) $n = 2\sigma + 2$. In this case, we have $K \in C^1(B_2)$. By Step 2 we know $\nabla K(0) = 0$. Similar to situation (1), Φ_{λ} also satisfies the estimates in Lemma 5.3. The rest of the proof is the same as that of situation (1).
- (3) $n>2\sigma+2$. By the assumption (K2) we have $K\in\mathcal{C}^{\alpha}(B_2)$ with $\alpha=\frac{n-2\sigma}{2}>1$. It follows from Step 2 that $|\nabla K(x_j)|\to 0$. Furthermore, using the moving sphere method as in Step 2 of Section 4 and the assumption (K2) we obtain $|\nabla K(x_j)|^{\frac{1}{\alpha-1}}M_j^{\frac{2}{n-2\sigma}}\to 0$ (see Lemma 5.4). This together with the assumption (K2) yields that the estimates for Φ_{λ} in Lemma 5.3 still hold. The rest of the proof is the same as that of situation (1).

Now we return to the detailed proof of Theorem 1.6 under the assumption (K2). We denote $M_i = u(x_i)$ and define

$$w_j(y) = M_j^{-1} u \left(x_j + M_j^{-\frac{2}{n-2\sigma}} y \right), \ h_j(y) = M_j^{-1} h \left(x_j + M_j^{-\frac{2}{n-2\sigma}} y \right) \text{ in } \Omega_j,$$
 (5.2)

where

$$\Omega_j = \left\{ y \in \mathbb{R}^n : x_j + M_j^{-\frac{2}{n-2\sigma}} y \in B_2 \backslash \Sigma \right\}.$$
 (5.3)

It follows from Step 1 and the classification results in [14] or [35] that, modulo a positive constant, $w_j(y)$ converges in C^2 norm to the standard bubble $U_0(y) = (1+|y|^2)^{-\frac{n-2\sigma}{2}}$ on every compact subset of \mathbb{R}^n . We also suppose that w_j is to be identically 0 outside Ω_j and K is to be identically 0 outside B_2 . For $\lambda > 0$, let

$$w_j^{\lambda}(y) = \left(\frac{\lambda}{|y|}\right)^{n-2\sigma} w_j \left(\frac{\lambda^2 y}{|y|^2}\right), \quad h_j^{\lambda}(y) = \left(\frac{\lambda}{|y|}\right)^{n-2\sigma} h_j \left(\frac{\lambda^2 y}{|y|^2}\right) \tag{5.4}$$

and let

$$K_j(y) = K\left(x_j + M_j^{-\frac{2}{n-2\sigma}}y\right).$$
 (5.5)

Then

$$w_{j}(y) = \int_{\mathbb{R}^{n}} \frac{K_{j}(z)w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz + h_{j}(y) \quad \text{for } y \in \Omega_{j}.$$
 (5.6)

By (3.7), w_i^{λ} satisfies

$$w_j^{\lambda}(y) = \int_{\mathbb{R}^n} \frac{K_j(z^{\lambda}) w_j^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}}}{|y-z|^{n-2\sigma}} dz + h_j^{\lambda}(y) \quad \text{for } y \in \Omega_j \backslash \overline{B}_{\lambda},$$
 (5.7)

where $z^{\lambda} = \frac{\lambda^2 z}{|z|^2}$ is the inversion of z with respect to ∂B_{λ} .

As Lemma 4.1 we also have

Lemma 5.1. Let $\lambda_0 = \frac{1}{2}$ and $\lambda_1 = 2$. Then there exist $\varepsilon_0 > 0$ and $j_0 > 1$ such that for all $j \ge j_0$,

$$w_j(y) - w_j^{\lambda_0}(y) \ge \varepsilon_0(|y| - \lambda_0)|y|^{2\sigma - 1 - n} + \varepsilon_0 M_j^{-1}(\lambda_0^{2\sigma - n} - |y|^{2\sigma - n}), \quad y \in \Omega_j \setminus \overline{B}_{\lambda_0}.$$
 (5.8)

Moreover, there exists $y^* \in B_{2\lambda_1} \backslash \overline{B}_{\lambda_1}$ such that for all $j \geq j_0$,

$$w_i(y^*) - w_i^{\lambda_1}(y^*) \le -\varepsilon_0. \tag{5.9}$$

For some small $\tau \in (0, \frac{1}{4})$ to be determined later, we denote

$$\Pi_j := \left\{ y \in \mathbb{R}^n : x_j + M_j^{-\frac{2}{n-2\sigma}} y \in B_\tau \backslash \Sigma \right\} \subset \Omega_j$$
 (5.10)

and

$$\Sigma_j := \left\{ y \in \mathbb{R}^n : x_j + M_j^{-\frac{2}{n-2\sigma}} y \in \Sigma \right\}. \tag{5.11}$$

Note that we have $\mathcal{L}^n(\Sigma_j) = 0$ due to $\mathcal{L}^n(\Sigma) = 0$. It follows from the same arguments as in Lemma 3.1 of [30] that for all large j, there holds

$$h_i^{\lambda}(y) \le h_i(y) \quad \forall \ y \in \Pi_i \backslash \overline{B}_{\lambda}, \ \lambda \in [1/2, 2].$$
 (5.12)

Let $\varphi_{\lambda}(y) = w_j(y) - w_j^{\lambda}(y)$, where we omit j in the notation for brevity. By (3.8), (5.12) and $K_j \equiv 0$ on $\Omega_j^c \setminus \Sigma_j$, we have for $\lambda \in [1/2, 2]$ and $y \in \Pi_j \setminus \overline{B}_{\lambda}$ that

$$\varphi_{\lambda}(y) \ge \int_{B_{\lambda}^{c}} G(0,\lambda;y,z) \left(K_{j}(z) w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - K_{j}(z^{\lambda}) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz$$

$$= \int_{\Pi_{j} \backslash B_{\lambda}} G(0,\lambda;y,z) b_{\lambda}(z) \varphi_{\lambda}(z) dz + J_{\lambda}(y) - \Phi_{\lambda}(y),$$
(5.13)

where

$$b_{\lambda}(y) = K_{j}(y) \frac{w_{j}(y)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(y)^{\frac{n+2\sigma}{n-2\sigma}}}{w_{j}(y) - w_{j}^{\lambda}(y)},$$
(5.14)

$$\Phi_{\lambda}(y) = \int_{\Omega_{j} \backslash B_{\lambda}} G(0, \lambda; y, z) \left(K_{j}(z^{\lambda}) - K_{j}(z) \right) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} dz$$
 (5.15)

and

$$J_{\lambda}(y) = \int_{\Omega_{j}\backslash\Pi_{j}} G(0,\lambda;y,z) K_{j}(z) \left(w_{j}(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz$$

$$- \int_{\Omega_{j}^{c}} G(0,\lambda;y,z) K_{j}(z^{\lambda}) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} dz.$$
(5.16)

Notice that $b_{\lambda}(y)$ is always non-negative. Thus, $\varphi_{\lambda} + \Phi_{\lambda}$ satisfies the integral inequality

$$\varphi_{\lambda}(y) + \Phi_{\lambda}(y) \ge \int_{\Pi_{j} \backslash B_{\lambda}} G(0, \lambda; y, z) b_{\lambda}(z) \varphi_{\lambda}(z) dz + J_{\lambda}(y), \quad y \in \Pi_{j} \backslash \overline{B}_{\lambda}, \tag{5.17}$$

where $\lambda \in [1/2, 2]$.

Now, we give an estimate for J_{λ} defined in (5.16). Suppose $\lambda_0 = 1/2$ and $\lambda_1 = 2$.

Lemma 5.2. There exists a constant $\beta_0 > 0$ independent of τ such that for any $\lambda_0 \le \lambda \le \lambda_1$ and for all large j,

$$J_{\lambda}(y) \ge \begin{cases} \beta_0(|y| - \lambda)u(x_j)^{-1}, & \text{if } \lambda \le |y| \le \lambda_1 + 1, \\ \beta_0 u(x_j)^{-1}, & \text{if } |y| > \lambda_1 + 1, \ y \in \Pi_j. \end{cases}$$
(5.18)

Proof. The proof is similar to that of Lemma 4.4. The main difference here is that the set Π_j defined in (5.10) depends on τ , but we require that the constant β_0 in (5.18) cannot depend on τ . For the sake of completeness, we also include the proof.

For any $z \in \mathbb{R}^n \setminus (\Pi_j \cup \Sigma_j)$ and $\lambda_0 \leq \lambda \leq \lambda_1$, we have $|z| \geq \frac{\tau}{2} u(x_j)^{\frac{2}{n-2\sigma}}$ for all large j and thus

$$w_j^{\lambda}(z) \le \left(\frac{\lambda}{|z|}\right)^{n-2\sigma} \max_{B_{\lambda_1}} w_j \le C\tau^{-(n-2\sigma)} u(x_j)^{-2},$$

where C > 0 depends only on n and σ . On the other hand, by the equation (1.12) we have

$$u(x) \ge 4^{2\sigma - n} \int_{B_2} K(y) u(y)^{\frac{n+2\sigma}{n-2\sigma}} dy =: A_0 > 0 \quad \text{ for all } x \in B_2 \backslash \Sigma, \tag{5.19}$$

and by the definition of w_i , we obtain

$$w_j(y) \ge \frac{A_0}{u(x_j)}$$
 for $y \in \Omega_j \backslash \Pi_j$. (5.20)

Therefore, for any $\tau \in (0, 1/4)$ there exists j_0 such that for all $j \geq j_0$ we have

$$w_j(z)^{\frac{n+2\sigma}{n-2\sigma}} - w_j^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \ge \frac{1}{2} w_j(z)^{\frac{n+2\sigma}{n-2\sigma}} \quad \text{in } \Omega_j \backslash \Pi_j.$$

As in Lemma 4.4, we still have for $\lambda_0 \le \lambda \le |y| \le \lambda_1 + 1$ and $|z| \ge \lambda_1 + 2$ that

$$\frac{\delta_1}{|y-z|^{n-2\sigma}}(|y|-\lambda) \le G(0,\lambda;y,z) \le \frac{\delta_2}{|y-z|^{n-2\sigma}}(|y|-\lambda)$$
(5.21)

where $\delta_1, \delta_2 > 0$ depend only on n and σ . Moreover, we can verify that for $|y| \ge \lambda_1 + 1$ and $|z| \ge \lambda_1 + 2$,

$$\frac{\delta_3}{|y-z|^{n-2\sigma}} \le G(0,\lambda;y,z) \le \frac{1}{|y-z|^{n-2\sigma}}$$
 (5.22)

for some $\delta_3 \in (0,1)$ depending only on n and σ .

Denote $\rho_j := u(x_j)^{\frac{2}{n-2\sigma}}$ and $A_1 := \min_{B_2} K > 0$. Then, for $\lambda \le |y| \le \lambda_1 + 1$ and for large j, we have

$$J_{\lambda}(y) \geq \frac{A_1}{2} \left(\frac{A_0}{u(x_j)}\right)^{\frac{n+2\sigma}{n-2\sigma}} \int_{\Omega_j \backslash \Pi_j} \frac{\delta_1}{|y-z|^{n-2\sigma}} (|y|-\lambda) dz$$

$$-C \int_{\Omega_j^c} \frac{\delta_2}{|y-z|^{n-2\sigma}} (|y|-\lambda) \left(\frac{\lambda}{|z|}\right)^{n+2\sigma} dz$$

$$\geq \frac{A_1 A_0^{\frac{n+2\sigma}{n-2\sigma}}}{2} (|y|-\lambda) u(x_j)^{-\frac{n+2\sigma}{n-2\sigma}} \int_{\left\{\frac{3\tau}{2}\rho_j \leq |z| \leq \frac{7}{4}\rho_j\right\} \backslash \Sigma_j} \frac{1}{|y-z|^{n-2\sigma}} dz$$

$$-C(|y|-\lambda) \int_{\left\{|z| \geq \frac{7}{4}\rho_j\right\} \cup \Sigma_j} \frac{1}{|y-z|^{n-2\sigma}} \left(\frac{1}{|z|}\right)^{n+2\sigma} dz$$

$$\geq C(|y|-\lambda) u(x_j)^{-1} - C(|y|-\lambda) u(x_j)^{-\frac{2n}{n-2\sigma}}$$

$$\geq \beta_0(|y|-\lambda) u(x_j)^{-1}$$

for some constant $\beta_0 > 0$ independent of τ , where we have used $\mathcal{L}^n(\Sigma_j) = 0$ and $u(x_j) \to \infty$. Similarly, for $|y| \ge \lambda_1 + 1$ and $y \in \Pi_j$, we have

$$J_{\lambda}(y) \ge Cu(x_j)^{-1} - Cu(x_j)^{-\frac{2n}{n-2\sigma}} \ge \beta_0 u(x_j)^{-1}$$

for another constant $\beta_0 > 0$ independent of τ . Lemma 5.2 is established.

To estimate Φ_{λ} with $\lambda \in [1/2, 2]$, we denote

$$Q_{\lambda}(z) = \left(K_{j}(z^{\lambda}) - K_{j}(z)\right) w_{j}^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \quad \text{for } z \in \Omega_{j} \backslash B_{\lambda}.$$
 (5.23)

Since $w_j(y)$ converges in $C^2_{loc}(\mathbb{R}^n)$ to $U_0(y)=(1+|y|^2)^{-\frac{n-2\sigma}{2}}$, we have for large j that

$$0 \le w_j^{\lambda}(z) \le 2\left(\frac{\lambda}{|z|}\right)^{n-2\sigma} \left(\frac{1}{1+|z^{\lambda}|^2}\right)^{\frac{n-2\sigma}{2}} \le C|z|^{2\sigma-n} \quad \text{for } z \in \Omega_j \backslash B_{\lambda}, \tag{5.24}$$

where C > 0 depends only on n and σ . In order to describe the behavior of Q_{λ} on $\Omega_j \backslash B_{\lambda}$ we define the following set

$$\mathcal{P}_j := \left\{ y \in \mathbb{R}^n : x_j + M_j^{-\frac{2}{n-2\sigma}} y \in B_{2\tau} \backslash \Sigma \right\} \subset \Omega_j.$$
 (5.25)

We next complete the proof of Theorem 1.6 under the assumption (K2) in three steps.

Step 3. We prove Theorem 1.6 under the assumption (K2) when $2\sigma < n < 2\sigma + 2$. By the assumption on K and (5.24) we have

$$|Q_{\lambda}(z)| \le \begin{cases} c(\tau)M_{j}^{-1}(|z| - \lambda)^{\alpha}|z|^{-2\sigma - n} & \text{for } z \in \mathcal{P}_{j} \backslash B_{\lambda}, \\ CM_{j}^{-1}|z|^{\alpha - 2\sigma - n} & \text{for } z \in \Omega_{j} \backslash \mathcal{P}_{j}, \end{cases}$$
(5.26)

where $c(\cdot)$ is a nonnegative continuous function with c(0)=0. Recall also that $\alpha=\frac{n-2\sigma}{2}$. By the definition of Q_{λ} , Φ_{λ} given in (5.15) can be written as

$$\Phi_{\lambda}(y) = \int_{\Omega_{j} \backslash B_{\lambda}} G(0, \lambda; y, z) Q_{\lambda}(z) dz.$$
 (5.27)

We have the following estimates for Φ_{λ} .

Lemma 5.3. For any $\lambda \in [1/2, 2]$, we have

$$|\Phi_{\lambda}(y)| \leq \begin{cases} \left(c(\tau)M_{j}^{-1} + \tau^{-\frac{n+2\sigma}{2}}M_{j}^{-1-\frac{n+2\sigma}{n-2\sigma}}\right)(|y| - \lambda) & \text{for } y \in B_{4\lambda} \backslash B_{\lambda}, \\ c(\tau)M_{j}^{-1}(|y|^{\alpha-n} + |y|^{2\sigma-n}\log|y|) + C\tau^{-\frac{n+2\sigma}{2}}M_{j}^{-1-\frac{n+2\sigma}{n-2\sigma}} & \text{for } y \in \Pi_{j} \backslash B_{4\lambda}, \end{cases}$$
(5.28)

where $c(\cdot)$ is a nonnegative continuous function satisfying c(0) = 0 and C > 0 is a constant depending only on n, σ and K.

Proof. It follows from (5.26) that

$$|\Phi_{\lambda}(y)| \le c(\tau) M_j^{-1} \int_{\mathcal{P}_j \setminus B_{\lambda}} G(0,\lambda;y,z) (|z| - \lambda)^{\alpha} |z|^{-2\sigma - n} dz$$

$$+ C M_j^{-1} \int_{\Omega_j \setminus \mathcal{P}_j} G(0,\lambda;y,z) |z|^{\alpha - 2\sigma - n} dz.$$
(5.29)

We consider the following two cases separately.

Case 1: $\lambda < |y| < 4\lambda$. As in the proof of Case 1 of Lemma 4.3, we consider three sets

$$\mathcal{A}_1 = \{ z \in \mathcal{P}_j \backslash B_\lambda : |z - y| < (|y| - \lambda)/3 \},$$

$$\mathcal{A}_2 = \{ z \in \mathcal{P}_j \backslash B_{\lambda} : |z - y| \ge (|y| - \lambda)/3 \text{ and } |z| \le 8\lambda \},$$

$$\mathcal{A}_3 = \{ z \in \mathcal{P}_j \backslash B_{\lambda} : |z| \ge 8\lambda \}.$$

Then, from Lemma 3.1 we have

$$G(0,\lambda;y,z) \le \begin{cases} C|y-z|^{2\sigma-n} & \text{if } z \in \mathcal{A}_{1}, \\ C\frac{(|y|-\lambda)(|z|^{2}-\lambda^{2})}{\lambda|y-z|^{n-2\sigma+2}} \le C\frac{(|y|-\lambda)(|z|-\lambda)}{|y-z|^{n-2\sigma+2}} & \text{if } z \in \mathcal{A}_{2}, \\ C\frac{(|y|-\lambda)(|z|^{2}-\lambda^{2})}{\lambda|y-z|^{n-2\sigma+2}} \le C(|y|-\lambda)|z|^{2\sigma-n} & \text{if } z \in \mathcal{A}_{3}. \end{cases}$$
(5.30)

Thus,

$$\int_{\mathcal{A}_1} G(0,\lambda;y,z)(|z|-\lambda)^{\alpha}|z|^{-2\sigma-n}dz \le C \int_{\mathcal{A}_1} |y-z|^{2\sigma-n}(|z|-\lambda)^{\alpha}|z|^{-2\sigma-n}dz$$

$$\le C(|y|-\lambda),$$

where we used $|z| - \lambda \le 4(|y| - \lambda)/3$ for $z \in \mathcal{A}_1$ and $\alpha + 2\sigma = \frac{n+2\sigma}{2} > 1$. For the integral over \mathcal{A}_2 , we have

$$\int_{\mathcal{A}_2} G(0,\lambda;y,z) (|z| - \lambda)^{\alpha} |z|^{-2\sigma - n} dz \le C \int_{\mathcal{A}_2} \frac{(|y| - \lambda)(|z| - \lambda)^{1+\alpha}}{|y - z|^{n-2\sigma + 2}} |z|^{-2\sigma - n} dz
\le C(|y| - \lambda),$$

where we used $|z| - \lambda \le 4|z-y|$ for $z \in A_2$ and $\alpha + 2\sigma > 1$. For the integral over A_3 , we have

$$\int_{\mathcal{A}_3} G(0,\lambda;y,z)(|z|-\lambda)^{\alpha}|z|^{-2\sigma-n}dz \le C(|y|-\lambda)\int_{\mathcal{A}_3} |z|^{\alpha-2n}dz$$

$$\le C(|y|-\lambda).$$

These estimates give that

$$\int_{\mathcal{P}_i \setminus B_\lambda} G(0,\lambda;y,z)(|z|-\lambda)^\alpha |z|^{-2\sigma-n} dz \le C(|y|-\lambda)$$
(5.31)

for some constant C > 0 depending only on n and σ . On the other hand, by Lemma 3.1 we obtain for large j that

$$\int_{\Omega_{j}\backslash \mathcal{P}_{j}} G(0,\lambda;y,z)|z|^{\alpha-2\sigma-n} dz \leq C(|y|-\lambda) \int_{|z|\geq \tau M_{j}^{\frac{2}{n-2\sigma}}} |z|^{\alpha-2n} dz$$

$$\leq C\tau^{-\frac{n+2\sigma}{2}} M_{j}^{-\frac{n+2\sigma}{n-2\sigma}} (|y|-\lambda). \tag{5.32}$$

This together with (5.29) and (5.31) implies that Lemma 5.3 holds for $y \in B_{4\lambda} \backslash B_{\lambda}$.

Case 2: $y \in \Pi_i \backslash B_{4\lambda}$. As in the proof of Case 2 of Lemma 4.3, we define four sets:

$$S_1 = \{ z \in \mathcal{P}_j \backslash B_\lambda : |z| < |y|/2 \},$$

$$S_2 = \{ z \in \mathcal{P}_i \backslash B_\lambda : |z| > 2|y| \},$$

$$\mathcal{S}_3 = \{z \in \mathcal{P}_j \backslash B_\lambda : |z - y| \le |y|/2\},$$

$$\mathcal{S}_4 = \{z \in \mathcal{P}_j \backslash B_\lambda : |z - y| \ge |y|/2 \text{ and } |y|/2 \le |z| \le 2|y|\}.$$

Notice that $G(0, \lambda; y, z) \leq C|y-z|^{2\sigma-n}$. By direct calculations we can get the following estimates:

$$\int_{\mathcal{S}_1} |y-z|^{2\sigma-n} |z|^{\alpha-2\sigma-n} dz \le \begin{cases} C|y|^{\alpha-n} & \text{if } \alpha > 2\sigma, \\ C|y|^{2\sigma-n} \log |y| & \text{if } \alpha = 2\sigma, \\ C|y|^{2\sigma-n} & \text{if } \alpha < 2\sigma, \end{cases}$$

$$\int_{\mathcal{S}_2} |y-z|^{2\sigma-n} |z|^{\alpha-2\sigma-n} dz \le C|y|^{\alpha-n},$$

$$\int_{\mathcal{S}_3} |y-z|^{2\sigma-n} |z|^{\alpha-2\sigma-n} dz \le C|y|^{\alpha-2\sigma-n} \int_{\mathcal{S}_3} |y-z|^{2\sigma-n} dz \le C|y|^{\alpha-n}$$

and

$$\int_{\mathcal{S}_4} |y-z|^{2\sigma-n} |z|^{\alpha-2\sigma-n} dz \leq C |y|^{\alpha-n}.$$

Therefore, we have

$$\int_{\mathcal{P}_j \setminus B_{\lambda}} G(0,\lambda;y,z)(|z|-\lambda)^{\alpha} |z|^{-2\sigma-n} dz \le C(|y|^{\alpha-n} + |y|^{2\sigma-n} \log|y|). \tag{5.33}$$

On the other hand, for $y \in \Pi_j \backslash B_\lambda$ we have $|y| \leq \frac{5\tau}{4} M_j^{\frac{2}{n-2\sigma}}$ and for $z \in \Omega_j \backslash \mathcal{P}_j$ we have $|z| \geq \frac{7\tau}{4} M_j^{\frac{2}{n-2\sigma}}$ when j is large. Thus,

$$\int_{\Omega_{j} \setminus \mathcal{P}_{j}} G(0,\lambda;y,z) |z|^{\alpha - 2\sigma - n} dz \le C \int_{|z| \ge \tau M_{j}^{\frac{2}{n - 2\sigma}}} |z|^{\alpha - 2n} dz \le C \tau^{-\frac{n + 2\sigma}{2}} M_{j}^{-\frac{n + 2\sigma}{n - 2\sigma}}.$$
 (5.34)

This together with (5.29) and (5.33) implies that Lemma 5.3 holds for $y \in \Pi_j \backslash B_{4\lambda}$. Thus the proof of Lemma 5.3 is finished.

For $\lambda \in [1/2, 2]$, we construct H_{λ} as

$$H_{\lambda}(y) = -\varepsilon_1 M_j^{-1} (\lambda^{2\sigma - n} - |y|^{2\sigma - n}) + \Phi_{\lambda}(y) \quad \text{for } y \in \Pi_j \backslash \overline{B}_{\lambda}. \tag{5.35}$$

Here ε_1 is chosen so that $0 < \varepsilon_1 \ll \varepsilon_0$ and

$$-\varepsilon_1 M_j^{-1}(\lambda^{2\sigma-n} - |y|^{2\sigma-n}) + J_{\lambda}(y) \ge \frac{1}{2} J_{\lambda}(y) \quad \text{for } y \in \Pi_j \backslash \overline{B}_{\lambda}, \tag{5.36}$$

where ε_0 is defined in Lemma 5.1. It follows from Lemma 5.2 that (5.36) is true by taking sufficiently small ε_1 . Notice that when τ is small, $c(\tau)$ is correspondingly small. Hence, by Lemma 5.3 we can choose τ to be small enough such that for any $\lambda \in [1/2, 2]$

$$H_{\lambda} < 0 \quad \text{in } \Pi_{i} \backslash \overline{B}_{\lambda}$$
 (5.37)

when j is sufficiently large. Moreover, by (5.17) and (5.36) we know that $\varphi_{\lambda} + H_{\lambda}$ satisfies the following integral inequality

$$\varphi_{\lambda}(y) + H_{\lambda}(y) \ge \int_{\Pi_{j} \backslash B_{\lambda}} G(0, \lambda; y, z) b_{\lambda}(z) \varphi_{\lambda}(z) dz + \frac{1}{2} J_{\lambda}(y), \quad y \in \Pi_{j} \backslash \overline{B}_{\lambda}$$
 (5.38)

with $\lambda \in [1/2, 2]$.

Now we apply the method of moving spheres to $\varphi_{\lambda} + H_{\lambda}$ for $\lambda \in [\lambda_0, \lambda_1]$ where $\lambda_0 = 1/2$ and $\lambda_1 = 2$. First, when $\lambda = \lambda_0$, by Lemmas 5.1 and 5.3 we have $\varphi_{\lambda_0} + H_{\lambda_0} \ge 0$ in $\Pi_j \backslash \overline{B}_{\lambda_0}$ for all large j. Define

$$\bar{\lambda} := \sup\{\mu \ge \lambda_0 \mid \varphi_{\lambda}(y) + H_{\lambda}(y) \ge 0, \ \forall \ y \in \Pi_j \setminus \overline{B}_{\lambda}, \ \forall \ \lambda_0 \le \lambda \le \mu\}.$$

Then, $\bar{\lambda}$ is well defined and $\bar{\lambda} \geq \lambda_0$ for all large j. Furthermore, by (5.37) and Lemma 5.1 we see that $\bar{\lambda} < \lambda_1$ for all large j. Then, as in Step 2 of Section 4, applying the moving sphere method one can derive a contradiction to the definition of $\bar{\lambda}$. Step 3 is established.

Step 4. We prove Theorem 1.6 under the assumption (K2) when $n=2\sigma+2$. In this case, $K \in C^1(B_2)$ and thus by Step 2 we have $\nabla K(0)=0$. For simplicity, we may assume $K \in C^1(\overline{B_2})$. Based on these properties of K we have for $\lambda \in [1/2,2]$ that

$$|K(x_j + M_j^{-\frac{2}{n-2\sigma}} z^{\lambda}) - K(x_j + M_j^{-\frac{2}{n-2\sigma}} z)| \le \begin{cases} c(\tau) M_j^{-1} (|z| - \lambda) & \text{for } z \in \mathcal{P}_j \backslash B_{\lambda}, \\ CM_j^{-1} |z| & \text{for } z \in \Omega_j \backslash \mathcal{P}_j. \end{cases}$$
(5.39)

Thus, by (5.24) we get

$$|Q_{\lambda}(z)| \le \begin{cases} c(\tau)M_{j}^{-1}(|z| - \lambda)|z|^{-2\sigma - n} & \text{for } z \in \mathcal{P}_{j} \backslash B_{\lambda}, \\ CM_{j}^{-1}|z|^{1 - 2\sigma - n} & \text{for } z \in \Omega_{j} \backslash \mathcal{P}_{j}, \end{cases}$$
(5.40)

where $c(\cdot)$ is a nonnegative continuous function with c(0) = 0. This indicates that Lemma 5.3 is still true. The rest of the proof of this step is the same as that of Step 3. Step 4 is established.

Step 5. We prove Theorem 1.6 under the assumption (K2) when $n>2\sigma+2$. Under the current case, our assumption is $K\in\mathcal{C}^{\alpha}(B_2)$ with $\alpha=\frac{n-2\sigma}{2}>1$. In particular, this implies that $K\in\mathcal{C}^1(B_2)$ and, by Step 2 we obtain $\nabla K(0)=0$. Thus 0 is a critical point of K, for $n\geq 2\sigma+4$, by the assumption on K there exists a neighborhood N of 0 such that (1.15) holds. Without loss of generality we may assume $N=B_2$, since otherwise we can consider the integral equation (1.12) on N and then make a rescaling argument. Namely, for $n\geq 2\sigma+4$ we assume that K satisfies

$$|\nabla^{i}K(y)| \le c(|y|)|\nabla K(y)|^{\frac{\alpha-i}{\alpha-1}}, \quad 2 \le i \le [\alpha], \quad y \in B_2$$

$$(5.41)$$

for some nonnegative continuous function $c(\cdot)$ with c(0) = 0.

Let $Z_j = |\nabla K(x_j)|$, then $\lim_{j\to\infty} Z_j = |\nabla K(0)| = 0$. Moreover, we have the following result.

Lemma 5.4.

$$Z_j^{\frac{1}{\alpha - 1}} M_j^{\frac{2}{n - 2\sigma}} \to 0.$$
 (5.42)

Proof. We first prove that there exists a constant C > 0 such that

$$Z_{j}^{\frac{1}{\alpha-1}} M_{j}^{\frac{2}{n-2\sigma}} \le C \quad \text{for all } j.$$
 (5.43)

If not, then up to a subsequence

$$Z_j^{\frac{1}{n-1}} M_j^{\frac{2}{n-2\sigma}} \to +\infty.$$
 (5.44)

The following proof is similar to that of Step 2 in Section 4, we mainly show the modifications that need to be made. Without loss of generality, we may assume

$$\lim_{j \to \infty} \frac{\nabla K(x_j)}{|\nabla K(x_j)|} = e = (1, 0, \dots, 0).$$
 (5.45)

As in Step 2 of Section 4, we define w_j and h_j as in (4.17) and define Ω_j as in (4.18). Then w_j still satisfies (4.21) on Ω_j where K_j is defined as in (4.20). Moreover, on every compact subset of \mathbb{R}^n , w_j converges in C^2 norm to $U_1 = U_0(\cdot - Re)$ by Step 1. We also extend w_j to be identically 0 outside Ω_j and K to be identically 0 outside B_2 .

Define w_j^{λ} and h_j^{λ} as in (4.19), then w_j^{λ} satisfies (4.22). It is clear that Lemma 4.1 still holds. Let $\varphi_{\lambda}(y) = w_j(y) - w_j^{\lambda}(y)$ with $\lambda \in [R-2,R+2]$. Recall that R>10 is a large positive constant to be determined later. Then φ_{λ} satisfies the following integral inequality for large j,

$$\varphi_{\lambda}(y) + \Phi_{\lambda}(y) \ge \int_{\Pi_{j} \backslash B_{\lambda}} G(0, \lambda; y, z) b_{\lambda}(z) \varphi_{\lambda}(z) dz + J_{\lambda}(y), \quad y \in \Pi_{j} \backslash \overline{B}_{\lambda}, \tag{5.46}$$

where Π_i , b_λ , Φ_λ and J_λ are respectively given as in (4.32), (4.36), (4.37) and (4.38).

Now we need to establish some estimates for Φ_{λ} on $\Pi_{j} \backslash \overline{B}_{\lambda}$. Define a special domain D_{λ} in $\Omega_{j} \backslash B_{\lambda}$ as follows,

$$D_{\lambda} = \{z \in \mathbb{R}^n : \lambda < |z| < 2\lambda, z_1 > 2|z'| \text{ where } z' = (z_2, \dots, z_n)\}.$$

Let $\tilde{l}_j := Z_j^{\frac{1}{\alpha-1}} M_j^{\frac{2}{n-2\sigma}}$, then by (5.44) we know $\tilde{l}_j \to +\infty$ and so $\tilde{l}_j \gg R$ when j is large. Under our current assumptions on K, we have

Claim 1: There exists a constant $C_0 > 0$ independent of λ such that for all large j,

$$K_j(z^{\lambda}) - K_j(z) \le -C_0 M_j^{-\frac{2}{n-2\sigma}} |\nabla K(x_j)| (|z| - \lambda), \quad z \in D_{\lambda}, \tag{5.47}$$

$$|K_j(z^{\lambda}) - K_j(z)| \le C_0 M_j^{-\frac{2}{n-2\sigma}} |\nabla K(x_j)| (|z| - \lambda), \quad z \in B_{2\tilde{l}_j} \setminus \overline{B}_{\lambda}, \tag{5.48}$$

$$|K_j(z^{\lambda}) - K_j(z)| \le C_0 M_j^{-1} |z|^{\alpha}, \quad z \in \Omega_j \backslash B_{2\tilde{l}_j}.$$
(5.49)

Proof. (1) For $z \in D_{\lambda}$ and $2\sigma + 2 < n < 2\sigma + 4$, by the mean value theorem and the assumptions on K we obtain that

$$K_{j}(z^{\lambda}) - K_{j}(z) = M_{j}^{-\frac{2}{n-2\sigma}} \nabla K \left(x_{j} + M_{j}^{-\frac{2}{n-2\sigma}} \left((1-\theta)z^{\lambda} + \theta z - Re \right) \right) \cdot (z^{\lambda} - z)$$

$$\leq M_{j}^{-\frac{2}{n-2\sigma}} \nabla K(x_{j}) \cdot (z^{\lambda} - z) + \sup_{0 \leq t \leq Z_{j}^{\frac{1}{\alpha}-1}} c(t) M_{j}^{-\frac{2}{n-2\sigma}} |\nabla K(x_{j})| |z^{\lambda} - z|$$

$$\leq -C_{0} M_{j}^{-\frac{2}{n-2\sigma}} |\nabla K(x_{j})| (|z| - \lambda)$$

for all large j, where $\theta \in (0,1)$ and $C_0 > 0$ is a constant.

For $z \in D_{\lambda}$ and $n \ge 2\sigma + 4$, by the mean value theorem and the assumptions on K we obtain

$$K_{j}(z^{\lambda}) - K_{j}(z) = M_{j}^{-\frac{2}{n-2\sigma}} \nabla K \left(x_{j} + M_{j}^{-\frac{2}{n-2\sigma}} \left((1-\theta)z^{\lambda} + \theta z - Re \right) \right) \cdot (z^{\lambda} - z)$$

$$\leq M_{j}^{-\frac{2}{n-2\sigma}} \nabla K(x_{j}) \cdot (z^{\lambda} - z) + C_{1} \sum_{i=2}^{[\alpha]} M_{j}^{-\frac{2i}{n-2\sigma}} |\nabla^{i}K(x_{j})| |z|^{i-1} |z^{\lambda} - z|$$

$$+ \sup_{0 \leq t \leq |x_{j}| + Z_{j}^{\frac{1}{\alpha} - 1}} c(t) M_{j}^{-\frac{2\alpha}{n-2\sigma}} |z|^{\alpha - 1} |z^{\lambda} - z|$$

$$\leq -C_{1} M_{j}^{-\frac{2}{n-2\sigma}} |\nabla K(x_{j})| (|z| - \lambda)$$

$$+ c(|x_{j}|) \sum_{i=2}^{[\alpha]} M_{j}^{-\frac{2i}{n-2\sigma}} |\nabla K(x_{j})|^{\frac{\alpha - i}{\alpha - 1}} |z|^{i-1} |z^{\lambda} - z|$$

$$\leq -C_{0} M_{j}^{-\frac{2}{n-2\sigma}} |\nabla K(x_{j})| (|z| - \lambda)$$

for all large j, where $\theta \in (0,1)$, $C_0, C_1, C_2 > 0$ are different constants.

(2) When $z \in B_{2\tilde{l}_i} \setminus \overline{B}_{\lambda}$, the proof is similar to that of (1), and we only need to notice that $M_j^{-\frac{2}{n-2\sigma}}|\nabla K(x_j)\cdot(z^{\lambda}-z)| \leq CM_j^{-\frac{2}{n-2\sigma}}|\nabla K(x_j)|(|z|-\lambda).$ (3) For $z\in\Omega_j\backslash B_{2\tilde{l}_j}$ and $2\sigma+2< n<2\sigma+4$, by the mean value theorem and the assumptions

on K we obtain that

$$|K_{j}(z^{\lambda}) - K_{j}(z)| \leq M_{j}^{-\frac{2}{n-2\sigma}} |\nabla K(x_{j})| (|z| - \lambda) + CM_{j}^{-\frac{2\alpha}{n-2\sigma}} |z|^{\alpha-1} (|z| - \lambda)$$

$$\leq C_{0} M_{j}^{-1} |z|^{\alpha}$$

for some constant $C_0 > 0$.

For $z\in\Omega_j\backslash B_{2\tilde{l}_j}$ and $n\geq 2\sigma+4$, by the mean value theorem and the assumptions on K we

$$|K_{j}(z^{\lambda}) - K_{j}(z)| \leq M_{j}^{-\frac{2}{n-2\sigma}} |\nabla K(x_{j})| (|z| - \lambda) + C \sum_{i=2}^{[\alpha]} M_{j}^{-\frac{2i}{n-2\sigma}} |\nabla^{i} K(x_{j})| |z|^{i-1} |z^{\lambda} - z|$$

$$+ C M_{j}^{-\frac{2\alpha}{n-2\sigma}} |z|^{\alpha-1} |z^{\lambda} - z|$$

$$\leq C_{0} M_{j}^{-1} |z|^{\alpha}$$

for some constant C_0 . Thus Claim 1 is proved.

It is also clear that Lemma 4.2 still holds. If we let

$$Q_{\lambda}(z) = \left(K_j(z^{\lambda}) - K_j(z) \right) w_j^{\lambda}(z)^{\frac{n+2\sigma}{n-2\sigma}} \quad \text{for } z \in \Omega_j \backslash B_{\lambda},$$

then by Claim 1 above and Lemma 4.2 we have

$$Q_{\lambda}(z) \le -CM_{j}^{-\frac{2}{n-2\sigma}} |\nabla K(x_{j})| (|z| - \lambda) \left(\frac{1}{1 + (z_{1} - \lambda)^{2} + |z'|^{2}} \right)^{\frac{n+2\sigma}{2}} \quad \text{for } z \in D_{\lambda} \quad (5.50)$$

and

$$|Q_{\lambda}(z)| \leq \begin{cases} CM_{j}^{-\frac{2}{n-2\sigma}} |\nabla K(x_{j})|(|z|-\lambda) & \text{for } z \in D_{\lambda}, \\ CM_{j}^{-\frac{2}{n-2\sigma}} |\nabla K(x_{j})|(|z|-\lambda)|z|^{-2\sigma-n} & \text{for } z \in B_{2\tilde{l}_{j}} \setminus (B_{\lambda} \cup D_{\lambda}), \\ CM_{j}^{-1} |z|^{\alpha-2\sigma-n} & \text{for } z \in \Omega_{j} \setminus B_{2\tilde{l}_{j}}. \end{cases}$$
(5.51)

We split $\Phi_{\lambda}(y)$ into two parts:

$$\Phi_{\lambda}(y) = \int_{B_{2\tilde{l}_{j}} \backslash B_{\lambda}} G(0, \lambda; y, z) Q_{\lambda}(z) dz + \int_{\Omega_{j} \backslash B_{2\tilde{l}_{j}}} G(0, \lambda; y, z) Q_{\lambda}(z) dz$$

$$=: \Phi_{1,\lambda}(y) + \Phi_{2,\lambda}(y). \tag{5.52}$$

By the estimates of $Q_{\lambda}(z)$ on $B_{2\tilde{l}_j}\backslash B_{\lambda}$, one can see easily that the estimates of $\Phi_{1,\lambda}$ are very similar to those of Φ_{λ} in Lemma 4.3 of Step 2 of Section 4. Hence we have the following:

Claim 2: There exists a constant C > 0 independent of λ such that for all large j,

$$\Phi_{1,\lambda}(y) \leq \begin{cases} -CM_j^{-\frac{2}{n-2\sigma}} |\nabla K(x_j)| (|y|-\lambda)\lambda^{-n} \log \lambda & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ -CM_j^{-\frac{2}{n-2\sigma}} |\nabla K(x_j)| |y|^{2\sigma-n} \lambda^{1-2\sigma} \log \lambda & \text{for } y \in \Pi_j \backslash \overline{B}_{4\lambda} \end{cases}$$

and

$$|\Phi_{1,\lambda}(y)| \leq \begin{cases} CM_j^{-\frac{2}{n-2\sigma}} |\nabla K(x_j)| (|y|-\lambda) \lambda^{2\sigma} & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ CM_j^{-\frac{2}{n-2\sigma}} |\nabla K(x_j)| |y|^{2\sigma-n} \lambda^{n+1} & \text{for } y \in \Pi_j \backslash \overline{B}_{4\lambda}. \end{cases}$$

Proof. We only need to replace $M_j^{-\frac{2}{n-2\sigma}}$ and $\Omega_j \backslash B_\lambda$ in Lemma 4.3 by $M_j^{-\frac{2}{n-2\sigma}} |\nabla K(x_j)|$ and $B_{2\tilde{l}_i} \backslash B_\lambda$, respectively.

Recall that under the current assumptions we have $1 < \alpha = (n-2\sigma)/2 < n$. For the estimates of $\Phi_{2,\lambda}$, by an argument similar to Lemma 4.6 of Section 4 we have

Claim 3: There exists a constant C > 0 independent of λ such that for all large j,

$$|\Phi_{2,\lambda}(y)| \leq \begin{cases} CM_j^{-1} \tilde{l}_j^{\alpha-n} (|y| - \lambda) \lambda^{-1} & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ CM_j^{-1} \tilde{l}_j^{\alpha-n} & \text{for } y \in \Pi_j \backslash B_{4\lambda}, \ \alpha \neq 2\sigma \\ CM_j^{-1} \tilde{l}_j^{2\sigma-n} \log \tilde{l}_j & \text{for } y \in \Pi_j \backslash B_{4\lambda}, \ \alpha = 2\sigma. \end{cases}$$

It is clear that Lemma 4.1 and Lemma 4.4 still hold. We construct a function H_{λ} which is non-positive on $\Pi_i \setminus \overline{B}_{\lambda}$. Let

$$H_{\lambda}(y) = -\varepsilon_1 M_j^{-1} (\lambda^{2\sigma - n} - |y|^{2\sigma - n}) + \Phi_{\lambda}(y), \quad y \in \Pi_j \backslash \overline{B}_{\lambda}$$
 (5.53)

for some small $\varepsilon_1 \in (0, \varepsilon_0/4)$ where ε_0 is defined in Lemma 4.1. By Lemma 4.4 we can choose $\varepsilon_1 > 0$ sufficiently small so that for any $\lambda \in [\lambda_0, \lambda_1]$,

$$J_{\lambda}(y) - \varepsilon_1 M_j^{-1}(\lambda^{2\sigma - n} - |y|^{2\sigma - n}) \ge \frac{1}{2} J_{\lambda}(y), \quad y \in \Pi_j \backslash \overline{B}_{\lambda}.$$
 (5.54)

Here we recall that $\lambda_0 = R - 2$ and $\lambda_1 = R + 2$. Then by Claim 2 and Claim 3 above we have for $\lambda \in [\lambda_0, \lambda_1]$ and for large j,

$$H_{\lambda}(y) < 0, \quad y \in \Pi_{i} \backslash \overline{B}_{\lambda}.$$
 (5.55)

Moreover, it follows from Lemma 4.1, Claim 2 and Claim 3 above that for large j,

$$\varphi_{\lambda_0}(y) + H_{\lambda_0}(y) \ge 0, \quad y \in \Pi_j \setminus \overline{B}_{\lambda_0}.$$

By (5.46) and (5.54), $\varphi_{\lambda} + H_{\lambda}$ satisfies

$$\varphi_{\lambda}(y) + H_{\lambda}(y) \ge \int_{\Pi_{j} \setminus B_{\lambda}} G(0, \lambda; y, z) b_{\lambda}(z) \varphi_{\lambda}(z) dz + \frac{1}{2} J_{\lambda}(y) \quad \text{for } y \in \Pi_{j} \setminus \overline{B}_{\lambda}, \quad (5.56)$$

where $\lambda \in [\lambda_0, \lambda_1]$. We define

$$\bar{\lambda} := \sup\{\mu \geq \lambda_0 \mid \varphi_{\lambda}(y) + H_{\lambda}(y) \geq 0, \ \forall \ y \in \Pi_i \setminus \overline{B}_{\lambda}, \ \forall \ \lambda_0 \leq \lambda \leq \mu\}.$$

Then, $\bar{\lambda}$ is well defined and $\bar{\lambda} \geq \lambda_0$ for all large j. Furthermore, by (5.55) and Lemma 4.1 we have $\bar{\lambda} < \lambda_1$ for all large j. Then, as in Step 2 of Section 4, applying the moving sphere method one can derive a contradiction to the definition of $\bar{\lambda}$. This proves that the sequence $Z_j^{\frac{1}{\alpha-1}} M_j^{\frac{2}{n-2\sigma}}$ is bounded.

Next we show that $Z_j^{\frac{1}{\alpha-1}}M_j^{\frac{2}{n-2\sigma}}\to 0$. If this does not hold, then there exists a constant $\varepsilon_2>0$ such that, after passing to a subsequence if necessary,

$$Z_j^{\frac{1}{\alpha-1}} M_j^{\frac{2}{n-2\sigma}} \ge \varepsilon_2 > 0. \tag{5.57}$$

The following proof is similar to that of (5.43), we will use the same notation and give only the corresponding modifications. Let $l_0 \gg R$ be a large positive constant which is to be determined after fixing R. Under our current assumptions on K, similar to Claim 1 we have

Claim 4: There exists a constant C > 0 independent of both l_0 and λ such that for all large j,

$$K_j(z^{\lambda}) - K_j(z) \le -CM_j^{-1}(|z| - \lambda), \quad z \in D_{\lambda}. \tag{5.58}$$

$$|K_j(z^{\lambda}) - K_j(z)| \le CM_j^{-1}(|z| - \lambda), \quad z \in B_{2l_0} \setminus B_{\lambda}.$$
 (5.59)

$$|K_j(z^{\lambda}) - K_j(z)| \le C_0 M_j^{-1} |z|^{\alpha}, \quad z \in \Omega_j \backslash B_{2l_0}.$$
 (5.60)

We split $\Phi_{\lambda}(y)$ into two parts:

$$\Phi_{\lambda}(y) = \int_{B_{2l_0} \backslash B_{\lambda}} G(0, \lambda; y, z) Q_{\lambda}(z) dz + \int_{\Omega_j \backslash B_{2l_0}} G(0, \lambda; y, z) Q_{\lambda}(z) dz$$

$$=: \Phi_{1,\lambda}(y) + \Phi_{2,\lambda}(y). \tag{5.61}$$

For the estimates of $\Phi_{1,\lambda}$, similar to Claim 2 we have

Claim 5: There exists a constant C > 0 independent of both l_0 and λ such that for all large j,

$$\Phi_{1,\lambda}(y) \le \begin{cases} -CM_j^{-1}(|y| - \lambda)\lambda^{-n}\log\lambda & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ -CM_j^{-1}|y|^{2\sigma - n}\lambda^{1 - 2\sigma}\log\lambda & \text{for } y \in \Pi_j \backslash \overline{B}_{4\lambda} \end{cases}$$

and

$$|\Phi_{1,\lambda}(y)| \le \begin{cases} CM_j^{-1}(|y| - \lambda)\lambda^{2\sigma} & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ CM_j^{-1}|y|^{2\sigma - n}\lambda^{n+1} & \text{for } y \in \Pi_j \backslash \overline{B}_{4\lambda}. \end{cases}$$

For the estimates of $\Phi_{2,\lambda}$, similar to Claim 3 we also have

Claim 6: There exists a constant C > 0 independent of both l_0 and λ such that for all large j,

$$|\Phi_{2,\lambda}(y)| \le \begin{cases} CM_j^{-1} l_0^{\alpha - n} (|y| - \lambda) \lambda^{-1} & \text{for } y \in B_{4\lambda} \backslash \overline{B}_{\lambda}, \\ CM_j^{-1} l_0^{\alpha - n} & \text{for } y \in \Pi_j \backslash B_{4\lambda}, \ \alpha \ne 2\sigma \\ CM_j^{-1} l_0^{2\sigma - n} \log l_0 & \text{for } y \in \Pi_j \backslash B_{4\lambda}, \ \alpha = 2\sigma. \end{cases}$$

It is clear that Lemma 4.1 and Lemma 4.4 still hold. We construct a function H_{λ} which is non-positive on $\Pi_{i} \setminus \overline{B}_{\lambda}$. Let

$$H_{\lambda}(y) = -\varepsilon_1 M_i^{-1} (\lambda^{2\sigma - n} - |y|^{2\sigma - n}) + \Phi_{\lambda}(y), \quad y \in \Pi_j \backslash \overline{B}_{\lambda}$$
 (5.62)

for some small $\varepsilon_1 \in (0, \varepsilon_0/4)$ where ε_0 is defined in Lemma 4.1. By Lemma 4.4 we can choose $\varepsilon_1 > 0$ sufficiently small so that for any $\lambda \in [\lambda_0, \lambda_1]$,

$$J_{\lambda}(y) - \varepsilon_1 M_j^{-1}(\lambda^{2\sigma - n} - |y|^{2\sigma - n}) \ge \frac{1}{2} J_{\lambda}(y), \quad y \in \Pi_j \backslash \overline{B}_{\lambda}.$$
 (5.63)

Recall that $\lambda_0 = R - 2$ and $\lambda_1 = R + 2$. Then by Claim 5 and Claim 6 above we can determine l_0 to be large enough so that for $\lambda \in [\lambda_0, \lambda_1]$ and for large j,

$$H_{\lambda}(y) < 0, \quad y \in \Pi_{j} \backslash \overline{B}_{\lambda}.$$
 (5.64)

Moreover, it follows from Lemma 4.1, Claim 5 and Claim 6 above that for large j,

$$\varphi_{\lambda_0}(y) + H_{\lambda_0}(y) \ge 0, \quad y \in \Pi_j \setminus \overline{B}_{\lambda_0}.$$

Now, as in Step 2 of Section 4, the method of moving spheres can be applied to $\varphi_{\lambda} + H_{\lambda}$ for $\lambda \in [\lambda_0, \lambda_1]$ to get a contradiction. Hence the proof of Lemma 5.4 is completed.

Once Lemma 5.4 holds, then by the assumptions on K we have the following estimates for $|K(x_j+M_j^{-\frac{2}{n-2\sigma}}z^\lambda)-K(x_j+M_j^{-\frac{2}{n-2\sigma}}z)|$ in $\Omega_j\backslash B_\lambda$. Here $\lambda\in[1/2,2]$.

Case 1: $2\sigma + 2 < n < 2\sigma + 4$. Using the mean value theorem and the assumptions on K yields that for large j,

$$\begin{split} & \left| K(x_j + M_j^{-\frac{2}{n-2\sigma}} z^{\lambda}) - K(x_j + M_j^{-\frac{2}{n-2\sigma}} z) \right| \\ & \leq M_j^{-\frac{2}{n-2\sigma}} |z - z^{\lambda}| \left| \nabla K \left(x_j + M_j^{-\frac{2}{n-2\sigma}} \left((1-\theta) z^{\lambda} + \theta z \right) \right) \right| \\ & \leq \sup_{0 \leq t \leq \tilde{M}_j |z|} c(t) M_j^{-\frac{2\alpha}{n-2\sigma}} |z|^{\alpha-1} |z - z^{\lambda}| + C M_j^{-\frac{2}{n-2\sigma}} |\nabla K(x_j)| |z - z^{\lambda}| \\ & \leq \begin{cases} c(\tau) M_j^{-1} |z|^{\alpha-1} (|z| - \lambda) & \text{for } z \in \mathcal{P}_j \backslash B_{\lambda}, \\ C M_j^{-1} |z|^{\alpha} & \text{for } z \in \Omega_j \backslash \mathcal{P}_j, \end{cases} \end{split}$$

where $\theta \in (0,1)$, $\tilde{M}_j := M_j^{-\frac{2}{n-2\sigma}}$ and Lemma 5.4 was used in the last inequality.

Case 2: $n \ge 2\sigma + 4$. Similar to Case 1, by the mean value theorem, the assumptions on K and Lemma 5.4 we obtain that for large j,

$$\begin{split} & \left| K(x_{j} + M_{j}^{-\frac{2}{n-2\sigma}} z^{\lambda}) - K(x_{j} + M_{j}^{-\frac{2}{n-2\sigma}} z) \right| \\ & \leq C \sum_{i=1}^{[\alpha]} M_{j}^{-\frac{2i}{n-2\sigma}} |\nabla^{i} K(x_{j})| |z|^{i-1} |z - z^{\lambda}| + \sup_{0 \leq t \leq |x_{j}| + \tilde{M}_{j}|z|} c(t) M_{j}^{-\frac{2\alpha}{n-2\sigma}} |z|^{\alpha-1} |z - z^{\lambda}| \\ & \leq C \left(M_{j}^{-\frac{2}{n-2\sigma}} |\nabla K(x_{j})| + c(|x_{j}|) \sum_{i=2}^{[\alpha]} M_{j}^{-\frac{2i}{n-2\sigma}} |\nabla K(x_{j})|^{\frac{\alpha-i}{\alpha-1}} |z|^{i-1} \right) |z - z^{\lambda}| \\ & + \sup_{0 \leq t \leq |x_{j}| + \tilde{M}_{j}|z|} c(t) M_{j}^{-\frac{2\alpha}{n-2\sigma}} |z|^{\alpha-1} |z - z^{\lambda}| \\ & \leq \begin{cases} c(\tau) M_{j}^{-1} |z|^{\alpha-1} (|z| - \lambda) & \text{for } z \in \mathcal{P}_{j} \backslash B_{\lambda}, \\ C M_{j}^{-1} |z|^{\alpha} & \text{for } z \in \Omega_{j} \backslash \mathcal{P}_{j}, \end{cases} \end{split}$$

where $\tilde{M}_j := M_j^{-\frac{2}{n-2\sigma}}$.

Thus, by combining with (5.24) we have for any $n > 2\sigma + 2$ that

$$|Q_{\lambda}(z)| \le \begin{cases} c(\tau)M_{j}^{-1}(|z| - \lambda)|z|^{\alpha - 2\sigma - n - 1} & \text{for } z \in \mathcal{P}_{j} \backslash B_{\lambda}, \\ CM_{j}^{-1}|z|^{\alpha - 2\sigma - n} & \text{for } z \in \Omega_{j} \backslash \mathcal{P}_{j}, \end{cases}$$
(5.65)

where $c(\cdot)$ is a nonnegative continuous function with c(0) = 0. This indicates that Lemma 5.3 is still true. The rest of the proof of this step is the same as that of Step 3. Step 5 is established. Therefore, the proof of Theorem 1.6 under the assumption (K2) is completely finished.

References

[1] J. H. Andrade, J. M. do Ó, Asymptotics for singular solutions to conformally invariant fourth order systems in the punctured ball. arXiv:2003.03487.

- [2] W. Ao, H. Chan, A. DelaTorre, M. Fontelos, M. González, J. Wei, On higher dimensional singularities for the fractional Yamabe problem: A nonlocal Mazzeo-Pacard program. *Duke Math. J.*, **168** (2019) no. 17, 3297-3411.
- [3] W. Ao, A. DelaTorre, M. González, J. Wei, A gluing approach for the fractional Yamabe problem with isolated singularities. *J. Reine Angew. Math.*, **763** (2020) 25-78.
- [4] W. Ao, M. González, A. Hyder, J. Wei, Removability of singularities and maximum principles for some fractional Laplacian equations. arXiv:2001.11683.
- [5] T. P. Branson, Differential operators canonically associated to a conformal structure. *Math. Scand.*, **57** (1985) 293-345.
- [6] L. Caffarelli, B. Gidas, J. Spruck, Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth. *Comm. Pure Appl. Math.*, 42 (1989) 271-297.
- [7] L. Caffarelli, T. Jin, Y. Sire, J. Xiong, Local analysis of solutions of fractional semi-linear elliptic equations with isolated singularities. *Arch. Ration. Mech. Anal.*, **213** (2014), no.1, 245-268.
- [8] S.-Y. A. Chang, F. Hang, P. C. Yang, On a class of locally conformally flat manifolds. *Int. Math. Res. Not.*, (2004), no. 4, 185-209.
- [9] S.-Y. A. Chang, Z.-C. Han and P. C. Yang, Some remarks on the geometry of a class of locally conformally flat metrics. Geometric analysis, 37–56, Progr. Math., 333, Birkhäuser/Springer, Cham, [2020].
- [10] C.-C. Chen, C.-S. Lin, Local behavior of singular positive solutions of semilinear elliptic equations with Sobolev exponent. *Duke Math. J.*, **78** (1995) 315-334.
- [11] C.-C. Chen, C.-S. Lin, Estimates of the conformal scalar curvature equation via the method of moving planes. *Comm. Pure Appl. Math.*, **50** (1997) 971-1017.
- [12] C.-C. Chen, C.-S. Lin, Estimates of the conformal scalar curvature equation via the method of moving planes, II. *J. Differential Geom.*, **49** (1998) 115-178.
- [13] C.-C. Chen, C.-S. Lin, On the asymptotic symmetry of singular solutions of the scalar curvature equations. *Math. Ann.*, **313** (1999) 229-245.
- [14] W. Chen, C. Li, B. Ou, Classification of solutions for an integral equation. *Comm. Pure Appl. Math.*, **59** (2006) 330-343.
- [15] A. DelaTorre, M. González, Isolated singularities for a semilinear equation for the fractional Laplacian arising in conformal geometry. *Rev. Mat. Iberoam.*, **34** (2018), no. 4, 1645-1678.
- [16] X. Du, H. Yang, Local behavior of positive solutions of higher order conformally invariant equations with a singular set. arXiv:2005.11998.
- [17] X. Du, H. Yang, Large singular solutions for conformal Q-curvature equations on \mathbb{S}^n . J. Differential Equations, **280** (2021) 618-643.
- [18] A. DelaTorre, M. del Pino, M. González, J. Wei, Delaunay-type singular solutions for the fractional Yamabe problem. *Math. Ann.*, **369** (2017) 597-626.
- [19] L. C. Evans, R. Gariepy, Measure Theory and Fine Properties of Functions, Studies in Advanced Mathematics. CRC Press, Boca Raton, FL (1992).

- [20] R. Frank, T. König, Classification of positive solutions to a nonlinear biharmonic equation with critical exponent. *Anal. PDE*, **12** (2019) 1101-1113.
- [21] M. González, R. Mazzeo, Y. Sire, Singular solutions of fractional order conformal Laplacians. *J. Geom. Anal.*, **22** (2012), no. 3, 845-863.
- [22] C.R. Graham, R. Jenne, L.J. Mason, G.A.J. Sparling, Conformally invariant powers of the Laplacian I. Existence. *J. Lond. Math. Soc.*, **46** (1992) 557-565.
- [23] M. J. Gursky, A. Malchiodi, A strong maximum principle for the Paneitz operator and a non-local flow for the *Q*-curvature. *J. Eur. Math. Soc. (JEMS)*, **17** (2015), no. 9, 2137-2173.
- [24] Q. Han, X. Li, Y. Li, Asymptotic expansions of solutions of the Yamabe equation and the σ_k -Yamabe equation near isolated singular points. *Comm. Pure Appl. Math.*, (2020), https://doi.org/10.1002/cpa.21943.
- [25] Z.-C. Han, Y.Y. Li, E. V. Teixeira, Asymptotic behavior of solutions to the σ_k -Yamabe equation near isolated singularities. *Invent. Math.*, **182** (2010), no. 3, 635-684.
- [26] F. Hang, P. C. Yang, *Q*-curvature on a class of manifolds with dimension at least 5. *Comm. Pure Appl. Math.*, **69** (2016), no. 8, 1452-1491.
- [27] A. Hyder, Y. Sire, Singular solutions for the constant *Q*-curvature problem. *J. Funct. Anal.*, **280** (2021), no. 3, 108819, 39 pp.
- [28] T. Jin, O. de Queiroz, Y. Sire, J. Xiong, On local behavior of singular positive solutions to nonlocal elliptic equations. *Calc. Var. Partial Differential Equations*, **56** (2017), no. 1, Art. 9, 25 pp.
- [29] T. Jin, Y.Y. Li, J. Xiong, The Nirenberg problem and its generalizations: A unified approach. *Math. Ann.*, **369** (2017), no,1-2, 109-151.
- [30] T. Jin, J. Xiong, Asymptotic symmetry and local behavior of solutions of higher order conformally invariant equations with isolated singularities. *Ann. Inst. Henri Poincaré Anal. Non Linéaire*, (2020). https://doi.org/10.1016/j.anihpc.2020.10.005.
- [31] A. Käenmäki, J. Lehrbäck, M. Vuorinen, Dimensions, Whitney covers, and tubular neighborhoods. *Indiana Univ. Math. J.*, **62** (2013), no. 6, 1861-1889.
- [32] N. Korevaar, R. Mazzeo, F. Pacard, R. Schoen, Refined asymptotics for constant scalar curvature metrics with isolated singularities. *Invent. Math.*, **135** (2) (1999) 233-272.
- [33] C. Li, Local asymptotic symmetry of singular solutions to nonlinear elliptic equations. *Invent. Math.*, **123** (1996) 221-231.
- [34] Y.Y. Li, Conformally invariant fully nonlinear elliptic equations and isolated singularities. *J. Funct. Anal.*, **233** (2006) 380-425.
- [35] Y.Y. Li, Remark on some conformally invariant integral equations: the method of moving spheres. *J. Eur. Math. Soc.* (JEMS), **6** (2004) 153-180.
- [36] Y.Y. Li, L. Zhang, Liouville-type theorems and Harnack-type inequalities for semilinear elliptic equations. *J. Anal. Math.*, **90** (2003) 27-87.
- [37] Y.Y. Li, M. Zhu, Uniqueness theorems through the method of moving spheres. *Duke Math. J.*, **80** (1995) 383-418.

- [38] C.-S. Lin, A classification of solutions of a conformally invariant fourth order equation in \mathbb{R}^n . *Comment. Math. Helv.*, **73** (1998) 206-231.
- [39] C.-S. Lin, Estimates of the scalar curvature equation via the method of moving planes III, *Comm. Pure Appl. Math.*, **53**(5) (2000) 611-646.
- [40] F. Marques, Isolated singularities of solutions to the Yamabe equation. *Calc. Var. Partial Differential Equations*, **32** (2008), no. 3, 349-371.
- [41] P. Mattila, Geometry of Sets and Measures in Euclidean Spaces: Fractals and rectifiability. Cambridge Studies in Advanced Mathematics, vol. 44, Cambridge University Press, Cambridge, 1995.
- [42] R. Mazzeo, F. Pacard, A construction of singular solutions for a semilinear elliptic equation using asymptotic analysis. *J. Differential Geom.*, **44** (1996), no. 2, 331-370.
- [43] D. Mitrea, Distributions, partial differential equations, and harmonic analysis, Springer, New York (2013).
- [44] F. Pacard, Solutions with high dimensional singular set, to a conformally invariant elliptic equation in \mathbb{R}^4 and in \mathbb{R}^6 . *Comm. Math. Phys.*, **159** (1994), no. 2, 423-432.
- [45] S. Paneitz, A quartic conformally covariant differential operator for arbitrary pseudo-Riemannian manifolds. Preprint, 1983, arXiv:0803.4331.
- [46] J. Ratzkin, On constant Q-curvature metrics with isolated singularities. arXiv:2001.07984.
- [47] R. Schoen, Conformal deformation of a Riemannian metric to constant scalar curvature. *J. Differential Geom.*, **20** (1984), no. 2, 479-495.
- [48] R. Schoen, The existence of weak solutions with prescribed singular behavior for a conformally invariant scalar equation. *Comm. Pure Appl. Math.*, **41** (1988), no. 3, 317-392.
- [49] R. Schoen, S.-T. Yau, Conformally flat manifolds, Kleinian groups and scalar curvature. *Invent. Math.*, **92** (1988) 47-71.
- [50] S. Taliaferro, Existence of large singular solutions of conformal scalar curvature equations in \mathbb{S}^n . *J. Funct. Anal.*, **224** (1) (2005) 192-216.
- [51] S. Taliaferro, L. Zhang, Asymptotic symmetries for conformal scalar curvature equations with singularity. *Calc. Var. Partial Differ. Equ.*, **26** (4) (2006) 401-428.
- [52] J. Xiong, L. Zhang, Isolated singularities of solutions to the Yamabe equation in dimension 6. arXiv:2006.13279v2.
- [53] H. Yang, Asymptotic behavior of positive solutions to a nonlinear biharmonic equation near isolated singularities. *Calc. Var. Partial Differential Equations*, **59** (2020), no. 4, Paper No. 130, 21 pp.
- [54] L. Zhang, Refined asymptotic estimates for conformal scalar curvature equation via moving sphere method. *J. Funct. Anal.*, **192** (2002), no. 2, 491-516.

T. Jin and H. Yang

Department of Mathematics, The Hong Kong University of Science and Technology

Clear Water Bay, Kowloon, Hong Kong, China

E-mail addresses: tianlingjin@ust.hk (T. Jin) mahuiyang@ust.hk (H. Yang)