

Nonlinearly stability of solutions on the outer Pressure Problem of Compressible Navier-Stokes System with Temperature-Dependent Heat Conductivity *

Guocai Cai^a, Canpei Chen^b, Yanfang Peng^{c†}

a. School of Mathematical Sciences, Xiamen University,
Xiamen 361005, P. R. China;

b. School of Mathematical Sciences,
University of Chinese Academy of Sciences, Beijing 100049, P. R. China

c. School of Mathematical Sciences, Guizhou Normal University,
Guiyang, 550001, P. R. China;

Abstract

In this paper, the one-dimensional compressible Navier-Stokes system with outer pressure boundary conditions is investigated. Under some suitable assumptions, we prove that the specific volume and the temperature are bounded from below and above independently of time, and then give the local and global existence of strong solutions. Furthermore, we also obtain the convergence of the global strong solution to a stationary state and the nonlinearly stability of its convergence. It is worth noticing that all the assumptions imposed on the initial data are the same as Takeyuki Nagasawa [Japan.J.Appl.Math.(1988)]. Therefore, our work can be regarded as an improvement of the results of Takeyuki.

Keywords: Compressible Navier-Stokes system; Temperature-dependent heat conductivity; Outer pressure problem; Stability

Math Subject Classification: 35Q35; 76N10.

1 Introduction

The one-dimensional compressible Navier-Stokes system describing the motion of a viscous heat-conducting perfect polytropic gas has the form in the Lagrangian mass coordinates (see [4, 22]):

$$v_t = u_x, \quad (1.1)$$

$$u_t + P_x = \left(\mu \frac{u_x}{v} \right)_x, \quad (1.2)$$

$$\left(e + \frac{1}{2}u^2 \right)_t + (Pu)_x = \left(\frac{\kappa\theta_x + \mu u u_x}{v} \right)_x, \quad (1.3)$$

*Partially supported by NNSFC Grant Nos. 11501143, 12071200, 11971401, 11871410 and 11871408.

†Email addresses: gotry@xmu.edu.cn (G.C. Cai), 20150008029@m.scnu.edu.cn (C.P. Chen), py-fang2005@sina.com (Y.F. Peng)

where $t > 0$ is time, $x \in \Omega = (0, 1)$ denotes the Lagrange mass coordinate, and the unknown functions $v > 0, u$ and P are the specific volume of the gas, fluid velocity and pressure respectively. In this paper, we focus on ideal polytropic gas, that is, P and e satisfy

$$P = R\theta/v, \quad e = c_v\theta + \text{const}, \quad (1.4)$$

where both specific gas constant R and heat capacity at constant volume c_v are positive constants. The viscosity μ and the heat conductivity κ are given by

$$0 < \mu = \tilde{\mu}\theta^\alpha, \quad \kappa = \tilde{\kappa}\theta^\beta, \quad (1.5)$$

with constants $\tilde{\mu}, \tilde{\kappa} > 0$ and $\alpha, \beta \geq 0$. The system (1.1)-(1.5) is supplemented with the initial data for any $x \in (0, 1)$,

$$(v, u, \theta)(x, 0) = (v_0, u_0, \theta_0)(x), \quad (1.6)$$

and the boundary conditions

$$\left(\frac{\mu}{v}u_x - P\right)(0, t) = \left(\frac{\mu}{v}u_x - P\right)(1, t) \triangleq -p(t) < 0, \quad \theta_x(0, t) = \theta_x(1, t) = 0. \quad (1.7)$$

where the first boundary conditions are called outer pressure boundary conditions. In addition, the initial data (1.6) should be compatible with the boundary conditions (1.7).

A large number of literatures have been devoted to the large-time existence and behavior of solutions to one-dimensional compressible Navier-Stokes equations. For constant coefficients ($\alpha = \beta = 0$) with large initial data, Kazhikhov and Shelukhin [16] first showed the global existence of solutions under the following boundary conditions:

$$u(0, t) = u(1, t) = 0, \quad \theta_x(0, t) = \theta_x(1, t) = 0, \quad (1.8)$$

which mean that the gas is confined into a fixed tube with impermeable gas. From then on, significant progress has been made on the mathematical aspect of the initial boundary value problems, see [1–3, 18–20] and the references therein. Motivated by the fact that in the case of isentropic flow a temperature dependence on the viscosity translates into a density dependence, there is a lot of research work (see [3, 6, 8, 9, 15, 21] and the references therein) concentrating on the case that μ is independent of θ , and heat conductivity is allowed to depend on temperature in a special way with a positive lower bound and balanced with corresponding constitution relations. In the case that $\alpha = 0, \beta > 0$ with boundary condition of either (1.7) with $p(t) \equiv 0$ or (1.8), Kawohl [15], Jiang [11, 12] and Wang [25] established the global existence of smooth solutions for (1.1)–(1.3), (1.6) under the assumption $\mu(v) \geq \mu_0 > 0$ for any $v > 0$ and κ may depend on both density and temperature. Moreover, when $\alpha = 0$ and $\beta \in (0, 3/2)$, Jenssen-Karper [10] proved the global existence of a weak solution to (1.1)–(1.7). Later on, for $\alpha = 0$ and $\beta \in (0, \infty)$, Pan-Zhang [21] obtained the global strong solutions under the non-slip and heat insulated boundary conditions. Recently, Duan-Guo-Zhu [5] obtained the global strong solutions of (1.1)–(1.7) under the condition that

$$(v_0, u_0, \theta_0) \in H^1 \times H^2 \times H^2. \quad (1.9)$$

And then, Huang-Shi-Sun [7] relaxed the initial condition (1.9) to

$$(v_0, u_0, \theta_0) \in H^1 \times H^1 \times H^1. \quad (1.10)$$

On the other hand, concerning the large-time behavior of the strong solutions to (1.1)-(1.6) and (1.8), Kazhikhov [13] first proved that for the case $\alpha = \beta = 0$, the strong solution is nonlinearly exponentially stable as time tends to infinity. More recently, Huang-Shi [9] extended the same result to $\alpha = 0, \beta > 0$. However, it should be pointed out here that the method used in [13] relies heavily on the non-degeneracy of both the viscosity μ and the heat conductivity κ , so that it can not be applied directly to the degenerate and nonlinear case ($\alpha = 0, \beta > 0$).

When it comes to the outer pressure problem (1.1)-(1.7) with $\alpha = \beta = 0$, Takeyuki Nagasawa [23] showed the convergence of the solutions to a stationary state and the rate of its convergence under some suitable assumptions. However, whether the same results are still hold for the system (1.1)-(1.7) with $\alpha = 0, \beta > 0$ remains open. One of our purposes is to give a positive answer to this problem. In fact, we establish the existence of the strong solutions for the problem (1.1)-(1.7) and study the large-time stability under the additional assumptions (1.14) and (1.15), more precisely, we have main results as follows.

Theorem 1.1. *Suppose that*

$$\alpha = 0, \quad \beta > 0, \quad 0 < p(t) \in C^1[0, +\infty),$$

and the initial data (v_0, u_0, θ_0) satisfies

$$(v_0, u_0, \theta_0) \in H^1(0, 1), \quad \inf_{x \in [0, 1]} v_0(x) > 0, \quad \inf_{x \in [0, 1]} \theta_0(x) > 0. \quad (1.11)$$

Then the initial-boundary-value problem (1.1)-(1.7) has a unique global strong solution (v, u, θ) such that for each $T > 0$,

$$\begin{cases} v, u, \theta \in L^\infty(0, T; H^1(0, 1)), \\ v_t \in L^\infty(0, T; L^2(0, 1)) \cap L^2(0, T; H^1(0, 1)), \\ u_x, \theta_x, u_t, \theta_t, v_{xt}, u_{xx}, \theta_{xx} \in L^2((0, T) \times (0, 1)). \end{cases} \quad (1.12)$$

Moreover,

$$C^{-1} \leq v(x, t) \leq C, \quad C^{-1} \leq \theta(x, t) \leq C, \quad (1.13)$$

where C is a positive constant depending on the initial data and T .

The following theorem indicates that the conclusion of Theorem 1.1 is uniform with respect to t and the global strong solution tends to a stationary state as time goes away, if one further assumes that $p(t)$ satisfies

$$m_p \triangleq \inf_{t \in [0, +\infty)} p(t) > 0 \quad (1.14)$$

and

$$I_p \triangleq \int_0^{+\infty} |p'(t)| dt < +\infty. \quad (1.15)$$

Theorem 1.2. *Under the assumptions of Theorem 1.1, suppose that $p(t) \in C^1[0, \infty)$ satisfies (1.14) and (1.15) in addition. Then the initial-boundary-value problem (1.1)-(1.7) has a unique global strong solution (v, u, θ) satisfying*

$$\begin{cases} v, u, \theta \in L^\infty(0, \infty; H^1(0, 1)), \\ v_t \in L^\infty(0, \infty; L^2(0, 1)) \cap L^2(0, \infty; H^1(0, 1)), \\ u_x, \theta_x, u_t, \theta_t, v_{xt}, u_{xx}, \theta_{xx} \in L^2((0, \infty) \times (0, 1)), \end{cases} \quad (1.16)$$

and for any $(x, t) \in (0, 1) \times (0, +\infty)$,

$$C^{-1} \leq v(x, t) \leq C, \quad C^{-1} \leq \theta(x, t) \leq C$$

with some positive constant C depending only on the initial data. Moreover, (v, u, θ) converges to a stationary state $(\hat{v}, \hat{u}, 0)$ in $H^1(0, 1)$ as $t \rightarrow +\infty$, where

$$\hat{v} \triangleq \frac{1}{2\bar{P}} \left\{ \int_0^1 \left(\frac{u_0^2}{2} + \theta_0 + p(0)v_0 \right) dx + \int_0^{+\infty} p'(\tau) \int_0^1 v(x, \tau) dx d\tau \right\} \quad (1.17)$$

and

$$\hat{\theta} \triangleq \frac{1}{2} \left\{ \int_0^1 \left(\frac{u_0^2}{2} + \theta_0 + p(0)v_0 \right) dx + \int_0^{+\infty} p'(\tau) \int_0^1 v(x, \tau) dx d\tau \right\} \quad (1.18)$$

with

$$\bar{P} \triangleq \lim_{t \rightarrow +\infty} p(t). \quad (1.19)$$

Similar to the proof of Theorem 3 in [23], we have the following results on rate of convergence of the strong solutions obtained by Theorem 1.2.

Theorem 1.3. *Assume that the assumptions of Theorem 1.2 hold. Then there exist positive constants $\lambda_1, \lambda_2 (\lambda_1 > \lambda_2), C$ depending on $\inf_{t \in [0, +\infty)} p(t), \sup_{t \in [0, +\infty)} p(t)$ and initial data such that the strong solution (v, u, θ) in Theorem 1.2 satisfies*

$$\int_0^1 \{u^2 + (v - \hat{v})^2 + (\theta - \hat{\theta})^2 + v_x^2\} dx \leq C \exp(-\lambda_1 t) \left\{ \int_0^t \exp(\lambda_1 \tau) F(\tau) d\tau + 1 \right\}, \quad (1.20)$$

$$\begin{aligned} \int_0^1 u_x^2 dx &\leq C \exp(-\lambda_1 t) \left\{ \int_0^t \exp(\lambda_1 \tau) F(\tau) d\tau + 1 \right\} \\ &+ C \left[\exp(-\lambda_2 t) \left\{ \int_0^t \exp(\lambda_2 \tau) (F(\tau) + |p'(\tau)|) d\tau + 1 \right\} + (p(t) - \bar{P})^2 \right], \end{aligned} \quad (1.21)$$

$$\int_0^1 \theta_x^2 dx \leq C \exp(-\lambda_2 t) \left\{ \int_0^t \exp(\lambda_2 \tau) (F(\tau) + |p'(\tau)|) d\tau + 1 \right\} \quad (1.22)$$

where

$$Y(t) \triangleq \exp \left\{ \int_0^t p(\tau) d\tau \right\}, \quad (1.23)$$

$$F(t) \triangleq \frac{1}{Y^2(t)} + \left(\frac{1}{Y(t)} \int_0^t Y(\tau) (\bar{P} - p(\tau)) d\tau \right)^2 + \int_t^{+\infty} |p'(\tau)| d\tau.$$

We now make some comments on the analysis of this paper. The key issue is to get the lower and upper bounds of v and θ (see (2.18), (2.19) and (2.50)). Compared with [7] and [23], the main difficulties come from the degeneracy and nonlinearity of the heat conductivity due to $\beta > 0$ and the outer pressure boundary conditions (1.7). Hence, in order to arrive at (2.50), some new ideas have to be involved here, which are based on three technical treatments. First, applying the standard energetic estimate (2.6) and the method in [7] yields that the specific volume v is bounded from above and below (see (2.12) or (2.18), (2.19)). Next, the lower bound of θ (see (2.22)) is taken into

account, and it suffices to show that $\|(\theta^{-1/2} - \alpha_1^{-1/2})_+\|_{L^\infty[0,1]} \leq C$. To this end, we multiply (2.10) by $(\theta^{-1/2} - \alpha_1^{-1/2})_+^p \theta^{-3/2}$ and apply Gronwall's inequality due to the observation that $\int_0^T (\theta^{-1/2} - \alpha_1^{-1/2})_+^2 dt$ can be bounded by $\int_0^T V(t) dt$ (see (2.24) and (2.25)). Finally, for $\beta \in (0, 1)$, multiplying (2.10) by $(\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+ \theta^{-(1+\beta)/2}$, and together with the standard energetic estimate (2.6), we obtain for any $\beta > 0$,

$$\int_0^T \int_0^1 \theta^{-1} \theta_x^2 dx dt \leq C,$$

which not only implies that

$$\int_0^T \max_{x \in [0,1]} \left| \theta(x, t) - \int_0^1 \theta(x, t) dx \right|^2 dt \leq C,$$

but also yields that the $L^2((0, T) \times (0, 1))$ -norm of θ_x^2 is bounded provided $\theta \leq \alpha_2$. In the case that $\theta > \alpha_2$, we multiply (2.10) by $(\theta - \alpha_2)_+$ so that the $L^2((0, T) \times (0, 1))$ -norm of θ_x can be bounded by the $L^4(0, T; L^2(0, 1))$ -norm of u_x which plays an important role in obtaining the bound on $L^2((0, T) \times (0, 1))$ -norm of both θ_x and u_{xx} (see Lemma 2.9). All these discussions will be carried out in the next section.

2 Proof of Theorem 1.1

We first state the following local existence result which can be proved by applying the contraction mapping principle (c.f. [14, 17, 24]).

Lemma 2.1. *Let (1.1)-(1.7) hold. Then there exists some $T > 0$ such that the initial-boundary-value problem (1.1)-(1.7) has a unique strong solution (v, u, θ) satisfying*

$$\begin{cases} v, u, \theta \in L^\infty(0, T; H^1(0, 1)), \\ v_t \in L^\infty(0, T; L^2(0, 1)) \cap L^2(0, T; H^1(0, 1)), \\ u_t, \theta_t, v_{xt}, u_{xx}, \theta_{xx} \in L^2((0, T) \times (0, 1)). \end{cases}$$

Then, when $p(t)$ satisfies (1.14) and (1.15) in addition, we obtain some priori estimates (see (2.12), (2.26), (2.41), (2.50), (2.55) below), where the constants depend only on the initial data of the problem so that we extend the local solution to the whole interval $[0, +\infty)$ and establish Theorem 1.2.

Without loss of generality, we assume that $\tilde{\mu} = \tilde{\kappa} = R = c_v = 1$. Moreover, integrating (1.2) over $[0, 1]$ and using (1.7), then

$$\int_0^1 u dx = \int_0^1 u_0 dx. \quad (2.1)$$

Hence, we may assume that

$$\int_0^1 u_0 dx = 0. \quad (2.2)$$

Unless otherwise stated, we follow the convention that C is an unspecified positive constant that may vary from expression to expression, even across an inequality (but not across an equality). Also C depends on initial data, β and $p(t)$ generally, and

the dependence of C on other parameters will be specified within parenthesis when necessary.

First, we derive the following representation of v which is essential to arrive at the lower and upper bounds of v .

Lemma 2.2. *It holds that*

$$v(x, t) = \frac{v_0(x)}{B(x, t)Y(t)} \left(1 + \frac{1}{v_0(x)} \int_0^t B(x, \tau)\theta(x, \tau)Y(\tau)d\tau \right), \quad (2.3)$$

where $Y(t)$ is defined in (1.23) and

$$B(x, t) \triangleq \exp \left\{ \int_0^x (u_0(y) - u(y, t))dy \right\}.$$

Proof. By (1.2), we get

$$u_t = \left(\frac{u_x}{v} - \frac{\theta}{v} \right)_x.$$

Integrating this over $(0, x)$ and using (1.7) give

$$\left(\int_0^x u dy \right)_t = \frac{u_x}{v} - \frac{\theta}{v} + p(t). \quad (2.4)$$

Then by (1.1),

$$\left(\int_0^x u dy \right)_t = (\ln v)_t - \frac{\theta}{v} + p(t). \quad (2.5)$$

Integrating (2.5) over $(0, t)$ yields

$$\ln v = \ln v_0 + \int_0^x (u - u_0)dy + \int_0^t \frac{\theta}{v} d\tau - \int_0^t p(\tau)d\tau,$$

which implies

$$v(x, t) = \frac{v_0(x)}{B(x, t)Y(t)} \exp \int_0^t \frac{\theta}{v}(x, \tau)d\tau.$$

This in particular gives (2.3) and finishes the proof of Lemma 2.2. \square

Next, we give the following energetic estimates.

Lemma 2.3. *Setting*

$$V(t) \triangleq \int_0^1 \left(\frac{\theta^\beta \theta_x^2}{v\theta^2} + \frac{u_x^2}{v\theta} \right) (x, t)dx,$$

we have

$$\sup_{0 \leq t \leq T} \int_0^1 (u^2/2 + v - \ln v + \theta - \ln \theta) dx + \int_0^T V(s)ds \leq C_0, \quad (2.6)$$

where and in what follows, C_0 and C are both positive constants depending only on β , $\|(v_0, u_0, \theta_0)\|_{H^1(0,1)}$, $\inf_{x \in [0,1]} v_0(x)$, $\inf_{x \in [0,1]} \theta_0(x)$, $m_{p,T}$, $M_{p,T}$, and $I_{p,T}$ with

$$m_{p,T} \triangleq \inf_{t \in [0,T]} p(t), \quad M_{p,T} \triangleq \sup_{t \in [0,T]} p(t), \quad I_{p,T} \triangleq \int_0^T |p'(t)|dt.$$

Proof. First, integrating (1.3) over $(0, 1)$ and using (1.1), (1.7) immediately give

$$\begin{aligned}
\left(\int_0^1 \left(\theta + \frac{u^2}{2} \right) dx \right)_t &= \left(\frac{\theta^\beta \theta_x}{v} + \frac{u(u_x - \theta)}{v} \right) \Big|_{x=0}^{x=1} \\
&= -p(t)(u(1) - u(0)) = -p(t) \int_0^1 u_x dx \\
&= - \left(\int_0^1 p(t) v dx \right)_t + \int_0^1 p'(t) v dx,
\end{aligned} \tag{2.7}$$

which implies

$$\begin{aligned}
\int_0^1 \left(\theta + \frac{u^2}{2} + p(t)v \right) dx &= \int_0^1 \left(\theta_0 + \frac{u_0^2}{2} + p(0)v_0 \right) dx + \int_0^t p'(\tau) \int_0^1 v dx d\tau \\
&\leq C + \int_0^t |(\ln p(\tau))'| \left(\int_0^1 p(\tau) v dx \right) d\tau.
\end{aligned} \tag{2.8}$$

Then by Gronwall's inequality, we have

$$\sup_{0 \leq t \leq T} \int_0^1 (u^2 + \theta + v) dx \leq C. \tag{2.9}$$

Next, by (1.2), we rewrite (1.3) as

$$\theta_t + \frac{\theta}{v} u_x = \left(\frac{\theta^\beta \theta_x}{v} \right)_x + \frac{u_x^2}{v}. \tag{2.10}$$

Multiplying (1.1) by $(1 - v^{-1})$, (1.2) by u , (2.10) by $1 - \theta^{-1}$, and adding them together, we obtain

$$\begin{aligned}
&\left(\frac{u^2}{2} + (v - \ln v) + (\theta - \ln \theta) \right)_t + \frac{u_x^2}{v\theta} + \frac{\theta^\beta \theta_x^2}{v\theta^2} \\
&= \left(u \left(\frac{u_x}{v} - \frac{\theta}{v} \right) \right)_x + u_x + \left((1 - \theta^{-1}) \frac{\theta^\beta \theta_x}{v} \right)_x.
\end{aligned} \tag{2.11}$$

Finally, integrating (2.11) over $(0, 1)$, and utilizing (1.1), (1.7) lead to

$$\begin{aligned}
&\left(\int_0^1 (u^2/2 + v - \ln v + \theta - \ln \theta) dx \right)_t + \int_0^1 \left(\frac{\theta^\beta \theta_x^2}{v\theta^2} + \frac{u_x^2}{v\theta} \right) dx \\
&= \left((1 - p(t)) \int_0^1 v dx \right)_t + \int_0^1 p'(t) v dx,
\end{aligned}$$

which together with (2.9) yields (2.6). \square

Lemma 2.4. *For any $T > 0$, it holds that*

$$C^{-1} e^{-M_{p,T} T} \leq v(x, t) \leq C, \tag{2.12}$$

for any $(x, t) \in [0, 1] \times [0, T]$.

Proof. First, denoting the average of a function f over $(0, 1)$ by

$$\bar{f} \triangleq \int_0^1 f dx.$$

Since the function $x - \ln x$ is convex, Jensen's inequality gives

$$\bar{\theta} - \ln \bar{\theta} \leq \int_0^1 (\theta - \ln \theta) dx,$$

which together with (2.6) implies

$$\bar{\theta}(t) \in [\alpha_1, \alpha_2], \quad (2.13)$$

where $0 < \alpha_1 < \alpha_2$ are two roots of

$$x - \ln x = C_0,$$

and C_0 is given by Lemma 2.3.

Thus, combining (2.13) with (2.9), we obtain

$$\begin{aligned} \left| \theta^{\frac{\beta+1}{2}} - \bar{\theta}^{\frac{\beta+1}{2}} \right| &\leq \frac{\beta+1}{2} \left(\int_0^1 \frac{\theta^\beta \theta_x^2}{\theta^2 v} dx \right)^{\frac{1}{2}} \left(\int_0^1 \theta v dx \right)^{\frac{1}{2}} \\ &\leq CV^{\frac{1}{2}}(t) \max_{x \in [0,1]} v^{\frac{1}{2}}(x, t), \end{aligned}$$

which leads to

$$\frac{\alpha_1}{4} - CV(t) \max_{x \in [0,1]} v(x, t) \leq \theta(x, t) \leq C + CV(t) \max_{x \in [0,1]} v(x, t), \quad (2.14)$$

for all $(x, t) \in [0, 1] \times [0, T]$.

Next, by (2.6) and Cauchy's inequality, we have

$$\left| \int_0^x u dy \right| \leq \int_0^1 |u| dy \leq \left(\int_0^1 u^2 dy \right)^{\frac{1}{2}} \leq C,$$

which shows that

$$C^{-1} \leq B(x, t) \leq C. \quad (2.15)$$

Hence, we deduce from (2.15), (2.3) and (2.14) that

$$\begin{aligned} v(x, t) &\leq C + C \int_0^t e^{-\int_\tau^t p(s) ds} \max_{x \in [0,1]} \theta(x, \tau) d\tau \\ &\leq C + C \int_0^t e^{-\int_\tau^t p(s) ds} \left(1 + V(\tau) \max_{x \in [0,1]} v(x, \tau) \right) d\tau \\ &\leq C + C \int_0^t e^{-m_p \tau(t-\tau)} \left(1 + V(\tau) \max_{x \in [0,1]} v(x, \tau) \right) d\tau \\ &\leq C + C \int_0^t V(\tau) \max_{x \in [0,1]} v(x, \tau) d\tau, \end{aligned}$$

which together with Gronwall's inequality indicates that

$$v(x, t) \leq C, \quad (2.16)$$

for all $(x, t) \in [0, 1] \times [0, T]$.

Finally, it follows from (2.3) and (2.15) that

$$v(x, t) \geq C e^{-M_p T}, \quad (2.17)$$

for all $(x, t) \in [0, 1] \times [0, T]$. Combining (2.16) and (2.17), we finish the proof of Lemma 2.4. \square

Moreover, if (1.14) and (1.15) hold, then all the constants C and C_0 stated in Lemmas 2.3 and 2.4 and even the lower bound of v are indeed independent of T . More precisely, we have the following further result.

Corollary 2.5. *If (1.14) and (1.15) hold, then there exists some positive constant \tilde{C} depending only on β , $\|(v_0, u_0, \theta_0)\|_{H^1(0,1)}$, $\inf_{x \in [0,1]} v_0(x)$, $\inf_{x \in [0,1]} \theta_0(x)$, m_p , M_p , and I_p such that*

$$M_{v,T} \triangleq \sup_{t \in [0,T]} \max_{x \in [0,1]} v(x, t) \leq \tilde{C}, \quad (2.18)$$

and

$$m_{v,T} \triangleq \inf_{t \in [0,T]} \min_{x \in [0,1]} v(x, t) \geq \tilde{C}^{-1}. \quad (2.19)$$

Proof. If (1.14) and (1.15) hold, then it is easy to see that there exist positive constants M_p and \bar{P} such that

$$\bar{P} \triangleq \lim_{t \rightarrow +\infty} p(t) \quad (2.20)$$

and

$$M_p \triangleq \sup_{T \in [0, +\infty)} M_{p,T} = \sup_{t \in [0, +\infty)} p(t) < \infty. \quad (2.21)$$

Thus, under such additional condition (1.14) and (1.15), an analysis completely parallel to those of Lemmas 2.3 and 2.4 except for some necessary modifications when it comes to $p(t)$ shows that all the constants C_0 and C in the proof of the above two lemmas are independent of T .

Next, it is clear that the inequality (2.18) is a direct consequence of (2.16), (1.14), (1.15), and (2.21). On the other hand, it follows from (2.3), (2.6), (2.14) and (2.21) that

$$\begin{aligned} v(x, t) &\geq C \int_0^t e^{-\int_\tau^t p(s) ds} \min_{x \in [0,1]} \theta(x, \tau) d\tau \\ &\geq C \int_0^t e^{-M_p(t-\tau)} \left(\frac{\alpha_1}{4} - CV(\tau) \right) d\tau \\ &= \frac{C\alpha_1}{4M_p} - \frac{C\alpha_1}{4M_p} e^{-M_p t} - C \int_0^t e^{-M_p(t-\tau)} V(\tau) d\tau. \end{aligned}$$

Notice that, by (2.6),

$$\begin{aligned} \int_0^t e^{-M_p(t-\tau)} V(\tau) d\tau &= \int_0^{\frac{t}{2}} e^{-M_p(t-\tau)} V(\tau) d\tau + \int_{\frac{t}{2}}^t e^{-M_p(t-\tau)} V(\tau) d\tau \\ &\leq e^{-\frac{M_p t}{2}} \int_0^\infty V(\tau) d\tau + \int_{\frac{t}{2}}^t V(\tau) d\tau \rightarrow 0, \quad \text{as } t \rightarrow \infty, \end{aligned}$$

as a result, there exists some $\tilde{T} > 0$ such that

$$v(x, t) \geq \frac{C\alpha_1}{8M_p},$$

for all $(x, t) \in [0, 1] \times [\tilde{T}, +\infty)$. Combining this and (2.16), (2.17) gives (2.18) and (2.19).
□

Now, we are in a position to show the lower bound of the temperature which is important for our further discussion.

Lemma 2.6. *It holds that for $T > 0$,*

$$\inf_{x \in [0,1]} \theta(x, t) \geq C^{-1}, \quad (2.22)$$

where C is a positive constant depending only on β , $\|(v_0, u_0, \theta_0)\|_{H^1(0,1)}$, $\inf_{x \in [0,1]} v_0(x)$, $\inf_{x \in [0,1]} \theta_0(x)$, $m_{p,T}$, $M_{p,T}$, $I_{p,T}$, $m_{v,T}$ and $M_{v,T}$.

Remark 2.1. *It should be note that in the case that $p(t) \in C^1[0, +\infty)$, all the constants C usually depend on β , $\|(v_0, u_0, \theta_0)\|_{H^1(0,1)}$, $\inf_{x \in [0,1]} v_0(x)$, $\inf_{x \in [0,1]} \theta_0(x)$, $m_{p,T}$, $M_{p,T}$, $I_{p,T}$, and T . However, under additional condition (1.14) and (1.15), Our discussion below still indicates that C is independent of T , which implies that C depends only on initial data, β , m_p , M_p , and I_p . In other words, all the estimates in this section will be uniform with respect to time provided (1.14) and (1.15) holds.*

Proof of Lemma 2.6. For $p > 2$, multiplying (2.10) by $(\theta^{-1/2} - \alpha_1^{-1/2})_+^p \theta^{-3/2}$ and integrating by parts give

$$\begin{aligned} & \frac{2}{p+1} \left(\int_0^1 (\theta^{-1/2} - \alpha_1^{-1/2})_+^{p+1} dx \right)_t + \int_0^1 \frac{u_x^2 (\theta^{-1/2} - \alpha_1^{-1/2})_+^p}{v \theta^{3/2}} dx \\ & \leq \int_0^1 \frac{u_x (\theta^{-1/2} - \alpha_1^{-1/2})_+^p}{v \theta^{1/2}} dx \\ & \leq \frac{1}{2} \int_0^1 \frac{u_x^2 (\theta^{-1/2} - \alpha_1^{-1/2})_+^p}{v \theta^{3/2}} dx + C \int_0^1 \theta^{1/2} (\theta^{-1/2} - \alpha_1^{-1/2})_+^p dx \\ & \leq \frac{1}{2} \int_0^1 \frac{u_x^2 (\theta^{-1/2} - \alpha_1^{-1/2})_+^p}{v \theta^{3/2}} dx \\ & \quad + C \max_{x \in [0,1]} (\alpha_1^{1/2} - \theta^{1/2})_+^2 \left(1 + \int_0^1 (\theta^{-1/2} - \alpha_1^{-1/2})_+^{p+1} dx \right), \end{aligned} \quad (2.23)$$

where we have used the following fact

$$\begin{aligned} & \int_0^1 \theta^{1/2} (\theta^{-1/2} - \alpha_1^{-1/2})_+^p dx \\ & = \alpha_1^{-1} \int_0^1 \theta^{-1/2} (\theta^{-1/2} - \alpha_1^{-1/2})_+^{p-2} (\alpha_1^{1/2} - \theta^{1/2})_+^2 dx \\ & \leq C \max_{x \in [0,1]} (\alpha_1^{1/2} - \theta^{1/2})_+^2 \left(1 + \int_0^1 (\theta^{-1/2} - \alpha_1^{-1/2})_+^{p+1} dx \right). \end{aligned}$$

On the other hand, it follows from (2.13) that for any $\beta \in (0, 1)$,

$$\begin{aligned} \max_{x \in [0,1]} (\alpha_1^{1/2} - \theta^{1/2})_+ & \leq C \int_0^1 \theta^{-\frac{1}{2}} |\theta_x| dx \\ & \leq C \left(\int_0^1 \theta^{\beta-2} \theta_x^2 dx \right)^{1/2} \left(\int_0^1 \theta^{1-\beta} dx \right)^{1/2} \\ & \leq C V^{1/2}(t) \end{aligned} \quad (2.24)$$

and that for $\beta \geq 1$,

$$\begin{aligned}
\max_{x \in [0,1]} (\alpha_1^{1/2} - \theta^{1/2})_+ &\leq C \max_{x \in [0,1]} (\alpha_1^{\beta/2} - \theta^{\beta/2})_+ \\
&\leq C \int_0^1 \theta^{\frac{\beta}{2}-1} |\theta_x| dx \\
&\leq C \left(\int_0^1 \frac{\theta^\beta \theta_x^2}{\theta^{2\nu}} dx \right)^{1/2} \\
&\leq CV^{1/2}(t).
\end{aligned} \tag{2.25}$$

Therefore, by (2.23)-(2.25) and Gronwall's inequality, we obtain

$$\|\theta^{-1/2} - \alpha_1^{-1/2}\|_{L^{p+1}[0,1]} \leq C,$$

with C independent of p . Letting $p \rightarrow +\infty$, we get (2.22). \square

Using Lemma 2.6, we have the following estimate on the $L^\infty(0, T; L^2(0, 1))$ -norm of v_x .

Lemma 2.7. *There exists a positive constant C such that*

$$\sup_{0 \leq t \leq T} \int_0^1 v_x^2 dx + \int_0^T \int_0^1 v_x^2 (\theta + 1) dx dt \leq C. \tag{2.26}$$

for any $T > 0$.

Proof. First, for $\beta \in (0, 1)$, multiplying (2.10) by $(\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+ \theta^{-(1+\beta)/2}$ gives

$$\begin{aligned}
&\frac{1}{1-\beta} \left(\int_0^1 (\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+^2 dx \right)_t \\
&\quad + \beta \int_0^1 \frac{\theta_x^2}{v\theta} 1_{(\theta > \alpha_2)} dx + \int_0^1 \frac{u_x^2 (\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+}{v\theta^{(1+\beta)/2}} dx \\
&= \frac{1+\beta}{2} \alpha_2^{\frac{1-\beta}{2}} \int_0^1 1_{(\theta > \alpha_2)} \frac{\theta^{\frac{\beta-3}{2}} \theta_x^2}{v} dx + \int_0^1 \frac{u_x \theta^{(1-\beta)/2} (\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+}{v} dx \\
&\leq \frac{\beta}{2} \int_0^1 \frac{\theta_x^2}{v\theta} 1_{(\theta > \alpha_2)} dx + C \int_0^1 \frac{\theta^\beta \theta_x^2}{v\theta^2} dx + \int_0^1 \frac{u_x \theta^{(1-\beta)/2} (\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+}{v} dx \\
&\leq \frac{\beta}{2} \int_0^1 \frac{\theta_x^2}{v\theta} 1_{(\theta > \alpha_2)} dx + C\varepsilon \int_0^1 \frac{u_x^2 (\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+}{v\theta^{(1+\beta)/2}} dx + C(\varepsilon)V(t),
\end{aligned} \tag{2.27}$$

where we have utilized the following two estimates

$$\begin{aligned}
&\int_0^1 \frac{|u_x| \theta^{(1-\beta)/2} (\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+}{v} dx \\
&\leq \varepsilon \int_0^1 \frac{u_x^2}{v\theta^\beta} 1_{(\theta > 2\alpha_2)} dx + C(\varepsilon) \max_{x \in [0,1]} (\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+^2 \int_0^1 \theta dx \\
&\quad + C \int_0^1 \frac{|u_x| (\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+}{v\theta^{1/2}} 1_{(\theta \leq 2\alpha_2)} dx \\
&\leq C\varepsilon \int_0^1 \frac{u_x^2 (\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+}{v\theta^{(1+\beta)/2}} dx + C(\varepsilon)V(t)
\end{aligned}$$

and

$$\begin{aligned} \max_{x \in [0,1]} (\theta^{(1-\beta)/2} - \alpha_2^{(1-\beta)/2})_+^2 &\leq C \max_{x \in [0,1]} (\theta^{1/2} - \alpha_2^{1/2})_+^2 \\ &\leq CV(t). \end{aligned} \quad (2.28)$$

Therefore, for $\beta \in (0, 1)$, choosing ε suitably small and integrating (2.27) over $[0, T]$, we have

$$\int_0^T \int_0^1 \theta^{-1} \theta_x^2 dx dt \leq C, \quad (2.29)$$

which still holds for $\beta \geq 1$ due to (2.6) and (2.22).

Moreover, by (2.29) and (2.13),

$$\int_0^T \max_{x \in [0,1]} (\theta(x, t) - \bar{\theta}(t))^2 dt \leq C \int_0^T \left(\int_0^1 \theta^{-1} \theta_x^2 dx \right) \left(\int_0^1 \theta dx \right) dt \leq C. \quad (2.30)$$

Next, by (1.1), one can rewrite (1.2) as

$$\left(u - \frac{v_x}{v} \right)_t - \frac{\theta v_x}{v^2} = -\frac{\theta_x}{v}. \quad (2.31)$$

Multiplying (2.31) by $u - \frac{v_x}{v}$ and integrating over $(0, 1)$, along with (2.13), we obtain

$$\begin{aligned} &\left(\frac{1}{2} \int_0^1 \left(u - \frac{v_x}{v} \right)^2 dx \right)_t + \int_0^1 \frac{\theta v_x^2}{v^3} dx \\ &= \int_0^1 \frac{u \theta v_x}{v^2} dx - \int_0^1 \frac{u \theta_x}{v} dx + \int_0^1 \frac{\theta_x v_x}{v^2} dx \\ &\leq C \int_0^1 \frac{u^2 \theta}{v} dx + \frac{1}{4} \int_0^1 \frac{\theta v_x^2}{v^3} dx + C \int_0^1 \frac{\theta_x^2}{\theta} dx \\ &\leq C \max_{x \in [0,1]} u^2 \int_0^1 \theta dx + \frac{1}{4} \int_0^1 \frac{\theta v_x^2}{v^3} dx + C \int_0^1 \frac{\theta_x^2}{\theta} dx \\ &\leq C \max_{x \in [0,1]} u^2 + \frac{1}{4} \int_0^1 \frac{\theta v_x^2}{v^3} dx + C \int_0^1 \frac{\theta_x^2}{\theta} dx, \end{aligned}$$

which directly leads to

$$\begin{aligned} &\int_0^1 \left(u - \frac{v_x}{v} \right)^2 dx + \int_0^t \int_0^1 \frac{\theta v_x^2}{v^3} dx dt \\ &\leq C + C \int_0^t \max_{x \in [0,1]} u^2 dt + C \int_0^t \int_0^1 \frac{\theta_x^2}{\theta} dx dt \\ &\leq C + C \int_0^t V(t) dt \\ &\leq C, \end{aligned} \quad (2.32)$$

where in the second inequality we have used (2.29),

$$|u| = \left| u - \int_0^1 u dx \right| \leq \int_0^1 |u_x| dx, \quad (2.33)$$

and

$$\begin{aligned} \int_0^1 |u_x| dx &\leq \left(\int_0^1 \frac{u_x^2}{v \theta} dx \right)^{1/2} \left(\int_0^1 v \theta dx \right)^{1/2} \\ &\leq C \left(\int_0^1 \frac{u_x^2}{v \theta} dx \right)^{1/2} \leq CV^{1/2}(t). \end{aligned} \quad (2.34)$$

Therefore, we have

$$\int_0^1 v_x^2 dx + \int_0^T \int_0^1 \theta v_x^2 dx dt \leq C, \quad (2.35)$$

due to the following simple fact

$$\sup_{0 \leq t \leq T} \int_0^1 v_x^2 dx \leq C \int_0^1 u^2 dx + C \sup_{0 \leq t \leq T} \int_0^1 \left(u - \frac{v_x}{v}\right)^2 dx \leq C.$$

Finally, together with (2.22) and (2.35), we derive

$$\int_0^T \int_0^1 v_x^2 dx dt \leq C.$$

Combining this with (2.35) gives (2.26), and we finish the proof of Lemma 2.7. \square

For further discussion, we need the following estimate on the $L^2((0, T) \times (0, 1))$ -norm of θ_x .

Lemma 2.8. *There exists a positive constant C such that*

$$\sup_{0 \leq t \leq T} \int_0^1 \theta^2 dx + \int_0^T \int_0^1 \theta_x^2 dx dt \leq C + C \int_0^T \left(\int_0^1 u_x^2 dx\right)^2 dt, \quad (2.36)$$

for any $T > 0$.

Proof. Multiplying (2.10) by $(1 - \alpha_2/\theta)_+$ and integrating by parts give

$$\begin{aligned} & \int_0^1 \frac{u_x^2 (\theta - \alpha_2)_+}{v \theta} dx - \left(\int_0^1 \int_{\alpha_2}^{\theta} \left(1 - \frac{\alpha_2}{s}\right)_+ ds dx \right)_t \\ &= \int_0^1 \frac{(\theta - \alpha_2)_+}{v} u_x dx + \alpha_2 \int_0^1 \frac{\theta^\beta \theta_x^2}{v \theta^2} 1_{(\theta > \alpha_2)} dx \\ &\leq \varepsilon \int_0^1 \frac{u_x^2}{v} dx + C(\varepsilon) \int_0^1 (\theta - \alpha_2)_+^2 dx + CV(t) \\ &\leq \varepsilon \int_0^1 \frac{u_x^2}{v} dx + C(\varepsilon) \max_{x \in [0,1]} (\theta^{1/2} - \alpha_2^{1/2})_+^2 + CV(t) \\ &\leq \varepsilon \int_0^1 \frac{u_x^2}{v} dx + C(\varepsilon) V(t), \end{aligned}$$

where in the third inequality we have used (2.28). Consequently, we get

$$\int_0^T \int_0^1 u_x^2 dx dt \leq C. \quad (2.37)$$

Multiplying (2.10) by $(\theta - \alpha_2)_+$ and integrating by parts, we check that

$$\begin{aligned}
& \left(\int_0^1 \frac{(\theta - \alpha_2)_+^2}{2} dx \right)_t + \int_0^1 \frac{\theta^\beta \theta_x^2}{v} 1_{(\theta > \alpha_2)} dx \\
&= - \int_0^1 \frac{\theta(\theta - \alpha_2)_+}{v} u_x dx + \int_0^1 \frac{(\theta - \alpha_2)_+ u_x^2}{v} dx \\
&\leq C \int_0^1 \frac{(\theta - \alpha_2)_+^2}{v} |u_x| dx + C \int_0^1 \frac{(\theta - \alpha_2)_+}{v} |u_x| dx \\
&\quad + \varepsilon \max_{x \in [0,1]} (\theta - \alpha_2)_+^2 + C(\varepsilon) \left(\int_0^1 u_x^2 dx \right)^2 \\
&\leq C\varepsilon \max_{x \in [0,1]} (\theta - \alpha_2)_+^2 + C \int_0^1 (\theta - \alpha_2)_+^2 dx \int_0^1 u_x^2 dx \\
&\quad + C(\varepsilon) \int_0^1 u_x^2 dx + C(\varepsilon) \left(\int_0^1 u_x^2 dx \right)^2 \\
&\leq C\varepsilon \int_0^1 \theta^\beta \theta_x^2 1_{(\theta > \alpha_2)} dx + C \int_0^1 (\theta - \alpha_2)_+^2 dx \int_0^1 u_x^2 dx \\
&\quad + C(\varepsilon) \int_0^1 u_x^2 dx + C(\varepsilon) \left(\int_0^1 u_x^2 dx \right)^2,
\end{aligned} \tag{2.38}$$

where we have used the following inequality

$$\max_{x \in [0,1]} (\theta - \alpha_2)_+^2 \leq C \int_0^1 \theta^\beta \theta_x^2 1_{(\theta > \alpha_2)} dx. \tag{2.39}$$

Thus, choosing ε suitably small and by Gronwall's inequality, we obtain

$$\begin{aligned}
& \sup_{t \in [0, T]} \left(\int_0^1 \frac{(\theta - \alpha_2)_+^2}{2} dx \right) + \int_0^T \int_0^1 \frac{\theta^\beta \theta_x^2}{v} 1_{(\theta > \alpha_2)} dx dt \\
&\leq C + C \int_0^T \left(\int_0^1 u_x^2 dx \right)^2 dt,
\end{aligned} \tag{2.40}$$

which together with (2.6) and (2.22) gives (2.36) and the proof is complete. \square

Now, we can derive the following uniform estimate on the $L^2((0, T) \times (0, 1))$ -norm of u_t and u_{xx} .

Lemma 2.9. *There exists a positive constant C such that*

$$\sup_{0 \leq t \leq T} \int_0^1 u_x^2 dx + \int_0^T \int_0^1 (u_t^2 + u_{xx}^2) dx dt \leq C, \tag{2.41}$$

for any $T > 0$.

Proof. First, by (1.7),

$$(u_x - (\theta - p(t)v))(0, t) = 0, \quad (u_x - (\theta - p(t)v))(1, t) = 0.$$

Multiplying (1.2) by $(u_x - (\theta - p(t)v))_x$ and integrating over $(0, 1)$, by (1.1) and (1.3), we find that

$$\begin{aligned}
& \left(\frac{1}{2} \int_0^1 u_x^2 dx \right)_t + \left(\frac{p(t)}{2} \int_0^1 \frac{v_x^2}{v} dx \right)_t + \int_0^1 \left(\frac{u_{xx}^2}{v} + \frac{\theta_x^2}{v} + p(t) \frac{\theta v_x^2}{v^2} \right) dx \\
&= \left(\int_0^1 (\theta - p(t)v) u_x dx \right)_t - \frac{p(t)}{2} \int_0^1 u_x \left(\frac{v_x}{v} \right)^2 dx + p(t) \int_0^1 u_x^2 dx \\
&+ \int_0^1 \frac{\theta^\beta \theta_x}{v} u_{xx} dx - \int_0^1 \frac{u_x^3}{v} dx + \int_0^1 \frac{\theta}{v} u_x^2 dx - p(t) \int_0^1 \frac{\theta_x v_x}{v} dx \\
&+ 2 \int_0^1 \frac{u_{xx} \theta_x}{v} dx + \int_0^1 \frac{u_{xx} u_x v_x - u_{xx} \theta v_x - u_x v_x \theta_x + \theta v_x \theta_x}{v^2} dx \\
&+ \frac{1}{2} \int_0^1 p'(t) v \left(\frac{v_x}{v} \right)^2 dx + \int_0^1 p'(t) u_x v dx \\
&\leq \left(\int_0^1 (\theta - p(t)v) u_x dx \right)_t + \frac{1}{2} \int_0^1 \frac{u_{xx}^2}{v} dx + C \int_0^1 (u_x^2 + v_x^2 + \theta_x^2) dx \\
&+ C \left(\int_0^1 u_x^2 dx \right)^2 + C |p'(t)| \left(1 + \int_0^1 u_x^2 dx \right) + C \int_0^1 \frac{\theta^{2\beta} \theta_x^2}{v} dx,
\end{aligned} \tag{2.42}$$

where we have used the following estimates

$$\begin{aligned}
& \left| \int_0^1 u_x \left(\frac{v_x}{v} \right)^2 dx \right| \\
&\leq C \int_0^1 u_x^2 v_x^2 dx + C \int_0^1 v_x^2 dx \\
&\leq C \max_{x \in [0,1]} u_x^2 \int_0^1 v_x^2 dx + C \int_0^1 v_x^2 dx \\
&\leq \frac{1}{16} \int_0^1 \frac{u_{xx}^2}{v} dx + C \int_0^1 u_x^2 dx + C \int_0^1 v_x^2 dx, \\
& \left| \int_0^1 \frac{\theta^\beta \theta_x}{v} u_{xx} dx \right| \leq \frac{1}{16} \int_0^1 \frac{u_{xx}^2}{v} dx + C \int_0^1 \frac{\theta^{2\beta} \theta_x^2}{v} dx, \\
& \left| \int_0^1 \frac{u_x^3}{v} dx \right| \leq C \max_{x \in [0,1]} |u_x| \int_0^1 u_x^2 dx \\
&\leq C \max_{x \in [0,1]} u_x^2 dx + C \left(\int_0^1 u_x^2 dx \right)^2 \\
&\leq \frac{1}{16} \int_0^1 \frac{u_{xx}^2}{v} dx + C \int_0^1 u_x^2 dx + C \left(\int_0^1 u_x^2 dx \right)^2
\end{aligned}$$

and

$$\begin{aligned}
& \left| \int_0^1 \frac{\theta}{v} u_x^2 dx \right| \leq C \max_{x \in [0,1]} u_x^2 \leq \frac{1}{16} \int_0^1 \frac{u_{xx}^2}{v} dx + C \int_0^1 u_x^2 dx, \\
& \left| p(t) \int_0^1 \frac{\theta_x v_x}{v} dx \right| \leq C \int_0^1 \theta_x^2 dx + C \int_0^1 v_x^2 dx, \\
& \left| \int_0^1 \frac{u_{xx} \theta_x}{v} dx \right| \leq \frac{1}{16} \int_0^1 \frac{u_{xx}^2}{v} dx + C \int_0^1 \theta_x^2 dx,
\end{aligned}$$

$$\begin{aligned}
& \left| \int_0^1 \frac{u_{xx}u_xv_x - u_{xx}\theta v_x - u_xv_x\theta_x + \theta v_x\theta_x}{v^2} dx \right| \\
& \leq C \int_0^1 |u_{xx}u_xv_x - u_{xx}\theta v_x - u_xv_x\theta_x + \theta v_x\theta_x| dx \\
& \leq \frac{1}{32} \int_0^1 \frac{u_{xx}^2}{v} dx + C \int_0^1 \theta_x^2 dx + C \int_0^1 u_x^2 v_x^2 dx + C \int_0^1 \theta^2 v_x^2 dx \\
& \leq \frac{1}{32} \int_0^1 \frac{u_{xx}^2}{v} dx + C \int_0^1 \theta_x^2 dx + C \left(\max_{x \in [0,1]} u_x^2 + \max_{x \in [0,1]} (\theta - \bar{\theta})^2 + 1 \right) \int_0^1 v_x^2 dx \\
& \leq \frac{1}{16} \int_0^1 \frac{u_{xx}^2}{v} dx + C \int_0^1 u_x^2 dx + C \int_0^1 \theta_x^2 dx + C \int_0^1 v_x^2 dx, \\
& \quad \left| \int_0^1 p'(t)v \left(\frac{v_x}{v} \right)^2 dx \right| \leq C|p'(t)| \int_0^1 v_x^2 dx \leq C|p'(t)|, \\
& \quad \left| \int_0^1 p'(t)u_x v dx \right| \leq C|p'(t)| + C|p'(t)| \int_0^1 u_x^2 dx
\end{aligned}$$

due to (2.26) and the following fact

$$\max_{x \in [0,1]} u_x^2 \leq \varepsilon \int_0^1 u_{xx}^2 dx + C(\varepsilon) \int_0^1 u_x^2 dx \quad (2.43)$$

for any $\varepsilon > 0$.

Now, it only remains to estimate the last term $\int_0^1 \frac{\theta^{2\beta}\theta_x^2}{v} dx$ on the right side of (2.42). To this end, multiplying (2.10) by $(\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}})_+ \theta^{\frac{\beta}{2}}$, we arrive at

$$\begin{aligned}
& \frac{1}{\beta+2} \left(\int_0^1 (\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}})_+^2 dx \right)_t + (\beta+1) \int_0^1 \frac{\theta^{2\beta}\theta_x^2}{v} 1_{\{\theta > \alpha_2\}} dx \\
& = \int_0^1 \frac{u_x^2 \theta^{\frac{\beta}{2}}}{v} (\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}})_+ dx - \int_0^1 \frac{u_x \theta^{\frac{\beta+2}{2}}}{v} (\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}})_+ dx \\
& \quad + \frac{\beta}{2} \alpha_2^{\frac{\beta+2}{2}} \int_0^1 \frac{\theta^{\frac{3\beta}{2}} \theta_x^2}{v \theta} 1_{\{\theta > \alpha_2\}} dx \\
& \leq \int_0^1 \frac{u_x^2 \theta^{\frac{\beta}{2}}}{v} (\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}})_+ dx - \int_0^1 \frac{u_x \theta^{\frac{\beta+2}{2}}}{v} (\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}})_+ dx \\
& \quad + \frac{\beta+1}{2} \int_0^1 \frac{\theta^{2\beta}\theta_x^2}{v} 1_{\{\theta > \alpha_2\}} dx + CV(t).
\end{aligned}$$

Consequently,

$$\begin{aligned}
& \frac{1}{\beta+2} \left(\int_0^1 (\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}})_+^2 dx \right)_t + \frac{\beta+1}{2} \int_0^1 \frac{\theta^{2\beta}\theta_x^2}{v} 1_{\{\theta > \alpha_2\}} dx \\
& \leq \int_0^1 \frac{u_x^2 \theta^{\frac{\beta}{2}}}{v} (\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}})_+ dx - \int_0^1 \frac{u_x \theta^{\frac{\beta+2}{2}}}{v} (\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}})_+ dx + CV(t) \quad (2.44) \\
& \triangleq I_1 + I_2 + CV(t).
\end{aligned}$$

On the other hand, a straightforward calculation indicates that

$$\begin{aligned}
I_1 &\leq \int_0^1 \frac{u_x^2}{v} (\theta^{\beta+1} - \alpha_2^{\beta+1})_+ dx \\
&\leq C \max_{x \in [0,1]} \left(\theta^{\beta+1} - \alpha_2^{\beta+1} \right)_+ \int_0^1 u_x^2 dx \\
&\leq C \int_0^1 |\theta^\beta \theta_x| 1_{(\theta > \alpha_2)} dx \int_0^1 u_x^2 dx \\
&\leq \frac{\beta+1}{4} \int_0^1 \frac{\theta^{2\beta} \theta_x^2}{v} 1_{(\theta > \alpha_2)} dx + C \left(\int_0^1 u_x^2 dx \right)^2
\end{aligned} \tag{2.45}$$

and

$$\begin{aligned}
I_2 &\leq \int_0^1 |u_x| (\theta^{\beta+2} - \alpha_2^{\beta+2})_+ dx \\
&\leq C \max_{x \in [0,1]} \left(\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}} \right)_+ \int_0^1 \left(\theta^{\frac{\beta+2}{2}} + \alpha_2^{\frac{\beta+2}{2}} \right) 1_{(\theta > \alpha_2)} |u_x| dx \\
&\leq C \int_0^1 |\theta^{\frac{\beta}{2}} \theta_x| 1_{(\theta > \alpha_2)} dx \int_0^1 \left(\theta^{\frac{\beta+2}{2}} + \alpha_2^{\frac{\beta+2}{2}} \right) 1_{(\theta > \alpha_2)} |u_x| dx \\
&\leq C \int_0^1 \theta^\beta \theta_x^2 1_{(\theta > \alpha_2)} dx + C \int_0^1 \left(\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}} \right)_+^2 dx \int_0^1 u_x^2 dx + C \int_0^1 u_x^2 dx.
\end{aligned} \tag{2.46}$$

Putting (2.45)-(2.46) into (2.44), together with (2.6), (2.37) and (2.40), and by Gronwall's inequality, we have

$$\begin{aligned}
&\sup_{0 \leq t \leq T} \left(\int_0^1 \left(\theta^{\frac{\beta+2}{2}} - \alpha_2^{\frac{\beta+2}{2}} \right)_+^2 dx \right) + \int_0^T \int_0^1 \frac{\theta^{2\beta} \theta_x^2}{v} dx dt \\
&\leq C + C \int_0^T \left(\int_0^1 u_x^2 dx \right)^2 dt.
\end{aligned} \tag{2.47}$$

Next, by (2.36),

$$\begin{aligned}
\left| \int_0^1 (\theta - p(t)v) u_x dx \right| &\leq \frac{1}{8} \int_0^1 u_x^2 dx + C \int_0^1 \theta^2 dx + C \\
&\leq \frac{1}{8} \int_0^1 u_x^2 dx + C \int_0^T \left(\int_0^1 u_x^2 dx \right)^2 dt + C,
\end{aligned}$$

which together with (2.42), (2.37), (2.36), (2.26) and Gronwall's inequality shows that

$$\int_0^1 u_x^2 dx + \int_0^T \int_0^1 u_{xx}^2 dx dt \leq C. \tag{2.48}$$

Hence, by (2.48), (2.36), (2.43) and (2.47),

$$\sup_{0 \leq t \leq T} \int_0^1 \theta^{\beta+2} dx + \int_0^T \left(\int_0^1 (1 + \theta^{2\beta}) \theta_x^2 dx + \max_{x \in [0,1]} u_x^2 \right) dt \leq C. \tag{2.49}$$

Finally, we rewrite (1.2) as

$$u_t = \frac{u_{xx}}{v} - \frac{u_x v_x}{v^2} - \frac{\theta_x}{v} + \frac{\theta v_x}{v^2},$$

which along with (2.48) and (2.49) implies

$$\begin{aligned}
\int_0^T \int_0^1 u_t^2 dx dt &\leq C \int_0^T \int_0^1 u_{xx}^2 dx dt + C \int_0^T \int_0^1 u_x^2 v_x^2 dx dt \\
&\quad + C \int_0^T \int_0^1 \theta_x^2 dx dt + C \int_0^T \int_0^1 \theta^2 v_x^2 dx dt \\
&\leq C + C \int_0^T \max_{x \in [0,1]} u_x^2 dt + C \int_0^T \left(\max_{x \in [0,1]} \theta^2 \int_0^1 v_x^2 dx \right) dt \\
&\leq C + C \int_0^T \max_{x \in [0,1]} |\theta - \bar{\theta}|^2 dt + C \int_0^T \int_0^1 v_x^2 dx dt \\
&\leq C.
\end{aligned}$$

Combining this and (2.48), we finish the proof of Lemma 2.9. \square

Lemma 2.10. *There exists a positive constant C such that*

$$C^{-1} \leq \theta(x, t) \leq C, \quad (2.50)$$

for any $(x, t) \in [0, 1] \times [0, T]$.

Proof. Multiplying (2.10) by $\theta^\beta \theta_t$ and integrating over $(0, 1)$, by (2.49), we give

$$\begin{aligned}
&\int_0^1 \theta^\beta \theta_t^2 dx + \frac{1}{2} \left(\int_0^1 \frac{(\theta^\beta \theta_x)^2}{v} dx \right)_t \\
&= -\frac{1}{2} \int_0^1 \frac{(\theta^\beta \theta_x)^2 u_x}{v^2} dx - \int_0^1 \frac{\theta^{\beta+1} \theta_t u_x}{v} dx + \int_0^1 \frac{\theta^\beta \theta_t u_x^2}{v} dx \\
&\leq C \max_{x \in [0,1]} |u_x| \int_0^1 (\theta^\beta \theta_x)^2 dx + \frac{1}{2} \int_0^1 \theta^\beta \theta_t^2 dx + C \int_0^1 \theta^{\beta+2} u_x^2 dx \\
&\quad + C \int_0^1 \theta^\beta u_x^4 dx \tag{2.51} \\
&\leq C \left(\int_0^1 \frac{(\theta^\beta \theta_x)^2}{v} dx \right)^2 + C \max_{x \in [0,1]} u_x^2 + \frac{1}{2} \int_0^1 \theta^\beta \theta_t^2 dx \\
&\quad + C \max_{x \in [0,1]} u_x^2 \int_0^1 \theta^{\beta+2} dx + C \max_{x \in [0,1]} u_x^4 \int_0^1 \theta^\beta dx \\
&\leq C \left(\int_0^1 \frac{(\theta^\beta \theta_x)^2}{v} dx \right)^2 + C \max_{x \in [0,1]} u_x^2 + C \max_{x \in [0,1]} u_x^4 + \frac{1}{2} \int_0^1 \theta^\beta \theta_t^2 dx.
\end{aligned}$$

On the other hand, by (2.48) and (2.49),

$$\begin{aligned}
&\int_0^T \max_{x \in [0,1]} u_x^4 dt \\
&\leq C \int_0^T \int_0^1 u_x^4 dx dt + C \int_0^T \int_0^1 |u_x^3 u_{xx}| dx dt \\
&\leq C \int_0^T \max_{x \in [0,1]} u_x^2 \left(\int_0^1 u_x^2 dx \right) dt + C \int_0^T \max_{x \in [0,1]} u_x^2 \left(\int_0^1 |u_x u_{xx}| dx \right) dt \\
&\leq C + \frac{1}{2} \int_0^T \max_{x \in [0,1]} u_x^4 dt + C \int_0^T \int_0^1 (u_x^2 + u_{xx}^2) dx dt \\
&\leq C + \frac{1}{2} \int_0^T \max_{x \in [0,1]} u_x^4 dt,
\end{aligned}$$

which implies that

$$\int_0^T \max_{x \in [0,1]} u_x^4 dt < C. \quad (2.52)$$

Therefore, together with (2.52), (2.49) and (2.51), by Gronwall's inequality, one gets

$$\sup_{0 \leq t \leq T} \int_0^1 (\theta^\beta \theta_x)^2 dx + \int_0^T \int_0^1 \theta^\beta \theta_t^2 dx dt \leq C. \quad (2.53)$$

In particular,

$$\begin{aligned} \max_{x \in [0,1]} |\theta^{\beta+1} - \bar{\theta}^{\beta+1}| &\leq (\beta + 1) \int_0^1 \theta^\beta |\theta_x| dx \\ &\leq C \left(\int_0^1 (\theta^\beta \theta_x)^2 dx \right)^{1/2} \\ &\leq C, \end{aligned} \quad (2.54)$$

which along with (2.22) gives (2.50) and we finish the proof of Lemma 2.10. \square

Finally, we obtain the following estimate on the $L^2((0,1) \times (0,T))$ -norm of θ_t and θ_{xx} .

Lemma 2.11. *There exists a positive constant C such that*

$$\sup_{0 \leq t \leq T} \int_0^1 \theta_x^2 dx + \int_0^T \int_0^1 (\theta_x^2 + \theta_t^2 + \theta_{xx}^2) dx dt \leq C. \quad (2.55)$$

for any $T > 0$.

Proof. By (2.50) and (2.53),

$$\sup_{0 \leq t \leq T} \int_0^1 \theta_x^2 dx + \int_0^T \int_0^1 \theta_t^2 dx dt \leq C. \quad (2.56)$$

Rewrite (2.10) as

$$\frac{\theta^\beta \theta_{xx}}{v} = -\frac{\beta \theta^{\beta-1} \theta_x^2}{v} + \frac{\theta^\beta \theta_x v_x}{v^2} - \frac{u_x^2}{v} + \frac{\theta u_x}{v} + \theta_t.$$

Hence,

$$\begin{aligned} \int_0^T \int_0^1 \theta_{xx}^2 dx dt &\leq C \int_0^T \int_0^1 (\theta_x^4 + \theta_x^2 v_x^2 + u_x^4 + u_x^2 + \theta_t^2) dx dt \\ &\leq C + C \int_0^T \max_{x \in [0,1]} \theta_x^2 dt + C \int_0^T \max_{x \in [0,1]} u_x^4 dt \\ &\leq C + C \int_0^T \int_0^1 |\theta_x \theta_{xx}| dx dt \\ &\leq C + C \int_0^T \int_0^1 \theta_x^2 dx dt + \frac{1}{2} \int_0^T \int_0^1 \theta_{xx}^2 dx dt \\ &\leq C + \frac{1}{2} \int_0^T \int_0^1 \theta_{xx}^2 dx dt, \end{aligned} \quad (2.57)$$

where we have used (2.26), (2.37), (2.50), (2.52) and (2.56). Therefore, together with (2.57) and (2.56), we give (2.55) and finish the proof of Lemma 2.11. \square

Proof of Theorem 1.1 With Lemmas 2.1, 2.4, 2.6, 2.9, 2.10 and 2.11 at hand, one can prove Theorem 1.1 by a standard argument. We omit it. \square

3 Stability of the strong solutions

In this section, we will concentrate on studying the stability of the strong solutions when (1.14) and (1.15) both hold. As indicated in Remark 2.1, all the constants C , \tilde{C} , and C_0 stated in Section 2 are independent of T .

Lemma 3.1. *It holds that*

$$\|(v - \hat{v}, u, \theta - \hat{\theta})(\cdot, t)\|_{H^1(0,1)} \rightarrow 0, \quad \text{as } t \rightarrow +\infty. \quad (3.1)$$

where $\hat{v}, \hat{\theta}$ are defined in (1.17), (1.18) respectively.

Proof. First, we check that

$$\int_0^{+\infty} \left(\left| \frac{d}{dt} \|v_x(\cdot, t)\|_{L^2}^2 \right| + \left| \frac{d}{dt} \|\theta_x(\cdot, t)\|_{L^2}^2 \right| \right) dt \leq C, \quad (3.2)$$

where we have taken advantage of the following facts

$$\int_0^1 v_x v_{xt} dx = \int_0^1 v_x u_{xx} dx$$

and

$$\int_0^1 \theta_x \theta_{xt} dx = - \int_0^1 \theta_{xx} \theta_t dx.$$

As a result, we deduce from (2.26), (2.55) and (3.2) that

$$\lim_{t \rightarrow +\infty} \int_0^1 (v_x^2 + \theta_x^2) dx = 0. \quad (3.3)$$

In particular,

$$\lim_{t \rightarrow +\infty} \int_0^1 ((v - \bar{v})^2 + (\theta - \bar{\theta})^2) dx \leq C \lim_{t \rightarrow +\infty} \int_0^1 (v_x^2 + \theta_x^2) dx = 0. \quad (3.4)$$

Multiplying (1.2) by u and integrating over $[0, 1]$ lead to

$$\frac{1}{2} \left(\int_0^1 u^2 dx \right)_t + \int_0^1 \frac{u_x^2}{v} dx = \int_0^1 \left(\frac{\theta}{v} - p(t) \right) u_x dx \leq C \int_0^1 |u_x| dx,$$

which yields that there exists some constant $C > 0$ such that for any $N > 0$ and $s, t \in [N, N + 1]$

$$\int_0^1 u^2(x, t) dx - \int_0^1 u^2(x, s) dx \leq C \int_N^{N+1} \int_0^1 |u_x| dx d\tau.$$

Integrating this with respect to s over $(N, N + 1)$, together with (2.33), we get

$$\sup_{N \leq t \leq N+1} \int_0^1 u^2(x, t) dx \leq C \left(\int_N^{N+1} \int_0^1 u_x^2 dx d\tau \right)^{1/2}.$$

Letting $N \rightarrow +\infty$ and by (2.37),

$$\lim_{t \rightarrow +\infty} \int_0^1 u^2(x, t) dx = 0. \quad (3.5)$$

Next, multiplying (2.4) by v and integrating over $[0, 1]$, we obtain

$$\bar{\theta} - p(t)\bar{v} = \int_0^1 u_x dx + \int_0^1 u_x \int_0^x u dy dx - \left(\int_0^1 v \int_0^x u dy dx \right)_t, \quad (3.6)$$

then

$$\begin{aligned} \int_0^t (\bar{\theta} - p(t)\bar{v})^2 dt &\leq \frac{1}{2} \int_0^t (\bar{\theta} - p(t)\bar{v})^2 dt + C \int_0^t \int_0^1 u_x^2 dx dt + C \\ &+ C \int_0^t \int_0^1 |u| dx (|\bar{\theta}'| + |p'(t)|\bar{v} + p(t)|\bar{v}'|) dt. \end{aligned} \quad (3.7)$$

On the other hand, Observe that, by (2.10),

$$\bar{\theta}' = - \int_0^1 \frac{\theta}{v} u_x dx + \int_0^1 \frac{u_x^2}{v} dx,$$

which along with (2.37), (2.41) and (2.50) yields

$$\int_0^{+\infty} (\bar{\theta}')^2 dt \leq C \quad (3.8)$$

and

$$\int_0^{+\infty} (\bar{v}')^2 dt = \int_0^{+\infty} \left(\int_0^1 u_x dx \right)^2 dt \leq C. \quad (3.9)$$

Therefore,

$$\int_0^{+\infty} (\bar{\theta} - p(t)\bar{v})^2 dt \leq C. \quad (3.10)$$

Moreover, by (3.8)- (3.10),

$$\begin{aligned} &\int_0^{+\infty} \left| \frac{d}{dt} (\bar{\theta} - p(t)\bar{v})^2 \right| dt \\ &\leq C \int_0^{+\infty} |\bar{\theta} - p(t)\bar{v}| (|\bar{\theta}'| + |p'(t)|\bar{v} + p(t)|\bar{v}'|) dt \\ &\leq C \int_0^{+\infty} (\bar{\theta} - p(t)\bar{v})^2 dt + C \int_0^{+\infty} ((\bar{\theta}')^2 + |p'(t)| + (\bar{v}')^2) dt \\ &\leq C, \end{aligned} \quad (3.11)$$

which together with (3.10) leads to

$$\lim_{t \rightarrow +\infty} (\bar{\theta} - p(t)\bar{v}) = 0.$$

In addition, by (2.8), we have

$$\bar{\theta} + p(t)\bar{v} = -\frac{1}{2} \int_0^1 u^2 dx + \int_0^1 \left(\theta_0 + \frac{u_0^2}{2} + p(0)v_0 \right) dx + \int_0^t p'(\tau) \int_0^1 v dx d\tau, \quad (3.12)$$

Hence, by (3.5) and (3.12),

$$\lim_{t \rightarrow +\infty} \bar{v} = \hat{v}, \quad \lim_{t \rightarrow +\infty} \bar{\theta} = \hat{\theta}. \quad (3.13)$$

Finally, introducing

$$\Phi(t) \triangleq \int_0^1 (u_x - \theta + p(t)v)^2 dx.$$

Multiplying (2.4) by v , one checks that

$$u_x - \theta + p(t)v = v \left(\int_0^x u dx \right)_t,$$

which together with (2.41) implies

$$\int_0^{+\infty} \Phi(t) dt \leq C \int_0^{+\infty} \int_0^1 u_t^2 dx dt \leq C. \quad (3.14)$$

On the other hand, by (1.7), (2.37), (2.41), (2.55) and (3.14), we conclude that

$$\begin{aligned} \int_0^{+\infty} |\Phi'(t)| dt &\leq C \int_0^{+\infty} \left| \int_0^1 (u_x - \theta + p(t)v) u_{xt} dx \right| dt \\ &\quad + C \int_0^{+\infty} \left| \int_0^1 (u_x - \theta + p(t)v) (-\theta + p(t)v)_t dx \right| dt \\ &= C \int_0^{+\infty} \left| \int_0^1 (u_x - \theta + p(t)v)_x u_t dx \right| dt \\ &\quad + C \int_0^{+\infty} \left| \int_0^1 (u_x - \theta + p(t)v) (-\theta + p(t)v)_t dx \right| dt \\ &\leq C \int_0^{+\infty} \int_0^1 u_{xx}^2 dx dt + C \int_0^{+\infty} \int_0^1 \theta_x^2 dx dt + C \int_0^{+\infty} \int_0^1 v_x^2 dx dt \\ &\quad + C \int_0^{+\infty} \int_0^1 u_t^2 dx dt + C \int_0^{+\infty} \int_0^1 u_x^2 dx dt \\ &\quad + C \int_0^{+\infty} \int_0^1 \theta_t^2 dx dt + C \int_0^{+\infty} |p'(t)| dt \\ &\leq C. \end{aligned}$$

Combining the above inequality with (3.14), we derive

$$\lim_{t \rightarrow +\infty} \Phi(t) = 0, \quad (3.15)$$

which together with (3.3) gives

$$\lim_{t \rightarrow +\infty} \int_0^1 u_x^2 dx = 0. \quad (3.16)$$

As a result, by virtue of (3.3), (3.4), (3.5), (3.13) and (3.16), we establish (3.1). \square

Proof of Theorem 1.2 Notice that all the conclusions in Section 2 are uniform with respect to time provided that (1.14) and (1.15) holds, by Theorem 1.1, the problem (1.1)-(1.7) has a unique global strong solution (v, u, θ) satisfying (1.12) and v, θ has a uniform positive below and upper bounds, Moreover, by Lemma 3.1, such (v, u, θ) converges to a stationary state $(\hat{v}, \hat{u}, 0)$ in $H^1(0, 1)$ as $t \rightarrow +\infty$. we finish the proof of Theorem 1.2.

References

- [1] Amosov, A. A., Zlotnik, A. A.: Global generalized solutions of the equations of the one-dimensional motion of a viscous heat-conducting gas. *Soviet Math. Dokl.* **38** (1989), 1-5.
- [2] Amosov, A. A., Zlotnik, A.A.: Solvability “in the large” of a system of equations of the one-dimensional motion of an inhomogeneous viscous heat-conducting gas. *Math. Notes* **52** (1992), 753-763.
- [3] Antontsev, S. N., Kazhikhov, A. V., Monakhov, V. N.: *Boundary Value Problems in Mechanics of Nonhomogeneous Fluids*. North-Holland, Amsterdam, 1990.
- [4] Batchelor, G. K.: *An Introduction to Fluid Dynamics*. Cambridge University Press, London, 1967.
- [5] Duan, R., Guo, A., Zhu, C. J.: Global strong solution to compressible Navier-Stokes equations with density dependent viscosity and temperature dependent heat conductivity. *J. Differential Equations* **262** (2017), 4314-4335.
- [6] Hsiao, L.: *Quasilinear Hyperbolic Systems and Dissipative Mechanisms*, World Scientific Publisher Co., 1997.
- [7] Huang B., Shi X. D., Sun Y.: Global strong solutions to compressible Navier-Stokes system with degenerate heat-conductivity and density-depending viscosity. *J. Differential Equations* **12** (2019), 4314-4335.
- [8] Hsiao, L., Jiang, S.: Nonlinear hyperbolic-parabolic coupled systems, *Handbook of Differential Equations: Evolutionary Equations*, Elsevier, **1** (2004), 287-384.
- [9] Huang, B., Shi X. D.: Nonlinearly exponential stability of compressible Navier-Stokes system with degenerate heat-conductivity. arXiv:1809.00467.
- [10] Jenssen, H. K., Karper, T. K.: One-dimensional compressible flow with temperature dependent transport coefficients, *SIAM Journal on Mathematical Analysis*, **42** (2010), 904-930.
- [11] Jiang, S.: On initial-boundary value problems for a viscous, heat-conducting, one-dimensional real gas, *J. Differential Equations* **110** (1994), 157-181.
- [12] Jiang, S.: On the asymptotic behavior of the motion of a viscous, heat-conducting, one-dimensional real gas, *Math. Z.* **216** (1994), 317-336.
- [13] Kazhikhov A. V.: To a theory of boundary value problems for equations of one-dimensional nonstationary motion of viscous heat-conduction gases, in: *Boundary Value Problem for Hydrodynamical Equations*, No.50, Institute of Hydrodynamics, Siberian Branch Acad. USSR, 1981,37-62, in Russian.
- [14] Kawashima, S., Nishida, T.: Global solutions to the initial value problem for the equations of one dimensional motion of viscous polytropic gases. *J. Math. Kyoto Univ.* **21** (1981), 825-837.
- [15] Kawohl, B.: Global existence of large solutions to initial-boundary value problems for a viscous, heat-conducting, one-dimensional real gas, *J. Differential Equations* **58** (1985), 76-103.

- [16] Kazhikhov, A. V., Shelukhin, V. V.: Unique global solution with respect to time of initial boundary value problems for one-dimensional equations of a viscous gas. *J. Appl. Math. Mech.* **41** (1977), 273-282.
- [17] Nash, J.: Le problème de Cauchy pour les équations différentielles d'un fluide général. *Bull. Soc. Math. France.* **90** (1962), 487-497.
- [18] Nagasawa, T.: On the asymptotic behavior of the one-dimensional motion of the poly-tropic ideal gas with stress-free condition. *Q. Appl. Math.* **46** (1988), 665-679.
- [19] Nagasawa, T.: On the one-dimensional motion of the polytropic ideal gas non-fixed on the boundary. *J. Differ. Equ.* **65** (1986), 49-67.
- [20] Nishida, T.: Equations of motion of compressible viscous fluids. *Pattern and Waves* (Nishida, T., Mimura, M., Fujii, H. Eds.). Kinokuniya/North-Holland, Amsterdam, pp. 97-128, 1986
- [21] Pan, R. H., Zhang, W. Z.: Compressible Navier-Stokes equations with temperature dependent heat conductivities, *Commun. Math. Sci.* **13** (2015), 401-425.
- [22] Serrin, J.: Mathematical principles of classical fluid mechanics. *Handbuch der Physik.* VIII/1 (Flügge, S., Truesdell, C. Eds.). Springer, Berlin, pp. 125-262, 1972.
- [23] Takeyuki, N.: On the outer pressure problem of the one-dimension of Polytropic ideal gas, *Japan J. Appl. Math.* **5** (1988), 53-85.
- [24] Tani, A.: On the first initial-boundary value problem of compressible viscous fluid motion, *Publications of the Research Institute for Mathematical Sciences*, **13** (1977), 193-253.
- [25] Wang, D. H.: Global solutions of the Navier-Stokes equations for viscous compressible flows, *Nonlinear Anal.* **52** (2003), 1867-1890.