

# CLAP: A New Algorithm for Promise CSPs\*

Lorenzo Ciardo<sup>†</sup>

Stanislav Živný<sup>†</sup>

25th October 2022

## Abstract

We propose a new algorithm for Promise Constraint Satisfaction Problems (PCSPs). It is a combination of the **C**onstraint **B**asic **L**P relaxation and the **A**ffine **I**P relaxation (CLAP). We give a characterisation of the power of CLAP in terms of a minion homomorphism. Using this characterisation, we identify a certain weak notion of symmetry which, if satisfied by infinitely many polymorphisms of PCSPs, guarantees tractability.

We demonstrate that there are PCSPs solved by CLAP that are not solved by any of the existing algorithms for PCSPs; in particular, not by the BLP + AIP algorithm of Brakensiek et al. [SICOMP’20] and not by a reduction to tractable finite-domain CSPs.

## 1 Introduction

**Constraint Satisfaction** Constraint Satisfaction Problems (CSPs) have driven some of the most influential developments in theoretical computer science, from NP-completeness to the PCP theorem [1, 2, 39] to semidefinite programming algorithms [63] to the Unique Games Conjecture [53].

A CSP over domain  $A$  is specified by a finite collection  $\mathbf{A}$  of relations over  $A$ , and is denoted by  $\text{CSP}(\mathbf{A})$ . Given on input a set of variables and a set of constraints, each of which uses relations from  $\mathbf{A}$ , the task is to decide the existence of an assignment of values from  $A$  to the variables that satisfies all the constraints. Classic examples of CSPs include 2-SAT, graph 3-colouring, and linear equations of fixed width over finite groups.

For Boolean CSPs, which are CSPs with  $|A| = 2$ , Schaefer proved that every such CSP is either solvable in polynomial time or is NP-complete [65]. Feder and Vardi famously conjectured that the same holds true for CSPs over arbitrary finite domains [41]. Furthermore, they realised the importance of considering closure properties of solution spaces of CSPs [41], which initiated the algebraic approach [26, 49, 50]. The key notion in the algebraic approach is that

---

\*An extended abstract of this work appeared in the Proceedings of the 2022 ACM-SIAM Symposium on Discrete Algorithms (SODA’22) [32]. The research leading to these results has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 714532). Stanislav Živný was supported by a Royal Society University Research Fellowship. The paper reflects only the authors’ views and not the views of the ERC or the European Commission. The European Union is not liable for any use that may be made of the information contained therein. This work was also supported by UKRI EP/X024431/1. For the purpose of Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. All data is provided in full in the results section of this paper.

<sup>†</sup>Department of Computer Science, University of Oxford, UK.

of *polymorphisms*, which are operations that take solutions to a CSP and are guaranteed to return, by a coordinatewise application, a solution to the same CSP. All CSPs admit projections (also known as dictators) as polymorphisms. However, the presence of less trivial polymorphisms, satisfying some notion of symmetry, is necessary for tractability. For instance, the set of solutions to 2-SAT is closed under the ternary majority operation  $\text{maj} : \{0, 1\}^3 \rightarrow \{0, 1\}$  that satisfies the following notion of symmetry:  $\text{maj}(a, a, b) = \text{maj}(a, b, a) = \text{maj}(b, a, a) = a$  for any  $a, b \in \{0, 1\}$ . Similarly, the set of solutions to Horn-SAT is closed under the binary minimum operation  $\text{min} : \{0, 1\}^2 \rightarrow \{0, 1\}$  that satisfies a different notion of symmetry:  $\text{min}(a, a) = a$ ,  $\text{min}(a, b) = \text{min}(b, a)$ , and  $\text{min}(a, \text{min}(b, c)) = \text{min}(\text{min}(a, b), c)$  for any  $a, b, c \in \{0, 1\}$ . The binary max operation – which is a polymorphism of dual Horn-SAT – has the same notion of symmetry, called semilattice [11]. Together with the ternary minority polymorphism, which captures linear equations on  $\{0, 1\}$ , this gives all non-trivial tractable cases from Schaefer’s dichotomy result.<sup>1</sup>

The polymorphisms of any CSP form a clone, in that they include all projections and are closed under composition. For instance, since Horn-SAT has  $\text{min}$  as a polymorphism, it also has the 4-ary minimum operation  $\text{min}_4(a, b, c, d) = \text{min}(a, \text{min}(b, \text{min}(c, d)))$  as a polymorphism. Building on the connection to universal algebra, the algebraic approach has been tremendously successful beyond decision CSPs, e.g. for robust satisfiability of CSPs [10, 35, 36], for exact optimisation of CSPs [54, 57, 66], and for characterising the power of algorithms [9, 14, 55, 56, 59, 67, 68]. The culmination of the algebraic approach is the positive resolution of the dichotomy conjecture by Bulatov [28] and Zhuk [71]. We refer the reader to [11] for a survey on the algebraic approach.

**Promise Constraint Satisfaction** In this paper, we study Promise Constraint Satisfaction Problems (PCSPs), whose systematic study was initiated by Austrin, Guruswami, and Håstad [5], and Brakensiek and Guruswami [20]. PCSPs form a vast generalisation of CSPs. In  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ , each constraint comes in two forms, a strict one in  $\mathbf{A}$  and a weak one in  $\mathbf{B}$ . The goal is to distinguish between (i) the case in which (the strong form of) the constraints can be simultaneously satisfied in  $\mathbf{A}$  and (ii) the case in which (even the weak form of) the constraints cannot be simultaneously satisfied in  $\mathbf{B}$ . The promise is that it is never the case that the PCSP is not satisfiable in the strict sense but is satisfiable in the weak sense. If the strict and weak forms coincide in every constraint (i.e., if  $\mathbf{A} = \mathbf{B}$ ) we get the (non-promise) CSPs. However, PCSPs include many fundamental problems that are inexpressible as CSPs.

The simplest example of strict vs. weak constraints is when the weak constraints are supersets of the strict constraints on the same domain (the first two examples below) or on a larger domain (the third example below); the notion of homomorphism from  $\mathbf{A}$  to  $\mathbf{B}$  formalises this for any PCSP.

First, can we distinguish a  $g$ -satisfiable  $k$ -SAT instance (in the sense that there is an assignment that satisfies at least  $g$  literals in each clause) from an instance that is not even 1-satisfiable? This problem was studied in [5], where it was shown to be solvable in polynomial time if  $\frac{g}{k} \geq \frac{1}{2}$  and NP-complete otherwise. Recently, this result has been generalised to arbitrary finite domains [23].

Second, can we distinguish a 3-SAT formula that admits an assignment satisfying exactly 1 literal in each clause (i.e., a satisfiable instance of **1-in-3-SAT**) from one that does not

---

<sup>1</sup>The trivial cases, called 0- and 1-valid, are captured by the constant-0 and constant-1 polymorphisms, respectively.

admit an assignment satisfying 1 or 2 literals in each clause (i.e., a non-satisfiable instance of **Not-All-Equal-3-SAT**)? Remarkably, while both **1-in-3** and **NAE** are NP-hard, this promise version is solvable in polynomial time [19, 20].

Third, can we distinguish a  $k$ -colourable graph from a graph that is not even  $\ell$ -colourable, where  $k \leq \ell$ ? This is the *approximate graph colouring* problem, which is believed to be NP-hard for any fixed  $3 \leq k \leq \ell$ , but has been elusive since the 1970s [43]. In particular, the larger the gap is between  $k$  and  $\ell$  the easier the problem could in principle be and, thus, the more challenging it is to prove NP-hardness. The current state of the art is NP-hardness for  $k = 3$  and  $\ell = 5$  [8], while already the case of  $k = 3$  and  $\ell = 6$  is open. For any  $k \geq 4$  and  $\ell = \ell(k) = \binom{k}{\lfloor k/2 \rfloor} - 1$ , NP-hardness has been established in [70].

While a systematic study of PCSPs was initiated only recently [5, 20], concrete PCSPs have been considered for a while, e.g. approximate graph [16, 43, 44, 51, 52, 69] and hypergraph colouring [40]. A highlight result is the dichotomy of Boolean symmetric PCSPs [42] (in which all constraint relations are symmetric), following an earlier classification of Boolean symmetric PCSPs with disequalities [20]. Very recent works have investigated certain Boolean non-symmetric PCSPs [24] and certain non-Boolean symmetric PCSPs [7]. Other recent results include, e.g., [4, 21, 45].

Most of the recent progress, including results on the approximate graph colouring problem [8, 70] and on the approximate graph homomorphism problem [58, 70], rely on the algebraic approach to PCSPs [8]. In particular, the breakthrough results in [8], building on [12], established that the complexity of PCSPs is captured by the polymorphism minions and certain types of symmetries these minions satisfy – these are non-nested identities on polymorphisms, such as the majority example but not the semilattice example. Crucially, minions are less structured than clones: A minion (of functions) is a set of operations closed under permuting coordinates, identifying coordinates, and introducing dummy coordinates, but not under composition.<sup>2</sup> Thus, unlike in our earlier CSP example (corresponding to Horn-SAT), a binary minimum polymorphism of a PCSP cannot in general be used to generate a 4-ary minimum polymorphism of the same PCSP.

Despite the momentous results in [8], there is a long way to go to classify all PCSPs, and it is not even clear whether a dichotomy for all PCSPs should be expected. When Feder and Vardi conjectured a CSP dichotomy [41], the Boolean case [65] and the graph case [46] had been fully classified. We seem quite far from these two cases being classified for PCSPs. Thus, further progress is needed on both the hardness and tractability part. This paper focusses on the latter.

**Finite tractability** Although PCSPs are (much) more general than CSPs, some PCSPs can be reduced to tractable CSPs. This idea was introduced in [19] under the name of homomorphic sandwiching (cf. Section 2 for a precise definition); PCSPs that are reducible to tractable (finite-domain) CSPs are called *finitely tractable*. Finite tractability is not sufficient to explain tractability of all tractable PCSPs. In particular, Barto et al. [8] showed that the above-mentioned example **1-in-3** vs. **NAE** is not finitely tractable, despite being a tractable PCSP [20]. We remark that it is not inconceivable (and in fact was conjectured in [19]) that every tractable (finite-domain) PCSP could be reducible to a tractable CSP possibly over an infinite domain; this is the case for the **1-in-3** vs. **NAE** problem [19]. However, while certain

---

<sup>2</sup>In this work, we shall use the more abstract notion of minion introduced in [22], cf. Definition 6.

infinite-domain CSPs are amenable to algebraic methods, the complexity of infinite-domain CSPs is far from understood, cf. [13, 17, 18] for recent work.

Since finite tractability does not capture all tractable PCSPs, there is need for other algorithmic tools. One possibility is to attempt to extend algorithmic techniques developed for CSPs.

There are two main algorithmic approaches to CSPs. On the one hand, there are local consistency methods [41], which have been studied in theoretical computer science but also in artificial intelligence, logic, and database theory. The power of local consistency for CSPs has been characterised in [9, 25], and it is known that the third level of consistency solves all so-called bounded-width CSPs [6]. On the other hand, there are CSPs solvable by algorithms based on generalisations of Gaussian elimination, most notably CSPs with a Mal'tsev polymorphism [29]. This method has been pushed to its limit, in a way, in [14, 48]. While the NP-hardness part of the CSP dichotomy has been known since [26], the challenge in proving the algorithmic part is the complicated interaction of these two very different algorithmic approaches. Although this interaction does not occur in Boolean CSPs, it occurs already in CSPs on three-element domains [27].

The characterisation of the power of the first level of the consistency methods, 1-consistency (also known as arc-consistency [61]), has been lifted from CSPs [41] to PCSPs in [8]. Rather than establishing 1-consistency combinatorially, one can employ convex relaxations.

**Relaxations** A canonical analogue of 1-consistency is the *basic linear programming relaxation* (BLP) [59], which in fact is stronger than 1-consistency [60]. The characterisation of the power of BLP has been lifted from CSPs [59] to PCSPs in [8], both in terms of a minion and a property of polymorphisms. The power of BLP is captured by a minion consisting of rational stochastic vectors<sup>3</sup> or, equivalently, by the presence of symmetric polymorphisms of all arities; these are polymorphisms invariant under any permutation of the coordinates. For example, we have seen that Horn-SAT, a classic CSP, has a binary symmetric polymorphism, namely min. We have also seen that min can generate a 4-ary operation  $\min_4$ , which is symmetric. Similarly, min can generate (via composition) symmetric operations of all arities, and thus Horn-SAT is solved by BLP.

A different relaxation of PCSPs is the *basic affine integer programming relaxation* (AIP) [19]. The power of AIP has been characterised, both in terms of a minion and a property of polymorphisms, in [8]. The minion capturing AIP consists of integer affine vectors.<sup>4</sup> Concerning polymorphisms, AIP is captured by polymorphisms of all odd arities that are invariant under permutations that only permute odd and even coordinates separately, and additionally satisfy that adjacent coordinates cancel each other out. The **1-in-3** vs. **NAE** problem is solved by AIP (cf. Example 5).

Brakensiek et al. [22] proposed a combination of the two above-mentioned relaxations, called BLP + AIP. Their algorithm has many interesting features. Firstly, it solves PCSPs that admit only infinitely many symmetric polymorphisms (i.e., not all arities are required as in the case of BLP). Secondly, it solves all tractable Boolean CSPs, thus demonstrating how research on PCSPs can shed new light on (non-promise) CSPs. In fact, [22] established the power of BLP + AIP in terms of a minion and (a property of) polymorphisms. The minion capturing BLP + AIP is essentially a product of the BLP and AIP minions [22]. Concerning

<sup>3</sup>A vector is *stochastic* if its entries are nonnegative and sum up to one.

<sup>4</sup>An integer vector is *affine* if its entries sum up to one.

polymorphisms, BLP + AIP is captured by polymorphisms of all odd arities that are invariant under permutations that only permute odd and even coordinates.

It may be that BLP + AIP is the only algorithm needed to solve all tractable *Boolean* PCSPs. However, as already observed in [22], BLP + AIP does not solve some rather simple, tractable, *non-Boolean* PCSPs. Motivated by this, we investigate algorithms that are stronger than BLP + AIP. We note that all PCSPs hitherto known to be tractable are solved by BLP + AIP or by finite tractability (i.e., by a reduction to a tractable finite-domain CSP). In this work, we provide an example of a PCSP that is tractable (through our algorithm) but is not solved by either of those two algorithmic techniques.

**Contributions** Building on the work of Brakensiek et al. [22], we study stronger relaxations for PCSPs and give three main contributions.

**(1) CLAP** Our first contribution is the introduction of CLAP to the study of PCSPs. Our goal was to design an algorithm that, unlike BLP + AIP, solves all CSPs of bounded width. While all bounded-width CSPs can be solved by 3-consistency [6], and thus also by the third level of the Sherali-Adams hierarchy for BLP (e.g., by [67]), Kozik showed that already (a special case of) the singleton arc-consistency (SAC) algorithm, introduced in [38] (cf. [15, 30]), solves all bounded-width CSPs [56]. Thus, we study the LP relaxation that we call the *singleton* BLP (SBLP), which is at least as strong as SAC. A special case of SBLP (without this name) implicitly appeared in the literature, e.g. in [5, 20] for Boolean PCSPs. The idea behind SBLP is essentially to run SAC but replace the arc-consistency check by the BLP; i.e., the algorithm repeatedly takes a variable-value pair  $(x, a)$  and tests the feasibility of the BLP with the requirement that  $x$  should be assigned the value  $a$ . If this LP is infeasible then  $a$  is removed from the domain of  $x$ . This is repeated until convergence. If any variable ends up with an empty domain then SBLP rejects, otherwise it accepts. Overall, the number of BLP calls occurring for an instance of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  with variable-set  $X$  is at most polynomial in the size of  $X$ . As mentioned above, this simple algorithm solves all bounded-width CSPs [56].

We adopt a modification of SBLP that turns out to be more naturally captured by a minion-oriented analysis: the *constraint* BLP (CBLP). This (possibly) stronger algorithm is a generalisation of SBLP in which we do not consider only variable-value pairs  $(x, a)$ , but rather the constraint-assignment pairs  $(\mathbf{x}, \mathbf{a})$  for every constraint in the instance. As in SBLP, if fixing a (local) assignment to a constraint yields an infeasible BLP then the assignment is removed from the constraint relation. Upon convergence, which takes at most polynomially many BLP calls, if any constraint ends up with an empty relation then CBLP rejects, otherwise it accepts.

Our algorithm CLAP first runs CBLP and then, upon termination, refines the solutions of CBLP by running (essentially) AIP. If one believes the suggestion in [22] that constantly many rounds of the Sherali-Adams hierarchy for BLP + AIP could solve all tractable (non-promise) CSPs, then it is not outrageous to believe that the same could be true for CLAP, and CLAP might be easier to analyse than such an algorithm.

**(2) Characterisation** Our second contribution is a minion characterisation of the power of CLAP, stated as Theorem 13. The objects in the minion are essentially matrices with a particular structure, which we call *skeletal* (cf. Definition 11). These matrices capture the CBLP part of CLAP and together with certain integer affine vectors form the minion (cf. Definition 12). Another, more conceptual contribution is the introduction of a minion of



matrices to the study of PCSPs.

**(3) H-symmetric polymorphisms** The minion characterisation is crucial to our third contribution: the identification of a sufficient condition for CLAP to work in terms of the symmetries of the polymorphisms. This is stated as Theorem 15, using the notion of  $H$ -symmetry. This condition can be more easily checked for concrete templates, thus allowing us to design a separating example that is not finitely tractable and is not solved by BLP + AIP (nor by local consistency methods, see [3]), but is solved by CLAP. It follows that our new algorithm is strictly more powerful than BLP + AIP (and separated by an interesting PCSP that is not reducible to a tractable finite-domain CSP via “gadget reductions”, which capture the algebraic approach to PCSPs [8]).

For a matrix  $H$ , a polymorphism  $f$  is  $H$ -symmetric if  $f$  is invariant under permutations of the coordinates but only on a specific set of inputs determined by  $H$  (cf. Definition 14). For instance, if  $H$  is a row vector then we obtain the requirement that  $f$  be symmetric on all inputs. If  $H$  is the identity matrix then we require that  $f$  be symmetric only on inputs in which different entries occur with different multiplicities. In general, the intuition is that we capture ‘symmetry with exceptions that depend on multiplicities’. We refer the reader to the discussion in Section 3 for details.

After necessary background material in Section 2, our algorithm CLAP and the main results are presented in Section 3; the proofs appear in Sections 4 and 5.

## 2 Preliminaries

We let  $\mathbb{N} = \{1, 2, \dots\}$  and  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ . The cardinality of  $\mathbb{N}$  shall be denoted by  $\aleph_0$ . For  $k \in \mathbb{N}$ ,  $[k]$  denotes the set  $\{1, \dots, k\}$ . For a set  $A$ ,  $\mathcal{P}(A)$  denotes the set of all subsets of  $A$ . We denote by  $\leq_p$  many-one polynomial-time reductions. We shall use standard notation for vectors and matrices. Vectors will be treated as column vectors and whenever convenient identified with the corresponding (row) tuples. Both tuples and vectors will be typed in bold font. We denote by  $\mathbf{e}_i$  the  $i$ -th standard unit vector of the appropriate size (which will be clear from the context); i.e.,  $\mathbf{e}_i$  is equal to 1 in the  $i$ -th coordinate and 0 elsewhere. We denote by  $\mathbf{0}_p$  and by  $\mathbf{1}_p$  the all-zero and all-one vector, respectively, of size  $p$ ; if the size is clear, we occasionally drop the subscript. The *support* of a vector  $\mathbf{v} = (v_i)$  of size  $p$  is the set  $\text{supp}(\mathbf{v}) = \{i \in [p] : v_i \neq 0\}$ .  $I_p$  denotes the identity matrix of order  $p$ , while  $O$  denotes an all-zero matrix of suitable size.

**Promise CSPs** A *signature*  $\sigma$  is a finite set of relation symbols  $R$ , each with its arity  $\text{ar}(R) \in \mathbb{N}$ . A *relational structure* over a signature  $\sigma$ , or a  $\sigma$ -structure, is a finite universe  $A$ , called the *domain* of  $\mathbf{A}$ , and a relation  $R^{\mathbf{A}} \subseteq A^{\text{ar}(R)}$  for each symbol  $R \in \sigma$ . For two  $\sigma$ -structures  $\mathbf{A}$  and  $\mathbf{B}$ , a mapping  $f : A \rightarrow B$  is called a *homomorphism* from  $\mathbf{A}$  to  $\mathbf{B}$ , denoted by  $f : \mathbf{A} \rightarrow \mathbf{B}$ , if  $f$  preserves all relations; that is, for every  $R \in \sigma$  and every tuple  $\mathbf{a} \in R^{\mathbf{A}}$ , we have  $f(\mathbf{a}) \in R^{\mathbf{B}}$ , where  $f$  is applied coordinatewise. The existence of a homomorphism from  $\mathbf{A}$  to  $\mathbf{B}$  is denoted by  $\mathbf{A} \rightarrow \mathbf{B}$ . A PCSP *template* is a pair  $(\mathbf{A}, \mathbf{B})$  of relational structures over the same signature such that  $\mathbf{A} \rightarrow \mathbf{B}$ . Without loss of generality, we will often assume that  $A$ , the domain of  $\mathbf{A}$ , is  $[n]$ .

**Definition 1.** Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template. The *decision version* of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  is the following problem: Given as input a relational structure  $\mathbf{X}$  over the same signature as  $\mathbf{A}$  and

$\mathbf{B}$ , output YES if  $\mathbf{X} \rightarrow \mathbf{A}$  and NO if  $\mathbf{X} \not\rightarrow \mathbf{B}$ . The *search version* of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  is the following problem: Given as input a relational structure  $\mathbf{X}$  over the same signature as  $\mathbf{A}$  and  $\mathbf{B}$  and such that  $\mathbf{X} \rightarrow \mathbf{A}$ , find a homomorphism from  $\mathbf{X}$  to  $\mathbf{B}$ .

For a relational structure  $\mathbf{A}$ , the *constraint satisfaction problem* (CSP) with template  $\mathbf{A}$  [41], denoted by  $\text{CSP}(\mathbf{A})$ , is  $\text{PCSP}(\mathbf{A}, \mathbf{A})$ .

**Example 2.** For  $k \geq 2$ , let  $\mathbf{K}_k$  be the structure with domain  $[k]$  and a binary relation  $\{(i, j) \in [k]^2 \mid i \neq j\}$ . Then,  $\text{CSP}(\mathbf{K}_k)$  is the standard graph  $k$ -colouring problem. For  $k \leq \ell$ ,  $\text{PCSP}(\mathbf{K}_k, \mathbf{K}_\ell)$  is the *approximate graph colouring problem* [43]. In the decision version, the task is to decide whether a graph is  $k$ -colourable or not even  $\ell$ -colourable. In the search version, given a  $k$ -colourable graph  $G$ , the task is to find an  $\ell$ -colouring of  $G$ . It is widely believed that for any fixed  $3 \leq k \leq \ell$ ,  $\text{PCSP}(\mathbf{K}_k, \mathbf{K}_\ell)$  is NP-hard; i.e., constantly many colours do not help. The current most general NP-hardness result is known for  $k = 3$  and  $\ell = 5$  by Bulín, Krokhin, and Opršal [8] and for  $k \geq 4$  and  $\ell = \ell(k) = \binom{k}{\lfloor k/2 \rfloor} - 1$  by Wrochna and Živný [70].

We call a PCSP template  $(\mathbf{A}, \mathbf{B})$  *tractable* if any instance of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  can be solved in polynomial time in the size of the input structure  $\mathbf{X}$ . It is easy to show that the decision version reduces to the search version [8] (but the converse is not known in general); for CSPs, the two versions are equivalent [26, 33]. Our results are for the decision version.

**Example 3.** Let **1-in-3** be the Boolean structure with domain  $\{0, 1\}$  and a single ternary relation  $\{(0, 0, 1), (0, 1, 0), (1, 0, 0)\}$ . Let **NAE** be the structure with domain  $\{0, 1\}$  and a single ternary relation  $\{0, 1\}^3 \setminus \{(0, 0, 0), (1, 1, 1)\}$ . Then,  $\text{CSP}(\mathbf{1-in-3})$  is the (positive) 1-in-3-SAT problem and  $\text{CSP}(\mathbf{NAE})$  is the (positive) Not-All-Equal-3-SAT problem. Since both of these problems are NP-hard [65], the PCSP templates  $(\mathbf{1-in-3}, \mathbf{1-in-3})$  and  $(\mathbf{NAE}, \mathbf{NAE})$  are both intractable. However, the PCSP template  $(\mathbf{1-in-3}, \mathbf{NAE})$  is tractable, as shown by Brakensiek and Guruswami [20].

**Definition 4.** Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template with signature  $\sigma$ . An operation  $f : A^L \rightarrow B$ , where  $L \in \mathbb{N}$ , is a *polymorphism* of arity  $L$  of  $(\mathbf{A}, \mathbf{B})$  if for every  $R \in \sigma$  of arity  $k = \text{ar}(R)$  and for any possible  $L \times k$  matrix whose rows are tuples in  $R^{\mathbf{A}}$ , the application of  $f$  on the columns of the matrix gives a tuple in  $R^{\mathbf{B}}$ . We denote by  $\text{Pol}(\mathbf{A}, \mathbf{B})$  the set of all polymorphisms of  $(\mathbf{A}, \mathbf{B})$ .

**Example 5.** The unary operation  $\neg : \{0, 1\} \rightarrow \{0, 1\}$  defined by  $\neg(a) = 1 - a$  is a polymorphism of  $(\mathbf{NAE}, \mathbf{NAE})$  but not a polymorphism of  $(\mathbf{1-in-3}, \mathbf{1-in-3})$ . For any odd  $L$ , the  $L$ -ary operation  $f : \{0, 1\}^L \rightarrow \{0, 1\}$  defined by  $f(a_1, \dots, a_L) = 1$  if  $a_1 - a_2 + a_3 - \dots + a_L > 0$  and  $f(a_1, \dots, a_L) = 0$  otherwise is a polymorphism of  $(\mathbf{1-in-3}, \mathbf{NAE})$ .

**Minions** Polymorphisms of CSPs form clones; i.e.,  $\text{Pol}(\mathbf{A}, \mathbf{A})$  contains all projections (also known as dictators) and is closed under composition [11]. Polymorphisms of the (more general) PCSPs form *minions*; i.e., they are closed under taking minors.<sup>5</sup> Formally, given an  $L$ -ary function  $f : A^L \rightarrow B$ , its *minor* relative to a map  $\pi : [L] \rightarrow [L']$  is the  $L'$ -ary function  $f_{/\pi} : A^{L'} \rightarrow B$  defined by

$$f_{/\pi}(a_1, \dots, a_{L'}) = f(a_{\pi(1)}, \dots, a_{\pi(L)}). \quad (1)$$

---

<sup>5</sup>We remark that clones are also closed under taking minors.

Equivalently, a minor of  $f$  is a function obtained from  $f$  by identifying variables, permuting variables, and introducing dummy variables. Rather than focussing on minions of functions, we consider here abstract minions, as described and used in [22].

**Definition 6.** A *minion*  $\mathcal{M}$  consists in the disjoint union of sets  $\mathcal{M}^{(L)}$  for  $L \in \mathbb{N}$  equipped with operations  $(\cdot)_{/\pi} : \mathcal{M}^{(L)} \rightarrow \mathcal{M}^{(L')}$  for all functions  $\pi : [L] \rightarrow [L']$ , which satisfy

- $(M_{/\pi})_{/\tilde{\pi}} = M_{/\tilde{\pi} \circ \pi}$  for  $\pi : [L] \rightarrow [L']$ ,  $\tilde{\pi} : [L'] \rightarrow [L'']$  and
- $M_{/\text{id}} = M$

for all  $M \in \mathcal{M}^{(L)}$ .

**Definition 7.** For two minions  $\mathcal{M}$  and  $\mathcal{N}$ , a *minion homomorphism*  $\xi : \mathcal{M} \rightarrow \mathcal{N}$  is a map that preserves arities and minors: Given  $M \in \mathcal{M}^{(L)}$  and  $\pi : [L] \rightarrow [L']$ ,  $\xi(M) \in \mathcal{N}^{(L')}$  and  $\xi(M_{/\pi}) = \xi(M)_{/\pi}$ .

For any PCSP template  $(\mathbf{A}, \mathbf{B})$ , the set  $\text{Pol}(\mathbf{A}, \mathbf{B})$  of its polymorphisms equipped with the operations described by (1) is a minion [8]. One of the results in [8] established that minion homomorphisms give rise to polynomial-time reductions: If there is a minion homomorphism from  $\text{Pol}(\mathbf{A}, \mathbf{B})$  to  $\text{Pol}(\mathbf{A}', \mathbf{B}')$ , then  $\text{PCSP}(\mathbf{A}', \mathbf{B}') \leq_p \text{PCSP}(\mathbf{A}, \mathbf{B})$ . Minions are also useful for characterising the power of algorithms, as we will discuss later.

**Remark 8.** Although we will not use this categorical view, we remark that a minion is nothing but a functor from the category of nonempty finite sets to the category of nonempty sets, and a minion homomorphism is a natural transformation.

**Existing algorithms** One way to establish tractability of PCSPs is to reduce to CSPs. Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template. A structure  $\mathbf{C}$  is called a (homomorphic) *sandwich* if  $\mathbf{A} \rightarrow \mathbf{C} \rightarrow \mathbf{B}$ . It is known that, in this case,  $\text{PCSP}(\mathbf{A}, \mathbf{B}) \leq_p \text{CSP}(\mathbf{C})$ .<sup>6</sup> Thus, if  $\mathbf{C}$  is a tractable CSP template then  $(\mathbf{A}, \mathbf{B})$  is a tractable PCSP template. If  $\mathbf{C}$  has a finite domain, we say that  $(\mathbf{A}, \mathbf{B})$  is *finitely tractable*.

**Example 9.** The PCSP template **(1-in-3, NAE)** from Example 3 is tractable [20] but not finitely tractable unless  $\text{P}=\text{NP}$ , as shown in [8].

Another way to establish tractability for PCSPs is to leverage convex relaxations. In Section 1, we mentioned three studied relaxations: BLP [59], AIP [20], and BLP + AIP [22]. Their powers have been characterised in [8, 22] in terms of certain minions and polymorphism identities. The details of these relaxations and the characterisations are provided in Appendix A.

All PCSPs hitherto known to be tractable are solved by finite tractability (i.e., by a reduction to a tractable finite-domain CSP) or by BLP + AIP. The next example identifies a simple PCSP template not captured by either of these two methods.

**Example 10.** Consider the relational structures  $\mathbf{A} = (A; R_1^{\mathbf{A}}, R_2^{\mathbf{A}})$  and  $\mathbf{B} = (B; R_1^{\mathbf{B}}, R_2^{\mathbf{B}})$  on the domain  $A = B = \{0, \dots, 6\}$  with the following relations:  $R_1^{\mathbf{A}} = \{(0, 0, 1), (0, 1, 0), (1, 0, 0)\}$  is **1-in-3** on  $\{0, 1\}$ ,  $R_1^{\mathbf{B}} = \{0, 1\}^3 \setminus \{(0, 0, 0), (1, 1, 1)\}$  is **NAE** on  $\{0, 1\}$ , and  $R_2^{\mathbf{A}} = R_2^{\mathbf{B}} = \{(2, 3), (3, 2), (4, 5), (5, 6), (6, 4)\}$ . The identity mapping is a homomorphism from  $\mathbf{A}$  to  $\mathbf{B}$ ,

---

<sup>6</sup>This is a special case of homomorphic relaxation [8], which we do not need here.



so  $(\mathbf{A}, \mathbf{B})$  is a PCSP template. Since the directed graph corresponding to  $R_2^{\mathbf{A}} = R_2^{\mathbf{B}}$  is a disjoint union of a directed 2-cycle and a directed 3-cycle, [22, Example 6.1] shows that the BLP + AIP algorithm does not solve  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ . We claim that the template  $(\mathbf{A}, \mathbf{B})$  is not finitely tractable. For contradiction, assume that there is a finite relational structure  $\mathbf{C} = (C; R_1^{\mathbf{C}}, R_2^{\mathbf{C}})$  such that  $\mathbf{A} \rightarrow \mathbf{C} \rightarrow \mathbf{B}$  and  $\text{CSP}(\mathbf{C})$  is tractable. We will argue that this would imply finite tractability of  $(\mathbf{1-in-3}, \mathbf{NAE})$ , which contradicts the result in [8] (unless  $\text{P}=\text{NP}$ ); cf. Example 9. Indeed, the existence of such  $\mathbf{C}$  gives the following chain of homomorphisms:

$$\mathbf{1-in-3} = (\{0, 1\}; R_1^{\mathbf{A}}) \rightarrow (A; R_1^{\mathbf{A}}) \rightarrow (C; R_1^{\mathbf{C}}) \rightarrow (B; R_1^{\mathbf{B}}) \rightarrow (\{0, 1\}; R_1^{\mathbf{B}}) = \mathbf{NAE} \quad (2)$$

where the first map is the inclusion of  $\{0, 1\}$  in  $A$ , the second and the third are the maps witnessing  $\mathbf{A} \rightarrow \mathbf{C} \rightarrow \mathbf{B}$ , and the fourth is any map  $g : B \rightarrow \{0, 1\}$  such that  $g(0) = 0$  and  $g(1) = 1$ . Let  $\tilde{\mathbf{C}} = (C; R_1^{\mathbf{C}})$ . Observe that  $\tilde{\mathbf{C}}$  is tractable since the inclusion map gives a minion homomorphism  $\text{Pol}(\mathbf{C}, \mathbf{C}) \rightarrow \text{Pol}(\tilde{\mathbf{C}}, \tilde{\mathbf{C}})$ , and thus  $\text{CSP}(\tilde{\mathbf{C}}) = \text{PCSP}(\tilde{\mathbf{C}}, \tilde{\mathbf{C}}) \leq_p \text{PCSP}(\mathbf{C}, \mathbf{C}) = \text{CSP}(\mathbf{C})$  by [8, Theorem 3.1]. This proves the claim, as (2) established  $\mathbf{1-in-3} \rightarrow \tilde{\mathbf{C}} \rightarrow \mathbf{NAE}$ .

Notice that the assignment  $f \mapsto g \circ f|_{\{0,1\}^L}$  (where  $f$  is a polymorphism of  $(\mathbf{A}, \mathbf{B})$  of arity  $L$  and  $g$  is the map considered above) yields a minion homomorphism from  $\text{Pol}(\mathbf{A}, \mathbf{B})$  to  $\text{Pol}(\mathbf{1-in-3}, \mathbf{NAE})$ . As established in [3, Corollary 4.2], the template  $(\mathbf{1-in-3}, \mathbf{NAE})$  does not have bounded width – i.e., is not solved by local consistency methods. It follows from [8, Lemma 7.5] that  $(\mathbf{A}, \mathbf{B})$  does not have bounded width either.

The template from Example 10 will be proved tractable later (in Example 16) using our new algorithm, which we will present next.

### 3 The CLAP algorithm

Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template with signature  $\sigma$  and let  $\mathbf{X}$  be an instance of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ . Without loss of generality, we assume that  $\sigma$  contains a unary symbol  $R_u$  such that  $R_u^{\mathbf{X}} = X$ ,  $R_u^{\mathbf{A}} = A$ , and  $R_u^{\mathbf{B}} = B$ . If this is not the case, the signature and the instance can be extended without changing the set of solutions. Our algorithm – the combined **CBLP**+**AIP** algorithm (CLAP), presented in Algorithm 1 and discussed below – builds on BLP [8] and BLP + AIP [22].

CLAP works in two stages. In the first stage, it runs CBLP; i.e., a modified version of the singleton arc-consistency algorithm (cf. [38]) where (i) the “arc-consistency” part is replaced by BLP, and (ii) the “singleton” part is boosted by requiring that every constraint-assignment pair (as opposed to every variable-value pair) is fixed at each iteration. In the second stage, it refines CBLP by doing an additional sanity check: At least one of the solutions computed by CBLP should be compatible with a solution of AIP. As in [22], this second stage requires that the AIP solution should only use those variables from the CBLP solution that have nonzero weight. There are two equivalent ways to enforce this requirement: Either by storing the nonzero variables at each iteration of CBLP in the first stage of the algorithm, or by simply running BLP + AIP as a black box in the second stage of the algorithm. We adopt the latter option to achieve a simpler presentation. Concretely, the first stage of CLAP is performed by initialising the sets  $S_{\mathbf{x}, R}$  of constraint-assignment pairs to the entire relation  $R^{\mathbf{A}}$ , and then progressively shrinking these sets by cycling over all constraint-assignment pairs and removing a pair whenever it yields an infeasible BLP. The second stage, that occurs if

all sets  $S_{\mathbf{x},R}$  are nonempty, is performed by cycling over each feasible constraint-assignment pair and running BLP + AIP on it. As soon as one constraint-assignment pair is accepted by BLP + AIP, the algorithm terminates and outputs YES. If no constraint-assignment pair is accepted, the algorithm outputs NO.

As in Appendix A, where BLP, AIP, and BLP + AIP are presented in full detail for completeness, by  $\lambda_{\mathbf{x},R}(\mathbf{a})$  we denote the variable of  $\text{BLP}(\mathbf{X}, \mathbf{A})$  associated with  $\mathbf{x} \in R^{\mathbf{X}}$  and  $\mathbf{a} \in R^{\mathbf{A}}$ , where  $R \in \sigma$ . The algorithm has polynomial time complexity in the size of the input instance: Letting  $g = \sum_{R \in \sigma} |R^{\mathbf{X}}| |R^{\mathbf{A}}|$ ,  $\mathcal{O}(g^2)$  BLP calls and  $\mathcal{O}(g)$  BLP + AIP calls occur. We say that CLAP *accepts* an instance  $\mathbf{X}$  of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  if Algorithm 1 returns YES. We say that CLAP *solves*  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  if, for every instance  $\mathbf{X}$  of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ , we have (i) if  $\mathbf{X} \rightarrow \mathbf{A}$  then CLAP accepts  $\mathbf{X}$ , and (ii) if  $\mathbf{X}$  is accepted by CLAP then  $\mathbf{X} \rightarrow \mathbf{B}$ .

---

**Algorithm 1:** The CLAP algorithm

---

**Input:** an instance  $\mathbf{X}$  of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  of signature  $\sigma$   
**Output:** YES if  $\mathbf{X} \rightarrow \mathbf{A}$  and NO if  $\mathbf{X} \not\rightarrow \mathbf{B}$

```

1 for  $R \in \sigma, \mathbf{x} \in R^{\mathbf{X}}$  do
2   | set  $S_{\mathbf{x},R} := R^{\mathbf{A}}$ ;
3 end
4 repeat
5   | for  $R \in \sigma, \mathbf{x} \in R^{\mathbf{X}}, \mathbf{a} \in S_{\mathbf{x},R}$  do
6     | if  $\text{BLP}(\mathbf{X}, \mathbf{A})$  with  $\lambda_{\mathbf{x},R}(\mathbf{a}) = 1$  and  $\lambda_{\mathbf{x}',R'}(\mathbf{a}') = 0$  for every  $R' \in \sigma, \mathbf{x}' \in R'^{\mathbf{X}}$ ,
7       |   and  $\mathbf{a}' \notin S_{\mathbf{x}',R'}$  is not feasible then
8         |   | remove  $\mathbf{a}$  from  $S_{\mathbf{x},R}$ ;
9         |   end
10    | end
11 until no set  $S_{\mathbf{x},R}$  is changed;
12 if some  $S_{\mathbf{x},R}$  is empty then
13   | return NO;
14 else
15   | for  $R \in \sigma, \mathbf{x} \in R^{\mathbf{X}}, \mathbf{a} \in S_{\mathbf{x},R}$  do
16     | if  $\text{BLP} + \text{AIP}(\mathbf{X}, \mathbf{A})$  with  $\lambda_{\mathbf{x},R}(\mathbf{a}) = 1$  and  $\lambda_{\mathbf{x}',R'}(\mathbf{a}') = 0$  for every  $R' \in \sigma,$ 
17       |    $\mathbf{x}' \in R'^{\mathbf{X}}$ , and  $\mathbf{a}' \notin S_{\mathbf{x}',R'}$  is feasible then
18         |   | return YES;
19         |   end
20   | end
21   | return NO;
22 end

```

---

**Characterisation** Our first main result – Theorem 13 – is a minion-theoretic characterisation of the power of the CLAP algorithm. In particular, we will introduce in Definition 12 a minion  $\mathcal{C}$  such that, for any PCSP template  $(\mathbf{A}, \mathbf{B})$ , the CLAP algorithm solves  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  if and only if there is a minion homomorphism from  $\mathcal{C}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$ . The two directions will be proved in Theorems 30 and 31, respectively, in Section 4. Combining Theorem 30 with our second main result – Theorem 15, proved in Section 5 – will then yield a sufficient condition

for CLAP to solve a given PCSP template, in terms of a weak notion of symmetry for the polymorphisms of the template.

The  $L$ -ary objects of the minion  $\mathcal{C}$  are pairs  $(M, \boldsymbol{\mu})$ , where  $M$  is a matrix with  $L$  rows and infinitely many columns encoding the BLP computations of CLAP and  $\boldsymbol{\mu}$  is an  $L$ -ary vector of integers encoding the AIP computation of CLAP. The matrices  $M$  in  $\mathcal{C}$  have a special structure, which we call ‘skeletal’.

**Definition 11.** Let  $M$  be a  $p \times \aleph_0$  matrix with  $p \in \mathbb{N}$ . We say that  $M$  is *skeletal* if, for each  $j \in [p]$ , either  $\mathbf{e}_j^T M = \mathbf{0}_{\aleph_0}^T$  or  $M\mathbf{e}_i = \mathbf{e}_j$  for some  $i \in \mathbb{N}$ .

In other words, either the  $j$ -th row of  $M$  is the zero vector or some column of  $M$  is the  $j$ -th standard unit vector. Equivalently,  $M$  is skeletal if there exist permutation matrices  $P \in \mathbb{R}^{p,p}$  and  $Q \in \mathbb{R}^{\aleph_0, \aleph_0}$  such that  $PMQ = \begin{bmatrix} I_k & \tilde{M} \\ O & O \end{bmatrix}$  for some  $k \leq p$  and some  $\tilde{M} \in \mathbb{R}^{k, \aleph_0}$ . The name indicates that the ‘body’ of a skeletal matrix (the nonzero rows) is completely supported by a ‘skeleton’ (the identity block).

We are now ready to define the minion  $\mathcal{C}$ . The  $L$ -ary objects of  $\mathcal{C}$  are pairs  $(M, \boldsymbol{\mu})$ , where  $M$  is a skeletal matrix of size  $L \times \aleph_0$  and  $\boldsymbol{\mu}$  is an affine vector (i.e., an integer vector whose entries sum up to one) of size  $L$ . We require that every column of  $M$  should be stochastic and  $M$  should have only finitely many different columns; the latter is formalised in  $(c_5)$  in Definition 12, which says that starting from some point all the columns are equal. We also require a particular relationship between  $M$  and  $\boldsymbol{\mu}$  formalised in  $(c_4)$ .

**Definition 12.** For  $L \in \mathbb{N}$ , let  $\mathcal{C}^{(L)}$  be the set of pairs  $(M, \boldsymbol{\mu})$  such that  $M \in \mathbb{Q}^{L, \aleph_0}$ ,  $\boldsymbol{\mu} \in \mathbb{Z}^L$ , and the following requirements are met:

- |  |   |
|--|---|
| $(c_1)$ $M$ is entrywise nonnegative;                  | $(c_4)$ $\text{supp}(\boldsymbol{\mu}) \subseteq \text{supp}(M\mathbf{e}_1)$ ;                        |
| $(c_2)$ $\mathbf{1}_L^T M = \mathbf{1}_{\aleph_0}^T$ ; | $(c_5)$ $\exists t \in \mathbb{N}$ such that $M\mathbf{e}_i = M\mathbf{e}_t \quad \forall i \geq t$ ; |
| $(c_3)$ $\mathbf{1}_L^T \boldsymbol{\mu} = 1$ ;        | $(c_6)$ $M$ is skeletal.  |

We define  $\mathcal{C}$  as the disjoint union of  $L$ -ary parts,  $\mathcal{C} := \bigcup_{L \geq 1} \mathcal{C}^{(L)}$ .

We defined  $\mathcal{C}$  as a set. For  $\mathcal{C}$  to be a minion, we need to define the minor operation on  $\mathcal{C}$  and verify that it preserves the structure of  $\mathcal{C}$ . This is easy and done in Section 4.1.

Our first result is the following characterisation of the power of CLAP.

**Theorem 13.** Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template. Then, CLAP solves  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  if and only if there is a minion homomorphism from  $\mathcal{C}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$ .

**H-symmetry** Our second main result is a sufficient condition on a PCSP template  $(\mathbf{A}, \mathbf{B})$  to guarantee that CLAP solves  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ . The condition is through symmetries satisfied by polymorphisms of the template. In particular, in Theorem 15 we will show that if  $\text{Pol}(\mathbf{A}, \mathbf{B})$  contains infinitely many operations that are ‘ $H$ -symmetric’ for a suitable matrix  $H$ , then there is a minion homomorphism from  $\mathcal{C}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$ , and thus CLAP solves  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  by Theorem 30.

In order to define the notion of  $H$ -symmetry, we need a few auxiliary definitions. A vector  $\mathbf{w} = (w_i) \in \mathbb{R}^p$  is *tieless* if, for any two indices  $i \neq i' \in [p]$ ,  $w_i \neq 0 \Rightarrow w_i \neq w_{i'}$ . A *tie matrix* is a matrix having integer nonnegative entries, each of whose columns is a tieless vector. Given an  $m \times p$  tie matrix  $H$ , we say that a vector  $\mathbf{v} \in \mathbb{R}^p$  is  *$H$ -tieless* if  $H\mathbf{v}$  is tieless.

Let  $A$  be a finite set, let  $L \in \mathbb{N}$ , and take  $\mathbf{a} = (a_1, \dots, a_L) \in A^L$ . We define the (multiplicity) vector  $\mathbf{a}^\#$  as the integer vector of size  $|A|$  whose  $a$ -th entry is  $|\{i \in [L] : a_i = a\}|$  for each  $a \in A$ .

**Definition 14.** Let  $A, B$  be finite sets, and consider a function  $f : A^L \rightarrow B$  for some  $L \in \mathbb{N}$ . Given an  $m \times |A|$  tie matrix  $H$ , we say that  $f$  is  $H$ -symmetric if

$$f_{/\pi}(\mathbf{a}) = f(\mathbf{a}) \quad \forall \pi : [L] \rightarrow [L] \text{ permutation, } \forall \mathbf{a} \in A^L \text{ such that } \mathbf{a}^\# \text{ is } H\text{-tieless.}$$

Our second result is the following sufficient condition for tractability of PCSPs.

**Theorem 15.** *Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template and suppose  $\text{Pol}(\mathbf{A}, \mathbf{B})$  contains  $H$ -symmetric operations of arbitrarily large arity for some  $m \times |A|$  tie matrix  $H$ ,  $m \in \mathbb{N}$ . Then there exists a minion homomorphism from  $\mathcal{C}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$ .*

Recall from Definition 11 the notion of a skeletal matrix. As it will be clear from the rest of the paper, the ‘skeleton’ represents the link between CLAP and the above-defined notion of  $H$ -symmetry. Indeed, on the one hand the presence of the identity block in a skeletal matrix captures the fact that each BLP solution computed by CLAP gives probability 1 to some constraint-assignment pair and probability 0 to all other constraint-assignment pairs for the same constraint (cf. line 6 of Algorithm 1). On the other hand, Lemma 35 (stated and proved in Section 5) shows that finitely many skeletal matrices can always be simultaneously reduced to  $H$ -tieless probability distributions – which are exactly the distributions on which  $H$ -symmetric functions are symmetric (cf. Definition 14).

We now mention some consequences of Theorem 15. First, observe that a vector of size 1 is always tieless. Hence, if we take any  $1 \times |A|$  integer nonnegative matrix as  $H$ , we have that  $H$  is a tie matrix and  $\mathbf{a}^\#$  is  $H$ -tieless for each tuple  $\mathbf{a}$  in the domain of  $f$ ; therefore, for such an  $H$ ,  $f$  being  $H$ -symmetric reduces to  $f$  being symmetric. On the other hand, having Definition 14 in mind, adding rows to  $H$  increases the chance that  $H\mathbf{a}^\#$  has some ties, in which case  $f$  is released from the requirement of being symmetric on  $\mathbf{a}$ . In this sense,  $H$  encodes the ‘exceptions to symmetry’ that  $f$  is allowed to have: The more rows  $H$  has, the stronger Theorem 15 becomes. If, for instance,  $H$  is the identity matrix of order  $|A|$ , then an  $H$ -symmetric operation needs to be symmetric only on those tuples where each entry occurs with a different multiplicity. A very special example of such an  $I_{|A|}$ -symmetric operation is a function  $f$  that returns (the homomorphic image of) the *most-frequent* entry in the input tuple whenever it is unique, and, in any other case,  $f$  is, say, (the homomorphic image of) a projection. Other, more creative choices for  $H$  allow capturing operations having more complex exceptions to symmetry, as shown in Example 16.

Theorems 13 and 15 together establish that the CLAP algorithm solves any PCSP template admitting arbitrarily large polymorphisms having some exceptions to symmetry that can be encoded via a tie matrix.

The importance of the next example lies in the fact that it provably separates CLAP from finite tractability and BLP + AIP; i.e., there are PCSP templates solvable by CLAP that are not finitely tractable and not solvable by BLP + AIP.

**Example 16.** Recall the PCSP template  $(\mathbf{A}, \mathbf{B})$  from Example 10, where it was shown that  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  is not finitely tractable and not solved by the BLP + AIP algorithm from [22]. We will show that  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  is solved by CLAP.

Take  $L \in \mathbb{N}$  and consider the function  $f : A^L \rightarrow B$  defined as follows: For  $\mathbf{a} = (a_1, \dots, a_L) \in A^L$ ,

- if  $\mathbf{a} \in \{0, 1\}^L$ , look at  $\mathbf{a}_1^\#$ , i.e., the multiplicity of  $1 \in A$  in the tuple  $\mathbf{a}$ ;
  - \* if  $\mathbf{a}_1^\# < \frac{L}{3}$ , set  $f(\mathbf{a}) = 0$ ;
  - \* if  $\mathbf{a}_1^\# > \frac{L}{3}$ , set  $f(\mathbf{a}) = 1$ ;
  - \* if  $\mathbf{a}_1^\# = \frac{L}{3}$ , set  $f(\mathbf{a}) = a_1$ ;
- if  $\mathbf{a} \in \{2, 3, 4, 5, 6\}^L$ ,
  - \* if there is a unique element  $a \in A$  having maximum multiplicity in  $\mathbf{a}$ , set  $f(\mathbf{a}) = a$ ;
  - \* if there is more than one element of  $A$  having maximum multiplicity in  $\mathbf{a}$ , set  $f(\mathbf{a}) = a_1$ ;
- otherwise, set  $f(\mathbf{a}) = 0$ .<sup>7</sup>

We claim that  $f \in \text{Pol}(\mathbf{A}, \mathbf{B})$ . To see that  $f$  preserves  $R_1$ , consider a tuple  $\boldsymbol{\rho} = (\mathbf{r}_1, \dots, \mathbf{r}_L)$  of elements of  $R_1^\mathbf{A}$ , where  $\mathbf{r}_i = (a_i, b_i, c_i)$  for  $i \in [L]$ . We shall let  $\mathbf{a} = (a_1, \dots, a_L)$ ,  $\mathbf{b} = (b_1, \dots, b_L)$ , and  $\mathbf{c} = (c_1, \dots, c_L)$ . Notice that

$$\mathbf{a}_1^\# + \mathbf{b}_1^\# + \mathbf{c}_1^\# = L. \quad (3)$$

If  $f(\mathbf{a}) = f(\mathbf{b}) = f(\mathbf{c}) = 0$ , then  $\mathbf{a}_1^\# \leq \frac{L}{3}$ ,  $\mathbf{b}_1^\# \leq \frac{L}{3}$ , and  $\mathbf{c}_1^\# \leq \frac{L}{3}$ ; by (3), this implies that  $\mathbf{a}_1^\# = \mathbf{b}_1^\# = \mathbf{c}_1^\# = \frac{L}{3}$ . Hence,  $(0, 0, 0) = (f(\mathbf{a}), f(\mathbf{b}), f(\mathbf{c})) = (a_1, b_1, c_1) = \mathbf{r}_1 \in R_1^\mathbf{A}$ , a contradiction. Similarly,  $f(\mathbf{a}) = f(\mathbf{b}) = f(\mathbf{c}) = 1$  would yield  $\mathbf{a}_1^\# \geq \frac{L}{3}$ ,  $\mathbf{b}_1^\# \geq \frac{L}{3}$ , and  $\mathbf{c}_1^\# \geq \frac{L}{3}$ ; again by (3), this implies that  $\mathbf{a}_1^\# = \mathbf{b}_1^\# = \mathbf{c}_1^\# = \frac{L}{3}$ , hence  $(1, 1, 1) = (f(\mathbf{a}), f(\mathbf{b}), f(\mathbf{c})) = (a_1, b_1, c_1) = \mathbf{r}_1 \in R_1^\mathbf{A}$ , also a contradiction. We conclude that  $f(\boldsymbol{\rho}) = (f(\mathbf{a}), f(\mathbf{b}), f(\mathbf{c})) \in R_1^\mathbf{B}$ , thus showing that  $f$  preserves  $R_1$ .

As for  $R_2$ , let  $\boldsymbol{\rho} = (\mathbf{r}_1, \dots, \mathbf{r}_L)$  be a tuple of elements of  $R_2^\mathbf{A}$ , where  $\mathbf{r}_i = (a_i, b_i)$  for  $i \in [L]$ , and let  $\mathbf{a} = (a_1, \dots, a_L)$  and  $\mathbf{b} = (b_1, \dots, b_L)$ . The directed graph having vertex set  $\{2, 3, 4, 5, 6\}$  and edge set  $R_2^\mathbf{A} = R_2^\mathbf{B}$  consists of the disjoint union of a directed 2-cycle and a directed 3-cycle and, hence, all of its vertices have in-degree and out-degree one. As a consequence, the multiplicity of a directed edge  $(a, b)$  in the tuple  $\boldsymbol{\rho}$  equals both the multiplicity of  $a$  in  $\mathbf{a}$  and the multiplicity of  $b$  in  $\mathbf{b}$ . Therefore, if the tuple  $\boldsymbol{\rho}$  has a unique element  $\mathbf{r} = (a, b)$  with maximum multiplicity, then  $f(\boldsymbol{\rho}) = (f(\mathbf{a}), f(\mathbf{b})) = (a, b) = \mathbf{r} \in R_2^\mathbf{B}$ . Otherwise,  $f(\boldsymbol{\rho}) = (a_1, b_1) = \mathbf{r}_1 \in R_2^\mathbf{B}$ . This shows that  $f$  preserves  $R_2$ , too, and is thus a polymorphism of  $(\mathbf{A}, \mathbf{B})$ .

Consider the matrix  $H = \text{diag}(1, 2, 1, 1, 1, 1)$ , and observe that  $H$  is a tie matrix. We claim that  $f$  is  $H$ -symmetric. Let  $\pi : [L] \rightarrow [L]$  be a permutation, and take a tuple  $\mathbf{a} = (a_1, \dots, a_L) \in A^L$  such that  $\mathbf{a}^\#$  is  $H$ -tieless; i.e., the vector  $H\mathbf{a}^\# = (\mathbf{a}_0^\#, 2\mathbf{a}_1^\#, \mathbf{a}_2^\#, \mathbf{a}_3^\#, \mathbf{a}_4^\#, \mathbf{a}_5^\#, \mathbf{a}_6^\#)$  is tieless. Write  $\tilde{\mathbf{a}} = (a_{\pi(1)}, \dots, a_{\pi(L)})$ , and observe that  $\tilde{\mathbf{a}}^\# = \mathbf{a}^\#$ .

- If  $\mathbf{a} \in \{0, 1\}^L$ , we get  $\mathbf{a}_0^\# \neq 2\mathbf{a}_1^\#$ ; since  $\mathbf{a}_0^\# + \mathbf{a}_1^\# = L$ , this gives  $2\mathbf{a}_1^\# \neq L - \mathbf{a}_1^\#$  so that  $\mathbf{a}_1^\# \neq \frac{L}{3}$ . As a consequence,  $f(\mathbf{a}) = f(\tilde{\mathbf{a}})$ .
- If  $\mathbf{a} \in \{2, 3, 4, 5, 6\}^L$ , the condition above implies that the tuple  $(\mathbf{a}_2^\#, \mathbf{a}_3^\#, \mathbf{a}_4^\#, \mathbf{a}_5^\#, \mathbf{a}_6^\#)$  has a unique maximum element and, hence, there is a unique element  $a$  of  $A$  having maximum multiplicity in  $\mathbf{a}$  (and in  $\tilde{\mathbf{a}}$ ). Therefore,  $f(\mathbf{a}) = a = f(\tilde{\mathbf{a}})$ .

<sup>7</sup>Assigning any value in  $\{0, \dots, 6\}$  to  $f(\mathbf{a})$  would work here.

- If  $\mathbf{a} \notin \{0, 1\}^L \cup \{2, 3, 4, 5, 6\}^L$ , then  $f(\mathbf{a}) = 0 = f(\tilde{\mathbf{a}})$ .

We conclude that, in each case,  $f(\mathbf{a}) = f(\tilde{\mathbf{a}}) = f_{/\pi}(\mathbf{a})$ , which means that  $f$  is  $H$ -symmetric. By Theorems 13 and 15, CLAP solves PCSP( $\mathbf{A}, \mathbf{B}$ ).

**Remark 17.** Consider the minion  $\mathcal{M}_{\text{BLP} + \text{AIP}}$  from [22] (cf. Appendix A.3). A direct consequence of Example 16, Theorem 13, and [22, Lemma 5.4] is that there is no minion homomorphism from  $\mathcal{M}_{\text{BLP} + \text{AIP}}$  to  $\mathcal{C}$ . On the other hand, the function

$$\begin{aligned} \vartheta : \mathcal{C} &\rightarrow \mathcal{M}_{\text{BLP} + \text{AIP}} \\ (M, \boldsymbol{\mu}) &\mapsto (M\mathbf{e}_1, \boldsymbol{\mu}) \end{aligned}$$

is readily seen to be a minion homomorphism. It follows that CLAP solves any PCSP template solved by BLP + AIP (as is also clear from the description of the two algorithms).

**Remark 18.** Similar to [22], the assumption in Theorem 15 can be weakened as follows: Instead of requiring  $H$ -symmetric polymorphisms of arbitrarily large arity, it turns out to be enough requiring  $H$ -block-symmetric polymorphisms of arbitrarily large width, where the definition of an  $H$ -block-symmetric operation mirrors that of a block-symmetric operation in [22]. The proof of this possibly stronger result is very similar to that of Theorem 15. For completeness, we include it in Appendix C. We point out that we do not know whether the condition in Theorem 15 (or the possibly weaker condition based on  $H$ -block-symmetric polymorphisms) is necessary for tractability via CLAP, but we suspect it is not.

**Remark 19.** A possibly stronger version of the CLAP algorithm consists in running BLP + AIP (instead of just BLP) at each iteration in the **for** loop in lines 5–9 of Algorithm 1, and then removing the additional **for** loop in lines 14–18. This algorithm can be called C(BLP + AIP). An analysis entirely analogous to the one presented in this paper shows that the power of C(BLP + AIP) is captured by the minion  $\tilde{\mathcal{C}}$  defined like  $\mathcal{C}$  with the following difference: The  $L$ -ary elements of  $\tilde{\mathcal{C}}$  are pairs  $(M, N)$ , where  $M$  is as in  $\mathcal{C}$  while  $N$  is an integer matrix of the same size as  $M$  taking the role of  $\boldsymbol{\mu}$  (in particular,  $N$  satisfies the ‘refinement condition’  $\text{supp}(N\mathbf{e}_i) \subseteq \text{supp}(M\mathbf{e}_i) \forall i \in \mathbb{N}$ , analogous to  $(c_4)$  in Definition 12). A possible direction for future research is to investigate whether the richer structure of  $\tilde{\mathcal{C}}$  can be exploited to obtain a stronger version of Theorem 15.

**Remark 20.** For CSPs, the characterisation of bounded width [9, 25] and its collapse [6] was preceded by a characterisation of width-1 CSPs [37, 41] and the collapse of width 2 to width 1 [34]. Thus the difference between width-1 CSPs and bounded-width CSPs is well understood. BLP and SBLP are the (convex relaxation) analogues of width 1 and SAC, respectively, and SAC solves all bounded-width CSPs [56]. Therefore, a natural question is whether a similar analysis can cast light on the difference in power between BLP on one side, and SBLP (and thus perhaps also of CBLP and CLAP) on the other side. We remark on two obstacles: Firstly, BLP is strictly more powerful than width 1 for CSPs [60]. Secondly, a good characterisation of the power of SBLP (and stronger algorithms studied in the present paper) would imply that these algorithms solve, in the special case of CSPs, all bounded-width CSPs – a non-trivial result implied by [56].



## 4 The power of the CLAP algorithm

The goal of this section is to prove Theorem 13. In Section 4.1, we will verify that  $\mathcal{C}$ , which appears in the statement of Theorem 13, is indeed a minion. In Sections 4.2 and 4.3, we will establish a compactness argument and present a condition that captures CLAP, respectively; both will be needed in the proof of Theorem 13. The two directions of Theorem 13 will be then proved in Section 4.4.

Minions are not only useful for capturing the complexity of PCSPs but also for characterising the power of algorithms. This will be done by using the concept of the free structure generated, for a given minion, by a relational structure [22] (cf. [8, Definition 4.1] for the definition in the special case of minions of functions).

**Definition 21.** Let  $\mathcal{M}$  be a minion and let  $\mathbf{A}$  be a (finite) relational structure with signature  $\sigma$ . The *free structure*  $\mathbb{F}_{\mathcal{M}}(\mathbf{A})$  is a relational structure with domain  $\mathcal{M}^{(|A|)}$  (potentially infinite) and signature  $\sigma$ . Given a relation  $R \in \sigma$  of arity  $k$ , a tuple  $(M_1, \dots, M_k)$  of elements of  $\mathcal{M}^{(|A|)}$  belongs to  $R^{\mathbb{F}_{\mathcal{M}}(\mathbf{A})}$  if and only if there is some  $Q \in \mathcal{M}^{(|R^{\mathbf{A}}|)}$  such that  $M_i = Q_{/\pi_i}$  for each  $i \in [k]$ , where  $\pi_i : R^{\mathbf{A}} \rightarrow A$  maps  $\mathbf{a} \in R^{\mathbf{A}}$  to its  $i$ -th coordinate  $a_i$ .

The next result will be useful to establish the connection between our algorithm CLAP, presented in Section 3, and the minion  $\mathcal{C}$ .

**Lemma 22.** *Let  $\mathcal{M}$  be a minion and let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template. Then there is a minion homomorphism from  $\mathcal{M}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$  if and only if  $\mathbb{F}_{\mathcal{M}}(\mathbf{A}) \rightarrow \mathbf{B}$ .*

The proof of Lemma 22 is based on that of [8, Lemma 4.4], which proves one-to-one correspondence but only for minions of functions. For completeness, we prove Lemma 22 in Appendix B.

### 4.1 $\mathcal{C}$ is a minion

The minor operation on  $\mathcal{C}$  is naturally defined via a matrix multiplication with a matrix that encodes the minor map. For a function  $\pi : [L] \rightarrow [L']$ , let  $P_\pi$  be the  $L' \times L$  matrix whose  $(i, j)$ -th entry is 1 if  $\pi(j) = i$ , and 0 otherwise. Note that  $P_\pi^T \mathbf{1}_{L'} = \mathbf{1}_L$  and, for each  $i \in [L']$ ,  $P_\pi^T \mathbf{e}_i = \sum_{j \in \pi^{-1}(i)} \mathbf{e}_j$ .

**Definition 23.** For  $(M, \boldsymbol{\mu}) \in \mathcal{C}^{(L)}$ , we define  $M_{/\pi} = P_\pi M$  and  $\boldsymbol{\mu}_{/\pi} = P_\pi \boldsymbol{\mu}$ , and we let the minor of  $(M, \boldsymbol{\mu})$  with respect to  $\pi$  be  $(M, \boldsymbol{\mu})_{/\pi} := (M_{/\pi}, \boldsymbol{\mu}_{/\pi})$ .

We remark that this definition is consistent with the minions  $\mathcal{Q}_{\text{conv}}$  and  $\mathcal{Z}_{\text{aff}}$  studied in [8], and the minion  $\mathcal{M}_{\text{BLP}+\text{AIP}}$  studied in [22], cf. Appendices A.1, A.2, and A.3.

**Proposition 24.**  *$\mathcal{C}$  is a minion.*

*Proof.* Write  $M = [m_{ij}]$  and  $\boldsymbol{\mu} = (\mu_i)$ . Observe that  $M_{/\pi} \in \mathbb{Q}^{L', \aleph_0}$  and  $\boldsymbol{\mu}_{/\pi} \in \mathbb{Z}^{L'}$ . The requirements  $(c_1)$ ,  $(c_2)$ ,  $(c_3)$ , and  $(c_5)$  are trivially satisfied by  $(M, \boldsymbol{\mu})_{/\pi}$ . As for  $(c_4)$ , suppose that  $\mathbf{e}_i^T P_\pi M \mathbf{e}_1 = 0$  but  $\mathbf{e}_i^T P_\pi \boldsymbol{\mu} \neq 0$ . It follows that  $\mu_j \neq 0$  for some  $j \in \pi^{-1}(i)$ . Hence,  $m_{j1} > 0$  and, then,

$$\mathbf{e}_i^T P_\pi M \mathbf{e}_1 = \sum_{j' \in \pi^{-1}(i)} \mathbf{e}_{j'}^T M \mathbf{e}_1 \geq \mathbf{e}_j^T M \mathbf{e}_1 > 0,$$

which is a contradiction. We now show that  $M_{/\pi}$  is skeletal. Choose  $j \in [L']$ , and suppose that  $\mathbf{e}_j^T M_{/\pi} \neq \mathbf{0}_{\mathbb{N}_0}^T$ . We obtain

$$\mathbf{0}_{\mathbb{N}_0} \neq M^T P_\pi^T \mathbf{e}_j = \sum_{\ell \in \pi^{-1}(j)} M^T \mathbf{e}_\ell$$

and, in particular,  $\exists \ell \in \pi^{-1}(j)$  such that  $\mathbf{e}_\ell^T M \neq \mathbf{0}_{\mathbb{N}_0}^T$ . Since  $M$  is skeletal, this implies that  $M\mathbf{e}_i = \mathbf{e}_\ell$  for some  $i \in \mathbb{N}$ . This yields

$$M_{/\pi} \mathbf{e}_i = P_\pi M \mathbf{e}_i = P_\pi \mathbf{e}_\ell = \mathbf{e}_{\pi(\ell)} = \mathbf{e}_j$$

as required. Hence,  $(c_6)$  is satisfied, too, and  $(M, \boldsymbol{\mu})_{/\pi} \in \mathcal{C}^{(L')}$ .

Finally, considering  $\tilde{\pi} : [L'] \rightarrow [L'']$  and the identity map  $\text{id} : [L] \rightarrow [L]$ , one readily checks that  $P_{\tilde{\pi} \circ \pi} = P_{\tilde{\pi}} P_\pi$  and  $P_{\text{id}} = I_L$ . Hence, the minor operations defined above satisfy the requirements of Definition 6.  $\square$

## 4.2 A compactness argument for $\mathcal{C}$

The set  $\mathcal{C}^{(L)}$  of the  $L$ -ary objects in  $\mathcal{C}$  is infinite unless  $L = 1$ . As a consequence, given a relational structure  $\mathbf{A}$  whose domain has size at least 2, the free structure  $\mathbb{F}_{\mathcal{C}}(\mathbf{A})$  has an infinite domain. We now describe a standard compactness argument analogous to [8, Remark 7.13] that will circumvent this inconvenience.

For  $D, L \in \mathbb{N}$ , consider the set

$$\mathcal{C}_D^{(L)} = \{(M, \boldsymbol{\mu}) \in \mathcal{C}^{(L)} : DM \text{ is entrywise integer, } M\mathbf{e}_i = M\mathbf{e}_D \ \forall i \geq D, \text{ and } \mathbf{1}_L^T |\boldsymbol{\mu}| \leq D\},$$

where  $|\boldsymbol{\mu}|$  denotes the vector whose entries are the absolute values of the entries of  $\boldsymbol{\mu}$ . Since  $\mathcal{C}_D^{(L)}$  is unambiguously determined by  $L \times (D + 1)$  integer numbers belonging to the set  $\{-D, \dots, D\}$ , it is finite. Observe that the set  $\mathcal{C}_D = \bigcup_{L \in \mathbb{N}} \mathcal{C}_D^{(L)}$  is closed under taking minors. Indeed, given  $(M, \boldsymbol{\mu}) \in \mathcal{C}_D^{(L)}$  and  $\pi : [L] \rightarrow [L']$ ,  $DP_\pi M = P_\pi DM$  is entrywise integer,  $P_\pi M \mathbf{e}_i = P_\pi M \mathbf{e}_D \ \forall i \geq D$ , and  $\mathbf{1}_{L'}^T |P_\pi \boldsymbol{\mu}| \leq \mathbf{1}_{L'}^T P_\pi |\boldsymbol{\mu}| = \mathbf{1}_L^T |\boldsymbol{\mu}| \leq D$ . Hence,  $\mathcal{C}_D$  is a sub-minion of  $\mathcal{C}$ . Observe also that  $\mathcal{C} = \bigcup_{D \in \mathbb{N}} \mathcal{C}_D$ . To see this, take  $(M, \boldsymbol{\mu}) \in \mathcal{C}^{(L)}$  and suppose that  $M\mathbf{e}_i = M\mathbf{e}_t \ \forall i \geq t$ . Let  $\tilde{D}$  be a common denominator of the finite set of rational numbers  $\{m_{ij} : i \in [L], j \in [t]\}$ , so that  $\tilde{D}M$  is entrywise integer. Let also  $\hat{D} = \mathbf{1}_L^T |\boldsymbol{\mu}|$ . Then,  $(M, \boldsymbol{\mu}) \in \mathcal{C}_{t\tilde{D}\hat{D}}$ .

**Proposition 25.** *Let  $\mathcal{M}$  be a minion such that  $\mathcal{M}^{(L)}$  is finite for each  $L \in \mathbb{N}$ , and suppose that there exist minion homomorphisms  $\xi_D : \mathcal{C}_D \rightarrow \mathcal{M}$  for each  $D \in \mathbb{N}$ . Then there exists a minion homomorphism  $\zeta : \mathcal{C} \rightarrow \mathcal{M}$ .*

*Proof.* For  $D \in \mathbb{N}$ , let  $\mathcal{C}_{\{D\}} = \bigcup_{L \leq D} \mathcal{C}_D^{(L)}$ . Observe that  $\mathcal{C}_{\{D\}}$  is a finite set and  $\mathcal{C}_{\{D\}} \subseteq \mathcal{C}_{\{D+1\}}$ . Moreover,  $\bigcup_{D \in \mathbb{N}} \mathcal{C}_{\{D\}} = \bigcup_{D \in \mathbb{N}} \mathcal{C}_D = \mathcal{C}$ . Indeed, given  $D' \in \mathbb{N}$ , we have that  $\mathcal{C}_{\{D'\}} \subseteq \mathcal{C}_{D'} \subseteq \bigcup_{D \in \mathbb{N}} \mathcal{C}_D$ , and, given  $L \in \mathbb{N}$ ,  $\mathcal{C}_{D'}^{(L)} \subseteq \mathcal{C}_{(D'L)}^{(L)} \subseteq \mathcal{C}_{\{D'L\}} \subseteq \bigcup_{D \in \mathbb{N}} \mathcal{C}_{\{D\}}$ . Consider an infinite rooted tree whose vertices are all the restrictions of the homomorphisms  $\xi_D$  to some  $\mathcal{C}_{\{D'\}}$ , whose root is the empty mapping, and the parent of a vertex corresponding to a function  $\mathcal{C}_{\{D'+1\}} \rightarrow \mathcal{M}$  is the vertex corresponding to the restriction of the function to  $\mathcal{C}_{\{D'\}}$ . This is an infinite connected tree. Moreover, since  $\mathcal{M}^{(L)}$  is finite for each  $L \in \mathbb{N}$

and since minion homomorphisms preserve the arities, there exist only finitely many distinct restrictions of minion homomorphisms to  $\mathcal{C}_{\{D\}}$ ; hence, the tree is locally finite. By König's Lemma, it contains an infinite path, which corresponds to an infinite chain of maps  $\zeta_i : \mathcal{C}_{\{i\}} \rightarrow \mathcal{M}$  such that  $\zeta_{i+1}$  extends  $\zeta_i \forall i \in \mathbb{N}$ . Their union  $\zeta : \mathcal{C} \rightarrow \mathcal{M}$  is then a minion homomorphism.  $\square$

### 4.3 The CLAP condition

Given a finite set  $C$ , consider the set  $\mathbb{S}(C)$  of the rational stochastic vectors of size  $|C|$ . Let  $U \subseteq C^k$ . For  $i \in [k]$ , consider the  $|C| \times |U|$  matrix  $E^{(U,i)}$  such that, for  $c \in C$  and  $\mathbf{c} = (c_1, \dots, c_k) \in U$ , the  $(c, \mathbf{c})$ -th entry of  $E^{(U,i)}$  is 1 if  $c_i = c$ , and 0 otherwise. Given  $\xi \in \mathbb{S}(U)$  and  $i \in [k]$ , we define the  $i$ -th *marginal* of  $\xi$  as

$$\xi^{(i)} = E^{(U,i)} \xi.$$

Observe that

$$\xi^{(i)T} \mathbf{1}_{|C|} = \xi^T E^{(U,i)T} \mathbf{1}_{|C|} = \xi^T \mathbf{1}_{|U|} = 1,$$

so that  $\xi^{(i)} \in \mathbb{S}(C)$ . We also define the set  $\mathbb{Z}(C)$  of the integer vectors of size  $|C|$  whose entries sum up to 1. Given  $U \subseteq C^k$ ,  $\zeta \in \mathbb{Z}(U)$ , and  $i \in [k]$ , we define

$$\zeta^{(i)} = E^{(U,i)} \zeta.$$

As before, observe that

$$\zeta^{(i)T} \mathbf{1}_{|C|} = \zeta^T E^{(U,i)T} \mathbf{1}_{|C|} = \zeta^T \mathbf{1}_{|U|} = 1,$$

so  $\zeta^{(i)} \in \mathbb{Z}(C)$ .

Let  $\mathbf{A}$  be a relational structure having domain  $A$  and signature  $\sigma$ . We define the relational structures  $\mathbb{S}(\mathbf{A})$  and  $\mathbb{Z}(\mathbf{A})$  as follows:

- $\mathbb{S}(\mathbf{A})$  has domain  $\mathbb{S}(A)$  and, for every symbol  $R \in \sigma$  of arity  $k$ ,  $R^{\mathbb{S}(\mathbf{A})} = \{(\xi^{(1)}, \dots, \xi^{(k)}) : \xi \in \mathbb{S}(R^{\mathbf{A}})\}$ ;
- $\mathbb{Z}(\mathbf{A})$  has domain  $\mathbb{Z}(A)$  and, for every symbol  $R \in \sigma$  of arity  $k$ ,  $R^{\mathbb{Z}(\mathbf{A})} = \{(\zeta^{(1)}, \dots, \zeta^{(k)}) : \zeta \in \mathbb{Z}(R^{\mathbf{A}})\}$ .

**Remark 26.**  $\mathbb{S}(\mathbf{A})$  and  $\mathbb{Z}(\mathbf{A})$  are denoted by  $\text{LP}(\mathbf{A})$  and  $\text{IP}(\mathbf{A})$  in [8], respectively. As noted in [8, Remarks 7.11 and 7.21],  $\mathbb{S}(\mathbf{A})$  coincides with the free structure of the minion  $\mathcal{Q}_{\text{conv}}$  generated by  $\mathbf{A}$  and, similarly,  $\mathbb{Z}(\mathbf{A})$  is the free structure of the minion  $\mathcal{Z}_{\text{aff}}$  generated by  $\mathbf{A}$ . (See Appendices A.1 and A.2 for the definitions of  $\mathcal{Q}_{\text{conv}}$  and  $\mathcal{Z}_{\text{aff}}$ , respectively.) In particular, given a relational structure  $\mathbf{X}$  with signature  $\sigma$ , BLP accepts  $\mathbf{X}$  as an instance of  $\text{CSP}(\mathbf{A})$  if and only if  $\mathbf{X} \rightarrow \mathbb{S}(\mathbf{A})$ ; similarly, AIP accepts  $\mathbf{X}$  as an instance of  $\text{CSP}(\mathbf{A})$  if and only if  $\mathbf{X} \rightarrow \mathbb{Z}(\mathbf{A})$ .

**Remark 27.** The assignment  $f : a \mapsto \mathbf{e}_a$  for each  $a \in A$  yields both a canonical homomorphism from  $\mathbf{A}$  to  $\mathbb{S}(\mathbf{A})$  and a canonical homomorphism from  $\mathbf{A}$  to  $\mathbb{Z}(\mathbf{A})$ . Indeed, for  $R \in \sigma$  of arity  $k$  and  $\mathbf{a} = (a_1, \dots, a_k) \in R^{\mathbf{A}}$ ,

$$f(\mathbf{a}) = (\mathbf{e}_{a_1}, \dots, \mathbf{e}_{a_k}) = (E^{(R^{\mathbf{A}},1)} \mathbf{e}_{\mathbf{a}}, \dots, E^{(R^{\mathbf{A}},k)} \mathbf{e}_{\mathbf{a}})$$

which belongs to both  $R^{\mathbb{S}(\mathbf{A})}$  and  $R^{\mathbb{Z}(\mathbf{A})}$  since  $\mathbf{e}_{\mathbf{a}} \in \mathbb{S}(R^{\mathbf{A}}) \cap \mathbb{Z}(R^{\mathbf{A}})$ .

In Proposition 29, we characterise the instances of a given PCSP template for which the CLAP algorithm returns YES in terms of the condition described in the following definition.

**Definition 28.** Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template, where  $\mathbf{A}$  and  $\mathbf{B}$  have signature  $\sigma$ . Given an instance  $\mathbf{X}$  of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ , we say that  $\mathbf{X}$  has the CLAP *condition* if the following holds:  $\forall R \in \sigma$  of arity  $k \exists s^R : R^{\mathbf{X}} \rightarrow \mathcal{P}(R^{\mathbf{A}}) \setminus \{\emptyset\}$  such that

(I)  $\forall \mathbf{x} = (x_1, \dots, x_k) \in R^{\mathbf{X}}, \forall \mathbf{a} = (a_1, \dots, a_k) \in s^R(\mathbf{x})$  there is a homomorphism  $h_{\mathbf{x}, \mathbf{a}} : \mathbf{X} \rightarrow \mathbb{S}(\mathbf{A})$  that satisfies:

1.  $h_{\mathbf{x}, \mathbf{a}}(x_i) = \mathbf{e}_{a_i} \forall i \in [k]$ ;
2.  $\forall \tilde{R} \in \sigma$  of arity  $\tilde{k}, \forall \tilde{\mathbf{x}} = (\tilde{x}_1, \dots, \tilde{x}_{\tilde{k}}) \in \tilde{R}^{\mathbf{X}} \exists \xi \in \mathbb{S}(\tilde{R}^{\mathbf{A}})$  such that
  - \*  $h_{\mathbf{x}, \mathbf{a}}(\tilde{x}_i) = E^{(\tilde{R}^{\mathbf{A}}, i)} \xi \quad \forall i \in [\tilde{k}]$ ;
  - \*  $\text{supp}(\xi) \subseteq s^{\tilde{R}}(\tilde{\mathbf{x}})$ .

(II)  $\exists \bar{R} \in \sigma, \bar{\mathbf{x}} \in \bar{R}^{\mathbf{X}}, \bar{\mathbf{a}} \in s^{\bar{R}}(\bar{\mathbf{x}})$  such that there is a homomorphism  $g : \mathbf{X} \rightarrow \mathbb{Z}(\mathbf{A})$  that satisfies:

- 1'.  $\forall \tilde{R} \in \sigma$  of arity  $\tilde{k}, \forall \tilde{\mathbf{x}} = (\tilde{x}_1, \dots, \tilde{x}_{\tilde{k}}) \in \tilde{R}^{\mathbf{X}} \exists \xi \in \mathbb{S}(\tilde{R}^{\mathbf{A}}), \exists \zeta \in \mathbb{Z}(\tilde{R}^{\mathbf{A}})$  such that
  - \*  $h_{\bar{\mathbf{x}}, \bar{\mathbf{a}}}(\tilde{x}_i) = E^{(\tilde{R}^{\mathbf{A}}, i)} \xi \quad \forall i \in [\tilde{k}]$ ;
  - \*  $g(\tilde{x}_i) = E^{(\tilde{R}^{\mathbf{A}}, i)} \zeta \quad \forall i \in [\tilde{k}]$ ;
  - \*  $\text{supp}(\zeta) \subseteq \text{supp}(\xi) \subseteq s^{\tilde{R}}(\tilde{\mathbf{x}})$ .

**Proposition 29.** *Given an instance  $\mathbf{X}$  of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ , CLAP accepts  $\mathbf{X}$  if and only if  $\mathbf{X}$  has the CLAP condition.*

*Proof.* Suppose that CLAP accepts  $\mathbf{X}$  and let  $\{S_{\mathbf{x}, R} : R \in \sigma, \mathbf{x} \in R^{\mathbf{X}}\}$  be the family of sets generated by the algorithm at termination. For each  $R \in \sigma$ , consider the map  $s^R : R^{\mathbf{X}} \rightarrow \mathcal{P}(R^{\mathbf{A}}) \setminus \{\emptyset\}$  defined by  $s^R(\mathbf{x}) = S_{\mathbf{x}, R}$ . For each  $\mathbf{x} \in R^{\mathbf{X}}, \mathbf{a} \in s^R(\mathbf{x})$ , consider the corresponding solution to  $\text{BLP}(\mathbf{X}, \mathbf{A})$  generated by the algorithm. Letting  $w_x$  be the probability distribution on  $A$  associated with  $x \in X$  in the linear program, we observe that the assignment  $x \mapsto w_x$  yields a homomorphism (call it  $h_{\mathbf{x}, \mathbf{a}}$ ) from  $\mathbf{X}$  to  $\mathbb{S}(\mathbf{A})$  that satisfies the requirement 1. Moreover, letting  $\xi$  be the probability distribution associated with a constraint  $\tilde{\mathbf{x}} \in \tilde{R}^{\mathbf{X}}$  for some  $\tilde{R} \in \sigma$ , observe that  $h_{\mathbf{x}, \mathbf{a}}$  also satisfies the requirement 2. Finally, let  $\bar{R} \in \sigma, \bar{\mathbf{x}} \in \bar{R}^{\mathbf{X}}, \bar{\mathbf{a}} \in S_{\bar{\mathbf{x}}, \bar{R}}$  be such that the condition in the **if** statement of line 15 of Algorithm 1 is met. Then 1' follows from the description of  $\text{BLP} + \text{AIP}$ .

The converse implication follows almost analogously, except for the following subtlety. The  $\text{BLP} + \text{AIP}$  algorithm requires that the BLP solution should be picked from the *relative interior* of the polytope of the feasible solutions (cf. Algorithm 2 in Appendix A.3). However, the homomorphism  $h_{\bar{\mathbf{x}}, \bar{\mathbf{a}}}$  whose existence witnesses part (II) of the CLAP condition may correspond to a BLP solution that is not in the relative interior of the polytope  $P$  of the feasible solutions of  $\text{BLP}(\mathbf{X}, \mathbf{A})$  satisfying  $\lambda_{\bar{\mathbf{x}}, \bar{R}}(\bar{\mathbf{a}}) = 1$  and  $\lambda_{\mathbf{x}', R'}(\mathbf{a}') = 0$  for every  $R' \in \sigma, \mathbf{x}' \in R'^{\mathbf{X}}$ , and  $\mathbf{a}' \notin S_{\mathbf{x}', R'}$ . If that is the case, the algorithm would not consider  $(h_{\bar{\mathbf{x}}, \bar{\mathbf{a}}}, g)$  as a solution for  $\text{BLP} + \text{AIP}$ . However, letting  $h'$  be a solution in the relative interior of  $P$ , the conditions (I) and (II) of CLAP are still satisfied if we let  $h'$  replace  $h_{\bar{\mathbf{x}}, \bar{\mathbf{a}}}$ ; and, in this case, the homomorphisms witnessing the CLAP condition do correspond to solutions found by the CLAP algorithm.<sup>8</sup> Hence, CLAP accepts  $\mathbf{X}$ .  $\square$

<sup>8</sup>Another way to phrase this is by saying that the existence of a pair  $(h, g)$  of homomorphisms such that

#### 4.4 Proof of Theorem 13

Our first goal is to prove the following.

**Theorem 30.** *If there is a minion homomorphism from  $\mathcal{C}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$  then CLAP solves  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ .*

*Proof.* Let  $\mathbf{X}$  be an instance of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ .

First we show that if  $\mathbf{X} \rightarrow \mathbf{A}$  then CLAP accepts  $\mathbf{X}$ , which is the easy direction. Consider a homomorphism  $f : \mathbf{X} \rightarrow \mathbf{A}$ . Given  $R \in \sigma$  of arity  $k$  and  $\mathbf{x} = (x_1, \dots, x_k) \in R^{\mathbf{X}}$ , let  $s^R(\mathbf{x}) = \{f(\mathbf{x})\}$ . For  $\mathbf{x} \in R^{\mathbf{X}}$  and  $\mathbf{a} = (a_1, \dots, a_k) = f(\mathbf{x}) \in s^R(\mathbf{x})$ , let  $h_{\mathbf{x}, \mathbf{a}} : \mathbf{X} \rightarrow \mathbb{S}(\mathbf{A})$  be the homomorphism obtained by composing  $f$  with the canonical homomorphism from  $\mathbf{A}$  to  $\mathbb{S}(\mathbf{A})$  of Remark 27 – i.e.,  $h_{\mathbf{x}, \mathbf{a}}(x) = \mathbf{e}_{f(x)} \forall x \in X$ . Observe that  $h_{\mathbf{x}, \mathbf{a}}(x_i) = \mathbf{e}_{f(x_i)} = \mathbf{e}_{a_i}$  for any  $i \in [k]$  and, given  $\tilde{R} \in \sigma$  of arity  $\tilde{k}$  and  $\tilde{\mathbf{x}} = (\tilde{x}_1, \dots, \tilde{x}_{\tilde{k}}) \in \tilde{R}^{\mathbf{X}}$ , setting  $\xi = \mathbf{e}_{f(\tilde{\mathbf{x}})}$  yields  $h_{\mathbf{x}, \mathbf{a}}(\tilde{x}_i) = \mathbf{e}_{f(\tilde{x}_i)} = E^{(\tilde{R}^{\mathbf{A}}, i)} \mathbf{e}_{f(\tilde{\mathbf{x}})} = E^{(\tilde{R}^{\mathbf{A}}, i)} \xi$  for any  $i \in [\tilde{k}]$ , and  $\text{supp}(\xi) = \text{supp}(\mathbf{e}_{f(\tilde{\mathbf{x}})}) = \{f(\tilde{\mathbf{x}})\} = s^{\tilde{R}}(\tilde{\mathbf{x}})$ . This shows that part (I) of Definition 28 is satisfied. As for part (II), choose any  $\bar{R} \in \sigma$  and  $\bar{\mathbf{x}} \in \bar{R}^{\mathbf{X}}$ , let  $\bar{\mathbf{a}} = f(\bar{\mathbf{x}})$ , and consider the homomorphism  $g : \mathbf{X} \rightarrow \mathbb{Z}(\mathbf{A})$  obtained by composing  $f$  with the canonical homomorphism from  $\mathbf{A}$  to  $\mathbb{Z}(\mathbf{A})$  of Remark 27 – i.e.,  $g(x) = \mathbf{e}_{f(x)} \forall x \in X$ . Given  $\tilde{R} \in \sigma$  of arity  $\tilde{k}$  and  $\tilde{\mathbf{x}} = (\tilde{x}_1, \dots, \tilde{x}_{\tilde{k}}) \in \tilde{R}^{\mathbf{X}}$ , setting  $\xi = \zeta = \mathbf{e}_{f(\tilde{\mathbf{x}})}$  yields  $g(\tilde{x}_i) = h_{\bar{\mathbf{x}}, \bar{\mathbf{a}}}(\tilde{x}_i) = \mathbf{e}_{f(\tilde{x}_i)} = E^{(\tilde{R}^{\mathbf{A}}, i)} \mathbf{e}_{f(\tilde{\mathbf{x}})} = E^{(\tilde{R}^{\mathbf{A}}, i)} \xi = E^{(\tilde{R}^{\mathbf{A}}, i)} \zeta$  for any  $i \in [\tilde{k}]$ , and  $\text{supp}(\zeta) = \text{supp}(\xi) = \{f(\tilde{\mathbf{x}})\} = s^{\tilde{R}}(\tilde{\mathbf{x}})$ . It follows that  $\mathbf{X}$  has the CLAP condition. By Proposition 29, CLAP accepts  $\mathbf{X}$ .

Second we show that if  $\mathbf{X}$  is accepted by CLAP then  $\mathbf{X} \rightarrow \mathbf{B}$ . So, suppose that  $\mathbf{X}$  is accepted by CLAP. By Proposition 29,  $\mathbf{X}$  has the CLAP condition. Using the terminology of Definition 28, consider the set  $\{h_1, \dots, h_t\} = \{h_{\mathbf{x}, \mathbf{a}} : R \in \sigma, \mathbf{x} \in R^{\mathbf{X}}, \mathbf{a} \in s^R(\mathbf{x})\}$ , where each  $h_{\mathbf{x}, \mathbf{a}}$  is a homomorphism from  $\mathbf{X}$  to  $\mathbb{S}(\mathbf{A})$  described in part (I) of the definition. We also consider the homomorphism  $g : \mathbf{X} \rightarrow \mathbb{Z}(\mathbf{A})$  of part (II) of the definition, corresponding to  $\bar{R} \in \sigma, \bar{\mathbf{x}} \in \bar{R}^{\mathbf{X}}, \bar{\mathbf{a}} \in s^{\bar{R}}(\bar{\mathbf{x}})$ . Without loss of generality, we set  $h_1 = h_{\bar{\mathbf{x}}, \bar{\mathbf{a}}}$ .

Let  $n = |A|$ . Given  $x \in X$ , consider the matrix  $M_x \in \mathbb{Q}^{n, \aleph_0}$  and the vector  $\mu_x \in \mathbb{Z}^n$  defined by

$$\begin{aligned} M_x \mathbf{e}_i &= h_i(x) \quad \forall i \in [t], \\ M_x \mathbf{e}_i &= h_t(x) \quad \forall i \in \mathbb{N} \setminus [t], \\ \mu_x &= g(x). \end{aligned}$$

We claim that  $(M_x, \mu_x) \in \mathcal{C}^{(n)}$ . The requirements  $(c_1), (c_2), (c_3)$ , and  $(c_5)$  in Definition 12 are easily seen to be satisfied. To check that  $M_x$  is skeletal, take  $a \in A$  and suppose that  $\mathbf{e}_a^T M_x \neq \mathbf{0}_{\aleph_0}^T$ . This means that  $\mathbf{e}_a^T M_x \mathbf{e}_d \neq 0$  for some  $d \in [t]$ . Hence,  $a \in \text{supp}(M_x \mathbf{e}_d) = \text{supp}(h_d(x))$ . Recall that we are assuming (with no loss of generality) that the signature  $\sigma$  of  $\mathbf{X}$ ,  $\mathbf{A}$ , and  $\mathbf{B}$  contains a unary symbol  $R_u$  such that  $R_u^{\mathbf{X}} = X$ ,  $R_u^{\mathbf{A}} = A$ , and  $R_u^{\mathbf{B}} = B$ . Notice that  $E^{(R_u^{\mathbf{A}}, 1)} = I_n$ . From part (I) of Definition 28, we deduce that  $\text{supp}(h_d(x)) \subseteq s^{R_u}(x)$  and, hence,  $a \in s^{R_u}(x)$ . We can then take the homomorphism  $h_i = h_{x, a}$ , which satisfies  $h_i(x) = \mathbf{e}_a$ , that is  $M_x \mathbf{e}_i = \mathbf{e}_a$ . So,  $M_x$  is skeletal and  $(c_6)$  is satisfied. Finally, to check  $(c_4)$ , choose  $a \in A$  and suppose that  $\mathbf{e}_a^T M_x \mathbf{e}_1 = 0$ . Since  $M_x \mathbf{e}_1 = h_1(x) = h_{\bar{\mathbf{x}}, \bar{\mathbf{a}}}(x)$ , this

---

each variable for  $g$  is zero whenever the corresponding variable for  $h$  is zero is equivalent to the existence, for any  $h'$  in the nonempty relative interior of the polytope of solutions of the BLP, of a solution  $g'$  of AIP that sets to zero any variable that is zero in  $h'$ . This is implicit in the analysis in [22].

implies that  $a \notin \text{supp}(h_{\tilde{\mathbf{x}}, \tilde{\mathbf{a}}}(x))$ . Choosing  $R_u$  as  $\tilde{R}$  and  $x$  as  $\tilde{x}$  in 1' of Definition 28, and using again the fact that  $E^{(R_u^{\mathbf{A}}, 1)} = I_n$ , we see that  $\text{supp}(g(x)) \subseteq \text{supp}(h_{\tilde{\mathbf{x}}, \tilde{\mathbf{a}}}(x))$ . Therefore,  $a \notin \text{supp}(g(x)) = \text{supp}(\mu_x)$ . Hence,  $(c_4)$  is satisfied, too, and the claim is proved.

Consider the map  $\gamma : X \rightarrow \mathcal{C}^{(n)}$  defined by  $x \mapsto (M_x, \mu_x)$ . We claim that  $\gamma$  is a homomorphism from  $\mathbf{X}$  to  $\mathbb{F}_{\mathcal{C}}(\mathbf{A})$ . With this claim, we can finish the proof. By assumption, there is a minion homomorphism from  $\mathcal{C}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$ . By Lemma 22 applied to  $\mathcal{C}$ , we have  $\mathbb{F}_{\mathcal{C}}(\mathbf{A}) \rightarrow \mathbf{B}$ . Composing  $\gamma$  with this homomorphism yields  $\mathbf{X} \rightarrow \mathbf{B}$ , as required. It remains to establish the claim.

**Claim:**  $\gamma$  is a homomorphism from  $\mathbf{X}$  to  $\mathbb{F}_{\mathcal{C}}(\mathbf{A})$ .

Take  $R \in \sigma$  of arity  $k$ , and let  $\mathbf{x} = (x_1, \dots, x_k) \in R^{\mathbf{X}}$ . We need to show that  $((M_{x_1}, \mu_{x_1}), \dots, (M_{x_k}, \mu_{x_k})) \in R^{\mathbb{F}_{\mathcal{C}}(\mathbf{A})}$ . For each  $i \in [t] \setminus \{1\}$ , consider a probability distribution  $\xi_i \in \mathbb{S}(R^{\mathbf{A}})$  corresponding to the homomorphism  $h_i$  and witnessing part 2 in Definition 28. Also, consider the probability distribution  $\xi_1 \in \mathbb{S}(R^{\mathbf{A}})$  and the integer distribution  $\zeta \in \mathbb{Z}(R^{\mathbf{A}})$  corresponding to  $h_1$  and  $g$ , respectively, and witnessing 1'. We introduce the matrix  $Q \in \mathbb{Q}^{|R^{\mathbf{A}}|, \mathbb{N}_0}$  and the vector  $\delta \in \mathbb{Z}^{|R^{\mathbf{A}}|}$  defined by

$$\begin{aligned} Q\mathbf{e}_i &= \xi_i \quad \forall i \in [t], \\ Q\mathbf{e}_i &= \xi_t \quad \forall i \in \mathbb{N} \setminus [t], \\ \delta &= \zeta. \end{aligned}$$

We claim that  $(Q, \delta) \in \mathcal{C}^{(|R^{\mathbf{A}}|)}$ . The requirements  $(c_1), (c_2), (c_3)$ , and  $(c_5)$  in Definition 12 are easily seen to be satisfied. Suppose  $\mathbf{e}_{\mathbf{a}}^T Q \neq \mathbf{0}_{\mathbb{N}_0}^T$  for some  $\mathbf{a} = (a_1, \dots, a_k) \in R^{\mathbf{A}}$ , so that there exists  $d \in [t]$  such that  $\mathbf{e}_{\mathbf{a}}^T Q \mathbf{e}_d \neq 0$ . Hence,  $\mathbf{a} \in \text{supp}(Q\mathbf{e}_d) = \text{supp}(\xi_d) \subseteq s^R(\mathbf{x})$ . Pick  $h_j = h_{\mathbf{x}, \mathbf{a}}$ . We have that

$$E^{(R^{\mathbf{A}}, p)} \xi_j = h_j(x_p) = h_{\mathbf{x}, \mathbf{a}}(x_p) = \mathbf{e}_{a_p} \quad \forall p \in [k].$$

Suppose that  $\xi_j \neq \mathbf{e}_{\mathbf{a}}$ . Then,  $\exists \mathbf{a}' = (a'_1, \dots, a'_k) \in R^{\mathbf{A}}$  such that  $\mathbf{a}' \neq \mathbf{a}$  and  $\mathbf{e}_{\mathbf{a}'}^T \xi_j > 0$ . Choose  $q \in [k]$  such that  $a'_q \neq a_q$ , and observe that

$$0 = \mathbf{e}_{a'_q}^T \mathbf{e}_{a_q} = \mathbf{e}_{a'_q}^T E^{(R^{\mathbf{A}}, q)} \xi_j \geq \mathbf{e}_{\mathbf{a}'}^T \xi_j > 0,$$

which is a contradiction. Hence,  $Q\mathbf{e}_j = \xi_j = \mathbf{e}_{\mathbf{a}}$ . We conclude that  $Q$  is skeletal and, therefore,  $(c_6)$  is satisfied. Finally, suppose that  $\mathbf{a} \notin \text{supp}(Q\mathbf{e}_1) = \text{supp}(\xi_1)$  for some  $\mathbf{a} \in R^{\mathbf{A}}$ . Recalling that  $\xi_1 \in \mathbb{S}(R^{\mathbf{A}})$  corresponds to the homomorphism  $h_1 = h_{\tilde{\mathbf{x}}, \tilde{\mathbf{a}}}$ , it follows from 1' that  $\text{supp}(\zeta) \subseteq \text{supp}(\xi_1)$ . Hence,  $\mathbf{a} \notin \text{supp}(\zeta) = \text{supp}(\delta)$ , so that  $(c_4)$  is satisfied, too. As a consequence,  $(Q, \delta) \in \mathcal{C}^{(|R^{\mathbf{A}}|)}$ , as claimed.

Now, we need to show that  $(M_{x_\alpha}, \mu_{x_\alpha}) = (Q, \delta)_{/\pi_\alpha}$  for each  $\alpha \in [k]$ , where  $\pi_\alpha : R^{\mathbf{A}} \rightarrow A$  maps  $\mathbf{a} \in R^{\mathbf{A}}$  to its  $\alpha$ -th coordinate. Observe first that, by definition,  $P_{\pi_\alpha} = E^{(R^{\mathbf{A}}, \alpha)}$  for each  $\alpha \in [k]$ . We see that

$$\begin{aligned} Q_{/\pi_\alpha} \mathbf{e}_i &= P_{\pi_\alpha} Q \mathbf{e}_i = E^{(R^{\mathbf{A}}, \alpha)} Q \mathbf{e}_i = E^{(R^{\mathbf{A}}, \alpha)} \xi_i = h_i(x_\alpha) = M_{x_\alpha} \mathbf{e}_i \quad \text{for } i \in [t] \quad \text{and} \\ Q_{/\pi_\alpha} \mathbf{e}_i &= P_{\pi_\alpha} Q \mathbf{e}_i = P_{\pi_\alpha} Q \mathbf{e}_t = M_{x_\alpha} \mathbf{e}_t = M_{x_\alpha} \mathbf{e}_i \quad \text{for } i \in \mathbb{N} \setminus [t], \end{aligned}$$

which yields  $Q_{/\pi_\alpha} = M_{x_\alpha}$ . Moreover,

$$\delta_{/\pi_\alpha} = P_{\pi_\alpha} \delta = E^{(R^{\mathbf{A}}, \alpha)} \delta = E^{(R^{\mathbf{A}}, \alpha)} \zeta = g(x_\alpha) = \mu_{x_\alpha}.$$



It follows that  $(M_{x_\alpha}, \mu_{x_\alpha}) = (Q_{/\pi_\alpha}, \delta_{/\pi_\alpha}) = (Q, \delta)_{/\pi_\alpha}$ . By Definition 21,  $((M_{x_1}, \mu_{x_1}), \dots, (M_{x_k}, \mu_{x_k})) \in R^{\mathbb{F}_{\mathcal{C}}(\mathbf{A})}$ , so  $\gamma : \mathbf{X} \rightarrow \mathbb{F}_{\mathcal{C}}(\mathbf{A})$  is a homomorphism.  $\square$

Our second goal is to prove the following.

**Theorem 31.** *If CLAP solves PCSP( $\mathbf{A}, \mathbf{B}$ ) then there is a minion homomorphism from  $\mathcal{C}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$ .*

**Remark 32.** The proof of Theorem 31 proceeds essentially by establishing that the free structure  $\mathbb{F}_{\mathcal{C}}(\mathbf{A})$  has the CLAP condition as an instance of PCSP( $\mathbf{A}, \mathbf{B}$ ). However, some care is needed when handling Proposition 29, which only applies to *finite* structures, while  $\mathbb{F}_{\mathcal{C}}(\mathbf{A})$  is not finite in general. To overcome this problem, we use a compactness argument tailored to our minion  $\mathcal{C}$  discussed in Section 4.2, which follows the ideas of [8].

We remark that the compactness argument for relational structures in the form stated in [22, Lemma A.6] does not entirely fit our proof structure, as the element  $(\mathbf{e}_1 \mathbf{1}_{\aleph_0}^T, \mathbf{e}_1)$  having the role of  $\bar{\mathbf{x}}$  in Definition 28 does not belong to every induced substructure of  $\mathbb{F}_{\mathcal{C}}(\mathbf{A})$ . A different option would have been to use the general compactness argument known as the (uncountable version of the) *compactness theorem of logic* [62], that applies to all *minion tests*<sup>9</sup> as derived in [31, Proposition 6] through [64].

*Proof of Theorem 31.* Let  $n = |A|$ . For  $D \in \mathbb{N}$ , denote  $\mathbb{F}_{\mathcal{C}_D}(\mathbf{A})$  by  $\mathbf{F}$  (where  $\mathcal{C}_D$  is the subminion of  $\mathcal{C}$  introduced in Section 4.2). Hence, the domain of  $\mathbf{F}$  is  $\mathcal{C}_D^{(n)}$ , which is finite. We claim that  $\mathbf{F}$  has the CLAP condition as an instance of PCSP( $\mathbf{A}, \mathbf{B}$ ).

For each  $R \in \sigma$  of arity  $k$  and for each  $\tau = ((M_1, \mu_1), \dots, (M_k, \mu_k)) \in R^{\mathbf{F}}$ , take  $(Q_\tau, \delta_\tau) \in \mathcal{C}_D^{(|R^{\mathbf{A}}|)}$  satisfying  $(M_j, \mu_j) = (Q_\tau, \delta_\tau)_{/\pi_j} \forall j \in [k]$ , where  $\pi_j : R^{\mathbf{A}} \rightarrow A$  maps  $\mathbf{a} \in R^{\mathbf{A}}$  to its  $j$ -th coordinate; i.e.,  $M_j = E^{(R^{\mathbf{A}}, j)} Q_\tau$  and  $\mu_j = E^{(R^{\mathbf{A}}, j)} \delta_\tau \forall j \in [k]$ . Given  $R \in \sigma$  of arity  $k$ , consider the map

$$\begin{aligned} s^R : R^{\mathbf{F}} &\rightarrow \mathcal{P}(R^{\mathbf{A}}) \setminus \{\emptyset\} \\ \tau &\mapsto \bigcup_{i \in \mathbb{N}} \text{supp}(Q_\tau \mathbf{e}_i). \end{aligned}$$

Let us first check part (I) of Definition 28. Pick  $\tau = ((M_1, \mu_1), \dots, (M_k, \mu_k)) \in R^{\mathbf{F}}$  and  $\mathbf{a} = (a_1, \dots, a_k) \in s^R(\tau)$ . We have that  $\mathbf{a} \in \text{supp}(Q_\tau \mathbf{e}_\alpha)$  for some  $\alpha \in \mathbb{N}$ , i.e.,  $\mathbf{e}_\alpha^T Q_\tau \mathbf{e}_\alpha \neq 0$ . Since  $Q_\tau$  is skeletal, the set  $L_{\tau, \mathbf{a}} = \{\ell \in \mathbb{N} : Q_\tau \mathbf{e}_\ell = \mathbf{e}_\mathbf{a}\}$  is nonempty; let  $\ell(\tau, \mathbf{a}) := \min(L_{\tau, \mathbf{a}})$ . Consider the map

$$\begin{aligned} h_{\tau, \mathbf{a}} : \mathcal{C}_D^{(n)} &\rightarrow \mathbb{S}(A) \\ (\hat{M}, \hat{\mu}) &\mapsto \hat{M} \mathbf{e}_{\ell(\tau, \mathbf{a})}. \end{aligned}$$

We claim that  $h_{\tau, \mathbf{a}}$  is a homomorphism from  $\mathbf{F}$  to  $\mathbb{S}(\mathbf{A})$ . Take  $\tilde{R} \in \sigma$  of arity  $\tilde{k}$ , and let  $\tilde{\tau} = ((\tilde{M}_1, \tilde{\mu}_1), \dots, (\tilde{M}_{\tilde{k}}, \tilde{\mu}_{\tilde{k}})) \in \tilde{R}^{\mathbf{F}}$ . Consider the pair  $(Q_{\tilde{\tau}}, \delta_{\tilde{\tau}}) \in \mathcal{C}_D^{(|\tilde{R}^{\mathbf{A}}|)}$ . We have that

$$h_{\tau, \mathbf{a}}(\tilde{\tau}) = (\tilde{M}_1 \mathbf{e}_{\ell(\tau, \mathbf{a})}, \dots, \tilde{M}_{\tilde{k}} \mathbf{e}_{\ell(\tau, \mathbf{a})}) = \left( E^{(\tilde{R}^{\mathbf{A}}, 1)} Q_{\tilde{\tau}} \mathbf{e}_{\ell(\tau, \mathbf{a})}, \dots, E^{(\tilde{R}^{\mathbf{A}}, \tilde{k})} Q_{\tilde{\tau}} \mathbf{e}_{\ell(\tau, \mathbf{a})} \right).$$

---

<sup>9</sup>Cf. Remark 33.

Since  $Q_{\tilde{\tau}}\mathbf{e}_{\ell(\tau, \mathbf{a})} \in \mathbb{S}(\tilde{R}^{\mathbf{A}})$ , we deduce that  $h_{\tau, \mathbf{a}}(\tilde{\tau}) \in \tilde{R}^{\mathbb{S}(\mathbf{A})}$ , as wanted. Therefore,  $h_{\tau, \mathbf{a}}$  is a homomorphism from  $\mathbf{F}$  to  $\mathbb{S}(\mathbf{A})$ . We now check that the requirements 1 and 2 in Definition 28 are met. The former follows from

$$h_{\tau, \mathbf{a}}((M_i, \boldsymbol{\mu}_i)) = M_i \mathbf{e}_{\ell(\tau, \mathbf{a})} = E^{(R^{\mathbf{A}}, i)} Q_{\tau} \mathbf{e}_{\ell(\tau, \mathbf{a})} = E^{(R^{\mathbf{A}}, i)} \mathbf{e}_{\mathbf{a}} = \mathbf{e}_{a_i} \quad \forall i \in [k].$$

To check the latter requirement, take  $\tilde{R} \in \sigma$  of arity  $\tilde{k}$  and  $\tilde{\tau} = ((\tilde{M}_1, \tilde{\boldsymbol{\mu}}_1), \dots, (\tilde{M}_{\tilde{k}}, \tilde{\boldsymbol{\mu}}_{\tilde{k}})) \in \tilde{R}^{\mathbf{F}}$ , and consider  $\boldsymbol{\xi} := Q_{\tilde{\tau}} \mathbf{e}_{\ell(\tau, \mathbf{a})}$ . Observe that

- $h_{\tau, \mathbf{a}}((\tilde{M}_i, \tilde{\boldsymbol{\mu}}_i)) = \tilde{M}_i \mathbf{e}_{\ell(\tau, \mathbf{a})} = E^{(\tilde{R}^{\mathbf{A}}, i)} Q_{\tilde{\tau}} \mathbf{e}_{\ell(\tau, \mathbf{a})} = E^{(\tilde{R}^{\mathbf{A}}, i)} \boldsymbol{\xi} \quad \forall i \in [\tilde{k}]$
- $\text{supp}(\boldsymbol{\xi}) = \text{supp}(Q_{\tilde{\tau}} \mathbf{e}_{\ell(\tau, \mathbf{a})}) \subseteq \bigcup_{i \in \mathbb{N}} \text{supp}(Q_{\tilde{\tau}} \mathbf{e}_i) = s^{\tilde{R}}(\tilde{\tau}).$

We now check part (II) of Definition 28. Take  $R_u$  as  $\bar{R}$ , and observe that

$$R_u^{\mathbf{F}} = \{(M, \boldsymbol{\mu}) \in \mathcal{C}_D^{(n)} : \exists (Q, \boldsymbol{\delta}) \in \mathcal{C}_D^{(n)} \text{ such that } M = E^{(R_u^{\mathbf{A}}, 1)} Q, \boldsymbol{\mu} = E^{(R_u^{\mathbf{A}}, 1)} \boldsymbol{\delta}\} = \mathcal{C}_D^{(n)},$$

where we have used that  $E^{(R_u^{\mathbf{A}}, 1)} = I_n$ . Consider the element  $\bar{\tau} = (\mathbf{e}_1 \mathbf{1}_{\mathbb{N}_0}^T, \mathbf{e}_1) \in \mathcal{C}_D^{(n)} = R_u^{\mathbf{F}}$ . Using again that  $E^{(R_u^{\mathbf{A}}, 1)} = I_n$ , we see that  $(Q_{\bar{\tau}}, \boldsymbol{\delta}_{\bar{\tau}}) = \bar{\tau}$ . We obtain

$$s^{R_u}(\bar{\tau}) = \bigcup_{i \in \mathbb{N}} \text{supp}(\mathbf{e}_1 \mathbf{1}_{\mathbb{N}_0}^T \mathbf{e}_i) = \bigcup_{i \in \mathbb{N}} \text{supp}(\mathbf{e}_1) = \{1\}.$$

Hence, we pick  $\bar{a} = 1$ . Notice that

$$\ell(\bar{\tau}, \bar{a}) = \min\{\ell \in \mathbb{N} : \mathbf{e}_1 \mathbf{1}_{\mathbb{N}_0}^T \mathbf{e}_\ell = \mathbf{e}_1\} = \min\{\ell \in \mathbb{N} : \mathbf{e}_1 = \mathbf{e}_1\} = \min \mathbb{N} = 1.$$

Consider the function

$$\begin{aligned} g : \mathcal{C}_D^{(n)} &\rightarrow \mathbb{Z}(\mathbf{A}) \\ (\hat{M}, \hat{\boldsymbol{\mu}}) &\mapsto \hat{\boldsymbol{\mu}}. \end{aligned}$$

Following the same procedure as for  $h_{\tau, \mathbf{a}}$ , we easily check that  $g$  is a homomorphism from  $\mathbf{F}$  to  $\mathbb{Z}(\mathbf{A})$ . We now verify that condition 1' of Definition 28 is satisfied. Given  $\tilde{R} \in \sigma$  of arity  $\tilde{k}$  and  $\tilde{\tau} = ((\tilde{M}_1, \tilde{\boldsymbol{\mu}}_1), \dots, (\tilde{M}_{\tilde{k}}, \tilde{\boldsymbol{\mu}}_{\tilde{k}})) \in \tilde{R}^{\mathbf{F}}$ , let  $\boldsymbol{\xi} := Q_{\tilde{\tau}} \mathbf{e}_1 \in \mathbb{S}(\tilde{R}^{\mathbf{A}})$  and  $\boldsymbol{\zeta} := \boldsymbol{\delta}_{\tilde{\tau}} \in \mathbb{Z}(\tilde{R}^{\mathbf{A}})$ . Then, given  $i \in [\tilde{k}]$ ,

- \*  $h_{\bar{\tau}, \bar{a}}((\tilde{M}_i, \tilde{\boldsymbol{\mu}}_i)) = \tilde{M}_i \mathbf{e}_{\ell(\bar{\tau}, \bar{a})} = \tilde{M}_i \mathbf{e}_1 = E^{(\tilde{R}^{\mathbf{A}}, i)} Q_{\tilde{\tau}} \mathbf{e}_1 = E^{(\tilde{R}^{\mathbf{A}}, i)} \boldsymbol{\xi};$
- \*  $g((\tilde{M}_i, \tilde{\boldsymbol{\mu}}_i)) = \tilde{\boldsymbol{\mu}}_i = E^{(\tilde{R}^{\mathbf{A}}, i)} \boldsymbol{\delta}_{\tilde{\tau}} = E^{(\tilde{R}^{\mathbf{A}}, i)} \boldsymbol{\zeta};$
- \*  $\text{supp}(\boldsymbol{\zeta}) = \text{supp}(\boldsymbol{\delta}_{\tilde{\tau}}) \subseteq \text{supp}(Q_{\tilde{\tau}} \mathbf{e}_1) = \text{supp}(\boldsymbol{\xi}) \subseteq \bigcup_{i \in \mathbb{N}} \text{supp}(Q_{\tilde{\tau}} \mathbf{e}_i) = s^{\tilde{R}}(\tilde{\tau})$

where, for the first inclusion in the third line, we have used  $(c_4)$  in Definition 12.

It follows that  $\mathbf{F}$  has the CLAP condition as an instance of PCSP( $\mathbf{A}, \mathbf{B}$ ), as claimed. Then, Proposition 29 implies that CLAP accepts  $\mathbf{F}$ . Since, by hypothesis, CLAP solves PCSP( $\mathbf{A}, \mathbf{B}$ ), we deduce that  $\mathbb{F}_{\mathcal{C}_D}(\mathbf{A}) = \mathbf{F} \rightarrow \mathbf{B}$ . By Lemma 22, there exists a minion homomorphism from  $\mathcal{C}_D$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$ . Finally, since the set of polymorphisms of  $(\mathbf{A}, \mathbf{B})$  of arity  $L$  is finite for every  $L \in \mathbb{N}$ , Proposition 25 allows us to conclude that there exists a minion homomorphism from  $\mathcal{C}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$ .  $\square$

**Remark 33.** It follows from the proofs of Theorems 30 and 31 that CLAP fits within the framework of *minion tests* recently introduced in [31]. More precisely,  $\text{CLAP} = \text{Test}_{\mathcal{C}}$ , which means that, for two  $\sigma$ -structures  $\mathbf{X}$  and  $\mathbf{A}$ ,  $\text{CLAP}(\mathbf{X}, \mathbf{A})$  accepts if and only if  $\mathbf{X} \rightarrow \mathbb{F}_{\mathcal{C}}(\mathbf{A})$ . Additionally, it follows from [31] that CLAP is a *conic* minion test, which essentially means that one can build a progressively tighter hierarchy of relaxations based on CLAP whose  $k$ -th level correctly classifies all instances of size  $k$ .

## 5 H-symmetric polymorphisms

This section contains the proof of Theorem 15. We remark that the machinery developed here can be extended to the more general setting of  $H$ -block-symmetric polymorphisms, at the only cost of dealing with a more cumbersome notation. This is done in Appendix C and results in Theorem 43 – a slightly stronger version of Theorem 15.

We shall need two helpful lemmas. The first lemma shows a property of  $H$ -symmetric functions that will be useful in the proof of Theorem 15. Throughout this section, without loss of generality, we consider  $A = [n]$ .

**Lemma 34.** *Let  $f : A^L \rightarrow B$  be  $H$ -symmetric for some  $m \times n$  tie matrix  $H$ , with  $m \in \mathbb{N}$ . Consider two maps  $\pi, \tilde{\pi} : [L] \rightarrow [n]$  such that  $P_{\pi} \mathbf{1}_L = P_{\tilde{\pi}} \mathbf{1}_L$  and the vector  $P_{\pi} \mathbf{1}_L$  is  $H$ -tieless. Then*

$$f_{/\pi}(1, \dots, n) = f_{/\tilde{\pi}}(1, \dots, n).$$

*Proof.* For  $a \in [n]$ , we have

$$|\pi^{-1}(a)| = \sum_{i \in [L]} (P_{\pi})_{ai} = \sum_{i \in [L]} \mathbf{e}_a^T P_{\pi} \mathbf{e}_i = \mathbf{e}_a^T P_{\pi} \mathbf{1}_L = \mathbf{e}_a^T P_{\tilde{\pi}} \mathbf{1}_L = |\tilde{\pi}^{-1}(a)|.$$

Hence, we can consider bijections  $\varphi_a : \pi^{-1}(a) \rightarrow \tilde{\pi}^{-1}(a)$  for each  $a \in [n]$ . Clearly, their union

$$\varphi = \bigcup_{a \in [n]} \varphi_a : [L] \rightarrow [L]$$

is also a bijection. For each  $i \in [L]$ , we have

$$(\tilde{\pi} \circ \varphi)(i) = \tilde{\pi}(\varphi(i)) = \tilde{\pi}(\varphi_{\pi(i)}(i)) = \pi(i)$$

and, hence,  $\tilde{\pi} \circ \varphi = \pi$ . Let  $\tilde{\mathbf{a}} = (\tilde{\pi}(1), \dots, \tilde{\pi}(L))$ . Notice that, for each  $a \in [n]$ ,

$$\mathbf{e}_a^T \tilde{\mathbf{a}}^{\#} = |\{i \in [L] : \tilde{\pi}(i) = a\}| = \mathbf{e}_a^T P_{\tilde{\pi}} \mathbf{1}_L$$

and, therefore,  $\tilde{\mathbf{a}}^{\#} = P_{\tilde{\pi}} \mathbf{1}_L = P_{\pi} \mathbf{1}_L$ , which is  $H$ -tieless. Using that  $f$  is  $H$ -symmetric, we find

$$f_{/\tilde{\pi}}(1, \dots, n) = f(\tilde{\mathbf{a}}) = f_{/\varphi}(\tilde{\mathbf{a}}) = (f_{/\varphi})_{/\tilde{\pi}}(1, \dots, n) = f_{/\tilde{\pi} \circ \varphi}(1, \dots, n) = f_{/\pi}(1, \dots, n),$$

as required.  $\square$

One intriguing property of skeletal matrices is that they can be simultaneously reduced to  $H$ -tieless vectors, in the sense of the next lemma. We say that a vector is *finitely supported* if it only has a finite number of nonzero entries.

**Lemma 35** (Tiebreak Lemma). *For  $k, p, m \in \mathbb{N}$ , let  $M_1, \dots, M_k \in \mathbb{Q}^{p, \aleph_0}$  be skeletal matrices, and let  $H$  be an  $m \times p$  tie matrix. Then there exists a stochastic finitely supported vector  $\mathbf{v} \in \mathbb{Q}^{\aleph_0}$  with  $\mathbf{e}_1^T \mathbf{v} > 0$  such that  $M_j \mathbf{v}$  is  $H$ -tieless for any  $j \in [k]$ .*

*Proof.* Let  $\Omega$  be the set of rational stochastic finitely supported vectors of size  $\aleph_0$  whose first entry is nonzero, and consider the map

$$f : \Omega \rightarrow \mathbb{N}_0$$

$$\hat{\mathbf{v}} \mapsto \sum_{j \in [k]} |\{(i, i') \in [m]^2 : i \neq i' \text{ and } \mathbf{e}_i^T H M_j \hat{\mathbf{v}} = \mathbf{e}_{i'}^T H M_j \hat{\mathbf{v}} \neq 0\}|.$$

In other words,  $f(\hat{\mathbf{v}})$  counts the total number of ties in the set of vectors  $\{H M_j \hat{\mathbf{v}} : j \in [k]\}$ . Let  $\mathbf{v}$  attain the minimum of  $f$  over  $\Omega$ . If  $f(\mathbf{v}) = 0$ , we are done. Otherwise, let  $j \in [k]$ ,  $i, i' \in [m]$  be such that  $i \neq i'$  and  $\mathbf{e}_i^T H M_j \mathbf{v} = \mathbf{e}_{i'}^T H M_j \mathbf{v} \neq 0$ . From  $\mathbf{e}_i^T H M_j \mathbf{v} \neq 0$ , we see that  $\exists \beta \in [p]$  such that  $\mathbf{e}_i^T H \mathbf{e}_\beta \neq 0$  and  $\mathbf{e}_\beta^T M_j \mathbf{v} \neq 0$ . In particular, we have  $\mathbf{e}_\beta^T M_j \neq \mathbf{0}_{\aleph_0}^T$ ; since  $M_j$  is skeletal, this implies that  $M_j \mathbf{e}_\alpha = \mathbf{e}_\beta$  for some  $\alpha \in \mathbb{N}$ . For  $\epsilon \in \mathbb{Q}$ ,  $0 < \epsilon < 1$ , consider the vector  $\mathbf{v}_\epsilon = (1 - \epsilon)\mathbf{v} + \epsilon \mathbf{e}_\alpha$ . Observe that  $\mathbf{v}_\epsilon \in \Omega$ . For  $g \in [k]$ , we have  $H M_g \mathbf{v}_\epsilon = (1 - \epsilon)H M_g \mathbf{v} + \epsilon H M_g \mathbf{e}_\alpha$ . By choosing  $\epsilon$  sufficiently small, we can assume that, for each  $g \in [k]$ ,  $H M_g \mathbf{v}_\epsilon$  does not have new ties other than those in  $H M_g \mathbf{v}$ . Moreover,

$$H M_j \mathbf{v}_\epsilon = (1 - \epsilon)H M_j \mathbf{v} + \epsilon H M_j \mathbf{e}_\alpha = (1 - \epsilon)H M_j \mathbf{v} + \epsilon H \mathbf{e}_\beta$$

and, hence,

$$\begin{aligned} \mathbf{e}_i^T H M_j \mathbf{v}_\epsilon &= (1 - \epsilon)\mathbf{e}_i^T H M_j \mathbf{v} + \epsilon \mathbf{e}_i^T H \mathbf{e}_\beta = (1 - \epsilon)\mathbf{e}_i^T H M_j \mathbf{v} + \epsilon \mathbf{e}_i^T H \mathbf{e}_\beta \\ &\neq (1 - \epsilon)\mathbf{e}_{i'}^T H M_j \mathbf{v} + \epsilon \mathbf{e}_{i'}^T H \mathbf{e}_\beta = \mathbf{e}_{i'}^T H M_j \mathbf{v}_\epsilon, \end{aligned}$$

where the disequality follows from  $\mathbf{e}_i^T H \mathbf{e}_\beta \neq 0$  and from the fact that  $H \mathbf{e}_\beta$  is a tieless vector by the definition of tie matrix. We conclude that  $f(\mathbf{v}_\epsilon) < f(\mathbf{v})$ , which contradicts our assumption.  $\square$

**Theorem** (Theorem 15 restated). *Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template and suppose  $\text{Pol}(\mathbf{A}, \mathbf{B})$  contains  $H$ -symmetric operations of arbitrarily large arity for some  $m \times |A|$  tie matrix  $H$ ,  $m \in \mathbb{N}$ . Then there exists a minion homomorphism from  $\mathcal{C}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$ .*

**Remark 36.** Before proving Theorem 15, we provide some intuition on the construction of the minion homomorphism whose existence shall establish the result. First, one fixes an  $H$ -symmetric polymorphism  $f$ . Then, the image of an  $L$ -ary element  $(M, \boldsymbol{\mu})$  of  $\mathcal{C}$  under the homomorphism is the function that (i) takes a tuple  $(a_1, \dots, a_L)$  of variables in  $A$  as input, (ii) deforms the tuple by changing the frequency of each variable according to the information carried by  $M$  and  $\boldsymbol{\mu}$ , and (iii) returns as output the evaluation of  $f$  on the deformed tuple. The deformation in step (ii) is encoded by the map  $\varphi$  defined in (4). Essentially,  $\varphi$  decides what frequency to assign to a variable  $a_i$  on the basis of the weight of  $i$  in the probability distribution  $M \mathbf{v}$  – where  $\mathbf{v}$  is the tie-breaking vector from Lemma 35. The integer distribution  $\boldsymbol{\mu}$  is also taken into account by  $\varphi$ , and its role is essentially to fill the gap between the size of the deformed tuple obtained above and the arity of  $f$ . If  $\text{Pol}(\mathbf{A}, \mathbf{B})$  is rich enough to provide  $H$ -symmetric polymorphisms of whichever arity we need,  $\boldsymbol{\mu}$  is inessential (cf. Remark 37).

*Proof of Theorem 15.* For  $D \in \mathbb{N}$ , consider the subminion  $\mathcal{C}_D$  of  $\mathcal{C}$  described in Section 4.2. Observe that  $S = \{M : (M, \boldsymbol{\mu}) \in \mathcal{C}_D^{(n)}\}$  is a finite set of skeletal matrices. Therefore, we can apply the Tiebreak Lemma 35 to find a stochastic finitely supported vector  $\mathbf{v} \in \mathbb{Q}^{\mathbb{N}_0}$  with  $\mathbf{e}_1^T \mathbf{v} > 0$  such that  $M\mathbf{v}$  is  $H$ -tieless for any  $M \in S$ . Since  $\mathbf{v}$  is finitely supported, we can find  $N' \in \mathbb{N}$  such that  $N'\mathbf{v}$  has integer entries. Let  $\sigma_1^H$  denote the largest singular value of  $H$  – i.e., the square root of the largest eigenvalue of  $H^T H$ . Set  $N = 2\lceil \sigma_1^H + 1 \rceil D^2 N'$ , and let  $f$  be an  $H$ -symmetric polymorphism of arity  $c \geq N^2$ . Write  $c = N\alpha + \beta$  with  $\alpha, \beta \in \mathbb{N}_0$ ,  $\beta \leq N - 1$ . Note that  $N^2 \leq c = N\alpha + \beta \leq N\alpha + N - 1 < N(\alpha + 1)$ , so  $N < \alpha + 1$  and, hence,  $\beta < \alpha$ .

Consider the function

$$\xi_D : \mathcal{C}_D \rightarrow \text{Pol}(\mathbf{A}, \mathbf{B})$$

defined as follows. Given  $L \in \mathbb{N}$  and  $(M, \boldsymbol{\mu}) \in \mathcal{C}_D^{(L)}$ , take the map  $\varphi : [c] \rightarrow [L]$  such that the corresponding  $L \times c$  matrix  $P_\varphi$  is

$$P_\varphi = \begin{pmatrix} \mathbf{1}_{\mathbf{e}_1^T(\alpha N M \mathbf{v} + \beta \boldsymbol{\mu})}^T & \mathbf{0}^T & \cdots & \mathbf{0}^T \\ \mathbf{0}^T & \mathbf{1}_{\mathbf{e}_2^T(\alpha N M \mathbf{v} + \beta \boldsymbol{\mu})}^T & \cdots & \mathbf{0}^T \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0}^T & \mathbf{0}^T & \cdots & \mathbf{1}_{\mathbf{e}_L^T(\alpha N M \mathbf{v} + \beta \boldsymbol{\mu})}^T \end{pmatrix}. \quad (4)$$

To verify that (4) is well defined, observe first that

$$\sum_{i=1}^L \mathbf{e}_i^T(\alpha N M \mathbf{v} + \beta \boldsymbol{\mu}) = \mathbf{1}_L^T(\alpha N M \mathbf{v} + \beta \boldsymbol{\mu}) = \alpha N \mathbf{1}_L^T M \mathbf{v} + \beta \mathbf{1}_L^T \boldsymbol{\mu} = \alpha N \mathbf{1}_{\mathbb{N}_0}^T \mathbf{v} + \beta = \alpha N + \beta = c.$$

Moreover, for each  $i \in [L]$ ,  $\mathbf{e}_i^T(\alpha N M \mathbf{v} + \beta \boldsymbol{\mu}) = \mathbf{e}_i^T(2\alpha\lceil \sigma_1^H + 1 \rceil D(DM)(N'\mathbf{v}) + \beta \boldsymbol{\mu})$  is an integer. If  $\mathbf{e}_i^T(\alpha N M \mathbf{v} + \beta \boldsymbol{\mu})$  was negative, then  $\mathbf{e}_i^T \boldsymbol{\mu} < 0$ . By the requirement  $(c_4)$  in Definition 12, this would imply that  $\mathbf{e}_i^T M \mathbf{e}_1 > 0$  and, hence,  $0 < \mathbf{e}_i^T M \mathbf{e}_1 \mathbf{e}_1^T \mathbf{v} \leq \mathbf{e}_i^T M \mathbf{v}$ . As a consequence,  $\mathbf{e}_i^T(DM)(N'\mathbf{v}) \geq 1$  so that

$$\mathbf{e}_i^T(\alpha N M \mathbf{v} + \beta \boldsymbol{\mu}) = 2\alpha\lceil \sigma_1^H + 1 \rceil D \mathbf{e}_i^T(DM)(N'\mathbf{v}) + \beta \mathbf{e}_i^T \boldsymbol{\mu} \geq 2\alpha\lceil \sigma_1^H + 1 \rceil D + \beta \mathbf{e}_i^T \boldsymbol{\mu} \geq \alpha D - \beta D > 0,$$

which is a contradiction. In conclusion, the numbers  $\mathbf{e}_i^T(\alpha N M \mathbf{v} + \beta \boldsymbol{\mu})$  are nonnegative integers summing up to  $c$ , so (4) is well defined.

We define  $\xi_D((M, \boldsymbol{\mu})) := f_{/\varphi}$ . Clearly,  $\xi_D((M, \boldsymbol{\mu})) \in \text{Pol}(\mathbf{A}, \mathbf{B})$ . We claim that the map  $\xi_D$  is a minion homomorphism. It is straightforward to check that  $\xi_D$  preserves arities so, to conclude, we need to show that it also preserves minors. Take  $L' \in \mathbb{N}$  and choose a map  $\pi : [L] \rightarrow [L']$ . Letting  $\tilde{\varphi} : [c] \rightarrow [L']$  be the map corresponding to the matrix

$$P_{\tilde{\varphi}} = \begin{pmatrix} \mathbf{1}_{\mathbf{e}_1^T(\alpha N P_\pi M \mathbf{v} + \beta P_\pi \boldsymbol{\mu})}^T & \mathbf{0}^T & \cdots & \mathbf{0}^T \\ \mathbf{0}^T & \mathbf{1}_{\mathbf{e}_2^T(\alpha N P_\pi M \mathbf{v} + \beta P_\pi \boldsymbol{\mu})}^T & \cdots & \mathbf{0}^T \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0}^T & \mathbf{0}^T & \cdots & \mathbf{1}_{\mathbf{e}_{L'}^T(\alpha N P_\pi M \mathbf{v} + \beta P_\pi \boldsymbol{\mu})}^T \end{pmatrix},$$

we see that  $\xi_D((M, \boldsymbol{\mu})/\pi) = f/\tilde{\varphi}$ . Moreover,  $\xi_D((M, \boldsymbol{\mu}))/\pi = (f/\varphi)/\pi = f/\pi \circ \varphi$ , where  $\varphi$  corresponds to the matrix  $P_\varphi$  in (4). Take  $\mathbf{a} = (a_1, \dots, a_{L'}) \in A^{L'}$ , and consider the map

$$\begin{aligned} \pi_{\mathbf{a}} : [L'] &\rightarrow [n] \\ i &\mapsto a_i. \end{aligned}$$

Observe that

$$\begin{aligned} f/\tilde{\varphi}(\mathbf{a}) &= (f/\tilde{\varphi})/\pi_{\mathbf{a}}(1, \dots, n) = f/\pi_{\mathbf{a} \circ \tilde{\varphi}}(1, \dots, n) \quad \text{and, similarly,} \\ f/\pi \circ \varphi(\mathbf{a}) &= (f/\pi \circ \varphi)/\pi_{\mathbf{a}}(1, \dots, n) = f/\pi_{\mathbf{a} \circ \pi \circ \varphi}(1, \dots, n). \end{aligned} \quad (5)$$

Notice that

$$\begin{aligned} P_{\pi_{\mathbf{a} \circ \tilde{\varphi}}} \mathbf{1}_c &= P_{\pi_{\mathbf{a}}} P_{\tilde{\varphi}} \mathbf{1}_c = P_{\pi_{\mathbf{a}}} (\alpha N P_{\pi} M \mathbf{v} + \beta P_{\pi} \boldsymbol{\mu}) \\ &= P_{\pi_{\mathbf{a}}} P_{\pi} (\alpha N M \mathbf{v} + \beta \boldsymbol{\mu}) = P_{\pi_{\mathbf{a}}} P_{\pi} P_{\varphi} \mathbf{1}_c = P_{\pi_{\mathbf{a} \circ \pi \circ \varphi}} \mathbf{1}_c. \end{aligned}$$

We claim that the vector  $P_{\pi_{\mathbf{a} \circ \tilde{\varphi}}} \mathbf{1}_c$  is  $H$ -tieless. Let  $\mathbf{u} = (u_i) = H P_{\pi_{\mathbf{a} \circ \tilde{\varphi}}} \mathbf{1}_c$ ; the claim is equivalent to  $\mathbf{u}$  being tieless. Let  $\mathbf{w} = (w_i) = \alpha N H P_{\pi_{\mathbf{a} \circ \pi}} M \mathbf{v}$  and  $\mathbf{z} = (z_i) = \beta H P_{\pi_{\mathbf{a} \circ \pi}} \boldsymbol{\mu}$ , so that  $\mathbf{u} = \mathbf{w} + \mathbf{z}$ . Choose  $i, i' \in [m]$  such that  $i \neq i'$  and  $u_i \neq 0$ . We need to show that  $u_i \neq u_{i'}$ . Suppose  $w_i = 0$ . We can write  $H^T \mathbf{e}_i = \sum_{g \in G} \lambda_g \mathbf{e}_g$  for  $G = \text{supp}(H^T \mathbf{e}_i)$ , where each  $\lambda_g$  is a positive integer (note that  $G \neq \emptyset$  since, otherwise,  $H^T \mathbf{e}_i = \mathbf{0}_n$ , which would imply  $u_i = 0$ ). Let  $F = (\pi_{\mathbf{a}} \circ \pi)^{-1}(G)$ . From  $w_i = 0$ , we obtain

$$0 = \mathbf{e}_i^T H P_{\pi_{\mathbf{a} \circ \pi}} M \mathbf{v} = (H^T \mathbf{e}_i)^T P_{\pi_{\mathbf{a} \circ \pi}} M \mathbf{v} = \sum_{g \in G} \lambda_g \mathbf{e}_g^T P_{\pi_{\mathbf{a} \circ \pi}} M \mathbf{v} = \sum_{g \in G} \lambda_g \sum_{j \in (\pi_{\mathbf{a} \circ \pi})^{-1}(g)} \mathbf{e}_j^T M \mathbf{v}$$

and, hence, the following chain of implications holds:

$$\begin{aligned} 0 &= \sum_{g \in G} \sum_{j \in (\pi_{\mathbf{a} \circ \pi})^{-1}(g)} \mathbf{e}_j^T M \mathbf{v} = \sum_{j \in F} \mathbf{e}_j^T M \mathbf{v} \quad \Rightarrow \quad \mathbf{e}_j^T M \mathbf{v} = 0 \quad \forall j \in F \\ \Rightarrow \quad \mathbf{e}_j^T M \mathbf{e}_1 &= 0 \quad \forall j \in F \quad \Rightarrow \quad \mathbf{e}_j^T \boldsymbol{\mu} = 0 \quad \forall j \in F \end{aligned}$$

(where the second implication follows from  $\mathbf{e}_1^T \mathbf{v} > 0$ , and the third follows from (c<sub>4</sub>) in Definition 12). Hence,

$$z_i = \beta \mathbf{e}_i^T H P_{\pi_{\mathbf{a} \circ \pi}} \boldsymbol{\mu} = \beta \sum_{g \in G} \lambda_g \mathbf{e}_g^T P_{\pi_{\mathbf{a} \circ \pi}} \boldsymbol{\mu} = \beta \sum_{g \in G} \lambda_g \sum_{j \in (\pi_{\mathbf{a} \circ \pi})^{-1}(g)} \mathbf{e}_j^T \boldsymbol{\mu} = 0,$$

so that  $u_i = w_i + z_i = 0$ , a contradiction. Hence,  $w_i > 0$ . Observe that  $(M/\pi_{\mathbf{a} \circ \pi}, \boldsymbol{\mu}/\pi_{\mathbf{a} \circ \pi}) \in \mathcal{C}_D^{(n)}$  and, hence,  $M/\pi_{\mathbf{a} \circ \pi} \in S$ . By the choice of  $\mathbf{v}$ , this implies that the vector  $P_{\pi_{\mathbf{a} \circ \pi}} M \mathbf{v} = M/\pi_{\mathbf{a} \circ \pi} \mathbf{v}$  is  $H$ -tieless; i.e.,  $H P_{\pi_{\mathbf{a} \circ \pi}} M \mathbf{v}$  is tieless. It follows that the vector  $H P_{\pi_{\mathbf{a} \circ \pi}} (DM)(N' \mathbf{v}) = \frac{1}{2\alpha[\sigma_1^H + 1]D} \mathbf{w}$  is also tieless; being it entrywise integer, and since  $\frac{1}{2\alpha[\sigma_1^H + 1]D} w_i > 0$ , we obtain

$$\left| \frac{1}{2\alpha[\sigma_1^H + 1]D} w_i - \frac{1}{2\alpha[\sigma_1^H + 1]D} w_{i'} \right| \geq 1 \quad \text{that yields} \quad |w_i - w_{i'}| \geq 2\alpha[\sigma_1^H + 1]D.$$

Denote the  $\ell_1$ -norm and the  $\ell_2$ -norm of a vector by  $\|\cdot\|_1$  and  $\|\cdot\|_2$ , respectively. Recall that the largest singular value of a matrix is its spectral operator norm – i.e.,  $\sigma_1^H = \max_{\mathbf{0} \neq \mathbf{x} \in \mathbb{R}^n} \frac{\|H\mathbf{x}\|_2}{\|\mathbf{x}\|_2}$



(see [47]). In particular,  $\|H\mathbf{x}\|_2 \leq \sigma_1^H \|\mathbf{x}\|_2$  for each vector  $\mathbf{x}$  of size  $n$ . Using the Cauchy-Schwarz inequality and the fact that the  $\ell_1$ -norm of a vector is greater than or equal to its  $\ell_2$ -norm, we find

$$\begin{aligned} |z_i - z_{i'}| &= \beta |(\mathbf{e}_i - \mathbf{e}_{i'})^T H P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu}| \leq \beta \|\mathbf{e}_i - \mathbf{e}_{i'}\|_2 \|H P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu}\|_2 \leq \beta \|\mathbf{e}_i - \mathbf{e}_{i'}\|_2 \sigma_1^H \|P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu}\|_2 \\ &\leq \beta \|\mathbf{e}_i - \mathbf{e}_{i'}\|_1 [\sigma_1^H + 1] \|P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu}\|_1 = 2\beta [\sigma_1^H + 1] \mathbf{1}_n^T P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu} \leq 2\beta [\sigma_1^H + 1] \mathbf{1}_n^T P_{\pi_{\mathbf{a}} \circ \pi} |\boldsymbol{\mu}| \\ &= 2\beta [\sigma_1^H + 1] \mathbf{1}_L^T |\boldsymbol{\mu}| \leq 2\beta [\sigma_1^H + 1] D < 2\alpha [\sigma_1^H + 1] D. \end{aligned}$$

We conclude the proof of the claim by noting that

$$|u_i - u_{i'}| = |(w_i - w_{i'}) - (z_{i'} - z_i)| \geq |w_i - w_{i'}| - |z_i - z_{i'}| > 2\alpha [\sigma_1^H + 1] D - 2\alpha [\sigma_1^H + 1] D = 0,$$

which implies  $u_i \neq u_{i'}$ . As a consequence, the vector  $P_{\pi_{\mathbf{a}} \circ \hat{\varphi}} \mathbf{1}_c$  is  $H$ -tieless. We can then apply Lemma 34 to conclude that  $f_{/\pi_{\mathbf{a}} \circ \hat{\varphi}}(1, \dots, n) = f_{/\pi_{\mathbf{a}} \circ \pi \circ \varphi}(1, \dots, n)$ . Hence, by (5),  $f_{/\hat{\varphi}} = f_{/\pi \circ \varphi}$ . Therefore,  $\xi_D((M, \boldsymbol{\mu})_{/\pi}) = \xi_D((M, \boldsymbol{\mu}))_{/\pi}$ , as required. It follows that  $\xi_D$  is a minion homomorphism.

Since the set of polymorphisms of  $(\mathbf{A}, \mathbf{B})$  of arity  $L$  is finite for every  $L \in \mathbb{N}$ , we can apply Proposition 25 to conclude that there exists a minion homomorphism  $\zeta : \mathcal{C} \rightarrow \text{Pol}(\mathbf{A}, \mathbf{B})$ .  $\square$

**Remark 37.** If  $\text{Pol}(\mathbf{A}, \mathbf{B})$  contains  $H$ -symmetric operations of *all* arities – as it happens for the PCSP template  $(\mathbf{A}, \mathbf{B})$  from Example 10, cf. Example 16 – the AIP part of CLAP is not required. Indeed, in that case, we can choose  $f$  in the proof of Theorem 15 to be an  $H$ -symmetric polymorphism of arity  $c = N^2$ , which implies  $\beta = 0$ . Therefore, the affine vector  $\boldsymbol{\mu}$  does not have any role in the definition of  $P_{\varphi}$  in (4), nor in the definition of the minion homomorphism  $\xi_D$ . It follows that, under this stronger hypothesis,  $\text{Pol}(\mathbf{A}, \mathbf{B})$  admits a minion homomorphism from a minion  $\hat{\mathcal{C}}$  whose  $L$ -ary elements are matrices in  $\mathbb{Q}^{L, \aleph_0}$  satisfying the requirements  $(c_1), (c_2), (c_5), (c_6)$  of Definition 12; notice that the projection  $(M, \boldsymbol{\mu}) \mapsto M$  yields a natural minion homomorphism from  $\mathcal{C}$  to  $\hat{\mathcal{C}}$ . The proofs of Theorems 30 and 31 can be straightforwardly modified to show that  $\hat{\mathcal{C}}$  captures the power of the algorithm CBLP – i.e., the simplified version of CLAP that does not run BLP + AIP at the end (cf. the discussion in Section 3).

## Acknowledgements

We would like to thank the anonymous referees of both the conference [32] and this full version of the paper.

## A Existing relaxations for PCSPs

Every CSP can be equivalently expressed as a 0–1 integer program in a standard way.

If the variables are allowed to take values in  $[0, 1]$ , we obtain the so-called basic linear programming relaxation (BLP) [59]. This naturally extends to PCSPs [8], as we describe in Appendix A.1.

If the variables are allowed to take integer values, we obtain the so-called basic affine integer programming relaxation (AIP) [20], studied in detail in [8], as we describe in Appendix A.2.

A combination of the two relaxations, called the BLP + AIP relaxation, was proposed in [22] and its power characterised in [22], as we describe in Appendix A.3.

Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template with signature  $\sigma$  and let  $\mathbf{X}$  be an instance of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ . In all three relaxations described below, we assume without loss of generality that  $\sigma$  contains a unary symbol  $R_u$  such that  $R_u^{\mathbf{X}} = X$ ,  $R_u^{\mathbf{A}} = A$ , and  $R_u^{\mathbf{B}} = B$ . If this is not the case, the signature and the instance can be extended without changing the set of solutions.

### A.1 BLP

The *basic linear programming relaxation* (BLP) of  $\mathbf{X}$ , denoted by  $\text{BLP}(\mathbf{X}, \mathbf{A})$ , is defined as follows.<sup>10</sup> The variables are  $\lambda_{\mathbf{x}, R}(\mathbf{a})$  for every  $R \in \sigma$ ,  $\mathbf{x} \in R^{\mathbf{X}}$ , and  $\mathbf{a} \in R^{\mathbf{A}}$ , and the constraints are given in Figure 1.

$$\begin{aligned}
0 &\leq \lambda_{\mathbf{x}, R}(\mathbf{a}) \leq 1 && \forall R \in \sigma, \forall \mathbf{x} \in R^{\mathbf{X}}, \forall \mathbf{a} \in R^{\mathbf{A}} \\
\sum_{\mathbf{a} \in R^{\mathbf{A}}} \lambda_{\mathbf{x}, R}(\mathbf{a}) &= 1 && \forall R \in \sigma, \forall \mathbf{x} \in R^{\mathbf{X}} \\
\sum_{\mathbf{a} \in R^{\mathbf{A}}, a_i = a} \lambda_{\mathbf{x}, R}(\mathbf{a}) &= \lambda_{x_i, R_u}(a) && \forall R \in \sigma, \forall \mathbf{x} \in R^{\mathbf{X}}, \forall a \in A, \forall i \in [\text{ar}(R)]
\end{aligned}$$

Figure 1: Definition of  $\text{BLP}(\mathbf{X}, \mathbf{A})$ .

We say that  $\text{BLP}(\mathbf{X}, \mathbf{A})$  accepts if the LP in Figure 1 is feasible, and rejects otherwise. By construction, if  $\mathbf{X} \rightarrow \mathbf{A}$  then  $\text{BLP}(\mathbf{X}, \mathbf{A})$  accepts. We say that BLP *solves*  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  if for every instance  $\mathbf{X}$  accepted by  $\text{BLP}(\mathbf{X}, \mathbf{A})$  we have  $\mathbf{X} \rightarrow \mathbf{B}$ .

We denote by  $\mathcal{Q}_{\text{conv}}$  the minion of stochastic vectors on  $\mathbb{Q}$  with the minor operation defined as in Section 4.1; i.e., if  $\mathbf{q} \in \mathcal{Q}_{\text{conv}}^{(L)}$  and  $\pi : [L] \rightarrow [L']$ , then  $\mathbf{q}_{/\pi} = P_{\pi} \mathbf{q}$ , where  $P_{\pi}$  is the  $L' \times L$  matrix whose  $(i, j)$ -th entry is 1 if  $\pi(j) = i$ , and 0 otherwise.

An  $L$ -ary operation  $f : A^L \rightarrow B$  is called *symmetric* if  $f(a_1, \dots, a_L) = f(a_{\pi(1)}, \dots, a_{\pi(L)})$  for every  $a_1, \dots, a_L \in A$  and every permutation  $\pi : [L] \rightarrow [L]$ .

The power of BLP for PCSPs is characterised in the following result.

**Theorem 38** ([8]). *Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template. The following are equivalent:*

- (1) BLP *solves*  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ .
- (2)  $\text{Pol}(\mathbf{A}, \mathbf{B})$  admits a minion homomorphism from  $\mathcal{Q}_{\text{conv}}$ .
- (3)  $\text{Pol}(\mathbf{A}, \mathbf{B})$  contains symmetric operations of all arities.

### A.2 AIP

The *basic affine integer programming relaxation* (AIP) of  $\mathbf{X}$ , denoted by  $\text{AIP}(\mathbf{X}, \mathbf{A})$ , is defined as follows. The variables are  $\tau_{\mathbf{x}, R}(\mathbf{a})$  for every  $R \in \sigma$ ,  $\mathbf{x} \in R^{\mathbf{X}}$ , and  $\mathbf{a} \in R^{\mathbf{A}}$ , and the constraints are given in Figure 2.

<sup>10</sup>The definition does not depend on  $\mathbf{B}$  and is the same as the BLP of an instance  $\mathbf{X}$  of  $\text{CSP}(\mathbf{A})$ ; the same holds for AIP and BLP + AIP.

$$\begin{array}{ll}
\tau_{\mathbf{x},R}(\mathbf{a}) \in \mathbb{Z} & \forall R \in \sigma, \forall \mathbf{x} \in R^{\mathbf{X}}, \forall \mathbf{a} \in R^{\mathbf{A}} \\
\sum_{\mathbf{a} \in R^{\mathbf{A}}} \tau_{\mathbf{x},R}(\mathbf{a}) = 1 & \forall R \in \sigma, \forall \mathbf{x} \in R^{\mathbf{X}} \\
\sum_{\mathbf{a} \in R^{\mathbf{A}}, a_i=a} \tau_{\mathbf{x},R}(\mathbf{a}) = \tau_{x_i, R_u}(a) & \forall R \in \sigma, \forall \mathbf{x} \in R^{\mathbf{X}}, \forall a \in A, \forall i \in [\text{ar}(R)]
\end{array}$$

Figure 2: Definition of  $\text{AIP}(\mathbf{X}, \mathbf{A})$ .

We say that  $\text{AIP}(\mathbf{X}, \mathbf{A})$  accepts if the affine program in Figure 2 is feasible, and rejects otherwise. By construction, if  $\mathbf{X} \rightarrow \mathbf{A}$  then  $\text{AIP}(\mathbf{X}, \mathbf{A})$  accepts. We say that  $\text{AIP}$  solves  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  if for every instance  $\mathbf{X}$  accepted by  $\text{AIP}(\mathbf{X}, \mathbf{A})$  we have  $\mathbf{X} \rightarrow \mathbf{B}$ .

We denote by  $\mathcal{Z}_{\text{aff}}$  the minion of affine vectors on  $\mathbb{Z}$  with the minor operation defined as in Section 4.1; i.e., if  $\mathbf{z} \in \mathcal{Z}_{\text{aff}}^{(L)}$  and  $\pi : [L] \rightarrow [L']$ , then  $\mathbf{z}_{/\pi} = P_{\pi}\mathbf{z}$ , where  $P_{\pi}$  is the  $L' \times L$  matrix whose  $(i, j)$ -th entry is 1 if  $\pi(j) = i$ , and 0 otherwise.

A  $(2L + 1)$ -ary operation  $f : A^{2L+1} \rightarrow B$  is called *alternating* if  $f(a_1, \dots, a_{2L+1}) = f(a_{\pi(1)}, \dots, a_{\pi(2L+1)})$  for every  $a_1, \dots, a_{2L+1} \in A$  and every permutation  $\pi : [2L + 1] \rightarrow [2L + 1]$  that preserves parity, and  $f(a_1, \dots, a_{2L-1}, a, a) = f(a_1, \dots, a_{2L-1}, a', a')$  for every  $a_1, \dots, a_{2L-1}, a, a' \in A$ . Intuitively, an alternating operation is invariant under permutations of its odd and even coordinates and has the property that adjacent coordinates cancel each other out.

The power of AIP for PCSPs is characterised in the following result.

**Theorem 39** ([8]). *Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template. The following are equivalent:*

- (1)  $\text{AIP}$  solves  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ .
- (2)  $\text{Pol}(\mathbf{A}, \mathbf{B})$  admits a minion homomorphism from  $\mathcal{Z}_{\text{aff}}$ .
- (3)  $\text{Pol}(\mathbf{A}, \mathbf{B})$  contains alternating operations of all odd arities.

### A.3 BLP+AIP

The *combined basic LP and affine IP algorithm* (BLP + AIP) is presented in Algorithm 2.

If  $\mathbf{X} \rightarrow \mathbf{A}$  then BLP + AIP accepts  $\mathbf{X}$  [22]. We say that BLP + AIP solves  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  if for every instance  $\mathbf{X}$  accepted by BLP + AIP we have  $\mathbf{X} \rightarrow \mathbf{B}$ .

We denote by  $\mathcal{M}_{\text{BLP}+\text{AIP}}$  the minion whose  $L$ -ary objects are pairs  $(\mathbf{q}, \mathbf{z})$ , where  $\mathbf{q} \in \mathbb{Q}^L$  is a stochastic vector and  $\mathbf{z} \in \mathbb{Z}^L$  is an affine vector, with the property that, for every  $i \in [L]$ ,  $q_i = 0$  implies  $z_i = 0$ . As before, the minor operation is defined as in Section 4.1; i.e., if  $(\mathbf{q}, \mathbf{z}) \in \mathcal{M}_{\text{BLP}+\text{AIP}}^{(L)}$  and  $\pi : [L] \rightarrow [L']$ , then  $(\mathbf{q}, \mathbf{z})_{/\pi} = (P_{\pi}\mathbf{q}, P_{\pi}\mathbf{z})$ , where  $P_{\pi}$  is the  $L' \times L$  matrix whose  $(i, j)$ -th entry is 1 if  $\pi(j) = i$ , and 0 otherwise.

A  $(2L + 1)$ -ary operation  $f : A^{2L+1} \rightarrow B$  is called *2-block symmetric* if  $f(a_1, \dots, a_{2L+1}) = f(a_{\pi(1)}, \dots, a_{\pi(2L+1)})$  for every  $a_1, \dots, a_{2L+1} \in A$  and every permutation  $\pi : [2L + 1] \rightarrow [2L + 1]$  that preserves parity.

The power of BLP + AIP for PCSPs is characterised in the following result.

**Theorem 40** ([22]). *Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template. The following are equivalent:*

---

**Algorithm 2:** The BLP + AIP algorithm

---

**Input:** an instance  $\mathbf{X}$  of  $\text{PCSP}(\mathbf{A}, \mathbf{B})$  of signature  $\sigma$   
**Output:** YES if  $\mathbf{X} \rightarrow \mathbf{A}$  and NO if  $\mathbf{X} \not\rightarrow \mathbf{B}$

- 1 find a relative interior point  $(\lambda_{\mathbf{x}, R}(\mathbf{a}))_{R \in \sigma, \mathbf{x} \in R^{\mathbf{X}}, \mathbf{a} \in R^{\mathbf{A}}}$  of  $\text{BLP}(\mathbf{X}, \mathbf{A})$ ;
- 2 **if** no relative interior point exists **then**
- 3     | return NO;
- 4 **end**
- 5 refine  $\text{AIP}(\mathbf{X}, \mathbf{A})$  by setting  $\tau_{\mathbf{x}, R}(\mathbf{a}) = 0$  if  $\lambda_{\mathbf{x}, R}(\mathbf{a}) = 0$ ;
- 6 **if** the refined  $\text{AIP}(\mathbf{X}, \mathbf{A})$  accepts **then**
- 7     | return YES;
- 8 **end**
- 9 return NO;

---

(1)  $\text{BLP} + \text{AIP}$  solves  $\text{PCSP}(\mathbf{A}, \mathbf{B})$ .

(2)  $\text{Pol}(\mathbf{A}, \mathbf{B})$  admits a minion homomorphism from  $\mathcal{M}_{\text{BLP} + \text{AIP}}$ .

(3)  $\text{Pol}(\mathbf{A}, \mathbf{B})$  contains 2-block-symmetric operations of all odd arities.

## B Proof of Lemma 22

In this section, we shall prove Lemma 22, which we restate below. The proof is based on that of [8, Lemma 4.4], which concerns minions of functions.

**Lemma** (Lemma 22 restated). *Let  $\mathcal{M}$  be a minion and let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template. Then there is a minion homomorphism from  $\mathcal{M}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$  if and only if  $\mathbb{F}_{\mathcal{M}}(\mathbf{A}) \rightarrow \mathbf{B}$ .*

*Proof.* Let  $A = [n]$ , and let  $\sigma$  be the signature of  $\mathbf{A}$  and  $\mathbf{B}$ . Suppose  $\xi : \mathcal{M} \rightarrow \text{Pol}(\mathbf{A}, \mathbf{B})$  is a minion homomorphism, and consider the function

$$\begin{aligned}
 f : \mathcal{M}^{(n)} &\rightarrow B \\
 M &\mapsto \xi(M)(1, \dots, n).
 \end{aligned}$$

For  $R \in \sigma$  of arity  $k$ , consider a tuple  $(M_1, \dots, M_k) \in R^{\mathbb{F}_{\mathcal{M}}(\mathbf{A})}$ . List the elements of  $R^{\mathbf{A}}$  as  $\mathbf{a}^{(1)}, \dots, \mathbf{a}^{(m)}$ . From Definition 21,  $\exists Q \in \mathcal{M}^{(m)}$  such that  $M_i = Q_{/\pi_i}$  for each  $i \in [k]$ , where  $\pi_i : [m] \rightarrow A$  maps  $j$  to the  $i$ -th coordinate of  $\mathbf{a}^{(j)}$ . It follows that, for each  $i \in [k]$ ,

$$f(M_i) = f(Q_{/\pi_i}) = \xi(Q_{/\pi_i})(1, \dots, n) = \xi(Q)_{/\pi_i}(1, \dots, n) = \xi(Q)(\pi_i(1), \dots, \pi_i(m)).$$

Hence,

$$f(M_1, \dots, M_k) = (\xi(Q)(\pi_1(1), \dots, \pi_1(m)), \dots, \xi(Q)(\pi_k(1), \dots, \pi_k(m))) = \xi(Q)(\mathbf{a}^{(1)}, \dots, \mathbf{a}^{(m)}) \in R^{\mathbf{B}}$$

since  $\xi(Q)$  is a polymorphism of  $(\mathbf{A}, \mathbf{B})$ . Therefore,  $f$  is a homomorphism from  $\mathbb{F}_{\mathcal{M}}(\mathbf{A})$  to  $\mathbf{B}$ .

Conversely, let  $f : \mathbb{F}_{\mathcal{M}}(\mathbf{A}) \rightarrow \mathbf{B}$  be a homomorphism, and consider the function  $\xi : \mathcal{M} \rightarrow \text{Pol}(\mathbf{A}, \mathbf{B})$  defined by  $\xi(M)(a_1, \dots, a_L) = f(M_{/\rho})$  for each  $L \in \mathbb{N}$ ,  $M \in \mathcal{M}^{(L)}$ ,  $(a_1, \dots, a_L) \in$

$A^L$ , where

$$\begin{aligned}\rho : [L] &\rightarrow [n] \\ i &\mapsto a_i.\end{aligned}$$

Let us first check that  $\xi$  is well defined – i.e., that  $\xi(M) \in \text{Pol}(\mathbf{A}, \mathbf{B})$ . For  $R \in \sigma$  of arity  $k$ , consider a matrix  $Z \in A^{L,k}$  such that each row of  $Z$  corresponds to a tuple in  $R^{\mathbf{A}}$ . We need to show that  $\xi(M)(Z) \in R^{\mathbf{B}}$ . Consider the maps

$$\begin{aligned}\tau : [L] &\rightarrow R^{\mathbf{A}} & \rho_j : [L] &\rightarrow [n] & \pi_j : R^{\mathbf{A}} &\rightarrow [n] \\ i &\mapsto Z^T \mathbf{e}_i, & i &\mapsto \mathbf{e}_i^T Z \mathbf{e}_j, & \mathbf{a} &\mapsto \mathbf{e}_j^T \mathbf{a},\end{aligned}$$

for  $j \in [k]$ . Observe that  $\rho_j = \pi_j \circ \tau$ , and set  $Q = M_{/\tau} \in \mathcal{M}(|R^{\mathbf{A}}|)$ . We obtain

$$\xi(M)(Z) = f(M_{/\rho_1}, \dots, M_{/\rho_k}) = f(M_{/\pi_1 \circ \tau}, \dots, M_{/\pi_k \circ \tau}) = f(Q_{/\pi_1}, \dots, Q_{/\pi_k}) \in R^{\mathbf{B}}$$

since  $(Q_{/\pi_1}, \dots, Q_{/\pi_k}) \in R^{\mathbb{F}\mathcal{M}(\mathbf{A})}$  and  $f$  is a homomorphism. Finally, we show that  $\xi$  is a minion homomorphism. Clearly,  $\xi$  preserves arities. To check that it preserves minors, let  $M \in \mathcal{M}^{(L)}$  and take a map  $\pi : [L] \rightarrow [L']$ . Given  $(a_1, \dots, a_{L'}) \in A^{L'}$ , consider the maps

$$\begin{aligned}\rho' : [L'] &\rightarrow [n] & \rho'' : [L] &\rightarrow [n] \\ i &\mapsto a_i, & i &\mapsto a_{\pi(i)},\end{aligned}$$

and observe that  $\rho'' = \rho' \circ \pi$ . We obtain

$$\begin{aligned}\xi(M_{/\pi})(a_1, \dots, a_{L'}) &= f((M_{/\pi})_{/\rho'}) = f(M_{/\rho' \circ \pi}) = f(M_{/\rho''}) = \xi(M)(a_{\pi(1)}, \dots, a_{\pi(L)}) \\ &= \xi(M)_{/\pi}(a_1, \dots, a_{L'}),\end{aligned}$$

which yields  $\xi(M_{/\pi}) = \xi(M)_{/\pi}$ , as desired.  $\square$

## C H-block-symmetric polymorphisms

Let  $\mathcal{C} = (\mathcal{C}_1, \dots, \mathcal{C}_\ell)$  be a partition of  $c \in \mathbb{N}$ ; i.e., the sets  $\mathcal{C}_i$  are pairwise disjoint and their union is  $[c]$ . Let  $c_i = |\mathcal{C}_i|$ , so that  $c = \sum_{i \in [\ell]} c_i$ . For each  $i \in [\ell]$ , we consider the unique monotonically increasing function  $\vartheta_i : [c_i] \rightarrow [c]$  such that  $\vartheta_i([c_i]) = \mathcal{C}_i$ . We also consider the function  $\chi_i : \mathcal{C}_i \rightarrow [c_i]$  such that  $\vartheta_i \circ \chi_i$  is the inclusion map of  $\mathcal{C}_i$  in  $[c]$ . Given  $c' \in \mathbb{N}$  and a map  $\pi : [c] \rightarrow [c']$ , we let  $\pi_{(i)} = \pi \circ \vartheta_i$ .

**Definition 41.** Let  $A, B$  be finite sets, and consider a function  $f : A^c \rightarrow B$  for some  $c \in \mathbb{N}$ . Given an  $m \times |A|$  tie matrix  $H$  and a partition  $\mathcal{C} = (\mathcal{C}_1, \dots, \mathcal{C}_\ell)$  of  $c$ , we say that  $f$  is *H-C-block-symmetric* if

$$\begin{aligned}f_{/\pi}(\mathbf{a}) &= f(\mathbf{a}) & \forall \pi : [c] &\rightarrow [c] \text{ permutation such that } \pi(\mathcal{C}_i) = \mathcal{C}_i \ \forall i \in [\ell], \\ & & \forall \mathbf{a} \in A^c &\text{ such that } (P_{\vartheta_i}^T \mathbf{a})^\# \text{ is } H\text{-tieless } \forall i \in [\ell].\end{aligned}$$

We say that  $f$  is *H-block-symmetric* with *width*  $W$  if  $W$  is the largest integer for which there is a partition  $\mathcal{C}$  of  $c$  such that each part of  $\mathcal{C}$  has size at least  $W$  and  $f$  is *H-C-block-symmetric*.<sup>11</sup> Without loss of generality, we consider  $A = [n]$ .

<sup>11</sup>The notion of *H-block-symmetric* operation is the *H*-analogue of that of *block-symmetric* operation in [22] (cf. Theorem 40).

**Lemma 42.** Let  $f : A^c \rightarrow B$  be  $H$ - $\mathcal{C}$ -block-symmetric for some  $m \times n$  tie matrix  $H$  ( $m \in \mathbb{N}$ ) and some partition  $\mathcal{C} = (\mathcal{C}_1, \dots, \mathcal{C}_\ell)$  of  $c$ . Consider two maps  $\pi, \tilde{\pi} : [c] \rightarrow [n]$  such that, for each  $i \in [\ell]$ ,  $P_{\pi(i)} \mathbf{1}_{c_i} = P_{\tilde{\pi}(i)} \mathbf{1}_{c_i}$  and the vector  $P_{\pi(i)} \mathbf{1}_{c_i}$  is  $H$ -tieless. Then

$$f_{/\pi}(1, \dots, n) = f_{/\tilde{\pi}}(1, \dots, n).$$

*Proof.* For  $i \in [\ell]$  and  $a \in [n]$ , we have

$$|\pi_{(i)}^{-1}(a)| = \mathbf{e}_a^T P_{\pi(i)} \mathbf{1}_{c_i} = \mathbf{e}_a^T P_{\tilde{\pi}(i)} \mathbf{1}_{c_i} = |\tilde{\pi}_{(i)}^{-1}(a)|.$$

Hence, we can consider bijections  $\varphi_{i,a} : \pi_{(i)}^{-1}(a) \rightarrow \tilde{\pi}_{(i)}^{-1}(a)$  for each  $i \in [\ell], a \in [n]$ . The union

$$\varphi_i = \bigcup_{a \in [n]} \varphi_{i,a} : [c_i] \rightarrow [c_i]$$

is also a bijection. Define  $\varphi : [c] \rightarrow [c]$  by letting  $\varphi|_{\mathcal{C}_i} = \vartheta_i \circ \varphi_i \circ \chi_i$  for each  $i \in [\ell]$ . Notice that  $\varphi(\mathcal{C}_i) = \mathcal{C}_i$  for each  $i \in [\ell]$ , so  $\varphi$  is a bijection. Take  $j \in [c]$  and suppose that  $j \in \mathcal{C}_i$ . We have

$$(\tilde{\pi} \circ \varphi)(j) = \tilde{\pi}(\varphi(j)) = \tilde{\pi}(\vartheta_i(\varphi_i(\chi_i(j)))) = \tilde{\pi}_{(i)}(\varphi_{i,\pi_{(i)}(\chi_i(j))}(\chi_i(j))) = \pi_{(i)}(\chi_i(j)) = (\pi \circ \vartheta_i \circ \chi_i)(j) = \pi(j)$$

and, hence,  $\tilde{\pi} \circ \varphi = \pi$ . Let  $\tilde{\mathbf{a}} = (\tilde{\pi}(1), \dots, \tilde{\pi}(c))$ . Notice that, for each  $i \in [\ell]$  and  $a \in [n]$ ,

$$\begin{aligned} \mathbf{e}_a^T (P_{\vartheta_i}^T \tilde{\mathbf{a}})^\# &= |\{j \in [c_i] : \mathbf{e}_j^T P_{\vartheta_i}^T \tilde{\mathbf{a}} = a\}| = |\{j \in [c_i] : \mathbf{e}_{\vartheta_i(j)}^T \tilde{\mathbf{a}} = a\}| = |\{j \in [c_i] : \tilde{\pi}(\vartheta_i(j)) = a\}| \\ &= |\{j \in [c_i] : \tilde{\pi}_{(i)}(j) = a\}| = \mathbf{e}_a^T P_{\tilde{\pi}(i)} \mathbf{1}_{c_i} \end{aligned}$$

and, therefore,  $(P_{\vartheta_i}^T \tilde{\mathbf{a}})^\# = P_{\tilde{\pi}(i)} \mathbf{1}_{c_i} = P_{\pi(i)} \mathbf{1}_{c_i}$ , which is  $H$ -tieless. Using that  $f$  is  $H$ - $\mathcal{C}$ -block-symmetric, we find

$$f_{/\tilde{\pi}}(1, \dots, n) = f(\tilde{\mathbf{a}}) = f_{/\varphi}(\tilde{\mathbf{a}}) = (f_{/\varphi})_{/\tilde{\pi}}(1, \dots, n) = f_{/\tilde{\pi} \circ \varphi}(1, \dots, n) = f_{/\pi}(1, \dots, n),$$

as required.  $\square$

**Theorem 43.** Let  $(\mathbf{A}, \mathbf{B})$  be a PCSP template and suppose  $\text{Pol}(\mathbf{A}, \mathbf{B})$  contains  $H$ -block-symmetric operations of arbitrarily large width for some  $m \times |A|$  tie matrix  $H$ ,  $m \in \mathbb{N}$ . Then there exists a minion homomorphism from  $\mathcal{C}$  to  $\text{Pol}(\mathbf{A}, \mathbf{B})$ .

*Proof.* For  $D \in \mathbb{N}$ , consider the subminion  $\mathcal{C}_D$  of  $\mathcal{C}$  described in Section 4.2. Observe that  $S = \{M : (M, \boldsymbol{\mu}) \in \mathcal{C}_D^{(n)}\}$  is a finite set of skeletal matrices. Therefore, we can apply the Tiebreak Lemma 35 to find a stochastic finitely supported vector  $\mathbf{v} \in \mathbb{Q}^{\mathbb{N}_0}$  with  $\mathbf{e}_1^T \mathbf{v} > 0$  such that  $M\mathbf{v}$  is  $H$ -tieless for any  $M \in S$ . Since  $\mathbf{v}$  is finitely supported, we can find  $N' \in \mathbb{N}$  such that  $N'\mathbf{v}$  has integer entries. Let  $\sigma_1^H$  denote the largest singular value of  $H$  – i.e., the square root of the largest eigenvalue of  $H^T H$ . Set  $N = 2\lceil \sigma_1^H + 1 \rceil D^2 N'$ , and let  $f$  be an  $H$ -block-symmetric polymorphism of width  $W \geq N^2$ . Letting  $c$  be the arity of  $f$ , consider a partition  $\mathcal{C} = (\mathcal{C}_1, \dots, \mathcal{C}_\ell)$  of  $c$  such that  $c_i = |\mathcal{C}_i| \geq W$  for each  $i \in [\ell]$  and  $f$  is  $H$ - $\mathcal{C}$ -block-symmetric. Write  $c_i = N\alpha_i + \beta_i$  with  $\alpha_i, \beta_i \in \mathbb{N}_0$ ,  $\beta_i \leq N - 1$ . Note that  $N^2 \leq W \leq c_i = N\alpha_i + \beta_i \leq N\alpha_i + N - 1 < N(\alpha_i + 1)$ , so  $N < \alpha_i + 1$  and, hence,  $\beta_i < \alpha_i$ .

Consider the function

$$\xi_D : \mathcal{C}_D \rightarrow \text{Pol}(\mathbf{A}, \mathbf{B})$$



defined as follows. Given  $L \in \mathbb{N}$  and  $(M, \boldsymbol{\mu}) \in \mathcal{C}_D^{(L)}$ , for each  $i \in [\ell]$  take the map  $\varphi_i : [c_i] \rightarrow [L]$  such that the corresponding  $L \times c_i$  matrix  $P_{\varphi_i}$  is

$$P_{\varphi_i} = \begin{pmatrix} \mathbf{1}_{\mathbf{e}_1^T(\alpha_i N M \mathbf{v} + \beta_i \boldsymbol{\mu})}^T & \mathbf{0}^T & \cdots & \mathbf{0}^T \\ \mathbf{0}^T & \mathbf{1}_{\mathbf{e}_2^T(\alpha_i N M \mathbf{v} + \beta_i \boldsymbol{\mu})}^T & \cdots & \mathbf{0}^T \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0}^T & \mathbf{0}^T & \cdots & \mathbf{1}_{\mathbf{e}_L^T(\alpha_i N M \mathbf{v} + \beta_i \boldsymbol{\mu})}^T \end{pmatrix}. \quad (6)$$

To verify that (6) is well defined, observe first that

$$\sum_{j=1}^L \mathbf{e}_j^T(\alpha_i N M \mathbf{v} + \beta_i \boldsymbol{\mu}) = \mathbf{1}_L^T(\alpha_i N M \mathbf{v} + \beta_i \boldsymbol{\mu}) = \alpha_i N \mathbf{1}_L^T M \mathbf{v} + \beta_i \mathbf{1}_L^T \boldsymbol{\mu} = \alpha_i N \mathbf{1}_{\mathbb{N}_0}^T \mathbf{v} + \beta_i = \alpha_i N + \beta_i = c_i.$$

Moreover, for each  $j \in [L]$ ,  $\mathbf{e}_j^T(\alpha_i N M \mathbf{v} + \beta_i \boldsymbol{\mu}) = \mathbf{e}_j^T(2\alpha_i[\sigma_1^H + 1]D(DM)(N'\mathbf{v}) + \beta_i \boldsymbol{\mu})$  is an integer. If  $\mathbf{e}_j^T(\alpha_i N M \mathbf{v} + \beta_i \boldsymbol{\mu})$  was negative, then  $\mathbf{e}_j^T \boldsymbol{\mu} < 0$ . By the requirement  $(c_4)$  in Definition 12, this would imply that  $\mathbf{e}_j^T M \mathbf{e}_1 > 0$  and, hence,  $0 < \mathbf{e}_j^T M \mathbf{e}_1 \mathbf{e}_1^T \mathbf{v} \leq \mathbf{e}_j^T M \mathbf{v}$ . As a consequence,  $\mathbf{e}_j^T(DM)(N'\mathbf{v}) \geq 1$  so that

$$\mathbf{e}_j^T(\alpha_i N M \mathbf{v} + \beta_i \boldsymbol{\mu}) = 2\alpha_i[\sigma_1^H + 1]D\mathbf{e}_j^T(DM)(N'\mathbf{v}) + \beta_i \mathbf{e}_j^T \boldsymbol{\mu} \geq 2\alpha_i[\sigma_1^H + 1]D + \beta_i \mathbf{e}_j^T \boldsymbol{\mu} \geq \alpha_i D - \beta_i D > 0,$$

which is a contradiction. In conclusion, the numbers  $\mathbf{e}_j^T(\alpha_i N M \mathbf{v} + \beta_i \boldsymbol{\mu})$  are nonnegative integers summing up to  $c_i$ , so (6) is well defined.

Consider the function  $\varphi : [c] \rightarrow [L]$  defined by  $\varphi|_{\mathcal{C}_i} = \varphi_i \circ \chi_i \ \forall i \in [\ell]$ , and let  $\xi_D((M, \boldsymbol{\mu})) := f_{/\varphi}$ . Clearly,  $\xi_D((M, \boldsymbol{\mu})) \in \text{Pol}(\mathbf{A}, \mathbf{B})$ . We claim that the map  $\xi_D$  is a minion homomorphism. It is straightforward to check that  $\xi_D$  preserves arities so, to conclude, we need to show that it also preserves minors. Take  $L' \in \mathbb{N}$  and choose a map  $\pi : [L] \rightarrow [L']$ . Letting  $\tilde{\varphi}_i : [c_i] \rightarrow [L']$  be the map corresponding to the matrix

$$P_{\tilde{\varphi}_i} = \begin{pmatrix} \mathbf{1}_{\mathbf{e}_1^T(\alpha_i N P_\pi M \mathbf{v} + \beta_i P_\pi \boldsymbol{\mu})}^T & \mathbf{0}^T & \cdots & \mathbf{0}^T \\ \mathbf{0}^T & \mathbf{1}_{\mathbf{e}_2^T(\alpha_i N P_\pi M \mathbf{v} + \beta_i P_\pi \boldsymbol{\mu})}^T & \cdots & \mathbf{0}^T \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0}^T & \mathbf{0}^T & \cdots & \mathbf{1}_{\mathbf{e}_{L'}^T(\alpha_i N P_\pi M \mathbf{v} + \beta_i P_\pi \boldsymbol{\mu})}^T \end{pmatrix}$$

for each  $i \in [\ell]$ , and considering  $\tilde{\varphi} : [c] \rightarrow [L']$  such that  $\tilde{\varphi}|_{\mathcal{C}_i} = \tilde{\varphi}_i \circ \chi_i \ \forall i \in [\ell]$ , we see that  $\xi_D((M, \boldsymbol{\mu})/\pi) = f_{/\tilde{\varphi}}$ . Moreover,  $\xi_D((M, \boldsymbol{\mu}))/\pi = (f_{/\varphi})/\pi = f_{/\pi \circ \varphi}$ , where  $\varphi$  is the map defined above. Take  $\mathbf{a} = (a_1, \dots, a_{L'}) \in A^{L'}$ , and consider the map

$$\begin{aligned} \pi_{\mathbf{a}} : [L'] &\rightarrow [n] \\ i &\mapsto a_i. \end{aligned}$$

Observe that

$$\begin{aligned} f_{/\tilde{\varphi}}(\mathbf{a}) &= (f_{/\tilde{\varphi}})_{/\pi_{\mathbf{a}}}(1, \dots, n) = f_{/\pi_{\mathbf{a}} \circ \tilde{\varphi}}(1, \dots, n) & \text{and, similarly,} \\ f_{/\pi \circ \varphi}(\mathbf{a}) &= (f_{/\pi \circ \varphi})_{/\pi_{\mathbf{a}}}(1, \dots, n) = f_{/\pi_{\mathbf{a}} \circ \pi \circ \varphi}(1, \dots, n). \end{aligned} \quad (7)$$

Notice that, for each  $i \in [\ell]$ ,  $\varphi \circ \vartheta_i = \varphi_i$  and  $\tilde{\varphi} \circ \vartheta_i = \tilde{\varphi}_i$ . Hence,

$$\begin{aligned} P_{(\pi_{\mathbf{a}} \circ \tilde{\varphi})_{(i)}} \mathbf{1}_{c_i} &= P_{\pi_{\mathbf{a}} \circ \tilde{\varphi} \circ \vartheta_i} \mathbf{1}_{c_i} = P_{\pi_{\mathbf{a}}} P_{\tilde{\varphi} \circ \vartheta_i} \mathbf{1}_{c_i} = P_{\pi_{\mathbf{a}}} P_{\tilde{\varphi}_i} \mathbf{1}_{c_i} = P_{\pi_{\mathbf{a}}} (\alpha_i N P_{\pi} M \mathbf{v} + \beta_i P_{\pi} \boldsymbol{\mu}) \\ &= P_{\pi_{\mathbf{a}}} P_{\pi} (\alpha_i N M \mathbf{v} + \beta_i \boldsymbol{\mu}) = P_{\pi_{\mathbf{a}}} P_{\pi} P_{\varphi_i} \mathbf{1}_{c_i} = P_{\pi_{\mathbf{a}} \circ \pi \circ \varphi \circ \vartheta_i} \mathbf{1}_{c_i} = P_{(\pi_{\mathbf{a}} \circ \pi \circ \varphi)_{(i)}} \mathbf{1}_{c_i}. \end{aligned}$$

We claim that the vector  $P_{(\pi_{\mathbf{a}} \circ \tilde{\varphi})_{(i)}} \mathbf{1}_{c_i} = P_{\pi_{\mathbf{a}} \circ \tilde{\varphi}_i} \mathbf{1}_{c_i}$  is  $H$ -tieless. Let  $\mathbf{u} = (u_t) = H P_{\pi_{\mathbf{a}} \circ \tilde{\varphi}_i} \mathbf{1}_{c_i}$ ; the claim is equivalent to  $\mathbf{u}$  being tieless. Let  $\mathbf{w} = (w_t) = \alpha_i N H P_{\pi_{\mathbf{a}} \circ \pi} M \mathbf{v}$  and  $\mathbf{z} = (z_t) = \beta_i H P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu}$ , so that  $\mathbf{u} = \mathbf{w} + \mathbf{z}$ . Choose  $t, t' \in [m]$  such that  $t \neq t'$  and  $u_t \neq 0$ . We need to show that  $u_t \neq u_{t'}$ . Suppose  $w_t = 0$ . We can write  $H^T \mathbf{e}_t = \sum_{g \in G} \lambda_g \mathbf{e}_g$  for  $G = \text{supp}(H^T \mathbf{e}_t)$ , where each  $\lambda_g$  is a positive integer (note that  $G \neq \emptyset$  since, otherwise,  $H^T \mathbf{e}_t = \mathbf{0}_n$ , which would imply  $u_t = 0$ ). Let  $F = (\pi_{\mathbf{a}} \circ \pi)^{-1}(G)$ . From  $w_t = 0$ , we obtain

$$0 = \mathbf{e}_t^T H P_{\pi_{\mathbf{a}} \circ \pi} M \mathbf{v} = (H^T \mathbf{e}_t)^T P_{\pi_{\mathbf{a}} \circ \pi} M \mathbf{v} = \sum_{g \in G} \lambda_g \mathbf{e}_g^T P_{\pi_{\mathbf{a}} \circ \pi} M \mathbf{v} = \sum_{g \in G} \lambda_g \sum_{j \in (\pi_{\mathbf{a}} \circ \pi)^{-1}(g)} \mathbf{e}_j^T M \mathbf{v}$$

and, hence, the following chain of implications holds:

$$\begin{aligned} 0 &= \sum_{g \in G} \sum_{j \in (\pi_{\mathbf{a}} \circ \pi)^{-1}(g)} \mathbf{e}_j^T M \mathbf{v} = \sum_{j \in F} \mathbf{e}_j^T M \mathbf{v} \Rightarrow \mathbf{e}_j^T M \mathbf{v} = 0 \quad \forall j \in F \\ \Rightarrow \mathbf{e}_j^T M \mathbf{e}_1 &= 0 \quad \forall j \in F \Rightarrow \mathbf{e}_j^T \boldsymbol{\mu} = 0 \quad \forall j \in F \end{aligned}$$

(where the second implication follows from  $\mathbf{e}_1^T \mathbf{v} > 0$ , and the third follows from  $(c_4)$  in Definition 12). Hence,

$$z_t = \beta_i \mathbf{e}_t^T H P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu} = \beta_i \sum_{g \in G} \lambda_g \mathbf{e}_g^T P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu} = \beta_i \sum_{g \in G} \lambda_g \sum_{j \in (\pi_{\mathbf{a}} \circ \pi)^{-1}(g)} \mathbf{e}_j^T \boldsymbol{\mu} = 0,$$

so that  $u_t = w_t + z_t = 0$ , a contradiction. Hence,  $w_t > 0$ . Observe that  $(M_{/\pi_{\mathbf{a}} \circ \pi}, \boldsymbol{\mu}_{/\pi_{\mathbf{a}} \circ \pi}) \in \mathcal{C}_D^{(n)}$  and, hence,  $M_{/\pi_{\mathbf{a}} \circ \pi} \in S$ . By the choice of  $\mathbf{v}$ , this implies that the vector  $P_{\pi_{\mathbf{a}} \circ \pi} M \mathbf{v} = M_{/\pi_{\mathbf{a}} \circ \pi} \mathbf{v}$  is  $H$ -tieless; i.e.,  $H P_{\pi_{\mathbf{a}} \circ \pi} M \mathbf{v}$  is tieless. It follows that the vector  $H P_{\pi_{\mathbf{a}} \circ \pi} (DM)(N^T \mathbf{v}) = \frac{1}{2\alpha_i \lceil \sigma_1^H + 1 \rceil D} \mathbf{w}$  is also tieless; being it entrywise integer, and since  $\frac{1}{2\alpha_i \lceil \sigma_1^H + 1 \rceil D} w_t > 0$ , we obtain

$$\left| \frac{1}{2\alpha_i \lceil \sigma_1^H + 1 \rceil D} w_t - \frac{1}{2\alpha_i \lceil \sigma_1^H + 1 \rceil D} w_{t'} \right| \geq 1 \quad \text{that yields} \quad |w_t - w_{t'}| \geq 2\alpha_i \lceil \sigma_1^H + 1 \rceil D.$$

Denote the  $\ell_1$ -norm and the  $\ell_2$ -norm of a vector by  $\|\cdot\|_1$  and  $\|\cdot\|_2$ , respectively. Recall that the largest singular value of a matrix is its spectral operator norm – i.e.,  $\sigma_1^H = \max_{\mathbf{x} \neq \mathbf{0}, \mathbf{x} \in \mathbb{R}^n} \frac{\|H\mathbf{x}\|_2}{\|\mathbf{x}\|_2}$  (see [47]). In particular,  $\|H\mathbf{x}\|_2 \leq \sigma_1^H \|\mathbf{x}\|_2$  for each vector  $\mathbf{x}$  of size  $n$ . Using the Cauchy-Schwarz inequality and the fact that the  $\ell_1$ -norm of a vector is greater than or equal to its  $\ell_2$ -norm, we find

$$\begin{aligned} |z_t - z_{t'}| &= \beta_i |(\mathbf{e}_t - \mathbf{e}_{t'})^T H P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu}| \leq \beta_i \|\mathbf{e}_t - \mathbf{e}_{t'}\|_2 \|H P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu}\|_2 \leq \beta_i \|\mathbf{e}_t - \mathbf{e}_{t'}\|_2 \sigma_1^H \|P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu}\|_2 \\ &\leq \beta_i \|\mathbf{e}_t - \mathbf{e}_{t'}\|_1 \lceil \sigma_1^H + 1 \rceil \|P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu}\|_1 = 2\beta_i \lceil \sigma_1^H + 1 \rceil \mathbf{1}_n^T P_{\pi_{\mathbf{a}} \circ \pi} \boldsymbol{\mu} \leq 2\beta_i \lceil \sigma_1^H + 1 \rceil \mathbf{1}_n^T P_{\pi_{\mathbf{a}} \circ \pi} |\boldsymbol{\mu}| \\ &= 2\beta_i \lceil \sigma_1^H + 1 \rceil \mathbf{1}_L^T |\boldsymbol{\mu}| \leq 2\beta_i \lceil \sigma_1^H + 1 \rceil D < 2\alpha_i \lceil \sigma_1^H + 1 \rceil D. \end{aligned}$$

We conclude the proof of the claim by noting that

$$|u_t - u_{t'}| = |(w_t - w_{t'}) - (z_{t'} - z_t)| \geq |w_t - w_{t'}| - |z_t - z_{t'}| > 2\alpha_i \lceil \sigma_1^H + 1 \rceil D - 2\alpha_i \lceil \sigma_1^H + 1 \rceil D = 0,$$

which implies  $u_t \neq u_{t'}$ . As a consequence, the vector  $P_{(\pi_{\mathbf{a}} \circ \tilde{\varphi})(i)} \mathbf{1}_{c_i}$  is  $H$ -tieless. We can then apply Lemma 42 to conclude that  $f_{/\pi_{\mathbf{a}} \circ \tilde{\varphi}}(1, \dots, n) = f_{/\pi_{\mathbf{a}} \circ \pi \circ \varphi}(1, \dots, n)$ . Hence, by (7),  $f_{/\tilde{\varphi}} = f_{/\pi \circ \varphi}$ . Therefore,  $\xi_D((M, \boldsymbol{\mu})_{/\pi}) = \xi_D((M, \boldsymbol{\mu}))_{/\pi}$ , as required. It follows that  $\xi_D$  is a minion homomorphism.

Since the set of polymorphisms of  $(\mathbf{A}, \mathbf{B})$  of arity  $L$  is finite for every  $L \in \mathbb{N}$ , we can apply Proposition 25 to conclude that there exists a minion homomorphism  $\zeta : \mathcal{C} \rightarrow \text{Pol}(\mathbf{A}, \mathbf{B})$ .  $\square$

## References

- [1] Sanjeev Arora, Carsten Lund, Rajeev Motwani, Madhu Sudan, and Mario Szegedy. Proof verification and the hardness of approximation problems. *J. ACM*, 45(3):501–555, 1998. doi:10.1145/278298.278306.
- [2] Sanjeev Arora and Shmuel Safra. Probabilistic checking of proofs: A new characterization of NP. *J. ACM*, 45(1):70–122, 1998. doi:10.1145/273865.273901.
- [3] Albert Atserias and Víctor Dalmau. Promise Constraint Satisfaction and Width. In *Proc. 2022 ACM-SIAM Symposium on Discrete Algorithms (SODA’22)*, pages 1129–1153, 2022. arXiv:2107.05886, doi:10.1137/1.9781611977073.48.
- [4] Per Austrin, Amey Bhangale, and Aditya Potukuchi. Improved inapproximability of rainbow coloring. In *Proc. 2020 ACM-SIAM Symposium on Discrete Algorithms (SODA’20)*, pages 1479–1495, 2020. arXiv:1810.02784, doi:10.1137/1.9781611975994.90.
- [5] Per Austrin, Venkatesan Guruswami, and Johan Håstad.  $(2+\epsilon)$ -Sat is NP-hard. *SIAM J. Comput.*, 46(5):1554–1573, 2017. arXiv:2013/159, doi:10.1137/15M1006507.
- [6] Libor Barto. The collapse of the bounded width hierarchy. *J. Log. Comput.*, 26(3):923–943, 2016. doi:10.1093/logcom/exu070.
- [7] Libor Barto, Diego Battistelli, and Kevin M. Berg. Symmetric Promise Constraint Satisfaction Problems: Beyond the Boolean Case. In *Proc. 38th International Symposium on Theoretical Aspects of Computer Science (STACS’21)*, volume 187 of *LIPIcs*, pages 10:1–10:16. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2021. arXiv:2010.04623, doi:10.4230/LIPIcs.STACS.2021.10.
- [8] Libor Barto, Jakub Bulín, Andrei A. Krokhin, and Jakub Opršal. Algebraic approach to promise constraint satisfaction. *J. ACM*, 68(4):28:1–28:66, 2021. arXiv:1811.00970, doi:10.1145/3457606.
- [9] Libor Barto and Marcin Kozik. Constraint Satisfaction Problems Solvable by Local Consistency Methods. *J. ACM*, 61(1), 2014. Article No. 3. doi:10.1145/2556646.
- [10] Libor Barto and Marcin Kozik. Robustly solvable constraint satisfaction problems. *SIAM J. Comput.*, 45(4):1646–1669, 2016. arXiv:1512.01157, doi:10.1137/130915479.
- [11] Libor Barto, Andrei Krokhin, and Ross Willard. Polymorphisms, and how to use them. In Andrei Krokhin and Stanislav Živný, editors, *The Constraint Satisfaction Problem: Complexity and Approximability*, volume 7 of *Dagstuhl Follow-Ups*, pages 1–44. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl, Germany, 2017. doi:10.4230/DFU.Vol7.15301.1.
- [12] Libor Barto, Jakub Opršal, and Michael Pinsker. The wonderland of reflections. *Isr. J. Math.*, 223(1):363–398, Feb 2018. arXiv:1510.04521, doi:10.1007/s11856-017-1621-9.
- [13] Libor Barto and Michael Pinsker. Topology is irrelevant (in a dichotomy conjecture for infinite domain constraint satisfaction problems). *SIAM J. Comput.*, 49(2):365–393, 2020. doi:10.1137/18M1216213.

- [14] Joel Berman, Pawel Idziak, Petar Marković, Ralph McKenzie, Matthew Valeriote, and Ross Willard. Varieties with few subalgebras of powers. *Trans. Am. Math. Soc.*, 362(3):1445–1473, 2010.
- [15] Christian Bessiere and Romuald Debruyne. Theoretical analysis of singleton arc consistency and its extensions. *Artif. Intell.*, 172(1):29–41, 2008. doi:[10.1016/j.artint.2007.09.001](https://doi.org/10.1016/j.artint.2007.09.001).
- [16] Avrim Blum. New approximation algorithms for graph coloring. *J. ACM*, 41(3):470–516, 1994. doi:[10.1145/176584.176586](https://doi.org/10.1145/176584.176586).
- [17] Manuel Bodirsky, Barnaby Martin, and Antoine Mottet. Discrete temporal constraint satisfaction problems. *J. ACM*, 65(2):9:1–9:41, 2018. arXiv:[1503.08572](https://arxiv.org/abs/1503.08572), doi:[10.1145/3154832](https://doi.org/10.1145/3154832).
- [18] Manuel Bodirsky, Antoine Mottet, Miroslav Olšák, Jakub Opršal, Michael Pinsker, and Ross Willard.  $\omega$ -categorical structures avoiding height 1 identities. *Trans. Am. Math. Soc.*, 374(1):327–350, 2021. arXiv:[2006.12254](https://arxiv.org/abs/2006.12254), doi:[10.1090/tran/8179](https://doi.org/10.1090/tran/8179).
- [19] Joshua Brakensiek and Venkatesan Guruswami. An algorithmic blend of LPs and ring equations for promise CSPs. In *Proc. 30th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA’19)*, pages 436–455, 2019. arXiv:[1807.05194](https://arxiv.org/abs/1807.05194), doi:[10.1137/1.9781611975482.28](https://doi.org/10.1137/1.9781611975482.28).
- [20] Joshua Brakensiek and Venkatesan Guruswami. Promise Constraint Satisfaction: Algebraic Structure and a Symmetric Boolean Dichotomy. *SIAM J. Comput.*, 50(6):1663–1700, 2021. arXiv:[1704.01937](https://arxiv.org/abs/1704.01937), doi:[10.1137/19M128212X](https://doi.org/10.1137/19M128212X).
- [21] Joshua Brakensiek, Venkatesan Guruswami, and Sai Sandeep. Conditional Dichotomy of Boolean Ordered Promise CSPs. In *Proc. 48th International Colloquium on Automata, Languages, and Programming (ICALP’21)*, volume 198 of *LIPIcs*, pages 37:1–37:15. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2021. arXiv:[2102.11854](https://arxiv.org/abs/2102.11854), doi:[10.4230/LIPIcs.ICALP.2021.37](https://doi.org/10.4230/LIPIcs.ICALP.2021.37).
- [22] Joshua Brakensiek, Venkatesan Guruswami, Marcin Wrochna, and Stanislav Živný. The power of the combined basic LP and affine relaxation for promise CSPs. *SIAM J. Comput.*, 49:1232–1248, 2020. arXiv:[1907.04383](https://arxiv.org/abs/1907.04383), doi:[10.1137/20M1312745](https://doi.org/10.1137/20M1312745).
- [23] Alex Brandts, Marcin Wrochna, and Stanislav Živný. The complexity of promise SAT on non-Boolean domains. *ACM Trans. Comput. Theory*, 13(4):26:1–26:20, 2021. arXiv:[1911.09065](https://arxiv.org/abs/1911.09065), doi:[10.1145/3470867](https://doi.org/10.1145/3470867).
- [24] Alex Brandts and Stanislav Živný. Beyond PCSP(1-in-3,NAE). *Information and Computation*, 2022. arXiv:[2104.12800](https://arxiv.org/abs/2104.12800), doi:[10.1016/j.ic.2022.104954](https://doi.org/10.1016/j.ic.2022.104954).
- [25] Andrei Bulatov. Bounded relational width. Unpublished manuscript, 2009. URL: <https://www2.cs.sfu.ca/~abulatov/papers/relwidth.pdf>.
- [26] Andrei Bulatov, Peter Jeavons, and Andrei Krokhin. Classifying the complexity of constraints using finite algebras. *SIAM J. Comput.*, 34(3):720–742, 2005. doi:[10.1137/S0097539700376676](https://doi.org/10.1137/S0097539700376676).
- [27] Andrei A. Bulatov. A dichotomy theorem for constraint satisfaction problems on a 3-element set. *J. ACM*, 53(1):66–120, 2006. doi:[10.1145/1120582.1120584](https://doi.org/10.1145/1120582.1120584).
- [28] Andrei A. Bulatov. A dichotomy theorem for nonuniform CSPs. In *Proc. 58th Annual IEEE Symposium on Foundations of Computer Science (FOCS’17)*, pages 319–330, 2017. arXiv:[1703.03021](https://arxiv.org/abs/1703.03021), doi:[10.1109/FOCS.2017.37](https://doi.org/10.1109/FOCS.2017.37).
- [29] Andrei A. Bulatov and Víctor Dalmau. A Simple Algorithm for Mal’tsev Constraints. *SIAM J. Comput.*, 36(1):16–27, 2006. doi:[10.1137/050628957](https://doi.org/10.1137/050628957).
- [30] Hubie Chen, Víctor Dalmau, and Berit Grußien. Arc consistency and friends. *J. Log. Comput.*, 23(1):87–108, 2013. doi:[10.1093/logcom/exr039](https://doi.org/10.1093/logcom/exr039).

- [31] Lorenzo Ciardo and Stanislav Živný. Hierarchies of minion tests for PCSPs through tensors. In *Proc. 2023 ACM-SIAM Symposium on Discrete Algorithms (SODA'23)*, 2023. To appear. [arXiv:2207.02277](#).
- [32] Lorenzo Ciardo and Stanislav Živný. CLAP: A New Algorithm for Promise CSPs. In *Proc. 2022 ACM-SIAM Symposium on Discrete Algorithms (SODA'22)*, pages 1057–1068, 2022. [arXiv:2107.05018](#), [doi:10.1137/1.9781611977073.46](#).
- [33] David A. Cohen. Tractable decision for a constraint language implies tractable search. *Constraints*, 9(3):219–229, 2004. [doi:10.1023/B:CONS.0000036045.82829.94](#).
- [34] Víctor Dalmau. There are no pure relational width 2 constraint satisfaction problems. *Inf. Process. Lett.*, 109(4):213–218, 2009. [doi:10.1016/j.ipl.2008.10.005](#).
- [35] Víctor Dalmau, Marcin Kozik, Andrei A. Krokhin, Konstantin Makarychev, Yury Makarychev, and Jakub Opršal. Robust algorithms with polynomial loss for near-unanimity CSPs. *SIAM J. Comput.*, 48(6):1763–1795, 2019. [arXiv:1607.04787](#), [doi:10.1137/18M1163932](#).
- [36] Víctor Dalmau and Andrei A. Krokhin. Robust Satisfiability for CSPs: Hardness and Algorithmic Results. *ACM Trans. Comput. Theory*, 5(4):15:1–15:25, 2013. [doi:10.1145/2540090](#).
- [37] Víctor Dalmau and Justin Pearson. Closure functions and width 1 problems. In *Proc. 4th International Conference on Principles and Practice of Constraint Programming (CP'99)*, volume 1713 of *Lecture Notes in Computer Science*, pages 159–173. Springer, 1999. [doi:10.1007/978-3-540-48085-3\\_12](#).
- [38] Romuald Debruyne and Christian Bessière. Some Practicable Filtering Techniques for the Constraint Satisfaction Problem. In *Proc. 15th International Joint Conference on Artificial Intelligence (IJCAI'97)*, pages 412–417. Morgan Kaufmann, 1997.
- [39] Irit Dinur. The PCP theorem by gap amplification. *J. ACM*, 54(3):12, 2007. [doi:10.1145/1236457.1236459](#).
- [40] Irit Dinur, Oded Regev, and Clifford Smyth. The hardness of 3-uniform hypergraph coloring. *Comb.*, 25(5):519–535, September 2005. [doi:10.1007/s00493-005-0032-4](#).
- [41] Tomás Feder and Moshe Y. Vardi. The Computational Structure of Monotone Monadic SNP and Constraint Satisfaction: A Study through Datalog and Group Theory. *SIAM J. Comput.*, 28(1):57–104, 1998. [doi:10.1137/S0097539794266766](#).
- [42] Miron Ficak, Marcin Kozik, Miroslav Olšák, and Szymon Stankiewicz. Dichotomy for Symmetric Boolean PCSPs. In *Proc. 46th International Colloquium on Automata, Languages, and Programming (ICALP'19)*, volume 132, pages 57:1–57:12. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2019. [arXiv:1904.12424](#), [doi:10.4230/LIPIcs.ICALP.2019.57](#).
- [43] M. R. Garey and David S. Johnson. The complexity of near-optimal graph coloring. *J. ACM*, 23(1):43–49, 1976. [doi:10.1145/321921.321926](#).
- [44] Venkatesan Guruswami and Sanjeev Khanna. On the hardness of 4-coloring a 3-colorable graph. *SIAM J. Discret. Math.*, 18(1):30–40, 2004. [doi:10.1137/S0895480100376794](#).
- [45] Venkatesan Guruswami and Sai Sandeep. d-To-1 Hardness of Coloring 3-Colorable Graphs with  $O(1)$  Colors. In *Proc. 47th International Colloquium on Automata, Languages, and Programming (ICALP'20)*, volume 168 of *LIPIcs*, pages 62:1–62:12. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2020. [doi:10.4230/LIPIcs.ICALP.2020.62](#).
- [46] Pavol Hell and Jaroslav Nešetřil. On the complexity of H-coloring. *J. Comb. Theory, Ser. B*, 48(1):92–110, 1990. [doi:10.1016/0095-8956\(90\)90132-J](#).
- [47] Roger A Horn and Charles R Johnson. *Matrix analysis*. Cambridge university press, 2012.

- [48] Paweł M. Idziak, Petar Marković, Ralph McKenzie, Matthew Valeriote, and Ross Willard. Tractability and learnability arising from algebras with few subpowers. *SIAM J. Comput.*, 39(7):3023–3037, 2010. doi:[10.1137/090775646](https://doi.org/10.1137/090775646).
- [49] Peter G. Jeavons. On the Algebraic Structure of Combinatorial Problems. *Theor. Comput. Sci.*, 200(1-2):185–204, 1998. doi:[10.1016/S0304-3975\(97\)00230-2](https://doi.org/10.1016/S0304-3975(97)00230-2).
- [50] Peter G. Jeavons, David A. Cohen, and Marc Gyssens. Closure Properties of Constraints. *J. ACM*, 44(4):527–548, 1997. doi:[10.1145/263867.263489](https://doi.org/10.1145/263867.263489).
- [51] Sanjeev Khanna, Nathan Linial, and Shmuel Safra. On the hardness of approximating the chromatic number. *Comb.*, 20(3):393–415, 2000. doi:[10.1007/s004930070013](https://doi.org/10.1007/s004930070013).
- [52] Subhash Khot. Improved Inapproximability Results for MaxClique, Chromatic Number and Approximate Graph Coloring. In *Proc. 42nd Annual IEEE Symposium on Foundations of Computer Science (FOCS'01)*, pages 600–609. IEEE Computer Society, 2001. doi:[10.1109/SFCS.2001.959936](https://doi.org/10.1109/SFCS.2001.959936).
- [53] Subhash Khot. On the power of unique 2-prover 1-round games. In *Proc. 34th Annual ACM Symposium on Theory of Computing (STOC'02)*, pages 767–775. ACM, 2002. doi:[10.1145/509907.510017](https://doi.org/10.1145/509907.510017).
- [54] Vladimir Kolmogorov, Andrei A. Krokhin, and Michal Rolínek. The complexity of general-valued CSPs. *SIAM J. Comput.*, 46(3):1087–1110, 2017. arXiv:[1502.07327](https://arxiv.org/abs/1502.07327), doi:[10.1137/16M1091836](https://doi.org/10.1137/16M1091836).
- [55] Vladimir Kolmogorov, Johan Thapper, and Stanislav Živný. The power of linear programming for general-valued CSPs. *SIAM J. Comput.*, 44(1):1–36, 2015. arXiv:[1311.4219](https://arxiv.org/abs/1311.4219), doi:[10.1137/130945648](https://doi.org/10.1137/130945648).
- [56] Marcin Kozik. Solving CSPs Using Weak Local Consistency. *SIAM J. Comput.*, 50(4):1263–1286, 2021. arXiv:[1605.00565](https://arxiv.org/abs/1605.00565), doi:[10.1137/18M117577X](https://doi.org/10.1137/18M117577X).
- [57] Marcin Kozik and Joanna Ochremiak. Algebraic properties of valued constraint satisfaction problem. In *Proc. 42nd International Colloquium on Automata, Languages, and Programming (ICALP'15)*, volume 9134 of *Lecture Notes in Computer Science*, pages 846–858. Springer, 2015. arXiv:[1403.0476](https://arxiv.org/abs/1403.0476), doi:[10.1007/978-3-662-47672-7\\_69](https://doi.org/10.1007/978-3-662-47672-7_69).
- [58] Andrei Krokhin and Jakub Opršal. The complexity of 3-colouring  $H$ -colourable graphs. In *Proc. 60th Annual IEEE Symposium on Foundations of Computer Science (FOCS'19)*, pages 1227–1239, 2019. arXiv:[1904.03214](https://arxiv.org/abs/1904.03214), doi:[10.1109/FOCS.2019.00076](https://doi.org/10.1109/FOCS.2019.00076).
- [59] Gábor Kun, Ryan O'Donnell, Suguru Tamaki, Yuichi Yoshida, and Yuan Zhou. Linear programming, width-1 CSPs, and robust satisfaction. In *Proc. 3rd Innovations in Theoretical Computer Science (ITCS'12)*, pages 484–495. ACM, 2012. doi:[10.1145/2090236.2090274](https://doi.org/10.1145/2090236.2090274).
- [60] Gábor Kun and Mario Szegedy. A new line of attack on the dichotomy conjecture. *Eur. J. Comb.*, 52:338–367, 2016. doi:[10.1016/j.ejc.2015.07.011](https://doi.org/10.1016/j.ejc.2015.07.011).
- [61] Alan K. Mackworth. Consistency in networks of relations. *Artif. Intell.*, 8(1):99–118, 1977. doi:[10.1016/0004-3702\(77\)90007-8](https://doi.org/10.1016/0004-3702(77)90007-8).
- [62] Anatolii Malcev. Untersuchungen aus dem gebiete der mathematischen logik. *Journal of Symbolic Logic*, 2(2), 1937.
- [63] Prasad Raghavendra. Optimal algorithms and inapproximability results for every CSP? In *Proc. 40th Annual ACM Symposium on Theory of Computing (STOC'08)*, pages 245–254, 2008. doi:[10.1145/1374376.1374414](https://doi.org/10.1145/1374376.1374414).
- [64] Danny Rorabaugh, Claude Tardif, and David L. Wehlau. Logical compactness and constraint satisfaction problems. *Log. Methods Comput. Sci.*, 13(1), 2017. doi:[10.23638/LMCS-13\(1:1\)2017](https://doi.org/10.23638/LMCS-13(1:1)2017).



- [65] Thomas Schaefer. The complexity of satisfiability problems. In *Proc. 10th Annual ACM Symposium on the Theory of Computing (STOC'78)*, pages 216–226, 1978. [doi:10.1145/800133.804350](#).
- [66] Johan Thapper and Stanislav Živný. The complexity of finite-valued CSPs. *J. ACM*, 63(4):37:1–37:33, 2016. [arXiv:1210.2987](#), [doi:10.1145/2974019](#).
- [67] Johan Thapper and Stanislav Živný. The power of Sherali-Adams relaxations for general-valued CSPs. *SIAM J. Comput.*, 46(4):1241–1279, 2017. [arXiv:1606.02577](#), [doi:10.1137/16M1079245](#).
- [68] Johan Thapper and Stanislav Živný. The limits of SDP relaxations for general-valued CSPs. *ACM Trans. Comput. Theory*, 10(3):12:1–12:22, 2018. [arXiv:1612.01147](#), [doi:10.1145/3201777](#).
- [69] Avi Wigderson. Improving the performance guarantee for approximate graph coloring. *J. ACM*, 30(4):729–735, 1983. [doi:10.1145/2157.2158](#).
- [70] Marcin Wrochna and Stanislav Živný. Improved hardness for  $H$ -colourings of  $G$ -colourable graphs. In *Proc. 2020 ACM-SIAM Symposium on Discrete Algorithms (SODA'20)*, pages 1426–1435, 2020. [arXiv:1907.00872](#), [doi:10.1137/1.9781611975994.86](#).
- [71] Dmitriy Zhuk. A proof of the CSP dichotomy conjecture. *J. ACM*, 67(5):30:1–30:78, 2020. [arXiv:1704.01914](#), [doi:10.1145/3402029](#).