GENERIC INJECTIVITY OF THE X-RAY TRANSFORM

MIHAJLO CEKIĆ AND THIBAULT LEFEUVRE

ABSTRACT. In dimensions ≥ 3 , we prove that the X-ray transform of symmetric tensors of arbitrary degree is generically injective with respect to the metric on closed Anosov manifolds and on manifolds with spherical strictly convex boundary, no conjugate points and a hyperbolic trapped set. Building on earlier work by Guillarmou, Knieper and the second author [GL19, GKL19], this solves locally the marked length spectrum rigidity conjecture in a neighborhood of a generic Anosov metric. This is the first work going beyond the negatively-curved assumption or dimension 2. Our method, initiated in [CL21a] and fully developed in the present paper, is based on a perturbative argument of the 0-eigenvalue of elliptic operators via microlocal analysis which turn the analytic problem of injectivity into an algebraic problem of representation theory. When the manifold is equipped with a Hermitian vector bundle together with a unitary connection, we also show that the twisted X-ray transform of symmetric tensors (with values in that bundle) is generically injective with respect to the connection. This property turns out to be crucial when solving the *holonomy inverse problem*, as studied in a subsequent article [CL21b].

Contents

1.	Introduction	2
2.	Technical preliminaries	10
3.	On spherical harmonics	17
4.	Pseudodifferential nature of perturbed generalized X-ray transforms	23
5.	Generic injectivity with respect to the connection	28
6.	Generic injectivity with respect to the metric	38
References		54

Date: November 7, 2021.

M. CEKIĆ AND T. LEFEUVRE

1. INTRODUCTION

Let M be a smooth closed *n*-dimensional manifold, with $n \ge 2$. Let \mathcal{M} (resp. \mathcal{M}^k , with $k \in \mathbb{Z}_{\ge 0}$) be the cone of smooth (resp. C^k) metrics on M. Recall that a metric $g \in \mathcal{M}$ is said to be *Anosov* if the geodesic flow $(\varphi_t)_{t\in\mathbb{R}}$ on its unit tangent bundle

$$SM := \{(x, v) \in TM \mid |v|_q = 1\}$$

is an Anosov flow (also called *uniformly hyperbolic* in the literature), in the sense that is there exists a continuous flow-invariant splitting of the tangent bundle of SM as:

$$T(SM) = \mathbb{R}X \oplus E_s \oplus E_u,$$

where X is the geodesic vector field, and such that:

$$\begin{aligned} \forall t \ge 0, \forall w \in E_s, \ |d\varphi_t(w)| \le Ce^{-t\lambda}|w|, \\ \forall t \le 0, \forall w \in E_u, \ |d\varphi_t(w)| \le Ce^{-|t|\lambda}|w|, \end{aligned}$$
(1.1)

the constants $C, \lambda > 0$ being uniform and the metric $|\bullet|$ arbitrary. We will denote by $\mathcal{M}_{\text{Anosov}}$ (resp. $\mathcal{M}^k_{\text{Anosov}}$) the space of smooth (resp. C^k) Anosov metrics on M and we will always assume in the following that it is not empty¹.

Historical examples of Anosov metrics were provided by metrics of negative sectional curvature [Ano67] but there are other examples as long as the metric carries "enough" zones of negative curvature, see [Ebe73, DP03]. As shown in [Con10], generic metrics have a non-trivial hyperbolic basic set, i.e. a compact invariant set, not reduced to a single periodic orbit, where (1.1) is satisfied (but this set may not be equal to the whole manifold though). Certain chaotic physical systems can also be described by Anosov Riemannian manifolds which are not globally negatively-curved: for instance, the *Sinaï billiards* which arise as a model in physics for the Lorentz gas (a gas of electrons in a metal) can be naturally approximated by Anosov surfaces but these surfaces have a lot of flat areas (they consist of two copies of a flat tori connected by negatively-curved cylinders which play the role of the obstacles), see [Kou15, Chapter 6] for instance.

1.1. Generic injectivity of the X-ray transform with respect to the metric: closed case. We let C be the set of free homotopy classes of loops on M. If $g \in \mathcal{M}_{\text{Anosov}}$, it is known [Kli74] that for all $c \in C$, there exists a unique g-geodesic $\gamma_g(c) \in c$. We will denote by L_g the marked length spectrum of g, defined as the map:

$$L_g \in \ell^{\infty}(\mathcal{C}), \quad L_g(c) := \ell_g(\gamma_g(c)),$$
(1.2)

where $\ell_g(\gamma)$ denotes the Riemannian length of a curve $\gamma \subset M$ computed with respect to the metric g.

¹Note that $\mathcal{M}_{Anosov}(\mathbb{S}^2) = \emptyset$ (see [MP11, Corollary 9.5] for instance), that is to say not all manifolds can carry Anosov metrics. It is also not known if manifolds carrying Anosov metrics also carry negatively-curved metrics (the converse being obviously true).

The closed curve $\gamma_g(c)$ on M can be lifted to SM to a periodic orbit $(\gamma_g(c), \dot{\gamma}_g(c))$ of $(\varphi_t)_{t \in \mathbb{R}}$, the geodesic flow of g. We then define the X-ray transform as the operator:

$$I^{g}: C^{0}(SM) \to \ell^{\infty}(\mathcal{C}), \ I^{g}f(c) := \frac{1}{L_{g}(c)} \int_{0}^{L_{g}(c)} f(\varphi_{t}(x,v)) \,\mathrm{d}\,t,$$
 (1.3)

where (x, v) is an arbitrary point of the lift of $\gamma_g(c)$. This operator (which can be defined more generally for any Anosov flow) plays an important role in dynamical systems. In restriction to Hölder-continuous functions, its kernel is given by *coboundaries*, namely for $\alpha \in (0, 1]$:

$$\ker I^g|_{C^{\alpha}(SM)} = \{Xu \mid u \in C^{\alpha}(SM), Xu \in C^{\alpha}(SM)\}^2$$

The restriction of this operator to symmetric tensors appears in some rigidity questions in Riemannian geometry, as we shall see. We introduce $\pi_m^* : C^{\infty}(M, \bigotimes_S^m T^*M) \to C^{\infty}(SM)$, the natural pullback of symmetric *m*-tensors, defined by $\pi_m^* f(x, v) := f_x(\bigotimes^m v)$. We then set

$$I_m^g := I^g \circ \pi_m^*. \tag{1.4}$$

Any symmetric tensor $f \in C^{\infty}(M, \otimes_{S}^{m}T^{*}M)$ admits a canonical decomposition f = Dp + h, where D is the symmetrized covariant derivative, $p \in C^{\infty}(M, \otimes_{S}^{m-1}T^{*}M)$, $h \in C^{\infty}(M, \otimes_{S}^{m}T^{*}M)$ and $D^{*}h = 0$, see §2.2.2 for further details. The Dp part is called *potential* whereas h is called solenoidal. Using the fundamental relation $X\pi_{m}^{*} = \pi_{m+1}^{*}D$, we directly see that:

$$\left\{Dp \mid p \in C^{\infty}(M, \bigotimes_{S}^{m-1}T^*M)\right\} \subset \ker I_m^g|_{C^{\infty}(M, \bigotimes_{S}^m T^*M)}.$$

If this equality holds, we say that the X-ray transform of symmetric *m*-tensors is *s*-injective (solenoidally injective, i.e. injective in restriction to solenoidal tensors). This is known to be true:

- for m = 0, 1 on all Anosov manifolds [DS03],
- for all $m \in \mathbb{Z}_{>0}$ on Anosov manifolds with non-positive curvature [GK80a, CS98],
- and for all $m \in \mathbb{Z}_{\geq 0}$ on all Anosov surfaces without any assumptions on the curvature by [Gui17a] (see also [PSU14] for the cases m = 0, 1, 2).

Although the s-injectivity of I_m^g is conjectured on Anosov manifolds of arbitrary dimension, it is still a widely open question. The main theorem of this article is a first step in this direction:

Theorem 1.1. There exists an integer $k_0 \gg 1$ such that the following holds. Let M be a smooth closed manifold of dimension ≥ 3 carrying Anosov metrics. For all $m \in \mathbb{Z}_{\geq 0}^3$, there exists an open dense set $\mathcal{R}_m \subset \mathcal{M}_{Anosov}^{k_0}$ such that for all metrics $g \in \mathcal{R}_m$, the X-ray transform I_m^g is s-injective. In particular, the space of metrics $\mathcal{R} := \bigcap_{m \geq 0} \mathcal{R}_m$ whose X-ray transforms are all s-injective is residual in $\mathcal{M}_{Anosov}^{k_0}$.

Here, open and dense is understood with respect to the C^{k_0} -topology. The integer $k_0 \gg 1$ is chosen large enough so that the framework of microlocal analysis developed below is welldefined (and this makes k_0 roughly proportional to n). One can also obtain a similar statement for smooth metrics: if $g \in \mathcal{M}_{\text{Anosov}}$ has injective X-ray transform I_m^g , then any smooth

²Of course, the geodesic vector field X depends on g. Note that when the context is clear, we try to avoid as much as possible the notation X_g in order not to burden the discussion.

³For m = 0, 1, the s-injectivity is already established [DS03].

perturbation of g (small enough in the C^{k_0} -topology) will also have injective X-ray transform; conversely, if $g \in \mathcal{M}_{\text{Anosov}}$ is not s-injective, then for any $\varepsilon > 0$, there is a smooth metric g'that is ε -close to g in the C^{k_0} -topology and such that its X-ray transform $I_m^{g'}$ is s-injective.

Observe that the sets \mathcal{R}_m and \mathcal{R} are invariant by the action (by pullback of metrics) of the group of diffeomorphisms with regularity $k_0 + 1$ that are isotopic to the identity, which we denote by $\text{Diff}_0^{k_0+1}(M)$ (the pullback of a C^{k_0} -metric by such a diffeomorphism is indeed C^{k_0} -regular).

As we shall see below, the generic s-injectivity of I_m^g is equivalent to the s-injectivity of an elliptic pseudodifferential operator Π_m^g introduced in [Gui17a], called the *generalized X*ray transform, which enjoys very good analytic properties. This operator will also naturally appear below when discussing the *twisted case*, i.e. when including a bundle $\mathcal{E} \to M$ in the discussion, see §1.4. In particular, this reduction to an elliptic Ψ DO will allow us to apply our technique of perturbation of the 0-eigenvalue of elliptic operators, see §1.5 for further details on the strategy of proof.

1.2. Generic injectivity on manifolds with boundary. We now turn to the case of a smooth Riemannian manifold (M, g) with boundary. We define the incoming (resp. outgoing) tail Γ_- (resp. Γ_+) as the set of points $(x, v) \in SM$ such that $\varphi_t(x, v)$ is defined for all $t \ge 0$ (resp. $t \le 0$); the trapped set is defined as $K = \Gamma_- \cap \Gamma_+$. We further assume that (M, g) satisfies the following assumptions:

- Spherical strictly convex boundary (SSCB): the boundary components of the manifold are diffeomorphic to spheres and the boundary is strictly convex in the sense that the second fundamental form is strictly positive;
- No conjugate points (NCP): the metric has no conjugate points;
- Hyperbolic trapped set (HTS): there exists a continuous flow-invariant Anosov decomposition as in (1.1) on the trapped set K.

We will use the short notation *spherical SNH* for such manifolds and also refer to *SNH* manifolds when the boundary components are not necessarily diffeomorphic to spheres (but satisfy all the previous assumptions). Typical and well-studied examples are provided by *simple manifolds* which are diffeomorphic to balls, without conjugate points and no trapped set; (spherical) SNH manifolds are a generalization of these, see [MP11, Gui17b] for further references. It was proved in a recent article [CEG20] that spherical SNH can always be isometrically embedded into closed Anosov manifolds and we shall see below that this has strong consequences.

Given a manifold M with spherical boundary, we let \mathcal{M}_{SNH} (resp. $\mathcal{M}_{\text{SNH}}^{k_0}$) be the set of all smooth (resp. C^{k_0}) metrics satisfying these properties. As in the closed case, this set is invariant by the action of a gauge group $\text{Diff}_0(M)$ which is the set of diffeomorphisms fixing the boundary ∂M .

Given $x \in \partial M$, we let $\nu(x)$ be the outward-pointing normal unit vector to the boundary and

$$\partial_{\pm}SM := \{(x,v) \in SM, \ x \in \partial M, \pm g(v,\nu(x)) \ge 0\}$$

be the incoming (-) and outgoing (+) boundary. The X-ray transform is defined as the operator

$$I^{g}: C^{\infty}(SM) \to C^{\infty}(\partial_{-}SM \setminus \Gamma_{-}), \ I^{g}f(x,v) := \int_{0}^{\ell_{+}(x,v)} f(\varphi_{t}(x,v)) \,\mathrm{d}\, t$$

where $\ell_+(x, v)$ denotes the *exit time* of $(x, v) \in SM$, namely the maximal positive time for which the geodesic flow is defined. As in the closed case, the X-ray transform of symmetric tensors is defined as $I_m^g = I^g \circ \pi_m^*$; a similar decomposition f = Dp + h between potential and solenoidal parts holds by requiring the extra condition $p|_{\partial M} = 0$. It is then easy to check that such potential tensors are in the kernel of I_m^g and it is conjectured that this should be the whole kernel of the X-ray transform. The s-injectivity is known to be true in a lot of cases but not in full generality:

- On simple manifolds: s-injectivity was proved for m = 0, 1 in any dimension [AR97]; further assuming non-positive sectional curvature, it was obtained for all $m \in \mathbb{Z}_{\geq 0}$ in [PS87]; and for all $m \in \mathbb{Z}_{\geq 0}$ on surfaces, without any curvature assumption, it was obtained in [PSU13]; generic s-injectivity was obtained for m = 2 in [SU05, SU08] (by proving s-injectivity for real analytic metrics);
- On SNH manifolds: in dimension $n \ge 3$, s-injectivity was proved on all SNH manifolds for m = 0, 1 and all SNH manifolds with non-positive sectional curvature for $m \ge 2$ in [Gui17b]; it was obtained on all SNH surfaces for all $m \in \mathbb{Z}_{\ge 0}$ in [Lef19a],
- On manifolds admitting a global foliation by convex hypersurfaces: s-injectivity was obtained in any dimension ≥ 3 for all $m \in \mathbb{Z}_{\geq 0}$ in [UV16, SUV18, dHUZ19].

We will then derive the following:

Corollary 1.2 (of Theorem 1.1 and [CEG20]). There exists an integer $k_0 \gg 1$ such that the following holds. Let M be a smooth manifold of dimension ≥ 3 with spherical boundary carrying SNH metrics. For all $m \in \mathbb{Z}_{\geq 0}^4$, there exists an open dense set $\mathcal{R}'_m \subset \mathcal{M}^{k_0}_{\text{SNH}}$ such that for all metrics $g \in \mathcal{R}'_m$, the X-ray transform I^g_m is s-injective. In particular, the space of metrics $\mathcal{R}' := \bigcap_{m \geq 0} \mathcal{R}'_m$ whose X-ray transforms are all s-injective is residual in $\mathcal{M}^{k_0}_{\text{SNH}}$.

Once again, the sets \mathcal{R}_m and \mathcal{R} are invariant by the action of $\operatorname{Diff}_0^{k_0+1}(M)$. We believe that the assumption that the boundary components are diffeomorphic to spheres could be removed (for that, one would need to avoid the use of [CEG20, Theorem 1] and follow directly the proof of Theorem 1.1 in the case of a manifold with boundary).

1.3. Application to rigidity problems. We now detail the consequences of Theorem 1.1 on three problems of rigidity.

1.3.1. The marked length spectrum rigidity conjecture. In the following, an isometry class, denoted by \mathfrak{g} , is defined as an orbit of metrics under the action of $\operatorname{Diff}_0(M)$, namely

$$\mathfrak{g} := \{ \phi^* g \mid \phi \in \mathrm{Diff}_0(M) \}$$
.

If M is closed, we let $\mathbb{M}_{\text{Anosov}} := \mathcal{M}_{\text{Anosov}}/\text{Diff}_0(M)$ (resp. $\mathbb{M}^{k_0}_{\text{Anosov}}$) be the moduli space of smooth (resp. C^{k_0}) Anosov metrics modulo the action of $\text{Diff}_0(M)$ (resp. $\text{Diff}_0^{k_0+1}(M)$). The

⁴For m = 0, 1, the s-injectivity is already established [Gui17b].

M. CEKIĆ AND T. LEFEUVRE

marked length spectrum introduced in (1.2) is invariant by the action of $\text{Diff}_0(M)$ and thus descends as a map

$$L: \mathbb{M}_{\text{Anosov}} \to \ell^{\infty}(\mathcal{C}), \ \mathfrak{g} \mapsto L_{\mathfrak{g}}.$$
 (1.5)

It is believed to parametrize entirely the moduli space of isometry classes.

Conjecture 1.3 (Burns-Katok [BK85]). Let M be a smooth n-dimensional closed manifold such that $\mathcal{M}_{Anosov}(M) \neq \emptyset$. Then the marked length spectrum map L in (1.5) is injective.

Originally, the conjecture was only phrased in the context of negatively-curved manifolds but it is believed to hold in the general Anosov case. Despite some partial results [GK80a, Kat88, CFF92, BCG95, Ham99, CS98, PSU14] and the proof of the conjecture in the twodimensional case for negatively-curved metrics [Cro90, Ota90a], this question is still widely open. Recently, Guillarmou, Knieper and the second author proved that the s-injectivity of $I_2^{g_0}$ implies that the conjecture holds true locally around g_0 (see [GL19] and [GKL19, Theorem 1.2]). In particular, by [CS98], this solves locally the conjecture around an Anosov metric with nonpositive curvature in any dimension (and without any assumptions on the curvature in dimension two by [PSU14, Gui17a]). A straightforward consequence of Theorem 1.1 is therefore the following:

Corollary 1.4 (of Theorem 1.1 and [GL19, GKL19]). There exists $k_0 \gg 1$ such that the following holds. Let M be a smooth n-dimensional closed manifold carrying Anosov metrics. There is an open dense set $\mathbb{R}_2 \subset \mathbb{M}^{k_0}_{\text{Anosov}}$ such that: for all $\mathfrak{g}_0 \in \mathbb{R}_2$, the marked length spectrum map L in (1.5) is locally injective near \mathfrak{g}_0 .

The set \mathbb{R}_2 is equal to $\mathcal{R}_2/\text{Diff}_0^{k_0+1}(M)$, where \mathcal{R}_2 is given by Theorem 1.1 (and this is well-defined since \mathcal{R}_2 is invariant by $\text{Diff}_0^{k_0+1}(M)$). By locally injective, we mean the following: there exists $k_1 \leq k_0$ such that for any $g_0 \in \mathfrak{g}_0$, there exists $\varepsilon_0 > 0$ such that the following holds: if $g_1, g_2 \in \mathcal{M}_{\text{Anosov}}$ are such that there exists $\phi_1, \phi_2 \in \text{Diff}_0^{k_0+1}(M)$ such that $\|\phi_1^*g_1 - g_0\|_{C^{k_1}} + \|\phi_2^*g_2 - g_0\|_{C^{k_1}} < \varepsilon_0$ and $L_{g_1} = L_{g_2}$, then g_1 and g_2 are isometric. Except in dimension two, this is the first result allowing to relax the negative curvature assumption.

1.3.2. Spectral rigidity. Since the celebrated paper of Kac [Kac66] "Can one hear the shape of a drum?", investigating the space of isospectral manifolds (i.e. manifolds with same spectrum for the Laplacian Δ_g on functions, counted with multiplicities) has been an important question in spectral geometry, see [Mil64, GK80a, GK80b, Vig80, Sar90, GWW92] for instance. It is known that there exist pairs of isospectral hyperbolic surfaces that are not isometric [Vig80]. On the other hand, by [GK80a], the s-injectivity of $I_2^{g_0}$ implies that (M, g_0) is spectrally rigid in the following sense: if $(g_s)_{s \in (-1,1)}$ is a smooth family of isospectral metrics, then they are isometric, i.e. there exists $(\phi_s)_{s \in (-1,1)}$ such that $g_0 = \phi_s^* g_s$. As a consequence, we obtain the following:

Corollary 1.5 (of Theorem 1.1). There exists $k_0 \gg 1$ such that the following holds. Let M be a n-dimensional closed manifold carrying Anosov metrics. Then, the open and dense set of isometry classes $\mathbb{R}_2 \subset \mathbb{M}^{k_0}_{\text{Anosov}}$ are spectrally rigid.

Once again, we conjecture that the previous corollary should actually hold for *all* Anosov metrics in any dimension.

1.3.3. Marked boundary distance function. Let (M, g) be an SNH manifold with boundary. For each pair of points $x, y \in \partial M$ and each homotopy class $[\gamma]$ of curves joining x to y, it is well-known that there exists a unique geodesic in that class. We let $d_g(x, y, [\gamma])$ be the length of that unique geodesic (it minimizes the length among all curves in the $[\gamma]$) and call the map d_g the marked boundary distance function. When M is simple (it is diffeomorphic to a ball), there is only a single geodesic joining x to y; we may then drop the $[\gamma]$ and we call d_g the boundary distance function. This function is invariant by the action of the gaugegroup $\text{Diff}_0(M)$ (it descends on the moduli space) and it is conjectured that this is the only obstruction to recovering the metric:

Conjecture 1.6 (Michel [Mic82]). Simple manifolds are boundary distance rigid and, more generally, SNH manifolds are marked boundary distance rigid in the sense that the marked boundary distance function:

$$d: \mathcal{M}_{\rm SNH}/{\rm Diff}_0(M) \ni \mathfrak{g} \mapsto d_{\mathfrak{g}} \tag{1.6}$$

is injective.

This conjecture is known in a certain number of cases but it still open in full generality. We refer to [Muk77, Muk81, MR78, Cro91, Mic82, Gro83, Ota90b, PU05, BI10, SUV18] for further details. Similarly to the closed case, it was shown in [SU05] that the local boundary distance rigidity of a metric g can be derived from the s-injectivity of its X-ray transform I_2^g . This was extended to SNH manifolds (not necessarily spherical) in [Lef20]. As a consequence, we obtain:

Corollary 1.7 (of Corollary 1.2 and [Lef20]). There exists $k_0 \gg 1$ such that the following holds. Let M be a smooth n-dimensional manifold with spherical boundary carrying SNH metrics. There is an open dense set $\mathbb{R}'_2 \subset \mathbb{M}^{k_0}_{\text{SNH}}$ such that: for all $\mathfrak{g}_0 \in \mathbb{R}'_2$, the marked boundary distance function d in (1.6) is locally injective near \mathfrak{g}_0 .

Here $\mathbb{R}'_2 = \mathcal{R}'_2 / \text{Diff}_0^{k_0+1}(M)$, where \mathcal{R}'_2 is given by Corollary 1.2.

1.4. Generic injectivity of the X-ray transform with respect to the connection. We now consider a smooth closed Anosov Riemannian manifold (M, g) and a smooth Hermitian vector bundle $\mathcal{E} \to M$. We let $\mathcal{A}_{\mathcal{E}}$ (resp. $\mathcal{A}_{\mathcal{E}}^k$) be the space of smooth (resp. C^k) unitary connections on the bundle \mathcal{E} . Contrary to the untwisted case (i.e. $\mathcal{E} = \mathbb{C} \times M$), (1.3) might not define a canonical notion of integration of sections along closed geodesics⁵. It is therefore more convenient to define a similar notion via microlocal analysis.

If $\nabla^{\mathcal{E}} \in \mathcal{A}_{\mathcal{E}}$ and $\pi : SM \to M$ denotes the projection, we can consider the pullback bundle $\pi^* \mathcal{E}$ equipped with the pullback connection $\pi^* \nabla^{\mathcal{E}}$ and define the operator $\mathbf{X} := (\pi^* \nabla^{\mathcal{E}})_X$ acting on $C^{\infty}(SM, \pi^* \mathcal{E})$. We then consider the meromorphic extension of the resolvent operators $\mathbf{R}_{\pm}(z) := (\mp \mathbf{X} - z)^{-1} : C^{\infty}(SM, \pi^* \mathcal{E}) \to \mathcal{D}'(SM, \pi^* \mathcal{E})$ to the whole complex plane \mathbb{C} (here \mathcal{D}' denotes the space of distributions), see §2.3 for further details on the Pollicott-Ruelle theory. It is known that there is an open and dense set of connections

⁵Actually, (1.3) defines an interesting notion if the bundle $\pi^* \mathcal{E} \to SM$ is transparent, i.e. the holonomy with respect to the connection $\pi^* \nabla^{\mathcal{E}}$ along closed geodesics is trivial, see [CL21b, Section 7.2]) for a discussion.

M. CEKIĆ AND T. LEFEUVRE

without resonances at 0 (see [CL21a]). When this is the case, we can define the generalized X-ray transform as:

$$\Pi_m^{\nabla^{z}} := \pi_{m*} (\mathbf{R}_+(0) + \mathbf{R}_-(0)) \pi_m^*, \tag{1.7}$$

acting on sections of $\otimes_S^m T^*M \otimes \mathcal{E} \to M$, see §2.3.3 for further details. This operator turns out to be pseudodifferential of order -1 (see [CL21b, Section 7]) and has some very good analytic properties (such as ellipticity), as we shall see.

Symmetric tensors with values in the bundle \mathcal{E} (also called twisted symmetric tensors in the following) also admit a canonical decomposition into a potential part and a solenoidal part, see §2.2.3. The twisted potential tensors are always contained in the kernel of $\Pi_m^{\nabla^{\mathcal{E}}}$ and we say that the operator is s-injective if this is an equality. We will prove the following:

Theorem 1.8. Let (M, g) be a smooth Anosov manifold of dimension ≥ 3 and let $\pi_{\mathcal{E}} : \mathcal{E} \to M$ be a smooth Hermitian vector bundle. There exists $k_0 \gg 1$ such that the following holds. For all $m \in \mathbb{Z}_{\geq 0}$, there exists an open dense set $\mathcal{S}_m \subset \mathcal{A}_{\mathcal{E}}^{k_0}$ of unitary connections with sinjective twisted generalized X-ray transform $\prod_m^{\nabla^{\mathcal{E}}}$. In particular, the space of connections $\mathcal{S} := \cap_{m \geq 0} \mathcal{S}_m$ whose twisted generalized X-ray transforms are all s-injective is residual in $\mathcal{A}_{\mathcal{E}}^{k_0}$.

We also point out here that a similar result holds for the induced connection on the endomorphism bundle, see Theorem 5.10. This plays a crucial role in the study of the *holonomy inverse problem* which consists in reconstructing a connection (up to gauge) from the knowledge of the trace of its holonomy along closed geodesics, see [CL21b] for further details. We believe that a similar result should hold in the boundary case and this is left for future investigation.

As we shall see, any section $f \in C^{\infty}(SM, \pi^* \mathcal{E})$ can be decomposed into spherical harmonics

$$f = \sum_{m \ge 0} f_m, \tag{1.8}$$

where $f_m \in C^{\infty}(SM, \Omega_m \otimes \mathcal{E})$ and $\Omega_m \to M$ is the bundle of spherical harmonics of degree $m \in \mathbb{Z}_{\geq 0}$. In particular, there is a well-defined notion of *degree* for smooth sections of $\pi^* \mathcal{E} \to SM$, which corresponds to the highest degree in the expansion (1.8) (note that this degree might be infinite). Sections of degree 0 are independent of the velocity variable v (also called spherical variable) and thus define sections of the bundle \mathcal{E} over the base manifold M.

The operator $\mathbf{X} := (\pi^* \nabla^{\mathcal{E}})_X$ has the remarkable property that it acts almost diagonally on this decomposition since

$$\mathbf{X}: C^{\infty}(M, \Omega_m \otimes \mathcal{E}) \to C^{\infty}(M, \Omega_{m-1} \otimes \mathcal{E}) \oplus C^{\infty}(M, \Omega_{m+1} \otimes \mathcal{E}),$$

that is it splits into $\mathbf{X} = \mathbf{X}_{-} + \mathbf{X}_{+}$, where \mathbf{X}_{\pm} corresponds to the projection onto the factor $\Omega_{m\pm 1} \otimes \mathcal{E}$. An important class of transport equations appearing in dynamical systems is that of *cohomological equations* which are equations of the form $\mathbf{X}u = f$, where $u, f \in C^{\infty}(SM, \pi^* \mathcal{E})$ (they can also be chosen in the space of distributions) and u and/or f have some prescribed properties. In particular, when both u and f are smooth, understanding how the degree of f controls that of u is an important question. It appears for instance when studying *transparent bundles* (see [Pat09, Pat12, Pat13, MP11, GPSU16, CL21b] for further details),

i.e. vector bundles with trivial holonomy along closed geodesics, or transparent manifolds (see [CL]) which is a particular case of the previous problem with $\mathcal{E} = TM$ (this is a real bundle though) and $\nabla^{\mathcal{E}} := \nabla^{\text{LC}}$, the Levi-Civita connection. We prove the following result:

Corollary 1.9 (of Theorem 1.8). Let (M, g) be a smooth Anosov manifold of dimension ≥ 3 and let $\pi_{\mathcal{E}} : \mathcal{E} \to M$ be a smooth Hermitian vector bundle. There exists $k_0 \gg 1$ and a residual set $\mathcal{S} \subset \mathcal{A}_{\mathcal{E}}^{k_0}$ such that if $\nabla^{\mathcal{E}} \in \mathcal{S}$, the following holds: let $f, u \in C^{\infty}(SM, \pi^* \mathcal{E})$ such that $\mathbf{X}u = f$ and $\deg(f) < \infty$; then $\deg(u) \leq \max(\deg(f) - 1, 0)$.

The set S is the same as in Theorem 1.8.

1.5. Strategy of proof, organization of the paper. The strategy of both Theorems 1.1 and 1.8 is the same, although the metric case (Theorem 1.1) is more involved due to complicated computations. The idea is also reminiscent of our previous work [CL21a], where a notion of operators of uniform divergence type was introduced. Let us discuss the metric case. If the X-ray transform $I_m^{g_0}$ is not s-injective for some $m \in \mathbb{Z}_{\geq 0}$ and $g_0 \in \mathcal{M}_{\text{Anosov}}$ (or $\mathcal{M}_{\text{Anosov}}^{k_0}$) then, equivalently, the generalized X-ray transform operator $\Pi_m^{g_0}$ is not s-injective. This operator is non-negative, pseudodifferential of order -1 and elliptic on ker $D_{g_0}^*$ (see §2.3.2): as a consequence, it has a well-defined spectrum when acting on the Hilbert space

$$\mathcal{H} := L^2(M, \otimes_S^2 T^*M) \cap \ker D^*_{q_0},$$

which lies in $\mathbb{R}_{\geq 0}$ and accumulates to z = 0. The fact that this operator is not s-injective is equivalent to the existence of an eigenvalue at z = 0. The accumulation of the spectrum at 0 (due to the compactness of the operator) is a slight difficulty and we first need to multiply $\Pi_m^{g_0}$ by a certain invertible Laplace-type operator Δ of order k > 1/2 to obtain $P_{g_0} = \pi_{\ker D_{g_0}^*} \Delta \Pi_m^{g_0} \Delta \pi_{\ker D_{g_0}^*}$ which is a pseudodifferential operator of positive order (hence the spectrum accumulates to $+\infty$) with same kernel as $\Pi_m^{g_0}$. The idea is to show that we can produce arbitrarily small perturbations g of the metric g_0 so that P_q has no eigenvalue at 0.

If γ denotes a small circle near 0 in \mathbb{C} (such that the interior of γ only contains the 0 eigenvalue of P_{g_0}) and λ_g is the sum of the eigenvalues of P_g inside γ , then by elementary spectral theory, we know that $C^{k_0} \ni g \mapsto \lambda_g$ is at least C^3 near g_0 when $k_0 \gg 1$ is large enough. Moreover, due to the non-negativity of the operators P_g , we have $d\lambda|_{g=g_0} = 0$. We then compute the second variation and show, using an elementary abstract perturbative Lemma 5.2, that for all $f \in C^{\infty}(M, \otimes_S^2 T^*M)$:

$$d^{2} \lambda|_{g=g_{0}}(f,f) = \sum_{i=1}^{a} \left(\left\langle d^{2} P|_{g=g_{0}}(f,f)u_{i}, u_{i} \right\rangle_{L^{2}} - 2 \left\langle P_{g_{0}}^{-1} d P|_{g=g_{0}}(f)u_{i}, d P|_{g=g_{0}}(f)u_{i} \right\rangle_{L^{2}} \right),$$
(1.9)

where d is the dimension of ker P_{g_0} and $(u_1, ..., u_d)$ is an L^2 -orthonormal basis of ker P_{g_0} . Writing the perturbation of the metric as $g_t = g_0 + tf$, we have $\lambda_{g_t} = t^2 d^2 \lambda_{g=g_0}(f, f) + t^3 \mathcal{O}(\|f\|_{C^{k_0}}^3)$ and it thus suffices to find $f \in C^{\infty}(M, \bigotimes_S^2 T^*M)$ such that $d^2 \lambda_{g=g_0}(f, f) > 0$. This means that one of the 0-eigenvalues was ejected for a small perturbation of g_0 , and iterating this process, one obtains a metric g close to g_0 such that P_q is injective.

For that, we assume that the contrary holds, namely that the second variation is always zero. We then consider the maps in (1.9), $f \mapsto \langle d^2 P |_{g=g_0}(f, f) u_i, u_i \rangle_{L^2}$ and $f \mapsto$

M. CEKIĆ AND T. LEFEUVRE

 $\langle P_{g_0}^{-1} dP|_{g=g_0}(f)u_i, dP|_{g=g_0}(f)u_i \rangle_{L^2}$. We show that these quantities can all be put in the form $\langle Af, f \rangle_{L^2}$, for some pseudodifferential operator A. The important point here is to evaluate the order of A and to compute exactly its principal symbol.

As a consequence, (1.9) can be put in the form $\langle Bf, f \rangle_{L^2} = 0$ for some Ψ DO denoted by $B \in \Psi^*(M, \bigotimes_S^2 T^*M \to \bigotimes_S^2 T^*M)^6$. Taking (real-valued) Gaussian states for the perturbations f, we then obtain by an elementary lemma that for all $(x_0, \xi_0) \in T^*M \setminus \{0\}$ and for all $f \in \bigotimes_S^2 T^*_{x_0} M$:

$$\langle \sigma_B(x_0,\xi_0)f,f\rangle_{\otimes^2_S T^*_{x_0}M} = 0,$$

where $\sigma_B(x_0, \xi_0) \in \operatorname{End}(\otimes_S^2 T_{x_0}^* M)$ denotes the principal symbol of B. In order to conclude, it is therefore sufficient to contradict the previous equality. This problem turns out to be of *purely algebraic nature* and relies on the representation theory of $\operatorname{SO}(n)$ via spherical harmonics, which is treated in the preliminary section §3. We also point out that the operator B is a priori not elliptic (see Remark 5.8), which prevents us from proving that, at least locally, there is only a finite-dimensional submanifold of isometry classes with non-injective X-ray transform.

The main technical ingredients are recalled in \S^2 but we assume that the reader is familiar with the basics of microlocal analysis. The proof of the genericity in the connection case is developed in \S^5 and the metric case is handled in \S^6 .

To conclude, let us mention that the approach initiated in [CL21a] and developed in the present paper to study generic properties of elliptic pseudodifferential operators seems new. It is at least very different from the historical approach of Uhlenbeck [Uhl76] and others.

Acknowledgement: M.C. has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 725967). The authors are grateful to Colin Guillarmou and Gabriel Paternain for their encouragement.

2. Technical preliminaries

2.1. Elementary Riemannian geometry. We refer to [Pat99] for further details on the content of this paragraph. Let (M, g) be a smooth Riemannian manifold. We denote by

$$SM := \{(x, v) \in TM \mid g_x(v, v) = 1\} \subset TM,$$

its unit tangent bundle. Let $(\varphi_t)_{t\in\mathbb{R}}$ be the geodesic flow generated by the vector field X. If π : $SM \to M$ denotes the projection, we define $\mathbb{V} := \ker d\pi$ to be the *vertical subspace*. Recall the definition of the *connection map* $\mathcal{K} : T(SM) \to TM$: consider $(x, v) \in SM, w \in T_{(x,v)}(SM)$ and a curve $(-\varepsilon, \varepsilon) \ni t \mapsto z(t) \in SM$ such that $z(0) = (x, v), \dot{z}(0) = w$; write z(t) = (x(t), v(t)); then $\mathcal{K}_{(x,v)}(w) := \nabla_{\dot{x}(t)}v(t)|_{t=0}$, where ∇ denotes the Levi-Civita connection of (M, g). The Sasaki metric g_{Sas} on SM is defined as follows:

$$g_{\text{Sas}}(w, w') := g(d\pi(w), d\pi(w')) + g(\mathcal{K}(w), \mathcal{K}(w')).$$

⁶If $\mathcal{E}, \mathcal{F} \to M$ are two vector bundles over M, we denote by $\Psi^*(M, \mathcal{E} \to \mathcal{F})$ the standard space of pseudodifferential operators (of all orders) obtained by quantizing symbols in the Kohn-Nirenberg class $\rho = 1, \delta = 0$, see [Shu01] for further details.

Write $\mathbb{H} := \ker \mathcal{K} \cap (\mathbb{R} \cdot X)^{\perp}$ for the *horizontal subspace* and $\mathbb{H}_{tot} := \mathbb{R}X \oplus \mathbb{H}$ for the *total horizontal space*. Then we have the following splitting:

$$T(SM) = \mathbb{R} \cdot X \oplus^{\perp} \mathbb{V} \oplus^{\perp} \mathbb{H}, \qquad (2.1)$$

and we will denote by $\pi_{\mathbb{H}}, \pi_{\mathbb{H}_{tot}}, \pi_{\mathbb{V}}$ the orthogonal projections onto the respective spaces $\mathbb{H}, \mathbb{H}_{tot}, \mathbb{V}$. We denote by ∇^{Sas} the gradient of the Sasaki metric. The splitting (2.1) gives rise to a decomposition of the gradient

$$\nabla^{\mathrm{Sas}}(f) = Xf \cdot X + \nabla_{\mathbb{H}}f + \nabla_{\mathbb{V}}f,$$

where $f \in C^{\infty}(SM)$, and $\nabla_{\mathbb{H}} f \in C^{\infty}(SM, \mathbb{H}), \nabla_{\mathbb{V}} f \in C^{\infty}(SM, \mathbb{V}).$

The geodesic vector field X is a contact vector field with contact 1-form α such that $\alpha(X) = 1$, $i_X d\alpha = 0$ and α has the expression:

$$\alpha_{(x,v)}(\xi) = g_x(\mathbf{d}_{(x,v)} \,\pi(\xi), v), \quad \xi \in T_{(x,v)}SM.$$
(2.2)

We have ker $\alpha = \mathbb{H} \oplus \mathbb{V}$ and $d\alpha$ is non-degenerate on ker α (it is a symplectic form). Moreover $d\alpha|_{\mathbb{H}\times\mathbb{H}} = d\alpha|_{\mathbb{V}\times\mathbb{V}} = 0$. The space ker α is equipped with a canonical almost complex structure J defined in the following way: if $Z \in C^{\infty}(SM, \ker \alpha)$, we write $Z = (Z_{\mathbb{H}}, Z_{\mathbb{V}})$ to denote its horizontal $Z_{\mathbb{H}} = d\pi(Z)$ and vertical $Z_{\mathbb{V}} = \mathcal{K}(Z)$ parts; then $JZ = (-Z_{\mathbb{V}}, Z_{\mathbb{H}})$, see [Pat99, Section 1.3.2]. For such Z, the following relation between the contact form and the Sasaki metric holds (see [Pat99, Proposition 1.24]):

$$\iota_Z d\alpha(\bullet) = -g_{\text{Sas}}(JZ, \bullet). \tag{2.3}$$

We will denote by $\operatorname{div}^{\operatorname{Sas}} := (\nabla^{\operatorname{Sas}})^*$ the divergence operator with respect to the Sasaki metric. When clear from context, we will drop the Sasaki superscript. Then we have:

$$\forall Z \in C^{\infty}(SM, T(SM)), \forall f \in C^{\infty}(SM), \quad \operatorname{div}(fZ) = f \operatorname{div}(Z) - Zf.$$
(2.4)

Equivalently, the divergence operator is defined by $\operatorname{div}(Z)d\operatorname{vol}_{g_{\operatorname{Sas}}} = -\mathcal{L}_Z(d\operatorname{vol}_{g_{\operatorname{Sas}}})$, where \mathcal{L}_Z is the Lie derivative along Z and $d\operatorname{vol}_{g_{\operatorname{Sas}}}$ is the volume form of the Sasaki metric. With our conventions, the following formal adjoint formula holds:

$$\forall Z \in C^{\infty}(SM, T(SM)), \quad Z^* = -Z + \operatorname{div}(Z).$$
(2.5)

The Sasaki volume form satisfies the property that (see [Pat99, Exercise 1.33])

$$d\operatorname{vol}_{g_{\operatorname{Sas}}} = \frac{1}{(n-1)!} \alpha \wedge (d\alpha)^{n-1}.$$
(2.6)

When the metric $g \in \mathcal{M}_{Anosov}$ is Anosov, the following crucial property is known [Kli74]:

$$E_s \cap \mathbb{V} = \{0\}, \ E_u \cap \mathbb{V} = \{0\}.$$
 (2.7)

As we shall see, this property is essential in proving the pseudodifferential nature of certain operators, see §4. This also implies that the manifold has no conjugate points, namely:

$$\forall t \neq 0, \quad \mathbb{V} \cap \mathrm{d}\,\varphi_t(\mathbb{V}) = \{0\}\,. \tag{2.8}$$

2.2. Symmetric tensors. This material is standard but it might be hard to locate a complete reference in the literature. Further details can be found in [DS10, GL21, Lef19b, CL21a].

2.2.1. Symmetric tensors in Euclidean space. Let (E, g_E) be a *n*-dimensional Euclidean space and $(\mathbf{e}_1, ..., \mathbf{e}_n)$ be an orthonormal basis. Let $\mathbf{e}_i^* := g_E(\mathbf{e}_i, \bullet)$ be the covector given by the musical isomorphism. We denote by $\otimes^m E^*$, the space of *m*-tensors and $\otimes^m_S E^*$ the space of symmetric *m*-tensors, namely $f \in \otimes^m_S E^*$ if and only if

$$\forall v_1, ..., v_m \in E, \ \forall \sigma \in \mathfrak{S}_m, \quad f(v_1, ..., v_m) = f(v_{\sigma(1)}, ..., v_{\sigma(m)}).$$

Here \mathfrak{S}_m denotes the permutation group of $\{1, \ldots, m\}$. Given $K = (k_1, \ldots, k_m)$, we write $\mathbf{e}_K^* := \mathbf{e}_{k_1}^* \otimes \ldots \otimes \mathbf{e}_{k_m}^*$. The metric g_E induces a natural inner product on $\otimes^m E^*$ given by:

$$\langle \mathbf{e}_K^*, \mathbf{e}_{K'}^* \rangle_{\otimes^m E^*} := \delta_{k_1 k_1'} \dots \delta_{k_m k_m'}.$$

The symmetrization operator $\mathcal{S}: \otimes^m E^* \to \otimes^m_S E^*$ defined by:

$$\mathcal{S}(\eta_1 \otimes \ldots \otimes \eta_m) := \frac{1}{m!} \sum_{\sigma \in \mathfrak{S}_m} \eta_{\sigma(1)} \otimes \ldots \otimes \eta_{\sigma(m)},$$

is the orthogonal projection onto $\otimes_S^m E^*$. We introduce the trace operator $\mathcal{T} : \otimes_S^m E^* \to \otimes_S^{m-2} E^*$:

$$\mathcal{T}f := \sum_{i=1}^{n} f(\mathbf{e}_i, \mathbf{e}_i, \bullet, ..., \bullet),$$

and this is formally taken to be equal to 0 for m = 0, 1. We say that a symmetric tensor is trace-free if its trace vanishes and denote by $\otimes_S^m E^*|_{0-\mathrm{Tr}}$ this subspace. We let $\mathcal{J} : \otimes_S^m E^* \to \otimes_S^{m+2} E^*$ be defined by $\mathcal{J}(f) := \mathcal{S}(g_E \otimes f)$ which is the adjoint of the trace map \mathcal{T} (with respect to the standard inner product previously defined on symmetric tensors). The operator $\mathcal{T} \circ \mathcal{J}$ is a scalar multiple of the identity on $\otimes_S^m E^*|_{0-\mathrm{Tr}}$. Moreover, the total space of symmetric tensors of degree m breaks up as the orthogonal sum:

$$\otimes_{S}^{m} E^{*} = \bigoplus_{k \ge 0} \mathcal{J}^{2k} \left(\otimes_{S}^{m-2k} E^{*}|_{0-\mathrm{Tr}} \right).$$

We define $\mathbf{P}_m(E)$ to be the set of homogeneous polynomials of degree m; $\mathbf{H}_m(E)$, the subset of harmonic homogeneous polynomials of degree m. There is a natural identification λ_m : $\otimes_S^m E^* \to \mathbf{P}_m(E)$ given by the evaluation map $\lambda_m f(v) := f(v, ..., v)$ (where $v \in E$). Moreover, $\lambda_m : \otimes_S^m E^*|_{0-\mathrm{Tr}} \to \mathbf{H}_m(E)$ is an isomorphism and

$$\lambda_m : \otimes_S^m E^* = \bigoplus_{k \ge 0} \mathcal{J}^{2k} \left(\otimes_S^{m-2k} E^* |_{0-\mathrm{Tr}} \right) \to \bigoplus_{k \ge 0} |v|^{2k} \mathbf{H}_{m-2k}(E),$$
(2.9)

is a graded isomorphism (it maps each summand to each summand isomorphically). We let $\mathbb{S}^{n-1} := \{v \in E \mid g_E(v,v) = 1\}$ be the unit sphere in E and $r : C^{\infty}(E) \to C^{\infty}(\mathbb{S}^{n-1})$ be the operator of restriction. Define $\pi_m^* := r \circ \lambda_m$, π_{m*} its adjoint, and denote by $\Omega_m(E)$ the spherical harmonics of degree m, namely

$$\Omega_m(E) := \ker(\Delta_{\mathbb{S}^{n-1}} + m(m+n-2)),$$

where $\Delta_{\mathbb{S}^{n-1}}$ denotes the induced Laplacian on the sphere, and $\mathbf{S}_m(E) := \bigoplus_{k \ge 0} \Omega_{m-2k}(E)$, where $\Omega_k(E) = \{0\}$ for k < 0. It is well-known that

$$\pi_m^* : \otimes_S^m E^* = \bigoplus_{k \ge 0} \mathcal{J}^{2k} \left(\otimes_S^{m-2k} E^* |_{0-\mathrm{Tr}} \right) \to \bigoplus_{k \ge 0} \Omega_{m-2k}(E) = \mathbf{S}_m(E),$$

is a graded isomorphism.

2.2.2. Symmetric tensors on Riemannian manifolds. We now consider a Riemannian manifold (M,g). Given $f \in C^{\infty}(M, \otimes_{S}^{m}T^{*}M)$, we define its symmetric derivative

$$Df := \mathcal{S} \circ \nabla f \in C^{\infty}(M, \otimes_{S}^{m+1}T^{*}M),$$

where ∇ is the Levi-Civita connection. The operator D is an elliptic differential operator of degree 1 and is of gradient type, i.e. its principal symbol is injective (see [CL21a, Section 3] for instance). When the geodesic flow is ergodic, its kernel is given by ker $(D) = \{0\}$ for m odd, and ker $(D) = \mathbb{R} \cdot S^{\frac{m}{2}}(g)$ for m even. Its adjoint is denoted by $D^*f = -\operatorname{Tr}(\nabla f) \in C^{\infty}(M, \otimes_{S}^{m-1}T^*M)$ and is of divergence type.

Any symmetric tensor $f \in C^{\infty}(M, \bigotimes_{S}^{m} T^{*}M)$ can be uniquely decomposed as

$$f = Dp + h,$$

where $h \in C^{\infty}(M, \bigotimes_{S}^{m}T^{*}M) \cap \ker D^{*}$ is the solenoidal part and $p \in C^{\infty}(M, \bigotimes_{S}^{m-1}T^{*}M)$ (and Dp is the potential part). We denote by $\pi_{\operatorname{ran} D}$ the L^{2} -orthogonal projection onto the first factor and by $\pi_{\ker D^{*}} = \mathbb{1} - \pi_{\operatorname{ran} D}$ the L^{2} -orthogonal projection onto the second factor. Both are pseudodifferential operators of order 0, namely in $\Psi^{0}(M, \bigotimes_{S}^{m}T^{*}M \to \bigotimes_{S}^{m}T^{*}M)$. The latter is given by the expression:

$$\pi_{\ker D^*} = \mathbb{1} - D(D^*D)^{-1}D^*.$$
(2.10)

More precisely, the principal symbol of D is given by $\sigma_D(x,\xi) = ij_{\xi}$ where $j_{\xi} = i\mathcal{S}(\xi \otimes \bullet)$, whereas that of D^* is given by $\sigma_{D^*}(x,\xi) = -ii_{\xi^{\sharp}}$, where i_w is the contraction by w. The space $\otimes_S^m T_x^* M$ breaks up as the orthogonal sum:

$$\otimes^m_S T^*_x M = \operatorname{ran}(j_{\xi}) \oplus^{\perp} \ker i_{\xi^{\sharp}}.$$

The principal symbol of $\pi_{\ker D^*}$ is then given by the orthogonal projection onto the second summand, namely

$$\sigma_{\pi_{\ker D^*}}(x,\xi) = \pi_{\ker \imath_{\xi^\sharp}}.$$
(2.11)

We have the important relation:

$$X\pi_m^* = \pi_{m+1}^* D. (2.12)$$

The spherical harmonics introduced previously in §2.2.1 allow to decompose smooth functions $f \in C^{\infty}(SM)$ as $f = \sum_{m\geq 0} f_m$, where $f_m \in C^{\infty}(M, \Omega_m)$ is the projection onto spherical harmonics of degree m and

$$\Omega_m(x) := \ker \left(\Delta_{\mathbb{V}}(x) + m(m+n-2) \right),$$

where $\Delta_{\mathbb{V}}(x) : C^{\infty}(S_x M) \to C^{\infty}(S_x M)$ denotes the vertical Laplacian acting on functions on $S_x M$ (i.e. the round Laplacian on the sphere). We call *degree* of f (denoted by deg(f)) the highest non-zero spherical harmonic in this expansion (which can take value $+\infty$) and say that f has finite Fourier content if its degree is finite. We will say that a function is *even* (resp. *odd*) if it contains only even (resp. odd) spherical harmonics in its expansion, i.e. $f_{2k+1} = 0$ for all $k \in \mathbb{Z}_{\geq 0}$ (resp. $f_{2k} = 0$ for all $k \in \mathbb{Z}_{\geq 0}$). The operator X acts on spherical harmonics as:

$$X: C^{\infty}(M, \Omega_m) \to C^{\infty}(M, \Omega_{m-1}) \oplus C^{\infty}(M, \Omega_{m+1}),$$

and therefore splits into $X = X_{-} + X_{+}$, where X_{\pm} denotes the projection onto the $\Omega_{m\pm 1}$ factor. The operator X_{+} is of gradient type and thus, for each $m \in \mathbb{N}$, ker $X_{+}|_{C^{\infty}(M,\Omega_{m})}$ is finite-dimensional, and we call elements in this kernel *Conformal Killing Tensors* (CKTs).

Eventually, we will also use another lift of symmetric tensors to the unit tangent bundle via the following map, which we call the *Sasaki lift*:

$$\pi_{m,\text{Sas}}^* : C^{\infty}(M, \otimes_S^m T^*M) \to C^{\infty}(SM, \otimes_S^m T^*(SM)), \pi_{m,\text{Sas}}^*f(\xi_1, ..., \xi_m) := f(\operatorname{d} \pi(\xi_1), ..., \operatorname{d} \pi(\xi_m)).$$
(2.13)

2.2.3. Twisted symmetric tensors. The previous discussion can be generalized in order to include a twist by a vector bundle $\mathcal{E} \to M$, see [CL21a, Section 2.3] for further details. We let $(e_1, ..., e_r)$ be a local orthonormal frame of \mathcal{E} (defined around a fixed point $x_0 \in M$). The smooth sections of the pullback bundle $\pi^* \mathcal{E} \to SM$ can also be decomposed into spherical harmonics, namely $f \in C^{\infty}(SM, \pi^* \mathcal{E})$ can be written as $f = \sum_{m\geq 0} f_m$, where $f_m \in C^{\infty}(M, \Omega_m \otimes \mathcal{E})$ and a similar notion of degree is defined, as well as the evenness/oddness of a section.

If $\nabla^{\mathcal{E}}$ is a unitary connection on \mathcal{E} given in a local patch of coordinates $U \ni x_0$ by $\nabla^{\mathcal{E}} = d + \Gamma$, where Γ is a connection 1-form with values in skew-Hermitian endomorphisms, $(\Gamma(\partial_{x_i}))_{k\ell} = \Gamma^k_{i\ell}$, and we consider a twisted symmetric tensor $f \in C^{\infty}(M, \otimes_S^m T^*M \otimes \mathcal{E})$, which we write locally as $f = \sum_{k=1}^r u_k \otimes e_k$ with $u_k \in C^{\infty}(U, \otimes_S^m T^*U)$, we can define its symmetric derivative

$$D_{\mathcal{E}}\left(\sum_{k=1}^{r} u_k \otimes e_k\right) = \sum_{k=1}^{r} \left(Du_k + \sum_{\ell=1}^{r} \sum_{i=1}^{n} \Gamma_{i\ell}^k \mathcal{S}(u_\ell \otimes dx_i) \right) \otimes e_k.$$
(2.14)

As in §2.2.2, twisted symmetric tensors can be uniquely decomposed as $f = D_{\mathcal{E}}p + h$, where $p \in C^{\infty}(M, \bigotimes_{S}^{m-1}T^{*}M \otimes \mathcal{E})$ and $h \in C^{\infty}(M, \bigotimes_{S}^{m}T^{*}M \otimes \mathcal{E})$ is solenoidal, i.e. in ker $(D_{\mathcal{E}})^{*}$.

The pullback operators extend as maps

$$\pi_m^*: C^{\infty}(M, \otimes_S^m T^*M \otimes \mathcal{E}) \to \oplus_{k \ge 0} C^{\infty}(M, \Omega_{m-2k} \otimes \mathcal{E}).$$

The bundle $\pi^* \mathcal{E} \to SM$ is naturally equipped with the pullback connection $\pi^* \nabla^{\mathcal{E}}$ and we set $\mathbf{X} := (\pi^* \nabla^{\mathcal{E}})_X$ which is a differential operator of order 1 acting on $C^{\infty}(SM, \pi^* \mathcal{E})$. We still have the relation:

$$\mathbf{X}\boldsymbol{\pi}_m^* = \boldsymbol{\pi}_{m+1}^* \boldsymbol{D}_{\mathcal{E}},\tag{2.15}$$

and \mathbf{X} decomposes as:

$$\mathbf{X}: C^{\infty}(M, \Omega_m \otimes \mathcal{E}) \to C^{\infty}(M, \Omega_{m-1} \otimes \mathcal{E}) \oplus C^{\infty}(M, \Omega_{m+1} \otimes \mathcal{E}),$$
(2.16)

that is **X** splits as $\mathbf{X} = \mathbf{X}_{-} + \mathbf{X}_{+}$, where \mathbf{X}_{+} is of gradient type. Elements in ker \mathbf{X}_{+} are called *twisted Conformal Killing Tensors*. Non-existence of twisted CKTs is a generic property of connections as proved in [CL21a].

2.3. **Pollicott-Ruelle theory.** The theory of Pollicott-Ruelle resonances which is briefly recalled below has been widely studied in the literature, see [Liv04, GL06, BL07, FRS08, FS11, FT13, DZ16]. We also refer to [Gui17a, CL21a, Lef19b] for further details on these paragraphs.

2.3.1. Meromorphic extension of the resolvents. Let $\mathcal{E} \to M$ be a Hermitian bundle over M equipped with a unitary connection $\nabla^{\mathcal{E}}$. We consider the pullback bundle $\pi^* \mathcal{E} \to SM$ equipped with the pullback connection $\pi^* \nabla^{\mathcal{E}}$ and set $\mathbf{X} := (\pi^* \nabla^{\mathcal{E}})_X$. Defining the domain

$$\mathcal{D}(\mathbf{X}) := \left\{ u \in L^2(SM, \pi^* \mathcal{E}) \mid \mathbf{X}u \in L^2(SM, \pi^* \mathcal{E}) \right\}^7,$$

the differential operator \mathbf{X} (of order 1) is skew-adjoint as an unbounded operator with dense domain $\mathcal{D}(X)$ and has absolutely continuous spectrum on $i\mathbb{R}$ (with possibly embedded eigenvalues).

We introduce the positive (resp. negative) $\mathbf{R}_+(z)$ (resp. $\mathbf{R}_-(z)$) resolvents, defined for for $\Re(z) > 0$ by:

$$\mathbf{R}_{\pm}(z) := (\mp \mathbf{X} - z)^{-1} = -\int_0^{\pm \infty} e^{\mp tz} e^{-t\mathbf{X}} \,\mathrm{d}\, t.$$

Note that given $z \in SM$ and $f \in \mathcal{E}_{\pi(\varphi_{-t}(z))}$, we have that $e^{-t\mathbf{X}}f(z) \in \mathcal{E}_{\pi(z)}$ is the parallel transport of f along the flowline $(\varphi_s(z))_{s\in[-t,0]}$ with respect to the connection $\pi^*\nabla^{\mathcal{E}}$.

These resolvents initially defined on $\{\Re(z) > 0\}$ can be meromorphically extended to \mathbb{C} by making **X** act on *anisotropic Sobolev spaces*. More precisely, there exists scale of Hilbert spaces \mathcal{H}^s_+ (where s > 0) and a constant c > 0 such that

$$\{\Re(z) > -cs\} \ni z \mapsto \mathbf{R}_{\pm}(z) \in \mathcal{H}^s_+,$$

are meromorphic families of operators with poles of finite rank. These spaces are defined so that $f \in \mathcal{H}^s_+$ (resp. \mathcal{H}^s_-) if and only if f is microlocally in H^s near E^*_s (resp. H^s near E^*_u) and microlocally H^{-s} near E^*_u (resp. H^{-s} near E^*_s). The poles are called the *Pollicott-Ruelle resonances*: they are intrinsic to the operators \mathbf{X} and do not depend on any choices made in the construction of the spaces. Moreover, these operators are holomorphic in $\{\Re(z) > 0\}$ and thus all the resonances are contained in $\{\Re(z) \leq 0\}$.

2.3.2. Generalized X-ray transform. When $\mathcal{E} = \mathbb{C} \times M$, $\mathbf{X} = X$ is nothing but the vector field and we use the notations $R_{\pm}(z)$ for the resolvents. In this case, there is a single resonance on $i\mathbb{R}$ located at z = 0. It is a pole of order 1 and the resolvents have the expansion near z = 0:

$$R_{\pm}(z) = -\frac{\Pi_{\pm}}{z} - R_{\pm,0} - R_{\pm,1}z + \mathcal{O}(z^2),$$

for some operators $R_{\pm,1}, R_{\pm,0} : \mathcal{H}^s_{\pm} \to \mathcal{H}^s_{\pm}$, bounded for any s > 0. Moreover, the spectral projection at 0 (i.e. the residue at z = 0) is

$$\Pi_{+} = \Pi_{-} = \langle \bullet, \mu \rangle \mathbf{1}, \tag{2.17}$$

where μ is the normalized Liouville measure i.e. so that $\langle \mathbf{1}, \mu \rangle = 1$, see [Gui17a] for instance. We record a few useful relations involving Π_{\pm} and $R_{\pm,0}$:

$$XR_{+,0} = R_{+,0}X = \mathbb{1} - \Pi_{+}, \quad XR_{-,0} = R_{-,0}X = -\mathbb{1} + \Pi_{-}, \quad R_{+,0}^* = R_{-,0},$$

$$XR_{+,1} = R_{+,1}X = -R_{+,0}, \quad XR_{-,1} = R_{-,1}X = R_{-,0}.$$

(2.18)

We introduce the operator

$$\mathcal{I} := R_{+,0} + R_{-,0} + \Pi_+ \tag{2.19}$$

⁷The scalar product on L^2 is given by $\langle f, f' \rangle_{L^2} := \int_{SM} h(f(x,v), f'(x,v))_{\mathcal{E}_x} d\mu(x,v)$, where h denotes the Hermitian metric on \mathcal{E} and μ is the Liouville measure on SM.

and define the generalized X-ray transform by:

$$\Pi_m := \pi_m * \mathcal{I} \pi_m^*, \tag{2.20}$$

which is an operator acting on sections of the bundle $\otimes_S^m T^*M \to M$ of symmetric tensors. We say that Π_m is s-injective if Π_m is injective when restricted to ker D^* . The following provides a relation with the X-ray transform I_m :

Lemma 2.1. Let $u \in C^{\infty}(M, \bigotimes_{S}^{m} T^{*}M)$. Then $\Pi_{m}u = 0$ if and only if there exists $v \in C^{\infty}(SM)$ such that $\pi_{m}^{*}u = X v$. Moreover, I_{m} is s-injective if and only if Π_{m} is s-injective.

Proof. The first part follows from [Gui17a, Theorem 1]; the second one is then the consequence of the smooth Livšic theorem (see eg. [Lef19b, Lemma 2.5.4]). \Box

By the preceding Lemma, the study of I_m is reduced to the study of Π_m and as we shall see in §4, Π_m enjoys very good analytic properties. In other words, in order to prove our main Theorem 1.1, it suffices to show that Π_m^g is generically s-injective with respect to the metric.

2.3.3. Twisted generalized X-ray transform. We now go back to the case of a Hermitian vector bundle $\mathcal{E} \to M$. For the sake of simplicity, we assume that ker $\mathbf{X} = \{0\}$ but the discussion could be generalized, see the footnote at the beginning of §5. We introduce:

$$\mathcal{I}_{\nabla}\varepsilon := \mathbf{R}_{+} + \mathbf{R}_{-},$$

where $\mathbf{R}_+ := -\mathbf{R}_+(z=0)$ and $\mathbf{R}_- := -\mathbf{R}_-(z=0)$, and define the twisted generalized X-ray transform by:

$$\Pi_m^{\nabla^{\mathcal{E}}} := \pi_{m*} \mathcal{I}_{\nabla^{\mathcal{E}}} \pi_m^* \tag{2.21}$$

Similarly to the first part of Lemma 2.1, the following was shown in [CL21a, Lemma 5.1]:

Lemma 2.2. Let $u \in C^{\infty}(M, \bigotimes_{S}^{m} T^{*}M \otimes \mathcal{E})$. Then $\Pi_{m}^{\nabla^{\mathcal{E}}} u = 0$ if and only if there exists $v \in C^{\infty}(SM, \pi^{*}\mathcal{E})$ such that $\pi_{m}^{*} u = \mathbf{X} v$.

2.4. Properties of the resolvent under perturbations. The generalized X-ray transform operators Π_m^g (we now add the index g to insist on the metric-dependence) and $\Pi_m^{\nabla^2}$ introduced in the previous paragraphs depend on a choice of metric g and/or connection $\nabla^{\mathcal{E}}$. In the following, we will consider perturbations of these operators with respect to these geometric data. For $\mathcal{E}, \mathcal{F} \to M$, two smooth Hermitian vector bundles over M, and $s \in \mathbb{R}$, the spaces of pseudodifferential operators $\Psi^s(M, \mathcal{E} \to \mathcal{F})$ are Fréchet spaces (see [GKL19, Section 2.1] for instance) where the seminorms are defined thanks to local coordinates by taking the seminorms of the full local symbol in the charts. Let us also mention that it is also possible to consider pseudodifferential operators obtained by quantizing symbols with limited regularity (see [Tay91]): actually, all the standard arguments of microlocal analysis (such as boundedness on Sobolev spaces for instance) involve only a finite number of derivatives of the full symbol, and this number depends linearly on the dimension. As a consequence, for $k \gg n$ (where n is the dimension of M), we can consider the space $\Psi_{(k)}^s(M, \mathcal{E} \to \mathcal{F})$ of pseudodifferential operators obtained by quantizing C^k -symbols (satisfying the usual symbolic rules of derivation). The remainder terms in this calculus are then operators with $C^{p(k)}$ -regular kernels (instead of smooth), where $p(k) \in \mathbb{Z}_{\geq 0}$.

Lemma 2.3. The following maps are smooth:

$$\mathcal{M}_{\text{Anosov}} \ni g \mapsto \Pi_m^g \in \Psi^{-1}(M, \otimes_S^m T^*M \to \otimes_S^m T^*M),$$
$$C^{\infty}(M, T^*M \otimes \text{End}_{\text{sk}}(\mathcal{E})) \ni \nabla^{\mathcal{E}} \mapsto \Pi_m^{\nabla^{\mathcal{E}}} \in \Psi^{-1}(M, \otimes_S^m T^*M \otimes \mathcal{E} \to \otimes_S^m T^*M \otimes \mathcal{E}).$$

Equivalently, for every $k \in \mathbb{Z}_{\geq 0}$, there are $k', k'' \gg \max(n, k)$ such that the following maps are C^k :

$$\mathcal{M}^{k'}_{\text{Anosov}} \ni g \mapsto \Pi^g_m \in \Psi^{-1}_{(k'')}(M, \otimes^m_S T^*M \to \otimes^m_S T^*M),$$
$$C^{k'}(M, T^*M \otimes \text{End}_{\text{sk}}(\mathcal{E})) \ni \nabla^{\mathcal{E}} \mapsto \Pi^{\nabla^{\mathcal{E}}}_m \in \Psi^{-1}_{(k'')}(M, \otimes^m_S T^*M \otimes \mathcal{E} \to \otimes^m_S T^*M \otimes \mathcal{E}).$$

Lemma 2.3 will be used in §5 and §6 in order to perturb the generalized X-ray transforms with respect to the connection/metric. For k = 0, Lemma 2.3 is precisely the content of [GKL19, Proposition 4.1]. Inspecting the proof, one can see that it also works for higher order derivatives. The heart of the proof is based on understanding the differentiability of the resolvent map

$$C^{\infty}(SM, T(SM)) \times \mathbb{C} \ni (Y, z) \mapsto (\mp Y - z)^{-1} \in \mathcal{L}(\mathcal{H}^s_{\pm}),$$

where \mathcal{H}^s_{\pm} is the scale of anisotropic Sobolev spaces (which can be made independent of the vector field by [Bon20]), and Y is a vector field close to X. This perturbation theory is now standard and we refer to [Bon20, DGRS20, CDKP20] for further details.

3. On spherical harmonics

We record here some facts about spherical harmonics. We keep the notation (E, g_E) for a Euclidean vector space of dimension n.

3.1. The restriction operator. In the following, we will need to understand how the degree of a function is changed when restricting to a hypersphere. For $\xi \in E^* \setminus \{0\}$, define $\vec{n}(\xi) := \frac{\xi^{\sharp}}{|\xi|}$. If \mathbb{S}^{n-1} denotes the unit sphere in E, we introduce $\mathbb{S}_{\xi}^{n-2} := \{v \in \mathbb{S}^{n-1} \mid \langle \xi, v \rangle = 0\}$. Any vector $v \in \mathbb{S}^{n-1} \setminus \{\pm \vec{n}(\xi)\}$ can be uniquely decomposed as $v = \cos(\varphi)\vec{n}(\xi) + \sin(\varphi)u$, where $\varphi \in (0,\pi), u \in \mathbb{S}_{\xi}^{n-2}$ (the diffeomorphism is singular at the extremal points $\varphi = 0$ and $\varphi = \pi$ but since they form a set of measure 0, this is harmless in what follows). The round measure d S(v) on \mathbb{S}^{n-1} is then given in these new coordinates by

$$d S(v) = \sin^{n-2}(\varphi) d \varphi d S_{\xi}(u), \qquad (3.1)$$

where $d S_{\xi}(u)$ denotes the canonical round measure on the (n-2)-dimensional sphere \mathbb{S}_{ξ}^{n-2} .

3.1.1. Standard restriction. We start with the following:

Lemma 3.1. Assume n > 2. Let $w \in C^{\infty}(\mathbb{S}^{n-1})$. Then w has degree $\leq m$ if and only if the restriction $w|_{\mathbb{S}^{n-2}_{\xi}}$ to any hypersphere \mathbb{S}^{n-2}_{ξ} has degree $\leq m$.

Proof. We start with the easy direction. If $w \in \Omega_m(\mathbb{S}^{n-1})$ is a spherical harmonic of degree m, then:

$$\|\mathbf{w}\|_{\mathbb{S}^{n-2}_{\xi}} \in \bigoplus_{k \ge 0} \Omega_{m-2k}(\mathbb{S}^{n-2}_{\xi}).$$

This fact follows from the following observation: if $\mathbf{w} \in \Omega_m(\mathbb{S}^{n-1})$, then w is the restriction of a harmonic homogeneous polynomial $P \in \mathbf{H}_m(E)$ defined on E. Let $H = \ker \xi$, then $P|_H$ is still a homogeneous polynomial of degree m (but it may not be harmonic, though) and thus its restriction to \mathbb{S}^{n-2}_{ξ} is a sum of spherical harmonics of degree $\leq m$ (and with same parity as m).

We now show the converse. The case m = 0 is obvious so we can always assume that $m \ge 1$. Note first that we may split w into odd and even terms, $w = w_{\text{odd}} + w_{\text{even}}$. We have $(w \mid_{\mathbb{S}_{\xi}^{n-2}})_{\text{odd}} = w_{\text{odd}} \mid_{\mathbb{S}_{\xi}^{n-2}}$ and $(w \mid_{\mathbb{S}_{\xi}^{n-2}})_{\text{even}} = w_{\text{even}} \mid_{\mathbb{S}_{\xi}^{n-2}}$, and so for every ξ , both $w_{\text{even}} \mid_{\mathbb{S}_{\xi}^{n-2}}$ and $w_{\text{odd}} \mid_{\mathbb{S}_{\xi}^{n-2}}$ are of degree $\le m$. Thus, we may assume w is either pure odd or pure even, and that moreover this is the parity of m (if m and w have distinct parities, then the hypothesis of the Lemma is true for m-1).

The conclusion is now implied by the following claim : let $w \in C^{\infty}(E \setminus \{0\})$ be an *m*-homogeneous function (since m > 0 it is at least continuous at x = 0) such that the restriction $w|_H$ to any hyperplane $H \subset E$ is a homogeneous polynomial of degree m. Then w is a homogeneous polynomial of degree m.

First of all, we start by proving that w is smooth at x = 0. Let $(\mathbf{e}_1, ..., \mathbf{e}_n)$ be an orthonormal basis of E and write $x = \sum_{i=1}^n x_i \mathbf{e}_i$. We claim that $\partial_{x_i}^{m+1} \mathbf{w} \equiv 0$ on E. Indeed, fix $i \in \{1, ..., n\}$, fix $x \in E$ and consider a hyperplane H containing both x and \mathbf{e}_i (which forces the condition n > 2). Then $f|_H$ is a polynomial of degree m. In particular, it is smooth and satisfies:

$$\partial_{x_i}^{m+1} \mathbf{w}(x) = \partial_t^{m+1} \mathbf{w}(x + t\mathbf{e}_i)|_{t=0} = 0,$$

since it is polynomial. Actually, $\partial_{x_i}^k w \equiv 0$ as long as $k \ge m+1$. In particular, P w = 0 on E, where $P = \sum_{i=1}^n \partial_{x_i}^{2m}$. As P is elliptic and w continuous, this gives that w is smooth on E.

We now write by Taylor's theorem:

$$\mathbf{w}(x) = \sum_{|\alpha| \le m} \frac{1}{\alpha!} x^{\alpha} \cdot \partial_x^{\alpha} \mathbf{w}(0) + R(x),$$

where $R(x) = \mathcal{O}(|x|^{m+1})$ as $|x| \to 0$ and define $S(x) := \sum_{|\alpha| \le m} \frac{1}{\alpha!} x^{\alpha} \cdot \partial_x^{\alpha} w(0)$. Taking any hyperplane H, we obtain:

$$w|_{H}(x) - S|_{H}(x) = \mathcal{O}(|x|^{m+1}).$$

The left-hand side is a polynomial of degree $\leq m$ so it implies that it is equal to 0. This gives that $R|_H \equiv 0$. Since this holds for every hyperplane H, this implies $R \equiv 0$ and w is a polynomial of degree $\leq m$. Using *m*-homogeneity of w, it is homogeneous of degree m.

Remark 3.2. We observe that the proof actually gives a stronger statement which is: assume $w \in C^{\infty}(E \setminus \{0\})$ is m-homogeneous and a polynomial of degree m in restriction to any plane (and not hyperplane), then it is a polynomial of degree m on E.

3.1.2. Differentiated restriction. The following lemma will be used for the generic s-injectivity with respect to the metric. We will denote by ∇ the gradient with respect to the spherical metric on \mathbb{S}^{n-1} .

Lemma 3.3. Assume $n \ge 3$ and let $m \in \mathbb{Z}_{\ge 0}$ and $w \in C^{\infty}(\mathbb{S}^{n-1})$ such that $\deg(w) \ge m+1$. Then, there exists $\xi \in E^* \setminus \{0\}$ such that $\langle \xi, \nabla w(\bullet) \rangle|_{\mathbb{S}^{n-2}_{\xi}}$ has degree $\ge m$ (seen as a function on \mathbb{S}^{n-2}_{ξ}).

This Lemma will be applied later in each fibre $E = T_x M$ and ∇ will be the vertical gradient $\nabla_{\mathbb{V}}$; we will take $\xi \in T_x^* M$. If m = 0, degree ≥ 0 also implies non-zero.

Proof. We assume the degree of $\langle \xi, \nabla w(\bullet) \rangle |_{\mathbb{S}^{n-2}_{\xi}}$ is always $\langle m$ (for all $\xi \neq 0$) and show that this forces w to be of degree $\leq m$. In fact, we may assume without loss of generality that w is either pure odd or pure even, and of the same parity as m + 1. Let us deal with the $m \geq 2$ case first.

First of all, we extend the smooth function w to an (m-1)-homogeneous function on E(which we still denote by w). In particular, this extension is smooth on $E \setminus \{0\}$. We now claim that for every ξ , $dw(\xi^{\sharp})$ is a homogeneous polynomial of degree $\leq m-2$ on ker ξ . Indeed, consider a point $v \in \ker \xi \setminus \{0\}$. The total gradient on E is

$$\nabla^{\text{tot}} f = \sum_{i=1}^{n} \partial_{x_i} f(v) . \partial_{x_i} = \nabla f + \mathrm{d} f(\vec{n}) . \vec{n},$$

where $\vec{n} := \frac{v}{|v|}$, ∇ denotes the gradient of f in restriction to the spheres and $(x_i)_{i=1}^n$ are the coordinates induced by an orthonormal basis $(\mathbf{e}_i)_{i=1}^n$ of E. Hence, for $v \in \ker \xi$, we have:

$$\langle \xi, \nabla \mathbf{w}(v) \rangle = \sum_{i=1}^{n} \xi_i \partial_{x_i} \mathbf{w}(v) - \mathbf{d} \mathbf{w}(\vec{n}) \cdot \langle \xi, \vec{n} \rangle = \mathbf{d} \mathbf{w}(\xi^{\sharp}).$$
(3.2)

Therefore, $\langle \xi, \nabla w(v) \rangle|_{\ker \xi}$ is a homogeneous function of degree m-2 whose restriction to the sphere \mathbb{S}^{n-2}_{ξ} is of same parity as m, and thus has degree $\leq m-2$. As a consequence, it is a homogeneous polynomial of degree m-2 on ker ξ .

We now fix an arbitrary $v_0 \in E \setminus \{0\}$ and consider the Taylor-expansion of w at this point:

$$w(v) = \underbrace{\sum_{|\alpha| \le m-1} (v - v_0)^{\alpha} (\alpha!)^{-1} \partial_x^{\alpha} f(v_0)}_{P(v):=} + R(v), \qquad (3.3)$$

where $R(v) = \mathcal{O}(|v - v_0|^m)$. We consider $v_1 \in E \setminus \{0\}$, $w \in \text{Span}(v_0, v_1)^{\perp}$. If we differentiate (3.3) in the *w*-direction and then restrict to the hyperplane w^{\perp} , then we know by the previous discussion that $dw(w)|_{w^{\perp}}$ is a polynomial of degree $\leq m - 2$, and so is $dP(w)|_{w^{\perp}}$. Moreover, from Taylor's theorem $dR(w) = \mathcal{O}(|v - v_0|^{m-1})$. As a consequence: $d(w - P)(w)|_{w^{\perp}} = dR(w)|_{w^{\perp}}$ is a polynomial of degree $\leq m - 2$ which vanishes to order m - 1 at v_0 ; it is therefore constant equal to 0. Evaluating at v_1 , this shows that dR(w) = 0 at v_1 .

We now introduce $G_{v_0} \subset SO(n)$, the isotropy subgroup of v_0 , i.e. the subgroup of rotations fixing the v_0 axis. By the previous discussion, R satisfies the following (see Figure 1): given a sphere $\mathbb{S}^{n-1}(r) := \{|v| = r\}, \gamma^* R = R$ for all $\gamma \in G_{v_0}$.

We restrict this equality to the unit sphere \mathbb{S}^{n-1} and observe that (3.3) implies $w|_{\mathbb{S}^{n-1}} = q_{v_0} + S_{v_0}$, where $q_{v_0} \in \Omega_{\leq m-1} = \bigoplus_{k\geq 0} \Omega_{m-1-k}$ is a sum of spherical harmonics of degree $\leq m-1$ and S_{v_0} is invariant by the action of G_{v_0} . Note that v_0 is arbitrary and taking



FIGURE 1. The function R is constant along the red orbits which correspond to the rotation around the v_0 -axis.

some other v_1 , we see that $S_{v_1} - S_{v_0} \in \Omega_{\leq m-1}$. As each Ω_k is a representation of SO(*n*) by pullback, in particular it is invariant by G_{v_1} . This gives that for all $\gamma \in G_{v_1}$, one has: $\gamma^* S_{v_1} - \gamma^* S_{v_0} = S_{v_1} - \gamma^* S_{v_0} \in \Omega_{\leq m-1}$. Hence, for all $\gamma \in G_{v_1}, \gamma^* S_{v_0} - S_{v_0} \in \Omega_{\leq m-1}$. Taking $\gamma' \in G_{v_2}$ for some other arbitrary v_2 , we see that $(\gamma')^* \gamma^* S_{v_0} - (\gamma')^* S_{v_0} \in \Omega_{\leq m-1}$ and since we also have $(\gamma')^* S_{v_0} - S_{v_0} \in \Omega_{\leq m-1}$, this gives that $(\gamma')^* \gamma^* S_{v_0} - S_{v_0} \in \Omega_{\leq m-1}$. By induction, for any $\gamma_1, \ldots, \gamma_\ell$ belonging to isotropy subgroups of SO(*n*), we have:

$$\gamma_1^* \dots \gamma_\ell^* S_{v_0} - S_{v_0} \in \Omega_{\leq m-1}.$$

As products of isotropy subgroups generate SO(n), we deduce that for all $\gamma \in SO(n)$, $\gamma^* S_{v_0} - S_{v_0} \in \Omega_{\leq m-1}$. Decomposing $S_{v_0} = \sum_{k \geq 0} (S_{v_0})_k$ into spherical harmonics, we then see that $\gamma^*(S_{v_0})_k = (S_{v_0})_k$ for all $k \geq m$ and $\gamma \in SO(n)$. As Ω_k is irreducible [Hel00, Theorem 3.1], this implies that $(S_{v_0})_k = 0$. Hence S_{v_0} is of degree $\leq m - 1$ and w $|_{\mathbb{S}^{n-1}} = q_{v_0} + S_{v_0}$ is also of degree $\leq m - 1$. This completes the proof of the $m \geq 2$ case.

Finally, if m = 0, 1, then (3.2) implies that $d w(\xi^{\sharp})|_{\ker \xi} \equiv 0$. Then taking R(v) := w(v) in (3.3), it is straightforward that d R(w) = 0 at v_1 where w, v_1 are as before. The remainder of the proof works the same to show that $w|_{\mathbb{S}^{n-1}}$ is invariant under the isotropy subgroups, hence constant, contradicting that deg(w) > 0.

3.2. The extension operator. In this paragraph, we study an operator of extension from a hypersphere \mathbb{S}_{ξ}^{n-2} to the whole sphere \mathbb{S}^{n-1} . First of all, for $m \in \mathbb{Z}_{\geq 1}$, we introduce the constant:

$$C_m := \int_0^\pi \sin^{m-1}(\varphi) \,\mathrm{d}\,\varphi = \sqrt{\pi} \frac{\Gamma(\frac{m}{2})}{\Gamma(\frac{m+1}{2})},\tag{3.4}$$

where Γ is the usual Gamma function. Given a smooth function $f \in C^{\infty}(\mathbb{S}^{n-2}_{\xi})$ or more generally, one can take a section $f \in C^{\infty}(\mathbb{S}^{n-2}_{\xi}, \pi^* \mathcal{E})$, where $\pi : \mathbb{S}^{n-1} \to \{0\}$ is the projection, we define its *extension of degree* $k \in \mathbb{N}$ to the whole sphere \mathbb{S}^{n-1} by the formula:

$$E_{\xi}^{k} f \in L^{2}(\mathbb{S}^{n-1}, \pi^{*} \mathcal{E}), \quad E_{\xi}^{k} f \big(\cos(\varphi) \vec{n}(\xi) + \sin(\varphi) u \big) := \sin^{k}(\varphi) f(u).$$
(3.5)

Note that E_{ξ}^k extends to $L^2(\mathbb{S}_{\xi}^{n-2}, \pi^* \mathcal{E})$ by continuity since by definition and (3.1)

$$\|E_{\xi}^{k}f\|_{L^{2}(\mathbb{S}^{n-1},\pi^{*}\mathcal{E})}^{2} = C_{2k+n-1}\|f\|_{L^{2}(\mathbb{S}^{n-2}_{\xi},\pi^{*}\mathcal{E})}^{2}.$$
(3.6)

Morever, we have (cf. [Lef19b, Lemma B.1.1]):

Lemma 3.4. For any $f \in C^{\infty}(\mathbb{S}^{n-2}_{\xi}, \pi^* \mathcal{E})$ and $f' \in \otimes^{m'}_{S} E^* \otimes \mathcal{E}$, and all $m \in \mathbb{Z}_{\geq 0}$, we have:

$$C_{m+m'+n-1} \int_{\mathbb{S}_{\xi}^{n-2}} \langle f(u), \pi_{m'}^* f'(u) \rangle_{\mathcal{E}} dS_{\xi}(u) = \langle \pi_{\ker_{i_{\xi}\sharp}} \pi_{m'*} E_{\xi}^m(f), f' \rangle_{\otimes_{S}^{m'} E^* \otimes \mathcal{E}}.$$
(3.7)

Proof. The left hand side of (3.7) equals, after using (3.4):

$$\begin{split} &\int_{\mathbb{S}_{\xi}^{n-2}} \int_{0}^{\pi} \left\langle \sin^{m}(\varphi) f(u), \pi_{m'}^{*} \pi_{\ker \imath_{\xi^{\sharp}}} f' \left(\cos(\varphi) . \vec{n}(\xi) + \sin(\varphi) . u \right) \right\rangle_{\mathcal{E}} . \sin^{n-2}(\varphi) . d\varphi . dS_{\xi}(u) \\ &= \int_{\mathbb{S}^{n-1}} \left\langle E_{\xi}^{m} f, \pi_{m'}^{*} \pi_{\ker \imath_{\xi^{\sharp}}} f' \right\rangle_{\mathcal{E}} . dS = \left\langle \pi_{\ker \imath_{\xi^{\sharp}}} \pi_{m'*} E_{\xi}^{m}(f), f' \right\rangle_{\otimes_{S}^{m'} E^{*} \otimes \mathcal{E}}, \end{split}$$

where in the first line we used that $\langle \xi, u \rangle = 0$ on \mathbb{S}_{ξ}^{n-2} by definition and in the second equality we used the Jacobian formula (3.1). This concludes the proof.

We have the following result on the degree:

Lemma 3.5. For all $\xi \in E^* \setminus \{0\}$, the following holds. Let $f \in C^{\infty}(\mathbb{S}^{n-2}_{\xi})$ such that $\deg(f) \ge m+1$. Then, $\deg(E^m_{\xi}(f)) \ge m+1$.

Proof. We argue by contradiction. We assume that $E_{\xi}^{m}(w)$ has degree $\leq m$. In particular, it is smooth. Moreover, observe that its (m-1)-th jet vanishes at the North pole $N := \{\varphi = 0\} = \vec{n}(\xi)$. We can therefore compute its differential of degree m. Let $0 \neq Z \in T_N \mathbb{S}^{n-1}$, Z = rZ', where |Z'| = 1. Note that in the (φ, u) -coordinates, Z' corresponds to ∂_{φ} at $(\varphi = 0, u)$ where $u = Z' \in \mathbb{S}_{\xi}^{n-2}$. Then:

$$d^{m} E_{\xi}^{m}(\mathbf{w})_{N}(Z,...,Z) = r^{m} \partial_{\varphi}^{m}(E_{\xi}^{m}(\mathbf{w}))(0,u) = r^{m} \partial_{\varphi}^{m} \big|_{\varphi=0}(\sin^{m}(\varphi) \,\mathbf{w} \,|_{\mathbb{S}_{\xi}^{n-2}}(u)) = m! \times r^{m} \,\mathbf{w} \,|_{\mathbb{S}_{\xi}^{n-2}}(u).$$
(3.8)

There is a natural identification between $T_N \mathbb{S}^{n-1}$ and $\ker \xi = \vec{n}(\xi)^{\perp}$, and with this identification $d^m E_{\xi}^m(\mathbf{w})_N$ defines a symmetric *m*-tensor on $\ker \xi$. Then (3.8) says $\mathbf{w}|_{\mathbb{S}_{\xi}^{n-2}} = \frac{1}{m!}\pi_m^*(d^m E_{\xi}^m(\mathbf{w})_N)$ has degree $\leq m$, which is a contradiction.

3.3. Multiplication of spherical harmonics. We end this section with standard results on multiplication of spherical harmonics:

Lemma 3.6. Let $m, k \in \mathbb{Z}_{\geq 0}$ and assume without loss of generality that $m \geq k$. If $f \in \Omega_m(E), f' \in \Omega_k(E)$, then:

$$f \times f' \in \bigoplus_{\ell=0}^k \Omega_{m+k-2\ell}(E).$$

Proof. First of all, extending f and f' by m- and k-homogeneity to E, respectively, we directly see that $f \times f'$ is a homogeneous polynomial of degree m + k and so by (2.9):

$$f \times f' \in \bigoplus_{\ell \ge 0} |v|^{2\ell} \mathbf{H}_{m+k-2\ell}(E).$$

The only non-trivial part is to show that the projection onto $\bigoplus_{\ell \ge k+1} |v|^{2\ell} \mathbf{H}_{m+k-2\ell}(E)$ is zero. For that it suffices to show that $\Delta^{\ell}(f \times f') = 0$ as long as $\ell \ge k+1$. Observe that:

$$\Delta(f \times f') = (\Delta f) \times f' + 2\nabla f \cdot \nabla f' + f \times (\Delta f') = 2\sum_{i=1}^{n} \partial_{v_i} f \times \partial_{v_i} f',$$

and thus by iteration:

$$\Delta^{\ell}(f \times f') = 2^{\ell} \sum_{|\alpha| = \ell} \partial_{v}^{\alpha} f \times \partial_{v}^{\alpha} f',$$

which clearly vanishes for $\ell \ge k+1$ as f' is a polynomial of degree k.

In the particular case where k = 1, the previous lemma shows that $f \in \Omega_1(E)$ gives rise to two operators f_{\pm} defined in the following way: if $w \in \Omega_m(E)$, then $f \times w = f_- w + f_+ w$ with $f_{\pm} w \in \Omega_{m\pm 1}(E)$. Moreover, by extending f and w as 1- and m-homogeneous harmonic polynomials denoted by the same letter, we get (∇ denotes the total gradient of E)

$$f_{-} w = \frac{1}{n + 2(m - 1)} (\nabla f \cdot \nabla w)|_{\mathbb{S}^{n-1}}.$$
(3.9)

In fact, for non-zero f the map $f_-: \Omega_m(E) \to \Omega_{m-1}(E)$ is surjective, implying also that $f_+: \Omega_m(E) \to \Omega_{m+1}(E)$ is injective (see [CL21a, Lemma 2.3]).

Lemma 3.7. Assume $n \ge 2$ and let $m, k \in \mathbb{Z}_{\ge 0}$. Consider $w \in C^{\infty}(\mathbb{S}^{n-1})$ such that $\deg(w) \ge m$. Then, there exists $f \in \mathbf{S}_k(E)$ such that $\deg(f, w) \ge m + k$.

Equivalently, there exists $f \in \bigotimes_{S}^{k} E^{*}$ such that $\deg(\pi_{k}^{*} f. \mathbf{w}) \geq m + k$.

Proof. We first prove the case k = 1; the cases m = 0 or k = 0 are trivial so we assume $m \ge 1$ from now on. We write $w = \sum_{j=0}^{\infty} w_j$, where $w_j \in \Omega_i$, and denote by the same letter the harmonic extension of w_j (as a *j*-homogeneous polynomial) to \mathbb{R}^n . Take $\ell \ge m$ such that $w_{\ell} \ne 0$, and assume for any $i = 1, \ldots, n$ that $(v_i)_+ w_{\ell} + (v_i)_- w_{\ell+2} = 0$, which by (3.9) is equivalent to:

$$(n+2(\ell+1))^{-1}\partial_{v_i} \mathbf{w}_{\ell+2} + v_i \mathbf{w}_{\ell} - |v|^2(n+2(\ell-1))^{-1}\partial_{v_i} \mathbf{w}_{\ell} = 0.$$

Multiplying by v_i and summing over i, we obtain using Euler's formula (i.e. homogeneity)

$$-\frac{\ell+2}{n+2(\ell+1)} \mathbf{w}_{\ell+2} = \frac{n+\ell-2}{n+2(\ell-1)} |v|^2 \mathbf{w}_{\ell}.$$

Applying Δ , this contradicts the fact that $w_{\ell} \neq 0$.

For general $k \in \mathbb{Z}_{\geq 1}$, by iteratively applying the case k = 1 above, there exist $f_1, ..., f_k \in \Omega_1(E)$ such that $\deg(f_k \cdots f_1 w) \geq m + k$. Since $f := f_k \cdots f_1 \in \mathbf{S}_k(E)$, this completes the proof.

Note that there is a straightforward extension to the bundle case (just by applying the previous lemma coordinate-wise), that is, when considering sections of a trivial bundle $\pi^* \mathcal{E} \to \mathbb{S}^{n-1}$, where $\pi : \mathbb{S}^{n-1} \to \{0\}$ is the constant map. We record it here and leave the proof as an exercise for the reader:

Lemma 3.8. Let $m, k \in \mathbb{Z}_{\geq 0}$. Consider $w \in C^{\infty}(\mathbb{S}^{n-1}, \pi^* \mathcal{E})$ such that $\deg(w) \geq m$. Then, there exists $f \in \mathbf{S}_k(E) \otimes \operatorname{End}_{\mathrm{sk}}(\mathcal{E})$ such that $\deg(f, w) \geq m + k$.

4. PSEUDODIFFERENTIAL NATURE OF PERTURBED GENERALIZED X-RAY TRANSFORMS

Under a weaker form, the results of this section can be found in [Gui17a, GL21] and in [Lef19b, Chapter 2] where the principal symbol of the generalized X-ray transform is computed in details. We here need a more general result where we "sandwich" (pseudo)differential operators:

Proposition 4.1. Let $P_R, P_L : C^{\infty}(SM, \pi^* \mathcal{E}) \to C^{\infty}(SM, \pi^* \mathcal{E})$ be differential operators of degree $m_R, m_L \ge 0$ and fix $m_1, m_2 \in \mathbb{N}_0$. Then the operator

$$A_{P_R,P_L} := \pi_{m_1*} P_L \mathcal{I}_{\nabla} \varepsilon P_R \pi_{m_2}^*,$$

is a classical⁸ pseudodifferential operator of order $m := m_R + m_L - 1$ in

$$A_{P_R,P_L} \in \Psi^m(M, \otimes_S^{m_2} T^*M \otimes \mathcal{E} \to \otimes_S^{m_1} T^*M \otimes \mathcal{E})$$

Moreover, its principal symbol satisfies, for any $f \in \bigotimes_{S}^{m_{2}}T_{x}^{*}M \otimes \mathcal{E}_{x}$ and $f' \in \bigotimes_{S}^{m_{1}}T_{x}^{*}M \otimes \mathcal{E}_{x}$:

$$\langle \sigma_{A_{P_R},P_L}(x,\xi)f,f' \rangle_{\otimes_S^{m_1}T^*M_x \otimes \mathcal{E}_x}$$

$$= \frac{2\pi}{|\xi|} \int_{\mathbb{S}_{\xi}^{n-2}} \left\langle \sigma_{P_R}\big((x,u),\xi_{\mathbb{H}}(x,u)\big)(\pi_{m_2}^*f(u)),\sigma_{P_L^*}\big((x,u),\xi_{\mathbb{H}}(x,u)\big)(\pi_{m_1}^*f'(u)) \right\rangle_{\mathcal{E}_x} \mathrm{d}\, S_{\xi}(u),$$

$$(4.1)$$

where $\xi_{\mathbb{H}}(x, u) := \xi(\mathbf{d}_{x,u} \pi(\bullet))$. More explicitly, the principal symbol of A_{P_R, P_L} is given by the formula, for any $m \in \mathbb{N}_0$:

$$\sigma_{A_{P_R,P_L}}(x,\xi)f = \frac{2\pi}{|\xi|} C_{m_1+m+n-1}^{-1} \pi_{\ker \iota_{\xi^{\sharp}}} \pi_{m_1*} E_{\xi}^m \Big[\sigma_{P_L P_R} \big(x, u, \xi_{\mathbb{H}}(x, u)\big) \big(\pi_{m_2}^* \pi_{\ker \iota_{\xi^{\sharp}}} f\big) \Big].$$
(4.2)

This was originally proved with $P_L = P_R = 1$ in [Gui17a], see also [Lef19b]. Note that one could actually take P_R, P_L to be pseudodifferential of arbitrary order (that only makes the proof slightly longer but the idea is the same).

In the following, we will refer to this result as the sandwich Proposition 4.1. In the case where $\mathcal{E} = \mathbb{C} \times M$, the formula reads (using that $\sigma_{P_L^*} = \overline{\sigma_{P_L}}$):

$$\langle \sigma_{A_{P_R},P_L}(x,\xi)f,f' \rangle_{\otimes_S^{m_1}T^*M_x \otimes \mathcal{E}_x}$$

= $\frac{2\pi}{|\xi|} \int_{\mathbb{S}_{\xi}^{n-2}} \sigma_{P_R}((x,u),\xi_{\mathbb{H}}(x,u)) \sigma_{P_L}((x,u),\xi_{\mathbb{H}}(x,u)) .\pi_{m_2}^*f(u).\overline{\pi_{m_1}^*f'(u)} \,\mathrm{d}\, S_{\xi}(u).$

We will only prove Proposition 4.1 in the case of the trivial line bundle with trivial connection in order to simplify the discussion; the generalization to the twisted case is straightforward modulo some tedious notations. We also make the following important remark:

Remark 4.2. Proposition 4.1 can also be generalized by considering differential operators

$$P_R: C^{\infty}(SM, \otimes_S^{m_2} T^*(SM)) \to C^{\infty}(SM), \quad P_L: C^{\infty}(SM) \to C^{\infty}(SM, \otimes_S^{m_1} T^*(SM)),$$

(of degree $m_R, m_L \ge 0$) and looking at the operator:

$$A_{P_R,P_L} := \pi_{m_1,\operatorname{Sas}_*} P_L \mathcal{I} P_R \pi_{m_2,\operatorname{Sas}}^*,$$

⁸See below (4.5) for a definition.

where $\pi_{m_2,Sas}^*$ denotes the Sasaki lift introduced in (2.13). The same proof shows that this operator is pseudodifferential of order $m_R + m_L - 1$ with principal symbol satisfying:

$$\langle \sigma_{A_{P_{R},P_{L}}}(x,\xi)f,f' \rangle_{\otimes_{S}^{m_{1}}T^{*}M_{x}}$$

$$= \frac{2\pi}{|\xi|} \int_{\mathbb{S}_{\xi}^{n-2}} \sigma_{P_{R}}((x,u),\xi_{\mathbb{H}}(x,u)) (\pi_{m_{2},\mathrm{Sas}}^{*}f(u)) \overline{\sigma_{P_{L}^{*}}((x,u),\xi_{\mathbb{H}}(x,u))} (\pi_{m_{1},\mathrm{Sas}}^{*}f'(u)) \mathrm{d} S_{\xi}(u),$$

$$(4.3)$$

where $\xi_{\mathbb{H}}(x, u) := \xi(d_{x,u} \pi(\bullet))$. We leave this claim as an exercise for the reader.

First of all, for some $\varepsilon > 0$ we set (here $e^{tX} = \varphi_t^*$):

$$\mathcal{I}_{\varepsilon} := \int_{-\varepsilon}^{+\varepsilon} e^{tX} \,\mathrm{d}\, t.$$

Lemma 4.3. The following property holds:

$$\pi_{m_1*}P_L\mathcal{I}P_R\pi_{m_2}^* - \pi_{m_1*}P_L\mathcal{I}_{\varepsilon}P_R\pi_{m_2}^* \in \Psi^{-\infty}(M, \otimes_S^{m_2}T^*M \to \otimes_S^{m_1}T^*M).$$

Proof. Recall that $\mathcal{I} = R_{+,0} + R_{-,0} + \Pi_+$ and the term Π_+ will only contribute to a smoothing operator. Observe that:

$$\pi_{m_{1}*}P_{L}(R_{+,0}+R_{-,0})P_{R}\pi_{m_{2}}^{*} - \pi_{m_{1}*}P_{L}\mathcal{I}_{\varepsilon}P_{R}\pi_{m_{2}}^{*} = \pi_{m_{1}*}P_{L}(e^{-\varepsilon X}R_{+,0} + e^{\varepsilon X}R_{-,0})P_{R}\pi_{m_{2}}^{*} - \underbrace{2\varepsilon.\pi_{m_{1}*}P_{L}\Pi_{+}P_{R}\pi_{m_{2}}^{*}}_{\in\Psi^{-\infty}}.$$

We want to prove that the first operator on the right hand side is smoothing. We deal with

$$K := \pi_{m_1} P_L e^{-\varepsilon X} R_{+,0} P_R \pi_{m_2}^*,$$

while the other term is dealt with similarly. It is sufficient to prove that if $f \in \mathcal{D}'(M, \otimes_S^{m_2} T^*M)$, one has $Kf \in C^{\infty}(M, \otimes_S^{m_1} T^*M)$. For that, we will use the wavefront set calculus of Hörmander [Hör03, Chapter 8].

Using the notation of §2.1, define the subbundles $\mathbb{H}^*, \mathbb{V}^* \subset T^*(SM)$ such that $\mathbb{H}^*(\mathbb{H} \oplus \mathbb{R} \cdot X) = 0, \mathbb{V}^*(\mathbb{V}) = 0$. Observe that since $\pi_{m_2}^*$ is a pullback operator, we have $WF(\pi_{m_2}^*f) \subset \mathbb{V}^*$ (see also [Lef19a, Lemma 2.1] for a detailed proof). Since P_R is a differential operator, we have $WF(P_R\pi_{m_2}^*f) \subset \mathbb{V}^*$. We then use the characterization of the wavefront set of the resolvent $R_{+,0}$ in [DZ16, Proposition 3.3], namely⁹:

$$WF'(R_{+,0}) \subset \Delta(T^*(SM)) \cup \Omega_+ \cup E_u^* \times E_s^*,$$
(4.4)

where $\Delta(T^*(SM))$ is the diagonal in $T^*(SM) \times T^*(SM)$, and

$$\Omega_{+} := \{ (\Phi_{t}(z,\xi), (z,\xi)) \mid t \ge 0, \ \langle \xi, X(z) \rangle = 0 \}$$

is the positive flow-out and $\Phi_t : T^*(SM) \to T^*(SM)$ is the symplectic lift of the geodesic flow $(\varphi_t)_{t\in\mathbb{R}}$, given by $\Phi_t(z,\xi) := (\varphi_t(z), \mathrm{d} \, \varphi_t^{-\top}(\xi))^{10}$. From (4.4) we obtain using [Hör03, Theorem 8.2.13]:

$$WF(R_{+,0}P_R\pi_{m_2}^*f) \subset E_u^* \cup_{t \ge 0} \Phi_t(\mathbb{V}^* \cap \ker \imath_X).$$

⁹We use the standard conventions, namely if $B : C^{\infty}(M) \to \mathcal{D}'(M)$ is a linear operator with kernel $K_B \in \mathcal{D}'(M \times M)$, we define $WF'(B) := \{(x, \xi, y, \eta) \in T^*M \times T^*M \mid (x, \xi, y, -\eta) \in WF(K_B)\}.$

¹⁰We use $^{-\top}$ to denote the inverse transpose.

By the behaviour of the wavefront set under pullbacks (see [Hör03, Theorem 8.2.4]):

$$WF(P_L e^{-\varepsilon X} R_{+,0} P_R \pi_{m_2}^* f) \subset WF(e^{-\varepsilon X} R_{+,0} P_R \pi_{m_2}^* f) \subset E_u^* \cup_{t \ge \varepsilon} \Phi_t(\mathbb{V}^* \cap \ker i_X).$$

Next, as $\pi_{m_{1*}}$ is a pushforward, we obtain (see [Mel03, Proposition 4.12]):

$$WF(Kf) \subset \left\{ (x,\xi) \in T^*M \mid \exists v \in S_x M, \, \xi(\mathsf{d}_{(x,v)} \, \pi(\bullet)) \in E^*_u \cup_{t \ge \varepsilon} \Phi_t(\mathbb{V}^* \cap \ker \imath_X) \right\},\$$

where we note that $\xi(d_{(x,v)} \pi(\bullet)) \in \mathbb{V}^*(x,v)$. As a consequence, the lemma is proved if we can show that $\mathbb{V}^* \cap (E_u^* \cup_{t \geq \varepsilon} \Phi_t(\mathbb{V}^* \cap \ker \iota_X)) = \{0\}$. But this follows from (2.7) and (2.8), completing the proof.

We now turn to the sandwich Proposition 4.1. For that, it is convenient to use the historical characterization of pseudodifferential operators [Hör65, Definition 2.1] which we now recall: P is a pseudodifferential operator of order $m \in \mathbb{R}$ if $P : C^{\infty}(M) \to C^{\infty}(M)$ is continuous, and there exists a sequence $s_0 = 0 < s_1 < \dots$ of real numbers converging to $+\infty$ such that for all $f \in C^{\infty}(M)$, $S \in C^{\infty}(M)$ such that $dS \neq 0$ on $\operatorname{supp}(f)$, there is an asymptotic expansion:

$$e^{-i\frac{S}{\hbar}}P(fe^{i\frac{S}{\hbar}}) \sim h^{-m} \sum_{j=0}^{+\infty} P_j(f,S)h^{s_j}.$$
 (4.5)

By this, we mean that for every integer N > 0, for every compact set ¹¹K of real-valued functions $S \in C^{\infty}(M)$ with $dS \neq 0$ on $\operatorname{supp}(f)$, for every 0 < h < 1, the following holds: the error term

$$h^{-s_N+m}\left(e^{-i\frac{S}{h}}P\left(fe^{i\frac{S}{h}}\right) - h^{-m}\sum_{j=0}^{N-1}P_j(f,S)h^{s_j}\right)$$
(4.6)

belongs to a bounded set in $C^{\infty}(M)$ with bound independent of h. In particular, P is *differential* if and only if the sum (4.5) ranges over a finite number of j's and P is *classical* if the s_j 's take integer values.

Proof of Proposition 4.1. We first note that the formula (4.2) is an immediate consequence of (4.1) and Lemma 3.4; henceforth we focus on (4.1). We divide the proof in two steps.

1. Principal symbol computation. For the moment, let us assume that the operator is pseudodifferential and compute its principal symbol. By Lemma 4.3, we can replace \mathcal{I} by $\mathcal{I}_{\varepsilon}$ in the definition of the operator, that is, it suffices to compute the principal symbol of $\pi_{m_1*}P_L\mathcal{I}_{\varepsilon}P_R\pi_{m_2}^*$.

Take a Lagrangian state $f_h := e^{i\frac{S}{h}}\widetilde{f}$, where $S \in C^{\infty}(M)$ is a real-valued, smooth phase such that $S(x_0) = 0$, $dS(x_0) = \xi_0$, $\widetilde{f} \in C^{\infty}(M, \bigotimes_S^{m_2} T^*M)$ and $\widetilde{f}(x_0) := f \in \bigotimes_S^{m_2} T^*_{x_0}M$, and further assume that dS does not vanish on the support of \widetilde{f} . As P_R is a differential operator, we have:

$$P_R \pi_{m_2}^* f_h = P_R \left(e^{i \frac{\pi_0^* S}{h}} \pi_{m_2}^* \widetilde{f} \right) = h^{-m_R} e^{i \frac{\pi_0^* S}{h}} \left(\sigma_{P_R} \left(\bullet, S_{\mathbb{H}}(\bullet) \right) . \pi_{m_2}^* \widetilde{f}(\bullet) + \mathcal{O}_{C^{\infty}}(h) \right),$$

¹¹A set $A \subset C^{\infty}(M)$ is bounded if there exists a sequence $(A_k)_{k \in \mathbb{Z}_{\geq 0}}$ such that for all $f \in A$, $||f||_{C^k(M)} \leq A_k$. It is known that A is compact if and only if it is closed and bounded. where $S_{\mathbb{H}}(x,v) := d_x S \circ d_{(x,v)} \pi$. Hence:

$$\begin{aligned} \mathcal{I}_{\varepsilon} P_{R} \pi_{m_{2}}^{*} f_{h}(x,v) &= \int_{-\varepsilon}^{+\varepsilon} h^{-m_{R}} e^{\frac{i}{h} S(\pi(\varphi_{t}(x,v)))} \\ &\times \left(\sigma_{P_{R}} \big(\varphi_{t}(x,v), S_{\mathbb{H}}(\varphi_{t}(x,v)) \big) \pi_{m_{2}}^{*} \widetilde{f}(\varphi_{t}(x,v)) + \mathcal{O}_{C^{\infty}}(h) \right) \mathrm{d} t, \end{aligned}$$

and thus:

$$P_{L}\mathcal{I}_{\varepsilon}P_{R}\pi_{m_{2}}^{*}f_{h}(x,v) = \int_{-\varepsilon}^{+\varepsilon} h^{-(m_{R}+m_{L})}e^{\frac{i}{h}S(\pi(\varphi_{t}(x,v)))} \Big(\sigma_{P_{R}}\big(\varphi_{t}(x,v),S_{\mathbb{H}}(\varphi_{t}(x,v))\big) \\ \times \sigma_{P_{L}}\big((x,v),S_{\mathbb{H}}^{(t)}(x,v)\big)\pi_{m_{2}}^{*}\widetilde{f}(\varphi_{t}(x,v)) + \mathcal{O}_{C^{\infty}}(h)\Big) \,\mathrm{d}\,t,$$

where $S_{\mathbb{H}}^{(t)}(x,v) := d_x S \circ d_{(x,v)} \pi \circ d_{(x,v)} \varphi_t$ (and $S_{\mathbb{H}}^{(0)} = S_{\mathbb{H}}$). This gives for $m = m_R + m_L - 1$ and any $f' \in \bigotimes_S^{m_1} T_{x_0}^* M$:

$$\begin{aligned} \langle \sigma_{A_{P_{R},P_{L}}}(x_{0},\xi_{0})f,f'\rangle_{\otimes_{S}^{m_{1}}T_{x_{0}}^{*}M} &= \lim_{h \to 0} h^{m} \langle (A_{P_{R},P_{L}}f_{h})(x_{0}),f'\rangle_{\otimes_{S}^{m_{1}}T_{x_{0}}^{*}M} \\ &= \lim_{h \to 0} h^{m} \int_{S_{x_{0}}M} (P_{L}\mathcal{I}_{\varepsilon}P_{R}\pi_{m_{2}}^{*}f_{h})(x_{0},v).\overline{\pi_{m_{1}}^{*}f'}(x_{0},v) \,\mathrm{d}\,S_{x_{0}}(v) \\ &= \lim_{h \to 0} h^{-1} \int_{S_{x_{0}}M} \int_{-\varepsilon}^{+\varepsilon} e^{\frac{i}{h}S(\pi(\varphi_{t}(x_{0},v)))}.\overline{\pi_{m_{1}}^{*}f'}(v) \\ &\times \left(\sigma_{P_{R}}\left(\varphi_{t}(x_{0},v),S_{\mathbb{H}}(\varphi_{t}(x_{0},v))\right).\sigma_{P_{L}}\left((x_{0},v),S_{\mathbb{H}}^{(t)}(x_{0},v)\right)\right) \cdot \sigma_{P_{L}}\left((x_{0},v),S_{\mathbb{H}}^{(t)}(x_{0},v)\right) + \mathcal{O}_{C^{\infty}}(h) \right) \,\mathrm{d}\,t \,\mathrm{d}\,S_{x_{0}}(v) , \end{aligned}$$

$$(4.7)$$

where $d S_{x_0}$ stands for the round measure on the sphere $S_{x_0}M$. As we shall see, the term h^{-1} comes from the fact that we will perform a stationary phase over a two-dimensional space.

We define the (real) phase $\Phi : (-\varepsilon, \varepsilon) \times S_{x_0}M \to \mathbb{R}$ by $\Phi(t, v) := S(\pi(\varphi_t(x_0, v)))$. We recall (see §3.1) the diffeomorphism, singular at the poles, $\mathbb{S}_{\xi_0}^{n-2} \times (0, \pi) \ni (u, \varphi) \mapsto v(u, \varphi) := \cos(\varphi)\vec{n}(\xi_0) + \sin(\varphi)u \in S_{x_0}M$. Observe that for fixed $u \in \mathbb{S}_{\xi_0}^{n-2}$, the phase $\Phi_u : (-\varepsilon, \varepsilon) \times (0, \pi) \to \mathbb{R}$ defined by $\Phi_u(t, \varphi) := \Phi(t, v(u, \varphi))$ has a critical point at $t = 0, \varphi = \frac{\pi}{2}$ and the determinant of the Hessian at this point is equal to $-|dS(x_0)|^2 = -|\xi_0|^2$ (see the proof of [GL21, Theorem 4.4] or [Lef19b, Theorem 2.5.1] for further details). Hence by the stationary phase lemma [Zwo12, Theorem 3.16], for any $u \in \mathbb{S}_{\xi_0}^{n-2}$, writing $v = v(u, \varphi)$:

$$\lim_{h \to 0} h^{-1} \int_{0}^{\pi} \int_{-\varepsilon}^{+\varepsilon} e^{\frac{i}{\hbar} \Phi_{u}(t,\varphi)} \cdot \overline{\pi_{m_{1}}^{*} f'}(v) \cdot \left(\mathcal{O}_{C^{\infty}}(h) + \sigma_{P_{R}}\left(\varphi_{t}(x_{0},v), S_{\mathbb{H}}(\varphi_{t}(x_{0},v))\right) \cdot \sigma_{P_{L}}\left((x_{0},v), S_{\mathbb{H}}^{(t)}(x_{0},v)\right) \cdot \pi_{m_{2}}^{*} \widetilde{f}(\varphi_{t}(x_{0},v))\right) \sin^{n-2}(\varphi) \, \mathrm{d} t \, \mathrm{d} \varphi$$

$$= \frac{2\pi}{|\xi_{0}|} \sigma_{P_{R}}\left((x_{0},u), S_{\mathbb{H}}(x_{0},u)\right) \cdot \sigma_{P_{L}}\left((x_{0},u), S_{\mathbb{H}}(x_{0},u)\right) \cdot \pi_{m_{2}}^{*} f(u) \cdot \overline{\pi_{m_{1}}^{*} f'}(u).$$
(4.8)

Using that this limit is uniform in u and integrating over $\mathbb{S}_{\xi_0}^{n-2}$, inserting into (4.7), as well as recalling the Jacobian formula (3.1), completes the proof.

2. Pseudodifferential nature. By the characterization (4.5) of Ψ DOs via the asymptotic expansion, the proof is very similar to the first point except that one needs to go to arbitrary order in the expansions. For the sake of simplicity, we assume that $m_1 = m_2 = 0$ (this does not change the nature of the proof). Consider an arbitrary $f \in C^{\infty}(M)$ and a compact set $K \subset C^{\infty}(M)$ of (real) phases S such that $dS \neq 0$ on $\operatorname{supp}(f)$. By compactness there is an $\varepsilon > 0$ such that for $S \in K$, $dS \neq 0$ on the 2ε -neighbourhood of $\operatorname{supp}(f)$. By Lemma 4.3, it suffices to show that $\pi_{0*}P_L\mathcal{I}_{\varepsilon}P_R\pi_0^*$ is a pseudodifferential operator of order m. In fact, for $d(x, \operatorname{supp}(f)) \geq \varepsilon$ by the definition of $\mathcal{I}_{\varepsilon}$ we have that $\pi_{0*}P_L\mathcal{I}_{\varepsilon}P_R\pi_0^*$ $\left(fe^{i\frac{S}{h}}\right) = 0$. Therefore, in what follows we will consider only $x \in N_{\varepsilon} := \{x \in M \mid d(x, \operatorname{supp}(f)) < \varepsilon\}$.

Since P_R is differential, we can write

$$P_R \pi_0^* \left(f e^{i\frac{S}{h}} \right) = h^{-m_R} e^{i\frac{\pi_0^* S}{h}} \underbrace{\sum_{j=0}^{m_R} P_R^{(j)}(\pi_0^* f, \pi_0^* S) h^j}_{=:f_h} = h^{-m_R} e^{i\frac{\pi_0^* S}{h}} f_h,$$

where $P_R^{(j)}(\pi_0^* f, \pi_0^* S)(x, v)$ depends on the jet of order $\leq j$ of f at x (and on the $(m_R - j)$ -th jet of the phase S). Then:

$$\mathcal{I}_{\varepsilon}P_{R}\pi_{0}^{*}\left(fe^{i\frac{S}{h}}\right)(x,v) = h^{-m_{R}}\int_{-\varepsilon}^{+\varepsilon} e^{\frac{i}{h}\pi_{0}^{*}S(\varphi_{t}(x,v))}f_{h}(\varphi_{t}(x,v))\,\mathrm{d}\,t,$$

which gives:

$$P_{L}\mathcal{I}_{\varepsilon}P_{R}\pi_{0}^{*}\left(fe^{i\frac{S}{h}}\right)(x,v) = h^{-(m_{R}+m_{L})}\int_{-\varepsilon}^{+\varepsilon} e^{\frac{i}{h}\pi_{0}^{*}S(\varphi_{t}(x,v))}\sum_{k=0}^{m_{L}}h^{k}P_{L}^{(k)}\left(e^{tX}f_{h},e^{tX}\pi_{0}^{*}S\right)(x,v)\,\mathrm{d}\,t,$$

and thus:

$$\pi_{0*} P_L \mathcal{I}_{\varepsilon} P_R \pi_0^* \left(f e^{i\frac{S}{h}} \right) (x)$$

= $h^{-(m_R + m_L)} \int_{S_x M} \int_{-\varepsilon}^{+\varepsilon} e^{\frac{i}{h} \pi_0^* S(\varphi_t(x, v))} \sum_{k=0}^{m_L} h^k P_L^{(k)} \left(e^{tX} f_h, e^{tX} \pi_0^* S \right) (x, v) \, \mathrm{d} t \, \mathrm{d} v.$

By introducing $\xi := d S(x) \neq 0$, and using the coordinates $(u, \varphi) \in \mathbb{S}_{\xi}^{n-2} \times (0, \pi)$ on $S_x M$ as in the previous step, as well as the formula (3.1), writing $v = v(u, \varphi)$ we obtain:

$$\pi_{0*}P_{L}\mathcal{I}_{\varepsilon}P_{R}\pi_{0}^{*}\left(fe^{i\frac{S}{h}}\right)(x) = h^{-(m_{R}+m_{L})}$$

$$\times \int_{\mathbb{S}_{\xi}^{n-2}} \int_{0}^{\pi} \int_{-\varepsilon}^{+\varepsilon} e^{\frac{i}{h}\pi_{0}^{*}S(\varphi_{t}(x,v))} \underbrace{\sum_{k=0}^{m_{L}} h^{k}P_{L}^{(k)}\left(e^{tX}f_{h}, e^{tX}\pi_{0}^{*}S\right)(x,v)\sin^{n-2}(\varphi) \,\mathrm{d}\,t \,\mathrm{d}\,\varphi \,\mathrm{d}\,S_{\xi}(u)}_{=:F(x,u,\varphi,t)}$$

$$= h^{-(m_{R}+m_{L})} \int_{\mathbb{S}_{\xi}^{n-2}} \left(\int_{0}^{\pi} \int_{-\varepsilon}^{+\varepsilon} e^{\frac{i}{h}\pi_{0}^{*}S(\varphi_{t}(x,v(u,\varphi)))}F(x,u,\varphi,t) \,\mathrm{d}\,t \,\mathrm{d}\,\varphi\right) \,\mathrm{d}\,S_{\xi}(u).$$

As in (4.8), for fixed $x \in M$, $u \in \mathbb{S}^{n-2}_{\xi}$, we apply the stationary phase lemma [Zwo12, Theorem 3.16] at t = 0, $\varphi = \frac{\pi}{2}$, which gives:

$$\pi_{0*} P_L \mathcal{I}_{\varepsilon} P_R \pi_0^* \left(f e^{i\frac{S}{h}} \right) (x) = h^{-(m_R + m_L - 1)} \int_{\mathbb{S}_{\xi}^{n-2}} e^{i\frac{S(x)}{h}} \left(\sum_{\ell=0}^{N-1} h^{\ell} A_{2\ell}(x, u, D_{\varphi}, D_t) F\left(x, u, \frac{\pi}{2}, 0\right) + h^N R'_N(x, u) \right) \mathrm{d} S_{\xi}(u) = h^{-(m_R + m_L - 1)} e^{i\frac{S(x)}{h}} \left(\sum_{\ell=0}^{N-1} h^{\ell} \int_{\mathbb{S}_{\xi}^{n-2}} A_{2\ell}(x, u, D_{\varphi}, D_t) F\left(x, u, \frac{\pi}{2}, 0\right) \mathrm{d} S_{\xi}(u) + h^N R_N(x) \right).$$

$$(4.9)$$

Here, for any $\ell \in \mathbb{N}$, $A_{2\ell}(x, u, D_{\varphi}, D_t)$ is a differential operator of degree $\leq 2\ell$ depending smoothly on x and u and R'_N satisfies the bound

$$||R'_N||_{C^0(N_{\varepsilon} \times \mathbb{S}^{n-2}_{\xi})} \le C_N ||F||_{C^{2N+3}(SN_{\varepsilon})},$$

where SU denotes the unit tangent bundle of U (where F is defined). The order 2N + 3 = 2N + 2 + 1 comes from the remainder term in [Zwo12, Theorem 3.16]. After integration in the variable u, i.e. setting $R_N(x) = \int_{\mathbb{S}_{\epsilon}^{n-2}} R'_N(x, u) \, \mathrm{d} S_{\xi}(u)$, this gives:

$$||R_N||_{C^0(N_{\varepsilon})} \le C_N ||F||_{C^{2N+3}(SN_{\varepsilon})}, \tag{4.10}$$

and one can control higher order derivatives of R_N in the same fashion (up to increasing the order of the norm on the right-hand side of (4.10)).

Next, observe that by definition:

$$F(x, u, \varphi, t) = \sin^{n-2}(\varphi) \sum_{k=0}^{m_L} P_L^{(k)} \left(e^{tX} \sum_{j=0}^{m_R} P_R^{(j)}(\pi_0^* f, \pi_0^* S) h^j, e^{tX} \pi_0^* S \right) (x, v(u, \varphi)) h^k,$$

which implies that the C^k -norms of F are controlled by the $C^{k'}$ -norms of f and S (for some $k' \gg k$), that is the remainder R_N is indeed negligible in the sense of (4.6). Also note that $A_{2\ell}(x, u, D_{\varphi}, D_t)F(x, u, \frac{\pi}{2}, 0)$ depends only on a finite number $K(\ell)$ of derivatives of the function f and the phase S at x. Hence (4.9) shows that the operator is pseudodifferential. \Box

5. GENERIC INJECTIVITY WITH RESPECT TO THE CONNECTION

We now prove Theorem 1.8 in this section. This case is much less technical than the metric case but still provides a good insight on the argument. In what follows, differentiation will be mostly carried out without recalling that the objects depend smoothly on the parameter and we refer the reader to $\S2.4$, Lemma 2.3 for further details.

5.1. **Preliminary remarks.** Consider an Anosov Riemannian manifold (M, g) with a Hermitian vector bundle $\mathcal{E} \to M$, equipped with a unitary connection $\nabla^{\mathcal{E}}$. Consider a linear perturbation $\nabla^{\mathcal{E}} + \tau\Gamma$ for some skew-Hermitian $\Gamma \in C^{\infty}(M, T^*M \otimes \operatorname{End}_{\operatorname{sk}}(\mathcal{E}))$ and $\tau \in \mathbb{R}$, and

the operator $\mathbf{X}_{\tau} := \pi^* (\nabla^{\mathcal{E}} + \tau \Gamma)_X$, where we recall that $\pi : SM \to M$ is the footpoint projection. We set $\mathbf{X} := \mathbf{X}_0$. For the sake of simplicity, let us assume that ker $\mathbf{X}|_{C^{\infty}(SM,\mathcal{E})} = \{0\}^{12}$, which is generically true by [CL21a]. We will consider the operator

$$P_{\tau} := \pi_{\ker D_0^*} \Delta_0 \Pi_m^{\tau} \Delta_0 \pi_{\ker D_0^*},$$

where $\Pi_m^{\tau} = \pi_{m*}(\mathbf{R}_+^{\tau} + \mathbf{R}_-^{\tau})\pi_m^*$, Δ_0 is an elliptic, formally self-adjoint, positive, pseudodifferential operator with diagonal principal symbol of order k > 1/2 that induces an isomorphism on Sobolev spaces H^s for any s and D_{τ} denotes the twisted (with respect to $\nabla^{\mathcal{E}} + \tau\Gamma$) symmetric derivative on tensors, as in §2.2.3. Here $\mathbf{R}_{\pm}^{\tau} := -\mathbf{R}_{\pm}^{\tau}(z=0)$ is the (opposite of) resolvent at zero of \mathbf{X}_{τ} as defined in §2.3.3. It is important to note that

$$\pi_{\ker D^*_{\tau}}\Pi^{\tau}_m = \Pi^{\tau}_m \pi_{\ker D^*_{\tau}} = \Pi^{\tau}_m,$$

and that by continuity we have:

Lemma 5.1. For $|\tau|$ small enough, the map $\tau \mapsto \pi_{\ker D^*_{\tau}} \in \Psi^0(M, \otimes^m_S T^*M \otimes \mathcal{E})$ is continuous. Moreover, for all $s \in \mathbb{R}$, there exists $\varepsilon = \varepsilon(s) > 0$ and C = C(s) > 0 such that for $|\tau| < \varepsilon$:

$$\forall f \in H^{s+k}(M, \otimes_S^m T^*M \otimes \mathcal{E}), \ \|\pi_{\ker D^*_\tau} \Delta_0 \pi_{\ker D^*_\tau} f\|_{H^s} \ge C \|\pi_{\ker D^*_\tau} f\|_{H^{s+k}}.$$
(5.1)

Hence for any $s \in \mathbb{R}$, for all $|\tau| < \varepsilon$, where $\varepsilon = \varepsilon(s) > 0$ is small enough, the following maps are isomorphisms:

$$\pi_{\ker D^*_{\tau}} \Delta_0 \pi_{\ker D^*_0} : \ker D^*_0 \cap H^{s+k} \to \ker D^*_{\tau} \cap H^s,$$

$$\pi_{\ker D^*_0} \Delta_0 \pi_{\ker D^*_{\tau}} : \ker D^*_{\tau} \cap H^{s+k} \to \ker D^*_0 \cap H^s.$$

Proof. The first claim is obvious from the formula (see (2.10)):

$$\pi_{\ker D^*_{\tau}} = \mathbb{1} - D_{\tau} (D^*_{\tau} D_{\tau})^{-1} D^*_{\tau}$$

Here it is important that ker $D_{\tau}|_{H^s} = \{0\}$ for any $s \in \mathbb{R}$ which follows from the fact that D_{τ} is elliptic and the following observation: if $D_0 f = 0$, then $\mathbf{X} \pi_m^* f = 0$ by (2.15) and this implies f = 0 by our assumptions, so the map $(D_{\tau}^* D_{\tau})^{-1}$ is continuous with respect to τ^{13} .

Next, since Δ_0 is an isomorphism on Sobolev spaces H^s , we have that $\|\Delta_0 f\|_{H^s} \ge C_s \|f\|_{H^{s+k}}$ for some $C_s > 0$. Using the identity

$$\pi_{\ker D_{\tau}^{*}} \Delta_{0} \pi_{\ker D_{\tau}^{*}} = [\pi_{\ker D_{\tau}^{*}} - \pi_{\ker D_{0}^{*}}, \Delta_{0}] \pi_{\ker D_{\tau}^{*}} + [\pi_{\ker D_{0}^{*}}, \Delta_{0}] \pi_{\ker D_{\tau}^{*}} + \Delta_{0} \pi_{\ker D_{\tau}^{*}},$$

as well as that $[\pi_{\ker D_0^*} - \pi_{\ker D_\tau^*}, \Delta_0] = o_{\Psi^{k-1}}(1)$, we obtain the estimate

$$\|\pi_{\ker D^*_{\tau}}f\|_{H^{s+k}} \le C_s \|\Delta_0 \pi_{\ker D^*_{\tau}}f\|_{H^s} \le C'(\|\pi_{\ker D^*_{\tau}}\Delta_0 \pi_{\ker D^*_{\tau}}f\|_{H^s} + \|\pi_{\ker D^*_{\tau}}f\|_{H^{s+k-1}})$$
(5.2)

¹²The following arguments can be generalized to the case where ker **X** consists of *stable* elements of degree 0 (equivalently, we will say that ker **X** is stably non-empty): by stable, we mean that any perturbation of the operator will still have the same resonant space at 0 and that this space only contains elements of degree 0. This is the case for the operator X acting on functions as it always has $\mathbb{C} \cdot \mathbf{1}$ (the constant sections) as resonant space at z = 0; this is also the case for $(\pi^* \nabla^{\operatorname{End}(\mathcal{E})})_X$ as it always contains $\mathbb{C} \cdot \mathbb{1}_{\mathcal{E}}$ and is generically equal to $\mathbb{C} \cdot \mathbb{1}_{\mathcal{E}}$ by [CL21a] (where $\nabla^{\operatorname{End}(\mathcal{E})}$ is the induced connection on $\operatorname{End}(\mathcal{E})$). Instead of taking the resolvent at 0, one needs to work with the holomorphic part of the resolvent. This is done in the metric case, see §6. For the sake of simplicity, we assume in this section that ker **X** is trivial.

 $^{^{13}}$ If ker X is not empty but stably non-empty, this also works.

for some constant C' = C'(s) > 0 independent of τ . To show (5.1), we argue by contradiction and assume there is a sequence $f_n \in \ker D^*_{\tau_n}$ with $||f_n||_{H^{s+k}} = 1$ and $||\pi^*_{\ker D_{\tau_n}} \Delta_0 f_n||_{H^s} \to 0$. We assume $\tau_n \to 0$ (but the same argument works if $\tau_n \to \tau$ for some $\tau \neq 0$). By compactness, we may assume f_n converges in H^{s+k-1} . In fact, we have:

$$\begin{split} \|f_n - f_m\|_{H^{s+k}} &\leq \|\pi_{\ker D^*_{\tau_n}}(f_n - f_m)\|_{H^{s+k}} + \|(\pi_{\ker D^*_{\tau_n}} - \pi_{\ker D^*_{\tau_m}})f_m\|_{H^{s+k}} \\ &\leq C'\|\pi_{\ker D^*_{\tau_n}}\Delta_0\pi_{\ker D^*_{\tau_n}}f_m\|_{H^s} + C'\|\pi_{\ker D^*_{\tau_n}}(f_n - f_m)\|_{H^{s+k-1}} + o(1) \\ &\leq \|\pi_{\ker D^*_{\tau_m}}\Delta_0f_m\|_{H^s} + \|(\pi_{\ker D^*_{\tau_n}} - \pi_{\ker D^*_{\tau_m}})\Delta_0f_m\|_{H^s} \\ &+ \|\pi_{\ker D^*_{\tau_n}}\Delta_0(\pi_{\ker D^*_{\tau_n}} - \pi_{\ker D^*_{\tau_m}})f_m\|_{H^s} + o(1) = o(1). \end{split}$$

as $m, n \to \infty$. In the second line, we used that $\pi_{\ker D_{\tau_n}^*} - \pi_{\ker D_{\tau_m}^*} = o_{\Psi^0}(1)$ and $||f_m||_{H^{s+k}} = 1$, (5.2), and the fact that $||\pi_{\ker D_{\tau_n}}^* \Delta_0 f_n||_{H^s} = o(1)$. In the last line, we also used the assumption that f_n converges in H^{s+k-1} . Therefore, $(f_n)_{n\in\mathbb{N}}$ is a Cauchy sequence in H^{s+k} and it converges to some $f \in H^{s+k}$ with $||f||_{H^{s+k}} = 1$, $D_0^*f = 0$ and $\pi_{\ker D_0^*}\Delta_0 f = 0$. Using that $[\pi_{\ker D_0^*}, \Delta_0]f = -\Delta_0 f$ and the fact that $[\pi_{\ker D_0^*}, \Delta_0] \in \Psi^{k-1}$ implies by elliptic regularity that $f \in H^{s+k+1}$. Bootstrapping we get $f \in C^{\infty}$ and so

$$0 = \langle \pi_{\ker D_0^*} \Delta_0 f, \Delta_0 f \rangle_{L^2} = \langle \Delta_0 f, f \rangle_{L^2},$$

which means that f = 0 as Δ_0 was chosen to be positive. This contradicts that $||f||_{H^{s+k}} = 1$ and proves (5.1).

Finally, by the first point we have $\|\pi_{\ker D_0^*} - \pi_{\ker D_\tau^*}\|_{H^s \to H^s} = o_s(1)$ as $\tau \to 0$, so by (5.1) for small $|\tau|$ depending on s we get

$$\begin{aligned} \|\pi_{\ker D_{\tau}^{*}}\Delta_{0}\pi_{\ker D_{0}^{*}}f\|_{H^{s}} &\geq \|\pi_{\ker D_{0}^{*}}\Delta_{0}\pi_{\ker D_{0}^{*}}f\|_{H^{s}} - \|(\pi_{\ker D_{0}^{*}} - \pi_{\ker D_{\tau}^{*}})\Delta_{0}\pi_{\ker D_{0}^{*}}f\|_{H^{s}} \\ &\geq \frac{C(s)}{2}\|\pi_{\ker D_{0}^{*}}f\|_{H^{s+k}}. \end{aligned}$$
(5.3)

Similarly, using (5.1) for $|\tau|$ small enough we obtain:

$$\|\pi_{\ker D^*_{\tau}}\Delta_0\pi_{\ker D^*_0}f\|_{H^s} \ge \frac{C(s)}{2}\|\pi_{\ker D^*_0}f\|_{H^{s+k}}.$$
(5.4)

Estimates (5.3) and (5.4) show that the operators $\pi_{\ker D_0^*} \Delta_0 \pi_{\ker D_\tau^*}$, $\pi_{\ker D_\tau^*} \Delta_0 \pi_{\ker D_0^*}$ are injective and have a closed range for $|\tau|$ small enough, and then the surjectivity follows since their L^2 -adjoints are injective. This completes the proof.

Next, using Proposition 4.1, the fact that Δ_0 acts diagonally to principal order and the equation (2.11), we obtain that for $\xi \in T_x^* M \setminus \{0\}$:

$$\sigma(P_{\tau})(x,\xi) = \frac{2\pi}{|\xi|} C_{2m+n-1}^{-1} \sigma(\Delta_0)^2(x,\xi) \pi_{\ker_{i_{\xi}\sharp}} \pi_{m*} \pi_m^* \pi_{\ker_{i_{\xi}\sharp}} \otimes \mathbb{1}_{\mathcal{E}_x}$$

Therefore, the symbol of P_{τ} at (x, ξ) is invertible on $\ker_{i_{\xi^{\sharp}}}$ and by standard microlocal analysis for each τ there exist pseudodifferential operators Q_{τ} and R_{τ} of respective orders 1 - 2k and $-\infty$ (cf. [Lef19b, Lemma 2.5.3]), such that

$$Q_{\tau}P_{\tau} = \pi_{\ker D_0^*} + R_{\tau}.$$

Using that $\Pi_m^{\tau} \ge 0$ we get $P_{\tau} \ge 0$ and it follows that $(P_{\tau} + 1)^{-1} : L^2 \cap \ker D_0^* \to L^2 \cap \ker D_0^*$ is compact and thus the spectrum of P_{τ} is well-defined. It is discrete, non-negative and accumulates at infinity, and the eigenfunctions of P_{τ} are smooth. Moreover, by Lemma 5.1 and again using that $\Pi_m^{\tau} \ge 0$, for small $|\tau|$ we have that

$$\pi_{\ker D^*_{\tau}} \Delta_0 \pi_{\ker D^*_0} : \ker P_{\tau}|_{C^{\infty} \cap \ker D^*_0} \xrightarrow{\sim} \ker \Pi^{\tau}_m|_{C^{\infty} \cap \ker D^*_0}$$

is an isomorphism. Therefore we see that for each τ , 0 is an eigenvalue of Π_m^{τ} if and only if 0 is an eigenvalue of P_{τ} .

5.2. Variations of the ground state. We assume that $\ker P_0|_{C^{\infty} \cap \ker D_0^*}$ is *d*-dimensional, for some $d \geq 1$, and spanned by $u_1, ..., u_d \in C^{\infty}(M, \bigotimes_S^m T^*M \otimes \mathcal{E}) \cap \ker D_0^*$ with $\langle u_i, u_j \rangle_{L^2} = \delta_{ij}$. Let Π_{τ} be the L^2 -orthogonal spectral projector

$$\Pi_{\tau} := \frac{1}{2\pi i} \oint_{\gamma'} (z - P_{\tau})^{-1} \,\mathrm{d}\,z, \tag{5.5}$$

where γ' is a small circle centred around 0 and not containing any other eigenvalue of P_0 in its interior. In particular, we have $\Pi_{\tau=0} = \sum_{i=1}^{d} \langle \bullet, u_i \rangle_{L^2} u_i$ and by ellipticity of P_0 on ker D_0^* we have the meromorphic expansion close to zero, valid on $S := L^2 \cap \ker D_0^*$

$$(z - P_0)^{-1} = \frac{\Pi_0}{z} - P_0^{-1} + zH_1 + \mathcal{O}(z^2), \qquad (5.6)$$

for some maps $P_0^{-1}, H_1: S \to S$. These maps satisfy the relations (cf. (2.18)):

$$P_0 P_0^{-1} = P_0^{-1} P_0 = \text{Id} - \Pi_0, \quad P_0 H_1 = -P_0^{-1}, \quad \Pi_0 P_0^{-1} = P_0^{-1} \Pi_0 = 0.$$
(5.7)

We introduce λ_{τ} as the sum of the eigenvalues of P_{τ} inside γ' :

$$\lambda_{\tau} := \operatorname{Tr}(P_{\tau} \Pi_{\tau}). \tag{5.8}$$

Note that both $\tau \mapsto \Pi_{\tau} \in \mathcal{L}(L^2)$ and $\tau \mapsto \lambda_{\tau} \in \mathbb{C}$ are smooth by standard elliptic theory (see [CL21a, Section 4]). Observe that $\lambda_{\tau=0} = 0$ and as $P_{\tau} \ge 0$, we have $\lambda_{\tau} \ge 0$. Our goal is to produce a small perturbation $\nabla^{\mathcal{E}} + \tau \Gamma$ (where Γ is skew-Hermitian) such that $\lambda_{\tau} > 0$ for $\tau \ne 0$. This will say that at least one of the eigenvalues was ejected from 0 and that ker P_{τ} is at most (d-1)-dimensional (for $\tau \ne 0$). Iterating the process, we will then obtain a perturbation of $\nabla^{\mathcal{E}}$ with injective (twisted) X-ray transform.

We make the easy observation that the first variation is zero, as $\lambda_{\tau=0} = 0$ is a local minimum of the smooth function $\tau \mapsto \lambda_{\tau}$:

$$\dot{\lambda}_{\tau=0} = 0. \tag{5.9}$$

Next, we note that since $P_{\tau}u_i = 0$ and $\Pi_m^{\tau} \ge 0$, we have $\Pi_m^{\tau}\Delta_0 u_i = 0$. Therefore, by Lemma 2.2, there exists $v_i \in C^{\infty}(SM, \mathcal{E})$ such that

$$\pi_m^* \Delta_0 u_i = \mathbf{X} \mathbf{v}_i, \quad \Pi_+ \mathbf{v}_i = 0.$$
(5.10)

By the mapping properties of \mathbf{X} (see (2.16)), we have that if m is even then v_i may be chosen odd and vice versa, if m is odd then v_i may be chosen even.

5.2.1. Second order variations. We now compute $\ddot{\lambda}$ at $\tau = 0$. We start with an abstract lemma, valid in a more general setting (this will also get used in the metric case, see Lemma 6.9 below):

Lemma 5.2. The following variational formula holds:

$$\ddot{\lambda} = \operatorname{Tr}(\ddot{P}_0 \Pi_0) - 2 \operatorname{Tr}(\Pi_0 \dot{P}_0 P_0^{-1} \dot{P}_0 \Pi_0).$$

Proof. We compute:

$$\begin{split} \dot{\Pi}_0 &= \frac{1}{2\pi i} \oint_{\gamma'} (z - P_0)^{-1} \dot{P}_0 (z - P_0)^{-1} dz = -\left(\Pi_0 \dot{P}_0 P_0^{-1} + P_0^{-1} \dot{P}_0 \Pi_0\right), \\ \ddot{\Pi}_0 &= 2 \times \frac{1}{2\pi i} \oint_{\gamma'} (z - P_0)^{-1} \dot{P}_0 (z - P_0)^{-1} \dot{P}_0 (z - P_0)^{-1} dz + \frac{1}{2\pi i} \oint_{\gamma'} (z - P_0)^{-1} \ddot{P}_0 (z - P_0)^{-1} dz. \end{split}$$

We expand the second formula using (5.6) to get:

$$\ddot{\Pi}_{0} = 2 \left[\Pi_{0} \dot{P}_{0} \Pi_{0} \dot{P}_{0} H_{1} + \Pi_{0} \dot{P}_{0} P_{0}^{-1} \dot{P}_{0} P_{0}^{-1} + \Pi_{0} \dot{P}_{0} H_{1} \dot{P}_{0} \Pi_{0} + P_{0}^{-1} \dot{P}_{0} \Pi_{0} \dot{P}_{0} P_{0}^{-1} + P_{0}^{-1} \dot{P}_{0} \Pi_{0} \dot{P}_{0} \Pi_{0} + H_{1} \dot{P}_{0} \Pi_{0} \dot{P}_{0} \Pi_{0} \right] - \left(\Pi_{0} \ddot{P}_{0} P_{0}^{-1} + P_{0}^{-1} \ddot{P}_{0} \Pi_{0} \right).$$
(5.11)

Therefore, we compute using (5.7):

$$P_0 \ddot{\Pi}_0 = 2(\mathrm{Id} - \Pi_0) \dot{P}_0 \left(\Pi_0 \dot{P}_0 P_0^{-1} + P_0^{-1} \dot{P}_0 \Pi_0 \right) - 2P_0^{-1} \dot{P}_0 \Pi_0 \dot{P}_0 \Pi_0 - (\mathrm{Id} - \Pi_0) \ddot{P}_0 \Pi_0, \quad (5.12)$$

which implies that, using the cyclicity of the trace and (5.7):

$$\operatorname{Tr}\left(P_{0}\ddot{\Pi}_{0}\right) = 2\operatorname{Tr}\left(\dot{P}_{0}\Pi_{0}\dot{P}_{0}P_{0}^{-1}\right) = 2\operatorname{Tr}\left(\Pi_{0}\dot{P}_{0}P_{0}^{-1}\dot{P}_{0}\Pi_{0}\right).$$

Finally, we obtain using once more the cyclicity of the trace:

$$\ddot{\lambda} = \text{Tr}\left(\ddot{P}_0 \Pi_0 + 2\dot{P}_0 \dot{\Pi}_0 + P_0 \ddot{\Pi}_0\right) = \text{Tr}\left(\ddot{P}_0 \Pi_0\right) - 2 \text{Tr}\left(\Pi_0 \dot{P}_0 P_0^{-1} \dot{P}_0 \Pi_0\right).$$

Next, we compute \dot{P}_0 , \ddot{P}_0 and apply the formula (5.14). Before doing that, note that by Lemma 5.1, on ker D_0^* we have

$$P_0^{-1} = (\pi_{\ker D_0^*} \Delta_0 \pi_{\ker D_0^*})^{-1} \Pi_m^{-1} (\pi_{\ker D_0^*} \Delta_0 \pi_{\ker D_0^*})^{-1},$$
(5.13)

where $-\Pi_m^{-1}$ stands for the holomorphic part of the resolvent at zero, i.e. if $(z - \Pi_m)^{-1} = \frac{\Pi_{m,0}}{z} + R_m(z)$, where $\Pi_{m,0}$ is the orthogonal projection onto ker $\Pi_m|_{\ker D_0^*}$ zero and $R_m(z)$ holomorphic close to z = 0 then $-\Pi_m^{-1} := R_m(0)$.

Lemma 5.3. We have:

$$\ddot{\lambda} = 2\sum_{i=1}^{d} \left(\left\langle \mathcal{I}_{\nabla} \varepsilon \left(\pi_{1}^{*} \Gamma. \mathbf{v}_{i} \right), \pi_{1}^{*} \Gamma. \mathbf{v}_{i} \right\rangle_{L^{2}} - \left\langle \Pi_{m}^{-1} \pi_{m*} \mathcal{I}_{\nabla} \varepsilon \left(\pi_{1}^{*} \Gamma. \mathbf{v}_{i} \right), \pi_{m*} \mathcal{I}_{\nabla} \varepsilon \left(\pi_{1}^{*} \Gamma. \mathbf{v}_{i} \right) \right\rangle_{L^{2}} \right).$$
(5.14)

Proof. We start with the first term in (5.14). For the variation of the resolvent, recalling that $\mathbf{R}_{-} = -\mathbf{X}^{-1}$ and $\mathbf{R}_{+} = \mathbf{X}^{-1}$ and the notation of §2.3.3 (γ is a small loop around zero), we have:

$$\ddot{\mathbf{R}}_{+} = \partial_{\tau}^{2}|_{\tau=0} \frac{1}{2\pi i} \oint_{\gamma} \frac{1}{z} (z + \mathbf{X}_{\tau})^{-1} dz = \partial_{\tau}^{2}|_{\tau=0} \mathbf{X}_{\tau}^{-1} = 2 \,\mathbf{R}_{+} \,\pi_{1}^{*} \Gamma \,\mathbf{R}_{+} \,\pi_{1}^{*} \Gamma \,\mathbf{R}_{+},$$

$$\ddot{\mathbf{R}}_{-} = (\ddot{\mathbf{R}}_{+})^{*} = \partial_{\tau}^{2}|_{\tau=0} \frac{1}{2\pi i} \oint_{\gamma} \frac{1}{z} (z - \mathbf{X}_{\tau})^{-1} dz = \partial_{\tau}^{2}|_{\tau=0} (-\mathbf{X}_{\tau})^{-1} = 2 \,\mathbf{R}_{-} \,\pi_{1}^{*} \Gamma \,\mathbf{R}_{-} \,\pi_{1}^{*} \Gamma \,\mathbf{R}_{-} \,.$$

(5.15)

Therefore

$$\ddot{P}_{0} = 2\pi_{\ker D_{0}^{*}}\Delta_{0}\pi_{m*}\left(\mathbf{R}_{+}\pi_{1}^{*}\Gamma\,\mathbf{R}_{+}\pi_{1}^{*}\Gamma\,\mathbf{R}_{+} + \mathbf{R}_{-}\pi_{1}^{*}\Gamma\,\mathbf{R}_{-}\pi_{1}^{*}\Gamma\,\mathbf{R}_{-}\right)\pi_{m}^{*}\Delta_{0}\pi_{\ker D_{0}^{*}}$$

and we obtain, using that $\mathbf{X} \mathbf{v}_i = \pi_m^* \Delta_0 u_i$ (see (5.10)):

$$\operatorname{Tr}(\ddot{P}_{0}\Pi_{0}) = 2\sum_{i=1}^{d} \left\langle \left(\mathbf{R}_{+} \pi_{1}^{*}\Gamma \,\mathbf{R}_{+} \pi_{1}^{*}\Gamma \,\mathbf{R}_{+} + \mathbf{R}_{-} \pi_{1}^{*}\Gamma \,\mathbf{R}_{-} \pi_{1}^{*}\Gamma \,\mathbf{R}_{-} \right) \mathbf{X} \,\mathbf{v}_{i}, \mathbf{X} \,\mathbf{v}_{i} \right\rangle_{L^{2}}$$

$$= 2\sum_{i=1}^{d} \left\langle \mathcal{I}_{\nabla}\varepsilon(\pi_{1}^{*}\Gamma \cdot \mathbf{v}_{i}), \pi_{1}^{*}\Gamma \cdot \mathbf{v}_{i} \right\rangle_{L^{2}},$$

$$(5.16)$$

using (2.18) in the second line, as well as that Γ is skew-Hermitian.

For the second term of (5.14), we first observe that similarly to (5.15)

$$\dot{\mathbf{R}}_{+} = \partial_{\tau}|_{\tau=0} \mathbf{X}_{\tau}^{-1} = -\mathbf{R}_{+} \pi_{1}^{*} \Gamma \mathbf{R}_{+}, \quad \dot{\mathbf{R}}_{-} = \partial_{\tau}|_{\tau=0} (-\mathbf{X}_{\tau})^{-1} = \mathbf{R}_{-} \pi_{1}^{*} \Gamma \mathbf{R}_{-}.$$

Therefore it holds that

$$\dot{P}_{0} = \pi_{\ker D_{0}^{*}} \Delta_{0} \pi_{m*} (-\mathbf{R}_{+} \pi_{1}^{*} \Gamma \mathbf{R}_{+} + \mathbf{R}_{-} \pi_{1}^{*} \Gamma \mathbf{R}_{-}) \pi_{m}^{*} \Delta_{0} \pi_{\ker D_{0}^{*}}, \qquad (5.17)$$

and we finally obtain that, using $(\dot{P}_0)^* = \dot{P}_0$:

$$\operatorname{Tr}\left(\Pi_{0}\dot{P}_{0}P_{0}^{-1}\dot{P}_{0}\Pi_{0}\right) = \sum_{i=1}^{d} \left\langle P_{0}^{-1}\dot{P}_{0}u_{i},\dot{P}_{0}u_{i}\right\rangle_{L^{2}}$$
$$= \sum_{i=1}^{d} \left\langle P_{0}^{-1}\pi_{\ker D_{0}^{*}}\Delta_{0}\mathcal{I}_{\nabla \mathcal{E}}\left(\pi_{1}^{*}\Gamma.\,\mathbf{v}_{i}\right),\pi_{\ker D_{0}^{*}}\Delta_{0}\mathcal{I}_{\nabla \mathcal{E}}\left(\pi_{1}^{*}\Gamma.\,\mathbf{v}_{i}\right)\right\rangle_{L^{2}} \quad (5.18)$$
$$= \sum_{i=1}^{d} \left\langle \Pi_{m}^{-1}\pi_{m*}\mathcal{I}_{\nabla \mathcal{E}}(\pi_{1}^{*}\Gamma.\,\mathbf{v}_{i}),\pi_{m*}\mathcal{I}_{\nabla \mathcal{E}}(\pi_{1}^{*}\Gamma.\,\mathbf{v}_{i})\right\rangle_{L^{2}}.$$

Here in the second line we used (5.17) and (5.10); in the final line we used (5.13). This proves the announced result.

5.2.2. Properties of the operators involved in the second variation. First of all we note that by Proposition 4.1, Π_m and hence $\Pi_m^{-1} \pi_{\ker D_0^*}$, are pseudodifferential operators of orders -1 and 1, respectively (recall Π_m^{-1} was defined just below (5.13)). It follows that, for $(x,\xi) \in T^*M \setminus 0$ (cf. [Lef19b, Lemma 2.5.3]):

$$\sigma_{\Pi_{m}}(x,\xi) = C_{n-1+2m}^{-1} \frac{2\pi}{|\xi|} \pi_{\ker \imath_{\xi^{\sharp}}} \pi_{m*} \pi_{m}^{*} \pi_{\ker \imath_{\xi^{\sharp}}} \otimes \mathbb{1}_{\mathcal{E}_{x}},$$

$$\sigma_{\Pi_{m}^{-1}\pi_{\ker D_{0}^{*}}}(x,\xi) = C_{n-1+2m} \frac{|\xi|}{2\pi} (\pi_{\ker \imath_{\xi^{\sharp}}} \pi_{m*} \pi_{m}^{*} \pi_{\ker \imath_{\xi^{\sharp}}})^{-1} \pi_{\ker \imath_{\xi^{\sharp}}} \otimes \mathbb{1}_{\mathcal{E}_{x}}.$$
(5.19)

We now fix $\mathbf{v} \in C^{\infty}(SM, \pi^* \mathcal{E})$ and introduce the multiplication map

$$M_{\mathbf{v}}: C^{\infty}(SM, \pi^* \operatorname{End}(\mathcal{E})) \to C^{\infty}(SM, \pi^* \mathcal{E}), \quad M_{\mathbf{v}}A := A. \mathbf{v}.$$

Its adjoint is given by $M_v^* w = \langle \bullet, v \rangle_{\mathcal{E}} \otimes w$, for any $w \in C^{\infty}(SM, \pi^* \mathcal{E})$. Next, we show that the terms appearing in the formula for $\ddot{\lambda}$ in Lemma 5.3 have pseudodifferential nature:

Lemma 5.4. For all $v \in C^{\infty}(SM, \pi^* \mathcal{E})$, the operator

$$Q_{\mathbf{v}} := \pi_{m*} \mathcal{I}_{\nabla} \varepsilon M_{\mathbf{v}} \pi_1^* \in \Psi^{-1} \left(M, T^* M \otimes \operatorname{End}(\mathcal{E}) \to \bigotimes_S^m T^* M \otimes \mathcal{E} \right)$$
(5.20)

is pseudodifferential of order -1 with principal symbol for $(x,\xi) \in T^*M \setminus \{0\}$

 $\sigma_{Q_{\mathbf{v}}}(x,\xi) \in \operatorname{Hom}(T_x^*M \otimes \operatorname{End}(\mathcal{E}_x), \otimes_S^m T_x^*M \otimes \mathcal{E}_x),$

given by, for $B \in T_x^* M \otimes \operatorname{End}(\mathcal{E}_x)$:

$$\sigma_{Q_{\mathbf{v}}}(x,\xi)B = C_{n-1+2m}^{-1} \frac{2\pi}{|\xi|} \pi_{\ker \imath_{\xi^{\sharp}}} \pi_{m*} E_{\xi}^{m} \left(\pi_{1}^{*}(\pi_{\ker \imath_{\xi^{\sharp}}}B). \mathbf{v}(x,\bullet) \right).$$

Proof. This follows directly from Proposition 4.1.

Lemma 5.5. For any $v \in C^{\infty}(SM, \pi^* \mathcal{E})$, the operator

$$L_{\mathbf{v}} := \pi_{1*} M_{\mathbf{v}}^* \mathcal{I}_{\nabla \mathcal{E}} M_{\mathbf{v}} \pi_1^* \in \Psi^{-1}(M, T^*M \otimes \operatorname{End}(\mathcal{E}) \to T^*M \otimes \operatorname{End}(\mathcal{E}))$$
(5.21)

is pseudodifferential of order -1 with principal symbol for $(x,\xi) \in T^*M \setminus \{0\}$

$$\sigma_{L_{\mathbf{v}}}(x,\xi) \in \operatorname{End}(T_x^*M \otimes \operatorname{End}(\mathcal{E}_x))$$

given by:

$$\sigma_{L_{v}}(x,\xi) = C_{n-1+2m}^{-1} \frac{2\pi}{|\xi|} \pi_{\ker \imath_{\xi^{\sharp}}} \pi_{1*} M_{v}^{*} (E_{\xi}^{m})^{*} E_{\xi}^{m} M_{v} \pi_{1}^{*} \pi_{\ker \imath_{\xi^{\sharp}}}$$

Proof. Once more, this follows from Proposition 4.1 and the formula, for $B \in T_x^* M \otimes \operatorname{End}(\mathcal{E}_x)$:

$$\langle \sigma_{L_{\mathbf{v}}}(x,\xi)B,B \rangle_{x} = \frac{2\pi}{|\xi|} \int_{\mathbb{S}_{\xi}^{n-2}} \left\langle \pi_{1}^{*}B(u).\,\mathbf{v}(x,u),\pi_{1}^{*}B(u).\,\mathbf{v}(x,u) \right\rangle_{x} \mathrm{d}\,S_{\xi}(u)$$

$$= C_{n-1+2m}^{-1} \frac{2\pi}{|\xi|} \left\langle E_{\xi}^{m}(\pi_{1}^{*}B.\,\mathbf{v}), E_{\xi}^{m}(\pi_{1}^{*}B.\,\mathbf{v}) \right\rangle_{L^{2}(\mathbb{S}^{n-1},\pi^{*}\mathcal{E}_{x})}$$

$$= C_{n-1+2m}^{-1} \frac{2\pi}{|\xi|} \left\langle \pi_{1*}M_{\mathbf{v}}^{*}(E_{\xi}^{m})^{*}E_{\xi}^{m}M_{\mathbf{v}}\pi_{1}^{*}B,B \right\rangle_{L^{2}}.$$

In the second line we used the Jacobian formula (3.1).

5.3. Assuming the second variation is zero. If $\ddot{\lambda} = 0$ for all linear variations $\nabla^{\mathcal{E}} + \tau \Gamma$ as in §5.2, where Γ is skew-Hermitian, by Lemma 5.3 and using the notation of (5.20), (5.21), this implies:

$$\forall \Gamma \in C^{\infty}(M, T^*M \otimes \operatorname{End}_{\operatorname{sk}}(\mathcal{E})), \quad \sum_{i=1}^d \langle L_{v_i}\Gamma, \Gamma \rangle_{L^2} = \sum_{i=1}^d \langle \Pi_m^{-1}Q_{v_i}\Gamma, Q_{v_i}\Gamma \rangle_{L^2}.$$
(5.22)

The idea is to apply the equality (5.22) (which is of *analytic* nature) to Gaussian states in order to derive an *algebraic* equality. Let $e_h(x_0, \xi_0)$ be a Gaussian state centered at $(x_0, \xi_0) \in T^*M \setminus \{0\}$, that is, a function which has the form in some local coordinates around x_0^{14} :

$$e_h(x_0,\xi_0)(x) = \frac{1}{(\pi h)^{\frac{n}{4}}} e^{\frac{i}{h}\xi_0 \cdot (x-x_0) - \frac{|x-x_0|^2}{2h}}.$$
(5.23)

We will use the following standard technical lemma:

Lemma 5.6. Let $P \in \Psi^m(M, E \to F)$ be a pseudodifferential operator of order $m \in \mathbb{R}$ acting on two Hermitian vector bundles $E, F \to M$. Let $e \in C^{\infty}(M, E)$ and $f \in C^{\infty}(M, F)$. Then:

$$\begin{split} \lim_{h \to 0} \Big\langle h^m P\Big(\Re(\mathbf{e}_h(x_0,\xi_0)).e \Big), \Re(\mathbf{e}_h(x_0,\xi_0)).f \Big\rangle_{L^2(M,F)} &= \frac{1}{2} \Big\langle \sigma_P(x_0,\xi_0)e(x_0), f(x_0) \big\rangle_{F_{x_0}} \\ &\quad + \frac{1}{2} \big\langle \sigma_P(x_0,-\xi_0)e(x_0), f(x_0) \big\rangle_{F_{x_0}}. \end{split}$$

We cannot directly apply (5.22) to $e_h(x_0, \xi_0)$. Γ where $\Gamma \in C^{\infty}(M, T^*M \otimes \operatorname{End}_{\operatorname{sk}}(\mathcal{E}))$ because $e_h(x_0, \xi_0)$. Γ is not skew-Hermitian. However, applying (5.22) to $\Re(e_h(x_0, \xi_0))$. Γ and using Lemma 5.6, as well as the principal symbol formulas in (5.19), Lemmas 5.4 and 5.5 (and that in particular these symbols are invariant under the antipodal map $(x, \xi) \mapsto (x, -\xi)$), taking $h \to 0$ we obtain:

$$\sum_{i=1}^{d} \left\langle E_{\xi_{0}}^{m}(\pi_{1}^{*}\Gamma.v_{i}), E_{\xi_{0}}^{m}(\pi_{1}^{*}\Gamma.v_{i}) \right\rangle_{L^{2}(\mathbb{S}^{n-1},\pi^{*}\mathcal{E}_{x_{0}})$$

$$= \sum_{i=1}^{d} \left\langle \pi_{m}^{*}\pi_{\ker\,i_{\xi_{0}^{\sharp}}} \left[\pi_{\ker\,i_{\xi_{0}^{\sharp}}} \pi_{m*}\pi_{m}^{*}\pi_{\ker\,i_{\xi_{0}^{\sharp}}} \right]^{-1} \pi_{\ker\,i_{\xi_{0}^{\sharp}}} \pi_{m*} E_{\xi_{0}}^{m}(\pi_{1}^{*}\Gamma.v_{i}), E_{\xi_{0}}^{m}(\pi_{1}^{*}\Gamma.v_{i}) \right\rangle_{L^{2}(\mathbb{S}^{n-1},\pi^{*}\mathcal{E}_{x_{0}})}.$$

$$(5.24)$$

Since $(\pi_m^* \pi_{\ker \imath_{\xi_0^{\sharp}}})^* = \pi_{\ker \imath_{\xi_0^{\sharp}}} \pi_{m*}$ in the L^2 sense, we have the orthogonal decomposition:

$$L^{2}(\mathbb{S}^{n-1}, \pi^{*} \mathcal{E}_{x_{0}}) = \pi^{*}_{m} \Big(\otimes^{m}_{S} \ker \imath_{\xi^{\sharp}_{0}} \otimes \mathcal{E}_{x_{0}} \Big) \oplus^{\perp} \ker \Big(\pi_{\ker \imath_{\xi^{\sharp}_{0}}} \pi_{m*} |_{L^{2}(\mathbb{S}^{n-1})} \Big).$$
(5.25)

In particular, if we define $w_i := E^m_{\xi_0}((\pi_1^*\Gamma, v_i)|_{\mathbb{S}^{n-2}_{\xi_0}})$, then we can write

$$\mathbf{w}_i = \pi_m^* T_i + h_i, \tag{5.26}$$

where $T_i \in \bigotimes_S^m \ker \imath_{\xi_0^{\sharp}} \otimes \mathcal{E}_{x_0}$ and $h_i \in \ker(\pi_{\ker \imath_{\xi_{\sharp}}} \pi_{m*}|_{L^2(\mathbb{S}^{n-1})})$. We also define

$$P_m := \pi_m^* \pi_{\ker \imath_{\xi_0^{\sharp}}} [\pi_{\ker \imath_{\xi_0^{\sharp}}} \pi_m * \pi_m^* \pi_{\ker \imath_{\xi_0^{\sharp}}}]^{-1} \pi_{\ker \imath_{\xi_0^{\sharp}}} \pi_m * : L^2(\mathbb{S}^{n-1}, \pi^* \mathcal{E}_{x_0}) \to L^2(\mathbb{S}^{n-1}, \pi^* \mathcal{E}_{x_0}),$$

¹⁴Alternatively, a Gaussian state is an *h*-dependent function whose semiclassical defect measure is a point $(x_0, \xi_0) \in T^*M$, see [Zwo12].

and observe that $P_m^2 = P_m$, $P_m^* = P_m$, so P_m is the orthogonal projection onto the first factor of (5.25). In particular $\pi_m^* T_i = P_m w_i$ and (5.24) reads:

$$\sum_{i=1}^{d} \| \mathbf{w}_{i} \|_{L^{2}(\mathbb{S}^{n-1},\pi^{*}\mathcal{E}_{x_{0}})}^{2} = \sum_{i=1}^{d} \| P_{m} \mathbf{w}_{i} \|_{L^{2}(\mathbb{S}^{n-1},\pi^{*}\mathcal{E}_{x_{0}})}^{2}.$$
(5.27)

As a consequence, in order to obtain a contradiction in (5.27), it is sufficient to exhibit a Γ such that $h_1 \neq 0$ (where h_1 is given in (5.26)). Since $h_1 \in \ker(\pi_{\ker i_{\xi^{\sharp}}} \pi_{m_*})$ and $\ker(\pi_{m_*}) \subset \ker(\pi_{\ker i_{\xi^{\sharp}}} \pi_{m_*})$, it is sufficient to show that the orthogonal projection of w_1 onto $\ker(\pi_{m_*}|_{L^2})$ is not zero, that is, it is sufficient to show that $w_1 = E_{\xi_0}^m(\pi_1^*\Gamma, v_1)$ has degree $\geq m + 1$:

Lemma 5.7. There exists $x_0 \in M$, $\xi_0 \in T^*_{x_0}M \setminus \{0\}$ and $\Gamma \in T^*_{x_0}M \otimes \operatorname{End}_{\operatorname{sk}}(\mathcal{E}_{x_0})$ such that $\operatorname{deg}(E^m_{\xi_0}(\pi_1^*\Gamma, v_1)) \geq m+1.$

Proof. For that, we will need the following claim:

$$\forall 1 \leq i \leq d$$
, we have $\deg(\mathbf{v}_i) \geq m+1$.

Indeed, assuming the contrary, since v_i has opposite parity as m this would force $\deg(v_i) \leq m-1$, that is, $v_i = \pi_{m-1}^* \tilde{v}_i$ for some $\tilde{v}_i \in C^{\infty}(M, \bigotimes_S^{m-1}T^*M)$. Recalling $\pi_m^* \Delta_0 u_i = \mathbf{X} v_i$ by (5.10) and using the relation (2.12), this implies $\Delta_0 u_i = D_{\mathcal{E}} \tilde{v}_i$. Hence $\pi_{\ker D_0^*} \Delta_0 u_i = 0$, implying $u_i = 0$ which is a contradiction.

Now we select x_0 according to v_1 , that is, we take x_0 such that at this point, the degree of v_1 in the fibre over x_0 is $\geq m+1$. This also implies that $\pi_1^*\Gamma$. v_1 has degree $\geq m+1$ (actually, the degree is at least m+2 but we do not need this) for some choice of $\Gamma \in T_{x_0}^*M \otimes \operatorname{End}_{\operatorname{sk}}(\mathcal{E}_{x_0})$ by Lemma 3.8 applied with k = 1. Then, by Lemma 3.1, we know that there exists a $\xi_0 \in T_{x_0}^*M \setminus \{0\}$ such that $\operatorname{deg}(\pi_1^*\Gamma. v_1|_{\mathbb{S}^{n-2}_{\xi_0}}) \geq m+1$ and then it suffices to apply the extension Lemma 3.5 to get that

$$\deg\left(E_{\xi_0}^m(\pi_1^*\Gamma.\operatorname{v}_1|_{\mathbb{S}_{\xi_0}^{n-2}})\right) \ge m+1$$

concluding the proof when $n \geq 3$.

This completes the proof of Theorem 1.8.

Remark 5.8. As mentioned in the introduction, Lemma 5.3 and (5.22) show that the second order derivative $d^2 \lambda(\Gamma, \Gamma)$ is of the form $\langle B\Gamma, \Gamma \rangle_{L^2}$, where *B* is some pseudodifferential operator (and the same will occur in the metric case). However, this operator is *a priori* not elliptic. More precisely, the proof only shows that there exists a point $x_0 \in M$ (where v_i has degree m + 1) where the principal symbol of *B* is non-zero. Had we been able to show the ellipticity of *B*, we would have obtained that locally the space of connections (up to gauge) with non-injective X-ray transform is finite-dimensional (and its tangent space would have been equal to the kernel of *B*).

Remark 5.9. Our proof does not give generic injectivity when n = 2. More precisely, Lemma 5.7 does not work in that case, since $E_{\xi_0}^m(\pi_1^*\Gamma, \mathbf{v}_1)$ always has degree equal to m. Therefore, the equality (5.27) always holds and our proof shows that the pseudodifferential operator $L_{\mathbf{v}_i} - Q_{\mathbf{v}_i}^* \prod_{m=1}^{n-1} Q_{\mathbf{v}_i}$ appearing in (5.22) is in fact of order -2, as opposed to the case $n \geq 3$

where we show that this operator is strictly of order -1. However, we believe that in the former case the second variation should also be non-zero, and that this should be provable using the method in §4 directly.

5.4. Endomorphism case. We conclude this section with a discussion of the endomorphism case. More precisely, if $\mathcal{E} \to M$ is a Hermitian vector bundle, we let $\operatorname{End}(\mathcal{E}) \to M$ be the vector bundle of endomorphisms. If $\nabla^{\mathcal{E}}$ is a unitary connection on \mathcal{E} , it induces a canonical connection $\nabla^{\operatorname{End}(\mathcal{E})}$ on $\operatorname{End}(\mathcal{E})$ defined so that it satisfies the Leibniz rule:

$$\left[\nabla^{\operatorname{End}(\mathcal{E})}u\right] \cdot f := \nabla^{\mathcal{E}}(u(f)) - u(\nabla^{\mathcal{E}}f),$$

for all $f \in C^{\infty}(M, \mathcal{E})$, $u \in C^{\infty}(M, \operatorname{End}(\mathcal{E}))$. Similarly to §2.3.3, one can define a twisted X-ray transform $\Pi_m^{\operatorname{End}(\mathcal{E})}$ with values in the endomorphism bundle $\operatorname{End}(\mathcal{E})$. More precisely, the operator $\pi^* \nabla_X^{\operatorname{End}(\mathcal{E})}$ always contains $\mathbb{C} \cdot \mathbb{1}_{\mathcal{E}}$ in its kernel and its kernel is generically reduced to $\mathbb{C}.\mathbb{1}_{\mathcal{E}}$ (see [CL21a], such a connection is also said to be *opaque*). We then set:

$$\Pi_m^{\operatorname{End}(\mathcal{E})} := \pi_{m*} (\mathbf{R}_+ + \mathbf{R}_- + \Pi_{\mathbb{C}.\mathbb{1}_{\mathcal{E}}}) \pi_m^*,$$

where $\Pi_{\mathbb{C},\mathbb{1}_{\mathcal{E}}}$ denotes the L^2 -orthogonal projection onto $\mathbb{C},\mathbb{1}_{\mathcal{E}}$ and \mathbf{R}_{\pm} is (the opposite of) the holomorphic part of the resolvents of $\pi^* \nabla_X^{\mathrm{End}(\mathcal{E})}$ at z = 0.

For m = 1, the solenoidal injectivity of the operator $\Pi_1^{\text{End}(\mathcal{E})}$ appears to be crucial when studying the *holonomy inverse problem* on Anosov manifolds, namely: to what extent does the trace of the holonomy of a connection along closed geodesics determine the connection? We proved in a companion paper [CL21b] that this problem is locally injective near a connection $\nabla^{\text{End}(\mathcal{E})}$ such that its induced operator $\Pi_1^{\text{End}(\mathcal{E})}$ is s-injective. Similarly to Theorem 1.8, this turns out to be a generic property:

Theorem 5.10. There exists $k_0 \gg 1$ such that the following holds. Let (M, g) be a smooth Anosov manifold of dimension $n \geq 3$, $\pi_{\mathcal{E}} : \mathcal{E} \to M$ be a smooth Hermitian vector bundle and let $m \in \mathbb{Z}_{\geq 0}$. Moreover, assume that the X-ray transform I_m with respect to (M, g) is s-injective. Then, there exists an open and dense set $\mathcal{S}_m \subset \mathcal{A}_{\mathcal{E}}^{k_0}$ of unitary connections with s-injective twisted generalized X-ray transform $\Pi_m^{\nabla^{\text{End}(\mathcal{E})}}$ on the endomorphism bundle.

Note that the main difference with Theorem 1.8 is that we need to assume that I_m is s-injective; this is known for m = 0, 1 on all Anosov manifolds [DS03] and this is a generic condition with respect to the metric by our Theorem 1.1.

Proof. We just point out the main differences with the proof of Theorem 1.8. If $\nabla^{\mathcal{E}}$ is a fixed unitary connection and $\nabla^{\mathcal{E}} + \Gamma$ (for $\Gamma \in C^{\infty}(M, T^*M \otimes \operatorname{End}_{\operatorname{sk}}(\mathcal{E}))$) is a perturbation, the induced connection on the endomorphism bundle is $\nabla^{\operatorname{End}(\mathcal{E})} + [\Gamma, \bullet]$. Then, in the computations of §5.2, each time that a term $\pi_1^*\Gamma$. v_i appears, it has to be replaced by $[\pi_1^*\Gamma, v_i]$ and the v_i 's are now elements of $C^{\infty}(SM, \pi^*\operatorname{End}(\mathcal{E}))$, where v_i satisfy a version of (5.10) for $\mathbf{X} := \pi^* \nabla_X^{\operatorname{End}}$.

Now, each v_i can be (uniquely) decomposed as $v_i = f_i.\mathbb{1}_{\mathcal{E}} + v_i^{\perp}$, where $f_i \in C^{\infty}(SM)$ and $v_i^{\perp} \in C^{\infty}(SM, \pi^* \text{End}(\mathcal{E}))$ is a (pointwise) trace-free endomorphism-valued section. One still has that $\mathbf{X} v_i$ is of degree m and v_i is of degree $\geq m + 1$. In fact, we claim that v_i^{\perp} is of

degree $\geq m + 1$. Indeed, assume that this is not the case, that is $\deg(\mathbf{v}_i^{\perp}) \leq m$; then f_i has to be of degree $\geq m + 1$ and

$$\mathbf{X} \mathbf{v}_i = (X f_i) . \mathbb{1}_{\mathcal{E}} + \mathbf{X} \mathbf{v}_i^{\perp}$$

is of degree m. As $(Xf_i).\mathbb{1}_{\mathcal{E}}$ and $\mathbf{X}\mathbf{v}_i^{\perp}$ are pointwise orthogonal as elements of $\pi^* \text{End}(\mathcal{E})$ (since $\mathbf{X}\mathbf{v}_i^{\perp}$ is trace-free), this forces each of them to be of degree $\leq m$ and thus Xf_i is of degree $\leq m$, and deg $(f_i) \geq m + 1$. But then the assumption that I_m is s-injective rules out this possibility.

Lemma 5.7 is then modified in the obvious way: one chooses a point x_0 such that $\deg(\mathbf{v}_i^{\perp}) \geq m+1$ and it suffices to find a $\Gamma \in T^*_{x_0} M \otimes \operatorname{End}_{\operatorname{sk}}(\mathcal{E}_{x_0})$ such that

$$[\pi_1^*\Gamma, \mathbf{v}_i] = [\pi_1^*\Gamma, \mathbf{v}_i^{\perp}]$$

has degree $\geq m + 1$. For that, we choose an orthonormal basis $(\mathbf{e}_1, ..., \mathbf{e}_r)$ of \mathcal{E}_{x_0} and write in that basis $\mathbf{v}_i^{\perp} = (m_{j\ell})_{1 \leq j,\ell \leq r}$. By assumption, there is an element $m_{j\ell}$ with degree $\geq m + 1$. Without loss of generality, we can assume it is in the first column $\ell = 1$. If it is obtained for some $j_0 \neq 1$, then taking a *real-valued* $\alpha \in \Omega_1$ such that $\alpha.m_{j_01}$ has degree $\geq m + 1$ (which is possible by Lemma 3.7), and setting $\Gamma := i\alpha \times \mathbf{e}_1 \otimes \mathbf{e}_1^*$, we get easily that $[\pi_1^*\Gamma, \mathbf{v}_i^{\perp}]$ has degree $\geq m + 1$.

If it is obtained for $j_0 = 1$, then we write

$$\mathbf{v}_i^{\perp} = \begin{pmatrix} m_{11} & b^{\top} \\ c & d \end{pmatrix},$$

where $m_{11} \in C^{\infty}(SM)$ has degree $\geq m + 1$, b, c are vectors of length r - 1 and $d \in C^{\infty}(S_{x_0}M, \operatorname{End}(\mathbb{C}^{r-1}))$. Note that $\operatorname{Tr}(\mathbf{v}_i^{\perp}) = 0 = m_{11} + \operatorname{Tr}(d)$. Moreover, writing $d = (m_{j\ell})_{2\leq j,\ell\leq r}$, there is an element $m_{j_0j_0}$ on the diagonal of d such that, if $f_{\geq m+1}$ denotes the projection of a function f onto Fourier modes of degree $\geq m + 1$, one has $(m_{j_0j_0})_{\geq m+1} \neq (m_{11})_{\geq m+1}$ (indeed, if not, this would contradict $\operatorname{Tr}(\mathbf{v}_i^{\perp}) = 0$). Without loss of generality, we can assume that $j_0 = 2$. Then, taking

$$\Gamma = \begin{pmatrix} 0 & \gamma^\top \\ -\gamma & 0 \end{pmatrix},$$

where γ is a vector of length r-1 and $\gamma^{\top} = (\alpha, 0, ..., 0)$ and $\alpha \in \Omega_1$ is real-valued, we obtain:

$$[\pi_1^*\Gamma, \mathbf{v}_i^{\perp}] = \begin{pmatrix} * & \gamma^{\top}d - m_{11}.\gamma^{\top} \\ * & * \end{pmatrix}, \quad \gamma^{\top}d - m_{11}.\gamma^{\top} = ((m_{22} - m_{11}).\alpha, *, ..., *).$$

By assumption, $m_{22} - m_{11}$ has degree $\geq m+1$ and it thus suffices to choose a real $\alpha \in \Omega_1$ such that $(m_{22} - m_{11}).\alpha$ has degree $\geq m+1$. The existence of such an α is once again guaranteed by Lemma 3.7. This completes the proof.

6. Generic injectivity with respect to the metric

We now prove Theorem 1.1. As we shall see, the computations follow from the same strategy as in the connection case, except that they are more involved.

6.1. **Preliminary computations.** A first point to address is that the unit tangent bundle now varies if we perturb the metric.

6.1.1. Scaling the geodesic vector fields on SM. The metric $g := g_0$ is fixed and we consider a smooth variation $(g_{\tau})_{\tau \in (-1,1)}$ of the metric. Each $\tau \in (-1,1)$ defines a unit tangent bundle

$$SM_{\tau} := \{(x, v) \in TM \mid g_{\tau}(v, v) = 1\} \subset TM,$$

and we write $SM := SM_0$. Each metric g_{τ} induces a geodesic vector field H_{τ} defined on the whole tangent bundle TM (which is tangent to SM_{τ} , for every $\tau \in (-1, 1)$). We let

$$\Phi_{\tau} : SM \to SM_{\tau}, \ (x,v) \mapsto \left(x, \frac{v}{|v|_{g_{\tau}}}\right),$$
(6.1)

be the natural projection onto SM_{τ} . We consider the family $X_{\tau} := \Phi_{\tau}^* H_{\tau} \in C^{\infty}(SM, T(SM))$, which depends smoothly on τ and is defined so that $X_{\tau=0} = X_0$ is the geodesic vector field of the metric g_0 . Note that $(X_{\tau})_{\tau \in (-1,1)}$ is a smooth family of Anosov vector fields on SM. In what follows, we will drop the sub-index zero when referring to quantities at $\tau = 0$. We will use $\sharp : T^*(SM) \to T(SM)$ to denote the musical isomorphism with respect to the Sasaki metric.

Lemma 6.1. We have:

$$\dot{X} = -\frac{1}{2}\pi_{2}^{*}\dot{g} \cdot X + \frac{1}{2}J\left[\nabla_{\mathbb{V}}(\pi_{2}^{*}\dot{g}) + \nabla_{\mathbb{H}}(\pi_{2}^{*}\dot{g})\right] - J\left(\left(\nabla_{X}^{\mathrm{Sas}}\pi_{2,\mathrm{Sas}}^{*}\dot{g}\right)(X,\pi_{\mathbb{H}}(\bullet))\right)^{\sharp} + \pi_{2,\mathrm{Sas}}^{*}\dot{g}(X,\pi_{\mathbb{H}}(\bullet))^{\sharp}.$$
(6.2)

Proof. Write $\beta_{\tau}(x,v)(\xi) = \langle v, d\pi(x,v)\xi \rangle_{g_{\tau}(x)}$ for the contact 1-form on SM_{τ} . Writing $\alpha_{\tau} := \Phi_{\tau}^* \beta_{\tau}$ and $\alpha := \beta_0$, and using $\pi \circ \Phi_t = \pi$, we obtain for any $\xi \in T_{(x,v)}SM$:

$$\alpha_{\tau}(x,v)(\xi) = |v|_{g_{\tau}}^{-1} g_{\tau}(v, \mathrm{d}\,\pi(x,v)(\xi)).$$

Differentiating, we obtain the relation:

$$\dot{\alpha} = -\frac{1}{2}\pi_2^* \dot{g} \cdot \alpha + \dot{g}(v, \mathrm{d}\,\pi(x, v)(\bullet)). \tag{6.3}$$

The pullback vector field X_{τ} is uniquely determined by the relations: $\iota_{X_{\tau}}\alpha_{\tau} = 1$ and $\iota_{X_{\tau}}d\alpha_{\tau} = 0$. Differentiating, we get $\iota_{\dot{X}}\alpha + \iota_X\dot{\alpha} = 0$ and thus using (6.3), $\alpha(\dot{X}) = -\frac{1}{2}\pi_2^*\dot{g}$. Since we can decompose $\dot{X} = \alpha(\dot{X})X + Y$ for some Y orthogonal to X this gives the first term in (6.2). It remains to compute Y.

For that, we introduce the 1-form, defined for $(x, v) \in SM$ as

$$A(x,v)(\bullet) := \dot{g}(v, \mathrm{d}\,\pi(x,v)(\bullet)) = \pi^*_{2,\mathrm{Sas}}\dot{g}(X,\bullet),$$

using the Sasaki lift introduced in (2.13). The first step is to compute $\iota_X dA$ and we claim:

$$\iota_X dA = -d(\pi_2^* \dot{g}) + (\nabla_X^{\operatorname{Sas}} \pi_{2,\operatorname{Sas}}^* \dot{g})(X, \pi_{\operatorname{\mathbb{H}_{tot}}}(\bullet)) - \pi_{2,\operatorname{Sas}}^* \dot{g}(X, J\pi_{\mathbb{V}}(\bullet)).$$
(6.4)

Recall here that $\mathbb{H}_{\text{tot}} = \mathbb{H} \oplus \mathbb{R} \cdot X$ denotes the total horizontal space, as explained in §2.1, and $\pi_{\mathbb{H}_{\text{tot}}}$ is the orthogonal projection onto this space (with respect to the Sasaki metric).

Note that A defines a 1-form on TM and we will first compute $\iota_X dA$ on TM. Then $\iota_X dA$ on SM is just the restriction. For $W, Z \in C^{\infty}(TM, T(TM))$, we have the formula:

$$dA(W,Z) = W \cdot \left(\pi_{2,\text{Sas}}^* \dot{g}(X,Z)\right) - Z \cdot \left(\pi_{2,\text{Sas}}^* \dot{g}(X,W)\right) - \pi_{2,\text{Sas}}^* \dot{g}(X,[W,Z]).$$
(6.5)

We now fix a point $p \in M$ and take a geodesic orthonormal frame $(E_1, ..., E_n)$ around p, i.e. such that $\nabla_{E_i} E_j(p) = 0$. Let X_i be the horizontal lift of E_i . We have that $\nabla_{X_i}^{\text{Sas}} X$ is vertical (see [Pat99, Lemma 1.25]) and $\nabla_X^{\text{Sas}} X_i$ is also vertical at p (as π is a Riemannian submersion). Hence by (6.5), at the point p:

$$\iota_X dA(X_i) = (\nabla_X^{\text{Sas}} \pi_{2,\text{Sas}}^* \dot{g})(X, X_i) + \pi_{2,\text{Sas}}^* \dot{g}(\nabla_X^{\text{Sas}} X, X_i) + \pi_{2,\text{Sas}}^* \dot{g}(X, \nabla_X^{\text{Sas}} X_i) - X_i \cdot \pi_2^* \dot{g} - \pi_{2,\text{Sas}}^* \dot{g}(X, \underbrace{[X, X_i]}_{=\nabla_X X_i - \nabla_{X_i} X}) = (\nabla_X^{\text{Sas}} \pi_{2,\text{Sas}}^* \dot{g})(X, X_i) - X_i \cdot \pi_2^* \dot{g}.$$
(6.6)

Introducing $Y_i := JX_i$, it can also be checked that $[X, Y_i] = JY_i = -X_i$ at the point p (this is an immediate consequence of the fact that $[X_i, Y_j] = 0$ at p, see [Pat99, Exercise 1.26]). Thus by (6.5), at p:

$$\iota_X dA(Y_i) = X \cdot \pi^*_{2,\text{Sas}} \dot{g}(X, Y_i) - Y_i \cdot \left(\pi^*_{2,\text{Sas}} \dot{g}(X, X)\right) - \pi^*_{2,\text{Sas}} \dot{g}(X, [X, Y_i]) = -Y_i \cdot \pi^*_2 \dot{g} - \pi^*_{2,\text{Sas}} \dot{g}(X, JY_i).$$
(6.7)

Combining (6.6) and (6.7) immediately yields (6.4) and proves the claim.

Hence, combining (6.4) with (6.3), we get

$$\iota_X d\dot{\alpha} = -\frac{1}{2} X(\pi_2^* \dot{g}) . \alpha - \frac{1}{2} d(\pi_2^* \dot{g}) + (\nabla_X^{\text{Sas}} \pi_{2,\text{Sas}}^* \dot{g}) (X, \pi_{\mathbb{H}_{\text{tot}}}(\bullet)) - \pi_{2,\text{Sas}}^* \dot{g} (X, J\pi_{\mathbb{V}}(\bullet)).$$

Using $\iota_X d\dot{\alpha} + \iota_{\dot{X}} d\alpha = \iota_X d\dot{\alpha} + \iota_Y d\alpha = 0$ together with (2.3), we get:

$$Y = -J(\iota_X d\dot{\alpha})^{\sharp} = \frac{1}{2}J\left(\nabla_{\mathbb{V}}\pi_2^* \dot{g} + \nabla_{\mathbb{H}}\pi_2^* \dot{g}\right) - J\left(\left(\nabla_X^{\mathrm{Sas}}\pi_{2,\mathrm{Sas}}^* \dot{g}\right)(X,\pi_{\mathbb{H}}(\bullet))\right)^{\sharp} + J\left(\pi_{2,\mathrm{Sas}}^* \dot{g}(X,J\pi_{\mathbb{V}}(\bullet))\right)^{\sharp}.$$

Note here that we used $(\nabla_X^{\text{Sas}}\pi^*\dot{g})(X,\pi_{\mathbb{H}}(\bullet)) = (\nabla_X^{\text{Sas}}\pi^*_{2,\text{Sas}}\dot{g})(X,\pi_{\mathbb{H}_{\text{tot}}}(\bullet)) - X(\pi_2^*\dot{g}).\alpha$, valid since $X(\pi_2^*\dot{g}) = (\nabla_X^{\text{Sas}}\pi^*_{2,\text{Sas}}\dot{g})(X,X)$. The last term can be simplified as $\pi^*_{2,\text{Sas}}\dot{g}(X,\pi_{\mathbb{H}}(\bullet))^{\sharp}$.

The last two terms of (6.2) vanish for a conformal perturbation. We introduce the differential operator $\Lambda^{\text{conf}} \in \text{Diff}^1(SM, \mathbb{C} \to T_{\mathbb{C}}(SM))$ of order one

$$\Lambda^{\operatorname{conf}}(f) := \frac{1}{2} \left(-fX + J \left[\nabla_{\mathbb{V}} f + \nabla_{\mathbb{H}} f \right] \right).$$
(6.8)

We also introduce $\Lambda^{aniso} \in \text{Diff}^1(SM, \otimes^2_S T^*(SM) \to T_{\mathbb{C}}(SM))$ of order one by:

$$\Lambda^{\text{aniso}}(f) := -J\left(\left(\nabla_X^{\text{Sas}}f\right)(X, \pi_{\mathbb{H}}(\bullet))\right)^{\sharp} + f(X, \pi_{\mathbb{H}}(\bullet))^{\sharp}$$
(6.9)

By construction $\dot{X} = \Lambda^{\text{conf}}(\pi_2^* f) + \Lambda^{\text{aniso}}(\pi_{2,\text{Sas}}^* f)$. In order to manipulate compact notations, we will write $\dot{X} := \Lambda(\pi_2^* f)$, although there is some abuse of notations here as there are two distinct lifts of f to SM.

We now compute the symbols of $\Lambda^{\text{conf,aniso}}$. They will be useful in the perturbation arguments in the following sections.

Lemma 6.2. For any $(x, v, \xi) \in T(SM)$, we have:

$$\sigma(\Lambda^{\text{conf}})(x,v,\xi) = \frac{i}{2}J(x,v)(\pi_{\mathbb{H}}\xi^{\sharp} + \pi_{\mathbb{V}}\xi^{\sharp}) = \frac{i}{2}J(x,v)(\xi^{\sharp} - \xi(X(x,v)).X(x,v)).$$

Proof. Consider the Lagrangian state $e^{i\frac{S}{\hbar}}f$, where S(x,v) = 0, $d S(x,v) = \xi$ and f(x,v) = 1. By (6.8), we compute

$$\Lambda^{\operatorname{conf}}(e^{i\frac{S}{h}}f) = -\frac{1}{2}e^{i\frac{S}{h}}\Big[fX - \frac{i}{h}fJ(\nabla_{\mathbb{H}}S + \nabla_{\mathbb{V}}S) - J(\nabla_{\mathbb{H}}f + \nabla_{\mathbb{V}}f)\Big].$$
(6.10)

We may directly read off the principal symbol from this expression:

$$\sigma(\Lambda^{\text{conf}})(x,v,\xi) = \lim_{h \to 0} h\Lambda(e^{i\frac{S}{h}}f)(x,v) = \frac{i}{2}J(\nabla_{\mathbb{H}}S + \nabla_{\mathbb{V}}S)(x,v) = \frac{i}{2}J(x,v)(\pi_{\mathbb{H}}\xi^{\sharp} + \pi_{\mathbb{V}}\xi^{\sharp}).$$

We have:

Lemma 6.3. For any $(x, v, \xi) \in T(SM)$ and $f \in \bigotimes_S^2 T^*_{(x,v)}(SM)$, we have:

$$\sigma(\Lambda^{\text{aniso}})(x,v,\xi)f = -i\langle\xi, X(x,v)\rangle J(x,v) (f(X(x,v),\pi_{\mathbb{H}}(\bullet)))^{\sharp}$$

Proof. We see that it suffices to compute the principal symbol of the first term in (6.9) as the second term is of lower order. We take a Lagrangian state $e^{i\frac{S}{\hbar}}f$ with S(x,v) = 0, $d S(x,v) = \xi$ and $f \in C^{\infty}(SM, \otimes_{S}^{2}T^{*}(SM)), f(x) =: f_{0}$. We have:

$$\sigma(\Lambda^{\text{aniso}})(x,v,\xi)f_0 = \lim_{h \to 0} h\Lambda^{\text{aniso}}(e^{i\frac{S}{h}}f)(x,v)$$

$$= \lim_{h \to 0} -hJ(x,v) \left(\frac{i}{h}XS.\left[f(X,\pi_{\mathbb{H}}(\bullet))\right]^{\sharp} + e^{i\frac{S}{h}}\left[(\nabla_X^{\text{Sas}}f)(X,\pi_{\mathbb{H}}(\bullet))\right]^{\sharp}\right)(x,v)$$

$$= -i\langle\xi, X(x,v)\rangle.J(x,v) \left(f_0(X(x,v),\pi_{\mathbb{H}}(\bullet))\right)^{\sharp}.$$

Eventually, we compute the divergence of \dot{X} in a geometric way. We prove:

Lemma 6.4. The following formula holds:

$$\operatorname{div}(\dot{X}) = X \left(\pi_0^* \operatorname{Tr}_{g_0}(\dot{g}) - \frac{n}{2} \pi_2^* \dot{g} \right).$$
(6.11)

In local coordinates $(x_i)_{1 \le i \le n}$ where g_0 and \dot{g} are identified with $n \times n$ symmetric matrices, we have $\operatorname{Tr}_{g_0}(\dot{g}) = \operatorname{Tr}(g_0^{-1}\dot{g})$.

Proof. Write $\Omega_{\tau} := d \operatorname{vol}_{\operatorname{Sas},g_{\tau}}$ for the Sasaki volume form of g_{τ} in SM_{τ} and $\Omega := \Omega_0$. Write \mathcal{J}_{τ} for the Jacobian of Φ_{τ} , i.e. $\Phi_{\tau}^*\Omega_{\tau} = \mathcal{J}_{\tau}\Omega$. Observe that

$$-\operatorname{div}(X_{\tau})\Omega = \mathcal{L}_{X_{\tau}}\Omega = \Phi_{\tau}^{*}(\mathcal{L}_{H_{\tau}}(\mathcal{J}_{\tau}^{-1} \circ \Phi_{\tau}^{-1}.\Omega_{\tau})) = \mathcal{J}_{\tau}.X_{\tau}(\mathcal{J}_{\tau}^{-1}).\Omega.$$

It follows that $\operatorname{div}(X_{\tau}) = \mathcal{J}_{\tau}^{-1} X_{\tau} \mathcal{J}_{\tau}$ and differentiating at $\tau = 0$ and using $\mathcal{J}_0 = 1$:

$$\operatorname{div}(\dot{X}) = X\dot{\mathcal{J}}.\tag{6.12}$$

In what follows we compute \mathcal{J}_{τ} . We will use that $d \operatorname{vol}_{g_{\tau}, \operatorname{Sas}}(x, v) = \pi^* d \operatorname{vol}_{g_{\tau}}(x, v) \wedge d \operatorname{vol}_{S_x M_{\tau}}(up \text{ to sign})$. We need the following auxiliary lemma:

Lemma 6.5. Let A_0 and A_1 be two symmetric, positive definite matrices, and denote by $\mathbb{S}_{A_i}^{n-1} = \{x \in \mathbb{R}^n \mid \langle A_i x, x \rangle = 1\} \subset \mathbb{R}^n$ the unit sphere with respect to the metric induced by A_i ; denote by $d \operatorname{vol}_{A_i}$ the induced volume form on $\mathbb{S}_{A_i}^{n-1}$. If $R(x) = \frac{x}{\sqrt{\langle A_1 x, x \rangle}}$ is the scaling map between the two spheres, then for $x \in \mathbb{S}_{A_0}^{n-1}$:

$$R^* d \operatorname{vol}_{A_1}(x) = \left(\sqrt{\frac{\det A_1}{\det A_0}} \cdot \langle A_1 x, x \rangle^{-\frac{n}{2}}\right) d \operatorname{vol}_{A_0}(x).$$
(6.13)

Proof. We first show the claim for $A_0 = \text{Id.}$ Denote $A_1 = A$ and write $R^* d \operatorname{vol}_A = j.d \operatorname{vol}_{\mathbb{S}^{n-1}}$ for some function j on \mathbb{S}^{n-1} . Observe that $d \operatorname{vol}_A(x) = \sqrt{\det A} \cdot i_{\sum_i x_i e_i} |dx|$ for $x \in \mathbb{S}^{n-1}_A$, where we write $|dx| = dx_1 \wedge \ldots \wedge dx_n$, e_i for the standard basis vectors of \mathbb{R}^n identified with ∂_i ; $\sum_i x_i e_i = x$ is the outer unit normal to \mathbb{S}^{n-1}_A at x. It is straightforward to compute:

$$\forall i = 1, \dots, n-1, \quad d_{e_n} R(e_i) = \partial_i R(e_n) = \frac{1}{\sqrt{A_{nn}}} \left(e_i - e_n \cdot \frac{A_{in}}{A_{nn}} \right).$$

This shows the claim at $x = e_n$:

$$R^*(d\operatorname{vol}_A)(e_n)(e_1,\ldots,e_{n-1}) = \sqrt{\det A} \cdot i_{\frac{e_n}{\sqrt{A_{nn}}}} dx_1 \wedge \ldots \wedge dx_n \left(\ldots,\frac{1}{\sqrt{A_{nn}}} \left(e_i - e_n \cdot \frac{A_{in}}{A_{nn}}\right),\ldots\right)$$
$$= (-1)^{n-1} \sqrt{\det A} \cdot \langle Ae_n,e_n \rangle^{-\frac{n}{2}}.$$

For general $x \in \mathbb{S}^{n-1}$, consider a $B \in SO(n)$ such that $B(e_n) = x$. Using that $B : \mathbb{S}_{B^T A B}^{n-1} \to \mathbb{S}_A^{n-1}$ is an isometry and the previous computation, the formula (6.13) for $A_0 = \text{Id}$ follows.

For general A_0 , simply consider a linear coordinate change given by B with $B^T A_0 B = \text{Id}$ and apply the previous result to $A = B^T A_1 B$. The completes the proof.

Using Lemma 6.5, we see that in local coordinates:

$$\begin{split} \Phi_{\tau}^{*}(\pi^{*}d\operatorname{vol}_{g_{\tau}}\wedge d\operatorname{vol}_{S_{x}M_{\tau}}) &= \pi^{*}d\operatorname{vol}_{g_{\tau}}\wedge \Phi_{\tau}^{*}(d\operatorname{vol}_{S_{x}M_{\tau}}) \\ &= \sqrt{\det g_{\tau}}|dx| \wedge \sqrt{\frac{\det g_{\tau}}{\det g_{0}}}.(\pi_{2}^{*}g_{\tau})^{-\frac{n}{2}}d\operatorname{vol}_{S_{x}M} \\ &= \frac{\det g_{\tau}}{\det g_{0}}.(\pi_{2}^{*}g_{\tau})^{-\frac{n}{2}}.\pi^{*}d\operatorname{vol}_{g_{0}}\wedge d\operatorname{vol}_{S_{x}M}. \end{split}$$

This shows that $\mathcal{J}_{\tau} = \frac{\det g_{\tau}}{\det g_0} . (\pi_2^* g_{\tau})^{-\frac{n}{2}}$, so taking the derivative at $\tau = 0$ and using (6.12) completes the proof and shows (6.11).

In what follows, we will frequently use the operators, defined for a distribution $v \in \mathcal{D}'(SM)$:

$$\forall u \in C^{\infty}(SM), \quad M_{\mathbf{v}}u := \mathbf{v} . u, \quad \forall Z \in C^{\infty}(SM, T(SM)), \quad N_{\mathbf{v}}Z := Z \mathbf{v} .$$
(6.14)

Sometimes u or Z will also be singular, in which case we will have to justify the extension of the corresponding operator to such functions.

6.1.2. Metric-dependent generalized X-ray transform. For each $\tau \in (-\varepsilon, \varepsilon)$ small enough, we can consider the positive resolvent $\mathbb{C} \ni z \mapsto (-X_{\tau} - z)^{-1}$ and the negative resolvent $\mathbb{C} \ni z \mapsto (X_{\tau} - z)^{-1}$. Since we have $X_{\tau} = (\Phi_{\tau})^* H_{\tau}$, the resolvents satisfy on $C^{\infty}(SM)$ that $(\pm X_{\tau} - z)^{-1} = (\Phi_{\tau})^* (\pm H_{\tau} - z)^{-1} (\Phi_{\tau})_*$ and therefore also

$$R_{\pm,0}^{X_{\tau}} = (\Phi_{\tau})^* R_{\pm,0}^{H_{\tau}} (\Phi_{\tau})_*, \quad \Pi_{\pm}^{X_{\tau}} = (\Phi_{\tau})^* \Pi_{\pm}^{H_{\tau}} (\Phi_{\tau})_*, \tag{6.15}$$

where the superscript denotes the vector field with respect to which the resolvent is taken, $R^{\bullet}_{\pm,0}$ is its holomorphic part at zero and Π^{\bullet}_{\pm} is the orthogonal projection to the resonant space at zero. From now on we drop the $\bullet = X_{\tau}$ superscript and simply write τ instead. By (2.17) we have

$$\Pi_{\pm}^{\tau} = \langle \bullet, \mu_{\tau} \rangle \mathbf{1}, \tag{6.16}$$

where **1** is the constant function and μ_{τ} is the pullback by Φ_{τ} of the normalised Liouville measure on SM_{τ} such that $\langle \mathbf{1}, \mu_{\tau} \rangle = 1$.

Let $(\pi_m^{g_\tau})^* : C^{\infty}(M, \otimes_S^m T^*M) \to C^{\infty}(SM_\tau)$ be the canonical pullback operator; we write $\pi_m^* = (\pi_m^{g_0})^*$. We may then compute, for a symmetric *m*-tensor *f*:

$$(\Phi_{\tau})^* (\pi_m^{g_{\tau}})^* f(x,v) = (\pi_m^{g_{\tau}})^* f\left(x, \frac{v}{|v|_{g_{\tau}}}\right) = f_x\left(\frac{v}{|v|_{g_{\tau}}}, \dots, \frac{v}{|v|_{g_{\tau}}}\right) = |v|_{g_{\tau}}^{-m} \pi_m^* f(x,v).$$

We denote $\chi_{\tau}(x,v) = |v|_{g_{\tau}}^{-m}$, so that by the previous equality:

$$\chi_{\tau}\pi_{m}^{*} = (\Phi_{\tau})^{*}(\pi_{m}^{g_{\tau}})^{*}, \quad (\pi_{m})_{*}\chi_{\tau} = (\pi_{m}^{g_{\tau}})_{*}(\Phi_{\tau})_{*}, \tag{6.17}$$

where the lower star denotes the pushforward, that is the L^2 adjoint of the pullback operator. We are in position to introduce the generalised X-ray transform with respect to H_{τ} and study its properties under re-scaling by (6.15) and (6.17):

$$\Pi_{m}^{\tau} := (\pi_{m}^{g_{\tau}})_{*} (R_{+,0}^{H_{\tau}} + R_{-,0}^{H_{\tau}} + \Pi_{+}^{H_{\tau}}) (\pi_{m}^{g_{\tau}})^{*} = \pi_{m*} \chi_{\tau} (\underbrace{R_{+,0}^{\tau} + R_{-,0}^{\tau} + \Pi_{+}^{\tau}}_{\mathcal{I}^{\tau} :=}) \chi_{\tau} \pi_{m}^{*}.$$
(6.18)

Moreover, the family

$$(-\varepsilon,\varepsilon) \ni \tau \mapsto \Pi_m^\tau \in \Psi^{-1}(M, \otimes_S^m T^*M \to \otimes_S^m T^*M)$$

depends smoothly on τ as stated in Lemma 2.3.

We keep the same strategy as in $\S5$ and define

$$P_{\tau} = \pi_{\ker D_0^*} \Delta_0 \Pi_m^{\tau} \Delta_0 \pi_{\ker D_0^*}, \qquad (6.19)$$

where Δ_0 is an elliptic, formally self-adjoint, positive, pseudodifferential operator of order k > 1/2 with diagonal principal symbol that induces an isomorphism on Sobolev spaces. As in Lemma 5.1, the maps:

$$\pi_{\ker D^*_{\tau}}\Delta_0\pi_{\ker D^*_0}: \ker D^*_0 \cap H^s(M, \otimes^m_S T^*M) \to \ker D^*_{\tau} \cap H^{s-k}(M, \otimes^m_S T^*M),$$

$$\pi_{\ker D^*_0}\Delta_0\pi_{\ker D^*_{\tau}}: \ker D^*_{\tau} \cap H^s(M, \otimes^m_S T^*M) \to \ker D^*_0 \cap H^{s-k}(M, \otimes^m_S T^*M),$$

are isomorphisms for τ small enough depending on $s \in \mathbb{R}$. In particular, as before, Π_m^{τ} is solenoidal injective (i.e. injective on symmetric tensors in ker D_{τ}^*) if and only if P_{τ} is solenoidal injective. In what follows we assume that $(u_i)_{i=1}^d$ is an L^2 -orthonormal basis of eigenstates of P_0 at 0. As in (5.8), we let λ_{τ} be the sum of the eigenvalues of P_{τ} inside a small contour near 0.

6.2. Variations of the ground state. We now compute the variations with respect to τ .

6.2.1. First order variations. As in the connection case, the first order variation of λ_{τ} at $\tau = 0$ is $\lambda_0 = 0$ and the second order variation is given by Lemma 5.2. We thus need to compute each term involved in the second derivative $\ddot{\lambda}_0$, namely \dot{P}_0 and \ddot{P}_0 . More precisely, the goal of this section is to compute $\sum_{i=1}^{d} \langle P_0^{-1} \dot{P}_0 u_i, \dot{P}_0 u_i \rangle_{L^2}$. We assume that $g_{\tau} = g + \tau f$, where $f \in C^{\infty}(M, \otimes_S^2 T^*M)$ is a perturbation such that

 $\dot{q} = f$. We write

$$-(-X_{\tau}-z)^{-1} = (z+X_{\tau})^{-1} = \frac{\Pi_{+}^{\tau}}{z} + R_{+,0}^{\tau} + zR_{+,1}^{\tau} + z^{2}R_{+,2}^{\tau} + \mathcal{O}(z^{3}),$$

$$-(X_{\tau}-z)^{-1} = (z-X_{\tau})^{-1} = \frac{\Pi_{-}^{\tau}}{z} + R_{-,0}^{\tau} + zR_{-,1}^{\tau} + z^{2}R_{-,2}^{\tau} + \mathcal{O}(z^{3}),$$
(6.20)

for the meromorphic expansion of the resolvents at zero. First of all, we compute the derivatives of χ_{τ} at $\tau = 0$:

$$\dot{\chi}_{\tau=0}(x,v) = -\frac{m}{2}\dot{g}(v,v) = -\frac{m}{2}\pi_2^*f(x,v),$$

$$\ddot{\chi}_{\tau=0}(x,v) = \frac{m}{2}\left(\frac{m}{2}+1\right)[\pi_2^*f]^2(x,v).$$
(6.21)

In the following we recall that $\Pi^{\tau}_{+} = \Pi^{\tau}_{-}$ (see (6.16)). We now turn to the first order variation of the resolvent:

Lemma 6.6. We have $\Pi_{\pm} = -\Pi_{\pm} X R_{\pm,0}$ and:

$$\dot{R}_{+,0} = -\left(R_{+,0}\dot{X}R_{+,0} + \Pi_{+}\dot{X}R_{+,1}\right), \quad \dot{R}_{-,0} = +\left(R_{-,0}\dot{X}R_{-,0} + \Pi_{-}\dot{X}R_{-,1}\right).$$

Proof. Let us deal with the second equality (the third one is similar). By (6.20), we have (recall that the contour γ around zero was defined in §2.4):

$$R_{+,0}^{\tau} = \frac{1}{2\pi i} \oint_{\gamma} (z + X_{\tau})^{-1} \frac{\mathrm{d}\,z}{z},\tag{6.22}$$

and differentiating with respect to τ , we get:

$$\dot{R}_{+,0} = -\frac{1}{2\pi i} \oint_{\gamma} (z+X)^{-1} \dot{X} (z+X)^{-1} \frac{\mathrm{d}\,z}{z} = -\left(R_{+,0} \dot{X} R_{+,0} + \Pi_{+} \dot{X} R_{+,1} + R_{+,1} \dot{X} \Pi_{+}\right).$$

To conclude, it suffices to observe that $\dot{X}\Pi_{+} = 0$ as \dot{X} is a vector field and the range of Π_{+} is always the constant functions $\mathbb{C} \cdot \mathbf{1}$.

As far as the derivative of the spectral projection is concerned, one starts with the equality

$$\Pi_{+}^{\tau} = \frac{1}{2\pi i} \oint_{\gamma} (z + X_{\tau})^{-1} \,\mathrm{d}\,z, \tag{6.23}$$

and then differentiates with respect to τ similarly as above; we omit the details.

Therefore, recalling (6.18), by (6.21) and Lemma 6.6 we obtain at $\tau = 0$:

$$\begin{aligned} \dot{\Pi}_{m} &= \pi_{m*} \left(-\frac{m}{2} \left(\pi_{2}^{*} f.\mathcal{I} + \mathcal{I} \pi_{2}^{*} f \right) + \dot{\mathcal{I}} \right) \pi_{m}^{*} \\ &= \pi_{m*} \left(-\frac{m}{2} \left(\pi_{2}^{*} f.\mathcal{I} + \mathcal{I} \pi_{2}^{*} f \right) - \left(R_{+,0} \dot{X} R_{+,0} + \Pi_{+} \dot{X} R_{+,1} \right) \\ &+ \left(R_{-,0} \dot{X} R_{-,0} + \Pi_{-} \dot{X} R_{-,1} \right) - \Pi_{+} \dot{X} R_{+,0} \right) \pi_{m}^{*}. \end{aligned}$$
(6.24)

Next, we observe that for m odd, $\pi_{m*}\mathbf{1} = 0$ and also $\pi_{m*}(\pi_2^*f) = 0$ and as a consequence, the terms in (6.24) involving Π_{\pm} disappear; when m is even, this simplification no longer occurs. Recalling the definition (6.19) of P_{τ} , we obtain from (6.24) the general expression (valid for m odd or even):

where $\varepsilon(m) = 0$ for m odd and $\varepsilon(m) = 1$ for m even. This last term is isolated on purpose because as we shall see, it only contributes to a smoothing remainder in the following argument and will therefore disappear in the principal symbol computations.

We let $u_i \in C^{\infty}(M, \bigotimes_S^m T^*M) \cap \ker D_0^*$ be one of the elements in the kernel of P_0 . Note that the operator Π_m being *real*, we can always assume that the u_i 's are real-valued. This implies using Lemma 2.1 that (as in §5)

$$\pi_m^* \Delta_0 \pi_{\ker D_0^*} u_i = X \, \mathbf{v}_i, \quad \text{for some} \quad \mathbf{v}_i \in C^\infty(SM) \quad \text{with} \quad \Pi_+ \, \mathbf{v}_i = 0, \tag{6.25}$$

which can also be chosen real-valued (since X is real). Hence, using (2.18) and the fact that $(R_{+,0} + R_{-,0})X = 0, X\Pi_{+} = \Pi_{+}X = \dot{X}\Pi_{+} = 0$, we obtain:

$$\dot{P}_{0}u_{i} = -\frac{m}{2}\pi_{\ker D_{0}^{*}}\Delta_{0}\pi_{m*}\left(\pi_{2}^{*}f.(R_{+,0}+R_{-,0})+(R_{+,0}+R_{-,0})\pi_{2}^{*}f\right)Xv_{i} + \pi_{\ker D_{0}^{*}}\Delta_{0}\pi_{m*}\left(-R_{+,0}\dot{X}\underbrace{R_{+,0}X}_{=1-\Pi_{+}}v_{i}+R_{-,0}\dot{X}\underbrace{R_{-,0}X}_{=-1+\Pi_{-}}v_{i}\right) + \varepsilon(m)\pi_{\ker D_{0}^{*}}\Delta_{0}\pi_{m*}\left(-\frac{m}{2}\pi_{2}^{*}f.\Pi_{+}+S(\pi_{2}^{*}f)\right)Xv_{i}$$

$$= -\pi_{\ker D_{0}^{*}}\Delta_{0}\pi_{m*}(R_{+,0}+R_{-,0})\left[\frac{m}{2}\pi_{2}^{*}f.Xv_{i}+\Lambda(\pi_{2}^{*}f)v_{i}\right] + \varepsilon(m)\pi_{\ker D_{0}^{*}}\Delta_{0}\pi_{m*}S(\pi_{2}^{*}f)Xv_{i}.$$

$$(6.26)$$

Let us introduce shorthand notation for the operators arising in (6.26), for any $v \in C^{\infty}(SM)$ and any symmetric 2-tensor $h \in C^{\infty}(M, \otimes_{S}^{2}T^{*}M)$:

$$B_{\mathbf{v}}h := \pi_{m*}S(\pi_2^*h)X\,\mathbf{v}_i, \quad Q_{\mathbf{v}}h := \pi_{m*}(R_{+,0} + R_{-,0})\left[\frac{m}{2}\pi_2^*hX\,\mathbf{v} + \Lambda(\pi_2^*h)\,\mathbf{v}\right]. \tag{6.27}$$

By using the facts that $P_0^{-1} = (\pi_{\ker D_0^*} \Delta_0 \pi_{\ker D_0^*})^{-1} \Pi_m^{-1} (\pi_{\ker D_0^*} \Delta_0 \pi_{\ker D_0^*})^{-1}$ on ker D_0^* and the equalities $D_0^* \pi_{m*}(R_{+,0} + R_{-,0}) = D_0^* \pi_{m*}S(\pi_2^*f) = 0$, we obtain from (6.26) (similarly to (5.18)):

$$\langle P_0^{-1} \dot{P}_0 u_i, \dot{P}_0 u_i \rangle_{L^2} = \left\langle \Pi_m^{-1} Q_{\mathbf{v}_i} f, Q_{\mathbf{v}_i} f \right\rangle_{L^2} + \varepsilon(m) \left\langle \Pi_m^{-1} B_{\mathbf{v}_i} f, B_{\mathbf{v}_i} f \right\rangle_{L^2} - 2\varepsilon(m) \Re \left(\langle Q_{\mathbf{v}_i} f, B_{\mathbf{v}_i} f \rangle_{L^2} \right)$$

$$(6.28)$$

We now prove that the operator B_v introduced in (6.27) is in fact smoothing:

Lemma 6.7. $B_{v_i} \in \Psi^{-\infty}(M, \otimes_S^2 T^*M \to \otimes_S^m T^*M)$ is a smoothing operator.

Proof. Similarly to the proof of Lemma 4.3, it suffices to show that B_{v_i} maps

$$\mathcal{D}'(M, \otimes_S^2 T^*M) \to C^{\infty}(M, \otimes_S^m T^*M)$$

boundedly. For that, let $h \in \mathcal{D}'(M, \otimes_S^2 T^*M)$. By [Lef19a, Lemma 2.1] we have WF $(\pi_2^*h) \subset \mathbb{V}^*$ and WF $(\pi_{2,\text{Sas}}^*h) \subset \mathbb{V}^*$ (as in the proof of Lemma 4.3). By resolvent identities (2.18) and using $\dot{X}\Pi_+ = 0$, we have:

$$S(\pi_2^*h)X\mathbf{v}_i = -\frac{m}{2}\Pi_+(\pi_2^*h.X\mathbf{v}_i) + \Pi_+\Lambda(\pi_2^*h)(R_{+,0} + R_{-,0})\mathbf{v}_i - \Pi_+\Lambda(\pi_2^*h)\mathbf{v}_i.$$
 (6.29)

By (2.17), the first term in (6.29) i.e. $-\frac{m}{2}\Pi_{+}(\pi_{2}^{*}h.X v_{i}) = -\frac{m}{2}\langle \pi_{2}^{*}h.X v_{i}, \mu \rangle \mathbf{1}$ is clearly smooth. Similarly, the last term $-\Pi_{+}\Lambda(\pi_{2}^{*}h) v_{i}$ is also obviously smooth.

We now deal with the remaining term. Define $v_i^u := (R_{+,0} + R_{-,0}) v_i$. Since v_i is smooth, this has wavefront set in E_u^* by the characterization of the wavefront set of the resolvent, see (4.4). Hence, the one-form dv_i^u has also wavefront set in E_u^* and thus by the multiplication rule of distributions (see [Hör03, Theorem 8.2.10] for instance), the inner product $\Lambda(\pi_2^*h) v_i^u = i_{\Lambda(\pi_2^*h)} dv_i^u$ is allowed (since $\mathbb{V}^* \cap E_u^* = \{0\}$ by the absence of conjugate points, see (2.8)). As a consequence

$$\Pi_{+}\Lambda(\pi_{2}^{*}h)(R_{+,0}+R_{-,0})\mathbf{v}_{i} = \langle \imath_{\Lambda(\pi_{2}^{*}h)} \operatorname{d} \mathbf{v}_{i}^{u}, \mu \rangle \mathbf{1}$$

is smooth, which proves the lemma.

We have an analogous statement to Lemma 5.4:

Lemma 6.8. For all $v \in C^{\infty}(SM)$,

$$Q_{\mathbf{v}} \in \Psi^0(M, \otimes_S^2 T^*M \to \otimes_S^m T^*M),$$

is a pseudodifferential operator of order 0 with principal symbol for $(x,\xi) \in T^*M \setminus \{0\}$, $\sigma_{Q_v}(x,\xi) \in \operatorname{Hom}(\otimes_S^2 T_x^*M, \otimes_S^m T_x^*M)$, given by:

$$\sigma_{Q_{\mathbf{v}}}(x,\xi)h = C_{n-1+2m}^{-1} \frac{i\pi}{|\xi|} \pi_{\ker \imath_{\xi^{\sharp}}} \pi_{m*} E_{\xi}^{m} \left(\langle \xi_{\mathbb{V}}, \nabla_{\mathbb{V}} \mathbf{v} \rangle . \pi_{2}^{*}h \right), \quad h \in \otimes_{S}^{2} T_{x}^{*} M$$

where $\xi_{\mathbb{V}}(x,v) \in \mathbb{H}^*(x,v)$ is defined by $\xi_{\mathbb{V}}(x,v)(\bullet) := \xi(\mathcal{K}_{x,v}(\bullet)).$

Observe that the main difference with Lemma 5.4 is that the operator is now of order 0 instead of -1: this is due to the fact that the operator Λ is of degree 1 (it costs derivatives of order 1 in f). Also note that the principal symbol of Q_v now satisfies $\sigma_{Q_v}(x, -\xi) = -\sigma_{Q_v}(x, \xi)$; this will not be a problem as Q_v will always appear twice in the brackets (hence the two minus signs will eventually give a plus).

Proof. We may re-write the operator $Q_{\rm v}$ in a sandwich form as follows:

$$Q_{\rm v} = \frac{m}{2} \pi_{m*} (R_{+,0} + R_{-,0}) M_{X\,\rm v} \pi_2^* + \pi_{m*} (R_{+,0} + R_{-,0}) N_{\rm v} (\Lambda^{\rm conf} \pi_2^* + \Lambda^{\rm aniso} \pi_{2,\rm Sas}^*), \quad (6.30)$$

where we use the notation of (6.14). By the sandwich Proposition 4.1 (and Remark 4.2 below), it follows that Q_v is pseudodifferential of order 0 and the first term of (6.30) is of order -1. By Lemma 6.2, the principal symbol of $N_v \Lambda^{\text{conf}}$ is given by, on co-vectors $\xi_{\mathbb{H}} = \xi \circ d_{x,v} \pi$:

$$\sigma(N_{\mathbf{v}}\Lambda^{\mathrm{conf}})(x,v,\xi_{\mathbb{H}}(x,v)) = \frac{i}{2}d_{x,v}\,\mathbf{v}\left(J(x,v)\left((\xi_{\mathbb{H}}(x,v))^{\sharp} - X(x,v).\langle\xi,v\rangle\right)\right)$$
$$= \frac{i}{2}\left\langle\nabla_{\mathbb{V}}\,\mathbf{v}(x,v),(0,\xi^{\sharp} - v.\langle\xi,v\rangle)\right\rangle = \frac{i}{2}\left\langle\xi_{\mathbb{V}}(x,v),\nabla_{\mathbb{V}}\,\mathbf{v}(x,v)\right\rangle.$$
(6.31)

Similarly, by Lemma 6.3, we have:

$$\sigma(N_{\mathbf{v}}\Lambda^{\mathrm{aniso}})(x,v,\xi_{\mathbb{H}}(x,v))h = -i\langle\xi_{\mathbb{H}}, X(x,v)\rangle.d_{x,v} \vee \left(J(h(X,\pi_{\mathbb{H}}(\bullet)))^{\sharp}\right)$$

$$= -i\langle\xi,v\rangle.d_{x,v} \vee \left(J(h(X,\pi_{\mathbb{H}}(\bullet)))^{\sharp}\right).$$
(6.32)

It then suffices to apply Proposition 4.1 and Remark 4.2 to conclude the proof. Note that the principal symbol of $N_{\rm v}\Lambda^{\rm aniso}$ does not appear as the expression in Remark 4.2 involves integration over the sphere $\{\langle \xi, v \rangle = 0\} = \mathbb{S}_{\xi}^{n-2}$.

To summarise the content of this section, we obtain from (6.28) using Lemmas 6.7 and 6.8:

$$\sum_{i=1}^{d} \langle P_0^{-1} \dot{P}_0 u_i, \dot{P} u_i \rangle_{L^2} = \sum_{i=1}^{d} \langle Q_{\mathbf{v}_i}^* \Pi_m^{-1} Q_{\mathbf{v}_i} f, f \rangle_{L^2(M, \otimes_S^2 T^* M)} + \langle \mathcal{O}_{\Psi^{-\infty}}(1) f, f \rangle_{L^2(M, \otimes_S^2 T^* M)}.$$
(6.33)

6.2.2. Second order variations. We now turn to the second variation of the resolvents which will allow us to compute $\sum_{i=1}^{d} \langle \ddot{P}_{0}u_{i}, u_{i} \rangle_{L^{2}}$. Similarly to Lemma 6.6, we have:

Lemma 6.9. We have $\ddot{\Pi}_{\pm} = 2\Pi_{\pm}\dot{X}R_{\pm,0}\dot{X}R_{\pm,0} \mp \Pi_{\pm}\ddot{X}R_{\pm,0}$ and:

$$\ddot{R}_{\pm,0} = 2 \left(\Pi_{\pm} \dot{X} R_{\pm,0} \dot{X} R_{\pm,1} + \Pi_{\pm} \dot{X} R_{\pm,1} \dot{X} R_{\pm,0} + R_{\pm,0} \dot{X} R_{\pm,0} \dot{X} R_{\pm,0} \right) \\ \mp \left(\Pi_{\pm} \ddot{X} R_{\pm,1} + R_{\pm,0} \ddot{X} R_{\pm,0} \right). \quad (6.34)$$

Proof. We prove the claims for the $(X_{\tau} + z)^{-1}$ resolvent, the other claim follows analogously. Differentiating the expression (6.22) twice at $\tau = 0$, we obtain the formula:

$$\ddot{R}_{+,0} = \frac{1}{\pi i} \oint_{\gamma} (X+z)^{-1} \dot{X} (X+z)^{-1} \dot{X} (X+z)^{-1} \frac{dz}{z} - \frac{1}{2\pi i} \oint_{\gamma} (X+z)^{-1} \ddot{X} (X+z)^{-1} \frac{dz}{z}.$$

The equation (6.34) now follows by the residue theorem and the expansion (6.20), using the relations $\dot{X}\Pi_{+} = \ddot{X}\Pi_{+} = 0$ to cancel the extra terms.

For the derivative of the spectral projector, by differentiating the formula (6.23) twice:

$$\ddot{\Pi}_{+} = \frac{1}{\pi i} \oint_{\gamma} (X+z)^{-1} \dot{X} (X+z)^{-1} \dot{X} (X+z)^{-1} dz - \frac{1}{2\pi i} \oint_{\gamma} (X+z)^{-1} \ddot{X} (X+z)^{-1} dz,$$

and the final result again follows from the expansion (6.20).

Next, differentiating (6.18) and inserting (6.21), we obtain at $\tau = 0$:

$$\ddot{\Pi}_{m} = \pi_{m*} \left(\frac{m}{2} \left(\frac{m}{2} + 1 \right) \left[(\pi_{2}^{*}f)^{2} \mathcal{I} + \mathcal{I}(\pi_{2}^{*}f)^{2} \right] - m \left[\pi_{2}^{*}f \mathcal{I} \dot{\mathcal{I}} + \dot{\mathcal{I}}\pi_{2}^{*}f \right] + \ddot{\mathcal{I}} + \frac{m^{2}}{2} \pi_{2}^{*}f \mathcal{I}\mathcal{I}\pi_{2}^{*}f \right] \pi_{m}^{*}.$$
(6.35)

We can already make some simplifications in the term $\langle \ddot{P}u_i, u_i \rangle_{L^2}$, where u_i is one of the eigenstates at 0 of P_0 . Recalling (6.25) and using $\mathcal{I}X = X\mathcal{I} = 0$, from (6.35) we get at $\tau = 0$:

$$\langle \ddot{P}_{0}u_{i}, u_{i}\rangle_{L^{2}} = \left\langle \pi_{\ker D_{0}^{*}}\Delta_{0}\ddot{\Pi}_{m}\Delta_{0}\pi_{\ker D_{0}^{*}}u_{i}, u_{i}\right\rangle_{L^{2}(M,\otimes_{S}^{m}T^{*}M)} = \left\langle \left(-m\left[\pi_{2}^{*}f.\dot{\mathcal{I}}+\dot{\mathcal{I}}\pi_{2}^{*}f\right]+\ddot{\mathcal{I}}+\frac{m^{2}}{2}\pi_{2}^{*}f.\mathcal{I}\pi_{2}^{*}f\right)Xv_{i}, Xv_{i}\right\rangle_{L^{2}(SM)}.$$
(6.36)

Similarly to §6.2.1, some terms will disappear when m is odd while in the case where m is even, they will only contribute up to a smoothing remainder. First of all, let us assume that m is odd. We make the simple observation that $\langle h_{\text{even}}, h_{\text{odd}} \rangle_{L^2} = 0$ for any $h_{\text{even}}, h_{\text{odd}} \in C^{\infty}(SM)$ with even and odd Fourier content, respectively. Therefore, the terms involving Π_{\pm} in (6.36) obtained after using Lemmas 6.6 and 6.9 to expand $\dot{\mathcal{I}}, \ddot{\mathcal{I}}$ vanish, and we get:

$$\begin{split} \langle \dot{P}_{0}u_{i}, u_{i}\rangle_{L^{2}} &= -m\left\langle \left[\pi_{2}^{*}f.\left(R_{-,0}\dot{X}R_{-,0} - R_{+,0}\dot{X}R_{+,0}\right) + \left(R_{-,0}\dot{X}R_{-,0} - R_{+,0}\dot{X}R_{+,0}\right)\pi_{2}^{*}f\right]Xv_{i}, Xv_{i}\right\rangle_{L^{2}} \\ &+ \left\langle \left(2\left(R_{+,0}\dot{X}R_{+,0}\dot{X}R_{+,0} + R_{-,0}\dot{X}R_{-,0}\dot{X}R_{-,0}\right) - R_{+,0}\ddot{X}R_{+,0} + R_{-,0}\ddot{X}R_{-,0}\right)Xv_{i}, Xv_{i}\right\rangle_{L^{2}} \\ &+ \frac{m^{2}}{2}\left\langle (R_{+,0} + R_{-,0})(\pi_{2}^{*}f.Xv_{i}), \pi_{2}^{*}f.Xv_{i}\right\rangle_{L^{2}}. \end{split}$$

Using the resolvent relations (2.18), the terms involving \ddot{X} cancel each other:

$$\left\langle \left(-R_{+,0}\ddot{X}R_{+,0} + R_{-,0}\ddot{X}R_{-,0} \right) X \mathbf{v}_i, X \mathbf{v}_i \right\rangle_{L^2(SM)} = 0.$$

$$= -m \left(- \left\langle \pi_{2}^{*}f.(R_{+,0} + R_{-,0})\dot{X}v_{i}, Xv_{i} \right\rangle_{L^{2}(SM)} + \left\langle \dot{X}(R_{+,0} + R_{-,0})(\pi_{2}^{*}f.Xv_{i}), v_{i} \right\rangle_{L^{2}(SM)} \right) - 2 \left\langle \dot{X}(R_{+,0} + R_{-,0})\dot{X}v_{i}, v_{i} \right\rangle_{L^{2}(SM)} + \frac{m^{2}}{2} \left\langle (R_{+,0} + R_{-,0})(\pi_{2}^{*}f.Xv_{i}), \pi_{2}^{*}f.Xv_{i} \right\rangle_{L^{2}(SM)} \\= \underbrace{2m \left\langle \pi_{2*} \left(Xv_{i}.(R_{+,0} + R_{-,0})\dot{X}v_{i} \right), f \right\rangle_{L^{2}} - m \left\langle \pi_{2*} \left(Xv_{i}.(R_{+,0} + R_{-,0})\left(\operatorname{div}(\dot{X}).v_{i}\right) \right), f \right\rangle_{L^{2}}}_{=:(I)} \\+ \underbrace{2 \left\langle (R_{+,0} + R_{-,0})\dot{X}v_{i}, \dot{X}v_{i} \right\rangle_{L^{2}}}_{=:(II)} - \underbrace{2 \left\langle (R_{+,0} + R_{-,0})\dot{X}v_{i}, \operatorname{div}(\dot{X}).v_{i} \right\rangle_{L^{2}}}_{=:(III)} \\+ \underbrace{\frac{m^{2}}{2} \left\langle \pi_{2*} \left(Xv_{i}.(R_{+,0} + R_{-,0})(\pi_{2}^{*}f.Xv_{i}) \right), f \right\rangle_{L^{2}}}_{=:(IV)}. \tag{6.37}$$

In the general case (where m is either even or odd), the previous equality still holds by adding on the right-hand side the extra term (V) which is equal to:

$$(\mathbf{V}) := \varepsilon(m) \left(-m \left\langle \pi_{2}^{*} f. \Pi_{+} \dot{X} \left(R_{+,0} + R_{-,0} - \mathbb{1} \right) \mathbf{v}_{i}, X \mathbf{v}_{i} \right\rangle_{L^{2}} + \frac{m^{2}}{2} \left\langle \pi_{2}^{*} f. \Pi_{+} (\pi_{2}^{*} f. X \mathbf{v}_{i}), X \mathbf{v}_{i} \right\rangle_{L^{2}} \right).$$
(6.38)

Here again we used (6.36) along with Lemmas 6.6 and 6.9 to expand the terms containing Π_{\pm} ; we also used (2.18) and $X\Pi_{\pm} = \dot{X}\Pi_{\pm} = 0$ to simplify the expression.

As before, we will see that each term in the previous equality can be written in the form $\langle Af, f \rangle_{L^2(M, \otimes_S^2 T^*M)}$, for some pseudodifferential operator A. In order to shorten the computations, it is important to understand the order of these operators: when taking Gaussian states, only the terms of highest order will remain. Also observe that (6.33) involves an operator of order 1: we therefore expect to find operators of order at most 1. First of all, we deal with the term (V) (appearing only when m is even):

Lemma 6.10. There exists
$$B'_{v_i} \in \Psi^{-\infty}(M, \bigotimes_S^2 T^*M \to \bigotimes_S^2 T^*M)$$
 such that:
 $(V) = \varepsilon(m) . \langle B'_{v_i} f, f \rangle_{L^2}.$
(6.39)

Proof. It follows from (6.38) that (6.39) holds with:

$$B_{\mathbf{v}_{i}}^{\prime} = \pi_{2*} \left(-m.M_{X\,\mathbf{v}_{i}} \Pi_{+} N_{(R_{+,0}+R_{-,0})\,\mathbf{v}_{i}} \Lambda + m.M_{X\,\mathbf{v}_{i}} \Pi_{+} N_{\mathbf{v}_{i}} \Lambda + \frac{m^{2}}{2} M_{X\,\mathbf{v}_{i}} \Pi_{+} M_{X\,\mathbf{v}_{i}} \right) \pi_{2}^{*}$$
(6.40)

The last two terms in (6.40) are obviously smoothing, while exactly the same wavefront set arguments as in Lemma 6.7 apply to show that the first term is smoothing.

We then prove (recall that $\xi_{\mathbb{V}}(x, v)(\bullet) = \xi(\mathcal{K}_{x,v}(\bullet))$):

Lemma 6.11. There exist pseudodifferential operators $A_{v_i}^{(I)}, A_{v_i}^{(II)}, A_{v_i}^{(III)}$ and $A_{v_i}^{(IV)}$ in the pseudodifferential algebra $\Psi^*(M, \otimes_S^2 T^*M \to \otimes_S^2 T^*M)$ of respective order 0, 1, 0 and -1 such that (recall (*) was defined in (6.37)):

$$(*) = \langle A_{v_i}^{(*)} f, f \rangle_{L^2}, \tag{6.41}$$

for $* \in \{I, II, III, IV\}$. Moreover, we have for $(x, \xi) \in T^*M \setminus \{0\}$ and $h \in \otimes_S^2 T_x^*M$:

$$\langle \sigma_{A^{(\mathrm{II})}}(x,\xi)h,h\rangle_{\otimes_{S}^{2}T_{x}^{*}M} = \frac{\pi}{|\xi|} C_{n-1+2m}^{-1} \left\| E_{\xi}^{m} \left(\pi_{2}^{*}h(\bullet).\langle \xi_{\mathbb{V}}(x,\bullet), \nabla_{\mathbb{V}} \mathbf{v}(x,\bullet) \rangle |_{\mathbb{S}_{\xi}^{n-2}} \right) \right\|_{L^{2}(S_{x}M)}^{2}.$$
(6.42)

Proof. Term (IV). Using the notation of (6.14), we observe that (6.41) holds for * = IV with:

$$A_{\mathbf{v}_{i}}^{(\mathrm{IV})} := \frac{m^{2}}{2} \pi_{2*} M_{X \, \mathbf{v}_{i}} (R_{+,0} + R_{-,0}) M_{X \, \mathbf{v}_{i}} \pi_{2}^{*}.$$

By the sandwich Proposition 4.1, this is a Ψ DO of order -1.

Term (I). Next, using the formula for the divergence in Lemma 6.4 together with $M_{v_i}Xu = X(M_{v_i}u) - M_{Xv_i}u$ and $(R_{+,0} + R_{-,0})X = 0 = X(R_{+,0} + R_{-,0})$, we see from (6.37) that (6.41) holds for * = I with:

$$A_{\mathbf{v}_{i}}^{(\mathrm{I})} := 2m.\pi_{2*}M_{X\,\mathbf{v}_{i}}(R_{+,0} + R_{-,0})N_{\mathbf{v}_{i}}\Lambda\pi_{2}^{*} + m.\pi_{2*}M_{X\,\mathbf{v}_{i}}(R_{+,0} + R_{-,0})M_{X\,\mathbf{v}_{i}}\left(\pi_{0}^{*}\operatorname{Tr}_{g_{0}}(\bullet) - \frac{n}{2}\pi_{2}^{*}(\bullet)\right).$$

$$(6.43)$$

By Proposition 4.1, the first term in (6.43) is of order 0 as Λ is of order 1, and the second term is of order -1.

Term (III). By Lemma 6.4 and using $M_{v_i}Xu = X(M_{v_i}u) - M_{Xv_i}u$, we get from (6.37):

$$A_{\mathbf{v}_i}^{(\mathrm{III})} = (2j_{g_0}\pi_{0*} - n.\pi_{2*})M_{X\,\mathbf{v}_i}(R_{+,0} + R_{-,0})N_{\mathbf{v}_i}\Lambda\pi_{2*}^*$$

where j_{g_0} denotes the multiplication by g_0 , that is $j_{g_0}u := u.g_0$ for $u \in C^{\infty}(M)$. By Proposition 4.1, this is a pseudodifferential operator of order 0.

Term (II). Finally, for the term (II), we write from (6.37):

$$A_{v_{i}}^{(\text{II})} := 2\left(\pi_{2*}\left(\Lambda^{\text{conf}}\right)^{*} + \pi_{2,\text{Sas}_{*}}\left(\Lambda^{\text{aniso}}\right)^{*}\right) N_{v_{i}}^{*}(R_{+,0} + R_{-,0}) N_{v_{i}}(\Lambda^{\text{conf}}\pi_{2}^{*} + \Lambda^{\text{aniso}}\pi_{2,\text{Sas}}^{*}),$$

where $(\Lambda^{\text{conf,aniso}})^*$ denotes the formal adjoint of the first order differential operators $\Lambda^{\text{conf,aniso}}$. Once more, applying Proposition 4.1 and Remark 4.2 shows that $A_{v_i}^{(\text{II})}$ is a Ψ DO of order 1 with the principal symbol given by (6.42), where we also use (6.31) and (6.32) to compute the principal symbols of $N_{v_i}\Lambda^{\text{conf}}$ and $N_{v_i}\Lambda^{\text{aniso}}$, and (3.6) to extend the formula to $S_x M$. Note that $N_{v_i}\Lambda^{\text{aniso}}$ does not contribute to the principal symbol as its symbol (see (6.32)) is proportional to $\langle \xi, v \rangle$ and the formula of Remark 4.2 involves integration over $\{\langle \xi, v \rangle = 0\} = \mathbb{S}_{\xi}^{n-2}$. As a consequence, (6.37) and the previous two lemmas show that:

$$\sum_{i=1}^{d} \langle \ddot{P}_0 u_i, u_i \rangle_{L^2} = \sum_{i=1}^{d} \langle R_{\mathbf{v}_i} f, f \rangle_{L^2} + \langle \mathcal{O}_{\Psi^{-\infty}}(1) f, f \rangle_{L^2}, \qquad (6.44)$$

where $R_{v_i} = A_{v_i}^{(I)} + A_{v_i}^{(II)} + A_{v_i}^{(III)} + A_{v_i}^{(IV)}$ is a pseudodifferential operator of order 1, whose principal symbol is given by that of $A_{v_i}^{(II)}$ and determined by (6.42).

6.3. Assuming the second variation is zero. We now assume that for any variation of the metric $g_{\tau} = g + \tau f$ with $f \in C^{\infty}(M, \otimes_{S}^{2}T^{*}M)$, the second derivative $\ddot{\lambda} = 0$ vanishes. Using Lemma 5.2, this implies that:

$$\sum_{i=1}^{d} \langle \ddot{P}_{0} u_{i}, u_{i} \rangle_{L^{2}(M, \otimes_{S}^{m} T^{*}M)} = 2 \sum_{i=1}^{d} \langle P_{0}^{-1} \dot{P}_{0} u_{i}, \dot{P}_{0} u_{i} \rangle_{L^{2}(M, \otimes_{S}^{m} T^{*}M)}.$$
(6.45)

In §6.2.1 and §6.2.2, we saw that both the right-hand side and the left-hand side can be written as quadratic forms in f, with pseudodifferential operators of order 1 acting on f. More precisely, combining (6.33) and (6.44), we have that (6.45) implies:

$$\forall f \in C^{\infty}(M, \otimes_{S}^{2} T^{*}M), \quad \sum_{i=1}^{d} \left\langle \left(R_{\mathbf{v}_{i}} - 2Q_{\mathbf{v}_{i}}^{*} \Pi_{m}^{-1} Q_{\mathbf{v}_{i}} \right) f, f \right\rangle_{L^{2}(M, \otimes_{S}^{2} T^{*}M)} = \langle \mathcal{O}_{\Psi^{-\infty}}(1) f, f \rangle_{L^{2}}, \tag{6.46}$$

where the remainder has this form by Lemmas 6.7 and 6.10 and appears only when m is even.

We then consider a real Gaussian state $f_h = \Re(e_h(x,\xi))\tilde{f}$ (see (5.23) for the notation), where $\tilde{f} \in C^{\infty}(M, \otimes_S^2 T^*M)$ is a smooth section such that $\tilde{f}(x) =: f$ and $\xi \in T_x^*M \setminus \{0\}$. Note that similarly to §5 we can only allow real perturbations of the metric, hence the need for the real part of the Gaussian state. Nevertheless, this will not be a problem insofar as the principal symbols of R_{v_i} and $Q_{v_i}^* \Pi_m^{-1} Q_{v_i}$ are preserved by the antipodal map in the fibres. We thus obtain by applying (6.46) to the Gaussian state $h.f_h$ and taking the limit as $h \to 0$, using Lemma 5.6:

$$\begin{split} C_{n-1+2m}^{-1} \frac{\pi}{|\xi|} &\sum_{i=1}^{d} \left\| E_{\xi}^{m} \left(\pi_{2}^{*}f(\bullet) \cdot \left\langle \xi_{\mathbb{V}}(x,\bullet), \nabla_{\mathbb{V}} \mathbf{v}_{i}(x,\bullet) \right\rangle |_{\mathbb{S}_{\xi}^{n-2}} \right) \right\|_{L^{2}(S_{x}M)}^{2} \\ &= 2 \sum_{i=1}^{d} \left\langle \sigma_{\Pi_{m}^{-1}\pi_{\ker D_{0}^{*}}}(x,\xi) \sigma_{Q_{\mathbf{v}_{i}}}(x,\xi) f, \sigma_{Q_{\mathbf{v}_{i}}}(x,\xi) f \right\rangle_{\otimes_{S}^{m}T_{x}^{*}M} \\ &= C_{n-1+2m}^{-1} \frac{\pi}{|\xi|} \sum_{i=1}^{d} \left\langle \left(\pi_{\ker \iota_{\xi}\sharp} \pi_{m*}\pi_{m}^{*}\pi_{\ker \iota_{\xi}\sharp} \right)^{-1} \pi_{\ker \iota_{\xi}\sharp} \pi_{m*}E_{\xi}^{m} \left(\pi_{2}^{*}f \cdot \left\langle \xi_{\mathbb{V}}, \nabla_{\mathbb{V}} \mathbf{v}_{i} \right\rangle |_{\mathbb{S}_{\xi}^{n-2}} \right), \\ &\pi_{\ker \iota_{\xi}\sharp} \pi_{m*}E_{\xi}^{m} \left(\pi_{2}^{*}f \cdot \left\langle \xi_{\mathbb{V}}, \nabla_{\mathbb{V}} \mathbf{v}_{i} \right\rangle |_{\mathbb{S}_{\xi}^{n-2}} \right) \right\rangle_{\otimes_{S}^{m}T_{x}^{*}M} \\ &= C_{n-1+2m}^{-1} \frac{\pi}{|\xi|} \sum_{i=1}^{d} \left\langle P_{m}E_{\xi}^{m} \left(\pi_{2}^{*}f \cdot \left\langle \xi_{\mathbb{V}}, \nabla_{\mathbb{V}} \mathbf{v}_{i} \right\rangle |_{\mathbb{S}_{\xi}^{n-2}} \right), E_{\xi}^{m} \left(\pi_{2}^{*}f \cdot \left\langle \xi_{\mathbb{V}}, \nabla_{\mathbb{V}} \mathbf{v}_{i} \right\rangle |_{\mathbb{S}_{\xi}^{n-2}} \right) \right\rangle_{\otimes_{S}^{m}T_{x}^{*}M}, \end{split}$$

where we used Lemmas 6.8 and 6.11 to compute the symbols of R_{v_i} and $Q_{v_i}^* \prod_m^{-1} Q_{v_i}$, and we introduced the following map:

$$P_m := \pi_m^* \pi_{\ker \imath_{\xi^\sharp}} \left(\pi_{\ker \imath_{\xi^\sharp}} \pi_m \ast \pi_m^* \pi_{\ker \imath_{\xi^\sharp}} \right)^{-1} \pi_{\ker \imath_{\xi^\sharp}} \pi_m \ast.$$

Note that P_m is the L^2 -orthogonal projection onto $\operatorname{ran}(\pi_m^* \pi_{\ker \imath_{\xi^{\sharp}}})$ in $L^2(S_x M)$ and so it holds that (cf. (5.25)):

$$L^{2}(S_{x}M) = \operatorname{ran}(\pi_{m}^{*}\pi_{\ker \imath_{\xi^{\sharp}}}) \oplus^{\perp} \ker(\pi_{\ker \imath_{\xi^{\sharp}}}\pi_{m*}).$$

Thus, we get cancelling the constant terms (similarly to (5.27)):

$$\sum_{i=1}^{d} \left\| E_{\xi}^{m} \left(\pi_{2}^{*} f.\langle \xi_{\mathbb{V}}, \nabla_{\mathbb{V}} v_{i} \rangle |_{\mathbb{S}_{\xi}^{n-2}} \right) \right\|_{L^{2}(S_{x}M)}^{2} = \sum_{i=1}^{d} \left\| P_{m} E_{\xi}^{m} \left(\pi_{2}^{*} f.\langle \xi_{\mathbb{V}}, \nabla_{\mathbb{V}} v_{i} \rangle |_{\mathbb{S}_{\xi}^{n-2}} \right) \right\|_{L^{2}(S_{x}M)}^{2}.$$
(6.47)

As $\ker(\pi_{m*}) \subset \ker(\pi_{\ker \imath_{\xi^{\sharp}}} \pi_{m*})$, to obtain a contradiction it is sufficient to show that the L^2 orthogonal projection onto $\ker(\pi_{m*})$ of the function $E_{\xi}^m\left(\pi_2^*f.\langle\xi_{\mathbb{V}},\nabla_{\mathbb{V}}v_1\rangle|_{\mathbb{S}_{\xi}^{n-2}}\right)$ is non-zero,
that is, it suffices to show the following:

Lemma 6.12. There exists $x \in M, \xi \in T_x^*M \setminus \{0\}$ and $f \in \otimes_S^2 T_x^*M$ such that:

$$\deg\left(E_{\xi}^{m}\left(\pi_{2}^{*}f.\langle\xi_{\mathbb{V}},\nabla_{\mathbb{V}}\,\mathbf{v}_{1}\rangle|_{\mathbb{S}_{\xi}^{n-2}}\right)\right) \geq m+1.$$
(6.48)

Proof. Observe that by Lemma 3.3, since v_1 has degree $\geq m + 1$ at some point $x \in M$, there exists $\xi \in T_x^*M$ such that $\langle \xi_{\mathbb{V}}, \nabla_{\mathbb{V}} v_1 \rangle |_{\mathbb{S}^{n-2}_{\xi}}$ has degree $\geq m$. Then, by Lemma 3.7 (observe here that the musical map is an isomorphism from $\ker\langle\xi,\bullet\rangle \subset T_xM$ to $\ker\iota_{\xi^{\sharp}} \subset T_x^*M$), there exists $f \in \bigotimes_S^2(\ker\iota_{\xi^{\sharp}})$ such that $\pi_2^*f.\langle\xi_{\mathbb{V}},\nabla_{\mathbb{V}} v_1\rangle|_{\mathbb{S}^{n-2}_{\xi}}$ has degree $\geq m+2$. Note that f can be naturally extended as a symmetric tensor in $\bigotimes_S^2 T_x^*M$ by setting $f(\xi^{\sharp},\bullet) = f(\bullet,\xi^{\sharp}) = 0$. Thus we can apply Lemma 3.5 to obtain (6.48).

This completes the proof of Theorem 1.1.

Remark 6.13. When n = 2, similarly as in Remark 5.9 our proof does not work. More precisely, the equality (6.47) always holds, which shows that the symbol of $R_{v_i} - 2Q_{v_i}^* \prod_m^{-1} Q_{v_i}$ appearing in (6.46) is zero, hence this operator is of order 0, as opposed to the case $n \ge 3$ where we show it is strictly of order 1.

6.4. Manifolds with boundary. This section is devoted to the proof of Corollary 1.2.

Proof of Corollary 1.2. We fix an integer $m \in \mathbb{Z}_{\geq 0}$. As in the closed case, the s-injectivity of I_m^g is equivalent to the s-injectivity of the normal operator $(I_m^g)^* I_m^g$ which has the same microlocal properties as the generalized X-ray transform Π_m^g , see [Gui17b] for instance. Hence the fact that \mathcal{R}'_m is open is an immediate consequence of elliptic theory so it suffices to show that \mathcal{R}'_m is dense in $\mathcal{M}^{k_0}_{\text{SNH}}$. We let $g \in \mathcal{M}^{k_0}_{\text{SNH}}$. By [CEG20], the manifold (M, g) can be embedded into a closed Anosov manifold $(M^{\text{ext}}, g_{\text{ext}})$ such that $g_{\text{ext}}|_M = g$ and by Theorem 1.1, we can perturb the metric g_{ext} (in the C^{k_0} -topology) to a new metric g'_{ext} such that this metric has injective X-ray transform $I_m^{g'_{\text{ext}}}$ on M^{ext} . In order to prove Corollary 1.2, it then suffices to show that $g'_{\text{ext}}|_M$ restricted to the manifold with boundary M has injective X-ray transform. In other words, it suffices to prove the following:

Lemma 6.14. Let (M, g) be a closed Anosov manifold and let $(N, g|_N) \subset (M, g)$ be an SNH manifold with boundary. If the X-ray transform I_m^g on the closed manifold is s-injective on M, then the X-ray transform $I_m^{g|_N}$ on the manifold with boundary is also s-injective on N.

Proof. We let $f \in C^{\infty}(N, \bigotimes_{S}^{m}T^{*}N)$ such that $I_{m}^{g|_{N}}f = 0$. First of all, by [Sha02, Lemma 2.2], we can write f = Dp + h, where $p|_{\partial N} = 0, \iota_{\nu}h = 0$ in a neighborhood of ∂N (where ν is the outward-pointing unit vector and is extended in the inner neighborhood of ∂N by flowing along the geodesics), ι_{ν} is the contraction by the vector ν and both tensors p, h are smooth. Observe that $I^{g|_{N}}f = 0 = I^{g|_{N}}h$. We claim that $\partial_{\nu}^{k}h = 0$ for all $k \geq 0$, that is h vanishes to infinite order on the boundary. For k = 0, this is contained in [Sha02, Lemma 2.3]. The proof is a simple observation: if $h_{x}(v, ..., v)$ is non-zero for some $x \in \partial M$ and $v \in T_{x}\partial N$ then it is also true in a small neighborhood of (x, v) and $\pi_{m}^{*}h$ has constant sign there; without loss of generality we can take it to be positive. Using short geodesics in a neighborhood of the boundary (with unit speed vector almost equal to v) we then get that $I_{m}h(x, v) > 0$, which is a contradiction. Hence we can write h = rh', where $r(x) := d(x, \partial N)$ is defined locally near the boundary and extended to an arbitrary positive function inside N. Then the same argument of positivity applies to h' and by iteration, we get that $h = \mathcal{O}(r^{\infty})$ at ∂N . Hence we can extend h by 0 outside N to get a smooth tensor (still denoted by h) in $C^{\infty}(M, \bigotimes_{S}^{m}T^{*}M)$.

We now claim that $I_m^g h = 0$ on M, that is the integral of $\pi_m^* h$ along closed geodesics in M is zero. Indeed, let $\gamma \subset SM$ be a closed orbit of the geodesic flow of length $\ell(\gamma)$, then:

$$I_m^g h(\gamma) = \frac{1}{\ell(\gamma)} \int_0^{\ell(\gamma)} \pi_m^* h(\varphi_t(x,v)) \,\mathrm{d}\, t = \frac{1}{\ell(\gamma)} \left(\int_I \pi_m^* h + \int_J \pi_m^* h \right),$$

where $I \subset [0, \ell(\gamma)]$ is the union of intervals of times t such that $\pi(\varphi_t(x, v)) \notin N, \pi : SM \to M$ denotes the projection and J is the complement of I. Observe that the integral over I is zero since h was extended by 0 outside N. Now, J splits as a union of subintervals, each of them corresponding to a segment of geodesic in N. By assumption, the integral of $\pi_m^* h$ over all these segments is 0. Hence $I_m^g h = 0$.

Since I_m^g is s-injective, we deduce that h = Du, for some $u \in C^{\infty}(M, \otimes^{m-1}T^*M)$, that is $\pi_m^* h = X\pi_{m-1}^* u$ is a coboundary. We now want to show that $u|_{\partial M} = 0$. We let $(x_0, v_0) \in SM$ be a point on the boundary ∂_+SN such that the orbit $(\varphi_t(x_0, v_0))_{t\in\mathbb{R}}$ is dense in SM and we let $c := \pi_{m-1}^* u(x_0, v_0)$. Observe that the following holds: if $t_1^- \ge 0$ denotes the first positive time such that $\varphi_{t_1^-}(x_0, v_0)$ intersects ∂_-SN° (that is, it is an inward pointing vector that is not tangent to the boundary of N), then $\pi_{m-1}^* u(\varphi_t(x_0, v_0))$ is constant equal to c for all $t \in [0, t_1^-]$. There is then a time $t_1^+ > t_1^-$ such that $\varphi_{t_1^+}(x_0, v_0) \in \partial_+SN$. For $t \in [t_1^-, t_1^+)$, the value of $\pi_{m-1}^* u(\varphi_t(x_0, v_0))$ is unknown but one has $\pi_{m-1}^* u(\varphi_{t_1^+}(x_0, v_0)) = c$ since:

$$I^{g|_{N}}h(\varphi_{t_{1}^{-}}(x_{0},v_{0})) = \int_{t_{1}^{-}}^{t_{1}^{+}} \pi_{m}^{*}h(\varphi_{t}(x_{0},v_{0})) \,\mathrm{d}\,t = 0 = \pi_{m-1}^{*}u(\varphi_{t_{1}^{+}}(x_{0},v_{0})) - \pi_{m-1}^{*}u(\varphi_{t_{1}^{-}}(x_{0},v_{0})).$$

Since the orbit of $\mathcal{O}(x_0, v_0)$ of (x_0, v_0) is dense in SM, the set $\mathcal{A} := \mathcal{O}(x_0, v_0) \cap (\partial_- SN \cup \partial_+ SN)$ is also dense in $\partial_- SN \cup \partial_+ SN$. Moreover, iterating the previous argument shows

that $\pi_{m-1}^* u|_{\mathcal{A}} = c$ and thus $\pi_{m-1}^* u|_{\partial_-SN \cup \partial_+SN} = c$. If m is even, m-1 is odd and this forces c to be 0 (just use the antipodal map $(x, v) \mapsto (x, -v)$); if m is odd, then changing at the very beginning u by $u + \lambda g^{\otimes (m-1)/2}$ for some $\lambda \in \mathbb{R}$ allows to take c = 0. Hence, we conclude that $u|_{\partial N} = 0$. This gives that f = Dp + h = D(p + u), where p + u vanishes on ∂N . \Box

This concludes the proof of Corollary 1.2.

References

- [Ano67] D. V. Anosov. Geodesic flows on closed Riemannian manifolds of negative curvature. Trudy Mat. Inst. Steklov., 90:209, 1967.
- [AR97] Yu. E. Anikonov and V. G. Romanov. On uniqueness of determination of a form of first degree by its integrals along geodesics. J. Inverse Ill-Posed Probl., 5(6):487–490 (1998), 1997.
- [BCG95] G. Besson, G. Courtois, and S. Gallot. Entropies et rigidités des espaces localement symétriques de courbure strictement négative. Geom. Funct. Anal., 5(5):731–799, 1995.
- [BI10] Dmitri Burago and Sergei Ivanov. Boundary rigidity and filling volume minimality of metrics close to a flat one. Ann. of Math. (2), 171(2):1183–1211, 2010.
- [BK85] K. Burns and A. Katok. Manifolds with nonpositive curvature. Ergodic Theory Dynam. Systems, 5(2):307–317, 1985.
- [BL07] Oliver Butterley and Carlangelo Liverani. Smooth Anosov flows: correlation spectra and stability. J. Mod. Dyn., 1(2):301–322, 2007.
- [Bon20] Yannick Guedes Bonthonneau. Perturbation of Ruelle resonances and Faure-Sjöstrand anisotropic space. Rev. Un. Mat. Argentina, 61(1):63–72, 2020.
- [CDKP20] Mihajlo Cekić, Semyon Dyatlov, Benjamin Küster, and Gabriel P. Paternain. The Ruelle zeta function at zero for nearly hyperbolic 3-manifolds. arXiv e-prints, page arXiv:2009.08558, September 2020.
- [CEG20] Dong Chen, Alena Erchenko, and Andrey Gogolev. Riemannian Anosov extension and applications. arXiv e-prints, page arXiv:2009.13665, September 2020.
- [CFF92] C. Croke, A. Fathi, and J. Feldman. The marked length-spectrum of a surface of nonpositive curvature. *Topology*, 31(4):847–855, 1992.
- [CL] Mihajlo Cekić and Thibault Lefeuvre. On transparent manifolds. in preparation.
- [CL21a] Mihajlo Cekić and Thibault Lefeuvre. Generic Dynamical Properties of Connections on Vector Bundles. International Mathematics Research Notices, 04 2021. rnab069.
- [CL21b] Mihajlo Cekić and Thibault Lefeuvre. The Holonomy Inverse Problem. arXiv e-prints, page arXiv:2105.06376, May 2021.
- [Con10] Gonzalo Contreras. Geodesic flows with positive topological entropy, twist maps and hyperbolicity. Ann. of Math. (2), 172(2):761–808, 2010.
- [Cro90] Christopher B. Croke. Rigidity for surfaces of nonpositive curvature. *Comment. Math. Helv.*, 65(1):150–169, 1990.
- [Cro91] Christopher B. Croke. Rigidity and the distance between boundary points. J. Differential Geom., 33(2):445–464, 1991.
- [CS98] Christopher B. Croke and Vladimir A. Sharafutdinov. Spectral rigidity of a compact negatively curved manifold. *Topology*, 37(6):1265–1273, 1998.
- [DGRS20] Nguyen Viet Dang, Colin Guillarmou, Gabriel Rivière, and Shu Shen. The Fried conjecture in small dimensions. Invent. Math., 220(2):525–579, 2020.
- [dHUZ19] Maarten V. de Hoop, Gunther Uhlmann, and Jian Zhai. Inverting the local geodesic ray transform of higher rank tensors. *Inverse Problems*, 35(11):115009, 27, 2019.
- [DP03] Victor J. Donnay and Charles C. Pugh. Anosov geodesic flows for embedded surfaces. Number 287, pages xviii, 61–69. 2003. Geometric methods in dynamics. II.

- [DS03] Nurlan S. Dairbekov and Vladimir A. Sharafutdinov. Some problems of integral geometry on Anosov manifolds. *Ergodic Theory Dynam. Systems*, 23(1):59–74, 2003.
- [DS10] N. S. Dairbekov and V. A. Sharafutdinov. Conformal Killing symmetric tensor fields on Riemannian manifolds. Mat. Tr., 13(1):85–145, 2010.
- [DZ16] Semyon Dyatlov and Maciej Zworski. Dynamical zeta functions for Anosov flows via microlocal analysis. Ann. Sci. Éc. Norm. Supér. (4), 49(3):543–577, 2016.
- [Ebe73] Patrick Eberlein. When is a geodesic flow of Anosov type? I,II. J. Differential Geometry, 8:437–463; ibid. 8 (1973), 565–577, 1973.
- [FRS08] Frédéric Faure, Nicolas Roy, and Johannes Sjöstrand. Semi-classical approach for Anosov diffeomorphisms and Ruelle resonances. Open Math. J., 1:35–81, 2008.
- [FS11] Frédéric Faure and Johannes Sjöstrand. Upper bound on the density of Ruelle resonances for Anosov flows. Comm. Math. Phys., 308(2):325–364, 2011.
- [FT13] Frédéric Faure and Masato Tsujii. Band structure of the Ruelle spectrum of contact Anosov flows. C. R. Math. Acad. Sci. Paris, 351(9-10):385–391, 2013.
- [GK80a] V. Guillemin and D. Kazhdan. Some inverse spectral results for negatively curved 2-manifolds. Topology, 19(3):301–312, 1980.
- [GK80b] Victor Guillemin and David Kazhdan. Some inverse spectral results for negatively curved nmanifolds. In Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), Proc. Sympos. Pure Math., XXXVI, pages 153–180. Amer. Math. Soc., Providence, R.I., 1980.
- [GKL19] Colin Guillarmou, Gerhard Knieper, and Thibault Lefeuvre. Geodesic stretch, pressure metric and marked length spectrum rigidity. *arXiv e-prints*, page arXiv:1909.08666, Sep 2019.
- [GL06] Sébastien Gouëzel and Carlangelo Liverani. Banach spaces adapted to Anosov systems. Ergodic Theory Dynam. Systems, 26(1):189–217, 2006.
- [GL19] Colin Guillarmou and Thibault Lefeuvre. The marked length spectrum of Anosov manifolds. Ann. of Math. (2), 190(1):321–344, 2019.
- [GL21] Sébastien Gouëzel and Thibault Lefeuvre. Classical and microlocal analysis of the x-ray transform on Anosov manifolds. Anal. PDE, 14(1):301–322, 2021.
- [GPSU16] Colin Guillarmou, Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. The X-ray transform for connections in negative curvature. Comm. Math. Phys., 343(1):83–127, 2016.
- [Gro83] Mikhael Gromov. Filling Riemannian manifolds. J. Differential Geom., 18(1):1–147, 1983.
- [Guil7a] Colin Guillarmou. Invariant distributions and X-ray transform for Anosov flows. J. Differential Geom., 105(2):177–208, 2017.
- [Gui17b] Colin Guillarmou. Lens rigidity for manifolds with hyperbolic trapped sets. J. Amer. Math. Soc., 30(2):561–599, 2017.
- [GWW92] C. Gordon, D. Webb, and S. Wolpert. Isospectral plane domains and surfaces via Riemannian orbifolds. *Invent. Math.*, 110(1):1–22, 1992.
- [Ham99] U. Hamenstädt. Cocycles, symplectic structures and intersection. Geom. Funct. Anal., 9(1):90–140, 1999.
- [Hel00] Sigurdur Helgason. Groups and geometric analysis. Integral geometry, invariant differential operators, and spherical functions, volume 83. Providence, RI: American Mathematical Society (AMS), 2000.
- [Hör65] Lars Hörmander. Pseudo-differential operators. Comm. Pure Appl. Math., 18:501–517, 1965.
- [Hör03] Lars Hörmander. The analysis of linear partial differential operators. I. Classics in Mathematics. Springer-Verlag, Berlin, 2003. Distribution theory and Fourier analysis, Reprint of the second (1990) edition [Springer, Berlin; MR1065993 (91m:35001a)].
- [Kac66] Mark Kac. Can one hear the shape of a drum? Amer. Math. Monthly, 73(4, part II):1–23, 1966.
- [Kat88] Anatole Katok. Four applications of conformal equivalence to geometry and dynamics. Ergodic Theory Dynam. Systems, 8* (Charles Conley Memorial Issue):139–152, 1988.

M. CEKIĆ AND T. LEFEUVRE

- [Kli74] Wilhelm Klingenberg. Riemannian manifolds with geodesic flow of Anosov type. Ann. of Math. (2), 99:1–13, 1974.
- [Kou15] Mickaël Kourganoff. *Geometry and dynamics of configuration spaces*. Theses, Ecole normale supérieure de lyon ENS LYON, December 2015.
- [Lef19a] Thibault Lefeuvre. On the s-injectivity of the X-ray transform on manifolds with hyperbolic trapped set. Nonlinearity, 32(4):1275–1295, 2019.
- [Lef19b] Thibault Lefeuvre. Sur la rigidité des variétés riemanniennes. PhD thesis, 2019. Thèse de doctorat dirigée par Guillarmou, Colin Mathématiques fondamentales Université Paris-Saclay (ComUE) 2019.
- [Lef20] Thibault Lefeuvre. Local marked boundary rigidity under hyperbolic trapping assumptions. J. Geom. Anal., 30(1):448–465, 2020.
- [Liv04] Carlangelo Liverani. On contact Anosov flows. Ann. of Math. (2), 159(3):1275–1312, 2004.
- [Mel03] Richard B. Melrose. Introduction to Microlocal Analysis. 2003. Available online: http://www-math.mit.edu/~rbm/iml90.pdf.
- [Mic82] René Michel. Sur la rigidité imposée par la longueur des géodésiques. Invent. Math., 65(1):71–83, 1981/82.
- [Mil64] J. Milnor. Eigenvalues of the Laplace operator on certain manifolds. Proc. Nat. Acad. Sci. U.S.A., 51:542, 1964.
- [MP11] Will Merry and Gabriel P. Paternain. *Inverse Problems in Geometry and Dynamics*. Lecture notes, 2011. Available online: https://www.dpmms.cam.ac.uk/~gpp24/ipgd(3).pdf.
- [MR78] R. G. Mukhometov and V. G. Romanov. On the problem of finding an isotropic Riemannian metric in an n-dimensional space. Dokl. Akad. Nauk SSSR, 243(1):41–44, 1978.
- [Muk77] R. G. Mukhometov. The reconstruction problem of a two-dimensional Riemannian metric, and integral geometry. Dokl. Akad. Nauk SSSR, 232(1):32–35, 1977.
- [Muk81] R. G. Mukhometov. On a problem of reconstructing Riemannian metrics. Sibirsk. Mat. Zh., 22(3):119–135, 237, 1981.
- [Ota90a] Jean-Pierre Otal. Le spectre marqué des longueurs des surfaces à courbure négative. Ann. of Math. (2), 131(1):151–162, 1990.
- [Ota90b] Jean-Pierre Otal. Sur les longueurs des géodésiques d'une métrique à courbure négative dans le disque. *Comment. Math. Helv.*, 65(2):334–347, 1990.
- [Pat99] Gabriel P. Paternain. Geodesic flows, volume 180 of Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 1999.
- [Pat09] Gabriel P. Paternain. Transparent connections over negatively curved surfaces. J. Mod. Dyn., 3(2):311–333, 2009.
- [Pat12] Gabriel P. Paternain. Transparent pairs. J. Geom. Anal., 22(4):1211–1235, 2012.
- [Pat13] Gabriel P. Paternain. Inverse problems for connections. In Inverse problems and applications: inside out. II, volume 60 of Math. Sci. Res. Inst. Publ., pages 369–409. Cambridge Univ. Press, Cambridge, 2013.
- [PS87] L. N. Pestov and V. A. Sharafutdinov. Integral geometry of tensor fields on a manifold of negative curvature. Dokl. Akad. Nauk SSSR, 295(6):1318–1320, 1987.
- [PSU13] Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. Tensor tomography on simple surfaces. Invent. Math., 193(1):229–247, 2013.
- [PSU14] Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. Tensor tomography: progress and challenges. Chin. Ann. Math. Ser. B, 35(3):399–428, 2014.
- [PU05] Leonid Pestov and Gunther Uhlmann. Two dimensional compact simple Riemannian manifolds are boundary distance rigid. Ann. of Math. (2), 161(2):1093–1110, 2005.
- [Sar90] Peter Sarnak. Determinants of Laplacians; heights and finiteness. In Analysis, et cetera, pages 601–622. Academic Press, Boston, MA, 1990.
- [Sha02] V. A. Sharafutdinov. A problem in integral geometry in a nonconvex domain. Sibirsk. Mat. Zh., 43(6):1430–1442, 2002.

56

- [Shu01] M. A. Shubin. Pseudodifferential operators and spectral theory. Springer-Verlag, Berlin, second edition, 2001. Translated from the 1978 Russian original by Stig I. Andersson.
- [SU05] Plamen Stefanov and Gunther Uhlmann. Boundary rigidity and stability for generic simple metrics. J. Amer. Math. Soc., 18(4):975–1003, 2005.
- [SU08] Plamen Stefanov and Gunther Uhlmann. Integral geometry on tensor fields on a class of non-simple Riemannian manifolds. *Amer. J. Math.*, 130(1):239–268, 2008.
- [SUV18] Plamen Stefanov, Gunther Uhlmann, and András Vasy. Inverting the local geodesic X-ray transform on tensors. J. Anal. Math., 136(1):151–208, 2018.
- [Tay91] Michael E. Taylor. Pseudodifferential operators and nonlinear PDE, volume 100 of Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 1991.
- [Uhl76] K. Uhlenbeck. Generic properties of eigenfunctions. Amer. J. Math., 98(4):1059–1078, 1976.
- [UV16] Gunther Uhlmann and András Vasy. The inverse problem for the local geodesic ray transform. Invent. Math., 205(1):83–120, 2016.
- [Vig80] Marie-France Vignéras. Variétés riemanniennes isospectrales et non isométriques. Ann. of Math. (2), 112(1):21–32, 1980.
- [Zwo12] Maciej Zworski. Semiclassical analysis, volume 138 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2012.

LABORATOIRE DE MATHÉMATIQUES D'ORSAY, UNIV. PARIS-SUD, CNRS, UNIVERSITÉ PARIS-SACLAY, 91405 ORSAY, FRANCE

Email address: mihajlo.cekic@u-psud.fr

UNIVERSITÉ DE PARIS AND SORBONNE UNIVERSITÉ, CNRS, IMJ-PRG, F-75006 PARIS, FRANCE. Email address: tlefeuvre@imj-prg.fr