SCATTERING THEORY FOR QUADRATIC NONLINEAR SCHRÖDINGER SYSTEM IN DIMENSION SIX

CHUANWEI GAO, FANFEI MENG, CHENGBIN XU, AND JIQIANG ZHENG

ABSTRACT. In this paper, we study the solutions to the energy-critical quadratic nonlinear Schrödinger system in $\dot{H}^1 \times \dot{H}^1$, where the sign of its potential energy can not be determined directly. If the initial data u_0 is radial or non-radial but satisfies the mass-resonance condition, and its energy is below that of the ground state, using the compactness/rigidity method, we give a complete classification of scattering versus blowing-up dichotomies depending on whether the kinetic energy of u_0 is below or above that of the ground state.

1. INTRODUCTION

We consider the quadratic nonlinear Schrödinger system:

$$\begin{cases} i\partial_t \mathbf{u} + A\mathbf{u} + \mathbf{f}(\mathbf{u}) = \mathbf{0}, & (t, x) \in \mathbb{R} \times \mathbb{R}^d, \\ \mathbf{u}(0, x) = \mathbf{u}_0(x), \end{cases}$$
(NLS system)

where \mathbf{u} , \mathbf{u}_0 , and $\mathbf{0}$ are all vector-valued complex functions with two components defined as follows

$$\mathbf{u} := \begin{pmatrix} u \\ v \end{pmatrix}, \ \mathbf{u}_{\mathbf{0}} := \begin{pmatrix} u_0 \\ v_0 \end{pmatrix}, \ \mathbf{0} := \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

and A is a 2×2 matrix, $\mathbf{f} : \mathbb{C}^2 \to \mathbb{C}^2$ as follows

$$A := \begin{pmatrix} \Delta & 0\\ 0 & \kappa \Delta \end{pmatrix}, \ \mathbf{f}(\mathbf{u}) := \begin{pmatrix} v\overline{u}\\ u^2 \end{pmatrix}, \tag{1.1}$$

 $u, v : \mathbb{R} \times \mathbb{R}^d \to \mathbb{C}$ are unknown functions, $\kappa \in (0, \infty)$ is a real number and Δ denotes the Laplacian in \mathbb{R}^d .

From a physical point view, (NLS system) is related to the Raman amplification in a plasma. This process is a nonlinear instability phenomenon (see [3, 4, 21, 22] for more details). Solutions to (NLS system) preserve the *mass*, *energy* and *momentum*, defined respectively by

$$M(\mathbf{u}) := \|u\|_{L^2}^2 + \|v\|_{L^2}^2 \equiv M(\mathbf{u}_0),$$

$$E(\mathbf{u}) := H(\mathbf{u}) - R(\mathbf{u}) \equiv E(\mathbf{u}_0),$$

$$P(\mathbf{u}) := \operatorname{Im} \int_{\mathbb{R}^d} \left(\overline{u}\nabla u + \frac{1}{2}\overline{v}\nabla v\right) \mathrm{d}x \equiv P(\mathbf{u}_0),$$

Date: March 19, 2022.

²⁰¹⁰ Mathematics Subject Classification. Primary 35Q55.

Key words and phrases. Energy-critical, quadratic nonlinear Schrödinger system, blow-up, scatter.

where

(kinetic energy)
$$H(\mathbf{u}) := \|u\|_{\dot{H}^1}^2 + \frac{\kappa}{2} \|v\|_{\dot{H}^1}^2$$

(potential energy)
$$R(\mathbf{u}) := \operatorname{Re} \int_{\mathbb{R}^d} \overline{v} u^2 \mathrm{d} x.$$

The equation (NLS system) is invariant under the scaling

$$\mathbf{u}_{\lambda}(t,x) = \lambda^2 \mathbf{u} \left(\lambda^2 t, \lambda x\right)$$

for any $\lambda > 0$. The critical regularity of Sobolev space is $\dot{\mathbf{H}}^{s_c}$, i.e. $\|\mathbf{u}_{\lambda}\|_{\dot{\mathbf{H}}^{s_c}_x(\mathbb{R}^d)} = \|\mathbf{u}\|_{\dot{\mathbf{H}}^{s_c}_x(\mathbb{R}^d)}$ where $s_c = \frac{d}{2} - 2$. Therefore, the equation (NLS system) is called masssubcritical if $d \leq 3$, mass-critical if d = 4, energy-subcritical if d = 5, and energycritical if d = 6. We say (NLS system) satisfies the mass-resonance condition if $\kappa = \frac{1}{2}$. A distinguished and exclusive feature of this case is that (NLS system) is left invariant under the following Galilean transformation

$$\begin{pmatrix} u(t,x) \\ v(t,x) \end{pmatrix} \to \begin{pmatrix} e^{ix \cdot \xi} e^{-it|\xi|^2} u(t,x-2t\xi) \\ e^{2ix \cdot \xi} e^{-2it|\xi|^2} v(t,x-2t\xi) \end{pmatrix}$$

for any $\xi \in \mathbb{R}^d$.

From a mathematical point view, even though the fact $A \leq 0$ (i.e. $\langle A\mathbf{u}, \mathbf{u} \rangle \leq 0$, $\forall \mathbf{u} \in \dot{\mathrm{H}}^{1}(\mathbb{R}^{d})$) superficially suggests that (NLS system) is similar to the classical nonlinear Schrödinger equation:

$$\partial_t u + \Delta u + \mu |u|^{p-1} u = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^d,$$
 (NLS)

the truth is quite different. One prominent difference among others is that there is no classification of the so-called focusing ($\mu = 1$) and defocusing ($\mu = -1$) in (NLS system), which can be seen from the fact the sign of potential energy $R(\mathbf{u})$ can not be judged directly from μ .

In this paper, we consider the energy-critical case of (NLS system) in the critical Sobolev space, that is, d = 6 and $\mathbf{u}_0 \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$. For any $s \ge 0$, we use $\dot{\mathrm{H}}^s$ to denote Hilbert space $\dot{H}^s \times \dot{H}^s$, and H^s to denote Hilbert space $H^s \times H^s$. For any $p \ge 1$, we use L^p to denote $L^p \times L^p$. We also denote the Schrödinger group $\mathbf{S}(t)$ as follows

$$\mathbf{S}(t) := \begin{pmatrix} e^{it\Delta} & 0\\ 0 & e^{\kappa it\Delta} \end{pmatrix}$$

By solution, we mean a function $\mathbf{u} \in C_t(I, \dot{\mathrm{H}}^1_x(\mathbb{R}^6)) \cap \mathrm{L}^4_{t,x}(I \times \mathbb{R}^6)$ on an interval $I \ni 0$ satisfying the Duhamel formula

$$\mathbf{u}(t) = \mathbf{S}(t)\mathbf{u}_0 + i\int_0^t \mathbf{S}(t-\tau)\mathbf{f}(\mathbf{u}(\tau))\mathrm{d}\tau$$

for $t \in I$. If the maximal interval of existence $I_{\max} = (T_{-}(\mathbf{u}), T_{+}(\mathbf{u})) = \mathbb{R}$, we call the solution \mathbf{u} is global.

We say that a solution **u** to (NLS system) blows up forward in time if there exists a time $t_1 \in I_{\text{max}} = (T_{-}(\mathbf{u}), T_{+}(\mathbf{u}))$ such that

$$S_{[t_1,T_+(\mathbf{u}))}(\mathbf{u}) = \infty$$

that **u** blows up backward in time if there exists a time $t_1 \in I_{\text{max}}$ such that

$$S_{(T_{-}(\mathbf{u}),t_{1}]}(\mathbf{u}) = \infty$$

 $\mathbf{2}$

and that a global solution **u** scatters if there exists $\mathbf{u}_{\pm} \in \dot{\mathrm{H}}^{1}(\mathbb{R}^{6})$ such that

$$\lim_{t \to \pm\infty} \|\mathbf{u}(t) - \mathbf{S}(t)\mathbf{u}_{\pm}\|_{\dot{\mathrm{H}}^{1}(\mathbb{R}^{6})} = 0$$

Definition 1.1 (Trivial scattering solution, [23]). We say a solution to a nonlinear equation is a *trivial scattering solution* if it also solves the corresponding linear equation. (In other words, the equation does not have nonlinear interaction.)

Remark 1.2. For (NLS), zero is the only trivial scattering solution. While for (NLS system), $(u, v) = (0, e^{\kappa i t \Delta} v_0)$ is a trivial scattering solution for any $v_0 \in \dot{H}^1(\mathbb{R}^6)$.

Definition 1.3 (Scattering size, [36]). The scattering size of a solution $\mathbf{u} = (u, v)^T$ to (NLS system) on the interval $I \ni 0$ is

$$S_{I}(\mathbf{u}) = \|\mathbf{u}\|_{\mathbf{L}^{4}_{t,x}(I \times \mathbb{R}^{6})}^{4} \sim \int_{I} \int_{\mathbb{R}^{6}} |u(t,x)|^{4} + |v(t,x)|^{4} dx dt.$$

Remark 1.4 ($\dot{\mathrm{H}}^1$ sacttering). It is easy to verify that the solution to (NLS system) will scatter in $\dot{\mathrm{H}}^1(\mathbb{R}^6)$ as $t \to +\infty$ if the solution $\mathbf{u}(t)$ to (NLS system) is global in $\dot{\mathrm{H}}^1(\mathbb{R}^6)$ with finite scattering size on \mathbb{R} , i.e., $S_{\mathbb{R}}(\mathbf{u}) < +\infty$.

The scattering solutions to (NLS) are attractive to many researchers, such as [7, 8, 13, 26] for the focusing case and [27, 35] for the defocusing case and so on. A quite interesting problem is the defocusing energy-critical case, i.e. $p = 1 + \frac{4}{d-2}$, one may refer to [1, 5, 19, 33] for more details.

For (NLS system) and its related models, there are amount of literature on those themes, such as [31, 32, 34] for three waves interaction and [12, 14, 29] for quadratic-type interaction. In [11, 16, 30], the authors have studied the problems of (NLS system) that arise in 2D. There are also many papers on the well-posedness and dynamics of the solutions to (NLS system) with data below the ground state \mathbf{Q} in dimensions, $1 \leq d \leq 5$, where $\mathbf{Q} := (\phi, \varphi)^T \in D_d$ is a pair of non-negative real-valued functions satisfying:

$$\begin{cases} \phi - \Delta \phi = \phi \varphi, \\ 2\varphi - \kappa \Delta \varphi = \phi^2, \end{cases} \quad x \in \mathbb{R}^d, \tag{1.2}$$

with $D_d := \{ \mathbf{G} \in \mathrm{H}^1(\mathbb{R}^d) \mid R(\mathbf{G}) > 0 \}$. Among others, up to translations, Hayashi, Li and Naumkin [11] proved the uniqueness of $\mathbf{Q} := (\phi_1, \frac{\phi_1}{\sqrt{2}})^T$ when $\kappa = 2$, where $\phi_1 \neq 0$ is the unique non-negative radial real-valued solution to

$$\phi_1 - \Delta \phi_1 = \frac{1}{\sqrt{2}}\phi_1^2.$$

With and without mass-resonance, the mass-critical case (d = 4) was studied in [15] and the energy-subcritical case (d = 5) was studied separately in [9, 10, 24, 28, 37].

In this paper, we concern about the energy-critical case (d = 6) of (NLS system). Our results show the classification of the solution as to scattering or blowing-up depending on the relationship between the initial data u_0 and the ground state $\mathbf{W} := (\phi, \varphi)^T \in D_6$, where $(\phi, \varphi)^t$ is a pair of non-negative real-valued functions satisfying:

$$\begin{cases} -\Delta\phi = \phi\varphi, \\ -\kappa\Delta\varphi = \phi^2, \end{cases} \quad x \in \mathbb{R}^6, \tag{1.3}$$

and $D_6 := \left\{ \mathbf{G} \in \dot{\mathrm{H}}^1(\mathbb{R}^6) \cap \mathrm{L}^3(\mathbb{R}^6) \mid R(\mathbf{G}) > 0 \right\}.$

At this point, some useful information of W are discussed. Define the operator $T: (\dot{H}^1 \cap L^3)_r(\mathbb{R}^6) \to H^1(\mathbb{R})$ as follows:

$$(Tf)(t) := e^{2t} f(e^t), \qquad t \in \mathbb{R}$$

By showing the map T gives a one-to-one correspondence between the ground states of (1.2) and those of (1.3), up to translations and dilations, [11] proved the uniqueness of \mathbf{W} for any $\kappa \in (0, \infty)$. Therefore, $\mathbf{W} = (\phi_0, \frac{\phi_0}{\sqrt{\kappa}})^T$, where $\phi_0 \neq 0$ is the unique non-negative radial real-valued solution to

$$-\Delta\phi_0 = \frac{1}{\sqrt{\kappa}}\phi_0^2. \tag{1.4}$$

Furthermore, we may write ϕ_0 explicitly as follows

$$\phi_0(x) = \frac{\sqrt{\kappa}}{\left(1 + \frac{|x|^2}{24}\right)^2} \in \dot{H}^1(\mathbb{R}^6)$$

one may refer to [18] for more details.

Our main results read as follows:

Theorem 1.5. For d = 6, $\mathbf{u}_0 \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$, and $E(\mathbf{u}_0) < E(\mathbf{W})$ in the (NLS system),

- (1) if $H(\mathbf{u}_0) < H(\mathbf{W})$, then \mathbf{u} scatters for \mathbf{u}_0 is non-radial with the massresonance condition or \mathbf{u}_0 is radial;
- (2) if $H(\mathbf{u_0}) > H(\mathbf{W})$, then \mathbf{u} blows up in finite time for u_0 is radial with $x\mathbf{u}_0 \in L^2(\mathbb{R}^6)$ or $\mathbf{u}_0 \in H^1(\mathbb{R}^6)$.

Remark 1.6. From the proof of Proposition 3.4, the restriction $E(\mathbf{u_0}) < E(\mathbf{W})$ implies $H(\mathbf{u_0}) \neq H(\mathbf{W})$, which indicates the completeness of our classification.

Remark 1.7 (Reason of the mass-resonance condition). The mass-resonance condition enables us to use the Galilean invariance of the critical solution \mathbf{u}_c to (NLS system) which is very useful in proving zero momentum $P(\mathbf{u}_c(t)) \equiv \mathbf{0}$ for the non-radial case and is only available when $\kappa = \frac{1}{2}$, that is the essential reason why we require $\kappa = \frac{1}{2}$ for the non-radial case in Theorem 1.5. One can refer to the proof of Proposition 6.5 in this paper or the statement of Remark 1.2 in [24] for more details. Under the assumption \mathbf{u}_0 is non-radial with the mass-resonance condition or \mathbf{u}_0 is radial, the scattering part of the results in Theorem 1.5 is consistent with that of the mass-critical case (d = 4) and the energy-subcritical case (d = 5). One can refer to [9, 10, 15, 24, 28, 37].

To prove the blowing-up result in Theorem 1.5, we exploit the Virial identity in Subsection 3.3. To prove the scattering result, we follow the strategies from [1, 18, 33] through analysing a kind of special solution, so called "critical solution", in Section 5 and Section 6. After a series of reductions, the "critical solution" shares the property of almost periodicity modulo symmetries:

Definition 1.8 (Almost periodic solution, [36]). We call a solution $\mathbf{u} = (u, v)^T$ to (NLS system) is almost periodic modulo symmetries with lifespan $I \ni 0$, if there exist functions $\lambda : I \to \mathbb{R}^+$, $x : I \to \mathbb{R}^6$, and $C : \mathbb{R}^+ \to \mathbb{R}^+$ such that

$$\int_{|x-x(t)| \ge \frac{C(\eta)}{\lambda(t)}} \left(|\nabla u(t,x)|^2 + \frac{\kappa}{2} |\nabla v(t,x)|^2 \right) \mathrm{d}x \le \eta$$

and

$$\int_{|\xi| \ge C(\eta)\lambda(t)} \left(|\xi|^2 |\hat{u}(t,\xi)|^2 + \frac{\kappa}{2} |\xi|^2 |\hat{v}(t,\xi)|^2 \right) \mathrm{d}\xi \leqslant \eta$$

for all $t \in I$ and $\eta > 0$. We refer to the function λ as the frequency scale function, x as the spatial center function, and C as the compactness modulus function.

From the Arzela-Ascoli Theorem, we have

Lemma 1.9 (Compactness in L^2). The set $\mathcal{F} \subset L^2(\mathbb{R}^d)$ is compact in $L^2(\mathbb{R}^d)$ if and only if the following conditions hold:

- (i) there exists A > 0 such that $||f||_{L^2(\mathbb{R}^d)} \leq A$, $\forall f \in \mathcal{F}$;
- (ii) for any $\varepsilon > 0$, there exists $R = R(\varepsilon) > 0$ such that

$$\int_{|x|\geqslant R} |f(x)|^2 \mathrm{d}x + \int_{|\xi|\geqslant R} |\hat{f}(\xi)|^2 \mathrm{d}\xi < \varepsilon, \qquad \forall \ f \in \mathcal{F}.$$

Remark 1.10. By Lemma 1.9, the set $\mathcal{F}_1 \subset \dot{H}^1(\mathbb{R}^d)$ is compact in $\dot{H}^1(\mathbb{R}^d)$ if and only if \mathcal{F}_1 is bounded in \dot{H}^1 uniformly and for any $\eta > 0$, there exists a compactness modulus function $C(\eta) > 0$ such that

$$\int_{|x| \ge C(\eta)} |\nabla f(x)|^2 \mathrm{d}x + \int_{|\xi| \ge C(\eta)} |\xi|^2 |\hat{f}(\xi)|^2 \mathrm{d}\xi < \eta, \qquad \forall \ f \in \mathcal{F}_1.$$

By using the Palais-Smale condition modulo symmetries, Proposition 4.9, we know that the critical solutions have to be almost periodic and minimal-kineticenergy blowing-up which is proven in Subsection 4.3 and stated as follows:

Theorem 1.11. If the scattering result of Theorem 1.5 fails, then there exists a solution $\mathbf{u}_c: I_c \times \mathbb{R}^6 \to \mathbb{C}^2$ with the maximal interval of existence I_c satisfying

$$\sup_{t \in I_c} H(\mathbf{u}_c(t)) < H(\mathbf{W})$$

which is almost periodic modulo symmetries and blows up in both directions. Furthermore, \mathbf{u}_c has the minimal kinetic energy among all the blowing-up solutions, that is,

$$\sup_{t\in I_c} H(\mathbf{u}_c(t)) \leqslant \sup_{t\in I} H(\mathbf{u}(t)),$$

where $\mathbf{u}: I \times \mathbb{R}^6 \to \mathbb{C}^2$ is a solution to (NLS system) with the maximal interval of existence I and blows up in at least one direction.

For the blow-up solutions \mathbf{u}_c to (NLS system) in Theorem 1.11, we can classify them by their different properties of the frequency scale function $\lambda(t) : I_c \to \mathbb{R}^+$.

Proposition 1.12 (Classification of critical solution, [19, 20]). If the scattering result of Theorem 1.5 fails, then there exists a minimal-kinetic-energy blowing-up solution $\mathbf{u}_c : I_c \times \mathbb{R}^6 \to \mathbb{C}^2$ which is almost periodic modulo symmetries with the maximal interval of existence I_c satisfying

$$S_{I_c}(\mathbf{u_c}) = \infty, \quad and \quad \sup_{t \in I_c} H(\mathbf{u}_c(t)) < H(\mathbf{W}).$$
 (1.5)

Furthermore, \mathbf{u}_c has to be one of the following three cases:

(1) a blowing-up solution in finite time if

 $|\inf I_c| < \infty, \quad or \quad \sup I_c < \infty;$

(2) a soliton-like solution if

$$I_c = \mathbb{R}, \quad and \quad \lambda(t) = 1, \ \forall \ t \in \mathbb{R};$$

(3) a low-to-high frequency cascade if

$$I_c = \mathbb{R}, \quad and \quad \inf_{t \in \mathbb{R}} \lambda(t) = 1, \quad \limsup_{t \to +\infty} \lambda(t) = \infty.$$

After clarifying our three enemies, we need to rule out them according to their respective properties. We will prove that each critical solution has to be **0** and eliminate the possibility of their existence one by one. For the first enemy, blowing-up solution in finite time, we analysis its compactness and use the Virial identity to obtain $\mathbf{u}_c \equiv \mathbf{0}$ in Section 5.

For the second and third enemies, by using the reduced Duhamel formulas(see Proposition 6.1 below), we will show that they enjoy the following negative regularity in Subsection 6.1.

Theorem 1.13 (Negative regularity). Let \mathbf{u}_c be as in Proposition 1.12 with $I_c = \mathbb{R}$. If

$$\sup_{t\in\mathbb{R}}H(\mathbf{u}_c(t))<\infty\quad and\quad \inf_{t\in\mathbb{R}}\lambda(t)\geq 1,$$

then there exists $\varepsilon > 0$ such that $\mathbf{u}_c \in \mathrm{L}^{\infty}_t(\mathbb{R}, \dot{\mathrm{H}}^{-\varepsilon}(\mathbb{R}^6)).$

It is easy to see that the momentum of any radial function is always zero. For the non-radial case, with additional mass-resonance condition, the above fact can also be recovered. This observation enables us to deduce $P(\mathbf{u}_c) \equiv \mathbf{0}$, if $\mathbf{u}_{c,0}$ is non-radial with the mass-resonance condition or $\mathbf{u}_{c,0}$ is radial, which is shown in Proposition 6.5. The zero momentum property helps us to obtain the control of spatial center function x(t) of the soliton-like solution in Corollary 6.6.

By Theorem 1.13 and Corollary 6.6, we use the Virial identity to eliminate the possibility of soliton-like solution. Finally, in order to exclude the last enemy, we divide its frequencies into low and high parts, and conquer them respectively. To be more precise, for the low frequency, we will use the negative regularity, and for the high frequency, we will use the property of low-to-high frequency cascade.

1.1. Outline of the paper. The following part of this paper is organized as follows: In Section 2, we clarify some preliminaries, including Bernstein inequalities, Strichartz estimate, local well-posedness, and long time perturbation theory of (NLS system). Section 3 are divided into three parts. We focus on ground state in Subsection 3.1, show the energy trapping in Subsection 3.2, and give the proof of blowing-up result of Theorem 1.5 in Subsection 3.3. Section 4 concentrates on the proof of Theorem 1.11 by the properties of the linear profile decomposition and the nonlinear profile. To obtain the scattering result, we rule out our three enemies arisen in Proposition 1.12 one by one in Section 5 and Section 6. Combining with the result of blowing-up in finite time in Theorem 3.10 and Theorem 3.11, we complete the proof of Theorem 1.5 finally.

2. Preliminaries

We mark $A \leq B$ to mean there exists a constant C > 0 such that $A \leq CB$. We indicate dependence on parameters via subscripts, e.g. $A \leq_x B$ indicates $A \leq CB$ for some C = C(x) > 0. We use $A \sim B$ to denote $A \leq B \leq A$.

 $\mathbf{6}$

Let $\phi(\xi)$ be a radial smooth function supported in the ball $\{\xi \in \mathbb{R}^6 : |\xi| \leq \frac{11}{10}\}$ and equal to 1 on the ball $\{\xi \in \mathbb{R}^6 : |\xi| \leq 1\}$. For each number N > 0, we define the Fourier multipliers

$$\begin{split} \widehat{P_{\leqslant N}\mathbf{g}}(\xi) &:= \phi\left(\frac{\xi}{N}\right) \hat{\mathbf{g}}(\xi), \\ \widehat{P_{>N}\mathbf{g}}(\xi) &:= \left(1 - \phi\left(\frac{\xi}{N}\right)\right) \hat{\mathbf{g}}(\xi), \\ \widehat{P_{N}\mathbf{g}}(\xi) &:= \left(\phi\left(\frac{\xi}{N}\right) - \phi\left(\frac{2\xi}{N}\right)\right) \hat{\mathbf{g}}(\xi), \end{split}$$

and similarly $P_{\leq N}$ and $P_{\geq N}$. We also define

$$P_{M < \cdot \leqslant N} := P_{\leqslant N} - P_{\leqslant M} = \sum_{M < N' \leqslant N} P_{N'}$$

whenever M < N. We will usually use these multipliers when M and N are *dyadic* numbers (that is, of the form 2^n for some integer n); in particular, all summations over N or M are understood to be over dyadic numbers. Nevertheless, it will occasionally be convenient to allow M and N to not be a power of 2. Like all Fourier multipliers, the Littlewood-Paley operators commute with the propagator $\mathbf{S}(t)$, as well as with differential operators such as $i\partial_t + A$.

Lemma 2.1 (Bernstein inequalities). For any $1 \leq p \leq q \leq \infty$ and $s \geq 0$,

$$\begin{split} \|P_{\geqslant N}\mathbf{g}\|_{\mathbf{L}^{p}_{x}(\mathbb{R}^{6})} &\lesssim N^{-s} \||\nabla|^{s} P_{\geqslant N}\mathbf{g}\|_{\mathbf{L}^{p}_{x}(\mathbb{R}^{6})} \lesssim N^{-s} \|\|\nabla|^{s}\mathbf{g}\|_{\mathbf{L}^{p}_{x}(\mathbb{R}^{6})},\\ \||\nabla|^{s} P_{\leqslant N}\mathbf{g}\|_{\mathbf{L}^{p}_{x}(\mathbb{R}^{6})} &\lesssim N^{s} \|P_{\leqslant N}\mathbf{g}\|_{\mathbf{L}^{p}_{x}(\mathbb{R}^{6})} \lesssim N^{s} \|\mathbf{g}\|_{\mathbf{L}^{p}_{x}(\mathbb{R}^{6})},\\ \||\nabla|^{\pm s} P_{N}\mathbf{g}\|_{\mathbf{L}^{p}_{x}(\mathbb{R}^{6})} &\lesssim N^{\pm s} \|P_{N}\mathbf{g}\|_{\mathbf{L}^{p}_{x}(\mathbb{R}^{6})} &\lesssim N^{\pm s} \|\mathbf{g}\|_{\mathbf{L}^{p}_{x}(\mathbb{R}^{6})},\\ \|P_{N}\mathbf{g}\|_{\mathbf{L}^{q}_{x}(\mathbb{R}^{6})} &\lesssim N^{\frac{6}{p} - \frac{6}{q}} \|P_{N}\mathbf{g}\|_{\mathbf{L}^{p}_{x}(\mathbb{R}^{6})}, \end{split}$$

where $|\nabla|^s$ is the classical fractional-order operator.

Lemma 2.2 (Gronwall-type inequality, [19]). Fix $\gamma > 0$. Given $0 < \eta < (1-2^{-\gamma})/2$ and $\{b_k\} \in l^{\infty}(\mathbb{N})$, let $\{x_k\} \in l^{\infty}(\mathbb{N})$ be a non-negative sequence obeying

$$x_k \leqslant b_k + \eta \sum_{l=0}^{\infty} 2^{-\gamma|k-l|} x_l \qquad for \quad all \quad k \ge 0.$$

$$(2.1)$$

Then

$$x_k \lesssim \sum_{l=0}^{\infty} r^{|k-l|} b_l \tag{2.2}$$

for some $r = r(\eta) \in (2^{-\gamma}, 1)$. Moreover, $r \downarrow 2^{-\gamma}$ as $\eta \downarrow 0$.

2.1. Strichartz estimate. Let Λ_s be the set of pairs (p,q) with $q \ge 2$ and satisfying

$$\frac{2}{q} = 6\left(\frac{1}{2} - \frac{1}{r}\right) - s.$$
 (2.3)

Define

$$\|\mathbf{u}\|_{\mathcal{S}(\dot{\mathrm{H}}^{s}(\mathbb{R}^{6}))} := \sup_{(q,r)\in\Lambda_{s}}\|\mathbf{u}\|_{\mathrm{L}^{q}_{t}(\mathbb{R},\mathrm{L}^{r}_{x}(\mathbb{R}^{6}))},$$

and

$$\|\mathbf{u}\|_{\mathcal{S}'(\dot{\mathrm{H}}^{s}(\mathbb{R}^{6}))} := \inf_{(q,r)\in\Lambda_{s}} \|\mathbf{u}\|_{\mathrm{L}^{q'}_{t}(\mathbb{R},\mathrm{L}^{r'}_{x}(\mathbb{R}^{6}))}$$

We extent our notation $\mathcal{S}(\dot{\mathrm{H}}^{s}(\mathbb{R}^{6})), \mathcal{S}'(\dot{\mathrm{H}}^{s}(\mathbb{R}^{6}))$ as follows: we write $\mathcal{S}(I, \dot{\mathrm{H}}^{s}(\mathbb{R}^{6}))$ or $\mathcal{S}'(I, \dot{\mathrm{H}}^{s}(\mathbb{R}^{6}))$ to indicate a restriction to a time subinterval $I \subset \mathbb{R}$.

Lemma 2.3 (Dispersive estimate). For any $t \neq 0$ and $r \ge 2$, we have

$$\begin{aligned} \|\mathbf{S}(t)\mathbf{g}\|_{\mathbf{L}^{\infty}_{x}(\mathbb{R}^{6})} \lesssim_{\kappa} |t|^{-3} \|\mathbf{g}\|_{\mathbf{L}^{1}(\mathbb{R}^{6})}, \\ \|\mathbf{S}(t)\mathbf{g}\|_{\mathbf{L}^{r}_{x}(\mathbb{R}^{6})} \lesssim_{\kappa} |t|^{\frac{6}{r}-3} \|\mathbf{g}\|_{\mathbf{L}^{r'}(\mathbb{R}^{6})} \end{aligned}$$

where r' is the Hölder conjugation index, that is, $\frac{1}{r} + \frac{1}{r'} = 1$.

By combining the endpoint results in [17] with Lemma 2.3, we can use $T - T^*$ method to obtain the following Strichartz estimates.

Theorem 2.4 (Strichartz estimates, [24]). The solution **u** to (NLS system) on an interval $I \ni t_0$ obeys

$$\|\mathbf{u}\|_{\mathcal{S}(I,\mathrm{L}^{2}(\mathbb{R}^{6}))} \leqslant C\left(\|\mathbf{u}(t_{0})\|_{\mathrm{L}^{2}(\mathbb{R}^{6})} + \|\mathbf{f}(\mathbf{u})\|_{\mathcal{S}'(I,\mathrm{L}^{2}(\mathbb{R}^{6}))}\right).$$
(2.4)

2.2. Local well-posedness. Standard argument can help us prove the following basic local well-posedness properties of the solution to (NLS system).

Theorem 2.5 (Local well-posedness, [2, 27]). For $\mathbf{u}_0 \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$, there exists a unique maximal-lifespan solution $\mathbf{u} : I \times \mathbb{R}^6 \to \mathbb{C}^2$ with the following properties holding

- (1) $0 \in I$ is an open interval.
- (2) If $\sup I$ is finite, then **u** blows-up forward in time. Similarly, if $\inf I$ is finite, then **u** blows-up backward in time.
- (3) If $\sup I = +\infty$ and **u** does not blow-up forward in time, then **u** scatters forward in time. Conversely, given $\mathbf{u}_+ \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$, there is a unique solution $\mathbf{u}(t)$ to (NLS system) in a neighborhood of $t = \infty$ such that

$$\lim_{t \to +\infty} \|\mathbf{u}(t) - \mathbf{S}(t)\mathbf{u}_+\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} = 0.$$

Analogous statements hold backward in time.

(4) There exists a small number $\delta_{sd} > 0$ satisfying that if $\|\mathbf{u}_0\|_{\mathcal{S}(I,\dot{\mathrm{H}}^1(\mathbb{R}^6))} \leq \delta_{sd}$, then \mathbf{u} is global and scatters with $S_{\mathbb{R}}(\mathbf{u}) \leq \delta_{sd}^4$.

Remark 2.6. The last result of Theorem 2.2 can be improved in the following manner: instead of using $\|\mathbf{u}_0\|_{\mathcal{S}(I,\dot{\mathrm{H}}^1(\mathbb{R}^6))} \leq \delta_{sd}$, we may use a weaker hypothesis $\|\mathbf{S}(t)\mathbf{u}_0\|_{\mathcal{S}(I,\dot{\mathrm{H}}^1(\mathbb{R}^6))} \leq \delta_{sd}$ to obtain the same result. And for any given $\mathbf{u}_0 \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$, one can utilize Strichartz estimate to deduce that there exists $I \ni 0$ such that $\|\mathbf{S}(t)\mathbf{u}_0\|_{\mathcal{S}(I,\dot{\mathrm{H}}^1(\mathbb{R}^6))} \leq \delta_{sd}$.

Inspired by [33], we can establish the following stability result under weak smallness conditions.

Proposition 2.7 (Long time perturbation theory). For each $M \gg 1$, there exist $\varepsilon = \varepsilon(M) \ll 1$ and $L = L(M) \gg 1$ such that the following holds. Let $\mathbf{u} = \mathbf{u}(t, x) \in \dot{\mathrm{H}}^{1}_{x}(\mathbb{R}^{6})$ for all t and solve

$$i\mathbf{u}_t + A\mathbf{u} + \mathbf{f}(\mathbf{u}) = \mathbf{0}.$$

Let $\tilde{\mathbf{u}} = \tilde{\mathbf{u}}(t, x) \in \dot{\mathrm{H}}^1_x(\mathbb{R}^6)$ for all t and define

 $\mathbf{e} := i\tilde{\mathbf{u}}_t + A\tilde{\mathbf{u}} + \mathbf{f}(\tilde{\mathbf{u}}).$

If

$$\begin{aligned} \|\tilde{\mathbf{u}}\|_{\mathcal{S}(\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} &\leq M, \qquad \|\nabla \mathbf{e}\|_{\mathcal{S}'(\mathrm{L}^{2}(\mathbb{R}^{6}))} &\leq \varepsilon, \\ \|\mathbf{S}(t-t_{0})(\mathbf{u}_{0}-\tilde{\mathbf{u}}(t_{0}))\|_{\mathcal{S}(\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} &\leq \varepsilon, \end{aligned}$$

then

$$\|\mathbf{u}\|_{\mathcal{S}(\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} \leq L.$$

Proof. Let $\mathbf{w} := \mathbf{u} - \tilde{\mathbf{u}} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$ for $\mathbf{u} = \begin{pmatrix} u \\ v \end{pmatrix}$ and $\tilde{\mathbf{u}} = \begin{pmatrix} \tilde{u} \\ \tilde{v} \end{pmatrix}$. Then \mathbf{w} solves the equation

$$i\mathbf{w}_t + A\mathbf{w} + \begin{pmatrix} \overline{\tilde{u}}w_2 + \tilde{v}\overline{w_1}\\ 2\tilde{u}w_1 \end{pmatrix} + \mathbf{f}(\mathbf{w}) + \mathbf{e} = \mathbf{0}.$$

Since $\|\tilde{\mathbf{u}}\|_{\mathcal{S}(\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} \leq M$, we can partition $[t_{0}, +\infty)$ into N = N(M) intervals $I_{j} = [t_{j}, t_{j+1})$ such that for each $0 \leq j \leq N-1$, the quantity $\|\tilde{\mathbf{u}}\|_{\mathcal{S}(I_{j}, \dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} \leq \delta$ is small enough. By Duhamel formula,

$$\mathbf{w}(t) = \mathbf{S}(t - t_j)\mathbf{w}(t_j) + i \int_{I_j} \mathbf{S}(t - s)\mathbf{N}(\mathbf{w}(s)) \mathrm{d}s,$$

where

$$\mathbf{N}(\mathbf{w}) = \begin{pmatrix} \overline{\tilde{u}}w_2 + \tilde{v}\overline{w_1} \\ 2\tilde{u}w_1 \end{pmatrix} + \mathbf{f}(\mathbf{w}) + \mathbf{e}.$$

By Theorem 2.4, we obtain

$$\|\mathbf{w}\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} \leq \|\mathbf{S}(t-t_j)\mathbf{w}(t_j)\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} + C\|\nabla\mathbf{N}(\mathbf{w})\|_{\mathcal{S}'(I_j,\mathrm{L}^2(\mathbb{R}^6))}.$$

Note that $\|\nabla \mathbf{e}\|_{\mathcal{S}'(\mathrm{L}^2(\mathbb{R}^6))} \leq \varepsilon$,

$$\begin{split} \|\nabla \mathbf{N}(\mathbf{w})\|_{\mathcal{S}'(I_{j}, \mathbf{L}^{2}(\mathbb{R}^{6}))} &\leqslant 3\|\nabla(\tilde{\mathbf{u}}\mathbf{w})\|_{\mathbf{L}_{t}^{2}(I_{j}, \mathbf{L}_{x}^{\frac{3}{2}}(\mathbb{R}^{6}))} + \|\nabla \mathbf{f}(\mathbf{w})\|_{\mathbf{L}_{t}^{2}(I_{j}, \mathbf{L}_{x}^{\frac{3}{2}}(\mathbb{R}^{6}))} \\ &+ \|\nabla \mathbf{e}\|_{\mathcal{S}'(\mathbf{L}^{2}(\mathbb{R}^{6}))} \\ &\leqslant 3\|\nabla \tilde{\mathbf{u}}\mathbf{w}\|_{\mathbf{L}_{t}^{2}(I_{j}, \mathbf{L}_{x}^{\frac{3}{2}}(\mathbb{R}^{6}))} + 3\|\tilde{\mathbf{u}}\nabla \mathbf{w}\|_{\mathbf{L}_{t}^{2}(I_{j}, \mathbf{L}_{x}^{\frac{3}{2}}(\mathbb{R}^{6}))} \\ &+ \|\nabla \mathbf{f}(\mathbf{w})\|_{\mathbf{L}_{t}^{2}(I_{j}, \mathbf{L}_{x}^{\frac{3}{2}}(\mathbb{R}^{6}))} + \varepsilon \\ &\leqslant 3\|\nabla \tilde{\mathbf{u}}\|_{\mathbf{L}_{t}^{4}(I_{j}, \mathbf{L}_{x}^{\frac{3}{2}}(\mathbb{R}^{6}))} \|\mathbf{w}\|_{\mathbf{L}_{t}^{4}(I_{j}, \mathbf{L}_{x}^{4}(\mathbb{R}^{6}))} \\ &+ 3\|\tilde{\mathbf{u}}\|_{\mathbf{L}_{t}^{4}(I_{j}, \mathbf{L}_{x}^{4}(\mathbb{R}^{6}))} \|\nabla \mathbf{w}\|_{\mathbf{L}_{t}^{4}(I_{j}, \mathbf{L}_{x}^{4}(\mathbb{R}^{6}))} \\ &+ 2\|\tilde{\mathbf{w}}\|_{\mathbf{L}_{t}^{4}(I_{j}, \mathbf{L}_{x}^{4}(\mathbb{R}^{6}))} \|\nabla \mathbf{w}\|_{\mathbf{L}_{t}^{4}(I_{j}, \mathbf{L}_{x}^{\frac{12}{5}}(\mathbb{R}^{6}))} + \varepsilon \\ &\leqslant 6\delta\|\mathbf{w}\|_{\mathcal{S}(I_{j}, \dot{\mathbf{H}}^{1}(\mathbb{R}^{6}))} + 2\|\mathbf{w}\|_{\mathcal{S}(I_{j}, \dot{\mathbf{H}}^{1}(\mathbb{R}^{6}))} + \varepsilon. \end{split}$$

Thus

$$\|\mathbf{w}\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} \leqslant \|\mathbf{S}(t-t_j)\mathbf{w}(t_j)\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} + 6C\delta \|\mathbf{w}\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} + 2C \|\mathbf{w}\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))}^2 + C\varepsilon.$$

$$(2.5)$$

Provided $\delta < \frac{1}{12C}$ and $\|\mathbf{S}(t-t_j)\mathbf{w}(t_j)\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} + C\varepsilon \leq \frac{1}{8C} - \frac{3}{2}\delta$, we can get

$$\|\mathbf{w}\|_{\mathcal{S}(I_{j},\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} \leqslant 6C\delta \|\mathbf{w}\|_{\mathcal{S}(I_{j},\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} + 2C \|\mathbf{w}\|_{\mathcal{S}(I_{j},\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))}^{2} + \frac{1}{8C} - \frac{3}{2}\delta,$$

which implies $\|\mathbf{w}\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} \leq \frac{1}{4C} - 3\delta$. Under this circumstance, we have

$$6C\delta \|\mathbf{w}\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} + 2C \|\mathbf{w}\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))}^2 \leqslant \frac{1}{2} \|\mathbf{w}\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))}.$$

Thus,

$$\|\mathbf{w}\|_{\mathcal{S}(I_{j},\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} \leq \frac{1}{2} \|\mathbf{w}\|_{\mathcal{S}(I_{j},\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} + \|\mathbf{S}(t-t_{j})\mathbf{w}(t_{j})\|_{\mathcal{S}(I_{j},\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} + C\varepsilon.$$

As a result,

$$\|\mathbf{w}\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} \leq 2\|\mathbf{S}(t-t_j)\mathbf{w}(t_j)\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} + 2C\varepsilon.$$
(2.6)

Now take $t = t_{j+1}$ in the above Duhamel formula and apply $S(t - t_{j+1})$ to both sides to obtain

$$\mathbf{S}(t-t_{j+1})\mathbf{w}(t_{j+1}) = \mathbf{S}(t-t_j)\mathbf{w}(t_j) + i \int_{I_j} \mathbf{S}(t-s)\mathbf{N}(\mathbf{w}(s)) \mathrm{d}s.$$

Similarly, we can also get

$$\|\mathbf{S}(t-t_{j+1})\mathbf{w}(t_{j+1})\|_{\mathcal{S}(\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} \\ \leq \|\mathbf{S}(t-t_{j})\mathbf{w}(t_{j})\|_{\mathcal{S}(\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} + 3C\delta\|\mathbf{w}\|_{\mathcal{S}(I_{j},\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))} + C\|\mathbf{w}\|_{\mathcal{S}(I_{j},\dot{\mathrm{H}}^{1}(\mathbb{R}^{6}))}^{2} + C\varepsilon,$$

and

$$\|\mathbf{S}(t-t_{j+1})\mathbf{w}(t_j)\|_{\mathcal{S}(\dot{\mathrm{H}}^1(\mathbb{R}^6))} \leq 2\|\mathbf{S}(t-t_j)\mathbf{w}(t_j)\|_{\mathcal{S}(\dot{\mathrm{H}}^1(\mathbb{R}^6))} + 2C\varepsilon.$$

Iterating the above procedure with the start $j = 0$, we obtain

$$\begin{aligned} \|\mathbf{S}(t-t_j)\mathbf{w}(t_{j+1})\|_{\mathcal{S}(\dot{\mathrm{H}}^1(\mathbb{R}^6))} &\leq 2^j \|\mathbf{S}(t-t_0)\mathbf{w}(t_0)\|_{\mathcal{S}(\dot{\mathrm{H}}^1(\mathbb{R}^6))} + (2^j-1)2C\varepsilon \\ &\leq 2^{j+2}C\varepsilon. \end{aligned}$$

To accommodate $\|\mathbf{S}(t-t_j)\mathbf{w}(t_j)\|_{\mathcal{S}(I_j,\dot{\mathrm{H}}^1(\mathbb{R}^6))} + C\varepsilon \leq \frac{1}{8C} - \frac{3}{2}\delta$ for all intervals I_j , $0 \leq j \leq N-1$, we require that

$$2^{N+2}C\varepsilon \leqslant \frac{1}{8C} - \frac{3}{2}\delta$$

which can be ensured by choosing $0 < \varepsilon \ll 1$.

3. VARIATIONAL CHARACTERIZATION

In this section, we are in the position to give the variational characterization for the sharp Gargliardo-Nirenberg inequality. Apart from stating the existence of the ground state, we will also show some relationships between the energy of solution and the energy of ground state, which facilitate us to obtain the finite time blow-up result in Theorem 1.5.

3.1. Ground state. We first recall some basic properties of ground state. As a quick application, we obtain the sharp Gargliardo-Nirenberg inequality. Then we will show the classification in Theorem 1.5 is complete.

Proposition 3.1 (Ground state, [12]). For $D_6 = \{ \mathbf{G} \in \dot{\mathrm{H}}^1 \cap \mathrm{L}^3 \mid R(\mathbf{G}) > 0 \}$, the minimal J_{\min} of the nonnegative functional

$$J(\mathbf{G}) := \left[H(\mathbf{G})\right]^3 \left[R(\mathbf{G})\right]^{-2}, \qquad \mathbf{G} \in D_6$$

are attained at $\mathbf{G} = (\Phi, \Psi)^T \in \mathbb{C}^2$, whose expression has to have the form of $(\Phi, \Psi) = (e^{i\theta}m\phi(nx), e^{2i\theta}m\varphi(nx))$, where m > 0, n > 0, $\theta \in \mathbb{R}$, and $\mathbf{W} := (\phi, \varphi)^T \in D_6$ is a pair of non-negative real-valued functions satisfying (1.3). The

function W is called a ground state with $J(W) = J_{\min}$. The set of all ground states is denoted as \mathcal{G} .

In addition, ground states for (1.3) are unique up to translations and dilations. The unique ground state is given by $\mathbf{W} = (\phi_0, \frac{\phi_0}{\sqrt{\kappa}})^T$ where ϕ_0 is a ground state of (1.4) in $\dot{H}^1(\mathbb{R}^6) \cap L^3(\mathbb{R}^6)$.

Remark 3.2. For the energy of ground state $\mathbf{W} = (\phi, \varphi)^T \in D_6$, we have the following scaling identity:

$$E\left(\lambda^{\alpha}\mathbf{W}(\lambda^{\beta} \cdot)\right) = \lambda^{2\alpha - 4\beta}H(\mathbf{W}) - \lambda^{3\alpha - 6\beta}R(\mathbf{W}), \qquad \forall \ \lambda \in (0, \infty).$$

Using variational derivatives and letting $\lambda=1$ in both sides of above identity, we can obtain

$$0 = \operatorname{Re} \int_{\mathbb{R}^{6}} \left(-2\Delta\phi - 2\phi\varphi \right) \cdot \overline{\left(\frac{\mathrm{d}}{\mathrm{d}\lambda}\Big|_{\lambda=1}\lambda^{\alpha}\phi(\lambda^{\beta}x)\right)} \\ + \left(-\kappa\Delta\varphi - \phi^{2}\right) \cdot \overline{\left(\frac{\mathrm{d}}{\mathrm{d}\lambda}\Big|_{\lambda=1}\lambda^{\alpha}\varphi(\lambda^{\beta}x)\right)} \mathrm{d}x \\ = (2\alpha - 4\beta)H(\mathbf{W}) - (3\alpha - 6\beta)R(\mathbf{W}) \\ = [2H(\mathbf{W}) - 3R(\mathbf{W})]\alpha - [4H(\mathbf{W}) - 6R(\mathbf{W})]\beta, \quad \forall \ \alpha \in \mathbb{R}, \ \beta \in \mathbb{R}.$$

This yields

$$H(\mathbf{W}): R(\mathbf{W}) = 3:2.$$
 (3.1)

This argument of scaling analysis produces the same effect as Pohozaev's identities.

Corollary 3.3 (Gagliardo-Nirenberg inequality). For any $\mathbf{g} \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$,

$$R(\mathbf{g}) \leqslant C_{GN}[H(\mathbf{g})]^{\frac{3}{2}},\tag{3.2}$$

where $C_{GN} = J_{\min}^{-\frac{1}{2}} > 0$ is a constant. Besides, $R(\mathbf{W}) = C_{GN}[H(\mathbf{W})]^{\frac{3}{2}}$ for any $\mathbf{W} \in \mathcal{G}$.

Proof. For $\mathbf{g} \in \dot{\mathrm{H}}^1(\mathbb{R}^6) \setminus D_6 = \left\{ \mathbf{g} \in \dot{\mathrm{H}}^1(\mathbb{R}^6) | R(\mathbf{g}) \leq 0 \right\}$, (3.2) holds obviously since $C_{GN} > 0$ and $H(\mathbf{g}) \geq 0$. For $\mathbf{g} \in D_6$, by Proposition 3.1, we have

$$[H(\mathbf{g})]^3 [R(\mathbf{g})]^{-2} = J(\mathbf{g}) \ge J_{\min},$$

which leads to (3.2) directly.

From the proof of Proposition 3.1, we can easily see that $R(\mathbf{W}) = C_{GN}[H(\mathbf{W})]^{\frac{3}{2}}$ for any $\mathbf{W} \in \mathcal{G}$.

Proposition 3.4 (Coercivity of energy). Let $0 < \delta < 1$. For the initial data $\mathbf{u}_0 \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$ and satisfying $E(\mathbf{u}_0) \leq (1-\delta)E(\mathbf{W})$ in (NLS system), if $H(\mathbf{u}_0) < H(\mathbf{W})$, then there exists $\delta'' = \delta''(\delta) > 0$ so that

$$2H(\mathbf{u}) - 3R(\mathbf{u}) \ge \delta'' H(\mathbf{u}); \tag{3.3}$$

if $H(\mathbf{u}_0) > H(\mathbf{W})$ then there exists $\tilde{\delta}'' = \tilde{\delta}''(\delta) > 0$ so that

$$2H(\mathbf{u}) - 3R(\mathbf{u}) \leqslant -\tilde{\delta}'' H(\mathbf{u}). \tag{3.4}$$

Proof. We start with the energy of \mathbf{u} , by (3.2),

$$E(\mathbf{u}) = H(\mathbf{u}) - R(\mathbf{u}) \ge H(\mathbf{u}) - C_{GN}[H(\mathbf{u})]^{\frac{3}{2}}.$$
(3.5)

Define $f(y) = y - C_{GN} y^{\frac{3}{2}}$ for $y = y(t) := H(\mathbf{u}(t)) \ge 0$. Then $f'(y) = 1 - \frac{3}{2}C_{GN} y^{\frac{1}{2}}$. Thus $f'(y) \ge 0$ only when $y \le y_0 := \frac{4}{9C_{GN}^2}$. From Remark 3.2 and Corollary 3.3, we may obtain $C_{GN} = \frac{2}{3[H(\mathbf{W})]^{\frac{1}{2}}}$ and $y_0 = H(\mathbf{W})$.

Therefore,

$$f_{\max}(y) = f(y_0) = f(H(\mathbf{W})) = H(\mathbf{W}) - C_{GN}[H(\mathbf{W})]^{\frac{3}{2}} = \frac{1}{3}H(\mathbf{W}) = E(\mathbf{W}).$$
(3.6)

For any time $t \in I$, by (3.5), (3.6), and the conversation of energy, we get

$$f(H(\mathbf{u}(t))) \leq E(\mathbf{u}(t)) = E(\mathbf{u_0}) < E(\mathbf{W}) = f(H(\mathbf{W})),$$

which means $H(\mathbf{u}(t)) \neq H(\mathbf{W})$ for any $t \in I$.

If $H(\mathbf{u}_0) < H(\mathbf{W})$, by the continuity of f(y) and y(t), there exists $\delta' = \delta'(\delta) > 0$ so that

$$H(\mathbf{u}(t)) \leq (1 - \delta')H(\mathbf{W}), \quad \forall \ t \in I.$$
 (3.7)

Similarly, if $H(\mathbf{u}_0) > H(\mathbf{W})$, there exists $\tilde{\delta}' = \tilde{\delta}'(\delta) > 0$ so that

$$H(\mathbf{u}(t)) \ge (1 + \tilde{\delta}')H(\mathbf{W}), \quad \forall \ t \in I.$$
(3.8)

Furthermore, there exist $\delta'' = \delta''(\delta') > 0$ and $\tilde{\delta}'' = \tilde{\delta}''(\delta, \tilde{\delta}') > 0$ so that

$$2 - 3\frac{R(\mathbf{u})}{H(\mathbf{u})} \ge 2 - 3C_{GN}[H(\mathbf{u})]^{\frac{1}{2}} = 2 - 2\left(\frac{H(\mathbf{u})}{H(\mathbf{W})}\right)^{\frac{1}{2}} \ge 2\left[1 - (1 - \delta')^{\frac{1}{2}}\right] := \delta'';$$

and

$$2 - 3\frac{R(\mathbf{u})}{H(\mathbf{u})} = 3\frac{E(\mathbf{u}_0)}{H(\mathbf{u})} - 1 \leqslant 3\frac{(1-\delta)E(\mathbf{W})}{(1+\delta')H(\mathbf{W})} - 1 = \frac{(1-\delta)}{(1+\delta')} - 1 := -\tilde{\delta}'',$$

that is,

$$2H(\mathbf{u}) - 3R(\mathbf{u}) \ge \delta'' H(\mathbf{u}); \tag{3.9}$$

and

$$H(\mathbf{u}) - 3R(\mathbf{u}) \leqslant -\tilde{\delta}'' H(\mathbf{u}). \tag{3.10}$$

So we complete the proof.

3.2. Energy trapping. For the solution to classical focusing (NLS), generally speaking, we can not determine the sign of its energy. However, under certain circumstance, we may determine the sign of a general function regardless of whether it is a solution to (NLS) or not. For us, we may generalize the above fact to our setting.

Lemma 3.5. For any
$$\mathbf{v} \in \dot{\mathrm{H}}^{1}(\mathbb{R}^{6})$$
, if $H(\mathbf{v}) \leq H(\mathbf{W})$, then
 $E(\mathbf{v}) \geq 0$.

2

Proof. We just need to show $H(\mathbf{v}) - C_{GN}[H(\mathbf{v})]^{\frac{3}{2}} \ge 0$, since

$$E(\mathbf{v}) = H(\mathbf{v}) - R(\mathbf{v}) \ge H(\mathbf{v}) - C_{GN}[H(\mathbf{v})]^{\frac{3}{2}}.$$

Considering $0 \leq H(\mathbf{v}) \leq H(\mathbf{W})$, we define $f(y) = y - C_{GN} y^{\frac{3}{2}}$ for $0 \leq y \leq H(\mathbf{W})$. Then $f'(y) = 1 - \frac{3}{2}C_{GN}y^{\frac{1}{2}}$. Thus $f'(y) \geq 0$ only when $y \leq y_0 := \frac{4}{9C_{GN}^2}$. The fact $C_{GN} = \frac{R(\mathbf{W})}{[H(\mathbf{W})]^{\frac{3}{2}}}$ and $H(\mathbf{W}) : R(\mathbf{W}) = 2 : 3$ implies

$$C_{GN} = \frac{2}{3[H(\mathbf{W})]^{\frac{1}{2}}}$$
 and $y_0 = H(\mathbf{W}).$

Therefore, we obtain $E(\mathbf{v}) \ge f(H(\mathbf{v})) \ge 0$ if $0 \le H(\mathbf{v}) \le H(\mathbf{W})$.

Next, we will establish the equivalence relation between the energy and the kinetic energy of the solution to (NLS system), that is, $H(\mathbf{u}) \sim E(\mathbf{u})$ under certain restrictions. For the classical (NLS), it is easy to obtain $E(\mathbf{u}) \leq H(\mathbf{u})$. Indeed, for the focusing case, we can use the fact that $R(\mathbf{u}) \geq 0$, which is missing in our setting; for the defocusing case, we can use the associated Gagliardo-Nirenberg inequality. However, for (NLS system), we do not have the natural comparison, $E(\mathbf{u}) \leq H(\mathbf{u})$. Our proof here is different from the argument in [18]. To obtain $|R(\mathbf{u})| \leq_{\kappa} [H(\mathbf{u})]^{\frac{3}{2}}$, the Arithmetic-Geometry Mean-value inequality is used. Combining with the coercivity of energy, we prove $H(\mathbf{u}) \sim E(\mathbf{u})$.

Lemma 3.6. For any $\mathbf{v} \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$,

$$|R(\mathbf{v})| \leqslant C(\kappa) [H(\mathbf{v})]^{\frac{3}{2}},\tag{3.11}$$

where $C(\kappa) = \sqrt{\frac{8}{27\kappa}} > 0.$

Proof. Using Hölder's inequality, Sobolev embedding, and the Arithmetic-Geometry Mean-value inequality,

$$\frac{1}{n}\sum_{i=1}^{n}a_i \geqslant \sqrt[n]{\prod_{i=1}^{n}a_i}, \quad \text{for } \forall \ 1 \leqslant i \leqslant n, a_i \geqslant 0,$$

we can compute directly, setting $\mathbf{v} := (\tilde{u}, \tilde{v})^T$,

$$\begin{aligned} |R(\mathbf{v})| &= \left| \operatorname{Re} \int_{\mathbb{R}^{6}} \overline{\tilde{v}} \tilde{u}^{2} dx \right| \leq \int_{\mathbb{R}^{6}} |\tilde{v}| |\tilde{u}|^{2} dx \\ &\leq \|\tilde{v}\|_{L^{3}(\mathbb{R}^{6})} \|\tilde{u}\|_{L^{3}(\mathbb{R}^{6})}^{2} \leq \|\tilde{v}\|_{\dot{H}^{1}(\mathbb{R}^{6})} \|\tilde{u}\|_{\dot{H}^{1}(\mathbb{R}^{6})}^{2} \\ &= \left[\frac{8}{\kappa} \left(\frac{\kappa}{2} \|\tilde{v}\|_{\dot{H}^{1}(\mathbb{R}^{6})}^{2} \right) \left(\frac{1}{2} \|\tilde{u}\|_{\dot{H}^{1}(\mathbb{R}^{6})}^{2} \right) \left(\frac{1}{2} \|\tilde{u}\|_{\dot{H}^{1}(\mathbb{R}^{6})}^{2} \right) \right]^{\frac{1}{2}} \\ &\leq \left\{ \frac{8}{\kappa} \left[\frac{\left(\frac{\kappa}{2} \|\tilde{v}\|_{\dot{H}^{1}(\mathbb{R}^{6})}^{2} \right) + \left(\frac{1}{2} \|\tilde{u}\|_{\dot{H}^{1}(\mathbb{R}^{6})}^{2} \right) + \left(\frac{1}{2} \|\tilde{u}\|_{\dot{H}^{1}(\mathbb{R}^{6})}^{2} \right) \right]^{3} \right\}^{\frac{1}{2}} \\ &= \sqrt{\frac{8}{27\kappa}} [H(\mathbf{v})]^{\frac{3}{2}}. \end{aligned}$$

Choosing $C(\kappa) = \sqrt{\frac{8}{27\kappa}}$, we obtain (3.11).

Using Lemma 3.6 and Proposition 3.4, we can establish the energy trapping result.

Corollary 3.7 (Energy Trapping). Let \mathbf{u} be a solution to (NLS system) with initial data \mathbf{u}_0 and maximal life-span $I_{\max} \ni 0$. If $E(\mathbf{u}_0) \leq (1-\delta)E(\mathbf{W})$ and $H(\mathbf{u}_0) \leq (1-\delta')H(\mathbf{W})$, then

$$H(\mathbf{u}(t)) \sim E(\mathbf{u}(t)), \quad \forall t \in I_{\max}.$$
 (3.12)

Proof. By (3.11) and (3.7), we may get the upper bound of energy $E(\mathbf{u})$,

$$\begin{split} E(\mathbf{u}(t)) &\leqslant H(\mathbf{u}(t)) + |R(\mathbf{u}(t))| \\ &\leqslant H(\mathbf{u}(t)) + C(\kappa) \left[H(\mathbf{u}(t))\right]^{\frac{3}{2}} \\ &\leqslant \left(1 + C(\kappa) \left[(1 - \delta')H(\mathbf{W})\right]^{\frac{1}{2}}\right) H(\mathbf{u}(t)). \end{split}$$

As for the lower bound of $E(\mathbf{u})$, the proof of Proposition 3.4 tells us

$$E(\mathbf{u}(t)) = \frac{1}{3}H(\mathbf{u}(t)) + \frac{1}{3}[2H(\mathbf{u}(t)) - 3R(\mathbf{u}(t))]$$

$$\geq \frac{1}{3}H(\mathbf{u}(t)) + \frac{1}{3}\delta''H(\mathbf{u}(t))$$

$$= \frac{1}{3}(1 + \delta'')H(\mathbf{u}(t)).$$

Combining the above two inequalities, we deduce the conclusion (3.12).

3.3. Virial identity and blowing-up. Despite being proved in [6], the part of blowing-up result of Theorem 1.5 will also be shown for completeness.

We aim to figure out how the mass is distributed in the spatial space. In particular, we try to make clear whether it concentrates at the origin or not, and how the distribution changes over time. So we consider a class of initial data satisfying $x\mathbf{u}_0 \in \mathrm{L}^2(\mathbb{R}^6)$ firstly. The law of conservation of mass suggests it useless to think about the change of $\|\mathbf{u}(t)\|_{\mathrm{L}^2}$ over time directly, so we redistribute the mass of \mathbf{u} to large radii and consider the behavior of $\|x\mathbf{u}(t)\|_{\mathrm{L}^2}$ when $t \in I_{max}$. For d = 6 in the (NLS system) we define a function

$$I(t) := \int_{\mathbb{R}^6} |x|^2 \left(2\kappa |u|^2 + |v|^2 \right) \mathrm{d}x.$$
(3.13)

Remark 3.8. The coefficients in (3.13) are chosen carefully to match with Propostion 3.4, one may refer to Remark 4.1 in [24] for details. More specifically, just by adjusting the rate of coefficients between $|u|^2$ and $|v|^2$ to be $2\kappa : 1$, we can obtain a factor $2H(\mathbf{u}) - 3R(\mathbf{u})$ in (3.14).

Lemma 3.9 (Virial identity). If **u** is a solution to (NLS system), for any real valued function $a \in C^{\infty}(\mathbb{R}^6)$, $t \in [0, T_+(\mathbf{u}))$, then (1)

$$\frac{\mathrm{d}}{\mathrm{d}t} \int_{\mathbb{R}^6} \left(2\kappa |u|^2 + |v|^2 \right) a(x) \mathrm{d}x = 2\kappa \operatorname{Im} \int_{\mathbb{R}^6} \left(2\overline{u}\nabla u + \overline{v}\nabla v \right) \cdot \nabla a(x) \mathrm{d}x.$$

(2)

$$\frac{\mathrm{d}^2}{\mathrm{d}t^2} \int_{\mathbb{R}^6} \left(2\kappa |u|^2 + |v|^2 \right) a(x) \mathrm{d}x$$

= $8\kappa \operatorname{Re} \int_{\mathbb{R}^6} \left(\overline{u_j} u_k + \frac{\kappa}{2} \overline{v_j} v_k \right) a_{jk}(x) \mathrm{d}x - 2\kappa \int_{\mathbb{R}^6} \left(|u|^2 + \frac{\kappa}{2} |v|^2 \right) \Delta \Delta a(x) \mathrm{d}x$
 $- 2\kappa \operatorname{Re} \int_{\mathbb{R}^6} \overline{v} u^2 \Delta a(x) \mathrm{d}x.$

Set $a(x) = x^2$, a simple calculation shows that

$$a(x) = |x|^2,$$
 $a_j(x) := \partial_{x_j} a = 2x_j,$ $\nabla a(x) = 2x,$

and

$$a_{jk}(x) := \partial_{x_i} \partial_{x_j} a = 2\delta_{jk}, \qquad \Delta a(x) = 2d, \qquad \Delta \Delta a(x) = 0.$$

As a result, by Lemma 3.9, we have

$$I'(t) = 4\kappa \operatorname{Im} \int_{\mathbb{R}^6} \left(2\overline{u}\nabla u + \overline{v}\nabla v \right) \cdot x \mathrm{d}x$$

and

$$I''(t) = 16\kappa H(\mathbf{u}) - 24\kappa R(\mathbf{u}) = 8\kappa [2H(\mathbf{u}) - 3R(\mathbf{u})].$$
(3.14)

By the Gagliardo-Nirenberg inequality (3.2), we know

$$\frac{1}{8\kappa} I''(t) = 2H(\mathbf{u}) - 3R(\mathbf{u})$$

$$\geq 2H(\mathbf{u}) - 2 \left[H(\mathbf{W})\right]^{-\frac{1}{2}} \left[H(\mathbf{u})\right]^{\frac{3}{2}}$$

$$= 2 \left[H(\mathbf{u})\right]^{\frac{3}{2}} \left(\left[H(\mathbf{u})\right]^{-\frac{1}{2}} - \left[H(\mathbf{W})\right]^{-\frac{1}{2}} \right).$$

So under the hypotheses of $E(\mathbf{u_0}) < E(\mathbf{W})$ and $H(\mathbf{u_0}) < H(\mathbf{W})$, we have I''(0) > 0, which implies I(t) is a convex function. We can expect \mathbf{u} is global and scatters.

Under the hypotheses of $E(\mathbf{u_0}) < E(\mathbf{W})$ and $H(\mathbf{u_0}) > H(\mathbf{W})$, Proposition 3.4 tells us that the right side of (3.14) is strictly negative, which implies I(t) is a concave function. Combing with I(0) > 0, we know the variance I(t) will tend to 0 in finite time, which means that all the mass of \mathbf{u} concentrates at the origin and \mathbf{u} has to blow up in finite time.

The coercivity of energy has showed some relations between the potential energy and the kinetic energy of solution in any time $t \in I_{\text{max}}$. But we need some uniform bounds sometimes.

Theorem 3.10. For $x\mathbf{u}_0 \in L^2(\mathbb{R}^6)$, we set I_{\max} be the maximal time interval of existence of $\mathbf{u}(t)$ solving (NLS system). If $E(\mathbf{u}_0) < E(\mathbf{W})$ and $H(\mathbf{u}_0) > H(\mathbf{W})$, then I_{\max} is finite and $H(\mathbf{u}(t)) > H(\mathbf{W})$, $\forall t \in I_{\max}$.

Proof. If $H(\mathbf{u}_0) > H(\mathbf{W})$, by the properties of f(y) as in the proof of Proposition 3.4 and (3.8), we can choose $\tilde{\delta}' > 0$ such that

$$H(\mathbf{u}(t)) \ge (1 + \tilde{\delta}')H(\mathbf{W}), \ \forall \ t \in I_{\max}.$$

And then,

$$2H(\mathbf{u}(t)) - 3R(\mathbf{u}(t)) = 3E(\mathbf{u}(t)) - H(\mathbf{u}(t))$$

= $3E(\mathbf{u}_0) - H(\mathbf{u}(t))$
 $\leqslant 3E(\mathbf{W}) - (1 + \tilde{\delta}')H(\mathbf{W})$
= $H(\mathbf{W}) - (1 + \tilde{\delta}')H(\mathbf{W}) = -\tilde{\delta}'H(\mathbf{W}).$

The Virial identity (3.14) shows that

$$I''(t) = 8\kappa [2H(\mathbf{u}) - 3R(\mathbf{u})] \leqslant -8\kappa \tilde{\delta}' H(\mathbf{W}), \ \forall \ t \in I_{\max}.$$

So, the time $t \in I_{\text{max}}$ must be finite.

Theorem 3.11. For $\mathbf{u}_0 \in \mathrm{H}^1(\mathbb{R}^6)$ radial instead of the assumption $x\mathbf{u}_0 \in \mathrm{L}^2(\mathbb{R}^6)$ in Theorem 3.10. If the other conditions keep the same as Theorem 3.10, then the same result holds as well.

Proof. We change the weighted function $|x|^2$ into a specific smooth cut-off function in (3.13), to be more specific,

$$\tilde{I}(t) := \int_{\mathbb{R}^6} \left(2\kappa |u|^2 + |v|^2 \right) a(x) \mathrm{d}x, \quad a(x) := R^2 \Gamma\left(\frac{|x|^2}{R^2}\right), \ \forall \ R > 0,$$

where $\Gamma(r)$ is a smooth concave function defined on $[0,\infty)$ satisfying

$$\Gamma(r) = \begin{cases} r, & r \leq 1, \\ 2, & r \geq 3, \end{cases} \quad \text{and} \quad \begin{cases} \Gamma''(r) \searrow, & r \leq 2, \\ \Gamma''(r) \nearrow, & r \geq 2. \end{cases}$$

By Lemma 3.9, we have

$$\begin{split} I''(t) &= 8\kappa \int_{\mathbb{R}^6} [2H(\mathbf{u}) - 3R(\mathbf{u})] \mathrm{d}x + \frac{1}{R^2} O\left(\int_{|x| \sim R} |\mathbf{u}|^2 \mathrm{d}x\right) \\ &+ 8\kappa \int_{\mathbb{R}^6} \left(\Gamma'\left(\frac{|x|^2}{R^2}\right) - 1 + \frac{2|x|^2}{R^2} \Gamma''\left(\frac{|x|^2}{R^2}\right)\right) [2H(\mathbf{u}) - 3R(\mathbf{u})] \mathrm{d}x \\ &- \frac{40}{3} \operatorname{Re} \int_{\mathbb{R}^6} \frac{2|x|^2}{R^2} \Gamma''\left(\frac{|x|^2}{R^2}\right) \overline{v} u^2 \mathrm{d}x. \end{split}$$

Because $\Gamma''(r) > 0$ and $\mathbf{u} \in L^2(\mathbb{R}^6)$, we can choose $R = R(M(\mathbf{u})))$ large enough to obtain

$$I''(t) \leqslant -4\kappa \tilde{\delta}' H(\mathbf{W}) - 8\kappa \int_{\mathbb{R}^6} \omega(x) [2H(\mathbf{u}) - 3R(\mathbf{u})] \mathrm{d}x,$$

where

$$\omega(x) = 1 - \Gamma'\left(\frac{|x|^2}{R^2}\right) - \frac{2|x|^2}{R^2}\Gamma''\left(\frac{|x|^2}{R^2}\right).$$

Observe that $0 \leq \Gamma \leq 1$ is radial, supp $(\Gamma) \subset \{x | |x| > R\}$, and

 $\Gamma(x) \lesssim \Gamma(y)$, uniformly for $|x| \leq |y|$,

we have

$$\left\| |x|^{\frac{5}{2}} \Gamma^{\frac{1}{4}} f \right\|_{L^{\infty}_{x}(\mathbb{R}^{6})}^{2} \lesssim \|f\|_{L^{2}_{x}(\mathbb{R}^{6})} \left\| \Gamma^{\frac{1}{2}} \nabla f \right\|_{L^{2}_{x}(\mathbb{R}^{6})}.$$

Furthermore, by the conservation of mass,

$$\operatorname{Re} \int_{\mathbb{R}^{6}} \Gamma(x) \overline{v} u^{2} \mathrm{d}x \lesssim \left\| \Gamma^{\frac{1}{4}} \mathbf{u}(t) \right\|_{\mathrm{L}_{x}^{\infty}(\mathbb{R}^{6})} \int_{\mathbb{R}^{6}} |\mathbf{u}(t,x)|^{2} \mathrm{d}x$$
$$\lesssim R^{-\frac{5}{2}} \left\| |x|^{\frac{5}{2}} \Gamma^{\frac{1}{4}} \mathbf{u}(t) \right\|_{\mathrm{L}_{x}^{\infty}(\mathbb{R}^{6})} M(\mathbf{u})$$
$$\lesssim R^{-\frac{5}{2}} \left\| \Gamma^{\frac{1}{2}} \nabla \mathbf{u}(t) \right\|_{\mathrm{L}_{x}^{2}(\mathbb{R}^{6})}^{\frac{1}{2}} [M(\mathbf{u})]^{\frac{5}{4}}.$$

This implies that $\tilde{I}''(t) < 0$ by choosing R > 0 large enough.

Theorem 3.10 and Theorem 3.11 have rigorously proved some original ideas in Subsection 3.3 and showed the blowing-up result in Theorem 1.5.

4. Proof of Theorem 1.11

In this section, we use the inverse Strichartz inequality to derive a linear profile decomposition for the Schrödinger propagator S(t). Combining with the nonlinear profile, we deduce the existence of critical solution.

4.1. Linear profile decomposition. In this subsection, we show the profile decomposition with scaling parameter of a radial uniformly bounded sequence in $\dot{H}^1(\mathbb{R}^6)$. We start by combining the Strichartz inequality for the Schrödinger propagator S(t), Theorem 2.4, and Sobolev embedding to obtain

$$\|\mathbf{S}(t)\mathbf{g}\|_{\mathbf{L}^{4}_{t,r}(\mathbb{R}\times\mathbb{R}^{6})} \lesssim \|\mathbf{S}(t)\nabla\mathbf{g}\|_{\mathbf{L}^{4}_{t,r}(\mathbb{R}\times\mathbb{R}^{6})} \lesssim \|\mathbf{g}\|_{\dot{\mathbf{H}}^{1}_{r}(\mathbb{R}^{6})}.$$
(4.1)

Our next result is a refinement of (4.1), which says that if the linear evolution of **g** is large in $L_{t,x}^4(\mathbb{R} \times \mathbb{R}^6)$, then the linear evolution of a single Littlewood-Paley piece of **g** is, at least partially, responsible.

Lemma 4.1 (Refined Strichartz estimate). For any function $\mathbf{g} \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$, we have

$$\|\mathbf{S}(t)\mathbf{g}\|_{\mathrm{L}^{4}_{t,x}(\mathbb{R}\times\mathbb{R}^{6})} \lesssim \|\mathbf{g}\|_{\mathrm{\dot{H}}^{1}(\mathbb{R}^{6})}^{\frac{1}{2}} \sup_{N\in2^{\mathbb{Z}}} \|\mathbf{S}(t)P_{N}\mathbf{g}\|_{\mathrm{L}^{4}_{t,x}(\mathbb{R}\times\mathbb{R}^{6})}^{\frac{1}{2}}$$

Proof. From the square function estimate and the Bernstein and Strichartz inequality, we can obtain the result above. \Box

The refined Strichartz estimate and Lemma 4.1 demonstrate that a linear solution to (NLS system) with a nontrivial space-time norm will be Fourier concentrated in at least one circular area. The following proposition will further manifest that a linear solution contains a concentrated "bubble" around a certain space-time.

Proposition 4.2 (Inverse Strichartz inequality). If the sequence $\{\mathbf{g}_n\}_{n=1}^{\infty} \subset \mathrm{H}^1(\mathbb{R}^6)$ satisfies

$$\lim_{n\to\infty} \|\mathbf{g}_n\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} = A \leqslant \infty, \quad and \quad \lim_{n\to\infty} \|\mathbf{S}(t)\mathbf{g}_n\|_{\mathrm{L}^4_{t,x}(\mathbb{R}\times\mathbb{R}^6)} = \varepsilon > 0.$$

Then there exists a subsequence of $\{n\}$ (still denoted by $\{n\}$), $\phi \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$, $\{\lambda_n\}_{n=1}^{\infty} \subset (0,\infty)$ and $\{(t_n, x_n)\}_{n=1}^{\infty} \subset \mathbb{R} \times \mathbb{R}^6$ such that

$$\lambda_n^2 \left[\mathbf{S}(t_n) \mathbf{g}_n \right] \left(\lambda_n x + x_n \right) \rightharpoonup \boldsymbol{\phi}(x) \quad weakly \ in \ \dot{\mathbf{H}}_x^1(\mathbb{R}^6), \tag{4.2}$$

$$\liminf_{n \to \infty} \left\{ \|\mathbf{g}_n\|_{\dot{\mathrm{H}}^1_x(\mathbb{R}^6)}^2 - \|\mathbf{g}_n - \boldsymbol{\phi}_n\|_{\dot{\mathrm{H}}^1_x(\mathbb{R}^6)}^2 \right\} = \|\boldsymbol{\phi}\|_{\dot{\mathrm{H}}^1_x(\mathbb{R}^6)}^2 \gtrsim A^{-10}\varepsilon^{12}, \tag{4.3}$$

$$\liminf_{n \to \infty} \left\{ \| \mathbf{S}(t) \mathbf{g}_n \|_{\mathbf{L}^4_{t,x}(\mathbb{R} \times \mathbb{R}^6)}^4 - \| \mathbf{S}(t) (\mathbf{g}_n - \boldsymbol{\phi}_n) \|_{\mathbf{L}^4_{t,x}(\mathbb{R} \times \mathbb{R}^6)}^4 \right\} \gtrsim A^{-20} \varepsilon^{24}, \qquad (4.4)$$

where

$$\phi_n(x) := \frac{1}{\lambda_n^2} \left[\mathsf{S}\left(-\frac{t_n}{\lambda_n^2} \right) \phi \right] \left(\frac{x - x_n}{\lambda_n} \right). \tag{4.5}$$

Remark 4.3. Under the hypotheses of Proposition 4.2 and passing to a further subsequence if necessary, the decoupling of potential energy holds (using Rellich-Kondrashov and refined Fatou), that is,

$$\liminf_{n\to\infty}\left\{\left\|\mathbf{g}_n\right\|_{\mathrm{L}^3_x(\mathbb{R}^6)}^3-\left\|\mathbf{g}_n-\boldsymbol{\phi}_n\right\|_{\mathrm{L}^3_x(\mathbb{R}^6)}^3-\left\|\mathsf{S}\left(-\frac{t_n}{\lambda_n^2}\right)\boldsymbol{\phi}\right\|_{\mathrm{L}^3_x(\mathbb{R}^6)}^3\right\}=0.$$

Using the above proposition, we can obtain the linear profile decomposition as follows:

Theorem 4.4 (Linear profile decomposition). Let $\phi_n = (\phi_n, \psi_n)$ be an uniformly bounded sequence in $\dot{\mathrm{H}}^1(\mathbb{R}^6)$ with $\|\phi_n\| \leq A$ for any $1 \leq n < +\infty$. Then there exists $M^* \in \{1, 2, \dots\} \cup \{\infty\}$ such that for each finite $1 \leq M \leq M^*$, (1) for each $1 \leq j \leq M$, there exist a sequence of space shifts $\{x_n^j\}_{n=1}^{\infty} \subset \mathbb{R}^6$, a sequence of time shifts $\{t_n^j\}_{n=1}^{\infty} \subset \mathbb{R}$, a sequence of scaling shifts $\{\lambda_n^j\}_{n=1}^{\infty} \subset (0, +\infty)$, and a profile $\phi^j := (\phi^j, \psi^j)$ (fixed in n) in $\dot{\mathrm{H}}^1(\mathbb{R}^6)$, with

$$\frac{\lambda_n^j}{\lambda_n^l} + \frac{\lambda_n^l}{\lambda_n^j} + \frac{|x_n^j - x_n^l|^2}{\lambda_n^j \lambda_n^l} + \frac{|t_n^j (\lambda_n^j)^2 - t_n^l (\lambda_n^l)^2|}{\lambda_n^j \lambda_n^l} \to +\infty, \quad \forall \ j \neq l \in \{1, 2, \cdots, M\}.$$
(4.6)

We may additionally assume that for each j either $t_n^j \equiv 0$ or $t_n^j \to \pm \infty$. Especially, for fixed $1 \leq j_0 \leq M$, there exists $\alpha(j_0) > 0$ such that

$$H\left(\boldsymbol{\phi}^{j_0}\right) \geqslant \alpha(j_0). \tag{4.7}$$

(2) There exists a sequence (in n) of remainders $\mathbf{\Phi}_n^M := (\Phi_n^M, \Psi_n^M)$ in $\dot{\mathrm{H}}^1(\mathbb{R}^6)$, such that

$$\boldsymbol{\phi}_{n} = \sum_{j=1}^{M} \mathcal{T}_{\lambda_{n}^{j}} \left[\mathbf{S} \left(\frac{-t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \right) \boldsymbol{\phi}^{j} \right] + \boldsymbol{\Phi}_{n}^{M}, \tag{4.8}$$

where $\mathcal{T}_{\lambda}\mathbf{u} = (T_{\lambda}u, T_{\lambda}v)^t$ for $\mathbf{u} = (u, v)^t$, with

$$\lim_{M \to M^*} \limsup_{n \to \infty} \left\| \mathbf{S}(t) \mathbf{\Phi}_n^M \right\|_{\mathbf{L}^4_{t,x}(\mathbb{R} \times \mathbb{R}^6)} = 0, \tag{4.9}$$

and

$$\mathbf{S}(-t_n^M) \left[\mathcal{T}_{\lambda_n^M}^{-1} \mathbf{\Phi}_n^M \right] \rightharpoonup 0 \qquad weakly \ in \ \dot{\mathbf{H}}^1(\mathbb{R}^6), \tag{4.10}$$

where
$$T_{\lambda}f(x) := \frac{1}{\lambda^2}f(\frac{x-x_n^j}{\lambda})$$
 and so $T_{\lambda}^{-1}f(x) := \lambda^2 f(\lambda x + x_n^j)$.

Besides, we have the asymptotic pythagorean expansion:

$$\|\phi_n\|_{\dot{H}^1(\mathbb{R}^6)}^2 = \sum_{j=1}^M \|\phi^j\|_{\dot{H}^1(\mathbb{R}^6)}^2 + \|\Phi_n^M\|_{\dot{H}^1(\mathbb{R}^6)}^2 + o_n(1),$$
(4.11)

$$\|\psi_n\|_{\dot{H}^1(\mathbb{R}^6)}^2 = \sum_{j=1}^M \|\psi^j\|_{\dot{H}^1(\mathbb{R}^6)}^2 + \|\Psi_n^M\|_{\dot{H}^1(\mathbb{R}^6)}^2 + o_n(1), \qquad (4.12)$$

$$\|\phi_n\|_{L^3(\mathbb{R}^6)}^3 = \sum_{j=1}^M \left\| e^{it_n^j \Delta} \phi^j \right\|_{L^3(\mathbb{R}^6)}^3 + \left\|\Phi_n^M\right\|_{L^3(\mathbb{R}^6)}^3 + o_n(1), \tag{4.13}$$

$$\|\psi_n\|_{L^3(\mathbb{R}^6)}^3 = \sum_{j=1}^M \left\| e^{\kappa i t_n^j \Delta} \psi^j \right\|_{L^3(\mathbb{R}^6)}^3 + \left\| \Psi_n^M \right\|_{L^3(\mathbb{R}^6)}^3 + o_n(1).$$
(4.14)

At the same, we have the orthogonality of the energy of linear profiles.

Corollary 4.5 (Energy pythagorean expansion). Under the assumptions of in Theorem 4.4, we have

$$H(\boldsymbol{\phi}_n) = \sum_{j=1}^M H\left(\boldsymbol{\phi}^j\right) + H\left(\boldsymbol{\Phi}_n^M\right) + o_n(1), \tag{4.15}$$

and

$$E(\boldsymbol{\phi}_n) = \sum_{j=1}^M E\left(\mathsf{S}\left(\frac{-t_n^j}{(\lambda_n^j)^2}\right)\boldsymbol{\phi}^j\right) + E\left(\boldsymbol{\Phi}_n^M\right) + o_n(1). \tag{4.16}$$

4.2. Nonlinear profile. First, we introduce the definition of nonlinear profile, which is associated with an initial data and a sequence of time, and then prove that it is well-defined in this subsection.

Definition 4.6. Let $\mathbf{v}_0 \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$, $\mathbf{v}(t, x) = \mathbf{S}(t)\mathbf{v}_0$ and let $\{t_n\}_{n=1}^{\infty}$ be a sequence with the limit $t_n \to t_\infty \in \mathbb{R} \cup \{\pm \infty\}$. We say that \mathbf{u} is a *nonlinear profile* associated with $(\mathbf{v}_0, \{t_n\}_{n=1}^{\infty})$ if there exists an interval $I \ni t_\infty$ (if $|t_\infty| = \infty$, $I = [a, +\infty)$ or $I = (-\infty, a]$) such that \mathbf{u} is a solution to (NLS system) in I and

$$\lim_{n \to \infty} \|\mathbf{u}(t_n, \cdot) - \mathbf{v}(t_n, \cdot)\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} = 0.$$

By the small data theory and Theorem 2.5, we can prove the existence and uniqueness of the nonlinear profile, which ensures the notation of the nonlinear profile is well-defined in Definition 4.6.

Proposition 4.7. There exists at least one nonlinear profile associated with $(\mathbf{v}_0, \{t_n\}_{n=1}^{\infty})$.

Proof. There are two cases to consider: either $|t_{\infty}| < \infty$ or $|t_{\infty}| = \infty$.

Case 1. If $|t_{\infty}| < \infty$, we take $\mathbf{u}_0 = \mathbf{v}(t_{\infty}, x) = \mathbf{S}(t_{\infty})\mathbf{v}_0$ and solve the initial value problem $\mathbf{u}|_{t=t_{\infty}} = \mathbf{u}_0$ in any time interval $I \ni t_{\infty}$ to obtain

$$\lim_{n \to \infty} \|\mathbf{u}(t_n, \cdot) - \mathbf{v}(t_n, \cdot)\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} \\
\leqslant \lim_{n \to \infty} \|\mathbf{S}(t_n - t_\infty)\mathbf{u}_0 - \mathbf{S}(t_n)\mathbf{v}_0\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} + \lim_{n \to \infty} \left\| \int_{t_\infty}^{t_n} \mathbf{S}(t_n - s)\mathbf{f}(\mathbf{u}(s))\mathrm{d}s \right\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} \\
= \lim_{n \to \infty} \|\mathbf{S}(t_n)\mathbf{v}_0 - \mathbf{S}(t_n)\mathbf{v}_0\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} + \lim_{n \to \infty} \left\| \int_{t_\infty}^{t_n} \mathbf{S}(t_n - s)\mathbf{f}(\mathbf{u}(s))\mathrm{d}s \right\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} \\
= 0.$$

Case 2. If $|t_{\infty}| = \infty$, we can assume $t_{\infty} = +\infty$ without loss of generality and solve the integral equation

$$\mathbf{u}(t) = \mathbf{S}(t)\mathbf{v}_0 + i\int_t^\infty \mathbf{S}(t-s)\mathbf{f}(\mathbf{u}(s))\mathrm{d}s$$

in $[t_N, +\infty) \times \mathbb{R}^6$ for $N \gg 1$ so large that $\|\mathbf{S}(t)\mathbf{v}_0\|_{\mathcal{S}([t_N, +\infty), \dot{\mathbf{H}}^1(\mathbb{R}^6))} \leq \delta_{sd}$, where δ_{sd} is as in Theorem 2.5. Then, \mathbf{u} is a solution in $I = [t_N, +\infty)$ with the terminal value \mathbf{v}_0 .

For large n, we have

$$\mathbf{u}(t_n) - \mathbf{v}(t_n) = i \int_{t_n}^{\infty} \mathbf{S}(t_n - s) \mathbf{f}(\mathbf{u}(s)) ds,$$

and

$$\|\nabla \mathbf{f}(\mathbf{u})\|_{\mathcal{S}'((t_n,+\infty),\mathrm{L}^2(\mathbb{R}^6))} < +\infty,$$

as in the proof of Theorem 2.5. So

$$\begin{split} &\lim_{n\to\infty} \|\mathbf{u}(t_n,\cdot) - \mathbf{v}(t_n,\cdot)\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} \leqslant \lim_{n\to\infty} \left\| \int_{t_n}^{\infty} \mathbf{S}(t_n-s) \mathbf{f}(\mathbf{u}(s)) \mathrm{d}s \right\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} \\ &\leqslant C \lim_{n\to+\infty} \|\nabla \mathbf{f}(\mathbf{u})\|_{\mathcal{S}'((t_n,\infty),\mathrm{L}^2(\mathbb{R}^6))} = 0. \end{split}$$

Combining the above two cases, we complete our proof.

Proposition 4.8. There exists at most one nonlinear profile associated with $(\mathbf{v}_0, \{t_n\}_{n=1}^{\infty})$ on an interval I with $t_{\infty} \in I$.

Proof. We can show it by contradiction. Assume $\mathbf{u}^{(1)}$ and $\mathbf{u}^{(2)}$ are both the non-linear profiles mentioned.

Case 1. If $|t_{\infty}| < \infty$, it is obvious from

$$\lim_{n \to \infty} \left\| \mathbf{u}^{(1)}(t_n, \cdot) - \mathbf{u}^{(2)}(t_n, \cdot) \right\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)}$$

$$\leqslant \lim_{n \to \infty} \left\| \mathbf{u}^{(1)}(t_n, \cdot) - \mathbf{v}(t_n, \cdot) \right\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} + \left\| \mathbf{u}^{(2)}(t_n, \cdot) - \mathbf{v}(t_n, \cdot) \right\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)}$$

$$= 0$$

and the well-posedness, the uniqueness in Duhamel formula, of **u**.

Case 2. If $|t_{\infty}| = \infty$, we can also assume $t_{\infty} = +\infty$. Because $\|\nabla \mathbf{u}^{(i)}\|_{\mathcal{S}(I, L^2(\mathbb{R}^6))} \leq \infty$ for i = 1, 2, for any $\varepsilon > 0$, there exists N > 0 such that $\forall n \ge N$

$$\left\|\nabla \mathbf{u}^{(i)}\right\|_{\mathcal{S}((t_n,+\infty),\mathrm{L}^2(\mathbb{R}^6))} < \varepsilon$$

By the proof of Theorem 2.5, for $M \gg N$, we have

$$\sup_{t \in (t_N, t_M)} \left\| \nabla \mathbf{u}^{(1)}(t, \cdot) - \nabla \mathbf{u}^{(2)}(t, \cdot) \right\|_{\mathbf{L}^2(\mathbb{R}^6)} \leqslant C \left\| \nabla \mathbf{u}^{(1)}(t_M, \cdot) - \nabla \mathbf{u}^{(2)}(t_M, \cdot) \right\|_{\mathbf{L}^2(\mathbb{R}^6)},$$

which implies that $\mathbf{u}^{(1)} \equiv \mathbf{u}^{(2)}$ on $(t_N, +\infty)$ and hence on I .

which implies that $\mathbf{u}^{(1)} \equiv \mathbf{u}^{(2)}$ on $(t_N, +\infty)$ and hence on I.

4.3. Existence of critical solution. In order to describe the properties of critical solution, especially compactness, we first briefly record a Palais-Smale condition modulo symmetries to describe the compactness.

Proposition 4.9 (Palais-Smale condition modulo symmetries, [36]). Let $\mathbf{u}_n : I_n \times \mathbb{R}^6 \to \mathbb{C}^2$ be a sequence of solutions to (NLS system) such that

$$\limsup_{n \to \infty} \sup_{t \in I_n} H(\mathbf{u}_n(t)) = H_c$$

and $\{t_n\}_{n=1}^{\infty} \subset I_n$ be a sequence of time such that

$$\lim_{n \to \infty} S_{\geq t_n}(\mathbf{u}_n) = \lim_{n \to \infty} S_{\leq t_n}(\mathbf{u}_n) = \infty.$$
(4.17)

Then there exists a subsequence of $\{\mathbf{u}_n(t_n)\}_{n=1}^{\infty}$ which converges in $\dot{\mathrm{H}}^1_x(\mathbb{R}^6)$ modulo symmetries.

Before the proof of Theorem 1.11, we introduce a notation L(H) by

$$L(H) := \sup \left\{ S_I(\mathbf{u}) \mid \mathbf{u} : I \times \mathbb{R}^6 \to \mathbb{C}^2, \sup_{t \in I} H(\mathbf{u}(t)) \leqslant H \right\}$$

for any $0 \leq H \leq H(\mathbf{W})$, and here the supremum is taken over all solutions \mathbf{u} : $I \times \mathbb{R}^6 \to \mathbb{C}^2$ to (NLS system) obeying $\sup_{t \in I} H(\mathbf{u}(t)) \leq H$. It is easy to see that $L : [0, H(\mathbf{W})] \to [0, \infty]$ is non-decreasing and satisfies $L(H(\mathbf{W})) = \infty$.

On the other hand, by the global existence and scattering result of small data, Theorem 2.5 and Proposition 1.4, we know

$$L(H) \leq CH^2, \quad \forall \ H < \delta_{sd},$$

where $\delta_{sd} > 0$ is the threshold arose in Theorem 2.5. By the long time perturbation theory and Proposition 2.7, we know that L is a continuous function, which suggests there exists a unique critical kinetic energy H_c such that

$$H_c \ge \delta_{sd} > 0,$$
 and $L(H) \begin{cases} < \infty, & H < H_c, \\ = \infty, & H \ge H_c. \end{cases}$ (4.18)

In particular, if $\mathbf{u} : I \times \mathbb{R}^6 \to \mathbb{C}^2$ is a maximal-lifespan solution to (NLS system) such that $\sup_{t \in I} H(\mathbf{u}(t)) < H_c$, then \mathbf{u} is global and

$$S_{\mathbb{R}}(\mathbf{u}) \leqslant L\left(\sup_{t\in\mathbb{R}} H(\mathbf{u}(t))\right) < \infty$$

Thus, the failure of Theorem 1.5 is equivalent to $0 < H_c < H(\mathbf{W})$.

Proof of Theorem 1.11. Because of the failure of Theorem 1.5, there exists a unique critical kinetic energy H_c satisfying (4.18). And we can find a sequence $\mathbf{u}_n : I_n \times \mathbb{R}^6 \to \mathbb{C}^2$ of solutions to (NLS system) with I_n compact, such that

$$\sup_{n \in \mathbb{N}} \sup_{t \in I_n} H(\mathbf{u}_n(t)) = H_c, \quad \text{and} \quad \lim_{n \to \infty} S_{I_n}(\mathbf{u}_n) = \infty.$$
(4.19)

Without loss of generality, we can take all $t_n \equiv 0$ in (4.17). Otherwise, if $t_n \in I_n$ be such that $S_{\geq t_n}(\mathbf{u}_n) = S_{\leq t_n}(\mathbf{u}_n)$, then let $\mathbf{u}_n(t) := \mathbf{u}(t_n - t)$.

By Proposition 4.9 and passing to a subsequence if necessary, we can find $\{\lambda_n\}_{n=1}^{\infty} \subset (0, +\infty)$ and a function $\mathbf{u}_{c,0} \in \dot{\mathrm{H}}^1(\mathbb{R}^6)$ such that

$$\lim_{n \to \infty} \|\mathcal{T}_{\lambda_n} \mathbf{u}_n(0) - \mathbf{u}_{c,0}\|_{\dot{\mathrm{H}}^1(\mathbb{R}^6)} = 0$$

Let $\mathbf{u}_c: I_c \times \mathbb{R}^6 \to \mathbb{C}^2$ be the maximal-lifespan solution to (NLS system) with initial data $\mathbf{u}_c(0) = \mathbf{u}_{c,0}$. Proposition 2.7 shows that $I_c \subset \liminf I_n$ and

$$\lim_{n \to \infty} \|\mathcal{T}_{\lambda_n} \mathbf{u}_n - \mathbf{u}_c\|_{L^{\infty}_t(K, \dot{\mathrm{H}}^1_x(\mathbb{R}^6))} = 0, \quad \text{for all compact } K \subset I_c.$$

Thus by (4.19),

$$\sup_{t \in I} H(\mathbf{u}(t)) \leqslant H_c. \tag{4.20}$$

We claim \mathbf{u}_c blows up both forward and backward in time. Indeed, if \mathbf{u}_c does not blow up forward in time, then $[0, \infty) \subset I_c$ and $S_{\geq 0}(\mathbf{u}_c) < \infty$. By Proposition 2.7, this implies $S_{\geq 0}(\mathbf{u}_n) = S_{\geq 0}(\mathcal{T}_{\lambda_n}\mathbf{u}_n) < \infty$ for sufficiently large n, which is a contradiction with (4.19). A similar argument proves that \mathbf{u}_c blows up backward in time, which completes the proof of our claim. Therefore, by (4.18),

$$\sup_{t \in I_c} H(\mathbf{u}_c(t)) \ge H_c. \tag{4.21}$$

Combining (4.21) with (4.20), we can obtain

$$\sup_{t \in I_c} H(\mathbf{u}_c(t)) = H_c. \tag{4.22}$$

It remains to show that \mathbf{u}_c is almost periodic modulo symmetries, which reminds us of Proposition 4.9. Considering an arbitrary sequence $\{\tau_n\}_{n=1}^{\infty} \subset I_c$, thanks to the fact that \mathbf{u}_c blows up in both time directions, we have

$$S_{\geqslant \tau_n}(\mathbf{u}_c) = S_{\leqslant \tau_n}(\mathbf{u}_c) = \infty.$$

By Proposition 4.9, there exists a subsequence of $\mathbf{u}_c(\tau_n)$, which is convergent in $\dot{\mathrm{H}}^1_x(\mathbb{R}^6)$ modulo symmetries. This implies the orbit $\mathcal{F}_c := \{\mathcal{T}_{\lambda_n}\mathbf{u}_c(\tau_n) \mid \tau_n \in I_c\}$ is precompact in $\dot{\mathrm{H}}^1_x(\mathbb{R}^6)$ modulo symmetries. By the precompactness of \mathcal{F}_c in $\dot{H}^1(\mathbb{R}^6)$ and the definition of \mathcal{T}_{λ} in Theorem 4.4,

$$\mathcal{T}_{\lambda_n} \mathbf{u}_c(\tau_n, x) = \left(\frac{1}{(\lambda_n)^2} u_c\left(\tau_n, \frac{x - x(\tau_n)}{\lambda_n}\right), \frac{1}{(\lambda_n)^2} v_c\left(\tau_n, \frac{x - x(\tau_n)}{\lambda_n}\right)\right)^t,$$

Recall Remark 1.10, we know for any $\eta > 0$, there exists a compactness modulus function $C(\eta) > 0$ such that

$$\int_{|x-x(\tau_n)| \ge \frac{C(\eta)}{\lambda_n}} \left(|\nabla u_c(\tau_n, x)|^2 + \frac{\kappa}{2} |\nabla v_c(\tau_n, x)|^2 \right) \mathrm{d}x \le \eta, \qquad \forall \ \tau_n \in I_c$$

and

$$\int_{|\xi| \ge C(\eta)\lambda_n} \left(|\xi|^2 |\widehat{u_c}(\tau_n, \xi)|^2 + \frac{\kappa}{2} |\xi|^2 |\widehat{v_c}(\tau_n, \xi)|^2 \right) \mathrm{d}\xi \leqslant \eta, \qquad \forall \ \tau_n \in I_c.$$

Choose the frequency scale function $\lambda(\tau_n) := \lambda_n$ in Definition 1.8 and we complete the proof of Theorem 1.11.

By Remark 1.10 and Theorem 1.11, we have established the compactness of $\{\mathbf{u}_c\} \subset \dot{\mathrm{H}}^1(\mathbb{R}^6)$. Therefore, we can describe them in a different way.

Remark 4.10. Similarly, one can use the strategy in [20] to prove Theorem 1.12. Define $\operatorname{osc}(T) := \inf_{t_0 \in J} \frac{\sup\{\lambda(t) \mid t \in J \text{ and } |t-t_0| \leq T[\lambda(t_0)]^{-2}\}}{\inf\{\lambda(t) \mid t \in J \text{ and } |t-t_0| \leq T[\lambda(t_0)]^{-2}\}}$ for any T > 0 and $a(t_0) := \frac{\lambda(t_0)}{\sup\{\lambda(t) \mid t \in J \text{ and } t \leq t_0\}} + \frac{\lambda(t_0)}{\sup\{\lambda(t) \mid t \in J \text{ and } t \geq t_0\}}$ for any $t_0 \in J$. Then the case of

$$\lim_{T \to \infty} \operatorname{ost}(T) = \infty, \ \inf_{t_0 \in J} a(t_0) = 0, \ I_c \neq \mathbb{R};$$

and

$$\lim_{T\to\infty}\operatorname{ost}(T)<\infty;$$

and

$$\lim_{T \to \infty} \operatorname{ost}(T) = \infty, \ \inf_{t_0 \in J} a(t_0) = 0, \ I_c = \mathbb{R}$$

or

$$\lim_{T \to \infty} osc(T) = \infty, \ \inf_{t_0 \in J} a(t_0) > 0$$

corresponds to the case of blowing-up solution in finite time, soliton-like solution, and low-to-high frequency cascade respectively.

5. NONEXISTENCE OF BLOWING-UP SOLUTION IN FINITE TIME

In the remaining part, we show the scattering result in Theorem 3.10. First of all, we prove the nonexistence of blowing-up solution.

Theorem 5.1. There dose not exist a solution to (NLS system) which blows up in finite time.

Proof. We argue by contradiction and suppose $\mathbf{u}_c : I_c \times \mathbb{R} \to \mathbb{C}^2$ is a finite-time blowing-up solution to (NLS system) with the maximal interval of existence $I_c = (-T_-(\mathbf{u}_c), T_+(\mathbf{u}_c))$. Without loss of generality, we may assume \mathbf{u}_c blows up forward in time, i.e. $T_+(\mathbf{u}_c) < \infty$.

First of all, we claim that

$$\liminf_{t \to T_{+}(\mathbf{u}_{c})} \lambda(t) = \infty.$$
(5.1)

Otherwise, if $\liminf_{t\to T_+(\mathbf{u}_c)} \lambda(t) < \infty$, we can choose $\{t_n\}_{n=1}^{\infty} \subset I_c$ such that $t_n \to T_+(\mathbf{u}_c)$ and define $\mathbf{u}_n : I_n \times \mathbb{R}^6 \to \mathbb{C}^2$ such that

$$\mathbf{u}_n(t,x) := \frac{1}{[\lambda(t_n)]^2} \mathbf{u}_c \left(t_n + \frac{t}{[\lambda(t_n)]^2}, x(t_n) + \frac{x}{\lambda(t_n)} \right),$$

where $I_n := \left\{ t \mid t_n \in I_c, t_n + \frac{t}{[\lambda(t_n)]^2} \in I_c \right\}$. It is easy to see that $0 \in I_n$ and $\{\mathbf{u}_n(t,x)\}_{n=1}^{\infty}$ is a sequence of solutions to (NLS system). Theorem 1.11 tells us the fact that \mathbf{u}_c is almost periodic modulo symmetries, combining with Remark 1.10, which implies $\{\mathbf{u}_n\}_{n=1}^{\infty} \subset \dot{\mathrm{H}}^1(\mathbb{R}^6)$ is precompact. Thus, after passing to a subsequence, there exists \mathbf{u}_0 such that

$$\lim_{n \to \infty} \|\mathbf{u}_n(0) - \mathbf{u}_0\|_{\dot{\mathbf{H}}^1_x(\mathbb{R}^6)} = 0.$$

We can get $\mathbf{u}_0 \neq \mathbf{0}$, by Sobolev embedding and the conservation of energy, from $H(\mathbf{u}_n(0)) = H(\mathbf{u}_c(t_n))$ and $\mathbf{u}_c \neq \mathbf{0}$. Let $\mathbf{u}(t, x)$ be the solution to (NLS system) associated to initial data $\mathbf{u}_0(x)$. On one hand, by Theorem 2.5 and Proposition 2.7, we can take n so large that there exists a compact interval $J \subset I_c$, on which \mathbf{u}_n is well-posed with finite scattering size. So, \mathbf{u}_c is well-posed with finite scattering size on the interval $\left\{t_n + \frac{t}{|\lambda(t_n)|^2} \mid t \in J\right\}$. On the other hand, we know that $t_n \to T_+(\mathbf{u}_c)$ and $\liminf_{n\to\infty} \lambda(t_n) = \liminf_{t\to T_+(\mathbf{u}_c)} \lambda(t) < \infty$, combining with $T_+(\mathbf{u}_c) < \infty$, which means \mathbf{u}_c is well-posed with finite scattering size on the interval $\left\{t_n + \frac{t}{|\lambda(t_n)|^2} \mid t \in J\right\} \cap (T_+(\mathbf{u}_c), +\infty)$, which is nonempty. That is impossible since \mathbf{u}_c blows up forward in time, i.e., $\exists t_1 \in I_c$ such that $S_{[t_1,T_+(\mathbf{u}_c))}(\mathbf{u}_c) = \infty$. As a consequence, (5.1) holds.

Set $\mathbf{u}_c := (u_c, v_c)^t$. For any $0 < \eta < 1$ and $t \in I_c$, by Hölder's inequality, we have

$$\begin{split} \int_{|x|\eta R} |u_c(t,x)|^2 \mathrm{d}x \\ &\leqslant \omega(6)\eta^2 R^2 \|u_c\|_{L^3_x(\mathbb{R}^6)}^2 + \omega(6) R^2 \left(\int_{|x-x(t)|>\eta R} |u_c(t,x)|^3 \mathrm{d}x \right)^{\frac{2}{3}} \end{split}$$

where $\omega(6)$ is a constant standing for the volume of unit ball in \mathbb{R}^6 . $\forall \varepsilon > 0$, let $\zeta = \zeta(\varepsilon, R) = 2^{-\frac{9}{4}} \varepsilon^{\frac{3}{2}} R^{-3} > 0$. Thanks to (5.1), there exists $\eta = \eta(\varepsilon, R, \zeta)$ such that $0 < \eta \leq 2^{-\frac{3}{4}} [H(\mathbf{W})]^{-\frac{1}{2}} \varepsilon^{\frac{1}{2}} R^{-1}$ and $C(\eta) \geq \lambda(t) C(\zeta)$. Thus,

$$\eta^2 R^2 \|u_c\|_{L^3_x(\mathbb{R}^6)}^2 \leqslant \eta^2 R^2 \|u_c\|_{\dot{H}^1_x(\mathbb{R}^6)}^2 < \eta^2 R^2 H(\mathbf{W}) \leqslant \frac{\varepsilon}{2}$$

and

$$\begin{aligned} R^2 \left(\int_{|x-x(t)|>\eta R} |u_c(t,x)|^3 \mathrm{d}x \right)^{\frac{2}{3}} &\leqslant R^2 \left(\int_{|x-x(t)|>C(\zeta)\frac{\eta R\lambda(t)}{C(\eta)}} |\nabla u_c(t,x)|^2 \mathrm{d}x \right)^{\frac{2}{3}} \\ &< R^2 \zeta^{\frac{2}{3}} = \frac{\varepsilon}{2}, \end{aligned}$$

which implies

$$\lim_{t \to \sup} \sup_{I_c} \int_{|x| < R} |u_c(t, x)|^2 \mathrm{d}x = 0, \quad \forall \ R > 0.$$
(5.2)

A similar way can help us to obtain

$$\limsup_{t \to \sup I_c} \int_{|x| < R} |v_c(t, x)|^2 \mathrm{d}x = 0, \quad \forall \ R > 0.$$
(5.3)

Let a(x) be a radial smooth function in Lemma 3.9 satisfying

$$a(x) = \begin{cases} 1, & |x| \le R, \\ 0, & |x| \ge 2R \end{cases}$$

and $|\nabla a(x)| \lesssim \frac{1}{|x|}$. Let

$$V_R(t) = \int_{\mathbb{R}^6} \left(2\kappa |u|^2 + |v|^2 \right) a(x) \mathrm{d}x.$$

One one hand, combining (5.2) with (5.3), we know

$$\limsup_{t \to \sup I_c} V_R(t) = \limsup_{t \to \sup I_c} \int_{\mathbb{R}^6} \left(2\kappa |u_c(t,x)|^2 + |v_c(t,x)|^2 \right) a(x) \mathrm{d}x = 0.$$
(5.4)

On the other hand, by Hardy's inequality and (1.5), we get

$$\begin{aligned} |V_{R}'(t)| &= 2\kappa \left| \int_{\mathbb{R}^{6}} (2\overline{u}_{c} \nabla u_{c} + \overline{v}_{c} \nabla v_{c}) \cdot \nabla a(x) \mathrm{d}x \right| \\ &\lesssim_{\kappa} \|\nabla \mathbf{u}_{c}\|_{\mathrm{L}^{2}(\mathbb{R}^{6})} \left\| \frac{\mathbf{u}_{c}}{|x|} \right\|_{\mathrm{L}^{2}(\mathbb{R}^{6})} \\ &\lesssim_{\kappa} [H(\mathbf{u}_{c})]^{2} < [H(\mathbf{W})]^{2}. \end{aligned}$$

So, Newton-Leibniz's formula tells us,

$$V_R(t_1) \lesssim_{\kappa} I_R(t_2) + |t_1 - t_2| [H(\mathbf{W})]^2, \quad \forall t_1, t_2 \in I_c.$$

Let $t_2 \to T_+(\mathbf{u}_c)$ and using (5.4), we obtain

$$V_R(t_1) \lesssim_{\kappa} |\sup I_c - t_1| [H(\mathbf{W})]^2, \quad \forall t_1 \in I_c$$

By the conservation of mass and the above estimate, we have

$$M(\mathbf{u}_{c,0}) = M(\mathbf{u}_{c}(t_{1})) \sim \int_{\mathbb{R}^{6}} 2\kappa |u_{c}(t_{1},x)|^{2} + |v_{c}(t_{1},x)|^{2} dx$$

= $\lim_{\mathbb{R}\to\infty} V_{R}(t_{1}) \lesssim_{\kappa} |\sup I_{c} - t_{1}|[H(\mathbf{W})]^{2}, \quad \forall t_{1} \in I_{c}$

Let $t_1 \to T_+(\mathbf{u}_c)$, we know $\mathbf{u}_{c,0} \equiv \mathbf{0}$. And thus, by the uniqueness of the solution to (NLS system), $\mathbf{u}_c \equiv \mathbf{0}$, which is a contradiction with (1.5).

6. NONEXISTENCE OF GLOBAL BLOWING-UP SOLUTION

Under the assumption of Theorem 1.5, we can prove $H_c = 0$ for any critical solution \mathbf{u}_c to (NLS system) and show the nonexistence of the ones. We prove the negative regularity, zero momentum and state the control of spatial center function x(t) of soliton-like solutions in Subsection 6.1, which are very powerful in excluding of our enemies. According to the unique properties of soliton-like solution and low-to-high frequency cascade, we make full use of Theorem 1.13 and Corollary 6.6 to rule out them one by one in Subsection 6.2 and Subsection 6.3.

6.1. Negative regularity. We show the proof of Theorem 1.13 in this subsection. To begin with, we present a special property of critical solution \mathbf{u}_c .

Proposition 6.1 (Reduced Duhamel formulas, [20, 25]). Let $\mathbf{u}_c : I_c \times \mathbb{R}^6 \to \mathbb{C}^2$ be a maximal-lifespan almost periodic modulo symmetries solution to (NLS system). Then $\mathcal{S}(-t)\mathbf{u}_c$ converges weakly to $\mathbf{0}$ in $\dot{\mathbf{H}}^1_x$ as $t \to \sup I_c$ or $t \to \inf I_c$. In particular, we have the 'reduced' Duhamel formulas

$$\mathbf{u}_{c}(t) = i \lim_{T \to \sup I_{c}} \int_{t}^{T} \mathbf{S}(t-s) \mathbf{f}(\mathbf{u}_{c}(s)) ds$$
$$= -i \lim_{T \to \inf I_{c}} \int_{T}^{t} \mathbf{S}(t-s) \mathbf{f}(\mathbf{u}_{c}(s)) ds$$

where the limits are to be understood in the weak $\dot{\mathrm{H}}^1_x$ topology.

Now, we turn to show the negative regularity of global almost periodic solution \mathbf{u}_c to (NLS system), which plays an important role in the preclusion of soliton-like solutions and low-to-high frequency cascades.

Proof of Theorem 1.13. Since $\mathbf{u}_c(t)$ is almost periodic modulo symmetries with $\lambda(t) \ge 1$, then there exists $N_0 = N_0(\eta)$ such that

$$\|\nabla (P_{\leqslant N_0} \mathbf{u}_c)\|_{\mathbf{L}^\infty_t(\mathbb{R}, \mathbf{L}^2_x(\mathbb{R}^6))} \leqslant \eta, \qquad \forall \ \eta > 0.$$
(6.1)

Denote A(N) by

$$A(N) := N^{-\frac{1}{2}} \| P_N \mathbf{u}_c \|_{\mathbf{L}^{\infty}_t(\mathbb{R}, \mathbf{L}^4_x(\mathbb{R}^6))},$$

for $N \leq 10N_0$.

Lemma 6.2 (Recurrence). Let A(N) be defined as above, for any $N \leq 10N_0$, we have

$$A(N) \lesssim_{\mathbf{u}_{c}} \left(\frac{N}{N_{0}}\right)^{\frac{1}{2}} + \eta \sum_{\frac{N}{10} \leqslant N_{1} \leqslant N_{0}} \left(\frac{N}{N_{1}}\right)^{\frac{1}{2}} A(N_{1}) + \eta \sum_{N_{1} < \frac{N}{10}} \left(\frac{N_{1}}{N}\right)^{\frac{1}{2}} A(N_{1}).$$
(6.2)

Proof. Fix $N \leq 10N_0$. By time-translation symmetry, it suffices to prove

$$N^{-\frac{1}{2}} \|\mathbf{u}_{c,0}\|_{\mathrm{L}^{4}} \lesssim \left(\frac{N}{N_{0}}\right)^{\frac{1}{2}} + \eta \sum_{\frac{N}{10} \leqslant N_{1} \leqslant N_{0}} \left(\frac{N}{N_{1}}\right)^{\frac{1}{2}} A(N_{1}) + \eta \sum_{N_{1} < \frac{N}{10}} \left(\frac{N_{1}}{N}\right)^{\frac{1}{2}} A(N_{1}).$$
(6.3)

By Lemma 2.1, Lemma 2.3, and Proposition 6.1, we get

$$N^{-\frac{1}{2}} \| P_N \mathbf{u}_{c,0} \|_{\mathrm{L}^4(\mathbb{R}^6)} \leqslant N^{-\frac{1}{2}} \left\| \int_0^{N^{-2}} \mathbf{S}(-\tau) P_N \mathbf{f}(\mathbf{u}_c(\tau)) \mathrm{d}\tau \right\|_{\mathrm{L}^4_x(\mathbb{R}^6)} + N^{-\frac{1}{2}} \int_{N^{-2}}^{\infty} \left\| \mathbf{S}(-\tau) P_N \mathbf{f}(\mathbf{u}_c(\tau)) \right\|_{\mathrm{L}^4_x(\mathbb{R}^6)} \mathrm{d}\tau \lesssim N^{\frac{1}{2}} \| P_N \mathbf{f}(\mathbf{u}_c) \|_{\mathrm{L}^\infty_t(\mathbb{R},\mathrm{L}^{\frac{4}{3}}_x(\mathbb{R}^6))}.$$

Let $\mathbf{f}(\mathbf{z}) = (f_1(\mathbf{z}), f_2(\mathbf{z}))^T$ for any $\mathbf{z} = (z_1, z_2)^T \in \mathbb{C}^2$, where $f_1(\mathbf{z}) = z_2 \bar{z}_1$ and $f_2(\mathbf{z}) = z_1^2$. Thus, for $\mathbf{u}_c = (u_c, v_c)^T$, we have

$$\begin{aligned} f_1(\mathbf{u}_c) - f_1(P_{>N_0}\mathbf{u}_c) &= v_c \overline{u_c} - (P_{>N_0}v_c)(P_{>N_0}\overline{u_c}) \\ &= (P_{\leqslant N}v_c + P_{>N}v_c)(P_{\leqslant N}\overline{u_c} + P_{>N}\overline{u_c}) - (P_{>N}v_c)(P_{>N}\overline{u_c}) \\ &= (P_{\leqslant N}v_c)(P_{\leqslant N}\overline{u_c}) + (P_{>N}v_c)(P_{\leqslant N}\overline{u_c}) + (P_{\leqslant N}v_c)(P_{>N}\overline{u_c}), \end{aligned}$$

and

$$f_{2}(\mathbf{u}_{c}) - f_{2}(P_{>N_{0}}\mathbf{u}_{c}) = u_{c}^{2} + (P_{>N_{0}}u_{c})^{2}$$

= $(P_{\leq N}u_{c} + P_{>N}u_{c})^{2} - (P_{>N}u_{c})^{2}$
= $(P_{\leq N}u_{c})^{2} + 2(P_{>N}u_{c})(P_{\leq N}u_{c}).$

By the Fundamental Theorem of Calculus, we obtain

$$f_i(\mathbf{z}^1) - f_i(\mathbf{z}^2) = (\mathbf{z}^1 - \mathbf{z}^2) \int_0^1 \partial_{\mathbf{z}} f_i(\mathbf{z}^1 + \theta(\mathbf{z}^1 - \mathbf{z}^2)) d\theta + \overline{(\mathbf{z}^1 - \mathbf{z}^2)} \int_0^1 \partial_{\mathbf{\bar{z}}} f_i(\mathbf{z}^1 + \theta(\mathbf{z}^1 - \mathbf{z}^2)) d\theta, \quad i = 1, 2.$$

where $\partial_{\mathbf{z}} f_1(\mathbf{z}) = (0, \bar{z}_1)^T$, $\partial_{\bar{\mathbf{z}}} f_1(\mathbf{z}) = (z_2, 0)^T$, $\partial_{\mathbf{z}} f_2(\mathbf{z}) = (2z_1, 0)^T$ and $\partial_{\bar{\mathbf{z}}} f_2(\mathbf{z}) = \mathbf{0}$. Then, we get

$$\begin{split} \mathbf{f}(\mathbf{u}_{c}) &= (f_{1}(\mathbf{u}_{c}), f_{2}(\mathbf{u}_{c}))^{T} \\ &= \left((P_{>N_{0}}v_{c})(P_{\leqslant N_{0}}\overline{u_{c}}) + (P_{\leqslant N_{0}}v_{c})(P_{>N_{0}}\overline{u_{c}}), 2(P_{>N_{0}}u_{c})(P_{\leqslant N_{0}}u_{c}) \right)^{T} \\ &+ \mathbf{f}(P_{>N_{0}}\mathbf{u}_{c}) + \mathbf{f}(P_{\frac{N}{10}\leqslant \cdot \leqslant N_{0}}\mathbf{u}_{c}) \\ &+ \left(P_{<\frac{N}{10}}\mathbf{u}_{c} \int_{0}^{1} \partial_{\mathbf{z}}f_{1}(P_{\frac{N}{10}\leqslant \cdot \leqslant N_{0}}\mathbf{u}_{c} + \theta P_{<\frac{N}{10}}\mathbf{u}_{c}) \mathrm{d}\theta \\ &+ \overline{P_{<\frac{N}{10}}}\mathbf{u}_{c} \int_{0}^{1} \partial_{\mathbf{z}}f_{2}(P_{\frac{N}{10}\leqslant \cdot \leqslant N_{0}}\mathbf{u}_{c} + \theta P_{<\frac{N}{10}}\mathbf{u}_{c}) \mathrm{d}\theta \\ &+ \overline{P_{<\frac{N}{10}}}\mathbf{u}_{c} \int_{0}^{1} \partial_{\mathbf{z}}f_{2}(P_{\frac{N}{10}\leqslant \cdot \leqslant N_{0}}\mathbf{u}_{c} + \theta P_{<\frac{N}{10}}\mathbf{u}_{c}) \mathrm{d}\theta \\ &+ \overline{P_{<\frac{N}{10}}}\mathbf{u}_{c} \int_{0}^{1} \partial_{\mathbf{z}}f_{2}(P_{\frac{N}{10}\leqslant \cdot \leqslant N_{0}}\mathbf{u}_{c} + \theta P_{<\frac{N}{10}}\mathbf{u}_{c}) \mathrm{d}\theta \\ &+ \overline{P_{<\frac{N}{10}}}\mathbf{u}_{c} \int_{0}^{1} \partial_{\mathbf{z}}f_{2}(P_{\frac{N}{10}\leqslant \cdot \leqslant N_{0}}\mathbf{u}_{c} + \theta P_{<\frac{N}{10}}\mathbf{u}_{c}) \mathrm{d}\theta \\ &+ \overline{P_{<\frac{N}{10}}}\mathbf{u}_{c} \int_{0}^{1} \partial_{\mathbf{z}}f_{2}(P_{\frac{N}{10}\leqslant \cdot \leqslant N_{0}}\mathbf{u}_{c} + \theta P_{<\frac{N}{10}}\mathbf{u}_{c}) \mathrm{d}\theta \right)^{T} \\ =: \mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}. \end{split}$$

Using Hölder and Bernstein,

$$N^{\frac{1}{2}} \| P_{N}(\mathcal{A} + \mathcal{B}) \|_{L^{\infty}_{t}(\mathbb{R}, L^{\frac{4}{3}}_{x}(\mathbb{R}^{6}))} \lesssim N^{\frac{1}{2}} \| P_{>N_{0}} \mathbf{u}_{c} \|_{L^{\infty}_{t}(\mathbb{R}, L^{\frac{12}{5}}_{x}(\mathbb{R}^{6}))} \| \mathbf{u}_{c} \|_{L^{\infty}_{t}(\mathbb{R}, L^{3}_{x}(\mathbb{R}^{6}))}$$
$$\lesssim N^{\frac{1}{2}} N^{\frac{1}{2}}_{0}.$$

Next, we turn to estimate the fourth term $\mathcal D$ and define $\mathcal D=\mathcal D_1+\mathcal D_2$ for

$$\mathcal{D}_{1} = \left(P_{\leq \frac{N}{10}} \mathbf{u}_{c} \int_{0}^{1} \partial_{\mathbf{z}} f_{1} (P_{\frac{N}{10} \leqslant \cdot \leqslant N_{0}} \mathbf{u}_{c} + \theta P_{\leq \frac{N}{10}} \mathbf{u}_{c}) \mathrm{d}\theta, + P_{\leq \frac{N}{10}} \mathbf{u}_{c} \int_{0}^{1} \partial_{\mathbf{z}} f_{2} (P_{\frac{N}{10} \leqslant \cdot \leqslant N_{0}} \mathbf{u}_{c} + \theta P_{\leq \frac{N}{10}} \mathbf{u}_{c}) \mathrm{d}\theta \right)^{T},$$

and

$$\mathcal{D}_{2} = \left(\overline{P_{<\frac{N}{10}}}\mathbf{u}_{c}\int_{0}^{1}\partial_{\bar{\mathbf{z}}}f_{1}(P_{\frac{N}{10}\leqslant\cdot\leqslant N_{0}}\mathbf{u}_{c} + \theta P_{<\frac{N}{10}}\mathbf{u}_{c})\mathrm{d}\theta, + \overline{P_{<\frac{N}{10}}}\mathbf{u}_{c}\int_{0}^{1}\partial_{\bar{\mathbf{z}}}f_{2}(P_{\frac{N}{10}\leqslant\cdot\leqslant N_{0}}\mathbf{u}_{c} + \theta P_{<\frac{N}{10}}\mathbf{u}_{c})\mathrm{d}\theta\right)^{T}.$$

It suffices to consider \mathcal{D}_1 since \mathcal{D}_2 can be handled similarly. Bernstein inequality yields

$$\|P_{\geq \frac{N}{10}}\partial_{\mathbf{z}}f_i(\mathbf{u}_c)\|_{\mathbf{L}_t^\infty(\mathbb{R},\mathbf{L}_x^2(\mathbb{R}^6))} \lesssim N^{-1}\|\nabla\mathbf{u}_c\|_{\mathbf{L}_t^\infty(\mathbb{R},\mathbf{L}_x^2(\mathbb{R}^6))}, \qquad i=1,2.$$

Thus, by Hölder's inequality and (6.1),

$$\begin{split} N^{\frac{1}{2}} \|P_{N}\mathcal{D}_{1}\|_{\mathrm{L}_{t}^{\infty}(\mathbb{R},\mathrm{L}_{x}^{\frac{4}{3}}(\mathbb{R}^{6}))} \\ &= N^{\frac{1}{2}} \sum_{i=1}^{2} \left\|P_{N}\left(P_{<\frac{N}{10}}\mathbf{u}_{c}\int_{0}^{1}\partial_{\mathbf{z}}f_{i}(P_{\frac{N}{10}\leqslant\cdot\leqslant N_{0}}\mathbf{u}_{c}+\theta P_{<\frac{N}{10}}\mathbf{u}_{c})\mathrm{d}\theta\right)\right\|_{\mathrm{L}_{t}^{\infty}(\mathbb{R},\mathrm{L}_{x}^{\frac{4}{3}}(\mathbb{R}^{6}))} \\ &\lesssim N^{\frac{1}{2}}\|P_{<\frac{N}{10}}\mathbf{u}_{c}\|_{\mathrm{L}_{t}^{\infty}(\mathbb{R},\mathrm{L}_{x}^{4}(\mathbb{R}^{6}))}\sum_{i=1}^{2}\left\|P_{>\frac{N}{10}}\left(\int_{0}^{1}\partial_{\mathbf{z}}f_{i}(P_{\frac{N}{10}\leqslant\cdot\leqslant N_{0}}\mathbf{u}_{c}+\theta P_{<\frac{N}{10}}\mathbf{u}_{c})\mathrm{d}\theta\right)\right\|_{\mathrm{L}_{t}^{\infty}(\mathbb{R},\mathrm{L}_{x}^{2}(\mathbb{R}^{6}))} \\ &\lesssim N^{-\frac{1}{2}}\|P_{<\frac{N}{10}}\mathbf{u}_{c}\|_{\mathrm{L}_{t}^{\infty}(\mathbb{R},\mathrm{L}_{x}^{4}(\mathbb{R}^{6}))}\|\nabla P_{\leqslant N_{0}}\mathbf{u}_{c}\|_{\mathrm{L}_{t}^{\infty}(\mathbb{R},\mathrm{L}_{x}^{2}(\mathbb{R}^{6}))} \\ &\lesssim \eta\sum_{N_{1}<\frac{N}{10}}\left(\frac{N_{1}}{N}\right)^{\frac{1}{2}}A(N_{1}). \end{split}$$

Then, we have

$$N^{\frac{1}{2}} \| P_N \mathcal{D} \|_{L_t^{\infty}(\mathbb{R}, L_x^{\frac{4}{3}}(\mathbb{R}^6))} = N^{\frac{1}{2}} \sum_{j=1}^2 \| P_N \mathcal{D}_j \|_{L_t^{\infty}(\mathbb{R}, L_x^{\frac{4}{3}}(\mathbb{R}^6))} \lesssim \eta \sum_{N_1 < \frac{N}{10}} \left(\frac{N_1}{N} \right)^{\frac{1}{2}} A(N_1).$$

We are left to estimate the contribution of \mathcal{C} . It remains to show

$$\|P_N \mathcal{C}\|_{\mathrm{L}^{\infty}_t(\mathbb{R},\mathrm{L}^{\frac{4}{3}}_x(\mathbb{R}^6))} \lesssim \eta \sum_{\frac{N}{10} \leqslant N_1 \leqslant N_0} N_1^{-\frac{1}{2}} A(N_1).$$

Using the triangle inequality, Bernstein, (6.1) and Hölder, we estimate as follow:

$$\begin{split} \|P_{N}\mathcal{C}\|_{L_{t}^{\infty}(\mathbb{R},L_{x}^{\frac{4}{3}}(\mathbb{R}^{6}))} &\lesssim \|\mathbf{f}(P_{\frac{N}{10}\leqslant\cdot\leqslant N_{0}}\mathbf{u}_{c})\|_{L_{t}^{\infty}(\mathbb{R},L_{x}^{\frac{4}{3}}(\mathbb{R}^{6}))} \\ &\lesssim \sum_{\frac{N}{10}\leqslant N_{1},N_{2}\leqslant N_{0}} \left(\|\overline{(P_{N_{1}}u_{c})}(P_{N_{2}}v_{c})\|_{L_{t}^{\infty}(\mathbb{R},L_{x}^{\frac{4}{3}}(\mathbb{R}^{6}))} + \|(P_{N_{1}}u_{c})(P_{N_{2}}u_{c})\|_{L_{t}^{\infty}(\mathbb{R},L_{x}^{\frac{4}{3}}(\mathbb{R}^{6}))} \right) \\ &\lesssim \eta \sum_{\frac{N}{10}\leqslant N_{1}\leqslant N_{2}} N_{2}^{-1} \|P_{N_{1}}u_{c}\|_{L_{t}^{\infty}(\mathbb{R},L_{x}^{2}(\mathbb{R}^{6}))} \\ &+ \sum_{\frac{N}{10}\leqslant N_{2}\leqslant N_{1}\leqslant N_{0}} \|P_{N_{1}}u_{c}\|_{L_{t}^{\infty}(\mathbb{R},L_{x}^{2}(\mathbb{R}^{6}))} \|P_{N_{2}}v_{c}\|_{L_{t}^{\infty}(\mathbb{R},L_{x}^{4}(\mathbb{R}^{6}))} \\ &\lesssim \eta \sum_{\frac{N}{10}\leqslant N_{1}\leqslant N_{0}} N_{1}^{-\frac{1}{2}}A(N_{1}) + \eta \sum_{\frac{N}{10}\leqslant N_{2}\leqslant N_{1}\leqslant N_{0}} \left(\frac{N_{2}}{N_{1}}\right) (N_{2}^{-\frac{1}{2}}A(N_{2})) \\ &\lesssim \eta \sum_{\frac{N}{10}\leqslant N_{1}\leqslant N_{0}} N_{1}^{-\frac{1}{2}}A(N_{1}). \end{split}$$

Finally, we put the above estimates together to obtain

$$\begin{aligned} A(N) &= N^{\frac{1}{2}} \| P_N(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}) \|_{L^{\infty}_t(\mathbb{R}, L^{\frac{4}{3}}_x(\mathbb{R}^6))} \\ &\lesssim_{\mathbf{u}_c} \left(\frac{N}{N_0} \right)^{\frac{1}{2}} + \eta \sum_{\frac{N}{10} \leqslant N_1 \leqslant N_0} \left(\frac{N}{N_1} \right)^{\frac{1}{2}} A(N_1) + \eta \sum_{N_1 < \frac{N}{10}} \left(\frac{N_1}{N} \right)^{\frac{1}{2}} A(N_1), \end{aligned}$$

which implies that (6.2) holds.

Proposition 6.3. Let \mathbf{u}_c be as in Theorem 1.13. Then

$$\mathbf{u}_c \in \mathcal{L}^{\infty}_t(\mathbb{R}, \mathcal{L}^p_x(\mathbb{R}^6)) \quad for \ \frac{14}{5} \leq p < 3.$$

In particular, by Hölder's inequality,

$$\nabla \mathbf{f}(\mathbf{u}_c) \in \mathcal{L}^{\infty}_t(\mathbb{R}, \mathcal{L}^r_x(\mathbb{R}^6)) \quad for \ \frac{7}{6} \leqslant r < \frac{6}{5}.$$

Proof. Combining Lemma 6.2 with Lemma 2.2, we deduce

$$\|P_N \mathbf{u}_c\|_{\mathbf{L}^{\infty}_t(\mathbb{R}, \mathbf{L}^{\frac{4}{3}}_x(\mathbb{R}^6))} \lesssim N \quad \text{for } N \leqslant 10N_0.$$
(6.4)

In fact, setting $N = 10 \cdot 2^{-k} N_0$, $x_k = A(10 \cdot 2^{-k} N_0)$ and $\gamma = 1$ in Lemma 2.2, we can obtain

$$A(10 \cdot 2^{-k} N_0) \lesssim \sum_{l=0}^{\infty} r^{|k-l|} (10 \cdot 2^{-l})^{\frac{1}{2}}.$$

Then taking $\eta > 0$ small enough such that $r < \sqrt{2}$, we have

$$A(10 \cdot 2^{-k} N_0) \lesssim (10 \cdot 2^{-k})^{\frac{1}{2}} \sum_{l=0}^{\infty} r^{|k-l|} (10 \cdot 2^{-l+k})^{\frac{1}{2}} \lesssim (10 \cdot 2^{-k})^{\frac{1}{2}}.$$

Observing the definition of A(N), we deduce (6.4) by

$$\|P_N \mathbf{u}_c\|_{\mathbf{L}^{\infty}_t(\mathbb{R},\mathbf{L}^{\frac{4}{3}}_x(\mathbb{R}^6))} \lesssim N_0^{-\frac{1}{2}} N^{\frac{1}{2}} (10 \cdot 2^{-k} N_0)^{\frac{1}{2}} \lesssim N.$$

By interpolation, we get

 $\|P_N \mathbf{u}_c\|_{\mathrm{L}^{\infty}_t(\mathbb{R},\mathrm{L}^p_x(\mathbb{R}^6))} \lesssim \|P_N \mathbf{u}_c\|_{\mathrm{L}^{\infty}_t(\mathbb{R},L^4_x(\mathbb{R}^6))}^{\frac{2(p-2)}{p}} \|P_N \mathbf{u}_c\|_{\mathrm{L}^{\infty}_t(\mathbb{R},\mathrm{L}^2_x(\mathbb{R}^6))}^{\frac{4}{p}-2} \lesssim N^{3-\frac{8}{p}} \lesssim N^{\frac{1}{7}}$ for all $N \leq 10N_0$. Using Lemma 2.1,

$$\begin{split} \|\mathbf{u}_{c}\|_{\mathcal{L}^{\infty}_{t}(\mathbb{R},\mathcal{L}^{p}_{x}(\mathbb{R}^{6}))} \leqslant \|P_{\leqslant N_{0}}\mathbf{u}_{c}\|_{\mathcal{L}^{\infty}_{t}(\mathbb{R},\mathcal{L}^{p}_{x}(\mathbb{R}^{6}))} + \|P_{>N_{0}}\mathbf{u}_{c}\|_{\mathcal{L}^{\infty}_{t}(\mathbb{R},\mathcal{L}^{p}_{x}(\mathbb{R}^{6}))} \\ \lesssim \sum_{N\leqslant N_{0}} N^{\frac{1}{7}} + \sum_{N>N_{0}} N^{2-\frac{6}{p}} \lesssim 1, \end{split}$$

which completes the proof of the proposition.

We can use the decomposition of frequency to deduce the following proposition.

Proposition 6.4 (Some negative regularity, [19]). Let \mathbf{u}_c be as in Theorem 1.13. If $|\nabla|^s \mathbf{f}(\mathbf{u}_c) \in \mathrm{L}^{\infty}_t(\mathbb{R}, \mathrm{L}^p_x(\mathbb{R}^6))$ for some $\frac{7}{6} \leq p < \frac{6}{5}$ and some $s \in [0, 1]$, then there exists $s_0 = s_0(p) > 0$ such that $\mathbf{u}_c \in \mathrm{L}^{\infty}_t(\mathbb{R}, \mathrm{\dot{H}}^{s-s_0+}_x(\mathbb{R}^6))$.

28

To end the proof of Theorem 1.13, we first apply Proposition 6.4 with s = 1 to show that $\mathbf{u} \in \mathrm{L}^{\infty}_{t}(\dot{\mathrm{H}}^{1-s_{0}+}_{x}(\mathbb{R}^{6}))$ for some $s_{0} > 0$. The fractional chain rule and Proposition 6.3 tell us that $|\nabla|^{1-s_{0}+}\mathbf{f}(\mathbf{u}_{c}) \in \mathrm{L}^{\infty}_{t}(\mathbb{R},\mathrm{L}^{p}_{x}(\mathbb{R}^{6}))$ for some $\frac{7}{6} \leq p < \frac{6}{5}$. Thus, another application of Proposition 6.4 facilitates us to obtain $\mathbf{u} \in \mathrm{L}^{\infty}_{t}(\dot{\mathrm{H}}^{1-2s_{0}+}_{x}(\mathbb{R}^{6}))$. Iterating this procedure finitely many times, we derive $\mathbf{u} \in \mathrm{L}^{\infty}_{t}(\dot{\mathrm{H}}^{-\varepsilon}_{x}(\mathbb{R}^{6}))$ for any $0 < \varepsilon < s_{0}$.

Proposition 6.5. If $\mathbf{u}_c := (u_c, v_c)^T$ be a minimal-kinetic-energy blowing-up solution to (NLS system) satisfying $\mathbf{u}_c \in \mathbf{L}_t^{\infty}(I_c, \mathbf{H}_x^1(\mathbb{R}^6))$ and its initial data $\mathbf{u}_{c,0}$ to be non-radial with the mass-resonance condition or $\mathbf{u}_{c,0}$ to be radial, then its momentum is zero, that is,

$$P(\mathbf{u}_c) := \operatorname{Im} \int_{\mathbb{R}^6} \left(\overline{u}_c \nabla u_c + \frac{1}{2} \overline{v_c} \nabla v_c \right) \mathrm{d}x \equiv \mathbf{0}.$$
(6.5)

Proof. First of all, we know that \mathbf{u}_c will be radial if its initial data $\mathbf{u}_{c,0}$ is radial. Combining with the fact that $\nabla f(x) = f'(r)\frac{x}{r}$ for radial function f(r) := f(x) when r = |x|, we have

$$P(\mathbf{u}_c) = \operatorname{Im} \int_{\mathbb{R}^6} \left(\overline{u}_c \nabla u_c + \frac{1}{2} \overline{v_c} \nabla v_c \right) dx$$
$$= \operatorname{Im} \int_{\mathbb{R}^6} \left(\overline{u_c(r)} u_c'(r) \frac{x}{r} + \frac{1}{2} \overline{v_c(r)} v_c'(r) \frac{x}{r} \right) dx$$
$$\equiv \mathbf{0},$$

where we have used the fact that $\int_{\mathbb{R}^6} g(x) dx \equiv 0$ for any odd function.

We are left to show the remaining case, i.e. $\mathbf{u}_{c,0}$ is non-radial with the massresonance condition. Under the mass-resonance condition, that is $\kappa = \frac{1}{2}$, (NLS system) enjoys the Galilean invariance property:

$$\mathbf{u}_{c}(t,x) = \begin{pmatrix} u_{c}(t,x) \\ v_{c}(t,x) \end{pmatrix} \to \begin{pmatrix} u_{c}^{\xi}(t,x) \\ v_{c}^{\xi}(t,x) \end{pmatrix} =: \begin{pmatrix} e^{ix\cdot\xi} e^{-it|\xi|^{2}} u_{c}(t,x-2t\xi) \\ e^{2ix\cdot\xi} e^{-2it|\xi|^{2}} v_{c}(t,x-2t\xi) \end{pmatrix}$$

for any $\xi \in \mathbb{R}^6$. Then, for $\mathbf{u}_c^{\xi} := (u_c^{\xi}, v_c^{\xi})^T$, we can compute

$$\begin{split} H(\mathbf{u}_{c}^{\xi}) &= \|\nabla u_{c}^{\xi}\|_{L^{2}(\mathbb{R}^{6})}^{2} + \frac{\kappa}{2} \|\nabla v_{c}^{\xi}\|_{L^{2}(\mathbb{R}^{6})}^{2} \\ &= \|i\xi u_{c} + \nabla u_{c}\|_{L^{2}(\mathbb{R}^{6})}^{2} + \frac{\kappa}{2} \|2i\xi v_{c} + \nabla v_{c}\|_{L^{2}(\mathbb{R}^{6})}^{2} \\ &= \int_{\mathbb{R}^{6}} |\operatorname{Re}(\nabla u_{c}) - \xi \operatorname{Im} u_{c}|^{2} + |\xi \operatorname{Re} u_{c} + \operatorname{Im}(\nabla u_{c})|^{2} dx \\ &+ \frac{\kappa}{2} \int_{\mathbb{R}^{6}} |\operatorname{Re}(\nabla v_{c}) - 2\xi \operatorname{Im} v_{c}|^{2} + |2\xi \operatorname{Re} v_{c} + \operatorname{Im}(\nabla v_{c})|^{2} dx \\ &= \int_{\mathbb{R}^{6}} |\nabla u_{c}|^{2} + |\xi|^{2} |u_{c}|^{2} + 2\xi (\operatorname{Re} u_{c} \operatorname{Im}(\nabla u_{c}) - \operatorname{Re}(\nabla u_{c}) \operatorname{Im} u_{c}) dx \\ &+ \frac{\kappa}{2} \int_{\mathbb{R}^{6}} |\nabla v_{c}|^{2} + 4|\xi|^{2} |v_{c}|^{2} + 4\xi (\operatorname{Re} v_{c} \operatorname{Im}(\nabla v_{c}) - \operatorname{Re}(\nabla v_{c}) \operatorname{Im} v_{c}) dx. \end{split}$$

Note that $\kappa = \frac{1}{2}$, we have

$$H(\mathbf{u}_{c}^{\xi}) - H(\mathbf{u}_{c}) - |\xi|^{2}M(\mathbf{u}_{c})$$

$$= 2\xi \cdot \int_{\mathbb{R}^{6}} \left(\operatorname{Re} u_{c} \operatorname{Im}(\nabla u_{c}) - \operatorname{Re}(\nabla u_{c}) \operatorname{Im} u_{c} \right) + \frac{1}{2} \left(\operatorname{Re} v_{c} \operatorname{Im}(\nabla v_{c}) - \operatorname{Re}(\nabla v_{c}) \operatorname{Im} v_{c} \right) dx$$

$$= 2\xi \cdot \int_{\mathbb{R}^{6}} \operatorname{Im}(\overline{u_{c}} \nabla u_{c}) + \frac{1}{2} \operatorname{Im}(\overline{v_{c}} \nabla v_{c}) dx = 2\xi \cdot P(\mathbf{u}_{c}).$$

Because $S_{I_c^{\xi}}(\mathbf{u}_c^{\xi}) = S_{I_c}(\mathbf{u}_c) = \infty$, \mathbf{u}_c^{ξ} is a blowing-up solution to (NLS system) as well. Combining with the fact that \mathbf{u}_c has the minimal kinetic energy among all the blowing-up solutions, we obtain

$$|\xi|^2 M(\mathbf{u}_c) + 2\xi \cdot P(\mathbf{u}_c) = H(\mathbf{u}_c^{\xi}) - H(\mathbf{u}_c) \ge 0$$

holds for any $\xi \in \mathbb{R}^6$. Setting $\xi = -\frac{P(\mathbf{u}_c)}{M(\mathbf{u}_c)}$, we find

$$0 \leq |\xi|^2 M(\mathbf{u}_c) + 2\xi \cdot P(\mathbf{u}_c) = -\frac{|P(\mathbf{u}_c)|^2}{M(\mathbf{u}_c)} \leq 0,$$

which implies (6.5).

Deduced from the property of negative regularity (Theorem 1.13) and zero momentum (Proposition 6.5) the movement of spatial center function x(t) of solitonlike solutions can be controlled, which can help us use Lemma 3.9 more effectively, combined with the finite mass of soliton-like solution ($\lambda(t) \ge 1$).

Corollary 6.6 (Control of spatial center function x(t), [19]). Let \mathbf{u}_c be a solitonlike solution as in Theorem 1.12. Then, for any $\eta > 0$, there exists $C(\eta) > 0$ such that

$$\sup_{t \in \mathbb{R}} \int_{|x-x(t)| \ge C(\eta)} |\mathbf{u}_c(t,x)|^2 \mathrm{d}x \lesssim \eta.$$

Furthermore, the spatial center function x(t) satisfies

 $|x(t)| = o(t), \qquad t \to \infty.$

6.2. Exclusion of soliton-like solution. Now, we use Virial identity to rule out the soliton-like solution. Its compactness has been described as an almost periodic solution. From the control of spatial center function x(t), Corollary 6.6, we know that x(t) = o(t) which means \mathbf{u}_c moves slower and slower. That implies the nonexistence of soliton-like solution for a large time.

Theorem 6.7. There exist no soliton-like solutions to (NLS system).

Proof. We argue by contradiction and suppose $\mathbf{u}_c : \mathbb{R} \times \mathbb{R}^6 \to \mathbb{C}^2$ is a soliton-like solution to (NLS system). By Definition 1.8 and $L^3(\mathbb{R}^6) \hookrightarrow \dot{H}^1(\mathbb{R}^6)$, for any $\eta > 0$, there exists $C(\eta) > 0$ such that

$$\sup_{t \in \mathbb{R}} \int_{|x-x(t)| \ge C(\eta)} \left(|\nabla \mathbf{u}_c|^2 + |\mathbf{u}_c|^3 \right) \mathrm{d}x \le \eta.$$
(6.6)

Using Corollary 6.6, we know there exists $T_0 = T_0(\eta) \in \mathbb{R}$ such that

$$|x(t)| \leqslant \eta t, \qquad \forall t \geqslant T_0. \tag{6.7}$$

Recalling Lemma 3.9, for a(x) be a radial smooth function satisfying

$$a(x) = \begin{cases} |x|^2, & |x| \le R, \\ 0, & |x| \ge 2R, \end{cases}$$

where R > 0 will be chosen later, we define

$$V_R(t) = \int_{\mathbb{R}^6} \left(2\kappa |u_c|^2 + |v_c|^2 \right) a(x) \mathrm{d}x.$$

By Theorem 1.13 and interpolation theorem, we know $\mathbf{u}_c := (u_c, v_c)^T \in \mathcal{L}^{\infty}_t(\mathbb{R}, \mathcal{L}^2_x(\mathbb{R}^6)).$ Then we get

$$|V_R'(t)| = \left| 2\kappa \operatorname{Im} \int_{\mathbb{R}^6} \left(2\overline{u_c} \nabla u_c + \overline{v_c} \nabla v_c \right) \cdot \nabla a(x) \mathrm{d}x \right| \lesssim_{\kappa} RH(\mathbf{u}_c) M(\mathbf{u}_c) \lesssim R, \quad (6.8)$$

and

$$\begin{split} V_R''(t) &= 8\kappa \operatorname{Re} \int_{\mathbb{R}^6} \left(\overline{u_{c,j}} u_{c,k} + \frac{\kappa}{2} \overline{v_{c,j}} v_{c,k} \right) a_{jk}(x) \mathrm{d}x \\ &- 2\kappa \int_{\mathbb{R}^6} \left(|u_c|^2 + \frac{\kappa}{2} |v_c|^2 \right) \Delta \Delta a(x) \mathrm{d}x - 2\kappa \operatorname{Re} \int_{\mathbb{R}^6} \overline{v_c} u_c^2 \Delta a(x) \mathrm{d}x \\ &= 8\kappa [2H(\mathbf{u}_c) - 3R(\mathbf{u}_c)] \\ &+ O\left(\int_{|x| \geqslant R} |\nabla \mathbf{u}_c|^2 + |\mathbf{u}_c|^3 \mathrm{d}x \right) + O\left(\int_{R \leqslant |x| \leqslant 2R} |\mathbf{u}_c|^3 \mathrm{d}x \right)^{\frac{2}{3}}. \end{split}$$

For $\eta > 0$ small enough, choosing

$$R := C(\eta) + \sup_{T_0 \leqslant t \leqslant T_1} |x(t)|$$

for any $T_1 > T_0$, by coercivity of energy (Proposition 3.4), energy trapping (Corollary 3.7), and (6.6), we have

$$V''(t) \gtrsim_{\kappa} E(\mathbf{u}_{c,0}). \tag{6.9}$$

Applying the Fundamental Theorem of Calculus on $[T_0, T_1]$, by (6.7), (6.8) and (6.9), we obtain

$$(T_1 - T_0)E(\mathbf{u}_{c,0}) \lesssim_{\kappa} R = C(\eta) + \sup_{T_0 \leqslant t \leqslant T_1} |x(t)| \leqslant C(\eta) + \eta T_1, \quad \forall T_1 > T_0.$$

Setting first $\eta \to 0$ and then $T_1 \to \infty$, we find $E(\mathbf{u}_{c,0}) = 0$. Using the conservation of energy and energy trapping again, we know $H(\mathbf{u}_c(t)) = 0, \forall t \in \mathbb{R}$. By (4.22), we get $H_c = 0$ which is a contradiction with $H_c \ge \delta_{sd} > 0$ in (4.18).

6.3. Exclusion of low-to-high frequency cascade. Our last enemy is the low-to-high cascade. An important ingredient is the negative regularity lemma, which indicates that u_c also falls into the L^2 space.

Theorem 6.8. There exist no low-to-high frequency cascade to (NLS system).

Proof. We argue by contradiction and suppose $\mathbf{u}_c : I_c \times \mathbb{R} \to \mathbb{C}^2$ is a low-to-high frequency cascade solution to (NLS system). Similar to the proof of Theorem 6.7, we know $\mathbf{u}_c \in \mathcal{L}^{\infty}_t(\mathbb{R}, \mathcal{L}^2_x(\mathbb{R}^6))$. Noticing the conservation of mass, we obtain

$$0 \leqslant M(\mathbf{u}_c) = M(\mathbf{u}_c(t)) := \int_{\mathbb{R}^6} |\mathbf{u}_c(t, x)|^2 \mathrm{d}x < \infty,$$

for any $t \in \mathbb{R}$.

On one hand, fixing $t \in \mathbb{R}$ and choosing $\eta > 0$ to be a small number, by Remark 1.10 and Theorem 1.11, we have

$$\int_{|\xi| \leqslant C(\eta)\lambda(t)} |\xi|^2 |\widehat{\mathbf{u}_c}(t,\xi)|^2 \mathrm{d}\xi < \eta.$$
(6.10)

At the same time, since $\mathbf{u}_c \in \mathrm{L}^{\infty}_t(\mathbb{R}, \mathcal{H}^{-\varepsilon}_x(\mathbb{R}^6))$, we know

$$\int_{|\xi| \leqslant C(\eta)\lambda(t)} |\xi|^{-2\varepsilon} |\widehat{\mathbf{u}_c}(t,\xi)|^2 \mathrm{d}\xi \lesssim 1.$$
(6.11)

Thus, interpolating (6.10) and (6.11) yields

$$\int_{|\xi| \leqslant C(\eta)\lambda(t)} |\widehat{\mathbf{u}_c}(t,\xi)|^2 \mathrm{d}\xi \lesssim \eta^{\frac{\varepsilon}{1+\varepsilon}}.$$
(6.12)

On the other hand, by Theorem 1.11,

$$\int_{|\xi|>C(\eta)\lambda(t)} |\widehat{\mathbf{u}_{c}}(t,\xi)|^{2} d\xi \leq [C(\eta)\lambda(t)]^{-2} \int_{\mathbb{R}^{6}} |\xi|^{2} |\widehat{\mathbf{u}_{c}}(t,\xi)|^{2} d\xi
\lesssim [C(\eta)\lambda(t)]^{-2} H(\mathbf{u}_{c}(t))
< [C(\eta)\lambda(t)]^{-2} H(\mathbf{W}).$$
(6.13)

Combining (6.12), (6.13), and Plancheral's theorem, we get the conclusion that

$$0 \leqslant M(\mathbf{u}_c) \lesssim \eta^{\frac{\varepsilon}{1+\varepsilon}} + [C(\eta)\lambda(t)]^{-2}, \qquad \forall \ t \in \mathbb{R}.$$

By Definition 1.12, there exists a time sequence $\{t_n\}_{n=1}^{\infty} \subset \mathbb{R}^+$ such that

$$\lim_{n \to \infty} t_n = +\infty, \text{ and } \lim_{n \to \infty} \lambda(t_n) = +\infty.$$

$$0 \leq \lim_{n \to \infty} M(\mathbf{u}_c(t_n)) \lesssim \eta^{\frac{\varepsilon}{1+\varepsilon}}.$$

Let $\eta \to 0$, we have $M(\mathbf{u}_c(t_n)) \to 0$, $n \to \infty$. Finally, by the conservation of mass, we know $\mathbf{u}_c \equiv \mathbf{0}$, which is a contradiction with (1.5) in Theorem 1.12.

References

- J. Bourgain, Global wellposedness of defocusing critical nonlinear Schrödinger equation in the radial case, J. Amer. Math. Soc., 12(1999), 145-171.
- [2] T. Cazenave, Semilinear Schrodinger Equations, Courant Lecture Notes in Mathematics, New York University, 2003.
- [3] M. Colin, Th. Colin and M. Ohta, Stability of solitary waves for a system of nonlinear Schrödinger equations with three wave interaction. Ann. Inst. H. Poincaré Anal. Non Linéaire, 26(2009), no. 6, 2211-2226.
- M. Colin, L. Di Menza and J. C. Saut, Solitons in quadratic media, Nonlinearity, 29(2016), no. 3, 1000-1035.
- [5] J. Colliander, M. Keel, G. Staffilani, H. Takaoka and T. Tao, Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in ℝ³, Ann. of Math., 167(2)(2008), no. 3, 767-865.
- [6] V. D. Dihn and L. Forcella, Blow-up results for systems of nonlinear Schrödinger equations with quadratic interaction, preprint, arXiv:2010.14595, 2020.
- [7] B. Dodson and J. Murphy, A new proof of scattering below the ground state for the 3D radial focusing cubic NLS, Proc. Amer. Math. Soc. 145(2017), no. 11, 4859-4867.
- [8] T. Duyckaerts and S. Roudenko, Threshold solutions for the focusing 3d cubic Schrödinger equation, Rev. Mat. Iberoam., 26(2010), no. 1, 1-56.
- [9] M. Hanamo, Global dynamics below the ground state for the quadratic Schrödinger system in 5d, preprint, arXiv:1805.12245, 2018.

32

,

NLS SYSTEM IN \mathbb{R}^6

- [10] M. Hanamo, T. Inui and K. Nishimura, Scattering for the quadratic nonlinear Schrödinger system in ℝ⁵ without mass-resonance condition, preprint, arXiv:1903.05880, 2019.
- [11] N. Hayashi, C. Li and P.I. Naumkin, On a system of nonlinear Schrödinger equations in 2D, Differential Integral Equations, 24(2011), no. 5/6, 417-434.
- [12] N. Hayashi, T. Ozawa and K. Tanaka, On a system of nonlinear Schrödinger equations with quadratic interaction, Ann. Inst. H. Poincaré Anal. Non Linéaire., 30(2013), no. 4, 661-690.
- [13] J. Holmer and S. Roudenko, A sharp condition of the radial 3D cubic nonlinear Schrödinger equation, Commun. Math. Phys., 282(2008), 435-467.
- [14] T. Inui, N. Kishimoto and K. Nishimura, Blow-up of the radially symmetric solutions for the quadratic nonlinear Schrödinger system without mass-resonance, Nonlinear Analysis 198(2020), 111895.
- [15] T. Inui, N. Kishimoto and K. Nishimura, Scattering for a mass critical NLS system below the ground state with and without mass-resonance condition, Discrete Contin. Dyn. Syst., 39(2019), no. 11, 6299-6353.
- [16] T. Iwabuchi, T. Ogawa and K. Uriya, Ill-posedness for a system of quadratic nonlinear Schrödinger equations in two dimensions, J. Funct. Anal., 271(2016), no. 1, 136-163.
- [17] M. Keel and T. Tao, Endpoint Strichartz estimate, Am. J. Math. 120(1998), no. 5, 955-980.
- [18] C. E. Kenig and F. Merle, Global well-posedness, scattering and blow-up for the energycritical, focusing, non-linear Schrödinger equation in the radial case, Invent. Math., 166(2006), no. 3, 645-675.
- [19] R. Killip and M. Visan, The focusing energy-critical nonlinear Schrödinger equation in dimensions five and higher, Amer. J. Math., 132(2010), no. 2, 361-424.
- [20] R. Killip and M. Visan, Nonlinear Schrödinger equations at critical regularity, Clay Mathematics Proceedings, 17(2013).
- [21] Y. S. Kivshar, A. A. Sukhorukova, E. A. Ostrovskayaa, T. J. Alexandera, O. Bang, S. M. Saltiel, C. B. Clausen and P. L. Christiansen, Multi-component optical solitary waves, Physica A: Statistical Mechanics and its Applications, 288(2000), no. 1-4, 152-173.
- [22] K. Koynov and S. Saltiel, Nonlinear phase shift via multistep (2) cascading, Optics Communications, 152(1998), no. 1, 96-100.
- [23] S. Masaki and M. Hamano, A sharp scattering threshold level for mass-subcritical nonlinear Schrödinger system, Discrete and Continuous Dynamical Systems 41(2021), no. 3, 1415-1447.
- [24] F. Meng and C. Xu, Scattering for mass-resonance nonlinear Schrödinger system in 5D, J. Differential Equations, 275(2021), 837-857.
- [25] F. Merle and L. Vega, Compactness at blow-up time for L² solutions of the critical nonlinear Schrödinger equation in 2D, Internat. Math. Res. Notices, (1998), no. 8, 399-425.
- [26] C. Miao, G. Xu, L. Zhao, The dynamics of the 3D radial NLS with the combined terms, Commun. Math. Phys., 318(2013), no. 1, 767-808.
- [27] C. Miao, J. Murphy and J. Zheng, The defocusing energy-supercritical NLS in four space dimensions, J. Functional Analysis, 267(2014), no. 2, 1662-1724.
- [28] N. Noguera and A. Pastor, On the dynamics of a quadratic Schrödinger system in dimension n = 5, Dyn. Partial Differ. Equ., 17(2020), no. 1, 1-17.
- [29] N. Noguera and A. Pastor, A system of Schrödinger equations with general quadratic-type nonlinearities, Commun. Contemp. Math., 23(2021), 2050023.
- [30] T. Ogawa and K. Uriya, Final state problem for a quadratic nonlinear Schrödinger system in two space dimensions with mass resonance , J. Differential Equations, 258(2015), no. 2, 483-503.
- [31] A. Pastor, Weak concentration and wave operator for a 3D coupled nonlinear Schrödinger system, J. Math. Phys., 56(2015), no. 2, 021507, 18 pp.
- [32] A. Pastor, On a three wave interaction Schrödinger systems with quadratic nonlinearities: Global wellposedness and standing waves, Commun. Pure Appl. Anal., 18(2019), no. 5, 2217-2242.
- [33] T. Tao and M. Visan, Stability of energy-critical nonlinear Schrödinger equations in high dimensions, Electron. J. Differential Equations, 118(2005), 1-28.
- [34] K. Uriya, Final state problem for a system of nonlinear Schrödinger equations with three wave interaction, J. Evol. Equ., 16(2016), no. 1, 173-191.
- [35] M. Visan, The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions, Duck Math. J., 138(2007), 281-374.

- [36] M. Visan, Global well-posedness and scattering for the defocusing cubic nonlinear Schrödinger equation in four dimensions, Int. Math. Res. Not. IMRN (2012), no. 5, 1037-1067.
- [37] H. Wang and Q. Yang, Scattering for the 5D quadratic NLS system without mass-resonance, Journal of Mathematical Physics, 60(2019), no. 12, 121508.

Beijing International Center for Mathematical Research, Peking University, Beijing, China, 100871

Email address: cwgao@pku.edu.cn

GRADUATE SCHOOL OF CHINA ACADEMY OF ENGINEERING PHYSICS, BEIJING, CHINA, 100088 *Email address*: mengfanfei17@gscaep.ac.cn

GRADUATE SCHOOL OF CHINA ACADEMY OF ENGINEERING PHYSICS, BEIJING, CHINA, 100088 Email address: xuchengbin190gscaep.ac.cn

Institute for Applied Physics and Computational Mathematics, Beijing, China, 100088

Email address: zhengjiqiang@gmail.com, zheng_jiqiang@iapcm.ac.cn