

DRAFT VERSION JANUARY 13, 2022
Typeset using L^AT_EX **modern** style in AASTeX62

Directly Determined Properties of HD 97658 from Interferometric Observations

TYLER G. ELLIS,¹ TABETHA BOYAJIAN,¹ KASPAR VON BRAUN,² ROXANNE LIGI,³
DENIS MOURARD,⁴ DIANA DRAGOMIR,⁵ GAIL H. SCHAEFER,⁶ AND
CHRISTOPHER D. FARRINGTON⁶

¹*Louisiana State University
202 Nicholson Hall
Baton Rouge, LA 70803*

²*Lowell Observatory
1400 W. Mars Hill Road
Flagstaff, AZ 86001, USA*

³*INAF-Osservatorio Astronomico di Brera
Via E. Bianchi 46
I-23807 Merate, Italy*

⁴*Université Côte d’Azur, Observatoire de la Côte d’Azur, CNRS, Laboratoire Lagrange, Parc
Valrose, 06108 Nice Cedex, France*

⁵*Department of Physics and Astronomy, University of New Mexico, 1919 Lomas Blvd NE,
Albuquerque, NM 87131, USA*

⁶*The CHARA Array of Georgia State University, Mount Wilson, CA 91023, USA*

(Received 6 May 2021; Revised 12 July 2021; Accepted 12 July 2021)

Submitted to ApJ

ABSTRACT

We conducted interferometric observations with the CHARA Array of transiting super-Earth host HD 97658 and measured its limb-darkened angular diameter to be $\theta_{\text{LD}} = 0.314 \pm 0.004$ mas. The combination of the angular diameter with the Gaia EDR3 parallax value with zero-point correction ($\pi = 46.412 \pm 0.022$ mas, $d = 21.546 \pm 0.011$ pc) yields a physical radius of $R_{\star} = 0.728 \pm 0.008 R_{\odot}$. We also measured the bolometric flux of the star to be $F_{\text{bol}} = 2.42 \pm 0.05 \times 10^{-8}$ erg s^{−1} cm^{−2}, which, together with angular size, allows a measurement of the effective temperature $T_{\text{eff}} = 5212 \pm 43$ K. Our directly determined physical stellar properties are in good agreement with previous estimates derived from spectroscopy. We used our measurements in combination with stellar evolutionary models and properties of the transit of HD 97658 b to determine the mass and age of HD 97658 as well as constrain the properties of the planet. Our results and our analysis of the TESS lightcurve

Corresponding author: Tyler Ellis
telli17@lsu.edu

on the planet (TOI-1821) corroborate previous studies of this system with tighter uncertainties.

Keywords: planetary systems — stars: individual (HD 97658) — stars: fundamental parameters (radii, temperatures, luminosities) — stars: late-type — techniques: interferometry — other: TOI-1821

1. INTRODUCTION

Interferometric observations of stars provide a unique opportunity to directly measure one of the most fundamental parameters of the star: its radius. Interferometry achieves the resolution of an extremely large telescope by combining the light from one or multiple pairs of telescopes across a variety of separations, or baselines. In particular, optical/near-infrared interferometry requires baselines of only tens of meters to achieve resolutions of milli-arcseconds (mas). Direct measurements of stellar radii at great precision will in turn reduce uncertainty in other derived stellar parameters (e.g. effective temperature, surface gravity, or density). Direct observations of stellar radii have highlighted a systematic discrepancy between evolutionary models and reality. [Boyajian et al. \(2012\)](#) has shown that stellar evolutionary models underestimate radii by $\sim 5\%$ and overestimate temperatures by $\sim 3\%$ for K and M dwarfs.

Observations with interferometric arrays play an important role in understanding as well as refining exoplanet system properties. In order to understand the properties of the exoplanet, the properties of the star must first be well constrained. In particular, transiting exoplanets provide a measure of the planet’s radius, but this measurement is in units of the host star’s radius. Any uncertainty or bias in stellar radius will propagate into estimates of the planet’s equilibrium temperature, density, habitability, and composition. Interferometry gives a direct measurement of the stellar radius with little or no dependence on stellar models. This technique has been used in the literature to refine the properties of several important systems, such as 55 Cancri which hosts five radial velocity exoplanets including another transiting super-Earth ([von Braun et al. 2011](#)) and transiting exoplanet host star GJ 436 where evolutionary models underestimated the stellar radius by $\sim 11\%$ ([von Braun et al. 2012](#)). There have also been multiple interferometric surveys of large numbers of exoplanet host stars such as [Baines et al. \(2008\)](#). The field of interferometry has also seen incredible developments in the field of imaging and astrometry with the ESO GRAVITY project ([Gravity Collaboration et al. 2017](#)). In 2019, the GRAVITY collaboration announced the first spectrum of an exoplanet observed with interferometry and refined the astrometric position with $100\mu\text{as}$ precision ([Gravity Collaboration et al. 2019](#)).

The exoplanet host star HD 97658 is of particular interest in this regard as it is the home of a transiting super-Earth ([Dragomir et al. 2013](#); [Howard et al. 2011](#)). HD 97658 is a bright, $m_V=7.78$ mag K1 dwarf with a moderately low iron content of $[\text{Fe}/\text{H}]=-0.23$ dex, which was discovered to have a Neptunian mass exoplanet by the

NASA Eta-Earth Keck-HIRES radial velocity survey (Howard et al. 2011). Follow-up time series observations with the *Spitzer* and *MOST* space telescopes detected a transit whose depth indicated an estimated planetary radius of a few Earth radii (Dragomir et al. 2013; van Grootel et al. 2014). These properties together make HD 97658 b a so-called super-Earth (planets with radii of 1–4 R_{\oplus} and masses of 1–10 M_{\oplus} ; Bryan et al. 2019). Super-Earths can take the form of water worlds with a smothering dense atmosphere or rocky behemoths with minimal atmospheres, both often consistent within the uncertainties of planetary mass and radius (Dragomir et al. 2013). Super-Earths captivate planetary scientists as they are the most populous of observed exoplanets (30–50% of Sun-like stars host one or more super-Earths (Bryan et al. 2019)), however they do not exist within our own solar system and must be studied solely as exoplanets.

We are interested in refining the properties of HD 97658 b by directly measuring the host star’s properties. In §2, we describe the interferometric observations of HD 97658. In §3, we report the resulting directly measured angular diameter, bolometric flux. In §4, we model and measure the mass of the star. We then derive updated properties of HD 97658 and its planet using the TESS lightcurve and our measured results. Lastly, in §5 we summarize and conclude this work.

2. INTERFEROMETRIC OBSERVATIONS

We observed HD 97658 over the course of several nights using the Georgia State University Center for High Angular Resolution Astronomy (CHARA) Array at the Mount Wilson Observatory using the Classic (near-IR), VEGA (optical), and PAVO (optical) beam combiners (ten Brummelaar et al. 2005; Mourard et al. 2009; Ireland et al. 2008). A summary of the observations is found in Table 1.

An interferometer measures visibilities (V), which quantifies the contrast of the dark and bright parts of the interference fringe pattern. In practice, this is the contrast of the time-averaged minimum and maximum power of the fringe pattern (see Lawson et al. (2000) for a full description of what an interferometer measures in Chapter 2.6).

As the visibilities measured at the time of observing include instrumental and atmospheric effects, it is necessary to observe stars with predictable visibilities to calibrate the data. These calibrator stars are observed in sequence with a science star and allow a measure of the combined systematic effects. Calibrator stars are of a known, and ideally unresolved, size.

Each observation consists of bracketed sequences of the form calibrator 1 - science target - calibrator 2 (or calibrator 1 again), and then reverse. One bracket is one observation of the target sandwiched by the calibrators.

As the beam combiners used in this work observe in different bandpasses, it is sometimes necessary to use different calibrators for each instrument. Wherever possible, we calibrated the raw square visibilities of the calibrators against each other (von Braun & Boyajian 2017). This provides insight into previously unknown duplicity,

activity, or other anomalous behaviors in a calibrator. No stars were expunged from our list of acceptable calibrators for any reason. The limb darkened angular size of the calibrators and magnitudes are summarized in Table 2.

We chose calibrators for this work using the JMMC Stellar Diameter Catalog (JSDC) version 2 (Chelli et al. 2016; Duvert 2016)¹. Ideally, we restrict our search for calibrators that are not resolved on the baselines used with beam combiner’s band-pass. Unresolved sources have predicted squared visibilities $V^2 \gtrsim 0.9$. Further, we discard potential calibrators that have known duplicity and/or have rapid rotation driving equatorial distention. Lastly, preference is given to calibrators of comparable brightness and with a minimal angular separation in the sky from the science target.

Calibrator selection is a nontrivial process which can affect the uncertainties in final calibrated data. van Belle & van Belle (2005) demonstrated that a 5% uncertainty in the calibrator diameter can propagate up to a $\sigma_{V^2} \sim 0.04$ uncertainty in the calibrated visibilities, though the amount depends on interferometer configuration, instrument, and calibrator size. Oftentimes, for small, faint targets such as HD 97658, most calibrators which are sufficiently smaller so as to be unresolved are often too faint to be observed. As such, it is sometimes necessary to compromise for partially resolved calibrators. The caveat with such a compromise is that the target’s diameter will only be known as well as the calibrators. In this work, the calibrators HD 101688 and HD 96738 are slightly resolved using the PAVO/VEGA beam combiner and S1-W1 baseline, $V^2 \simeq 0.6$. Thus, the predicted angular size of these calibrators are of concern to the error budget. However, within this size regime, $\theta < 0.45$ mas, the dominant source of error is in the measurement of the visibilities rather than assumptions of calibrator size (van Belle & van Belle 2005).

We used the *isoclassify*² program to estimate the angular diameters of these calibrators independently of the JMMC’s surface brightness relationship method (Huber et al. 2017; Berger et al. 2020). We find the *isoclassify* and JMMC diameters to be consistent, though the adopted *isoclassify* diameters had larger, more conservative uncertainties (an average of $1.9\times$ larger). All of differences in size are less than 1σ , with the average statistical tension $Avg\left((|\theta_{\text{iso}} - \theta_{\text{JMMC}}|)/\sqrt{\sigma_{\text{iso}}^2 + \sigma_{\text{JMMC}}^2}\right) = 0.33$.

As previously mentioned, the greater uncertainty in the calibrated visibilities from using resolved calibrators is propagated forward into the uncertainty of the fitted angular diameter. We also note that data taken with the VEGA on several nights are calibrated with only a single star. However, our analysis finds consistency in the calibrated results from night to night and between PAVO and VEGA, assuring confidence in our choice of calibrators and calibration methods (Ligi, R. et al. 2016; Baines et al. 2018; Lachaume et al. 2019).

In order to complete an observation, the interferometric fringes must be found by equalizing the optical path length from the stars to the beam combiner through the

¹ http://www.jmmc.fr/catalogue_jsdc.htm

² <https://github.com/danxhuber/isoclassify>

two telescopes. The fringes are found by scanning the additional path length up and down for one of the telescopes until fringes are detected. In good conditions, scanning initially takes about 5–15 minutes per star and about twice that in difficult observational conditions. Finding the fringe packet for subsequent observations of the same star typically goes much faster, at most a few minutes.

Observations with the Classic beam combiner consist of approximately 2.5 minutes of integration in the H band. Shutter sequences proceed and follow integration. Classic data are reduced using the REDFLUOR package to produce raw squared visibilities which are then calibrated using the CALIBIR package — both of these software routines are provided as binary executables from CHARA³. Classic observes in a single spectral channel at a time and gathers a single data point per bracket.

Observations with the PAVO beam combiner consist of around 2 minutes integration followed by approximately 3 minutes of shutter sequences and dark integration for reduction and calibration. PAVO data are reduced and calibrated using IDL routines which are also provided by CHARA (Ireland et al. 2008)⁴. As PAVO is an integral field unit, it collects a spectra of fringes with each bracket between 630–950 nm bandwidth (resolution $R = 30$) (Ireland et al. 2008).

Observations with VEGA require more time. Calibrators are observed during ~ 15 min while the science star is generally observed during ~ 30 min to ensure enough signal. The data are then reduced using the *vegadrs* pipeline (Mourard et al. 2009, 2011). VEGA’s bandpass is broken into 20 nm bins and creates two data points per baseline at a time.

Date [UT]	Baseline	Combiner	Brackets	Calibrators
2015-02-04	S1/W1	Classic	5	HD 93152, HD 99267
2015-02-05	S1/W1	Classic	7	HD 95804, HD 99267
2015-02-11	S1/W1	PAVO	4	HD 101688, HD 96738
2017-03-13	W1/W2	VEGA	2	HD 89239
2017-03-17	S2/W2	VEGA	2	HD 107168
2018-04-28	E2/W2	VEGA	2	HD 97638
2019-05-05	S1/W2	VEGA	3	HD 96738
2020-03-05	S2/E2	VEGA	4	HD 96738, HD 107168

Table 1. Summary of observations of HD 97658. Both PAVO and Classic instruments were used in the single baseline mode of the CHARA array, while VEGA uses multiple baselines simultaneously. See ten Brummelaar et al. (2005, Table 1) for a complete description of the available baselines. Each bracket corresponds to one observation of the target. See § 2 for further details

3. DIRECTLY DETERMINED STELLAR PROPERTIES

3.1. *Stellar Diameter*

³ http://www.astro.gsu.edu/~theo/chara_reduction/index.html

⁴ <https://gitlab.chara.gsu.edu/fabien/pavo.git>

Calibrator	V mag	H mag	θ_{LD} [mas]
HD 89239	6.530	6.599	0.159±0.008
HD 93152	5.285	5.384	0.279±0.012
HD 95804	6.766	6.288	0.208±0.005
HD 96738	5.593	5.442	0.269±0.033
HD 99267	6.606	6.091	0.241±0.006
HD 101688	6.291	5.730	0.281±0.011
HD 107168	6.220	5.969	0.241±0.011

Table 2. Summary of calibrator stars used in all observing campaigns. All photometry data are taken from the JMMC Stellar Diameters Catalog v2. The cited angular diameters are limb darkened and estimated using *isoclassify*.

The calibrated squared visibilities V^2 can be fit to the radial profile of the 2D Fourier pair of a uniformly illuminated disk or a limb darkened disk, θ_{UD} and θ_{LD} , respectively. The resulting profile is a function of the projected baseline B , observational wavelength λ , and most importantly the angular size of the object (Hanbury Brown et al. 1974). The functional form, shown in Eq. 1 is a combination of Bessel functions of the first kind, $J_\alpha(x)$, scaled with the linear limb darkening coefficient μ .

$$V^2 = \left[\left(\frac{1-\mu}{2} + \frac{\mu}{3} \right)^{-1} \cdot \left((1-\mu) \cdot \frac{J_1(x)}{x} + \mu \sqrt{\frac{\pi}{2}} \cdot \frac{J_{3/2}(x)}{x^{3/2}} \right) \right]^2 \quad (1)$$

where $x = \frac{\pi B \theta}{\lambda}$.

We fit the uniform disk model, Eq. 1 with $\mu = 0$, for the combined datasets using the *Scipy curve_fit* nonlinear least-squares minimization routine (Jones et al. 2001). As part of the fitting process, we find the optimized fit for many realizations of the dataset by sampling the wavelength solution uncertainty. The resulting distribution of fits for the uniform disk is $\theta_{UD} = 0.296 \pm 0.004$ mas. We performed the fitting routine for the VEGA and PAVO datasets separately as a check for consistency and found the best fit uniform disk angular diameters of $\theta_{UD,VEGA} = 0.282 \pm 0.024$ mas and $\theta_{UD,PAVO} = 0.296 \pm 0.004$ mas. As seen in Fig. 1, there is more internal spread amongst the VEGA data than the Classic or PAVO data, which drives the uncertainty up slightly. The enlarged uncertainties are likely caused by dividing the starlight of this faint target amongst the multiple simultaneous baselines of VEGA. We opt not to perform this same fit with the Classic data as HD 97658 is not well resolved by the instrument, but can still act as a sanity check for the other instruments. The uniform disk angular diameter and other stellar properties are summarized in Table 3.

We fit the limb darkened disk model using the same technique as above, adding in Monte Carlo realizations of the limb darkening parameter μ as part of the fitting process. As all visibility curves are near unity at low spatial frequencies, it is safe to combine the Classic data with the VEGA/PAVO data. We estimated the limb darkening parameter μ using the Limb Darkening Toolkit (LDTK) (Parviainen & Aigrain 2015; Husser et al. 2013). Throughout this work we find the limb darkening

coefficient and associated limb darkened diameter in the Bessel R filter. We provide the LDTK module with estimates of T_{eff} , $\log(g) = 4.5 \pm 0.1$, $Z = 0.03 \pm 0.01$. We iterate the fitting process twice to reflect our refined measurements in T_{eff} which goes into the estimation of the limb darkening coefficient. The first run uses the fit of θ_{UD} with our measurement of the bolometric flux to estimate T_{eff} , as is discussed in the following section § 3.2. Then we use the results from the first fit of θ_{LD} to estimate T_{eff} and find our new estimation of μ . With the final estimation of μ , we then find our final fit of θ_{LD} . We scale the uncertainties in μ by a factor of 5 during each iteration of the fitting process as the LDTK distribution seems unrealistically tight when compared to Claret & Bloemen (2011), though this does not significantly contribute to the final error budget. Our final estimation of the linear limb darkening coefficient in the Bessel R filter is $\mu_{\text{R}} = 0.629 \pm 0.014$.

The iteration process resulted in a best fit limb darkened angular diameter of $\theta_{\text{LD}} = 0.314 \pm 0.004$ mas. The reduced chi-square for the linear limb darkened model is $\chi^2_{\nu=113} = 0.934$ ($p = 0.68$), indicating good agreement between theory and observation. This fit and the calibrated squared visibilities are shown in Figure 1 with the uncertainties scaled to fix χ^2_{ν} at unity.

We calculate the physical radius of HD 97658 as $0.728 \pm 0.008 R_{\odot}$ using the Gaia EDR3, zero-point corrected parallax measurement (Table 3). Applying the zero-point correction, the Gaia parallax of 46.412 ± 0.022 mas yields a corresponding distance of $d = 21.546 \pm 0.011$ pc (Gaia Collaboration et al. 2021; Lindegren et al. 2021). Previously published radius estimates include Dragomir et al. (2013) who obtained an estimate of $0.703^{+0.035}_{-0.030} R_{\odot}$ from evolutionary models fit within EXOFAST (Eastman et al. (2013)) and van Grootel et al. (2014) who derived $0.741^{+0.024}_{-0.023} R_{\odot}$ from the spectroscopic temperature, bolometric correction, and Hipparcos parallax. It is of interest that the estimation of the radius using the evolutionary models in EXOFAST is lower than the spectroscopic radius and the directly measured radius in this paper. As has been explored in Boyajian et al. (2012), evolutionary models systematically underestimate the radius by a few percent. The high resolution capabilities of the interferometer complemented with the exquisite parallax measurements from Gaia allow us to report a physical radius with about 1/3 the uncertainty of prior works. The uncertainty in our physical radius predominantly comes from the fit of the angular diameter, with the parallax contribution essentially negligible.

3.2. Bolometric Flux and Temperature

We fit an interpolated K0.5 Pickles (Pickles 1998) template spectrum to collected literature measurements of broadband photometry to measure the bolometric flux. Photometry used in this fit include measurements in Johnson *UBV*, Cousins *R_c*, *I_c*, 2Mass *JHK*. The photometric measurements are from van Leeuwen (2007); Skrutskie et al. (2006); Kotoneva et al. (2002); Koen et al. (2010); Bessell (2000); Kharchenko (2001); Droege et al. (2006); Mermilliod (1994).




Figure 1. Top panel shows the calibrated squared visibilities and uncertainties from CHARA observations of HD 97658. We binned and averaged the data by equal spacing in of 10^7 rad^{-1} for clarity, and plot these points over all of the calibrated data shown in transparent grey. Classic observations are shown as red circles grouped on the left at lower spatial frequencies, VEGA observations are green diamonds in the middle, and PAVO observations are blue squares grouped on the right at higher spatial frequencies. The fitted visibility curve for a limb darkened disk is shown as a dashed line with the parameters found in §3.1. We show the uncertainty in the angular diameter as the grey region around the best fit curve. The residuals for the fit are shown in the bottom panel. See §3.1 for details.

Interstellar extinction was fixed at 0 in the fitting routine as the proximity of the star should render any extinction effects negligible. The fit was also performed with A_V as a free parameter. This modification found a value of $A_V = 0.027 \pm 0.015$, so we accept our original assumption of no extinction for this work.

The template spectra was then scaled to fit to the photometric measurements and integrated to obtain a bolometric flux $F_{\text{bol}} = 2.42 \pm 0.05 \times 10^{-8} \text{ erg s}^{-1} \text{ cm}^{-2}$. We use the updated filter profiles and zero-point corrections as discussed in Mann & von Braun (2015). Further, we account for unknown systematics by applying a 2% addition in quadrature to the bolometric flux uncertainty as is suggested in Bohlin et al. (2014). Our flux measurement and the parallax distance gives a stellar luminosity

of $L = 0.351 \pm 0.007 L_{\odot}$. The assembled photometry and spectral fit are shown in Figure 2. This model has a goodness of fit of $\chi^2_{\nu=29} = 0.79$ ($p = 0.78$). Further details of this technique are available in [van Belle et al. \(2007\)](#); [von Braun & Boyajian \(2017\)](#). We extended the infrared portion of the Spectral Energy Distribution (SED) ($\lambda > 12.5 \mu\text{m}$) with the WISE W 1–4 data to check for an infrared excess, but did not observe any such excess ([Wright et al. 2010](#)). We note that HD 97658 is saturated in W1 and W2, so we opted to exclude all of these data from the SED fit.




Figure 2. The template spectrum shown in blue is an interpolated Pickles K0.5V stellar spectrum with nominal metallicity. We used literature broadband photometry and associated uncertainties and bandpass widths to scale the spectrum. The photometry is shown as the black points. The horizontal bars on the photometry data points represent the width of the filter bandpass. This scaled spectrum was then integrated to yield the bolometric flux. We calculate the scaled model flux values at the center of the bandpasses and show them as red crosses here. The bottom panel shows the percent residual of this fit. See § 3.2 for discussion.

The Stefan-Boltzmann equation can be rewritten to express the temperature in terms of the observables — bolometric flux and angular diameter:

$$T_{\text{eff}} = 2341 \cdot \left(\frac{F_{\text{bol}}}{\theta_{\text{LD}}^2} \right)^{1/4} \text{ K}, \quad (2)$$

where the bolometric flux is in units of $10^{-8} \text{ erg s}^{-1} \text{ cm}^{-2}$ and the angular diameter is in mas. We determined an effective temperature of $T_{\text{eff}} = 5212 \pm 43 \text{ K}$ given the limb darkened diameter and above bolometric flux. Howard et al. (2011) measured the effective temperature of HD 97658 as $5170 \pm 44 \text{ K}$ which was determined from spectroscopy. This temperature is lower than what we found, but consistent with our result at the 1σ level.

4. DERIVED STELLAR AND PLANETARY PROPERTIES

Using the measured properties above, we compute here other properties of the star and its planet. As the host star plays an important role in determining many of the properties of the planet, we can also update several of the estimated properties of the planet with our improved stellar parameters. We also include new observations of the planet’s transit with TESS. A summary of all the following results is shown in Table 3.

4.1. Age and Mass Estimation with Isochrones

We use two stellar evolutionary models to estimate the stellar age and mass. The Garstec and YaPSI stellar evolutionary models (Spada et al. 2017; Weiss & Schlattl 2008) were fit using the **bagemass** Fortran program (Maxted et al. 2015). These models use our directly measured temperature, the luminosity inferred from the parallax and bolometric flux, the spectroscopically determined $[\text{Fe}/\text{H}]$ from Howard et al. (2011), and the density inferred from the transit discussed later in this section as predictors of goodness of fit. We assumed uniform priors on age, mass, and surface $[\text{Fe}/\text{H}]$. One of the issues with estimating age using isochrones for this range of stellar masses is a lower mass star afforded a longer time to evolve can have the same observable properties as a higher mass star that is younger. This bias is reflected as the elongated distribution in age and mass. We show the posterior distribution for the YaPSI and Garstec models in Figure 3.

We then checked the consistency of the two model’s predicted age and mass using a χ^2 -test. We first combined (i.e. summed) the two posterior distributions for the ages/masses and computed the resultant median age and mass. We then compute:

$$\chi^2 = \sum_{i=1}^2 \frac{x_i - \bar{x}}{\sigma_i^2} \quad (3)$$

where x_i is the median mass/age from the two models, \bar{x} is the median from the combined distribution, and σ is their associated uncertainties. Assuming that these

are drawn from a χ^2 distribution, the confidence in the agreement of the two posterior distributions can be computed with:

$$p = \int_{\chi^2}^{\infty} f(\chi^2) d\chi^2 \quad (4)$$

The χ^2 distribution, $f(\chi^2)$, for this case has $2 - 1 = 1$ degrees of freedom, for the two samples of the age/mass. The computed p-value for the ages is $p=0.62$ and $p=0.27$ for the masses. Both the mass and age distributions agree within 95% confidence, so we conclude that the median and 1σ confidence interval of the combined distribution is representative of both. The combined isochrone models estimated the age of HD 97658 as $3.9^{+2.6}_{-2.03}$ Gyr and the mass as $0.773^{+0.015}_{-0.018} M_{\odot}$.

HD 97658 has a chromospheric Ca II H and K activity index of $\log(R'_{HK}) = -4.971$ (Isaacson & Fischer 2010). Isaacson & Fischer (2010) then used the relationship from Mamajek & Hillenbrand (2008, Eq. 3) to estimate an age of 6.06 ± 0.91 Gyr⁵. Using the Gaia $G_{BP} - G_{RP} = 0.843$ and the rotation period of 34 ± 2 days from Guo et al. (2020), we find another estimate of the age of HD 97658 of 6.25 ± 0.56 Gyr using gyrochronology with stardate (Angus et al. 2019). Both of these techniques agree within $\sim 1\sigma$ of the combined isochrone model age from this work.




Figure 3. Posterior distribution of age and mass from the *bagemass* Bayesian evolutionary track fitting program as discussed in § 4. The YaPSI model output is shown on the left and the Garstec model on right. Higher posterior density is shown as the darker regions in the bottom left plot. The posterior population distribution for the ages is shown on the top and the posterior population distribution for the mass is shown on the right. Black dashed vertical/horizontal lines indicate the median for the age (3.2 ± 1.9 Gyr, 4.7 ± 2.5 Gyr) and mass ($0.782 \pm 0.011 M_{\odot}$, $0.763 \pm 0.011 M_{\odot}$) for the YaPSI and Garstec models, respectively. We combine the solutions to derive a final estimated age $3.9^{+2.6}_{-2.03}$ Gyr and mass $0.773^{+0.015}_{-0.018} M_{\odot}$ as described in § 4.

⁵ In order to estimate uncertainty on this age, we applied the suggested RMS of 0.07 dex on $\log \tau$ from Mamajek & Hillenbrand (2008)

4.2. Exoplanet Modeling with TESS

The *TESS* mission (Ricker et al. 2015) observed HD 97658 during Sector 22 for a total duration of approximately 23 days (the full Sector duration of ~ 27 days, minus ~ 4 day gap from d13 to d17). The normalized PDCSAP lightcurve is presented in Figure 4. We first use the short cadence (2 minute) mission processed PDCSAP time series data to look for signs of rotation due to starspots rotating in and out of view on the surface of the star. We find that the average brightness of HD 97658 is stable with a RMS (root mean square) of 385 ppm, having no evidence of long-term variability due to spots during this observation period. This conclusion is consistent with *TESS* observation period being shorter than the derived rotation period $P_{\text{rot}} = 34 \pm 2$ d from the spectroscopic analysis of the Calcium H and K lines (S_{HK}) (Guo et al. 2020).




Figure 4. The TESS PSCSAP 2 min lightcurve binned into 10 minute bins. The known planet HD 97658 b has transits at days 1905 and 1924, as well as a third transit during the gap.

Next, we search for transits in the *TESS* data by computing a Box Least-Squares (BLS) periodogram (Kovács et al. 2002), shown in Figure 5, using the *astropy* package⁶. We identify the known transiting planet HD 97658 b, with an approximate period of 9.474 d. We note that only two transits of HD 97658 b are detected in *TESS* data, while the third falls within the data gap. We use EXOFASTv2 (Eastman 2017) to simultaneously fit the orbital parameters to the *TESS* time series along with the full radial velocity (RV) data series from Dragomir et al. (2013). The program also simultaneously fit the MIST evolutionary models to estimate stellar properties (Choi et al. 2016).

EXOFASTv2 reported a transit depth of HD 97658 b of 712 ± 38 ppm with a duration of 2.80 ± 0.04 hours, centered at BJD 2458904.9366 ± 0.0008 . With our measurement of the stellar radius and the transit depth, we compute a planet radius of $2.12 \pm 0.06 R_{\oplus}$.

⁶ <https://docs.astropy.org/en/stable/timeseries/bls.html>




Figure 5. The Box Least-Squares periodogram of the TESS lightcurve of HD 97658. The strongest peak coincides with the known planet HD 97658 b with a period of ~ 9.5 days. Most of the other peaks are harmonics of HD 97658 b, but there is a notable spike at 1.05 days which is explored as a potential planet candidate. There is a peak at ~ 3.8 days which proved insignificant upon further inspection. The other peaks are harmonics of the 9.5 and 1.05 day signals.

The resultant temperature of HD 97658 b can then be found as $T_{\text{eq}} = T_{\star} \sqrt{R_{\star}/2a}$, neglecting albedo. We find the equilibrium temperature for planet b of 750 ± 13 K. Our estimation of the equilibrium temperature is in line with the estimate from [van Grootel et al. \(2014\)](#). Using the planetary mass found with EXOFASTv2, $7.5 \pm 0.9 M_{\oplus}$, and the planetary radius derived from our direct measurement of the stellar radius and the TESS transit depth, we determine a density of $\rho_p = 3.7 \pm 0.5 \text{ g cm}^{-3}$. The density is consistent within 1σ of [Dragomir et al. \(2013\)](#) and [van Grootel et al. \(2014\)](#). This fitting of the RV and transit data yielded a semi-amplitude of $2.8 \pm 0.3 \text{ m/s}$, which is also within $1-\sigma$ of [Dragomir et al. \(2013\)](#). The low amplitude RV signal largely limits the uncertainty in the measurements of planet b’s density and any improvements are due to the decreased uncertainty in the transit depth and stellar radius.

Finally, we use the *TESS* time series to look for any additional transiting planets. Interestingly enough, the BLS periodogram analysis comes up with a signal at 1.054 d. We use EXOFASTv2 to model this candidate signal as an additional planet in the system at the same time as planet b. If such a planet candidate exists, EXOFASTv2 indicates it would have a period of $1.05443179^{+0.00000011}_{-0.00000018}$ days and cause a transit depth of $88 \pm 17 \text{ ppm}$ lasting 1.36 hours. The T_0 for the model was found as BJD 2458907.1 ± 0.3 . Given this depth and our measurement of the stellar radius, the planet candidate would have a radius of $0.74 R_{\oplus}$. The planet would be located at 0.019 au with an equilibrium temperature of 1565 K, found using the same method as for planet b. The transit model overlaps with the known planet transit and reduces the depth for planet b to 674 ± 38 and the corresponding radius to $2.06 \pm 0.06 R_{\oplus}$. The folded lightcurve for this planet candidate is shown in Figure 6 right panel. The

10 minute binned out of transit photometry has a root mean square deviation of 39 ppm, about half of the transit depth.




Figure 6. All of the TESS photometry is shown as transparent grey points. The folded and 10 minute binned TESS photometry with associated errors are shown as blue points. The transit models from EXOFASTv2 are the dashed black line. **Left:** The best fit model for planet b with period 9.4896 days. **Right:** The EXOFASTv2 transit model for the planet candidate “c” identified in Figure 5 with period 1.054 days is shown as the blue dashed line. See § 4.2 for details. Note that the depth is approximately an order of magnitude less than the left plot.

In attempts to verify this signal with existing data, we searched the RV data for a signal with a period of 1.05 days, but did not find evidence to support it. However, we note that a transiting planet with this period and transit depth would not be massive enough to induce reflex motions detectable with current RV instruments/observations. We compared the Bayesian Information Criterion (BIC), a measure of the goodness of the fit that penalizes over-parameterization, of the single planet model to the two planet model (Schwarz 1978). We find there is a small preference to the two planet model $\Delta BIC \simeq -8$. However, we still present this signal as only a candidate due to the low strength of the transit signal and lack of RV corroboration.

While beyond the scope of this paper, we propose that further investigation of this candidate would benefit from including archival time series over much longer time baselines than the TESS Sector 22 data presented here. In particular, enabling transit timing variation analysis in the orbital fit for planet b in the model could provide additional independent evidence for the putative companion “c”.

4.3. Star and Planet Properties from Transit Observables

Seager & Mallén-Ornelas (2003) demonstrated that combining the transit depth ΔF , duration τ and period P derived from the exoplanet transit lightcurve analysis yield the stellar density. Thus, with a direct determination of the stellar radius through interferometry, we can then directly obtain the stellar mass. This method

has been applied to 55 Cnc (Crida et al. 2018b,a) and HD 219134 (Ligi et al. 2019), for which the joint likelihood of the stellar mass and radius ($\mathcal{L}_{MR_\star} = \mathcal{L}_{MR_\star}(\rho_\star, R_\star)$) is expressed through the probability density function (PDF) of the density and radius (see Eq. (2) of Ligi et al. 2019). The PDF of the radius is itself expressed as a function of the PDF of the observables θ_{LD} (angular diameter) and d (distance), considered as Gaussian.

This yields the PDF of the planetary mass and radius, that depends on ΔF , P , the semi-amplitude of the RV measurement K , and \mathcal{L}_{MR_\star} . Importantly, this method also allows computation of the correlation between parameters. This prevents, for example, determining absurd planetary densities that would not correspond to a realistic planetary mass.

Using this technique, and considering only planet b , we obtain $\rho_\star = 3.1 \pm 0.3 \text{ g cm}^{-3}$ which yields $M_\star = 0.85 \pm 0.08 \text{ M}_\odot$, with a correlation of $\text{Corr}(R_\star, M_\star) = 0.41$. This low correlation is due to the high uncertainty on the stellar density. This direct determination of the mass is higher but consistent with those obtained with the different stellar evolutionary models.

Applying the stellar mass derived from the transit model, the interferometrically derived stellar radius, transit model, and the RV semi-amplitude we find the planetary mass $M_p = 8.3 \pm 1.1 \text{ M}_\oplus$ and radius $R_p = 2.12 \pm 0.06 \text{ R}_\oplus$, with the corresponding density $\rho_p = 4.8 \pm 0.7 \text{ g cm}^{-3}$. These measurements are in good agreement with those found from the EXOFASTv2 analysis. We note that the correlation between the planet's mass and radius is very low ($\text{Corr}(R_p, M_p) = 0.09$). This is explained by $\text{Corr}(R_\star, M_\star)$, which is already low, and the high uncertainty on the transit and RV measurements parameters. Observations with higher precision are needed to reduce these uncertainties and increase the correlation between the parameters.

5. CONCLUSION

In this work we use the GSU CHARA interferometric array to obtain a highly precise measurement of the angular diameter of HD 97658. We combine measurements from both the infrared Classic instrument as well as the optical PAVO and VEGA instruments for more complete coverage of the UV plane, which helps ensure a well defined angular diameter. We also combine photometric measurements from a panoply of sources to find bolometric flux with SED fitting. These two measurements allow an estimation of temperature independent of distance measurements.

We provide the most direct measurement of the star's radius which paired with the Gaia parallax produce a $\sim 1\%$ uncertainty in the physical radius and a $\sim 0.5\%$ uncertainty in effective temperature. Previous works exploring the properties of HD 97658 were able to obtain estimates of the stellar radius and temperatures which are in good agreement with the measurements performed in this work. Because of this we cannot report substantially different composition and properties of HD 97658 b, but we can provide greater certainty in the previous results.

Table 3. Summarized Properties of the HD 97658 System

Property	Value	Source
Measured Stellar Properties		
Parallax [mas]	46.412 ± 0.022	Gaia Collaboration et al. (2021); Lindegren et al. (2021)
Distance [pc]	21.546 ± 0.011	Gaia Collaboration et al. (2021); Lindegren et al. (2021)
[Fe/H] [dex]	-0.23 ± 0.03	Howard et al. (2011)
$\theta_{\text{UD-R}}$ [mas]	0.296 ± 0.004	§3.1 Interferometry
θ_{LD} [mas]	0.314 ± 0.004	§3.1 Interferometry
Linear Limb Darkening μ_{R}	0.629 ± 0.014	§3.1 Parviainen & Aigrain (2015); Husser et al. (2013)
R_{\star} [R_{\odot}]	0.728 ± 0.008	§3.1 Interferometry, Parallax
F_{Bol} [erg s $^{-1}$ cm $^{-2}$]	$2.42 \pm 0.05 \times 10^{-8}$	§3.2 SED Templates
T_{eff} [K]	5212 ± 43	§3.2 Interferometry, SED
L_{\star} [L_{\odot}]	0.351 ± 0.007	§3.2 F_{Bol} , Parallax
Isochrone Properties — § 4.1		
Age [Gyr]	$3.9^{+2.6}_{-2.03}$	Combined Isochrone Models
M_{\star} [M_{\odot}] ^a	$0.773^{+0.015}_{-0.018}$	Combined Isochrone Models
EXOFASTv2 Model Derived Properties — § 4.2		
Transit Depth [ppm]	712 ± 38	EXOFASTv2
Period [days]	9.4897116 ± 0.0000008	EXOFASTv2
T_0 [BJD]	2458904.9366 ± 0.0008	EXOFASTv2
R_{p}/R_{\star}	0.0267 ± 0.0007	EXOFASTv2
Inclination [deg]	$89.05^{+0.41}_{-0.24}$	EXOFASTv2
Impact Parameter	$0.39^{+0.11}_{-0.18}$	EXOFASTv2
Eccentricity	$0.05^{+0.04}_{-0.03}$	EXOFASTv2
M_{\star} [M_{\odot}] ^a	0.75 ± 0.02	EXOFASTv2, MIST
M_{p} [M_{\oplus}] ^a	7.5 ± 0.9	EXOFASTv2, MIST, K
RV Semi-Amplitude K [m/s]	2.8 ± 0.3	EXOFASTv2, RVs from (Dragomir et al. 2013)
a/R_{\star}	24.2 ± 0.7	EXOFASTv2
R_{p} [R_{\oplus}]	2.12 ± 0.06	Transit Depth, Interferometric R_{\star}
ρ_{p} [g cm $^{-3}$]	3.7 ± 0.5	Transit Derived R_{p} , M_{p}
T_{Eq} [K]	751 ± 12	EXOFASTv2, a/R_{\star} , T_{eff}
Stellar and Planetary Properties from Transit Observables — § 4.3		
ρ_{\star} [g cm $^{-3}$]	3.1 ± 0.3	Transit Observed Properties
M_{\star} [M_{\odot}] ^a	0.85 ± 0.08	Interferometric R_{\star} , Transit Derived ρ_{\star}
$\log(g)$ [cgs]	4.64 ± 0.04	Interferometric R_{\star} , Transit Derived ρ_{\star}
$\text{Corr}(R_{\star}, M_{\star})$	0.41	
M_{p} [M_{\oplus}] ^a	8.3 ± 1.1	Transit Derived $M_{\star}(\rho_{\star}, R_{\star})$ and K
ρ_{p} [g cm $^{-3}$]	4.8 ± 0.7	Transit Derived M_{p} , R_{p}
$\text{Corr}(R_{\text{p}}, M_{\text{p}})$	0.09	

^aThe table reflects the computed mass of the star and planet with two different methods. See § 4 for more details.

Follow-up observations of HD 97658 with JWST will allow a more precise and accurate measurement of the transit depth. This is a particularly interesting measurement to pin down as current best measurements of the transit depth are accurate to only $\sim 5\%$, which complicates more accurate analysis of the planet. These follow up observations would provide further exciting insight into this nearby super-Earth planet. We also eagerly await the Magdalena Ridge Observatory interferometer which will enable observations of fainter targets and baseline bootstrapping which will ease optimal sampling of the UV plane (Creech-Eakman et al. 2018).

ACKNOWLEDGEMENTS

We offer our sincere appreciation to the observing team, scientists, and support staff at the CHARA Array. This work is based upon observations obtained with the Georgia State University Center for High Angular Resolution Astronomy Array at Mount Wilson Observatory. The CHARA Array is supported by the National Science Foundation under Grant No. AST-1636624 and AST-1715788. Institutional support has been provided from the GSU College of Arts and Sciences and the GSU Office of the Vice President for Research and Economic Development. The VEGA time at the CHARA Array was granted through the NOIR Lab community access program (PI: Ligi; 2017A-0162, 2018A-0178, 2018B-0019, 2020A-0172).

R.L. has received funding from the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement n. 664931 and from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement CoG #683029)

We thank Dan Huber and Tim White for their time so willingly given to make this project better. We thank Pierre Maxted for their assistance in compiling and running **bagemass**. We thank Jason Eastman for their assistance with **EXOFASTv2**. Finally, we offer our appreciation to the anonymous reviewer whose thorough comments helped make this article its best possible realization.

This work uses results from the TESS mission. Funding for the TESS mission is provided by NASA’s Science Mission directorate.

This research made use of the Jean-Marie Mariotti Center JSDC catalogue⁷.

This research made use of the SIMBAD literature database, operated at CDS, Strasbourg, France, and of NASA’s Astrophysics Data System.

This work presents results from the European Space Agency (ESA) space mission Gaia. Gaia data are being processed by the Gaia Data Processing and Analysis Consortium (DPAC). Funding for the DPAC is provided by national institutions, in particular the institutions participating in the Gaia MultiLateral Agreement (MLA). The Gaia mission website is <https://www.cosmos.esa.int/gaia>. The Gaia archive website is <https://archives.esac.esa.int/gaia>.

⁷ available at http://www.jmmc.fr/catalogue_jsdc.htm

This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration.

This research made use of Lightkurve, a Python package for Kepler and TESS data analysis (Lightkurve Collaboration, 2018).

Facilities: CHARA (PAVO, VEGA, Classic)

Software: Scipy (Jones et al. 2001), astropy (Astropy Collaboration et al. 2018), Red-fluor & Calibir, PAVO (Ireland et al. 2008), vegadrs (Mourard et al. 2009, 2011), bagemass (Maxted et al. 2015), EXOFAST v2 (Eastman 2017), Limb Darkening Toolkit (Parviainen & Aigrain 2015; Husser et al. 2013), Lightkurve (and dependencies) (Lightkurve Collaboration et al. 2018), celerite (Foreman-Mackey et al. 2017), Astrocut (Brasseur et al. 2019), Astroquery (Ginsburg et al. 2019)

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author. All interferometric data are available in the CHARA archive ⁸.

REFERENCES

- | | |
|--|---|
| <p>Angus, R., Morton, T. D.,
Foreman-Mackey, D., et al. 2019, AJ,
158, 173</p> <p>Astropy Collaboration, Price-Whelan,
A. M., Sipőcz, B. M., et al. 2018, AJ,
156, 123</p> <p>Baines, E. K., Armstrong, J. T., Schmitt,
H. R., et al. 2018, AJ, 155, 30</p> <p>Baines, E. K., McAlister, H. A., ten
Brummelaar, T. A., et al. 2008, ApJ,
680, 728</p> <p>Berger, T. A., Huber, D., van Saders,
J. L., et al. 2020, AJ, 159, 280</p> <p>Bessell, M. S. 2000, PASP, 112, 961</p> | <p>Bohlin, R. C., Gordon, K. D., &
Tremblay, P. E. 2014, Publications of
the Astronomical Society of the Pacific,
126, 711</p> <p>Boyajian, T. S., von Braun, K., van Belle,
G., et al. 2012, ApJ, 757, 112</p> <p>Brasseur, C. E., Phillip, C., Fleming,
S. W., Mullally, S. E., & White, R. L.
2019, Astrocut: Tools for creating
cutouts of TESS images, ASCL,
ascl:1905.007</p> <p>Bryan, M. L., Knutson, H. A., Lee, E. J.,
et al. 2019, AJ, 157, 52</p> <p>Chelli, A., Duvert, G., Bourguès, L., et al.
2016, A&A, 589, A112</p> <p>Choi, J., Dotter, A., Conroy, C., et al.
2016, ApJ, 823, 102</p> <p>Claret, A., & Bloemen, S. 2011, A&A,
529, A75</p> |
|--|---|
- ⁸ <http://www.chara.gsu.edu/observers/data-policy-ac529>

- Creech-Eakman, M. J., Romero, V. D., Payne, I., et al. 2018, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 10701, Proc. SPIE, 1070106
- Crida, A., Ligi, R., Dorn, C., Borsa, F., & Lebreton, Y. 2018a, *Research Notes of the American Astronomical Society*, 2, 172
- Crida, A., Ligi, R., Dorn, C., & Lebreton, Y. 2018b, *ApJ*, 860, 122
- Dragomir, D., Matthews, J. M., Eastman, J. D., et al. 2013, *The Astrophysical Journal Letters*, 772, L2. <http://stacks.iop.org/2041-8205/772/i=1/a=L2>
- Droege, T. F., Richmond, M. W., Sallman, M. P., & Creager, R. P. 2006, *PASP*, 118, 1666
- Duvert, G. 2016, *VizieR Online Data Catalog*, 2345
- Eastman, J. 2017, EXOFASTv2: Generalized publication-quality exoplanet modeling code, ASCL, ascl:1710.003
- Eastman, J., Gaudi, B. S., & Agol, E. 2013, *PASP*, 125, 83
- Foreman-Mackey, D., Agol, E., Angus, R., & Ambikasaran, S. 2017, *ArXiv*. <https://arxiv.org/abs/1703.09710>
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2021, *A&A*, 649, A1
- Ginsburg, A., Sipőcz, B. M., Brasseur, C. E., et al. 2019, *AJ*, 157, 98
- Gravity Collaboration, Abuter, R., Accardo, M., et al. 2017, *A&A*, 602, A94
- Gravity Collaboration, Lacour, S., Nowak, M., et al. 2019, *A&A*, 623, L11
- Guo, X., Crossfield, I. J. M., Dragomir, D., et al. 2020, *AJ*, 159, 239
- Hanbury Brown, R., Davis, J., Lake, R. J., & Thompson, R. J. 1974, *MNRAS*, 167, 475
- Howard, A. W., Johnson, J. A., Marcy, G. W., et al. 2011, *ApJ*, 730, 10
- Huber, D., Zinn, J., Bojsen-Hansen, M., et al. 2017, *ApJ*, 844, 102
- Husser, T.-O., Wende-von Berg, S., Dreizler, S., et al. 2013, *A&A*, 553, A6
- Ireland, M. J., Mérand, A., ten Brummelaar, T. A., et al. 2008, in *Optical and Infrared Interferometry*, Vol. 7013, 701324
- Isaacson, H., & Fischer, D. 2010, *ApJ*, 725, 875
- Jones, E., Oliphant, T., Peterson, P., et al. 2001, *SciPy: Open source scientific tools for Python*, Online, [Online]. <http://www.scipy.org/>
- Kharchenko, N. V. 2001, *Kinematika i Fizika Nebesnykh Tel*, 17, 409
- Koen, C., Kilkenney, D., van Wyk, F., & Marang, F. 2010, *MNRAS*, 403, 1949
- Kotoneva, E., Flynn, C., Chiappini, C., & Matteucci, F. 2002, *MNRAS*, 336, 879
- Kovács, G., Zucker, S., & Mazeh, T. 2002, *A&A*, 391, 369
- Lachaume, R., Rabus, M., Jordán, A., et al. 2019, *MNRAS*, 484, 2656
- Lawson, P., et al. 2000, *Principles of Long Baseline Stellar Interferometry*
- Lightkurve Collaboration, Cardoso, J. V. d. M., Hedges, C., et al. 2018, *Lightkurve: Kepler and TESS time series analysis in Python*, *Astrophysics Source Code Library*, ASCL, ascl:1812.013
- Ligi, R., Dorn, C., Crida, A., et al. 2019, *A&A*, 631, A92
- Ligi, R., Creevey, O., Mourard, D., et al. 2016, *A&A*, 586, A94. <https://doi.org/10.1051/0004-6361/201527054>
- Lindgren, L., Bastian, U., Biermann, M., et al. 2021, *A&A*, 649, A4
- Mamajek, E. E., & Hillenbrand, L. A. 2008, *ApJ*, 687, 1264
- Mann, A. W., & von Braun, K. 2015, *Publications of the Astronomical Society of the Pacific*, 127, 102
- Maxted, P. F. L., Serenelli, A. M., & Southworth, J. 2015, *A&A*, 575, A36
- Mermilliod, J.-C. 1994, *VizieR Online Data Catalog*, 2193
- Mourard, D., Clausse, J. M., Marcotto, A., et al. 2009, *A&A*, 508, 1073
- Mourard, D., Bério, P., Perraut, K., et al. 2011, *A&A*, 531, A110

- Parviainen, H., & Aigrain, S. 2015, MNRAS, 453, 3821.
<http://mnras.oxfordjournals.org/lookup/doi/10.1093/mnras/stv1857>
- Pickles, A. J. 1998, PASP, 110, 863
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, Journal of Astronomical Telescopes, Instruments, and Systems, 1, 014003
- Schwarz, G. 1978, The Annals of Statistics, 6, 461 . <https://doi.org/10.1214/aos/1176344136>
- Seager, S., & Mallén-Ornelas, G. 2003, ApJ, 585, 1038
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, The Astronomical Journal, 131, 1163.
<https://doi.org/10.1086%2F498708>
- Spada, F., Demarque, P., Kim, Y.-C., Boyajian, T. S., & Brewer, J. M. 2017, ApJ, 838, 161
- ten Brummelaar, T. A., McAlister, H. A., Ridgway, S. T., et al. 2005, ApJ, 628, 453
- van Belle, G. T., Ciardi, D. R., & Boden, A. F. 2007, ApJ, 657, 1058
- van Belle, G. T., & van Belle, G. 2005, PASP, 117, 1263
- van Grootel, V., Gillon, M., Valencia, D., et al. 2014, The Astrophysical Journal, 786, 2. <http://stacks.iop.org/0004-637X/786/i=1/a=2>
- van Leeuwen, F. 2007, A&A, 474, 653
- von Braun, K., & Boyajian, T. 2017, arXiv e-prints, arXiv:1707.07405
- von Braun, K., Boyajian, T. S., ten Brummelaar, T. A., et al. 2011, ApJ, 740, 49
- von Braun, K., Boyajian, T. S., Kane, S. R., et al. 2012, ApJ, 753, 171
- Weiss, A., & Schlattl, H. 2008, Ap&SS, 316, 99
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, AJ, 140, 1868