

UNIQUENESS OF A POSITIVE SOLUTION FOR THE LAPLACE EQUATION WITH INDEFINITE SUPERLINEAR BOUNDARY CONDITION

KENICHIRO UMEZU

ABSTRACT. In this paper, we consider the Laplace equation with a class of indefinite superlinear boundary conditions and study the uniqueness of positive solutions that this problem possesses. Superlinear elliptic problems can be expected to have multiple positive solutions under certain situations. To our end, by conducting spectral analysis for the linearized eigenvalue problem at an unstable positive solution, we find sufficient conditions for ensuring that the implicit function theorem is applicable to the unstable positive one. An application of our results to the logistic boundary condition arising from population genetics is given.

1. INTRODUCTION AND MAIN RESULTS

Let Ω be a bounded domain of \mathbb{R}^N , $N \geq 1$, with smooth boundary $\partial\Omega$. Consider positive solutions of the Laplace equation with indefinite superlinear boundary condition

$$\begin{cases} -\Delta v = 0 & \text{in } \Omega, \\ \frac{\partial v}{\partial \nu} = \lambda g(x)(v + v^p) & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where $p > 1$ is a given exponent, $\lambda > 0$ is a parameter, $g \in C^{1+\theta}(\partial\Omega)$ changes sign, and ν is the unit outer normal to $\partial\Omega$. Throughout this paper, unless stated otherwise, we assume that $p > 1$ is *subcritical*, i.e.,

$$1 < p < \frac{N}{N-2} \quad \text{if } N > 2. \quad (1.2)$$

A nonnegative solution $u \in C^{2+\alpha}(\overline{\Omega})$, $\alpha \in (0, 1)$, of (1.1) is called *positive* if $u \not\equiv 0$. Using the strong maximum principle (SMP) and Hopf's boundary point lemma (BPL) ([26]), a positive solution of (1.1) is positive in $\overline{\Omega}$. For the existence of a positive solution, we can show that if $\int_{\partial\Omega} g \geq 0$, no positive solution of (1.1) exists for any $\lambda > 0$ (Proposition 2.7). Therefore, in terms of the existence, we focus our consideration of (1.1) on the case when $\int_{\partial\Omega} g < 0$.

2020 *Mathematics Subject Classification.* 35J20, 35J25, 35J65, 35B32, 35P30, 92D40.

Key words and phrases. Laplace equation, indefinite superlinear boundary condition, positive solution, spectral analysis, variational approach, population genetics.

This work was supported by JSPS KAKENHI Grant Number JP18K03353.

When $\int_{\partial\Omega} g < 0$, let $\lambda_1(g) > 0$ be the positive principal eigenvalue (smallest positive eigenvalue) of the linear eigenvalue problem

$$\begin{cases} -\Delta\varphi = 0 & \text{in } \Omega, \\ \frac{\partial\varphi}{\partial\nu} = \lambda g(x)\varphi & \text{on } \partial\Omega. \end{cases} \quad (1.3)$$

Here, an eigenvalue of (1.3) is called *principal* if the eigenfunctions associated with it have constant sign. In fact, (1.3) has exactly two principal eigenvalues $\lambda = 0, \lambda_1(g)$ if $\int_{\partial\Omega} g < 0$, which are both simple and possess eigenfunctions that are positive in $\overline{\Omega}$ ([32]). We call $\varphi_1(g)$ a positive eigenfunction associated with $\lambda_1(g)$ (nonzero constants are the eigenfunctions associated with the principal eigenvalue $\lambda = 0$). We note that the smallest eigenvalue of $(-\Delta, \frac{\partial}{\partial\nu} - \lambda g(x))$ is positive for $\lambda \in (0, \lambda_1(g))$ (Lemma 2.1). For the case of $\int_{\partial\Omega} g \geq 0$, we know that (1.3) has no positive principal eigenvalue (i.e., zero is a unique nonnegative principal eigenvalue), thus, it is understood that $\lambda_1(g) = 0$, and $\varphi_1(g)$ is a positive constant.

In this paper, we aim to give a class of g satisfying $\int_{\partial\Omega} g < 0$ in which (1.1) has a *unique* positive solution v_λ for each $\lambda \in (0, \lambda_1(g))$ and no positive solution for any $\lambda \geq \lambda_1(g)$. Moreover, we show that $v_\lambda \in C^2(\overline{\Omega})$ is parametrized smoothly by $\lambda \in (0, \lambda_1(g))$, bifurcates from the *constant line* $\{(\lambda, 0)\}$ at $\lambda = \lambda_1(g)$, satisfies $\|v_\lambda\|_{C(\overline{\Omega})} \rightarrow \infty$ as $\lambda \searrow 0$, and is unstable (Figure 1). In Section 4, we apply this result to the Laplace equation with the logistic boundary condition, (4.1).




FIGURE 1. Positive solution set $\{(\lambda, v_\lambda)\}$ of (1.1) as a smooth curve.

For our aim, the change of variables

$$w = \lambda^{\frac{1}{p-1}} v \quad (1.4)$$

is useful, because it transforms (1.1) into the problem

$$\begin{cases} -\Delta w = 0 & \text{in } \Omega, \\ \frac{\partial w}{\partial\nu} = \lambda g(x)w + g(x)w^p & \text{on } \partial\Omega. \end{cases} \quad (1.5)$$

Before discussing the uniqueness issue, we consider the general case of g , and use the fibering map method to establish the existence of positive solutions and their properties for (1.5) as follows:

Theorem 1.1. *Assume that $\int_{\partial\Omega} g(x) < 0$. Then, (1.5) possesses a positive solution w_λ for every $\lambda \in (0, \lambda_1(g))$ and no positive solution for any $\lambda \geq \lambda_1(g)$. Furthermore, w_λ satisfies the following:*

- (i) w_λ is unstable,
- (ii) $\|w_\lambda\|_{C^2(\bar{\Omega})} \rightarrow 0$ as $\lambda \nearrow \lambda_1(g)$,
- (iii) $\sup_{\lambda \in (0, \lambda_1(g))} \|w_\lambda\|_{C(\bar{\Omega})} < \infty$, and
- (iv) $\inf_{\lambda \in (0, \bar{\lambda})} \|w_\lambda\|_{C(\bar{\Omega})} > 0$ for any $\bar{\lambda} \in (0, \lambda_1(g))$.

We refer to [34] for the existence of positive solutions to a similar type of nondivergence elliptic problem with indefinite superlinear boundary conditions. For nonlinear elliptic equations with definite superlinear boundary conditions, we refer to the survey article [29]. We refer to [25, Theorem 3] for a similar existence result for positive solutions of the Neumann problem

$$\begin{cases} -\Delta w = \lambda m(x)w + m(x)w^p & \text{in } \Omega, \\ \frac{\partial w}{\partial \nu} = 0 & \text{on } \partial\Omega. \end{cases} \quad (1.6)$$

Here, $p > 1$ is subcritical, i.e., $p < \frac{N+2}{N-2}$ if $N > 2$, $\lambda > 0$ is a parameter, $m \in C^\theta(\bar{\Omega})$ changes sign such that $\int_{\Omega} m < 0$, and $(-\Delta - \lambda m(x), \frac{\partial}{\partial \nu})$ has a positive smallest eigenvalue. See also [2, 4, 5, 6, 12, 24, 30] for the existence and related issues for positive solutions of the analogous indefinite superlinear elliptic equations with linear Dirichlet or Neumann boundary conditions.

Next, we consider a special case of g that is central to this paper. For a sign changing $g \in C^{1+\theta}(\partial\Omega)$, we set

$$\begin{aligned} \Gamma_{\pm} &:= \{x \in \partial\Omega : g(x) \gtrless 0\}, \\ \Gamma_0 &:= \{x \in \partial\Omega : g(x) = 0\}, \\ \Gamma_{-,0} &:= \Gamma_- \cup \Gamma_0. \end{aligned} \quad (1.7)$$

It should be noted that Γ_+ is open in the relative topology of $\partial\Omega$. We then introduce the following condition for Γ_+ (Figure 2):

$$\Gamma_+ \text{ is a compact submanifold of } \partial\Omega \text{ with dimension } N-1. \quad (1.8)$$

For $g \in C^{1+\theta}(\partial\Omega)$ equipped with (1.8), we define $g_\delta \in C^{1+\theta}(\partial\Omega)$ by the formula

$$g_\delta := g^+ - \delta g^-, \quad \delta > \delta_0 := \frac{\int_{\partial\Omega} g^+}{\int_{\partial\Omega} g^-}. \quad (1.9)$$

We observe that $\int_{\partial\Omega} g_\delta < 0$, and $\int_{\partial\Omega} g_\delta \nearrow \int_{\partial\Omega} g_{\delta_0} = 0$ as $\delta \searrow \delta_0$, and know that $\lambda_1(g_\delta) \rightarrow 0$ as $\delta \searrow \delta_0$ (see (3.1)).




FIGURE 2. Situations admitting (1.8): (i) $\Gamma_+ = \bigcup_{j=1}^k \gamma_j$, (ii) $\Gamma_+ = \Gamma_1 \cup \left(\bigcup_{j=2}^k \gamma_j \right)$, and (iii) $\Gamma_{-,0} = \gamma_1 \cup \gamma_2$.

We then present our main result for (1.5), where a precise description of the positive solution set $\{(\lambda, v_\lambda)\}$ of (1.5) is given if δ is sufficiently close to δ_0 , i.e., if the situation of g_δ is assumed to be near the critical case $\int_{\partial\Omega} g_{\delta_0} = 0$.

Theorem 1.2. *Let $g_\delta \in C^{1+\theta}(\partial\Omega)$ be introduced by (1.9). If δ is sufficiently close to δ_0 , then the positive solution set of (1.5) with $g = g_\delta$ for $\lambda \in (0, \lambda_1(g_\delta))$ is given as follows (Figure 3):*

- (i) (1.5) with $g = g_\delta$ possesses a unique positive solution w_λ for every $\lambda \in (0, \lambda_1(g_\delta))$, and the positive solution set $\{(\lambda, w_\lambda) : \lambda \in (0, \lambda_1(g_\delta))\}$ is represented by a smooth curve,
- (ii) $w_\lambda \rightarrow 0$ in $C^2(\overline{\Omega})$ as $\lambda \nearrow \lambda_1(g_\delta)$, i.e., bifurcation from $\{(\lambda, 0)\}$ at $(\lambda_1(g_\delta), 0)$ occurs subcritically, and
- (iii) $w_\lambda \rightarrow w_0$ in $C^2(\overline{\Omega})$ as $\lambda \searrow 0$ for some $w_0 \in C^2(\overline{\Omega})$, where w_0 is a unique positive solution of (1.5) with $g = g_\delta$ for $\lambda = 0$ (actually the smooth positive solution curve is extended slightly to $\lambda < 0$).

As a byproduct of assertion (iii), we obtain that if δ is close to δ_0 , then the uniqueness of a positive solution holds for (1.5) with $g = g_\delta$ for $\lambda = 0$:

$$\begin{cases} -\Delta w = 0 & \text{in } \Omega, \\ \frac{\partial w}{\partial \nu} = g(x)w^p & \text{on } \partial\Omega. \end{cases}$$




FIGURE 3. Positive solution set $\{(\lambda, w_\lambda)\}$ of (1.5) with $g = g_\delta$ as a smooth curve in the case that δ is close to δ_0 .

We refer to [25, Theorem 2] for the existence result for this problem. We also refer to [8] for high multiplicity of positive solutions of (1.6) with $\lambda = 0$ when the negative part of m is large (see [7] for the Dirichlet case).

Going back to (1.1) by (1.4), Theorems 1.1 and 1.2 are analogous to Theorems 2.12 and 3.5 in Sections 2 and 3, respectively. Theorem 3.5 provides us with a precise description of the positive solution set for (1.1), as shown in Figure 1.

To conclude the Introduction, it should be noted that our approach employed in the proofs of Theorem 1.1 and 1.2 remains valid for the following more general situation:

$$\begin{cases} -\Delta w = 0 & \text{in } \Omega, \\ \frac{\partial w}{\partial \nu} = \lambda g(x)w + f(x)w^p & \text{on } \partial\Omega, \end{cases} \quad (1.10)$$

where $p > 1$ is subcritical, $\lambda > 0$, and $g, f \in C^{1+\theta}(\partial\Omega)$ both change signs (see Theorems 2.13 and 3.6 in Sections 2 and 3, respectively). In addition, similar results to Theorems 1.1 and 1.2 can be established for (1.6). For high multiplicity of positive solutions to a similar type of (1.6), we refer to [31].

The remainder of this paper is organized as follows. Section 2 is devoted to the proof of Theorem 1.1. In Subsection 2.1, we prove the existence assertion by employing the variational approach based on the Nehari manifold and fibering map with (1.5). In Subsection 2.2, we use a Picone type identity to verify the nonexistence assertion. In Subsection 2.3, we prove assertion (i) with the aid of an idea from [9]. In Subsection 2.4, we prove assertions (ii) to (iv) using a variational approach.

Section 3 is devoted to the proof of Theorem 1.2, where a crucial step is the establishment of Proposition 3.3. We conduct spectral analysis for the linearized eigenvalue problem at each positive solution of (1.5).

In Section 4, we present an application of Theorems 1.1 and 1.2 to logistic boundary conditions. Our main results in this section are Theorems 4.1, 4.2, and 4.4. In Subsection 4.5, an analysis for the one-dimensional space case is given, where a numerical observation is added.

2. PROOF OF THEOREM 1.1

In this section, we prove Theorem 1.1.

2.1. Existence. This subsection assumes that $\int_{\partial\Omega} g < 0$. We define the functional associated with (1.5)

$$J_\lambda(w) := \frac{1}{2}E_\lambda(w) - \frac{1}{p+1}G(w), \quad w \in H^1(\Omega),$$

where

$$E_\lambda(w) := \int_{\Omega} |\nabla w|^2 - \lambda \int_{\partial\Omega} g(x)w^2, \quad G(w) := \int_{\partial\Omega} g(x)|w|^{p+1}.$$

Using the divergence theorem, it is easy to deduce that

$$\int_{\partial\Omega} g(x)\varphi_1(g)^2 > 0 \quad \text{and} \quad G(\varphi_1(g)) > 0,$$

which are used repeatedly in the following.

The next lemma implies that $E_\lambda(\cdot)$ is coercive in $H^1(\Omega)$ for $\lambda \in (0, \lambda_1(g))$. Here, $\|\cdot\|$ denotes the usual norm of $H^1(\Omega)$.

Lemma 2.1. *Let $I \subset (0, \lambda_1(g))$ be a compact interval. Then, there exists a constant $c_I > 0$ such that*

$$E_\lambda(w) \geq c_I \|w\|^2, \quad w \in H^1(\Omega) \quad \text{and} \quad \lambda \in I.$$

For our procedure, we use the fibering map for J_λ , $\lambda \in (0, \lambda_1(g))$, which is introduced as follows: for $w \neq 0$, we set

$$j_w(t) := J_\lambda(tw) = \frac{t^2}{2}E_\lambda(w) - \frac{t^{p+1}}{p+1}G(w), \quad t > 0.$$

The associated Nehari manifold is also introduced:

$$\mathcal{N}_\lambda := \{w \in H^1(\Omega) \setminus \{0\} : j'_w(1) = 0\} = \{w \in H^1(\Omega) \setminus \{0\} : E_\lambda(w) = G(w)\}.$$

We split \mathcal{N}_λ into three parts as follows:

$$\begin{aligned} \mathcal{N}_\lambda^\pm &:= \{w \in \mathcal{N}_\lambda : j''_w(1) \geq 0\} \\ &= \{w \in H^1(\Omega) \setminus \{0\} : E_\lambda(w) = G(w), E_\lambda(w) \geq pG(w)\}, \\ \mathcal{N}_\lambda^0 &:= \{w \in \mathcal{N}_\lambda : j''_w(1) = 0\} \\ &= \{w \in H^1(\Omega) \setminus \{0\} : E_\lambda(w) = G(w), E_\lambda(w) = pG(w)\}. \end{aligned}$$

However, from Lemma 2.1, we deduce that if $\lambda \in (0, \lambda_1(g))$, then for any $w \in \mathcal{N}_\lambda$, we obtain

$$E_\lambda(w) - pG(w) = (1-p)E_\lambda(w) < 0.$$

The next lemma is thus proved.

Lemma 2.2. $\mathcal{N}_\lambda = \mathcal{N}_\lambda^- \neq \emptyset$ and $\mathcal{N}_\lambda^+ \cup \mathcal{N}_\lambda^0 = \emptyset$ for $\lambda \in (0, \lambda_1(g))$.

Proof. It remains to prove $\mathcal{N}_\lambda^- \neq \emptyset$. We use a positive eigenfunction $\varphi_1 = \varphi_1(g)$ associated with $\lambda_1(g)$ to verify the assertion. We infer from Lemma 2.1 that $E_\lambda(\varphi_1) > 0$. Since $G(\varphi_1) > 0$, the fibering map $j_{\varphi_1}(\cdot)$ has a unique global maximum point $t_0 > 0$. This implies that $j'_{\varphi_1}(t_0) = 0 > j''_{\varphi_1}(t_0)$. Thus, $t_0\varphi_1 \in \mathcal{N}_\lambda^-$. \square

The next lemma asserts that $J_\lambda(\cdot)$ is positive in \mathcal{N}_λ^- for each $\lambda \in (0, \lambda_1(g))$.

Lemma 2.3. Let $\lambda \in (0, \lambda_1(g))$. Then, $J_\lambda(w) > 0$ for all $w \in \mathcal{N}_\lambda^-$.

Proof. Let $\lambda \in (0, \lambda_1(g))$, and $w \in \mathcal{N}_\lambda^-$. Then, $E_\lambda(w) > 0$ from Lemma 2.1, which implies that $G(w) > 0$. Hence, $J_\lambda(w) = j_w(1) = \max_{t>0} j_w(t) > 0$. \square

We then prove the existence of a minimizer in \mathcal{N}_λ^- for J_λ . Let $\lambda \in (0, \lambda_1(g))$. From Lemma 2.3, we can say $\gamma := \inf_{\mathcal{N}_\lambda^-} J_\lambda(w) \geq 0$, and we take a minimizing sequence $\{w_n\} \subset \mathcal{N}_\lambda^-$ such that $J_\lambda(w_n) \searrow \gamma$. We then obtain the following lemma.

Lemma 2.4. $\sup_{n \geq 1} \|w_n\| < \infty$.

Proof. By contradiction, we assume $\|w_n\| \rightarrow \infty$. Say $\eta_n = \frac{w_n}{\|w_n\|}$, and $\|\eta_n\| = 1$. Since $J_\lambda(w_n)$ is bounded by the choice of w_n , we infer the existence of $C > 0$ such that

$$\left(\frac{1}{2} - \frac{1}{p+1}\right) E_\lambda(w_n) = J_\lambda(w_n) \leq C.$$

This implies that $\overline{\lim}_{n \rightarrow \infty} E_\lambda(\eta_n) \leq 0$. Thus, by Lemma 2.1, $\eta_n \rightarrow 0$ in $H^1(\Omega)$, which is a contradiction. \square

From Lemma 2.4, we may infer that for some $w_0 \in H^1(\Omega)$, $w_n \rightharpoonup w_0$, and $w_n \rightarrow w_0$ in $L^2(\Omega)$ and $L^{p+1}(\partial\Omega)$. We then have the following.

Lemma 2.5.

- (i) $E_\lambda(w_0) > 0$ and $G(w_0) > 0$.
- (ii) $w_n \rightarrow w_0$ in $H^1(\Omega)$.

Proof. We verify assertion (i). We claim that $\gamma > 0$. Assume by contradiction that $J_\lambda(w_n) \rightarrow 0$, and then, from $w_n \in \mathcal{N}_\lambda^-$, we deduce

$$E_\lambda(w_n) = \left(\frac{1}{2} - \frac{1}{p+1}\right)^{-1} J_\lambda(w_n) \rightarrow 0.$$

From Lemma 2.1, $\|w_n\| \rightarrow 0$. However, $\eta_n = \frac{w_n}{\|w_n\|}$ satisfies

$$E_\lambda(\eta_n) = G(\eta_n) \|w_n\|^{p-1} \rightarrow 0,$$

and using Lemma 2.1 again, we obtain $\|\eta_n\| \rightarrow 0$, which is contradictory for $\|\eta_n\| = 1$, as desired. Since $w_n \in \mathcal{N}_\lambda^-$ and $\gamma > 0$, we see

$$G(w_0) = \lim_{n \rightarrow \infty} G(w_n) = \lim_{n \rightarrow \infty} \left(\frac{1}{2} - \frac{1}{p+1} \right)^{-1} J_\lambda(w_n) = \left(\frac{1}{2} - \frac{1}{p+1} \right)^{-1} \gamma > 0.$$

In particular, $w_0 \neq 0$, thus, Lemma 2.1 ensures $E_\lambda(w_0) > 0$. Assertion (i) is now verified.

We then verify assertion (ii). Assume to the contrary that

$$w_n \not\rightarrow w_0 \text{ in } H^1(\Omega). \quad (2.1)$$

Since $E_\lambda(w_0) > 0$ and $G(w_0) > 0$, j_{w_0} has a unique global maximum point $t_0 > 0$, and consequently, $t_0 w_0 \in \mathcal{N}_\lambda^-$. By taking a suitable subsequence of $\{w_n\}$, denoted by the same notation, (2.1) infers

$$J_\lambda(t_0 w_0) = j_{w_0}(t_0) < \lim_{n \rightarrow \infty} j_{w_n}(t_0) \leq \lim_{n \rightarrow \infty} j_{w_n}(1) = \lim_{n \rightarrow \infty} J_\lambda(w_n) = \gamma,$$

which is contradictory for $t_0 w_0 \in \mathcal{N}_\lambda^-$. Here, we have used the fact that j_{w_n} has the unique global maximum point $t = 1$. Assertion (ii) has been verified. \square

From Lemma 2.5, we derive the following existence result.

Proposition 2.6. *Assume that $\int_{\partial\Omega} g < 0$. Then, (1.5) has a variational positive solution for every $\lambda \in (0, \lambda_1(g))$.*

Proof. By Lemma 2.5 (i), there exists $t_0 > 0$ such that $j'_{w_0}(t_0) = 0 > j''_{w_0}(t_0)$, thus, $t_0 w_0 \in \mathcal{N}_\lambda^-$. We claim that $t_0 = 1$. Once this is verified, Lemma 2.5 (ii) shows that

$$J_\lambda(w_0) = \lim_{n \rightarrow \infty} J_\lambda(w_n) = \gamma \quad \text{and} \quad w_0 \in \mathcal{N}_\lambda^-,$$

thus, Lemma 2.3 provides

$$J_\lambda(w_0) = \min_{w \in \mathcal{N}_\lambda^-} J_\lambda(w) > 0.$$

Without loss of generality, we may assume that $w_0 \geq 0$. Since $w_0 \neq 0$, the SMP and BPL apply, and we obtain that $w_0 > 0$ in $\overline{\Omega}$. By a similar argument as in [11, Theorem 2.3], w_0 is a critical point for J_λ . The desired conclusion thus follows.

It remains to show that $t_0 = 1$. Note that if $j'_{w_0}(t) = 0 \geq j''_{w_0}(t)$, then $t = t_0$. Since $j'_{w_n}(1) = 0 > j''_{w_n}(1)$ from $w_n \in \mathcal{N}_\lambda^-$, passing to the limit provides that $j'_{w_0}(1) = 0 \geq j''_{w_0}(1)$, using Lemma 2.5 (ii). Thus, $t_0 = 1$. \square

2.2. Nonexistence. This subsection deals with the general case of g such that $g \in C^{1+\theta}(\partial\Omega)$ changes sign. For $\lambda \in \mathbb{R}$, let $\sigma_1 = \sigma_1(\lambda) \in \mathbb{R}$ be the smallest eigenvalue of the eigenvalue problem

$$\begin{cases} -\Delta\phi = \sigma\phi & \text{in } \Omega, \\ \frac{\partial\phi}{\partial\nu} = \lambda g(x)\phi & \text{on } \partial\Omega. \end{cases}$$

We can characterize σ_1 by the following variational formula ([32]):

$$\sigma_1 = \inf \left\{ \int_{\Omega} |\nabla \phi|^2 - \lambda \int_{\partial\Omega} g(x) \phi^2 : \int_{\Omega} \phi^2 = 1 \right\}. \quad (2.2)$$

We know that σ_1 is simple and has positive eigenfunctions, and it satisfies

$$\begin{cases} \sigma_1(0) = \sigma_1(\lambda_1(g)) = 0, \\ \sigma_1(\lambda) > 0 & \text{for } \lambda \in (0, \lambda_1(g)), \\ \sigma_1(\lambda) < 0 & \text{for } \lambda > \lambda_1(g). \end{cases} \quad (2.3)$$

We call $\Phi_1 \in C^{2+\alpha}(\overline{\Omega})$ a positive eigenfunction of (2.2) that is associated with σ_1 , satisfying that $\Phi_1 > 0$ in $\overline{\Omega}$. Note that Φ_1 is characterized by $\varphi_1(g)$ multiplied by a positive constant when $\lambda = \lambda_1(g)$.

We then obtain the following nonexistence result.

Proposition 2.7. *Let $p > 1$ be arbitrary (without (1.2)). Then, (1.5) has no positive solution for any $\lambda \geq \lambda_1(g)$.*

Proof. First, we consider the case when $\int_{\partial\Omega} g \geq 0$ and $\lambda = 0$. It should be noted that $\lambda_1(g) = 0$ in this case. If w is a positive solution of (1.5) for $\lambda = 0$, then w is not a constant, and the divergence theorem gives

$$0 = \int_{\Omega} -\Delta w w^{-p} = \int_{\Omega} \nabla w \nabla(w^{-p}) - \int_{\partial\Omega} g < - \int_{\partial\Omega} g \leq 0.$$

The assertion thus follows.

Next, we consider the case of $\lambda > 0$, and prove the nonexistence assertion only for the case of $\int_{\partial\Omega} g < 0$. The case of $\int_{\partial\Omega} g \geq 0$ is similar, so we omit the proof. Let w be a positive solution of (1.5) for $\lambda > 0$. We then use the Picone type identity

$$\begin{aligned} & -\sigma_1 \Phi_1^{p+1} w^{1-p} \\ &= \left(\frac{\Phi_1}{w} \right)^p (\Delta \Phi_1 w - \Phi_1 \Delta w) = \left(\frac{\Phi_1}{w} \right)^p \sum_{j=1}^N \frac{\partial}{\partial x_j} \left(w^2 \frac{\partial}{\partial x_j} \left(\frac{\Phi_1}{w} \right) \right). \end{aligned}$$

The divergence theorem shows that

$$\begin{aligned} & -\sigma_1 \int_{\Omega} \Phi_1^{p+1} w^{1-p} \\ &= \int_{\Omega} \left(\frac{\Phi_1}{w} \right)^p \sum_{j=1}^N \frac{\partial}{\partial x_j} \left(w^2 \frac{\partial}{\partial x_j} \left(\frac{\Phi_1}{w} \right) \right) \\ &= -p \int_{\Omega} \left(\frac{\Phi_1}{w} \right)^{p-1} w^2 \left| \nabla \left(\frac{\Phi_1}{w} \right) \right|^2 + \int_{\partial\Omega} \left(\frac{\Phi_1}{w} \right)^p \left(\frac{\partial \Phi_1}{\partial \nu} w - \Phi_1 \frac{\partial w}{\partial \nu} \right) \\ &= -p \int_{\Omega} \left(\frac{\Phi_1}{w} \right)^{p-1} w^2 \left| \nabla \left(\frac{\Phi_1}{w} \right) \right|^2 - \int_{\partial\Omega} g \Phi_1^{p+1} \\ &\leq - \int_{\partial\Omega} g \Phi_1^{p+1} \end{aligned}$$

$$= \lambda^{-1} \left(\sigma_1 \int_{\Omega} \Phi_1^{p+1} - p \int_{\Omega} \Phi_1^{p-1} |\nabla \Phi_1|^2 \right).$$

Here, the last equality is deduced from the computation of $\int_{\Omega} -\Delta \Phi_1 \Phi_1^p$ via the divergence theorem. Thus, using (2.3), we infer $\lambda \in (0, \lambda_1(g))$. \square

2.3. Instability. Let w be a positive solution of (1.5) for $\lambda \geq 0$. We call $\gamma_1(\lambda, w) \in \mathbb{R}$ the smallest eigenvalue of the eigenvalue problem

$$\begin{cases} -\Delta \phi = \gamma \phi & \text{in } \Omega, \\ \frac{\partial \phi}{\partial \nu} = \lambda g(x) \phi + p g(x) w^{p-1} \phi + \gamma \phi & \text{on } \partial \Omega. \end{cases} \quad (2.4)$$

It is well known that γ_1 is simple, and the corresponding eigenfunctions possess constant signs. For a nonnegative eigenfunction $\phi_1(\lambda, w)$ associated with γ_1 , from the SMP and BPL, we obtain that $\phi_1 > 0$ in $\bar{\Omega}$. The positive solution w is called *asymptotically stable* and *unstable* if $\gamma_1 > 0$ and $\gamma_1 < 0$, respectively.

In view of Propositions 2.6, we then prove the following instability result.

Proposition 2.8. *Let $p > 1$ be arbitrary (without (1.2)). If $\int_{\partial \Omega} g < 0$, then every positive solution of (1.5) for $\lambda \in [0, \lambda_1(g))$ is unstable.*

Proof. The proof is inspired by that of [9, Theorem 1]. Let w be a positive solution of (1.5) for $\lambda \in [0, \lambda_1(g))$. Say $h(t) := \lambda t + t^p$, $t > 0$, and then, for any $\lambda > 0$, we see that $h(w) > 0$ in $\bar{\Omega}$, and $h''(w) = p(p-1)w^{p-2} > 0$ in $\bar{\Omega}$. We also see that γ_1 and ϕ_1 satisfy

$$\begin{cases} -\Delta \phi_1 = \gamma_1 \phi_1 & \text{in } \Omega, \\ \frac{\partial \phi_1}{\partial \nu} = g(x) h'(w) \phi_1 + \gamma_1 \phi_1 & \text{on } \partial \Omega. \end{cases}$$

The divergence theorem shows that

$$\begin{aligned} \gamma_1 \int_{\Omega} h(w) \phi_1 &= \int_{\Omega} (-\Delta \phi_1 h(w) + \Delta w h'(w) \phi_1) \\ &= - \int_{\Omega} h''(w) |\nabla w|^2 \phi_1 - \gamma_1 \int_{\partial \Omega} h(w) \phi_1. \end{aligned}$$

Therefore,

$$\gamma_1 = - \frac{\int_{\Omega} h''(w) |\nabla w|^2 \phi_1}{\int_{\Omega} h(w) \phi_1 + \int_{\partial \Omega} h(w) \phi_1} < 0,$$

as desired. \square

2.4. Bounds and Asymptotics. We complete the proof of Theorem 1.1 in this subsection. The next proposition asserts that the variational positive solution w_λ is bounded in $H^1(\Omega)$ for $\lambda \in (0, \lambda_1(g))$.

Proposition 2.9. *Assume that $\int_{\partial \Omega} g < 0$. Let w_λ be the variational positive solution of (1.5) for $\lambda \in (0, \lambda_1(g))$, constructed by Proposition 2.6. Then,*

$$\sup_{\lambda \in (0, \lambda_1(g))} \|w_\lambda\| < \infty.$$

Proof. We verify the existence of $C > 0$ satisfying that if $w_n := w_{\lambda_n}$ is the positive solution of (1.5) for $\lambda = \lambda_n \in (0, \lambda_1(g))$, then

$$J_{\lambda_n}(w_n) = \inf_{w \in \mathcal{N}_{\lambda_n}^-} J_{\lambda_n}(w) \leq C. \quad (2.5)$$

To this end, we take a smooth function \hat{w} in $\overline{\Omega}$ such that $G(\hat{w}) > 0$. Then, Lemma 2.1 gives $E_{\lambda_n}(\hat{w}) > 0$. Therefore, the mapping

$$t \mapsto \frac{t^2}{2} E_{\lambda_n}(\hat{w}) - \frac{t^{p+1}}{p+1} G(\hat{w})$$

has a global maximum point $t = t_n > 0$, implying that $t_n \hat{w} \in \mathcal{N}_{\lambda_n}^-$. Thus, $J_{\lambda_n}(w_n) \leq J_{\lambda_n}(t_n \hat{w})$. Observing now that

$$\begin{aligned} J_{\lambda_n}(t_n \hat{w}) &\leq \frac{t_n^2}{2} \left(\int_{\Omega} |\nabla \hat{w}|^2 + \lambda_1(g) \int_{\partial\Omega} g^-(x) \hat{w}^2 \right) - \frac{t_n^{p+1}}{p+1} G(\hat{w}) \\ &\leq \sup_{t>0} \left\{ \frac{t^2}{2} \left(\int_{\Omega} |\nabla \hat{w}|^2 + \lambda_1(g) \int_{\partial\Omega} g^-(x) \hat{w}^2 \right) - \frac{t^{p+1}}{p+1} G(\hat{w}) \right\} < \infty, \end{aligned}$$

assertion (2.5) is thus verified.

We then prove this proposition. Assume by contradiction that $w_n := w_{\lambda_n}$ is the positive solution of (1.5) for $\lambda = \lambda_n \in (0, \lambda_1(g))$ with the condition that $\lambda_n \rightarrow \lambda_{\infty} \in [0, \lambda_1(g)]$ and $\|w_n\| \rightarrow \infty$. Letting $\eta_n = \frac{w_n}{\|w_n\|}$, we may deduce that for some $\eta_0 \in H^1(\Omega)$, $\eta_n \rightarrow \eta_0$, and $\eta_n \rightarrow \eta_0$ in $L^2(\Omega)$ and $L^{p+1}(\partial\Omega)$. From $w_n \in \mathcal{N}_{\lambda_n}^-$, we see that

$$E_{\lambda_n}(w_n) = \left(\frac{1}{2} - \frac{1}{p+1} \right)^{-1} J_{\lambda_n}(w_n). \quad (2.6)$$

Combining (2.5) and (2.6) shows that $\overline{\lim}_{n \rightarrow \infty} E_{\lambda_n}(\eta_n) \leq 0$. Because $\|\eta_n\| = 1$, Lemma 2.1 shows that $\lambda_{\infty} = 0$ or $\lambda_1(g)$, and

$$0 \leq E_{\lambda_{\infty}}(\eta_0) \leq \underline{\lim}_{n \rightarrow \infty} E_{\lambda_n}(\eta_n) \leq \overline{\lim}_{n \rightarrow \infty} E_{\lambda_n}(\eta_n) \leq 0.$$

This implies that $\eta_n \rightarrow \eta_0$ in $H^1(\Omega)$, $\|\eta_0\| = 1$, and $\eta_0 \geq 0$. Moreover, if $\lambda_{\infty} = 0$, then η_0 is a positive constant, whereas $\eta_0 = c\varphi_1(g)$ for some $c > 0$ if $\lambda_{\infty} = \lambda_1(g)$. However, since $w_n \in \mathcal{N}_{\lambda_n}^-$, we infer

$$G(\eta_0) = \lim_{n \rightarrow \infty} G(\eta_n) = \lim_{n \rightarrow \infty} E_{\lambda_n}(\eta_n) \|w_n\|^{1-p} = 0,$$

thus,

$$0 = G(\eta_0) = \begin{cases} \eta_0^{p+1} \int_{\partial\Omega} g(x) < 0, & \lambda_{\infty} = 0, \\ c^{p+1} G(\varphi_1(g)) > 0, & \lambda_{\infty} = \lambda_1(g), \end{cases}$$

which is a contradiction. \square

Corollary 2.10. *Under the conditions of Proposition 2.9, $w_{\lambda} \rightarrow 0$ in $C^2(\overline{\Omega})$ as $\lambda \nearrow \lambda_1(g)$.*

Proof. By elliptic regularity ([29, Theorem 2.2]), it follows from Proposition 2.9 that $\sup_{\lambda \in (0, \lambda_1(g))} \|w_\lambda\|_{W^{1,r}(\Omega)} < \infty$ for $r > N$. Combining the Schauder estimate and the compactness argument shows that, up to a subsequence, $w_{\lambda_n} \rightarrow w_\infty \geq 0$ in $C^2(\overline{\Omega})$ as $\lambda_n \nearrow \lambda_1(g)$, and w_∞ is a nonnegative solution of (1.5) for $\lambda = \lambda_1(g)$. Proposition 2.7 completes the proof with $w_\infty = 0$. \square

The following *a priori* lower bound can be derived for the positive solutions of (1.5):

Proposition 2.11. *Assume $\int_{\partial\Omega} g < 0$. Then, for $\bar{\lambda} \in (0, \lambda_1(g))$, there exists $C > 0$ such that if w is a positive solution of (1.5) for $\lambda \in (0, \bar{\lambda})$, then $\|w\|_{C(\overline{\Omega})} > C$.*

Proof. If not, then we take a positive solution w_n of (1.5) for $\lambda = \lambda_n \searrow 0$ such that $\|w_n\|_{C(\overline{\Omega})} \rightarrow 0$ because $(\lambda_1(g), 0)$ is a unique bifurcation point from $\{(\lambda, 0)\}$ for the positive solutions of (1.5). Then, $\|w_n\| \rightarrow 0$, and additionally, Lemma 2.1 shows that $0 < E_{\lambda_n}(w_n) = G(w_n)$. Set $\eta_n = \frac{w_n}{\|w_n\|}$, and we may deduce that for some $\eta_\infty \in H^1(\Omega)$, $\eta_n \rightarrow \eta_\infty$, and $\eta_n \rightarrow \eta_\infty$ in $L^2(\Omega)$ and $L^{p+1}(\partial\Omega)$. Then, η_n admits

$$\begin{aligned} \int_{\Omega} |\nabla \eta_n|^2 &= \lambda_n \int_{\partial\Omega} g(x) \eta_n^2 + \|w_n\|^{p-1} \int_{\partial\Omega} g(x) \eta_n^{p+1} \rightarrow 0, \\ G(\eta_n) &\rightarrow \int_{\partial\Omega} g(x) \eta_\infty^{p+1} \geq 0. \end{aligned}$$

Therefore, $\eta_n \rightarrow \eta_\infty$ in $H^1(\Omega)$, η_∞ is a positive constant, and thus, $\int_{\partial\Omega} g(x) \geq 0$, which is a contradiction. \square

End of proof of Theorem 1.1. The existence and nonexistence assertions follow from Propositions 2.6 and 2.7, respectively. Proposition 2.8 implies assertion (i). Assertion (ii) is verified by Corollary 2.10. Assertion (iii) is obtained from Proposition 2.9 by the bootstrap argument developed in the proof of [29, Theorem 2.2]. Finally, Proposition 2.11 implies assertion (iv). \square

Recalling (1.4), we convert Theorem 1.1 to (1.1) with $v_\lambda = \lambda^{-\frac{1}{p-1}} w_\lambda$ to formulate the following theorem:

Theorem 2.12. *Assume that $\int_{\partial\Omega} g(x) < 0$. Then, (1.1) possesses a positive solution v_λ for $\lambda \in (0, \lambda_1(g))$ and no positive solution for any $\lambda \geq \lambda_1(g)$. Moreover, v_λ satisfies the following four conditions:*

- (i) v_λ is unstable.
- (ii) $\|v_\lambda\|_{C^2(\overline{\Omega})} \rightarrow 0$ as $\lambda \nearrow \lambda_1(g)$.
- (iii) $\sup_{\lambda \in (\underline{\lambda}, \lambda_1(g))} \|v_\lambda\|_{C(\overline{\Omega})} < \infty$ for any $\underline{\lambda} \in (0, \lambda_1(g))$, and additionally, the following asymptotics holds:

$$c_1 \lambda^{-\frac{1}{p-1}} \leq \|v_\lambda\|_{C(\overline{\Omega})} \leq c_2 \lambda^{-\frac{1}{p-1}} \quad \text{as } \lambda \searrow 0$$

for some $c_2 > c_1 > 0$.

(iv) $\inf_{\lambda \in (0, \bar{\lambda})} \|v_\lambda\|_{C(\bar{\Omega})} > 0$ for any $\bar{\lambda} \in (0, \lambda_1(g))$.

Similar results to those of Theorem 1.1 can be presented for (1.10), which is evaluated in the same way as employed in this section:

Theorem 2.13. *Assume that $\int_{\partial\Omega} g(x) < 0$ and $\int_{\partial\Omega} f(x)\varphi_1(g)^{p+1} > 0$. Then, (1.10) has a positive solution W_λ for every $\lambda \in (0, \lambda_1(g))$ and no positive solution for $\lambda = \lambda_1(g)$, satisfying*

- (i) $\sup_{\lambda \in (\underline{\lambda}, \lambda_1(g))} \|W_\lambda\|_{C(\bar{\Omega})} < \infty$ for any $\underline{\lambda} \in (0, \lambda_1(g))$ and
- (ii) $\|W_\lambda\|_{C^2(\bar{\Omega})} \rightarrow 0$ as $\lambda \nearrow \lambda_1(g)$.

If additionally $\int_{\partial\Omega} f \neq 0$, then it is possible to take $\underline{\lambda} = 0$. Moreover,

- (iii) $\int_{\partial\Omega} f < 0$ implies that $\inf_{\lambda \in (0, \bar{\lambda})} \|W_\lambda\|_{C(\bar{\Omega})} > 0$ for $\bar{\lambda} \in (0, \lambda_1(g))$, whereas
- (iv) $\int_{\partial\Omega} f > 0$ implies that $\inf_{\lambda \in (\underline{\lambda}, \bar{\lambda})} \|W_\lambda\|_{C(\bar{\Omega})} > 0$ for $0 < \underline{\lambda} < \bar{\lambda} < \lambda_1(g)$, and $\|W_\lambda\|_{C^2(\bar{\Omega})} \rightarrow 0$ as $\lambda \searrow 0$.

The case of $\int_{\partial\Omega} f = 0$ must be treated delicately. For instance, let f satisfy (1.8) additionally. Then, we obtain assertion (i) of Theorem 2.13 with $\bar{\lambda} = 0$, which is derived from the argument in the first paragraph of the proof of Lemma 3.1 below. Consequently, we obtain conclusion (iv) of the theorem because (1.10) does not have any positive solution for $\lambda = 0$ under $\int_{\partial\Omega} f = 0$. In case $\int_{\partial\Omega} f = 0$, it is an open question to solve how W_λ behaves as $\lambda \searrow 0$. We cannot yet exclude the possibility that $\|W_\lambda\|_{C(\bar{\Omega})} \rightarrow \infty$ as $\lambda \searrow 0$.

3. PROOF OF THEOREM 1.2

In this section, we consider (1.5) with $g = g_\delta$ introduced by (1.9), and prove Theorem 1.2. Note that $\int_{\partial\Omega} g_\delta < 0$.

3.1. Vanishing positive solutions. Let $g_\delta \in C^{1+\theta}(\partial\Omega)$, $\delta > \delta_0$, satisfy (1.9). We then deduce

$$\lambda_1(g_\delta) \longrightarrow 0 \quad \text{as } \delta \searrow \delta_0. \quad (3.1)$$

Indeed, (3.1) is verified in a similar manner as in the proof of [28, Lemma 6.6], using the condition that $\int_{\partial\Omega} g_\delta \nearrow \int_{\partial\Omega} g_{\delta_0} = 0$ as $\delta \searrow \delta_0$. Letting $\varphi_1(g_\delta)$ be the positive eigenfunction associated with $\lambda_1(g_\delta)$ such that $\|\varphi_1(g_\delta)\| = 1$, we additionally derive from (3.1) that

$$\int_{\Omega} |\nabla \varphi_1(g_\delta)|^2 = \lambda_1(g_\delta) \int_{\partial\Omega} g_\delta(x) \varphi_1(g_\delta)^2 \longrightarrow 0 \quad \text{as } \delta \searrow \delta_0.$$

This implies that $\varphi_1(g_\delta)$ converges to a constant $c > 0$ in $H^1(\Omega)$ as $\delta \searrow \delta_0$, and by elliptic regularity,

$$\varphi_1(g_\delta) \longrightarrow c \quad \text{in } C(\overline{\Omega}) \quad \text{as } \delta \searrow \delta_0. \quad (3.2)$$

The next lemma asserts that the positive solutions of (1.5) with $g = g_\delta$ vanish uniformly in $\lambda \in [0, \lambda_1(g_\delta))$ as $\delta \searrow \delta_0$.

Lemma 3.1. *Setting*

$$C_\delta := \sup\{\|w\|_{C(\overline{\Omega})} : w \text{ is a positive solution of (1.5)} \\ \text{with } g = g_\delta \text{ for } \lambda \in [0, \lambda_1(g_\delta))\},$$

we obtain that $C_\delta \rightarrow 0$ as $\delta \searrow \delta_0$.

Proof. Let $\bar{\delta} > \delta_0$ be fixed as close to δ_0 . First, we prove

$$\sup_{\delta \in (\delta_0, \bar{\delta})} C_\delta < \infty. \quad (3.3)$$

If not, then there exist $\delta_n \in (\delta_0, \bar{\delta})$ and positive solutions w_n of (1.5) with $g = g_{\delta_n}$ for $\lambda_n \in [0, \lambda_1(g_{\delta_n}))$ such that $\|w_n\|_{C(\overline{\Omega})} \rightarrow \infty$. Since w_n is harmonic in Ω , and $g_{\delta_n} = g^+ - \delta_n g^-$, we see

$$\|w_n\|_{C(\overline{\Omega})} = \max_{\partial\Omega} w_n = w_n(x_n), \quad x_n \in \Gamma_+. \quad (3.4)$$

Here, we have used the fact that $\frac{\partial w_n}{\partial \nu}(x_n) > 0$ by applying the SMP and BPL. Since Γ_+ is compact, up to a subsequence, $x_n \rightarrow x_\infty$ for some $x_\infty \in \Gamma_+$, as well as x_∞ is an interior point of Γ_+ . This leads us to a contradiction using the blow up argument as in the proof of Lemma 3.4 by Kim, Liang, and Shi [18, Sect. 5]. Assertion (3.3) is thus proved. Immediately, we have (3.3) with $\|\cdot\|_{C(\overline{\Omega})}$ replaced by $\|\cdot\|$ because of (3.1).

We then consider $\delta \searrow \delta_0$, and take a positive solution w of (1.5) with $g = g_\delta$ for $\lambda \in [0, \lambda_1(g_\delta))$. The boundedness of w in $H^1(\Omega)$ infers that there exists $\delta_n \searrow \delta_0$ and $\lambda_n \in [0, \lambda_1(g_{\delta_n}))$ such that for some $w_0 \in H^1(\Omega)$, $w_n \rightharpoonup w_0$, and $w_n \rightarrow w_0$ in $L^2(\Omega)$ and $L^{p+1}(\partial\Omega)$. We claim that $w_n \rightarrow 0$ in $H^1(\Omega)$. From the definition of w_n , we deduce

$$\int_{\Omega} \nabla w_n \nabla \psi = \lambda_n \int_{\partial\Omega} g_{\delta_n}(x) w_n \psi + \int_{\partial\Omega} g_{\delta_n}(x) w_n^p \psi, \quad \psi \in H^1(\Omega),$$

thus, because of (3.1), passing to the limit yields

$$\int_{\Omega} \nabla w_0 \nabla \psi = \int_{\partial\Omega} g_{\delta_0}(x) w_0^p \psi.$$

This implies that w_0 is a nonnegative weak solution of (1.5) with $g = g_{\delta_0}$ for $\lambda = 0$. Therefore, $w_0 \equiv 0$ because $\int_{\partial\Omega} g_{\delta_0} = 0$; thus, $w_n \rightarrow 0$ in $L^{p+1}(\partial\Omega)$. Finally, we deduce

$$\int_{\Omega} |\nabla w_n|^2 = \lambda_n \int_{\partial\Omega} g_{\delta_0} w_n^2 + \int_{\partial\Omega} g_{\delta_n}(x) w_n^{p+1} \longrightarrow 0,$$

as desired.

By using the bootstrap argument with elliptic regularity, as in the proof of [29, Theorem 2.2], we then deduce that $\|w_n\|_{W^{1,r}(\Omega)} \rightarrow 0$ for $r > N$. Sobolev's embedding theorem shows that $\|w_n\|_{C(\overline{\Omega})} \rightarrow 0$, as desired. \square

Remark 3.2. A positive solution of (1.5) with $g = g_\delta$ is unique for λ less than but close to $\lambda_1(g_\delta)$. This is verified by the combination of Proposition 2.7, the fact that the positive constant C_δ introduced by Lemma 3.1 is finite for a fixed $\delta > \delta_0$, and a unique local bifurcation curve of positive solutions from the simple eigenvalue $\lambda_1(g_\delta)$. It should be noted that this assertion is unconditional for $\delta > \delta_0$.

3.2. Our strategy. Let w be a positive solution of (1.5) with $g = g_\delta$ for $\lambda \in [0, \lambda_1(g_\delta))$. Proposition 2.8 tells us that w is unstable, meaning that the smallest eigenvalue of (2.4) with $g = g_\delta$ for (λ, w) is negative. We then look for a certain condition of δ under which it does *not* have a zero eigenvalue. More precisely, for the application of the implicit function theorem (IFT) to all the (λ, w) , we find such δ that is uniform in (λ, w) , based on the Fredholm alternative. Thus, we deduce assertions (i) to (iii) in Theorem 1.2, using the existence assertion of Theorem 1.1 and the uniqueness assertion of Remark 3.2.

We then present our strategy more precisely in the general setting of (1.5). We consider the eigenvalue problem

$$\begin{cases} -\Delta\phi = 0 & \text{in } \Omega, \\ \frac{\partial\phi}{\partial\nu} = \lambda g(x)\phi + \mu g(x)w^{p-1}\phi & \text{on } \partial\Omega, \end{cases} \quad (3.5)$$

where w is a positive solution of (1.5) for $\lambda \geq 0$. We deduce that (2.4) has a zero eigenvalue if and only if $\mu = p$ is an eigenvalue of (3.5). Therefore, let us study the distribution of the eigenvalues of (3.5). As in (1.3), an eigenvalue of (3.5) is called principal if the eigenfunctions associated with it have constant sign. We observe that (3.5) possesses exactly two principal eigenvalues $\mu_1^\pm \in \mathbb{R}$, which are both simple, satisfying

$$\begin{cases} \mu_1^- = 0 < 1 = \mu_1^+, & \lambda = 0, \\ \mu_1^- < 0 < 1 = \mu_1^+, & \lambda \in (0, \lambda_1(g)). \end{cases}$$

Indeed, $\mu = 1$ is always a principal eigenvalue of (3.5) with the positive eigenfunction $\phi = w$ if $\lambda \in [0, \lambda_1(g))$. It is clear that $\mu_1^- = 0$ if $\lambda = 0$. From Lemma 2.1, we infer that $\mu_1^- < 0$. Moreover, by a similar argument as in [10], we obtain that an eigenvalue μ larger than μ_1^+ is not principal. From these observations, we deduce that if $p < \mu_2^+$ for a second positive eigenvalue μ_2^+ of (3.5), then (2.4) has no zero eigenvalue, which is our desired situation.

3.3. Analysis of the second eigenvalue. We complete the proof of Theorem 1.2 by proving the following result.

Proposition 3.3. *Let $g_\delta \in C^{1+\theta}(\partial\Omega)$ be introduced by (1.9). For $\delta > \delta_0$, let $m_\delta \geq 1$ be given by the formula*

$$m_\delta = \inf\{\mu_2^+ : \mu_2^+ \text{ is the second positive eigenvalue of (3.5)}$$

with a positive solution w of (1.5) with $g = g_\delta$ for $\lambda \in [0, \lambda_1(g_\delta))\}$.

Then, $m_\delta \rightarrow \infty$ as $\delta \searrow \delta_0$.

Proof. We call ϕ_2 the eigenfunction associated with μ_2^+ and satisfying $\|\phi_2\| = 1$, and ϕ_2 changes sign. Assume by contradiction that $m_\delta < C$ as $\delta \searrow \delta_0$. We then obtain a sequence $\{(\delta_n, \lambda_n, w_n)\}$ such that $\delta_n \searrow \delta_0$, $\lambda_n \in [0, \lambda_1(g_{\delta_n}))$, $\mu_{2,n}^+ := \mu_2^+(\lambda_n, w_n) < C$, and w_n is a positive solution of (1.5) with $g = g_{\delta_n}$ for $\lambda_n \in [0, \lambda_1(g_{\delta_n}))$. Thus, $\|w_n\|_{C(\bar{\Omega})} \leq C_{\delta_n}$ using C_{δ_n} given by Lemma 3.1 with $\delta = \delta_n$. We then deduce

$$\begin{aligned} \int_{\Omega} |\nabla \phi_{2,n}|^2 &= \lambda_n \int_{\partial\Omega} g_n(x) \phi_{2,n}^2 + \mu_{2,n}^+ \int_{\partial\Omega} g_n(x) w_n^{p-1} \phi_{2,n}^2 \\ &\leq \lambda_n \int_{\partial\Omega} g_n(x) \phi_{2,n}^2 + CC_{\delta_n}^{p-1} \int_{\partial\Omega} g_n^+(x) \phi_{2,n}^2, \end{aligned}$$

where $\phi_{2,n} = \phi_2(\mu_{2,n}^+)$ and $g_n = g_{\delta_n}$. By Lemma 3.1 and (3.1), passing to the limit shows that $\int_{\Omega} |\nabla \phi_{2,n}|^2 \rightarrow 0$. Since $\|\phi_{2,n}\| = 1$, we infer that up to a subsequence and for a constant $c_\infty \neq 0$,

$$\begin{aligned} \phi_{2,n} &\rightarrow c_\infty \quad \text{in } H^1(\Omega) \quad (\text{implying that this is the case in } L^{p+1}(\partial\Omega)), \\ \phi_{2,n} &\rightarrow c_\infty \quad \text{a.e. in } \Omega \text{ and } \partial\Omega. \end{aligned}$$

We then deduce that

$$\phi_{2,n} \rightarrow c_\infty \quad \text{in } C(\bar{\Omega}). \quad (3.6)$$

Once this is done, we obtain that $\phi_{2,n}$ has constant sign if n is sufficiently large, which is the desired contradiction. Let us show how to deduce (3.6). Since c_∞ is a constant, we infer

$$\begin{cases} (-\Delta + 1)(\phi_{2,n} - c_\infty) = \phi_{2,n} - c_\infty & \text{in } \Omega, \\ \frac{\partial}{\partial \nu}(\phi_{2,n} - c_\infty) = \lambda_n g_n(x) \phi_{2,n} + \mu_{2,n}^+ g_n(x) w_n^{p-1} \phi_{2,n} & \text{on } \partial\Omega. \end{cases} \quad (3.7)$$

It should be noted that $\|\phi_{2,n}\|_{C(\bar{\Omega})}$ is bounded, based on the bootstrap argument used in the proof of Lemma 3.1. Thus, Lebesgue's dominated convergence theorem applies, and for a fixed $r > N$, we deduce

$$\phi_{2,n} - c_\infty \rightarrow 0 \quad \text{in } L^r(\Omega).$$

Similarly, using Lemma 3.1 and the assumption that $\lambda_n \rightarrow 0$ and $\mu_{2,n}^+$ is bounded, we deduce that

$$\lambda_n g_n(x) \phi_{2,n} + \mu_{2,n}^+ g_n(x) w_n^{p-1} \phi_{2,n} \rightarrow 0 \quad \text{in } L^r(\partial\Omega).$$

The $W^{1,r}$ -estimate ([3, (3.3)Proposition]) applies to (3.7), and we infer that

$$\phi_{2,n} - c_\infty \rightarrow 0 \quad \text{in } W^{1,r}(\Omega),$$

from which (3.6) follows by Sobolev's embedding theorem. \square

End of proof of Theorem 1.2. From Proposition 3.3, we find that given $p > 1$ satisfying (1.2), it is possible to choose $\bar{\delta} > \delta_0$ such that if $\delta \in (\delta_0, \bar{\delta})$, then $p < \mu_2^+$ for a positive solution w of (1.5) with $\lambda \in [0, \lambda_1(g_\delta)]$. The proof of Theorem 1.2 is now complete. \square

Remark 3.4. A similar strategy as in the proof of Theorem 1.2 was applied in [17] to an indefinite sublinear elliptic equation with a Robin boundary condition.

We then use (1.4) to convert Theorem 1.2 to (1.1), which is our main result for (1.1).

Theorem 3.5. *Let $g_\delta \in C^{1+\theta}(\partial\Omega)$ be given by (1.9). If we take δ sufficiently close to δ_0 , then the positive solution set of (1.5) with $g = g_\delta$ for $\lambda \in (0, \lambda_1(g_\delta))$ is described as follows (see Figure 1):*

- (i) (1.5) with $g = g_\delta$ possesses a unique positive solution v_λ for each $\lambda \in (0, \lambda_1(g_\delta))$, and the positive solution set $\{(\lambda, v_\lambda) : \lambda \in (0, \lambda_1(g_\delta))\}$ is represented by a smooth curve.
- (ii) $v_\lambda \rightarrow 0$ in $C^2(\bar{\Omega})$ as $\lambda \nearrow \lambda_1(g_\delta)$, i.e., bifurcation from $\{(\lambda, 0)\}$ at $(\lambda_1(g_\delta), 0)$ occurs subcritically.
- (iii) $\lambda^{\frac{1}{p-1}} v_\lambda \rightarrow w_0$ in $C^2(\bar{\Omega})$ as $\lambda \searrow 0$ for some $w_0 \in C^2(\bar{\Omega})$, where w_0 is a unique positive solution of (1.5) with $g = g_\delta$ for $\lambda = 0$.

From Theorem 2.13 (iv), similar results as in Theorem 1.2 are deduced for (1.10). Using (1.7), we introduce for a sign changing function $f \in C^{1+\theta}(\partial\Omega)$ the condition

$$\partial\Omega = \Gamma_+(f) \cup \Gamma_-(f) \quad \text{and} \quad \overline{\Gamma_+(f)} \cap \overline{\Gamma_-(f)} = \emptyset. \quad (3.8)$$

Then, we have the following:

Theorem 3.6. *Let $f \in C^{1+\theta}(\partial\Omega)$ be a sign changing function that satisfies (3.8) and the condition $\int_{\partial\Omega} f > 0$. Let $g_\delta \in C^{1+\theta}(\partial\Omega)$ be given by (1.9). If δ is sufficiently close to δ_0 , then the positive solution set of (1.10) with $g = g_\delta$ for $\lambda \in (0, \lambda_1(g_\delta))$ is given as follows (see Figure 4):*

- (i) (1.10) with $g = g_\delta$ possesses a unique positive solution w_λ for every $\lambda \in (0, \lambda_1(g_\delta))$, and the positive solution set $\{(\lambda, w_\lambda) : \lambda \in (0, \lambda_1(g_\delta))\}$ is represented by a smooth curve,
- (ii) $w_\lambda \rightarrow 0$ in $C^2(\bar{\Omega})$ as $\lambda \nearrow \lambda_1(g_\delta)$ and $\lambda \searrow 0$.

Proof. Because of (3.2), we find that the condition $\int_{\partial\Omega} f > 0$ is sufficient for getting $\int_{\partial\Omega} f(x)\varphi_1(g_\delta)^{p+1} > 0$ for δ close to δ_0 , thus ensuring the existence of the positive solution W_λ of (1.10) with $g = g_\delta$ for $\lambda \in (0, \lambda_1(g_\delta))$ by Theorem 2.13.

Assertion (ii) is a direct consequence of assertions (ii) and (iv) of Theorem 2.13.

To verify assertion (i), we show that the assertion similar to (3.4) holds. We can deduce that $x_n \in \Gamma_+(f)$ for n large enough where $w_n(x_n) = \max_{\partial\Omega} w_n \rightarrow \infty$ as in (3.4), which is essential for our procedure. It should be noted that $\frac{\partial w_n}{\partial \nu}(x_n) > 0$ by the SMP and BPL. We observe from (3.8) that if we assume $x_n \in \Gamma_-(f)$ by contradiction, then $f(x_n) < -c_0$ for some $c_0 > 0$, thus, for n that is large enough,

$$\frac{\partial w_n}{\partial \nu}(x_n) = (\lambda_n g_{\delta_n}(x_n) + f(x_n)(w_n(x_n))^{p-1})w_n(x_n) < 0,$$

which is a contradiction. The rest of the proof of assertion (i) follows the same line of argument in this section. \square




FIGURE 4. Positive solution set $\{(\lambda, w_\lambda)\}$ of (1.10) with $g = g_\delta$ as a smooth curve in the case that δ is close to δ_0 .

4. APPLICATIONS TO INDEFINITE LOGISTIC BOUNDARY CONDITIONS

Let Ω be a bounded domain of \mathbb{R}^N , $N \geq 1$, with smooth boundary $\partial\Omega$. Consider nonnegative solutions of the problem

$$\begin{cases} -\Delta u = 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = \lambda r(x)u(1-u) & \text{on } \partial\Omega, \end{cases} \quad (4.1)$$

where $r \in C^{1+\theta}(\partial\Omega)$, $\theta \in (0, 1)$, changes sign, $\lambda > 0$ is a parameter, and ν is the unit outer normal to $\partial\Omega$. A motivation for our study of (4.1) arises in population genetics ([15, 9]). For previous works on the boundary version, we refer to [21, 22, 18, 23]. Clearly, $u \equiv 0, 1$ satisfies (4.1) for all $\lambda > 0$, which are called *constant solutions*, and

$$\begin{aligned} \{(\lambda, 0)\} &:= \{(\lambda, u) : \lambda > 0, u \equiv 0\}, \\ \{(\lambda, 1)\} &:= \{(\lambda, u) : \lambda > 0, u \equiv 1\} \end{aligned}$$

are said to be the *constant lines*. A positive solution $u \in C^{2+\alpha}(\overline{\Omega})$, $\alpha \in (0, 1)$, of (4.1) is defined similarly. We may regard (λ, u) as a positive solution of (4.1). It should be noted that $(\lambda, 1)$ is a (constant) positive solution.

In this section, we discuss the existence and uniqueness of *nonconstant* positive solutions u of (4.1), which have been well studied for the case that $u \leq 1$ in $\overline{\Omega}$ ([21, 22, 18, 23]). In turn, our objective is to discuss the case of $u \not\leq 1$ in $\overline{\Omega}$, i.e., the case that $u > 1$ somewhere in $\overline{\Omega}$. Such positive solutions are called *large positive solutions*. Similar studies on large positive solutions are [19, 14, 1, 27], where the logistic type equation $-\Delta u = \lambda r(x)(u - u^p)$ in Ω , $p > 1$, is considered under linear Dirichlet, Neumann or Robin boundary conditions. For $\lambda < 0$, (λ, u) is a positive solution of (4.1) if and only if $(-\lambda, u)$ is a positive solution of (4.1) with r replaced by $-r$, because of the symmetry $\lambda r(x) = (-\lambda)(-r(x))$. For $\lambda = 0$, it is clear that $\{(0, c)\} := \{(\lambda, u) : \lambda = 0, u \equiv c > 0\}$ is the positive solution set.

4.1. Known results for positive solutions $u \leq 1$. In this subsection, we focus our consideration on nonnegative solutions u of (4.1) such that $u \leq 1$ in $\overline{\Omega}$, and summarize known results for them from [22, 23], as illustrated by Figure 5. By applying the SMP and BPL, a nonconstant positive solution u implies that $0 < u < 1$ in $\overline{\Omega}$.

When $\int_{\partial\Omega} r \neq 0$, in view of positive solutions $u(x) \in (0, 1)$ in $\overline{\Omega}$, it suffices to consider the case of $\int_{\partial\Omega} r < 0$ because all the results obtained for this case can be converted automatically into the case of $\int_{\partial\Omega} r > 0$. Indeed, using the change $U = 1 - u$, we transform (4.1) into the problem

$$\begin{cases} -\Delta U = 0 & \text{in } \Omega, \\ \frac{\partial U}{\partial \nu} = \lambda(-r(x))U(1 - U) & \text{on } \partial\Omega. \end{cases}$$

Let us now assume that $\int_{\partial\Omega} r < 0$. In terms of bifurcation from the constant line $\{(\lambda, 0)\}$, an important role is played by the positive principal eigenvalue $\lambda_1(r) > 0$ of (1.3) with $g = r$. The local bifurcation theory from simple eigenvalues by Crandall and Rabinowitz [13] shows that a nonconstant positive solution u of (4.1) bifurcates from $\{(\lambda, 0)\}$ uniquely at $(\lambda_1(r), 0)$, which is *supercritical*, i.e., in the direction $\lambda > \lambda_1(r)$ (see Figure 5 (i)). Then, by employing the IFT, $u = u_\lambda$ is parametrized smoothly by $\lambda \in (\lambda_1(r), \infty)$ ([22, Theorems 2.5 and 2.8]). Since the bifurcating positive solution curve at $(\lambda_1(r), 0)$ is unique, we see that (λ, u_λ) , $\lambda \in (\lambda_1(r), \infty)$, is a unique nonconstant positive solution of (4.1) ([22, Theorem 2.7]). For $\lambda \leq \lambda_1(r)$, (4.1) has no nonconstant positive solution ([22, Theorem 2.6]). For the stability of the nonnegative solutions, $(\lambda, 0)$ is asymptotically stable and unstable for $\lambda \leq \lambda_1(r)$ and $\lambda > \lambda_1(r)$, respectively, $(\lambda, 1)$ is unstable for $\lambda > 0$, and (λ, u_λ) is asymptotically stable for $\lambda > \lambda_1(r)$ ([22, Theorem 3.1, 3.2 and 3.3]).

When $\int_{\partial\Omega} r = 0$, it is proved by a bifurcation approach from $\{(0, c)\}$ that (4.1) has a unique nonconstant positive solution u_λ for each $\lambda > 0$, which bifurcates from $\{(0, c)\}$ at $c = \frac{1}{2}$ and is parametrized smoothly by $\lambda \in (0, \infty)$ ([23, Theorem 1.1, Lemma 2.2]). For the stability of the nonnegative

solutions, $(\lambda, 0)$ and $(\lambda, 1)$ are both unstable for $\lambda > 0$, whereas (λ, u_λ) is asymptotically stable for $\lambda > 0$ ([23, Theorem 1.2]).




FIGURE 5. Cases (i) $\int_{\partial\Omega} r < 0$, (ii) $\int_{\partial\Omega} r > 0$, and (iii) $\int_{\partial\Omega} r = 0$.

To conclude this subsection, let us demonstrate our developing scenario presented in the next subsection for large positive solutions of (4.1). When $\int_{\partial\Omega} r > 0$, using the change $v = u - 1$, another bifurcation approach from the constant line $\{(\lambda, 1)\}$ can be employed. We then transform (4.1) into (1.1) with $g = -r$ and $p = 2$:

$$\begin{cases} -\Delta v = 0 & \text{in } \Omega, \\ \frac{\partial v}{\partial \nu} = \lambda(-r(x))(v + v^2) & \text{on } \partial\Omega. \end{cases} \quad (4.2)$$

The local bifurcation theory also applies to (4.2) at $(\lambda_1(-r), 0)$, and (4.2) possesses a smooth solution curve $\{(\lambda(\sigma), v(\sigma)) : \sigma \in (-\bar{\sigma}, \bar{\sigma})\}$, $\bar{\sigma} > 0$, which uniquely bifurcates at $(\lambda_1(-r), 0)$, such that $(\lambda(0), v(0)) = (\lambda_1(-r), 0)$, $\lambda(\sigma) > \lambda_1(-r)$ and $v(\sigma) < 0$ in $\bar{\Omega}$ for $\sigma \in (-\bar{\sigma}, 0)$, and $\lambda(\sigma) < \lambda_1(-r)$ and $v(\sigma) > 0$ in $\bar{\Omega}$ for $\sigma \in (0, \bar{\sigma})$. It should be noted that the curve $\{(\lambda(\sigma), v(\sigma)) : \sigma \in (-\bar{\sigma}, 0)\}$ corresponds to positive solutions of (4.1) satisfying $u < 1$ in $\bar{\Omega}$, whereas the curve $\{(\lambda(\sigma), v(\sigma)) : \sigma \in (0, \bar{\sigma})\}$ corresponds to large positive solutions of (4.1). The latter result will be strengthened by Theorems 2.12 and 3.5.

4.2. Main results for large positive solutions $u \not\leq 1$. In this subsection, we consider large positive solutions of (4.1). The large positive solution u is divided into the two cases:

- (a) $u \geq 1$ in $\overline{\Omega}$, and
- (b) $u \not\geq 1$ in $\overline{\Omega}$ ($u < 1$ somewhere in $\overline{\Omega}$).

Here, case (a) implies that $u > 1$ in $\overline{\Omega}$, i.e., (λ, v) with $v = u - 1$ is a positive solution of (4.2), whereas case (b) implies that (λ, v) is a sign changing solution of (4.2).

When $\int_{\partial\Omega} r > 0$, the positive solution curve $\{(\lambda(\sigma), v(\sigma)) : \sigma \in (0, \overline{\sigma})\}$ of (4.2) added to $(\lambda_1(-r), 0)$ is extended globally in $\mathbb{R} \times C^2(\overline{\Omega})$ as a subcontinuum \mathcal{C}_0 , using the global bifurcation result ([20, Theorem 6.4.3], [33, Theorem 1.1]). Indeed, $\mathcal{C}_0 \setminus \{(\lambda_1(-r), 0)\}$ does not meet any point on $\{(\lambda, 0)\}$ by the uniqueness of $(\lambda_1(-r), 0)$. Moreover, equipped with $g = -r$, Lemma 2.7 and Proposition 2.11 show that

$$\{\lambda \geq 0 : (\lambda, v) \in \mathcal{C}_0 \setminus \{(\lambda_1(-r), 0)\}\} \subset (0, \lambda_1(-r)),$$

implying that \mathcal{C}_0 is unbounded, and bifurcates from infinity at some $\lambda \in [0, \lambda_1(-r)]$. We then deduce that (4.1) possesses the unbounded subcontinuum

$$\tilde{\mathcal{C}}_0 = \{(\lambda, u) : u = 1 + v, (\lambda, v) \in \mathcal{C}_0\}$$

for which $\tilde{\mathcal{C}}_0$ bifurcates at $(\lambda_1(-r), 1)$, and $\tilde{\mathcal{C}}_0 \setminus \{(\lambda_1(-r), 1)\}$ consists of large positive solutions with condition (a). One of our aims is to strengthen this global bifurcation result by Theorems 2.12 and 3.5.

We then present our main results for large positive solutions of (4.1) in the case of $\int_{\partial\Omega} r > 0$, where (1.2) is assumed with $p = 2$, i.e., $N = 1, 2, 3$.

Theorem 4.1. *Suppose that $\int_{\partial\Omega} r > 0$. Then, (4.1) possesses at least one large positive solution $u_\lambda \in C^2(\overline{\Omega})$ with condition (a) for each $\lambda \in (0, \lambda_1(-r))$. Moreover, there is no large positive solution of (4.1) with condition (a) for any $\lambda \geq \lambda_1(-r)$ nor with condition (b) for any $\lambda \in (0, \lambda_1(-r)]$. Additionally, the following five assertions hold:*

- (i) u_λ is unstable.
- (ii) $\|u_\lambda\|_{C(\overline{\Omega})} \rightarrow \infty$ as $\lambda \searrow 0$. More precisely, there exists $0 < c_1 < c_2$ such that
$$c_1 \lambda^{-1} \leq \|u_\lambda\|_{C(\overline{\Omega})} \leq c_2 \lambda^{-1} \quad \text{as } \lambda \searrow 0. \quad (4.3)$$
- (iii) $\sup_{\lambda \in (\underline{\lambda}, \lambda_1(-r))} \|u_\lambda\|_{C(\overline{\Omega})} < \infty$ for any $\underline{\lambda} \in (0, \lambda_1(-r))$.
- (iv) $\|u_\lambda - 1\|_{C^2(\overline{\Omega})} \rightarrow 0$ as $\lambda \nearrow \lambda_1(-r)$. Particularly, (λ, u_λ) is the bifurcating positive solution on $\tilde{\mathcal{C}}_0$ for λ that is close to $\lambda_1(-r)$.
- (v) $\inf_{\lambda \in (0, \overline{\lambda})} \|u_\lambda - 1\|_{C(\overline{\Omega})} > 0$ for any $\overline{\lambda} \in (0, \lambda_1(-r))$.

The next theorem is a direct consequence of Theorems 3.5 and 4.1.

Theorem 4.2. *Let $r \in C^{1+\theta}(\partial\Omega)$ be under (1.8) with $g = -r$, and let $r_\delta = \delta r^+ - r^-$, $\delta > \delta_0 := \frac{\int_{\partial\Omega} r^-}{\int_{\partial\Omega} r^+}$, be a function in $C^{1+\theta}(\partial\Omega)$. Then, the large*

positive solution (λ, u_λ) of (4.1) with $r = r_\delta$, given by Theorem 4.1, satisfies the following three properties, provided that δ is sufficiently close to δ_0 (see Figure 6):

- (i) u_λ is a unique large positive solution (and consequently a unique nonconstant positive solution), and the positive solution set $\{(\lambda, u_\lambda) : \lambda \in (0, \lambda_1(-r_\delta))\}$ is represented by a smooth curve.
- (ii) $u_\lambda \rightarrow 1$ in $C^2(\overline{\Omega})$ as $\lambda \nearrow \lambda_1(-r_\delta)$.
- (iii) $\lambda u_\lambda \rightarrow w_0$ in $C^2(\overline{\Omega})$ as $\lambda \searrow 0$ for some $w_0 \in C^2(\overline{\Omega})$, where w_0 is a unique positive solution of (1.5) with $g = -r_\delta$ and $p = 2$ for $\lambda = 0$.

Remark 4.3. Without any smallness condition on $\delta - \delta_0$, (4.1) with $r = r_\delta$ has a unique nonconstant positive solution u_λ for λ around, but not equal to $\lambda_1(-r_\delta)$, satisfying $0 < u_\lambda < 1$ in $\overline{\Omega}$ for $\lambda > \lambda_1(-r_\delta)$ and $u_\lambda > 1$ in $\overline{\Omega}$ for $\lambda < \lambda_1(-r_\delta)$. Moreover, there is no nonconstant positive solution of (4.1) for $\lambda = \lambda_1(-r_\delta)$ (cf. Remark 3.2).




FIGURE 6. Positive solution curve of (4.1) for $r = r_\delta$ with δ larger than but close to δ_0 .

When $\int_{\partial\Omega} r < 0$, motivated by (4.3), the following asymptotics condition for positive solutions of (4.1) is introduced when $\lambda \searrow 0$: there exists $c_0 > 0$ such that

$$\|u_n\|_{C(\overline{\Omega})} \leq c_0 \lambda_n^{-1}, \quad n \rightarrow \infty, \quad (4.4)$$

for a sequence $\{(\lambda_n, u_n)\}$ of positive solutions of (4.1) that satisfies $\|u_n\|_{C(\overline{\Omega})} \rightarrow \infty$ as $\lambda_n \searrow 0$.

We then state our main result for the case of $\int_{\partial\Omega} r \leq 0$.

Theorem 4.4. *Let $N \geq 1$ be arbitrary. Assume that $\int_{\partial\Omega} r \leq 0$. Then, (4.1) has no large positive solution with condition (a) for any $\lambda > 0$. If $\int_{\partial\Omega} r < 0$, then additionally assuming $N = 1, 2, 3$ and (4.4) provides that $(\lambda, 1)$ is the only positive solution of (4.1) for small $\lambda > 0$.*

Remark 4.5. To obtain (4.4), it is sufficient that (1.8) with $g = -r$ holds, which follows from the same argument as in the proof of Lemma 3.1.

4.3. Proof of Theorem 4.1. The assertions in Theorem 4.1 are deduced directly from Theorem 2.12, except for the nonexistence assertion for the large positive solutions with condition (b). We verify it using the sub and supersolution method.

Let u be a large positive solution with condition (b) for some $\lambda \in (0, \lambda_1(-r)]$. Note that $u - 1$ changes sign. We then set $\underline{u} := \min(u, 1)$, and \underline{u} is a supersolution of (4.1) such that $0 < \underline{u} \leq 1$ and $\underline{u} \not\equiv 1$ in $\overline{\Omega}$. Note that $u \equiv 0$ is an unstable constant solution of (4.1) for every $\lambda > 0$, thus, we can construct a positive subsolution of (4.1) smaller than \underline{u} in $\overline{\Omega}$. By the sub and supersolution method, we obtain a nonconstant positive solution u_1 of (4.1) such that $u_1 \leq 1$ in $\overline{\Omega}$, which is contradictory for [22, Theorem 2.6]. The proof of Theorem 4.1 is complete. \square

4.4. Proof of Theorem 4.4. The first assertion comes from Proposition 2.7 with $g = -r$. We next prove the second assertion. It suffices to verify that (4.1) has no large positive solution with condition (b) for $\lambda > 0$ small. Assume by contradiction that (4.1) has positive solutions $\{(\lambda_n, u_n)\}$ with $\lambda_n \searrow 0$ such that $u_n - 1$ changes sign, and we deduce that $\|u_n\|_{C(\overline{\Omega})} \rightarrow \infty$. Indeed, if $\|u_n\|_{C(\overline{\Omega})}$ is bounded, then $\|u_n\|$ is also bounded, and so is $\|u_n\|_{C^\alpha(\overline{\Omega})}$ by elliptic regularity. Hence, up to a subsequence, $u_n \rightarrow u_\infty$ in $C(\overline{\Omega})$. We then infer that $u_\infty \equiv 1$ because $u_n - 1$ changes sign and u_∞ is a constant. However, this is contradictory, because we can show that $\{(\lambda, 1)\}$ is a unique bifurcation curve from $\{(0, c)\}$ at $(\lambda, u) = (0, 1)$, as in the proof of [16, Proposition 3.1].

From (4.4), we obtain that $\|\lambda_n u_n\|_{C(\overline{\Omega})} \leq c_0$. Say $v_n = \lambda_n u_n$, and (λ_n, v_n) is a positive solution of the problem:

$$\begin{cases} -\Delta v = 0 & \text{in } \Omega, \\ \frac{\partial v}{\partial \nu} = \lambda r(x)v - r(x)v^2 & \text{on } \partial\Omega. \end{cases}$$

We observe that $\|v_n\|$ is bounded, thus, for some $v_\infty \in H^1(\Omega)$ and up to a subsequence, $v_n \rightharpoonup v_\infty \geq 0$, and $v_n \rightarrow v_\infty$ in $L^2(\Omega)$ and $L^3(\partial\Omega)$. The argument then proceeds by dividing it into the two cases:

(i) Case $v_\infty \not\equiv 0$ on $\partial\Omega$: By definition,

$$\int_{\Omega} \nabla v_n \nabla \varphi - \lambda_n \int_{\partial\Omega} r(x)v_n \varphi + \int_{\partial\Omega} r(x)v_n^2 \varphi = 0, \quad \varphi \in H^1(\Omega). \quad (4.5)$$

Passing to the limit, we deduce

$$\int_{\Omega} \nabla v_\infty \nabla \varphi + \int_{\partial\Omega} r(x)v_\infty^2 \varphi = 0,$$

thus, $v_\infty > 0$ in $\overline{\Omega}$ by the SMP and BPL, which is contradictory for assumption $\int_{\partial\Omega} r < 0$.

(ii) Case $v_\infty \equiv 0$ on $\partial\Omega$: We then deduce that $v_n \rightarrow 0$ in $H^1(\Omega)$ because (4.5) with $\varphi = v_n$ implies

$$\int_{\Omega} |\nabla v_n|^2 = \lambda_n \int_{\partial\Omega} r(x)v_n^2 - \int_{\partial\Omega} r(x)v_n^3 \rightarrow 0.$$

Say $\psi_n = \frac{v_n}{\|v_n\|}$, and $\|\psi_n\| = 1$. Up to a subsequence and for some $\psi_\infty \in H^1(\Omega)$, $\psi_n \rightharpoonup \psi_\infty$, and $\psi_n \rightarrow \psi_\infty$ in $L^2(\Omega)$ and $L^3(\partial\Omega)$. Since

$$\int_{\Omega} |\nabla \psi_n|^2 = \lambda_n \int_{\partial\Omega} r\psi_n^2 - \|v_n\| \int_{\partial\Omega} r\psi_n^3 \rightarrow 0,$$

we infer that $\psi_n \rightarrow \psi_\infty$ in $H^1(\Omega)$, and ψ_∞ is a positive constant. From (4.5) with $\varphi = 1$, we infer that

$$\int_{\partial\Omega} \lambda_n r(x)\psi_n = \int_{\partial\Omega} r(x)\psi_n^2 \|v_n\|.$$

Using $v_n = \lambda_n u_n$, we obtain that

$$\int_{\partial\Omega} r(x)\psi_n = \|u_n\| \int_{\partial\Omega} r\psi_n^2.$$

It should be noted that $\|u_n\| \rightarrow \infty$ from the condition $\|u_n\|_{C(\bar{\Omega})} \rightarrow \infty$ by elliptic regularity. Passing to the limit, we deduce

$$\begin{aligned} \int_{\partial\Omega} r\psi_n &\rightarrow \psi_\infty \int_{\partial\Omega} r < 0, \\ \|u_n\| \int_{\partial\Omega} r\psi_n^2 &\rightarrow -\infty, \end{aligned}$$

which is a contradiction. The proof of Theorem 4.4 is now complete. \square

4.5. Large positive solutions in the one dimensional case. In this subsection, we consider the one dimensional case of (4.1):

$$\begin{cases} -u''(x) = 0, & x \in I = (0, 1), \\ -u'(0) = \lambda r_0 u(0)(1 - u(0)), \\ u'(1) = \lambda r_1 u(1)(1 - u(1)), \end{cases} \quad (4.6)$$

where $\lambda > 0$ is a parameter, and $r_0 < 0 < r_1$. Then, the positive solutions u are determined necessarily as linear functions $u = cx + d$ with the constants c, d that satisfy $d > 0$ and $c + d > 0$.

Proposition 4.6. *Problem (4.6) does not have any large positive solution with condition (b).*

Proof. We argue by contradiction. Let u be a positive solution of (4.6) for some $\lambda > 0$ such that

$$u(x_-) := \min_{x=0,1} u(x) < 1 < \max_{x=0,1} u(x) =: u(x_+).$$

Then, it follows that $(x_-, x_+) = (0, 1), (1, 0)$. However, this is contradictory for the boundary condition. \square

Consequently, any nonconstant positive solution (λ, u) of (4.6) fulfills that either $u > 1$ in $[0, 1]$ or $0 < u < 1$ in $[0, 1]$. Combining this assertion with Theorems 4.1 and 4.2 then provides the following corollary. Set $r(0) = r_0$ and $r(1) = r_1$, and deduce that $\lambda_1(r) = 0$ if $r_0 + r_1 = 0$.

Corollary 4.7. *The following two assertions hold:*

- (i) *If $r_0 + r_1 \leq 0$, then (4.6) has a unique nonconstant positive solution u_λ for $\lambda > \lambda_1(r)$, satisfying $0 < u_\lambda < 1$ in $[0, 1]$, and there is no nonconstant positive solution of (4.6) for any $\lambda \in (0, \lambda_1(r))$.*
- (ii) *If $r_0 + r_1 > 0$, then (4.6) has a nonconstant positive solution u_λ for $\lambda \neq \lambda_1(-r)$ and no nonconstant positive solution for $\lambda = \lambda_1(-r)$ such that $0 < u_\lambda < 1$ in $[0, 1]$ for $\lambda > \lambda_1(-r)$, whereas $u_\lambda > 1$ in $[0, 1]$ for $\lambda \in (0, \lambda_1(-r))$. Moreover, u_λ is unique for $\lambda > \lambda_1(-r)$. Particularly, given $r_0 < 0$, choose $r_1 > 0$ such that $r_0 + r_1 > 0$ is sufficiently close to zero, and then, u_λ is also unique for $\lambda \in (0, \lambda_1(-r))$.*

A numerical observation of (4.6) assists the understanding of the nonconstant positive solution set. This observation suggests that a nonconstant positive solution u_λ is always unique if it exists, and one obtains the following limiting behaviors of u_λ both as $\lambda \rightarrow \infty$ and $\lambda \searrow 0$:

(i) in each case of r_0, r_1 , the limiting function of u_λ as $\lambda \rightarrow \infty$ is given by $u_\infty(x) = x$, which is consistent with [22, Theorem 4.1] and [23, Theorem 1.3];

(ii) in the case when $r_0 + r_1 > 0$, the limiting function of λu_λ as $\lambda \searrow 0$ is given by

$$w_0(x) = -\frac{\left(1 - \sqrt{\frac{-r_0}{r_1}}\right)^2}{(-r_0)}x + \frac{1 - \sqrt{\frac{-r_0}{r_1}}}{(-r_0)},$$

which is the unique positive solution of the problem

$$\begin{cases} -w'' = 0 & \text{in } (0, 1), \\ -w'(0) = -r_0 w(0)^2, \\ w'(1) = -r_1 w(1)^2, \end{cases}$$

and thus, this is consistent with Theorem 4.2 (iii).

To conclude this paper, we mention that when $N > 1$, it is an open question to solve whether (4.1) has bifurcating positive solutions from $\{(\lambda, 1)\}$ at a positive *non principal* eigenvalues of (1.3) with $g = -r$. If yes, then Proposition 2.7 tells us that these bifurcating positive solutions should be large positive solutions with condition (b). Otherwise, Corollary 4.7 could be extended to $N > 1$ in a certain class of sign changing weights r . In the case of $r < 0$ on $\partial\Omega$ (sign definite superlinear case), the existence of such a bifurcating positive solution of (4.1) from $\{(\lambda, 1)\}$ was discussed in [18].

REFERENCES

- [1] G. A. Afrouzi, K. J. Brown, K. J. *Positive mountain pass solutions for a semilinear elliptic equation with a sign-changing weight function*, *Nonlinear Anal.* **64** (2006), 409–416.
- [2] S. Alama, G. Tarantello, *On semilinear elliptic equations with indefinite nonlinearities*, *Calc. Var. Partial Differential Equations* **11** (1993), 439–475.
- [3] H. Amann, *Nonlinear elliptic equations with nonlinear boundary conditions*, *New developments in differential equations (Proc. 2nd Scheveningen Conf., Scheveningen, 1975)*, pp. 43–63, *North-Holland Math. Studies*, Vol. 21, North-Holland, Amsterdam, 1976.
- [4] H. Amann, J. López-Gómez, *A priori bounds and multiple solutions for superlinear indefinite elliptic problems*, *J. Differential Equations* **146** (1998), 336–374.
- [5] H. Berestycki, I. Capuzzo-Dolcetta, L. Nirenberg, *Superlinear indefinite elliptic problems and nonlinear Liouville theorems*, *Top. Methods Nonlinear Anal.* **4** (1994), 59–78.
- [6] H. Berestycki, I. Capuzzo-Dolcetta, L. Nirenberg, *Variational methods for indefinite superlinear homogeneous elliptic problems*, *Nonlinear Differential Equations Appl.* **2** (1995), 553–572.
- [7] D. Bonheure, J. M. Gomes, P. Habets, *Multiple positive solutions of superlinear elliptic problems with sign-changing weight*, *J. Differential Equations* **214** (2005), 36–64.
- [8] A. Boscaggin, *A note on a superlinear indefinite Neumann problem with multiple positive solutions*, *J. Math. Anal. Appl.* **377** (2011), 259–268.
- [9] K. J. Brown, P. Hess, *Stability and uniqueness of positive solutions for a semilinear elliptic boundary value problem*, *Differential Integral Equations* **3** (1990), 201–207.
- [10] K. J. Brown, S. S. Lin, *On the existence of positive eigenfunctions for an eigenvalue problem with indefinite weight function*, *J. Math. Anal. Appl.* **75** (1980), 112–120.
- [11] K. J. Brown, Y. Zhang, *The Nehari manifold for a semilinear elliptic equation with a sign-changing weight function*, *J. Differential Equations* **193** (2003), 481–499.
- [12] W. Chen, C. Li, *Indefinite elliptic problems in a domain*, *Discrete Contin. Dyn. Systems* **3** (1997), 333–340.
- [13] M. G. Crandall, P. H. Rabinowitz, *Bifurcation from simple eigenvalues*, *J. Funct. Anal.* **8** (1971), 321–340.
- [14] M. Delgado, A. Suárez, *On the existence and multiplicity of positive solutions for some indefinite nonlinear eigenvalue problem*, *Proc. Amer. Math. Soc.* **132** (2004), 1721–1728.
- [15] W. H. Fleming, *A selection-migration model in population genetics*, *J. Math. Biol.* **2** (1975), 219–233.
- [16] U. Kaufmann, H. Ramos Quoirin, K. Umezu, *Nonnegative solutions of an indefinite sublinear Robin problem I: positivity, exact multiplicity, and existence of a subcontinuum*, *Ann. Mat. Pura Appl. (4)* **199** (2020), 2015–2038.

- [17] U. Kaufmann, H. Ramos Quoirin, K. Umezu, *Nonnegative solutions of an indefinite sublinear Robin problem II: local and global exactness results*, (2020), Israel J. Math., accepted. arXiv:2001.09315
- [18] C.-G. Kim, Z.-P. Liang, J.-P. Shi, *Existence of positive solutions to a Laplace equation with nonlinear boundary condition*, Z. Angew. Math. Phys. **66** (2015), 3061–3083.
- [19] B. Ko, K. Brown, *The existence of positive solutions for a class of indefinite weight semilinear elliptic boundary value problems*, Nonlinear Anal. **39** (2000), 587–597.
- [20] J. López-Gómez, Spectral theory and nonlinear functional analysis, Research Notes in Mathematics 426, Chapman & Hall/CRC, Boca Raton, FL, 2001.
- [21] G. F. Madeira, *Existence and regularity for a nonlinear boundary flow problem of population genetics*, Nonlinear Anal. **70** (2009), 974–981.
- [22] G. F. Madeira, A. S. do Nascimento, *Bifurcation of stable equilibria and nonlinear flux boundary condition with indefinite weight*, J. Differential Equations **251** (2011), 3228–3247.
- [23] G. F. Madeira, A. S. do Nascimento, *Bifurcation of stable equilibria under nonlinear flux boundary condition with null average weight*, J. Math. Anal. Appl. **441** (2016), 121–139.
- [24] T. Ouyang, *On positive solutions of semilinear equations $\Delta u + \lambda u + hu^p = 0$ on compact manifolds, Part II*, Indiana Univ. Math. J. **40** (1991), 1083–1141.
- [25] K. Pflüger, *On indefinite nonlinear Neumann problems*, Partial differential and integral equations (Newark, DE, 1997), 335–346, Int. Soc. Anal. Appl. Comput., **2**, Kluwer Acad. Publ., Dordrecht, 1999.
- [26] M. H. Protter, H. F. Weinberger, Maximum principles in differential equations. Prentice-Hall, Inc., Englewood Cliffs, N.J. 1967.
- [27] H. Ramos Quoirin, A. Suárez, *Positive solutions for some indefinite nonlinear eigenvalue elliptic problems with Robin boundary conditions*, Nonlinear Anal. **114** (2015), 74–86.
- [28] H. Ramos Quoirin, K. Umezu, *An indefinite concave-convex equation under a Neumann boundary condition I*, Israel J. Math. **220** (2017), 103–160.
- [29] J. D. Rossi, *Elliptic problems with nonlinear boundary conditions and the Sobolev trace theorem*, Stationary partial differential equations. Vol. II, 311–406.
- [30] H. Tehrani, *On indefinite superlinear elliptic equations*, Calc. Var. Partial Differential Equations **4** (1996), 139–153.
- [31] A. Tellini, *High multiplicity of positive solutions for superlinear indefinite problems with homogeneous Neumann boundary conditions*, J. Math. Anal. Appl. **467** (2018), 673–698.
- [32] K. Umezu, *On eigenvalue problems with Robin type boundary conditions having indefinite coefficients*, Appl. Anal. **85** (2006), 1313–1325.
- [33] K. Umezu, *Global bifurcation results for semilinear elliptic boundary value problems with indefinite weights and nonlinear boundary conditions*, NoDEA Nonlinear Differential Equations Appl. **17** (2010), 323–336.
- [34] M. Zhu, *On elliptic problems with indefinite superlinear boundary conditions*, J. Differential Equations **193** (2003), 180–195.

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, IBARAKI UNIVERSITY,
MITO 310-8512, JAPAN

Email address: `kenichiro.umezu.math@vc.ibaraki.ac.jp`