THE STRUCTURE OF ALGEBRAIC BAER *-ALGEBRAS

ZSOLT SZŰCS AND BALÁZS TAKÁCS

ABSTRACT. The purpose of this note is to describe when a general complex algebraic *-algebra is pre- C^* -normed, and to investigate their structure when the *-algebras are Baer *-rings in addition to algebraicity. As a main result we prove the following theorem for complex algebraic Baer *-algebras: every *-algebra of this kind can be decomposed as a direct sum $M \oplus B$, where M is a finite dimensional Baer *-algebra and B is a commutative algebraic Baer *-algebra. The summand M is *-isomorphic to a finite direct sum of full complex matrix algebras of size at least 2×2 . The commutative summand B is *-isomorphic to the linear span of the characteristic functions of the clopen sets in a Stonean topological space.

As an application we show that a group G is finite exactly when the complex group algebra $\mathbb{C}[G]$ is an algebraic Baer *-algebra.

Published in Pacific Journal of Mathematics, Vol. 316 (2022), No. 2, 431-452. DOI: https://doi.org/10.2140/pjm.2022.316.431

1. INTRODUCTION AND PRELIMINARIES

The authors investigated complex algebraic *-algebras in the recently published article [13]. The present paper is a continuation of this research, and our goal is that to derive new and significant properties of this class of *-algebras (Theorems 2.3, 3.6 and 4.1).

First we recall the notions in question. All rings and algebras are associative and not necessarily unital. An element a of an algebra Aover a field \mathbb{F} is said to be *algebraic*, if a generates a finite dimensional subalgebra, or equivalently, a is a root of a polynomial with coefficients from \mathbb{F} . The algebra A is *algebraic*, if every element of A is algebraic. It is obvious that every subalgebra of an algebraic algebra is algebraic as well. For general properties of algebraic algebras we refer the reader to [5].

Date: August 14, 2021.

²⁰²⁰ Mathematics Subject Classification. Primary 16W10; Secondary 22D15, 46L99.

Key words and phrases. Algebraic algebra; Baer *-ring; von Neumann regular algebra; AW^* -algebra; group algebra.

An *involution* of a ring A is a mapping $*: A \to A$ such that

$$(a^*)^* = a; \ (a+b)^* = a^* + b^*; \ (ab)^* = b^*a^* \qquad (a,b \in A).$$

A *-ring is a ring with involution. We say that A is a *-algebra, if it is an algebra over the complex numbers \mathbb{C} , and there is an involution * : $A \to A$ with the additional property

$$(\lambda a)^* = \overline{\lambda} a^* \qquad (\lambda \in \mathbb{C}, \ a \in A).$$

So our convention in the paper is that a *-algebra is always complex and not necessarily unital. The set $\{a \in A \mid a^* = a\}$ of *self-adjoint elements* is denoted by A_{sa} , while A_+ is the set of *positive elements* of A, i.e., $a \in A_+$ iff $a = \sum_{j \in J} a_j^* a_j$ for some finite system $(a_j)_{j \in J}$ in A.

For a *-algebra A and an $a \in A$, the notation $\text{Sp}_A(a)$ stands for the *spectrum* of a. That is,

$$\operatorname{Sp}_A(a) := \{\lambda \in \mathbb{C} | \lambda \mathbf{1} - a \text{ has no inverse in } A\}$$

when A has unit 1, and if A is non-unital, then $\operatorname{Sp}_A(a)$ denotes the spectrum of (0, a) in the standard unitization $A^1 := \mathbb{C} \oplus A$ of A. The *-algebra A is said to be *hermitian*, if every selfadjoint element in A has real spectrum. A is symmetric (resp. completely symmetric), if $\operatorname{Sp}_A(a^*a) \subseteq \mathbb{R}_+$ for any $a \in A$ (resp. $\operatorname{Sp}_A(x) \subseteq \mathbb{R}_+$ for any $x \in A_+$), see 9.8. in [8]. Note here that a unital *-algebra A is symmetric iff for all $a \in A$ the element $\mathbf{1} + a^*a$ has inverse in A. For other general properties of *-algebras and C*-algebras see [3], [7] and [8].

The presence of algebraic *-algebras is natural in the theories of operator algebras and harmonic analysis. Namely, the *-algebra of finite rank operators on a complex Hilbert space ([13], Example 4.2) and the convolution *-algebra of finitely supported complex functions defined on a discrete locally finite group ([13], Theorem 6.5) are algebraic *algebras. Moreover, algebraic *-algebras appear in the important theory of approximately finite dimensional C^* -algebras. According to one of the definitions ([3], III.2), a C^* -algebra \mathscr{A} is approximately finite dimensional if it has an increasing sequence $(A_n)_{n\in\mathbb{N}}$ of finite dimensional *-subalgebras such that $A := \bigcup_{n\in\mathbb{N}}A_n$ is norm-dense in \mathscr{A} . It is obvious that A is an algebraic *-algebra.

The examples above explain that it might be of interest to study general properties of these *-algebras. The authors' recent paper [13] contains many results in this context (Theorems 3.8 and 4.1, Proposition 3.10). For instance, in section 3 we introduced the concept of an E^* -algebra, that is, a *-algebra A with the following extension property: every representable positive functional defined on an arbitrary *-subalgebra has a representable positive linear extension to A. Among other results, we showed that a *-algebra is an E^* -algebra if and only if it is hermitian and every selfadjoint element is algebraic ([13], Theorem 3.8). In fact, more was proved: the class of E^* -algebras coincides with the class of completely symmetric algebraic *-algebras ([13], Corollary 4.5).

To continue the investigation in this direction, the recent paper focuses on two conditions which an algebraic *-algebra A may fulfil:

- (1) a pre- C^* -norm exists on A;
- (2) A is a Baer *-algebra.

In the subsections 1.1 and 1.2, we make a short survey on these properties. Our main result related to (1) is Theorem 2.3, which is a characterization theorem. In relation to (2) we prove Theorem 3.6, which is a structure theorem for algebraic Baer *-algebras. As an application of the latter theorem we show that for a group G the group algebra $\mathbb{C}[G]$ is an algebraic Baer *-algebra iff G is finite (Theorem 4.1).

1.1. Algebraic pre- C^* -algebras. Let A be a *-algebra. A submultiplicative seminorm $\sigma: A \to \mathbb{R}_+$ is a C^* -seminorm ([8], 9.5), if

(1.1)
$$\sigma(a^*a) = \sigma(a)^2 \quad (a \in A).$$

The kernel of σ , i.e., the *-ideal $\{a \in A | \sigma(a) = 0\}$ is denoted by ker σ . If ker $\sigma = \{0\}$, that is, σ is a (possibly incomplete) norm, then we say that σ is a *pre-C**-*norm* and (A, σ) is a *pre-C**-*algebra*.

One of the most important concepts in the general theory of *algebras is the following ([8], Definition 10.1.1). For every $a \in A$, let (1.2)

 $\gamma_A(a) := \sup\{\|\pi(a)\| | \pi \text{ is a }^*\text{-representation of } A \text{ on a Hilbert space}\}.$

If γ_A is finite valued, then it is called the *Gelfand-Naimark seminorm* of A, and it is the greatest C^{*}-seminorm on A ([8], Theorem 10.1.3).

The concrete examples of algebraic *-algebras above have a common property: they all admit a pre- C^* -norm. Of course, in general, an algebraic *-algebra does not have a pre- C^* -norm. For example, Theorem 9.7.22 in [8] produces many finite dimensional (hence algebraic) *algebras which not possess a pre- C^* -norm. So a natural question arises: if A is a given algebraic *-algebra, then what kind of assumptions guarantee the existence of a pre- C^* -norm on A? In section 2, we give purely algebraic answers to this question (Theorem 2.3), and in fact, each of these conditions actually characterize the existence of a pre- C^* -norm on A. We recall them here.

Definition 1.1. The involution of a *-ring A is proper ([8], page 990; [1], §2), if for any $a \in A$ the equation $a^*a = 0$ implies that a = 0.

By (1.1), it is immediate that every pre- C^* -algebra has proper involution. In Theorem 2.3, we show for an algebraic *-algebra that the properness of the involution guarantees that the *-algebra admits a pre- C^* -norm.

The following concept was first studied by J. von Neumann (see the Introduction in [4]); for general properties see [2] and [4].

Definition 1.2. A ring A is said to be *von Neumann regular*, if for every $a \in A$ there is an $x \in A$ such that axa = a.

Von Neumann regularity and hermicity imply for an algebraic *-algebra that the *-algebra has a pre- C^* -norm (Theorem 2.3). In subsection 3.1, which deals with Baer *-algebras without Abelian summand, von Neumann regularity is very important.

Before the next definition we recall the notion of annihilators ([1], page 12): if A is a ring, $S \subseteq A$ is an arbitrary subset, then the right annihilator of S is

$$\operatorname{ann}_{rA}(S) := \{ y \in A | ay = 0 \ \forall a \in S \},\$$

which is obviously a right ideal of A. The *left annihilator* $\operatorname{ann}_{lA}(S)$ of S can be defined similarly (which is a left ideal); a non-trivial ideal of A is an *annihilator ideal*, if it is a left annihilator of some set $S \subseteq A$. (We drop A from the index, when it is clear from the context in which algebra the annihilator is considered.)

Definition 1.3. A *-ring A is said to be a

(a) Rickart *-ring ([1], §3), if for any $a \in A$ the right annihilator of $\{a\}$ is a principal right ideal generated by a projection f, that is,

$$\operatorname{ann}_r(\{a\}) = fA := \{fy | y \in A\}$$

with an $f \in A$ such that $f = f^* = f^2$.

- (b) weakly Rickart *-ring ([1], §5), if for every $a \in A$ there is a projection $e \in A$ with the following properties:
 - (1) ae = a;

(2) ay = 0 implies ey = 0 for every $y \in A$.

If A is a weakly Rickart *-ring, the conditions (1) and (2) uniquely determines the projection e, and it is called *the right projection* of a; in notation: $e = \operatorname{RP}(a)$. The *left projection* LP(a) can be obtained similarly.

A (weakly) Rickart *-algebra is a *-algebra which is also a (weakly) Rickart *-ring.

By §3 and §5 in [1], we have the following properties. A *-ring A is a Rickart *-ring if and only if A is a weakly Rickart *-ring with

4

unity; in this case the definitions above give $\operatorname{RP}(a) = \mathbf{1} - f$. A very important observation is that the involution of a weakly Rickart *-ring is proper, hence algebraic *-algebras which are also weakly Rickart *-algebras admit pre- C^* -norms (Theorem 2.3).

For a *-ring A let $\mathbf{P}(A)$ be the set of projections in A; let \leq denote the natural ordering, i.e., $p \leq q$ iff p = pq = qp $(p, q \in \mathbf{P}(A))$. Note here that there is an ordering on the set of self-adjoint elements ([1], Definition 8 on page 70 and §50). In the results of this paper these orderings coincide on $\mathbf{P}(A)$, hence we use the same notation: for $x, y \in$ $A_{sa} x \leq y$ means that $y - x \in A_+$.

The projections of a weakly Rickart *-algebra form a lattice with the following operations ([1], Proposition 7 on page 29):

$$e \lor f = f + \operatorname{RP}(e - ef); \ e \land f = e - \operatorname{LP}(e - ef) \quad (e, f \in \operatorname{P}(A)).$$

1.2. Algebraic Baer *-algebras. As we mentioned in the preceding subsection, we prove that algebraic weakly Rickart *-algebras have pre- C^* -norms (Theorem 2.3). In Section 3, we investigate algebraic *algebras with the following stronger assumption, which was introduced by I. Kaplansky to give an algebraic axiomatization of von Neumann algebras. (See the introductions of [1] and [4].)

Definition 1.4. Let A be a *-ring. We say that A is a *Baer* *-*ring* ([1], §4), if for every non-empty subset $S \subseteq A$ the right annihilator $\operatorname{ann}_r(S)$ of S is a principal right ideal generated by a projection. A *Baer* *-*algebra* is a *-algebra which is also a Baer *-ring.

For Baer *-rings the reader is referred to the fundamental works of S. K. Berberian [1] and [2].

Our main result in the context of algebraic Baer *-algebras is Theorem 3.6, which gives a complete description for this class. To obtain this theorem we collect some indispensable properties of Baer *-rings here (see [1]).

By the definitions, it is obvious that a Baer *-ring A is a Rickart *-ring, hence the involution of A is proper and A is unital. A very important property of the projection lattice of a Baer *-ring is contained in the following statement ([1], Proposition 1 in §4):

Proposition 1.5. The following conditions on a *-ring A are equivalent.

- (i) A is a Baer *-ring;
- (ii) A is a Rickart *-ring whose projections form a complete lattice.

From a given Baer *-ring or *-algebra we can easily obtain "new" Baer *-rings ([1], §4). If A is a Baer *-ring (resp. *-algebra), then eAe

is a Baer *-subring (resp. *-subalgebra) for every projection $e \in \mathbf{P}(A)$, as well as the *bicommutant* $S'' := \{a \in A | as = sa \forall s \in S\}$ of every *-closed subset S of A. It is obvious that the image of a Baer *-ring via a *-isomorphism (i.e., injective *-homomorphism) is also a Baer *-ring.

The more concrete examples of Baer *-rings are the so-called AW^* algebras, that is, C^* -algebras which are Baer *-algebras (see [1] and [12]). For instance, every von Neumann algebra is an AW^* -algebra ([1], Proposition 9 on page 24), but for us the following two examples are the most important.

Example 1.6. The class of finite dimensional Baer *-algebras is precisely the class of finite dimensional C^* -algebras. Indeed, a Baer *-algebra A has proper involution, hence finite dimensionality implies that A is a C^* -algebra ([8], Theorem 9.7.22). Moreover, A can be written as a finite direct sum of standard complex matrix *-algebras (see also [3], page 74):

(1.3)
$$A \cong \bigoplus_{k=1}^{m} M_{n_k}(\mathbb{C}).$$

Furthermore, it is clear that full matrix *-algebras over \mathbb{C} are Baer *-algebras, as well as their finite direct sums.

Example 1.7. By Theorem 1 in §7 of [1] and the well-known Gelfand-Naimark theorem, a commutative C^* -algebra \mathscr{B} is an AW^* -algebra if and only if it is *-isomorphic to the C^* -algebra $\mathscr{C}(T; \mathbb{C})$ of the complex valued continuous functions on a *Stonean topological space* T (that is, T is compact Hausdorff and *extremally disconnected*: every open set has open closure).

For a Stonean topological space T, the C^* -algebra $\mathscr{C}(T; \mathbb{C})$ naturally contains a norm-dense algebraic Baer *-subalgebra. Indeed, the projections of $\mathscr{C}(T; \mathbb{C})$ are exactly the characteristic functions of the clopen (closed and open) sets in T ([1], page 40 and 41), and these form a complete lattice. Hence, if \widehat{B} denotes the linear span of the characteristic functions of the clopen sets, then it is a dense ([1], Proposition 1 in §8) unital *-subalgebra with complete projection lattice. From the commutativity, it is easy to see that \widehat{B} is algebraic, moreover the supremumnorm is a pre- C^* -norm on \widehat{B} . Our result (Theorem 2.3) implies that \widehat{B} is a Rickart *-algebra, thus \widehat{B} is a Baer *-algebra by Proposition 1.5.

Now one may ask the following question: are there algebraic Baer *-algebras that have different structure than the ones in the previous examples? The answer is in our result, Theorem 3.6. It states that every algebraic Baer *-algebra is a direct sum of a finite dimensional Baer *algebra and a commutative algebraic Baer *-algebra as in Example 1.7. 2. Existence of pre- C^* -norms on algebraic *-algebras

Our recently published paper [13] contains an equivalent condition for a pre- C^* -algebra to be algebraic (Theorem 4.1). As it was mentioned in 1.1 of the Introduction, we assume algebraicity in this section and we study conditions which imply that an algebraic *-algebra has a pre- C^* -norm (Theorem 2.3). This is our main result in this part.

The following statement about algebraic algebras is elementary (see Lemma 3.2 in [13] and Lemma 3.12 in [10]).

Lemma 2.1. Let A be a complex algebraic algebra. For an element $b \in A$, denote by A_b the subalgebra generated by $\mathbf{1}$ and b (resp. b) if A is unital (resp. A is non-unital).

Then for every $b \in A$, the spectrum $\text{Sp}_A(b)$ is non-empty, finite and the following is true:

(2.1)
$$\operatorname{Sp}_{A}(b) = \begin{cases} \operatorname{Sp}_{A_{b}}(b), & \text{when } A \text{ is unital,} \\ \operatorname{Sp}_{A_{b}}(b) \cup \{0\}, & \text{when } A \text{ is non-unital.} \end{cases}$$

Furthermore, for any $a, b \in A$

(2.2)
$$\operatorname{Sp}_A(ab) = \operatorname{Sp}_A(ba)$$

holds.

If A is a complex algebra, then A_J stands for the Jacobson radical of A. Following Palmer's terminology in [7] and [8], A is semisimple iff $A_J = \{0\}$.

The next statement collects several results from the authors' paper [13].

Theorem 2.2. Let A be a *-algebra such that every selfadjoint element of A is algebraic. Then the following are equivalent.

(i) A is hermitian.

(ii) $A_J = \ker \gamma_A$.

If these properties hold, then A is a completely symmetric algebraic *-algebra such that the elements of the Jacobson radical A_J are nilpotent. Furthermore, the Gelfand-Naimark seminorm γ_A in (1.2) is finite valued, hence it is the greatest C*-seminorm on A.

Proof. See [13]: the equivalence comes from Theorem 3.8 (iia) and (iic); the properties follow from Proposition 3.10. (b), (g), (f) and Theorem 3.8 (iii).

Our main result on general algebraic *-algebras is the following statement. We use it often in the Baer *-algebra case, Section 3. Note here that the properties listed below (with the exception of (i)) are purely algebraic conditions. **Theorem 2.3.** Let A be an algebraic *-algebra. The following assertions are equivalent.

- (i) There exists a pre- C^* -norm on A.
- (ii) The involution of A is proper.
- (iii) A is hermitian and semisimple.
- (iv) Every non-zero selfadjoint $b \in A_{sa}$ can be written in a finite sum

$$b = \sum_{j \in J} \lambda_j e_j,$$

where $(\lambda_j)_{j \in J}$ is a finite system of non-zero distinct real numbers and $(e_j)_{j \in J}$ is a finite system of non-zero orthogonal projections in A.

- (v) A is a weakly Rickart *-ring.
- (vi) A is hermitian and von Neumann regular.

Proof. We prove the equivalence of (i), (ii) and (iii) first.

(i) \Rightarrow (ii): By the C^{*}-property (1.1), the involution is proper on any *-algebra which admits a pre-C^{*}-norm.

(ii) \Rightarrow (iii): We show that A is hermitian if the involution of A is proper. Let us fix a selfadjoint $b \in A_{sa}$; we have to prove that $\operatorname{Sp}_A(b) \subseteq \mathbb{R}$.

The equation in (2.1) implies that it is enough to see the inclusion $\operatorname{Sp}_{A_b}(b) \subseteq \mathbb{R}$. Since *b* is algebraic, the subalgebra A_b generated by *b* (which is trivially a *-subalgebra) is finite dimensional, and has proper involution from the assumption. So Theorem 9.7.22 in [8] forces A_b to be hermitian, hence $\operatorname{Sp}_{A_b}(b) \subseteq \mathbb{R}$ holds.

Now we conclude the semisimplicity. The Jacobson radical A_J is a *-subalgebra of A, hence $A_J = (A_J)_{sa} + \mathbf{i}(A_J)_{sa}$ is true (\mathbf{i} is the complex imaginary unit). By properness, $(A_J)_{sa} = \{0\}$. As we recalled in Theorem 2.2 above, every element of A_J is nilpotent. Thus for an $a \in (A_J)_{sa}$, let $n \in \mathbb{N}$ be the minimal positive integer such that $a^n = 0$. If $n \geq 2$, then $2n - 2 \geq n$ and $n - 1 \geq 1$. So $a^{2n-2} = 0$, and since $a^* = a$, it follows that $(a^{n-1})^*(a^{n-1}) = 0$. The involution is proper, which implies $a^{n-1} = 0$. This contradicts the minimality of n, thus n = 1, i.e., a = 0.

(iii) \Rightarrow (i): This follows from Theorem 2.2, since γ_A is a pre-C^{*}-norm on A.

Now we show the implications $(i) \Rightarrow (iv) \Rightarrow (iii)$.

 $(i) \Rightarrow (iv)$: Let $\|\cdot\|$ be a pre- C^* -norm on A. Fix a non-zero $b \in A_{sa}$ selfadjoint element, and let B be the *-subalgebra generated by b, which is finite dimensional since b is algebraic. Hence $(B, \|\cdot\||_B)$ is a nonzero finite dimensional commutative C^* -algebra. Since b is selfadjoint, the well-known commutative Gelfand-Naimark theorem and spectral theory imply the existence of a finite system $(\lambda_j)_{j\in J}$ of non-zero distinct real numbers and a finite system $(e_j)_{j\in J}$ of non-zero orthogonal projections in B such that

$$b = \sum_{j \in J} \lambda_j e_j.$$

(iv) \Rightarrow (iii): Let $b \in A_{sa}$ be an arbitrary selfadjoint element. From the assumption (iv) write $b = \sum_{j \in J} \lambda_j e_j$ with a finite system $(\lambda_j)_{j \in J}$ of non-zero distinct real numbers and a finite system $(e_j)_{j \in J}$ of non-zero orthogonal projections in A. Then an easy calculation shows that b is a root of the polynomial

$$p(z) = z \prod_{j \in J} (z - \lambda_j).$$

Hence the spectral mapping property implies that

$$\operatorname{Sp}_A(b) \subseteq \{\lambda_j | j \in J\} \cup \{0\} \subseteq \mathbb{R},\$$

which means that A is hermitian.

To show semisimplicity, assume that $b \in (A_J)_{sa} \setminus \{0\}$. The Jacobson radical is an ideal, hence for any $j_0 \in J$ from the orthogonality we have

$$e_{j_0} = \left(\frac{1}{\lambda_{j_0}}e_{j_0}\right)b \in A_J.$$

But this is absurd, since the radical does not contain non-zero idempotent elements. From this it follows that $(A_J)_{sa} = \{0\}$, thus $A_J = \{0\}$. The equivalence of (i), (ii), (iii) and (iv) has been proved; let us proceed to the properties (v) and (vi).

(ii) & (iv) \Rightarrow (v): According to Definition 1.3, we have to prove for any $a \in A$ that there is a projection $e \in A$ with the following properties:

- (1) ae = a;
- (2) ay = 0 implies ey = 0 for every $y \in A$.

We may suppose that $a \neq 0$. From the assumption (ii) we infer that $a^*a \neq 0$, thus by (iv) we may write

$$a^*a = \sum_{j \in J} \lambda_j e_j$$

with a finite system $(\lambda_j)_{j \in J}$ of non-zero distinct real numbers and a finite system $(e_j)_{j \in J}$ of non-zero orthogonal projections in A. Define the projection e by the following equation:

(2.3)
$$e := \sum_{j \in J} e_j.$$

By orthogonality, it is indeed a projection.

Let us prove property (1), that is, ae = a. The involution is proper, hence

$$ae = a \iff ae - a = 0 \iff (ae - a)^*(ae - a) = 0 \Leftrightarrow$$

 $ea^*ae - a^*ae - ea^*a + a^*a = 0,$

and the last equation is true, since the obvious equalities

$$ea^*ae = a^*ae = ea^*a = a^*a$$

hold.

(2): Let $y \in A$ be arbitrary with ay = 0. Then $a^*ay = 0$, that is,

$$\left(\sum_{j\in J}\lambda_j e_j\right)y=0.$$

Since every $\lambda_j \neq 0$, by orthogonality we conclude that

$$0 = \left(\sum_{j \in J} \frac{1}{\lambda_j} e_j\right) \left(\sum_{j \in J} \lambda_j e_j\right) y = \left(\sum_{j \in J} e_j\right) y = ey,$$

hence A is a weakly Rickart *-ring.

 $(v) \Rightarrow (ii)$: Any weakly Rickart *-ring has a proper involution (subsection 1.1 in the Introduction).

(ii), (iii) & (iv) \Rightarrow (vi): Hermicity follows from (iii). To see the von Neumann regularity, let $a \in A$ be an arbitrary non-zero element. We seek an $x \in A$ such that axa = a. Similar to the preceding proofs, by (ii) and (iv) we may write

$$a^*a = \sum_{j \in J} \lambda_j e_j$$

with a finite system $(\lambda_j)_{j \in J}$ of non-zero distinct real numbers and a finite system $(e_j)_{j \in J}$ of non-zero orthogonal projections in A. Define the element x by the equation

$$x := \left(\sum_{j \in J} \frac{1}{\lambda_j} e_j\right) a^*.$$

Then

$$axa = a\left[\left(\sum_{j\in J} \frac{1}{\lambda_j} e_j\right) a^*\right] a = a\left(\sum_{j\in J} \frac{1}{\lambda_j} e_j\right) \left(\sum_{j\in J} \lambda_j e_j\right) = a\left(\sum_{j\in J} e_j\right) = a.$$

10

(The last equation is property (1) in the proof of the implication (ii) & $(iv) \Rightarrow (v)$.)

 $(vi) \Rightarrow (iii)$: A von Neumann regular algebra is always semisimple: if $a \in A_J, x \in A$ and axa = a, then axax = ax, that is, ax is an idempotent in the radical. Thus, ax = 0, which implies a = 0.

Remark 2.4. We note here that if there is a complete pre- C^* -norm on the algebraic *-algebra A, then A is finite dimensional. In fact, by Kaplansky's Lemma 7 in [6], an infinite dimensional semisimple Banach algebra has an element with infinite spectrum. Thus, by Lemma 2.1, every semisimple algebraic Banach algebra is finite dimensional. Furthermore, a von Neumann regular Banach algebra is finite dimensional as well (this is also Kaplansky's theorem: [7], Theorem 2.1.18).

Remark 2.5. Let A be an algebraic *-algebra.

- Semisimplicity does not imply that A is hermitian (see Theorem 9.7.22 in [8]), and the same is true for von Neumann regularity (semisimple finite dimensional *-algebras are von Neumann regular, including the non-hermitian ones).
- (2) If A satisfies the properties of the previous theorem (hence it is a weakly Rickart *-ring), then for an element $a \in A$ the right projection RP(a) of a is the projection e in (2.3). That is, if

$$a^*a = \sum_{j \in J} \lambda_j e_j$$

for a finite system $(\lambda_j)_{j \in J}$ of non-zero distinct real numbers and a finite system $(e_j)_{j \in J}$ of non-zero orthogonal projections in A, then

$$\operatorname{RP}(a) = \sum_{j \in J} e_j = \operatorname{RP}(a^*a).$$

Furthermore, from the proof of the theorem it can be easily seen that the projections e_j $(j \in J)$ are in the subalgebra generated by a^*a . It follows that the right projection of any $a \in A$ is in the *-subalgebra generated by a.

(3) By means of Theorem 2.2, one can show that property (iv) actually characterizes algebraic pre- C^* -algebras in the class of *-algebras. Namely, a *-algebra A is an algebraic *-algebra with a pre- C^* -norm if and only if every non-zero selfadjoint $b \in A_{sa}$ can be written in a finite sum

(2.4)
$$b = \sum_{j \in J} \lambda_j e_j,$$

where $(\lambda_j)_{j \in J}$ is a finite system of non-zero distinct complex numbers and $(e_j)_{j \in J}$ is a finite system of non-zero orthogonal projections in A. In this case every normal element b of A has the form (2.4). Moreover,

$$Sp_A(b) \cup \{0\} = \{\lambda_j | j \in J\} \cup \{0\}$$

holds for the spectrum of b. The numbers λ_j are real when b is selfadjoint, and they are non-negative if b is a positive element, since A is completely symmetric by Theorem 2.2.

Now we list further properties of general algebraic pre- C^* -algebras (some of them were shown in the authors' paper [13]). These are indispensable in Section 3.

Proposition 2.6. If A is an algebraic pre- C^* -algebra, then the following assertions hold.

- (1) The pre- C^* -norm on A is unique.
- (2) Denote by $C^*(A)$ the completion of A with respect to the unique pre- C^* -norm. Then for every closed ideal $I \subseteq C^*(A)$ the equation $I = \overline{A \cap I}$ holds, where the bar means the closure in the norm of $C^*(A)$.
- (3) A satisfies the **(EP)-axiom** ([1], Definition 1 on page 43), that is, for any $a \in A \setminus \{0\}$ there exists $a w \in \{a^*a\}''$ such that $w = w^*$ and $(a^*a)w^2$ is a non-zero projection.
- (4) A satisfies the (UPSR)-axiom ([1], Definition 10 on page 70), that is, for any x ∈ A₊ there is a unique y ∈ A₊ such that y² = x and in addition y ∈ {x}".
- (5) A is the linear span of its projections.

Moreover, if A has a unit element, then:

- (6) $a^*a \leq ||a||^2 \mathbf{1}$ holds for any $a \in A$.
- (7) A is a **directly finite** algebra, that is, for any $a, b \in A$ the equation ab = 1 implies that ba = 1.

Proof. (1): Theorem 4.1 (a) in [13] shows this.

(2): This statement is Theorem 4.1 (f) in [13].

(3): Let $a \in A$ be a non-zero element. Since $a^*a \ge 0$, by Theorem 2.3 and Remark 2.5 (3), we may write

$$a^*a = \sum_{j \in J} \lambda_j e_j$$

with a finite system $(\lambda_j)_{j \in J}$ of non-zero distinct positive numbers and a finite system $(e_j)_{j \in J}$ of non-zero orthogonal projections in A. Then the element

$$w = \sum_{j \in J} \frac{1}{\sqrt{\lambda_j}} e_j$$

is selfadjoint (in fact, positive), moreover orthogonality implies

$$(a^*a)w^2 = \left(\sum_{j\in J}\lambda_j e_j\right)\left(\sum_{j\in J}\frac{1}{\lambda_j}e_j\right) = \sum_{j\in J}e_j = \operatorname{RP}(a),$$

which is a non-zero projection, as desired. The property $w \in \{a^*a\}''$ for the bicommutant also holds, since w is an element of the *-subalgebra generated by a^*a according to Remark 2.5 (2).

(4): Let $x \in A_+$ be a non-zero positive element. Then x is of the form

$$x = \sum_{j \in J} \lambda_j e_j$$

with a finite system $(\lambda_j)_{j \in J}$ of non-zero distinct positive numbers and a finite system $(e_j)_{j \in J}$ of non-zero orthogonal projections in A. Obviously the element

$$y := \sum_{j \in J} \sqrt{\lambda_j} e_j$$

is positive and orthogonality shows that $y^2 = x$.

To see the uniqueness, let $z \in A$ be a positive element such that $z^2 = x$. Let A_z (resp. A_x) denote the *-subalgebra generated by z (resp. x). Then $A_x \subseteq A_z$, moreover these are finite dimensional C^* -algebras by the algebraicity. From Remark 2.5 (2), it follows that $y \in A_x$, hence $y \in A_z$. But $y^2 = x = z^2$ and it is well-known that every positive element has a *unique* positive square root in a C^* -algebra, thus y = z. The inclusion $y \in \{x\}''$ is also clear from $y \in A_x$.

(5): The equality $A = A_{sa} + \mathbf{i}A_{sa}$ holds for the *-algebra A. Thus, Theorem 2.3 (iv) shows that every element of A can be written as a linear combination of the projections in A.

For the rest of the proof, assume that A is unital.

(6): For an arbitrary $a \in A$, the *-subalgebra generated by a^*a and **1** is a finite dimensional C^* -algebra. Hence spectral theory and the C^* -property imply the inequality $a^*a \leq ||a||^2 \mathbf{1}$.

(7): This follows from (2.2) in Lemma 2.1. \Box

As we noted in the Introduction 1.2, every Baer *-ring is a Rickart *-ring. Hence for an algebraic *-algebra which is also a Baer *-ring, the properties in Theorem 2.3 and Proposition 2.6 come true. We formulate this in the next statement.

Corollary 2.7. If A is an algebraic Baer *-algebra, then A is a directly finite, von Neumann regular and completely symmetric *-algebra with a unique pre-C*-norm $\|\cdot\|$, satisfying the (EP)- and (UPSR)-axioms. Moreover, A is the linear span of its projections, and for any $a \in A$ the inequality $a^*a \leq \|a\|^2 \mathbf{1}$ holds.

To close this section we present the following lemma which characterizes finite dimensionality of algebraic *-algebras with pre- C^* -norms. It is used in the proof of Theorem 3.3.

Lemma 2.8. An algebraic pre- C^* -algebra A is finite dimensional if and only if every orthogonal system of non-zero projections in A is finite.

Proof. If A is finite dimensional, then it is obvious that every orthogonal system of non-zero projections in A is finite, since such a system is linearly independent.

For the converse, suppose that every orthogonal system of non-zero projections in A is finite. Note first that it is enough to prove finite dimensionality in the case of unital A. Indeed, if A is not unital, then let $A^1 := \mathbb{C} \oplus A$ be the standard unitization of the *-algebra A. Now it is clear that A^1 is also an algebraic *-algebra with a pre- C^* -norm ([8], Proposition 9.1.13 (b)). Assume that $(\lambda_k, p_k)_{k \in \mathbb{N}}$ is an infinite orthogonal system of projections in A^1 . Then orthogonality shows for every $k, m \in \mathbb{N}, k \neq m$ that

$$(0,0) = (\lambda_k, p_k)(\lambda_m, p_m) = (\lambda_k \lambda_m, \lambda_k p_m + \lambda_m p_k + p_k p_m),$$

hence $\lambda_k \neq 0$ holds for at most one $k \in \mathbb{N}$. Dropping out this possible exception, the system has the form $(0, p_k)_{k \in \mathbb{N}}$. Now it is clear that $(p_k)_{k \in \mathbb{N}}$ would be an infinite orthogonal system of projections in A, which is impossible according to the assumption. Thus, all of the assertions we made for A are true for A^1 . Moreover if A^1 is finite dimensional, then A is also finite dimensional.

Suppose that A is a unital algebraic pre- C^* -algebra such that every orthogonal system of non-zero projections in A is finite. Then it is easy to see that A does not contain a strictly increasing sequence of non-zero projections. Indeed, if $(q_k)_{k\in\mathbb{N}}$ is such sequence, then the projections $p_0 := q_0, p_{k+1} := q_{k+1} - q_k \ (k \in \mathbb{N})$ would be form an infinite orthogonal system of non-zero projections.

By Theorem 2.3 (vi), the *-algebra A is von Neumann regular, hence every finitely generated right ideal I of A is of the form eA with an idempotent $e \in A$ (see Lemma 1.3 on page 68 in [10] or Theorem 1.1 in [4]). Furthermore, since the involution of A is proper (Theorem 2.3 (ii)), Proposition 3 on page 229 in [1] implies that we may assume

14

that e is a projection. This and the preceding paragraph imply that A satisfies the ascending chain condition for finitely generated right ideals. Hence, by Lemma 2.9 on page 80 in [10], we obtain that A is Artinian. In fact, the proof of this lemma and the arguments above imply that every right ideal of A is generated by a projection. Thus, the unital A is a modular annihilator algebra ([7], Definition 8.4.6), because it is semisimple (hence semiprime: [7], Definition 4.4.1) and obviously satisfies property (a) of Theorem 8.4.5 in [7]. Now Proposition 8.4.14 of [7] implies that A is finite dimensional, since it is a normed algebra. \Box

3. Algebraic Baer *-Algebras

The main purpose of this section is to answer the following questions. Theorem 3.6 answers them.

- (1) In the light of Corollary 2.7, what other properties an algebraic Baer *-algebra has?
- (2) Are there examples of algebraic Baer *-algebras different than the ones given in Examples 1.6 and 1.7?

The lattice of projections of a Baer *-ring is complete ([1], §15) and these rings have well-developed structure theory. We recall some relevant definitions. We say that a *-ring is *Abelian* if all of its projections are central. A *-ring A is properly non-Abelian, if the only Abelian central projection is 0, i.e., if for a central projection $e \in A$ the *-ring eAeis Abelian, then e = 0.

By Corollary 2.7, an algebraic Baer *-algebra A is Abelian iff A is commutative, since it is the linear span of its projections (the same equivalence is true for AW^* -algebras: [1], Examples on page 90).

The key to our structure Theorem 3.6 is the following decomposition $([1], \text{ Theorem 1 } (2) \text{ in } \S15)$:

Theorem 3.1. Let A be a Baer *-ring. Then there exists a unique central projection $h \in A$ such that in the decomposition

$$A = (\mathbf{1} - h)A + hA$$

the summand $M := (\mathbf{1} - h)A$ is a properly non-Abelian Baer *-subring, while B := hA is an Abelian Baer *-subring.

If $B = \{0\}$, then we say that A has no Abelian summand.

By this theorem and the observations above, an algebraic Baer *algebra A can be decomposed as a sum of algebraic Baer *-subalgebras

$$(3.1) A = M \oplus B,$$

where M has no Abelian summand and B is commutative. Thus, if we fully analyze the properly non-Abelian and the commutative case separately, we get a complete description of general algebraic Baer *algebras. This has been done in the following subsections.

3.1. Algebraic Baer *-algebras without Abelian summands. The following statement is a reformulation of Corollary 3 on page 231 in [1] (the assumptions for A can be found in the statement of this corollary and at the beginning of §51). Note here that a unital *-ring A is finite iff $x^*x = \mathbf{1}$ implies $xx^* = \mathbf{1}$ for any $x \in A$.

Lemma 3.2. Let A be a finite, von Neumann regular and symmetric Baer *-ring without Abelian summand, satisfying the (EP)- and (UPSR)-axioms. Assume that A contains a central element $i \in A$ with the properties $i^2 = -1$ and $i^* = -i$.

If, for any $a \in A$, there is a positive integer k such that $a^*a \leq k\mathbf{1}$, then every system of non-zero orthogonal projections in A is finite.

We are ready to prove our result on algebraic Baer *-algebras that have no Abelian summand.

Theorem 3.3. Let M be a *-algebra. The following are equivalent.

- (i) M is an algebraic Baer *-algebra without Abelian summand.
- (ii) M is a finite dimensional Baer *-algebra without Abelian summand.

If one (hence all) of the properties holds for M, then M is a finite direct sum of full complex matrix algebras of size at least 2×2 .

Proof. (i) \Rightarrow (ii): By Corollary 2.7, M satisfies all of the assertions of the previous lemma (with $i := \mathbf{i1}$), hence every system of non-zero orthogonal projections in M is finite. Now Lemma 2.8 implies the finite dimensionality of M.

(ii) \Rightarrow (i): Finite dimensional algebras are algebraic.

As we mentioned in Example 1.6, finite dimensional Baer *-algebras are finite direct sum of full matrix *-algebras over \mathbb{C} . Since M has no Abelian summand, then the size of every matrix algebra in the decomposition (1.3) of Example 1.6 must be at least 2×2 .

3.2. Commutative algebraic Baer *-algebras. Our first result shows that the completion with respect to the unique pre- C^* -norm (Corollary 2.7) is also a Baer *-algebra in the commutative case.

Lemma 3.4. Let B be a commutative Baer *-algebra which is algebraic. If $C^*(B)$ stands for the completion of B with respect to its unique pre-C*-norm, then $C^*(B)$ is a commutative AW*-algebra, and every projection of $C^*(B)$ is in B.

16

Proof. We prove that $C^*(B)$ is a Baer *-algebra by Definition 1.4. Since the algebras in question are commutative, the left and right attributes for the annihilators are redundant, so we have to prove for a fixed non-empty set $S \subseteq C^*(B)$ that

$$\operatorname{ann}_{C^*(B)}(S) = \{a \in C^*(B) | Sa = \{0\}\} = fC^*(B)$$

for some projection $f \in C^*(B)$. We show that such a projection exists in B.

For the non-empty set $B \cap \operatorname{ann}_{C^*(B)}(S)$ the Baer property implies that there is projection $e \in B$ such that

$$eB = \operatorname{ann}_B \left(B \cap \operatorname{ann}_{C^*(B)}(S) \right).$$

We state that for the projection $f := \mathbf{1} - e \in B$ the equation

$$\operatorname{ann}_{C^*(B)}(S) = C^*(B)f$$

is true. To see this, it is enough to claim the following equality:

(3.2)
$$eC^*(B) = \operatorname{ann}_{C^*(B)} \left(\operatorname{ann}_{C^*(B)}(S) \right).$$

Indeed, if (3.2) holds, then taking annihilators on both sides gives by Proposition 1 (3) in §3 of [1] that

$$C^*(B)(\mathbf{1}-e) = \operatorname{ann}_{C^*(B)}(eC^*(B)) = \operatorname{ann}_{C^*(B)}(S).$$

Now we prove (3.2). Let $a \in \operatorname{ann}_{C^*(B)}(S)$ be an arbitrary element. Note first that every annihilator is obviously a closed ideal in the commutative C^* -algebra $C^*(B)$. Thus, $B \cap \operatorname{ann}_{C^*(B)}(S)$ is dense in $\operatorname{ann}_{C^*(B)}(S)$ by Proposition 2.6 (2), since B is algebraic. So there is a sequence $(a_n)_{n \in \mathbb{N}}$ in $B \cap \operatorname{ann}_{C^*(B)}(S)$ such that $a_n \to a$. Since $e \in$ $\operatorname{ann}_B(B \cap \operatorname{ann}_{C^*(B)}(S))$ we infer that $ea_n = 0$ for every $n \in \mathbb{N}$. This implies $0 = ea_n \to ea$, that is,

$$e \in \operatorname{ann}_{C^*(B)} \left(\operatorname{ann}_{C^*(B)}(S) \right),$$

thus the inclusion

$$eC^*(B) \subseteq \operatorname{ann}_{C^*(B)} \left(\operatorname{ann}_{C^*(B)}(S) \right)$$

follows.

For the reversed inclusion, let $x \in \operatorname{ann}_{C^*(B)}(\operatorname{ann}_{C^*(B)}(S))$ be arbitrary. The latter set is a closed ideal in $C^*(B)$, so using Proposition 2.6 (2) again we get a sequence $(x_n)_{n\in\mathbb{N}}$ in $B\cap\operatorname{ann}_{C^*(B)}(\operatorname{ann}_{C^*(B)}(S))$ such that $x_n \to x$. Thus, $x_n \in B$ and $x_n\operatorname{ann}_{C^*(B)}(S) = \{0\}$ for every $n \in \mathbb{N}$. In particular, $x_n(B\cap\operatorname{ann}_{C^*(B)}(S)) = \{0\}$. But this exactly shows that $x_n \in \operatorname{ann}_B(B\cap\operatorname{ann}_{C^*(B)}(S)) = eB$ for every $n \in \mathbb{N}$. Taking the limit of the sequence $(x_n)_{n\in\mathbb{N}}$, we have $x \in \overline{eB} = eC^*(B)$, so (3.2) has been proved. To see that every projection of $C^*(B)$ is in B, let $p \in \mathbf{P}(C^*(B))$ be arbitrary. The preceding proof shows that

$$\operatorname{ann}_{C^*(B)}(\{p\}) = qC^*(B)$$

with a projection q in B. But in a unital commutative ring it is well known that the annihilator of an idempotent p is the ideal generated by 1-p, hence q = 1-p ([1], Proposition 1 in §1), so $p = 1-q \in B$. \Box

Our next theorem characterizes algebraic Baer *-algebras among commutative *-algebras.

Theorem 3.5. Let B be a commutative *-algebra. The following statements are equivalent.

- (i) B is an algebraic Baer *-algebra.
- (ii) There exists a Stonean topological space T such that B is *isomorphic to the linear span of the characteristic functions of the clopen sets in T.

Proof. (i) \Rightarrow (ii): Suppose that B is an algebraic Baer *-algebra. Then from the previous lemma, it follows that $C^*(B)$ is an AW^* -algebra. Hence, by Theorem 1 in §7 of [1] and the Gelfand-Naimark theorem, $C^*(B)$ is *-isomorphic to the C^* -algebra $\mathscr{C}(T; \mathbb{C})$ of the complex valued continuous functions on a Stonean topological space T. If \widehat{B} denotes the image of B via the Gelfand-transform, the lemma also concludes that all of the projections in $\mathscr{C}(T; \mathbb{C})$ are actually contained in \widehat{B} . These projections are the characteristic functions of the clopen sets in T ([1], page 40 and 41). Since an algebraic Baer *-algebra is the linear span of its projections (Corollary 2.7), we obtain (ii).

(ii) \Rightarrow (i): The proof of this direction was discussed in Example 1.7. If \widehat{B} denotes the linear span of the characteristic functions of the clopen sets in a Stonean topological space T, then by the arguments in the example we get that \widehat{B} is an algebraic Baer *-algebra. Hence, if B is *-isomorphic to \widehat{B} , then B is an algebraic Baer *-algebra as well. \Box

3.3. General algebraic Baer *-algebras. Now we are in position to prove our structure theorem for algebraic Baer *-algebras.

Theorem 3.6. The following conditions are equivalent for a *-algebra A.

- (i) A is an algebraic Baer *-algebra.
- (ii) A can decomposed as a sum M⊕B, where M is *-isomorphic to a finite direct sum of full complex matrix algebras of size at least 2 × 2, while the summand B is *-isomorphic to the linear span

of the characteristic functions of the clopen sets in a Stonean topological space T.

Proof. (i) \Rightarrow (ii): According to (3.1), we can decompose A into a sum

 $A = M \oplus B,$

where M has no Abelian summand and B is commutative. Now from Theorems 3.3 and 3.5, the properties in (ii) immediately follow.

(ii) \Rightarrow (i): If $A = M \oplus B$ as in (ii), then M and B are algebraic Baer *-algebras by Theorems 3.3 and 3.5. It is clear that their direct sum is an algebraic Baer *-algebra as well.

Corollary 3.7. The C^{*}-algebras which contain a dense algebraic Baer *-algebra are exactly the AW^{*}-algebras with finite codimensional Abelian summand.

Proof. Suppose that \mathscr{A} is a C^* -algebra containing a dense algebraic Baer *-algebra A. By Corollary 2.7, there is a unique pre- C^* -norm on A. The density and the uniqueness imply that the completion $C^*(A)$ of A with respect to this norm is actually \mathscr{A} . According to Theorem 3.6, we may write $A = M \oplus B$, where M is a finite direct sum of full complex matrix algebras of size at least 2×2 , while the summand B is *-isomorphic to the linear span of the characteristic functions of the clopen sets in a Stonean topological space T. Hence this form and the finite dimensionality of M clearly imply that

$$C^*(A) = M \oplus C^*(B).$$

We infer that $C^*(B)$ is an Abelian AW^* -algebra from Lemma 3.4. Since M is a finite dimensional AW^* -algebra without commutative ideals, it follows that $C^*(A)$ is an AW^* -algebra with finite codimensional Abelian summand.

Now let \mathscr{A} be an AW^* -algebra with finite codimensional Abelian summand. By Theorem 3.1, \mathscr{A} can be decomposed as a sum

$$\mathscr{A} = M \oplus \mathscr{B},$$

where M is a finite dimensional AW^* -algebra without Abelian summand and \mathscr{B} is an Abelian (commutative) AW^* -algebra. By Example 1.7 and Theorem 3.5, \mathscr{B} contains a dense algebraic Baer *-algebra B, hence the algebraic Baer *-algebra $A := M \oplus B$ is dense in \mathscr{A} .

4. An application to complex group algebras

Let G be a group and let \mathbb{F} be a field. The group algebra $\mathbb{F}[G]$ consists of all formal finite sums of the form $\sum_{g \in G} \lambda_g g$, where $\lambda_g \in \mathbb{F}$. For $a, b \in \mathbb{F}[G], a = \sum_{g \in G} \lambda_g g, b = \sum_{h \in G} \mu_h h$ and $\lambda \in \mathbb{F}$, the operations are defined by

$$a+b:=\sum_{g\in G}(\lambda_g+\mu_g)g;\ ab:=\sum_{g,h\in G}(\lambda_g\mu_h)gh;\ \lambda a:=\sum_{g\in G}(\lambda\lambda_g)g.$$

If $\mathbb{F} = \mathbb{C}$, then an involution can be defined by

$$a^* = \left(\sum_{g \in G} \lambda_g g\right)^* := \sum_{g \in G} \overline{\lambda_g} g^{-1}.$$

It is easy to see that $\mathbb{C}[G]$ is a *-algebra with proper involution, moreover it admits a pre- C^* -norm. (For the theory of group rings and locally compact groups we refer the reader to [10], [7] and [8].)

Our result specifies which groups have the property that their group algebras over \mathbb{C} are algebraic Baer *-algebras.

Theorem 4.1. If G is a group, then the following statements are equivalent.

- (i) The complex group algebra $\mathbb{C}[G]$ is an algebraic Baer *-algebra.
- (ii) G is finite.

Proof. (i) \Rightarrow (ii): The elements of G constitute an algebraic basis for $\mathbb{C}[G]$, hence we need to prove that $\mathbb{C}[G]$ is finite dimensional. From Theorem 3.6, we obtain that $\mathbb{C}[G]$ splits into a direct sum $M \oplus B$, where M is a finite dimensional Baer *-algebra, B is a commutative algebraic Baer *-algebra which is *-isomorphic to the linear span of the characteristic functions of the clopen sets in a Stonean topological space T. We examine two cases:

- (I) $M = \{0\};$
- (II) $M \neq \{0\}.$

Case (I): if $M = \{0\}$, then $\mathbb{C}[G] = B$, i. e., $\mathbb{C}[G]$ is commutative. It is well known that this occurs exactly when G is commutative. Regarding G as a discrete group, its dual group \hat{G} is a compact group (see [3], Chapter VII.1). The Fourier-Gelfand transform (which is a *-homomorphism) maps $\mathbb{C}[G]$ injectively onto a dense *-subalgebra of $\mathscr{C}(\hat{G};\mathbb{C})$. Thus, the latter C^* -algebra contains a dense algebraic Baer *-algebra. Corollary 3.7 implies that $\mathscr{C}(\hat{G};\mathbb{C})$ is an AW^* -algebra. From this, it follows that \hat{G} is a Stonean topological space (Example 1.7), that is, compact and extremally disconnected. Now, from Theorem 1 in [11], we obtain that \hat{G} is discrete. As \hat{G} is compact and discrete, \hat{G} is finite. Case (II): if $M \neq \{0\}$, then B is an annihilator ideal of $\mathbb{C}[G]$ (since B is the left/right annihilator of M), which is finite codimensional. By Passman's Theorem 3.1 in [9], G is finite.

(ii) \Rightarrow (i): If G is finite, then $\mathbb{C}[G]$ is a finite dimensional *-algebra with a C*-norm, thus it is an algebraic Baer *-algebra.

An analogous statement can be proved in the case of locally compact groups.

Corollary 4.2. Let G be a locally compact group and let β be a left Haar-measure of G. If $\mathscr{K}_{\beta}(G; \mathbb{C})$ stands for the convolution *-algebra of compactly supported complex valued functions on G, then the following statements are equivalent.

(i) $\mathscr{K}_{\beta}(G; \mathbb{C})$ is an algebraic Baer *-algebra.

Proof. (i)⇒(ii): The conditions in (i) imply that G is discrete. Indeed, Theorem 6.5 in [13] or the presence of the unit element in the Baer *-algebra $\mathscr{K}_{\beta}(G; \mathbb{C})$ ensures that G is discrete. Now the group algebra $\mathbb{C}[G]$ can be identified with $\mathscr{K}_{\beta}(G; \mathbb{C})$, hence Theorem 4.1 applies. (ii)⇒(i): Obvious. □

Remark 4.3. Let G be a group with unit $\mathbf{1}_G$.

(1) The conditions "algebraic" and "being a Baer *-algebra" are independent for group algebras over \mathbb{C} . For example, if G is a finitely generated torsion-free commutative group, then for an arbitrary field \mathbb{F} the group algebra $\mathbb{F}[G]$ has no zero-divisors ([10], Lemma 1.1 in §1). Thus, in the case of $\mathbb{F} = \mathbb{C}$, the group algebra $\mathbb{C}[G]$ is obviously a Baer *-algebra. Moreover, for a $g \in$ $G \setminus \{\mathbf{1}_G\} \subseteq \mathbb{C}[G]$, we have that g has infinite order. So $\{g^n | n \in$ $\mathbb{Z}\}$ is a linearly independent system in the group algebra, hence g is not an algebraic element.

In fact, a theorem of Herstein ([10], Theorem 3.11 in §3) states that if \mathbb{F} is a field with zero characteristic, then $\mathbb{F}[G]$ is algebraic if and only if G is locally finite. Thus, for a non-finite locally finite group G, our Theorem 4.1 implies that $\mathbb{C}[G]$ is an algebraic pre- C^* -algebra which is not a Baer *-algebra.

(2) In the absence of an involution, we say that a ring A is Baer, if the right annihilator $\operatorname{ann}_r(S)$ of every non-empty subset $S \subseteq A$ is a principal right ideal generated by an idempotent ([2]). By definition, a Baer *-ring is a Baer ring, but the converse is not true in general (9.1.42 in [8]).

⁽ii) G is finite.

However, if a *-ring is a Baer ring which is von Neumann regular with proper involution ([2], Proposition 1.13) or symmetric ([1], Ex. 5A in page 25), then it is a Baer *-ring. Since complex group algebras $\mathbb{C}[G]$ have proper involution, the properties "Baer ring" and "Baer *-ring" are equivalent for the algebraic complex group algebras (Theorems 2.2 and 2.3 imply that such algebras are symmetric and von Neumann regular). So, by the above mentioned theorem of Herstein and Theorem 4.1, a locally finite group G is finite if and only if $\mathbb{C}[G]$ is a Baer ring.

Acknowledgement

The authors thank the anonymous referee for the insightful comments that improved the presentation the paper.

References

- S. K. Berberian, *Baer *-rings*. Springer-Verlag, Berlin-Heidelberg, 1972, 2nd printing 2011.
- 2. S. K. Berberian, *Baer rings and Baer* *-*rings*, The University of Texas at Austin, 2003.
- 3. K. R. Davidson, C*-algebras by example, American Mathematical Society, 1996.
- 4. K. R. Goodearl, Von Neumann regular rings. Pitman, London-California, 1979.
- 5. N. Jacobson, Structure of rings, American Mathematical Society, 1968.
- I. Kaplansky, Ring isomorphism of Banach algebras, Canad. J. Math., 6 (1954), 374-381.
- 7. T. W. Palmer, Banach algebras and the general theory of *-algebras I. Cambridge University Press, 1994.
- 8. T. W. Palmer, Banach algebras and the general theory of *-algebras II. Cambridge University Press, 2001.
- D. S. Passman, Minimal ideals in group rings, Proc. Amer. Math. Soc., 31 (1972), 81-86.
- D. S. Passman, The algebraic structure of group rings, John Wiley and Sons, Inc., 1977.
- M. Rajagopalan, Fourier transform in locally compact groups, Acta Sci. Math. (Szeged), 25 (1964), 86-89.
- K. Saitô, J. D. M. Wright, Monotone complete C*-algebras and generic dynamics. Springer-Verlag, London, 2015.
- Zs. Szűcs, B. Takács, Hermitian and algebraic *-algebras, representable extensions of positive functionals, Studia Math. 256 (2021), 311-343.

Department of Differential Equations, Budapest University of Technology and Economics, Műegyetem rakpart 3., 1111 Budapest, Hungary

Email address: szucszs@math.bme.hu

DEPARTMENT OF ANALYSIS, BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS, MŰEGYETEM RAKPART 3., 1111 BUDAPEST, HUNGARY

Email address: takacsb@math.bme.hu