

# Global existence of weak solutions to the drift-flux system for general pressure laws

Hai-Liang Li <sup>\*a,b</sup> and Ling-Yun Shou <sup>†a,b</sup>

<sup>a</sup>*School of Mathematical Sciences, Capital Normal University, Beijing 100048, P.R. China*

<sup>b</sup>*Academy for Multidisciplinary Studies, Capital Normal University, Beijing 100048, P.R. China*

## Abstract

The initial value problem to the multi-dimensional drift-flux model for two-phase flow is investigated in this paper, and the global existence of weak solutions with finite energy is established for general pressure-density functions without the monotonicity assumption.

**Key words:** two-phase flow, drift-flux model, global weak solutions, non-monotone pressure laws, quantitative regularity estimate

## 1 Introduction

The drift-flux model for two-phase flow can be used widely in many applied scientific areas, such as nuclear, chemical-process, petroleum, cryogenic, bio-medical, oil-and-gas, microtechnology, sprays [3, 17, 20, 27, 31, 32, 40, 52, 59], and so on. In this paper, we consider the initial value problem (IVP) for the drift-flux system in the periodic domain  $\mathbb{T}^d := \mathbb{R}^d / \mathbb{Z}^d$  ( $d \geq 2$ ):

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t n + \operatorname{div}(n u) = 0, \\ \partial_t((\rho + n)u) + \operatorname{div}((\rho + n)u \otimes u) + \nabla P(\rho, n) = \mu \Delta u + (\mu + \lambda) \nabla \operatorname{div} u, \end{cases} \quad x \in \mathbb{T}^d, \quad t > 0, \quad (1.1)$$

with the initial data

$$(\rho, n, (\rho + n)u)(x, 0) = (\rho_0, n_0, m_0)(x), \quad x \in \mathbb{T}^d, \quad (1.2)$$

---

\*E-mail: hailiang.li.math@gmail.com (H.-L. Li).

†Corresponding author. E-mail: shoulingyun11@gmail.com (L.-Y. Shou).

where  $\rho = \rho(x, t) \in \mathbb{R}_+$  and  $n = n(x, t) \in \mathbb{R}_+$  denote the densities of two fluids respectively,  $u = u(x, t) \in \mathbb{R}^d$  stands for the mixed velocity, the shear viscosity coefficient  $\mu$  and the bulk viscosity coefficient  $\lambda$  satisfy

$$\mu > 0, \quad 2\mu + \lambda > 0, \quad (1.3)$$

and the pressure  $P(\rho, n)$  is assumed to be the general laws

$$\begin{cases} P(\rho, n) \in C^1(\mathbb{R}_+ \times \mathbb{R}_+), \\ \frac{1}{C_0}(\rho^\gamma + n^\alpha) - C_1 \leq P(\rho, n) \leq C_0(\rho^\gamma + n^\alpha) + C_1, \\ |\partial_\rho P(\rho, n)| \leq C_2(\rho^{\tilde{\gamma}-1} + 1), \quad |\partial_n P(\rho, n)| \leq C_2(n^{\tilde{\alpha}-1} + 1), \\ |P(\rho, n)| \leq C_3(\rho^\beta + n^\beta) \quad \text{for } 0 \leq \rho, n \leq 1, \end{cases} \quad (1.4)$$

with the constants  $C_i$  ( $i = 0, 1, 2, 3$ ),  $\tilde{\gamma}, \tilde{\alpha}$ , and  $\beta$  satisfying

$$C_0 \geq 1, \quad C_1 \geq 0, \quad C_2 > 0, \quad C_3 > 0, \quad \tilde{\gamma}, \tilde{\alpha} \geq 1, \quad 0 < \beta < 1. \quad (1.5)$$

There are many significant progress made recently on the mathematical analysis of global solutions to the drift-flux system (1.1) [12, 18–21, 28, 30, 42, 45, 46, 50, 51, 53, 54, 54, 55, 55–58]. Among them, for the pressure

$$P(\rho, n) = C(-b(\rho, n) + \sqrt{b^2(\rho, n) + c(\rho, n)}),$$

the global existence and asymptotical behaviors of solutions to (1.1) have been established in [18, 21] for one-dimensional general initial data and in [28, 30, 53, 56, 57] for multi-dimensional small initial data around the constant equilibrium state. For the pressure

$$P(\rho, n) = C\rho_l^\gamma \left( \frac{n}{\rho_l} - \rho \right)^\gamma,$$

the well-posedness and dynamics of global weak solutions to one-dimensional free boundary value problem were studied in [19, 20, 54, 55]. As for the pressure

$$P(\rho, n) = \rho^\gamma + n^\alpha,$$

the existence of global weak solutions with finite energy to the three-dimensional drift-flux equations (1.1) was investigated by Bresch-Mucha-Zatorska [12] for  $\gamma, \alpha > 1$  in a semi-stationary Stokes regime, by Vasseur-Wen-Yu [45] for  $\gamma > \frac{9}{5}$  and  $\alpha$  satisfying that either  $\alpha$  is close to  $\gamma$  enough or  $\alpha \geq 1$  if

$$\underline{c}\rho_0 \leq n_0 \leq \bar{c}\rho_0, \quad (1.6)$$

with two constants  $0 < \underline{c} \leq \bar{c} < \infty$ , and then by Wen [50] for two independent adiabatic constants  $\gamma, \alpha \geq \frac{9}{5}$ . For more general pressures laws  $P(\rho, n)$  which can be non-monotone on a compact set of the

two variables  $\rho$  and  $n$ , the global weak solutions to (1.1) has been obtained by Novotný-Pokorný [42] in 3D under the assumption (1.6) with two constants  $0 \leq \underline{c} \leq \bar{c} < \infty$  and by Wen-Zhu [51] in 1D without the assumption (1.6). In addition, there are many interesting studies on the global existence of weak solutions to some models related to (1.1), such as compressible Oldroyd-B model [1], two-dimensional non-resistive compressible MHD equations [35], compressible Navier-Stokes equations with entropy transport [38], compressible Navier-Stokes-Vlasov-Fokker-Planck system [40], etc. The reader can refer to the review papers [49, 51].

The drift-flux system (1.1) can be viewed as a simplified model of the compressible nonconservative two-fluid equations (cf. [8, 11, 32, 49]):

$$\begin{cases} \alpha^+ + \alpha^- = 1, \\ \partial_t(\alpha^\pm \rho^\pm) + \operatorname{div}(\alpha^\pm \rho^\pm u^\pm) = 0, \\ \partial_t(\alpha^\pm \rho^\pm u^\pm) + \operatorname{div}(\alpha^\pm \rho^\pm u^\pm \otimes u^\pm) + \alpha^\pm \nabla P^\pm(\rho^\pm) = \operatorname{div}(\alpha^\pm \tau^\pm) + \sigma^\pm \alpha^\pm \rho^\pm \nabla \Delta(\alpha^\pm \rho^\pm), \end{cases} \quad (1.7)$$

where  $\alpha^\pm$ ,  $\rho^\pm$ ,  $u^\pm$ ,  $P^\pm(\rho^\pm)$ , and  $\tau^\pm$  denote the volume fractions, densities, velocities, pressures, and stress tensors of two fluids respectively. The global existence of weak solutions to (1.7) with degenerate viscosity coefficients was proved in [8, 9] for  $P^+(\rho^+) = P^-(\rho^-)$ , and the global well-posedness and optimal time-decay rates of strong solutions near the constant equilibrium state to (1.7) were studied in [22, 47] for  $P^+(\rho^+) \neq P^-(\rho^-)$ .

The drift-flux system (1.1) for either  $\rho = 0$  or  $\rho = n$  can reduce to the barotropic compressible Navier-Stokes equations:

$$\begin{cases} \partial_t \varrho + \operatorname{div}(\varrho v) = 0, \\ \partial_t(\varrho v) + \operatorname{div}(\varrho v \otimes v) + \nabla \pi(\varrho) = \mu \Delta v + (\mu + \lambda) \nabla \operatorname{div} v, \end{cases} \quad (1.8)$$

where  $\varrho$ ,  $v$ , and  $\pi(\varrho)$  denote the density, velocity, and pressure respectively. There are many important investigations about global weak solutions to compressible Navier-Stokes equations (1.8) with constant viscosity coefficients [10, 23, 24, 29, 33, 37, 43]. For instance, concerning the  $\gamma$ -law  $\pi(\varrho) = \varrho^\gamma$ , the existence of global weak solutions with finite energy to (1.8) has been established by Lions [37] for  $\gamma \geq \frac{3d}{d+2}$  ( $d = 2, 3$ ) and  $\gamma > \frac{d}{2}$  ( $d \geq 4$ ), then by Feireisl-Novotný-Petzeltová [23] for  $\gamma > \frac{d}{2}$  ( $d \geq 2$ ), by Jiang-Zhang [33] for  $\gamma > 1$  subject to spherically symmetric initial data, and by Plotnikov-Weigant [43] for  $\gamma = 1$  in 2D. Indeed, the concentration phenomenon of the convective term  $\rho u \otimes u$  may occur for  $\gamma \in [1, \frac{d}{2}]$  [29]. Feireisl [24] proved the global existence of weak solutions to (1.8) for the pressure laws allowed to be non-monotone on a compact set. Bresch-Jabin [10] developed new compactness tools to obtain global weak solutions to (1.8) for more general stress tensors, including thermodynamically unstable pressure laws and anisotropic viscosity coefficients. Moreover, some important progress has been made about global weak solutions to compressible Navier-Stokes equations with degenerate viscosity coefficients, refer to [5–7, 13, 34, 39, 44] and references therein.

However, there are no any results about the global existence problem to the multi-dimensional drift-flux model (1.1) without the monotonicity assumption of the pressure as we know so far. The purpose of this paper is to extend the compactness techniques in [10] to the two-phase case and establish the global existence of weak solutions to the IVP (1.1)-(1.2) for the general pressure laws (1.4).

First, we give the definition of global weak solutions to the IVP (1.1)-(1.2) as follows.

**definition 1.1.**  $(\rho, n, (\rho + n)u)$  is said to be a global weak solution to the IVP (1.1)-(1.2) if for any time  $T > 0$ , the following properties hold:

(1). It holds

$$\begin{cases} 0 \leq \rho \in C([0, T]; L^\gamma_{weak}(\mathbb{T}^d)), & 0 \leq n \in C([0, T]; L^\alpha_{weak}(\mathbb{T}^d)), \\ u \in L^2(0, T; H^1(\mathbb{T}^d)), & \sqrt{\rho + n}u \in L^\infty(0, T, L^2(\mathbb{T}^d)), & (\rho + n)u \in C([0, T]; L^{\frac{2 \min\{\gamma, \alpha\}}{\min\{\gamma, \alpha\} + 1}}(\mathbb{T}^d)), \\ (\rho, n, (\rho + n)u)(x, 0) = (\rho_0, n_0, m_0)(x), & \text{a.e. } x \in \mathbb{T}^d. \end{cases}$$

(2). The continuity equations (1.1)<sub>1</sub>-(1.1)<sub>2</sub> are satisfied in the sense of renormalized solutions:

$$\begin{cases} \partial_t b(\rho) + \operatorname{div}(b(\rho)u) + [b'(\rho)\rho - b(\rho)] \operatorname{div} u = 0 & \text{in } \mathcal{D}(\mathbb{T}^d \times (0, T)), \\ \partial_t b(n) + \operatorname{div}(b(n)u) + [b'(n)n - b(n)] \operatorname{div} u = 0 & \text{in } \mathcal{D}(\mathbb{T}^d \times (0, T)), \end{cases}$$

where  $b \in C^1(\mathbb{R})$  satisfies  $b'(z) = 0$  for all  $z \in \mathbb{R}$  large enough.

(3). The momentum equation (1.1)<sub>3</sub> is satisfied in  $\mathcal{D}'(\mathbb{T}^d \times (0, T))$ .

(4). The energy inequality holds for a.e.  $t \in (0, T)$ :

$$\begin{aligned} & \int_{\mathbb{T}^d} [\frac{1}{2}(\rho + n)|u|^2 + G(\rho, n)] dx + \int_0^t \int_{\mathbb{T}^d} [\mu|\nabla u|^2 + (\mu + \lambda)(\operatorname{div} u)^2] dx d\tau \\ & \leq \int_{\mathbb{T}^d} [\frac{1}{2} \frac{|m_0|^2}{\rho_0 + n_0} + G(\rho_0, n_0)] dx, \end{aligned} \tag{1.9}$$

where the Helmholtz free energy function  $G(\rho, n)$  is defined by

$$G(\rho, n) := \begin{cases} \rho \int_1^\rho P(s, \frac{n}{\rho} s) s^{-2} ds, & \text{if } 0 \leq n \leq \bar{c}\rho, \quad 0 < \bar{c} < \infty, \\ \rho \int_0^\rho P(s, \frac{n}{\rho} s) s^{-2} ds, & \text{if } \rho \geq 0, \quad n \geq 0, \quad C_1 = 0. \end{cases} \tag{1.10}$$

Then, under the assumption (1.6), we have the global existence of weak solutions to the IVP (1.1)-(1.2) below.

**Theorem 1.1.** Suppose that the initial data  $(\rho_0, n_0, m_0)$  satisfies

$$\begin{cases} 0 \leq \underline{c}\rho_0(x) \leq n_0(x) \leq \bar{c}\rho_0(x), & \text{a.e. } x \in \mathbb{T}^d, \\ \frac{m_0(x)}{\sqrt{(\rho_0 + n_0)(x)}} = 0, & \text{if } (\rho_0 + n_0)(x) = 0, \quad \text{a.e. } x \in \mathbb{T}^d, \\ (\rho_0, n_0, \frac{m_0}{\sqrt{\rho_0 + n_0}}) \in L^\gamma(\mathbb{T}^d) \times L^\alpha(\mathbb{T}^d) \times L^2(\mathbb{T}^d), \end{cases} \tag{1.11}$$

with two constants  $0 < \underline{c} \leq \bar{c} < \infty$ . Then, if it holds for the pressure  $P(\rho, n)$  given by (1.4) that

$$\begin{cases} \gamma > \frac{d}{2}, & \gamma \geq \frac{3d}{d+2}, & \alpha \geq 1, \\ \tilde{\gamma}, \tilde{\alpha} \leq \max\{\gamma, \alpha\} + \theta, & \theta := \frac{2}{d} \max\{\gamma, \alpha\} - 1 > 0, \end{cases} \quad (1.12)$$

the IVP (1.1)-(1.2) admits a global weak solution  $(\rho, n, (\rho + n)u)$  in the sense of Definition 1.1.

**Remark 1.1.** The assumption (1.4)<sub>4</sub> implies  $P(0, 0) = 0$  and the continuity of  $G(\rho, n)$  near  $(0, 0)$ .

**Remark 1.2.** There are many interesting pressure laws satisfying (1.4), for instance,

(1) We can take

$$P(\rho, n) = \rho^\gamma + n^\alpha + \sum_{i=1}^N c_i \rho^{\gamma_i} n^{\alpha_i},$$

where the constants  $N$ ,  $\gamma_i$ , and  $\alpha_i$  ( $i = 1, \dots, N$ ) satisfy  $N \geq 1$ ,  $0 \leq \gamma_i < \gamma$ ,  $0 \leq \alpha_i < \alpha$ , and  $\gamma_i + \alpha_i < \max\{\gamma, \alpha\}$ , and the constants  $c_i$  ( $i = 1, \dots, N$ ) are allowed to be negative such that  $P(\rho, n)$  can be non-monotone. In particular, one can choose the monotone pressure

$$P(\rho, n) = \rho^\gamma + n^\alpha.$$

(2) Let  $\Pi_1(\rho)$  and  $\Pi_2(n)$  be two non-monotone pressure laws for compressible Navier-Stokes equations given in [10], for instance, the virial expansion with high order terms, which can be thermodynamically unstable. Then the pressure

$$P(\rho, n) = \Pi_1(\rho) + \Pi_2(n)$$

satisfies (1.4). In particular, the pressure function  $P(\rho, n) = \Pi_1(\rho) + n^2$  is corresponding to the two-dimensional non-resistive compressible MHD [35].

(3) An example of pressure laws is

$$P(\rho, n) = \rho^\gamma(1 + 2 \cos \rho) + n^\alpha(1 + 2 \cos n),$$

which is oscillatory even for large  $\rho$  and  $n$ .

**Remark 1.3.** If (1.6) holds, then one can show  $0 \leq \underline{c}n(x, t) \leq \rho(x, t) \leq \bar{c}\rho(x, t)$  for a.e.  $(x, t) \in \mathbb{T}^d \times (0, T)$  due to the comparison principle for (1.1)<sub>1</sub>-(1.1)<sub>2</sub> so that  $G(\rho, n)$  satisfies

$$\frac{1}{C_{\underline{c}, \bar{c}}}(\rho + n)^{\max\{\gamma, \alpha\}} - C_{\underline{c}, \bar{c}} \leq G(\rho, n) \leq C_{\underline{c}, \bar{c}}(\rho + n)^{\max\{\gamma, \alpha\}} + C_{\underline{c}, \bar{c}}, \quad (1.13)$$

where  $C_{\underline{c}, \bar{c}} \geq 1$  is a constant. Indeed, we can prove under the assumptions (1.11)-(1.12) that both  $\rho$  and  $n$  are bounded in  $L^{\max\{\gamma, \alpha\} + \theta}(0, T; L^{\max\{\gamma, \alpha\} + \theta}(\mathbb{T}^d))$  for  $\max\{\gamma, \alpha\} + \theta \geq 2$  so as to apply the arguments of renormalized solutions.

Without the condition (1.6), we also have the global existence of weak solutions to the IVP (1.1)-(1.2) as follows.

**Theorem 1.2.** *Suppose that the initial data  $(\rho_0, n_0, m_0)$  satisfies*

$$\begin{cases} \rho_0(x) \geq 0, & n_0(x) \geq 0, & \text{a.e. } x \in \mathbb{T}^d, \\ \frac{m_0(x)}{\sqrt{(\rho_0 + n_0)(x)}} = 0, & \text{if } (\rho_0 + n_0)(x) = 0, & \text{a.e. } x \in \mathbb{T}^d, \\ (\rho_0, n_0, \frac{m_0}{\sqrt{\rho_0 + n_0}}) \in L^\gamma(\mathbb{T}^d) \times L^\alpha(\mathbb{T}^d) \times L^2(\mathbb{T}^d). \end{cases} \quad (1.14)$$

Then, if it holds for the pressure  $P(\rho, n)$  given by (1.4) that

$$\begin{cases} C_1 = 0, \\ \gamma, \alpha > \frac{d}{2}, & \gamma, \alpha \geq \frac{3d}{d+2}, \\ \tilde{\gamma} \leq \gamma + \theta_1, & \theta_1 := \frac{2}{d}\gamma - \frac{\gamma}{\min\{\gamma, \alpha\}} > 0, \\ \tilde{\alpha} \leq \alpha + \theta_2, & \theta_2 := \frac{2}{d}\alpha - \frac{\alpha}{\min\{\gamma, \alpha\}} > 0, \end{cases} \quad (1.15)$$

the IVP (1.1)-(1.2) admits a global weak solution  $(\rho, n, (\rho + n)u)$  in the sense of Definition 1.1.

**Remark 1.4.** *Theorem 1.2 is the first result on the global existence of weak solutions to the multi-dimensional drift-flux model (1.1) for non-monotonous pressure laws without any requirement on the relation between  $\rho_0$  and  $n_0$  as in (1.6). As pointed in [21, 50], the system (1.1) without the condition (1.6) is more realistic in some physical situations and more "two-phase" properties from mathematical points of view.*

**Remark 1.5.** *The adiabatic constants  $\gamma, \alpha$  can take  $\frac{3d}{d+2}$  for  $d = 2, 3$ , which extends the previous work by Bresch-Jabin [10] about global existence of weak solutions to compressible Navier-Stokes equations (1.8) for thermodynamically unstable pressure laws.*

**Remark 1.6.** *If (1.4) holds with  $C_1 = 0$ , then we have*

$$\frac{1}{C_0} \left( \frac{\rho^\gamma}{\gamma-1} + \frac{n^\alpha}{\alpha-1} \right) \leq G(\rho, n) \leq C_0 \left( \frac{\rho^\gamma}{\gamma-1} + \frac{n^\alpha}{\alpha-1} \right). \quad (1.16)$$

*This implies under the assumptions (1.14)-(1.15) that  $\rho$  and  $n$  belong to  $L^{\gamma+\theta_1}(0, T; L^{\gamma+\theta_1}(\mathbb{T}^d))$  for  $\gamma + \theta_1 \geq 2$  and  $L^{\alpha+\theta_2}(0, T; L^{\alpha+\theta_2}(\mathbb{T}^d))$  for  $\alpha + \theta_2 \geq 2$  respectively.*

**Remark 1.7.** *By (1.10), the condition  $C_1 = 0$  in Theorem 1.2 can be removed provided  $n_0 \leq \bar{c}\rho_0$  for some constant  $\bar{c} > 0$ .*

We explain the main strategies to prove Theorem 1.1 and Theorem 1.2. By the Faedo-Galerkin approximation and vanishing artificial viscosity, one can construct the approximate sequence  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  for  $\delta \in (0, 1)$  to the IVP (1.1)-(1.2) with the artificial pressure.

Then, we pass to the limit as  $\delta \rightarrow 0$ . As emphasized in many related papers [42, 45, 51], the key point is to show the strong convergence of the two densities  $\rho_\delta$  and  $n_\delta$ . Due to the possible non-monotonicity on  $P(\rho, n)$  in the two variables  $\rho$  and  $n$ , it is difficult to apply the techniques by Bresch-Jabin [10] to estimate  $\rho_\delta$  and  $n_\delta$  separately. To overcome this difficulty, we first make use of the ideas inspired by [45] to deduce

$$\begin{aligned} (\rho_\delta, n_\delta) &\rightarrow (\rho, n) \quad \text{in } L^1(0, T; L^1(\mathbb{T}^d)) \times L^1(0, T; L^1(\mathbb{T}^d)) \\ \iff \rho_\delta + n_\delta &\rightarrow \rho + n \quad \text{in } L^1(0, T; L^1(\mathbb{T}^d)) \quad \text{as } \delta \rightarrow 0. \end{aligned}$$

By virtue of the compactness criterion introduced in [2, 10], one needs to derive the quantitative regularity estimate of the sum  $\vartheta_\delta := \rho_\delta + n_\delta$ :

$$\frac{1}{\|\mathcal{K}_h\|_{L^1}} \int_0^T \int_{\mathbb{T}^{2d}} \mathcal{K}_h(x-y) |\vartheta_\delta^x - \vartheta_\delta^y| dx dy dt \rightarrow 0 \quad \text{uniformly in } \delta \quad \text{as } h \rightarrow 0,$$

with  $\mathcal{K}_h$  the periodic symmetric kernel given by (4.15). By the structures of the mass equations, we turn to estimate the term of  $\operatorname{div} u_\delta^x - \operatorname{div} u_\delta^y$ , which can be decomposed into the pressure part and the effective viscous flux part.

The key pressure part can be analyzed as follows:

$$\begin{aligned} P_\delta(\rho_\delta^x, n_\delta^x) - P_\delta(\rho_\delta^y, n_\delta^y) &= P_\delta(\rho_\delta^x, n_\delta^x) - P_\delta(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) \\ &\quad + P_\delta(A_{\rho^y, n^y} \vartheta_\delta^y, B_{\rho^y, n^y} \vartheta_\delta^y) - P_\delta(\rho_\delta^y, n_\delta^y) \\ &\quad + P_\delta(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y) - P_\delta(A_{\rho^y, n^y} \vartheta_\delta^x, B_{\rho^y, n^y} \vartheta_\delta^x) \\ &\quad + P_\delta(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P_\delta(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y), \end{aligned} \tag{1.17}$$

where  $(A_{\rho, n}, B_{\rho, n})$  is defined by

$$(A_{\rho, n}, B_{\rho, n}) := \begin{cases} \left( \frac{\rho}{\rho+n}, \frac{n}{\rho+n} \right), & \text{if } \rho + n > 0, \\ (0, 0), & \text{if } \rho + n = 0. \end{cases} \tag{1.18}$$

We can prove that for some domain  $\mathbb{Q} \subset \mathbb{T}^{2d} \times (0, T)$  with  $\mathbb{T}^{2d} \times (0, T)/\mathbb{Q}$  arbitrarily small, the first three terms on the right-hand side of (1.17) on  $\mathbb{Q}$  tend to 0 as  $h \rightarrow 0$  uniformly with respect to  $\delta$ , and the last term of  $P_\delta(A_{\rho, n} \vartheta_\delta, B_{\rho, n} \vartheta_\delta)$  in the one variable  $\vartheta_\delta$  can be estimated by the arguments as used in [10] with some modifications.

Meanwhile, to deal with the effective viscous flux part, we make use of the structures of the momentum equation and the commutator estimates of Riesz operator, which is different from the previous analysis [10] and may be applicable to other related models in fluid dynamics.

The rest part of this paper is arranged as follows. The global existence of weak solutions for the approximate system with two small parameters  $\varepsilon \in (0, 1)$  and  $\delta \in (0, 1)$  is obtained in Section 2. In

Section 3, we pass to the limit as  $\varepsilon \rightarrow 0$  by the compactness theorem of Lions-Feireisl. In Section 4, we prove the strong convergence of two densities by deriving quantitative regularity estimate of their sum and show the convergence of approximate sequence to an expected weak solution as  $\delta \rightarrow 0$ . In Appendix, we give some technical lemmas which are used in this paper.

## 2 Faedo–Galerkin approximation

We are ready to solve the following approximate system for  $\varepsilon \in (0, 1)$  and  $\delta \in (0, 1)$ :

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = \varepsilon \Delta \rho, \\ \partial_t n + \operatorname{div}(n u) = \varepsilon \Delta n, \\ \partial_t((\rho + n)u) + \operatorname{div}((\rho + n)u \otimes u) + \nabla P_\delta(\rho, n) + \varepsilon \nabla u \cdot \nabla(\rho + n) \\ = \mu \Delta u + (\mu + \lambda) \nabla \operatorname{div} u, \quad x \in \mathbb{T}^d, \quad t > 0, \end{cases} \quad (2.1)$$

with the initial data

$$(\rho, n, u)(x, 0) = (\rho_{0,\delta}, n_{0,\delta}, u_{0,\delta})(x), \quad x \in \mathbb{T}^d, \quad (2.2)$$

where  $P_\delta(\rho, n)$  is the artificial pressure

$$P_\delta(\rho, n) := \mathbf{1}_{\rho+n \geq \delta} P(\rho, n) + \delta(\rho + n)^{p_0}, \quad p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1, \quad (2.3)$$

and  $(\rho_{0,\delta}, n_{0,\delta}, u_{0,\delta})$  is the regularized initial data

$$(\rho_{0,\delta}, n_{0,\delta}, u_{0,\delta})(x) := (\rho_0 * j_\delta + \delta, n_0 * j_\delta + \delta, \frac{\frac{m_0}{\sqrt{\rho_0 + n_0}} * j_\delta}{\sqrt{\rho_0 * j_\delta + n_0 * j_\delta + 2\delta}})(x). \quad (2.4)$$

Here  $\mathbf{1}_{s \geq k}$  is a cut-off function satisfying

$$\mathbf{1}_{s \leq k} := \begin{cases} 1, & \text{if } 0 \leq s \leq k, \\ \text{smooth}, & \text{if } k \leq s \leq 2k, \\ 0, & \text{if } s \geq 2k, \end{cases} \quad \mathbf{1}_{s \geq k} := 1 - \mathbf{1}_{s \leq k},$$

and  $j_\delta$  is a function such that

$$\begin{cases} j_\delta \in C^\infty(\mathbb{T}^d), \quad \|j_\delta\|_{L^1} = 1, \quad 0 \leq j_\delta \leq \delta^{-\frac{1}{2p_0}}, \\ j_\delta * f \rightarrow f \quad \text{in } L^p(\mathbb{T}^d) \quad \text{as } \delta \rightarrow 0, \quad \forall f \in L^p(\mathbb{T}^d), \quad p \in [1, \infty). \end{cases} \quad (2.5)$$

It is easy to verify under the assumptions of either Theorem 1.1 or Theorem 1.2 that  $(\rho_{0,\delta}, n_{0,\delta}, (\rho_{0,\delta} + n_{0,\delta})u_{0,\delta})$  converges to  $(\rho_0, u_0, m_0)$  strongly in  $L^\gamma(\mathbb{T}^d) \times L^\alpha(\mathbb{T}^d) \times L^{\frac{2 \min\{\gamma, \alpha\}}{\min\{\gamma, \alpha\} + 1}}(\mathbb{T}^d)$  and satisfies

$$\begin{cases} \|\rho_{0,\delta}\|_{L^\gamma} \leq \|\rho_0\|_{L^\gamma}, \quad \|n_{0,\delta}\|_{L^\alpha} \leq \|n_0\|_{L^\alpha}, \quad \|\sqrt{\rho_{0,\delta} + n_{0,\delta}} u_{0,\delta}\|_{L^2} \leq \|\frac{m_0}{\sqrt{\rho_0 + n_0}}\|_{L^2}, \\ 0 < \delta \leq \rho_{0,\delta}(x), n_{0,\delta}(x) \leq C\delta^{-\frac{1}{2p_0}}, \quad \frac{1}{c_{*,\delta}} \leq \frac{n_{0,\delta}(x)}{\rho_{0,\delta}(x)} \leq c_{*,\delta}, \quad x \in \mathbb{T}^d, \quad c_{*,\delta} := C\delta^{-\frac{1}{2p_0}-1} > 1, \\ \underline{c}\rho_{0,\delta}(x) \leq n_{0,\delta}(x) \leq \bar{c}\rho_{0,\delta}(x), \quad \text{if } \underline{c}\rho_0(x) \leq n_0(x) \leq \bar{c}\rho_0(x), \quad x \in \mathbb{T}^d. \end{cases} \quad (2.6)$$



Without loss of generalization, we can assume  $P(\rho, n) \in C^2(\mathbb{R}_+ \times \mathbb{R}_+)$ . Indeed, if this is not the case, one may approximate  $P(\rho, n)$  by a regular sequence. We can compute a constant  $c_\delta \rightarrow \infty$  as  $\delta \rightarrow 0$  such that both  $\partial_\rho P_\delta(\rho, n) > 0$  and  $\partial_n P_\delta(\rho, n) > 0$  hold for any  $(\rho, n) \in \mathbb{R}_+ \times \mathbb{R}_+$  satisfying  $\rho + n \geq c_\delta > 0$ , and then set

$$\begin{cases} P_{1,\delta}(\rho, n) := P_\delta(\rho, n) + C_* \mathbf{1}_{\rho+n \leq C_* c_\delta} ((\rho + n)^{\tilde{\gamma} + \tilde{\alpha}} + \rho + n), \\ P_{2,\delta}(\rho, n) := C_* \mathbf{1}_{\rho+n \leq C_* c_\delta} ((\rho + n)^{\tilde{\gamma} + \tilde{\alpha}} + \rho + n), \end{cases} \quad (2.7)$$

with  $C_* > 1$  a sufficiently large constant. Then the pressure  $P_\delta(\rho, n)$  can be re-written as

$$P_\delta(\rho, n) = P_{1,\delta}(\rho, n) - P_{2,\delta}(\rho, n). \quad (2.8)$$

Note that  $P_{1,\delta}(\rho, n) \in C^2(\mathbb{R}_+ \times \mathbb{R}_+)$  is monotone increasing with respect to both  $\rho \geq 0$  and  $n \geq 0$ , and  $P_{2,\delta}(\rho, n) \in C^\infty(\mathbb{R}_+ \times \mathbb{R}_+)$  satisfies  $P_{2,\delta}(\rho, n) \geq 0$  and  $P_{2,\delta}(\rho, n) = 0$  for  $\rho + n \geq 2C_* c_\delta$ , so we can employ the classical Lions-Feireisl approach [24, 37] to solve the approximate system (2.1).

Let  $\psi_\ell \in C^\infty(\mathbb{T}^d)$  ( $\ell = 1, \dots$ ) be an orthonormal basis in  $L^2$ . Denote the finite dimensional spaces  $X_\ell \subset L^2$  and its projector  $\mathbf{P}_\ell : L^2 \rightarrow X_\ell$  by

$$X_\ell := \text{span}\{\psi_j\}_{j=1}^\ell, \quad \mathbf{P}_\ell f := \sum_{i=1}^\ell \psi_i \int_{\mathbb{T}^d} \psi_i f dx, \quad \forall f \in X_\ell.$$

**Proposition 2.1.** *Let  $p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1$ ,  $\delta \in (0, 1)$ ,  $\varepsilon \in (0, 1)$ , and integer  $\ell \geq 1$ . Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, there exists a unique global regular solution  $(\rho_\ell, n_\ell, u_\ell)$  for (2.1) with the initial data  $(\rho_{0,\delta}, n_{0,\delta}, \mathbf{P}_\ell u_{0,\delta})$ . In addition, it holds for any  $T > 0$  that*

$$\begin{cases} \|(\rho_\ell, n_\ell)\|_{L^\infty(0,T;L^{p_0})} + \varepsilon^{\frac{1}{2}} \|(\nabla \rho_\ell, \nabla n_\ell)\|_{L^2(0,T;L^2)} \leq C_\delta, \\ \|\sqrt{\rho_\ell + n_\ell} u_\ell\|_{L^\infty(0,T;L^2)} + \|u_\ell\|_{L^2(0,T;H^1)} \leq C_\delta, \\ 0 \leq \frac{1}{c_{*,\delta}} \rho_\ell(x, t) \leq n_\ell(x, t) \leq c_{*,\delta} \rho_\ell(x, t), \quad \text{a.e. } (x, t) \in \mathbb{T}^d \times (0, T), \\ 0 \leq \underline{c} \rho_\ell(x, t) \leq n_\ell(x, t) \leq \bar{c} \rho_\ell(x, t), \quad \text{if } \underline{c} \rho_0(x) \leq n_0(x) \leq \bar{c} \rho_0(x), \quad \text{a.e. } (x, t) \in \mathbb{T}^d \times (0, T), \end{cases} \quad (2.9)$$

where  $C_\delta > 0$  is a constant independent of  $\varepsilon$  and  $\ell$ , and  $c_{*,\delta} > 1$  is the constant given by (2.6)<sub>2</sub>.

Furthermore, we have

$$\begin{aligned} & \int_{\mathbb{T}^d} \left[ \frac{1}{2} (\rho_\ell + n_\ell) |u_\ell|^2 + G_\delta(\rho_\ell, n_\ell) \right] dx + \int_0^t \int_{\mathbb{T}^d} [\mu |\nabla u_\ell|^2 + (\mu + \lambda) (\text{div } u_\ell)^2] dx d\tau \\ & \leq \int_{\mathbb{T}^d} \left[ \frac{1}{2} (\rho_{0,\delta} + n_{0,\delta}) |u_{0,\delta}|^2 + G_\delta(\rho_{0,\delta}, n_{0,\delta}) \right] dx + C_\delta \varepsilon, \end{aligned} \quad (2.10)$$

where  $G_\delta(\rho, n)$  is defined by

$$G_\delta(\rho, n) := \begin{cases} \rho \int_1^\rho P(s, \frac{n}{\rho} s) s^{-2} ds, & \text{under the assumptions of Theorem 1.1,} \\ \rho \int_0^\rho P(s, \frac{n}{\rho} s) s^{-2} ds, & \text{under the assumptions of Theorem 1.2.} \end{cases} \quad (2.11)$$

**Proof.** By the Faedo-Galerkin approximation and fixed point arguments [23, 25, 41], there is a time  $T_\ell \in (0, T]$  such that the system (2.1) with the initial data  $(\rho_{0,\delta}, n_{0,\delta}, \mathbf{P}_\ell u_{0,\delta})$  can be solved uniquely on  $[0, T_\ell]$ . The details are omitted here.

To show  $T_\ell = T$ , we need to establish the a-priori estimates (2.9) on  $\mathbb{T}^d \times (0, T_\ell)$  uniformly in  $\ell$ . Note that  $u_\ell$  is regular since all the Sobolev norms in the finite dimensional space  $X_\ell$  are equivalent. By the comparison principle for (2.1)<sub>1</sub>-(2.1)<sub>2</sub>, we have

$$0 \leq \frac{1}{c_{*,\delta}} \rho_\ell(x, t) \leq n_\ell(x, t) \leq c_{*,\delta} \rho_\ell(x, t), \quad (x, t) \in \mathbb{T}^d \times (0, T_\ell). \quad (2.12)$$

Multiplying (1.1)<sub>1</sub>, (1.1)<sub>2</sub>, and (1.1)<sub>3</sub> by  $\partial_{\rho_\ell} G_\ell(\rho_\ell, n_\ell)$ ,  $\partial_\ell G_{n_\ell}(\rho_\ell, n_\ell)$ , and  $u_\ell$  respectively and using the fact

$$\rho \partial_\rho G_\delta(\rho, n) + n \partial_n G_\delta(\rho, n) - P_\delta(\rho, n) = 0,$$

we show for  $t \in [0, T_\ell]$  that

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{T}^d} \left( \frac{1}{2} (\rho_\ell + n_\ell) |u_\ell|^2 + G_\delta(\rho_\ell, n_\ell) \right) dx \\ & + \int_{\mathbb{T}^d} \left( \mu |\nabla u_\ell|^2 + (\mu + \lambda) (\operatorname{div} u_\ell)^2 + \varepsilon \delta p_0 \rho_\ell^{p_0-2} |\nabla \rho_\ell|^2 + \varepsilon \delta p_0 n_\ell^{p_0-2} |\nabla n_\ell|^2 \right) dx \\ & = -\varepsilon \int_{\mathbb{T}^d} \left( \partial_{\rho_\ell \rho_\ell}^2 G_\delta(\rho_\ell, n_\ell) |\nabla \rho_\ell|^2 + 2 \partial_{\rho_\ell n_\ell}^2 G_\delta(\rho_\ell, n_\ell) \nabla \rho_\ell \cdot \nabla n_\ell + \partial_{n_\ell n_\ell}^2 G_\delta(\rho_\ell, n_\ell) |\nabla n_\ell|^2 \right) dx. \end{aligned} \quad (2.13)$$

It follows from (1.13), (2.3), and (2.12) that

$$\begin{cases} G_\delta(\rho_\ell, n_\ell) \geq \frac{\delta}{p_0 - 1} (\rho_\ell + n_\ell)^{p_0} - C_\delta, \\ (|\partial_{\rho_\ell \rho_\ell}^2 G_\delta| + |\partial_{\rho_\ell n_\ell}^2 G_\delta| + |\partial_{n_\ell n_\ell}^2 G_\delta|)(\rho_\ell, n_\ell) \leq C_\delta (\rho_\ell + n_\ell)^{p_0-2} + C_\delta. \end{cases} \quad (2.14)$$

By (2.14)<sub>2</sub> and the Young inequality, the right-hand side of (2.13) can be estimates as

$$\begin{aligned} & -\varepsilon \int_{\mathbb{T}^d} \left( \partial_{\rho_\ell \rho_\ell}^2 G_\delta(\rho_\ell, n_\ell) |\nabla \rho_\ell|^2 + 2 \partial_{\rho_\ell n_\ell}^2 G_\delta(\rho_\ell, n_\ell) \nabla \rho_\ell \cdot \nabla n_\ell + \partial_{n_\ell n_\ell}^2 G_\delta(\rho_\ell, n_\ell) |\nabla n_\ell|^2 \right) dx \\ & \leq \frac{\varepsilon \delta p_0}{2} \int_{\mathbb{T}^d} (\rho_\ell^{p_0-2} |\nabla \rho_\ell|^2 + n_\ell^{p_0-2} |\nabla n_\ell|^2) dx + C_\delta \varepsilon \int_{\mathbb{T}^d} (|\nabla \rho_\ell|^2 + |\nabla n_\ell|^2) dx + C_\delta \varepsilon. \end{aligned} \quad (2.15)$$

To control the second term on the right-hand side of (2.15), one can have by (2.1)<sub>1</sub>-(2.1)<sub>2</sub> that

$$\begin{aligned} & C_\delta \frac{d}{dt} \int_{\mathbb{T}^d} (|\rho_\ell|^2 + |n_\ell|^2) dx + 2C_\delta \varepsilon \int_{\mathbb{R}^d} (|\nabla \rho_\ell|^2 + |\nabla n_\ell|^2) dx \\ & = -2C_\delta \int_{\mathbb{T}^d} (\rho_\ell + n_\ell) \operatorname{div} u_\ell dx \\ & \leq \frac{1}{2} \int_{\mathbb{T}^d} (\mu |\nabla u_\ell|^2 + (\mu + \lambda) (\operatorname{div} u_\ell)^2) dx + C_\delta \int_{\mathbb{T}^d} (|\rho_\ell|^2 + |n_\ell|^2) dx. \end{aligned} \quad (2.16)$$

Adding (2.13) and (2.15)-(2.16) together and making use of (1.3), (2.6), (2.14)<sub>1</sub>, and the Grönwall inequality, we obtain

$$\begin{aligned} & \sup_{t \in [0, T]} \int_{\mathbb{T}^d} [(\rho_\ell + n_\ell) |u_\ell|^2 + \rho_\ell^{p_0} + n_\ell^{p_0}] dx \\ & + \int_0^T \int_{\mathbb{T}^d} [|\nabla u_\ell|^2 + \varepsilon (1 + \rho_\ell^{p_0-2} + n_\ell^{p_0-2}) (|\nabla \rho_\ell|^2 + |\nabla n_\ell|^2)] dx dt \leq C_\delta, \end{aligned} \quad (2.17)$$

which together with [25, Lemma 3.2] leads to

$$\|u_\ell\|_{L^2(0,T;L^2)} \leq C_\delta. \quad (2.18)$$

By the Sobolev inequality, it also holds that

$$\begin{aligned} \|\rho_\ell + n_\ell\|_{L^{p_0+1}(0,T;L^{p_0+1})}^{p_0-1} &= \|(\rho_\ell + n_\ell)^{p_0-1}\|_{L^{\frac{p_0+1}{p_0-1}}(0,T;L^{\frac{p_0+1}{p_0-1}})} \\ &\leq \|(\rho_\ell + n_\ell)^{p_0-1}\|_{L^1(0,T;L^1)} + C\|\nabla(\rho_\ell + n_\ell)^{p_0-1}\|_{L^2(0,T;L^2)}. \end{aligned} \quad (2.19)$$

The combination of (1.3), (2.12), and (2.17)-(2.19) gives rise to  $T_\ell = T$  and (2.9). By (2.13), (2.15), and (2.17), one can show (2.10). The proof of Proposition 2.1 is completed.  $\square$

With the help of Proposition 2.1 and standard arguments (cf. [42, 45]), we have the following global existence of weak solutions to the IVP (2.1)-(2.2).

**Proposition 2.2.** *Let  $p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1$ ,  $\delta \in (0, 1)$ , and  $\varepsilon \in (0, 1)$ . Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, there exists a global weak solution  $(\rho_\varepsilon, n_\varepsilon, (\rho_\varepsilon + n_\varepsilon)u_\varepsilon)$  to the IVP (2.1)-(2.2) satisfying for any  $T > 0$  that*

$$\left\{ \begin{aligned} &\|(\rho_\varepsilon, n_\varepsilon)\|_{L^\infty(0,T;L^{p_0})} + \varepsilon^{\frac{1}{2}}\|(\nabla\rho_\varepsilon, \nabla n_\varepsilon)\|_{L^2(0,T;L^2)} \leq C_\delta, \\ &\|\sqrt{\rho_\varepsilon + n_\varepsilon}u_\varepsilon\|_{L^\infty(0,T;L^2)} + \|u_\varepsilon\|_{L^2(0,T;H^1)} \leq C_\delta, \\ &\|(\rho_\varepsilon, n_\varepsilon)\|_{L^{p_0+1}(0,T;L^{p_0+1})} \leq C_\delta, \\ &0 \leq \frac{1}{c_{*,\delta}}\rho_\varepsilon(x, t) \leq n_\varepsilon(x, t) \leq c_{*,\delta}\rho_\varepsilon(x, t), \quad \text{a.e. } (x, t) \in \mathbb{T}^d \times (0, T), \\ &0 \leq \underline{c}\rho_\varepsilon(x, t) \leq n_\varepsilon(x, t) \leq \bar{c}\rho_\varepsilon(x, t), \quad \text{if } \underline{c}\rho_0(x) \leq n_0(x) \leq \bar{c}\rho_0(x), \quad \text{a.e. } (x, t) \in \mathbb{T}^d \times (0, T), \end{aligned} \right. \quad (2.20)$$

where  $C_\delta > 0$  is a constant independent of  $\varepsilon$ , and  $c_{*,\delta} > 1$  is the constant given by (2.6)<sub>2</sub>.

Furthermore, we have

$$\begin{aligned} &\int_{\mathbb{T}^d} \left[ \frac{1}{2}(\rho_\varepsilon + n_\varepsilon)|u_\varepsilon|^2 + G_\delta(\rho_\varepsilon, n_\varepsilon) \right] dx + \int_0^t \int_{\mathbb{T}^d} [\mu|\nabla u_\varepsilon|^2 + (\mu + \lambda)(\operatorname{div} u_\varepsilon)^2] dx d\tau \\ &\leq \int_{\mathbb{T}^d} \left[ \frac{1}{2}(\rho_{0,\delta} + n_{0,\delta})|u_{0,\delta}|^2 + G_\delta(\rho_{0,\delta}, n_{0,\delta}) \right] dx + C_\delta \varepsilon, \quad \text{a.e. } t \in (0, T), \end{aligned} \quad (2.21)$$

with  $G_\delta(\rho, n)$  denoted by (2.11).

### 3 Vanishing artificial viscosity

In this section, we aim to study the following IVP:

$$\left\{ \begin{aligned} &\partial_t \rho + \operatorname{div}(\rho u) = 0, \\ &\partial_t n + \operatorname{div}(n u) = 0, \\ &\partial_t((\rho + n)u) + \operatorname{div}((\rho + n)u \otimes u) + \nabla P_\delta(\rho, n) = \mu \Delta u + (\mu + \lambda) \nabla \operatorname{div} u, \quad x \in \mathbb{T}^d, \quad t > 0, \end{aligned} \right. \quad (3.1)$$

with the initial data

$$(\rho, n, u)(x, 0) = (\rho_{0,\delta}, n_{0,\delta}, u_{0,\delta})(x), \quad x \in \mathbb{T}^d, \quad (3.2)$$

where  $P_\delta(\rho, n)$  and  $(\rho_{0,\delta}, n_{0,\delta}, u_{0,\delta})$  are given by (2.3) and (2.4) respectively.

First, by (2.20) and standard compactness arguments, we have the following convergence of weak solutions to the IVP (3.1)-(3.2).

**Lemma 3.1.** *Let  $T > 0$ ,  $p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1$ , and  $(\rho_\varepsilon, n_\varepsilon, (\rho_\varepsilon + n_\varepsilon)u_\varepsilon)$  be the weak solution to the IVP (3.1)-(3.2) for  $\varepsilon \in (0, 1)$  given by Proposition 2.2. Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, there is a limit  $(\rho, n, (\rho + n)u)$  such that as  $\varepsilon \rightarrow 0$ , it holds up to a subsequence (still denoted by  $(\rho_\varepsilon, n_\varepsilon, (\rho_\varepsilon + n_\varepsilon)u_\varepsilon)$ ) that*

$$\left\{ \begin{array}{ll} (\rho_\varepsilon, n_\varepsilon) \rightharpoonup (\rho, n) & \text{in } L^{p_0+1}(0, T; L^{p_0+1}(\mathbb{T}^d)) \times L^{p_0+1}(0, T; L^{p_0+1}(\mathbb{T}^d)), \\ \varepsilon(\nabla \rho_\varepsilon, \nabla n_\varepsilon) \rightarrow 0 & \text{in } L^2(0, T; L^2(\mathbb{T}^d)), \\ u_\varepsilon \rightharpoonup u & \text{in } L^2(0, T; H^1(\mathbb{T}^d)), \\ (\rho_\varepsilon, n_\varepsilon) \rightarrow (\rho, n) & \text{in } C([0, T]; L_{weak}^{p_0}(\mathbb{T}^d)) \times C([0, T]; L_{weak}^{p_0}(\mathbb{T}^d)), \\ (\rho_\varepsilon, n_\varepsilon) \rightarrow (\rho, n) & \text{in } C([0, T]; H^{-1}(\mathbb{T}^d)) \times C([0, T]; H^{-1}(\mathbb{T}^d)), \\ (\rho_\varepsilon + n_\varepsilon)u_\varepsilon \rightarrow (\rho + n)u & \text{in } C([0, T]; L_{weak}^{\frac{2p_0}{p_0+1}}(\mathbb{T}^d)) \cap C([0, T]; H^{-1}(\mathbb{T}^d)). \end{array} \right. \quad (3.3)$$

Before showing the strong convergence of two densities, we need to have the following lemma.

**Lemma 3.2.** *Let  $T > 0$ ,  $(\rho_\varepsilon, n_\varepsilon, (\rho_\varepsilon + n_\varepsilon)u_\varepsilon)$  be the weak solution to the IVP (2.1)-(2.2) for  $\varepsilon \in (0, 1)$  given by Proposition 2.2, and  $(\rho, n, (\rho + n)u)$  be the limit obtained by Lemma 3.1. Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, it holds*

$$\left\{ \begin{array}{ll} \lim_{\varepsilon \rightarrow 0} \int_0^T \int_{\mathbb{T}^d} (\rho_\varepsilon + n_\varepsilon) |A_{\rho_\varepsilon, n_\varepsilon} - A_{\rho, n}|^p dx dt = 0, & p \in [1, \infty), \\ \lim_{\varepsilon \rightarrow 0} \int_0^T \int_{\mathbb{T}^d} (\rho_\varepsilon + n_\varepsilon) |B_{\rho_\varepsilon, n_\varepsilon} - B_{\rho, n}|^p dx dt = 0, & p \in [1, \infty), \end{array} \right. \quad (3.4)$$

where  $(A_{\rho, n}, B_{\rho, n})$  is defined by (1.18).

Furthermore, as  $\varepsilon \rightarrow 0$ , we have

$$\begin{aligned} & (\rho_\varepsilon, n_\varepsilon) \rightarrow (\rho, n) \quad \text{in } L^1(0, T; L^1(\mathbb{T}^d)) \times L^1(0, T; L^1(\mathbb{T}^d)) \\ & \iff \rho_\varepsilon + n_\varepsilon \rightarrow \rho + n \quad \text{in } L^1(0, T; L^1(\mathbb{T}^d)). \end{aligned} \quad (3.5)$$

**Proof.** The proof of (3.4) can be found in [45]. Denote  $F_\sigma(\rho, n) := \frac{\rho^2}{\rho+n+\sigma}$  for any  $\sigma > 0$  so that it holds

$$\left\{ \begin{array}{l} |\rho \partial_\rho F_\sigma(\rho, n) + n \partial_n F_\sigma(\rho, n) - F_\sigma(\rho, n)| = \frac{\sigma \rho^2}{(\rho + n + \sigma)^2} \leq \sigma, \\ \partial_{\rho\rho}^2 F_\sigma(\rho, n) |\nabla \rho|^2 + 2 \partial_{\rho n}^2 F_\sigma(\rho, n) \nabla \rho \cdot \nabla n + \partial_{nn}^2 F_\sigma(\rho, n) |\nabla n|^2 \geq 0, \\ |\partial_\rho F_\sigma(\rho, n)| + |\partial_n F_\sigma(\rho, n)| \leq 4. \end{array} \right. \quad (3.6)$$

Let  $\eta_\sigma \in C_c^\infty(\mathbb{T}^d)$  for any  $\sigma > 0$  be the Friedrichs mollifier. Applying  $\eta_\sigma*$  to both sides of (2.1)<sub>1</sub>-(2.1)<sub>2</sub>, we obtain

$$\begin{cases} \partial_t \eta_\sigma * \rho_\varepsilon + \operatorname{div}((\eta_\sigma * \rho_\varepsilon)u_\varepsilon) = \varepsilon \Delta(\eta_\sigma * \rho_\varepsilon) + r_{1,\sigma}, \\ r_{1,\sigma} := \operatorname{div}((\eta_\sigma * \rho_\varepsilon)u_\varepsilon) - \eta_\sigma * \operatorname{div}(\rho_\varepsilon u_\varepsilon), \\ \partial_t \eta_\sigma * n_\varepsilon + \operatorname{div}((\eta_\sigma * n_\varepsilon)u_\varepsilon) = \varepsilon \Delta(\eta_\sigma * n_\varepsilon) + r_{2,\sigma}, \\ r_{2,\sigma} := \operatorname{div}((\eta_\sigma * n_\varepsilon)u_\varepsilon) - \eta_\sigma * \operatorname{div}(n_\varepsilon u_\varepsilon). \end{cases} \quad (3.7)$$

By the commutator estimates of Friedrichs mollifier (cf. [16, 36]), the terms  $r_{1,\sigma}$  and  $r_{2,\sigma}$  converge to 0 strongly in  $L^1(0, T; L^1(\mathbb{T}^d))$ . Multiplying (3.7)<sub>1</sub> and (3.7)<sub>3</sub> by  $\partial_{\eta_\sigma * \rho_\varepsilon} F_\sigma(\eta_\sigma * \rho_\varepsilon, \eta_\sigma * n_\varepsilon)$  and  $\partial_{\eta_\sigma * n_\varepsilon} F_\sigma(\eta_\sigma * \rho_\varepsilon, \eta_\sigma * n_\varepsilon)$  respectively, adding the resulted equations together, and then making use of (3.6), we have

$$\begin{aligned} & \partial_t F_\sigma(\eta_\sigma * \rho_\varepsilon, \eta_\sigma * n_\varepsilon) + \operatorname{div}(F_\sigma(\eta_\sigma * \rho_\varepsilon, \eta_\sigma * n_\varepsilon)u_\varepsilon) \\ & \leq \sigma |\operatorname{div} u_\varepsilon| + \varepsilon \Delta F_\sigma(\eta_\sigma * \rho_\varepsilon, \eta_\sigma * n_\varepsilon) + 4(|r_{1,\sigma}| + |r_{2,\sigma}|), \end{aligned}$$

which gives rise to

$$\begin{aligned} & \int_0^T \int_{\mathbb{T}^d} F_\sigma(\eta_\sigma * \rho_\varepsilon, \eta_\sigma * n_\varepsilon) dx dt \\ & \leq T \int_{\mathbb{T}^d} F_\sigma(\eta_\sigma * \rho_{0,\delta}, \eta_\sigma * n_{0,\delta}) dx + \sigma T^{\frac{3}{2}} \|\operatorname{div} u_\varepsilon\|_{L^2(0,T;L^2)} + 4T \| (r_{1,\sigma}, r_{2,\sigma}) \|_{L^1(0,T;L^1)}. \end{aligned} \quad (3.8)$$

By the dominated convergence theorem, we take the limit as  $\sigma \rightarrow 0$  in (3.8) to derive

$$\int_0^T \int_{\mathbb{T}^d} \frac{\rho_\varepsilon^2}{\rho_\varepsilon + n_\varepsilon} dx dt \leq T \int_{\mathbb{T}^d} \frac{\rho_{0,\delta}^2}{\rho_{0,\delta} + n_{0,\delta}} dx. \quad (3.9)$$

Similarly, one has

$$\int_0^T \int_{\mathbb{T}^d} \frac{\rho^2}{\rho + n} dx dt = T \int_{\mathbb{T}^d} \frac{\rho_{0,\delta}^2}{\rho_{0,\delta} + n_{0,\delta}} dx. \quad (3.10)$$

By using (3.3), (3.9)-(3.10), and

$$\begin{cases} \rho = (\rho + n)A_{\rho,n}, & \frac{\rho^2}{\rho + n} = \rho A_{\rho,n} = (\rho + n)A_{\rho,n}^2, \\ n = (\rho + n)B_{\rho,n}, & \frac{n^2}{\rho + n} = n B_{\rho,n} = (\rho + n)B_{\rho,n}^2, \end{cases} \quad (3.11)$$

we obtain

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \int_0^T \int_{\mathbb{T}^d} (\rho_\varepsilon + n_\varepsilon) |A_{\rho_\varepsilon, n_\varepsilon} - A_{\rho,n}|^2 dx dt \\ & = \lim_{\varepsilon \rightarrow 0} \int_0^T \int_{\mathbb{T}^d} \left( \frac{\rho_\varepsilon^2}{\rho_\varepsilon + n_\varepsilon} - 2\rho_\varepsilon A_{\rho,n} + (\rho_\varepsilon + n_\varepsilon) A_{\rho,n}^2 \right) dx dt \\ & \leq \int_0^T \int_{\mathbb{T}^d} \left( \frac{\rho^2}{\rho + n} - 2\rho A_{\rho,n} + (\rho + n) A_{\rho,n}^2 \right) dx dt = 0, \end{aligned}$$

which together with the Hölder inequality and the fact

$$0 \leq A_{\rho,n}, B_{\rho,n} \leq 1 \quad (3.12)$$

leads to (3.4)<sub>1</sub>. Similarly, one can verify (3.4)<sub>2</sub>.

Then, we are ready to prove (3.5). Obviously, as  $\varepsilon \rightarrow 0$ , the convergence of both  $\rho_\varepsilon$  and  $n_\varepsilon$  strongly in  $L^1(0, T; L^1(\mathbb{T}^d))$  implies the convergence of  $\rho_\varepsilon + n_\varepsilon$  strongly in  $L^1(0, T; L^1(\mathbb{T}^d))$ . Assume  $\rho_\varepsilon + n_\varepsilon \rightarrow \rho + n$  in  $L^1(0, T; L^1(\mathbb{T}^d))$  as  $\varepsilon \rightarrow 0$ . By virtue of (3.4)<sub>1</sub> for  $p = 1$  and (3.11)-(3.12), we have

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \|\rho_\varepsilon - \rho\|_{L^1(0, T; L^1)} \\ &= \lim_{\varepsilon \rightarrow 0} \|(\rho_\varepsilon + n_\varepsilon)A_{\rho_\varepsilon, n_\varepsilon} - (\rho_\varepsilon + n_\varepsilon)A_{\rho, n} + (\rho_\varepsilon + n_\varepsilon)A_{\rho, n} - (\rho + n)A_{\rho, n}\|_{L^1(0, T; L^1)} \\ &\leq \lim_{\varepsilon \rightarrow 0} (\|\rho_\varepsilon - (\rho_\varepsilon + n_\varepsilon)A_{\rho, n}\|_{L^1(0, T; L^1)} + \|\rho_\varepsilon + n_\varepsilon - \rho - n\|_{L^1(0, T; L^1)}) = 0. \end{aligned}$$

Similarly, one can show that  $n_\varepsilon$  converges to  $n$  strongly in  $L^1(0, T; L^1(\mathbb{T}^d))$  as  $\varepsilon \rightarrow 0$ . The proof of Lemma 3.2 is completed.  $\square$

We consider the cut-off function

$$T_k(s) := \begin{cases} s, & \text{if } 0 \leq s \leq k, \\ \text{smooth and concave,} & \text{if } k \leq s \leq 3k, \\ 2k, & \text{if } s \geq 3k, \end{cases} \quad (3.13)$$

which was introduced in [23] and references therein. As in [23, 37], we can prove the following property of effect viscous flux.

**Lemma 3.3.** *Let  $T > 0$ ,  $\delta \in (0, 1)$ ,  $(\rho_\varepsilon, n_\varepsilon, (\rho_\varepsilon + n_\varepsilon)u_\varepsilon)$  be the weak solution to the IVP (3.1)-(3.2) for  $\varepsilon \in (0, 1)$  given by Proposition 2.2, and  $(\rho, u, (\rho + n)u)$  be the limit obtained by Lemma 3.1. Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, for any  $k > 0$ , it holds up to a subsequence (still labelled by  $(\rho_\varepsilon, n_\varepsilon, (\rho_\varepsilon + n_\varepsilon)u_\varepsilon)$ ) that*

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \int_0^T \int_{\mathbb{T}^d} (P_\delta(\rho_\varepsilon, n_\varepsilon) - (2\mu + \lambda) \operatorname{div} u_\varepsilon) T_k(\rho_\varepsilon + n_\varepsilon) dx dt \\ &= \int_0^T \int_{\mathbb{T}^d} (\overline{P_\delta(\rho, n)} - (2\mu + \lambda) \operatorname{div} u) \overline{T_k(\rho + n)} dx dt, \end{aligned} \quad (3.14)$$

where  $\overline{f}$  denotes the weak limit of  $f_\varepsilon$  as  $\varepsilon \rightarrow 0$ .

Finally, we prove the strong convergence of two densities.

**Lemma 3.4.** *Let  $T > 0$ ,  $p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1$ ,  $(\rho_\varepsilon, n_\varepsilon, (\rho_\varepsilon + n_\varepsilon)u_\varepsilon)$  be the weak solution to the IVP (2.1)-(2.2) for  $\varepsilon \in (0, 1)$  given by Proposition 2.2, and  $(\rho, n, (\rho + n)u)$  be the limit obtained by Lemma 3.1. Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, as  $\varepsilon \rightarrow 0$ , it holds up to a subsequence (still denoted by  $(\rho_\varepsilon, n_\varepsilon)$ ) that*

$$(\rho_\varepsilon, n_\varepsilon) \rightarrow (\rho, n) \quad \text{in } L^p(0, T; L^p(\mathbb{T}^d)) \times L^p(0, T; L^p(\mathbb{T}^d)), \quad p \in [1, p_0 + 1). \quad (3.15)$$

**Proof.** Let  $\vartheta_\varepsilon := \rho_\varepsilon + n_\varepsilon$ ,  $\vartheta := \rho + n$ , and  $b_\sigma(s) := (s + \sigma) \log(s + \sigma)$  for  $s \geq 0$  and  $\sigma > 0$ . By (3.7), one has

$$\partial_t(\eta_\sigma * \vartheta_\varepsilon) + \operatorname{div}((\eta_\sigma * \vartheta_\varepsilon)u_\varepsilon) = \varepsilon \Delta(\eta_\sigma * \vartheta_\varepsilon) + r_{1,\sigma} + r_{2,\sigma}, \quad (3.16)$$

where  $\eta_\sigma \in C_c^\infty(\mathbb{T}^d)$  for  $\sigma > 0$  is the Friedrichs mollifier, and  $r_{i,\sigma}$  ( $i = 1, 2$ ) are given by (3.7). Multiplying (3.16) by  $b'(\eta_\sigma * \vartheta_\varepsilon) = 1 + \log(\eta_\sigma * \vartheta_\varepsilon + \sigma)$ , we get

$$\begin{aligned} \partial_t(b(\eta_\sigma * \vartheta_\varepsilon)) + \operatorname{div}(b(\eta_\sigma * \vartheta_\varepsilon)u_\varepsilon) + (b'(\eta_\sigma * \vartheta_\varepsilon)\eta_\sigma * \vartheta_\varepsilon - b(\eta_\sigma * \vartheta_\varepsilon)) \operatorname{div} u_\varepsilon \\ \leq \varepsilon \Delta(b(\eta_\sigma * \vartheta_\varepsilon)) + b'(\eta_\sigma * \vartheta_\varepsilon)(r_{1,\sigma} + r_{2,\sigma}). \end{aligned} \quad (3.17)$$

We integrate (3.17) over  $\mathbb{T}^d \times [0, t]$  for  $t \in [0, T]$  and take the limit as  $\sigma \rightarrow 0$  to derive

$$\int_{\mathbb{T}^d} \vartheta_\varepsilon \log \vartheta_\varepsilon dx \leq \int_{\mathbb{T}^d} (\rho_{0,\delta} + n_{0,\delta}) \log(\rho_{0,\delta} + n_{0,\delta}) dx - \int_0^t \int_{\mathbb{T}^d} \vartheta_\varepsilon \operatorname{div} u_\varepsilon dx d\tau, \quad \text{a.e. } t \in (0, T). \quad (3.18)$$

Similarly, one has

$$\int_{\mathbb{T}^d} \vartheta \log \vartheta dx = \int_{\mathbb{T}^d} (\rho_{0,\delta} + n_{0,\delta}) \log(\rho_{0,\delta} + n_{0,\delta}) dx - \int_0^t \int_{\mathbb{T}^d} \vartheta \operatorname{div} u dx d\tau, \quad \text{a.e. } t \in (0, T). \quad (3.19)$$

Let  $\bar{f}$  denote the weak limit of  $f_\varepsilon$  as  $\varepsilon \rightarrow 0$ , and  $P_{1,\delta}(\rho, n)$ ,  $P_{2,\delta}(\rho, n)$ ,  $(A_{\rho,n}, B_{\rho,n})$ , and  $T_k(f)$  be defined by (2.7)<sub>1</sub>, (2.7)<sub>2</sub>, (1.18), and (3.13) respectively. It can be shown by (2.8), (3.11), (3.14), and (3.18)-(3.19) that

$$\begin{aligned} \int_{\mathbb{T}^d} (\overline{\vartheta \log \vartheta} - \vartheta \log \vartheta) dx \\ \leq \lim_{\varepsilon \rightarrow 0} \int_0^t \int_{\mathbb{T}^d} (\vartheta \operatorname{div} u - \vartheta_\varepsilon \operatorname{div} u_\varepsilon) dx d\tau = \sum_{i=1}^4 I_i^1, \quad \text{a.e. } t \in (0, T), \end{aligned} \quad (3.20)$$

where  $I_i^1$  ( $i = 1, 2, 3, 4$ ) are given by

$$\begin{aligned} I_1^1 &:= \lim_{\varepsilon \rightarrow 0} \int_0^t \int_{\mathbb{T}^d} [(\vartheta - \overline{T_k(\vartheta)}) \operatorname{div} u - (\vartheta_\varepsilon - T_k(\vartheta_\varepsilon)) \operatorname{div} u_\varepsilon] dx d\tau, \\ I_2^1 &:= \frac{1}{2\mu + \lambda} \lim_{\varepsilon \rightarrow 0} \int_0^t \int_{\mathbb{T}^d} [P_\delta(\rho_\varepsilon, n_\varepsilon) - P_\delta(A_{\rho,n}\vartheta_\varepsilon, B_{\rho,n}\vartheta_\varepsilon)] (\overline{T_k(\vartheta)} - T_k(\vartheta_\varepsilon)) dx d\tau, \\ I_3^1 &:= \frac{1}{2\mu + \lambda} \lim_{\varepsilon \rightarrow 0} \int_0^t \int_{\mathbb{T}^d} P_{1,\delta}(A_{\rho,n}\vartheta_\varepsilon, B_{\rho,n}\vartheta_\varepsilon) (\overline{T_k(\vartheta)} - T_k(\vartheta_\varepsilon)) dx d\tau, \\ I_4^1 &:= \frac{1}{2\mu + \lambda} \lim_{\varepsilon \rightarrow 0} \int_0^t \int_{\mathbb{T}^d} P_{2,\delta}(A_{\rho,n}\vartheta_\varepsilon, B_{\rho,n}\vartheta_\varepsilon) (T_k(\vartheta_\varepsilon) - \overline{T_k(\vartheta)}) dx d\tau. \end{aligned}$$

It follows from (2.20), (3.3), and lower semi-continuity of the weak limit  $\vartheta - \overline{T_k(\vartheta)}$  that

$$\begin{aligned} I_1^1 &\leq \sup_{\varepsilon \rightarrow 0} \lim (\|\vartheta - \overline{T_k(\vartheta)}\|_{L^2(0,T;L^2)} \|\operatorname{div} u\|_{L^2(0,T;L^2)} + \|\vartheta_\varepsilon - T_k(\vartheta_\varepsilon)\|_{L^2(0,T;L^2)} \|\operatorname{div} u_\varepsilon\|_{L^2(0,T;L^2)}) \\ &\leq C_\delta \sup_{\varepsilon \rightarrow 0} \lim \|\vartheta_\varepsilon - T_k(\vartheta_\varepsilon)\|_{L^2(0,T;L^2)} \\ &\leq \frac{C_\delta \|\vartheta_\delta\|_{L^{\frac{p_0+1}{2}}(0,T;L^{p_0+1})}}{k^{\frac{p_0-1}{2}}} \leq \frac{C_\delta}{k^{\frac{p_0-1}{2}}}. \end{aligned}$$

For any  $x_1, x_2, y_1, y_2 \geq 0$  and  $F(x, y) \in C^1(\mathbb{R}_+ \times \mathbb{R}_+)$ , we have

$$\begin{aligned} F(x_1, y_1) - F(x_2, y_2) &= (x_1 - x_2) \int_0^1 \partial_x F(x_2 + \theta(x_1 - x_2), y_2 + \theta(y_1 - y_2)) d\theta \\ &\quad + (y_1 - y_2) \int_0^1 \partial_y F(x_2 + \theta(x_1 - x_2), y_2 + \theta(y_1 - y_2)) d\theta, \end{aligned} \quad (3.21)$$

which together with (1.4), (2.3), (2.20)<sub>4</sub>, and (3.4) yields

$$\begin{aligned} I_2^1 &\leq Ck \lim_{\varepsilon \rightarrow 0} \int_0^t \int_{\mathbb{T}^d} (\vartheta_\varepsilon + 1)^{p_0-1} \vartheta_\varepsilon^{1-\frac{1}{p_0}} \vartheta_\varepsilon^{\frac{1}{p_0}} (|A_{\rho_\varepsilon, n_\varepsilon} - A_{\rho, n}| + |B_{\rho_\varepsilon, n_\varepsilon} - B_{\rho, n}|) dx d\tau \\ &\leq Ck \sup_{\varepsilon \rightarrow 0} \lim \left( \int_0^t \int_{\mathbb{T}^d} (\vartheta_\varepsilon + 1)^{p_0+1} dx d\tau \right)^{\frac{p_0-1}{p_0}} \\ &\quad \times \lim_{\varepsilon \rightarrow 0} \left( \int_0^t \int_{\mathbb{T}^d} \vartheta_\varepsilon (|A_{\rho_\varepsilon, n_\varepsilon} - A_{\rho, n}|^{p_0} + |B_{\rho_\varepsilon, n_\varepsilon} - B_{\rho, n}|^{p_0}) dx d\tau \right)^{\frac{1}{p_0}} = 0. \end{aligned} \quad (3.22)$$

Noticing that both  $T_k(s)$  and  $P_{1,\delta}(A_{\rho,n}s, B_{\rho,n}s)$  are monotone increasing with respect to  $s \geq 0$ , we apply [26, Theorem 10.19] to have

$$I_3^1 \leq 0. \quad (3.23)$$

Substituting the above estimates of  $I_i^1$  ( $i = 1, 2, 3$ ) into (3.20) and taking the limit as  $k \rightarrow \infty$ , we deduce

$$\int_{\mathbb{T}^d} (\overline{\vartheta \log \vartheta} - \vartheta \log \vartheta) dx \leq \frac{1}{2\mu + \lambda} \lim_{\varepsilon \rightarrow 0} \int_0^t \int_{\mathbb{T}^d} P_{2,\delta}(A_{\rho,n}\vartheta_\varepsilon, B_{\rho,n}\vartheta_\varepsilon) (\vartheta_\varepsilon - \vartheta) dx d\tau. \quad (3.24)$$

Since  $P_{2,\delta}(sA_{\rho,n}, sB_{\rho,n})$  has compact support in  $\{s \in \mathbb{R}_+ \mid s \leq 2C_*c_\delta\}$  due to (2.7)<sub>2</sub>, there is a constant  $C_{*,\delta} > 0$  such that both  $C_{*,\delta}s \log s - sP_{2,\delta}(A_{\rho,n}s, B_{\rho,n}s)$  and  $C_{*,\delta}s \log s + P_{2,\delta}(A_{\rho,n}s, B_{\rho,n}s)$  are strictly convex for any  $s \geq 0$ , and therefore we get

$$\begin{aligned} &\lim_{\varepsilon \rightarrow 0} \int_0^t \int_{\mathbb{T}^d} [P_{2,\delta}(A_{\rho,n}\vartheta_\varepsilon, B_{\rho,n}\vartheta_\varepsilon)\vartheta_\varepsilon - P_{2,\delta}(\rho, n)\vartheta] dx d\tau \\ &\leq C_{*,\delta} \int_0^t \int_{\mathbb{T}^d} (\overline{\vartheta \log \vartheta} - \vartheta \log \vartheta) dx d\tau, \end{aligned} \quad (3.25)$$

and

$$\begin{aligned} &\lim_{\varepsilon \rightarrow 0} \int_0^t \int_{\mathbb{T}^d} [P_{2,\delta}(\rho, n) - P_{2,\delta}(A_{\rho,n}\vartheta_\varepsilon, B_{\rho,n}\vartheta_\varepsilon)] \vartheta dx d\tau \\ &\leq \lim_{\varepsilon \rightarrow 0} \int_{\{(x,\tau) \in \mathbb{T}^d \times [0,t] \mid \vartheta \leq 2C_*c_\delta\}} [P_{2,\delta}(A_{\rho,n}\vartheta, B_{\rho,n}\vartheta) - P_{2,\delta}(A_{\rho,n}\vartheta_\varepsilon, B_{\rho,n}\vartheta_\varepsilon)] \vartheta dx d\tau \\ &\leq C_{*,\delta} \lim_{\varepsilon \rightarrow 0} \int_{\{(x,\tau) \in \mathbb{T}^d \times [0,t] \mid \vartheta \leq 2C_*c_\delta\}} (\vartheta_\varepsilon \log \vartheta_\varepsilon - \vartheta \log \vartheta) \vartheta dx d\tau \\ &\leq 2C_{*,\delta} C_*c_\delta \int_0^t \int_{\mathbb{T}^d} (\overline{\vartheta \log \vartheta} - \vartheta \log \vartheta) dx d\tau, \end{aligned} \quad (3.26)$$

where one has used (3.11),  $P_{2,\delta}(sA_{\rho,n}, sB_{\rho,n}) \geq 0$ , the compact support of  $P_{2,\delta}(sA_{\rho,n}, sB_{\rho,n})$ , and the convexity of  $C_{*,\delta}s \log s - P_{2,\delta}(sA_{\rho,n}, sB_{\rho,n})$ . By (3.24)-(3.26), we have

$$\int_{\mathbb{T}^d} (\overline{\vartheta \log \vartheta} - \vartheta \log \vartheta) dx \leq \frac{C_{*,\delta}(2C_*c_\delta + 1)}{2\mu + \lambda} \int_0^t \int_{\mathbb{T}^d} (\overline{\vartheta \log \vartheta} - \vartheta \log \vartheta) dx d\tau. \quad (3.27)$$



The combination of (2.20)<sub>4</sub>, (3.5), (3.27), the Grönwall inequality, and the convexity of  $s \log s$  gives rise to (3.15). The proof of Lemma 3.4 is completed.  $\square$

With the help of Proposition 2.2 and Lemmas 3.1-3.4, we have the global existence of weak solutions to the IVP (3.1)-(3.2).

**Proposition 3.1.** *Let  $p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1$  and  $\delta \in (0, 1)$ . Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, there exists a global weak solution  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  to the IVP (3.1)-(3.2) satisfying for any  $T > 0$  that*

$$\begin{cases} \|\rho_\delta\|_{L^\infty(0,T;L^\gamma)} + \|n_\delta\|_{L^\infty(0,T;L^\alpha)} + \delta^{\frac{1}{p_0}} \|(\rho_\delta, n_\delta)\|_{L^\infty(0,T;L^{p_0})} \leq C, \\ \|\sqrt{\rho_\delta + n_\delta}u_\delta\|_{L^\infty(0,T;L^2)} + \|u_\delta\|_{L^2(0,T;H^1)} \leq C, \\ \rho_\delta(x, t) \geq 0, \quad n_\delta(x, t) \geq 0, \quad \text{a.e. } (x, t) \in \mathbb{T}^d \times (0, T), \\ \underline{\rho}_\delta(x, t) \leq n_\delta(x, t) \leq \bar{c}\rho_\delta(x, t), \quad \text{if } \underline{\rho}_0(x) \leq n_0(x) \leq \bar{c}\rho_0(x), \quad \text{a.e. } (x, t) \in \mathbb{T}^d \times (0, T), \end{cases} \quad (3.28)$$

where  $C > 0$  is a constant independent of  $\delta$ .

Furthermore, it holds

$$\begin{aligned} & \int_{\mathbb{T}^d} \left[ \frac{1}{2}(\rho_\delta + n_\delta)|u_\delta|^2 + G_\delta(\rho_\delta, n_\delta) \right] dx + \int_0^t \int_{\mathbb{T}^d} [\mu|\nabla u_\delta|^2 + (\mu + \lambda)(\operatorname{div} u_\delta)^2] dx d\tau \\ & \leq \int_{\mathbb{T}^d} \left[ \frac{1}{2}(\rho_{0,\delta} + n_{0,\delta})|u_{0,\delta}|^2 + G_\delta(\rho_{0,\delta}, n_{0,\delta}) \right] dx, \quad \text{a.e. } t \in (0, T), \end{aligned} \quad (3.29)$$

where  $G_\delta(\rho, n)$  is given by (2.11).

## 4 Vanishing artificial pressure

### 4.1 Higher integrability of densities

In this section, we prove Theorem 1.1 and Theorem 1.2 by taking the limit in (3.1) as  $\delta \rightarrow 0$ . Since the uniform estimate (3.28) are not enough to show the convergence of  $P_\delta(\rho_\delta, n_\delta)$  to  $P(\rho, n)$  as  $\delta \rightarrow 0$ , we need the higher integrability of  $\rho_\delta$  and  $n_\delta$  uniformly in  $\delta$  as follows.

**Lemma 4.1.** *Let  $T > 0$ ,  $p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1$ ,  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  be the weak solution to the IVP (3.1)-(3.2) for  $\delta \in (0, 1)$  given by Proposition 3.1, and  $\vartheta_\delta := \rho_\delta + n_\delta$ . Then, under the assumptions of Theorem 1.1, it holds*

$$\int_0^T \int_{\mathbb{T}^d} (\vartheta_\delta^{\max\{\gamma, \alpha\} + \theta} + \delta \vartheta_\delta^{p_0 + \theta}) dx dt \leq C, \quad (4.1)$$

with  $C > 0$  a constant independent of  $\delta$  and

$$\theta := \frac{2}{d} \max\{\gamma, \alpha\} - 1 > 0.$$

In addition, under the assumptions of Theorem 1.2, we have

$$\int_0^T \int_{\mathbb{T}^d} (\rho_\delta^{\gamma+\theta_1} + n_\delta^{\alpha+\theta_2} + \delta \vartheta_\delta^{p_0+\theta_1} + \delta \vartheta_\delta^{p_0+\theta_2}) dx dt \leq C, \quad (4.2)$$

with

$$\theta_1 := \frac{2}{d}\gamma - \frac{\gamma}{\min\{\gamma, \alpha\}} > 0, \quad \theta_2 := \frac{2}{d}\alpha - \frac{\alpha}{\min\{\gamma, \alpha\}} > 0.$$

**Proof.** We are ready to show (4.2), and (4.1) can be proved by similar arguments. Denote by  $F_\delta$  the effect viscous flux as

$$F_\delta := P_\delta(\rho_\delta, n_\delta) - \frac{1}{|\mathbb{T}^d|} \int_{\mathbb{T}^d} P_\delta(\rho_\delta, n_\delta) dx - (2\mu + \lambda) \operatorname{div} u_\delta. \quad (4.3)$$

One deduces from the equation (3.1)<sub>3</sub> that

$$F_\delta = (-\Delta)^{-1} \operatorname{div} [\partial_t (\vartheta_\delta u_\delta) + \operatorname{div} (\vartheta_\delta u_\delta \otimes u_\delta)], \quad (4.4)$$

where the operator  $(-\Delta)^{-1} : W^{k-2,p}(\mathbb{T}^d) \rightarrow W^{k,p}(\mathbb{T}^d)$  for  $k \in \mathbb{R}$  and  $p \in (1, \infty)$  is defined by

$$(-\Delta)^{-1} f = g, \quad g \text{ is the solution of the problem } -\Delta g = f, \quad \int_{\mathbb{T}^d} f = 0. \quad (4.5)$$

It can be verified by the Sobolev inequality and  $L^p(\mathbb{T}^d)$ - $L^p(\mathbb{T}^d)$  boundedness of the operator  $\nabla(-\Delta)^{-1} \operatorname{div}$  that

$$\|(-\Delta)^{-1} \operatorname{div} f\|_{L^{\frac{pd}{d-p}}} \leq C \|\nabla(-\Delta)^{-1} \operatorname{div} f\|_{L^p} \leq C \|f\|_{L^p}, \quad \forall f \in L^p, \quad p \in (1, d). \quad (4.6)$$

By (3.1) and (4.4), we obtain

$$\int_0^T \int_{\mathbb{T}^d} \rho_\delta^\theta P_\delta(\rho_\delta, n_\delta) dx dt = \sum_{i=1}^4 I_i^2, \quad (4.7)$$

with

$$\begin{aligned} I_1^2 &:= \int_0^T \int_{\mathbb{T}^d} \rho_\delta^\theta \left( \frac{1}{|\mathbb{T}^d|} \int_{\mathbb{T}^d} P_\delta(\rho_\delta, n_\delta) dx + (2\mu + \lambda) \operatorname{div} u_\delta \right) dx dt, \\ I_2^2 &:= \int_{\mathbb{T}^d} \rho_\delta^\theta (-\Delta)^{-1} \operatorname{div} (\vartheta_\delta u_\delta) dx \Big|_0^T, \\ I_3^2 &:= (\theta - 1) \int_0^T \int_{\mathbb{T}^d} \rho_\delta^\theta \operatorname{div} u_\delta (-\Delta)^{-1} \operatorname{div} (\vartheta_\delta u_\delta) dx dt, \\ I_4^2 &:= \sum_{i,j=1}^d \int_0^T \int_{\mathbb{T}^d} \rho_\delta^\theta [\mathcal{R}_i \mathcal{R}_j (\vartheta_\delta u_\delta^i u_\delta^j) - u_\delta^j \mathcal{R}_i \mathcal{R}_j (\vartheta_\delta u_\delta^i)] dx dt, \quad R_i := (-\Delta)^{-\frac{1}{2}} \partial_i, \quad i = 1, \dots, d. \end{aligned}$$

First, one deduces by (3.28) and  $2\theta_1 < \gamma + \theta_1$  that

$$\begin{aligned} I_1^2 &\leq T \|\rho_\delta^\theta\|_{L^\infty(0,T;L^1)} \|P_\delta(\rho_\delta, n_\delta)\|_{L^\infty(0,T;L^1)} + (2\mu + \lambda) T^{\frac{1}{2}} \|\rho_\delta^\theta\|_{L^2(0,T;L^2)} \|\operatorname{div} u_\delta\|_{L^2(0,T;L^2)} \\ &\leq C \left(1 + \frac{1}{\varepsilon_0}\right) + \varepsilon_0 \int_0^T \int_{\mathbb{T}^d} \rho_\delta^{\gamma+\theta_1} dx dt, \end{aligned}$$

with the constant  $\varepsilon_0 > 0$  to be chosen later. By (3.28), one has

$$\|\vartheta_\delta\|_{L^\infty(0,T;L^{\gamma_0})} + \|\vartheta_\delta u_\delta\|_{L^\infty(0,T;L^{\frac{2\gamma_0}{\gamma_0+1}})} \leq C, \quad (4.8)$$

where we used the convenient notation

$$\gamma_0 := \min\{\gamma, \alpha\}.$$

For the term  $I_2^2$ , making use of (2.6), (3.28), (4.6), (4.8), and  $\frac{1}{d} + 1 - \frac{\theta_1}{\gamma} > \frac{\gamma_0+1}{2\gamma_0}$ , we have

$$\begin{aligned} I_2^2 &\leq \|\rho_\delta^{\theta_1}\|_{L^\infty(0,T;L^{\frac{\gamma}{\theta_1}})} \|(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)\|_{L^\infty(0,T;L^{\frac{2\gamma_0 d}{(d-2)\gamma_0+d}}} \\ &\quad + \|\rho_{0,\delta}\|_{L^{\frac{\gamma}{\theta_1}}} \|(-\Delta)^{-1} \operatorname{div}((\rho_{0,\delta} + n_{0,\delta})u_{0,\delta})\|_{L^{\frac{2\gamma_0 d}{(d-2)\gamma_0+d}}} \\ &\leq C\|\rho_\delta\|_{L^\infty(0,T;L^\gamma)}^{\theta_1} \|\vartheta_\delta u_\delta\|_{L^\infty(0,T;L^{\frac{2\gamma_0}{\gamma_0+1}}} + C\|\rho_{0,\delta}\|_{L^\gamma}^\theta \|(\rho_{0,\delta} + n_{0,\delta})u_{0,\delta}\|_{L^{\frac{2\gamma_0}{\gamma_0+1}}} \leq C. \end{aligned}$$

As in [37], we analyse the terms  $I_i^2$  ( $i = 3, 4$ ) for two cases  $d \geq 3$  and  $d = 2$  separately:

- Case 1:  $d \geq 3$ .

Since it holds  $\frac{\theta_1}{\gamma} + \frac{1}{\gamma_0} + \frac{d-2}{2d} + \frac{d-2}{2d} = 1$ , we obtain by (3.28), (4.8), the Sobolev embedding  $H^1(\mathbb{R}^d) \hookrightarrow L^{\frac{2d}{d-2}}(\mathbb{R}^d)$ , and the boundedness of Riesz operator that

$$I_4^2 \leq C\|\rho_\delta^{\theta_1}\|_{L^\infty(0,T;L^{\frac{\gamma}{\theta_1}})} \|\vartheta_\delta\|_{L^\infty(0,T;L^{\gamma_0})} \|u_\delta\|_{L^2(0,T;L^{\frac{2d}{d-2}}}^2 \leq C.$$

Similarly, the term  $I_3^2$  for  $d \geq 4$  can be estimated by

$$\begin{aligned} I_3^2 &\leq \|\rho_\delta^{\theta_1}\|_{L^\infty(0,T;L^{\frac{\gamma}{\theta_1}})} \|\operatorname{div} u_\delta\|_{L^2(0,T;L^2)} \|(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)\|_{L^2(0,T;L^{\frac{2\gamma_0 d}{(d-4)\gamma_0+2d}}} \\ &\leq C\|\rho_\delta\|_{L^\infty(0,T;L^\gamma)}^{\theta_1} \|\operatorname{div} u_\delta\|_{L^2(0,T;L^2)} \|\vartheta_\delta\|_{L^\infty(0,T;L^{\gamma_0})} \|u_\delta\|_{L^2(0,T;L^{\frac{2d}{d-2}}} \leq C. \end{aligned}$$

For  $d = 3$ , it follows from (3.28), (4.6), (4.8), and the Sobolev embedding  $H^1(\mathbb{T}^3) \hookrightarrow L^6(\mathbb{T}^3)$  that

$$\begin{aligned} I_3^2 &\leq C\|\rho_\delta^{\theta_1}\|_{L^{\frac{5\gamma_0-3}{2\gamma_0-3}}(0,T;L^{\frac{5\gamma_0-3}{2\gamma_0-3}})} \|\operatorname{div} u_\delta\|_{L^2(0,T;L^2)} \|\vartheta_\delta u_\delta\|_{L^{\frac{2(5\gamma_0-3)}{\gamma_0+3}}(0,T;L^{\frac{6(5\gamma_0-3)}{13\gamma_0+3}}} \\ &\leq C\|\rho_\delta^{\theta_1}\|_{L^{\frac{\gamma+\theta_1}{\theta_1}}(0,T;L^{\frac{\gamma+\theta_1}{\theta_1}})} \|\operatorname{div} u_\delta\|_{L^2(0,T;L^2)} \|\vartheta_\delta\|_{L^\infty(0,T;L^{\gamma_0})}^{\frac{\gamma_0+3}{5\gamma_0-3}} \|u_\delta\|_{L^2(0,T;L^6)}^{\frac{\gamma_0+3}{5\gamma_0-3}} \|\vartheta_\delta u_\delta\|_{L^\infty(0,T;L^{\frac{2\gamma_0}{\gamma_0+1}}} \\ &\leq C\left(1 + \frac{1}{\varepsilon_0}\right) + \varepsilon_0 \int_0^T \int_{\mathbb{T}^3} \rho_\delta^{\gamma+\theta_1} dx dt, \end{aligned}$$

- Case 2:  $d = 2$ .

One can show by (3.28) and (4.6) that

$$\begin{aligned} I_3^2 &\leq C\|\rho_\delta^{\theta_1}\|_{L^{\frac{2\gamma_0-1}{\gamma_0-1}}(0,T;L^{\frac{2\gamma_0-1}{\gamma_0-1}})} \|\operatorname{div} u_\delta\|_{L^2(0,T;L^2)} \|\vartheta_\delta u_\delta\|_{L^{2(2\gamma_0-1)}(0,T;L^{\frac{2\gamma_0}{\gamma_0+1}}} \\ &\leq C\|\rho_\delta\|_{L^{\gamma+\theta_1}(0,T;L^{\gamma+\theta_1})}^{\theta_1} \|\operatorname{div} u_\delta\|_{L^2(0,T;L^2)} \|\sqrt{\vartheta_\delta}\|_{L^{2(2\gamma_0-1)}(0,T;L^{2(2\gamma_0-1)}} \|\sqrt{\vartheta_\delta} u_\delta\|_{L^\infty(0,T;L^2)} \\ &\leq C\left(1 + \frac{1}{\varepsilon_0}\right) + \varepsilon_0 \int_0^T \int_{\mathbb{T}^2} (\rho_\delta^{\gamma+\theta_1} + n_\delta^{\alpha+\theta_2}) dx dt. \end{aligned}$$

In addition, one deduces by the critical Sobolev embedding  $H^1(\mathbb{T}^2) \hookrightarrow \mathcal{BMO}(\mathbb{T}^2)$  for  $d = 2$  and the commutator estimate (A.11) that

$$\begin{aligned}
I_4^2 &\leq \sum_{i,j=1}^2 \|\rho_\delta^{\theta_1}\|_{L^{\frac{2\gamma_0-1}{\gamma_0-1}}(0,T;L^{\frac{2\gamma_0-1}{\gamma_0-1}})} \|\mathcal{R}_i \mathcal{R}_j (\vartheta_\delta u_\delta^i u_\delta^j) - u_\delta^j \mathcal{R}_i \mathcal{R}_j (\vartheta_\delta u_\delta^i)\|_{L^{\frac{2\gamma_0-1}{\gamma_0}}(0,T;L^{\frac{2\gamma_0-1}{\gamma_0}})} \\
&\leq C \|\rho_\delta\|_{L^{\gamma+\theta_1}(0,T;L^{\gamma+\theta_1})}^{\theta_1} \|u_\delta\|_{L^2(0,T;\mathcal{BMO})} \|\vartheta_\delta u_\delta\|_{L^{2(2\gamma_0-1)}(0,T;L^{\frac{2\gamma_0-1}{\gamma_0-1}})} \\
&\leq C \|\rho_\delta\|_{L^{\gamma+\theta_1}(0,T;L^{\gamma+\theta_1})}^{\theta_1} \|\sqrt{\vartheta_\delta}\|_{L^{2(2\gamma_0-1)}(0,T;L^{2(2\gamma_0-1)})} \|u_\delta\|_{L^2(0,T;H^1)} \|\sqrt{\vartheta_\delta} u_\delta\|_{L^\infty(0,T;L^2)} \\
&\leq C(1 + \frac{1}{\varepsilon_0}) + \varepsilon_0 \int_0^T \int_{\mathbb{T}^2} (\rho_\delta^{\gamma+\theta_1} + n_\delta^{\alpha+\theta_2}) dxdt.
\end{aligned}$$

Substituting the above estimates of  $I_i^2$  ( $i = 1, 2, 3, 4$ ) into (4.7), we obtain

$$\int_0^T \int_{\mathbb{T}^d} \rho_\delta^{\theta_1} P_\delta(\rho_\delta, n_\delta) dxdt \leq C(1 + \frac{1}{\varepsilon_0}) + 3\varepsilon_0 \int_0^T \int_{\mathbb{T}^d} (\rho_\delta^{\gamma+\theta_1} + n_\delta^{\alpha+\theta_2}) dxdt. \quad (4.9)$$

Similarly, one can show

$$\int_0^T \int_{\mathbb{T}^d} n_\delta^{\theta_2} P_\delta(\rho_\delta, n_\delta) dxdt \leq C(1 + \frac{1}{\varepsilon_0}) + 3\varepsilon_0 \int_0^T \int_{\mathbb{T}^d} (\rho_\delta^{\gamma+\theta_1} + n_\delta^{\alpha+\theta_2}) dxdt. \quad (4.10)$$

By (1.4) and (2.3), it holds

$$\begin{aligned}
&(\rho_\delta^{\theta_1} + n_\delta^{\theta_2}) P_\delta(\rho_\delta, n_\delta) \\
&\geq (\rho_\delta^{\theta_1} + n_\delta^{\theta_2}) (P(\rho_\delta, n_\delta) + \delta(\rho_\delta + n_\delta)^{p_0}) - \mathbf{1}_{\vartheta_\delta \leq \delta} (\rho_\delta^{\theta_1} + n_\delta^{\theta_2}) P(\rho_\delta, n_\delta) \\
&\geq \frac{1}{2C_1} (\rho_\delta^{\gamma+\theta_1} + n_\delta^{\alpha+\theta_2}) + \delta(\rho_\delta^{p_0+\theta_1} + n_\delta^{p_0+\theta_2}) - C.
\end{aligned} \quad (4.11)$$

Combining (4.9)-(4.11) together and choosing  $\varepsilon_0 = \frac{1}{24C_1}$ , we derive (4.2). The proof of Lemma 4.1 is completed.  $\square$

By (3.28) and (4.1), we have the convergence of weak solutions to the IVP (3.1)-(3.2).

**Lemma 4.2.** *Let  $T > 0$ ,  $p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1$ , and  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  be the weak solution to the IVP (3.1)-(3.2) for  $\delta \in (0, 1)$  given by Proposition 3.1. Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, there is a limit  $(\rho, n, (\rho + n)u)$  such that as  $\delta \rightarrow 0$ , it holds up to a subsequence (still denoted by  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$ ) that*

$$\left\{ \begin{array}{ll}
(\rho_\delta, n_\delta) \xrightarrow{*} (\rho, n) & \text{in } L^\infty(0, T; L^\gamma(\mathbb{T}^d)) \times L^\infty(0, T; L^\alpha(\mathbb{T}^d)), \\
\delta(\rho_\delta + n_\delta)^{p_0} \rightarrow 0 & \text{in } L^1(0, T; L^1(\mathbb{T}^d)), \\
u_\delta \rightharpoonup u & \text{in } L^2(0, T; H^1(\mathbb{T}^d)), \\
(\rho_\delta, n_\delta) \rightarrow (\rho, n) & \text{in } C([0, T]; L_{weak}^\gamma(\mathbb{T}^d)) \times C([0, T]; L_{weak}^\alpha(\mathbb{T}^d)), \\
(\rho_\delta, n_\delta) \rightarrow (\rho, n) & \text{in } C([0, T]; H^{-1}(\mathbb{T}^d)) \times C([0, T]; H^{-1}(\mathbb{T}^d)), \\
(\rho_\delta + n_\delta)u_\delta \rightarrow (\rho + n)u & \text{in } C([0, T]; L_{weak}^{\frac{2 \min\{\gamma, \alpha\}}{\min\{\gamma, \alpha\} + 1}}(\mathbb{T}^d)) \cap C([0, T]; H^{-1}(\mathbb{T}^d)).
\end{array} \right. \quad (4.12)$$

## 4.2 Strong convergence of densities

By similar arguments as used in Lemma 3.2, the strong convergence of two densities  $\rho_\delta$  and  $n_\delta$  and the strong convergence of their sum  $\rho_\delta + n_\delta$  are equivalent.

**Lemma 4.3.** *Let  $T > 0$ ,  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  be the weak solution to the IVP (3.1)-(3.2) for  $\delta \in (0, 1)$  given by Proposition 3.1, and  $(\rho, n, (\rho + n)u)$  be the limit obtained by Lemma 4.2. Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, as  $\delta \rightarrow 0$ , it holds*

$$\begin{cases} \rho_\delta - (\rho_\delta + n_\delta)A_{\rho, n} \rightarrow 0 & \text{in } L^1(0, T; L^1(\mathbb{T}^d)), \\ n_\delta - (\rho_\delta + n_\delta)B_{\rho, n} \rightarrow 0 & \text{in } L^1(0, T; L^1(\mathbb{T}^d)), \end{cases} \quad (4.13)$$

where  $(A_{\rho, n}, B_{\rho, n})$  is defined by (1.18).

Furthermore, as  $\delta \rightarrow 0$ , we have

$$\begin{aligned} (\rho_\delta, n_\delta) &\rightarrow (\rho, n) && \text{in } L^1(0, T; L^1(\mathbb{T}^d)) \times L^1(0, T; L^1(\mathbb{T}^d)) \\ \iff \rho_\delta + n_\delta &\rightarrow \rho + n && \text{in } L^1(0, T; L^1(\mathbb{T}^d)). \end{aligned} \quad (4.14)$$

As in [2, 10, 12], introduce the symmetric periodic kernel

$$\mathcal{K}_h(x) := \begin{cases} \frac{1}{(h+|x|)^d}, & \text{if } 0 \leq |x| \leq \frac{1}{2}, \\ \text{smooth}, & \text{if } \frac{1}{2} \leq |x| \leq \frac{2}{3}, \\ \text{smooth and independent of } h, & \text{if } \frac{2}{3} \leq |x| \leq \frac{3}{4}, \\ 0, & \text{if } \frac{3}{4} \leq |x| \leq 1. \end{cases} \quad (4.15)$$

It is easy to show for a constant  $C > 1$  independent of  $h$  and a suitably small constant  $h_0 \in (0, 1)$  that

$$|x| |\nabla \mathcal{K}_h(x)| \leq C \mathcal{K}_h(x), \quad \frac{1}{C} |\log h| \leq \|\mathcal{K}_h\|_{L^1} \leq C |\log h|, \quad h \in (0, h_0). \quad (4.16)$$

Define the functional

$$L_{h,p}(f) := \int_{\mathbb{T}^{2d}} \overline{\mathcal{K}}_h(x-y) |\Lambda[f]|^p dx dy, \quad (4.17)$$

where we use the convenient notations

$$\begin{cases} \Lambda[f] := f^x - f^y, \\ f^x := f(x, t), \\ \overline{\mathcal{K}}_h := \frac{\mathcal{K}_h}{\|\mathcal{K}_h\|_{L^1}}. \end{cases} \quad (4.18)$$

By virtue of Lemma A.3 below, to derive the strong convergence of  $\vartheta_\delta := \rho_\delta + n_\delta$ , we need

$$\lim_{h \rightarrow 0} \limsup_{\delta \rightarrow 0} \operatorname{ess\,sup}_{t \in [0, T]} L_{h,1}(\vartheta_\delta) = 0.$$

It is shown in Lemma 4.4 below that the above estimate is the same thing as

$$\lim_{h \rightarrow 0} \limsup_{\delta \rightarrow 0} \operatorname{ess\,sup}_{t \in [0, T]} \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) (w_\delta^x + w_\delta^y) dx dy = 0,$$

where  $w_\delta$  is given by (4.21) below, and  $\chi$  is the function

$$\chi(s) := \begin{cases} |s|^2, & \text{if } |s| \leq 1, \\ \text{smooth}, & \text{if } 1 \leq |s| \leq 2, \\ |s|, & \text{if } |s| \geq 2, \end{cases} \quad (4.19)$$

which satisfies

$$\begin{cases} |\chi(s) - \frac{1}{2}\chi'(s)s| \leq \frac{1}{2}\chi'(s)s, & 0 \leq \chi'(s)s \leq C\chi(s) \leq C|s|, \quad s \in \mathbb{R}, \\ \chi(s) \geq \frac{1}{C}|s|, & |s| \geq 1. \end{cases} \quad (4.20)$$

**Lemma 4.4.** *Let  $T > 0$ ,  $h \in (0, h_0)$ ,  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  be the weak solution to the IVP (3.1)-(3.2) for  $\delta \in (0, 1)$  given by Proposition 3.1,  $\vartheta_\delta := \rho_\delta + n_\delta$ , and  $w_\delta$  be the solution to the auxiliary problem*

$$\begin{cases} \partial_t w_\delta + u_\delta \cdot \nabla w_\delta + \lambda_0 \Xi_\delta w_\delta = 0, & x \in \mathbb{T}^d, \quad t \in (0, T], \\ \Xi_\delta := \vartheta_\delta |\operatorname{div} u_\delta| + |\operatorname{div} u_\delta| + M |\nabla u_\delta| + \rho_\delta^\gamma + \rho_\delta^{\tilde{\gamma}} + n_\delta^\alpha + n_\delta^{\tilde{\alpha}} + 1, \\ w_\delta(x, 0) = e^{-\lambda_0(\rho_{0,\delta} + n_{0,\delta})(x)}, & x \in \mathbb{T}^d, \end{cases} \quad (4.21)$$

where  $\lambda_0 \geq 1$  is a constant to be chosen, and  $M$  is the localized maximal operator defined by (A.9). Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, we have

$$\begin{cases} 0 \leq w_\delta \leq e^{-\lambda_0 \vartheta_\delta} \leq 1, \\ \operatorname{ess\,sup}_{t \in [0, T]} \int_{\mathbb{T}^d} \vartheta_\delta \log(1 + |\log w_\delta|) dx \leq C(1 + \lambda_0), \end{cases} \quad (4.22)$$

and for  $\sigma_* > 0$  to be chosen later,

$$\operatorname{ess\,sup}_{t \in [0, T]} (L_{h,1}(\vartheta_\delta))^2 \leq \frac{C(1 + \lambda_0)}{\log(1 + |\log \sigma_*|)} + \frac{1}{\sigma_*} \operatorname{ess\,sup}_{t \in [0, T]} \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) (w_\delta^x + w_\delta^y) dx dy, \quad (4.23)$$

where  $L_{h,1}(f)$ ,  $\Lambda[f]$ ,  $f^x$ ,  $\bar{\mathcal{K}}_h$ , and  $\chi$  are given by (4.17), (4.18)<sub>1</sub>, (4.18)<sub>2</sub>, (4.18)<sub>3</sub>, and (4.19) respectively, and  $C > 0$  is a constant independent of  $\delta$ ,  $h$ , and  $\sigma_*$ .

**Proof.** First, it follows from the maximal principle for the transport equation (4.21)<sub>1</sub> that

$$0 \leq w_\delta \leq 1, \quad (4.24)$$

which leads to  $|\log w_\delta| = -\log w_\delta$ . Since there holds

$$\partial_t \vartheta_\delta + u_\delta \cdot \nabla \vartheta_\delta + \vartheta_\delta \operatorname{div} u_\delta = 0 \quad \text{in } \mathcal{D}'(\mathbb{T}^d \times (0, T)), \quad (4.25)$$

one derives by (4.21)<sub>2</sub> and the argument of renormalized solutions as in (3.16)-(3.18) that

$$\partial_t e^{-\lambda_0 \vartheta_\delta} + u_\delta \cdot \nabla e^{-\lambda_0 \vartheta_\delta} + \lambda_0 \Xi_\delta e^{-\lambda_0 \vartheta_\delta} = \lambda_0 (\Xi_\delta + \vartheta_\delta \operatorname{div} u_\delta) e^{-\lambda_0 \vartheta_\delta} \geq 0 \quad \text{in } \mathcal{D}'(\mathbb{T}^d \times (0, T)). \quad (4.26)$$

From the comparison principle for (4.21)<sub>1</sub> and (4.26), we have (4.22)<sub>1</sub>.

Next, by (4.22)<sub>1</sub> and the argument of renormalized solutions (formally multiplying (4.21)<sub>1</sub> by  $-\vartheta_\delta(1 + |\log w_\delta|)w_\delta^{-1}$ ), we obtain

$$\begin{aligned} & \partial_t (\vartheta_\delta \log(1 + |\log w_\delta|)) + \operatorname{div} (u_\delta \vartheta_\delta \log(1 + |\log w_\delta|)) \\ &= \frac{\lambda_0 \vartheta_\delta \Xi_\delta}{1 + |\log w_\delta|} \leq \frac{\lambda_0 \vartheta_\delta \Xi_\delta}{1 + \lambda_0 \vartheta_\delta} \leq \Xi_\delta \quad \text{in } \mathcal{D}'(\mathbb{T}^d \times (0, T)). \end{aligned} \quad (4.27)$$

Since we have

$$\int_{\mathbb{T}^d} \vartheta_\delta \log(1 + |\log w_\delta|)|_{t=0} dx \leq C \lambda_0,$$

and the term  $\Xi_\delta$  on the right-hand side of (4.27) is uniformly bounded in  $L^1(0, T; L^1(\mathbb{T}^d))$  due to (1.12)<sub>2</sub>, (1.15)<sub>2</sub>-(1.15)<sub>3</sub>, (3.28), and (4.1)-(4.2), one shows (4.22)<sub>2</sub> after integrating (4.27) over  $\mathbb{T}^d \times [0, t]$ .

Finally, one has by (4.19) and (4.20)<sub>2</sub> that

$$\begin{aligned} L_{h,1}(\vartheta_\delta) &= \left( \int_{\{(x,y) \in \mathbb{T}^{2d} \mid |\Lambda[\vartheta_\delta]| \leq 1\}} + \int_{\{(x,y) \in \mathbb{T}^{2d} \mid |\Lambda[\vartheta_\delta]| \geq 1\}} \right) \bar{\mathcal{K}}_h(x-y) |\Lambda[\vartheta_\delta]| dx dy \\ &\leq C \left( \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) dx dy \right)^{\frac{1}{2}}. \end{aligned} \quad (4.28)$$

Owing to (4.19)<sub>1</sub>, the Young inequality of convolution type, and  $w_\delta^x + w_\delta^y \geq \sigma_*$  for  $w_\delta^x \geq \sigma_*$  or  $w_\delta^y \geq \sigma_*$ , we obtain

$$\begin{aligned} & \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) dx dy \\ &= \left( \int_{\{(x,y) \in \mathbb{T}^{2d} \mid w_\delta^x \leq \sigma_* \text{ and } w_\delta^y \leq \sigma_*\}} + \int_{\{(x,y) \in \mathbb{T}^{2d} \mid w_\delta^x \geq \sigma_* \text{ or } w_\delta^y \geq \sigma_*\}} \right) \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) dx dy \\ &\leq \frac{C}{\log(1 + |\log \sigma_*|)} \int_{\mathbb{T}^d} \vartheta_\delta \log(1 + |\log w_\delta|) dx + \frac{1}{\sigma_*} \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) (w_\delta^x + w_\delta^y) dx dy, \end{aligned}$$

which together with (4.22)<sub>2</sub> and (4.28) leads to (4.23).  $\square$

With the help of the continuity equation (4.25), we have the quantitative regularity estimate with weight.

**Lemma 4.5.** *Let  $T > 0$ ,  $h \in (0, h_0)$ ,  $\lambda_0 \geq 1$ ,  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  be the weak solution to the IVP (3.1)-(3.2) for  $\delta \in (0, 1)$  given by Proposition 3.1,  $\vartheta_\delta := \rho_\delta + n_\delta$ , and  $w_\delta$  be the solution to the IVP (4.21).*

*Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, it holds for  $t \in [0, T]$  that*

$$\begin{aligned} & \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) (w_\delta^x + w_\delta^y) dx dy \\ & \leq L_{h,1}(\rho_{0,\delta} + n_{0,\delta}) + \frac{C}{|\log h|^{\frac{1}{2}}} + (C - 2\lambda_0) \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) \Xi_\delta^x w_\delta^x dx dy d\tau \\ & \quad - 2 \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[\operatorname{div} u_\delta] (\chi'(\Lambda[\vartheta_\delta]) \vartheta_\delta^y + \chi(\Lambda[\vartheta_\delta])) w_\delta^x dx dy d\tau, \end{aligned} \quad (4.29)$$

where  $L_{h,1}(f)$ ,  $\Lambda[f]$ ,  $f^x$ ,  $\bar{\mathcal{K}}_h$ ,  $\chi$ , and  $\Xi_\delta$  are defined by (4.17), (4.18)<sub>2</sub>, (4.18)<sub>1</sub>, (4.18)<sub>3</sub>, (4.19), and (4.21)<sub>2</sub> respectively, and  $C > 0$  is a constant independent of  $\delta$ ,  $h$ , and  $\lambda_0$ .

**Proof.** It can be verified by (4.25) that

$$\begin{aligned} & \partial_t \Lambda[\vartheta_\delta] + \operatorname{div}_x(\Lambda[\vartheta_\delta]u_\delta^x) + \operatorname{div}_y(\Lambda[\vartheta_\delta]u_\delta^y) \\ &= -\vartheta_\delta^y \operatorname{div}_x u_\delta^x + \vartheta_\delta^x \operatorname{div}_y u_\delta^y \\ &= \frac{1}{2} \Lambda[\vartheta_\delta](\operatorname{div}_x u_\delta^x + \operatorname{div}_y u_\delta^y) - \frac{1}{2}(\vartheta_\delta^x + \vartheta_\delta^y) \Lambda[\operatorname{div} u_\delta] \quad \text{in } \mathcal{D}'(\mathbb{T}^d \times (0, T)). \end{aligned} \quad (4.30)$$

By virtue of (4.21), (4.30), and the argument of renormalized solutions as in (3.16)-(3.18) (formally multiplying (4.30) by  $\bar{\mathcal{K}}(x-y)\chi'(\Lambda[\vartheta_\delta])(w_\delta^x + w_\delta^y)$  and integrating the resulted equation over  $\mathbb{T}^{2d} \times [0, t]$ ), one can show

$$\begin{aligned} & \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y)\chi(\Lambda[\vartheta_\delta])(w_\delta^x + w_\delta^y) dx dy \Big|_0^t \\ &= \int_0^t \int_{\mathbb{T}^{2d}} \nabla \bar{\mathcal{K}}_h(x-y) \cdot \Lambda[u_\delta]\chi(\Lambda[\vartheta_\delta])(w_\delta^x + w_\delta^y) dx dy d\tau \\ &+ \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y)\chi(\Lambda[\vartheta_\delta])(\partial_t w_\delta^x + u_\delta^x \cdot \nabla_x w_\delta^x + \partial_t w_\delta^y + u_\delta^y \cdot \nabla_y w_\delta^y) dx dy d\tau \\ &+ \frac{1}{2} \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y)(2\chi(\Lambda[\vartheta_\delta]) - \chi'(\Lambda[\vartheta_\delta])\Lambda[\vartheta_\delta])(\operatorname{div}_x u_\delta^x + \operatorname{div}_y u_\delta^y)(w_\delta^x + w_\delta^y) dx dy d\tau \\ &- \frac{1}{2} \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y)\chi'(\Lambda[\vartheta_\delta])(\vartheta_\delta^x + \vartheta_\delta^y)\Lambda[\operatorname{div} u_\delta](w_\delta^x + w_\delta^y) dx dy d\tau. \end{aligned}$$

Making use of symmetry, we have

$$\int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y)\chi(\Lambda[\vartheta_\delta])(w_\delta^x + w_\delta^y) dx dy \Big|_0^t = \sum_{i=1}^4 I_i^3, \quad (4.31)$$

where  $I_i^3$  ( $i = 1, 2, 3, 4$ ) are given by

$$\begin{aligned} I_1^3 &:= 2 \int_0^t \int_{\mathbb{T}^{2d}} \nabla \bar{\mathcal{K}}_h(x-y) \cdot \Lambda[u_\delta]\chi(\Lambda[\vartheta_\delta])w_\delta^x dx dy d\tau, \\ I_2^3 &:= 2 \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y)\chi(\Lambda[\vartheta_\delta])(\partial_t w_\delta^x + u_\delta^x \cdot \nabla_x w_\delta^x) dx dy d\tau, \\ I_3^3 &:= \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y)(2\chi(\Lambda[\vartheta_\delta]) - \chi'(\Lambda[\vartheta_\delta])\Lambda[\vartheta_\delta])(\operatorname{div}_x u_\delta^x + \operatorname{div}_y u_\delta^y)w_\delta^x dx dy d\tau, \\ I_4^3 &:= - \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y)(\vartheta_\delta^x + \vartheta_\delta^y)\chi'(\Lambda[\vartheta_\delta])\Lambda[\operatorname{div} u_\delta]w_\delta^x dx dy d\tau. \end{aligned}$$



By (4.16), (4.20)<sub>1</sub>, (4.22)<sub>1</sub>, (A.6), and (A.10), it holds

$$\begin{aligned}
I_1^3 &\leq C \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) (D_{|x-y|} u_\delta^x + D_{|x-y|} u_\delta^y) \chi(\Lambda[\vartheta_\delta]) w_\delta^x dx dy d\tau \\
&\leq C \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) (|\Lambda[D_{|x-y|} u_\delta]| + 2D_{|x-y|} u_\delta^x) (\vartheta_\delta^x + \vartheta_\delta^y) w_\delta^x dx dy d\tau \\
&\leq C \int_0^t \|\vartheta_\delta(\tau)\|_{L^2} \int_{\mathbb{T}^d} \bar{\mathcal{K}}_h(z) \|(D_{|z|} u_\delta^{-z} - D_{|z|} u_\delta)(\tau)\|_{L^2} dz d\tau \\
&\quad + C \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) M |\nabla_x u_\delta^x| w_\delta^x dx dy d\tau \\
&\leq \frac{C \|\vartheta_\delta\|_{L^2(0,T;L^2)} \|u_\delta\|_{L^2(0,T;H^1)}}{|\log h|^{\frac{1}{2}}} + C \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) M |\nabla_x u_\delta^x| w_\delta^x dx dy d\tau,
\end{aligned}$$

where  $D_{|x|} f$  and  $Mf$  are defined by (A.7) and (A.9) respectively. For  $I_2^3$ , it follows from (4.21)<sub>1</sub> that

$$I_2^3 = -2\lambda_0 \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) \Xi_\delta^x w_\delta^x dx dy d\tau.$$

In addition, by (4.20)<sub>1</sub> and the fact

$$\begin{aligned}
&(2\chi(\Lambda[\vartheta_\delta]) - \chi'(\Lambda[\vartheta_\delta])\Lambda[\vartheta_\delta]) (\operatorname{div}_x u_\delta^x + \operatorname{div}_y u_\delta^y) + (\vartheta_\delta^x + \vartheta_\delta^y) \chi'(\Lambda[\vartheta_\delta]) \Lambda[\operatorname{div} u_\delta] \\
&= 2(2\chi(\Lambda[\vartheta_\delta]) - \chi'(\Lambda[\vartheta_\delta])\Lambda[\vartheta_\delta]) \operatorname{div}_x u_\delta^x - 2(\chi'(\Lambda[\vartheta_\delta])\vartheta_\delta^y + \chi(\Lambda[\vartheta_\delta])) \Lambda[\operatorname{div} u_\delta],
\end{aligned}$$

we infer

$$\begin{aligned}
I_3^3 + I_4^3 &\leq C \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) |\operatorname{div}_x u_\delta^x| w_\delta^x dx dy d\tau \\
&\quad - 2 \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[\operatorname{div} u_\delta] (\chi'(\Lambda[\vartheta_\delta])\vartheta_\delta^y + \chi(\Lambda[\vartheta_\delta])) w_\delta^x dx dy d\tau.
\end{aligned}$$

One obtain (4.29) after substituting the above estimates of  $I_i^3$  ( $i = 1, 2, 3, 4$ ) into (4.31).  $\square$

Next, the truncated part of  $\Lambda[\operatorname{div} u_\delta]$  can be decomposed the pressure part and the effect viscous part.

**Lemma 4.6.** *Let  $T > 0$ ,  $p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1$ ,  $h \in (0, h_0)$ ,  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  be the weak solution to the IVP (3.1)-(3.2) for  $\delta \in (0, 1)$  given by Proposition 3.1,  $\vartheta_\delta := \rho_\delta + n_\delta$ , and  $w_\delta$  be the solution to the IVP (4.21). Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, it holds for  $k \geq 1$  and  $t \in [0, T]$  that*

$$\begin{aligned}
&- \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[\operatorname{div} u_\delta] (\chi'(\Lambda[\vartheta_\delta])\vartheta_\delta^y + \chi(\Lambda[\vartheta_\delta])) w_\delta^x dx dy d\tau \\
&\leq \frac{C}{k} - \frac{1}{2\mu + \lambda} \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[P_\delta] \Psi_{\delta,k} dx dy d\tau \\
&\quad + \frac{1}{2\mu + \lambda} \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[F_\delta] \Psi_{\delta,k} dx dy d\tau,
\end{aligned} \tag{4.32}$$

with

$$\Psi_{\delta,k} := (\chi'(\Lambda[\vartheta_\delta])\vartheta_\delta^y + \chi(\Lambda[\vartheta_\delta])) \mathbf{1}_{\vartheta_\delta^x \leq k} \mathbf{1}_{\vartheta_\delta^y \leq k} w_\delta^x, \tag{4.33}$$

where  $P_\delta$ ,  $F_\delta$ ,  $L_{h,1}(f)$ ,  $\Lambda[f]$ ,  $f^x$ ,  $\bar{\mathcal{K}}_h$ , and  $\chi$  are given by (2.3), (4.3), (4.17), (4.18)<sub>1</sub>, (4.18)<sub>2</sub>, (4.18)<sub>3</sub>, and (4.19) respectively, and  $C > 0$  is a constant independent of  $\delta$ ,  $h$ ,  $k$ , and  $\lambda_0$ .

**Proof.** By (3.28), (4.1)-(4.2), (4.19)<sub>1</sub>, and (4.22)<sub>1</sub>, we have

$$\begin{aligned}
& - \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[\operatorname{div} u_\delta] (\chi'(\Lambda[\vartheta_\delta]) \vartheta_\delta^y + \chi(\Lambda[\vartheta_\delta])) (1 - \mathbf{1}_{\vartheta_\delta^x \leq k} \mathbf{1}_{\vartheta_\delta^y \leq k}) w_\delta^x dx dy d\tau \\
& \leq C \int_{\{(x,y,\tau) \in \mathbb{T}^{2d} \mid \vartheta_\delta^x \geq k \text{ or } \vartheta_\delta^y \geq k\}} \bar{\mathcal{K}}_h(x-y) (|\operatorname{div}_x u_\delta^x| + |\operatorname{div}_y u_\delta^y|) (\vartheta_\delta^x w_\delta^x + 1) dx dy d\tau \\
& \leq \frac{C}{k} \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) (|\operatorname{div}_x u_\delta^x| + |\operatorname{div}_y u_\delta^y|) (\vartheta_\delta^x + \vartheta_\delta^y) dx dy d\tau \\
& \leq \frac{C}{k} \|\operatorname{div} u_\delta\|_{L^2(0,T;L^2)} \|\vartheta_\delta\|_{L^2(0,T;L^2)} \leq \frac{C}{k},
\end{aligned} \tag{4.34}$$

where one has used the facts  $|\chi'(\Lambda[\vartheta_\delta])| \vartheta_\delta^y \leq C(\vartheta_\delta^x + 1)$  if  $|\lambda[\vartheta_\delta]| \leq 2$ ,  $\chi'(\Lambda[\vartheta_\delta]) \vartheta_\delta^y + \chi(\Lambda[\vartheta_\delta]) = \vartheta_\delta^x \operatorname{sign} \Lambda[\vartheta_\delta]$  for  $|\lambda[\vartheta_\delta]| \geq 2$ ,  $\vartheta_\delta^x + \vartheta_\delta^y \geq k$  for either  $\vartheta_\delta^x \geq k$  or  $\vartheta_\delta^y \geq k$ , and

$$\vartheta_\delta w_\delta \leq \vartheta_\delta e^{-\lambda_0 \vartheta_\delta} \leq \frac{1}{\lambda_0} \leq 1.$$

By (4.3) and (4.34), (4.32) holds.  $\square$

We are ready to estimate the key pressure part.

**Lemma 4.7.** *Let  $T > 0$ ,  $p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1$ ,  $h \in (0, h_0)$ ,  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  be the weak solution to the IVP (3.1)-(3.2) for  $\delta \in (0, 1)$  given by Proposition 3.1,  $\vartheta_\delta := \rho_\delta + n_\delta$ , and  $w_\delta$  be a solution to the IVP (4.21). Then, for any  $\zeta > 0$ , there is a constant  $\delta_1(\zeta) \in (0, 1)$  such that it holds for  $\delta \in (0, \delta_1(\zeta))$ ,  $k \geq 1$ , and  $t \in [0, T]$  that*

$$\begin{aligned}
& - \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[P_\delta] \Psi_{\delta,k} dx dy d\tau \\
& \leq C k^{p_0} \left( \delta^\beta + \zeta + \int_0^t (L_{h,1}(A_{\rho,n}) + L_{h,1}(B_{\rho,n})) d\tau \right) \\
& \quad + C \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) \left( (\rho_\delta^x)^\gamma + (\rho_\delta^x)^{\tilde{\gamma}} + (n_\delta^x)^\alpha + (n_\delta^x)^{\tilde{\alpha}} + 1 \right) w_\delta^x dx dy d\tau,
\end{aligned} \tag{4.35}$$

where  $(A_{\rho,n}, B_{\rho,n})$ ,  $L_{h,1}(f)$ ,  $\Lambda[f]$ ,  $f^x$ ,  $\bar{\mathcal{K}}_h$ ,  $\chi$ , and  $\Psi_{\delta,k}$  are given by (1.18), (4.17), (4.18)<sub>1</sub>, (4.18)<sub>2</sub>, (4.18)<sub>3</sub>, (4.19), and (4.39) respectively, and  $C > 0$  is a constant independent of  $\delta$ ,  $h$ ,  $k$ , and  $\lambda_0$ .

**Proof.** To begin with, as  $\delta \rightarrow 0$ , we deduce by (4.13) that

$$\begin{cases} \rho_\delta - A_{\rho,n} \vartheta_\delta \rightarrow 0 & \text{a.e. in } \mathbb{T}^d \times (0, T), \\ n_\delta - B_{\rho,n} \vartheta_\delta \rightarrow 0 & \text{a.e. in } \mathbb{T}^d \times (0, T), \end{cases}$$

which together with the Egorov theorem implies that for any  $\zeta > 0$ , there exist a constant  $\delta_1(\zeta) \in (0, 1)$  and a domain  $Q_{\zeta,T} \subset \mathbb{T}^d \times (0, T)$  such that

$$\begin{cases} |\mathbb{T}^d \times (0, T) / Q_{\zeta,T}| \leq \zeta, \\ |(\rho_\delta - A_{\rho,n} \vartheta_\delta)(x, t)| + |(n_\delta - B_{\rho,n} \vartheta_\delta)(x, t)| \leq \zeta, \quad (x, t) \in Q_{\zeta,T}, \quad \delta \in (0, \delta_1(\zeta)). \end{cases} \tag{4.36}$$

Denoting

$$\mathbb{Q}_{\zeta,t} = \{(x, y, \tau) \in \mathbb{T}^{2d} \times (0, t) \mid (x, \tau) \in Q_{\zeta,t} \text{ and } (y, \tau) \in Q_{\zeta,t}\}, \quad t \in [0, T],$$

we obtain by (2.3), (4.22)<sub>1</sub>, and (4.36) that

$$\begin{aligned} & - \int_{\mathbb{T}^{2d} \times (0,t)/\mathbb{Q}_{\zeta,t}} \bar{\mathcal{K}}_h(x-y) \Lambda[P_\delta(\rho_\delta, n_\delta)] \Psi_{\delta,k} dx dy d\tau \\ & \leq C(k^\gamma + k^\alpha + \delta k^{p_0} + 1) \int_{\mathbb{T}^{2d} \times (0,t)/\mathbb{Q}_{\zeta,t}} \bar{\mathcal{K}}_h(x-y) dx dy d\tau \leq Ck^{p_0} \zeta. \end{aligned}$$

According to (1.17), the integration on  $\mathbb{Q}_{\zeta,t}$  can be decomposed by

$$- \int_{\mathbb{Q}_{\zeta,t}} \bar{\mathcal{K}}_h(x-y) \Lambda[P_\delta(\rho_\delta, n_\delta)] \Psi_{\delta,k} dx dy d\tau = \sum_{i=1}^6 I_i^4, \quad (4.37)$$

where  $I_i^4$  ( $i = 1, \dots, 6$ ) are given by

$$\begin{aligned} I_1^4 &:= - \int_{\mathbb{Q}_{\zeta,t}} \bar{\mathcal{K}}_h(x-y) (P_\delta(\rho_\delta^x, n_\delta^x) - P_\delta(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x)) \Psi_{\delta,k} dx dy d\tau, \\ I_2^4 &:= - \int_{\mathbb{Q}_{\zeta,t}} \bar{\mathcal{K}}_h(x-y) (P_\delta(A_{\rho^y, n^y} \vartheta_\delta^y, B_{\rho^y, n^y} \vartheta_\delta^y) - P_\delta(\rho_\delta^y, n_\delta^y)) \Psi_{\delta,k} dx dy d\tau, \\ I_3^4 &:= - \int_{\mathbb{Q}_{\zeta,t}} \bar{\mathcal{K}}_h(x-y) (P_\delta(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y) - P_\delta(A_{\rho^y, n^y} \vartheta_\delta^y, B_{\rho^y, n^y} \vartheta_\delta^y)) \Psi_{\delta,k} dx dy d\tau, \\ I_4^4 &:= - \int_{\mathbb{Q}_{\zeta,t}} \bar{\mathcal{K}}_h(x-y) (\mathbf{1}_{\vartheta_\delta^x \leq \delta} P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - \mathbf{1}_{\vartheta_\delta^y \leq \delta} P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)) \Psi_{\delta,k} dx dy d\tau, \\ I_5^4 &:= -\delta \int_{\mathbb{Q}_{\zeta,t}} \bar{\mathcal{K}}_h(x-y) ((A_{\rho^x, n^x} \vartheta_\delta^x)^{p_0} - (A_{\rho^x, n^x} \vartheta_\delta^y)^{p_0} + (B_{\rho^x, n^x} \vartheta_\delta^x)^{p_0} - (B_{\rho^x, n^x} \vartheta_\delta^y)^{p_0}) \Psi_{\delta,k} dx dy d\tau, \\ I_6^4 &:= - \int_{\mathbb{Q}_{\zeta,t}} \bar{\mathcal{K}}_h(x-y) (P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)) \Psi_{\delta,k} dx dy d\tau. \end{aligned}$$

It follows from (1.4), (3.11)-(3.12), (3.21), (4.22)<sub>1</sub>, and (4.36) that

$$\begin{aligned} I_1^4 &\leq C \int_{\mathbb{Q}_{\zeta,t}} \bar{\mathcal{K}}_h(x-y) [((\rho_\delta^x + A_{\rho^x, n^x} \vartheta_\delta^x)^{\tilde{\alpha}-1} + \delta(\rho_\delta^x + A_{\rho^x, n^x} \vartheta_\delta^x)^{p_0-1} + 1) |\rho_\delta^x - A_{\rho^x, n^x} \vartheta_\delta^x| \\ &\quad + ((n_\delta^x + B_{\rho^x, n^x} \vartheta_\delta^x)^{\tilde{\alpha}-1} + \delta(n_\delta^x + B_{\rho^x, n^x} \vartheta_\delta^x)^{p_0-1} + 1) |n_\delta^x - B_{\rho^x, n^x} \vartheta_\delta^x|] (\vartheta_\delta^x + \vartheta_\delta^y) \mathbf{1}_{\vartheta_\delta^x \leq k} \mathbf{1}_{\vartheta_\delta^y \leq k} dx dy d\tau \\ &\leq Ck^{p_0} \zeta. \end{aligned}$$

Similarly, one has

$$I_2^4 \leq Ck^{p_0} \zeta.$$

For the term  $I_3^4$ , we have by (1.4), (3.12), (3.21), and (4.22)<sub>1</sub> that

$$I_3^4 \leq Ck^{p_0} (L_{h,1}(A_{\rho,n}) + L_{h,1}(B_{\rho,n})).$$

Additionally, it is easy to prove

$$I_4^4 + I_5^4 \leq Ck^{p_0} \delta^\beta.$$

The key term  $I_6^4$  can be controlled by the damping term in (4.21)<sub>1</sub> with respect to  $x$ , so we need to overcome the difficulties caused by the possible dependence of  $I_6^5$  in terms of  $y$ . To this end, in the spirit of [10], we analyze  $I_6^4$  in the following three cases:

- Case 1:  $(P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)) \Lambda[\vartheta_\delta] \geq 0$ .

Note that if  $P(\rho, n)$  is monotone increasing in both  $\rho$  and  $n$ , then all the cases reduce to Case 1. In this case, since  $P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)$ ,  $\Lambda[\vartheta_\delta]$ , and  $\chi'(\Lambda[\vartheta_\delta])$  have a same sign, it holds by (4.20)<sub>1</sub> that

$$\begin{aligned} & - (P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)) \Psi_{\delta, k} \\ & \leq | (P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)) | \\ & \quad \times \left( -\frac{1}{2} |\chi'(\Lambda[\vartheta_\delta])| (\vartheta_\delta^x + \vartheta_\delta^y) + |\chi(\Lambda[\vartheta_\delta]) - \frac{1}{2} \chi'(\Lambda[\vartheta_\delta]) \Lambda[\vartheta_\delta]| \right) \leq 0, \quad (x, y, \tau) \in \mathbb{Q}_{\zeta, t}. \end{aligned}$$

- Case 2:  $(P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)) \Lambda[\vartheta_\delta] < 0$  and  $\vartheta_\delta^y \leq 2\vartheta_\delta^x$ .

We make use of (1.4), (3.12), (3.21), and (4.36)<sub>2</sub> to obtain

$$\begin{aligned} & - (P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)) \Psi_{\delta, k} \\ & \leq C((A_{\rho^x, n^x} \vartheta_\delta^x + A_{\rho^x, n^x} \vartheta_\delta^y)^{\tilde{\gamma}-1} + 1) A_{\rho^x, n^x} |\Lambda[\vartheta_\delta]| (\chi'(\Lambda[\vartheta_\delta]) \vartheta_\delta^y + \chi(\Lambda[\vartheta_\delta])) w_\delta^x \\ & \quad + C((B_{\rho^x, n^x} \vartheta_\delta^x + B_{\rho^x, n^x} \vartheta_\delta^y)^{\tilde{\alpha}-1} + 1) B_{\rho^x, n^x} |\Lambda[\vartheta_\delta]| (\chi'(\Lambda[\vartheta_\delta]) \vartheta_\delta^y + \chi(\Lambda[\vartheta_\delta])) w_\delta^x \\ & \leq C((A_{\rho^x, n^x} \vartheta_\delta^x)^{\tilde{\gamma}} + A_{\rho^x, n^x} \vartheta_\delta^x + (B_{\rho^x, n^x} \vartheta_\delta^x)^{\tilde{\alpha}} + B_{\rho^x, n^x} \vartheta_\delta^x) (\chi'(\Lambda[\vartheta_\delta]) \Lambda[\vartheta_\delta] + \chi(\Lambda[\vartheta_\delta])) w_\delta^x \\ & \leq C((\rho_\delta^x + \zeta)^{\tilde{\gamma}} + \rho_\delta^x + \zeta + (n_\delta^x + \zeta)^{\tilde{\alpha}} + n_\delta^x + \zeta) \chi(\Lambda[\vartheta_\delta]) w_\delta^x, \quad (x, y, \tau) \in \mathbb{Q}_{\zeta, t}. \end{aligned}$$

- Case 3:  $(P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)) \Lambda[\vartheta_\delta] < 0$  and  $\vartheta_\delta^y > 2\vartheta_\delta^x$ .

In this case, one has

$$\begin{cases} |\Lambda[\vartheta_\delta]| = \vartheta_\delta^y - \vartheta_\delta^x > \vartheta_\delta^x \geq 0, & \chi'(\Lambda[\vartheta_\delta]) < 0, & \vartheta_\delta^y = |\Lambda[\vartheta_\delta]| + \vartheta_\delta^x < 2|\Lambda[\vartheta_\delta]|, \\ P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) > P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y), \end{cases}$$

which gives rise to

$$\begin{aligned} & - (P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)) \Psi_{\delta, k} \\ & = (P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) - P(A_{\rho^x, n^x} \vartheta_\delta^y, B_{\rho^x, n^x} \vartheta_\delta^y)) (|\chi'(\Lambda[\vartheta_\delta])| \vartheta_\delta^y - \chi(\Lambda[\vartheta_\delta])) w_\delta^x \\ & \leq (P(A_{\rho^x, n^x} \vartheta_\delta^x, B_{\rho^x, n^x} \vartheta_\delta^x) + C_1) (2|\chi'(\Lambda[\vartheta_\delta])| |\Lambda[\vartheta_\delta]| + \chi(\Lambda[\vartheta_\delta])) w_\delta^x \\ & \leq C((\rho_\delta^x + \zeta)^\gamma + (n_\delta^x + \zeta)^\alpha + 1) \chi(\Lambda[\vartheta_\delta]) w_\delta^x, \quad (x, y, \tau) \in \mathbb{Q}_{\zeta, t}. \end{aligned}$$

Combining the above three cases together, we get

$$I_6^4 \leq C \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) ((\rho_\delta^x)^\gamma + (\rho_\delta^x)^{\tilde{\gamma}} + (n_\delta^x)^\alpha + (n_\delta^x)^{\tilde{\alpha}} + 1) \chi(\Lambda[\vartheta_\delta]) w_\delta^x dx dy d\tau.$$

Substituting the above estimates of  $I_i^4$  ( $i = 1, \dots, 6$ ) into (4.37), we derive (4.35). The proof of Lemma 4.7 is completed.  $\square$

**Remark 4.1.** Let  $\Pi(\rho)$  be the general pressure laws for compressible Navier-Stokes equations given in [10]. For  $P(\rho, n) = \Pi(\rho + n)$ , one can repeat same arguments as in [10] to estimate the pressure part (4.35). However, even for  $P(\rho, n) = \Pi(\rho) + \Pi(n)$ , we have to reduce the two variables  $\rho_\delta$  and  $n_\delta$  into the one variable  $\vartheta_\delta$  by applying the decomposition (1.17).

Finally, different from the previous work by Bresch-Jabin [10], we make use of the structures of the equations (3.1) and the commutator estimates (A.11)-(A.12) of the Riesz operator  $\mathcal{R}_i = (-\Delta)^{-1}\partial_i$  to control the effect viscous flux part.

**Lemma 4.8.** Let  $T > 0$ ,  $p_0 > \gamma + \tilde{\gamma} + \alpha + \tilde{\alpha} + 1$ ,  $h \in (0, h_0)$ ,  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  be the weak solution to the IVP (3.1)-(3.2) for  $\delta \in (0, 1)$  given by Proposition 3.1,  $\vartheta_\delta := \rho_\delta + n_\delta$ , and  $w_\delta$  be the solution to the IVP (4.21). Then, under the assumptions of either Theorem 1.1 or Theorem 1.2, it holds for  $k \geq 1$  and  $t \in [0, T]$  that

$$\int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[F_\delta] \Psi_{\delta,k} dx dy d\tau \leq \frac{C \lambda_0 k^{p_0}}{|\log h|^{\frac{1}{2} \min\{\frac{1}{2}, \frac{2\gamma_0-d}{\gamma_0+d}\}}}, \quad \gamma_0 := \min\{\gamma, \alpha\}, \quad (4.38)$$

where  $F_\delta$ ,  $\Lambda[f]$ ,  $f^x$ ,  $\bar{\mathcal{K}}_h$ , and  $\Psi_{\delta,k}$  are given by (4.3), (4.18)<sub>1</sub>, (4.18)<sub>2</sub>, (4.18)<sub>3</sub>, and (4.39) respectively, and  $C > 0$  is a constant independent of  $\delta$ ,  $h$ ,  $k$ , and  $\lambda_0$ .

**Proof.** By the argument of renormalized solutions for (4.21)<sub>1</sub> and (4.25), one has

$$\partial_t \Psi_{\delta,k} + \operatorname{div}_x(\Psi_{\delta,k} u_\delta^x) + \operatorname{div}_y(\Psi_{\delta,k} u_\delta^y) = R_{\delta,k} \quad \text{in } \mathcal{D}'(\mathbb{T}^{2d} \times (0, T)), \quad (4.39)$$

with

$$R_{\delta,k} := (\Psi_{\delta,k} - \partial_{\vartheta_\delta^x} \Psi_{\delta,k}) \operatorname{div}_x u_\delta^x + (\Psi_{\delta,k} - \partial_{\vartheta_\delta^y} \Psi_{\delta,k}) \operatorname{div}_y u_\delta^y - \lambda_0 \partial_{w_\delta^x} \Psi_{\delta,k} \Xi_\delta^x w_\delta^x.$$

By (4.21)<sub>2</sub> and (4.22)<sub>1</sub>, it is easy to verify

$$\begin{cases} \|\Psi_{\delta,k}\|_{L^\infty(0,T;L^\infty(\mathbb{T}^{2d}))} \leq Ck, \\ \|R_{\delta,k}\|_{L^\infty(0,T;L^\infty(\mathbb{T}^{2d}))} \leq C\lambda_0 k^{p_0} (|\operatorname{div}_x u_\delta^x| + |\operatorname{div}_y u_\delta^y| + M|\nabla_x u_\delta^x| + 1), \end{cases} \quad (4.40)$$

where the constant  $C > 0$  is independent of  $\delta, k$ , and  $\lambda_0$ , and the localized maximal operator  $M : L^p(\mathbb{T}^d) \rightarrow L^p(\mathbb{T}^d)$  for any  $p \in [1, \infty)$  is defined by (A.7). From (4.4), we have

$$\Lambda[F_\delta] = \partial_t \Lambda[(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)] + \Lambda[(-\Delta)^{-1} \operatorname{div} \operatorname{div}(\vartheta_\delta u_\delta \otimes u_\delta)]. \quad (4.41)$$

One deduces by (4.39), (4.41), and integration by parts that

$$\int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[F_\delta] \Psi_{\delta,k} dx dy d\tau = \sum_{i=1}^4 I_i^5, \quad (4.42)$$

where  $I_i^5$  ( $i = 1, 2, 3, 4$ ) are given by

$$\begin{aligned} I_1^5 &:= \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}(x-y) \Lambda[(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)] \Psi_{\delta,k} dx dy \Big|_0^t, \\ I_2^5 &:= - \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)] R_{\delta,k} dx dy d\tau, \\ I_3^5 &:= \int_0^t \int_{\mathbb{T}^{2d}} \nabla \bar{\mathcal{K}}(x-y) \cdot \Lambda[u_\delta] \Lambda[(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)] \Psi_{\delta,k} dx dy d\tau, \\ I_4^5 &:= \sum_{i,j=1}^d \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[\mathcal{R}_i \mathcal{R}_j ((\vartheta_\delta u_\delta^i) u_\delta^j) - u_\delta^j \mathcal{R}_i \mathcal{R}_j (\vartheta_\delta u_\delta^i)] \Psi_{\delta,k} dx dy d\tau. \end{aligned}$$

First, we estimate the term  $I_1^5$ . It follows from (4.8), (A.3), and the boundedness of the operator  $\nabla(-\Delta)^{-1} \operatorname{div}$  that

$$\begin{aligned} &L_{h,p}((-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)) \\ &\leq \frac{C \|(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)\|_{L^p}}{|\log h|^{\frac{1}{2}}} + \frac{C \|\nabla(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)\|_{L^{\frac{2\gamma_0}{\gamma_0+1}}}}{|\log h|^{pd \min\{0, \frac{1}{p} - \frac{\gamma_0+1}{2\gamma_0}\} + p}} \\ &\leq C \left( \frac{1}{|\log h|^{\frac{1}{2}}} + \frac{1}{|\log h|^{pd \min\{0, \frac{1}{p} - \frac{\gamma_0+1}{2\gamma_0}\} + p}} \right), \quad p \in [1, \frac{2\gamma_0 d(\gamma_0 + 1)}{d(\gamma_0 + 1) - 2\gamma_0}], \end{aligned} \quad (4.43)$$

which together with (2.6), (4.8), and (4.40)<sub>2</sub> implies

$$\begin{aligned} I_1^5 &\leq Ck \operatorname{ess\,sup}_{t \in [0, T]} L_{h,1}((-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)) + Ck L_{h,1}((-\Delta)^{-1} \operatorname{div}((\rho_{0,\delta} + n_{0,\delta}) u_{0,\delta})) \\ &\leq \frac{Ck}{|\log h|^{\frac{1}{2}}}. \end{aligned}$$

To control the term  $I_2^5$ , we obtain by (3.28), (4.8), (4.40)<sub>2</sub>, and (4.43) that

$$\begin{aligned} I_2^5 &\leq Ck^{p_0} \lambda_0 \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) |Er[(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)]| (|\operatorname{div}_x u_\delta^x| + |\operatorname{div}_y u_\delta^y| + M|\nabla_x u_\delta^x| + 1) dx dy d\tau \\ &\leq Ck^{p_0} \lambda_0 \left( \|\nabla u_\delta\|_{L^2(0, T; L^2)} \operatorname{ess\,sup}_{t \in [0, T]} L_{h,2}((-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta))^{\frac{1}{2}} + \operatorname{ess\,sup}_{t \in [0, T]} L_{h,1}((-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)) \right) \\ &\leq \frac{Ck^{p_0} \lambda_0}{|\log h|^{\min\{\frac{1}{4}, \frac{2\gamma_0-d}{2\gamma_0}\}}}. \end{aligned}$$

Similarly to the estimate of  $I_1^3$  in (4.31), since  $(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)$  is uniformly bounded in  $L^2(0, T; L^2(\mathbb{T}^d))$  due to  $\gamma_0 > \frac{d}{2}$ , one can prove

$$I_3^5 \leq Ck \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) |\Lambda[(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)]| M |\nabla_x u_\delta^x| dx dy d\tau + \frac{Ck}{|\log h|^{\frac{1}{2}}}. \quad (4.44)$$

By (3.28), (4.8), and (4.43), we have

$$\begin{aligned} &\int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) |\Lambda[(-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta)]| M |\nabla_x u_\delta^x| dx dy d\tau \\ &\leq C \|M |\nabla u_\delta|\|_{L^2(0, T; L^2)} \operatorname{ess\,sup}_{t \in [0, T]} L_{h,2}((-\Delta)^{-1} \operatorname{div}(\vartheta_\delta u_\delta))^{\frac{1}{2}} \\ &\leq \frac{C}{|\log h|^{\min\{\frac{1}{4}, \frac{2\gamma_0-d}{2\gamma_0}\}}}. \end{aligned} \quad (4.45)$$

We obtain from (4.44)-(4.45) that

$$I_3^5 \leq \frac{Ck}{|\log h|^{\min\{\frac{1}{4}, \frac{2\gamma_0-d}{2\gamma_0}\}}}.$$

Finally, we are ready to estimate the last term  $I_4^5$ . Inspired by [37], we make use of the commutator estimate (A.12) to have

$$\|\nabla(\mathcal{R}_i\mathcal{R}_j(\vartheta_\delta u_\delta^i u_\delta^j) - u_\delta^j \mathcal{R}_i\mathcal{R}_j(\vartheta_\delta u_\delta^i))(t)\|_{L^p} \leq C\|\nabla u_\delta^j(t)\|_{L^2}\|\vartheta_\delta u_\delta^i(t)\|_{L^q}, \quad \text{a.e. } t \in (0, T), \quad (4.46)$$

where the constants  $p \in (1, \infty)$  and  $q \in (2, \infty)$  satisfy  $\frac{1}{2} + \frac{1}{q} = \frac{1}{p} < 1$ . Note that one may not apply (4.46) directly for  $\gamma_0 \leq d$ . For example,  $\vartheta_\delta u_\delta$  is uniformly bounded in  $L^2(0, T; L^{\frac{6\gamma_0}{6+\gamma_0}}(\mathbb{T}^d))$  for  $d = 3$ , but we have  $\frac{1}{2} + \frac{6+\gamma_0}{6\gamma_0} \geq 1$  provided  $\gamma_0 \leq 3$ . To overcome this difficulty, we truncate by using  $1 = \mathbf{1}_{\vartheta_\delta \leq L} + \mathbf{1}_{\vartheta_\delta \geq L}$  for  $L \geq 1$  leading to  $I_4^5 = I_1^6 + I_2^6$  with

$$I_1^6 := \sum_{i,j=1}^d \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[\mathcal{R}_i\mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \leq L} u_\delta^i u_\delta^j) - u_\delta^j \mathcal{R}_i\mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \leq L} u_\delta^i)] \Psi_{\delta,k} dx dy d\tau,$$

$$I_2^6 := \sum_{i,j=1}^d \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \Lambda[\mathcal{R}_i\mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \geq L} u_\delta^i u_\delta^j) - u_\delta^j \mathcal{R}_i\mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \geq L} u_\delta^i)] \Psi_{\delta,k} dx dy d\tau.$$

The terms  $I_1^6$  and  $I_2^6$  can be estimated in the case of  $d \geq 3$  and the case of  $d = 2$  as follows:

- Case 1:  $d \geq 3$ .

We show after using (4.40)<sub>1</sub>, (A.3), (A.12), and  $H^1(\mathbb{T}^d) \hookrightarrow L^{\frac{2d}{d-2}}(\mathbb{T}^d)$  that

$$\begin{aligned} I_1^6 &\leq Ck \sum_{i,j=1}^d \int_0^t L_{h,1}(\mathcal{R}_i\mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \leq L} u_\delta^i u_\delta^j) - u_\delta^j \mathcal{R}_i\mathcal{R}_j(\mathbf{1}_{\vartheta_\delta \leq L} \vartheta_\delta u_\delta^i)) d\tau \\ &\leq Ck \left( \frac{1}{|\log h|^{\frac{1}{2}}} + \frac{1}{|\log h|^2} \sum_{i,j=1}^d \|\nabla[\mathcal{R}_i\mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \leq L} u_\delta^i u_\delta^j) - u_\delta^j \mathcal{R}_i\mathcal{R}_j(\mathbf{1}_{\vartheta_\delta \leq L} \vartheta_\delta u_\delta^i)]\|_{L^1(0,T;L^{\frac{d}{d-1}})} \right) \\ &\leq \frac{Ck}{|\log h|^{\frac{1}{2}}} \left( 1 + \|\nabla u_\delta\|_{L^2(0,T;L^2)} \|\vartheta_\delta \mathbf{1}_{\vartheta_\delta \leq L} u_\delta\|_{L^2(0,T;L^{\frac{2d}{d-2}})} \right) \\ &\leq \frac{CkL}{|\log h|^{\frac{1}{2}}}. \end{aligned}$$

For the term  $I_2^6$ , it holds

$$\begin{aligned} I_2^6 &\leq Ck \sum_{i,j=1}^d \left( \|\mathcal{R}_i\mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \geq L} u_\delta^i u_\delta^j)\|_{L^1(0,T;L^{p_2})} + \|u_\delta^j \mathcal{R}_i\mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \geq L} u_\delta^i)\|_{L^1(0,T;L^{p_2})} \right) \\ &\leq Ck \|\vartheta_\delta \mathbf{1}_{\vartheta_\delta \geq L}\|_{L^\infty(0,T;L^{p_3})} \|u_\delta\|_{L^2(0,T;L^{\frac{2d}{d-2}}}^2 \\ &\leq \frac{Ck \|\vartheta_\delta\|_{L^\infty(0,T;L^{\gamma_0})}^{\frac{\gamma_0}{p_3}} \|u_\delta\|_{L^2(0,T;H^1)}^2}{L^{\frac{1}{d}(\gamma_0 - \frac{d}{2})}} \leq \frac{Ck}{L^{\frac{1}{d}(\gamma_0 - \frac{d}{2})}}, \end{aligned}$$

with the constants  $p_2 \in (1, \infty)$  and  $p_3 \in (\frac{d}{2}, \gamma_0)$  satisfying

$$\frac{1}{p_2} + \frac{d-2}{d} = \frac{1}{p_2}, \quad \frac{\gamma_0 - p_3}{p_3} = \frac{1}{d}(\gamma_0 - \frac{d}{2}).$$

- Case 2:  $d = 2$ .

By (4.40)<sub>1</sub>, (A.3), (A.12), and  $H^1(\mathbb{T}^2) \hookrightarrow L^p(\mathbb{T}^2)$  for any  $p \in [1, \infty)$ , the term  $I_1^6$  can be estimated by

$$\begin{aligned} I_1^6 &\leq Ck \sum_{i,j=1}^2 \int_0^t L_{h,1}(\mathcal{R}_i \mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \leq L} u_\delta^i u_\delta^j) - u_\delta^j \mathcal{R}_i \mathcal{R}_j(\mathbf{1}_{\vartheta_\delta \leq L} \vartheta_\delta u_\delta^i)) d\tau \\ &\leq Ck \left( \frac{1}{|\log h|^{\frac{1}{2}}} + \frac{1}{|\log h|^{\frac{3}{2}}} \sum_{i,j=1}^2 \|\nabla[\mathcal{R}_i \mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \leq L} u_\delta^i u_\delta^j) - u_\delta^j \mathcal{R}_i \mathcal{R}_j(\mathbf{1}_{\vartheta_\delta \leq L} \vartheta_\delta u_\delta^i)]\|_{L^1(0,T;L^{\frac{4}{3}})} \right) \\ &\leq \frac{Ck}{|\log h|^{\frac{1}{2}}} (1 + \|\nabla u_\delta\|_{L^2(0,T;L^2)} \|\vartheta_\delta \mathbf{1}_{\vartheta_\delta \leq L} u_\delta\|_{L^2(0,T;L^4)}) \\ &\leq \frac{Ck(1 + L\|u_\delta\|_{L^2(0,T;H^1)})}{|\log h|^{\frac{1}{2}}} \leq \frac{CkL}{|\log h|^{\frac{1}{2}}}. \end{aligned}$$

For the term  $I_2^6$ , we make use of (4.40)<sub>1</sub>, (A.3), (A.11), and  $H^1(\mathbb{T}^2) \hookrightarrow \mathcal{BMO}(\mathbb{T}^2) \hookrightarrow L^p(\mathbb{T}^2)$  for any  $p \in [1, \infty)$  to obtain

$$\begin{aligned} I_2^6 &\leq Ck \sum_{i,j=1}^2 \|\mathcal{R}_i \mathcal{R}_j(\vartheta_\delta \mathbf{1}_{\vartheta_\delta \leq L} u_\delta^i u_\delta^j) - u_\delta^j \mathcal{R}_i \mathcal{R}_j(\mathbf{1}_{\vartheta_\delta \leq L} \vartheta_\delta u_\delta^i)\|_{L^1(0,T;L^{\frac{4\gamma_0}{3\gamma_0+1})}} \\ &\leq Ck \|u_\delta\|_{L^2(0,T;\mathcal{BMO})} \|\vartheta_\delta \mathbf{1}_{\vartheta_\delta \geq L} u_\delta\|_{L^2(0,T;L^{\frac{4\gamma_0}{3\gamma_0+1})}} \\ &\leq Ck \|u_\delta\|_{L^2(0,T;H^1)} \|\vartheta_\delta \mathbf{1}_{\vartheta_\delta \geq L}\|_{L^\infty(0,T;L^{\frac{2\gamma_0}{\gamma_0+1})}} \|u_\delta\|_{L^2(0,T;L^{\frac{4\gamma_0}{\gamma_0-1})}} \\ &\leq \frac{Ck}{L^{\frac{\gamma_0-1}{2}}} \|u_\delta\|_{L^2(0,T;H^1)}^2 \|\vartheta_\delta\|_{L^\infty(0,T;L^{\gamma_0})}^{\frac{\gamma_0+1}{2}} \leq \frac{Ck}{L^{\frac{\gamma_0-1}{2}}} \end{aligned}$$

Combining the above two cases about the estimates of  $I_i^6$  ( $i = 1, 2$ ) together, we derive

$$I_4^5 \leq Ck \left( \frac{L}{|\log h|^{\frac{1}{2}}} + \frac{1}{L^{\frac{1}{d}(\gamma_0 - \frac{d}{2})}} \right), \quad d \geq 2. \quad (4.47)$$

Thence one can choose  $L = |\log h|^{\frac{1}{2(\frac{\gamma_0}{d} + \frac{1}{2})}}$  in (4.47) to have

$$I_4^5 \leq \frac{Ck}{|\log h|^{\frac{1}{2} \min\{1, \frac{2\gamma_0-d}{2\gamma_0+d}\}}}.$$

Substituting the above estimates of  $I_i^5$  ( $i = 1, 2, 3, 4$ ) into (4.42), we prove (4.38). The proof of Lemma 4.8 is completed.  $\square$

**Remark 4.2.** *It should be noted that Lemma 4.8 about the compactness of effect viscous flux indeed holds for any adiabatic constants  $\gamma, \alpha > \frac{d}{2}$ .*

**Proof of Theorem 1.1 and Theorem 1.2:** Let  $T > 0$ ,  $(\rho_\delta, n_\delta, (\rho_\delta + n_\delta)u_\delta)$  be the weak solution to the IVP (3.1)-(3.2) for  $\delta \in (0, 1)$  given by Proposition 3.1,  $\vartheta_\delta := \rho_\delta + n_\delta$ ,  $w_\delta$  be the solution to the



IVP (4.21), and  $(\rho, u, (\rho + n)u)$  be the limit obtained by Lemma 4.2. Then, with the help of Lemmas 4.5-4.8, for any  $\zeta > 0$ , there exists a constant  $\delta_1(\zeta) \in (0, 1)$  such that for  $\delta \in (0, \delta_1(\zeta))$ ,  $h \in (0, h_0)$ ,  $k \geq 1$ , and  $\lambda_0 \geq 1$ , we have

$$\begin{aligned} & \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) (w_\delta^x + w_\delta^y) dx dy \\ & \leq \frac{C}{k} + C\lambda_0 k^{p_0} h_1(\delta, \zeta, h) + (C - 2\lambda_0) \int_0^t \int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) |\Lambda[\vartheta_\delta]| \Xi_\delta^x w_\delta^x dx dy d\tau, \end{aligned}$$

where  $C > 0$  is a constant independent of  $\delta$ ,  $h$ ,  $\zeta$ , and  $\lambda_0$ , and  $h_1(\delta, \zeta, h) \in (0, 1)$  is given by

$$\begin{aligned} h_1(\delta, \zeta, h) & := L_{h,1}(\rho_{0,\delta} + n_{0,\delta}) + \frac{1}{|\log h|^{\frac{1}{2} \min\{\frac{1}{2}, \frac{2\gamma_0-d}{\gamma_0+d}\}}} \\ & + \delta^\beta + \zeta + \int_0^T (L_{h,1}(A_{\rho,n}) + L_{h,1}(B_{\rho,n})) dt, \quad \gamma_0 := \min\{\gamma, \alpha\}. \end{aligned}$$

Choosing

$$\lambda_0 = \frac{C}{2} + 1, \quad k = h_1(\delta, \zeta, h)^{-\frac{1}{p_0+1}},$$

we have

$$\int_{\mathbb{T}^{2d}} \bar{\mathcal{K}}_h(x-y) \chi(\Lambda[\vartheta_\delta]) (w_\delta^x + w_\delta^y) dx dy \leq C h_1(\delta, \zeta, h)^{\frac{1}{p_0+1}},$$

which together with (4.23) implies for any  $\sigma_* > 0$  that

$$(L_{h,1}(\vartheta_\delta))^2 \leq \frac{C}{\log(1 + |\log \sigma_*|)} + \frac{C h_1(\delta, \zeta, h)^{\frac{1}{p_0+1}}}{\sigma_*}.$$

Thus, one can choose  $\sigma_* = h_1(\delta, \zeta, h)^{\frac{1}{2(p_0+1)}}$  to obtain

$$(L_{h,1}(\vartheta_\delta))^2 \leq \frac{C}{\log(1 + |\log h_1(\delta, \zeta, h)|)} + h_1(\delta, \zeta, h)^{\frac{1}{2(p_0+1)}}.$$

Since it follows  $\lim_{h \rightarrow 0} \limsup_{\delta \rightarrow 0} h_1(\delta, \zeta, h) = \zeta$  for any small  $\zeta > 0$ , we gain

$$\lim_{h \rightarrow 0} \limsup_{\delta \rightarrow 0} \operatorname{ess\,sup}_{t \in [0, T]} L_{h,1}(\vartheta_\delta) = 0. \quad (4.48)$$

Due to (4.14), (4.48), Lemma A.3 below, and the fact that  $\partial_t \vartheta_\delta$  is uniformly bounded in  $L^\infty(0, T; W^{-1, \frac{2\gamma_0}{\gamma_0+1}}(\mathbb{T}^d))$ , it holds up to a subsequence (still denoted by  $(\rho_\delta, n_\delta)$ ) that

$$(\rho_\delta, n_\delta) \rightarrow (\rho, n) \quad \text{in } L^1(0, T; L^1(\mathbb{T}^d)) \times L^1(0, T; L^1(\mathbb{T}^d)) \quad \text{as } \delta \rightarrow 0, \quad (4.49)$$

which together with Lemma 4.1 and the Egorov theorem gives rise to

$$P_\delta(\rho_\delta, n_\delta) \rightarrow P(\rho, n) \quad \text{in } L^1(0, T; L^1(\mathbb{T}^d)) \quad \text{as } \delta \rightarrow 0. \quad (4.50)$$

By (4.12) and (4.50), one can show that  $(\rho, n, (\rho + n)u)$  satisfies the properties (1)-(3) in Definition 1.1.

Finally, we show the energy inequality (1.9) under the assumptions of Theorem 1.1, and the case of Theorem 1.2 can be proved similarly. For any  $L \geq 1$  and the smooth function  $j_\delta$  satisfying (2.5), it is easy to verify for  $(\rho_{0,\delta}^{1,L}, n_{0,\delta}^{1,L}) := ((\rho_0 \mathbf{1}_{\rho_0 \leq L}) * j_\delta + \delta, (n_0 \mathbf{1}_{\rho_0 \leq L}) * j_\delta + \delta)$  that

$$\left| \int_{\mathbb{T}^d} G(\rho_{0,\delta}^{1,L}, n_{0,\delta}^{1,L}) dx \right| \rightarrow 0 \quad \text{as } \delta \rightarrow 0, \quad (4.51)$$

and for  $(\rho_{0,\delta}^{2,L}, n_{0,\delta}^{2,L}) := ((\rho_0 \mathbf{1}_{\rho_0 \geq L}) * j_\delta + \delta, (n_0 \mathbf{1}_{n_0 \geq L}) * j_\delta + \delta)$  that

$$\begin{aligned} & \left| \int_{\mathbb{T}^d} G(\rho_{0,\delta}^{2,L}, n_{0,\delta}^{2,L}) dx \right| \\ & \leq \int_{\mathbb{T}^d} \rho_{0,\delta}^{2,L} \int_1^{\rho_{0,\delta}^{2,L}} (P_0 s^{\gamma-2} + P_0 \frac{(n_{0,\delta}^{2,L})^\Gamma}{(\rho_{0,\delta}^{2,L})^\Gamma} s^{\Gamma-2} + s^{-2}) ds dx \\ & \leq \int_{\mathbb{T}^d} \rho_{0,\delta}^{2,L} \left( \int_1^{\rho_{0,\delta}^{2,L}} (P_0 s^{\gamma-2} + P_0 \frac{(n_{0,\delta}^{2,L})^\Gamma}{(\rho_{0,\delta}^{2,L})^\Gamma} s^{\Gamma-2}) ds + 1 \right) dx \\ & \leq C \int_{\mathbb{T}^d} \left( (\rho_0 \mathbf{1}_{\rho_0 \geq L})^\gamma + (n_0 \mathbf{1}_{n_0 \geq L})^\Gamma + \rho_0 \mathbf{1}_{\rho_0 \geq L} \right) dx \rightarrow 0 \quad \text{as } L \rightarrow \infty \text{ uniformly in } \delta, \end{aligned} \quad (4.52)$$

where one has used (1.4), (1.10)-(1.11), and the Young inequality for convolutions. By (2.6), we also infer

$$\left| \int_{\mathbb{T}^d} \rho_{0,\delta} \int_1^{\rho_{0,\delta}} \left( \mathbf{1}_{s + \frac{n_{0,\delta}}{\rho_{0,\delta}} s \leq \delta} P(s, \frac{n_{0,\delta}}{\rho_{0,\delta}} s) + \delta (s + \frac{n_{0,\delta}}{\rho_{0,\delta}} s)^{p_0} \right) s^{-2} \right| \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \quad (4.53)$$

By (4.51)-(4.53), it follows

$$\int_{\mathbb{T}^d} G(\rho_0, n_0) dx = \lim_{\delta \rightarrow 0} \int_{\mathbb{T}^d} G_\delta(\rho_{0,\delta}, n_{0,\delta}) dx. \quad (4.54)$$

By (4.1)-(4.2), (4.49), a similar argument of truncation, one gets

$$\int_0^T \psi \int_{\mathbb{T}^d} G(\rho, n) dx dt = \lim_{\delta \rightarrow 0} \int_0^T \psi \int_{\mathbb{T}^d} G_\delta(\rho_\delta, n_\delta) dx dt, \quad \forall \psi \in C_c^\infty(0, T), \quad (4.55)$$

By (2.21), (3.29), (4.12), (4.54)-(4.55), and the lower semi-continuity of weak limits, the solution  $(\rho, u, (\rho + n)u)$  to the IVP (1.1)-(1.2) satisfies the energy inequality (1.9) for a.e.  $t \in (0, T)$ . Theorem 1.1 and Theorem 1.2 are proved.

## A

In this section, we present some technical lemmas which are useful for our analysis.

**Lemma A.1.** *Let  $p \in [1, \infty)$ ,  $\varepsilon > 0$ , and  $f_\varepsilon$  be uniformly bounded in  $L^p(\mathbb{T}^d)$ . Then, for any  $h \in (0, h_0)$ , it holds*

$$\sup_{|z| < h} \int_{\mathbb{T}^d} |f_\varepsilon(x+z) - f_\varepsilon(x)|^p dx \leq C \left( \frac{\|f_\varepsilon\|_{L^1}^p}{|\log h|^p} + L_{h^{\frac{1}{d+1}}, p}(f_\varepsilon) \right), \quad (A.1)$$

$$L_{h,p}(f_\varepsilon) \leq C \left( \frac{\|f_\varepsilon\|_{L^p}^p}{|\log h|^{\frac{1}{2}}} + \sup_{|z| < \frac{1}{|\log h|}} \int_{\mathbb{T}^d} |f_\varepsilon(x+z) - f_\varepsilon(x)|^p dx \right), \quad (A.2)$$

where  $L_{h,p}(f)$  is defined by (4.17), and  $C > 0$  is a constant independent of  $h$  and  $\varepsilon$ .

Furthermore, if  $f_\varepsilon$  is uniformly bounded in  $W^{1,q}(\mathbb{T}^d)$  for  $q \in [1, d)$ , then we have

$$L_{h,p}(f_\varepsilon) \leq C \left( \frac{\|f_\varepsilon\|_{L^p}^p}{|\log h|^{\frac{1}{2}}} + \frac{\|\nabla f_\varepsilon\|_{L^q}^p}{|\log h|^{pd \min\{0, \frac{1}{p} - \frac{1}{q}\} + p}} \right), \quad p \in [1, \frac{qd}{d-q}). \quad (\text{A.3})$$

**Proof.** First, for any  $|z| < h$  and  $\sigma \in (0, 1)$ , it can be shown by (4.16) and the fact  $\|\nabla \mathcal{K}_\sigma\|_{L^\infty} \leq \frac{C}{\sigma^{d+1}}$  that

$$\begin{aligned} & \int_{\mathbb{T}^d} |f_\varepsilon(x+z) - f_\varepsilon(x)|^p dx \\ & \leq C \int_{\mathbb{T}^d} \left| \frac{\mathcal{K}_\sigma}{\|\mathcal{K}_h\|_{L^1}} * f_\varepsilon(x+z) - \frac{\mathcal{K}_\sigma}{\|\mathcal{K}_h\|_{L^1}} * f_\varepsilon(x) \right|^p dx + C \int_{\mathbb{T}^d} \left| f_\varepsilon(x) - \frac{\mathcal{K}_\sigma}{\|\mathcal{K}_h\|_{L^1}} * f_\varepsilon(x) \right|^p dx \\ & \leq \frac{C}{\|\mathcal{K}_\sigma\|_{L^1}^p} \int_{\mathbb{T}^d} \left| \int_{\mathbb{T}^d} (\mathcal{K}_\sigma(x+z-y) - \mathcal{K}_\sigma(x-y)) f_\varepsilon(y) dy \right|^p dx \\ & \quad + \frac{C}{\|\mathcal{K}_\sigma\|_{L^1}^p} \int_{\mathbb{T}^d} \left| \int_{\mathbb{T}^d} \mathcal{K}_\sigma(x-y) (f_\varepsilon(x) - f_\varepsilon(y)) dy \right|^p dx \\ & \leq \frac{C|z|^p}{|\log \sigma|^p} \|\nabla \mathcal{K}_\sigma\|_{L^\infty}^p \|f_\varepsilon\|_{L^1}^p + \frac{C}{|\log \sigma|} \int_{\mathbb{T}^{2d}} \mathcal{K}_\sigma(x-y) |f_\varepsilon(x) - f_\varepsilon(y)|^p dx dy \\ & \leq \frac{Ch^p \|f_\varepsilon\|_{L^1}^p}{\sigma^{p(d+1)} |\log \sigma|^p} + L_{\sigma,p}(f_\varepsilon). \end{aligned}$$

One may choose  $\sigma = h^{\frac{1}{d+1}}$  in the above inequality to prove (A.2).

Next, we show (A.1). It follows

$$L_{h,p}(f_\varepsilon) = \sum_{i=1}^3 I_i^7,$$

where  $I_i^7$  ( $i = 1, 2, 3$ ) are given by

$$\begin{aligned} I_1^7 & := \int_{\{|x-y| \geq \frac{1}{2}\}} \bar{\mathcal{K}}_h(x-y) |f_\varepsilon(x) - f_\varepsilon(y)|^p dx dy, \\ I_2^7 & := \int_{\{\frac{1}{|\log h|} \leq |x-y| < \frac{1}{2}\}} \bar{\mathcal{K}}_h(x-y) |f_\varepsilon(x) - f_\varepsilon(y)|^p dx dy, \\ I_3^7 & := \int_{\{0 \leq |x-y| < \frac{1}{|\log h|}\}} \bar{\mathcal{K}}_h(x-y) |f_\varepsilon(x) - f_\varepsilon(y)|^p dx dy. \end{aligned}$$

Due to (4.16) and  $\mathcal{K}_h(x) \leq C$  for any  $|x| \geq \frac{1}{2}$ , one can bound  $I_1^8$  by

$$I_1^7 \leq \frac{C \|f_\varepsilon\|_{L^p}^p}{|\log h|}.$$

For the term  $I_2^7$ , we have

$$\begin{aligned} I_2^7 & \leq C \sum_{n=0}^{\log |\log h| - 1} \frac{1}{|\log h|} \int_{\{\frac{1}{2} e^{-n-1} \leq |z| \leq \frac{1}{2} e^{-n}\}} e^{nd} \int_{\mathbb{T}^d} |f_\varepsilon(x) - f_\varepsilon(x-z)|^p dx dz \\ & \leq \frac{C \|f_\varepsilon\|_{L^p}^p}{|\log h|^{\frac{1}{2}}} \sum_{n=0}^{\infty} e^{-\frac{n}{2}} \leq \frac{C \|f_\varepsilon\|_{L^p}^p}{|\log h|^{\frac{1}{2}}}, \end{aligned}$$

where we have used the facts  $\mathcal{K}_h(z) \leq \frac{C}{|z|^d}$  for any  $|z| > 0$  and  $|\log h|^{-\frac{1}{2}} \leq e^{-\frac{1}{2}(n+1)}$  for any integer  $0 \leq n \leq \log |\log h| - 1$ . For the term  $I_3^7$ , it is easy to obtain

$$I_3^7 = \int_{\{0 \leq |z| < \frac{1}{|\log h|}\}} \overline{\mathcal{K}}_h(z) dz \int_{\mathbb{T}^d} |f_\varepsilon(x) - f_\varepsilon(x-z)|^p dx \leq \sup_{|z| < \frac{1}{|\log h|}} \int_{\mathbb{T}^d} |f_\varepsilon(x+z) - f_\varepsilon(x)|^p dx.$$

Combing the above estimates of  $I_i^7$  ( $i = 1, 2, 3$ ) together, we derive (A.2).

Moreover, for any  $|z| < h$ , due to the Sobolev inequality and

$$f_\varepsilon(x+z) - f_\varepsilon(x) = \int_0^1 z \cdot \nabla f_\varepsilon(x + \omega z) d\omega,$$

it holds

$$\left( \int_{\mathbb{T}^d} |f_\varepsilon(x+z) - f_\varepsilon(x)|^p dx \right)^{\frac{1}{p}} \leq \left( \int_{\mathbb{T}^d} |f_\varepsilon(x+z) - f_\varepsilon(x)|^q dx \right)^{\frac{1}{q}} \leq |z| \|\nabla f_\varepsilon\|_{L^q}, \quad p \in [1, q],$$

and

$$\begin{aligned} & \left( \int_{\mathbb{T}^d} |f_\varepsilon(x+z) - f_\varepsilon(x)|^p dx \right)^{\frac{1}{p}} \\ & \leq \left( \int_{\mathbb{T}^d} |f_\varepsilon(x+z) - f_\varepsilon(x)|^{\frac{qd}{d-q}} dx \right)^{\frac{(d-q)\theta}{qd}} \left( \int_{\mathbb{T}^d} \int_0^1 |z|^q |\nabla f_\varepsilon(x + \omega z)|^q d\omega dx \right)^{\frac{1-\theta}{q}} \\ & \leq C |z|^{1-\theta} \|f_\varepsilon - \frac{1}{|\mathbb{T}^d|} \int_{\mathbb{T}^d} f dx\|_{L^{\frac{qd}{d-q}}}^\theta \|\nabla f_\varepsilon\|_{L^q}^{1-\theta} \\ & \leq C |z|^{1-\theta} \|\nabla f_\varepsilon\|_{L^q}, \quad p \in (q, \frac{qd}{d-q}), \end{aligned}$$

where the constant  $\theta \in (0, 1)$  satisfies  $\frac{1}{p} = \frac{d-q}{qd}\theta + \frac{1}{q}(1-\theta)$ . Thus, (A.3) follows by (A.2) and the above two estimates. The proof of Lemma A.1 is completed.  $\square$

The following compactness criterion is a consequence of Lemma A.1 and the Riesz-Fréchet-Kolmogorov criterion (cf. [4, page 111]).

**Lemma A.2.** *Let  $\varepsilon > 0$ , and  $f_\varepsilon$  be uniformly bounded in  $L^p(\mathbb{T}^d)$  for  $p \in [1, \infty)$ . Then the sequence  $f_\varepsilon$  is strongly compact in  $L^p(\mathbb{T}^d)$  if and only if*

$$\lim_{h \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} L_{h,p}(f_\varepsilon) = 0, \quad (\text{A.4})$$

where  $L_{h,p}(f_\varepsilon)$  is defined by (4.17).

For the compactness in time and space, we have the following lemma of Lions-Aubin type.

**Lemma A.3.** *Let  $T > 0$ ,  $\varepsilon > 0$ ,  $f_\varepsilon$  be uniformly bounded in  $L^p(0, T; L^p(\mathbb{T}^d))$  for  $p \in [1, \infty)$ , and  $\partial_t f_\varepsilon$  be uniform bounded in  $L^q(0, T; W^{-m,1}(\mathbb{T}^d))$  for  $q > 1$  and  $m \geq 0$ . Then  $f_\varepsilon$  is strongly compact in  $L^p(0, T; L^p(\mathbb{T}^d))$  if and only if*

$$\lim_{h \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} \int_0^T L_{h,p}(f_\varepsilon) dt = 0, \quad (\text{A.5})$$

where  $L_{h,p}(f_\varepsilon)$  is defined by (4.17).

**Proof.** For any  $|z| < h < \frac{T}{2}$ , it holds

$$\|f_\varepsilon(x+z, t+h) - f_\varepsilon(x, t)\|_{L^p(0, \frac{T}{2}; L^p)} = \sum_{i=1}^3 I_i^8,$$

with

$$I_1^8 := \|f_\varepsilon(x+z, t+h) - f_\varepsilon(x, t+h)\|_{L^p(0, \frac{T}{2}; L^p)},$$

$$I_2^8 := \|f_\varepsilon * \eta_\delta(x, t+h) - f_\varepsilon * \eta_\delta(x, t)\|_{L^p(0, \frac{T}{2}; L^p)},$$

$$I_3^8 := \|f_\varepsilon(x, t+h) - f_\varepsilon * \eta_\delta(x, t+h)\|_{L^p(0, \frac{T}{2}; L^p)} + \|f_\varepsilon(x, t) - f_\varepsilon * \eta_\delta(x, t)\|_{L^p(0, \frac{T}{2}; L^p)},$$

where  $\eta_\delta \in C_c^\infty(\mathbb{T}^d)$  for  $\delta \in (0, 1)$  is the Friedrichs mollifier. It can be verified by (A.1) and (A.5) that

$$\lim_{h \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} I_1^8 = 0,$$

$$\lim_{h \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} I_2^8 \leq \lim_{h \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} h^{1-\frac{1}{q}} \|\partial_t f_\varepsilon * \eta_\delta\|_{L^q(0, T; L^p)} = 0, \quad \delta \in (0, 1),$$

$$\lim_{\delta \rightarrow 0} \lim_{h \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} I_3^8 \leq 2 \lim_{\delta \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} \sup_{|z| < \delta} \|f_\varepsilon(x+z, t) - f_\varepsilon(x, t)\|_{L^p(0, T; L^p)} = 0.$$

By the above estimates and the Riesz-Fréchet-Kolmogorov criterion,  $f_\varepsilon$  is strongly compact in  $L^p(0, \frac{T}{2}; L^p(\mathbb{T}^d))$ .

Repeating same arguments for  $\tilde{f}_\varepsilon(x, t) := f_\varepsilon(x, T-t)$ , we derive the compactness of  $f_\varepsilon$  in  $L^p(\frac{T}{2}, T; L^p(\mathbb{T}^d))$ .  $\square$

The next lemma is useful in Lemmas 4.6 and 4.8.

**Lemma A.4** ([2, 12]). *For any  $f \in H^1(\mathbb{T}^d)$ , it holds*

$$|f(x) - f(y)| \leq C|x-y|(D_{|x-y|}f(x) + D_{|x-y|}f(y)), \quad (\text{A.6})$$

where  $C > 0$  is a constant depending only on  $d$ , and  $D_r f(x)$  is denoted as

$$D_r f(x) := \frac{1}{r} \int_{|z| \leq r} \frac{|\nabla f(x+z)|}{|z|^{d-1}} dz, \quad (\text{A.7})$$

which satisfies

$$D_r f(x) \leq CM|\nabla f|(x), \quad (\text{A.8})$$

where the localized maximal operator  $M : L^p(\mathbb{T}^d) \rightarrow L^p(\mathbb{T}^d)$  for any  $p \in [1, \infty)$  is defined by

$$Mf(x) := \sup_{r \in (0, 1]} \frac{1}{|B(0, r)|} \int_{B(0, r)} f(x+z) dz. \quad (\text{A.9})$$

In addition, we have

$$\int_{\mathbb{T}^d} \mathcal{K}_h(z) \|D_{|z|}f - D_{|z|}f(\cdot+z)\|_{L^2} dz \leq C\|f\|_{H^1} |\log h|^{\frac{1}{2}}, \quad (\text{A.10})$$

where the kernel  $\mathcal{K}_h$  is given by (4.15).

The following lemma about the commutator estimates of the Riesz operator  $\mathcal{R}_i := (-\Delta)^{-\frac{1}{2}}\partial_i$  was introduced by Coifman-Rochberg-Weiss [15] and Coifman-Meyer [14], and applied to the global existence of weak solution for isentropic compressible Navier-Stokes equations by Lions [37].

**Lemma A.5** ([14, 15]). *For  $p \in (1, \infty)$ , it holds*

$$\|f\mathcal{R}_i\mathcal{R}_jg - \mathcal{R}_i\mathcal{R}_j(fg)\|_{L^p} \leq C\|f\|_{\mathcal{BMO}}\|g\|_{L^p}, \quad f \in \mathcal{BMO}(\mathbb{T}^d), \quad g \in L^p(\mathbb{T}^d), \quad (\text{A.11})$$

where  $\mathcal{BMO}(\mathbb{T}^d)$  denotes the bounded mean oscillation space, and  $C > 0$  is a constant depending only on  $p$  and  $d$ .

Moreover, for  $q_i \in (1, \infty)$  ( $i = 1, 2, 3$ ) satisfying  $\frac{1}{q_1} = \frac{1}{q_2} + \frac{1}{q_3}$ , it holds

$$\|\nabla[f\mathcal{R}_i\mathcal{R}_jg - \mathcal{R}_i\mathcal{R}_j(fg)]\|_{L^{q_1}} \leq C\|\nabla f\|_{L^{q_2}}\|g\|_{L^{q_3}} \quad f \in W^{1,q_2}(\mathbb{T}^d), \quad g \in L^{q_3}(\mathbb{T}^d), \quad (\text{A.12})$$

where  $C > 0$  is a constant depending only on  $q_i$  ( $i = 1, 2$ ) and  $d$ .

**Acknowledgments.** The authors would like to thank the referees for their valuable suggestions and comments on the manuscript. The research of the paper is supported by National Natural Science Foundation of China (No.11931010, 11671384 and 11871047) and by the key research project of Academy for Multidisciplinary Studies, Capital Normal University, and by the Capacity Building for Sci-Tech Innovation-Fundamental Scientific Research Funds (No.007/20530290068).

## References

- [1] J.-W. Barrett, Y. Lu, E. Süli. Existence of large-data finite-energy global weak solutions to a compressible Oldroyd-B model. *Commun. Math. Sci*, 2017, 15: 1265-1323.
- [2] F.B. Belgacem, P.-E. Jabin. Compactness for nonlinear continuity equations. *J. Funct. Anal*, 2013, 264: 139-168.
- [3] C. E. Brennen. *Fundamentals of Multiphase Flow*. Cambridge Univ. Press, 2005.
- [4] H. Brezis, *Functional analysis, Sobolev spaces and partial differential equations*. Universitext. Springer, New York, 2011.
- [5] D. Bresch, B. Desjardins. Existence of global weak solutions for a 2D viscous shallow water equations and convergence to the quasi-geostrophic model. *Commun. Math. Phys*, 2003, 238: 211-223.
- [6] D. Bresch, B. Desjardins, D. Gérard-Varet. On compressible Navier-Stokes equations with density dependent viscosities in bounded domains. *J. Math. Pures Appl*, 2007, 87: 227-235.

- [7] D. Bresch, B. Desjardins, C.-K. Lin. On some compressible fluid models: Korteweg, lubrication, and shallow water systems. *Commun. Partial Differential Equations*, 2003, 28: 843-868.
- [8] D. Bresch, B. Desjardins, J.-M. Ghidaglia, E. Grenier. Global weak solutions to a generic two-fluid model. *Arch. Ration. Mech. Anal.*, 2010, 196: 599–629.
- [9] D. Bresch, X. Huang, J. Li. Global weak solutions to one-dimensional non-conservative viscous compressible two-phase system. *Comm. Math. Phys.*, 2012, 309: 737–755.
- [10] D. Bresch, P.-E Jabin. Global existence of weak solutions for compressible Navier-Stokes equations: thermodynamically unstable pressure and anisotropic viscous stress tensor. *Ann. of Math. (2)*, 2018, 188: 577-684.
- [11] D. Bresch, B. Desjardins, J.M. Ghidaglia, E. Grenier, M. Hilliaret. Multifluid models including compressible fluids. *Handbook of Mathematical Analysis in Mechanics of Viscous Fluids*, Eds. Y. Giga et A. Novotný, 2018, 52: 2927-2978.
- [12] D. Bresch, P.-B Mucha, E. Zatorska. Finite-energy solutions for compressible two-fluid Stokes system. *Arch. Ration. Mech. Anal.*, 2019, 232: 987-1029.
- [13] D. Bresch, A. Vasseur, C. Yu. Global existence of entropy-weak solutions to the compressible Navier-Stokes equations with non-linear density dependent viscosities. *arXiv preprint arXiv:1905.02701*, 2019.
- [14] R.R. Coifman, Y. Meyer. On commutators of singular integrals and bilinear singular integrals, *Trans. Amer. Math. Soc*, 1975, 212: 315-331.
- [15] R.R. Coifman, R. Rochberg, G. Weiss. Factorization theorems for Hardy spaces in several variables. *Ann. of Math. (2)*, 1976, 103: 611-635.
- [16] R.J. DiPerna, P.-L. Lions. Ordinary differential equations, transport theory and Sobolev spaces, *Invent. Math.* 1989, 98: 511-547.
- [17] L. Desvillettes. Some aspects of the modeling at different scales of multiphase flows, *Comput. Methods Appl. Mech. Engrg*, 2010, 199: 1265-1267.
- [18] S. Evje, K.H. Karlsen. Global existence of weak solutions for a viscous two-phase model, *J. Differential Equations*, 2008, 245: 2660-2703.
- [19] S. Evje, Steinar, K.H. Karlsen. Global weak solutions for a viscous liquid-gas model with singular pressure law. *Commun. Pure Appl. Anal.*, 2009, 8: 1867-1894.

- [20] S. Evje. Weak solutions for a gas-liquid model relevant for describing gas-kick in oil wells. *SIAM J. Math. Anal.*, 2011, 43: 1887-1922.
- [21] S. Evje, H. Wen, C. Zhu. On global solutions to the viscous liquid-gas model with unconstrained transition to single-phase flow. *Math. Models Methods Appl. Sci.*, 2017, 27: 323-346.
- [22] S. Evje, W. Wang, H. Wen. Global well-posedness and decay rates of strong solutions to a non-conservative compressible two-fluid model. *Arch. Ration. Mech. Anal.*, 2016, 221: 1285–1316.
- [23] E. Feireisl, A. Novotný, H. Petzeltová. On the existence of globally defined weak solutions to the Navier-Stokes equations. *J. Math. Fluid Mech.*, 2001, 3: 358-392.
- [24] E. Feireisl. Compressible Navier-Stokes equations with a non-monotone pressure law. *J. Differential Equations*, 2002, 184: 97-108.
- [25] E. Feireisl. Dynamics of viscous compressible fluids. *Oxford Lecture Series in Mathematics and its Applications*, 26. Oxford University Press, Oxford, 2004.
- [26] E. Feireisl. A. Novotny, *Singular Limits in Thermodynamics of Viscous Fluids*, Birkhäuser Verlag, Basel, 2009.
- [27] D. Gidaspow. *Multiphase flow and fluidization*, Boston, MA: Academic Press Inc, 1994.
- [28] Z. Guo, J. Yang, L. Yao. Global strong solution for a three-dimensional viscous liquid-gas two-phase flow model with vacuum. *J. Math. Phy.*, 2001, 52: 093102.
- [29] X. Hu. Hausdorff dimension of concentration for isentropic compressible Navier-Stokes equations, *Arch. Ration. Mech. Anal.*, 2019, 234: 375-416.
- [30] C. Hao, H.-L Li. Well-posedness for a multidimensional viscous liquid-gas two-phase flow model, *SIAM J. Math. Anal.*, 2012, 44: 1304-1332.
- [31] M. Ishii. One-dimensional drift-flux model and constitutive equations for relative motion between phases in various two-fluid flow regimes, Tech. report, Argonne National Laboratory Report ANL,77-47, 1977.
- [32] M. Ishii, T. Hibiki. *Thermo-fluid dynamics of two-phase flow*. With a foreword by Lefteri H. Tsoukalas, Springer, New York, 2006.
- [33] S. Jiang, P. Zhang. On spherically symmetric solutions of the compressible isentropic Navier-Stokes equations. *Comm. Math. Phys.*, 2001, 215, 559-581.



- [34] J. Li, Z. Xin. Global existence of weak solutions to the barotropic compressible Navier-Stokes flows with degenerate viscosities, arXiv:1504.06826, 2015.
- [35] Y. Li, Y. Sun. Global weak solutions to a two-dimensional compressible MHD equations of viscous non-resistive fluids. *J. Differential Equations*, 2019, 267: 3827-3851.
- [36] P.-L. Lions. *Mathematical topics in fluid mechanics. Vol. 1. Incompressible models.* Oxford Lecture Series in Mathematics and its Applications, 3. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1996.
- [37] P.-L. Lions. *Mathematical topics in fluid mechanics. Vol. 2. Compressible models.* Oxford Lecture Series in Mathematics and its Applications, 10. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1998.
- [38] D. Maltese, M. Michálek, P. Mucha, A. Novotný, M. Pokorný, E. Zatorska. Existence of weak solutions for compressible Navier-Stokes equations with entropy transport. *J. Differential Equations*, 2016, 261: 4448-4485.
- [39] A. Mellet, A. Vasseur. On the isentropic compressible Navier-Stokes equation. *Commun. Partial Differential Equations*, 2007, 32: 431-452.
- [40] A. Mellet, A. Vasseur. Asymptotic analysis for a Vlasov-Fokker-Planck/compressible Navier-Stokes system of equations. *Comm. Math. Phys*, 2008, 281: 573-596.
- [41] A. Novotný, I. Straškraba. *Introduction to the mathematical theory of compressible flow*, Oxford Lecture Series in Mathematics and its Applications, 27. Oxford University Press, Oxford, 2004.
- [42] A. Novotný, M. Pokorný. Weak solutions for some compressible multicomponent fluid models. *Arch. Rational Mech. Anal*, 2020, 235: 355-403.
- [43] P. I. Plotnikov, W. Weigant. Isothermal Navier-Stokes equations and Radon transform. *SIAM J. Math. Anal*, 2015, 47: 626-653.
- [44] A. Vasseur, C. Yu. Existence of global weak solutions for 3D degenerate compressible Navier-Stokes equations. *Inventiones mathematicae*, 2016, 206: 1–40.
- [45] A. Vasseur, H. Wen, C. Yu. Global weak solution to the viscous two-fluid model with finite energy, *J. Math. Pures Appl.* (9), 2019, 125: 247-282.
- [46] W. Wang, W. Wang. Large time behavior for the system of a viscous liquid-gas two-phase flow model in  $\mathbb{R}^3$ , *Journal of Differential Equations*, 2016, 261: 5561-5589.

- [47] Y. Wang, H. Wen, L. Yao. On a non-conservative compressible two-fluid model in a bounded domain: global existence and uniqueness. *Journal of Mathematical Fluid Mechanics*, 2021, 23: 1-24.
- [48] H. Wen, L. Zhu. Global well-posedness and decay estimates of strong solutions to a two-phase model with magnetic field. *J. Differential Equations*, 2018, 264: 2377–2406.
- [49] H. Wen, L. Yao, C. Zhu. Review on mathematical analysis of some two-phase flow models. *Acta Mathematica Scientia*, 2018, 38: 1617-1636.
- [50] H. Wen. On global solutions to a viscous compressible two-fluid model with unconstrained transition to single-phase flow in three dimensions, *Calculus of Variations and Partial Differential Equations*, 2021, 60: 1-38.
- [51] H. Wen, C. Zhu. Remarks on global weak solutions to a two-fluid type model, *Communications on Pure Applied Analysis*, 2021.
- [52] G.B. Wallis. *One-dimensional two-fluid flow*, McGraw-Hill, New York, 1979.
- [53] L. Yao, T. Zhang, C. Zhu. Existence of asymptotic behavior of global weak solutions to a 2D viscous liquid-gas two-phase flow model. *SIAM J. Math. Anal.*, 2010, 42: 1874-1897.
- [54] L. Yao, C. Zhu. Free boundary value problem for a viscous two-phase model with mass-dependent viscosity. *J. Differential Equations*, 2009, 247: 2705-2739.
- [55] L. Yao, C. Zhu. Existence and uniqueness of global weak solution to a two-phase flow model with vacuum. *Math. Ann.*, 2011, 349: 903-928.
- [56] L. Yao, J. Yang, Z. Guo. Global classical solution for a three-dimensional viscous liquid-gas two-fluid flow model with vacuum. *Acta Math. Appl. Sin. Engl. Ser.*, 2014, 30: 989–1006.
- [57] H. Yu. Global strong solutions to the 3D viscous liquid-gas two-phase flow model. *Journal of Differential Equations*, 2021, 272: 732-759.
- [58] Y. Zhang, C. Zhu. Global existence and optimal convergence rates for the strong solutions in  $H^2$  to the 3D viscous liquid-gas two-phase flow model. *Journal of Differential Equations*, 2015, 258: 2315-2338.
- [59] N. Zuber. On the dispersed two-phase flow in the laminar flow regime. *Chem. Engrg. Sci.*, 1964, 19: 897-917.