On a formula that is not in "Grothendieck Topologies in Posets"


Eduardo Ochs

November 18, 2021

Abstract

The paper [Lin14] shows that when \mathbf{P} is an Artinian poset and \mathbf{E} is the topos $\mathbf{Set}^{\mathbf{P}}$ then there are bijections between the set of subsets of \mathbf{P} , the set of Grothendieck topologies on \mathbf{E} , and the set of nuclei on the Heyting Algebra $\mathrm{Sub}(1_{\mathbf{E}})$. It also shows that there are nice formulas for converting between subsets, Grothendieck topologies, and nuclei, but the formula for converting a nucleus to a subset is not spelled out explicitly. These notes fix that gap.

Let \mathbf{P} be a downward-directed poset, that we will also regard as a category. Then $\mathbf{E} := \mathbf{Set^P}$ is a topos and $H := \mathrm{Sub}(1_{\mathbf{E}})$ is a Heyting Algebra. Let's denote the set of all Grothendieck topologies on \mathbf{E} by $\mathsf{GrTops}(\mathbf{P})$, the set of all nuclei on H by $\mathsf{Nucs}(\mathbf{P})$, and the set of subsets of (the set of points of) \mathbf{P} by $\mathcal{P}(\mathbf{P})$. In [Lin14] Bert Lindenhovius shows that when \mathbf{P} is Artinian we have bijections between $\mathcal{P}(\mathbf{P})$, $\mathsf{Nucs}(\mathbf{P})$, and $\mathsf{GrTops}(\mathbf{P})$. He uses notational conventions in which J always denotes a Grothendieck topology, j always denotes a nucleus, and X always denotes a subset of \mathbf{P} , and he writes the components these bijections as $(j \mapsto J_j)$, $(J \mapsto j_J)$, and so on; we will also write them here as $(j \mapsto J)$, $(J \mapsto j)$, etc, and we will write the bijections as $(X \leftrightarrow j)$, $(X \leftrightarrow J)$, and $(j \leftrightarrow J)$. Let's put all this in a diagram:



He defines the components of these bijections as:

The annotations like "(C.4.2, C.2)" indicate where these components are defined. Note that one of the annotations says "(e-mail)"; this is because that formula doesn't appear explicitly in [Lin14], and so I (Eduardo) asked him (Bert) if that formula was what I guessed it would be, and he replied with a formula slightly shorter than my guess, and a proof...

These notes are just to make his formula and his proof available in a public place. All the mathematical content here is by Bert Lindenhovius, and all the typesetting was done by Eduardo Ochs, who found Bert's proof hard to follow and decided to typeset it in Natural Deduction form. *Note:* when I wrote the first version of these notes I listed Bert as the main author and me as the coauthor, but he told me that he preferred to be credited only in the text. I insisted, and explained that I would have never been able to find these proofs by myself — but he insisted more.

If we combine $(j \mapsto J)$ and $(J \mapsto X)$ we get this:

$$\begin{array}{lll} (j \mapsto J): & J_{j}(p) & = & \left\{S \in \mathcal{D}(\downarrow p) : p \in j(S)\right\} \\ (J \mapsto X): & X_{J} & = & \left\{p \in \mathbf{P} : J(p) = \left\{\downarrow p\right\}\right\} \\ (j \mapsto J \mapsto X): & X_{J_{j}} & = & \left\{p \in \mathbf{P} : \left\{S \in \mathcal{D}(\downarrow p) : p \in j(S)\right\} = \left\{\downarrow p\right\}\right\} \\ & = & \left\{p \in \mathbf{P} : \forall S \in \mathcal{D}(\downarrow p). \left((p \in j(S)) \leftrightarrow (S = \downarrow p)\right)\right\} \\ (j \mapsto X): & X_{j} & = & \left\{p \in \mathbf{P} : p \not\in j(\downarrow p \setminus \{p\})\right\} \end{array}$$

It is not obvious at all that $X_{J_j} = X_j$. We will prove that $p \in X_j$ iff $p \in X_{J_j}$, where:

$$(p \in X_j) = (p \notin j(\downarrow p \setminus \{p\}))$$

$$(p \in X_{J_j}) = \forall S \in \mathcal{D}(\downarrow p). (p \in j(S)) \leftrightarrow (S = \downarrow p)$$

Look:

Look:
$$\frac{p \in X_{J_j}}{\downarrow p \setminus \{p\} \in \mathcal{D}(\downarrow p)} \qquad \overline{\forall S \in \mathcal{D}(\downarrow p).(p \in j(S)) \leftrightarrow (S = \downarrow p)} \\
\underline{(p \in j(\downarrow p \setminus \{p\})) \leftrightarrow (\downarrow p \setminus \{p\} = \downarrow p)} \\
\underline{(p \notin j(\downarrow p \setminus \{p\})) \leftrightarrow (\downarrow p \setminus \{p\} \neq \downarrow p)} \qquad \overline{\downarrow p \setminus \{p\} \neq \downarrow p} \\
\underline{p \notin j(\downarrow p \setminus \{p\})} \\
\underline{p \notin j(\downarrow p \setminus \{p\})} \\
\underline{p \in X_j}$$

$$S \in \mathcal{D}(\downarrow p)]^2 \\
\underline{S \in J(p)} \qquad [S \neq \lfloor p \rfloor^1 \qquad [S \in \mathcal{D}(\lfloor p)]^2]$$

$$F \in X_{j}$$

$$\underline{S \subset \downarrow p} \qquad [S \neq \downarrow p]^{1} \qquad \underline{[S \in \mathcal{D}(\downarrow p)]^{2}}$$

$$\underline{P \notin S} \qquad \underline{S \subset \downarrow p}$$

$$\underline{S \subset \downarrow p \setminus \{p\}} \qquad p \in X_{j}$$

$$\underline{j(S) \subset j(\downarrow p \setminus \{p\})} \qquad p \notin j(\downarrow p \setminus \{p\})$$

$$\underline{P \notin j(S)} \qquad 1$$

$$\underline{(S \neq \downarrow p) \to (p \notin j(S))} \qquad (S = \downarrow p) \qquad (S = \downarrow p) \to (p \in j(S))$$

$$\underline{(p \in j(S)) \leftrightarrow (S = \downarrow p)} \qquad 2$$

$$\underline{P \notin J(S)} \qquad P \notin J(S) \qquad P$$

This proves that X_j and X_{J_j} are equal.

Now let's check that X_j and X'_j are equal, where X'_j is defined as:

$$X_{j} = \{ p \in \mathbf{P} : p \notin j(\downarrow p \setminus \{p\}) \}$$

$$X'_{j} = \{ p \in \mathbf{P} : j(\downarrow p) \neq j(\downarrow p \setminus \{p\}) \}$$

Proof:

$$\frac{[p \in j(\downarrow p \setminus \{p\})]^{1}}{\downarrow p \subseteq j(\downarrow p \setminus \{p\})} \frac{\downarrow p \setminus \{p\} \subseteq \downarrow p}{j(\downarrow p \setminus \{p\}) \subseteq j(\downarrow p)} \frac{\downarrow p \subseteq j(\downarrow p)}{j(\downarrow p) \subseteq (j \circ j)(\downarrow p \setminus \{p\})} \frac{\downarrow p \subseteq j(\downarrow p)}{j(\downarrow p) \subseteq j(\downarrow p \setminus \{p\})} \frac{\downarrow p \subseteq j(\downarrow p)}{\downarrow p \subseteq j(\downarrow p \setminus \{p\})} \frac{\downarrow p \subseteq j(\downarrow p) = j(\downarrow p \setminus \{p\})}{\downarrow p \subseteq j(\downarrow p \setminus \{p\})} \frac{j(\downarrow p) = j(\downarrow p \setminus \{p\})}{p \in j(\downarrow p \setminus \{p\})} 1$$

$$\frac{(p \in j(\downarrow p \setminus \{p\})) \leftrightarrow (j(\downarrow p) = j(\downarrow p \setminus \{p\}))}{(p \notin j(\downarrow p \setminus \{p\})) \leftrightarrow (j(\downarrow p) \neq j(\downarrow p \setminus \{p\}))} 1$$

$$p \in X_{j} \leftrightarrow p \in X'_{j}$$

The formula of X'_j above is the one that I asked if it was correct; Bert answered that yes, and showed that it is equivalent to the slighty shorter formula for X_j .

References

[Lin14] A.J. Lindenhovius. "Grothendieck Topologies on Posets". https://arxiv.org/pdf/1405.440 2014.