

# Amplitudes in persistence theory

Barbara Giunti<sup>\*</sup>, John S. Nolan<sup>†</sup>, Nina Otter<sup>‡</sup>, Lukas Waas<sup>§</sup>

## Abstract

The use of persistent homology in applications is justified by the validity of certain stability results. At the core of such results is a notion of distance between the invariants that one associates with data sets. Here we introduce a general framework to study stability questions in multiparameter persistence, where there is no natural choice of invariants and distances between them. We define amplitudes, monotone, and subadditive invariants that arise from assigning a non-negative real number to objects of an abelian category. We then present different ways to associate distances to such invariants, and we provide a classification of classes of amplitudes relevant for topological data analysis. We study distances associated with such amplitudes by investigating their discriminative power as well as relationships between different choices of distances, and show with examples how this framework can help in proving stability results in topological data analysis.

## 1 Introduction

One-parameter persistent homology (PH) is, arguably, the most successful method in Topological Data Analysis (TDA), since it has been used in a variety of applications, ranging from nanoporous materials [LBD<sup>+</sup>17] to pulmonary diseases [BPC<sup>+</sup>18], with new preprints being released on a weekly basis<sup>1</sup>.

In one-parameter PH, one associates a 1-parameter family of spaces to a data set, and by computing the homology of this family one then extracts algebraic invariants, called “barcodes”, which give a summary of how topological invariants, such as the number of connected components, holes, and voids, evolve as the parameter value changes. PH can meaningfully be used for data analysis thanks to stability results, which, roughly, guarantee that the pipeline from the input data to the barcodes is Lipschitz, with respect to appropriate notions of distances on the input data and barcodes. Such distances are well-understood from a theoretical point of view, since they rely on matchings between intervals in barcodes, and they differ in how the cost between matchings is computed. In what is called the “bottleneck distance”, one considers the cost of matching the largest intervals, while in what are generally called “ $p$ -Wasserstein distances” one takes the weighted sum of the costs of the matching over all the intervals. Of these distances, the bottleneck distance has been most widely used in applications [BW20, HKNU17] and has been studied the most from a theoretical point of view. However, for many applications, the bottleneck distance is too coarse, and there have recently been renewed efforts in studying stability results for  $p$ -Wasserstein distances [ST20].

---

<sup>\*</sup>Institute of Geometry, Graz University of Technology, Graz, Austria, *email*: bgiunti@albany.edu

<sup>†</sup>Department of Mathematics, UC Berkeley, Berkeley, CA, US, *email*:

<sup>‡</sup>Department of Mathematics, UCLA, Los Angeles, CA, US, *email*: n.otter@qmul.ac.uk

<sup>§</sup>Department of Mathematics, Heidelberg University, Germany, *email*: lwaas@mathi.uni-heidelberg.de

<sup>1</sup>A database of applications of methods in TDA, including persistent homology, is being maintained at [GLR22], and at the time of writing it includes more than two hundred applications of persistent homology. The entries in the database can be filtered by application area, data format, and topological methods.

In many applications, one is interested in studying data sets across multiple parameters, and thus in what is called “multiparameter persistent homology” one wishes to study the homology of a multiparameter family of spaces associated to a data set. It is well-known that there is no generalization of the barcode for multiparameter families of spaces [CZ09]. In practice, one thus chooses invariants that are meaningful for a specific problem, but which do not give a complete characterization of the resulting persistence module.

Choosing an appropriate invariant in multiparameter persistence requires defining what it means to be “persistent”, so as to be able to distinguish between topological signal and noise. To do this, one could first focus either on finding an appropriate notion of signal, or of noise. Several different approaches, focused on finding an appropriate notion of signal, have been taken in the literature. The first invariant that was suggested is the “rank invariant” [CZ09], which encodes the information on the ranks of structure morphisms in the module. More recently, new invariants have been proposed, thanks to the introduction of new techniques into the field, stemming from representation theory [BE18, BBLO20, BBCB20], and commutative algebra [HOST19, Tho19, Mil20a, BBOO21]. To be able to study the stability of such invariants, one usually needs to first associate suitable distances to them. Studying distances associated to such choices of invariants is currently one of the most active areas of research in TDA. Several different notions have been proposed, including the interleaving distance which, in an appropriate sense, generalizes the bottleneck distance for multiparameter persistence, see [Les15]. More recently, generalizations of Wasserstein-type distances and  $\ell^p$ -distances have been studied in [BSS23] and [BBL21], respectively.

In a complementary line of work, the authors of [SCL<sup>+</sup>17] proposed an approach to study notions of “noise” for multiparameter persistence modules, with the idea of stabilizing invariants by minimizing them on disks, rather than proving a stability result for each of them. Such an approach to minimizing invariants was also taken, in the 1-parameter setting, in [BMSW17].

In our work, we take a different perspective: we focus on classes of invariants that capture a certain notion of “size” or “measure” of persistence, and on ways of associating distances with such invariants. Specifically, given an abelian category  $\mathcal{A}$ , we are interested in studying real-valued maps  $\alpha: \text{ob } \mathcal{A} \rightarrow [0, \infty]$  that mimic the properties that we would expect a size function to satisfy. Namely, we would like the map to be monotone with respect to subobjects and quotients, and to be subadditive with respect to short exact sequences (see Section 2).

Since our motivation stems from studying invariants arising from  $n$ -parameter persistence modules, we then proceed to investigate specific properties of such smaller classes of invariants. To an amplitude  $\alpha$  defined for  $n$ -parameter persistence modules, we may associate a function  $\mu_\alpha$  defined on intervals in  $\mathbb{R}^n$ , by evaluating  $\alpha$  on interval modules. We may then attempt to study  $\alpha$  in terms of this function  $\mu_\alpha$ . For a persistence module  $M$ , satisfying appropriate finiteness conditions, we obtain lower and upper bounds for  $\alpha$  entirely in terms of  $\mu_\alpha$  (see Proposition 4.3 and the following section). Furthermore, the amplitudes for which either one of these bounds are strict may be classified entirely in terms of their function on intervals  $\mu_\alpha$  (Theorems 4.11 and 4.18). These are respectively for the lower and the upper bound: “quasi-simple” amplitudes which behave like a maximum under direct sums and provide a generalization of the simple amplitudes of [SCL<sup>+</sup>17]; “additive” which are additive under short exact sequences (see Section 4.1 and Section 4.2, respectively).

We then proceed to study distances associated to amplitudes on categories of persistence modules. In general, such distances are extended pseudometrics. We show that under appropriate assumptions, such a pseudometric is a metric exactly when it is induced by an amplitude whose value is 0 only for zero objects (see Section 5). We then study relationships between distances arising from relationships between different amplitudes. In particular, we focus on stability results, and we show how with our framework we can deduce new stability results from known ones (see Section 6). We believe this to be an important contribution for applications of persistent homology

to the study of data, as in such applications one often seeks the best way to capture persistence, depending on the specific problem. Therefore one often is interested in several amplitudes at once and in studying how to relate these different amplitudes in a stable way.

Finally, we also show how our framework can be used to study the stability of invariants that fail to be amplitudes. This is the case for the  $c_p$ -rank, which is not an amplitude as it fails to be monotone, but whose stability can nevertheless be studied in our framework (see Section 7).

We conclude this introduction with an example illustrating the usefulness of our framework for applications in which PH is used as a representation of data and fed into a Machine Learning pipeline. In such applications, one usually associates a vector to the barcode obtained. All such vectorizations usually depend on a parameter, which in turn becomes a hyperparameter to tune in the Machine Learning pipeline. With our framework, we can study how vectorizations corresponding to different choices of parameters are related, and thus it may help in optimizing a grid search during hyperparameter tuning.

**Example 1.1.** Several vectorizations are considered in PH [AAJ<sup>+</sup>23], such as persistence landscapes [Bub15], persistence images [AEK<sup>+</sup>17], and tropical coordinates [Kal19]. In this example, we showcase how our framework allows to relate the different choices for the hyperparameter of the vectorization. We choose as a vectorization the tropical coordinate given by the sum of the lengths of the  $k$  longest intervals, where the natural number  $k$  is the hyperparameter.

Suppose that we are given two different persistence modules  $M$  and  $M'$ , and their corresponding persistence diagrams PD and PD', which one might think of as representations of two different data points. We then may aim to associate different vectorizations to such persistence diagrams, for instance, corresponding to taking the  $k$  or  $l$  longest intervals. We then have the following relationship between such vectorizations:

$$d_l(M, M') \leq \max \left\{ 1, \frac{l}{k} \right\} d_k(M, M')$$

(see Corollary 6.5), where  $d_l$  and  $d_k$  are the distances on the persistence modules induced by taking the  $l$  longest or  $k$  longest intervals, respectively. Therefore, even if the entire Machine Learning pipeline might not be stable, we have control over how different vectorizations are related. In Figure 1 we give an illustration of such a pipeline.




Figure 1: We consider a persistence module and its persistence diagram obtained by computing persistent homology in a specific degree for a given data point. We then aim to associate a vectorization to such a persistence diagram, for instance by associating to it the vector of  $k$ -longest intervals, for different values of  $k$ , e.g.,  $k = 1$  or  $k = 3$ . Given such a vectorization for all data points in the data set, we then proceed to a classification or regression task using a Machine Learning method, such as SVM.

## 2 Amplitudes on abelian categories

We begin with the definition of amplitudes, which are real-valued maps that behave well with respect to the structure of abelian categories.

**Definition 2.1.** For an abelian category  $\mathcal{A}$ , a class function  $\alpha: \text{ob } \mathcal{A} \rightarrow [0, \infty]$  is called an **amplitude** if  $\alpha(0) = 0$  and for every short exact sequence

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

in  $\mathcal{A}$  the following inequalities hold:

$$\begin{aligned} \text{(Monotonicity)} \quad & \alpha(A) \leq \alpha(B) \\ & \alpha(C) \leq \alpha(B) \quad , \end{aligned} \tag{1}$$

$$\text{(Subadditivity)} \quad \alpha(B) \leq \alpha(A) + \alpha(C) \quad . \tag{2}$$

Moreover, if (2) is an equality, the amplitude is said to be **additive**, and if  $\alpha(A) < \infty$  for all  $A$  in  $\text{ob } \mathcal{A}$ , then the amplitude is said to be **finite**.

We note that as a direct consequence of the monotonicity property, an amplitude is an isomorphism invariant.

**Example 2.2.** (Max degree) Consider the category  $\text{Ch}$  of non-negatively-graded chain complexes over an abelian category, and chain maps between them. Define an amplitude  $\delta$  on  $\text{Ch}$  as

$$\delta(C) := \max\{n \in \mathbb{N} \mid C_n \neq 0\}$$

for all  $C$  in  $\text{ob } \text{Ch}$ . It is clear that  $\delta(0) = 0$ . Since a short sequence in  $\text{Ch}$  is exact if and only if it is exact degreewise, we also have the properties of monotonicity (1) and subadditivity (2). Moreover,  $\delta$  is finite. However, it is immediate to see that, in general, (2) is a strict inequality for this amplitude, and thus  $\delta$  is not additive.

**Example 2.3.** In [ST20], the authors consider  $p$ -norms of finitely presented 1-parameter persistence modules and use them to define distances isometric to the  $p$ -Wasserstein distances. Here, we note that  $p$ -norms are amplitudes. Let  $\oplus_{j \in J} \{\mathbb{I}(b_j, d_j)\}_{j \in J}$  be the finite barcode decomposition of a persistence module  $M$  (see, for example, [Oud15] for the precise definition of these terms, which we will introduce in Section 3). The angular brackets denote that the interval can be open or closed on each endpoint. Then the  $p$ -norm of  $M$ , for  $p \in [1, \infty]$ , is

$$\rho_p(M) := \begin{cases} \left( \sum_{j \in J} |d_j - b_j|^p \right)^{\frac{1}{p}} & \text{if } p < \infty \\ \max_{j \in J} |d_j - b_j| & \text{if } p = \infty \end{cases} .$$

Using the results of [BL15], the authors of [ST20] show that such  $p$ -norms satisfy the monotonicity and subadditivity conditions (1)-(2). In particular,  $\rho_1$  is an additive amplitude.

### 2.1 Span metrics

We now introduce metrics associated with amplitudes. A classical method to define a metric is to minimize the cost of a path between two objects. By properties of amplitudes, the path can be shortened into a span, following the technique of [SCL<sup>+</sup>17].

**Definition 2.4.** A **cost function** is a function  $f: [0, \infty]^4 \rightarrow [0, \infty]$  satisfying the following conditions:

- Monotonicity:  $f(v) \leq f(w)$  whenever  $v \leq w$ .
- Subadditivity:  $f(v + w) \leq f(v) + f(w)$  for any  $v, w$ .
- Behavior at zero:  $f(0, 0, 0, 0) = 0$ .
- Exchange law:  $f(x_1, x_2, x_3, x_4) = f(x_3, x_2, x_1, x_4) = f(x_1, x_4, x_3, x_2)$ .

**Example 2.5.** The 1-norm  $\| - \|_1: [0, \infty]^4 \rightarrow [0, \infty]$ , taking  $(x_1, x_2, x_3, x_4)$  to  $\sum_{i=1}^4 x_i$ , is a cost function.

**Definition 2.6.** Let  $\mathcal{A}$  be an abelian category with amplitude  $\alpha$ . Given a span  $\mathbf{c}: A \xleftarrow{\varphi} C \xrightarrow{\psi} B$  in  $\mathcal{A}$  and a cost function  $f$ , the  **$f$ -cost of  $\mathbf{c}$**  is

$$f_\alpha(\mathbf{c}) := f(\alpha(\ker \varphi), \alpha(\operatorname{coker} \varphi), \alpha(\ker \psi), \alpha(\operatorname{coker} \psi)).$$

One can similarly define the  $f$ -cost of a cospan.

**Remark 2.7.** For every cost function  $f$ , we have that, given a span  $\mathbf{c}: A \xleftarrow{\varphi} C \xrightarrow{\psi} B$ , there exists a cospan  $\mathbf{c}': A \xrightarrow{\psi'} C \xleftarrow{\varphi'} B$  with  $f_\alpha(\mathbf{c}') \leq f_\alpha(\mathbf{c})$ , and similarly with the roles of spans and cospans swapped. The proof is in [SCL<sup>+</sup>17, Cor. 8.3] for  $f = \| - \|_1$ , but generalizes immediately to our setting.

We use the  $f$ -cost to define a (pseudo, extended) metric:

**Definition 2.8.** Given an amplitude  $\alpha$  on an abelian category  $\mathcal{A}$  and a cost function, the **span metric associated to  $\alpha$**  is:

$$d_{f_\alpha}(A, B) = \inf \left\{ f_\alpha(\mathbf{c}) \mid \mathbf{c}: A \xleftarrow{\varphi} C \xrightarrow{\psi} B \right\},$$

with the convention that  $\inf \emptyset = \infty$ . When  $f = \| - \|_1$ , we say simply **span metric associated to  $\alpha$** , denoted by  $d_\alpha$ .

**Proposition 2.9.** Let  $\mathcal{A}$  be an abelian category with amplitude  $\alpha$ , and let  $f$  be a cost function. Then the assignment  $d_{f_\alpha}$  is a pseudometric on  $\operatorname{ob} \mathcal{A}$ .

*Proof.* It is clear that  $d_{f_\alpha}(A, A) = 0$  and that  $d_{f_\alpha}(A, B) = d_{f_\alpha}(B, A)$  for every  $A, B$  in  $\operatorname{ob} \mathcal{A}$ . For the triangle inequality, consider two spans  $\mathbf{c}_1: A \xleftarrow{\varphi_1} H_1 \xrightarrow{\psi_1} B$  and  $\mathbf{c}_2: B \xleftarrow{\varphi_2} H_2 \xrightarrow{\psi_2} C$ . By Remark 2.7 applied to the span  $H_1 \xrightarrow{\psi_1} B \xleftarrow{\varphi_2} H_2$ , we obtain a span

$$\mathbf{c}_3: A \xleftarrow{\varphi_1} H_1 \xleftarrow{\varphi_3} H_3 \xrightarrow{\psi_3} H_2 \xrightarrow{\psi_2} C.$$

Thus, it is left to show that  $f_\alpha(\mathbf{c}_3) \leq f_\alpha(\mathbf{c}_1) + f_\alpha(\mathbf{c}_2)$ . This follows by the argument presented in [SCL<sup>+</sup>17, Prop. 8.2] and the properties of  $f$ .  $\square$

**Remark 2.10.** This result is not true in the more general case of weight functions [BSS23] and therefore provides motivation for the use of amplitudes or noise systems. Indeed, one of the motivations for the introduction of the axioms of noise systems [SCL<sup>+</sup>17] was to allow for studying distances using only (co)spans instead of arbitrary zigzags.

**Example 2.11.** An interval module is an object in  $\text{Fun}(\mathbb{R}, \text{Vect})$  that assumes value  $\mathbb{F}$  over an interval of  $\mathbb{R}$  of the form  $[a, b)$ , with the identity as a transition morphism, and 0 otherwise. Consider two interval modules  $\mathbb{I}[a, a+r)$  and  $\mathbb{I}[b, b+r)$ , for  $r > 0 \in \mathbb{R}$  and  $a, b \in \mathbb{R}$  with  $a+r < b$ , and  $\alpha = \rho_p$ . We have  $d_{\rho_p}(\mathbb{I}[a, a+r), \mathbb{I}[b, b+r)) = 2r$ , and taking  $f = \max$ ,  $d_{f_{\rho_p}}(\mathbb{I}[a, a+r), \mathbb{I}[b, b+r)) = r$ .

**Lemma 2.12.** *Let  $\mathcal{A}$  be an abelian category with amplitude  $\alpha$ . Then, for any  $A, B$  in  $\text{ob } \mathcal{A}$ , the following inequality holds:*

$$|\alpha(A) - \alpha(B)| \leq d_\alpha(A, B).$$

In other words,  $\alpha$  is 1-Lipschitz with respect to  $d_\alpha$ .

*Proof.* By the triangle inequality and by definition of  $d_\alpha$ , it suffices to show that whenever there is a morphism  $f: A \rightarrow B$  with  $c_\alpha(f) \leq \varepsilon$ , then

$$|\alpha(A) - \alpha(B)| \leq \varepsilon.$$

Again by the triangle inequality and epi-mono factorization, we may restrict to the case where  $f$  is either an epi- or monomorphism. If  $f$  is a mono, we then have a short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow \text{coker } f \rightarrow 0$  with  $\alpha(\text{coker}(f)) \leq \varepsilon$ . Hence, by monotonicity and subadditivity of amplitudes follows that,

$$\alpha(A) \leq \alpha(B) \leq \alpha(A) + \varepsilon,$$

showing the required inequality. The case of  $f$  an epimorphism follows dually.  $\square$

**Lemma 2.13.** *Let  $\mathcal{A}$  be an abelian category and  $\alpha$  an amplitude on  $\mathcal{A}$ . Then, for each  $A$  in  $\text{ob } \mathcal{A}$ ,*

$$\alpha(A) = d_\alpha(A, 0).$$

*Proof.* We observe that  $\alpha(A) = c_\alpha(A \rightarrow 0)$ , and thus  $d_\alpha(A, 0) \leq \alpha(A)$ .

For the converse inequality, consider a cospan  $A \xrightarrow{\varphi} B \leftarrow 0$ . It suffices to show that  $c_\alpha(A \rightarrow 0) \leq c_\alpha(\varphi) + c_\alpha(0 \rightarrow B)$  to obtain the missing inequality by passing to the infimum. Since  $0 = 0 \circ \varphi$ , we have

$$c_\alpha(A \rightarrow 0) \leq c_\alpha(\varphi) + c_\alpha(B \rightarrow 0) = c_\alpha(\varphi) + c_\alpha(0 \rightarrow B). \quad \square$$

In data analysis, stability is a crucial property as it guarantees the robustness of the analysis: the extracted information (aka, the amplitude) does not vary excessively if the input data is only slightly perturbed. In more detail, we want to have that the span metric associated with a certain cost function of an amplitude is bounded by some (meaningful) metric on the input. This property is in general difficult to prove. To ease the process, we now present results bounding span metrics with each other. Using this technique, one can then prove a stability result for a certain span metric and then transfer it to other span metrics.

**Definition 2.14.** A **cost-processing function** is a function  $g: [0, \infty] \rightarrow [0, \infty]$  satisfying:

- Monotonicity:  $g(v) \leq g(w)$  whenever  $v \leq w$ .
- Subadditivity:  $g(v + w) \leq g(v) + g(w)$ .
- Behavior at zero:  $g(0) = 0$ .

**Example 2.15.** For any positive integer  $n$ , the function  $\sqrt[n]{\phantom{x}}$  is a cost-processing function.

If  $f: [0, \infty]^4 \rightarrow [0, \infty]$  is a cost function, and  $g: [0, \infty] \rightarrow [0, \infty]$  is a cost-processing function, then  $g \circ f: [0, \infty]^4 \rightarrow [0, \infty]$  is a cost function. Furthermore, we have the following bound on span metrics:

**Proposition 2.16.** *Let  $\mathcal{A}$  be an abelian category with amplitude  $\alpha$ ,  $f$  a cost function, and  $g$  a cost-processing function. Then*

$$d_{(g \circ f)_\alpha}(A, B) \leq 4g(d_{f_\alpha}(A, B))$$

for all  $A, B \in \text{ob } \mathcal{A}$ .

*Proof.* Consider a (co)span  $\mathbf{c}: A \xleftarrow{\varphi} C \xrightarrow{\psi} B$ . For ease of notation, write  $\alpha_1 = \alpha(\ker \varphi)$ ,  $\alpha_2 = \alpha(\text{coker } \varphi)$ ,  $\alpha_3 = \alpha(\ker \psi)$ , and  $\alpha_4 = \alpha(\text{coker } \psi)$ . Write  $f_i = f(0, \dots, \alpha_i, \dots, 0)$ , where  $\alpha_i$  is the  $i$ -th input. Then

$$f(\alpha_1, \alpha_2, \alpha_3, \alpha_4) \leq \sum_{i=1}^4 f_i \leq 4 \max_i f_i \leq 4f_\alpha(\mathbf{c}).$$

Therefore  $g(f(\alpha_1, \alpha_2, \alpha_3, \alpha_4)) \leq 4g(f_\alpha(\mathbf{c}))$ . Taking infima over all possible cospans  $\mathbf{c}$  yields the claim.  $\square$

**Proposition 2.17.** *Let  $g: [0, \infty] \rightarrow [0, \infty]$  be a cost-processing function. Let  $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$  be an exact functor that satisfies  $\alpha'(F(A)) \leq g(\alpha(A))$  for all  $A \in \text{ob } \mathcal{A}$ . Then we have*

$$d_{\alpha'}(F(A), F(B)) \leq 4g(d_\alpha(A, B))$$

for all  $A, B \in \text{ob } \mathcal{A}$ . In particular, if  $g$  is continuous at 0,  $F$  induces a continuous function  $\text{ob } \mathcal{A} \rightarrow \text{ob } \mathcal{A}'$  with respect to the topologies induced by  $d_\alpha$  and  $d_{\alpha'}$ .

*Proof.* Let  $f = \|\cdot\|_1$ . Consider a (co)span  $\mathbf{c}: A \xleftarrow{\varphi} C \xrightarrow{\psi} B$ . As above, write  $\alpha_1 = \alpha(\ker \varphi)$ ,  $\alpha_2 = \alpha(\text{coker } \varphi)$ ,  $\alpha_3 = \alpha(\ker \psi)$ , and  $\alpha_4 = \alpha(\text{coker } \psi)$ . Then

$$\begin{aligned} d_{f_\alpha}(F(A), F(B)) &\leq f_{\alpha'}(F(\mathbf{c})) \leq \sum_{i=1}^4 g(\alpha_i) \leq 4 \max_i g(\alpha_i) = 4g\left(\max_i \alpha_i\right) \\ &\leq 4g\left(\sum_{i=1}^4 \alpha_i\right) = 4(g \circ f)_\alpha(\mathbf{c}). \end{aligned}$$

Now take the infimum over all  $\mathbf{c}$  and apply Proposition 2.9.  $\square$

## 2.2 Noise systems and amplitudes

The idea of amplitude is closely related to that of a noise system, introduced in [SCL<sup>+</sup>17]. While the concept of amplitude arises from the aim to capture a notion of size for objects in an abelian category, noise systems were introduced to define stable approximations of persistence modules, and thus a noise system can be seen as a way to hierarchically organize persistence modules according to how small they are. Here we define noise systems, introduced in [SCL<sup>+</sup>17] for tame persistence modules  $\mathbb{Q}^n \rightarrow \text{Vect}$ , for arbitrary abelian categories and show that they are related to amplitudes by an equivalence between appropriately defined categories.

**Definition 2.18.** Let  $\mathcal{A}$  be an abelian category and  $[0, \infty)$  the poset of non-negative reals. A **noise system** in  $\mathcal{A}$  is a collection  $\mathcal{S} = \{\mathcal{S}_\varepsilon\}_{\varepsilon \in [0, \infty)}$  of collections of objects of  $\mathcal{A}$ , indexed by non-negative reals  $\varepsilon$ , such that:

- (i) the zero object belongs to  $\mathcal{S}_\varepsilon$  for any  $\varepsilon$ ;
- (ii) if  $0 \leq \tau < \varepsilon$ , then  $\mathcal{S}_\tau \subseteq \mathcal{S}_\varepsilon$ ;
- (iii) if  $0 \rightarrow F \rightarrow G \rightarrow H \rightarrow 0$  is an exact sequence in  $\mathcal{A}$ , then
  - if  $G$  is in  $\mathcal{S}_\varepsilon$ , then so are  $F$  and  $H$  (subobject property);
  - if  $F$  is in  $\mathcal{S}_\varepsilon$  and  $H$  is in  $\mathcal{S}_\tau$ , then  $G$  is in  $\mathcal{S}_{\varepsilon+\tau}$  (additivity property).

In [SCL<sup>+</sup>17], the authors provide several examples of noise systems. Among these examples, they discuss the property for a noise system of being closed under direct sum. Given a noise system  $\mathcal{S}$  on an abelian category  $\mathcal{A}$ ,  $\mathcal{S}$  is closed under direct sum if, for all  $\varepsilon \in \mathbb{R}$  and for any pair of objects  $M, N$  in  $\mathcal{S}_\varepsilon$ , then  $M \oplus N$  is also in  $\mathcal{S}_\varepsilon$ . Later works [GC17, RC18] describe a machinery called **contour** that generates noise systems that are in particular closed under direct sums. Since every noise system can be seen as an amplitude (see Proposition 2.22), the theory described in [GC17, RC18] can be used to generate amplitudes that on direct sums take as value the maximum of the evaluations on single elements. We call such amplitudes quasi-simple, and classify them in the applied topology frameworks in Section 4.1.

**Definition 2.19.** Let  $C$  be a cone in  $\mathbb{R}^n$ , i.e. the set of all linear combinations with non-negative coefficients of a finite collection of non-negative elements in  $\mathbb{R}^n$ , and  $\| - \|$  a norm on  $\mathbb{R}^n$ . Then the shift amplitude  $\varsigma$  on the functor category  $\text{Fun}(\mathbb{R}^n, \text{Vect})$  is defined as follows: for any  $M$  in  $\text{ob Fun}(\mathbb{R}^n, \text{Vect})$ ,

$$\varsigma(M) := \inf \{ \|c\| \mid c \in C \text{ and } M(t \leq t + c) = 0 \text{ for all } t \in \mathbb{R}^n \}.$$

The fact that this indeed is an amplitude follows from the properties of a noise system and the definition of standard noise in the direction of a cone in [SCL<sup>+</sup>17, Par. 6.2].

If the cone is generated by one element, then we get a simpler description of  $\varsigma$ , which also motivates the name. Let  $c \in C$  be the only vector in  $C$  with  $\|c\| = 1$ . Then, for any  $M$  in  $\text{ob Fun}(\mathbb{R}^n, \text{Vect})$ ,

$$\varsigma(M) := \inf \{ \varepsilon \in \mathbb{R} \mid M(t \leq t + \varepsilon c) = 0 \}.$$

In what follows, we often refer to the shift amplitude along a cone generated by one element as **shift along a vector**  $v$ . Note that the shift amplitude  $\varsigma$  does not need to be finite, in general. To see this it is enough to consider a module that persists forever in the direction of the cone.

We now study in detail the relation between amplitudes and noise systems. In particular, we will show that noise systems and amplitudes are the same by way of an adjunction.

**Definition 2.20.** Let  $(\mathcal{A}, \mathcal{S})$  and  $(\mathcal{A}', \mathcal{S}')$  be abelian categories with noise systems. An additive functors  $F: (\mathcal{A}, \mathcal{S}) \rightarrow (\mathcal{A}', \mathcal{S}')$  is **noise-bounding with constant**  $K \in (0, \infty)$  if there exists a non-negative real number  $K \geq 0$  such that, for any  $A$  in  $\text{ob } \mathcal{A}$ , if  $A \in \mathcal{S}_\varepsilon$ , then  $F(A) \in \mathcal{S}_{K\varepsilon}$ . The **category of noise systems**  $\text{Ns}$  is the category whose objects are pairs  $(\mathcal{A}, \mathcal{S})$  consisting of an abelian category  $\mathcal{A}$  and a noise system  $\mathcal{S}$ , and whose morphisms are noise-bounding functors<sup>2</sup>.

**Definition 2.21.** Let  $(\mathcal{A}, \alpha)$  and  $(\mathcal{A}', \alpha')$  be abelian categories with amplitudes. An additive functor  $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$  is **amplitude-bounding with constant**  $K \in (0, \infty)$  if  $\alpha'(FA) \leq K\alpha(A)$  for all objects  $A$  in  $\text{ob } \mathcal{A}$ . The **category of amplitudes**  $\text{Amp}$  is the category whose objects are pairs  $(\mathcal{A}, \alpha)$  of abelian categories and amplitudes, and whose morphisms are amplitude-bounding functors.

---

<sup>2</sup>As we are dealing with categories of large categories here, we do not expect  $\text{Ns}$  to be locally small.



For our applications to stability, we will be especially interested in the case in that  $\mathcal{A}$  is an abelian category with two amplitudes  $\alpha$  and  $\alpha'$ , and we consider identity functor  $\text{id}_{\mathcal{A}}: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}, \alpha')$ . The name is a slight abuse of notation, as this functor behaves as the identity only on part of the structure, but the symbol is more precise: the subscript contains only the category and not the amplitude. In this context, the following is worth noting:

- $\text{id}_{\mathcal{A}}$  does not need be a morphism in  $\text{Amp}$ .
- If  $\text{id}_{\mathcal{A}}$  is a morphism in  $\text{Amp}$ , it does not need be invertible.
- If  $\text{id}_{\mathcal{A}}$  is invertible, it is not necessarily the case that  $\alpha = \alpha'$ . Rather,  $\alpha'$  allows us to infer upper and lower bounds on  $\alpha$ , and vice versa.

We now establish the relationship between noise systems and amplitudes. Essentially, this relationship comes down to characterizing a norm on a vector space by the set of  $\varepsilon$ -balls around the origin.

**Proposition 2.22.** *Given an abelian category  $\mathcal{A}$  with a noise system  $\mathcal{S} = \{\mathcal{S}_{\varepsilon}\}_{\varepsilon \geq 0}$ , the function  $\alpha_{\mathcal{S}}$  on  $\text{ob } \mathcal{A}$  defined by*

$$\alpha_{\mathcal{S}}(A) := \inf \{ \varepsilon \mid A \in \mathcal{S}_{\varepsilon} \},$$

*with the convention that  $\inf \emptyset = \infty$ , is an amplitude on  $\mathcal{A}$ . This construction gives rise to a functor  $\text{NA}: \text{Ns} \rightarrow \text{Amp}$  defined by sending:*

- An object  $(\mathcal{A}, \mathcal{S})$  of  $\text{Ns}$  to the object  $(\mathcal{A}, \alpha_{\mathcal{S}})$  of  $\text{Amp}$ , and
- A noise-bounding functor  $F: (\mathcal{A}, \mathcal{S}) \rightarrow (\mathcal{A}', \mathcal{S}')$  to an amplitude-bounding functor  $F: (\mathcal{A}, \alpha_{\mathcal{S}}) \rightarrow (\mathcal{A}', \alpha_{\mathcal{S}'})$ .

*Proof.* All of the verifications are straightforward. □

Having constructed amplitudes from noise systems in Proposition 2.22, we are led to consider the converse:

**Definition 2.23.** Let  $\alpha$  be an amplitude on an abelian category  $\mathcal{A}$ . The **noise system associated to  $\alpha$** ,  $\mathcal{S}(\alpha)$ , is defined by

$$A \in \mathcal{S}(\alpha)_{\varepsilon} \iff \alpha(A) \leq \varepsilon,$$

for all  $\varepsilon \in \mathbb{R}_{\geq 0}$ . This construction induces a functor  $\text{AN}: \text{Amp} \rightarrow \text{Ns}$  as in Proposition 2.22.

**Proposition 2.24.** *The functors  $\text{AN}: \text{Amp} \leftrightarrow \text{Ns}: \text{NA}$  induce an equivalence of categories. The natural isomorphisms of the equivalence are all given by the identity on the underlying abelian categories, and are noise (amplitude) bounding for any  $K > 1$ .*

*Proof.* Both natural isomorphisms  $\text{NA} \circ \text{AN} \cong \text{id}_{\text{Amp}}$  and  $\text{AN} \circ \text{NA} = \text{id}_{\text{Ns}}$  are induced by the identity. The former is easily seen to fulfill  $\text{NA}(\text{AN}(\alpha)) = \alpha$  directly. For the latter, note first that  $\mathcal{S}_{\varepsilon} \subset \text{AN}(\text{NA}(\mathcal{S}))_{\varepsilon}$ . Hence, the identity in one direction is noise bounding with constant 1. Conversely, for any  $A \in \text{AN}(\text{NA}(\mathcal{S}))_{\varepsilon}$ , we have  $A \in \mathcal{S}_{K\varepsilon}$  for any  $K > 1$ . Hence, the converse direction is noise bounding with constant  $K > 1$ . □

We can now describe a class of amplitude-bounding functors that induces Lipschitz functions between metric spaces.

**Proposition 2.25.** *Let  $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$  be an additive amplitude-bounding functor with constant  $K$  satisfying the following two properties for any short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  in  $\mathcal{A}$ :*

1. If  $A$  has finite amplitude, then the sequence  $F(A) \rightarrow F(B) \rightarrow F(C) \rightarrow 0$  is  $\alpha'$ -exact;
2. If  $C$  has finite amplitude, then the sequence  $0 \rightarrow F(A) \rightarrow F(B) \rightarrow F(C)$  is  $\alpha'$ -exact.

Then the induced map  $F: (\text{ob } \mathcal{A}, d_\alpha) \rightarrow (\text{ob } \mathcal{A}', d_{\alpha'})$  is  $K$ -Lipschitz.

**Remark 2.26.** In particular, conditions 1 and 2 of Proposition 2.25 are satisfied if  $F$  is exact.

*Proof.* Without loss of generality (by passing to the appropriate quotient categories if necessary), we may assume that the sequences above are actually exact, and not just  $\alpha'$ -exact. Next, observe that, for the computation of the metric, only the morphisms that have finite cost are relevant. Thus, it suffices to show that for any  $\phi: A \rightarrow B$  with finite cost, we have  $c_{\alpha'}(F(\phi)) \leq K c_\alpha(\phi)$ .

Consider first the case where  $\phi$  is a monomorphism. In this case, we have a short exact sequence

$$0 \rightarrow A \xrightarrow{\phi} B \rightarrow \text{coker } \phi \rightarrow 0.$$

By assumption,  $F$  sends this short exact sequence into an exact sequence  $0 \rightarrow F(A) \xrightarrow{F(\phi)} F(B) \rightarrow F(\text{coker } \phi)$ . In particular,  $\text{coker}(F(\phi))$  is a subobject of  $F(\text{coker } \phi)$ . Hence,

$$c_{\alpha'}(F(\phi)) = \alpha'(\text{coker}(F(\phi))) \leq \alpha'(F(\text{coker } \phi)) \leq K \alpha(\text{coker } \phi) = K c_\alpha(\phi)$$

which finishes the proof in this case. By duality, the result also holds when  $\phi$  is an epimorphism.

In general, if  $\phi: A \rightarrow B$  has finite cost, we may consider its epi-mono factorization  $\phi = \psi \circ \eta$ , where  $\eta: A \rightarrow C$  is an epimorphism and  $\psi: C \rightarrow B$  is a monomorphism, and both  $\psi$  and  $\eta$  have finite cost. Our conditions imply that  $F(\phi) = F(\psi) \circ F(\eta)$  is also an epi-mono factorization. Then

$$c_{\alpha'}(F(\phi)) = c_{\alpha'}(F(\psi)) + c_{\alpha'}(F(\eta)) \leq K c_\alpha(\psi) + K c_\alpha(\eta) = K c_\alpha(\phi). \quad \square$$

As a partial converse, we can show that an additive functor that induces Lipschitz maps on metric spaces is necessarily amplitude-bounding:

**Proposition 2.27.** Let  $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$  be an additive functor such that the induced map  $F: (\text{ob } \mathcal{A}, d_\alpha) \rightarrow (\text{ob } \mathcal{A}', d_{\alpha'})$  is  $K$ -Lipschitz. Then  $F$  is amplitude-bounding with constant  $K$ .

*Proof.* By Lemma 2.13 we know that  $\alpha'(F(A)) = d_{\alpha'}(F(A), 0)$ . Since  $F$  is additive, we have that  $F0 \cong 0$ . Thus, we obtain  $\alpha'(F(A)) = d_{\alpha'}(F(A), 0) \leq K d_\alpha(A, 0) = \alpha(A)$ .  $\square$

## 2.3 Distances from noise systems vs. amplitudes

In this section, we relate distances obtained from noise systems and amplitudes, respectively. Noise systems and amplitudes induce metrics on the underlying abelian categories (see Definition 2.8), and we show that passing from a noise system to its corresponding amplitude does not affect the resulting metric. This all assembles into a commutative diagram of functors between various categories of noise systems, amplitudes, and pseudometric spaces.

We begin by defining the relevant categories.

**Definition 2.28.** Let **Pseudom** be the **category of pseudometric spaces** and Lipschitz maps, i.e.  $K$ -Lipschitz maps for some  $K \geq 0$ .

Here, we use the usual definition of “pseudometric space”, except that we do not require the underlying objects of our pseudometric spaces to be sets, and we allow pseudometrics to take values in  $[0, \infty]$ .

To ensure that the functors of interest between abelian categories induce Lipschitz maps between the corresponding pseudometric spaces, we restrict our attention to exact functors, as in the following definition.

**Definition 2.29.** We denote by  $\text{Ns}^{ex}$  be the subcategory of  $\text{Ns}$  where morphisms are exact noise-bounding functors, and by  $\text{Amp}^{ex}$  be the subcategory of  $\text{Amp}$  where morphisms are exact amplitude-bounding functors.

We are now ready to investigate the functoriality of the standard metric constructions.

**Proposition 2.30.** *There exists a functor  $\text{Ns}^{ex} \rightarrow \text{Pseudom}$*

- (i) *Sending a pair  $(\mathcal{A}, \mathcal{S})$  to the pseudometric space  $(\text{ob } \mathcal{A}, d_{\mathcal{S}})$ , where  $d_{\mathcal{S}}$  is the metric defined by*

$$d_{\mathcal{S}}(A, B) = \inf\{\varepsilon \mid A \text{ and } B \text{ are } \varepsilon\text{-close}\},$$

*where as usual  $\inf \emptyset = \infty$ , and the definition of  $\varepsilon$ -closeness is given in [SCL<sup>+</sup>17, Def. 8.4];*

- (ii) *Sending an exact noise-bounding functor  $F: (\mathcal{A}, \mathcal{S}) \rightarrow (\mathcal{A}', \mathcal{S}')$  to the map on objects  $F: (\text{ob } \mathcal{A}, d_{\mathcal{S}}) \rightarrow (\text{ob } \mathcal{A}', d_{\mathcal{S}'})$ .*

*Furthermore, if  $F: (\mathcal{A}, \mathcal{S}) \rightarrow (\mathcal{A}', \mathcal{S}')$  is exact and noise-bounding with constant  $K$ , then the map on objects is  $K$ -Lipschitz.*

*Proof.* The fact that  $d_{\mathcal{S}}$  is a pseudometric is proved, in a special case, in [SCL<sup>+</sup>17, Prop. 8.7] and their arguments generalize immediately to the case we are considering. Hence, the only nontrivial part of the proposition is the last claim, since this implies the functor  $\text{Ns}^{ex} \rightarrow \text{Pseudom}$  is well-defined on morphisms. So, suppose we are given an exact functor  $F: (\mathcal{A}, \mathcal{S}) \rightarrow (\mathcal{A}', \mathcal{S}')$  which is noise-bounding with constant  $K$ . Furthermore, suppose  $d_{\mathcal{S}}(A, B) < \varepsilon$  for some  $A, B \in \text{ob } \mathcal{A}$ . Then we can choose a span  $A \xleftarrow{f_0} C \xrightarrow{f_1} B$  with  $\ker f_0 \in \mathcal{S}_{\varepsilon_1}$ ,  $\ker f_1 \in \mathcal{S}_{\varepsilon_2}$ ,  $\text{coker } f_0 \in \mathcal{S}_{\varepsilon_3}$ , and  $\text{coker } f_1 \in \mathcal{S}_{\varepsilon_4}$ , for some  $\varepsilon_i$  satisfying  $\sum_{i=1}^4 \varepsilon_i < \varepsilon$ . Since  $F$  is noise bounding and exact, the span  $FA \xleftarrow{Ff_0} FC \xrightarrow{Ff_1} FB$  serves as a witness to the inequality  $d_{\mathcal{S}'}(FA, FB) < K\varepsilon$ . Because this holds for any  $\varepsilon > d_{\mathcal{S}}(A, B)$ , we see that  $d_{\mathcal{S}'}(FA, FB) \leq Kd_{\mathcal{S}}(A, B)$ , i.e.  $F$  is  $K$ -Lipschitz.  $\square$

**Corollary 2.31.** *There exists a functor  $\text{Amp}^{ex} \rightarrow \text{Pseudom}$*

- (i) *Sending an object  $(\mathcal{A}, \alpha)$  to the pseudometric space  $(\text{ob } \mathcal{A}, d_{\alpha})$ , where  $d_{\alpha}$  is as defined in Definition 2.8;*
- (ii) *Sending an exact amplitude-bounding functor  $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$  to the map on objects  $F: (\text{ob } \mathcal{A}, d_{\alpha}) \rightarrow (\text{ob } \mathcal{A}', d_{\alpha'})$ .*

*Furthermore, if  $F: (\mathcal{A}, \alpha) \rightarrow (\mathcal{A}', \alpha')$  is exact and amplitude bounding with constant  $K$ , then  $F: (\text{ob } \mathcal{A}, d_{\alpha}) \rightarrow (\text{ob } \mathcal{A}', d_{\alpha'})$  is  $K$ -Lipschitz.*

*Proof.* By Proposition 2.9, such a functor is uniquely given by the composition  $\text{Amp}^{ex} \xrightarrow{\text{AN}} \text{Ns}^{ex} \rightarrow \text{Pseudom}$   $\square$

Applying Proposition 2.24, we obtain from this:

**Corollary 2.32.** *The functors of Proposition 2.30, Corollary 2.31, Proposition 2.22, and Definition 2.23 assemble into on-the-nose commutative triangles*

$$\begin{array}{ccc} \text{Amp}^{ex} & \xrightarrow{\text{AN}} & \text{Ns}^{ex} \\ & \searrow & \swarrow \\ & \text{Pseudom} & \end{array}, \quad \begin{array}{ccc} \text{Ns}^{ex} & \xrightarrow{\text{NA}} & \text{Amp}^{ex} \\ & \searrow & \swarrow \\ & \text{Pseudom} & \end{array}.$$

Let us end this section, with a remark on another class of metrics for homological algebra setting which was defined in [BCZ21].

**Remark 2.33.** In [BCZ21] several notions of metrics for triangulated (persistence) categories were constructed. These rely on the additional data of specifying a class of objects  $\mathcal{F}$ . Instead of assigning weights to morphisms, these distances assign weights to triangles. In particular, the metrics  $\underline{d}_{\mathcal{F}}$  of [BCZ21, Rem. 2.0.5] may seem conceptually very close to amplitude distances. Roughly,  $\underline{d}_{\mathcal{F}}$  measure how many exact triangles with lower tip in  $\mathcal{F}$  it takes to transform one object into another, and weights this value by the weight of the triangles. At first glance, this makes amplitude distances appear to be a special case of the distances in [BCZ21]. Given an abelian category  $\mathcal{A}$  with amplitude  $\alpha$ , the obvious route of comparison would be to consider the (bounded) derived category of  $\mathcal{A}$ . One may then take  $\mathcal{F} = \{C \mid \sum_{i \in \mathbb{Z}} \alpha(H_i(C)) < \infty\}$  and assign to each exact triangle  $T$

$$\begin{array}{ccc} A & \xrightarrow{\quad} & B \\ & \nwarrow \quad \nearrow & \\ & C & \end{array}$$

the weight  $\omega(T) = \sum_{i \in \mathbb{Z}} \alpha(H_i(C))$ .

We note, however, that these types of weights do not fulfill the axioms of [BCZ21, Def. 2.0.1]. In particular, they fail normalization. Furthermore, in the definition of  $\underline{d}_{\mathcal{F}}$ , [BCZ21] require all triangles in a sequence to be oriented in the same way, and then symmetrize the metric by taking a maximum. This roughly corresponds to setting

$$\begin{aligned} d(A, B) = \max \Big\{ & \inf \{ \alpha(\ker f) + \alpha(\operatorname{coker} f) \mid f \in \operatorname{Hom}_{\mathcal{A}}(A, B) \}, \\ & \inf \{ \alpha(\ker f) + \alpha(\operatorname{coker} f) \mid f \in \operatorname{Hom}_{\mathcal{A}}(B, A) \} \Big\}, \end{aligned}$$

which differs from what we defined. For instance, in the case of 1-parameter persistence modules and the shift amplitude, this value will be infinite for any two interval modules that are not isomorphic.

Additionally, in [BCZ21, Sec. 4] the authors define metrics that take persistence into account by adding an additional persistence structure to the triangulated category and generalizing from exact triangles to triangles which are exact up to an  $\varepsilon$ -isomorphism. However, in these scenarios, they only weigh such almost exact triangles by the necessary amount of shifting. In particular, there seems to be no way to represent arbitrary amplitude distances in this framework, and therefore, although their approach seems close to ours, they are different in practice.

### 3 Persistence modules and finiteness conditions

In this section, we first cover some finiteness notions for multiparameter persistence modules and then we discuss some properties of multiparameter persistence modules on intervals.

**Notation 3.1.** We denote by  $\operatorname{Vect}$  the abelian category of finite-dimensional vector spaces over some fixed field  $\mathbb{F}$ . We use  $\mathcal{Q}$  to denote an arbitrary poset, and by abuse of notation, we denote by  $\mathcal{Q}$  also the canonical posetal category associated with  $\mathcal{Q}$ .

**Definition 3.2.** A functor  $\mathcal{Q} \rightarrow \operatorname{Vect}$  is a **persistence module**. When  $\mathcal{Q} = P^n$  for  $P$  a totally ordered set and  $n \geq 1$  a natural number, we also call such a functor an  **$n$ -parameter persistence module**. Furthermore, when  $n \geq 2$ , we alternatively call such a module a **multiparameter persistence module**.

Persistence modules are one of the main objects of study in TDA: by computing simplicial homology with coefficients in a field of an  $n$ -parameter family of simplicial complexes, one obtains for each homology degree an  $n$ -parameter persistence module.

**Remark 3.3.** We will constantly make use of the fact that persistence modules over  $\mathcal{Q} = \mathbb{Z}^n, \mathbb{R}^n$ , or more generally any monoidal poset, admit an equivalent formulation as algebraic objects. In Definition 3.2 we define persistence modules as functors  $\mathcal{Q} \rightarrow \text{Vect}$ . The category with objects such as functors, and morphisms given by natural transformations between them, can be easily seen to be isomorphic to the category with objects  $\mathcal{Q}$ -graded modules over the monoidal ring  $\mathbb{F}[\mathcal{Q}_{\geq 0}]$ , and morphisms given by homogeneous module homomorphisms. Using the latter perspective, we will frequently denote the application of  $M(u \leq v)$  to an element  $m \in M_u$  in the form

$$x^{u-v}m := M(u \leq v)m,$$

for  $M \in \text{Vect}^{\mathcal{Q}}$ ,  $u \leq v \in \mathcal{Q}$ .

### 3.1 Finiteness conditions

To balance computability and generality, we focus on persistence modules that admit a *finite encoding* [Mil20b]. Roughly, a finitely encoded module is a module that can be described using a finite poset (Definition 3.11). However, this level of generality is not entirely suitable to our setting since the category of finitely encoded persistence modules is not abelian [Mil20b, Ex. 4.25]. Instead, as a consequence of [Waa23, Thm 1.1], we restore abelianity by only allowing for encodings of certain types. Particular attention will be paid to the class of staircase modules (Example 3.8).




Figure 2: Examples illustrating the difference between staircase module and finitely encoded module. To every point is associated a copy of the field  $\mathbb{F}$  or 0 if it is in or outside a blue region, respectively. The thick line denotes a closed boundary, and the lack of it an open boundary. The structure morphisms are given by the identity on  $\mathbb{F}$  whenever possible.

We begin by introducing some general notation and language.

**Definition 3.4.** An **upset** of a poset  $\mathcal{Q}$  is a subset  $U \subseteq \mathcal{Q}$  such that for any  $u \in U$  and  $q \in \mathcal{Q}$  whenever  $u \leq q$  then  $q \in U$ . Similarly, a **downset** of a poset  $\mathcal{Q}$  is a subset  $D \subseteq \mathcal{Q}$  such that for any  $d \in D$  and  $q \in \mathcal{Q}$  whenever  $d \geq q$  then  $q \in D$ . An **interval** in  $\mathcal{Q}$  is the intersection of an upset and a downset of  $\mathmathcal{Q}$ .

**Notation 3.5.** For a poset  $\mathcal{Q}$ , and  $p, q \in \mathcal{Q}$ , we denote

1. The upset given by all elements greater or equal to  $q$ , by  $[q, \infty)$ .

2. The downset given by all elements lesser or equal to  $q$ , by  $(-\infty, p]$ .
3. The interval given by the intersection of  $[p, \infty)$  and  $(-\infty, q]$  by  $[p, q]$ .

For  $A \subset B$  subsets of a poset  $\mathcal{Q}$ , we write  $A \subseteq B$  to indicate that  $A$  is a downset in  $B$  and  $A \subseteq$  to indicate that  $A$  is an upset in  $B$ .

**Definition 3.6.** Recall that an **algebra** on a set  $S$ , is a subset of the power set of  $S$ , which contains  $S$  and is closed under unions and complements.

As already mentioned above, it is useful to have a general framework that specifies what class of persistence modules one allows for. This is taken care of by the following definition:

**Definition 3.7.** Let  $\mathcal{Q}$  be a poset. An **encoding structure**  $\mathfrak{X}$  on  $\mathcal{Q}$  is a subset of the powerset of  $\mathcal{Q}$ , such that

1.  $\mathfrak{X}$  is an algebra (i.e. closed under finite unions and complements, and contains  $\mathcal{Q}$ );
2. Every element of  $\mathfrak{X}$  is a finite union of intervals, which are themselves contained in  $\mathfrak{X}$ ;
3. If  $I \in \mathfrak{X}$  is an interval of  $\mathcal{Q}$ , then there exist upsets  $U, V \in \mathfrak{X}$ , such that  $I = U \cap V^c$ ;

**Example 3.8.** Let  $\mathbb{T}$  be a totally ordered set. A **cube** in  $\mathbb{T}^n$  is a subset of the form  $I_1 \times \cdots \times I_n$ , where the  $I_j$  are intervals in  $\mathbb{T}$ . A finite union of cubes that are unbounded above is called a **staircase**. The set

$$\mathfrak{C} := \{S \subset \mathbb{T}^n \mid S \text{ is a finite union of cubes in } \mathbb{T}^n\}$$

is an encoding structure, which we call the **staircase structure**  $\mathbb{T}^n$ , as the upsets of  $\mathfrak{C}$  are precisely the staircases. We denote by  $\underline{\mathfrak{C}}$  the subalgebra of  $\mathfrak{C}$  generated by the closed staircases. Again,  $\underline{\mathfrak{C}}$  is an encoding structure on  $\mathbb{T}$ .

Furthermore, observe that the intervals of  $\mathfrak{C}$  are all given by finite unions of cubes in  $\mathbb{T}^n$  and that the intervals of  $\underline{\mathfrak{C}}$  are given by finite unions of cubes  $I_1 \times \cdots \times I_n$  where the  $I_j$  are closed at the left and open at the right.

**Example 3.9.** The algebra generated by piecewise linear upsets, by semi-algebraic upsets or alternatively by any o-minimal structure in the sense of [vdD98], defines an encoding structure on  $\mathbb{R}^n$ . Similarly, one may consider the subalgebra of the latter examples generated by only the topologically closed upsets, called the closed below encoding structures.

**Definition 3.10.** An **encoding** of an object  $M \in \text{ob Vect}^{\mathcal{Q}}$  by a poset  $\mathcal{Q}'$  is a morphism  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  together with an object  $M' \in \text{ob Vect}^{\mathcal{Q}'}$  and an isomorphism  $\eta: M \xrightarrow{\sim} e^*M'$  between the object and the pullback of  $M'$  along  $e$ . Moreover, the encoding is **finite** if  $\mathcal{Q}$  is finite and  $M'$  is pointwise finite-dimensional. Let  $\mathfrak{X}$  be an encoding structure on  $\mathcal{Q}$  and  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$ ,  $\eta: M \xrightarrow{\sim} e^*M'$  a finite encoding of  $M$ . If  $e$  is such that for every upset  $U \subset \mathcal{Q}'$ ,  $e^{-1}(U)$  is an element of  $\mathfrak{X}$ , or equivalently that  $e^{-1}(p) \in \mathfrak{X}^{\text{int}}$ , for every  $p \in \mathcal{Q}'$ , then we say that the encoding is **of class  $\mathfrak{X}$**  or an  **$\mathfrak{X}$ -encoding**.

**Definition 3.11.** An object  $M$  in  $\text{ob Vect}^{\mathcal{Q}}$  is **finitely encoded** if it admits a finite encoding. Moreover,  $M$  is  **$\mathfrak{X}$ -encoded**, or **encoded by  $\mathfrak{X}$** , if it admits a  $\mathfrak{X}$ -encoding. If  $\mathcal{Q} = \mathbb{T}$  is a totally ordered poset, and  $\mathfrak{X} = \mathfrak{C}$  is the staircase structure, we say that  $M$  is a **staircase module**.

**Notation 3.12.** For a poset  $\mathcal{Q}$  and an encoding structure  $\mathfrak{X}$  on  $\mathcal{Q}$  we denote by  $\text{PerM}(\mathcal{Q})$  the full subcategory of  $\text{Vect}^{\mathcal{Q}}$  given by finitely encoded persistence modules, and by  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  the full subcategory given by the objects that are finitely  $\mathfrak{X}$ -encoded. Note that, when  $n = 1$ , the categories  $\text{PerM}(\mathbb{R}^n)$  and  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  coincide.

**Example 3.13.** Let  $I \subset \mathcal{Q}$  be an interval of an encoding structure  $\mathfrak{X}$  on  $\mathcal{Q}$ . We denote  $\mathbb{F}[I]$ , the **interval module on  $I$** , the module obtained as follows. Consider the finite poset with two elements 0 and 1 and the unique non-trivial relation given by  $0 \leq 1$ . We denote this by  $\{0 \leq 1\}$ , and similarly we denote by  $\{0 \leq 1\}^2$  the poset given by the cartesian product  $\{0, 1\}^2$  with partial order  $(a, b) \leq (a', b')$  if and only if  $a \leq a'$  and  $b \leq b'$ .

Consider the map of partially ordered sets  $e: \mathcal{Q} \rightarrow \{0 \leq 1\}^2$  induced by the two maps  $e_D: \mathcal{Q} \rightarrow \{0 \leq 1\}$  and  $e_U: \mathcal{Q} \rightarrow \{0 \leq 1\}$  defined as follows:

$$e_D(q) := \begin{cases} 0 & \text{if } q \in D \\ 1 & \text{otherwise} \end{cases} \quad e_U(q) := \begin{cases} 1 & \text{if } q \in U \\ 0 & \text{otherwise} \end{cases}.$$

Let  $M' \in \text{Vect}^{\{0 \leq 1\}^2}$  be the unique persistence module with  $M'_{(0,1)} = \mathbb{F}$  and 0 otherwise. Finally, define  $\mathbb{F}[I] = e^*M'$ . By construction,  $\mathbb{F}[I]$  is  $\mathfrak{X}$ -encoded. Explicitly,  $\mathbb{F}[I]$  is given by  $\mathbb{F}$  on  $I$ , 0 everywhere else, and with identity as structure morphism for all relations in  $I$ .

A useful property of finitely encoded modules is that there always exists a common finite encoding.

**Lemma 3.14.** *Let  $M_1, \dots, M_n$  be finitely encodable persistence modules over  $\mathcal{Q}$ . Then there exist an encoding map  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  which is part of an encoding for  $M_1, \dots, M_n$  simultaneously. Furthermore, if  $M_1, \dots, M_n$  are  $\mathfrak{X}$  encodable, for some encoding structure  $\mathfrak{X}$  on  $\mathcal{Q}$ , then  $e$  can also be taken of class  $\mathfrak{X}$ .*

*Proof.* We only show the case  $n = 2$ . The general case is entirely analogous. Let  $e_M: \mathcal{Q} \rightarrow \mathcal{P}_M$  and  $e_N: \mathcal{Q} \rightarrow \mathcal{P}_N$  be finite encodings of  $M$  and  $N$ , respectively, with modules  $M'$  and  $N'$ . Consider the induced map

$$e: \mathcal{Q} \longrightarrow \mathcal{P}_M \times \mathcal{P}_N \\ q \mapsto (e_M(q), e_N(q)).$$

Define two modules  $M''$  and  $N''$  as the pullback along the canonical projections  $\pi_M: \mathcal{P}_M \times \mathcal{P}_N \rightarrow \mathcal{P}_M$ ,  $\pi_N: \mathcal{P}_M \times \mathcal{P}_N \rightarrow \mathcal{P}_N$  of  $M'$  and  $N'$ , respectively. It follows that there exist two isomorphisms  $M \cong e^*M''$  and  $N \cong e^*N''$ , proving the claim. Furthermore, the fibers of  $e$  are, by construction, given by the pairwise intersections of the fibers of  $e_M$  and  $e_N$ . Hence, if the latter two are of class  $\mathfrak{X}$ , so is  $e$ .  $\square$

**Remark 3.15.** *Let  $I \subset \mathcal{Q}$  be an interval. Then, for any encoding map  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  of  $\mathbb{F}[I]$ ,  $I$  is saturated under  $e$ , i.e. we have  $e^{-1}(e(I)) = I$ . To see this, fix some module  $M$  over  $\mathcal{Q}'$  such that  $e^*(M) \cong \mathbb{F}[I]$ . Then, by construction,  $I = e^{-1}(S)$  where  $S$  is the support of  $M$ , and thus  $I$  is saturated. A convenient consequence of this property is that intersections of intervals are compatible under common encodings, in the sense that*

$$e\left(\bigcap I_j\right) = \bigcap e(I_j),$$

*whenever  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  is a common encoding map for a family of interval modules  $\mathbb{F}[I_j]$ .*

As an immediate consequence of Lemma 3.14 and the additivity of pullback functors, we obtain that the sum of two finitely encoded persistence modules (of a certain class  $\mathfrak{X}$ ) is again finitely encoded (and of class  $\mathfrak{X}$ ). In other words, we have:

**Corollary 3.16.** *For any encoding structure  $\mathfrak{X}$  on  $\mathcal{Q}$ , the category  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  is an additive subcategory of  $\text{Vect}^{\mathcal{Q}}$ .*

For our purposes, we are particularly interested in the cases where  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  is a full *abelian* subcategory of  $\text{Vect}^{\mathcal{Q}}$ . Even if, as we have already recalled, it is in general not true, under certain connectivity assumptions, which are fulfilled for the algebra of cubes and certain subalgebras of the PL and semialgebraic sets, this turns out to be the case. We provide now a brief introduction to the topic and refer to [Waa23] for more details and the full proofs.

**Definition 3.17.** We say that a poset  $\mathcal{Q}$  is  **$\leq$ -connected** if it is connected as a category. In other words, if, for every  $q, q' \in \mathcal{Q}$ , there exists a zigzag

$$q \leq q_1 \leq \cdots \leq q_k \leq q'$$

of inequalities between the two elements, where  $\leq$  denotes either  $\leq$  or  $\geq$ . The maximal  $\leq$ -connected subsets of  $\mathcal{Q}$  are called the  **$\leq$ -components** of  $\mathcal{Q}$ . We denote the set of these by  $\pi_0(\mathcal{Q})$ .

**Remark 3.18.** Every cube in  $C \subset \mathbb{R}^n$  is  $\leq$ -connected. To see this, note that a zigzag between any two elements  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  of the cube is given by the elements

$$x \leq (y_1, x_2, \dots, x_n) \leq (y_1, y_2, x_3, \dots, x_n) \leq \cdots \leq (y_1, \dots, y_{n-1}, x_n) \leq y.$$

As a cube is a product of subsets of  $\mathbb{R}$ , all of these elements are again in  $C$ .

More generally, every element of the closed below encoding structures described in Example 3.9 has finitely many  $\leq$ -connected components.

We now describe how the connected components of intervals of an encoding structure  $\mathfrak{X}$  influence the categorical properties of  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ .

**Definition 3.19.** An encoding structure  $\mathfrak{X}$  on a poset  $\mathcal{Q}$  is **connective** if every interval  $I$  in  $\mathfrak{X}$  has only finitely many  $\leq$ -components and these components are again elements of  $\mathfrak{X}$ .

**Remark 3.20.** The encoding structure of cubes  $\mathfrak{C}$  is connective. Indeed, let  $I \subset \mathbb{R}^n$  be an interval given by a finite union of cubes. First, note that we can write  $I$  as a finite disjoint union of cubes  $C_i$ . By Example 3.18, each of these cubes is  $\leq$ -connected. Hence, all of the  $\leq$ -components of  $I$  can be written as a finite disjoint union of  $C_i$ . As the components are pairwise disjoint and there are only finitely many  $C_i$ , there can only be finitely many  $\leq$ -components.

For connective encoding structures  $\mathfrak{X}$ , the category  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  is a full abelian subcategory of  $\text{Vect}^{\mathcal{Q}}$  (Theorem 3.22). However, the analogous argument as in Remark 3.20 does not work in the general piecewise linear or semialgebraic setting. In fact, the antidiagonal  $\{(x, y) \mid x = -y\}$  already is a PL interval with infinitely many  $\leq$ -connected components. For many applications, it can be of interest to allow for PL or semialgebraic encodings (compare [Mil20b]) while still retaining the abelianity of the underlying category.

**Example 3.21.** [Waa23, Rem. 3.18] For any of the encoding structures described in Example 3.9, the closed below encoding structure is connective.

For connective encoding structures, we have the following theorem:

**Theorem 3.22.** ([Waa23, Thm. 3.7]) Let  $\mathfrak{X}$  be a connective encoding structure on the poset  $\mathcal{Q}$ . Then the category  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  is a full abelian subcategory of  $\text{Vect}^{\mathcal{Q}}$ .

**Corollary 3.23.** The category  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  is a full abelian subcategory of  $\text{Vect}^{\mathbb{R}^n}$ .

The connective components of an interval are the key part to understanding morphisms between interval modules (see [BL15] for many examples). We will make extensive use of this understanding in Section 4. As a direct consequence of [Mil20b, Prop. 3.10], morphisms between interval modules can be explicitly classified as follows.



**Corollary 3.24.** Let  $I, I' \subset \mathcal{Q}$  be intervals,  $C \in \pi_0(I \cap I')$  and  $p, q \in C$ . Set  $S := \{J \in \pi_0(I \cap I') \mid J \subseteq I, J \subseteq I'\}$ . Then

1. for every morphism  $f: \mathbb{F}[I] \rightarrow \mathbb{F}[I']$ , we have  $f_p = f_q$  for every  $p, q \in C$ . Hence, we can denote by  $f_C$  the canonical element of  $\mathbb{F}$  specifying  $f_p: \mathbb{F} \rightarrow \mathbb{F}$ ;
2. The linear map  $\text{Hom}_{\text{PerM}(\mathcal{Q})}(\mathbb{F}[I], \mathbb{F}[I']) \rightarrow \mathbb{F}^S$  that associates  $f \mapsto (f_C)_{C \in S}$  is an isomorphism.

**Definition 3.25.** Let  $\mathfrak{X}$  be an encoding structure on a poset  $\mathcal{Q}$ . The **category of encoding intervals**  $\mathfrak{X}^{\text{int}}$  has as objects intervals  $I$  in  $\mathfrak{X}$  and morphisms given by

$$\text{Hom}_{\mathfrak{X}^{\text{int}}}(I, I') := \{J \subset I \cap I' \mid J \in \mathfrak{X}^{\text{int}}, J \subseteq I, J \subseteq I'\}$$

with  $\text{id}_I = I$  and composition given by  $J' \circ J := J \cap J'$ .

**Remark 3.26.** One can use Corollary 3.24 to make the construction in Example 3.13 functorial:

$$\begin{aligned} \mathbb{F}[-]: \mathfrak{X}^{\text{int}} &\rightarrow \text{PerM}_{\mathfrak{X}}(\mathcal{Q}) \\ J \subset I \cap I' &\mapsto g \end{aligned}$$

where  $g$  is the unique morphism in  $\text{Hom}_{\text{PerM}(\mathcal{Q})}(\mathbb{F}[I], \mathbb{F}[I'])$  specified by  $(f_C)_{C \in S} \in \mathbb{F}^S$ , with  $f_C = 1$  if  $C \subset J$  and zero otherwise.

**Remark 3.27.** Recall that a subquotient of  $A \in \mathcal{A}$ , in an abelian category  $\mathcal{A}$ , is an object  $B$  together with an epimorphism  $f: A \twoheadrightarrow C$  and a monomorphism  $B \hookrightarrow C$ . We say that  $B$  admits the structure of a subquotient of  $A$  if such a pair of arrows exists. By considering the pullback square

$$\begin{array}{ccc} C' & \hookrightarrow & A \\ \downarrow & & \downarrow \\ B & \hookrightarrow & C \end{array}$$

the cospan can be equivalently replaced by a span given by a monomorphism  $C' \hookrightarrow A$  and an epimorphism  $C' \twoheadrightarrow B$ .

Remark 3.26 induces the following dictionary between algebraic and poset language.

**Corollary 3.28.** Let  $\mathfrak{X}$  be a connective encoding structure,  $I \in \mathfrak{X}^{\text{int}}$ , and  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ . We denote by  $\text{sub}(M)$ ,  $\text{quot}(M)$  and  $\text{subquot}(M)$  the isomorphism classes of subobjects, quotient objects, and subquotient objects of  $M$ , respectively. The functor  $I \mapsto \mathbb{F}[I]$  described in Remark 3.26 together with Corollary 3.24 induces natural bijections

$$\begin{aligned} \{J \in \mathfrak{X}^{\text{int}} \mid J \subseteq I\} &\cong \text{sub}(\mathbb{F}[I]) \\ \{J \in \mathfrak{X}^{\text{int}} \mid J \subsetneq I\} &\cong \text{quot}(\mathbb{F}[I]) \\ \{J \in \mathfrak{X}^{\text{int}} \mid J \subset I\} &\cong \text{subquot}(\mathbb{F}[I]). \end{aligned}$$

Furthermore, unions can be translated into sums and extensions as follows. Let  $I_0, I_1 \in \mathfrak{X}^{\text{int}}$ . Then

1.  $I = I_0 \cup I_1$ , for  $I_0, I_1 \subseteq I$ , if and only if there exists an epimorphism

$$\mathbb{F}[I_0] \oplus \mathbb{F}[I_1] \twoheadrightarrow \mathbb{F}[I]$$

which is componentwise a monomorphism.

2.  $I = I_0 \cup I_1$ , for  $I_0, I_1 \subseteq I$ , if and only if there exists a monomorphism

$$\mathbb{F}[I] \hookrightarrow \mathbb{F}[I_0] \oplus \mathbb{F}[I_1]$$

which is componentwise an epimorphism.

3.  $I = I_0 \cup I_1$ , for  $I_0 \subseteq I$ ,  $I_1 \subseteq I$  with  $I_0 \cap I_1 = \emptyset$  if and only if there is a short exact sequence

$$0 \rightarrow \mathbb{F}[I_0] \rightarrow \mathbb{F}[I] \rightarrow \mathbb{F}[I_1] \rightarrow 0.$$

Finally, we describe how one may restrict and extend finitely encoded persistence modules from and to intervals.

**Proposition 3.29.** *Let  $I \in \mathfrak{X}$ . Denote by  $\mathfrak{X}|_I$  the algebra on  $I$  given by elements of the form  $S \cap I$ , for  $S \in \mathfrak{X}$ . Then the following holds:*

1.  $\mathfrak{X}|_I$  is an encoding structure on  $I$ .
2. If  $\mathfrak{X}$  is connective, then so is  $\mathfrak{X}|_I$ .
3. The extension by 0 functor

$$\begin{aligned} -|^\mathcal{Q}: \text{PerM}_{\mathfrak{X}|_I}(I) &\rightarrow \text{PerM}_{\mathfrak{X}}(\mathcal{Q}) \\ M &\mapsto \{p \mapsto \begin{cases} M_p & p \in I \\ 0 & p \notin I \end{cases}, \end{aligned}$$

with the obvious structure morphisms and definition of morphism, is well-defined and fully faithful. If  $\mathfrak{X}$  is connective, then  $-|^\mathcal{Q}$  defines an exact fully faithful embedding of abelian categories.

4. For  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ , the module  $(M|_I)|^\mathcal{Q}$  is naturally a subquotient of  $M$ .

*Proof.* 1. and 2. are immediate from the fact that every interval of  $I$  is also an interval of  $\mathcal{Q}$ .

To show the remaining two points, we restrict without loss of generality to the case where  $I$  is either an upset or a downset in  $\mathfrak{X}$ . Indeed, the general case is a composition of the latter two. By dualizing, it suffices to show the case where  $I$  is an upset. Denote by  $i$  the inclusion  $I \hookrightarrow \mathcal{Q}$ . Consider the left adjoint to the restriction functor  $i^*: \text{Vect}^\mathcal{Q} \rightarrow \text{Vect}^I$ , namely  $i_!: \text{Vect}^I \rightarrow \text{Vect}^\mathcal{Q}$ .  $i_!$  is given by left Kan-extension, i.e. we have  $(i_!M)_p = \varinjlim_{q \leq p, q \in I} M_q$ , with the induced structure morphisms. Since  $I$  is an upset, it follows that  $(i_!M) = M|^\mathcal{Q}$ , as defined above. The counit of adjunction  $i_!: i^*M \rightarrow M$  is given by the identity at  $p \in I$  and by  $0 \hookrightarrow M_p$  otherwise. In particular, it is given by a monomorphism and  $(M|_I)|^\mathcal{Q}$  is naturally a submodule of  $M$ , which shows point 4. The unit map  $N \rightarrow i^*i_!N$  is an isomorphism, indeed for every  $p \in I$  the indexing category  $\{q \leq p, q \in I\}$  has terminal object  $p$ . It follows that  $i_!$  is fully faithful. By construction, it is exact. It remains to see that  $i_!$  restricts to a functor of the subcategories of  $\mathfrak{X}|_I$  and  $\mathfrak{X}$  encodable submodules. Let  $M \in \text{PerM}_{\mathfrak{X}|_I}(I)$  with an encoding map  $e: I \rightarrow \mathcal{Q}'$ , and encoding module  $N \in \text{PerM}(\mathcal{Q}')$ . We extend  $e$  to a map of posets  $e_{-\infty}: \mathcal{Q} \rightarrow \mathcal{Q}'_{-\infty}$ , by sending every point not in  $I$  to  $-\infty$ . Since  $I$  is an upset, this construction defines morphism of posets, which has fibers in  $\mathfrak{X}$ . Denote by  $N_{-\infty}$  the extension of  $N$  by 0 at  $-\infty$ . Then

$$M|^\mathcal{Q} \cong (e^*N)|^\mathcal{Q} \cong e_{-\infty}^*(N_{-\infty}).$$

It follows that  $M|^\mathcal{Q}$  is  $\mathfrak{X}$  encoded, as was to be shown.  $\square$

## 4 Classifications of amplitudes in TDA

In this section, we show how every amplitude on staircase persistence modules is bounded above and below by two classes of functions, and we show that these two classes are in bijection with two classes of amplitudes. To do so, we study amplitudes with respect to their values on interval modules. We then obtain a classification result for the classes of additive amplitudes, which are amplitudes that are additive on short exact sequences, as well as what we call “quasi-simple” amplitudes, which are amplitudes that behave like a maximum on direct sums. Our classification of quasi-simple amplitudes provides a generalization of [GC17].

Throughout this section, we assume that  $\mathcal{Q}$  is a fixed poset and  $\mathfrak{X}$  is a connective encoding structure on  $\mathcal{Q}$ .

**Definition 4.1.** Given an amplitude  $\alpha$  on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ , the **function on intervals associated to  $\alpha$**  is given by

$$\begin{aligned} \mu_{\alpha}: \mathfrak{X}^{\text{int}} &\rightarrow [0, \infty] \\ I &\mapsto \alpha(\mathbb{F}[I]). \end{aligned}$$

We first note some properties of  $\mu_{\alpha}$  which may be immediately verified from the defining properties of an amplitude and Corollary 3.28.

**Lemma 4.2.** *Let  $\alpha$  be an amplitude on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ . Then, for all  $I, I_0, I_1 \in \mathfrak{X}^{\text{int}}$  the following hold:*

1.  $\mu_{\alpha}(\emptyset) = 0$  ;
2. If  $I_0 \subset I_1$  then  $\mu_{\alpha}(I_0) \leq \mu_{\alpha}(I_1)$  ;
3. If  $I = I_0 \cup I_1$ , where, for each  $i = 0, 1$ ,  $I_i \subseteq I$  or  $I_i \subsetneq I$ , then  $\mu_{\alpha}(I) \leq \mu_{\alpha}(I_0) + \mu_{\alpha}(I_1)$ .

The function on intervals provides lower and upper bounds for the amplitude it is associated with:

**Proposition 4.3.** *Let  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  and  $\alpha$  an amplitude on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ . Consider any  $\mathfrak{X}$ -encoding of  $M$ , given by  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  and  $M' \in \text{PerM}(\mathcal{Q}')$ . We have the following inequalities:*

$$\max_{p \leq q, M'_p \rightarrow M'_q \neq 0} \mu_{\alpha}(e^{-1}[p, q]) \stackrel{(3.1)}{\leq} \alpha(M) \stackrel{(3.2)}{\leq} \sum_{q \in \mathcal{Q}'} \dim(M'_q) \mu_{\alpha}(e^{-1}(q)). \quad (3)$$

*Proof.* We begin by proving inequality (3.1). Let  $p \leq q$ . If  $M'_p \rightarrow M'_q$  is nonzero, then there exists a morphism  $\phi: \mathbb{F}[p, \infty) \rightarrow M'$ , such that  $\phi(t)$  is nonzero for all  $t \in [p, q]$ . Hence, the image of  $\phi$  is an interval module  $\mathbb{F}[I]$ , for some  $I \subset [p, \infty)$ , such that  $[p, q] \subseteq I$ . It follows that  $\mathbb{F}[p \leq q]$  is a quotient of  $\mathbb{F}[I]$ , and we obtain the left-hand-side diagram in (4). By applying  $e^*$ , we obtain the right-hand-side diagram in (4).

$$\begin{array}{ccc} \mathbb{F}[I] & \hookrightarrow & M' \\ \downarrow & & \\ \mathbb{F}[p \leq q] & & \end{array} \quad \begin{array}{ccc} e^*(\mathbb{F}[I]) & \hookrightarrow & e^*(M') \\ \downarrow & & \\ e^*(\mathbb{F}[p \leq q]) & & \end{array} \quad (4)$$

Since  $e^*M' \cong M$ , inequality (3.1) follows by monotonicity of amplitudes.

We now prove inequality (3.2). We proceed by induction over the number  $n$  of elements  $q \in \mathcal{Q}'$  for which  $M'_q \neq 0$ . If  $n = 0$ , the inequality is trivially satisfied. If  $n = 1$ , and  $q \in \mathcal{Q}'$  is such that  $M'_q \neq 0$  is of dimension  $k$ , then

$$M \cong e^*(M') \cong e^*(\mathbb{F}[q])^k \cong \mathbb{F}[e^{-1}(q)]^k.$$

It follows from the subadditivity of amplitudes that  $\alpha(M) = \alpha(\mathbb{F}[e^{-1}(q)]^k) \leq k\alpha(\mathbb{F}[e^{-1}(q)])$ . We now assume the claim to be true for  $n$ , and show that then it holds for  $n + 1$ . Take  $q \in \max(\{p \in \mathcal{Q}' \mid M'_p \neq 0\})$ . Denote by  $M'' \in \text{PerM}(\mathcal{Q}')$  the persistence module with  $M''_p = M'_p$  if  $p = q$  and  $M''_p = 0$  otherwise. By maximality, there is a short exact sequence

$$0 \rightarrow M'' \rightarrow M' \rightarrow M'/M'' \rightarrow 0. \quad (5)$$

By construction,  $|\{p \in \mathcal{Q}' \mid (M'/M'')_p \neq 0\}| = n$ , since  $M'/M''$  and  $M$  differ only for their evaluation at  $q$ . In particular, the inductive assumption holds for  $e^*(M'/M'')$ . Applying  $e^*$  to (5), we obtain a short exact sequence

$$0 \rightarrow e^*(M'') \rightarrow M \rightarrow e^*(M'/M'') \rightarrow 0.$$

From it, by subadditivity of amplitudes and inductive assumption, we obtain the claim:

$$\begin{aligned} \alpha(M) &\leq \alpha(e^*(M'')) + \alpha(e^*(M'/M'')) \\ &= \sum_{q' \in \mathcal{Q}'} \dim(M''_{q'}) \mu_\alpha(e^{-1}(q')) + \sum_{q' \in \mathcal{Q}'} \dim((M'/M'')_{q'}) \mu_\alpha(e^{-1}(q')) \\ &= \sum_{q' \in \mathcal{Q}'} \dim(M''_{q'} + (M'/M'')_{q'}) \mu_\alpha(e^{-1}(q')) = \sum_{q' \in \mathcal{Q}'} \dim(M'_{q'}) \mu_\alpha(e^{-1}(q')). \quad \square \end{aligned}$$

Next we aim to characterize the amplitudes for which one of the two bounds in Equation 3 is attained (for all  $M$ ): in Subsection 4.1 we classify the class of amplitudes that realizes the lower bound (3.1), while in Subsection 4.2 we classify the one that realizes the upper bound (3.2).

#### 4.1 Quasi-simple amplitudes

The goal of this subsection is to show that the amplitudes for which inequality (3.1) is an equality, for any encoding, behave like a maximum on direct sums of persistence modules. We call this type of amplitudes *quasi-simple*. To reach this goal we first describe some characterizing properties of quasi-simple amplitudes (Proposition 4.4 and Definition 4.5). Then we define a class of functions over intervals that behave essentially like taking the diameter of an interval in the direction of the poset and are therefore called *directed diameters* (Definition 4.8). Finally, we prove that quasi-simple amplitudes are in bijection with directed diameters (Theorem 4.11).

**Proposition 4.4.** *Let  $\alpha$  be an amplitude on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ . Then the following conditions are equivalent:*

1. *For all  $M, N \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ ,  $\alpha(M \oplus N) = \max\{\alpha(M), \alpha(N)\}$ .*
2. *Let  $I_i \in \mathfrak{X}^{\text{int}}$ . For all  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ , the following holds: If there exists an epimorphism  $\bigoplus_{i=1, \dots, n} \mathbb{F}[I_i] \twoheadrightarrow M$  such that the resulting morphism  $\mathbb{F}[I_i] \rightarrow M$  is a monomorphism for every  $i$ , then*

$$\alpha(M) = \max\{\mu_\alpha(I_i) \mid i = 1, \dots, n\}.$$

3. Let  $I_i \in \mathfrak{X}^{\text{int}}$ . For all  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ , the following holds: If there exists a monomorphism  $M \hookrightarrow \bigoplus_{i=1, \dots, n} \mathbb{F}[I_i]$  such that the resulting morphism  $M \rightarrow \mathbb{F}[I_i]$  is an epimorphism for every  $i$ , then

$$\alpha(M) = \max\{\mu_{\alpha}(I_i) \mid i = 1, \dots, n\}.$$

4. For every  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  and every encoding of  $M$  with  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  and  $M' \in \text{PerM}(\mathcal{Q}')$ , we have:

$$\alpha(M) = \max_{p \leq q, M'_p \rightarrow M'_q \neq 0} \mu_{\alpha}(e^{-1}([p, q])).$$

*Proof.* The implications (1.  $\implies$  2.) and (1.  $\implies$  3.) follow directly from the amplitudes' axioms. We now show (2.  $\implies$  1.). For  $M, N \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  choose upsets  $U_i, V_j \in \mathfrak{X}$ , with  $i = 1, \dots, n$  and  $j = 1, \dots, m$ , and epimorphisms  $\oplus \phi_i: \bigoplus \mathbb{F}[U_i] \twoheadrightarrow M$  and  $\oplus \psi_j: \bigoplus \mathbb{F}[V_j] \twoheadrightarrow N$ , see [Mil20b, Theorem 6.12]. Let  $I_i \subseteq U_i$  and  $J_j \subseteq V_j$  be the subsets given by the points in  $\mathcal{Q}$  where  $\phi_i$  and  $\psi_j$  are nonzero, respectively. We then have induced natural isomorphisms  $\mathbb{F}[I_i] \cong \text{im}(\phi_i)$  and  $\mathbb{F}[J_j] \cong \text{im}(\psi_j)$ . In particular, we obtain induced epimorphisms  $\bigoplus \mathbb{F}[I_i] \twoheadrightarrow M$  and  $\bigoplus \mathbb{F}[J_j] \twoheadrightarrow N$ , which are componentwise monomorphisms, and consequently also an epimorphism

$$\bigoplus \mathbb{F}[I_i] \bigoplus \bigoplus \mathbb{F}[J_j] \twoheadrightarrow M \oplus N,$$

which is componentwise a monomorphism. Then, applying the assumption twice, we have

$$\begin{aligned} \alpha(M \oplus N) &= \max_{I \in \{I_1, \dots, I_n, J_1, \dots, J_m\}} \{\mu_{\alpha}(I)\} = \max \left\{ \max_{i=1, \dots, n} \{\mu_{\alpha}(I_i)\}, \max_{j=1, \dots, m} \{\mu_{\alpha}(J_j)\} \right\} \\ &= \max\{\alpha(M), \alpha(N)\}. \end{aligned}$$

The proof of (3.  $\implies$  1.) follows a similar argument by duality.

To see (2.  $\implies$  4.), choose some encoding  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$ ,  $M' \in \text{PerM}(\mathcal{Q}')$  of  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ . We have an epimorphism  $\phi: \bigoplus_{p \in \mathcal{Q}'} (\mathbb{F}[p, \infty))^{\dim(M'_p)} \twoheadrightarrow M'$ , which at some fixed  $p \in \mathcal{Q}'$  is an isomorphism when restricted to the  $p$ -component. Next, we note that by Corollary 3.28, the image of such a monomorphism  $\phi_p$  is isomorphic to an interval module  $\mathbb{F}[I_p]$  where  $I_p \subseteq \mathcal{Q}'$  is an interval having a unique minimal element  $p$ . Thus, by taking the direct sum of these interval modules, we have an epimorphism  $\bigoplus (\mathbb{F}[I_p]) \twoheadrightarrow M'$ , which is componentwise a monomorphism. Diagrammatically, we illustrate the argument below.

$$\begin{array}{ccccc} \bigoplus_{p \in \mathcal{Q}'} (\mathbb{F}[p, \infty))^{\dim(M'_p)} & \xrightarrow{\phi} & M' & \leftarrow & \bigoplus \text{im} \phi_p \cong \bigoplus (\mathbb{F}[I_p]) \\ & \nearrow \phi_p & \uparrow & & \nearrow \\ \mathbb{F}[p, \infty) & \xrightarrow{\quad} & \text{im} \phi_p & & \end{array}$$

Since  $e^*$  is exact, we obtain an epimorphism  $\bigoplus (\mathbb{F}[e^{-1}(I_p)]) \twoheadrightarrow M$ , which is also a monomorphism componentwise. It hence follows by the assumption that there exists an  $s$  such that

$$\alpha(M) = \mu_{\alpha}(e^{-1}(I_s)).$$

Let  $p$  be the minimal element of  $I_s$  and  $q_0, \dots, q_n$  the finite set of maximal elements of  $I_s$ . Then  $I_s = \bigcup_{i=0, \dots, n} [p, q_i]$ . Thus, we have the canonical monomorphism  $\mathbb{F}[I_s] \hookrightarrow \bigoplus_{i=1, \dots, n} \mathbb{F}[p, q_i]$ , which is componentwise an epimorphism. This induces a monomorphism  $\mathbb{F}[e^{-1}(I_0)] \hookrightarrow \bigoplus_{i=1, \dots, n} \mathbb{F}[e^{-1}[p, q_i]]$

which is componentwise an epimorphism. Since we already showed that (2.  $\iff$  3.), from the hypothesis we have

$$\alpha(M) = \mu_\alpha(e^{-1}(I_s)) = \max_{i=1,\dots,n} \mu_\alpha(e^{-1}[p, q_i]).$$

Since  $[p, q_i] \subseteq I_s$ , for each  $i = 1, \dots, n$  it follows that  $M'_p \rightarrow M'_{q_i}$  is nonzero. Hence we obtain

$$\alpha(M) = \max_{p \leq q, M'_p \rightarrow M'_q \neq 0} \mu_\alpha(e^{-1}([p, q])).$$

Finally, we show (4.  $\implies$  1.). Let  $M, N \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  and let  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  be a common encoding map of  $M$  and  $N$ , with encoding modules  $M'$  and  $N'$ . We thus have  $e^*(M' \oplus N') \cong M \oplus N$ . We note that  $(M' \oplus N')_p \rightarrow (M' \oplus N')_q$  is nonzero if and only if one of the two components is nonzero. Thus,

$$\left\{ p \leq q \mid (M' \oplus N')_p \rightarrow (M' \oplus N')_q \neq 0 \right\} = \left\{ p \leq q \mid M'_p \rightarrow M'_q \neq 0 \right\} \cup \left\{ p \leq q \mid N'_p \rightarrow N'_q \neq 0 \right\}$$

and the claim follows.  $\square$

**Definition 4.5.** An amplitude  $\alpha$  on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  is **quasi-simple** if it satisfies any of the equivalent properties of Proposition 4.4.

For example, the shift amplitude (Definition 2.19) is quasi-simple amplitude.

**Remark 4.6.** We note that quasi-simple amplitudes generalize the simple noise systems defined in [SCL<sup>+</sup>17]. Under the equivalence of noise systems and amplitudes (see Proposition 2.24), we have that noise systems can be thought of as corresponding to a specific kind of quasi-simple amplitudes. Namely, if one replaces finitely encoded by finitely presented persistence modules, simple noise systems correspond to quasi-simple amplitudes that additionally satisfy the following two properties: (1) for every persistence module  $M$  and every  $\varepsilon \geq 0$ , there exists a minimal subobject  $M'$  with the property that  $\alpha(M/M') \leq \varepsilon$ ; and (2) the subobject  $M'$  has rank less or equal to  $M$ .

As an immediate corollary of the characterization of quasi-simple amplitudes in Proposition 4.4 we obtain:

**Lemma 4.7.** If  $\alpha$  is quasi-simple then for all  $I_0, I_1, I \in \mathfrak{X}$ , where  $I_0$  and  $I_1$  are both upsets or both downsets of  $I$ , we have

$$\mu_\alpha(I) = \max\{\mu_\alpha(I_0), \mu_\alpha(I_1)\}.$$

**Definition 4.8.** A function  $\vec{\mu}: \mathfrak{X}^{\text{int}} \rightarrow [0, \infty]$  is a **directed diameter**, if for all  $I, I', I_0, I_1 \in \mathfrak{X}^{\text{int}}$  it fulfills the following properties:

1.  $\vec{\mu}(\emptyset) = 0$ ;
2. If  $I' \subset I$ , then  $\vec{\mu}(I') \leq \vec{\mu}(I)$ ;
3. If  $I = I_0 \cup I_1$ ,  $I_0 \subseteq I$ , and  $I_1 \subseteq I$ , then  $\vec{\mu}(I) \leq \vec{\mu}(I_0) + \vec{\mu}(I_1)$ ;
4. If  $I = I_0 \cup I_1$  and  $I_0, I_1 \subseteq I$  or  $I_0, I_1 \subseteq I$ , then  $\vec{\mu}(I) = \max\{\vec{\mu}(I_0), \vec{\mu}(I_1)\}$ .

**Example 4.9.** Let  $\mathfrak{C}$  be the staircase encoding structure and  $C$  a cone in  $\mathbb{R}^n$ , and  $\varsigma$  the shift amplitude (Definition 2.19). Then the function on interval associated to  $\varsigma$ ,  $\mu_\varsigma$ , is a directed diameter on  $\mathfrak{C}^{\text{int}}$ . Explicitly

$$I \mapsto \sup\{\|c\| \mid c \in C \text{ and } t + c \in I \text{ for some } t \in I\}$$

or equivalently

$$I \mapsto \inf\{||c|| \mid c \in C \text{ and } t + c \notin I \text{ for all } t \in I\}.$$

The value  $\mu_c(I)$  measures the diameter of  $I$  in the direction of the cone  $C$ .

Before proving that the above example is not just a coincidence, i.e. that the function on interval associated to any quasi-simple amplitude is a directed diameter, we first note the following symmetry property of directed diameters:

**Proposition 4.10.** *Let  $\vec{\mu}$  be a directed diameter on  $\mathfrak{X}^{\text{int}}$ . Then, for every  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ , we have the following equalities*

$$\begin{aligned} \sup\{\vec{\mu}(I) \mid I \in \mathfrak{X}^{\text{int}}, \mathbb{F}[I] \hookrightarrow M\} &= \sup\{\vec{\mu}(I) \mid I \in \mathfrak{X}^{\text{int}}, M \twoheadrightarrow \mathbb{F}[I]\} \\ &= \sup\{\vec{\mu}(I) \mid I \in \mathfrak{X}^{\text{int}}, \mathbb{F}[I] \text{ is a subquotient of } M\} \end{aligned}$$

*Proof.* It is straightforward to see that the first two expressions are lesser than or equal to the final one. By dualizing, it thus suffices to show that

$$\sup\{\vec{\mu}(I) \mid I \text{ is a subquotient of } M\} \leq \sup\{\vec{\mu}(I) \mid \mathbb{F}[I] \hookrightarrow M\}.$$

Suppose we are given a subquotient zigzag  $M \hookleftarrow N \twoheadrightarrow \mathbb{F}[I]$ . As in the proof of Proposition 4.4 we may choose interval modules  $I_1, \dots, I_n \in \mathfrak{X}^{\text{int}}$  together with an epimorphism

$$\bigoplus_{i=1, \dots, n} \mathbb{F}[I_i] \twoheadrightarrow N$$

which is componentwise a monomorphism. Consequently, we also have an epimorphism

$$\bigoplus_{i=1, \dots, n} \mathbb{F}[I_i] \twoheadrightarrow \mathbb{F}[I].$$

By replacing  $I_i$  with the interval corresponding to the image of  $\mathbb{F}[I_i] \twoheadrightarrow \mathbb{F}[I]$ , denoted  $\hat{I}_i$ , we obtain an epimorphism

$$\bigoplus_{i=1, \dots, n} \mathbb{F}[\hat{I}_i] \twoheadrightarrow \mathbb{F}[I],$$

which is componentwise a monomorphism. It follows by Corollary 3.28 that  $I = \bigcup_{i=1, \dots, n} \hat{I}_i$  and  $\hat{I}_i \subseteq I$ . Consequently, by assumption,  $\vec{\mu}(I) = \vec{\mu}(\hat{I}_k)$ , for some  $k$ . By construction, there is an epimorphism  $\mathbb{F}[I_k] \twoheadrightarrow \mathbb{F}[\hat{I}_k]$ . Hence, again by Corollary 3.28, it follows that  $\hat{I}_k \subseteq I_k$  and consequently that  $\vec{\mu}(I) = \vec{\mu}(\hat{I}_k) \leq \vec{\mu}(I_k)$ . The composition of monomorphisms

$$\mathbb{F}[I_k] \hookrightarrow N \hookrightarrow M,$$

show that  $\mathbb{F}[I_k]$  is a submodule of  $M$ , with  $\vec{\mu}(I_k) \geq \vec{\mu}(I)$ , as claimed.  $\square$

**Theorem 4.11.** *The assignment  $\alpha \mapsto \mu_\alpha$  induces a bijection between quasi-simple amplitudes on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  and directed diameters on  $\mathfrak{X}^{\text{int}}$ . The inverse map is given by  $\vec{\mu} \mapsto \alpha_{\vec{\mu}}$ , where  $\alpha_{\vec{\mu}}(M) := \sup\{\vec{\mu}(I) \mid I \in \mathfrak{X}^{\text{int}}, \mathbb{F}[I] \hookrightarrow M\}$ .*

*Proof.* By Lemma 4.7 the image of  $\alpha \mapsto \mu_\alpha$  is a directed diameter.

We now show that  $\alpha_{\vec{\mu}}$  is an amplitude. Note that by Proposition 4.10, we may freely replace the monomorphism in the definition of  $\alpha_{\vec{\mu}}$  by an epimorphism or subquotient. From the assumption  $\mu(\emptyset) = 0$ , we immediately have  $\alpha_{\vec{\mu}}(0) = 0$ . Monotonicity under monomorphisms

(epimorphisms) follows directly from the monomorphism (epimorphism) description of  $\alpha_\mu$ , respectively. Finally, to prove the subadditivity, consider the diagram

$$\begin{array}{ccccccc} & & \mathbb{F}[I] & & & & \\ & & \downarrow i & & & & \\ 0 & \longrightarrow & M' & \longrightarrow & M & \xrightarrow{g} & M'' \longrightarrow 0 \end{array},$$

with the lower row exact. We can complete the diagram into a commutative diagram with exact rows as follows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & i^{-1}M' & \longrightarrow & \mathbb{F}[I] & \longrightarrow & \text{img} \circ i \longrightarrow 0 \\ & & \downarrow & & \downarrow i & & \downarrow \\ 0 & \longrightarrow & M' & \longrightarrow & M & \xrightarrow{g} & M'' \longrightarrow 0 \end{array}.$$

Since  $i^{-1}M'$  and  $\text{img} \circ i$  are, respectively, a submodule and a quotient of  $\mathbb{F}[I]$ , by Corollary 3.28, they are of the form  $\mathbb{F}[I_0]$  with  $I_0 \subseteq I$  and  $\mathbb{F}[I_1]$  with  $I_1 \subseteq I$ , respectively, and  $I = I_0 \cup I_1$ . It follows that  $\vec{\mu}(I) \leq \sum \vec{\mu}(I_0) + \vec{\mu}(I_1)$ , which shows subadditivity.

Next, we show that  $\alpha_{\vec{\mu}}$  is quasi-simple. It is enough to prove that, for every  $M, N \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ ,

$$\alpha_{\vec{\mu}}(M_0 \oplus M_1) \geq \alpha_{\vec{\mu}}(M_0) + \alpha_{\vec{\mu}}(M_1).$$

Given a monomorphism  $\mathbb{F}[I] \hookrightarrow M_0 \oplus M_1$  with  $I \in \mathfrak{X}^{\text{int}}$ , we consider the image of the two components of this morphism. Since they are given by quotient objects of  $\mathbb{F}[I]$  by Corollary 3.28, they are of the form  $I_0, I_1 \subseteq I$ , for  $I_0, I_1 \in \mathfrak{X}^{\text{int}}$ . Furthermore, by construction, we have a monomorphism  $\mathbb{F}[I] \hookrightarrow \mathbb{F}[I_0] \oplus \mathbb{F}[I_1]$ . It follows, again by Corollary 3.28, that  $I = I_0 \cup I_1$  and therefore that  $\vec{\mu}(I) = \max\{\mu(I_0), \mu(I_1)\}$ . Without loss of generality, assume that the maximum is attained for  $I_0$ . Then, since by construction we have an embedding  $\mathbb{F}[I_0] \hookrightarrow M_0$ , it follows that  $\alpha_{\vec{\mu}}(M) \leq \alpha_{\vec{\mu}}(M_0)$ , i.e.  $\alpha_{\vec{\mu}}$  is quasi-simple.

Finally, let us verify that the two constructions are inverse to each other. By Proposition 4.4, in particular from the second and third characterization, it follows that if we start with an amplitude, pass to a directed diameter, and then pass back to amplitudes, we re-obtain the original amplitude. Conversely, by Corollary 3.28, every interval  $I$  is always maximal among intervals  $J$ , with the property that there exists a monomorphism  $\mathbb{F}[J] \hookrightarrow \mathbb{F}[I]$ , it also follows that starting with a directed diameter, passing to an amplitude, and then taking the function on intervals is the identity.  $\square$

As an immediate corollary of Proposition 4.10 and we Theorem 4.11 obtain:

**Corollary 4.12.** *Let  $\alpha$  be a quasi-simple amplitude on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ . Then the following equalities hold for any  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ :*

$$\begin{aligned} \sup\{\alpha(\mathbb{F}[I]) \mid I \in \mathfrak{X}^{\text{int}}, \mathbb{F}[I] \hookrightarrow M\} &= \sup\{\alpha(\mathbb{F}[I]) \mid I \in \mathfrak{X}^{\text{int}}, M \twoheadrightarrow \mathbb{F}[I]\} \\ &= \sup\{\alpha(\mathbb{F}[I]) \mid I \in \mathfrak{X}^{\text{int}}, \mathbb{F}[I] \text{ is a subquotient of } M\}. \end{aligned}$$

## 4.2 Additive amplitudes

Next, we classify the amplitudes for which inequality (3.2) is strict. We show that these are the *additive amplitudes* (see Definition 2.1), and can be described in terms of integrating, in some sense, the Hilbert function of a persistence module<sup>3</sup>. As a consequence, we have that additive

<sup>3</sup>Such a characterization may be understood as a Riesz-(Markov)-Representation type theorem for additive amplitudes (see [Rud87, Thm. 2.14] for a reference of the latter).



amplitudes take the same values on persistence modules that have the same Hilbert function but potentially different structure morphisms.

**Proposition 4.13.** *Let  $\alpha$  be an amplitude on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ . Then the following conditions are equivalent:*

1.  $\alpha$  is additive.
2. For  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  and  $M' \in \text{PerM}(\mathcal{Q}')$ ,  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  part of an  $\mathfrak{X}$ -encoding of  $M$ , we have

$$\alpha(M) = \sum_{q \in \mathcal{Q}'} \dim(M'_q) \mu_{\alpha}(e^{-1}(q)).$$

*Proof.* The implication (1.  $\implies$  2.) may be proved with an inductive argument, analogously to the proof of Proposition 4.3, inequality (3.2), if one replaces the appropriate inequalities by equalities.

For (2.  $\implies$  1.), we need to show that, given a short exact sequence  $S = 0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  in  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ ,  $\alpha(B) = \alpha(A) + \alpha(C)$ . By [Waa23, Prop. 3.6], we can take a sequence  $S' = 0 \rightarrow A' \rightarrow B' \rightarrow C' \rightarrow 0$  in  $\text{PerM}(\mathcal{Q}')$ , such that by pulling back  $S'$  along  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  we obtain  $S$ . Moreover,  $e$  can be chosen such that  $e^*$  is fully faithful, and thus  $S'$  is also exact. Therefore we have for each  $q \in \mathcal{Q}'$

$$\dim B'_q = \dim A'_q + \dim C'_q.$$

Consequently, we obtain:

$$\begin{aligned} \alpha(B) &= \sum_{q \in \mathcal{Q}'} \dim(B'_q) \mu_{\alpha}(e^{-1}(q)) = \sum_{q \in \mathcal{Q}'} \dim(A'_q) \mu_{\alpha}(e^{-1}(q)) + \sum_{q \in \mathcal{Q}'} \dim(C'_q) \mu_{\alpha}(e^{-1}(q)) \\ &= \alpha(A) + \alpha(C). \end{aligned} \quad \square$$

A direct consequence of the above characterization of additive amplitudes is that the associated function on intervals is completely determined by its behavior on the connected components of each interval.

**Lemma 4.14.** *Let  $\alpha$  be an additive amplitude on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ . Then, for all  $I, I_1, \dots, I_n \in \mathfrak{X}^{\text{int}}$  with  $I = \bigcup_{i=1}^n I_i$  and  $I_i \cap I_j = \emptyset$  for all  $i \neq j$ , it holds that*

$$\mu_{\alpha}(I) = \sum_{i=1}^n \mu_{\alpha}(I_i).$$

*Proof.* Choose a common encoding map  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  of class  $\mathfrak{X}$ , as well as encoding modules  $F$  and  $F_i$ , for  $\mathbb{F}[I]$  and  $\mathbb{F}[I_i]$  respectively. Then, using the definition of function on intervals, Lemma 4.2, Proposition 4.13, and Remark 3.15, we have

$$\mu_{\alpha}(I) = \alpha(\mathbb{F}[I]) = \sum_{q \in e(I)} \mu_{\alpha}(e^{-1}(q)) = \sum_{i=1}^n \sum_{q \in e(I_i)} \mu_{\alpha}(e^{-1}(q)) = \sum_{i=1}^n \alpha(\mathbb{F}[I_i]) = \sum_{i=1}^n \mu_{\alpha}(I_i). \quad \square$$

**Definition 4.15.** A **content** on an algebra  $\mathfrak{A}$  is a function  $\overset{\circ}{\mu}: \mathfrak{A} \rightarrow [0, \infty]$  such that:

1.  $\overset{\circ}{\mu}(\emptyset) = 0$ ;
2. For all  $S_1, S_2 \in \mathfrak{A}$  with  $S_1 \cap S_2 = \emptyset$ ,  $\overset{\circ}{\mu}(S_1 \cup S_2) = \overset{\circ}{\mu}(S_1) + \overset{\circ}{\mu}(S_2)$ .

By a **content on  $\mathfrak{X}^{\text{int}}$**  we mean a function  $\overset{\circ}{\mu}: \mathfrak{X} \rightarrow [0, \infty]$  which extends to a content on  $\mathfrak{X}$ .

**Remark 4.16.** Note that as  $\mathfrak{X}$  is generated by  $\mathfrak{X}^{\text{int}}$  under disjoint unions, a function  $\overset{\circ}{\mu}: \mathfrak{X} \rightarrow [0, \infty]$  extends to a content on  $\mathfrak{X}$  if, and only if for all  $I, I_1, \dots, I_n \in \mathfrak{X}^{\text{int}}$ , for some  $n \geq 0$ , with  $I = \bigcup_{i=1}^n I_i$  and  $I_i \cap I_j = \emptyset$  if  $i \neq j$ , it holds that  $\overset{\circ}{\mu}(I) = \sum_{i=1}^n \overset{\circ}{\mu}(I_i)$ . Furthermore, this extension is unique, so we really will not distinguish between contents on  $\mathfrak{X}^{\text{int}}$  and on  $\mathfrak{X}$ .

By the Lemma 4.14, we have that, for every additive amplitude  $\alpha$ , its function on intervals defines a content on  $\mathfrak{X}$ .

Recall that a function  $f: S \rightarrow \mathbb{N}$  is called measurable with respect to an algebra  $\mathfrak{A}$  on  $S$  if each of its fibers lies in  $\mathfrak{A}$ . Furthermore, measurable functions  $f: S \rightarrow \mathbb{N}$  can be integrated with respect to a content  $\overset{\circ}{\mu}$  on  $\mathcal{A}$  via the formula  $\int f d\overset{\circ}{\mu} = \sum_{n \in \mathbb{N}} f(n) \overset{\circ}{\mu}(f^{-1}(n))$ .

**Remark 4.17.** If  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ , then its Hilbert function

$$\begin{aligned} \dim_M: \mathcal{Q} &\rightarrow \mathbb{N} \\ q &\mapsto \dim M_q \end{aligned}$$

is  $\mathfrak{X}$ -measurable. Indeed, choose any encoding function  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  for  $M$ . Then every fiber of  $\dim_M$  is given by a finite union of fibers of  $e$ . As any such fiber is an interval of  $\mathfrak{X}$ , measurability follows.

We can then phrase the relationship between contents and additive amplitudes as follows.

**Theorem 4.18.** The assignment  $\alpha \mapsto \mu_\alpha$  induces a bijection between additive amplitudes of  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  and contents on  $\mathfrak{X}$ . The inverse map is given by

$$\overset{\circ}{\mu} \mapsto \{M \mapsto \int \dim_M d\overset{\circ}{\mu}\}.$$

*Proof.* By Lemma 4.14, we know that for any additive amplitude  $\alpha$  its function on intervals  $\mu_\alpha$  is a content on  $\mathfrak{X}^{\text{int}}$ .

Conversely, we denote by  $\overset{\circ}{\mu}$  any content on  $\mathfrak{X}$ . We have that the integral  $M \mapsto \int \dim_M d\overset{\circ}{\mu}$  is well defined by Remark 4.17. It is immediate from the additivity of Hilbert functions under short exact sequences that this expression does indeed define an amplitude on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ .

It remains to see that the two constructions are inverse to each other. Since  $\langle \mathfrak{X}^{\text{int}} \rangle = \mathfrak{X}$ , it is enough to show that given a content  $\overset{\circ}{\mu}$  on  $\mathfrak{X}$  we can recover the content on  $\mathfrak{X}^{\text{int}}$ . Indeed, for any  $I \in \mathfrak{X}^{\text{int}}$  we have

$$\int \dim_{\mathbb{F}[I]} d\overset{\circ}{\mu} = \mu(I).$$

On the other hand, if we begin with an amplitude  $\alpha$ , then for any  $M \in \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  and  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$  part of an  $\mathfrak{X}$ -encoding of  $M$  using Proposition 4.13, we have

$$\begin{aligned} \alpha(M) &= \sum_{q \in \mathcal{Q}'} \dim(M'_q) \mu_\alpha(e^{-1}(q)) = \sum_{n \in \mathbb{N}} n \sum_{q \in \mathcal{Q}', \dim(M'_q)=n} \mu_\alpha(e^{-1}(q)) \\ &= \sum_{n \in \mathbb{N}} n \mu_\alpha(\dim_M^{-1}(n)) = \int \dim_M d\mu_\alpha. \end{aligned} \quad \square$$

**Example 4.19.** Let  $\mathfrak{X}$  be a connective encoding structure on  $\mathbb{R}^n$ . Every upset in  $\mathbb{R}^n$  is Lebesgue measurable<sup>4</sup>. Let  $\mu$  be the restriction of the Lebesgue measure to  $\mathfrak{X}$ . Then we can generalize the construction in Theorem 4.18 to define the  $(p)$ -Hilbert amplitude of a module  $M$  in  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ :

$$\text{Hilb}^p(M) = \left( \int \dim_M d\mu \right)^{\frac{1}{p}}$$

<sup>4</sup>See [GG06, Thm. 4.4] for the case of  $[0, 1]^n$ . The proof there easily generalizes to  $\mathbb{R}^n$ .

If  $p = 1$ , we simply write  $\text{Hilb}(-)$ . If  $n = 1$  and  $p = 1$ , this coincides with  $\rho_1$  (Example 2.3).

**Remark 4.20.** *In a sense, Theorem 4.18 is a (monoid and) persistence modules version of the classical sheaf function correspondence of [KS90, Thm. 9.7.1]. Indeed, in the setting of constructible persistence modules in the sense of [Ber22], which correspond to constructible sheaves fulfilling certain microsupport conditions (see [BP21]) similar results were proven in [Ber22]. Hence, one could have also obtained Theorem 4.18 by computing the Grothendieck monoid of  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ . Recall that the Grothendieck monoid of an Abelian category  $\mathcal{A}$  (first defined in [BG16]) is a commutative monoid  $\mathcal{M}_{\mathcal{A}}$ , together with a map  $\text{ob } \mathcal{A} \rightarrow \mathcal{M}_{\mathcal{A}}$  which sends 0 to 0, is additive under short exact sequences, and is universal amongst such monoid valued maps.  $\mathcal{M}_{\mathcal{A}}$  may explicitly be constructed by taking the free monoid on  $\text{ob } \mathcal{A}$  (with 0 as neutral element), and quotienting out by  $A+C \sim B$ , whenever there is a short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ . It follows from the universal property of the Grothendieck monoid that additive amplitudes on  $\mathcal{A}$  correspond to monoid morphisms  $\mathcal{M}_{\mathcal{A}} \rightarrow [0, \infty]$ .*

*If  $\mathcal{A} = \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ , the Grothendieck monoid is the monoid of bounded maps  $f: \mathcal{Q} \rightarrow \mathbb{N}$  which are  $\mathfrak{X}$ -measurable. Denote the latter monoid by  $\mathcal{M}_{\mathfrak{X}}^n(\mathcal{Q}, \mathbb{N})$ . The bijection is induced by the Hilbert function. Indeed, by the same inductive argument as in the proof of Proposition 4.3, one shows that the map sending  $M$  to its Hilbert function is injective. Conversely, to see that bounded  $\mathfrak{X}$ -measurable  $\mathbb{N}$  valued function may be obtained as a Hilbert function, one can show that the constant regions of such a function may further be subdivided into intervals of  $\mathfrak{X}^{\text{int}}$ . Since indicator functions of intervals lie in the image of the Hilbert function construction, we obtain surjectivity. Finally, one may verify that the set of monoid morphisms*

$$\text{Hom}_{\text{Grpd}}(\mathcal{M}_{\mathfrak{X}}^n(\mathcal{Q}, \mathbb{N}), [0, \infty])$$

*is equivalently the set of contents on  $\mathfrak{X}$ .*

## 5 Discriminativity of amplitude metrics in TDA

In Section 2.1 we introduced span metrics associated to an amplitude on a category of persistence modules. As we have seen, the span metric is an extended pseudometric. In this section, we investigate under which conditions the span metric is an extended metric on the set of isomorphism classes of persistence modules. In other words, when are two modules with distance 0 isomorphic? To answer this question, we start with a first assumption: we only allow for amplitudes that do not send non-zero objects to 0.

**Definition 5.1.** An amplitude  $\alpha$  is **strict** if  $\alpha(A) = 0 \implies A \cong 0$  for all  $A \in \mathcal{A}$ .

**Example 5.2.** Many classical examples of amplitudes are not strict, even in the one-parameter setting. Consider for example the shift amplitude on  $\text{PerM}_{\mathcal{C}}(\mathbb{R})$  induced by the directed diameter sending an interval to its length. Its metric is equivalent to the interleaving distance of 1-parameter persistence modules. For interval modules of the shape  $\mathbb{F}[p, p]$ , the shift amplitude takes the value 0. More generally, it is 0 for all ephemeral persistence modules (see [CCBdS16, BP21]). We may restore strictness by quotienting out by the subcategory of ephemeral persistence modules. This is equivalent to instead restricting to the encoding structure  $\underline{\mathcal{G}}$ , i.e. only allowing for half-open intervals closed at the left and open at the right (compare with [BP21]).

The question of discriminativity of the span metric can be phrased as a convergence question for spans. Specifically, does the existence of spans  $A \leftarrow C_{\varepsilon} \rightarrow B$  with arbitrarily small cost allow one to conclude that there is a span  $A \leftarrow C_0 \rightarrow B$  with cost 0? In general, it is difficult to answer if one does not require some type of tameness assumption of the objects in use. In the

finitely generated case, for example, the diagonal shift amplitudes are known to induce a metric on isomorphism classes (see [Les15]). In the framework of constructible sheaves in [PSW21] it is shown that interleaving type distances on categories of appropriately constructible sheaves have this property.

In this section, we show such a result for general amplitude distances over persistence modules under certain topological boundary assumptions. We begin by setting up the framework:

**Definition 5.3.** We say that a subspace  $\mathbb{P} \subset \mathbb{R}^n$  is **locally open above** if for all  $x \in \mathbb{P}$  there exists neighborhood  $U \subset \mathbb{R}^n$ , such that  $(x + \mathbb{R}_{\geq 0}^n) \cap U \subset \mathbb{P}$ .

Clearly,  $\mathbb{R}^n$  is locally open above, and so is every interval given by the intersection of an upset and an open downset.

**Theorem 5.4.** Let  $\mathbb{P} \subset \mathbb{R}^n$  be locally open above. Let  $\mathfrak{X}$  be a connective encoding structure on  $\mathbb{P}$  fulfilling the following two properties:

1. Every upset  $U \in \mathfrak{X}$  is closed;
2. Every interval  $[p, q] \subset \mathbb{P}$ , with  $p, q \in (\mathbb{P}_{-\infty, \infty})$ , is an interval of  $\mathfrak{X}$ .

Let  $\alpha$  be an amplitude on  $\text{PerM}_{\mathfrak{X}}(\mathbb{P})$ . Then  $\alpha$  is strict if and only if the induced span metric on  $\text{PerM}_{\mathfrak{X}}(\mathbb{P})$  induces a metric on isomorphism classes of  $\text{PerM}_{\mathfrak{X}}(\mathbb{P})$ .

**Remark 5.5.** The assumptions on  $\mathfrak{X}$  in Theorem 5.4 are fulfilled, for example, for the encoding structure of closed staircases  $\underline{\mathfrak{C}}$  on  $\mathbb{R}^n$ , as well as for any of the closed below encoding structures in Example 3.9.

For the remainder of the section, we assume  $\mathbb{P}$  and  $\mathfrak{X}$  as in the statement of Theorem 5.4.

**Remark 5.6.** It is possible to drop the strictness assumption on  $\alpha$  and locally openness above on  $\mathbb{P}$  by passing to the quotient category in which objects with amplitude 0 are identified with 0, thus forcefully making  $\alpha$  strict. In many cases, the latter is of the shape  $\text{PerM}_{\mathfrak{X}'}(\mathcal{Q})$  for some smaller algebra and  $\mathcal{Q}$  a locally open above subset of  $\mathbb{R}^n$ .

For example, if  $\mathbb{P} = [0, 1]^n$ ,  $\mathfrak{X}$  is the encoding structure of cubes and we are considering the Hilbert amplitude with respect to the Lebesgue measure on  $\mathbb{R}^n$ , then said quotient category is equivalently given by persistence modules encoded by (open above) half open cubes on the half-open cube  $[0, 1)^n$  (compare [BP21]). Furthermore, one should note that the following proof of Theorem 5.4 also works in the case when  $\mathbb{P}$  is discrete, with minor alterations.

We begin with several remarks on the conditions of Theorem 5.4 before proving a series of intermediate results, which will result in the proof of Theorem 5.4. First, note that under the connectedness assumptions on  $\mathfrak{X}$ , whether two modules are isomorphic can generally be checked at finitely many points. This is made formal by the following proposition.

**Proposition 5.7.** Let  $\mathfrak{X}$  be a connective encoding structure over a poset  $\mathcal{Q}$ . Let  $M, N$  be objects in  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$ . Then there exists a finite subset of  $S \subset \mathcal{Q}$ , such that  $M$  and  $N$  are isomorphic if and only if  $i_S^*(M) \cong i_S^*(N)$ , where  $i_S: S \hookrightarrow \mathcal{Q}$  is the inclusion of posets.

*Proof.* Choose a common  $\mathfrak{X}$ -encoding  $e: \mathcal{Q} \rightarrow \mathcal{Q}'$ , with  $M', N' \in \text{Vect}^{\mathcal{Q}'}$ , of  $M$  and  $N$ . By [Waa23, Lem. 3.7], we may without loss of generality assume that  $e$  has  $\leq$ -connected fibers and that the relations in  $\mathcal{Q}'$  are generated by the images of relations under  $e$ . Denote by  $S'$  the set of elements of  $\mathcal{Q}$  given by choosing, for every generating relation in  $\mathcal{Q}'$ , a pair of elements  $s \leq s' \in \mathcal{Q}$  mapping to it. As  $\mathcal{Q}'$  is finite, so is  $S'$ . Furthermore, denote by  $S$  the set given by  $S'$  together with, for each pair  $s \leq s' \in S'$  which are contained in a common fiber of  $e$ , the elements of a

zigzag connecting  $s$  and  $s'$  in  $S'$ . Then  $e' := e|_S$  fulfills the requirements of [Waa23, Prop. 3.6] and hence  $e'^*$  is fully faithful. In particular, every isomorphism  $e'^*M' \cong i_S^*M \xrightarrow{\sim} i_S^*N \cong e'^*N$  extends to an isomorphism  $M' \xrightarrow{\sim} N'$ . The latter then pullback to an isomorphism of  $M$  and  $N$  under  $e$ .  $\square$

We call a subset  $S \subset \mathcal{Q}$  as in Proposition 5.7 a **comparison set**. The following lemma shows that in our framework the property of being a comparison set is stable under small perturbations.

**Lemma 5.8.** *Let  $\mathbb{P} \subset \mathbb{R}^n$  be locally open above,  $\mathfrak{X}$  an encoding structure given by closed sets on  $\mathbb{P}$ , and  $S \subset \mathbb{P}$  a comparison set for  $M, N$  in  $\text{ob PerM}_{\mathfrak{X}}(\mathbb{P})$ . Then, for  $\varepsilon > 0$  small enough,  $S + \varepsilon$  is again a comparison set.*

*Proof.* Using that  $S$  is finite, this is immediate from the fact that, as  $\mathfrak{X}$  is given by closed upsets and  $\mathbb{P}$  is locally open for each  $s \in S$ , the morphisms  $M_s \rightarrow M_{s+\varepsilon}$  is an isomorphism, for  $\varepsilon > 0$  small enough.  $\square$

Next, we need the following two technical tools.

**Lemma 5.9.** *Let  $\mathfrak{X}$  be any connective encoding structure on a poset  $\mathcal{Q}$ , such that every interval  $[p, \infty)$ ,  $(-\infty, p]$ , for  $p \in \mathcal{Q}_{-\infty, \infty}$ , is an interval of  $\mathfrak{X}$ . Let  $\alpha$  an amplitude on  $\text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  and  $M, N$  in  $\text{ob PerM}_{\mathfrak{X}}(\mathcal{Q})$ , and  $I \in \mathfrak{X}^{\text{int}}$  be such that  $M, N$  are constant (i.e. of the form  $\mathbb{F}^k[I]$ ) when restricted to  $I$ . Let  $p \in I$  and let*

$$\varepsilon := \min\{\alpha(\mathbb{F}[[p, \infty) \cap I]), \alpha(\mathbb{F}[(-\infty, p] \cap I])\}.$$

*Consider a span  $M \xleftarrow{f} L \xrightarrow{g} N$  with cost  $c_\alpha(f) + c_\alpha(g) < \varepsilon$ , and the induced cospan  $g': M \rightarrow \hat{L} \leftarrow N$ :  $f'$  obtained by taking the pushout of the former span. Then both  $f'$  and  $g'$  are given by isomorphisms at  $p$ .*

*Proof.* Let us first assume  $I = \mathcal{Q}$ . Then we can consider the following diagram

$$\begin{array}{ccccccc} & & \ker(f) & \longrightarrow & \ker(f') & & \\ & & \downarrow & & \downarrow & & \\ \ker(g) & \longrightarrow & L & \xrightarrow{g} & \mathbb{F}^{k'}[I] & \longrightarrow & \text{coker}(g) \\ & & \downarrow f & & \downarrow f' & & \\ \ker(g') & \longrightarrow & \mathbb{F}^k[I] & \xrightarrow{g'} & \hat{L} & \longrightarrow & \text{coker}(g') \\ & & \downarrow & & \downarrow & & \\ & & \text{coker}(f) & \longrightarrow & \text{coker}(f') & & \end{array}$$

with the inner square given by a pushout. As a consequence of the monotonicity of  $\alpha$ , we obtain that  $\alpha(\text{coker}(f)), \alpha(\text{coker}(g)), \alpha(\ker(f')), \alpha(\ker(g')) < \varepsilon$ . Note that, by the epimorphisms in the diagram, for any point that does not lie in the support of the latter four modules,  $f'$  and  $g'$  are isomorphisms. Furthermore, all kernels are submodules of a module  $\mathbb{F}^l[I]$ , which has all structure morphisms given by monomorphisms. In particular, any element at  $m \in \ker(f)_p$  induces a submodule of the shape  $\mathbb{F}[[p, \infty) \cap I]$ . Hence, by monotonicity of the amplitude, it follows that  $p$  is not contained in the support of the kernels. An essentially dual argument for the cokernels shows that  $p$  is not contained in the support of the cokernels either, and the claim is proved.

It remains to show that it is always possible to reduce to the case of  $\mathcal{Q} = I$ . To see this, consider the extension by 0 functor  $(-)|^{\mathcal{Q}}: \text{PerM}_{\mathfrak{X}|_I}(I) \hookrightarrow \text{PerM}_{\mathfrak{X}}(\mathcal{Q})$  of Proposition 3.29. Since

$(-)|^{\mathcal{Q}}$  is exact, we can transport  $\alpha$  to  $\text{PerM}_{\mathfrak{X}|_I}(I)$  along it. Denote the inclusion  $I \hookrightarrow \mathcal{Q}$  by  $i$ . Then  $i^*$  is an amplitude bounding functor (with constant 1). To see this, note that  $(i^*M)|^{\mathcal{Q}}$  is always a subquotient of  $M$ . Hence, by applying  $i^*$  to the setup of the theorem and using that it preserves all finite limits and colimits, we can always reduce to the case  $I = \mathcal{Q}$ .  $\square$

We now have everything necessary available to finish the proof of Theorem 5.4.

*Proof of Theorem 5.4.* The sufficient condition is immediate as  $\alpha(M) = d_\alpha(M, 0)$ .

For the converse, assume  $M, N$  in  $\text{ob PerM}_{\mathfrak{X}}(\mathbb{P})$  with  $d_\alpha(M, N) = 0$ . Choose a common  $\mathfrak{X}$ -encoding map  $e: \mathbb{P} \rightarrow \mathcal{Q}$  of  $M$  and  $N$  and a comparison set  $S \subset \mathbb{P}$ . Since  $\mathbb{P}$  is locally open above,  $S$  is finite, and  $\mathfrak{X}$  is given by closed sets, we can find  $\varepsilon > 0$  such that each set  $s + [0, 2\varepsilon)^n = [s, s + 2\varepsilon) \subset \mathbb{R}^n$  is contained in a fiber of  $e$ , for  $s \in S$ . Thus,  $M$  and  $N$  are both constant when restricted to any of these intervals. In particular, let  $\varepsilon$  be small enough, such that (the proof of) Lemma 5.8 applies, and hence  $S' := S + \vec{\varepsilon}$  is still a comparison set for  $M$  and  $N$ . Next, choose  $0 < \delta \leq \min\{\alpha(\mathbb{F}[s, s + \varepsilon), \mathbb{F}[s + \varepsilon, s + 2\varepsilon)) \mid s \in S\}$ . Such a  $\delta$  exists, as  $\alpha$  is strict and  $S$  is finite. As  $d_\alpha(M, N) = 0$ , there exists a span  $M \leftarrow L \rightarrow N$  of cost  $< \delta$ . Then, by Lemma 5.9, the pushout-induced cospan  $M \rightarrow \hat{L} \leftarrow N$  is given by isomorphisms on  $S'$ , i.e. we have an isomorphism  $i_{S'}^*M \cong i_{S'}^*N$ . Since  $S'$  is a comparison set, the claim is proved.  $\square$

## 6 Stability for amplitudes in TDA

The typical TDA workflow goes from data to invariants passing through parametrized (typically, filtered) simplicial complexes and persistent modules, and one can study the stability for each of these passages, or for some composition of them. Here we present a new possibility, naturally suggested by our approach. Namely, we consider functors from the category of staircase persistence modules to itself, possibly changing the source poset and the amplitudes. Then, studying the stability of these functors, we can infer new stability results from old ones.

Recall from the concept of amplitude-bounding functors (Definition 2.21). To obtain continuity and stability results, one can consider an amplitude-bounding functor

$$F: (\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n), \alpha) \rightarrow (\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n), \alpha')$$

with the additional property of being Lipschitz with respect to the distances induced by the amplitudes:

$$d_{\alpha'}(F(M), F(N)) \leq K d_\alpha(M, N)$$

for all staircase persistence modules  $M, N$  and for some  $K > 0$ . Then any stability result of the form

$$d_\alpha(H_k(D), H_k(E)) \leq K' d(D, E)$$

where  $D, E$  and  $d$  are appropriate input data sets and distances between these for the persistent homology pipeline, such as finite metric spaces together with the Gromow-Hausdorff distance, or Lipschitz functions on a topological space together with the sup-norm, would give us a stability result for  $\alpha'$ :

$$d_{\alpha'}(H_k(D), H_k(E)) \leq K d_\alpha(H_k(D), H_k(E)) \leq K K' d(D, E).$$

The proof of new stability results usually requires substantial work, and we therefore believe this to be a very powerful application of the change in perspective that our framework gives. In this section, we present some proofs of concept for 1-parameter persistence modules. In Section 6.1, we describe the  $k$ -longest-intervals amplitude (for  $k \in \mathbb{N}$ ) on  $\text{PerM}_{\mathfrak{C}}(\mathbb{R})$  and prove a bound relating the induced span metrics for varying  $k$ . We discuss the relationship between these span metrics and Wasserstein distances in Section 6.2. As a second example, we discuss in Section 6.3 how the theory of cost-processing functions allows us to relate the interleaving distance to the span metric coming from the Hilbert amplitude.

## 6.1 Stability results for the $k$ -longest-intervals amplitude

Often, in practice one only expects a certain number of bars to carry relevant data. However, this number may vary in different situations, and one needs not only to be able to choose how many bars to consider but also how to relate these different choices. We now discuss how to tackle these issues.

**Definition 6.1.** Let  $k \in \mathbb{N}$ . Given  $M$  in  $\text{ob PerM}_{\mathfrak{C}}(\mathbb{R})$ , write its interval-decomposition  $\oplus_{i=1}^n \mathbb{I}\langle b_i, b_i + \ell_i \rangle$ , where each interval can be closed or open at endpoints, and assume (after reordering as necessary) that  $\ell_1 \geq \ell_2 \geq \dots \geq \ell_n$ . Then the  **$k$ -longest-intervals amplitude** is defined as follows:

$$k\text{LI}(M) := \sum_{i=1}^k \ell_i.$$

**Remark 6.2.** The  $k$ -longest-intervals amplitude is discussed as a key example of a “tropical coordinate” in [Kal19]. Other tropical coordinates do not typically give amplitudes on  $\text{PerM}_{\mathfrak{C}}(\mathbb{R})$ .

**Proposition 6.3.** For any  $k \in \mathbb{N}$ , the function  $k\text{LI}$  is an amplitude on  $\text{PerM}_{\mathfrak{C}}(\mathbb{R})$ .

The proof relies on the matching theorem of [BL15, Theorem 4.2], which we recall for convenience. For a persistence module  $M$  in  $\text{PerM}_{\mathfrak{C}}(\mathbb{R})$ , we denote by  $\mathcal{D}_M = \{\langle b_i^M, d_i^M \rangle \mid i = 1, \dots, n\}$  the multiset of its barcode. Suppose we are given a short exact sequence of interval-decomposable persistence modules  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ . Then, by possibly adding empty intervals to the barcodes, we can find a common indexing of  $\mathcal{D}_A$ ,  $\mathcal{D}_B$ , and  $\mathcal{D}_C$  such that for all indices  $i$ , we have  $d_i^A = d_i^B$ ,  $b_i^B = b_i^C$ ,  $b_i^A \geq b_i^B$ , and  $d_i^B \geq d_i^C$ . Graphically, for every bar in  $B$ , we can truncate the bar on the left to obtain a bar in  $A$ , and truncate the original bar on the right to obtain a bar in  $C$ , and every bar in  $A$  and  $C$  arises in this way.

*Proof of Proposition 6.3.* Clearly  $k\text{LI}(0) = 0$ . Further, note that the monotonicity of  $k\text{LI}$  is an immediate consequence of the preceding discussion, as every bar in a submodule or quotient of a module  $B$  corresponds injectively to a (potentially) longer bar in  $B$ .

We are left with showing subadditivity. As ephemeral modules (i.e. modules that are supported on points) do not contribute to  $k\text{LI}$ , we may as well assume all intervals involved to be left closed right open, to simplify the proof.

Given a short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  of interval decomposable modules, let us first assume that  $B$  has at most  $k$  interval components. Then, by the discussion preceding this proof, so do  $A$  and  $C$ . Therefore, for  $M = A, B$ , or  $C$ , we have that  $k\text{LI}(M)$  is given by the sum of the lengths of all intervals. In particular  $k\text{LI}(B) = k\text{LI}(A) + k\text{LI}(C)$ .

For the general case, let  $B'$  denote the submodule of  $B$  given by the  $k$  longest bars (in case there are multiple such modules, choose any). We then have a commutative diagram with vertical monomorphisms and exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & A' & \longrightarrow & B' & \longrightarrow & C' \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C \longrightarrow 0 \end{array}$$

where  $A'$  is the intersection of  $B'$  with the kernel of  $B \rightarrow C$ , and  $C'$  is the image of  $B'$  in  $C$ . By the special case above, together with monotonicity of  $k\text{LI}$ , we obtain the claimed inequality:

$$k\text{LI}(B) = k\text{LI}(B') = k\text{LI}(A') + k\text{LI}(C') \leq k\text{LI}(A) + k\text{LI}(C). \quad \square$$

We now show that we can always compare  $k$ -longest-interval amplitudes for different  $k$  values.

**Lemma 6.4.** *Let  $k, \ell \in \mathbb{N}$  and let  $\text{id}_{\text{PerM}_{\mathfrak{C}}(\mathbb{R})} : (\text{PerM}_{\mathfrak{C}}(\mathbb{R}), k\text{LI}) \rightarrow (\text{PerM}_{\mathfrak{C}}(\mathbb{R}), \ell\text{LI})$  be given by the identity on objects.*

1. *If  $k \geq \ell$ , then  $\text{id}$  is amplitude-bounding with constant  $K = 1$ .*
2. *If  $1 \leq k \leq \ell$ , then  $\text{id}$  is amplitude-bounding with constant  $K = \ell/k$ .*

*Proof.* 1. This is immediate.

2. For a family of real numbers  $(x_i)_{1 \leq i \leq N}$ , we have the inequality

$$\max_{S \subset \{1, \dots, N\}, |S|=\ell} \left( \frac{\sum_{i \in S} x_i}{\ell} \right) \leq \max_{S' \subset \{1, \dots, N\}, |S'|=k} \left( \frac{\sum_{i \in S'} x_i}{k} \right),$$

Given  $M$  in  $\text{ob PerM}_{\mathfrak{C}}(\mathbb{R})$ , let  $(x_i)_{1 \leq i \leq N}$  be the lengths of the intervals in the barcode of  $M$  (in arbitrary order). Then

$$k\text{LI}(A) = \max_{S \subset \{1, \dots, N\}, |S|=k} \sum_{i \in S} x_i,$$

and similarly for  $\ell\text{LI}(A)$ . The stated inequality then implies the result.  $\square$

**Corollary 6.5.** *Let  $k, \ell \geq 1$  and let  $\text{id}_{\text{PerM}_{\mathfrak{C}}(\mathbb{R})} : (\text{PerM}_{\mathfrak{C}}(\mathbb{R}), k\text{LI}) \rightarrow (\text{PerM}_{\mathfrak{C}}(\mathbb{R}), \ell\text{LI})$  be given by the identity. Then*

$$\text{d}_{\ell\text{LI}}(M, N) \leq \max \left\{ 1, \frac{\ell}{k} \right\} \text{d}_{k\text{LI}}(M, N)$$

for any  $M, N \in \text{ob PerM}_{\mathfrak{C}}(\mathbb{R})$ .

*Proof.* This follows from Lemma 6.4 and Remark 2.26.  $\square$

## 6.2 Wasserstein distances versus $k$ -longest-intervals span metrics

The Wasserstein distance is a commonly used metric in TDA [CSEHM10, RT16, ST20]. Here, we present a simple result relating it to the span metric of the  $p$ -norm amplitude.

**Definition 6.6.** Let  $M, N$  be objects in  $\text{PerM}_{\mathfrak{C}}(\mathbb{R})$  and  $p \in [1, \infty)$ . The  **$p$ -Wasserstein distance** between  $M$  and  $N$  is given by

$$W_p(\text{Dgm}(M), \text{Dgm}(N)) = \inf_L \left( \sum_{x \in \text{Dgm}(M)} \|x - L(x)\|_p \right)^{\frac{1}{p}}$$

where the infimum is taken over all bijections  $L : \text{Dgm}(M) \rightarrow \text{Dgm}(N)$  (after possibly adding points on the diagonal to  $\text{Dgm}(M)$  or  $\text{Dgm}(N)$  so that such bijections exist).

As shown in [ST20], the Wasserstein distance can be reinterpreted as an  $f$ -span metric with  $f$  given by  $\| - \|_p$  and  $\alpha = \rho_p$ .

**Definition 6.7.** [ST20] Let  $M$  and  $N$  be two objects in  $\text{PerM}_{\mathfrak{C}}(\mathbb{R})$ , and  $p \in [1, \infty]$ . The algebraic  $p$ -Wasserstein distance between  $M$  and  $N$  is given by

$$W_p^{\text{alg}}(M, N) = \text{d}_{f_\alpha}(M, N) = \inf_{(C, \varphi, \psi)} \|(\rho_p(\ker \varphi), \rho_p(\text{coker } \varphi), \rho_p(\ker \psi), \rho_p(\text{coker } \psi))\|_p$$

where the infimum is taken over the spans  $M \xleftarrow{\varphi} C \xrightarrow{\psi} N$ , for  $C \in \text{ob PerM}_{\mathfrak{C}}(\mathbb{R})$  and  $f$  is the cost function  $\| - \|_p$ .



In [ST20, Thm 7.27 and 7.28], the authors prove that Definition 6.7 is indeed an alternative definition for the Wasserstein distance of persistence modules which are pointwise finite-dimensional with bounded  $p$ -energy. We use the result here, with a slightly stronger hypothesis: we require the persistence modules to be staircase modules. The proof of the result in this case is simpler, and now relies almost entirely on the theorem about the structure of persistence submodules and quotients of [BL15].

Using the equivalences of norms in  $\mathbb{R}^4$  and the fact that  $W_p^{alg} = d_{f_{\rho_p}}$ , where  $f$  is given by the  $p$ -norm on  $\mathbb{R}^4$ , we obtain the following equivalence:

**Corollary 6.8.** *For  $p \in [1, \infty]$  and  $M, N \in \text{PerM}_{\mathcal{C}}(\mathbb{R})$ , the following equivalence of metrics holds:*

$$W_p^{alg}(M, N) \leq d_{\rho_p}(M, N) \leq 4^{1-\frac{1}{p}} W_p^{alg}(M, N).$$

*In particular, for  $p = 1$ , we obtain  $W_1^{alg} = d_{\rho_1}$ .*

**Remark 6.9.** *In [BSS23, Th. 5.16], a version of the case  $p = 1$  of Corollary 6.8 is proven under slightly different assumptions.*

We can now discuss the relationship between Wasserstein distances and span metrics arising from the  $k$ -longest intervals amplitude. Corollary 6.5 allows for a different perspective on the distances  $d_{k\text{LI}}$ : they interpolate (in a monotone manner) between the interleaving and 1-Wasserstein distances. Intuitively, it is quite clear: the interleaving distance is equal to the bottleneck distance [CCSG<sup>+</sup>09, Les15], i.e. it is measuring the cost of the longest interval, while the 1-Wasserstein is adding the costs of all intervals. When  $k$  increases,  $d_{k\text{LI}}$  considers the cost of more and more intervals, thus “getting closer” to the 1-Wasserstein. Precisely, we have

$$d_I(M, N) \leq d_{k\text{LI}}(M, N) \xrightarrow{k \rightarrow \infty} W^1(M, N).$$

where the first inequality follows by [GC17, Prop. 12.2] and the limits is given by Corollary 6.8 since, for any  $M, N \in \text{PerM}_{\mathcal{C}}(\mathbb{R})$ ,  $d_{k\text{LI}}(M, N) = W^1(M, N)$  for  $k$  sufficiently large.

From the above observations, we have that the 1-Wasserstein distance bounds the span metric of the  $p$ -norms and of the  $k$ -longest interval amplitudes. Thus, we can use [ST20, Theorem 4.8] to obtain a stability result. We present the result in the notation of [ST20] and refer to [ST20] for the appropriate definitions.

**Corollary 6.10.** *(Stability of  $\rho_p$  and  $k\text{LI}$ ) Denote by  $T$  either  $\rho_p$  or  $k\text{LI}$ . Let  $K$  be any finite CW complex, and  $f, g: K \rightarrow \mathbb{R}$  monotone maps. Let  $H_k(f)$  and  $H_k(g)$  be the  $k$ -th persistent homology computed with respect to the corresponding sublevel set filtrations. Then, for all  $p \in [1, \infty]$  and  $k \in \mathbb{N}$ , we have:*

$$d_T(H_k(f), H_k(g)) \leq \|f - g\|_1.$$

In [Kal19, Section 5], there are stability results relating the absolute-value distance of  $k\text{LI}$  (i.e. the metric  $d'(A, B) := |k\text{LI}(A) - k\text{LI}(B)|$ ) to the  $p$ -Wasserstein distances and the bottleneck distance. This double stability property is also a consequence of the fact the absolute-value distance does not discriminate much. By contrast, span metrics are better at distinguishing persistence modules, at the price of not being as stable as the absolute-value distance.

### 6.3 The interleaving distance and the $\rho_1$ span metric

Proposition 2.17 allows for a simple proof of the following stability result between interleaving distance and the measure distances defined in [BSS23].

**Proposition 6.11.** *Let  $d_I$  be the interleaving distance on  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  with respect to the vector  $v = (1, \dots, 1) \in \mathbb{R}^n$ ,  $d_{\varsigma}$  the distance coming from the shift amplitude  $\varsigma$  with respect to  $v$  and the supremum norm, and  $d_{\rho_1}$  the path distance on  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  induced by the Hilbert amplitude. Then we have*

$$d_I(M, N) \leq d_{\varsigma}(M, N) \leq 4(d_{\rho_1}(M, N))^{\frac{1}{n}}$$

for all  $M, N \in \text{ob PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ .

*Proof.* The first inequality is [GC17, Prop. 12.2]. Hence, by Proposition 2.17, it suffices to show that  $\sqrt[n]{\cdot}$  defines a function  $g$  as in Proposition 2.17 taking  $F$  to be the identity functor,  $\alpha = \text{Hilb}(-)$ , and  $\alpha' = \varsigma$ . So, let  $M \in \text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  and, for any  $u \in \mathbb{R}^n$  and  $m \in M_u$ , consider the unique morphism  $\mathbb{F}[u, \infty) \rightarrow M$  such that  $1 \mapsto m$ . The image of this morphism is of the form  $\mathbb{F}[I]$  where  $I = [u, \infty) \cap D$  for  $D$  a downset in  $\mathbb{R}^n$ . By monotonicity of  $\text{Hilb}(-)$ , we have that  $\text{Hilb}(M) \geq \lambda(S)$ . Note that if  $x^{\varepsilon v} m \neq 0$  then  $S$  contains at least the cube of volume  $\varepsilon$  with lower left corner at  $u$ . The latter has volume  $\varepsilon^n$ . Hence,

$$\text{Hilb}(M) \geq \lambda(S) \geq \sup\{\varepsilon \mid x^{\varepsilon v} m \neq 0 \text{ for some } m \in M_w, w \in \mathbb{R}^n\}^n = \varsigma(M)^n.$$

Taking the  $n$ -th square root of the resulting inequality concludes the proof.  $\square$

**Remark 6.12.** *Proposition 6.11 holds more generally with  $\mathfrak{C}$  replaced by any connective encoding structure  $\mathfrak{X}$  that is closed under shifts and contains all upsets of the form  $[u, \infty)$ .*

## 7 Application: the $c_{\tau}$ -rank

In this section, we apply the techniques we presented to the study of the  $c_{\tau}$ -rank, showing that it is often not continuous.

This invariant is the continuous version of the  $c_{\mathfrak{p}}$ -rank described in [HOST19] and, intuitively, measures the number of ways in which a module goes to infinity in certain directions (given by  $\mathfrak{p}$ ). Since  $\mathfrak{p}$  can be assumed to be  $\langle x_1, \dots, x_t \rangle$  without loss of generality, we assume so throughout this section. The  $c_{\mathfrak{p}}$ -rank fails to be an amplitude (Example 7.2). However, it is given by the composition of two functors and an amplitude (Definition 7.3 and Proposition 7.4). This formulation allows us to extend the definition to the continuous setting (Definition 7.11) and then prove its (in)stability (Proposition 7.16) by studying the behavior of the two functors in the composition (Propositions 7.14 and 7.15).

In [HOST19, Definition 4.19], the  $c_{\mathfrak{p}}$ -rank of a finitely generated  $N^n$ -graded module  $M$  is defined to be the rank of the 0-th local cohomology of  $M$  as a module over  $\mathbb{F}[x_{t+1}, \dots, x_n]$ . Here, we give a more general definition of  $c_{\mathfrak{p}}$ -rank using homogeneous prime ideals of  $\mathbb{F}[x_1, \dots, x_n]$ , which, by [HOST19, Lemma 4.6], are generated by a subset of the variables.

**Definition 7.1.** Consider the abelian category  $\mathcal{A}$  of finitely generated  $\mathbb{N}^n$ -graded modules  $M$  over the polynomial ring  $\mathbb{F}[x_1, \dots, x_n]$  with coefficients in a field  $\mathbb{F}$ . Let  $\mathfrak{p}$  be any homogeneous prime ideal in  $\mathbb{F}[x_1, \dots, x_n]$ , and without loss of generality assume for simplicity that  $\mathfrak{p} = \langle x_1, \dots, x_t \rangle$ . The  **$c_{\mathfrak{p}}$ -rank**  $c_{\mathfrak{p}}(M)$  of a module  $M \in \text{ob } \mathcal{A}$  is the rank of  $H_{\mathfrak{p}}^0(M)$ , the 0-th local cohomology of  $M$ , as a module over  $\mathbb{F}[x_{t+1}, \dots, x_n]$ :

$$c_{\mathfrak{p}}(M) := \text{rank}(H_{\mathfrak{p}}^0(M)) = \text{rank}(\{m \in M \mid \mathfrak{p}^k \cdot m = 0 \text{ for all } k \text{ large enough}\}).$$

The above definition is well-posed as, for any finitely generated module  $M$ , the module  $H_{\mathfrak{p}}^0(M)$  is finitely generated over  $\mathbb{F}[x_{t+1}, \dots, x_n]$  since the statement in Lemma 4.18 in [HOST19] holds more generally for any homogeneous prime ideal  $\mathfrak{p}$ .

The  $c_{\mathfrak{p}}$ -rank of a module can be interpreted as a measure of the size of its submodule generated by elements living along the coordinate directions orthogonal to those spanned by  $\mathfrak{p}$ , see [HOST19, Proposition 4.20]. This interpretation is valid for general homogeneous primes  $\mathfrak{p}$ , and not just associated primes; if  $\mathfrak{p}$  is a homogeneous prime that is not an associated prime, then the  $c_{\mathfrak{p}}$ -rank is zero.

**Example 7.2** (The  $c_{\mathfrak{p}}$ -rank is not an amplitude). Consider the  $\mathbb{N}^2$ -graded modules  $B = \mathbb{F}[x_1, x_2]$  and  $C = \mathbb{F}[x_1, x_2]/\langle x_1 \rangle$  over  $\mathbb{F}[x_1, x_2]$ , with the canonical projection  $B \rightarrow C$ . Then  $c_{\langle x_2 \rangle}(B) = 0$  but  $c_{\langle x_2 \rangle}(C) = 1$ , and the monotonicity axiom fails.

We now discuss two constructions — the continuous 0-th local cohomology and the quotient restriction — which allow us to frame the  $c_{\mathfrak{p}}$ -rank in the amplitude setting and extend it to staircase modules.

Given a persistence module  $M$  over  $\mathbb{Z}^n$ , we can express its 0-th local cohomology with respect to  $\mathfrak{p}$  as

$$H_{\mathfrak{p}}^0(M) = \ker(M \rightarrow \prod_{0 \leq i \leq t} M_{x_i}).$$

Here,  $M_{x_i}$  is the persistence module obtained by localizing  $x_i$ . An immediate application of [Mil20a, Thm. 5.2] shows that this construction indeed preserves staircase encodings and hence descends to a functor  $\text{PerM}_{\mathfrak{C}}(\mathbb{Z}^n)$

$$H_{\mathfrak{p}}^0: \text{PerM}_{\mathfrak{C}}(\mathbb{Z}^n) \rightarrow \text{PerM}_{\mathfrak{C}}(\mathbb{Z}^n).$$

Next, we consider the functor

$$(-)/\mathfrak{J}_{\mathfrak{p}}: \text{PerM}_{\mathfrak{C}}(\mathbb{Z}^n) \rightarrow \text{PerM}_{\mathfrak{C}}(\mathbb{Z}^{n-t})$$

obtained by taking the quotient by the ideal  $\mathfrak{J}_{\mathfrak{p}} = \langle x_{t+1} - 1, \dots, x_n - 1 \rangle$ . Finally, we equip  $\mathbb{Z}^{n-t}$  with the counting measure. We are now ready to give the following definition:

**Definition 7.3.** The  **$c_{\mathfrak{p}}$ -rank** for staircase  $\mathbb{Z}^n$ -graded modules is defined as the following composition:

$$\text{ob}(\text{PerM}_{\mathfrak{C}}(\mathbb{Z}^n)) \xrightarrow{H_{\mathfrak{p}}^0} \text{ob}(\text{PerM}_{\mathfrak{C}}(\mathbb{Z}^n)) \xrightarrow{(-)/\mathfrak{J}_{\mathfrak{p}}} \text{ob}(\text{PerM}_{\mathfrak{C}}(\mathbb{Z}^r)) \xrightarrow{\text{Hilb}(-)} [0, \infty].$$

In other words, the  $c_{\mathfrak{p}}$ -rank is given by the 1-Hilbert amplitude of  $(H_{\mathfrak{p}}^0(M))/\mathfrak{J}_{\mathfrak{p}}$ , for  $M$  in  $\text{PerM}_{\mathfrak{C}}(\mathbb{Z}^n)$ .

This formulation agrees with the usual one in our case of interest.

**Proposition 7.4.** Let  $M$  be a staircase persistence module over  $\mathbb{Z}^n$  and  $\mathfrak{p}$  a homogeneous prime ideal of  $\mathbb{F}[x_1, \dots, x_n]$ . Then Definitions 7.1 and 7.3 of  $c_{\mathfrak{p}}$ -rank of  $M$  coincide.

*Proof.* Let  $M$  be a module in  $\text{PerM}_{\mathfrak{C}}(\mathbb{Z}^n)$ . To simplify notation, we set  $N = H_{\mathfrak{p}}^0(M)$ , considered as  $\mathbb{F}[x_{t+1}, \dots, x_n]$ -module. The goal is to show that the rank of  $N$  is equivalently given by  $\text{Hilb}(N/\mathfrak{J}_{\mathfrak{p}}N)$ . Note that  $\text{Hilb}(N/\mathfrak{J}_{\mathfrak{p}}N) = \dim_{\mathbb{F}}(N/\mathfrak{J}_{\mathfrak{p}}N)$ , where by  $\dim_{\mathbb{F}}$  we mean the dimension of the  $\mathbb{Z}^{n-t}$ -graded objects considered as an  $\mathbb{F}$ -vector space. As in Definition 7.3,  $N/\mathfrak{J}_{\mathfrak{p}}N$  can be computed by first localizing at the elements  $x_{t+1}, \dots, x_n$  and then restricting the resulting module to  $\mathbb{Z}^t \times \{0\} \subset \mathbb{Z}^n$  (compare [Mil20a, Rem. 2.23] for the analogous real cases). The localized module  $N_{x_{t+1}, \dots, x_n}$  is a  $\mathbb{Z}^n$  graded module over  $\mathbb{F}[x_1, \dots, x_n, x_{t+1}^{-1}, \dots, x_n^{-1}]$ . The quantity  $\dim_{\mathbb{F}}(N/\mathfrak{J}_{\mathfrak{p}}N)$  is independent from the multiplicative structure in direction of  $x_1, \dots, x_t$ . Hence, we may instead consider  $N_{x_{t+1}, \dots, x_n}$  as a  $\mathbb{Z}^{n-t}$ -graded  $R = \mathbb{F}[x_{t+1}^{\pm 1}, \dots, x_n^{\pm 1}]$ -module, denoted by  $N'$ , given by

$$N'(v) = \bigoplus_{u \in \mathbb{Z}^t} N(u, v).$$

$N'$  is a free  $R$ -module of rank  $\dim_{\mathbb{F}}(N'(0))$ . Since the restriction to  $\mathbb{Z}^t \times \{0\} \subset \mathbb{Z}^n$  corresponds to evaluating at 0 in this setting, we have:

$$\text{Hilb}(N/\mathfrak{J}_{\mathbf{p}}N) = \dim_{\mathbb{F}}(N/\mathfrak{J}_{\mathbf{p}}N) = \dim_{\mathbb{F}}(N'(0)) = \text{rank}_R(N').$$

To conclude we need to show that  $\text{rank}_R(N') = \text{rank}(N)$ . By definition,  $\text{rank}(N)$  is given by the  $\text{Quot}(\mathbb{F}[x_{t+1}, \dots, x_n])$ -dimension of  $N \otimes_{\mathbb{F}[x_{t+1}, \dots, x_n]} \text{Quot}(\mathbb{F}[x_{t+1}, \dots, x_n])$ . By standard commutative algebra, the latter admits the alternative description:

$$\begin{aligned} N \otimes_{\mathbb{F}[x_{t+1}, \dots, x_n]} \text{Quot}(\mathbb{F}[x_{t+1}, \dots, x_n]) &\cong (N \otimes_{\mathbb{F}[x_{t+1}, \dots, x_n]} R) \otimes_R \text{Quot}(\mathbb{F}[x_{t+1}, \dots, x_n]) \\ &\cong N' \otimes_R \text{Quot}(\mathbb{F}[x_{t+1}, \dots, x_n]). \end{aligned}$$

As  $N'$  is, by construction, a free module of rank  $\dim_{\mathbb{F}}(N'(0))$ , the dimension of the right-hand side expression is  $\text{rank}_R(N')$ . This finishes the proof.  $\square$

We can now construct the  $c_p$ -rank for  $\mathbb{R}^n$ -modules by defining the continuous version of the 0-th local cohomology. The role of homogeneous prime ideals for  $\mathbb{Z}^n$ -modules is taken by the faces of the positive cone for  $\mathbb{R}^n$ -modules, with a corresponding notion of localizations. For more detail about these two concepts, see [Mil20a, Sec. 2, in particular Def. 2.17].

**Definition 7.5.** A **face** of the positive cone  $\mathbb{R}^n$  is a set of the form

$$\tau = \left\{ \sum_{i \in I} \alpha_i e_i \mid \alpha_i \geq 0 \right\},$$

where  $I$  is a subset of  $\{1, \dots, n\}$  and  $e_i$  denotes the  $i$ -th canonical unit vector in  $\mathbb{R}^n$ . The **dimension of  $\tau$**  is  $|I|$ . Moreover,  $\tau^c$  denotes the face **orthogonal** to  $\tau$ , i.e. it is the cone spanned by the canonical basis vectors in  $\mathbb{R}^n$  which are not in  $\tau$ .

**Definition 7.6.** Given a face  $\tau$ , the **localization of a module  $M$  in  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  along  $\tau$** , denoted by  $M_{\tau}$  is defined as the tensor product

$$M_{\tau} := M \otimes_{\mathbb{F}[\mathbb{R}_{\geq 0}^n]} \mathbb{F}[\mathbb{R}_{\geq 0}^n + \mathbb{Z}\tau],$$

viewing  $M$  as a  $\mathbb{R}^n$ -graded  $\mathbb{F}[\mathbb{R}_{\geq 0}^n]$ -module.

Note that localization in this sense sends the indicator modules of cubes into indicator modules of cubes and is exact [Mil20a, Lem. 2.24]. Hence, as an application of [Mil20b, Thm. 6.12], localisation preserves staircase encodings. Thus, the following persistent module  $H_{\tau}^0$  is again a staircase module, since  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  is abelian and the product is finite:

**Definition 7.7.** Let  $\tau$  be a face of the positive cone in  $\mathbb{R}^n$  and  $M$  in  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ . We define the following persistence module

$$H_{\tau}^0(M) := \ker(M \rightarrow \prod_{0 \neq \rho \subset \tau} M_{\rho}),$$

where the product is taken over the nontrivial faces of  $\tau$ . The functor  $H_{\tau}^0: \text{PerM}_{\mathfrak{C}}(\mathbb{R}^n) \rightarrow \text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  induced by this construction is the **0-th local cohomology with respect to  $\tau$** .

**Remark 7.8.** Note that, in Definition 7.7, we might have equivalently just taken the product over the 1-dimensional faces of  $\tau$ . Indeed, let  $\{v_0, \dots, v_k\} \subset \tau_i$  be any set of generators for a face  $\tau_i$  of  $\tau$ . Then  $m \in M$  being mapped to 0 in  $M_{\mathbb{R}_{\geq 0} v_j}$ , for all  $0 \leq j \leq k$ , already implies that  $m$  is mapped to 0 in  $M_{\tau}$ .

The role of modding out the ideal  $\mathfrak{J}_{\mathbf{p}}$  in Definition 7.3 is taken by the quotient restriction functor from [Mil20a, Def. 2.22].

**Definition 7.9.** Let  $\tau$  be a face of the positive cone in  $\mathbb{R}^n$ . Given a persistence module  $M$  in  $\text{Vect}^{\mathbb{R}^n}$ , the **quotient restriction**  $M/\tau$  of  $M$  along  $\tau$  is defined as the  $\mathbb{R}^n/\mathbb{R}\tau$ -module given by

$$(M/\tau)_{v+\mathbb{R}\tau} := \varinjlim_{u \in v+\tau} M_u$$

with the induced structure maps.

In other words,  $M$  can be thought as the quotient by equivalence relation  $m \sim x^v m$ , for  $m \in M$  and  $v \in \tau$ , of an  $\mathbb{R}^n$ -graded module.

**Lemma 7.10.** *The quotient restriction preserves staircase encodings. Moreover, the functor*

$$(-)/\tau: \text{PerM}_{\mathfrak{C}}(\mathbb{R}^n) \rightarrow \text{PerM}_{\mathfrak{C}}(\mathbb{R}^{n-t})$$

*induced by this construction is exact.*

In the above statement, we have identified  $\mathbb{R}^n/\mathbb{R}\tau$  with  $\mathbb{R}^{\tau^c}$ , which in turn is naturally identified with  $\mathbb{R}^{n-t}$  via the canonical basis of  $\mathbb{R}^n$ .

*Proof.* Quotient restriction is equivalently defined as first localizing at  $\tau$  and then restricting along  $\mathbb{R}\tau^c$  (see [Mil20b, Rem. 2.23]). As both the localization and the restriction to a subposet are exact, the functor  $\text{Vect}^{\mathbb{R}^n} \rightarrow \text{Vect}^{\mathbb{R}^n}$  defined in this fashion is exact. Furthermore, it sends interval modules of cubes to interval modules of cubes. More specifically, if  $\mathbb{F}[C]$  is the interval module of an upset-cube  $C$ , then  $\mathbb{F}[C]_{\tau} = \mathbb{F}[C + \mathbb{R}\tau]$ . The restriction to  $\mathbb{R}\tau$  acts as  $\mathbb{F}[C + \mathbb{R}\tau] \mapsto \mathbb{F}[(C + \mathbb{R}\tau) \cap \mathbb{R}\tau^c]$ . The latter is again the indicator module of a cube. By an application of [Mil20b, Thm. 6.12] and using exactness, we thus obtain that quotient restriction indeed sends staircase modules into staircase modules.  $\square$

We now have all it is needed to define the continuous  $c_{\tau}$ -rank.

**Definition 7.11.** Let  $\tau$  be a face of the positive cone in  $\mathbb{R}^n$  of dimension  $t$ . The  **$c_{\tau}$ -rank**  $c_{\tau}: \text{PerM}_{\mathfrak{C}}(\mathbb{R}^n) \rightarrow [0, \infty]$  is defined as the composition

$$\text{ob}(\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)) \xrightarrow{H_{\tau}^0} \text{ob}(\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)) \xrightarrow{(-)/\tau^c} \text{ob}(\text{PerM}_{\mathfrak{C}}(\mathbb{R}^{n-t})) \xrightarrow{\text{Hilb}(-)} [0, \infty].$$

This definition is compatible with the discrete one in the following sense:

**Proposition 7.12.** *Let  $f_n: \mathbb{R}^n \rightarrow \mathbb{Z}^n$  the map of posets given by taking the floor map (i.e. the map that sends a real to the largest integer smaller than it) in each component. Let  $\mathbf{p}$  be a homogeneous prime of  $\mathbb{Z}[x_1, \dots, x_n]$  with  $t$  generators and  $\tau$  the corresponding face of the positive cone of dimension  $t$ . Then, for each  $M$  in  $\text{PerM}_{\mathfrak{C}}(\mathbb{Z}^n)$ , we have*

$$c_{\mathbf{p}}(M) = c_{\tau}(f_n^* M).$$

*Proof.* A straightforward verification shows that  $H_{\tau}^0 \circ f_n^* \cong f_n^* \circ H_{\mathbf{p}}^0$  and  $(-)/\tau^c \circ f_n^* \cong f_{n-t}^* \circ (-)/\mathfrak{J}_{\mathbf{p}}$ . Next, note that the map  $f_{n-t}$  is measure-preserving with respect to the Lebesgue measure and the counting measure, as the inverse of a point is given by a cube of volume one, with parts of its boundary removed. In particular,  $\text{Hilb}(f_{n-t}^* N) = \text{Hilb}(N)$  for all  $N$  in  $\text{PerM}_{\mathfrak{C}}(\mathbb{Z}^{n-t})$ . Piecing these equalities together gives the result.  $\square$

## 7.1 Stability of the $c_\tau$ -rank

Having defined the  $c_\tau$ -rank, we are now ready to study its stability. The idea is to prove the stability of the two functors composing the  $c_\tau$ -rank, and then to use the techniques of Section 6 to study the stability of the 1-Hilbert amplitude. The concatenation of these results gives us (in)stability results for the  $c_\tau$ -rank. We begin studying the quotient restriction by proving an ancillary lemma.

In what follows, without loss of generality, we assume  $\tau$  is spanned by the first  $t$  canonical vectors, and we identify  $\mathbb{R}^n/\mathbb{R}\tau$  with  $\mathbb{R}^{n-t}$ .

**Lemma 7.13.** *Let  $\tau$  be a face of the positive cone in  $\mathbb{R}^n$  of dimension  $t$ , with  $0 < t \leq n-1$ ,  $1 \leq q < \infty$ , and  $M \in \text{ob PerM}_\mathbb{C}(\mathbb{R}^n)$ . Consider  $\mathbb{R}^n$  and  $\mathbb{R}^{n-t}$  equipped with the Lebesgue measure. If  $\text{Hilb}^p(M) < \infty$ , then  $\text{Hilb}^q(M/\tau) = 0$  for all  $1 \leq p < \infty$ .*

*Proof.* Fix  $q \in \mathbb{N}_{\geq 1}$ . For  $M \in \text{ob PerM}_\mathbb{C}(\mathbb{R}^n)$ , assume that  $\text{Hilb}^q(M/\tau) > 0$ . Choose a staircase encoding of  $M/\tau$  given by a map  $e: \mathbb{R}^{n-t} \rightarrow Q$ . By assumption, at least one of the fibers  $F$  of this map has non-zero measure and the dimension of  $M/\tau$  over this fiber is nonzero. As  $\lambda(F) > 0$ , there exists a closed cube  $Q \subset F$ , with positive volume  $\lambda(Q) > 0$  and minimal element  $u \in Q$ , which is fully contained in  $F$ . As all structure morphisms of  $M/\tau$  which are going from points in  $F$  to points in  $F$  are isomorphisms, we have that  $x^v \hat{m} \neq 0$ , for  $v \in -u + Q$  and  $\hat{m} \in M/\tau(u)$ . For any representative  $m \in M$  mapping to  $\hat{m}$ , it thus holds that  $x^v m \neq 0$  for  $v \in \tau \times (-u + Q)$ . In particular,  $\dim(M(u')) \geq 1$  holds for all  $u'$  in a cube of shape  $(\tau + v_0) \times Q$ , which has Lebesgue measure  $\infty \cdot \lambda(Q) = \infty$ . Consequently we have  $\text{Hilb}^p(M) = \infty$ , for all  $1 \leq p < \infty$ .  $\square$

**Proposition 7.14.** *Let  $\tau$  be a face of the positive cone of  $\mathbb{R}^n$  of dimension  $t$ , where  $1 \leq t \leq n-1$ . Then we have that, for all  $M, N$  in  $\text{PerM}_\mathbb{C}(\mathbb{R}^n)$ ,*

$$d_{\alpha'}(M/\tau, N/\tau) \leq K d_\alpha(M, N) \quad \text{for}$$

1. all  $K > 0$  and  $\alpha = \alpha' = \text{Hilb}^p(-)$ , for all  $1 \leq p < \infty$ ;
2.  $K = 1$ ,  $\alpha = \varsigma$ , and  $\alpha' = \varsigma'$ , where the  $\varsigma, \varsigma'$  are shift amplitudes in the direction of a fixed vector in  $\mathbb{R}_{\geq 0}^n$  and its image under the projection to the first  $n-t$  components, respectively;
3.  $K = 1$ ,  $\alpha = I$ , and  $\alpha' = I'$ , where  $d_I, d_{I'}$  are interleaving distances in the direction of a fixed vector in  $\mathbb{R}_{\geq 0}^n$  and its image under the projection to the first  $n-t$  components, respectively.

Moreover,  $(-)/\tau$  is not continuous with respect to the span metric induced by  $\text{Hilb}^p(-)$  on target and source category, for any choice of finite measure  $\mu$  on the Lebesgue  $\sigma$ -algebra on  $\mathbb{R}^n$  and any nonzero measure  $\mu'$  on the one on  $\mathbb{R}^{n-t}$ , for  $1 \leq p < \infty$ .

*Proof.* By Lemma 7.10 and Remark 2.26, to prove 1.–2. it suffices to show that quotient restriction is amplitude bounding in the appropriate sense.

1.  $\text{Hilb}^p(-)$  is, trivially, amplitude bounding for any  $K > 0$  by Lemma 7.13.
2. Using the equivalent algebraic description of persistence modules, we have

$$x^{\varepsilon v'} m = x^{\varepsilon v'} (x^{\varepsilon(v-v')} m) = x^{\varepsilon v} m$$

for each  $m \in M/\tau$  and  $\varepsilon > 0$ . Thus, if multiplying by  $x^{\varepsilon v}$  annihilates  $M$ , multiplying by  $x^{\varepsilon v'}$  annihilates  $M/\tau$ . By Definition 2.19 of shift amplitude, this means that the quotient restriction functor  $(-)/\tau: (\text{PerM}_\mathbb{C}(\mathbb{R}^n), \varsigma) \rightarrow (\text{PerM}_\mathbb{C}(\mathbb{R}^{n-t}), \varsigma')$  is amplitude bounding with constant  $K = 1$ .

3. It is enough to observe that the quotient restriction sends  $\varepsilon$ -interleavings into  $\varepsilon'$ -interleavings, with  $\varepsilon' \leq \varepsilon$ .

Finally, consider any staircase upset  $U \subset \mathbb{R}^{n-t}$  with  $\mu'(U) \neq 0$ , which, if  $\mu' \neq 0$ , exists by continuity from below of  $\mu'$ . Further, define  $U_n := ne_1 + \mathbb{R}_{\geq 0}^t \times U$ , for all  $n \in \mathbb{N}$ . Consider  $\mathbb{F}[U_n]$ . By our assumption on  $\tau$ , we have that  $\mathbb{R}_{\geq 0}^t = \tau$ , and therefore  $\mathbb{F}[U_n]/\tau \cong \mathbb{F}[U]$ . Thus,  $\text{Hilb}^p(\mathbb{F}[U_n]/\tau) = c > 0$  independently of  $n$ . Furthermore, we have  $\cap_{n \in \mathbb{N}} U_n = \emptyset$ . As finite measures are continuous from above, we have that  $\text{Hilb}^p(\mathbb{F}[U_n]) = \mu(U_n)^p$  converges to 0. Hence,  $\mathbb{F}[U_n]$  converges to 0 in the span metric, showing that  $(-)/\tau$  is not continuous.  $\square$

The statement 1. should be understood more in terms of a general incompatibility between the  $L^p$ -amplitudes and the changes of dimension of the underlying poset than in terms of a stability result. To be more precise, topologically, Lemma 7.13 translates into the statement that  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  (as a space) is the topologically disjoint union of the fibers of quotient restriction. From this perspective, while quotient restriction is continuous with respect to the respective  $L^p$ -distances, this is because it is in fact locally constant. Therefore, we can interpret Proposition 7.14 as saying that while quotient restriction interacts well with shift amplitudes, it is essentially incompatible with any of the  $L^p$ -amplitudes.

**Proposition 7.15.** *Let  $\tau$  be some face of the positive cone in  $\mathbb{R}^n$  with positive dimension. The 0-th local cohomology  $H_{\tau}^0: (\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n), \alpha) \rightarrow (\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n), \alpha')$  is*

1. *Amplitude bounding with  $K = 1$  if  $\alpha = \alpha'$ , for any amplitude  $\alpha$ ;*
2. *1-Lipschitz continuous with respect to the interleaving distance  $d_I$ ;*
3. *Not continuous with respect to  $d_{\text{Hilb}^p(-)}$ , for  $1 \leq p < \infty$ , with respect to any finite, nonzero measure on the Lebesgue  $\sigma$ -algebra on  $\mathbb{R}^n$ ;*
4. *1-Lipschitz continuous with respect to  $d_{\text{Hilb}^p(-)}$  with the Lebesgue measure on  $\mathbb{R}^n$ , for  $1 \leq p < \infty$ .*

*Proof.* 1. Since  $H_{\tau}^0$  is not exact, we cannot use Remark 2.26. Nonetheless, the claim is an immediate consequence of the monotonicity of amplitudes, since  $H_{\tau}^0(M)$  is given by a subobject of  $M$ .

2. Kernels, products, and localisations all commute with shifts, and therefore, so does  $H_{\tau}^0$ . In particular,  $H_{\tau}^0$  carries  $\varepsilon$ -interleavings into  $\varepsilon$ -interleavings, and the claim follows.

3. Consider closed cubes  $Q_k := [0, k]^n \subset \mathbb{R}^n$  with side-length  $k$ , for  $k \in \mathbb{N}$ . Then  $M^k := \mathbb{F}[Q_k]$  defines a sequence of finitely presented modules converging to  $\mathbb{F}[\mathbb{R}_{\geq 0}^n]$  in  $d_{\text{Hilb}^p(-)}$  with respect to any finite measure. As  $\tau \neq 0$ , we have that  $H_{\tau}^0(\mathbb{F}[\mathbb{R}_{\geq 0}^n]) = 0$ . At the same time, it holds that  $H_{\tau}^0(M^k) = M^k$ , as  $Q_k$  is bounded in all directions. The latter sequence does not converge to 0, and the claim follows.

4. To prove the fourth claim, we show that  $H_{\tau}^0$  fulfills conditions 1 and 2 of Proposition 2.25. Condition 2 is satisfied by the left exactness of  $H_{\tau}^0$ . For condition 1, let  $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$  be an exact sequence in  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  such that  $\text{Hilb}^p(M') < \infty$ . Consider the following diagram with exact horizontal rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M'' \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \prod_{0 \neq \rho \subset \tau} M'_{\rho} & \longrightarrow & \prod_{0 \neq \rho \subset \tau} M_{\rho} & \longrightarrow & \prod_{0 \neq \rho \subset \tau} M''_{\rho} \longrightarrow 0 \end{array}$$

Assume  $\text{Hilb}^p(\prod_{0 \neq \rho \subset \tau} M'_{\rho}) = 0$ . Then, by the snake lemma, we have a short exact sequence  $0 \rightarrow H_{\tau}^0(M') \rightarrow H_{\tau}^0(M) \rightarrow H_{\tau}^0(M'') \rightarrow N$ , where  $N$  is a quotient of  $\prod_{0 \neq \rho \subset \tau} M'_{\rho}$ . In particular,  $\text{Hilb}^p(N) = 0$ , i.e.  $N \cong 0$  in  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)/0_{\text{Hilb}^p(-)}$ . Hence, the former sequence with

$N$  replaced by 0 is  $\text{Hilb}^p(-)$ -exact, and condition 1 is satisfied. It remains to prove that  $\text{Hilb}^p\left(\prod_{0 \neq \rho \subset \tau} M'_\rho\right) = 0$ . As  $\prod_{0 \neq \rho \subset \tau} M'_\rho$  is a finite product, it suffices to show that  $M'_\rho$  has amplitude 0, for  $\rho \subset \tau$  a face of positive dimension. Without loss of generality, we assume that  $\rho = \mathbb{R}_{\geq 0}\langle e_1, \dots, e_t \rangle$  for some  $t$  smaller than the dimension of  $\tau$ . Note that, for a staircase module  $N$  to have  $L^p$ -amplitude 0, it suffices that each submodule  $N' \subset N$  generated by a homogeneous element  $x \in N$  has  $L^p$ -amplitude 0. Indeed, this condition already implies that the fibers of the encoding over which the module is nontrivial are all 0-sets. So, we assume that  $m \in M_\rho$  is supported on a set  $S$  with nonzero measure. As all transition morphisms in direction  $\rho$  are isomorphisms, we may assume  $m$  to come from some homogeneous  $m' \in M'$  with  $x^v m' \neq 0$  for all  $v \in \rho$ . By shifting, we may assume the degree of  $m'$  to be zero. By assumption, there exists some  $\varepsilon > 0$  such that  $x^u m \neq 0$  for all  $u$  in the complement  $\rho^c$  of  $\rho$  whose norm is lesser or equal to  $\varepsilon$ . Thus, also  $x^{u+v} m' \neq 0$  for  $v \in \rho$  and  $u \in \rho^c$  with  $\|u\| \leq \varepsilon$ . But then,  $m'$  is supported on  $(\mathbb{R}^t \times [0, \varepsilon)^{n-t}) \cap \mathbb{R}_{\geq 0}^n$ . Hence,  $m'$  is supported on a set with infinite Lebesgue measure, in contradiction to the amplitude finiteness of  $M'$ .  $\square$

We can now conclude by proving some (in)stability results for the  $c_\tau$ -rank.

**Proposition 7.16.** *The  $c_\tau$ -rank as a function of pseudo metric spaces  $c_\tau: (\text{ob PerM}_\mathcal{C}(\mathbb{R}^n), d) \rightarrow [0, \infty]$  is:*

1. *Not continuous when  $d = d_I$ , the interleaving distance along a diagonal vector;*
2. *Not continuous when  $d = d_{\text{Hilb}^p(-)}$ , where  $\text{Hilb}^p(-)$  is with respect to any finite, nonzero measure on the Lebesgue  $\sigma$ -algebra on  $\mathbb{R}^n$ , for  $1 \leq p < \infty$ ;*
3.  *$K$ -Lipschitz continuous when  $d = d_{\text{Hilb}^p(-)}$ , where  $\text{Hilb}^p(-)$  is with respect to the Lebesgue measure, for any  $K > 0$  for  $1 \leq p < \infty$ .*

*Proof.* 1. For  $k \in \mathbb{N}$ , consider the sequence of persistence modules

$$M_k := \mathbb{F}\left[\left[0, \frac{1}{k}\right] \times [0, 1]^{t-1} \times \mathbb{R}_{\geq 0}^{n-t}\right]^k.$$

For  $k \rightarrow \infty$ ,  $M_k$  converges to 0 in the interleaving distance. At the same time, we have

$$H_\tau^0(M_k)/\tau^c = \mathbb{F}\left[\left[0, \frac{1}{k}\right] \times [0, 1]^{t-1}\right]^k.$$

Thus,

$$c_\tau(M_k) = \text{Hilb}\left(\mathbb{F}\left[\left[0, \frac{1}{k}\right] \times [0, 1]^{t-1}\right]^k\right) = k\lambda\left(\left[0, \frac{1}{k}\right] \times [0, 1]^{t-1}\right) = 1$$

showing that  $c_\tau$ -rank is not continuous in 0 with respect to the interleaving distance.

2. Using the same technique as the previous claim, define  $M_k := \mathbb{F}[ke_{k+1} + [0, 1]^t \times \mathbb{R}_{\geq 0}^{n-t}]$ , for all  $k \in \mathbb{N}$ . By an analogous computation, we have  $c_\tau\text{-rank}(M_k) = 1$ . At the same time,  $M_k$  converges to 0, by the continuity from above of finite measures.

3. By Propositions 7.14 and 7.15,  $(-)/\tau^c \circ H_\tau^0$  is amplitude bounding, for any constant  $K > 0$ , for the amplitude  $\text{Hilb}^p(-)$  with respect to the Lebesgue measure both on target and source. Since any amplitude is 1-Lipschitz with respect to the span metric induced by it (see Lemma 2.12), the result follows by composition of Lipschitz functions.  $\square$

Note that the Lipschitz-continuity with respect to any  $K$  and  $L^p$ -amplitudes essentially translates to the fact that the fibers of the  $c_\tau$ -rank all have infinite distance from each other. Hence, with respect to these distances the  $c_\tau$ -rank is actually locally constant. Summarizing, the  $c_\tau$ -rank



does not fulfill a meaningful continuity result with respect to any of the amplitude distances of Proposition 7.16.

There are two possible ways to amend the difficulties we described in the previous paragraph. The first one is to consider metrics specifically depending on  $\tau$ . In applications, different parameter directions might often carry very different information. Hence, one could for example consider  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$  as  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^{n-t})$  valued persistence modules over  $\mathbb{R}^t$  and then study  $L^p$ -distance with respect to different amplitudes on  $\text{PerM}_{\mathfrak{C}}(\mathbb{R}^n)$ . Alternatively, when  $\tau$  is a face of codimension 1, one may consider the barcode of  $H_{\tau}^0(M)/\tau^c$  as an invariant instead. This contains more information than the  $c_{\tau}$ -rank. At the same time, by the isometry theorem in [Les15, Thm. 3.4], together with Proposition 7.14 and Proposition 7.15, this gives an invariant that is Lipschitz continuous with respect to the interleaving distance.

## Acknowledgments

We would like to thank the organizers of the 2020 Applied Category Theory Adjoint School, for bringing us together in this collaboration. Part of this work was initiated during a postdoctoral fellowship of NO and an internship of LW at the Max Planck Institute for Mathematics in the Sciences, and NO and LW would like to thank the Max Planck Institute for their support. We thank Håvard B. Bjerkevik, Wojciech Chachólski, Adrian Clough, Nicolò De Ponti, Oliver Gäfvert, and Ezra Miller for useful discussions. BG was supported by the grant FAR2019-UniMORE and by the Austrian Science Fund (FWF) grant numbers P 29984-N35 and P 33765-N. NO was supported by Royal Society Research Grant RGS\R2\212169.

## References

- [AAJ<sup>+</sup>23] Dashti Ali, Aras Asaad, Maria-Jose Jimenez, Vidit Nanda, Eduardo Paluzo-Hidalgo, and Manuel Soriano-Trigueros. A survey of vectorization methods in topological data analysis. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, pages 1–14, 2023.
- [AEK<sup>+</sup>17] Henry Adams, Tegan Emerson, Michael Kirby, Rachel Neville, Chris Peterson, Patrick Shipman, Sofya Chepushtanova, Eric Hanson, Francis Motta, and Lori Ziegelmeier. Persistence images: A stable vector representation of persistent homology. *Journal of Machine Learning Research*, 18(8):1–35, 2017.
- [BBCB20] Magnus Bakke Botnan and William Crawley-Boevey. Decomposition of persistence modules. *Proceedings of the AMS*, 148:4581–4596, 2020.
- [BBL21] Håvard Bakke Bjerkevik and Michael Lesnick.  $\ell^p$ -distances on multiparameter persistence modules, 2021. Preprint, available at arXiv:2106.13589.
- [BBLO20] Magnus Bakke Botnan, Vadim Lebovici, and Steve Oudot. On rectangle-decomposable 2-parameter persistence modules. In *36th International Symposium on Computational Geometry, SoCG 2020, June 23–26, 2020, Zürich, Switzerland*, volume 164 of *LIPIcs*, pages 22:1–22:16, 2020.
- [BBOO21] Magnus Bakke Botnan, Steffen Oppermann, and Steve Oudot. Signed barcodes for multi-parameter persistence via rank decompositions and rank-exact resolutions, 2021. Preprint, available at arXiv:2107.06800.

- [BCZ21] Paul Biran, Octav Cornea, and Jun Zhang. Triangulation and persistence: Algebra 101, 2021. Preprint, available at arXiv:2104.12258.
- [BE18] Mickaël Buchet and Emerson G. Escolar. Realizations of indecomposable persistence modules of arbitrarily large dimension. In *34th International Symposium on Computational Geometry (SoCG 2018)*, volume 99 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 15:1–15:13, Dagstuhl, Germany, 2018.
- [Ber22] Nicolas Berkouk. Persistence and the sheaf-function correspondence, 2022. Preprint, available at arXiv:2207.06335.
- [BG16] Arkady Berenstein and Jacob Greenstein. Primitively generated hall algebras. *Pacific Journal of Mathematics*, 281, 2016.
- [BL15] Ulrich Bauer and Michael Lesnick. Induced matchings of barcodes and the algebraic stability of persistence. *Journal of Computational Geometry*, 6(2):162–191, 2015.
- [BMSW17] Ulrich Bauer, Axel Munk, Hannes Sieling, and Max Wardetzky. Persistence barcodes versus Kolmogorov signatures: Detecting modes of one-dimensional signals. *Foundations of Computational Mathematics*, 17(1):1–33, Feb 2017.
- [BP21] Nicolas Berkouk and François Petit. Ephemeral persistence modules and distance comparison. *Algebraic & Geometric Topology*, 21(1):247–277, Feb 2021.
- [BPC<sup>+</sup>18] Francisco Belchi, Mariam Pirashvili, Joy Conway, Michael Bennett, Ratko Djukanovic, and Jacek Brodzki. Lung topology characteristics in patients with chronic obstructive pulmonary disease. *Scientific Reports*, 8(1):5341, 2018.
- [BSS23] Peter Bubenik, Jonathan Scott, and Donald Stanley. Exact weights, path metrics, and algebraic wasserstein distances. *Journal of Applied and Computational Topology*, 7(2):185–219, Jun 2023.
- [Bub15] Peter Bubenik. Statistical topological data analysis using persistence landscapes. *Journal of Machine Learning Research*, 16(3):77–102, 2015.
- [BW20] David Bramer and Guo-Wei Wei. Atom-specific persistent homology and its application to protein flexibility analysis. *Computational and Mathematical Biophysics*, 8(1):1–35, 2020.
- [CCBdS16] Frederic Chazal, William Crawley-Boevey, and Vin de Silva. The observable structure of persistence modules. *Homology, Homotopy and Applications*, 18(2):247–265, 2016.
- [CCSG<sup>+</sup>09] Frédéric Chazal, David Cohen-Steiner, Marc Glisse, Leonidas J. Guibas, and Steve Y. Oudot. Proximity of persistence modules and their diagrams. In *Proceedings of the Twenty-Fifth Annual Symposium on Computational Geometry*, SCG '09, pages 237–246, New York, NY, USA, 2009. Association for Computing Machinery.
- [CSEHM10] David Cohen-Steiner, Herbert Edelsbrunner, John Harer, and Yuriy Mileyko. Lipschitz functions have  $l_p$ -stable persistence. *Foundations of Computational Mathematics*, 10(2):127–139, 2010.
- [CZ09] Gunnar Carlsson and Afra J. Zomorodian. The theory of multidimensional persistence. *Discrete & Computational Geometry*, 42(1):71–93, Jul 2009.

- [GC17] Oliver Gäfvert and Wojciech Chachólski. Stable invariants for multidimensional persistence, 2017. Preprint, available at arXiv:1703.03632.
- [GG06] Benjamin T. Graham and Geoffrey R. Grimmett. Influence and sharp-threshold theorems for monotonic measures. *The Annals of Probability*, 34(5):1726–1745, 2006.
- [GLR22] Barbara Giunti, Jānis Lazovskis, and Bastian Rieck. DONUT: Database of Original & Non-Theoretical Uses of Topology, 2022. <https://donut.topology.rocks>.
- [HKNU17] Christoph Hofer, Roland Kwitt, Marc Niethammer, and Andreas Uhl. Deep learning with topological signatures. *Advances in Neural Information Processing Systems*, 30:1633–1643, 2017.
- [HOST19] Heather A. Harrington, Nina Otter, Hal Schenck, and Ulrike Tillmann. Stratifying multiparameter persistent homology. *SIAM Journal on Applied Algebra and Geometry*, 3(3):439–471, 2019.
- [Kal19] Sara Kališnik. Tropical coordinates on the space of persistence barcodes. *Foundations of Computational Mathematics*, 19(1):101–129, 2019.
- [KS90] Masaki Kashiwara and Pierre Schapira. *Sheaves on manifolds*, volume 292 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1990. With a chapter in French by Christian Houzel.
- [LBD<sup>+</sup>17] Yongjin Lee, Senja D. Barthel, Paweł Dłotko, S. Mohamad Moosavi, Kathryn Hess, and Berend Smit. Quantifying similarity of pore-geometry in nanoporous materials. *Nature communications*, 8:15396, 2017.
- [Les15] Michael Lesnick. The theory of the interleaving distance on multidimensional persistence modules. *Foundations of Computational Mathematics*, 15(3):613–650, 2015.
- [Mil20a] Ezra Miller. Essential graded algebra over polynomial rings with real exponents, 2020. Preprint, available at arXiv:2008.03819.
- [Mil20b] Ezra Miller. Homological algebra of modules over posets, 2020. Preprint, available at arXiv:2008.00063.
- [Oud15] Steve Y. Oudot. *Persistence theory: from quiver representations to data analysis*, volume 209 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015.
- [PSW21] Francois Petit, Pierre Schapira, and Lukas Waas. A property of the interleaving distance for sheaves, 2021. Preprint, available at arXiv:2108.13018.
- [RC18] Henri Riihimäki and Wojciech Chachólski. Generalized persistence analysis based on stable rank invariant, 2018. Preprint, available at arXiv:1807.01217.
- [RT16] Andrew Robinson and Katharine Turner. Hypothesis testing for topological data analysis, 2016. Preprint, available at arXiv:1310.7467.
- [Rud87] Walter Rudin. *Real and Complex Analysis, 3rd Ed.* McGraw-Hill, Inc., USA, 1987.
- [SCL<sup>+</sup>17] Martina Scalamiero, Wojciech Chachólski, Anders Lundman, Ryan Ramanujam, and Sebastian Öberg. Multidimensional persistence and noise. *Foundations of Computational Mathematics*, 17:1367–1406, 2017.

- [ST20] Primož Skraba and Katharine Turner. Wasserstein stability for persistence diagrams, 2020. Preprint, available at [arXiv:2006.16824](https://arxiv.org/abs/2006.16824).
- [Tho19] Ashleigh L. Thomas. Invariants and metrics for multiparameter persistent homology, 2019. PhD thesis, Duke University.
- [vdD98] Lou van den Dries. *Tame topology and o-minimal structures*, volume 248. Cambridge university press, 1998.
- [Waa23] Lukas Waas. Abelian categories of finitely encodable multiparameter persistence modules, 2023. Available at [www.mathi.uni-heidelberg.de/~lwaas/work/pers\\_notes\\_01.pdf](http://www.mathi.uni-heidelberg.de/~lwaas/work/pers_notes_01.pdf).