

# $O_M(\mathbb{R}^n)$ as locally convex Orlicz space

Jan Kiszyński

the date of receipt and acceptance should be inserted later

**Abstract** The original M.Valdivia proof of his theorem on representation of the space  $O_M$  uses results of S.Rolewicz concerning metric linear spaces and results of A.Grothendieck from his theory of topological tensor product. We present a more direct proof.

**Keywords** Multiplication · convolution · the M.Valdivia representation of  $O_M$

## 1 Introduction

Usually the L.Schwartz set  $S(\mathbb{R}^n)$  of infinitely differentiable functions on  $\mathbb{R}^n$  rapidly decreasing at infinity is treated as a topological space with distinguished topology. But in fact  $S(\mathbb{R}^n)$  admits at least two important locally convex topologies. These topologies are determined by systems of seminorms  $\{\rho_{\alpha,\beta}; \alpha \in \mathbb{N}, \beta \in \mathbb{N}_0^n\}$  and  $\{\tilde{\rho}_{\alpha,\beta}; \alpha \in \mathbb{N}, \beta \in \mathbb{N}_0^n\}$  whose exact definition will be given in Section 2.

Let  $S(\mathbb{R}^n)$  be the L.Schwartz set of infinitely differentiable functions on  $\mathbb{R}^n$  rapidly decreasing at infinity and let  $S'(\mathbb{R}^n)$  be its dual set. In 1981 M.Valdivia has shown in [12] that for every element  $\Phi$  the space  $O_M(\mathbb{R}^n)$  there is a distribution  $T$  on  $\mathbb{R}^n$  rapidly decreasing at infinity such that

$$\Phi \cdot \psi = T * \psi \quad \text{for every } \psi \in S(\mathbb{R}^n)$$

where  $T * \psi := T_{r_0}(\tilde{\psi})$  is a distinguished regularization of T.M.Valdivia used theory of linear metric spaces developed in the book [6] of S.Rolewicz and used also the theory of topological tensor products of A.Grothendieck [4]. Our aim is to prove the Valdivia theorem more directly, remaining in the framework of  $(S(\mathbb{R}^n), \tilde{\rho})$  and  $(S(\mathbb{R}^n), \tilde{\rho})'$ . We use the operators  $(m \cdot)$  of multiplication of elements of  $(S(\mathbb{R}^n), \tilde{\rho})$  by multipliers  $m$  of  $(S(\mathbb{R}^n), \tilde{\rho})$ , and we distinguish

the operators  $(m \cdot)$  from the multipliers themselves. Moreover, we use the elementary convolution  $(S'(\mathbb{R}^n), \tilde{\rho})' \times (S(\mathbb{R}^n), \tilde{\rho}) \ni (T, \psi) \rightarrow T(\tilde{\psi}) \in (S(\mathbb{R}^n), \tilde{\rho})$ . Notice that elementary convolutions are deeply related to regularization, the last being described in Sect.4.10 of John Horváth book [6]. Distributions  $T \in (S(\mathbb{R}^n), \tilde{\rho})'$  rapidly decreasing at  $\infty$  are in [3, Sect.125.0] characterized by their regularizations. The distinguished regularization  $T_{r_0}$  occurring in the proof of 4.1 has the value at  $\tilde{\psi} \in (S(\mathbb{R}^n), \tilde{\rho})$  equal to  $F(\psi)$ , the Fourier image of  $\psi$ . This resembles the Fourier-Schwartz transformation.

## 2 Locally convex topologies in the L.Schwartz set $S(\mathbb{R}^n)$

Except of the L.Schwartz set  $S(\mathbb{R}^n)$  of interest is a larger set  $\tilde{S}(\mathbb{R}^n)$  whose element are these functions  $\psi \in C^\infty(\mathbb{R}^n; \mathbb{C})$  for which  $\lim_{|x| \rightarrow \infty} [\partial^\beta \psi](x) = 0$  for every  $\beta \in \mathbb{N}_0^n$  in sense of the usual convergence, not necessarily rapid. The set  $\tilde{S}(\mathbb{R}^n)$  admits the locally convex topology  $\rho$  determined by system of seminorms  $\{\rho_{\alpha, \beta}; \alpha \in \mathbb{N}, \beta \in \mathbb{N}_0^n\}$  such that

$$[\rho_{\alpha, \beta}(\psi)](x) = \sup_{x \in \mathbb{R}^n} (1 + |x|^2)^{-\frac{\alpha}{2}} |[\partial^\beta \psi](x)| \text{ for every } x \in \mathbb{R}^n$$

and every  $\psi \in \tilde{S}(\mathbb{R}^n)$ .

The set  $\tilde{S}(\mathbb{R}^n)$  admits also the locally convex topology  $\tilde{\rho}$ , determined by the system of seminorms  $\{\tilde{\rho}_{\alpha, \beta}; \alpha \in \mathbb{N}, \beta \in \mathbb{N}_0^n\}$  such that

$$[\tilde{\rho}_{\alpha, \beta}(\psi)](x) = \int_{\mathbb{R}^n} (1 + |x|^2)^{-\frac{\alpha}{2}} |[\partial^\beta \psi](x)| dx \text{ for every } \psi \in \tilde{S}(\mathbb{R}^n).$$

where the integral over  $\mathbb{R}^n$  is an absolutely convergent integral in sense of Lebesgue. Indeed, from [2, Lemma 4.15] it follows that there holds

**Theorem 2.1** *For every  $\alpha \in \mathbb{N}, \beta \in \mathbb{N}_0^n$  and  $\psi \in \tilde{S}(\mathbb{R}^n)$  there is  $\mu \in [0, \infty]$  such that*

$$\tilde{\rho}_{\alpha, \beta}(\psi) \leq \mu \rho_{\alpha, \beta}(\psi).$$

*Theorem 2.1 implies the*

**Corollary 2.1** *In the set  $\tilde{S}(\mathbb{R}^n)$  the topology  $\tilde{\rho}$  is no stronger than the topology  $\rho$ .*

## 3 Multipliers of $(S(\mathbb{R}^n), \tilde{\rho})$

The function  $m \in C^\infty(\mathbb{R}; \mathbb{C})$  is called multiplier of the space  $(S(\mathbb{R}^n), \tilde{\rho})$ , if  $m \cdot (S(\mathbb{R}^n), \tilde{\rho}) \subset (S(\mathbb{R}^n), \tilde{\rho})$ . The set of all multipliers of  $(S(\mathbb{R}^n), \tilde{\rho})$  is denoted by  $m(\mathbb{R}^n)$  and constitutes a complex multiplication algebra. Let  $O_M(\mathbb{R}^n)$  be the set of functions belonging to  $C^\infty(\mathbb{R}^n; \mathbb{C})$ , growing at the most polynomially at  $\infty$ . The above assumption about growth means exactly that for every  $\Phi \in$

$O_M(\mathbb{R}^n)$  and every multiindex  $\beta \in \mathbb{N}_0^n$  there is  $\alpha \in \mathbb{N}$  such that  $\lim_{|x| \rightarrow \infty} \sup (1 + |x|^2)^{-\frac{\alpha}{2}} |[\partial^\beta \Phi](x)| < \infty$ , in usual sense, i.e. non necessarily in sense of rapid convergence.

**Theorem 3.1** ( $m(\mathbb{R}^n) = O_M(\mathbb{R}^n)$ .) *The inclusion  $O_M(\mathbb{R}^n) \subset m(\mathbb{R}^n)$  is almost evident. Indeed, if  $\Phi \in O_M(\mathbb{R}^n)$  and  $\psi \in S(\mathbb{R}^n)$ , the the product  $\Phi \cdot \psi$  is an infinitely differentiable complex valued function on  $\mathbb{R}^n$  rapidly decreasing to zero at infinity, so that  $\Phi \in m(\mathbb{R}^n)$ . Proof of the inclusion  $m(\mathbb{R}^n) \subset O_M(\mathbb{R}^n)$  is more complicated and is postponed to the Appendix.*

**Theorem 3.2** *If  $m \in m(\mathbb{R}^n)$ , then the mapping  $(S(\mathbb{R}^n), \tilde{\rho}) \ni \psi \rightarrow m \cdot \psi \in (S(\mathbb{R}^n), \tilde{\rho})$  is continuous.*

*Proof* Since the locally convex space  $(S(\mathbb{R}^n), \tilde{\rho})$  is metrizable, its topology is characterized whenever characterized are the convergent countable sequences  $(\psi_k)_{k \in \mathbb{N}}$  of functions belonging to  $(S(\mathbb{R}^n), \tilde{\rho})$ . We can characterize the convergence of such sequences is as follows:

**Lemma 3.1** *The countable sequence  $(\psi_k)_{k \in \mathbb{N}} \subset (S(\mathbb{R}^n), \tilde{\rho})$  is convergent to zero if and only if for every  $\beta \in \mathbb{N}_0^n$ , the following two conditions are satisfied:*

$$\lim_{k \rightarrow \infty} \partial^\beta \psi_k = 0 \text{ almost uniformly on } \mathbb{R}^n, \quad (3.1)$$

$$\lim_{|x| \rightarrow \infty} |x|^\alpha |\partial^\beta \psi_k(x)| = 0 \text{ uniformly with respect to } \alpha \in \mathbb{N}. \quad (3.2)$$

*The limits occurring in these conditions are understood in usual sense of Calculus, i.e. not in sense of rapid convergence.*

*Proof* The assertions (3.1) and (3.2) express in terms of the Calculus the fact that the sequence  $(\psi_k)_{k \in \mathbb{N}}$  converges to zero in sense of  $(S(\mathbb{R}^n), \tilde{\rho})$ , i.e. that  $\lim_{k \rightarrow \infty} \tilde{\rho}_{\alpha, \beta}(\psi_k) = 0$  for every  $(\alpha, \beta) \in \mathbb{N} \times \mathbb{N}_0^n$  where  $\tilde{\rho}_{\alpha, \beta}(\psi_k) = \int_{\mathbb{R}^n} (1 + |x|^2)^{\frac{\alpha}{2}} |(\partial^\beta \psi_k)(x)| dx$ . Therefore the countable sequence  $(\psi_k)_{k \in \mathbb{N}}$  converges to zero in the topology of  $(S(\mathbb{R}^n), \tilde{\rho})$  if and only if both the conditions (3.1) and (3.2) are satisfied.

*Proof (of Theorem 3.2)* We have to prove that if  $m \in m(\mathbb{R}^n)$  and both the conditions (3.1) and (3.2) are satisfied, then analogous condition, say (3.1)\* and (3.2)\*, are satisfied for the sequence  $(m \cdot \psi_k)_{k \in \mathbb{N}}$ . Indeed, the Leibniz rule (see [6, Sect.2.5, Proposition 3]) says that

$$\partial^\alpha (m \cdot \psi_k) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} (\partial^{\alpha-\beta}(m)) \cdot (\partial^\beta \psi_k).$$

Hence (3.1)\* is a consequence of (3.1) and (3.2)\* is a consequence of (3.2).

**Remark 3.1** If  $m \in m(\mathbb{R}^n)$  is given, then the mapping  $(S(\mathbb{R}^n), \tilde{\rho}) \ni \psi \rightarrow m \cdot \psi \in (S(\mathbb{R}^n), \tilde{\rho})$  is continuous.

#### 4 Appendix. Proof of the inclusion $m(\mathbb{R}^n) \subset O_M(\mathbb{R}^n)$ and proof of the M.Valdivia representation of the space $O_M(\mathbb{R}^n)$ by rapidly decreasing distributions

We shall divide this Appendix onto several subsections.

##### 4.1 $O_M(\mathbb{R}^n)$ as a locally convex Orlicz space.

Let the elements of  $O_M(\mathbb{R}^n)$  be denoted by  $\Phi$ . Then

$$O_M(\mathbb{R}^n) = \{\Phi \in C^\infty(\mathbb{R}^n; \mathbb{C}) : \text{for every } n \in \mathbb{N} \text{ the integral} \\ \int_{\mathbb{R}^n} (1 + |x|^2)^{-\frac{n}{2}} |[\partial^\beta \Phi](x)| dx \text{ is convergent}\}.$$

The above follows from the fact that elements of OM are operators of multiplication acting in the space  $(S(\mathbb{R}^n), \tilde{\rho})$  which admits the locally convex topology determined by the system of seminorms  $\{\tilde{\rho}_{\alpha,\beta}; \alpha \in \mathbb{N}, \beta \in \mathbb{N}_0^n\}$  such that for every  $\psi \in S(\mathbb{R}^n)$  the Lebesgue integral

$$\tilde{\rho}_{\alpha,\beta}(\psi) = \int_{\mathbb{R}^n} (1 + |x|^2)^{-\frac{n}{2}} |[\partial^\beta \Phi](x)| dx \text{ is convergent.}$$

It follows that  $O_M(\mathbb{R}^n)$  is uniquely determined by the system of spaces  $\{S_{-n} : n \in \mathbb{N}\}$  consisting of convergent integrals

$$\int_{\mathbb{R}^n} (1 + |x|^2)^{-\frac{n}{2}} \sup_{|\beta| \leq b_n} |[\partial^\beta \Phi](x)| dx \quad (4.1)$$

where  $b_n \in \mathbb{N}$  is fixed for every space  $S_{-n}$ . Each  $S_{-n}$  is a locally convex space with topology determined by system of seminorms equal to integrals occurring in (4.1). The system of spaces  $\{S_{-n} : n \in \mathbb{N}\}$  can be treated as a single locally convex Orlicz space denoted by  $\tilde{O}$  whose topology is determined by the system of seminorms  $\{s_\beta : \beta \in \mathbb{N}_0^n\}$  such that  $s_\beta(\Phi) = \int_{\mathbb{R}^n} (1 + |x|^2)^{-\frac{n}{2}} |[\partial^\beta \Phi](x)| dx$  for

every  $\Phi \in \tilde{O}$ , and the set of convergence factors is equal to  $\{(1 + |x|^2)^{-\frac{n}{2}} : n \in \mathbb{N}\}$ . In definition of this locally convex Orlicz space the above seminorms and the above convergence factors are equally important.

##### 4.2 The locally convex space $((m(\mathbb{R}^n)\cdot), op_m)$ of operators of multiplication

For every  $m \in m(\mathbb{R}^n)$  denote by  $(m\cdot)$  the operator of multiplication the elements of  $S(\mathbb{R}^n)$  by  $m$ . The set  $(m(\mathbb{R}^n)\cdot)$  of multiplication operators admits the locally convex topology determined by the set of seminorms  $\{s_{\alpha,\beta,\psi}; \alpha \in \mathbb{N}, \beta \in \mathbb{N}_0^n, \psi \in S(\mathbb{R}^n)\}$  such that

$$s_{\alpha,\beta,\psi}((m\cdot)) = \tilde{\rho}_{\alpha,\beta}(m \cdot \psi) \text{ for every } (m(\mathbb{R}^n)\cdot).$$

The so defined locally convex topology in the set  $(m(\mathbb{R}^n)\cdot)$  will be denoted by  $op_m$ . Existence of this topology is crucial for Subsection 4.5. Notice that for compact subsets of  $\mathbb{R}^n$  the topology  $op_m$  coincides with the topology introduced in [6, Sect.2.4, Example 11 and Sect.2.5, Example 8]. The elementary convolution  $T * \psi$  where  $\psi \in S(\mathbb{R}^n)$  and  $T$  is a distribution on  $(S(\mathbb{R}^n), \tilde{\rho})$  was analysed in [13, Sect.VI.3].

#### 4.3 The openworkness of the set $(m(\mathbb{R}^n)\cdot)$

The set  $(m(\mathbb{R}^n)\cdot)$  is openwork (in sense of this adjective explained in common words in [12, p.724]) because for every  $k \in \mathbb{N}$  there exists in  $(m(\mathbb{R}^n)\cdot)$  the inner automorphism ([2, Sect.190.D], [7, Sect.I.7])

$$A_k : (m(\mathbb{R}^n)\cdot) \ni (m\cdot) \rightarrow F_o^k \circ (m\cdot) \circ F_o^{-k} \in (m(\mathbb{R}^n)\cdot).$$

In the above formula  $F_o$  denotes the Fourier transformation restricted to  $S(\mathbb{R}^n)$ . Let us stress that using the bracket-less symbol of superposition of operators we admit the rule that the operators act in order from right to left. For our proof of Theorem 3.1 the openworkness of  $(m(\mathbb{R}^n)\cdot)$  is only nonessential curiosity.

#### 4.4 Proof of the inclusion $m(\mathbb{R}^n) \subset O_M(\mathbb{R}^n)$

**Theorem 4.1** *The inclusion*

$$m(\mathbb{R}^n) \subset O_M(\mathbb{R}^n) \tag{4.2}$$

*is true*

*Proof* Usual multiplication of complex numbers implies that the inclusion (4.2) is equivalent to the inclusion

$$m(\mathbb{R}^n) \cdot \psi \subset O_M(\mathbb{R}^n) \cdot \psi \text{ for every } \psi \in (S(\mathbb{R}^n), \tilde{\rho}). \tag{4.3}$$

Left side of 4.3 is equal to the set of values of the operators  $(m(\mathbb{R}^n)\cdot)$  acting in  $S(\mathbb{R}^n)$ . Right side of 4.3 is equal to the set of values operators  $\{\phi \in O_M(\mathbb{R}^n)\}$  which, by definition of  $O_M(\mathbb{R}^n)$ , is the maximal set of multiplication operators acting in  $S(\mathbb{R}^n)$ .  $\square$

#### 4.5 Proof of M.Valdivia representation of the space $O_M(\mathbb{R}^n)$ by rapidly decreasing distributions

We shall consider actions by multiplication. All the used properties of multiplication are consequences of the usual properties of multiplication of complex numbers. Recall that a function  $m \in C^\infty(\mathbb{R}^n; \mathbb{C})$  is called a multiplier of the

space  $(S(\mathbb{R}^n), \tilde{\rho})$  if  $m \cdot (S(\mathbb{R}^n), \tilde{\rho}) \in (S(\mathbb{R}^n), \tilde{\rho})$ . The above (very popular) definition concerns a single element of  $C^\infty(\mathbb{R}^n; \mathbb{C})$  and is not useful if one wants to speak about action of a set of operators of multiplication in a set of functions.

We define the action of the set  $(m(\mathbb{R}^n) \cdot)$  of operators of multiplication in the set  $m(\mathbb{R}^n)$  of multipliers of the space  $(S(\mathbb{R}^n), \tilde{\rho})$  by the equality

$$(m(\mathbb{R}^n) \cdot) \cdot m(\mathbb{R}^n) := m(\mathbb{R}^n) \cdot (S(\mathbb{R}^n), \tilde{\rho}). \quad (4.4)$$

Using theory of regularization of distributions exposed by John Horváth in Section 4.10 of his book [6], and using the definition of distribution on  $\mathbb{R}^n$  rapidly decreasing at  $\infty$  due in [3, Sect.125.0], one concludes that the representation theorem of M.Valdivia is equivalent to the following Theorem 4.2.

**Theorem 4.2** *The distribution  $T \in S(\mathbb{R}^n)$  rapidly decreases at  $\infty$  if and only if the set  $C^\infty(\mathbb{R}^n; \mathbb{C})$  admits the locally convex topology determined by the system of seminorms  $\{s_\tau : \tau \in \Xi\}$  such that if  $\psi \in C^\infty(\mathbb{R}^n; \mathbb{C})$ , then  $s_\tau(\psi) \in F(\psi)$  for every  $\tau \in \Xi$*

*Proof* The formulas occurring in Theorem 4.2 mean that there is a unique regularization  $\hat{R}$  defined on  $(S(\mathbb{R}^n), \tilde{\rho})'$  such that  $\hat{R}(T) = T(\psi)$  for every  $T \in (S(\mathbb{R}^n), \tilde{\rho})'$  and every  $\psi \in (S(\mathbb{R}^n), \tilde{\rho})$ . Existence and uniqueness of such regularization are obvious.  $\square$

#### 4.6 A remark concerning the Gagliardo-Nirenberg equality

Consider the metrizable linear space  $C_B \subset C^\infty(\mathbb{R}^n; \mathbb{C})$  and its imbedding into Lebesgue space  $L(\mathbb{R}^n)$ . In their book [2] R.A.Adams and J.J.F.Fournier observed that existence of the imbedding of  $C_B$  into  $L(\mathbb{R}^n)$  is equivalent to the Lemma 4.15 of [2] which, by turns, is equivalent to the simple behaviour of  $C_B$  described on p.79 of [2]. This observation (as well as sole Lemma 4.15) is surprising. Also its connections with famous Gagliardo-Nirenberg equality are interesting. For that matter we shall limit ourselves to superficial presentation of the related literature.

The papers I and II of E.Gagliardo and III of L.Nirenberg are quoted in the books [10] and [11].

- I. E.Gagliardo, *Careatteristiche della tracce sulla frontiera relative ad alcune di funzioni in  $n$  variabili*,  
Rend. Sem. Mat. Padova 27 (1957), 284-305.
- II. E.Gagliardo, *Proprietà di alcune classi di funzioni in più variabili*,  
Ricerche Mat. Napoli 7 (1958), 102-137.
- III. L.Nirenberg, *On elliptic partial differential equations*,  
Ann. della Scuole Normale Superiore, Pisa 13 (1959), 116-169.

Russian Academy of Sciences published the translations of E.Gagliardo papers I and II in *Matematika* 5:4 (1961), 87-116.

Haim Brezis and Petru Mironescu published in HaL a survey of papers related to Gagliardo- Nirenberg equality.

## References

1. Adams, R.A.: *Sobolev Spaces*, Academic Press, 1975.
2. Adams, R.A., and Fournier, J.J.F. : *Sobolev Spaces*, 2nd ed., Academic Press, 2003.
3. *Encyclopedic Dictionary of Mathematics*, by Mathematical Society of Japan, second edition of the English translation, MIT Press, 1987.
4. Grothendieck, A.: *Topological Tensor Products and Nuclear Spaces*, Mem. Amer. Math. Soc., No. 16, 1955 (in French).
5. Harjulehto, P. and Hästö, P.: *Orlicz Spaces and Generalized Orlicz Spaces*, Springer, 2019.
6. Horváth, J.: *Topological Vector Spaces and Distributions*, Addison-Wesley, 1966, Dover Publications, 2012.
7. Lang, S.: *Algebra*, Addison Wesley, 1965 ; Russian transl., Moscow, 1968.
8. Rolewicz, S.: *Metric Linear Spaces*, PWN, Warszawa, 1972.
9. Schwartz, L.: *Théorie des Distributions*, new edition, Hermann, Paris, 1966 (in French).
10. Stein, E.M.: *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970 ; Russian tranns.: Moscow, 1973.
11. Stein, E.M. and Weiss, G.: *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press 1971 ; Russian transl.: Moscow, 1974.
12. Valdivia, M.: *A representation of the space  $O_M$* , Math.Z. 177 (4), 463-478 (1981).
13. Yosida, K.: *Functional Analysis*, Sixth Ed., Springer, 1980.