Symbols of compact truncated Toeplitz operators

Ryan O'Loughlin

E-mail address: R.OLoughlin@leeds.ac.uk

School of Mathematics, University of Leeds, Leeds, LS2 9JT, U.K.

Abstract

This paper characterises the dual of the model space K_I^1 , where I is an inner function, intersected with the shifted Hardy space, zH^1 . With this duality result, it is then shown that every bounded truncated Toeplitz operator on the model space K_I^2 has a bounded symbol if and only if every compact truncated Toeplitz operator on K_I^2 has a symbol which is of the form I multiplied by a continuous function.

Keywords: Hardy space, truncated Toeplitz operator, model space.

MSC: 30H10, 47B35, 46E20.

1 Introduction

In this paper we provide two new overlapping results. We first characterise the dual space of any given backward shift invariant subspace of the Hardy space H^1 intersected with zH^1 . We then use our duality result to study the question: when does a bounded truncated Toeplitz operator have a bounded symbol? This question has generated much research interest and is one of the most fundamental problems concerning truncated Toeplitz operators. The main result of this paper shows that every bounded truncated Toeplitz operator on K_I^2 has a bounded symbol if and only if every compact truncated Toeplitz operator on K_I^2 has a symbol which is of the form $I\phi$ where ϕ is a continuous function on the unit circle.

Throughout we let $1 \leq p < \infty$ and we denote the unit circle by \mathbb{T} . All L^p spaces, and L^{∞} , are assumed to be on the unit circle, \mathbb{T} . We denote the Hardy space on the unit circle by H^p and we refer the reader to [8, 15] for background theory on the Hardy space. Throughout we let I be an inner function (i.e. a function in H^2 which is unimodular on \mathbb{T}). It is shown in Chapter 5 of [7] that all backward shift invariant subspaces of H^p are of the form $K_I^p := H^p \cap IH_0^p$, where $\overline{H_0^p} := \{\overline{f} : f \in zH^p\}$. We call the space K_I^p a model space, and we refer the reader to [7] for background theory on model spaces. Integrals will be evaluated with respect to m, where mis the normalised Lebesgue measure on \mathbb{T} . We define $C(\mathbb{T})$ to be the continuous functions on the unit circle. We let BMOA denote the set of all analytic functions of bounded mean oscillation, i.e., $f \in BMOA$ means $f \in H^2$ and

$$\sup_{A} \frac{1}{|A|} \int_{A} |f - f_A| dm < \infty,$$

where the supremum is taken over all arcs $A \subset \mathbb{T}$ and

$$f_A := \frac{1}{|A|} \int_A f dm.$$

It can be checked that BMOA is a linear vector space and an easy adaptation of Proposition 2.5 in [13] shows that when equipped with the norm

$$||f||_* := \left| \int_{\mathbb{T}} f dm \right| + \sup_A \frac{1}{|A|} \int_A |f - f_A| dm,$$

BMOA becomes a Banach space.

When $1 we let <math>P : L^p \to H^p$ be the Riesz projection, and $P_I = PI(Id - P)\overline{I}$ be the projection from L^p to K_I^p . When p = 2, P and P_I are in fact orthogonal projections. As noted in Section 2 of [16], these projections can be expressed as integral operators independent of $p \in (1, \infty)$. With these projections, for $1 we can write <math>L^p = \overline{H_0^p} \oplus H^p$ or $L^p = \overline{H_0^p} \oplus K_I^p \oplus IH^p$ (see page 20 of [17] for details) and furthermore for p = 2 the summations are orthogonal summations.

For $\phi \in L^2$, we define the truncated Toeplitz operator A_{ϕ} on bounded functions from K_I^2 by

$$A_{\phi}(f) = P_I(\phi f), \qquad f \in K_I^2 \cap L^{\infty},$$

where here $P_I : L^2 \to K_I^2$. In contrast to the Toeplitz operators on H^2 the operator A_{ϕ} may be extended to a bounded operator on K_I^2 even for some unbounded symbols.

We use the notation $\mathcal{T}(I)$ to denote the space of bounded truncated Toeplitz operators on K_I^2 and $\mathcal{T}_c(I)$ to denote the space of compact truncated Toeplitz operators on K_I^2 . Previous results in [4] identify the dual space of $K_I^1 \cap zH^1$ for a certain class of inner functions.

The results of Section 2 give an alternative description of the space dual to $K_I^1 \cap zH^1$, and furthermore this description is valid for all inner functions. Specifically, the description of the space dual to $K_I^1 \cap zH^1$ given in Section 2 initially realises $(K_I^1 \cap zH^1)^*$ as a quotient space and then we further show that $(K_I^1 \cap zH^1)^*$ may be realised as a space of analytic functions.

In Section 3 we will see that this realisation of $(K_I^1 \cap zH^1)^*$ lays the foundation for us to prove the main result of this paper, which is every bounded truncated Toeplitz operator on K_I^2 has a bounded symbol if and only if every compact truncated Toeplitz operator on K_I^2 has a symbol lying in $IC(\mathbb{T})$.

2 Duality results

In this section we describe the dual and predual of the space $K_I^1 \cap zH^1$. This description is a crucial tool needed in Section 2, where we study the symbols of truncated Toeplitz operators.

For not necessarily closed subspaces $M \subseteq X$ and $N \subseteq X^*$ of a Banach space X we define the annihilator

$$\mathcal{M}^{\perp} := \{ \ell \in X^* : \ell(x) = 0 \quad \forall x \in \mathcal{M} \}$$

and the pre-annihilator

$${}^{\perp}\mathcal{N} := \{ x \in X : \ell(x) = 0 \quad \forall \ell \in \mathcal{N} \}.$$

We note that with the above definitions the Hahn-Banach separation Theorem implies that $^{\perp}(M^{\perp})$ is the norm closure of M.

Lemma 2.1. $K_I^2 \cap zH^2$ is dense in $K_I^1 \cap zH^1$.

Proof. Clearly we have $I\overline{zH_0^2} \cap H^2 \subset I\overline{zH_0^1} \cap H^1$. If we show

$$(I\overline{zH_0^2}\cap H^2)^{\perp} \subseteq (I\overline{zH_0^1}\cap H^1)^{\perp},$$

then as ${}^{\perp}(M^{\perp})$ is the norm closure of M for a subspace M we obtain that $I\overline{zH_0^2} \cap H^2$ is dense in $I\overline{zH_0^1} \cap H^1$.

A result originally due to Fefferman-Stein [10], which can also be found as Theorem 2.2 in Chapter 9 of [1], shows that the dual space of H^1 is BMOA. By duality in the H^2 sense (which agrees with the H^1 -BMOA duality pairing) we have if $g \in (I\overline{z}H_0^2 \cap H^2)^{\perp}$ then $g \in BMOA$ and $\langle I\overline{z}H_0^2, g \rangle = 0$, which implies $g \in \frac{I}{z}H^2 \cap BMOA$ so $gz \in BMOA \cap IH^2$. If we define $G := \frac{gz}{I}$, then as the space H^1 is invariant by multiplication by I, we can deduce that BMOA (being the dual of H^1) is invariant under the Toeplitz operator $T_{\overline{I}}$ and so $G \in BMOA$. If $f \in I\overline{z}H_0^1 \cap H^1$ then $\frac{fz}{I} = \overline{F}$ for some $F \in H_0^1$, and if we denote \langle , \rangle to be the H^1 -BMOA duality pairing then we have

$$\langle f,g\rangle = \langle \frac{fz}{I}, \frac{gz}{I} \rangle = \langle \overline{F}, G \rangle = 0.$$

Thus, we have established $I\overline{zH_0^2} \cap H^2$ is dense in $I\overline{zH_0^1} \cap H^1$. Now clearly as multiplication by z is an isometry we have $K_I^2 \cap zH^2$ is dense in $K_I^1 \cap zH^1$.

Theorem 2.2. $l \in (K_I^1 \cap zH^1)^*$ if and only if there is a unique $v + Q \in L^{\infty}/Q$, where $Q = L^{\infty} \cap (H^2 + \overline{IH^2})$, such that

$$l(f) = l_{v+Q}(f) := \int_{\mathbb{T}} f(\zeta)v(\zeta) \, dm(\zeta).$$

Furthermore $||l_{v+Q}|| = ||v+Q||_{L^{\infty}/Q}$.

Proof. Clearly as $v \in L^{\infty}$ this will define a bounded linear functional. This functional is well defined as if v_2 is another representative of v then $v - v_2 \in Q$ but for all f in $K_I^2 \cap zH^2$ and any $q \in Q$ by orthogonality in the H^2 sense we have $\int_{\mathbb{T}} f(\zeta)q(\zeta) dm(\zeta) = 0$ and so by the above lemma we must indeed have $\int_{\mathbb{T}} f(\zeta)(v(\zeta) - v_2(\zeta)) dm(\zeta) = 0$ for all $f \in K_I^1 \cap zH^1$.

To prove the converse implication we must find the annihilator of $K_I^1 \cap z H^1$. The annihilator of $K_I^1 \cap z H^1$ (viewed as a subspace of L^1) is given by $(K_I^1 \cap z H^1)^{\perp} = \{b \in L^{\infty} : \int_{\mathbb{T}} f(\zeta)b(\zeta) dm(\zeta) = 0$ for all $f \in K_I^1 \cap z H^1\}$, but as a result of our previous lemma this is equal to $\{b \in L^{\infty} : \int_{\mathbb{T}} f(\zeta)b(\zeta) dm(\zeta) = 0$ for all $f \in K_I^2 \cap z H^2\}$.

We now argue that

$$\{b \in L^{\infty} : \int_{\mathbb{T}} f(\zeta)b(\zeta) \, dm(\zeta) = 0 \text{ for all } f \in K_I^2 \cap zH^2\} = Q := L^{\infty} \cap (H^2 + \overline{IH^2}).$$

A straightforward computation verifies the \supseteq inclusion. Before we show the \subseteq inclusion we first note that $f \in K_I^2 \cap zH^2$ if and only if $f \in K_I^2$ and f(0) = 0, which is equivalent to $f \in K_I^2$ and $f \perp k_0^I$, where $1 - \overline{I(0)}I(z) = k_0^I(z) \in K_I^2$ is the reproducing kernel at 0, (see Chapter 5 of [11] for details on the reproducing kernels for model spaces). Hence $K_I^2 \cap zH^2 = K_I^2 \ominus \mathbb{C}k_0^I$, giving $K_I^2 \ominus (K_I^2 \cap zH^2) = \mathbb{C}k_0^I$. Now to prove the \subseteq inclusion, take $b \in L^\infty$ such that $\langle f, \overline{b} \rangle_2 = 0$ for every $f \in K_I^2 \cap zH^2$. Decompose $b = g + \overline{I}\varphi_1 + \varphi_2$, with $g, \varphi_1 \in H^2$ and $\varphi_2 \in K_I^2$. Hence, for every $f \in K_I^2 \cap zH^2$, we have

$$0 = \langle f, \bar{g} \rangle_2 + \langle f, I\varphi_1 \rangle_2 + \langle f, \varphi_2 \rangle_2.$$

The first term $\langle f, \bar{g} \rangle_2$ is zero because $f \in zH^2$ and $g \in H^2$, the second term $\langle f, I\varphi_1 \rangle_2$ is also zero because $f \in K_I^2$ and $I\varphi_1 \in IH^2$. Thus we obtain that, for every $f \in K_I^2 \cap zH^2, \langle f, \varphi_2 \rangle_2 = 0$. That means that $\varphi_2 \in K_I^2 \ominus (K_I^2 \ominus zH^2)$, and as previously pointed out $K_I^2 \ominus (K_I^2 \cap zH^2) = \mathbb{C}k_0^I$, which means $\varphi_2 = c(1 - \overline{I(0)}I)$ for some $c \in \mathbb{C}$. Hence

$$b = g + \overline{I\varphi_1} + \overline{c(1 - \overline{I(0)}I)} = g + \overline{c} + \overline{I\left(\varphi_1 - c\overline{I(0)}\right)}.$$

We then deduce that $b \in L^{\infty} \cap \left(H^2 + \overline{IH^2}\right) = Q.$

Now given $l \in (K_I^1 \cap zH^1)^*$, the Hahn-Banach Theorem shows there is a unique v + Q where $v \in L^{\infty}$ such that $l(f) = l_{v+Q}(f) = \int_{\mathbb{T}} f(\zeta)v(\zeta) dm(\zeta)$ and furthermore that $||l_{v+Q}|| = ||v + Q||_{L^{\infty}/Q}$. We refer the reader to page 97 Theorem 4.9 of [18] for details.

One can also obtain a set theoretic description of $(K_I^1 \cap zH^1)^*$ which more so resembles the classical duality result for model spaces, which is $(K_I^p)^* = K_I^q$ where $1 and <math>\frac{1}{p} + \frac{1}{q} = 1$ (see Theorem 5.10.1 in [7]).

We let $P_{I,z}$ denote the orthogonal projection $L^2 \to K_I^2 \cap zH^2$ and we let $E : L^{\infty}/Q \to P_{I,z}(L^{\infty})$ be the well defined map $v + Q \mapsto P_{I,z}(\overline{v})$. If we equip $P_{I,z}(L^{\infty})$ with the norm $||P_{I,z}(f)|| := \inf\{||g||_{L^{\infty}} : P_{I,z}(g) = P_{I,z}(f)\}$ then E is an isometric antilinear isomorphism. Now as $P_{I,z}P = P_{I,z}$ and $P(L^{\infty}) = BMOA$ (see Theorem 3.5.11 in [7]), we may view E as a map from L^{∞}/Q to $P_{I,z}(BMOA)$. Furthermore, we can write $P_{I,z}(BMOA)$ as

$$\{k \in K_I^2 \cap zH^2 : \exists h \in H^2 \text{ and } \exists c \in K_I^2 \ominus (K_I^2 \cap zH^2) \text{ with } c+k+Ih \in \text{BMOA}\} := K.$$

Noting that, as in the proof of Theorem 2.2, we have that $K_I^2 \ominus (K_I^2 \cap zH^2) = \mathbb{C}k_0^I \subseteq H^{\infty} \subseteq BMOA$, this allows us to further write K as

$$\{k \in K_I^2 \cap zH^2 : \exists h \in H^2 \text{ with } k + Ih \in BMOA\}.$$

As mentioned previously, because the space H^1 is invariant by multiplication by I, and as BMOA is the dual space of H^1 (see Theorem 2.2 in Chapter 9 of [1]), we

can deduce that BMOA is invariant under the Toeplitz operator $T_{\overline{I}}$. So, if $k \in K$ is such that $k + Ih = b \in BMOA$ with $h \in H^2$, then $T_{\overline{I}}(k + Ih) = T_{\overline{I}}(b)$, and so as $k \in \ker T_{\overline{I}}$, we have $h = T_{\overline{I}}(b) \in BMOA$. Thus, we can in fact write K as

$$\{k \in K_I^2 \cap zH^2 : \exists h \in BMOA \text{ with } k + Ih \in BMOA\}.$$

The above description of K is clearly equal to span(BMOA, $I(BMOA)) \cap K_I^2 \cap zH^2$, so we conclude

$$P_{I,z}(L^{\infty}) = \text{span}(\text{BMOA}, I(\text{BMOA})) \cap K_I^2 \cap zH^2.$$

Thus via E and the above expression for $P_{I,z}(L^{\infty})$ we can isometrically identify $(K_I^1 \cap zH^1)^*$ as span(BMOA, $I(BMOA)) \cap K_I^2 \cap zH^2$, where for $k \in K$ we have $||k|| = \inf\{||g||_{L^{\infty}} : P_{I,z}(g) = k\}.$

Remark. We remark that adapting the above theorem and reasoning we can similarly deduce $(K_I^1)^* = P_I(BMOA) = \text{span}(BMOA, I(BMOA)) \cap K_I^2$ where for $k \in \text{span}(BMOA, I(BMOA)) \cap K_I^2$ we have $||k|| = \inf\{||g||_{L^{\infty}} \colon P_I(g) = k\}$.

We note that in general *IBMOA* $\not\subseteq$ BMOA. In fact, the conditions for when *IBMOA* \subseteq BMOA can be found as Theorem 1 in [9].

If one wants to consider the predual of the space $K_I^1 \cap zH^1$, we have the following result which may be found as Lemma 3.1 in [5].

Theorem 2.3. Let \mathcal{F}_I be the closure of the set $\overline{IH^{\infty}} + H^{\infty}$ in the weak * topology of the space L^{∞} . Then $(C(\mathbb{T})/(\mathcal{F}_I \cap C(\mathbb{T})))^* = K_I^1 \cap zH^1$.

Specifically, in the above theorem there is an isometric isomorphism given by $h \mapsto l_h$ where $h \in K_I^1 \cap zH^1$ and $l_h \in (C(\mathbb{T})/(\mathcal{F}_I \cap C(\mathbb{T})))^*$ is defined by $l_h(\phi) = \int_{\mathbb{T}} \phi h \, dm$ where ϕ is any representative of an element in $(C(\mathbb{T})/(\mathcal{F}_I \cap C(\mathbb{T})))$.

3 Application to truncated Toeplitz operators

In this section after giving a brief overview on some of the previously known results about the symbols of truncated Toeplitz operators, use our duality results from our previous section to deduce the main result of this paper, which is Theorem 3.2.

The question of whether every bounded truncated Toeplitz operator has a bounded symbol is an interesting one. This question has led to much research activity within the community with many questions being answered and many new questions being posed. Here we give the reader a brief background on this topic. In Sarason's seminal work of 2007 [19] he initiated a systematic study of truncated Toeplitz operators with symbols in L^2 . In this paper, one of the most natural questions posed was whether every bounded truncated Toeplitz operator has a bounded symbol. This question was then shown to be negative in 2009 (see [3]). In fact, the authors actually constructed a bounded rank one truncated Toeplitz operator which was shown to have no bounded symbol. To build on this work, in [2] the authors gave two conditions on the inner function, I, which are equivalent to every bounded truncated Toeplitz operator on K_I^2 having a bounded symbol. (See Theorem 3.1 below.)

Motivated by these findings, a similar study in to the symbols of compact truncated Toeplitz operators was initiated. Section 5 of [6] gives an overview of many results in this area. In particular, the role played by bounded symbols in the case of bounded truncated Toeplitz operators seems to be replaced by symbols of the form $IC(\mathbb{T})$ when we are considering compact truncated Toeplitz operators. Specifically, Proposition 5.4 of [6] shows that if $\phi \in IC(\mathbb{T})$ then A_{ϕ}^{I} is compact, however using the example of the rank one truncated Toeplitz operator which does not possess a bounded symbol, we know there exists a compact truncated Toeplitz operator without a symbol in $IC(\mathbb{T})$, as trivially $IC(\mathbb{T}) \subseteq L^{\infty}$. One question posed in [6] was whether there was a compact truncated Toeplitz operator with a symbol in $IC(\mathbb{T}) + IH^{\infty}$ that has no continuous symbol. This question was then answered affirmatively in [14] when a compact truncated Toeplitz operator with this property was then constructed.

Following the results in [2] one may suspect that there are conditions on the inner function I which are equivalent to every compact truncated Toeplitz operator on K_I^2 having a symbol in $IC(\mathbb{T})$. We may further suspect that these conditions may be similar in nature to the conditions on the inner function I which are equivalent to every bounded truncated Toeplitz operator on K_I^2 having a bounded symbol. In fact, in this section we will prove that every compact truncated Toeplitz operator on K_I^2 has a symbol in $IC(\mathbb{T})$ if and only if every bounded truncated Toeplitz operator on K_I^2 has a bounded symbol. We show this with Theorem 3.2 below.

In the following we define $\mathcal{C}_p(I)$ to be the set of all finite complex Borel measures μ on the unit circle such that the embedding $K_I^p \to L^p(|\mu|)$ is continuous.

Theorem 3.1. [2]

The following are equivalent:

- 1. any bounded truncated Toeplitz operator on K_I^2 admits a bounded symbol;
- 2. $C_1(I^2) = C_2(I^2);$
- 3. for any $f \in H^1 \cap I^2 \overline{zH_0^1}$ there exists $x_i, y_i \in K_I^2$ with $\sum_i ||x_i||_2 ||y_i||_2 < \infty$ such that $f = \sum_i x_i y_i$.

The inner function I is said to be one-component if and only if there exists an $\eta \in (0, 1)$ such that

$$\{z \in \mathbb{D} : |I(z)| < \eta\}$$

is connected. We remark that by Corollary 2.5 in [2] the equivalent conditions of the theorem below are fulfilled when I is a one component inner function.

Throughout the rest of the paper we will focus on proving the following theorem.

Theorem 3.2. The equivalent conditions of Theorem 3.1 are satisfied if and only if any compact truncated Toeplitz operator on K_I^2 has a symbol in $IC(\mathbb{T})$.

We note that the forward implication of the above theorem has already been noticed. Indeed, as noted in Section 3.2 of [6], we have $C_2(I^2) = C_2(I)$, and thus in view of Theorem 3.1, Theorem 5.2 in [6] states if every bounded truncated Toeplitz operator on K_I^2 has a bounded symbol then every compact truncated Toeplitz operator on K_I^2 is of the form $A_{I\phi}$ where $\phi \in C(\mathbb{T})$.

The main theorem of this paper is the backwards implication of the above theorem, that is, if any compact truncated Toeplitz operator on K_I^2 has a symbol in $IC(\mathbb{T})$, then the equivalent conditions of Theorem 3.1 are satisfied. As the workings for the proof of the backwards implication of Theorem 3.2 are quite long and detailed we first outline the idea of the proof for the reader.

We first consider the map $L: X \to K^1_{I^2} \cap zH^1$, given by $f \mapsto If$ (where X is defined as in (1) below). Then using the results of Section 1 as well as Theorems 3.6 and 3.3, which identify the predual and dual space of X as $\mathcal{T}_c(I)$ and $\mathcal{T}(I)$ respectively, we can describe the pre-adjoint, *L, and adjoint, L^* , of L,

where $Q_2 = L^{\infty} \cap (H^2 + \overline{I^2 H^2})$. The above description of L^* is given by Theorem 3.4 and the description of *L is given by Proposition 3.7.

Then for an inner function I, by Proposition 3.8 we notice that any compact truncated Toeplitz operator on K_I^2 having a symbol in $IC(\mathbb{T})$, is actually equivalent to L being isomorphic. Then when L is isomorphic, L must also be isomorphic and in particular the image of L^* must be equal to $\mathcal{T}(I)$. Then under the assumption Lis surjective, by Corollary 3.5 (which shows the image of L^* is all truncated Toeplitz operators on K_I^2 with a bounded symbol) we are able to deduce that every bounded truncated Toeplitz operator on K_I^2 has a bounded symbol.

Following the results of [2], we define the Banach spaces

$$X = \{ \sum x_i \overline{y_i} : x_i, y_i \in K_I^2, \sum ||x_i||_2 ||y_i||_2 < \infty \},$$
(1)

and

$$X_a = \{ \sum x_i y_i : x_i, y_i \in K_I^2, \sum ||x_i||_2 ||y_i||_2 < \infty \}.$$

The norm in the space of X and X_a is defined as the infimum of $\sum ||x_i||_2 ||y_i||_2$ over all possible representations. We note there is an isometric isomorphism from X to X_a given by

$$f \mapsto \overline{z}If,$$
 (2)

and one can also show that the forward shift from $X_a \to K_{I^2}^1 \cap zH^1$ is bounded. The following theorem can be found as Theorem 2.3 in [2].

Theorem 3.3. The dual space of X can be naturally identified with $\mathcal{T}(I)$. Namely, continuous linear functionals over X are of the form

$$\Phi_A(f) = \sum_i \langle Ax_i, y_i \rangle, \qquad f = \sum_i x_i \overline{y_i} \in X,$$

with $A \in \mathcal{T}(I)$, and the correspondence between X and $\mathcal{T}(I)$ is one to one and isometric.

We define a bounded linear map

$$L: X \to K^1_{I^2} \cap zH^1,$$

given by

$$f \mapsto If.$$

Now taking into account the results of Section 1 and Theorem 3.3, when considering the adjoint, L^* , of L we obtain a bounded map

$$L^*: L^\infty/Q_2 \to \mathcal{T}(I),$$

where $Q_2 = L^{\infty} \cap (H^2 + \overline{I^2 H^2})$ Explicitly, L^* is the unique map satisfying

$$\langle L(k), g + Q_2 \rangle = \langle k, L^*(g + Q_2) \rangle, \tag{3}$$

for each $k \in X$ and $g + Q_2 \in L^{\infty}/Q_2$ (here the duality pairings, denoted \langle , \rangle , are given by Theorem 2.2 and 3.3 respectively). If we denote A to be the truncated Toeplitz operator $L^*(g + Q_2)$, then for $k = x\overline{y}$ with $x \in K_I^{\infty}, y \in K_I^2$ equating both sides of equation (3) gives us

$$\int_{\mathbb{T}} x \overline{y} Ig \ dm = \int_{\mathbb{T}} A(x) \overline{y} \ dm,$$

and so as $y \in K_I^2$, and by orthogonality we know $\langle f, y \rangle_2 = \langle P_I(f), y \rangle_2$ for each $f \in L^2$ we have

$$\int_{\mathbb{T}} (A(x) - xIg)\overline{y} \ dm = \int_{\mathbb{T}} (A(x) - P_I(xIg))\overline{y} \ dm = 0.$$

Now by density of K_I^{∞} in K_I^2 (see Section 2.5 of [12]) and non-degeneracy of the integral we can deduce

$$A = A_{Ig}^I$$

We conclude the following result;

Theorem 3.4. There is a bounded linear map $L^* : L^{\infty}/Q_2 \to \mathcal{T}(I)$, given by

$$g + Q_2 \mapsto A_{Ig}.\tag{4}$$

Corollary 3.5. The image of L^* is exactly all elements of $\mathcal{T}(I)$ which possess a bounded symbol.

Proof. Clearly the image of L^* is contained the set of all bounded truncated Toeplitz operators with a bounded symbol. Conversely, if A_{ϕ}^I has a bounded symbol g then $\overline{I}g \in L^{\infty}$, so $\overline{I}g + Q_2 \in L^{\infty}/Q_2$ and maps to A_{ϕ}^I through L^* .

In the case when every bounded truncated Toeplitz operator has a bounded symbol the map L^* is isomorphic, and so the norm of A_{Ig} is equivalent to $||g + Q_2||_{L^{\infty}/Q_2}$.

We now seek to describe the pre-adjoint of L. The second part of Theorem 2.3 in [2] states the following.

Theorem 3.6. The dual space of all compact truncated Toeplitz operators can be identified with X, via the duality pairing $\sum_i x_i \overline{y_i} \mapsto L_{\sum_i x_i \overline{y_i}}$, where

$$L_{\sum_{i} x_{i} \overline{y_{i}}}(A) = \sum_{i} \langle Ax_{i}, y_{i} \rangle,$$

for each compact truncated Toeplitz operator A. Furthermore the duality pairing $\sum_i x_i \overline{y_i} \mapsto L_{\sum_i x_i \overline{y_i}}$ is one-to-one and isometric map between X and $(\mathcal{T}_c(I))^*$.

In general the pre-adjoint of a bounded linear map may not exist. Nonetheless we may define the map $^*L: C(\mathbb{T})/(\mathcal{F}_{I^2} \cap C(\mathbb{T})) \to \mathcal{T}_c(I)$, where

$$g + (\mathcal{F}_{I^2} \cap C(\mathbb{T})) \mapsto A_{Ig}.$$

Proposition 3.7. The map *L is a well defined bounded, linear, injective map and $(*L)^* = L$ (i.e. *L is the pre-adjoint of L).

Proof. The map ${}^{*}L$ is clearly linear. Proposition 5.4 in [6] ensures that the specified A_{Ig} is indeed a compact truncated Toeplitz operator. Noting that $A_{\phi}^{I} = 0$ if and only if $\phi \in \overline{IH^2} + IH^2$ (see Section 3 of [19]), we see that the map is well defined and injective. By non-uniqueness of the symbol we have that

$$|A_{Ig}|| \leq \inf\{\|\overline{Ih_1} + Ig + Ih_2\|_{L^{\infty}}\} = \inf\{\|\overline{I^2h_1} + g + h_2\|_{L^{\infty}}\}\$$

taken over all $\overline{I^2h_1} + h_2 \in (\mathcal{F}_{I^2} \cap C(\mathbb{T}))$. We see the above expression is actually equal to $\|g + (\mathcal{F}_{I^2} \cap C(\mathbb{T}))\|$, and thus *L is bounded

We now argue $({}^{*}L)^{*} = L$. Proposition 4.1 in [2] states that every element of X can be expressed as a sum of four elements of the form $x\overline{y}$ for $x, y \in K_{I}^{2}$. So for an arbitrary element $\sum_{i=1}^{4} x_{i}\overline{y_{i}} \in X$, we know $({}^{*}L)^{*}$ is the linear map satisfying

$$\langle *L(g + (\mathcal{F}_{I^2} \cap C(\mathbb{T}))), \sum_{i=1}^4 x_i \overline{y_i} \rangle = \langle g + (\mathcal{F}_{I^2} \cap C(\mathbb{T})), (*L)^* (\sum_{i=1}^4 x_i \overline{y_i}) \rangle,$$

for every $g + (\mathcal{F}_{I^2} \cap C(\mathbb{T})) \in C(\mathbb{T})/(\mathcal{F}_{I^2} \cap C(\mathbb{T}))$ and every $\sum_{i=1}^4 x_i \overline{y_i} \in X$. Here the duality pairing on the left hand side is understood by the duality described in Theorem 3.6 and on the right hand side the duality is described by Theorem 2.3. Explicitly, this means $(*L)^*$ is a linear map satisfying

$$\int_{\mathbb{T}} Ig \sum_{i=1}^{4} x_i \overline{y_i} \, dm = \int_{\mathbb{T}} g(^*L)^* (\sum_{i=1}^{4} x_i \overline{y_i}) \, dm, \tag{5}$$

i.e.

$$\int_{\mathbb{T}} g \underbrace{\left(I \sum_{i=1}^{4} x_i \overline{y_i} - (^*L)^* (\sum_{i=1}^{4} x_i \overline{y_i}) \right)}_{:=u} dm = 0 \tag{6}$$

for all $g + (\mathcal{F}_{I^2} \cap C(\mathbb{T})) \in C(\mathbb{T})/(\mathcal{F}_{I^2} \cap C(\mathbb{T}))$. Now, u defined as above lies in $K^1_{I^2} \cap zH^1$, so by Theorem 2.3 the map l_u , where $l_u(g + (\mathcal{F}_{I^2} \cap C(\mathbb{T}))) \mapsto \int_{\mathbb{T}} gu \, dm$ defines a continuous linear functional on $C(\mathbb{T})/(\mathcal{F}_{I^2} \cap C(\mathbb{T}))$. So when (6) holds we must have l_u is the zero functional, however by Theorem 2.3 we know $u \mapsto l_u$ is an isometric isomorphism and so we must have u = 0 and so $(^*L)^*(\sum_{i=1}^4 x_i\overline{y_i}) = I\sum_{i=1}^4 x_i\overline{y_i}$ and therefore $(^*L)^* = L$.

We can make a result which is analogous to Corollary 3.5 but in the case of continuous symbols.

Proposition 3.8. The image of *L is all truncated Toeplitz operators of the form A_{ϕ} where $\phi \in IC(\mathbb{T})$

As the map L is clearly injective, L must have dense range and we can make the following corollary which has been previously noticed in the proof of Lemma 3.5 in [5].

Corollary 3.9. Truncated Toeplitz operators of the form A_{ϕ} where $\phi \in IC(\mathbb{T})$ are dense in $\mathcal{T}_c(I)$.

We now can prove our main result, which is a one way implication of Theorem 3.2.

Theorem 3.10. If every compact truncated Toeplitz operator on K_I^2 is of the form $A_{I\phi}$ where $\phi \in C(\mathbb{T})$ then every bounded truncated Toeplitz operator has a bounded symbol.

Proof. If every compact truncated Toeplitz operator on K_I^2 is of the form $A_{I\phi}$ where $\phi \in C(\mathbb{T})$ then by Proposition 3.8 we know that *L is surjective (and hence isomorphic). Now by Proposition 3.7 we must also have that $(*L)^* = L$ is isomorphic, and hence L^* is isomorphic. Now by Corollary 3.5 this must mean every bounded truncated Toeplitz operator on K_I^2 has a bounded symbol.

Remark. Examining the above proof we can obtain a concise alternate proof of a result already known from Theorem 3.1, which is L is isomorphic (or equivalently condition 3 holds in Theorem 3.1) implies every bounded truncated Toeplitz operator on K_I^2 has a bounded symbol.

We now easily state the proof of Theorem 3.2.

Proof of Theorem 3.2. The forward implication is already known by previous results (see the reasoning laid out after the statement of Theorem 3.2) and the backwards implication is Theorem 3.10.

A long standing open conjecture regarding symbols of bounded truncated Toeplitz operators is the following.

Conjecture 3.1. Every bounded truncated Toeplitz operator on K_I^2 has a bounded symbol if and only if I is one-component.

We can now state an equivalent formulation of the above conjecture, which is the following.

Conjecture 3.2. Every compact truncated Toeplitz operator on K_I^2 has a symbol in $IC(\mathbb{T})$ if and only if I is one-component.

Acknowledgements

The author is grateful to EPSRC for financial support. The author would like to thank Professor Partington for his guidance. The author is grateful to the referee for their valuable comments.

Declarations

Declarations of interest: none.

Competing interests: none.

This research was supported by the EPSRC (grant number 1972662).

The sponsor (EPSRC) had no role in study design; in the collection, analysis and interpretation of data or in writing of the report; and in the decision to submit the article for publication.

References

- [1] M. Andersson. Topics in complex analysis. Springer New York, 1997.
- [2] A. Baranov, R. Bessonov, and V. Kapustin. Symbols of truncated Toeplitz operators. J. Funct. Anal., 261(12):3437–3456, 2011.
- [3] A. Baranov, I. Chalendar, E. Fricain, J. Mashreghi, and D. Timotin. Bounded symbols and reproducing kernel thesis for truncated Toeplitz operators. *Journal* of Functional Analysis, 259(10):2673–2701, 2010.
- [4] R. V. Bessonov. Duality theorems for coinvariant subspaces of H¹. Adv. Math., 271:62–90, 2015.
- [5] R. V. Bessonov. Fredholmness and compactness of truncated Toeplitz and Hankel operators. Integral Equations Operator Theory, 82(4):451–467, 2015.
- [6] I. Chalendar, E. Fricain, and D. Timotin. A survey of some recent results on truncated Toeplitz operators. In *Recent progress on operator theory and approximation in spaces of analytic functions*, volume 679 of *Contemp. Math.*, pages 59–77. Amer. Math. Soc., Providence, RI, 2016.
- [7] J. A. Cima and W. T. Ross. The backward shift on the Hardy space. American Mathematical Soc., 2000.
- [8] P. L. Duren. Theory of H^p spaces. Pure Appl. Math, 38:74, 1970.
- K. M. Dyakonov. Division and multiplication by inner functions and embedding theorems for star-invariant subspaces. *American Journal of Mathematics*, 115(4):881–902, 1993.
- [10] C. Fefferman and E. M. Stein. H^p spaces of several variables. Acta mathematica, 129:137–193, 1972.
- [11] S. R. Garcia, J. Mashreghi, and W. T. Ross. Introduction to model spaces and their operators, volume 148 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2016.
- [12] S.R Garcia and W.T. Ross. Recent progress on truncated Toeplitz operators. Blaschke Products and Their Applications, 65:265–319, 2013.

- [13] D. Girela. Analytic functions of bounded mean oscillation. In Complex Function Spaces (Mekrijärvi, 1999) Univ. Joensuu Dept. Math. Rep. Ser.4, pages 61–170. Univ. Joensuu, Joensuu, 2001.
- [14] P. Gorkin and J. R. Partington. Norms of truncated Toeplitz operators and numerical radii of restricted shifts. *Comput. Methods Funct. Theory*, 19(3):487– 508, 2019.
- [15] N. K. Nikolski. Operators, Functions, and Systems-An Easy Reading: Hardy, Hankel, and Toeplitz, volume 1. American Mathematical Soc., 2002.
- [16] R. O'Loughlin. Matrix-valued truncated Toeplitz operators: unbounded symbols, kernels and equivalence after extension. arXiv preprint arXiv:2012.00654, 2020.
- [17] R. O'Loughlin. Multidimensional Toeplitz and truncated Toeplitz operators. https://etheses.whiterose.ac.uk/29284/, University of Leeds, 2021.
- [18] Walter Rudin. Functional analysis. International Series in Pure and Applied Mathematics. McGraw-Hill, Inc., New York, second edition, 1991.
- [19] D. Sarason. Algebraic properties of truncated Toeplitz operators. Oper. Matrices, 1(4):491–526, 2007.