

Strain Tensors and Matching Property on Degenerated Hyperbolic Surfaces

Liang-Biao Chen and Peng-Fei Yao

Key Laboratory of Systems and Control

Institute of Systems Science, Academy of Mathematics and Systems Science

Chinese Academy of Sciences, Beijing 100190, P. R. China

School of Mathematical Sciences

University of Chinese Academy of Sciences, Beijing 100049, China

e-mail: pfyao@iss.ac.cn

Abstract We prove the regularity of solutions to the strain tensor equation on degenerated hyperbolic surfaces S where the Gauss curvature is zero on a part of boundary. Furthermore, we obtain the density property that smooth infinitesimal isometries are dense in the $W^{2,2}(S, \mathbb{R}^3)$ infinitesimal isometries. Finally, the matching property is established. Those results are important tools in obtaining recovery sequences (Γ -lim sup inequality) for dimensionally-reduced shell theories in elasticity.

Keywords shell, nonlinear elasticity, Riemannian geometry, tensor analysis

Mathematics Subject Classifications (2010) 74K20(primary), 74B20(secondary).

1 Introduction and Main Results

Let $M \subset \mathbb{R}^3$ be a surface with a normal \vec{n} and let the middle surface of a shell be an open set $S \subset M$. Let $T^k S$ denote all the k -order tensor fields on S for an integer $k \geq 0$. Let $T_{\text{sym}}^2 S$ be all the 2-order symmetrical tensor fields on S . For $y \in W^{1,2}(S, \mathbb{R}^3)$, we decompose it into $y = W + w\vec{n}$, where $w = \langle y, \vec{n} \rangle$ and $W \in TS$. For $U \in T_{\text{sym}}^2 S$ given, linear strain tensor of a displacement $y \in W^{1,2}(S, \mathbb{R}^3)$ of the middle surface S takes the form

$$\text{sym } DW + w\Pi = U \quad \text{for } x \in S, \quad (1.1)$$

This work is supported by the National Science Foundation of China, grants no. 12071463 and Key Research Program of Frontier Sciences, CAS, no. QYZDJ-SSW-SYS011.

where D is the connection of the induced metric in M , $2\text{sym } DW = DW + D^T W$, and Π is the second fundamental form of M . Equation (1.1) plays a fundamental role in the theory of thin shells, see [2, 4, 3, 5, 7, 9] and many others. When $U = 0$, a solution y to (1.1) is referred to as an *infinitesimal isometry*.

The type of equation (1.1) depends on the sign of the curvature on the region S : It is elliptic if S has positive curvature; it is parabolic if the curvature is zero but $\Pi \neq 0$ on S ; it is hyperbolic if S has negative curvature.

Here we establish the regularity of solutions to (1.1) when S is a non-characteristic region where its curvature is zero on a part of boundary, that will be specified below. Then it is proved that smooth infinitesimal isometries are dense in the $W^{2,2}(S, \mathbb{R}^3)$ infinitesimal isometries on the region S . Finally, the matching property is derived that smooth enough infinitesimal isometries can be matched with higher order infinitesimal isometries. Those results are important tools in obtaining recovery sequences (Γ -lim sup inequality) for dimensionally-reduced shell theories in elasticity, when the elastic energy density scales like h^β , $\beta \in (2, 4)$, that is, intermediate regime between pure bending ($\beta = 2$) and the von-Kármán regime ($\beta = 4$). Such results have been obtained for elliptic surfaces [5], developable surfaces [2], and hyperbolic surfaces [7, 9]. A survey on this topic is presented in [3].

In this paper we study the degenerated hyperbolic equation (1.1), which is equivalent to a degenerated hyperbolic scalar equation of the form

$$\langle D^2 w, Q^* \Pi \rangle + \frac{1}{\kappa} \langle Dw, X_0 \rangle + \kappa (\text{tr}_g \Pi) w = \kappa f + \frac{1}{\kappa} \langle X_0, F \rangle + \langle DF, Q^* \Pi \rangle, \quad (1.2)$$

for $x \in S$, $\kappa \neq 0$, where $w \in L^2(S)$ is the unknown, $f \in L^2(S)$ and $F \in L^2(S, TS)$ are given, and κ is the Gauss curvature. When $\kappa = 0$, there are two terms to degenerate in (1.2): The coefficient of $\langle Dw - F, X_0 \rangle$ becomes infinite; the second derivative of w along the direction of the nonzero principal curvature is zero in $\langle D^2 w, Q^* \Pi \rangle$. Those situations challenge the analysis of (1.2).

Here we employ the Bochner technique and the tensor analysis to cope with the degenerates in (1.2), where some priori estimates near the zero curvature curve in Section 2 play the key role.

We state our main results as follows. Let $S \subset M$ be given by

$$S = \{ \alpha(t, s) \mid (t, s) \in [0, a) \times (0, b) \}, \quad a > 0, b > 0,$$

where $\alpha : [0, a) \times [0, b] \rightarrow M$ is an imbedding map which is a family of regular curves with two parameters t, s such that

$$\Pi(\alpha_t(t, s), \alpha_t(t, s)) > 0 \quad \text{for all } (t, s) \in [0, a) \times [0, b], \quad (1.3)$$

where $\alpha(\cdot, s)$ is a closed curve with period a for each $s \in [0, b]$ and

$$\{ \alpha(t, 0) \mid t \in [0, a) \}$$

is a closed curve or just one point.

Curvature assumptions Let κ be the Gaussian curvature function on M . We assume that S satisfies the following curvature conditions:

$$\kappa(x) < 0 \quad \text{for } x \in S \cup \Gamma_0; \quad (1.4)$$

$$\kappa = 0, \quad D\kappa(x) \neq 0 \quad \text{for } x \in \Gamma_b, \quad (1.5)$$

where

$$\Gamma_b = \{ \alpha(t, b) \mid t \in [0, a] \}, \quad \Gamma_0 = \{ \alpha(t, 0) \mid t \in [0, a] \}.$$

Our main results are the following.

Theorem 1.1. *Let S be of class $C^{2,1}$. For $U \in W^{2,2}(S, T_{\text{sym}}^2 S)$, there exists a solution $y = W + w\vec{n} \in W^{1,2}(S, \mathbb{R}^3)$ to equation (1.1) satisfying the bounds*

$$\|W\|_{W^{2,1}(S, TS)}^2 + \|w\|_{W^{1,2}(S)}^2 \leq C \|U\|_{W^{2,2}(S, T_{\text{sym}}^2 S)}^2. \quad (1.6)$$

If, in addition, $S \in C^{m+1,1}$, $U \in W^{m+1,2}(S, T_{\text{sym}}^2 S)$ for some $m \geq 2$, then

$$\|W\|_{W^{m+1,2}(S, TS)}^2 + \|w\|_{W^{m,2}(S)}^2 \leq C \|U\|_{W^{m+1,2}(S, T_{\text{sym}}^2 S)}^2.$$

By the imbedding theorem [1, P. 158], the following corollary is immediate.

Corollary 1.1. *Let $m \geq 0$ be an integer and let S be of $S \in C^{m+2,1}$. Then problem (1.1) admits a solution $y = W + w\vec{n} \in C_B^m(S, \mathbb{R}^3)$ satisfying*

$$\|W\|_{C_B^{m+1}(S, TS)} + \|w\|_{C_B^m(S)} \leq C \|U\|_{C_B^{m+3}(S, T_{\text{sym}}^2 S)},$$

where

$$C_B^m(S, \mathbb{R}^3) = \{ y \in C^m(S, \mathbb{R}^3) \mid D^\alpha y \in L^\infty(S, \mathbb{R}^3) \text{ for } |\alpha| \leq m \}.$$

For $y \in W^{1,2}(S, \mathbb{R}^3)$, we denote the left hand side of equation (1.1) by $\text{sym } \nabla y$. Let

$$\mathcal{V}(S, \mathbb{R}^3) = \{ y \in W^{2,2}(S, \mathbb{R}^3) \mid \text{sym } \nabla y = 0 \}.$$

Theorem 1.2. *Let S be of class $C^{m+3,1}$ for some integer $m \geq 0$. Then, for every $y \in \mathcal{V}(S, \mathbb{R}^3)$ there exists a sequence $\{y_k\} \subset \mathcal{V}(S, \mathbb{R}^3) \cap C_B^m(S, \mathbb{R}^3)$ such that*

$$\lim_{k \rightarrow \infty} \|y - y_k\|_{W^{2,2}(S, \mathbb{R}^3)} = 0.$$

A one parameter family $\{y_\varepsilon\}_{\varepsilon > 0} \subset C_B^1(\overline{S}, \mathbb{R}^3)$ is said to be a (generalized) m th order infinitesimal isometry if the change of metric induced by y_ε is of order ε^{m+1} , that is,

$$\|\nabla^T y_\varepsilon \nabla y_\varepsilon - g\|_{L^\infty(S, T^2)} = \mathcal{O}(\varepsilon^{m+1}) \quad \text{as } \varepsilon \rightarrow 0,$$

where g is the induced metric of M from \mathbb{R}^3 , see [2]. A given m th order infinitesimal isometry can be modified by higher order corrections to yield an infinitesimal isometry of order $m_1 > m$, a property to which we refer to by *matching property of infinitesimal isometries*, [2, 5]. This property plays an important role in the construction of a recover sequence in the Γ -limit for thin shells.

Theorem 1.3. *Let S be of class $C^{4m,1}$. Given $y \in \mathcal{V}(S, \mathbb{R}^3) \cap C_B^{4m-2}(S, \mathbb{R}^3)$, there exists a family $\{z_\varepsilon\}_{\varepsilon>0} \subset C_B^2(S, \mathbb{R}^3)$, equi-bounded in $C_B^2(S, \mathbb{R}^3)$, such that for all small $\varepsilon > 0$ the family:*

$$y_\varepsilon = \text{id} + \varepsilon y + \varepsilon^2 z_\varepsilon$$

is a m th order infinitesimal isometry of class $C_B^2(S, \mathbb{R}^3)$.

2 Some priori estimates near the zero curvature curve

Let ∇ and D denote the connection of \mathbb{R}^3 in the Euclidean metric and the one of M in the reduced metric, respectively. We have to treat the relationship between ∇ and D carefully.

Let $m \geq 1$ be an integer. Let $T \in T^m M$ be a m th order tensor field on M . We define a $m - 1$ th order tensor field by

$$i_Y T(Y_1, \dots, Y_{m-1}) = T(Y, Y_1, \dots, Y_{m-1}) \quad \text{for } Y_1, \dots, Y_{m-1} \in TM,$$

which is called an *inner product* of T with Y . For any $T \in T^2 S$ and $\alpha \in T_x M$,

$$\text{tr}_g i_\alpha DT$$

is a linear functional on $T_x M$, where $\text{tr}_g i_\alpha DT$ is the trace of the 2-order tensor field $i_\alpha DT$ in the induced metric g . Thus there is a vector, denoted by $\text{div}_g T$, such that

$$\langle \text{div}_g T, \alpha \rangle = \text{tr}_g i_\alpha DT \quad \text{for } \alpha \in T_x M, x \in M.$$

Clearly, the above formula defines a vector field $\text{div}_g T \in TM$.

We need a linear operator Q ([7], [9]) as follows. For each point $p \in M$, the Riesz representation theorem implies that there exists an isomorphism $Q : T_p M \rightarrow T_p M$ such that

$$\langle \alpha, Q\beta \rangle = \det(\alpha, \beta, \vec{n}(p)) \quad \text{for } \alpha, \beta \in T_p M. \quad (2.1)$$

Let e_1, e_2 be an orthonormal basis of $T_p M$ with positive orientation, that is,

$$\det(e_1, e_2, \vec{n}(p)) = 1.$$

Then Q can be expressed explicitly by

$$Q\alpha = \langle \alpha, e_2 \rangle e_1 - \langle \alpha, e_1 \rangle e_2 \quad \text{for all } \alpha \in T_p M. \quad (2.2)$$

Clearly, Q satisfies

$$Q^T = -Q, \quad Q^2 = -\text{Id}.$$

The operator Q plays an important role in our analysis.

In the present section, we consider problem

$$\begin{cases} \operatorname{div}_g Q \nabla \vec{n} V = f_1, \\ \operatorname{div}_g V = f_2, \end{cases} \quad (2.3)$$

where $V \in TS$ and f_i are functions on S .

For given $\varepsilon > 0$, set

$$S_{b-\varepsilon} = \{ \alpha(t, s) \mid (t, s) \in [0, a) \times (b - \varepsilon, b) \}, \quad \Gamma_{b-\varepsilon} = \{ \alpha(t, b - \varepsilon) \mid t \in [0, a) \}.$$

The main results of this section are the following.

Theorem 2.1. *Let $S_{b-\varepsilon}$ be of class $C^{3,1}$. For given $\varepsilon > 0$ small, there is $\sigma_\varepsilon > 0$ such that for any solution $V \in TS_{b-\varepsilon}$ to (2.3), the following estimates hold true.*

$$\sigma_\varepsilon \|V\|_{L^2(S_{b-\varepsilon}, TS_{b-\varepsilon})}^2 \leq \|f_1\|_{L^2(S_{b-\varepsilon})}^2 + \|f_2\|_{L^2(S_{b-\varepsilon})}^2 + \|V\|_{L^2(\Gamma_{b-\varepsilon}, TM)}^2, \quad (2.4)$$

$$\begin{aligned} \sigma_\varepsilon \|V\|_{W^{1,2}(S_{b-\varepsilon}, TS_{b-\varepsilon})}^2 &\leq \|f_1\|_{W^{1,2}(S_{b-\varepsilon})}^2 + \|f_2\|_{W^{1,2}(S_{b-\varepsilon})}^2 + \|i_\nu DV\|_{L^2(\Gamma_b)}^2 \\ &\quad + \|V\|_{L^2(\Gamma_\varepsilon \cup \Gamma_b, TM)}^2 + \|DV\|_{L^2(\Gamma_{-\varepsilon}, T\Gamma_{-\varepsilon})}^2, \end{aligned} \quad (2.5)$$

where ν is the outside normal of $S_{b-\varepsilon}$.

We make some preparations first. The proof of Theorem 2.1 will be given in the end of this section.

Lemma 2.1. *We have*

$$QD_X Y = D_X(QY) \quad \text{for } X, Y \in TM. \quad (2.6)$$

Proof. It follows from (2.1) that

$$\begin{aligned} \langle Z, QD_X Y \rangle &= \det(Z, \nabla_X Y, \vec{n}) \\ &= X \det(Z, Y, \vec{n}) - \det(\nabla_X Z, Y, \vec{n}) - \det(Z, Y, \nabla_X \vec{n}) \\ &= X \langle Z, QY \rangle - \langle \nabla_X Z, QY \rangle = \langle Z, \nabla_X(QY) \rangle \quad \text{for } X, Y, Z \in TM, \end{aligned} \quad (2.7)$$

this gives (2.6). □

Lemma 2.2. *The following formulas are true.*

$$\langle X, Y \rangle Z = \langle Z, Y \rangle X + \langle Z, QX \rangle QY \quad \text{for } X, Y, Z \in TS. \quad (2.8)$$

Proof. Let $p \in S$ be given. If $Y = 0$, then (2.8) holds. We assume that $|Y| = 1$. Then QY , Y forms an orthonormal basis of T_pS . Thus

$$\langle\langle Z, Y \rangle X + \langle Z, QX \rangle QY, Y \rangle = \langle\langle Z, Y \rangle X, Y \rangle = \langle\langle X, Y \rangle Z, Y \rangle,$$

$$\begin{aligned} \langle\langle Z, Y \rangle X + \langle Z, QX \rangle QY, QY \rangle &= \langle Z, Y \rangle \langle X, QY \rangle + \langle Z, QX \rangle \\ &= \langle Z, Y \rangle \langle X, QY \rangle + \langle Z, Y \rangle \langle Y, QX \rangle + \langle Z, QY \rangle \langle QY, QX \rangle \\ &= \langle\langle X, Y \rangle Z, QY \rangle. \end{aligned}$$

Thus (2.8) follows. \square

Lemma 2.3. *Let $P \in T^2S$. Let X and Y be vector fields and f be a function. Then*

$$\operatorname{div}_g(PX) = \langle P, DX \rangle + \langle \operatorname{div}_g P, X \rangle, \quad (2.9)$$

$$\operatorname{div}_g(fP) = f \operatorname{div}_g P + P^T Df.$$

Proof. Let $\{e_1, e_2\}$ be a normal frame field at p . We have

$$\begin{aligned} \operatorname{div}_g(PX) &= \sum \langle D_{e_i} PX, e_i \rangle = \sum e_i P(X, e_i) = \sum [(D_{e_i} P)(X, e_i) + \langle P D_{e_i} X, e_i \rangle] \\ &= \sum_i DP(X, e_i, e_i) + \sum_{i,j} \langle P e_j, e_i \rangle DX(e_j, e_i) \\ &= \langle P, DX \rangle + \langle \operatorname{div}_g P, X \rangle \quad \text{at } p, \end{aligned}$$

and

$$\begin{aligned} \langle \operatorname{div}_g(fP), X \rangle &= \sum D(fP)(X, e_i, e_i) = \sum D_{e_i}(fP)(X, e_i) \\ &= \sum [e_i(f)P(X, e_i) + f(D_{e_i} P)(X, e_i)] \\ &= \langle X, P^T Df + f \operatorname{div}_g P \rangle \quad \text{at } p. \end{aligned}$$

\square

Lemma 2.4. *Let $P \in T^2S$ and let $p \in S$ be given. Then the following identities hold.*

$$\langle Qv, w \rangle P - \langle Pv, w \rangle Q = Qv \otimes Pw - Pv \otimes Qw, \quad (2.10)$$

$$\langle Qv, w \rangle QP + \langle Pv, w \rangle \operatorname{id} = Qv \otimes QPw + Pv \otimes w \quad \text{for } v, w \in T_pS. \quad (2.11)$$

Proof. Set

$$\Lambda(v, w, P) = \langle Qv, w \rangle P - \langle Pv, w \rangle Q - Qv \otimes Pw + Pv \otimes Qw.$$

Let $\{e_1, e_2\}$ be an orthonormal basis of T_pS with positive orientation. Since $\Lambda(v, w, P)$ is linear with respect to v , w , and P , respectively, for (2.10) it suffices to prove $\Lambda(v, w, P) = 0$ for $v = e_i$, $w = e_j$, and $P = e_k \otimes e_l$. From (2.2), we obtain

$$\langle Qe_i, e_j \rangle = -\langle e_1 \wedge e_2, e_i \otimes e_j \rangle = \langle e_1 \wedge e_2, e_j \otimes e_i \rangle \quad \text{for } 1 \leq i, j \leq 2.$$

Then we compute

$$\begin{aligned}
\Lambda(e_i, e_j, e_k \otimes e_l)(e_m, e_n) &= \langle \Lambda(e_i, e_j, e_k \otimes e_l)e_m, e_n \rangle \\
&= \langle Qe_i, e_j \rangle \delta_{km} \delta_{ln} - \delta_{ki} \delta_{lj} \langle Qe_m, e_n \rangle - \langle Qe_i, e_m \rangle \delta_{kj} \delta_{ln} + \delta_{ki} \delta_{lm} \langle Qe_j, e_n \rangle \\
&= -\langle e_1 \wedge e_2, (\delta_{km} \delta_{ln} e_i \otimes e_j - \delta_{ki} \delta_{lj} e_m \otimes e_n - \delta_{kj} \delta_{ln} e_i \otimes e_m + \delta_{ki} \delta_{lm} e_j \otimes e_n) \rangle \\
&= -\langle e_1 \wedge e_2, [\delta_{ln} e_i \otimes (\delta_{km} e_j - \delta_{kj} e_m) - \delta_{ki} e_n \otimes (\delta_{lm} e_j - \delta_{lj} e_m)] \rangle. \tag{2.12}
\end{aligned}$$

One observes (2.12) to have

$$\Lambda(e_i, e_j, e_k \otimes e_l)(e_m, e_n) = -\Lambda(e_i, e_m, e_k \otimes e_l)(e_j, e_n).$$

It immediately follows that

$$\Lambda(e_i, e_j, e_k \otimes e_l)(e_j, e_n) = 0 \quad \text{for } j = m.$$

Let $i = j \neq m$. From (2.12), we have

$$\begin{aligned}
\Lambda(e_i, e_i, e_k \otimes e_l)(e_m, e_n) &= -e_1 \wedge e_2 [\delta_{ln} e_i \otimes (\delta_{km} e_i - \delta_{ki} e_m) - \delta_{ki} e_n \otimes (\delta_{lm} e_i - \delta_{li} e_m)] \\
&= \delta_{ki} e_1 \wedge e_2 [\delta_{ln} e_i \otimes e_m + e_n \otimes (\delta_{lm} e_i - \delta_{li} e_m)].
\end{aligned}$$

If $n = i$, then

$$\Lambda(e_i, e_i, e_k \otimes e_l)(e_m, e_i) = \delta_{ki} \delta_{lm} e_1 \wedge e_2 (e_i \otimes e_i) = 0.$$

If $n \neq i$, then $n = m$ and

$$\Lambda(e_i, e_i, e_k \otimes e_l)(e_m, e_m) = \delta_{ki} e_1 \wedge e_2 [\delta_{lm} (e_i \otimes e_m + e_m \otimes e_i) - \delta_{li} e_m \otimes e_m] = 0.$$

Let $i \neq j$ and $j \neq m$. Then $i = m$. It follows from (2.12) that

$$\begin{aligned}
&\Lambda(e_i, e_j, e_k \otimes e_l)(e_m, e_n) \\
&= -\delta_{km} \langle e_1 \wedge e_2, \delta_{ln} e_m \otimes e_j - e_n \otimes (\delta_{lm} e_j - \delta_{lj} e_m) \rangle \\
&= \begin{cases} -\delta_{km} \delta_{lj} \langle e_1 \wedge e_2, e_m \otimes e_j + e_j \otimes e_m \rangle = 0 & \text{if } n = j, \\ -\delta_{km} \delta_{lm} \langle e_1 \wedge e_2, e_m \otimes e_j - e_m \otimes e_j \rangle = 0 & \text{otherwise } n = m. \end{cases}
\end{aligned}$$

Using (2.10), we have

$$\Lambda(Qv, Qw, -QPQ) = -\langle Qv, w \rangle QPQ - \langle Pv, w \rangle Q + v \otimes QPw - QPv \otimes w = 0.$$

Thus (2.11) follows from $\Lambda(Qv, Qw, -QPQ)Q = 0$. \square

We further assume that

$$\det(\alpha_t, \alpha_s, \vec{n}) > 0 \quad \text{for } \vec{S}. \tag{2.13}$$

For otherwise, we replace $\alpha(t, s)$ with $\alpha(-t, s)$.

Let $x \in \Gamma_b$. Since $\kappa(x) = 0$ and $D\kappa(x) \neq 0$, from [8, Lemma 2.6], there exist vector fields X_1, X_2 in a neighborhood of x satisfying $\nabla \vec{n} X_i = \lambda_i X_i$, where λ_i are the principal curvatures. Clearly we may extend the vector fields X_i to the region $\overline{S}_{b-\varepsilon}$ when $\varepsilon > 0$ is given small. We assume that X_i are vector fields such that

$$\nabla \vec{n} X_i = \lambda_i X_i, \quad |X_i| = 1, \quad \langle X_1, X_2 \rangle = 0 \quad \text{for } x \in \overline{S}_{b-\varepsilon}, \quad (2.14)$$

where

$$\begin{aligned} \lambda_1 &> 0 \quad \text{for } x \in \overline{S}_{b-\varepsilon}, \\ \lambda_2 &< 0 \quad \text{for } x \in S_{b-\varepsilon}, \quad \lambda_2 = 0 \quad \text{for } x \in \Gamma_b. \end{aligned}$$

Lemma 2.5. *For given $\varepsilon > 0$ small*

$$X_2(\lambda_2) \neq 0 \quad \text{for } x \in S_{b-\varepsilon}.$$

Proof. It will suffice to prove

$$X_2(\lambda_2) \neq 0 \quad \text{for } x \in \Gamma_b.$$

First, we claim that

$$\langle Q\alpha_t, X_2 \rangle(x) \neq 0 \quad \text{for } x \in \Gamma_b.$$

If not, then $\langle Q\alpha_t, X_2 \rangle(x) = 0$ implies that $X_2 = \eta \alpha_t$ with $\eta \neq 0$, and thus

$$\eta^2 \Pi(\alpha_t, \alpha_t) = \langle \nabla_{X_2} \vec{n}, X_2 \rangle = \lambda_2(x) = 0,$$

which contradicts the assumption $\Pi(\alpha_t, \alpha_t) \neq 0$.

In addition, assumption (1.5) implies

$$\langle D\kappa, \alpha_t \rangle = 0, \quad D\kappa = \eta Q\alpha_t \quad \text{for } x \in \Gamma_b, \quad (2.15)$$

for some $\eta \neq 0$. Thus we obtain

$$X_2(\lambda_2) = \frac{1}{\lambda_1} X_2(\kappa) = \frac{\eta}{\lambda_1} \langle X_2, Q\alpha_t \rangle \neq 0 \quad \text{for } x \in \Gamma_b. \quad (2.16)$$

□

We assume that

$$X_2(\lambda_2) > 0 \quad \text{for } x \in \overline{S}_{b-\varepsilon}. \quad (2.17)$$

For otherwise, we replace X_2 with $-X_2$. Furthermore, we assume that X_1, X_2 has positive orientation. For otherwise, we replace X_1 with $-X_1$. Thus

$$QX_2 = X_1, \quad QX_1 = -X_2 \quad \text{for } x \in \overline{S}_{b-\varepsilon}. \quad (2.18)$$

Let $Y \in TS_{b-\varepsilon}$ be given. Define

$$\mathcal{L}_Y V = e^{-s\lambda_2}[(\operatorname{div}_g Q\nabla\bar{n}V + \langle V, Y \rangle)X_1 + \lambda_2(\operatorname{div}_g V)X_2], \quad (2.19)$$

for $V \in TS_{b-\varepsilon}$ and $s > 0$. For given $V, W \in TS_{b-\varepsilon}$, we have

$$\begin{aligned} \langle W, \mathcal{L}_Y V \rangle &= e^{-s\lambda_2}[(\operatorname{div}_g Q\nabla\bar{n}V + \langle V, Y \rangle)\langle W, X_1 \rangle + \lambda_2(\operatorname{div}_g V)\langle W, X_2 \rangle] \\ &= \operatorname{div}_g(e^{-s\lambda_2}\langle W, X_1 \rangle Q\nabla\bar{n}V + e^{-s\lambda_2}\lambda_2\langle W, X_2 \rangle V) - Q\nabla\bar{n}V[e^{-s\lambda_2}\langle W, X_1 \rangle] \\ &\quad - V[e^{-s\lambda_2}\lambda_2\langle W, X_2 \rangle] + e^{-s\lambda_2}\langle V, Y \rangle\langle W, X_1 \rangle \\ &= \langle \mathcal{L}_Y^* W, V \rangle + \operatorname{div}_g(e^{-s\lambda_2}\langle W, X_1 \rangle Q\nabla\bar{n}V + e^{-s\lambda_2}\lambda_2\langle W, X_2 \rangle V) \end{aligned} \quad (2.20)$$

for $x \in S_{b-\varepsilon}$, where

$$\begin{aligned} e^{s\lambda_2}\langle \mathcal{L}_Y^* W, V \rangle &= -\langle DW, X_1 \otimes Q\nabla\bar{n}V + \lambda_2 X_2 \otimes V \rangle + s\langle Q\nabla\bar{n}V, D\lambda_2 \rangle\langle W, X_1 \rangle \\ &\quad + s\lambda_2\langle V, D\lambda_2 \rangle\langle W, X_2 \rangle - \langle W, D_{Q\nabla\bar{n}V} X_1 \rangle - \langle V, D\lambda_2 \rangle\langle W, X_2 \rangle \\ &\quad - \lambda_2\langle W, D_V X_2 \rangle + \langle V, Y \rangle\langle W, X_1 \rangle \quad \text{for } x \in S_{b-\varepsilon}, \end{aligned} \quad (2.21)$$

where the following formulas have been used

$$DW(Z_1, Z_2) = \langle DW, Z_1 \otimes Z_2 \rangle \quad \text{for } Z_1, Z_2 \in TS_{b-\varepsilon}.$$

On the other hand, from (2.19) and Lemma 2.3, we obtain

$$\begin{aligned} e^{s\lambda_2}\langle V, \mathcal{L}_Y W \rangle &= \langle V, (\operatorname{div}_g Q\nabla\bar{n}W + \langle W, Y \rangle)X_1 + \lambda_2(\operatorname{div}_g W)X_2 \rangle \\ &= \langle DW, \langle V, X_1 \rangle Q\nabla\bar{n} + \lambda_2\langle V, X_2 \rangle \operatorname{id} \rangle \\ &\quad + \langle \operatorname{div}_g Q\nabla\bar{n} + Y, W \rangle\langle V, X_1 \rangle \quad \text{for } x \in S_{b-\varepsilon}. \end{aligned} \quad (2.22)$$

Proposition 2.1. *For given $\varepsilon > 0$ small, the following identities hold true.*

(i) *For any $x \in S_{b-\varepsilon}$,*

$$\langle W, -\mathcal{L}_Y V \rangle = \langle -\mathcal{L}_Y^* W, V \rangle - \operatorname{div}_g(e^{-s\lambda_2}\langle W, X_1 \rangle Q\nabla\bar{n}V + e^{-s\lambda_2}\lambda_2\langle W, X_2 \rangle V); \quad (2.23)$$

(ii) *For any $X \in TS_{b-\varepsilon}$ with $|X| = 1$,*

$$\begin{aligned} &\langle X_2, X \rangle \langle (\langle W, X_1 \rangle Q\nabla\bar{n}V + \lambda_2\langle W, X_2 \rangle V), X \rangle \\ &= \lambda_2\langle V, X \rangle\langle W, X \rangle - \Pi(QX, QX)\langle V, X_1 \rangle\langle W, X_1 \rangle \quad \text{for } x \in S_{b-\varepsilon}. \end{aligned} \quad (2.24)$$

Proof. (2.23) follows from (2.20).

Next, we prove (2.24). Using (2.8) where $X = X_2$, $Y = X$, and $Z = V$, we have

$$\langle X_2, X \rangle V = \langle V, X \rangle X_2 + \langle V, X_1 \rangle QX \quad \text{for } x \in S_{b-\varepsilon}.$$

Thus we obtain

$$\begin{aligned}
& \langle X_2, X \rangle (\langle W, X_1 \rangle Q \nabla \bar{n} V + \lambda_2 \langle W, X_2 \rangle V) \\
&= \langle W, X_1 \rangle Q \nabla \bar{n} (\langle V, X \rangle X_2 + \langle V, X_1 \rangle Q X) + \lambda_2 \langle W, X_2 \rangle (\langle V, X \rangle X_2 + \langle V, X_1 \rangle Q X) \\
&= \lambda_2 \langle W, X_1 \rangle \langle V, X \rangle X_1 + \langle W, X_1 \rangle \langle V, X_1 \rangle Q \nabla \bar{n} Q X + \lambda_2 \langle W, X_2 \rangle (\langle V, X \rangle X_2 + \langle V, X_1 \rangle Q X) \\
&= \lambda_2 \langle V, X \rangle W + \langle W, X_1 \rangle \langle V, X_1 \rangle Q \nabla \bar{n} Q X + \lambda_2 \langle W, X_2 \rangle \langle V, X_1 \rangle Q X \quad \text{for } x \in S_{b-\varepsilon}.
\end{aligned}$$

It follows by $\langle QX, X \rangle = 0$ that

$$\begin{aligned}
& \langle X_2, X \rangle (\langle W, X_1 \rangle Q \nabla \bar{n} V + \lambda_2 \langle W, X_2 \rangle V), X \rangle \\
&= \lambda_2 \langle V, X \rangle \langle W, X \rangle - \Pi(QX, QX) \langle V, X_1 \rangle \langle W, X_1 \rangle \quad \text{for } x \in S_{b-\varepsilon}.
\end{aligned}$$

□

Proposition 2.2. *Let $Y \in TS$ and let \mathcal{L}_Y be given in (2.19). Then there exists a constant $\sigma_s > 0$ such that*

$$-e^{s\lambda_2} \langle \mathcal{L}_Y W + L_Y^* W, W \rangle \geq \sigma_s |W|^2 \quad \text{for } W \in TS_{b-\varepsilon}, \quad x \in S_{b-\varepsilon} \quad (2.25)$$

for $s > 0$ large and $\varepsilon > 0$ small enough.

Proof. Using Lemma 2.3, (2.19), (2.18), and (2.11), we obtain

$$\begin{aligned}
e^{s\lambda_2} \langle \mathcal{L}_Y W, W \rangle &= (\operatorname{div}_g Q \nabla \bar{n} W + \langle W, Y \rangle) \langle W, X_1 \rangle + \lambda_2 (\operatorname{div}_g W) \langle W, X_2 \rangle \\
&= \langle DW, \langle W, X_1 \rangle Q \nabla \bar{n} + \lambda_2 \langle W, X_2 \rangle \operatorname{id} \rangle + \langle W, \operatorname{div}_g(Q \nabla \bar{n}) + Y \rangle \langle W, X_1 \rangle \\
&= \langle DW, X_1 \otimes Q \nabla \bar{n} W + \lambda_2 X_2 \otimes W \rangle + \langle W, \operatorname{div}_g(Q \nabla \bar{n}) + Y \rangle \langle W, X_1 \rangle. \quad (2.26)
\end{aligned}$$

It follows from (2.21) and (2.26) that

$$\begin{aligned}
e^{s\lambda_2} \langle \mathcal{L}_Y W + L_Y^* W, W \rangle &= s \langle Q \nabla \bar{n} W, D\lambda_2 \rangle \langle W, X_1 \rangle + s \lambda_2 \langle W, D\lambda_2 \rangle \langle W, X_2 \rangle \\
&\quad - \langle W, D_{Q \nabla \bar{n} W} X_1 \rangle - \langle W, D\lambda_2 \rangle \langle W, X_2 \rangle - \lambda_2 \langle W, D_W X_2 \rangle + \langle W, Y \rangle \langle W, X_1 \rangle \\
&\quad + \langle W, \operatorname{div}_g(Q \nabla \bar{n}) + Y \rangle \langle W, X_1 \rangle \\
&= -s X_2(\lambda_2) [\lambda_1 \langle W, X_1 \rangle^2 - \lambda_2 \langle W, X_2 \rangle^2] - X_2(\lambda_2) \langle W, X_2 \rangle^2 \\
&\quad + 2s \lambda_2 X_1(\lambda_2) \langle W, X_1 \rangle \langle W, X_2 \rangle - X_1(\lambda_2) \langle W, X_1 \rangle \langle W, X_2 \rangle \\
&\quad - \langle W, D_{Q \nabla \bar{n} W} X_1 \rangle - \lambda_2 \langle W, D_W X_2 \rangle + \langle W, Y \rangle \langle W, X_1 \rangle \\
&\quad + \langle W, \operatorname{div}_g(Q \nabla \bar{n}) + Y \rangle \langle W, X_1 \rangle. \quad (2.27)
\end{aligned}$$

Since $\lambda_2 = \mathcal{O}(\varepsilon)$ on $S_{b-\varepsilon}$, (2.25) follows from (2.17) and (2.27), when $s > 0$ is large enough and $\varepsilon > 0$ is small enough, respectively. □

Lemma 2.6. For any $X, Y \in TS$, the following holds.

$$\nabla \vec{n}[X, Y] = D_X \nabla \vec{n} Y - D_Y \nabla \vec{n} X, \quad (2.28)$$

$$\operatorname{div}_g[X, Y] = X \operatorname{div}_g Y - Y \operatorname{div}_g X, \quad (2.29)$$

Proof. A direct calculation yields

$$D_X \nabla \vec{n} Y - D_Y \nabla \vec{n} X = \nabla_X \nabla_Y \vec{n} - \nabla_Y \nabla_X \vec{n} - \langle \nabla_X \nabla_Y \vec{n} - \nabla_Y \nabla_X \vec{n}, \vec{n} \rangle = \nabla_{[X, Y]} \vec{n}.$$

For $v \in C_0^1(S)$, we have

$$\begin{aligned} - \int_S v \operatorname{div}_g[X, Y] dg &= \int_S [X, Y] v dg = \int_S (XYv - YXv) dg \\ &= \int_S [-(Yv) \operatorname{div}_g X + (Xv) \operatorname{div}_g Y] dg = \int_S v (Y \operatorname{div}_g X - X \operatorname{div}_g Y) dx. \end{aligned}$$

Then (2.29) follows. \square

Let $\Phi \in TS$ be given such that

$$\Pi(\Phi, \Phi) \neq 0 \quad \text{for } p \in \overline{S}_{b-\varepsilon}.$$

Define

$$RV = D_V Q \nabla \vec{n} \Phi - D_{Q \nabla \vec{n} V} \Phi \quad \text{for } V \in TS_{b-\varepsilon}. \quad (2.30)$$

Set

$$h_1 = \frac{\langle R\Phi, Q\Phi \rangle}{\Pi(\Phi, \Phi)}, \quad h_2 = \frac{\langle RQ\Phi, Q\Phi \rangle - h_1 \langle \nabla \vec{n} Q\Phi, \Phi \rangle}{|\Phi|^2}, \quad (2.31)$$

$$Z = \frac{1}{|\Phi|^2} (R - h_1 Q \nabla \vec{n} - h_2 \operatorname{id})^T \Phi. \quad (2.32)$$

Lemma 2.7. The following formula is true.

$$R = h_1 Q \nabla \vec{n} + h_2 \operatorname{id} + Z \otimes \Phi. \quad (2.33)$$

Proof. We have

$$\langle (R - h_1 Q \nabla \vec{n} - h_2 \operatorname{id}) \Phi, Q\Phi \rangle = h_1 \Pi(\Phi, \Phi) - h_1 \Pi(\Phi, \Phi) = 0.$$

Since $\frac{Q\Phi}{|\Phi|}, \frac{\Phi}{|\Phi|}$ forms an orthonormal frame, it follows that

$$\begin{aligned} (R - h_1 Q \nabla \vec{n} - h_2 \operatorname{id}) W &= \frac{1}{|\Phi|^2} \langle (R - h_1 Q \nabla \vec{n} - h_2 \operatorname{id}) W, \Phi \rangle \Phi \\ &\quad + \frac{1}{|\Phi|^4} \langle (R - h_1 Q \nabla \vec{n} - h_2 \operatorname{id}) W, Q\Phi \rangle Q\Phi \\ &= \langle W, Z \rangle \Phi \quad \text{for } W \in TS_{b-\varepsilon}, \end{aligned}$$

where $\langle \Phi, Q\Phi \rangle = 0$. Thus (2.33) follows. \square

Proposition 2.3. *Let $V \in TS_{b-\varepsilon}$ be a solution to problem (2.3). Let $\Phi \in TS_{b-\varepsilon}$ be given. Then*

$$\mathcal{L}_Z[\Phi, V] = e^{-s\lambda_2}[(\Phi f_1 - h_1 f_1 - h_2 f_2 + \langle H, V \rangle)X_1 + \lambda_2(\Phi f_2 - \langle D \operatorname{div}_g \Phi, V \rangle)X_2], \quad (2.34)$$

where Z is given in (2.32) and

$$H = \nabla \vec{n} Q D \operatorname{div}_g \Phi + h_1 \operatorname{div}_g Q \nabla \vec{n} - \operatorname{div}_g R - i_Z D \Phi.$$

Proof. From (2.28), (2.30), (2.29), (2.9), and (2.33), we have

$$\begin{aligned} \operatorname{div}_g Q \nabla \vec{n}[\Phi, V] &= \operatorname{div}_g Q (D_\Phi \nabla \vec{n} V - D_V \nabla \vec{n} \Phi) = \operatorname{div}_g (D_\Phi Q \nabla \vec{n} V - D_V Q \nabla \vec{n} \Phi) \\ &= \operatorname{div}_g [\Phi, Q \nabla \vec{n} V] - \operatorname{div}_g R V \\ &= \Phi \operatorname{div}_g Q \nabla \vec{n} V - Q \nabla \vec{n} V \operatorname{div}_g \Phi - \langle h_1 Q \nabla \vec{n} + h_2 \operatorname{id} + Z \otimes \Phi, DV \rangle \\ &\quad - \langle \operatorname{div}_g R, V \rangle \\ &= \Phi f_1 - Q \nabla \vec{n} V \operatorname{div}_g \Phi - h_1 (\langle Q \nabla \vec{n}, DV \rangle + \langle \operatorname{div}_g Q \nabla \vec{n}, V \rangle) - h_2 \operatorname{div}_g V \\ &\quad - \langle D_\Phi V - D_V \Phi, Z \rangle + \langle h_1 \operatorname{div}_g Q \nabla \vec{n}, V \rangle - D \Phi(Z, V) \\ &= -\langle [\Phi, V], Z \rangle + \langle \nabla \vec{n} Q D \operatorname{div}_g \Phi + h_1 \operatorname{div}_g Q \nabla \vec{n} - i_Z D \Phi - \operatorname{div}_g R, V \rangle \\ &\quad + \Phi f_1 - h_2 f_2, \end{aligned}$$

and

$$\operatorname{div}_g[\Phi, V] = \Phi \operatorname{div}_g V - V \operatorname{div}_g \Phi = \Phi f_2 - \langle D \operatorname{div}_g \Phi, V \rangle.$$

Thus (2.34) follows. \square

Lemma 2.8. $\langle X_2, Q \alpha_t \rangle < 0$ for $x \in \Gamma_b$, where X_2 is given in (2.14).

Proof. Since $Q \alpha_t / |\alpha_t|$, $\alpha_t / |\alpha_t|$ forms an orthonormal vector basis long Γ_0 with positive orientation, the assumption (2.13) and the curvature conditions (1.4) and (1.5) imply

$$D \kappa = \eta Q \alpha_t, \quad \eta \neq 0, \quad \langle D \kappa, -Q \alpha_t \rangle > 0 \quad \text{for } x \in \Gamma_b. \quad (2.35)$$

Then

$$\eta = \langle D \kappa, Q \alpha_t \rangle / |\alpha_t|^2 < 0 \quad \text{for } x \in \Gamma_b.$$

It follows from (2.17) that

$$\langle X_2, Q \alpha_t \rangle = \frac{1}{\eta} \langle X_2, D \kappa \rangle = \frac{\lambda_1}{\eta} X_2(\lambda_2) < 0 \quad \text{for } x \in \Gamma_b.$$

\square

Proof of Theorem 2.1. Step 1. Let $V \in TS_{b-\varepsilon}$ satisfy (2.3). Taking $Y = 0$ and $W = V$ in (2.19), (2.23), and (2.24), respectively, we have, by (2.3),

$$\begin{aligned}\mathcal{L}_0 V &= e^{-s\lambda_2}(f_1 X_1 + \lambda_2 f_2 X_2), \\ \langle V, -\mathcal{L}_0 V \rangle &= \langle -\mathcal{L}_0^* V, V \rangle - \operatorname{div}_g(e^{-s\lambda_2} \langle V, X_1 \rangle Q \nabla \vec{n} V + e^{-s\lambda_2} \lambda_2 \langle V, X_2 \rangle V),\end{aligned}\quad (2.36)$$

and

$$\begin{aligned}\langle X_2, X \rangle \langle (\langle V, X_1 \rangle Q \nabla \vec{n} V + \lambda_2 \langle V, X_2 \rangle V), X \rangle \\ = \lambda_2 \langle V, X \rangle^2 - \Pi(QX, QX) \langle V, X_1 \rangle^2 \quad \text{for } X \in TS_{b-\varepsilon},\end{aligned}\quad (2.37)$$

respectively. Since $Q\alpha_t/|\alpha_t|$, $\alpha_t/|\alpha_t|$ forms an orthonormal vector basis along $\Gamma_{b-\varepsilon}$ and Γ_b , respectively, it follows from (2.13) that

$$\nu = -Q\alpha_t/|\alpha_t| \quad \text{for } x \in \Gamma_b; \quad \nu = Q\alpha_t/|\alpha_t| \quad \text{for } x \in \Gamma_{b-\varepsilon}.$$

When given $\varepsilon > 0$ is small, from Lemma 2.8, we have

$$\langle X_2, \nu \rangle > 0 \quad \text{for } x \in \Gamma_b; \quad \langle X_2, \nu \rangle < 0 \quad \text{for } x \in \Gamma_{b-\varepsilon}.\quad (2.38)$$

We integrate (2.36) over $S_{b-\varepsilon}$ and use (2.37), where $X = \nu$, Proposition 2.2, and (2.38) to obtain

$$\begin{aligned}2(V, -\mathcal{L}_0 V)_{L^2(S_{b-\varepsilon}, TS_{b-\varepsilon})} &= (V, -\mathcal{L}_0^* V - \mathcal{L}_0 V)_{L^2(S_{b-\varepsilon}, TS_{b-\varepsilon})} \\ &+ \int_{\Gamma_{b-\varepsilon} \cup \Gamma_b} \frac{e^{-\lambda_2 s}}{\langle X_2, \nu \rangle} [\Pi(Q\nu, Q\nu) \langle V, X_1 \rangle^2 - \lambda_2 \langle V, \nu \rangle^2] d\Gamma \\ &\geq \sigma_s \|V\|_{L^2(S_{b-\varepsilon}, TS_{b-\varepsilon})}^2 + \int_{\Gamma_b} \frac{e^{-\lambda_2 s}}{\langle X_2, \nu \rangle} \Pi(Q\nu, Q\nu) \langle V, X_1 \rangle^2 d\Gamma \\ &+ \int_{\Gamma_{b-\varepsilon}} \frac{e^{-\lambda_2 s}}{\langle X_2, \nu \rangle} [\Pi(Q\nu, Q\nu) \langle V, X_1 \rangle^2 d\Gamma - \lambda_2 \langle V, \nu \rangle^2] d\Gamma.\end{aligned}\quad (2.39)$$

Thus (2.4) follows.

Step 2. We prove (2.5). Let

$$\Phi_1 = \alpha_t, \quad \Phi_2 = \Phi_1 + \eta Q\Phi_1,\quad (2.40)$$

where $\eta > 0$ is given small such that

$$\Pi(\Phi_i, \Phi_i) > 0 \quad \text{for } x \in S_{b-\varepsilon}, \quad i = 1, 2.\quad (2.41)$$

Clearly, Φ_1, Φ_2 forms a basis of vector fields. It follows from (2.34) that

$$\mathcal{L}_{Z_i}[\Phi_i, V] = e^{-s\lambda_2}[(\Phi_i f_1 - h_{i1} f_1 - h_{i2} f_2 + \langle H_i, V \rangle)X_1 + \lambda_2(\Phi_i f_2 - \langle D \operatorname{div}_g \Phi_i, V \rangle)X_2],\quad (2.42)$$

where Z_i is given in (2.32) with $\Phi = \Phi_i$, and

$$H_i = \nabla \vec{n} Q D \operatorname{div}_g \Phi_i + h_{i1} \operatorname{div}_g Q \nabla \vec{n} - \operatorname{div}_g R_i - i_{Z_i} D \Phi_i.$$

We repeat the produce in Step 1 with $\mathcal{L}_0 V$ replaced by $\mathcal{L}_{Z_i}[\Phi_i, V]$ to obtain (2.5). \square

3 The degenerated hyperbolic regions

In the sequel we assume that $\Gamma_0 = \{ \alpha(t, 0) \mid t \in [0, a] \}$ is a non-degenerated curve, that does not contain just one point. For otherwise, we may replace Γ_0 with any curve

$$\Gamma_s = \{ \alpha(t, s) \mid t \in [0, a] \} \quad \text{for } s \in (0, b).$$

Let

$$X_0 = Q\nabla\vec{n}QD\kappa \quad \text{for } x \in S.$$

Let $F \in W^{1,2}(S, TS)$ and $f \in L^2(S)$. We consider a degenerated hyperbolic problem

$$\begin{cases} \langle D^2w, Q^*\Pi \rangle + \frac{1}{\kappa}X_0w + \kappa(\text{tr}_g\Pi)w = \kappa f + \frac{1}{\kappa}\langle X_0, F \rangle + \langle DF, Q^*\Pi \rangle & x \in S, \\ w = q_0, \quad \langle Dw, Q\nabla\vec{n}\alpha_t \rangle = q_1, & x \in \Gamma_0. \end{cases} \quad (3.1)$$

Theorem 3.1. *Let $m \geq 0$ be an integer. Let S be of class $C^{m,1}$. Let $F \in W^{m+1,2}(S, TS)$, $f \in W^{m,2}(S)$, $q_0 \in W^{m+1,2}(\Gamma_0)$, and $q_1 \in W^{m,2}(\Gamma_0)$. Then problem (3.1) admits a unique solution $w \in W^{m+1,2}(S)$ satisfying*

$$\begin{aligned} \|w\|_{W^{m+1,2}(S)}^2 + \|w\|_{W^{m+1,2}(\Gamma_b)}^2 &\leq C(\|f\|_{W^{m,2}(S)}^2 + \|F\|_{W^{m+1,2}(S,TS)}^2 \\ &\quad + \|q_0\|_{W^{m+1,2}(\Gamma_0)}^2 + \|q_1\|_{W^{m,2}(\Gamma_0)}^2), \end{aligned} \quad (3.2)$$

$$\langle Dw, X_0 \rangle = \langle F, X_0 \rangle \quad \text{for } x \in \Gamma_0. \quad (3.3)$$

Let

$$\Gamma_2(w, S) = \int_0^a (|D^2w|^2 + |Dw|^2 + |w|^2) \circ \alpha(t, 0) dt. \quad (3.4)$$

Theorem 3.2. *Let S be of class $C^{2,1}$. Let w solve problem (3.1). Then there are constants $C > c > 0$ such that*

$$\begin{aligned} c\Gamma_2(w, S) &\leq \|w\|_{W^{2,2}(S)}^2 + \|f\|_{W^{1,2}(S)}^2 + \|F\|_{W^{2,2}(S,TS)}^2 \\ &\leq C(\Gamma_2(w, S) + \|f\|_{W^{1,2}(S)}^2 + \|F\|_{W^{2,2}(S,TS)}^2). \end{aligned} \quad (3.5)$$

The proofs of Theorems 3.1 and 3.2 will be given in the end of this section.

Fix $\varepsilon_0 > 0$ small, such that (2.17), Proposition 2.2, and

$$\langle X_2, Q\alpha_t \rangle < 0 \quad \text{for } x \in \overline{S}_{b-\varepsilon_0} \quad (\text{by Lemma 2.8}), \quad (3.6)$$

hold true. The curve $\Gamma_{b-\varepsilon_0}$ divides S into two regions:

$$S = \Sigma_1 \cup \Sigma_2 \cup \Gamma_{b-\varepsilon_0},$$

where

$$\Sigma_1 = \{ \alpha(t, s) \mid (t, s) \in [0, a] \times (0, b - \varepsilon_0) \}, \quad \Sigma_2 = \{ \alpha(t, s) \mid (t, s) \in [0, a] \times (b - \varepsilon_0, b) \}.$$

We shall obtain solutions to the boundary-value problems on the regions Σ_1 and Σ_2 , separately. Then paste them together to have solutions to problem (3.1) on the region S .

To apply some existence results in [7, 9] to the boundary-value problem on the regions Σ_i , we recall some boundary operators in [7]. Let $x \in \Gamma_{b-\varepsilon_0}$ be given. $\mu \in T_x S$ with $|\mu| = 1$ is said to be the *noncharacteristic normal* outside Σ_2 if there is a curve $\zeta : (0, \varepsilon) \rightarrow \Sigma_2$ such that

$$\zeta(0) = x, \quad \zeta'(0) = -\mu, \quad \Pi(\mu, \beta) = 0 \quad \text{for } \beta \in T_x \Gamma_{b-\varepsilon_0}.$$

Let μ be the the noncharacteristic normal field along Γ_0 outside Σ_2 . We define the boundary operators $\mathcal{T}_i : T_x M \rightarrow T_x M$ by

$$\mathcal{T}_i \beta = \frac{1}{2} \left[\beta + (-1)^i \chi(\mu, \beta) \rho(\beta) Q \nabla \vec{n} \beta \right] \quad \text{for } \beta \in T_x M, \quad i = 1, 2,$$

where

$$\chi(\mu, \beta) = \text{sign det}(\mu, \beta, \vec{n}), \quad \varrho(\beta) = \frac{1}{\sqrt{-\kappa}} \text{sign } \Pi(\beta, \beta),$$

and sign is the sign function. Noting that

$$\mu = \frac{Q \nabla \vec{n} \alpha_t}{|\nabla \vec{n} \alpha_t|} \quad \text{for } x \in \Gamma_{b-\varepsilon_0},$$

it follows from (1.3) and (2.13) that

$$\chi(\mu, \alpha_t) \circ \alpha(t, b - \varepsilon_0) = 1, \quad \varrho(\alpha_t) \circ \alpha(t, b - \varepsilon_0) = \frac{1}{\sqrt{-\kappa}} \quad \text{for } t \in [0, a).$$

Thus

$$(\mathcal{T}_2 - \mathcal{T}_1) \alpha_t \circ \alpha(t, 0) = -\frac{1}{\sqrt{-\kappa}} Q \nabla \vec{n} \alpha_t \quad \text{for } t \in [0, a).$$

Similarly, as Γ_0 is a part of the boundary of the region Σ_1 , then

$$(\mathcal{T}_2 - \mathcal{T}_1) \alpha_t \circ \alpha(t, 0) = \frac{1}{\sqrt{-\kappa}} Q \nabla \vec{n} \alpha_t \quad \text{for } t \in [0, a).$$

Since Σ_1 is non-degenerated hyperbolic region, by similar arguments as for [7, Theorems 4.2 and 4.3] (or [9]), we have the following. The details are omitted.

Proposition 3.1. *Problem*

$$\begin{cases} \langle D^2 w, Q^* \Pi \rangle + \frac{1}{\kappa} X_0 w + \kappa (\text{tr}_g \Pi) w = \kappa f + \frac{1}{\kappa} \langle X_0, F \rangle + \langle DF, Q^* \Pi \rangle, & x \in \Sigma_1, \\ w = q_0, \quad \langle Dw, Q \nabla \vec{n} \alpha_t \rangle = q_1 & \text{for } x \in \Gamma_0 \end{cases} \quad (3.7)$$

admits a unique solution w satisfying

$$\begin{aligned} \|w\|_{\mathbb{W}^{m+1,2}(\Sigma_1)}^2 + \sum_{i=0}^{m+1} \|D^i w\|_{L^2(\Gamma_{b-\varepsilon_0})}^2 &\leq C(\|f\|_{\mathbb{W}^{m,2}(\Sigma_1)}^2 + \|F\|_{\mathbb{W}^{m+1,2}(\Sigma_1, T\Sigma_1)}^2) \\ &\quad + \|q_0\|_{\mathbb{W}^{m+1,2}(\Gamma_0)}^2 + \|q_1\|_{\mathbb{W}^{m,2}(\Gamma_0)}^2. \end{aligned}$$

Consider a degenerated problem

$$\begin{cases} \langle D^2w, Q^*\Pi \rangle + \frac{1}{\kappa}X_0w + \kappa(\operatorname{tr}_g\Pi)w = \kappa f + \frac{1}{\kappa}\langle X_0, F \rangle + \langle DF, Q^*\Pi \rangle, & x \in \Sigma_2, \\ w = q_0, \quad \langle Dw, Q\nabla\vec{n}\alpha_t \rangle = q_1 & \text{for } x \in \Gamma_{b-\varepsilon_0}. \end{cases} \quad (3.8)$$

Consier the regions

$$\Sigma_{2\varepsilon} = \{ \alpha(t, s) \mid (t, s) \in [0, a) \times (b - \varepsilon_0, b - \varepsilon) \} \quad \text{for } 0 < \varepsilon < \varepsilon_0. \quad (3.9)$$

Since $\Sigma_{2\varepsilon}$ are also non-degenerated hyperbolic regions for all given $0 < \varepsilon < \varepsilon_0$, for the same reasons as for Σ_1 , problem (3.8) has solutions in $W^{m+1,2}(\Sigma_{2\varepsilon})$ on the regions $\Sigma_{2\varepsilon}$. Because $0 < \varepsilon < \varepsilon_0$ can be arbitrarily small, problem (3.8) actually admits a unique solution w on the region Σ_2 , which satisfies $w \in W^{m+1,2}(\Sigma_{2\varepsilon})$ for all $0 < \varepsilon < \varepsilon_0$. Then Theorem 3.1 follows from Propositions 3.1 and 3.2 later.

Lemma 3.1. *We have*

$$-\kappa(\nabla\vec{n})^{-1}\beta = Q\nabla\vec{n}Q\beta \quad \text{for } \beta \in T_xM, \quad x \in M, \quad \kappa(x) \neq 0.$$

Proof. Let $\kappa(x) \neq 0$. Let e_1, e_2 be an orthonormal basis of T_xS with positive orientation such that

$$\nabla\vec{n}e_i = \lambda_i e_i \quad \text{for } x,$$

where λ_i are the principal curvatures. We have

$$\begin{aligned} Q\nabla\vec{n}Q\beta &= Q\nabla\vec{n}Q(\langle\beta, e_1\rangle e_1 + \langle\beta, e_2\rangle e_2) = Q\nabla\vec{n}(-\langle\beta, e_1\rangle e_2 + \langle\beta, e_2\rangle e_1) \\ &= Q(-\langle\beta, e_1\rangle\lambda_2 e_2 + \langle\beta, e_2\rangle\lambda_1 e_1) = -\langle\beta, e_1\rangle\lambda_2 e_1 - \langle\beta, e_2\rangle\lambda_1 e_2 \\ &= -\kappa(\nabla\vec{n})^{-1}\beta. \end{aligned}$$

□

Lemma 3.2. *For $V \in W^{1,2}(S, TS)$, we have*

$$\langle D(\nabla\vec{n}V), Q^*\Pi \rangle = \operatorname{div}_g \kappa V \quad \text{for } x \in S.$$

Proof. Let $x \in S$ be fixed. Let e_1, e_2 be an orthonormal basis of T_xS with positive orientation such that

$$\nabla\vec{n}e_1 = \lambda_1 e_1.$$

Suppose that E_1, E_2 be a frame field normal at x such that

$$E_i = e_i \quad \text{at } x.$$

We have

$$\begin{aligned}
\langle D(\nabla \vec{n}V), Q^*\Pi \rangle &= \lambda_2 E_1 \langle \nabla \vec{n}V, E_1 \rangle + \lambda_1 E_2 \langle \nabla \vec{n}V, E_2 \rangle \\
&= \lambda_2 E_1 \langle V, \nabla \vec{n}E_1 \rangle + \lambda_1 E_2 \langle V, \nabla \vec{n}E_2 \rangle \\
&= \lambda_2 \langle D_{E_1}V, \nabla \vec{n}E_1 \rangle + \lambda_1 \langle D_{E_2}V, \nabla \vec{n}E_2 \rangle + \lambda_2 \langle V, D_{E_1}(\nabla \vec{n}E_1) \rangle + \lambda_1 \langle V, D_{E_2}(\nabla \vec{n}E_2) \rangle \\
&= \kappa \operatorname{div}_g V + (\Pi_{22}\Pi_{111} + \Pi_{11}\Pi_{122}) \langle V, E_1 \rangle + (\Pi_{22}\Pi_{121} + \Pi_{11}\Pi_{222}) \langle V, E_2 \rangle \\
&= \kappa \operatorname{div}_g V + (\Pi_{22}\Pi_{111} + \Pi_{11}\Pi_{221} - 2\Pi_{12}\Pi_{121}) \langle V, E_1 \rangle \\
&\quad + (\Pi_{22}\Pi_{112} + \Pi_{11}\Pi_{222} - 2\Pi_{12}\Pi_{122}) \langle V, E_2 \rangle \\
&= \kappa \operatorname{div}_g V + E_1(\kappa) \langle V, E_1 \rangle + E_2(\kappa) \langle V, E_2 \rangle = \operatorname{div}_g \kappa V \quad \text{at } x,
\end{aligned}$$

where $\Pi_{ij} = \Pi(E_i, E_j)$ and $\Pi_{ijk} = D\Pi(E_i, E_j, E_k)$. \square

Lemma 3.3. For $v, w \in W^{2,2}(S)$, we have

$$\langle D^2v, Q^*\Pi \rangle = \operatorname{div}_g i_{Dv}Q^*\Pi, \quad (3.10)$$

$$\Pi(QDv, QDw) + w \langle D^2v, Q^*\Pi \rangle = \operatorname{div}_g (w i_{Dv}Q^*\Pi). \quad (3.11)$$

Proof. Let $x \in S$ be fixed. Suppose that E_1, E_2 is a frame field normal at x with positive orientation such that

$$\Pi(E_1, E_2) = 0 \quad \text{at } x.$$

Then

$$QE_1 = -E_2, \quad QE_2 = E_1 \quad \text{in a neighborhood of } x.$$

Using the above formulas, we compute

$$\begin{aligned}
\operatorname{div}_g i_{Dv}Q^*\Pi &= E_1 \langle i_{Dv}Q^*\Pi, E_1 \rangle + E_2 \langle i_{Dv}Q^*\Pi, E_2 \rangle = E_1[\Pi(QDv, QE_1)] + E_2[\Pi(QDv, QE_2)] \\
&= E_1(v_1\Pi_{22} - v_2\Pi_{12}) + E_2(-v_1\Pi_{12} + v_2\Pi_{11}) \\
&= v_{11}\Pi_{22} + v_1\Pi_{221} - v_2\Pi_{121} - v_1\Pi_{122} + v_{22}\Pi_{11} + v_2\Pi_{112} \\
&= \langle D^2v, Q^*\Pi \rangle,
\end{aligned}$$

where $v_i = E_i v$, $\Pi_{ijk} = D\Pi(E_i, E_j, E_k)$, and the following formulas have been used

$$\Pi_{221} = \Pi_{122}, \quad \Pi_{121} = \Pi_{112}.$$

Since $\langle i_{Dv}Q^*\Pi, Dw \rangle = \Pi(QDv, QDw)$, (3.11) follows from (3.10). \square

Proposition 3.2. Problem (3.8) admits a unique solution w satisfying

$$\begin{aligned}
\|w\|_{W^{1,2}(\Sigma_2)}^2 + \|w\|_{W^{1,2}(\Gamma_b)}^2 &\leq C(\|f\|_{L^2(\Sigma_2)}^2 + \|F\|_{W^{1,2}(\Sigma_2, T\Sigma_2)}^2 \\
&\quad + \|q_0\|_{W^{1,2}(\Gamma_{b-\varepsilon_0})}^2 + \|q_1\|_{L^2(\Gamma_{b-\varepsilon_0})}^2), \quad (3.12)
\end{aligned}$$

$$\langle Dw, X_0 \rangle = \langle F, X_0 \rangle \quad \text{for } x \in \Gamma_0. \quad (3.13)$$

Furthermore, for $m \geq 1$, the following estimates hold.

$$\begin{aligned} \|w\|_{\mathbb{W}^{m+1,2}(\Sigma_2)}^2 + \|w\|_{\mathbb{W}^{m+1,2}(\Gamma_b)}^2 &\leq C(\|f\|_{\mathbb{W}^{m,2}(\Sigma_2)}^2 + \|F\|_{\mathbb{W}^{m+1,2}(\Sigma_2, T\Sigma_2)}^2 \\ &\quad + \|q_0\|_{\mathbb{W}^{m+1,2}(\Gamma_{b-\varepsilon_0})}^2 + \|q_2\|_{\mathbb{W}^{m,2}(\Gamma_{b-\varepsilon_0})}^2). \end{aligned} \quad (3.14)$$

Proof. Our task is to establish (3.12)-(3.14). Let w solve problem (3.8). As before, we define

$$W = (\nabla \vec{n})^{-1}(Dw - F) \quad \text{for } x \in \Sigma_2. \quad (3.15)$$

Then

$$Dw = \nabla \vec{n} W + F \quad \text{for } x \in \Sigma_2.$$

Let E_1, E_2 be an orthonormal frame on $S_{b-\varepsilon_0}$ with positive orientation. From Lemma 2.1, we have

$$\begin{aligned} \operatorname{div}_g QDw &= \langle D_{E_1}(QDw), E_1 \rangle + \langle D_{E_2}(QDw), E_2 \rangle = \langle QD_{E_1}Dw, E_1 \rangle + \langle QD_{E_2}Dw, E_2 \rangle \\ &= -\langle D_{E_1}Dw, QE_1 \rangle - \langle D_{E_2}Dw, QE_2 \rangle = \langle D_{E_1}Dw, E_2 \rangle - \langle QD_{E_2}Dw, E_1 \rangle = 0. \end{aligned}$$

Thus we obtain

$$\operatorname{div}_g Q\nabla \vec{n} W = -\operatorname{div}_g QF \quad \text{for } x \in \Sigma_2.$$

On the other hand, from Lemma 3.1, we have

$$X_0 = -\kappa(\nabla \vec{n})^{-1}D\kappa \quad \text{for } x \in \Sigma_2.$$

Then

$$\langle W, D\kappa \rangle = \langle Dw - F, (\nabla \vec{n})^{-1}D\kappa \rangle = \frac{1}{\kappa} \langle F - Dw, X_0 \rangle \quad \text{for } x \in \Sigma_2.$$

In addition, it follows from Lemma 3.2 and the first equation in (3.8) that

$$\begin{aligned} \kappa \operatorname{div}_g W + \langle W, D\kappa \rangle &= \operatorname{div}_g \kappa W = \langle D(\nabla \vec{n} W), Q^* \Pi \rangle = \langle D^2 w - DF, Q^* \Pi \rangle \\ &= \kappa[f - (\operatorname{tr}_g \Pi)w] + \frac{1}{\kappa} \langle X_0, F - Dw \rangle \quad \text{for } x \in \Sigma_2. \end{aligned}$$

Thus we obtain

$$\operatorname{div}_g W = f - (\operatorname{tr}_g \Pi)w \quad \text{for } x \in \Sigma_2.$$

That is, $W \in T\Sigma_2$ solves problem

$$\begin{cases} \operatorname{div}_g Q\nabla \vec{n} W = -\operatorname{div}_g QF & \text{for } x \in \Sigma_2, \\ \operatorname{div}_g W = -(\operatorname{tr}_g \Pi)w + f & \text{for } x \in \Sigma_2. \end{cases} \quad (3.16)$$

Step 1. Consider the case of $m = 0$.

We first prove (3.13). Let $\mathcal{L}_0 W$ be given by (2.19) where $V = W$ and $Y = 0$. It follows from (3.16) that

$$\mathcal{L}_0 W = e^{-s\lambda_2} \{ -(\operatorname{div}_g QF)X_2 + \lambda_2 [-(\operatorname{tr}_g \Pi)w + f] X_2 \}. \quad (3.17)$$

From (2.23) and (2.25), we have

$$2\langle W, -\mathcal{L}_0 W \rangle \geq \sigma_{\varepsilon_0} |W|^2 - \operatorname{div}_g (e^{-s\lambda_2} \langle W, X_2 \rangle Q \nabla \vec{n} W + e^{-s\lambda_2} \lambda_2 \langle W, X_2 \rangle W), \quad (3.18)$$

for $x \in \Sigma_2$. Let $\Sigma_{2\varepsilon}$ be given in (3.9) for $0 < \varepsilon < \varepsilon_0$. We integrate (3.18) over $\Sigma_{2\varepsilon}$ and use (2.24) to have

$$\begin{aligned} 2(W, -\mathcal{L}_0 W)_{L^2(\Sigma_{2\varepsilon}, T\Sigma_{2\varepsilon})} &\geq \sigma_{\varepsilon_0} \|W\|_{L^2(\Sigma_{2\varepsilon}, T\Sigma_{2\varepsilon})}^2 \\ &+ \int_{\Gamma_{b-\varepsilon}} \frac{e^{-s\lambda_2}}{\langle X_2, \nu \rangle} \left[\frac{\Pi(\alpha_t, \alpha_t)}{|\alpha_t|^2} \langle W, X_1 \rangle^2 + |\lambda_2| \langle W, \nu \rangle^2 \right] d\Gamma \\ &+ \int_{\Gamma_{b-\varepsilon_0}} \frac{e^{-s\lambda_2}}{\langle X_2, \nu \rangle} \left[\frac{\Pi(\alpha_t, \alpha_t)}{|\alpha_t|^2} \langle W, X_1 \rangle^2 + |\lambda_2| \langle W, \nu \rangle^2 \right] d\Gamma, \end{aligned} \quad (3.19)$$

for $0 < \varepsilon < \varepsilon_0$.

Next, we deal with the first boundary integration in the right hand side of (3.19). Since

$$\langle W, \nu \rangle = \langle \nu, X_2 \rangle \langle W, X_1 \rangle + \frac{\langle \nu, X_2 \rangle}{\lambda_2} \langle Dw - F, X_2 \rangle \quad \text{for } x \in \Sigma_{2\varepsilon},$$

it follows that

$$\begin{aligned} &\frac{\Pi(\alpha_t, \alpha_t)}{|\alpha_t|^2} \langle W, X_1 \rangle^2 + |\lambda_2| \langle W, \nu \rangle^2 \\ &\geq \frac{\langle \nu, X_2 \rangle^2}{2|\lambda_2|} \langle Dw - F, X_2 \rangle^2 + \left[\frac{\Pi(\alpha_t, \alpha_t)}{|\alpha_t|^2} - |\lambda_2| \langle \nu, X_1 \rangle^2 \right] \langle W, X_1 \rangle^2 \quad \text{for } x \in \Gamma_{b-\varepsilon}. \end{aligned}$$

Noting that

$$|\lambda_2| = \mathcal{O}(\varepsilon), \quad \langle X_2, \nu \rangle > 0 \quad \text{for } x \in \Gamma_{b-\varepsilon},$$

we have

$$\begin{aligned} &\int_{\Gamma_{b-\varepsilon}} \frac{e^{-s\lambda_2}}{\langle X_2, \nu \rangle} \left[\frac{\Pi(\alpha_t, \alpha_t)}{|\alpha_t|^2} \langle W, X_1 \rangle^2 + |\lambda_2| \langle W, \nu \rangle^2 \right] d\Gamma \\ &\geq \sigma \int_{\Gamma_{b-\varepsilon}} \left(\frac{1}{|\lambda_2|} \langle Dw - F, X_2 \rangle^2 + \langle W, X_1 \rangle^2 \right) d\Gamma \quad \text{for } \varepsilon > 0 \text{ small}, \end{aligned} \quad (3.20)$$

which imply that

$$\langle Dw, X_2 \rangle = \langle F, X_2 \rangle \quad \text{for } x \in \Gamma_b. \quad (3.21)$$

On the other hand, from (2.35),

$$\begin{aligned} X_0 &= Q \nabla \vec{n} Q D \kappa = -\eta Q \nabla \vec{n} \alpha_t = \eta \langle \nabla \vec{n} \alpha_t, Q X_1 \rangle X_1 + \eta \langle \nabla \vec{n} \alpha_t, Q X_2 \rangle X_2 \\ &= -\eta \langle \nabla \vec{n} \alpha_t, X_2 \rangle X_1 + \eta \langle \nabla \vec{n} \alpha_t, X_1 \rangle X_2 = \eta \lambda_1 \langle \alpha_t, X_1 \rangle X_2 \quad \text{for } x \in \Gamma_b. \end{aligned} \quad (3.22)$$

Thus (3.13) follows.

We claim that

$$\langle X_1, \tau \rangle \neq 0 \quad \text{for } x \in \Gamma_b, \quad (3.23)$$

where $\tau = \alpha_t/|\alpha_t|$. For otherwise, $\langle X_1, \tau \rangle = 0$ implies that $\tau = \pm X_2$ and

$$\Pi(\tau, \tau) = \Pi(X_2, X_2) = 0,$$

which contradicts (1.3). Thus

$$\begin{aligned} \langle Dw, \tau \rangle &= \langle \tau, X_1 \rangle \langle Dw, X_1 \rangle + \langle \tau, X_2 \rangle \langle Dw, X_2 \rangle \\ &= \langle \tau, X_1 \rangle \langle Dw, X_1 \rangle + \langle \tau, X_2 \rangle \langle F, X_2 \rangle, \end{aligned}$$

that is

$$\langle Dw, X_1 \rangle = \frac{1}{\langle X_1, \tau \rangle} (\langle Dw, \tau \rangle - \langle X_2, \tau \rangle \langle F, X_2 \rangle) \quad \text{for } x \in \Gamma_b.$$

Then

$$\begin{aligned} \langle W, X_1 \rangle &= \frac{1}{\lambda_1} \langle Dw - F, X_1 \rangle = \frac{1}{\lambda_1 \langle X_1, \tau \rangle} \langle Dw, \tau \rangle \\ &\quad - \frac{1}{\lambda_1} \langle F, X_1 \rangle - \frac{\langle X_2, \tau \rangle}{\lambda_1 \langle X_1, \tau \rangle} \langle F, X_2 \rangle \quad \text{for } x \in \Gamma_b. \end{aligned} \quad (3.24)$$

We now integrate (3.18) over Σ_2 and use (2.24) to obtain

$$\begin{aligned} 2(W, -\mathcal{L}_0 W)_{L^2(\Sigma_2, T\Sigma_2)} &\geq \sigma_{\varepsilon_0} \|W\|_{L^2(\Sigma_2, T\Sigma_2)}^2 + \int_{\Gamma_b} \frac{e^{-s\lambda_2} \Pi(\alpha_t, \alpha_t)}{\langle X_2, \nu \rangle |\alpha_t|^2} \langle W, X_1 \rangle^2 d\Gamma \\ &\quad + \int_{\Gamma_{b-\varepsilon_0}} \frac{e^{-s\lambda_2}}{\langle X_2, \nu \rangle} \left[\frac{\Pi(\alpha_t, \alpha_t)}{|\alpha_t|^2} \langle W, X_1 \rangle^2 - \lambda_2 \langle W, \nu \rangle^2 \right] d\Gamma. \end{aligned} \quad (3.25)$$

It follows from (3.25), (3.17), and (3.24) that

$$\begin{aligned} \|Dw\|_{L^2(\Sigma_2, T\Sigma_2)}^2 &+ \int_{\Gamma_b} \langle Dw, \tau \rangle^2 d\Gamma \leq C(\|w\|_{L^2(\Sigma_2)}^2 + \|f\|_{L^2(\Sigma_2)}^2 + \|F\|_{W^{1,2}(\Sigma_2, T\Sigma_2)}^2 \\ &\quad + \|F\|_{L^2(\Gamma_b, TM)}^2 + \|F\|_{L^2(\Gamma_{b-\varepsilon_0}, TM)}^2 + \|Dw\|_{L^2(\Gamma_{b-\varepsilon_0}, TM)}^2) \\ &\leq C(\|w\|_{L^2(\Sigma_2)}^2 + \|f\|_{L^2(\Sigma_2)}^2 + \|F\|_{W^{1,2}(\Sigma_2, T\Sigma_2)}^2 + \|Dw\|_{L^2(\Gamma_{b-\varepsilon_0}, TM)}^2). \end{aligned} \quad (3.26)$$

Thus (3.12) follows from (3.26) and the boundary data in (3.8).

Step 2. We prove (3.14) in the case of $m = 1$. The case of $m \geq 2$ can be treated by the similar arguments.

Let W be given in (3.15). Suppose that Φ_2 and Φ_2 are given in (2.40) such that

$$\Pi(\Phi_i, \Phi_i) > 0 \quad \text{for } x \in \Sigma_2$$

and Φ_2, Φ_2 forms a basis of vector fields on Σ_2 . By a similar computation as in (2.42), we have

$$\begin{aligned} \mathcal{L}_{Z_i}[\Phi_i, W] &= e^{-s\lambda_2}[(\Phi_i f_2 - h_{i1} f_2 - h_{i2} f_2 + \langle H_i, W \rangle)X_2 \\ &\quad + \lambda_2(\Phi_i f_2 - \langle D \operatorname{div}_g \Phi_i, W \rangle)X_2], \end{aligned} \quad (3.27)$$

where Z_i, h_{2i} , and H_i are the same as in (2.42), and

$$f_2 = -\operatorname{div}_g QF, \quad f_2 = -(\operatorname{tr}_g \Pi)w + f \quad \text{for } x \in \Sigma_2.$$

By a similar argument as in Step 1 yields

$$\begin{aligned} &2([\Phi_i, W], -\mathcal{L}_{Z_i}[\Phi_i, W])_{L^2(\Sigma_2, T\Sigma_2)} \\ &\geq \sigma_{\varepsilon_0} \|[\Phi_i, W]\|_{L^2(\Sigma_2, T\Sigma_2)}^2 + \int_{\Gamma_b \cup \Gamma_{b-\varepsilon_0}} e^{-s\lambda_2} P_i(W_\varepsilon) d\Gamma, \end{aligned} \quad (3.28)$$

where

$$P_i(W) = \frac{1}{\langle X_2, \nu \rangle} [\Pi(\tau, \tau) \langle [\Phi_i, W], X_1 \rangle^2 - \lambda_2 \langle [\Phi_i, W], \nu \rangle^2], \quad i = 1, 2. \quad (3.29)$$

Since

$$\langle X_2, \nu \rangle > 0 \quad \text{for } x \in \Gamma_b, \quad \langle X_2, \nu \rangle < 0 \quad \text{for } x \in \Gamma_{b-\varepsilon_0},$$

from (3.27)-(3.29), we obtain

$$\begin{aligned} \|[\Phi_i, W]\|_{L^2(\Sigma_2, T\Sigma_2)}^2 &+ \int_{\Gamma_b} \langle D\Phi_i W, X_1 \rangle^2 d\Gamma \leq C(\|f\|_{W^{1,2}(\Sigma_2)}^2 + \|F\|_{W^{2,2}(\Sigma_2, T\Sigma_2)}^2 \\ &+ \|w\|_{W^{1,2}(\Sigma_2)}^2 + \|w\|_{L^2(\Sigma_2)}^2 + \|Dw\|_{L^2(\Sigma_2, T\Sigma_2)}^2), \end{aligned} \quad (3.30)$$

for $i = 1, 2$, which imply that

$$W \in W^{1,2}(\Sigma_2, T\Sigma_2).$$

Thus, by the trace theorem,

$$W \in L^2(\Gamma_b, TM). \quad (3.31)$$

It is easy to check that $\tau = \alpha_t/|\alpha_t|$, X_2 forms a vector basis along Γ_b by (3.23). From (3.21), we have

$$Dw = p_1 \tau + p_2 X_2 \quad \text{for } x \in \Gamma_b,$$

where

$$p_1 = \frac{1}{1 - \langle \tau, X_2 \rangle^2} [\tau(w) - \langle \tau, X_2 \rangle \langle F, X_2 \rangle], \quad p_2 = \frac{1}{1 - \langle \tau, X_2 \rangle^2} [-\langle \tau, X_2 \rangle \tau(w) + \langle F, X_2 \rangle].$$

Thus

$$\begin{aligned} |\langle D_\tau Dw, X_1 \rangle| &\geq |\langle \tau, X_1 \rangle| |\tau(p_1)| - C(|Dw| + |F|) \\ &\geq |\langle \tau, X_1 \rangle| |\tau\tau(w)| - C(|\tau(w)| + |F| + |DF|) \quad \text{for } x \in \Gamma_b. \end{aligned} \quad (3.32)$$

Next, we compute

$$\begin{aligned} \langle D_{\Phi_1}(\nabla \vec{n}W), X_1 \rangle &= \Phi_1(\lambda_1 \langle W, X_1 \rangle) - \langle \nabla \vec{n}W, D_{\Phi_1} X_1 \rangle \\ &= \Phi_1(\lambda_1) \langle W, X_1 \rangle + \langle \nabla \vec{n} D_{\Phi_1} W, X_1 \rangle + \lambda_1 \langle W, D_{\Phi_1} X_1 \rangle - \langle \nabla \vec{n}W, D_{\Phi_1} X_1 \rangle. \end{aligned}$$

Thus

$$\begin{aligned} \lambda_1 \langle D_{\Phi_1} W, X_1 \rangle &= \langle \nabla \vec{n} D_{\Phi_1} W, X_1 \rangle = \langle D_{\Phi_1}(Dw - F), X_1 \rangle \\ &\quad - \Phi_1(\lambda_1) \langle W, X_1 \rangle - \lambda_1 \langle W, D_{\Phi_1} X_1 \rangle + \langle \nabla \vec{n}W, D_{\Phi_1} X_1 \rangle \end{aligned}$$

which yields, by (3.31) and (3.32),

$$\begin{aligned} |\langle D_{\Phi_1} W, X_1 \rangle| &\geq |\alpha_t| |\langle D_\tau Dw, X_1 \rangle| - C(|W| + |DF|) \\ &\geq \sigma |\tau\tau(w)| - C(|W| + |\tau(w)| + |F| + |DF|) \quad \text{for } x \in \Gamma_b, \end{aligned}$$

since $\Phi_1 = |\alpha_t| \tau$. Thus (3.14) follows from (3.30) in the case of $m = 1$. \square

Proof of Theorem 3.1 We solve problem (3.7) on the region Σ_1 to have a solution w_2 and then solve problem (3.8) on Σ_2 with the data

$$q_0 = w_2, \quad q_2 = \langle Dw_2, Q \nabla \vec{n} \alpha_t \rangle \quad \text{for } x \in \Gamma_{b-\varepsilon_0},$$

to obtain the solution w_2 . Furthermore, paste two solutions together to have a solution to problem (3.1) on the region S . Clearly, the solution meets our needs. \square

Proof of Theorem 3.2 From Theorem 3.1 and [7, Theorem 4.3], we have

$$\begin{aligned} c\Gamma_2(w, S) &\leq \|w\|_{W^{2,2}(\Sigma_1)}^2 + \|f\|_{W^{1,2}(\Sigma_1)}^2 + \|F\|_{W^{2,2}(\Sigma_1, T\Sigma_1)}^2 \\ &\leq \|w\|_{W^{2,2}(S)}^2 + \|f\|_{W^{1,2}(S)}^2 + \|F\|_{W^{2,2}(S, TS)}^2 \\ &\leq C(\Gamma_2(w, S) + \|f\|_{W^{1,2}(S)}^2 + \|F\|_{W^{2,2}(S, TS)}^2). \end{aligned}$$

\square

4 Proofs of Theorems 1.1-1.3 in Section 1

Let

$$U \in T_{\text{sym}}^2 S.$$

Consider problem

$$\begin{cases} Dv = \nabla \vec{n}V + F & \text{for } x \in S, \\ \operatorname{div}_g V + v \operatorname{tr}_g \Pi = f & \text{for } x \in S, \end{cases} \quad (6.1)$$

where $(V, v) \in W^{1,2}(S, T) \times W^{1,2}(S)$ is the unknown and

$$F = Q[D(\operatorname{tr}_g U) - \operatorname{div}_g U], \quad f = -\operatorname{tr}_g U(Q\nabla \vec{n}\cdot, \cdot) \quad \text{for } x \in S. \quad (6.2)$$

For $y \in W^{1,2}(S, \mathbb{R}^3)$, let

$$2v = \nabla y(e_2, e_2) - \nabla y(e_2, e_2) \quad \text{for } x \in S, \quad (6.3)$$

$$V = (\nabla \vec{n})^{-1}(Dv - F) \quad \text{for } x \in S, \quad (6.4)$$

where e_2, e_2 is an orthonormal basis of $T_x S$ with positive orientation.

By [7, Section 2], there is a $y \in W^{1,2}(S, \mathbb{R}^3)$ to solve problem (1.1) if and only if (v, V) , being given in (6.3) and (6.4), solves problem (6.1). In that case, we have

$$\begin{cases} \nabla_{e_1} y = U(e_1, e_1)e_1 + [v + U(e_1, e_2)]e_2 - \langle QV, e_1 \rangle \vec{n}, \\ \nabla_{e_2} y = [-v + U(e_1, e_2)]e_1 + U(e_2, e_2)e_2 - \langle QV, e_2 \rangle \vec{n}, \end{cases} \quad \text{for } x \in S. \quad (6.5)$$

Moreover, from [7, Theorem 2.1], (V, v) is a solution to problem (6.1) if and only if v solves problem

$$\langle D^2 v, Q^* \Pi \rangle + \frac{1}{\kappa} X_0 v + v \kappa \operatorname{tr}_g \Pi = \kappa f + \frac{1}{\kappa} \langle X_0, F \rangle + \langle DF, Q^* \Pi \rangle \quad \text{for } x \in S, \quad (6.6)$$

where f and F are given in (6.2).

Proof of Theorem 1.1 We solve the degenerated hyperbolic problem

$$\begin{cases} \langle D^2 v, Q^* \Pi \rangle + \frac{1}{\kappa} X_0 v + \kappa (\operatorname{tr}_g \Pi) v = \kappa f + \frac{1}{\kappa} \langle X_0, F \rangle + \langle DF, Q^* \Pi \rangle & x \in S, \\ v = \langle Dv, Q\nabla \vec{n} \alpha_t \rangle = 0 & \text{for } x \in \Gamma_0, \end{cases}$$

by Theorem 3.1, to have the solution $v \in W^{m+1,2}(S)$. Then, set

$$V = (\nabla \vec{n})^{-1}(Dw - F) \quad \text{for } x \in S.$$

It follows from [7, Theorem 2.1] that problem (1.1) admits a solution $y \in W^{m,2}(S, \mathbb{R}^3)$ that satisfies (6.5) on the region S .

Let $y = W + w\vec{n}$, where $w = \langle y, \vec{n} \rangle \in W^{m,2}(S)$. Since

$$\operatorname{sym} DW = U - w\Pi \quad \text{for } x \in S,$$

it follows from [6, Lemma 4.3] that

$$W \in W^{m+1,2}(S, TS).$$

□

Proof of Theorem 1.2 Let $y \in W^{2,2}(S, \mathbb{R}^3)$ be an infinitesimal isometry. Let

$$2v = \nabla y(e_2, e_2) - \nabla y(e_2, e_2), \quad V = (\nabla \vec{n})^{-1} Dv \quad \text{for } x \in S,$$

where e_2, e_2 is an orthonormal basis of $T_x S$ with positive orientation. It follows from (6.5) and (6.6) that

$$\begin{cases} \nabla_{e_2} y = ve_2 - \langle QV, e_2 \rangle \vec{n}, \\ \nabla_{e_2} y = -ve_2 - \langle QV, e_2 \rangle \vec{n} \end{cases} \quad \text{for } x \in S. \quad (6.7)$$

and

$$\langle D^2 v, Q^* \Pi \rangle + \frac{1}{\kappa} X_0 v + v \kappa \operatorname{tr}_g \Pi = 0 \quad \text{for } x \in S. \quad (6.8)$$

From (6.7), (6.8), Theorems 3.1, and 3.2, we have

$$\|y\|_{W^{2,2}(S)}^2 \leq C(\|v\|_{W^{1,2}(S)}^2 + \|V\|_{W^{1,2}(S, TS)}^2) \leq C\Gamma_2(v, S) \quad \text{for } y \in \mathcal{V}(S, \mathbb{R}^3), \quad (6.9)$$

where $\Gamma_2(v, S)$ is given in (3.4) with w replaced by v .

For given $\varepsilon > 0$, we take $q_{0\varepsilon} \in W^{m+2,2}(\Gamma_0)$ and $q_{1\varepsilon} \in W^{m+1,2}(\Gamma_0)$ such that

$$\|v - q_{0\varepsilon}\|_{W^{2,2}(\Gamma_0)}^2 + \|\langle Dv, Q\nabla \vec{n} \alpha_t \rangle - q_{1\varepsilon}\|_{W^{1,2}(\Gamma_0)}^2 \leq \varepsilon. \quad (6.10)$$

Then we solve problem

$$\begin{cases} \langle D^2 v, Q^* \Pi \rangle + \frac{1}{\kappa} X_0 v + \kappa(\operatorname{tr}_g \Pi)v = 0 & \text{for } x \in S, \\ v = q_{0\varepsilon}, \quad \langle Dw, Q\nabla \vec{n} \alpha_t \rangle = q_{1\varepsilon} & x \in \Gamma_0 \end{cases}$$

to obtain the solution v_ε . Thus there is $y_\varepsilon \in \mathcal{V}(S, \mathbb{R}^3)$ with $2v_\varepsilon = \nabla y_\varepsilon(e_2, e_2) - \nabla y_\varepsilon(e_2, e_2)$. By Theorem 3.1, $y_\varepsilon \in W^{m+2,2}(S, \mathbb{R}^3)$, and, thus, by the imbedding theorem (see [1, P. 158])

$$y_\varepsilon \in \mathcal{V}(S, \mathbb{R}^3) \cup C_B^m(S, \mathbb{R}^3).$$

Finally, it follows from Theorem 3.2, (6.9), and (6.10) that

$$\|y - y_\varepsilon\|_{W^{2,2}(S)}^2 \leq C\varepsilon.$$

The proof is complete. □

Proof of Theorem 1.3 As in [2] we conduct in $2 \leq i \leq m$. Let

$$y_\varepsilon = \sum_{j=0}^{i-1} \varepsilon^j z_j$$

be an $(i-1)$ th order isometry of class $C_B^{2+4(m-i+1)}(S, \mathbb{R}^3)$, where $z_0 = \operatorname{id}$ and $z_2 = y$ for some $i \geq 2$. Then

$$\sum_{j=0}^k \nabla^T z_j \nabla z_{k-j} = 0 \quad \text{for } 1 \leq k \leq i-1.$$

Next, we shall find out $z_i \in C_B^{2+4(m-i)}(S, \mathbb{R}^3)$ such that

$$\phi_\varepsilon = y_\varepsilon + \varepsilon^i z_i$$

is an i th order isometry. By Corollary 1.1 there exists a solution $z_i \in C_B^{2+4(m-i)}(S, \mathbb{R}^3)$ to problem

$$\text{sym } \nabla z_i = -\frac{1}{2} \text{sym} \sum_{j=1}^{i-1} \nabla^T z_j \nabla z_{i-j}$$

which satisfies

$$\begin{aligned} \|z_i\|_{C_B^{2+4(m-i)}(S, \mathbb{R}^3)} &\leq C \left\| \sum_{j=1}^{i-1} \text{sym} \nabla^T z_j \nabla z_{i-j} \right\|_{C_B^{2+4(m-i)+3}(S, \mathbb{R}^3)} \\ &\leq C \sum_{j=1}^{i-1} \|z_j\|_{C_B^{2+4(m-i+1)}(S, \mathbb{R}^3)} \|z_{i-j}\|_{C_B^{2+4(m-i+1)}(S, \mathbb{R}^3)}. \end{aligned}$$

The conduction completes. □

Compliance with Ethical Standards

Conflict of Interest: The author declares that there is no conflict of interest.

Ethical approval: This article does not contain any studies with human participants or animals performed by the authors.

References

- [1] D. Gilbary, Neil S. Trudinger, Elliptic partial differential equations of second order. Reprint of the 1998 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2001.
- [2] P. Hornung, M. Lewicka, M. R. Pakzad, Infinitesimal isometries on developable surfaces and asymptotic theories for thin developable shells. *J. Elasticity* 111 (2013), no. 1, 1-19.
- [3] M. Lewicka, M. R. Pakzad, The infinite hierarchy of elastic shell models: some recent results and a conjecture. *Infinite dimensional dynamical systems*, 407-420, Fields Inst. Commun., 64, Springer, New York, 2013.
- [4] M. Lewicka, M. G. Mora, M. R. Pakzad, Shell theories arising as low energy Γ -limit of 3d nonlinear elasticity. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 9 (2010), no. 2, 253-295.
- [5] —, The matching property of infinitesimal isometries on elliptic surfaces and elasticity of thin shells. *Arch. Ration. Mech. Anal.* 200 (2011), no. 3, 1023-1050.
- [6] P. F. Yao, Modeling and control in vibrational and structural dynamics. A differential geometric approach. Chapman & Hall/CRC Applied Mathematics and Nonlinear Science Series. CRC Press, Boca Raton, FL, 2011.

- [7] —, Linear Strain Tensors on Hyperbolic Surfaces and Asymptotic Theories for Thin Shells, *SIAM J. Math. Anal.*, Vol. 51(2019), No. 2, pp. 1387-1435.
- [8] —, Optimal exponentials of thickness in Korn's inequalities for parabolic and elliptic shells *Annali di Matematica Pura ed Applicata (1923 -)* <https://doi.org/10.1007/s10231-020-01000-6>.
- [9] —, Strain tensors on hyperbolic surfaces and their applications, *J. Funct. Anal.* (2021), 108986, doi: <https://doi.org/10.1016/j.jfa.2021.108986>.