

ON BIRKHOFF ANGLES IN NORMED SPACES

HENDRA GUNAWAN, MUHAMAD JAMALUDIN, AND MAS DAFFA PRATAMADIRDJA

ABSTRACT. Associated to Birkhoff orthogonality, we study Birkhoff angles in a normed space and present some of their basic properties. We also discuss how to decide whether an angle is more acute or more obtuse than another. In addition, given two vectors x and y in a normed space, we study the formula for Birkhoff ‘cosine’ of the angle from x to y from which we can, in principal, compute the angle. Some examples will be presented.

1. INTRODUCTION

Let $(X, \|\cdot\|)$ be a real normed space. (Throughout this note, X will be assumed so, unless otherwise stated.) Unlike in an inner product space, there is not a standard definition of orthogonality, let alone the definition of angles, in X . We know, for instance, the notion of Pythagorean orthogonality (denoted by \perp_P), which states that

$$x \perp_P y \text{ if and only if } \|x - y\|^2 = \|x\|^2 + \|y\|^2,$$

and the notion of isosceles orthogonality (denoted by \perp_I), which states that

$$x \perp_I y \text{ if and only if } \|x + y\| = \|x - y\|,$$

for $x, y \in X$, as introduced by R.C. James in [6]. Both of these definition coincides with the usual definition of orthogonality when X is an inner product space and the norm is induced by the inner product.

In 1935, G. Birkhoff [3] introduced a different notion of orthogonality in a normed space, inspired by the property of the tangent to a circle in the Euclidean space (see Figure 1.1). A vector $x \in X$ is said to be B -orthogonal to another vector $y \in X$, denoted by $x \perp_B y$, if and only if

$$(1.1) \quad \|x + \lambda y\| \geq \|x\|$$

for every $\lambda \in \mathbb{R}$. This notion of orthogonality is then known as Birkhoff orthogonality, as studied in [7]. One may observe that this definition also coincides with the usual orthogonality when X is an inner product space and the norm is induced by the inner product. However, unlike Pythagorean orthogonality and isosceles orthogonality, Birkhoff orthogonality is not symmetric: $x \perp_B y$ does not imply $y \perp_B x$. For discussions on various notions of orthogonality in normed spaces, see [1, 2, 9].




Figure 1.1. A vector x is B -orthogonal to another vector y in ℓ_2^2 (\mathbb{R}^2 with the usual norm). The circle has radius $\|x\|$.

2010 Mathematics Subject Classification. 46B20.

Key words and phrases. Birkhoff angles, Birkhoff orthogonality, normed spaces.




Figure 1.2. A vector x is B -orthogonal to another vector y in ℓ_2^∞ (\mathbb{R}^2 with the supremum norm). The square is the set of vectors with length $\|x\|$. Note that here the vector y is not B -orthogonal to x .

Once we define an orthogonality criterion in X , we may ask how we can define angles in X that are compatible with the orthogonality defined there. If we use Pythagorean orthogonality, then we can define the angle *between* two nonzero vectors x and y in X by

$$A_P(x, y) := \arccos \frac{\|x\|^2 + \|y\|^2 - \|x - y\|^2}{2\|x\|\|y\|}.$$

If we adopt isosceles orthogonality, then we can define the angle between two nonzero vectors x and y in X by

$$A_I(x, y) := \arccos \frac{\|x + y\|^2 - \|x - y\|^2}{4\|x\|\|y\|}.$$

(See [4] for some properties of Pythagorean angles and isosceles angles.) Now, if we opt to define Birkhoff orthogonality in X , how should we define the angles *from* a vector *to* another vector in X ? To answer this question, we must dig deeper to the case where X is an inner product space. This paper presents some results that might be of interest to the readers, **especially undergraduate students who are interested in geometry of normed spaces**. For further readings, we refer the readers to [2, 8, 11].

2. PRELIMINARY OBSERVATION: ACUTE AND OBTUSE B-ANGLES

For the moment, let $(X, \langle \cdot, \cdot \rangle)$ be an inner product space, and $\|x\| := \langle x, x \rangle^{1/2}$ be the induced norm on X . Let $x \in X$ and $y \in X \setminus \{0\}$. By simple calculations, one may observe that

- when $\langle x, y \rangle \geq 0$ (that is, x and y form an acute angle), the inequality (1.1) holds for $\lambda \leq -\frac{2\langle x, y \rangle}{\|y\|^2}$ or $\lambda \geq 0$.
- when $\langle x, y \rangle \leq 0$ (that is, x and y form an obtuse angle), the inequality (1.1) holds for $\lambda \leq 0$ or $\lambda \geq -\frac{2\langle x, y \rangle}{\|y\|^2}$.
- when $\langle x, y \rangle = 0$ (that is, x and y are orthogonal to each other), the inequality (1.1) holds for every $\lambda \in \mathbb{R}$.

The sets of the values of λ that satisfies the inequality (1.1) are visualized in the following figure:




Figure 2.1. The sets of the values of λ for which the inequality (1.1) holds when (a) $\langle x, y \rangle \geq 0$, (b) $\langle x, y \rangle \leq 0$, and (c) $\langle x, y \rangle = 0$, where $\gamma := \frac{2|\langle x, y \rangle|}{\|y\|^2}$.

Note that when $\langle x, y \rangle \neq 0$ (that is, x and y form a proper acute or obtuse angle), the value of $\gamma := \frac{2|\langle x, y \rangle|}{\|y\|^2}$ is strictly positive. This means that, there exist some values of λ for which the inequality (1.1) fails to hold.

We now go back to normed spaces in general. Based on the above observation, we can define acute and obtuse angle in a normed space $(X, \|\cdot\|)$ through the sets of the values of λ satisfying the inequality (1.1). Let $x, y \in X$. We say that

- x forms an **acute B-angle** to y , denoted by $x A_B y$, if the inequality (1.1) holds for every $\lambda \geq 0$.
- x forms an **obtuse B-angle** to y , denoted by $x O_B y$, if the inequality (1.1) holds for every $\lambda \leq 0$.
- x is **B-orthogonal** to y , denoted by $x \perp_B y$, if x forms an acute B-angle and an obtuse B-angle to y simultaneously.

As studied in [5, 10], we have the following propositions. We leave the proof of the first proposition to the readers.

Proposition 2.1. *Let $x, y \in X \setminus \{0\}$ and $a, b \in \mathbb{R} \setminus \{0\}$.*

- (1) *Suppose that $x A_B y$. If $ab > 0$ (that is, a and b have the same sign), then $ax A_B by$. If $ab < 0$, then $ax O_B by$.*
- (2) *Suppose that $x O_B y$. If $ab > 0$, then $ax O_B by$. If $ab < 0$, then $ax A_B by$.*

Proposition 2.2. *Let $x, y \in X$. Then we have*

- (1) *$x A_B y$ if and only if there exists $\delta > 0$ such that the inequality (1.1) holds for every $\lambda \in [0, \delta)$.*
- (2) *$x O_B y$ if and only if there exists $\delta > 0$ such that the inequality (1.1) holds for every $\lambda \in (-\delta, 0]$.*
- (3) *$x \perp_B y$ if and only if there exists $\delta > 0$ such that the inequality (1.1) holds for every $\lambda \in (-\delta, \delta)$ simultaneously.*

Proof. We shall only prove the first statement, as the second one can be proven in a similar way and the third one is a consequence of the first and the second ones. Now, the ‘only if’ part is immediate, and so we only need to prove the ‘if’ part. Suppose that there exists $\delta > 0$ such that the inequality $\|x + \lambda y\| \geq \|x\|$ holds for every $\lambda \in [0, \delta)$. If x or y equals 0, then the inequality obviously holds for every $\lambda \geq 0$. So assume that $x, y \neq 0$. Suppose that, to the contrary, there exists $\lambda' \geq \delta$ such that $\|x + \lambda' y\| < \|x\|$. Choose $n \in \mathbb{N}$ such that $\frac{\lambda'}{n} \in (0, \delta)$. By the hypothesis, we have $\|x + \frac{\lambda'}{n} y\| \geq \|x\|$. But, by the triangle inequality, we find that

$$\begin{aligned} \left\| x + \frac{\lambda'}{n} y \right\| &= \left\| \frac{n-1}{n} x + \frac{1}{n} (x + \lambda' y) \right\| \\ &\leq \frac{n-1}{n} \|x\| + \frac{1}{n} \|x + \lambda' y\| \\ &< \frac{n-1}{n} \|x\| + \frac{1}{n} \|x\| \\ &= \|x\|. \end{aligned}$$

Thus we obtain a contradiction. Therefore, we conclude that the inequality $\|x + \lambda y\| \geq \|x\|$ must hold for every $\lambda \geq 0$. \square

3. PROPER ACUTE AND PROPER OBTUSE B-ANGLES

If X is equipped with an inner product $\langle \cdot, \cdot \rangle$ and the induced norm $\|x\| := \langle x, x \rangle^{1/2}$, we find that two nonzero vectors x and y form a proper acute angle if and only if there exists $\gamma > 0$ such that $\|x + \lambda y\| < \|x\|$ precisely for every $\lambda \in (-\gamma, 0)$. Likewise, x and y form a proper obtuse angle if and only if there exists $\gamma > 0$ such that $\|x + \lambda y\| < \|x\|$ precisely for every $\lambda \in (0, \gamma)$. The value of γ in both cases is given by $\gamma := \frac{2|\langle x, y \rangle|}{\|y\|^2}$. Our aim now is to formulate the criteria for proper acute and proper obtuse B-angles in a normed space $(X, \|\cdot\|)$.

Let $x, y \in X$. We define that

- x forms a **proper acute B-angle** to y , denoted by $x P A_B y$, if $x A_B y$ but $x \not\perp_B y$.
- x forms a **proper obtuse B-angle** to y , denoted by $x P O_B y$, if $x O_B y$ but $x \not\perp_B y$.

Note that, given $x, y \in X$, we now have three exclusive possibilities: $x P A_B y$, $x P O_B y$, or $x \perp_B y$.

As in inner product spaces, we obtain analogous results in normed spaces, as stated in the following theorem.

Theorem 3.1. *Let $x, y \in X$. Then there are only three (exclusive) possibilities for the set of values of λ for which the inequality (1.1) holds (or fails to hold), namely:*

- (1) *There exists $\gamma > 0$ such that the inequality (1.1) fails to hold precisely for every $\lambda \in (-\gamma, 0)$.*
- (2) *There exists $\gamma > 0$ such that the inequality (1.1) fails to hold precisely for every $\lambda \in (0, \gamma)$.*
- (3) *The inequality (1.1) holds for every $\lambda \in \mathbb{R}$.*

To prove the theorem, we consider the following set

$$S(x, y) := \{\lambda \in \mathbb{R} : \|x + \lambda y\| < \|x\|\},$$

for $x, y \in X$. The above theorem is a consequence of the following statements.

Theorem 3.2. *Let $x, y \in X$. Then each of the following statements hold:*

- (1) $0 \notin S(x, y)$.
- (2) *If there exists $\lambda > 0$ such that $\lambda \in S(x, y)$, then $(0, \lambda] \subseteq S(x, y)$ and $\mu \notin S(x, y)$ for every $\mu \leq 0$.*
- (3) *If there exists $\lambda < 0$ such that $\lambda \in S(x, y)$, then $[\lambda, 0) \subseteq S(x, y)$ and $\mu \notin S(x, y)$ for every $\mu \geq 0$.*
- (4) $S(x, y)$ is bounded.
- (5) *If $S(x, y) \neq \emptyset$, then $0 = \inf S(x, y) < \sup S(x, y)$ or $\inf S(x, y) < \sup S(x, y) = 0$.*
- (6) *If $S(x, y) \neq \emptyset$, then $\inf S(x, y) \notin S(x, y)$ and $\sup S(x, y) \notin S(x, y)$.*

Proof. (1) Obvious.

- (2) Let $\lambda > 0$ and $\lambda \in S(x, y)$. It follows from the first statement that $0 \notin S(x, y)$. Now, for $0 < \lambda' < \lambda$, we observe that

$$\begin{aligned} \|x + \lambda' y\| &= \left\| \frac{\lambda - \lambda'}{\lambda} x + \frac{\lambda'}{\lambda} x + \lambda' y \right\| \\ &\leq \frac{\lambda - \lambda'}{\lambda} \|x\| + \frac{\lambda'}{\lambda} \|x + \lambda y\| \\ &< \frac{\lambda - \lambda'}{\lambda} \|x\| + \frac{\lambda'}{\lambda} \|x\| \\ &= \|x\|, \end{aligned}$$

whence $\lambda' \in S(x, y)$. This proves that $(0, \lambda] \subseteq S(x, y)$. Next, suppose that there exists $\mu < 0$ such that $\mu \in S(x, y)$. Notice that for $t_0 = \frac{-\mu}{\lambda - \mu} \in (0, 1)$, we have

$$0 = (1 - t_0)\mu + t_0\lambda.$$

Hence

$$\begin{aligned} \|x\| &= \|x + [(1 - t_0)\mu + t_0\lambda]y\| \\ &\leq (1 - t_0)\|x + \mu y\| + t_0\|x + \lambda y\| \\ &< (1 - t_0)\|x\| + t_0\|x\| \\ &= \|x\|, \end{aligned}$$

which cannot be true. Therefore we conclude that $\mu \notin S(x, y)$ for every $\mu \leq 0$.

- (3) Similar to the proof of (2).
- (4) Assuming that $S(x, y) \neq \emptyset$, let $\lambda \in S(x, y)$. We observe that

$$\left| \|x\| - |\lambda| \|y\| \right| \leq \|x + \lambda y\| < \|x\|.$$

The assumption that $S(x, y) \neq \emptyset$ means that $x, y \neq 0$. Using the above inequality, we obtain

$$0 < |\lambda| < \frac{2\|x\|}{\|y\|}.$$

Hence $S(x, y)$ is bounded.

- (5) It follows from (2), (3), and (4).
- (6) Suppose that $S(x, y) \neq \emptyset$. Assume that there exists $\lambda^* > 0$ such that $\lambda^* \in S(x, y)$. By (2), (4), and (5), $\inf S(x, y) = 0 \notin S(x, y)$ and $\sup S(x, y)$ exists. Let $s := \sup S(x, y)$. Now the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(\lambda) = \|x + \lambda y\|$ is continuous everywhere, with

$$\begin{aligned} f(\lambda) &< \|x\| \text{ if } \lambda \in (0, s), \text{ and} \\ f(\lambda) &\geq \|x\| \text{ if } \lambda \leq 0 \text{ or } \lambda > s. \end{aligned}$$

Hence

$$\|x\| \leq \lim_{\lambda \rightarrow s^+} f(\lambda) = f(s) = \lim_{\lambda \rightarrow s^-} f(\lambda) \leq \|x\|.$$

Thus $\|x + sy\| = f(s) = \|x\|$, which implies that $\sup S(x, y) \notin S(x, y)$. By similar arguments, if there exists $\lambda^* < 0$ such that $\lambda^* \in S(x, y)$, then $\inf S(x, y) \notin S(x, y)$. \square

Remark 3.3. In view of Theorem 3.2, the value of γ in Theorem 3.1, part (1), equals $-\inf S(x, y)$; while that in part (2) equals $\sup S(x, y)$.

From Theorem 3.1, parts (1) and (2), we have the following corollary.

Corollary 3.4. Let $x, y \in X$. Then we have

- (1) $x PA_B y$ if and only if there exists $\gamma > 0$ such that (1.1) fails to hold precisely for every $\lambda \in (-\gamma, 0)$.
- (2) $x PO_B y$ if and only if there exists $\gamma > 0$ such that (1.1) fails to hold precisely for every $\lambda \in (0, \gamma)$.

As for acute and obtuse B -angles, we also have the following results for proper acute and proper obtuse B -angles.

Proposition 3.5. Let $x, y \in X \setminus \{0\}$ and $a, b \in \mathbb{R} \setminus \{0\}$.

- (1) Suppose that $x PA_B y$. If $ab > 0$, then $ax PA_B by$. If $ab < 0$, then $ax PO_B by$.
- (2) Suppose that $x PO_B y$. If $ab > 0$, then $ax PO_B by$. If $ab < 0$, then $ax PA_B by$.

4. COMPARING TWO B-ANGLES

Let $x, y \in X$. Throughout this section, we shall consider the case where $x PA_B y$. (The discussion for the case where $x PO_B y$ is similar.) For the purpose of our discussion here, we write $\gamma(x, y)$ for the largest value of γ for which the inequality (1.1) fails to hold precisely for every $\lambda \in (-\gamma, 0)$.

Let $x, y_1, y_2 \in X \setminus \{0\}$. If $x PA_B y_1$ and $x PO_B y_2$ (which includes the possibility that $x \perp_B y_2$), then it is safe to say that the B -angle from x to y_1 is more acute than that from x to y_2 . The question now is: what can we say when $x PA_B y_1$ and $x PA_B y_2$? How can we tell that one B -angle is more acute than the other? Suppose that $\gamma(x, y_1) < \gamma(x, y_2)$ (see the figure below).




Figure 4.1. The blue rays indicated the sets of the values of λ for which $\|x + \lambda y_1\| \geq \|x\|$ and $\|x + \lambda y_2\| \geq \|x\|$ hold, respectively.

Intuitively, we might want to conclude that the B -angle from x to y_2 is more acute than that from x to y_1 (that is, the larger the value of $\gamma(x, y)$, the more acute the B -angle from x to y). However, this conclusion is too early to make. Observe the following example in ℓ_2^∞ .

Let $x := (1, 0)$, $y_c := (c, c) \in \ell_2^\infty$, where $c > 0$. By observation, we see that $x PA_B y_c$ (see the figure below).




Figure 4.2. The vector x forms a proper acute B -angle to the vector y in ℓ_2^∞ .

Let us now take two different values of c , say $c_1 = 1$ and $c_2 = 2$, so that we have $y_1 := (1, 1)$ and $y_2 := (2, 2)$. By simple calculation, we obtain $\gamma(x, y_1) = 1$ and $\gamma(x, y_2) = \frac{1}{2}$. Thus $\gamma(x, y_2) < \gamma(x, y_1)$. However, we would **not** say that the B-angle from x to y_1 is more acute than that from x to y_2 since $y_2 = 2y_1$. In such a case, we would want to have the B-angle from x to y_1 **equal to** that from x to y_2 . To compare two proper B-acute angles, we need to find a number which in general depends on the two vectors but invariant under positive scalar multiplications. In order to do so, we go back to inner product spaces, to get an inspiration.

Suppose, for the moment, X is equipped with an inner product $\langle \cdot, \cdot \rangle$ and its induced norm $\|x\| := \langle x, x \rangle^{1/2}$. Let $x, y \in X \setminus \{0\}$, and consider the expression $\frac{\langle x, y \rangle}{\|x\| \|y\|}$, which is nothing but the cosine of the angle between x and y . We note that the value of this expression, and so is the angle, is invariant under positive scalar multiplications. Indeed, if $x' := ax$ and $y' := by$ with $a, b > 0$, then

$$\frac{\langle x', y' \rangle}{\|x'\| \|y'\|} = \frac{\langle ax, by \rangle}{\|ax\| \|by\|} = \frac{ab \langle x, y \rangle}{ab \|x\| \|y\|} = \frac{\langle x, y \rangle}{\|x\| \|y\|}.$$

Thus, normalizing both vectors, we get

$$\langle \hat{x}, \hat{y} \rangle = \frac{\langle x, y \rangle}{\|x\| \|y\|}$$

where $\hat{x} := \frac{x}{\|x\|}$ and $\hat{y} := \frac{y}{\|y\|}$.

We turn back to our normed space $(X, \|\cdot\|)$. Let $x, y \in X \setminus \{0\}$. Our goal is to find a number $\gamma^* = \gamma^*(x, y)$ such that $\gamma^*(ax, by) = \gamma^*(x, y)$ whenever $a, b > 0$. The key is the lemma below.

Lemma 4.1. *Let $x, y \in X \setminus \{0\}$ and suppose that $x PA_B y$. If $a, b > 0$, then $ax PA_B by$ and*

$$\gamma(ax, by) = \frac{a}{b} \gamma(x, y).$$

Proof. Let $a, b > 0$. By Proposition 3.5, we have $ax PA_B by$. Now suppose that $\lambda \in (-\gamma(x, y), 0)$, where we have $\|x + \lambda y\| < \|x\|$. By simple manipulations, we find that

$$\|ax + \frac{a}{b} \lambda (by)\| < \|ax\|.$$

Thus $\frac{a}{b} \lambda \in (-\gamma(ax, by), 0)$. Since we also have $\frac{a}{b} \lambda \in (-\frac{a}{b} \gamma(x, y), 0)$, we conclude that $\gamma(ax, by) \geq \frac{a}{b} \gamma(x, y)$. On the other hand, since $x = \frac{1}{a}(ax)$ and $y = \frac{1}{b}(by)$, we obtain $\gamma(ax, by) \leq \frac{a}{b} \gamma(x, y)$ via similar arguments. As a consequence, we arrive at the conclusion that

$$\gamma(ax, by) = \frac{a}{b} \gamma(x, y),$$

as claimed. \square

We are now ready to define the number γ^* with the desired property. Let $x, y \in X$ such that $x PA_B y$. Here of course $x, y \neq 0$, for otherwise x will be B-orthogonal to y per definition. We then define

$$(4.1) \quad \gamma^*(x, y) := \frac{\|y\|}{\|x\|} \gamma(x, y).$$

Accordingly, we have the following lemma.

Lemma 4.2. *Let $x, y \in X \setminus \{0\}$ and suppose that $x PA_B y$. If $a, b > 0$, then $\gamma^*(ax, by) = \gamma^*(x, y)$.*

Proof. Let $a, b > 0$. By (4.1), we have

$$\gamma^*(ax, by) = \frac{\|by\|}{\|ax\|} \gamma(ax, by) = \frac{b\|y\|}{a\|x\|} \frac{a}{b} \gamma(x, y) = \frac{\|y\|}{\|x\|} \gamma(x, y) = \gamma^*(x, y),$$

as desired. \square

Based on the above lemma, given $x, y \in X$ such that $x PA_B y$, we have $\gamma^*(x', y') = \gamma^*(x, y)$ for every $x' = ax$ and $y' = by$ with $a, b > 0$. Using this fact, we can now consider the unit vectors $\hat{x} := \frac{x}{\|x\|}$ and $\hat{y} := \frac{y}{\|y\|}$, where we have

$$\gamma^*(x, y) = \gamma^*(\hat{x}, \hat{y}) = \gamma(\hat{x}, \hat{y}).$$

This equality tells us that to compare acute B-angles, it suffices for us to compare the values of $\gamma(\cdot, \cdot)$ among the associated unit vectors. The same is also true for $x, y \in X$ for which $x PO_B y$. To be precise, we define the following.

Let $x, y_1, y_2 \in X \setminus \{0\}$ such that $x PA_B y_1$ and $x PA_B y_2$. Let \hat{x} , \hat{y}_1 , and \hat{y}_2 be the unit vectors associated to x, y_1 , and y_2 , respectively. We say that

- the B-angle from x to y_1 is **more acute** than that from x to y_2 if $\gamma(\hat{x}, \hat{y}_1) > \gamma(\hat{x}, \hat{y}_2)$.
- the B-angle from x to y_1 is **the same** as that from x to y_2 if $\gamma(\hat{x}, \hat{y}_1) = \gamma(\hat{x}, \hat{y}_2)$.
- the B-angle from x to y_1 is **more obtuse** than that from x to y_2 if $\gamma(\hat{x}, \hat{y}_1) < \gamma(\hat{x}, \hat{y}_2)$.

(Remark that similar definitions can be formulated for $x, y_1, y_2 \in X \setminus \{0\}$ such that $x PO_B y_1$ and $x PO_B y_2$.)

In the same spirit, we can also compare the B-angles for $x_1, x_2, y \in X \setminus \{0\}$ such that $x_1 PA_B y$ and $x_2 PA_B y$. Here we say that

- the B-angle from x_1 to y is **more acute** than that from x_2 to y if $\gamma(\hat{x}_1, \hat{y}) > \gamma(\hat{x}_2, \hat{y})$.
- the B-angle from x_1 to y is **the same** as that from x_2 to y if $\gamma(\hat{x}_1, \hat{y}) = \gamma(\hat{x}_2, \hat{y})$.
- the B-angle from x_1 to y is **more obtuse** than that from x_2 to y if $\gamma(\hat{x}_1, \hat{y}) < \gamma(\hat{x}_2, \hat{y})$.

(As in the previous case, similar definitions can be formulated for $x_1, x_2, y \in X \setminus \{0\}$ such that $x_1 PO_B y$ and $x_2 PO_B y$.)

Example 4.3. Let $x := (1, 0), y := (y_1, y_2) \in \ell_2^\infty$. Firstly, let us consider the case where $y := (a, 1)$ with $0 < a < 1$.




Figure 4.3. The vectors $x = (1, 0)$ and $y = (a, 1)$ with $0 < a < 1$ in ℓ_2^∞ .

Note that $\hat{x} = x$ and $\hat{y} = y$, and $x PA_B y$. Indeed, there exists $\gamma(x, y) > 0$ such that $\|x + \lambda y\| < \|x\| = 1$ precisely for every $\lambda \in (-\gamma(x, y), 0)$. To find the value of $\gamma(x, y)$, we solve the inequality

$$\max\{|1 + \lambda a|, |\lambda|\} = \|x + \lambda y\| < \|x\| = 1$$

for λ . We obtain that λ must satisfy $-2/a < \lambda < 0$ and $-1 < \lambda < 0$. Since $0 < a < 1$, we have $1/a > 1$, and so we find that $\lambda \in (-1, 0)$. Therefore $\gamma(x, y) = 1$, independent of the value of $a \in (0, 1)$. In this case, the B-angle from $x = (1, 0)$ to $y_1 = (a_1, 1)$ is the same as that from $x = (1, 0)$ to $y_2 = (a_2, 1)$ whenever $0 < a_1, a_2 < 1$.

Secondly, let us consider the case where $x := (1, 0), y := (1, a)$ with $0 < a \leq 1$.




Figure 4.4. The vectors $x = (1, 0)$ and $y = (1, a)$ with $0 < a \leq 1$ in ℓ_2^∞ .

Note that $\hat{y} = y$ and $x PA_B y$. The value of $\gamma(x, y) > 0$ for which $\|x + \lambda y\| < \|x\| = 1$ precisely for every $\lambda \in (-\gamma(x, y), 0)$ can be found by solving the inequality

$$\max\{|1 + \lambda|, |a\lambda|\} = \|x + \lambda y\| < \|x\| = 1$$

for λ . Here we obtain that $-2 < \lambda < 0$ and $-\frac{1}{a} < \lambda < \frac{1}{a}$, which yields $\lambda \in (-\min\{2, \frac{1}{a}\}, 0)$. For $0 < a \leq \frac{1}{2}$, we have $\frac{1}{a} \geq 2$, and so $\min\{2, \frac{1}{a}\} = 2$. For $\frac{1}{2} < a \leq 1$, we have $1 \leq \frac{1}{a} < 2$, which gives $\min\{2, \frac{1}{a}\} = \frac{1}{a}$. Therefore $\gamma(x, y) = 2$ for $a \in (0, 1/2]$ and $\gamma(x, y) = \frac{1}{a}$ for $a \in (\frac{1}{2}, 1]$. In this example, the B-angle from $x = (1, 0)$ to $y_1 = (1, a_1)$ is more acute than that from $x = (1, 0)$ to $y_2 = (1, a_2)$ provided that $0 < a_1 \leq \frac{1}{2} < a_2 \leq 1$.

Example 4.4. Let $x := (1, 0), y = (a, \sqrt{1 - a^2}) \in \ell_2^2$ with $0 < a < 1$.




Figure 4.5. The vectors $x = (1, 0)$ and $y = (a, \sqrt{1 - a^2})$ with $0 < a < 1$ in ℓ_2^2 .

Observe that $\|x\| = \|y\| = 1$ and $x PA_B y$. We can find the value of $\gamma(x, y) > 0$ for which $\|x + \lambda y\| < \|x\| = 1$ precisely for every $\lambda \in (-\gamma(x, y), 0)$. By simple computations, one may obtain that

$$\gamma(x, y) = \frac{2\langle x, y \rangle}{\|y\|^2} = \frac{2a}{a^2 + (1 - a^2)} = 2a.$$

Thus here $\gamma(x, y)$ gets larger as a tends to 1, that is, as y is approaching x .

5. CONCLUDING REMARKS: THE ANALOG OF THE COSINE OF B-ANGLES

We have defined acute and obtuse B-angles in a normed space $(X, \|\cdot\|)$ via the sets of values of λ for which the inequality (1.1) holds (or fails to hold), and defined the criteria for proper acute and proper obtuse B-angles from a vector x to another vector y in X . We have also defined a number $\gamma^*(\cdot, \cdot)$ which can be used to compare the B-angle from x to y_1 and that from x to y_2 , or from x_1 to y and that from x_2 to y . It is then tempting to have a formula for the B-angle from a vector to another vector in X , as in an inner product space.

As the readers might have guessed by now, the formula is only one step away from the formula of $\gamma(\cdot, \cdot)$ on the unit sphere. Let $x, y \in X \setminus \{0\}$. Define

$$k(x, y) = \begin{cases} \frac{1}{2}\gamma(\hat{x}, \hat{y}), & \text{if } x PA_B y; \\ 0, & \text{if } x \perp_B y; \\ -\frac{1}{2}\gamma(\hat{x}, \hat{y}), & \text{if } x PO_B y. \end{cases}$$

Notice that if X is an inner product space which is also equipped with the induced norm, then $k(x, y) = \langle \hat{x}, \hat{y} \rangle$, which is nothing but the cosine of the angle between x and y .

The following proposition gives basic properties of the function $k(x, y)$, which may be viewed as the **cosine of the B-angle** from x to y .

Proposition 5.1. *Let $x, y \in X \setminus \{0\}$ and $a, b \neq 0$. Then the following statements hold:*

- (1) $|k(x, y)| \leq 1$.
- (2) *If $ab > 0$, then $k(ax, by) = k(x, y)$. If $ab < 0$, then $k(ax, by) = -k(x, y)$.*

Proof. (1) Suppose that we have some values of λ satisfying $\|\hat{x} + \lambda\hat{y}\| < 1$. Then

$$\|\hat{x}\| - |\lambda| \|\hat{y}\| \leq \|\hat{x} + \lambda\hat{y}\| < 1.$$

Hence we have $|1 - |\lambda|| < 1$, whence $0 < |\lambda| < 2$. This implies that $\gamma(\hat{x}, \hat{y}) \leq 2$, and therefore $|k(x, y)| = \frac{1}{2}\gamma(\hat{x}, \hat{y}) \leq 1$.

(2) We shall only prove it for the case where $ab < 0$ and $x PA_B y$, and leave the other cases to the readers. Without loss of generality, assume that $a > 0 > b$. The hypothesis that $x PA_B y$ means that there exists $\gamma(\hat{x}, \hat{y}) > 0$ such that $\|\hat{x} + \lambda\hat{y}\| < 1$ precisely for every $\lambda \in (-\gamma(\hat{x}, \hat{y}), 0)$. Since $a > 0 > b$, we have $ax PO_B by$, which means that there exists $\gamma(\widehat{ax}, \widehat{by}) > 0$ such that $\|\widehat{ax} + \lambda\widehat{by}\| < 1$ precisely for every $\lambda \in (0, \gamma(\widehat{ax}, \widehat{by}))$. Now the assumption that $a > 0 > b$ implies that $\widehat{ax} = \hat{x}$ and $\widehat{by} = -\hat{y}$. For $0 < \lambda' < \gamma(\hat{x}, \hat{y})$, set $\lambda = -\lambda'$. Then $-\gamma(\hat{x}, \hat{y}) < \lambda < 0$, and so we obtain

$$\|\widehat{ax} + \lambda'(\widehat{by})\| = \|\hat{x} + \lambda'(-\hat{y})\| = \|\hat{x} + \lambda\hat{y}\| < 1.$$

This implies that $\gamma(\hat{x}, \hat{y}) \leq \gamma(\widehat{ax}, \widehat{by})$. Conversely, by setting $x' = ax, y' = by$, we have $x = \frac{1}{a}x', y = \frac{1}{b}y'$, so that $\gamma(\hat{x}, \hat{y}) \geq \gamma(\widehat{ax}, \widehat{by})$. Therefore $\gamma(\widehat{ax}, \widehat{by}) = \gamma(\hat{x}, \hat{y})$. Consequently, we obtain

$$k(x, y) = \frac{1}{2}\gamma(\hat{x}, \hat{y}) = \frac{1}{2}\gamma(\widehat{ax}, \widehat{by}) = -k(ax, by),$$

as expected. □

Both properties in the above proposition are basic properties of the cosine of the B-angles in X . We miss, however, the symmetric property $k(x, y) = k(y, x)$. To see that we do not have this property, take for an example $x := (1, 0)$ and $y = (1, 1)$ in ℓ_2^∞ . Here $x PA_B y$ and $y \perp_B x$, and thus $k(x, y) > 0 = k(y, x)$. This example tells us that, in general, $k(x, y) \neq k(y, x)$. (This is why we do not use the phrase the B-angle **between** x **and** y , but the B-angle **from** x **to** y .)

Furthermore, given a vector $x \in X$, the set of vectors y such that $k(x, y) = 1$ may consists more than just $y = x$. We have seen this in Example 4.3 where $X = \ell_2^\infty$, $x := (1, 0)$ and $y := (1, a)$ with $a \in [0, 1/2)$. The same also happens with the set of vectors y such that $k(x, y) = -1$.

Similar to Example 4.3, we can also compute the values of $k(x, y)$ for $x := (1, 0)$ and $y := (\cos \theta, \sin \theta) \in \ell_2^1$ with $\theta \in (-\pi, \pi]$. Here k can be seen as a function of θ , say $k = f(\theta)$, where

$$f(\theta) := \begin{cases} 1, & -\frac{\pi}{4} < \theta < \frac{\pi}{4}; \\ 0, & -\frac{3\pi}{4} \leq \theta \leq -\frac{\pi}{4} \text{ or } \frac{\pi}{4} \leq \theta \leq \frac{3\pi}{4}; \\ -1, & -\pi < \theta < -\frac{3\pi}{4} \text{ or } \frac{3\pi}{4} < \theta \leq \pi. \end{cases}$$

The graph of $k = f(\theta)$ is presented below:




Figure 5.1. The graph of k as a function of θ for $\theta \in (-\pi, \pi]$.

Acknowledgements. This work is supported by P2MI-ITB 2021 Program.

REFERENCES

- [1] J. Alonso, H. Martini, and S. Wu, “On Birkhoff orthogonality and isosceles orthogonality in normed linear spaces”, *Aequat. Math.* **83**, No. 2 (2012), 153–189.
- [2] V. Balestro, A.G. Horvath, H. Martini, and R. Teixeira, “Angles in normed spaces”, *arXiv:1607.06938v1* [math.MG]
- [3] G. Birkhoff, “Orthogonality in linear metric spaces”, *Duke Math. J.* **1**, No. 2 (1935), 169–172.

- [4] H. Gunawan, J. Lindiarni, and O. Neswan, “ P -, I -, g -, and D - angles in normed spaces”, *ITB J. Sci.* **40A**, No. 1 (2008), 24–32.
- [5] M. Jamaludin, *Sudut Birkhoff pada Ruang Bernorma* (in Indonesian), Final Project Report, Mathematics Undergraduate Program, Bandung Institute of Technology, 2021.
- [6] R.C. James, “Orthogonality in normed linear spaces”, *Duke Math. J.* **12** (1945), 291–302.
- [7] R.C. James, “Orthogonality and linear functionals in normed linear spaces”, *Trans. Amer. Math. Soc.* **61** (1947), 265–292.
- [8] P. Miličić, “On the B-angle and g -angle in normed spaces”, *JIPAM* Vol. **8**, Iss. 4, art. 99 (2007).
- [9] J.R. Partington, “Orthogonality in normed spaces”, *Bull. Austral. Math. Soc.* **33** (1986), 449–455.
- [10] M.D. Pratamadirdja, *Ortogonalitas dan Karakterisasi Sudut Birkhoff di ℓ_N^1* (in Indonesian), Final Project Report, Mathematics Undergraduate Program, Bandung Institute of Technology, 2020.
- [11] C. Zhi-Zhi, L. Wei, L. Lü-lin, and J. You-qing, “Projections, Birkhoff orthogonality and angles in normed spaces”, *Comm. Math. Res.* **27**, No. 4 (2011).

FACULTY OF MATHEMATICS AND NATURAL SCIENCES, BANDUNG INSTITUTE OF TECHNOLOGY, BANDUNG 40132, INDONESIA

Email address: hgunawan@math.itb.ac.id

FACULTY OF MATHEMATICS AND NATURAL SCIENCES, BANDUNG INSTITUTE OF TECHNOLOGY, BANDUNG 40132, INDONESIA

Email address: mjamaludin@students.itb.ac.id

FACULTY OF MATHEMATICS AND NATURAL SCIENCES, BANDUNG INSTITUTE OF TECHNOLOGY, BANDUNG 40132, INDONESIA

Email address: pratamadirdja@students.itb.ac.id