# AN OBSERVATION ON UNBOUNDED LINEAR OPERATORS WITH AN ARBITRARY SPECTRUM

MARAT V. MARKIN

ABSTRACT. We furnish a simple way of constructing an unbounded closed linear operator in a complex Banach space, whose spectrum is an arbitrary nonempty closed, in particular compact, subset of the complex plane.

## 1. INTRODUCTION

By *Gelfand's spectral radius theorem* (see, e.g., [3]), the spectrum of a bounded linear operator on a complex Banach space is a nonempty compact subset of the complex plane. Conversely, an arbitrary nonempty compact set in  $\mathbb{C}$  is the spectrum of a such an operator (cf. [3, Theorem 5.7]).

While the spectrum of an unbounded closed linear operator in a complex Banach space is known to be a closed subset of the complex plane, including  $\emptyset$  and  $\mathbb{C}$ , (see, e.g., [1,3]), the natural question is whether an arbitrary closed set in  $\mathbb{C}$  is the spectrum of a such an operator. Of particular interest are unbounded closed linear operators with nonempty compact spectra.

We give a simple way of constructing an unbounded closed linear operator in a complex Banach space, whose spectrum is any given nonempty closed, in particular compact, subset of the complex plane.

### 2. Preliminaries

For a closed linear operator A in a complex Banach space X, the set

$$\rho(A) := \{\lambda \in \mathbb{C} \mid \exists R(\lambda, A) := (A - \lambda I)^{-1} \in L(X) \}$$

1

(*I* is the *identity operator* on *X*, L(X) is the space of bounded linear operators on *X*) and its complement  $\sigma(A) := \mathbb{C} \setminus \rho(A)$  are called the operator's *resolvent set* and *spectrum*, respectively.

The spectrum is partitioned into three pairwise disjoint subsets,  $\sigma_p(A)$ ,  $\sigma_c(A)$ , and  $\sigma_r(A)$ , called the *point*, *continuous*, and *residual spectrum* of A, respectively, as follows:

$$\begin{split} \sigma_p(A) &:= \left\{ \lambda \in \mathbb{C} \, | \, A - \lambda I \text{ is not injective, i.e., } \lambda \text{ is an eigenvalue of } A \right\},\\ \sigma_c(A) &:= \left\{ \lambda \in \mathbb{C} \, \Big| \, A - \lambda I \text{ is injective, not surjective, and } \overline{R(A - \lambda I)} = X \right\},\\ \sigma_r(A) &:= \left\{ \lambda \in \mathbb{C} \, \Big| \, A - \lambda I \text{ is injective and } \overline{R(A - \lambda I)} \neq X \right\} \end{split}$$

<sup>2020</sup> Mathematics Subject Classification. Primary 47A10; Secondary 47A08, 47B15.

Key words and phrases. Bounded linear operator, closed linear operator, normal operator, spectrum.

 $(R(\cdot))$  is the range of an operator and  $\overline{\cdot}$  is the closure of a set) (see, e.g., [1,3]).

In the complex Hilbert space  $l_2$  of square-summable sequences, the operator of multiplication by a sequence  $(m_n)_{n \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}}$ 

$$M\left(x_{n}\right)_{n\in\mathbb{N}}:=\left(m_{n}x_{n}\right)_{n\in\mathbb{N}}$$

with maximal domain

$$D(M) := \{ (x_n)_{n \in \mathbb{N}} \in l_2 \mid (m_n x_n)_{n \in \mathbb{N}} \in l_2 \}$$

is a (densely defined) closed linear operator, which is bounded  $i\!f\!f$  the multiplier sequence is bounded, i.e.,

$$\sup_{n\in\mathbb{N}}|m_n|<\infty.$$

In fact, the operator M is *normal* (see, e.g., [2]) and

$$\sigma(M) = \overline{\{m_n\}}$$

with

$$\sigma_p(M) = \{m_n\}, \ \sigma_c(M) = \{m_n\} \setminus \{m_n\}, \ \text{and} \ \sigma_r(M) = \emptyset,$$

where  $\{m_n\}$  is the set of values of  $(m_n)_{n \in \mathbb{N}}$  [3].

Thus, by choosing such multiplier sequence  $(m_n)_{n \in \mathbb{N}}$  that  $\{m_n\}_{n \in \mathbb{N}}$  is a countable dense subset of an arbitrary nonempty closed set  $\sigma$  in the complex plane, we obtain the multiplication operator M in  $l_2$  with

$$\sigma(M) = \sigma$$

the normal operator M being bounded whenever the set  $\sigma$  is compact (see, e.g., [2]).

In the complex Banach space  $Y := L_p(0,1)$   $(1 \le p < \infty)$  or Y := C[0,1], the latter equipped with the maximum norm

$$C[0,1] \ni x \mapsto ||x||_{\infty} := \max_{0 \le t \le 1} |x(t)|,$$

the differentiation operator

$$Dx := x'$$

with domain

$$D(D) := \{ x \in L_p(0,1) \, | \, x(\cdot) \in AC[a,b], \ x' \in L_p(0,1), \ x(0) = 0 \}$$

or

$$D(D):=\left\{x\in C^1[0,1]\,\big|\,x(0)=0\right\},$$

respectively, is an unbounded closed linear operator with

 $\sigma(D) = \emptyset$ 

and

$$[R(\lambda, D)y](t) = \int_0^t e^{\lambda(t-s)}y(s) \, ds, \ \lambda \in \mathbb{C}, y \in Y,$$

(cf. [3]), the operator being densely defined in  $L_p(0,1)$   $(1 \le p < \infty)$  but not in C[0,1].

## 3. UNBOUNDED LINEAR OPERATOR WITH AN ARBITRARY SPECTRUM

Let  $\sigma$  be arbitrary nonempty closed set in the complex plane, M be the multiplication operator in the complex space  $l_2$  relative to a multiplier sequence  $(m_n)_{n \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}}$ such that  $\{m_n\}$  is a countable dense subset of  $\sigma$ , and hence,

$$\sigma(M) = \overline{\{m_n\}} = \sigma,$$

and D be the differentiation operator in the complex space  $L_p(0,1)$   $(1 \le p < \infty)$  or C[0,1] with  $\sigma(D) = \emptyset$  (see Preliminaries).

In the complex Banach space  $X \oplus Y$ , where  $X := l_2$  and  $Y := L_p(0,1)$   $(1 \le p < \infty)$  or Y := C[0,1], equipped with the norm

$$X \oplus Y \in (x, y) \mapsto ||(x, y)|| := ||x||_X + ||y||_Y,$$

the operator matrix

$$\begin{bmatrix} M & 0 \\ 0 & D \end{bmatrix}$$

defines an unbounded closed linear operator A, whose domain

$$D(A) := D(M) \oplus D(D)$$

is dense in  $X \oplus Y$  whenever D(D) is dense in Y, with

$$\sigma(A) = \sigma(M) \cup \sigma(D) = \sigma(M) = \overline{\{m_n\}} = \sigma$$

and, as is easily seen,

$$\sigma_p(A) = \sigma_p(M) = \{m_n\}, \ \sigma_c(A) = \sigma_c(M) = \overline{\{m_n\}} \setminus \{m_n\}, \ \text{and} \ \sigma_r(A) = \emptyset$$

(see Preliminaries).

# Remarks 3.1.

- Thus, the spectrum of an unbounded closed linear operator in a complex Banach space can be an arbitrary closed subset of the complex plane, including Ø and C, (cf. [3, Theorem 5.7]).
- If, in the prior example,  $Y := L_2(0, 1), X \oplus Y$  is a Hilbert space relative to the inner product

$$X \oplus Y \in (x_1, y_1), (x_2, y_2) \mapsto \langle (x_1, y_1), (x_2, y_2) \rangle := \langle x_1, x_2 \rangle_X + \langle y_1, y_2 \rangle_Y.$$

### References

- N. Dunford and J.T. Schwartz with the assistance of W.G. Bade and R.G. Bartle, *Linear Operators. Part I: General Theory*, Interscience Publishers, New York, 1958.
- [2] \_\_\_\_\_, Linear Operators. Part II: Spectral Theory. Self Adjoint Operators in Hilbert Space, Interscience Publishers, New York, 1963.
- [3] M.V. Markin, *Elementary Operator Theory*, De Gruyter Graduate, Walter de Gruyter GmbH, Berlin/Boston, 2020.

DEPARTMENT OF MATHEMATICS CALIFORNIA STATE UNIVERSITY, FRESNO 5245 N. BACKER AVENUE, M/S PB 108 FRESNO, CA 93740-8001, USA

 $Email \ address: mmarkin@csufresno.edu$