

REGULARITY AND ENERGY OF SECOND ORDER HYPERBOLIC BOUNDARY VALUE PROBLEMS ON NON-TIMELIKE HYPERSURFACES

SHIQI MA

ABSTRACT. We study the second order hyperbolic equations with initial conditions, a nonhomogeneous Dirichlet boundary condition and a source term. We prove the solution possesses H^1 regularity on any piecewise C^1 -smooth non-timelike hypersurfaces. We generalize the notion of energy to these hypersurfaces, and establish an estimate of the difference between the energies on the hypersurface and on the initial plane where the time $t = 0$. The energy is shown to be conserved when the source term and the boundary datum are both zero. We also obtain an L^2 estimate for the normal derivative of the solution. In the proofs we first show these results for C^2 -smooth solutions by using the multiplier methods, and then we go back to the original results by approximation.

Keywords: Hyperbolic equations, regularity, non-timelike hypersurface, energy estimates, normal derivative.

2020 Mathematics Subject Classification: 35L10, 35L20, 35B65, 35B38, 83A05.

1. INTRODUCTION

Throughout the article we assume that T is a fixed positive constant, Ω is an open domain in \mathbb{R}^n ($n \geq 1$) with C^∞ -smooth boundary, and $A(x) = (a_{ij}(x)) \in C^{n+4}(\bar{\Omega}; \mathbb{R}^{n^2})$ is a real-valued symmetric $n \times n$ matrix function. We focus on the following second order hyperbolic initial/boundary value problem,

$$\begin{cases} \partial_t^2 u - \nabla \cdot (A(x)\nabla u) = G & \text{in } Q := \Omega \times (0, T), \\ u = f & \text{on } \Sigma := \partial\Omega \times (0, T), \\ u = u_0, u_t = u_1 & \text{on } \Omega \times \{t = 0\}, \end{cases} \quad (1.1)$$

under the condition

$$\begin{cases} G \in L^2(Q), f \in H^1(\Sigma) := L^2(0, T; H^1(\partial\Omega)) \cap H^1(0, T; L^2(\partial\Omega)), \\ u_0 \in H^1(\Omega), u_1 \in L^2(\Omega), u_0(x) = f(x, 0) \text{ on } \partial\Omega. \end{cases} \quad (1.2)$$

The existence and regularity results of the solution of (1.1) under variant initial/boundary data can be found in the literature. The existence of the unique solution $u \in C([0, T]; H^1(\Omega))$ of (1.1) under the condition (1.2) is given in [LLT86, (3.5)]. In [LT81] the authors proved another existence result for $f \in L^2(0, T; L^2(\partial\Omega))$ and $G = 0$ by using the cosine operators technique, and they also showed the map from f to u is continuous. Then, in [LT83] the same authors improved the regularity of u from $L^2(0, T; L^2(\Omega))$ to $C([0, T]; L^2(\Omega))$. In [Sak70a, Sak70b] Sakamoto studied the problem (1.1) with higher order regularities by using pseudo-differential operator. The book [LM72] contains a comprehensive treatments of the non-homogeneous boundary value problems, including hyperbolic equations. See also [Ava97, LT88] for related applications. These studies mentioned above treat the solution u as maps $[0, T] \rightarrow X$ where X are function spaces defined in Ω , namely, the cylinder Q are foliated horizontally and u is defined in every horizontal slice $\Omega \times \{t = \tau\}$ for $\forall \tau \in [0, T]$.




FIGURE 1. An example of the non-timelike hypersurfaces Γ_S .

In this article, we investigate the properties of the solution of (1.1)-(1.2) on *non-horizontal* hypersurfaces Γ_S described by

$$\Gamma_S := \{(x, S(x)); x \in \overline{\Omega}\}.$$

Check Fig. 1 as an example. The restrictions of the solution on slanted hypersurfaces have already appeared in the literature. In [RU14], the authors studied the equation $(\partial_t^2 - \Delta + q)U = 0$ incited by an incident wave $\delta(t - x_1)$. When encountered with the potential q , the incident wave generates a scattered wave u such that

$$U(x, t) = \delta(t - x_1) + u(x, t)H(t - x_1)$$

where H is the Heaviside function. [RU14, Theorem 1] proves that the scattered wave u is also a solution of the equation $(\partial_t^2 - \Delta + q)u = 0$ and

$$u(t, x', t) = -\frac{1}{2} \int_{-\infty}^t q(s, x') ds, \quad \forall (x', t) \in \mathbb{R}^n.$$

The expression above involves the restriction of u on the lightlike hypersurface $\{(t, x', t)\}$ of \mathbb{R}^{n+1} . Similar situations also appeared in [RS20a, RS20b, MPMS21], and in [MS21] in the presence of a Riemannian metric.

When regarding Γ_S as a submanifold of Q , the upward normal vector ν of Γ_S is given by

$$\nu := (\nu_x, \nu_t) = \left(\frac{-\nabla S}{\sqrt{1 + |\nabla S|^2}}, \frac{1}{\sqrt{1 + |\nabla S|^2}} \right). \quad (1.3)$$

We say a hypersurface is timelike (resp. lightlike, spacelike) with respect to A at a point p if and only if the normal vector ν at p satisfies

$$|\nu_x|_A < |\nu_t|, \quad (\text{resp. } |\nu_x|_A = |\nu_t|, \quad |\nu_x|_A > |\nu_t|), \quad \text{where } |\xi|_A := \bar{\xi}^T A(x)\xi.$$

And the hypersurface is said to be timelike (resp. lightlike, spacelike) if it is timelike (resp. lightlike, spacelike) at every point. In this article we restrict our attentions only to non-timelike (i.e. lightlike or spacelike) hypersurfaces. Therefore, we put the following assumption.

Assumption 1.1. $S \in C^1(\overline{\Omega}, [0, T])$ piecewisely¹, and Γ_S is a non-timelike hypersurface, i.e. and $|\nabla S(x)|_A \leq 1$ for $\forall x \in \overline{\Omega}$.

We also assume that $-\nabla \cdot (A(x)\nabla)$ is an elliptic operator, namely,

Assumption 1.2. There exist two positive constants c_1, c_2 such that for $\forall \xi \in \mathbb{C}^n$ and $\forall x \in \overline{\Omega}$,

$$c_1|\xi|^2 \leq |\xi|_A^2 := \bar{\xi}^T A(x)\xi \leq c_2|\xi|^2.$$

¹For convenience, we shall say “ $S \in C^1(\overline{\Omega}, [0, T])$ piecewisely” if S is piecewisely C^1 -smooth in $\overline{\Omega}$ and its range is contained in $[0, T]$.

1.1. Main results. In this work we establish H^1 regularity and energy estimates for the solution of (1.1)-(1.2) on any non-timelike hypersurfaces. The energy is estimated not directly on the value of the energy itself, but on the *difference* between the energy on the hypersurface and the energy at time $t = 0$. The estimate of the difference is sharper than the estimate of the value of the energy itself, see Theorem 1.1 and the succeeding discussion below for details.

To state the main results, we introduce several notations. We define the energy $\mathcal{E}(u; \Gamma_S)$ of u on Γ_S whenever the following expression can be well-defined:

$$\mathcal{E}(u; \Gamma_S) := \int_{\Omega} [|\nabla(u(x, S(x)))|_A^2 + (1 - |\nabla S(x)|_A^2)|u_t(x, S(x))|^2] dx. \quad (1.4)$$

When $S(x) = \text{constant}$, (1.4) coincides with the classical energy definition. The first result involves (1.1) underneath Γ_S , so let us introduce the following notations (see Fig. 2):

$$\begin{cases} Q_{\tau} := \{(x, t); x \in \Omega, \tau \leq t \leq S(x)\} \subset Q, & \Sigma_{\tau} := \{(x, t); x \in \partial\Omega, \tau \leq t \leq S(x)\} \subset \Sigma, \\ H_{\tau} := Q_0 \cap \{(x, \tau); x \in \Omega\}, & \Gamma_{S, \tau} := \{(x, S(x)); \tau \leq S(x) \leq T_2\} \\ T_1 := \inf\{S(x); x \in \Omega\}, & T_2 := \sup\{S(x); x \in \Omega\}. \end{cases}$$

Note that when $\tau \leq T_1$, $\Gamma_{S, \tau} = \Gamma_S$. And when Γ_S is not horizontal, Q_0 and Σ_0 will be strict subsets of Q and Σ , respectively.




FIGURE 2. Left: the cylinder Q . Right: the part of Q underneath Γ_S .

In what follows we use the notation $\langle T_2 \rangle := (1 + |T_2|^2)^{1/2}$ for simplicity.

Theorem 1.1. *Given Assumptions 1.1 and 1.2, then in the system (1.1)-(1.2), the restriction of the solution u on Γ_S is in $H^1(\Gamma_S)$, and $\mathcal{E}(u; \Gamma_S)$ is well-defined. Moreover, there holds*

$$\begin{aligned} |\mathcal{E}(u; \Gamma_S) - \int_{\Omega} (|\nabla u_0|_A^2 + |u_1|^2) dx| \\ \leq C \langle T_2 \rangle^{1/2} [\|\nabla u_0\|_{L^2(\Omega)} + \|u_1\|_{L^2(\Omega)} + \langle T_2 \rangle^{1/2} (\|f\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)})] \\ \times (\|f\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)}), \end{aligned}$$

for some constant C depending only on A and the dimension n .

Remark 1.1. When Γ_S is not horizontal, we have $Q_0 \subsetneq Q$ and $\Sigma_0 \subsetneq \Sigma$, so the estimate given in Theorem 1.1 only requires parts of the data f and G . When Γ_S is strictly spacelike, i.e. $|\nabla S(x)|_A \leq C < 1$ for all $x \in \bar{\Omega}$ for some constant C , Theorem 1.1 implies $u_t|_{\Gamma_S} \in L^2(\Gamma_S)$ as well.

In [Tay11, Chapter 2 Fig. 6.1, 6.2, 8.1] Taylor studied the spacelike cases where the lateral boundary is an empty set. The lightlike case is not covered by [Tay11]. Also, the results given in [Tay11] is given in a form of (1.5) below, and the inequality in Theorem 1.1 seems

to be new. In some applications it is more convenient to use the following estimate, which is a direct consequence of Theorem 1.1,

$$\mathcal{E}(u; \Gamma_S) \leq C(\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) + C\langle T_2 \rangle (\|f\|_{H^1(\Sigma_0)}^2 + \|G\|_{L^2(Q_0)}^2). \quad (1.5)$$

But readers should note that the estimate given in Theorem 1.1 is sharper than (1.5).

The energy can also be defined on different non-timelike hypersurfaces. Denote $S_0(x) \equiv 0$ and $\Gamma_0 := \{(x, S_0(x)); x \in \bar{\Omega}\}$ so that $\Gamma_0 = \bar{\Omega} \times \{t = 0\}$. And we define a family of hypersurfaces $\{\Gamma_\tau\}_{\tau>0}$ (see Fig. 3) by

$$\Gamma_\tau := \{(x, S_\tau(x)); x \in \bar{\Omega}\},$$

with S_τ satisfying the following requirement,

$$S_\tau \in C^1(\bar{\Omega}; [0, T]) \text{ piecewisely, such that for } \tau_1 < \tau_2, S_{\tau_1}(x) \leq S_{\tau_2}(x) \forall x \in \bar{\Omega}. \quad (1.6)$$




FIGURE 3. Non-timelike hypersurfaces

If for every fixed $x \in \bar{\Omega}$, the function $S_\tau(x)$ is continuous with respect to τ , then the family $\{\Gamma_\tau\}_{\tau \geq 0}$ will form another foliation of \bar{Q} comparing to the standard foliation $\Omega \times \{t = \tau\}$ with $\tau \in [0, T]$. We define the corresponding energy on Γ_τ whenever it can be well-defined:

$$\mathcal{E}(u; \Gamma_\tau) := \int_{\Omega} [|\nabla(u(x, S_\tau(x)))|_A^2 + (1 - |\nabla S_\tau(x)|_A^2)|u_t(x, S_\tau(x))|^2] dx. \quad (1.7)$$

By Theorem 1.1, the following corollary about $\mathcal{E}(u; \Gamma_\tau)$ is an immediate result.

Corollary 1.2. *Given Assumptions 1.1 and 1.2, and assuming the family $\{\Gamma_\tau\}_{\tau \geq 0}$ satisfies (1.6), then in the system (1.1)-(1.2), when $G = 0$ and $f = 0$, the energy $\mathcal{E}(u; \Gamma_\tau)$ is well-defined and is conserved, i.e.*

$$\mathcal{E}(u; \Gamma_\tau) = \int_{\Omega} (|\nabla u_0|_A^2 + |u_1|^2) dx, \text{ for } \tau \geq 0.$$

Corollary 1.2 generalizes the classical energy conservation law, which says the energy is conserved on every horizontal surface $\Omega \times \{t = \tau\}$. It also generalizes the energy conservation in [Tay11, §2.5 & §2.6] to the complex-valued case.

We also obtain an estimate of the conormal derivative (with respect to A) $u_{\nu, A} := \nu_\Sigma \cdot A(x)\nabla u$, where ν_Σ signifies the outer unit normal vector to Σ .

Theorem 1.3. *Under the same assumptions as in Theorem 1.1, we have*

$$\|u_{\nu, A}\|_{L^2(\Sigma_0)}^2 \leq C\langle T_2 \rangle (\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) + C\langle T_2 \rangle^2 (\|G\|_{L^2(Q_0)}^2 + \|f\|_{H^1(\Sigma_0)}^2),$$

for some constant C depending only on A and the dimension n .

Remark 1.2. When $S(x) = \text{constant}$, the hypersurface Γ_S will be horizontal and so $Q_0 = Q$ and $\Sigma_0 = \Sigma$. In this case, the estimate about $u_{\nu,A}$ is proved in [LLT86, (4.7)] (note that $|u_{\nu,A}| \leq C|u_\nu|$). However, when Γ_S is not horizontal, Theorem 1.3 seems to be new.

The proofs of Theorems 1.1, 1.3 and Corollary 1.2 are presented in Section 4.

1.2. The motivation. The classical result [LLT86, Remark 2.10] says that the solution u of (1.1)-(1.2) satisfies

$$u \in C([0, T]; H^1(\Omega)), \quad \partial_t u \in C([0, T]; L^2(\Omega)),$$

which implies

$$u \in H^1(Q).$$

Therefore by the trace theorem, the restriction of u to Γ_S has regularity $H^{1/2}$. However, when Γ_S is horizontal, e.g. when $\Gamma_S = \Omega \times \{t = \tau\}$ for some τ , the restriction of u on Γ_S has H^1 regularity due to [LLT86, Remark 2.10]. From the point of view of the relativity of simultaneity in the theory of relativity [Wal84], $\Omega \times \{t = \tau\}$ should not be more special than other non-timelike hypersurfaces Γ_S . Therefore, one would expect that the trace theorem for the solution u is not sharp and u shall also enjoy the same H^1 regularity on slanted Γ_S as on $\Omega \times \{t = \tau\}$. This is the motivations of the work.




FIGURE 4. The observers X and Y stand at the origin O at the time $\tau = 0$. X stands still while Y possesses an instantaneous velocity v pointing to the right. $\{t, x_1\}$ is the instantaneous coordinate of X and $\{\tilde{t}, \tilde{x}_1\}$ is that of Y . The time and space axes are perpendicular to each other for both $\{t, x_1\}$ and $\{\tilde{t}, \tilde{x}_1\}$, under the Minkowski metric $-dt^2 + dx_1^2$.

Let us explain the idea by a simplified example. Assume there is a sound wave $u(x, t)$ propagating inside a domain Ω which satisfies the wave equation $(\partial_t^2 - \Delta)u = 0$, and the sound speed is normalized to 1 in the medium, i.e. $A(x) = 1$ in Ω . An observer X is located inside Ω and he/she stands still relative to Ω . Another observer Y is also located inside Ω but is moving in the direction of the first axis, say, x_1 , at a constant speed v which is slower than that sound speed, $|v| < 1$. We assume the Ω is large enough such that all the events mentioned here take place inside $\Omega \times [0, T]$ for T large enough. Then, from the perspective of X , the simultaneity at time τ is $\Omega \times \{t = \tau\}$, while from the perspective of Y the simultaneity is a slanted plane Γ_S with $S(x) = vx$, see Fig. 4. Let (t, x) be the spacetime coordinate of X and we denote $\gamma = (1 - v^2)^{-1/2}$ as the Lorentz factor. Then according to the theory of relativity, the spacetime coordinate (\tilde{t}, \tilde{x}) of Y should satisfy

$$\tilde{t} = \gamma(t - vx_1), \quad \tilde{x}_1 = \gamma(x_1 - vt), \quad \tilde{x}_j = x_j \quad (j = 2, \dots, n).$$

The sound wave \tilde{u} that Y experienced should be $\tilde{u}(\tilde{x}, \tilde{t}) := u(x, t)$. The wave equation is preserved under the Lorentz transformation, namely,

$$(\partial_{\tilde{t}}^2 - \Delta_{\tilde{x}})\tilde{u}(\tilde{x}, \tilde{t}) = (\partial_t^2 - \Delta_x)u(x, t).$$

This means that what Y heard is also a wave which satisfies the same physical law with the sound that X heard. Theorem 1.1 tells us the profile of the sound (the wave shape across the space at a fixed time) that Y heard has the same H^1 spacial regularity with the sound that X heard.

This article is organized as follows. In Section 2 we make some preparations which are necessary for the subsequent analysis. Section 3 is devoted to the proof of an intermediate result in which the solution is assumed to have C^2 -smoothness. Then the C^2 -smoothness constraint is lifted in Section 4 by dealing with a compatibility issue and a regularity issue consecutively. We conclude this work in Section 5 with some remarks about possible future directions.

2. SOME PREPARATIONS

Throughout the article we denote by C a generic constant whose value may varies from line to line. We use $\operatorname{div}_{x,t}(\vec{a}, b)$ to signify $\nabla \cdot \vec{a} + \partial_t b$. The following lemma shall be used in Section 3 where the constraint $u \in C^2$ is stipulated.

Lemma 2.1. *Assume $u \in C^2$, and $\vec{\varphi}(x) = (\varphi_1(x), \dots, \varphi_n(x)) \in C^1(\mathbb{R}^n; \mathbb{R}^n)$ piecewisely, then at every C^1 continuous point of $\vec{\varphi}$, we have the following identities:*

$$\begin{aligned} 2 \operatorname{Re}\{\bar{u}_t[\partial_t^2 u - \nabla \cdot (A(x)\nabla u)]\} \\ = \operatorname{Re} \operatorname{div}_{x,t} [-2\bar{u}_t A(x)\nabla u, |u_t|^2 + |\nabla u|_A^2], \end{aligned} \quad (2.1)$$

$$\begin{aligned} 2 \operatorname{Re}\{(\vec{\varphi} \cdot \nabla \bar{u})[\partial_t^2 u - \nabla \cdot (A(x)\nabla u)]\} \\ = \operatorname{Re} \operatorname{div}_{x,t} [\vec{\varphi}(|\nabla u|_A^2 - |u_t|^2) - 2(\vec{\varphi} \cdot \nabla \bar{u})A\nabla u, 2(\vec{\varphi} \cdot \nabla \bar{u})u_t] \\ + (\nabla \cdot \vec{\varphi})(|u_t|^2 - |\nabla u|_A^2) - (\vec{\varphi} \cdot \nabla A)(\nabla u, \nabla u) + 2(\partial_j \varphi_k) \operatorname{Re}(\bar{u}_k a_{jl} u_l), \end{aligned} \quad (2.2)$$

where Re stands for the real part, and the summation convention is called for the last term.

Proof. A straightforward computation shows

$$2 \operatorname{Re}\{\bar{u}_t \partial_t^2 u\} = (|u_t|^2)', \quad 2 \operatorname{Re}\{\bar{u}_t \nabla \cdot (A\nabla u)\} = 2 \operatorname{Re} \nabla \cdot (\bar{u}_t A\nabla u) - (|\nabla u|_A^2)',$$

which give (2.1).

Similarly, we can show

$$\begin{aligned} 2 \operatorname{Re}\{(\vec{\varphi} \cdot \nabla \bar{u}) \partial_t^2 u\} &= \operatorname{Re} \partial_t [2(\vec{\varphi} \cdot \nabla \bar{u})u_t] - \vec{\varphi} \cdot 2 \operatorname{Re}(u_t \nabla \bar{u}_t) \\ &= \operatorname{Re} \partial_t [2(\vec{\varphi} \cdot \nabla \bar{u})u_t] - \vec{\varphi} \cdot \nabla (|u_t|^2) \\ &= \operatorname{Re} \partial_t [2(\vec{\varphi} \cdot \nabla \bar{u})u_t] - \nabla \cdot (\vec{\varphi}|u_t|^2) + (\nabla \cdot \vec{\varphi})|u_t|^2. \end{aligned} \quad (2.3)$$

$\vec{\varphi}$ is a real-valued vector function, so we also have

$$\begin{aligned} 2 \operatorname{Re}\{(\vec{\varphi} \cdot \nabla \bar{u}) \nabla \cdot (A\nabla u)\} &= \operatorname{Re} \nabla \cdot [2(\vec{\varphi} \cdot \nabla \bar{u})A\nabla u] - 2 \operatorname{Re}[\nabla(\vec{\varphi} \cdot \nabla \bar{u}) \cdot A\nabla u] \\ &= \operatorname{Re} \nabla \cdot [2(\vec{\varphi} \cdot \nabla \bar{u})A\nabla u] - 2 \operatorname{Re}[(\partial_j \varphi_k) \bar{u}_k a_{jl} u_l] - 2 \operatorname{Re}[\varphi_k \bar{u}_{jk} a_{jl} u_l]. \end{aligned} \quad (2.4)$$

We compute $\varphi_k \bar{u}_{jk} a_{jl} u_l$ as follows,

$$\begin{aligned} \varphi_k \bar{u}_{jk} a_{jl} u_l &= \varphi_k \partial_k (\bar{u}_j a_{jl} u_l) - \varphi_k \bar{u}_j (\partial_k a_{jl}) u_l - \varphi_k \bar{u}_j a_{jl} u_{lk} \\ &= \vec{\varphi} \cdot \nabla (|\nabla u|_A^2) - (\vec{\varphi} \cdot \nabla A)(\nabla u, \nabla u) - \varphi_k u_{jk} a_{lj} \bar{u}_l \\ &= \nabla \cdot (\vec{\varphi} |\nabla u|_A^2) - (\nabla \cdot \vec{\varphi}) |\nabla u|_A^2 - (\vec{\varphi} \cdot \nabla A)(\nabla u, \nabla u) - \varphi_k u_{jk} a_{lj} \bar{u}_l. \end{aligned} \quad (2.5)$$

Because A is symmetric and $A, \vec{\varphi}$ are real-valued, we can conclude from (2.5) that

$$2 \operatorname{Re}[\varphi_k \bar{u}_{jk} a_{jl} u_l] = \nabla \cdot (\vec{\varphi} |\nabla u|_A^2) - (\nabla \cdot \vec{\varphi}) |\nabla u|_A^2 - (\vec{\varphi} \cdot \nabla A)(\nabla u, \nabla u). \quad (2.6)$$

Combining (2.4) and (2.6), we arrive at

$$2 \operatorname{Re}\{(\bar{\varphi} \cdot \nabla \bar{u}) \nabla \cdot (A \nabla u)\} = \operatorname{Re} \nabla \cdot [2(\bar{\varphi} \cdot \nabla \bar{u}) A \nabla u] - 2(\partial_j \varphi_k) \operatorname{Re}(\bar{u}_k a_j u_l) \\ - \nabla \cdot (\bar{\varphi} |\nabla u|_A^2) + (\nabla \cdot \bar{\varphi}) |\nabla u|_A^2 + (\bar{\varphi} \cdot \nabla A)(\nabla u, \nabla u). \quad (2.7)$$

Subtracting (2.7) from (2.3), we arrive at (2.2). The proof is done. \square

In the following lemma we abbreviate $\mathcal{E}(u; \Gamma_S)$ as $\mathcal{E}(u)$ for simplicity, and we show the energy possesses a similar triangle inequality property.

Lemma 2.2. *For any u_1, u_2 such that $\mathcal{E}(u_1), \mathcal{E}(u_2)$ are well-defined, we have $\mathcal{E}(u_1 - u_2)$ is also well-defined and*

$$|\mathcal{E}(u_1) - \mathcal{E}(u_2)| \leq \mathcal{E}(u_1 - u_2) + 2\sqrt{\mathcal{E}(u_1 - u_2)}\sqrt{\mathcal{E}(u_j)}, \quad j = 1, 2.$$

Proof. First, similar to $|a + b| \leq |a| + |b|$, we also have $|a + b|_A \leq |a|_A + |b|_A$. This is because

$$|a + b|_A^2 = |a|_A^2 + |b|_A^2 + 2 \operatorname{Re}\{\bar{a} \cdot Ab\} = |a|_A^2 + |b|_A^2 + 2 \operatorname{Re}\{\overline{A^{1/2}a} \cdot A^{1/2}b\} \\ \leq |a|_A^2 + |b|_A^2 + 2|a|_A|b|_A = (|a|_A + |b|_A)^2.$$

By this triangle inequality, it is straightforward to check $\mathcal{E}(u_1 - u_2) \leq 2[\mathcal{E}(u_1) + \mathcal{E}(u_2)]$, so $\mathcal{E}(u_1 - u_2)$ is well-defined.

The triangle inequality obtained above also gives $|a|_A - |b|_A \leq |a - b|_A$. Hence

$$|\nabla(u_1)|_A^2 - |\nabla(u_2)|_A^2 = [|\nabla(u_1)|_A - |\nabla(u_2)|_A]^2 + 2[|\nabla(u_1)|_A - |\nabla(u_2)|_A]|\nabla(u_2)|_A \\ \leq |\nabla(u_1 - u_2)|_A^2 + 2|\nabla(u_1 - u_2)|_A|\nabla(u_2)|_A.$$

Similarly,

$$|\partial_t u_1|^2 - |\partial_t u_2|^2 \leq |\partial_t(u_1 - u_2)|^2 + 2|\partial_t(u_1 - u_2)||\partial_t u_2|.$$

Hence,

$$\mathcal{E}(u_1) - \mathcal{E}(u_2) \\ \leq \mathcal{E}(u_1 - u_2) + 2 \int [|\nabla(u_1 - u_2)|_A |\nabla(u_2)|_A + (1 - |\nabla S(x)|_A^2) |\partial_t(u_1 - u_2)| |\partial_t u_2|] \, dx \\ \leq \mathcal{E}(u_1 - u_2) + 2 \int \sqrt{|\nabla(u_1 - u_2)|_A^2 + (1 - |\nabla S(x)|_A^2) |\partial_t(u_1 - u_2)|^2} \\ \times \sqrt{|\nabla(u_2)|_A^2 + (1 - |\nabla S(x)|_A^2) |\partial_t u_2|^2} \, dx \\ \leq \mathcal{E}(u_1 - u_2) + 2\sqrt{\mathcal{E}(u_1 - u_2)}\sqrt{\mathcal{E}(u_2)}.$$

Similar arguments also imply

$$\mathcal{E}(u_1) - \mathcal{E}(u_2) \leq \mathcal{E}(u_1 - u_2) + 2\sqrt{\mathcal{E}(u_1 - u_2)}\sqrt{\mathcal{E}(u_1)}.$$

Therefore,

$$|\mathcal{E}(u_1) - \mathcal{E}(u_2)| \leq \mathcal{E}(u_1 - u_2) + 2\sqrt{\mathcal{E}(u_1 - u_2)}\sqrt{\mathcal{E}(u_j)}, \quad j = 1, 2.$$

The proof is done. \square

2.1. The decomposition of the gradient. This part is devoted to the analysis of the relation between $|\nu_\Sigma|_A^2 |\nabla u|_A^2$ and $|u_{\nu,A}|^2$ which will appear in (3.9). Here ν_Σ stands for the outer unit normal vector to Σ , and $u_{\nu,A}$ signifies the conormal derivative with respect to A , i.e. $u_{\nu,A} := \nu_\Sigma \cdot A \nabla u$. Readers can skip this part for the first time.

Recall that $\Sigma = \partial\Omega \times (0, T)$. When Ω is the unit ball and $A(x)$ is the identity matrix, by straightforward computations it can be checked that on Σ we have the following identity:

$$|\nu_\Sigma|_A^2 |\nabla u|_A^2 = |u_{\nu,A}|^2 + \frac{1}{2} \sum_{i \neq j} |X_{ij} u|^2, \quad \text{where } X_{ij} = x_i \partial_j - x_j \partial_i, \quad |x| = 1, \quad (2.8)$$

see e.g. [RS20b, (1.19)]. Note that here X_{ij} are vector fields tangential to the sphere of the unit ball. Therefore, the square norm of the gradient $|\nabla u|^2$ is decomposed into the desired term $|u_{\nu,A}|^2$ along with other terms which are tangential gradients on Σ . For general domain Ω and general matrix $A(x)$, we can show a similar decomposition result.

Lemma 2.3. *Assume Ω is a C^1 -smooth domain, and the matrix $A(x)$ satisfies Assumption 1.2 with constants c_1 and c_2 . Then for a C^1 -smooth function u , there exist two constants C depending only on c_1, c_2 and the dimension n such that on Σ we have*

$$|\nu_\Sigma|_A^2 |\nabla u|_A^2 \leq 1 \times |u_{\nu,A}|^2 + C |u_{\nu,A}| |\nabla_\Sigma u| + C |\nabla_\Sigma u|^2, \quad (2.9)$$

where $\nabla_\Sigma u$ represents the tangential gradient of u on Σ .

Remark 2.1. In Lemma 2.3 we emphasize the coefficient of the term $|u_{\nu,A}|^2$ is exactly 1.

Proof of Lemma 2.3. Denote $e_1 = \nu_\Sigma$ and fix an orthonormal basis $\{e_2, \dots, e_{n+1}\}$ of a local chart of Σ . Then $\{e_1, e_2, \dots, e_{n+1}\}$ is also an locally orthonormal basis of \mathbb{R}^{n+1} . The conclusion (2.9) is a local estimate so local arguments are enough for the proof.

For simplicity we denote

$$E(x) = (Ae_1, e_2, \dots, e_{n+1}).$$

Assumption 1.2 guarantees that $E(x)$ is always invertible; this is because $e_1 \cdot Ae_1 \geq c_1 |e_1|^2 = c_1 > 0$, so Ae_1 always has nonzero component in e_1 direction, and so $\{Ae_1, e_2, \dots, e_{n+1}\}$ is always linearly independent. Hence, thanks to the existence of $E^{-1}(x)$, we can compute

$$\begin{aligned} |\nabla u|_A^2 &= (\nabla \bar{u})^T A \nabla u = (\nabla \bar{u})^T E(x) [A^{-1/2} E(x)]^{-1} [E^T(x) A^{-1/2}]^{-1} E^T(x) \nabla u \\ &= (\nabla \bar{u})^T E(x) [E^{-1}(x) A E^{-1,T}(x)] E^T(x) \nabla u \\ &= (\bar{u}_{\nu,A}, e_2 \cdot \nabla \bar{u}, \dots, e_{n+1} \cdot \nabla \bar{u}) [E^{-1}(x) A E^{-1,T}(x)] (u_{\nu,A}, e_2 \cdot \nabla u, \dots, e_{n+1} \cdot \nabla u)^T. \end{aligned}$$

Here $E^{-1,T}(x)$ signifies the transpose of the inverse of the matrix $E(x)$. Let us denote

$$M(x) = E^{-1}(x) A(x) E^{-1,T}(x)$$

and use M_{ij} to signify the elements of M . Note that M is symmetric, then we have

$$|\nabla u|_A^2 = M_{11} |u_{\nu,A}|^2 + 2 \operatorname{Re} \{ \bar{u}_{\nu,A} \sum_{j=2}^n M_{1j} (e_j \cdot \nabla u) \} + \sum_{k,l=2}^n M_{kl} (e_k \cdot \nabla \bar{u}) (e_l \cdot \nabla u). \quad (2.10)$$

We claim that $e_l \cdot \nabla u$ is a component of the tangential gradient of u on Σ , which is similar to the $X_{ij} u$ term in (2.8). To see that, we choose a C^1 -smooth curve $\gamma: (-1, 1) \rightarrow \Sigma$ on Σ satisfying $\gamma(0) = x$ and $\dot{\gamma}(0) = e_l$, then $e_l \cdot \nabla u(x)$ can be represented as $\frac{d}{dt} |_{t=0} (u(\gamma(t)))$. This justifies our claim. Hence, we have

$$|e_l \cdot \nabla u| \leq |\nabla_\Sigma u|, \quad l = 2, \dots, n+1,$$

and so we can continue (2.10) as

$$|\nu_\Sigma|_A^2 |\nabla u|_A^2 = |\nu_\Sigma|_A^2 (M_{11} |u_{\nu,A}|^2 + C |u_{\nu,A}| |\nabla_\Sigma u| + C |\nabla_\Sigma u|^2)$$

$$\leq (|\nu_\Sigma|_A^2 M_{11})|u_{\nu,A}|^2 + C|u_{\nu,A}||\nabla_\Sigma u| + C|\nabla_\Sigma u|^2, \quad (2.11)$$

for some constant C . From the definition of the matrix M , it can be checked that C depends only on c_1 , c_2 and n .

It is left to show

$$|\nu_\Sigma|_A^2 M_{11} = 1. \quad (2.12)$$

To see this, let us represent the inverse matrix $E^{-1}(x)$ as

$$E^{-1}(x) = (f_1(x), f_2(x), \dots, f_{n+1}(x))^T,$$

then the matrix identity $E^{-1}(x)E(x) = I$ gives

$$f_1(x) \cdot e_j(x) = 0, \quad j = 2, \dots, n+1. \quad (2.13)$$

$$f_1(x) \cdot A(x)e_1(x) = 1. \quad (2.14)$$

Because $\{e_1, e_2, \dots, e_{n+1}\}$ is an orthonormal basis in a chart, from (2.13) we see $f_1(x)$ is parallel to $e_1(x)$, i.e. $f_1(x) = \lambda(x)e_1(x)$ for some function $\lambda(x)$. Substitute this into (2.14), we see $\lambda(x) = |e_1(x)|_A^{-2} = |\nu_\Sigma|_A^{-2}$. Therefore, we have

$$M_{11}|\nu_\Sigma|_A^2 = f_1^T(x)A(x)f_1(x)|\nu_\Sigma|_A^2 = \lambda(x)e_1^T(x)A(x)\lambda(x)e_1(x)|\nu_\Sigma|_A^2 = \lambda^2(x)|\nu_\Sigma|_A^4 = 1,$$

which is (2.12). Combining (2.11) and (2.12), we can complete the proof. \square

3. THE C^2 SMOOTH CASE

Recall $\langle T_2 \rangle := (1 + |T_2|^2)^{1/2}$. In this section we aim to prove the following result.

Proposition 3.1. *Given Assumptions 1.1 and 1.2. Assume $u \in C^2(\overline{Q})$ solves (1.1)-(1.2). Then we have*

$$\begin{aligned} |\mathcal{E}(u; \Gamma_S) - \int_\Omega (|\nabla u_0|_A^2 + |u_1|^2) dx| \\ \leq C \langle T_2 \rangle^{1/2} [\|\nabla u_0\|_{L^2(\Omega)} + \|u_1\|_{L^2(\Omega)} + \langle T_2 \rangle^{1/2} (\|f\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)})] \\ \times (\|f\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)}), \end{aligned}$$

for some constant C depending only on A and the dimension n . Especially, when $f = 0$ and $G = 0$ in (1.2), we have

$$\mathcal{E}(u; \Gamma_S) = \|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2. \quad (3.1)$$

This section and a major portion of Section 4 involve the same hypersurface Γ_S , so we abbreviate $\mathcal{E}(u; \Gamma_S)$ as $\mathcal{E}(u)$ for short in these parts. Also, for technical reasons we introduce the following functional $\mathcal{E}_\tau(u)$ which we shall call it *partial energy* and which takes a real number τ as its parameter,

$$\mathcal{E}_\tau(u) := \int_{\pi(\Gamma_{S,\tau})} [|\nabla(u(x, S(x)))|_A^2 + (1 - |\nabla S(x)|_A^2)|u_t(x, S(x))|^2] dx, \quad (3.2)$$

where $\pi: (x, t) \mapsto x$ is a projection map. Readers can compare (3.2) with (1.4), and shall distinguish the notation $\mathcal{E}_\tau(u)$ with $\mathcal{E}(u; \Gamma_\tau)$ defined on different hypersurfaces Γ_τ . When $\tau \leq T_1$, we have $\pi(\Gamma_{S,\tau}) = \Omega$, so $\mathcal{E}_\tau(u)$ will be a constant with respect to τ when $\tau \leq T_1$. Under Assumption 1.1, we always have $\mathcal{E}_\tau(u) \leq \mathcal{E}(u)$; and when $\tau \leq T_1$ we have $\mathcal{E}_\tau(u) = \mathcal{E}(u)$.

The relationship between $\|u\|_{H^1(\Gamma_S)}$ and $\mathcal{E}(u)$ are given below,

$$\|u\|_{H^1(\Gamma_S)}^2 = \int_\Omega |\nabla(u(x, S(x)))|^2 \frac{dx}{(1 + |\nabla S(x)|^2)^{1/2}} \leq c_1^{-1} \int_\Omega |\nabla(u(x, S(x)))|_A^2 dx$$

$$\begin{aligned}
 &= c_1^{-1} \int_{\Omega} (|\nabla u|_A^2 + |\nabla S|_A^2 |u_t|^2 + 2 \operatorname{Re}\{\nabla S \cdot \bar{u}_t A \nabla u\}) \, dx \\
 &\leq c_1^{-1} \int_{\Omega} (|\nabla u|_A^2 + c_2 |u_t|^2 + 2 \operatorname{Re}\{\nabla S \cdot \bar{u}_t A \nabla u\}) \, dx \\
 &\leq c_1^{-1} \max\{1, c_2\} \mathcal{E}(u),
 \end{aligned} \tag{3.3}$$

where we used Assumptions 1.1 and 1.2. For readers convenience we also record the following identity,

$$|\nabla(u)|_A^2 + (1 - |\nabla S|_A^2) |u_t|^2 = |\nabla u|_A^2 + |u_t|^2 + 2 \operatorname{Re}\{\bar{u}_t \nabla S \cdot A \nabla u\}.$$

The arguments of proving Proposition 3.1 are divided into several steps.

Lemma 3.2. *Under the same condition as in Proposition 3.1, we have*

$$\mathcal{E}_{\tau}(u) \leq 2 \int_{H_{\tau}} (|\nabla u(x, \tau)|_A^2 + |u_t(x, \tau)|^2) \, dx + 6 \|f_t\|_{L^2(\Sigma_{\tau})} \|u_{\nu, A}\|_{L^2(\Sigma_{\tau})} + 8(T_2 - \tau) \|G\|_{L^2(Q_{\tau})}^2.$$

Proof. Equation (1.1) gives $\partial_t^2 u - \nabla \cdot (A(x) \nabla u) = G$ in Q_{τ} . Hence, integrating the identity (2.1) in Q_{τ} , we can have

$$\begin{aligned}
 &\operatorname{Re} \int_{Q_{\tau}} 2\bar{u}_t G = \operatorname{Re} \int_{\partial Q_{\tau}} \nu_{Q_{\tau}} \cdot (-2\bar{u}_t A \nabla u, |u_t|^2 + |\nabla u|_A^2) \\
 &= \operatorname{Re} \int_{H_{\tau}} (\vec{0}, -1) \cdot (-2\bar{u}_t A \nabla u, |u_t|^2 + |\nabla u|_A^2) + \operatorname{Re} \int_{\Sigma_{\tau}} \nu_{\Sigma} \cdot (-2\bar{u}_t A \nabla u, |u_t|^2 + |\nabla u|_A^2) \\
 &\quad + \operatorname{Re} \int_{\Gamma_{S, \tau}} \nu \cdot (-2\bar{u}_t A \nabla u, |u_t|^2 + |\nabla u|_A^2) \quad (\nu \text{ is defined in (1.3)}) \\
 &= - \int_{H_{\tau}} (|u_t|^2 + |\nabla u|_A^2) \, d\sigma - 2 \operatorname{Re} \int_{\Sigma_{\tau}} \bar{u}_t u_{\nu, A} \, d\sigma \quad (u_{\nu, A} := \nu_{\Sigma} \cdot A \nabla u) \\
 &\quad + \operatorname{Re} \int_{\Gamma_{S, \tau}} (2\nabla S \cdot \bar{u}_t A \nabla u + |u_t|^2 + |\nabla u|_A^2) \frac{d\sigma}{\sqrt{1 + |\nabla S|^2}} \\
 &= - \int_{H_{\tau}} (|u_t|^2 + |\nabla u|_A^2) \, d\sigma - 2 \operatorname{Re} \int_{\Sigma_{\tau}} \bar{u}_t u_{\nu, A} \, d\sigma \\
 &\quad + \int_{\pi(\Gamma_{S, \tau})} (2 \operatorname{Re}\{\nabla S \cdot \bar{u}_t A \nabla u\} + |u_t|^2 + |\nabla u|_A^2) \, dx \\
 &= - \int_{H_{\tau}} (|u_t|^2 + |\nabla u|_A^2) \, d\sigma - 2 \operatorname{Re} \int_{\Sigma_{\tau}} \bar{u}_t u_{\nu, A} \, d\sigma + \mathcal{E}_{\tau}(u),
 \end{aligned} \tag{3.4}$$

where $\nu_{Q_{\tau}}$ signifies the outer unit normal of Q_{τ} , ν_{Σ} is the outer unit normal vector to Σ , and $u_{\nu, A}$ signifies the conormal derivative with respect to A , i.e. $u_{\nu, A} := \nu_{\Sigma} \cdot A \nabla u$.

We denote

$$e(\tau) := \int_{H_{\tau}} (|\nabla u|_A^2 + |u_t|^2) \, dx. \tag{3.5}$$

Substituting (3.5) into (3.4), we have

$$\begin{aligned}
 e(\tau) &= - \operatorname{Re} \int_{Q_{\tau}} 2\bar{u}_t G + \mathcal{E}_{\tau}(u) - 2 \operatorname{Re} \int_{\Sigma_{\tau}} \bar{u}_t u_{\nu, A} \, d\sigma \\
 &\leq \int_{Q_{\tau}} \frac{1}{K} |u_t|^2 + \int_{Q_{\tau}} K |G|^2 + \mathcal{E}_{\tau}(u) - 2 \operatorname{Re} \int_{\Sigma_{\tau}} \bar{u}_t u_{\nu, A} \, d\sigma \\
 &\leq \frac{1}{K} \int_{\tau}^{T_2} e(s) \, ds + K \|G\|_{L^2(Q_{\tau})}^2 + \mathcal{E}_{\tau}(u) + 2 \int_{\Sigma_{\tau}} |u_t u_{\nu, A}| \, d\sigma,
 \end{aligned}$$

where K can be any positive number. Recall that under Assumption 1.1 we have $|\nabla S|_A \leq 1$, so $\mathcal{E}_\tau(u)$ is always non-increasing with respect to τ . Hence, by Grönwall's inequality we can obtain

$$e(\tau) \leq e^{(T_2-\tau)/K} (K \|G\|_{L^2(Q_\tau)}^2 + \mathcal{E}_\tau(u) + 2 \int_{\Sigma_\tau} |u_t u_{\nu,A}| \, d\sigma), \quad \forall \tau \in [0, T_2]. \quad (3.6)$$

On the other hand, when $\tau \leq T_1$, from (3.4) we also obtain

$$\begin{aligned} \mathcal{E}_\tau(u) &= e(\tau) + 2 \operatorname{Re} \int_{\Sigma_\tau} \bar{u}_t u_{\nu,A} \, d\sigma + \operatorname{Re} \int_{Q_\tau} 2\bar{u}_t G \\ &\leq e(\tau) + 2 \int_{\Sigma_\tau} |u_t u_{\nu,A}| \, d\sigma + \frac{1}{\epsilon} \int_{Q_\tau} |G|^2 + \epsilon \int_\tau^{T_2} e(s) \, ds \\ &\leq e(\tau) + 2 \int_{\Sigma_\tau} |u_t u_{\nu,A}| \, d\sigma + \frac{1}{\epsilon} \|G\|_{L^2(Q_\tau)}^2 \\ &\quad + \epsilon \int_\tau^{T_2} e^{(T_2-s)/K} [K \|G\|_{L^2(Q_s)}^2 + \mathcal{E}_s(u) + 2 \int_{\Sigma_s} |u_t u_{\nu,A}| \, d\sigma] \, ds \quad (\text{by (3.6)}) \\ &\leq e(\tau) + 2 \int_{\Sigma_\tau} |u_t u_{\nu,A}| \, d\sigma + \frac{1}{\epsilon} \|G\|_{L^2(Q_\tau)}^2 \\ &\quad + \epsilon K (e^{(T_2-\tau)/K} - 1) [K \|G\|_{L^2(Q_\tau)}^2 + \mathcal{E}_\tau(u) + 2 \int_{\Sigma_\tau} |u_t u_{\nu,A}| \, d\sigma] \\ &\leq e(\tau) + 2[1 + \epsilon K (e^{(T_2-\tau)/K} - 1)] \int_{\Sigma_\tau} |u_t u_{\nu,A}| \, d\sigma \\ &\quad + \left[\frac{1}{\epsilon} + \epsilon K^2 (e^{(T_2-\tau)/K} - 1) \right] \|G\|_{L^2(Q_\tau)}^2 + \epsilon K (e^{(T_2-\tau)/K} - 1) \mathcal{E}_\tau(u). \end{aligned}$$

By setting $\epsilon = [2K(e^{(T_2-\tau)/K} - 1)]^{-1}$ and absorbing $\mathcal{E}_\tau(u)$ on the right-hand side (RHS) by the left-hand side (LHS), we obtain

$$\begin{aligned} \frac{1}{2} \mathcal{E}_\tau(u) &\leq e(\tau) + 3 \int_{\Sigma_\tau} |u_t u_{\nu,A}| \, d\sigma + [2K(e^{(T_2-\tau)/K} - 1) + \frac{K}{2}] \|G\|_{L^2(Q_\tau)}^2 \\ &\leq e(\tau) + 3 \int_{\Sigma_\tau} |u_t u_{\nu,A}| \, d\sigma + (T_2 - \tau) \left[2 \frac{K e^{(T_2-\tau)/K}}{T_2 - \tau} - \frac{3}{2} \frac{K}{T_2 - \tau} \right] \|G\|_{L^2(Q_\tau)}^2. \quad (3.7) \end{aligned}$$

When we treat the coefficient in front of $\|G\|_{L^2(Q_\tau)}^2$ as a function of K , then elementary calculus shows the minimum value of the coefficient is in between 3.5 and 4, i.e.,

$$3.5 < \min_{K>0} \left\{ 2 \frac{K e^{(T_2-\tau)/K}}{T_2 - \tau} - \frac{3}{2} \frac{K}{T_2 - \tau} \right\} < 4, \quad \text{provided } \tau < T_2.$$

Hence, by choosing the value of K according to T_2 and τ properly, (3.7) can be improved to

$$\mathcal{E}_\tau(u) \leq 2e(\tau) + 6 \|f_t\|_{L^2(\Sigma_\tau)} \|u_{\nu,A}\|_{L^2(\Sigma_\tau)} + 8(T_2 - \tau) \|G\|_{L^2(Q_\tau)}^2,$$

which is the conclusion. \square

Remark 3.1. The piecewise smoothness of S does not cause any trouble to the proof of Lemma 3.2. This is because the integration operation does not require the integrand to be smooth everywhere.

The RHS of the inequality in Lemma 3.2 involves a norm of $u_{\nu,A}$, and to achieve an *a-priori* estimate of u on Γ_S , we also need to estimate the term $\|u_{\nu,A}\|_{L^2(\Sigma_0)}$. This can be done by playing with (2.2). The multiplier “ $\vec{\varphi} \cdot \nabla u$ ” in (2.2) is a generalization of “ $x \cdot \nabla u$ ” which has been seen in the literature, e.g. [Mor61, (12)], [Ike05, Lemma 2.1], [RS20b, Lemma 3.3].

Lemma 3.3. *Under the same condition as in Proposition 3.1, for vector-valued function $\vec{\varphi} \in C^1(\bar{\Omega}, \mathbb{R}^n)$ satisfying $\vec{\varphi}(x) = A(x)\nu_\Sigma(x)$ on $\partial\Omega$, we have*

$$\begin{aligned} \|u_{\nu,A}\|_{L^2(\Sigma_\tau)}^2 &\leq C\|f\|_{H^1(\Sigma_\tau)}^2 + C(\sup_\Omega |\vec{\varphi}|) [\|\nabla u(\cdot, \tau)\|_{L^2(\Omega)}^2 + \|u_t(\cdot, \tau)\|_{L^2(\Omega)}^2] \\ &\quad + C[T_2 - \tau + (\sup_\Omega |\vec{\varphi}|)] [(T_2 - \tau)\|G\|_{L^2(Q_\tau)}^2 + \mathcal{E}_\tau(u)] \\ &\quad + C[T_2 - \tau + (\sup_\Omega |\vec{\varphi}|)]^2 \|f_t\|_{L^2(\Sigma_\tau)}^2, \end{aligned}$$

for some constant C depending only on $\vec{\varphi}$, A and n .

Proof. Integrating (2.2) in Q_τ , we have

$$\begin{aligned} 2 \operatorname{Re} \int_{Q_\tau} (\vec{\varphi} \cdot \nabla \bar{u}) G &= 2 \operatorname{Re} \int_{Q_\tau} (\vec{\varphi} \cdot \nabla \bar{u}) [\partial_t^2 u - \nabla \cdot (A(x) \nabla u)] \\ &= \operatorname{Re} \int_{Q_\tau} \operatorname{div}_{x,t} [\vec{\varphi} (|\nabla u|_A^2 - |u_t|^2) - 2(\vec{\varphi} \cdot \nabla \bar{u}) A \nabla u, 2(\vec{\varphi} \cdot \nabla \bar{u}) u_t] \\ &\quad + \int_{Q_\tau} [(\nabla \cdot \vec{\varphi})(|u_t|^2 - |\nabla u|_A^2) - (\vec{\varphi} \cdot \nabla A)(\nabla u, \nabla u) + 2(\partial_j \varphi_k) \operatorname{Re}(\bar{u}_k a_{jl} u_l)] \\ &= \operatorname{Re} \int_{\Sigma_\tau} [(\nu_\Sigma \cdot \vec{\varphi})(|\nabla u|_A^2 - |u_t|^2) - 2(\vec{\varphi} \cdot \nabla \bar{u}) u_{\nu,A}] \\ &\quad + \operatorname{Re} \int_{\Gamma_{S,\tau}} [(\nu_x \cdot \vec{\varphi})(|\nabla u|_A^2 - |u_t|^2) + 2(\vec{\varphi} \cdot \nabla \bar{u})(\nu_t u_t - \nu_x \cdot A \nabla u)] \\ &\quad - \operatorname{Re} \int_{\Omega \times \{\tau\}} 2(\vec{\varphi} \cdot \nabla \bar{u}) u_t \\ &\quad + \int_{Q_\tau} [(\nabla \cdot \vec{\varphi})(|u_t|^2 - |\nabla u|_A^2) - (\vec{\varphi} \cdot \nabla A)(\nabla u, \nabla u) + 2(\partial_j \varphi_k) \operatorname{Re}(\bar{u}_k a_{jl} u_l)] \\ &=: I_1 + I_2 + I_3 - \operatorname{Re} \int_{\Omega \times \{\tau\}} 2(\vec{\varphi} \cdot \nabla \bar{u}) u_t, \end{aligned} \tag{3.8}$$

where I_1 , I_2 and I_3 represent the integrals on Σ_τ , $\Gamma_{S,\tau}$ and Q_τ , respectively. We estimate I_1 , I_2 and I_3 separately.

Recall that $\vec{\varphi}(x) = A(x)\nu_\Sigma(x)$ on $\partial\Omega$, so I_1 can be simplified as

$$I_1 = \int_{\Sigma_\tau} [|\nu_\Sigma|_A^2 (|\nabla u|_A^2 - |u_t|^2) - 2|u_{\nu,A}|^2] \, d\sigma. \tag{3.9}$$

By Lemma 2.3 we can obtain

$$|\nu_\Sigma|_A^2 |\nabla u|_A^2 \leq \frac{3}{2} |u_{\nu,A}|^2 + C |\nabla_\Sigma u|^2 \text{ on } \Sigma, \tag{3.10}$$

for some constant C depending only on n and c_1, c_2 of Assumption 1.2, where $\nabla_\Sigma u$ represents the tangential gradient of u on Σ . Substituting (3.10) into (3.9), we can continue

$$\begin{aligned} I_1 &\leq \int_{\Sigma_\tau} \left[\left(\frac{3}{2} |u_{\nu,A}|^2 + C |\nabla_\Sigma u|^2 - |u_t|^2 \right) - 2|u_{\nu,A}|^2 \right] \, d\sigma \\ &\leq -\frac{1}{2} \|u_{\nu,A}\|_{L^2(\Sigma_\tau)}^2 + C \|u\|_{H^1(\Sigma_\tau)}^2, \end{aligned} \tag{3.11}$$

for some constant C depending on c_1, c_2, n .

The integral I_2 is on Γ_S , and on Γ_S we have

$$\nabla u(x, S(x)) = \nabla(u(x, S(x))) - \nabla S(x) u_t(x, S(x)).$$

For simplicity we abbreviate $\nabla(u(x, S(x)))$ as $\nabla(u)$. Then we compute

$$\begin{aligned}
 & (\nu_x \cdot \vec{\varphi})(|\nabla u|_A^2 - |u_t|^2) \\
 &= (\nu_x \cdot \vec{\varphi})[|\nabla(u) - \nabla S u_t|_A^2 - |u_t|^2] \\
 &= (\nu_x \cdot \vec{\varphi})[|\nabla(u)|_A^2 - 2\operatorname{Re}\{\nabla S \bar{u}_t \cdot A \nabla(u)\} - (1 - |\nabla S|_A^2)|u_t|^2], \tag{3.12}
 \end{aligned}$$

and

$$\begin{aligned}
 & 2(\vec{\varphi} \cdot \nabla \bar{u})(\nu_t u_t - \nu_x \cdot A \nabla u) \times (1 + |\nabla S|^2)^{1/2} \\
 &= 2[\vec{\varphi} \cdot \nabla(\bar{u}) - \vec{\varphi} \cdot \nabla S \bar{u}_t][\nu_t u_t - \nu_x \cdot A \nabla(u) + \nu_x \cdot A \nabla S u_t] \times (1 + |\nabla S|^2)^{1/2} \\
 &= 2[\vec{\varphi} \cdot \nabla(\bar{u}) - \vec{\varphi} \cdot \nabla S \bar{u}_t][u_t + \nabla S \cdot A \nabla(u) - |\nabla S|_A^2 u_t] \quad (\text{recall (1.3)}) \\
 &= 2\vec{\varphi} \cdot \nabla(\bar{u})(1 - |\nabla S|_A^2)u_t - 2(\vec{\varphi} \cdot \nabla S)(1 - |\nabla S|_A^2)|u_t|^2 \\
 &\quad + 2\vec{\varphi} \cdot \nabla(\bar{u})\nabla S \cdot A \nabla(u) - 2(\vec{\varphi} \cdot \nabla S)\bar{u}_t \nabla S \cdot A \nabla(u). \tag{3.13}
 \end{aligned}$$

Combining (3.12) with (3.13), we obtain

$$\begin{aligned}
 & \operatorname{Re}[(\nu_x \cdot \vec{\varphi})(|\nabla u|_A^2 - |u_t|^2) + 2(\vec{\varphi} \cdot \nabla \bar{u})(\nu_t u_t - \nu_x \cdot A \nabla u)] \times (1 + |\nabla S|^2)^{1/2} \\
 &= -(\vec{\varphi} \cdot \nabla S)[|\nabla(u)|_A^2 + (1 - |\nabla S|_A^2)|u_t|^2] \\
 &\quad + 2\operatorname{Re}\{[\vec{\varphi} \cdot \nabla(\bar{u})][\nabla S \cdot A \nabla(u) + (1 - |\nabla S|_A^2)u_t]\}. \tag{3.14}
 \end{aligned}$$

With the help of (3.14), we can estimate I_2 in the following way,

$$\begin{aligned}
 |I_2| &= \left| \operatorname{Re} \int_{\Gamma_{S,\tau}} [(\nu_x \cdot \vec{\varphi})(|\nabla u|_A^2 - |u_t|^2) + 2(\vec{\varphi} \cdot \nabla \bar{u})(\nu_t u_t - \nu_x \cdot A \nabla u)] d\sigma \right| \\
 &\leq \int_{\Gamma_{S,\tau}} |\vec{\varphi} \cdot \nabla S| [|\nabla(u)|_A^2 + (1 - |\nabla S|_A^2)|u_t|^2] \frac{d\sigma}{\sqrt{1 + |\nabla S|^2}} \\
 &\quad + 2 \int_{\Gamma_{S,\tau}} |\vec{\varphi} \cdot \nabla(\bar{u})| [|\nabla S \cdot A \nabla(u)| + (1 - |\nabla S|_A^2)|u_t|] \frac{d\sigma}{\sqrt{1 + |\nabla S|^2}} \\
 &\leq (c_1^{-1/2} + 2)(\sup_{\Omega} |\vec{\varphi}|) \int_{\Gamma_{S,\tau}} [|\nabla(u)|_A^2 + (1 - |\nabla S|_A^2)|u_t|^2] \frac{d\sigma}{\sqrt{1 + |\nabla S|^2}} \\
 &= (c_1^{-1/2} + 2)(\sup_{\Omega} |\vec{\varphi}|) \mathcal{E}_{\tau}(u). \tag{3.15}
 \end{aligned}$$

For I_3 , we have

$$\begin{aligned}
 |I_3| &\leq \int_{Q_{\tau}} |(\nabla \cdot \vec{\varphi})(|u_t|^2 - |\nabla u|_A^2) - (\vec{\varphi} \cdot \nabla A)(\nabla u, \nabla u) + 2(\partial_j \varphi_k) \operatorname{Re}(\bar{u}_k a_{j l} u_l)| \\
 &\leq C \int_{Q_{\tau}} (|\nabla u|_A^2 + |u_t|^2) dx dt = C \int_{\tau}^{T_2} e(s) ds,
 \end{aligned}$$

for some constant C depending only on $\vec{\varphi}$, A and n , and the $e(s)$ is defined in (3.5). Hence by (3.6) which requires Assumption 1.1, we can have

$$\begin{aligned}
 |I_3| &\leq C \int_{\tau}^{T_2} e^{(T_2-s)/K} [K \|G\|_{L^2(Q_s)}^2 + \mathcal{E}_s(u) + 2 \int_{\Sigma_s} |u_t u_{\nu, A}| d\sigma] ds \\
 &\leq C \int_{\tau}^{T_2} e^{(T_2-s)/K} ds \cdot [K \|G\|_{L^2(Q_{\tau})}^2 + \mathcal{E}_{\tau}(u) + 2 \int_{\Sigma_{\tau}} |u_t u_{\nu, A}| d\sigma] \\
 &\leq CK(e^{(T_2-\tau)/K} - 1) [K \|G\|_{L^2(Q_{\tau})}^2 + \mathcal{E}_{\tau}(u) + \frac{1}{\epsilon} \|u_t\|_{L^2(\Sigma_{\tau})}^2 + \epsilon \|u_{\nu, A}\|_{L^2(\Sigma_{\tau})}^2]. \tag{3.16}
 \end{aligned}$$

In (3.8), there is a term $\operatorname{Re} \int_{Q_\tau} (\vec{\varphi} \cdot \nabla \bar{u}) G$, and similar to the estimation of I_3 , we can also estimate this integral as follows,

$$\begin{aligned}
 |\operatorname{Re} \int_{Q_\tau} (\vec{\varphi} \cdot \nabla \bar{u}) G| &\leq \frac{1}{2} (\sup_\Omega |\vec{\varphi}|) \int_{Q_\tau} (c_1^{-1} K^{-1} |\nabla u|_A^2 + K |G|^2) \\
 &\leq \frac{1}{2c_1} (\sup_\Omega |\vec{\varphi}|) \frac{1}{K} \int_\tau^{T_2} e(s) \, ds + \frac{1}{2} (\sup_\Omega |\vec{\varphi}|) K \|G\|_{L^2(Q_\tau)}^2 \\
 &\leq \frac{1}{2c_1} (\sup_\Omega |\vec{\varphi}|) (e^{(T_2-\tau)/K} - 1) [K \|G\|_{L^2(Q_\tau)}^2 + \mathcal{E}_\tau(u) + \frac{1}{\epsilon} \|u_t\|_{L^2(\Sigma_\tau)}^2 + \epsilon \|u_{\nu,A}\|_{L^2(\Sigma_\tau)}^2] \\
 &\quad + \frac{1}{2} (\sup_\Omega |\vec{\varphi}|) K \|G\|_{L^2(Q_\tau)}^2 \\
 &\leq \frac{1}{2} (\sup_\Omega |\vec{\varphi}|) \cdot K (e^{(T_2-\tau)/K} - 1 + c_1) \cdot \|G\|_{L^2(Q_\tau)}^2 \\
 &\quad + \frac{1}{2c_1} (\sup_\Omega |\vec{\varphi}|) (e^{(T_2-\tau)/K} - 1) [\mathcal{E}_\tau(u) + \frac{1}{\epsilon} \|u_t\|_{L^2(\Sigma_\tau)}^2 + \epsilon \|u_{\nu,A}\|_{L^2(\Sigma_\tau)}^2]. \tag{3.17}
 \end{aligned}$$

Combining (3.16) and (3.17) and setting $K = T_2 - \tau$, we have

$$\begin{aligned}
 |I_3| + |\operatorname{Re} \int_{Q_\tau} (\vec{\varphi} \cdot \nabla \bar{u}) G| \\
 \leq C [T_2 - \tau + (\sup_\Omega |\vec{\varphi}|)] [(T_2 - \tau) \|G\|_{L^2(Q_\tau)}^2 + \mathcal{E}_\tau(u) + \frac{1}{\epsilon} \|u_t\|_{L^2(\Sigma_\tau)}^2 + \epsilon \|u_{\nu,A}\|_{L^2(\Sigma_\tau)}^2], \tag{3.18}
 \end{aligned}$$

for some constant C depending only on $\vec{\varphi}$, A and n .

Now, by combining (3.11), (3.15) and (3.18) with (3.8), we obtain

$$\begin{aligned}
 &\frac{1}{2} \|u_{\nu,A}\|_{L^2(\Sigma_\tau)}^2 \\
 &\leq C \|u\|_{H^1(\Sigma_\tau)}^2 + |I_2| + 2 \left| \int_{Q_\tau} (\vec{\varphi} \cdot \nabla \bar{u}) G \right| + |I_3| + \left| \int_{\Omega \times \{\tau\}} 2(\vec{\varphi} \cdot \nabla \bar{u}) u_t \right| \\
 &\leq C \|u\|_{H^1(\Sigma_\tau)}^2 + (c_1^{-1/2} + 2) (\sup_\Omega |\vec{\varphi}|) \mathcal{E}_\tau(u) + (\sup_\Omega |\vec{\varphi}|) [\|\nabla u(\cdot, \tau)\|_{L^2(\Omega)}^2 + \|u_t(\cdot, \tau)\|_{L^2(\Omega)}^2] \\
 &\quad + C [T_2 - \tau + (\sup_\Omega |\vec{\varphi}|)] [(T_2 - \tau) \|G\|_{L^2(Q_\tau)}^2 + \mathcal{E}_\tau(u) + \frac{1}{\epsilon} \|u_t\|_{L^2(\Sigma_\tau)}^2 + \epsilon \|u_{\nu,A}\|_{L^2(\Sigma_\tau)}^2].
 \end{aligned}$$

By setting $\epsilon = \{4C [T_2 - \tau + (\sup_\Omega |\vec{\varphi}|)]\}^{-1}$, we arrive at

$$\begin{aligned}
 \frac{1}{4} \|u_{\nu,A}\|_{L^2(\Sigma_\tau)}^2 &\leq C \|u\|_{H^1(\Sigma_\tau)}^2 + C (\sup_\Omega |\vec{\varphi}|) [\|\nabla u(\cdot, \tau)\|_{L^2(\Omega)}^2 + \|u_t(\cdot, \tau)\|_{L^2(\Omega)}^2] \\
 &\quad + C [T_2 - \tau + (\sup_\Omega |\vec{\varphi}|)] [(T_2 - \tau) \|G\|_{L^2(Q_\tau)}^2 + \mathcal{E}_\tau(u)] \\
 &\quad + C [T_2 - \tau + (\sup_\Omega |\vec{\varphi}|)]^2 \|u_t\|_{L^2(\Sigma_\tau)}^2.
 \end{aligned}$$

The proof is done. \square

Recall $\langle T_2 \rangle := (1 + |T_2|^2)^{1/2}$. With the help of Lemmas 3.2 and 3.3, we are able to bound $\mathcal{E}(u)$ and $\|u_{\nu,A}\|_{L^2(\Sigma_0)}$ by the initial/boundary data and the source term.

Lemma 3.4. *Under the same condition as in Proposition 3.1, we have*

$$\begin{aligned}
 \mathcal{E}(u) &\leq C (\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) + C \langle T_2 \rangle (\|G\|_{L^2(Q_0)}^2 + \|f\|_{H^1(\Sigma_0)}^2), \\
 \|u_{\nu,A}\|_{L^2(\Sigma_0)}^2 &\leq C \langle T_2 \rangle (\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) + C \langle T_2 \rangle^2 (\|f\|_{H^1(\Sigma_0)}^2 + \|G\|_{L^2(Q_0)}^2),
 \end{aligned}$$

for some constant C depending only on A and the dimension n .

Proof. Using the inequality in Lemma 3.2 with $\tau = 0$, and noting that $u(\cdot, 0) = u_0$ and $u_t(\cdot, 0) = u_1$, we have

$$\mathcal{E}(u) \leq 2 \int_{\Omega} (|\nabla u_0|_A^2 + |u_1|^2) dx + \frac{3}{\epsilon} \|f_t\|_{L^2(\Sigma_0)}^2 + \epsilon \|u_{\nu,A}\|_{L^2(\Sigma_0)}^2 + 8T_2 \|G\|_{L^2(Q_0)}^2. \quad (3.19)$$

Substituting the inequality in Lemma 3.3 with $\tau = 0$ into (3.19), and setting ϵ in (3.19) to be $\{2C[T_2 + (\sup_{\Omega} |\bar{\varphi}|)]\}^{-1}$, we obtain

$$\mathcal{E}(u) \leq C(\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) + C[T_2 + (\sup_{\Omega} |\bar{\varphi}|) + 1](\|f\|_{H^1(\Sigma_0)}^2 + \|G\|_{L^2(Q_0)}^2), \quad (3.20)$$

for some constant C depending only on $\bar{\varphi}$, A and n . Recall that in Lemma 3.3 we have fixed the value of $\bar{\varphi}$ on $\partial\Omega$. But we still have the freedom to choose the value of $\bar{\varphi}$ in the interior of Ω . We can choose $\bar{\varphi}$ in such a way that $\sup_{\Omega} |\bar{\varphi}| \leq 2\sup_{\partial\Omega} |\bar{\varphi}|$, and this choice guarantees

$$\sup_{\Omega} |\bar{\varphi}| \leq 2 \sup_{\partial\Omega} |A \cdot \nu_{\Sigma}| \leq 2c_2,$$

for the constant c_2 given in Assumption 1.2. Combining this with (3.20), we arrive at the first inequality of the lemma.

For the second inequality, we substitute the inequality in Lemma 3.2 into the inequality in Lemma 3.3 with $\tau = 0$, and this gives

$$\begin{aligned} \|u_{\nu,A}\|_{L^2(\Sigma_0)}^2 &\leq C[T_2 + (\sup_{\Omega} |\bar{\varphi}|)] [\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2] + C[T_2 + (\sup_{\Omega} |\bar{\varphi}|)]^2 \|f\|_{H^1(\Sigma_0)}^2 \\ &\quad + C[T_2 + (\sup_{\Omega} |\bar{\varphi}|)] T_2 \|G\|_{L^2(Q_0)}^2 \\ &\quad + C[T_2 + (\sup_{\Omega} |\bar{\varphi}|)] \left[\frac{1}{\epsilon} \|f_t\|_{L^2(\Sigma_0)}^2 + \epsilon \|u_{\nu,A}\|_{L^2(\Sigma_0)}^2 + T_2 \|G\|_{L^2(Q_0)}^2 \right]. \end{aligned}$$

By letting $\epsilon = \{2C[T_2 + (\sup_{\Omega} |\bar{\varphi}|)]\}^{-1}$ and absorbing the $\|u_{\nu,A}\|_{L^2(\Sigma_0)}^2$ -term on the RHS by the LHS, we obtain

$$\begin{aligned} \|u_{\nu,A}\|_{L^2(\Sigma_0)}^2 &\leq C[T_2 + (\sup_{\Omega} |\bar{\varphi}|)] (\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) \\ &\quad + C[T_2 + (\sup_{\Omega} |\bar{\varphi}|)]^2 (\|f\|_{H^1(\Sigma_0)}^2 + \|G\|_{L^2(Q_0)}^2). \end{aligned}$$

Again, by using $\sup_{\Omega} |\bar{\varphi}| \leq 2c_2$, we obtain the second inequality. The proof is complete. \square

Remark 3.2. When the hypersurface Γ_S is horizontal, the corresponding estimate of the Neumann data $u_{\nu,A}$ given in Lemma 3.4 will be a quantitative version of [LLT86, (2.7)].

Now we are ready to prove Proposition 3.1.

Proof of Proposition 3.1. In (3.4), when $\tau = 0$, we have

$$\begin{aligned} |\mathcal{E}(u) - e(0)| &= |2 \operatorname{Re} \int_{\Sigma_0} \bar{f}_t u_{\nu,A} d\sigma + \operatorname{Re} \int_{Q_0} 2\bar{u}_t G| \\ &\leq 2 \int_{\Sigma_0} |f_t| |u_{\nu,A}| d\sigma + 2 \int_{Q_0} |u_t| |G| \\ &\leq 2 \|f_t\|_{L^2(\Sigma_0)} \|u_{\nu,A}\|_{L^2(\Sigma_0)} + 2 \|u_t\|_{L^2(Q_0)} \|G\|_{L^2(Q_0)}. \end{aligned} \quad (3.21)$$

For $\|u_t\|_{L^2(Q_0)}$, noticing $\|u_t\|_{L^2(H_{\tau})}^2 \leq e(\tau)$, so by (3.6) with K set to be T_2 , we have

$$\begin{aligned} \|u_t\|_{L^2(Q_0)}^2 &= \int_0^{T_2} \|u_t\|_{L^2(H_{\tau})}^2 d\tau \leq \int_0^{T_2} e(\tau) d\tau \\ &\leq T_2 (e^{T_2/T_2} - 1) (\mathcal{E}(u) + T_2 \|G\|_{L^2(Q_0)}^2) + 2 \int_{\Sigma_0} |u_t u_{\nu,A}| d\sigma \end{aligned}$$

$$\leq CT_2(\mathcal{E}(u) + T_2\|G\|_{L^2(Q_0)}^2 + 2\|f_t\|_{L^2(\Sigma_0)}\|u_{\nu,A}\|_{L^2(\Sigma_0)}).$$

Combining this with the estimates of $\mathcal{E}(u)$ and $\|u_{\nu,A}\|_{L^2(\Sigma_0)}$ given in Lemma 3.4, we obtain

$$\begin{aligned} \|u_t\|_{L^2(Q_0)}^2 &\leq CT_2(\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) + C(T_2 + 1)^2(\|G\|_{L^2(Q_0)}^2 + \|f\|_{H^1(\Sigma_0)}^2) \\ &\quad + CT_2^2\|G\|_{L^2(Q_0)}^2 + CT_2^2\|f_t\|_{L^2(\Sigma_0)}^2 + C(T_2 + 1)(\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) \\ &\quad + C(T_2 + 1)^2(\|f\|_{H^1(\Sigma_0)}^2 + \|G\|_{L^2(Q_0)}^2) \\ &\leq C\langle T_2 \rangle(\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) + C\langle T_2 \rangle^2(\|G\|_{L^2(Q_0)}^2 + \|f\|_{H^1(\Sigma_0)}^2), \end{aligned} \quad (3.22)$$

for some constant C depending only on A and the dimension n . Substituting (3.22) into (3.21), we obtain

$$\begin{aligned} |\mathcal{E}(u) - e(0)| &\leq 2\|f_t\|_{L^2(\Sigma_0)}\|u_{\nu,A}\|_{L^2(\Sigma_0)} + C\langle T_2 \rangle^{1/2}(\|\nabla u_0\|_{L^2(\Omega)} + \|u_1\|_{L^2(\Omega)})\|G\|_{L^2(Q_0)} \\ &\quad + C\langle T_2 \rangle(\|G\|_{L^2(Q_0)} + \|f\|_{H^1(\Sigma_0)})\|G\|_{L^2(Q_0)}. \end{aligned}$$

Again, by the estimate of $\|u_{\nu,A}\|_{L^2(\Sigma_0)}$ given in Lemma 3.4, we arrive at

$$\begin{aligned} |\mathcal{E}(u) - e(0)| &\leq C\langle T_2 \rangle^{1/2}[\|\nabla u_0\|_{L^2(\Omega)} + \|u_1\|_{L^2(\Omega)} + \langle T_2 \rangle^{1/2}(\|f\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)})] \\ &\quad \times (\|f\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)}) \end{aligned}$$

The proof is complete. \square

4. APPROXIMATION OF THE SOLUTION

The results in Section 3 are based on the prerequisite $u \in C^2$. This is not true for the system (1.1) when:

- (1) only the compatibility condition up to order zero is satisfied;
- (2) and the initial/boundary data are merely H^1 .

However, these two issues can be overcome by approximation. That is to say, we can find smooth sequences $u_{0,\epsilon}$, $u_{1,\epsilon}$, f_ϵ , G_ϵ which converge to u_0 , u_1 , f , G , respectively, and under these approximate data, we obtain approximate solutions u_ϵ , which will converge in $H^1(\Gamma_S)$. For the compatibility conditions issue, we shall modify f_ϵ . And for the regularity issue, with the help of the estimate given in Lemma 3.4, we are able to approximate the system (1.1) with smooth enough initial/boundary data and to show the corresponding approximate solution has a limit in $H^1(\Gamma_S)$ and the corresponding energy is well-defined.

For readers convenience we reproduce [LLT86, Remark 2.10] below.

Lemma 4.1. *Assume A is a C^1 -smooth real-valued symmetric matrix function, and $\partial\Omega$ is C^2 . Let u be a solution of the system (1.1) with (G, f, u_0, u_1) satisfying the regularity assumptions (m is a non-negative integer)*

$$\begin{cases} G \in L^1(0, T; H^m(\Omega)), \quad \frac{d^m G}{dt^m} \in L^1(0, T; L^2(\Omega)), \\ f \in H^{m+1}(\Sigma) := L^2(0, T; H^{m+1}(\partial\Omega)) \cap H^{m+1}(0, T; L^2(\partial\Omega)), \\ u_0 \in H^{m+1}(\Omega), \quad u_1 \in H^m(\Omega), \end{cases}$$

and satisfying all necessary compatibility conditions up to order m . Then

$$u \in C([0, T]; H^{m+1}(\Omega)), \quad \frac{d^{(m+1)}u}{dt^{(m+1)}} \in C([0, T]; L^2(\Omega)), \quad \text{and} \quad \frac{\partial u}{\partial \nu} \in H^m(\Sigma).$$

Remark 4.1. The case of Lemma 4.1 when $\nabla \cdot (A(x)\nabla)$ is replaced by Δ is covered by [LLT86, Remark 2.10]. And in [LLT86, Section 4] the authors discussed how to generalize from Δ to $\nabla \cdot (A(x)\nabla)$ when $m = 0, 1$. Actually, by following the same steps in the proof

of [LLT86, Theorem 2.2], we can generalize the scenario to any integer m not only 0 and 1, and the proof is straightforward so we omit it.

Readers should note that, in this work, the function space for the source Q is set to be $L^2(Q)$, which is a subset of the space $L^1(0, T; H^m(\Omega))$ with $m = 0$ used in [LLT86, Remark 2.10].

4.1. The compatibility issue. We first deal with the compatibility condition issue. Recall $\mathcal{E}(u)$ stands for $\mathcal{E}(u; \Gamma_S)$.

Lemma 4.2. *Given Assumptions 1.1 and 1.2. In system (1.1), assume $u_0, u_1 \in C^\infty(\overline{\Omega})$, $f \in C^\infty(\overline{\Sigma})$, $G \in C^\infty(\overline{Q})$, with $u_0(x) = f(x, 0)$ on $\partial\Omega$. Then we have $u \in H^1(\Gamma_S)$ and*

$$\begin{aligned} |\mathcal{E}(u) - \int_{\Omega} (|\nabla u_0|_A^2 + |u_1|^2) dx| \\ \leq C \langle T_2 \rangle^{1/2} [\|\nabla u_0\|_{L^2(\Omega)} + \|u_1\|_{L^2(\Omega)} + \langle T_2 \rangle^{1/2} (\|f\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)})] \\ \times (\|f\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)}). \end{aligned}$$

and

$$\|u_{\nu, A}\|_{L^2(\Sigma_0)}^2 \leq C \langle T_2 \rangle (\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) + C \langle T_2 \rangle^2 (\|G\|_{L^2(Q_0)}^2 + \|f\|_{H^1(\Sigma_0)}^2).$$

Proof. We shall find a sequence of C^2 -smooth solutions $\{u_\epsilon\}_{\epsilon>0}$ of

$$\begin{cases} \partial_t^2 u_\epsilon - \nabla \cdot (A(x) \nabla u_\epsilon) = G & \text{in } Q, \\ u_\epsilon = f_\epsilon & \text{on } \Sigma, \\ u_\epsilon = u_0, \partial_t u_\epsilon = u_1 & \text{on } \Omega \times \{t = 0\}, \end{cases} \quad (4.1)$$

by using the regularity result in Lemma 4.1. For this, we need smoothness of u_0, u_1, f and G , which are already assumed in this lemma, and we also need certain higher order compatibility conditions to be satisfied on $\Omega \times \{t = 0\}$. To guarantee the compatibility conditions, we modify the Dirichlet boundary datum f .

Let us construct a series of Dirichlet boundary data $\{f_\epsilon\}_{\epsilon>0}$ in the following way. First, we define u_k in $\overline{\Omega}$ iteratively for $k \geq 2$ from u_0 and u_1 which is already given as the initial data,

$$u_k(x) := \partial_t^{k-2} G(x, 0) + \nabla \cdot (A(x) \nabla u_{k-2}(x)), \quad x \in \overline{\Omega}.$$

recall that $u_0, u_1 \in C^\infty(\overline{\Omega})$. Recall the smoothness of A stipulated at the beginning of the article, i.e. $A \in C^{m+4}(\overline{\Omega})$. This guarantees

$$u_{2k}, u_{2k+1} \in C^{m+5-2k}(\overline{\Omega}), \quad \forall 1 \leq k \leq \frac{K-1}{2}, \quad (4.2)$$

for certain integer K which shall be determined later. Then, we fix a cutoff function $\chi \in C_c^\infty(\mathbb{R})$ satisfying $\chi(t) = 1$ when $|t| \leq 1$ and $\chi(t) = 0$ when $|t| \geq 2$, and we set

$$f_\epsilon(x, t) := \chi(t/\epsilon) \sum_{k=0}^K \frac{t^k}{k!} u_k(x) + (1 - \chi(t/\epsilon)) f(x, t), \quad \forall (x, t) \in \overline{\Sigma}. \quad (4.3)$$

By (4.2) and (4.3), $f_\epsilon \in C^{n+5-K}(\overline{\Sigma})$. Now the compatibility conditions up to the order K required by Lemma 4.1, i.e.,

$$u_k = \partial_t^k f_\epsilon \text{ on } \partial\Omega \times \{t = 0\}, \quad \forall k : 0 \leq k \leq K, \quad (4.4)$$

is satisfied (see also [Eva10, §7.2.3 (62)]). Therefore, we can use Lemma 4.1 up to order $N := \min\{n + 5 - K, K\}$ to conclude that the corresponding solution u_ϵ satisfies

$$u_\epsilon \in C([0, T]; H^N(\Omega)), \quad \partial_t u_\epsilon \in C([0, T]; H^{N-1}(\Omega)).$$

By the Sobolev embedding theorem we know $H^N(\Omega) \subset C^2(\overline{\Omega})$ when $N - n/2 \geq 2$, so we set $K = \lceil n/2 \rceil + 2$, and thus

$$u_\epsilon \in C([0, T]; C^2(\overline{\Omega})), \quad \partial_t u_\epsilon \in C([0, T]; C^1(\overline{\Omega})), \quad (4.5)$$

so $\nabla \cdot (A \nabla u_\epsilon(\cdot, t)) \in C(\overline{\Omega})$ for each t . Therefore, the equation $(\partial_t^2 - \Delta)u_\epsilon = 0$ implies

$$\partial_t^2 u_\epsilon(x, t) \text{ is continuous with respect to } t. \quad (4.6)$$

By (4.5) and (4.6) we obtain

$$u_\epsilon \in C^2(\overline{Q}). \quad (4.7)$$

The C^2 -smoothness of u_ϵ guarantees us to use Proposition 3.1 to conclude

$$\begin{aligned} |\mathcal{E}(u_\epsilon) - \int_{\Omega} (|\nabla u_0|_A^2 + |u_1|^2) dx| \\ \leq C \langle T_2 \rangle^{1/2} [\|\nabla u_0\|_{L^2(\Omega)} + \|u_1\|_{L^2(\Omega)} + \langle T_2 \rangle^{1/2} (\|f_\epsilon\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)})] \\ \times (\|f_\epsilon\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)}), \end{aligned} \quad (4.8)$$

for some constant C depending only on A and n . Further, by the linearity of (4.1) we obtain

$$\mathcal{E}(u_{\epsilon_1} - u_{\epsilon_2}) \leq C \langle T_2 \rangle \|f_{\epsilon_1} - f_{\epsilon_2}\|_{H^1(\Sigma_0)}^2, \quad \forall \epsilon_1, \epsilon_2 > 0. \quad (4.9)$$

Besides the compatibility conditions, the construction (4.3) also guarantees $f_\epsilon \rightarrow f$ in $H^1(\Sigma)$. Indeed, it is straightforward to check that

$$\|f - f_\epsilon\|_{L^2(\Sigma)} \rightarrow 0, \quad \|\nabla|_{\Sigma, x}(f - f_\epsilon)\|_{L^2(\Sigma)} \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0^+, \quad (4.10)$$

where $\nabla|_{\Sigma, x}$ stands for the spatial tangential gradient on Σ . Therefore, to show $f_\epsilon \rightarrow f$ in $H^1(\Sigma)$, it is left to show $\|\partial_t(f - f_\epsilon)\|_{L^2(\Sigma)} \rightarrow 0$ as $\epsilon \rightarrow 0^+$. By (4.3) one can compute

$$\begin{aligned} \partial_t(f_\epsilon - f)(x, t) &= \frac{1}{\epsilon} \chi'(\frac{t}{\epsilon}) \left[\sum_{k=0}^K \frac{t^k}{k!} u_k(x) - f(x, t) \right] + \chi(\frac{t}{\epsilon}) \left[\sum_{k=0}^{K-1} \frac{t^k}{k!} u_k(x) - f'(x, t) \right] \\ &= \frac{1}{\epsilon} \chi'(\frac{t}{\epsilon}) \left[\sum_{k=0}^K \frac{t^k}{k!} u_k(x) - u_0(x) - f'(x, \xi_t) t \right] + \chi(\frac{t}{\epsilon}) \left[\sum_{k=0}^{K-1} \frac{t^k}{k!} u_k - f' \right] \\ &= \frac{t}{\epsilon} \chi'(\frac{t}{\epsilon}) \left[\sum_{k=1}^K \frac{t^{k-1}}{k!} u_k(x) - f'(x, \xi_t) \right] + \chi(\frac{t}{\epsilon}) \left[\sum_{k=0}^{K-1} \frac{t^k}{k!} u_k(x) - f'(x, t) \right], \end{aligned}$$

where we used the compatibility condition “ $u_0(x) = f(x, 0)$ on $\partial\Omega$ ” and the Taylor’s expansion with Lagrange remainder, and the ξ_t comes from the Lagrange remainder. Note that for every $\epsilon > 0$, $\frac{(\cdot)}{\epsilon} \chi'(\frac{\cdot}{\epsilon})$ and $\chi(\frac{\cdot}{\epsilon})$ are uniformly bounded by $\max\{2 \sup |\chi'|, 1\}$ in $[0, T]$, so $|\partial_t(f_\epsilon - f)|^2$ is dominated by an integrable function F given as follows,

$$F(x, t) := \max\{2 \sup |\chi'|, 1\}^2 \left(\left| \sum_{k=1}^K \frac{t^{k-1}}{k!} u_k(x) - f'(x, \xi_t) \right| + \left| \sum_{k=0}^{K-1} \frac{t^k}{k!} u_k(x) - f'(x, t) \right| \right)^2.$$

Also, from (4.3) we see $f_\epsilon = f$ when $t \geq 2\epsilon$, so $\partial_t(f_\epsilon - f) = 0$ when $t \geq 2\epsilon$ and hence $\partial_t(f_\epsilon - f) \rightarrow 0$ almost everywhere in Σ as $\epsilon \rightarrow 0^+$. Therefore, by Lebesgue’s dominated convergence theorem, we obtain

$$\|\partial_t(f - f_\epsilon)\|_{L^2(\Sigma)} \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0^+. \quad (4.11)$$

Combining (4.10) with (4.11), we arrive at

$$\|f - f_\epsilon\|_{H^1(\Sigma)} \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0^+. \quad (4.12)$$

Now, combining (4.12) with (4.9), we see $\mathcal{E}(u_{\epsilon_1} - u_{\epsilon_2})$ goes to zero as $\epsilon_1, \epsilon_2 \rightarrow 0^+$, which implies $\{\nabla_{\Gamma_S} u_\epsilon\}$ is a Cauchy sequence in $L^2(\Gamma_S)$ and $\{\partial_t u_\epsilon\}$ is a Cauchy sequence in $L^2(\Gamma_S)$.

We claim that “ $\{\nabla_{\Gamma_S} u_\epsilon\}$ being a Cauchy sequence in $L^2(\Gamma_S)$ ” is enough to conclude $\{u_\epsilon\}$ is a Cauchy sequence in $H^1(\Gamma_S)$. This is due to the reason that every u_ϵ has the same trace on $\partial\Gamma_S$. Hence, we can define $u|_{\Gamma_S}$ as the limit of $u_\epsilon|_{\Gamma_S}$, i.e.,

$$u|_{\Gamma_S} := \lim_{\epsilon \rightarrow 0^+} u_\epsilon|_{\Gamma_S},$$

and the estimate (4.9) implies the limit $u|_{\Gamma_S}$ is in $H^1(\Gamma_S)$. Moreover, the energy $\mathcal{E}(u)$ is well-defined and by Lemma 2.2 and we see $\mathcal{E}(u_\epsilon) \rightarrow \mathcal{E}(u)$ as $\epsilon \rightarrow 0^+$. Now, taking the limit of (4.8), we obtain

$$\begin{aligned} & |\mathcal{E}(u) - \int_{\Omega} (|\nabla u_0|_A^2 + |u_1|^2) dx| \\ & \leq C \langle T_2 \rangle^{1/2} [\|\nabla u_0\|_{L^2(\Omega)} + \|u_1\|_{L^2(\Omega)} + \langle T_2 \rangle^{1/2} (\|f\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)})] \\ & \quad \times (\|f\|_{H^1(\Sigma_0)} + \|G\|_{L^2(Q_0)}). \end{aligned}$$

For $\|u_{\nu,A}\|_{L^2(\Sigma_0)}$, by combining the arguments above with Lemma 3.4, we can see $u_{\nu,A}$ is also well-define on Σ_0 and

$$\|u_{\nu,A}\|_{L^2(\Sigma_0)}^2 \leq C \langle T_2 \rangle (\|\nabla u_0\|_{L^2(\Omega)}^2 + \|u_1\|_{L^2(\Omega)}^2) + C \langle T_2 \rangle^2 (\|G\|_{L^2(Q_0)}^2 + \|f\|_{H^1(\Sigma_0)}^2).$$

The proof is complete. \square

4.2. The regularity issue. The C^∞ -smooth regularities requirements for u_0 , u_1 , f and G in Lemma 4.2 can be further improved. Recall the statement of Theorem 1.1..

Proof of Theorem 1.1. We shall utilize Lemma 4.2. One can find sequences $\{u_{0,\epsilon}\}_{\epsilon>0}$, $\{u_{1,\epsilon}\}_{\epsilon>0}$, $\{\tilde{f}_\epsilon\}_{\epsilon>0}$, $\{G_\epsilon\}_{\epsilon>0}$, satisfying the following requirements:

$$\begin{cases} \{u_{0,\epsilon}\}_{\epsilon>0} \subset C^\infty(\overline{\Omega}) \text{ such that } \|u_{0,\epsilon} - u_0\|_{H^1(\Omega)} \leq \epsilon, \\ \{u_{1,\epsilon}\}_{\epsilon>0} \subset C^\infty(\overline{\Omega}) \text{ such that } u_{1,\epsilon} \rightarrow u_1 \text{ in } L^2(\Omega), \\ \{\tilde{f}_\epsilon\}_{\epsilon>0} \subset C^\infty(\overline{\Sigma}) \text{ such that } \|\tilde{f}_\epsilon - f\|_{H^1(\Sigma)} \leq \epsilon, \\ \{G_\epsilon\}_{\epsilon>0} \subset C^\infty(\overline{Q}) \text{ such that } G_\epsilon \rightarrow G \text{ in } L^2(Q). \end{cases} \quad (4.13)$$

The condition “ $u_0(x) = f(x, 0)$ on $\partial\Omega$ ” mentioned in (1.2) does not guarantee $u_{0,\epsilon}(x) = \tilde{f}_\epsilon(x, 0)$ on $\partial\Omega$. Hence we need to modify \tilde{f}_ϵ to f_ϵ so that $(u_{0,\epsilon}, u_{1,\epsilon}, f_\epsilon, G_\epsilon)$ meet the requirements of Lemma 4.2.

Similar to (4.3), we fix a cutoff function $\chi \in C_c^\infty(\mathbb{R})$ satisfying $\chi(t) = 1$ when $|t| \leq 1$ and $\chi(t) = 0$ when $|t| \geq 2$, and set

$$f_\epsilon(x, t) := \chi(t/\epsilon) u_{0,\epsilon}(x) + (1 - \chi(t/\epsilon)) \tilde{f}_\epsilon(x, t), \quad \forall (x, t) \in \overline{\Sigma}. \quad (4.14)$$

Then,

$$f_\epsilon \in C^\infty(\overline{\Omega}), \quad \text{and} \quad u_{0,\epsilon}(x) = f_\epsilon(x, 0) \text{ on } \partial\Omega. \quad (4.15)$$

It can be seen that (see (4.10))

$$\|f - f_\epsilon\|_{L^2(\Sigma)} \rightarrow 0, \quad \|\nabla|_{\Sigma,x}(f - f_\epsilon)\|_{L^2(\Sigma)} \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0^+, \quad (4.16)$$

so, to guarantee $f_\epsilon \rightarrow f$ in $H^1(\Sigma)$, it is left to show $\|\partial_t(f - f_\epsilon)\|_{L^2(\Sigma)} \rightarrow 0$ as $\epsilon \rightarrow 0^+$. By (4.14) we have

$$\begin{aligned} & \partial_t(f_\epsilon - f)(x, t) = \partial_t(\tilde{f}_\epsilon - f) - \chi(t/\epsilon) \tilde{f}'_\epsilon + \frac{1}{\epsilon} \chi'(t/\epsilon) (u_{0,\epsilon}(x) - \tilde{f}_\epsilon(x, t)) \\ & = \partial_t(\tilde{f}_\epsilon - f) - \chi(t/\epsilon) \tilde{f}'_\epsilon \\ & \quad + \frac{t}{\epsilon} \chi'(t/\epsilon) \frac{[u_{0,\epsilon}(x) - u_0(x)] + [f(x, 0) - \tilde{f}_\epsilon(x, 0)] + [\tilde{f}_\epsilon(x, 0) - \tilde{f}_\epsilon(x, t)]}{t}, \end{aligned} \quad (4.17)$$

where we used the compatibility condition $u_0(x) = f(x, 0)$ on $\partial\Omega$.

The function $\frac{(\cdot)}{\epsilon}\chi'(\frac{\cdot}{\epsilon})$ is supported in the interval $[\epsilon, 2\epsilon]$ and is bounded by $2\max|\chi'|$, and $\|u_{0,\epsilon} - u_0\|_{H^1(\Omega)} \leq \epsilon$, so

$$\begin{aligned} \int_{\Sigma} \left| \frac{t}{\epsilon}\chi'(\frac{t}{\epsilon}) \frac{u_{0,\epsilon}(x) - u_0(x)}{t} \right|^2 &\leq C \int_{\epsilon}^{2\epsilon} \left(\int_{\partial\Omega} \left| \frac{u_{0,\epsilon}(x) - u_0(x)}{\epsilon} \right|^2 \right) dt \\ &= C\epsilon^{-1} \|u_{0,\epsilon} - u_0\|_{L^2(\partial\Omega)}^2 \leq C\epsilon^{-1} \|u_{0,\epsilon} - u_0\|_{H^1(\Omega)}^2 \\ &\leq C\epsilon, \end{aligned} \quad (4.18)$$

where we used the trace theorem. Similarly, we have

$$\int_{\Sigma} \left| \frac{t}{\epsilon}\chi'(\frac{t}{\epsilon}) \frac{f(x, 0) - \tilde{f}_{\epsilon}(x, 0)}{t} \right|^2 \leq C\epsilon^{-1} \|f - \tilde{f}_{\epsilon}\|_{L^2(\partial\Omega)}^2 \leq C\epsilon^{-1} \|f - \tilde{f}_{\epsilon}\|_{H^1(\Sigma)}^2 \leq C\epsilon. \quad (4.19)$$

We can also estimate the last term in (4.17). Note that \tilde{f}_{ϵ} is a smooth function, so we have

$$\begin{aligned} \int_{\Sigma} \left| \frac{t}{\epsilon}\chi'(\frac{t}{\epsilon}) \frac{\tilde{f}_{\epsilon}(x, 0) - \tilde{f}_{\epsilon}(x, t)}{t} \right|^2 d\sigma &= \int_{\Sigma} \left| \frac{t}{\epsilon}\chi'(\frac{t}{\epsilon}) \right|^2 |\partial_t \tilde{f}_{\epsilon}(x, t) + \mathcal{O}(t)|^2 d\sigma \\ &\leq 2 \int_{\Sigma} \left| \frac{t}{\epsilon}\chi'(\frac{t}{\epsilon}) \right|^2 |\partial_t \tilde{f}_{\epsilon}(x, t)|^2 d\sigma + \mathcal{O}(\epsilon^2). \end{aligned}$$

The function $\left| \frac{t}{\epsilon}\chi'(\frac{t}{\epsilon}) \right|^2 |\partial_t \tilde{f}_{\epsilon}(x, t)|^2$ is dominated by $C|\partial_t \tilde{f}_{\epsilon}(x, t)|^2$ for certain constant C , whose Lebesgue integral in Σ is bounded by $C\|\tilde{f}_{\epsilon}\|_{H^1(\Sigma)}$, and hence bounded by $C\|f\|_{H^1(\Sigma)} + 1$ when ϵ is small enough. Also, the function $\left| \frac{t}{\epsilon}\chi'(\frac{t}{\epsilon}) \right|^2 |\partial_t \tilde{f}_{\epsilon}(x, t)|^2$ converges to 0 for $\forall(x, t) \in \bar{\Sigma}$ as $\epsilon \rightarrow 0^+$, which is because $\left| \frac{t}{\epsilon}\chi'(\frac{t}{\epsilon}) \right|^2$ converges to 0 for $\forall t \in [0, T]$ as $\epsilon \rightarrow 0^+$. Therefore, by Lebesgue's dominated convergence theorem we have

$$\lim_{\epsilon \rightarrow 0^+} \int_{\Sigma} \left| \frac{t}{\epsilon}\chi'(\frac{t}{\epsilon}) \frac{\tilde{f}_{\epsilon}(x, 0) - \tilde{f}_{\epsilon}(x, t)}{t} \right|^2 d\sigma \leq 2 \lim_{\epsilon \rightarrow 0^+} \int_{\Sigma} \left| \frac{t}{\epsilon}\chi'(\frac{t}{\epsilon}) \right|^2 |\partial_t \tilde{f}_{\epsilon}(x, t)|^2 d\sigma = 0. \quad (4.20)$$

Combining (4.18), (4.19), (4.20) with (4.17), we conclude $\|\partial_t(f_{\epsilon} - f)\|_{L^2(\Sigma)} \rightarrow 0$, so

$$\|f_{\epsilon} - f\|_{H^1(\Sigma)} \rightarrow 0, \quad \epsilon \rightarrow 0^+. \quad (4.21)$$

By (4.13) and (4.15), we see $(u_{0,\epsilon}, u_{1,\epsilon}, f_{\epsilon}, G_{\epsilon})$ meet the requirement of Lemma 4.2, so the corresponding solution u_{ϵ} satisfies

$$\begin{aligned} |\mathcal{E}(u_{\epsilon}) - \int_{\Omega} (|\nabla u_{0,\epsilon}|_A^2 + |u_{1,\epsilon}|^2) dx| \\ \leq C\langle T_2 \rangle^{1/2} [\|\nabla u_{0,\epsilon}\|_{L^2(\Omega)} + \|u_{1,\epsilon}\|_{L^2(\Omega)} + \langle T_2 \rangle^{1/2} (\|f_{\epsilon}\|_{H^1(\Sigma_0)} + \|G_{\epsilon}\|_{L^2(Q_0)})] \\ \times (\|f_{\epsilon}\|_{H^1(\Sigma_0)} + \|G_{\epsilon}\|_{L^2(Q_0)}), \end{aligned} \quad (4.22)$$

and

$$\begin{aligned} |\mathcal{E}(u_{\epsilon_1} - u_{\epsilon_2}) - \int_{\Omega} (|\nabla(u_{0,\epsilon_1} - u_{0,\epsilon_2})|_A^2 + |u_{1,\epsilon_1} - u_{1,\epsilon_2}|^2) dx| \\ \leq C\langle T_2 \rangle^{1/2} [\|\nabla(u_{0,\epsilon_1} - u_{0,\epsilon_2})\|_{L^2(\Omega)} + \|u_{1,\epsilon_1} - u_{1,\epsilon_2}\|_{L^2(\Omega)} + \langle T_2 \rangle^{1/2} (\|f_{\epsilon_1} - f_{\epsilon_2}\|_{H^1(\Sigma_0)} \\ + \|G_{\epsilon_1} - G_{\epsilon_2}\|_{L^2(Q_0)})] \times (\|f_{\epsilon_1} - f_{\epsilon_2}\|_{H^1(\Sigma_0)} + \|G_{\epsilon_1} - G_{\epsilon_2}\|_{L^2(Q_0)}). \end{aligned} \quad (4.23)$$

Combining the limits given in (4.13) and (4.21) with (4.23), we see the limit of $\{u_{\epsilon}\}$ and $\{\partial_t u_{\epsilon}\}$ exist in $H^1(\Gamma_S)$ and in $L^2(\Sigma_S)$, respectively, and the limit coincides with the solution u . Therefore, by taking the limit of (4.22), we arrive at the conclusion. \square

Proof of Corollary 1.2. Recall the notation $\mathcal{E}(u; \Gamma_\tau)$ defined in (1.7). When $f = 0$ and $G = 0$, by Theorem 1.1 we have

$$\mathcal{E}(u; \Gamma_\tau) = \int_{\Omega} (|\nabla u_0|_A^2 + |u_1|^2) dx,$$

which is true for any τ . The proof is done. \square

We can borrow the arguments in the proof of Theorem 1.1 to show Theorem 1.3.

Proof of Theorem 1.3. The sequences $(u_{0,\epsilon}, u_{1,\epsilon}, f_\epsilon, G_\epsilon)$ constructed in the proof of Theorem 1.1 meet the requirement of Lemma 4.2, so the corresponding solution u_ϵ satisfies

$$\|u_{\epsilon,\nu,A}\|_{L^2(\Sigma_0)}^2 \leq C\langle T_2 \rangle (\|\nabla u_{0,\epsilon}\|_{L^2(\Omega)}^2 + \|u_{1,\epsilon}\|_{L^2(\Omega)}^2) + C\langle T_2 \rangle^2 (\|G_\epsilon\|_{L^2(Q_0)}^2 + \|f_\epsilon\|_{H^1(\Sigma_0)}^2).$$

By taking the limit, we obtain the result. \square

5. CONCLUDING REMARK

One of the possible future directions succeeding this work is to study the case where the initial/boundary data possess higher order regularities. The corresponding results are classical for horizontal planes, i.e. for $\Gamma_S = \Omega \times \{t = \tau\}$, see e.g. [Eva10, §7.2.3] for Dirichlet boundary condition cases. However, the method used for horizontal planes seems not to directly apply to the non-timelike case. More careful investigations are needed.

Another direction is to look at the timelike case. Physically speaking, the timelike case corresponds to supersonic waves. In this case the shock wave phenomenon could happen, which could cause additional singularities to the solution. Hence, a reasonable expectation is that the regularity of the solution on timelike hypersurfaces will not be as good as on non-timelike hypersurfaces.

Acknowledgements. The author was supported by the Academy of Finland (Finnish Centre of Excellence in Inverse Modelling and Imaging, grant numbers 312121 and 309963).

REFERENCES

- [Ava97] G. Avalos, *Sharp regularity estimates for solutions of the wave equation and their traces with prescribed Neumann data*, Appl. Math. Optim. **35** (1997), no. 2, 203–219. MR1424790
- [Eva10] L. C. Evans, *Partial differential equations*, Second, Graduate Studies in Mathematics, vol. 19, American Mathematical Society, Providence, RI, 2010. MR2597943
- [Ike05] R. Ikehata, *Local energy decay for linear wave equations with variable coefficients*, J. Math. Anal. Appl. **306** (2005), no. 1, 330–348. MR2132904
- [LLT86] I. Lasiecka, J.-L. Lions, and R. Triggiani, *Nonhomogeneous boundary value problems for second order hyperbolic operators*, J. Math. Pures Appl. (9) **65** (1986), no. 2, 149–192. MR867669
- [LM72] J.-L. Lions and E. Magenes, *Non-homogeneous boundary value problems and applications. Vol. I–III*, Die Grundlehren der mathematischen Wissenschaften, Band 181, Springer-Verlag, New York-Heidelberg, 1972–1973. MR0350177
- [LT81] I. Lasiecka and R. Triggiani, *A cosine operator approach to modeling $L_2(0, T; L_2(\Gamma))$ –boundary input hyperbolic equations*, Appl. Math. Optim. **7** (1981), no. 1, 35–93. MR600559
- [LT83] I. Lasiecka and R. Triggiani, *Regularity of hyperbolic equations under $L_2(0, T; L_2(\Gamma))$ -Dirichlet boundary terms*, Appl. Math. Optim. **10** (1983), no. 3, 275–286. MR722491
- [LT88] I. Lasiecka and R. Triggiani, *A lifting theorem for the time regularity of solutions to abstract equations with unbounded operators and applications to hyperbolic equations*, Proc. Amer. Math. Soc. **104** (1988), no. 3, 745–755. MR964851
- [Mor61] C. S. Morawetz, *The decay of solutions of the exterior initial-boundary value problem for the wave equation*, Comm. Pure Appl. Math. **14** (1961), 561–568. MR132908
- [MPMS21] C. J. Meroño, L. Potenciano-Machado, and M. Salo, *The fixed angle scattering problem with a first order perturbation*, Ann. Henri Poincaré (2021).
- [MS21] S. Ma and M. Salo, *Fixed angle inverse scattering in the presence of a riemannian metric*, J. Inverse Ill-Posed Probl. (2021). Pre-published online. doi: 10.1515/jiip-2020-0119.

- [RS20a] Rakesh and M. Salo, *Fixed angle inverse scattering for almost symmetric or controlled perturbations*, SIAM J. Math. Anal. **52** (2020), no. 6, 5467–5499. MR4170189
- [RS20b] Rakesh and M. Salo, *The fixed angle scattering problem and wave equation inverse problems with two measurements*, Inverse Problems **36** (2020), no. 3, 035005, 42. MR4068234
- [RU14] Rakesh and G. Uhlmann, *Uniqueness for the inverse backscattering problem for angularly controlled potentials*, Inverse Problems **30** (2014), no. 6, 065005, 24. MR3224125
- [Sak70a] R. Sakamoto, *Mixed problems for hyperbolic equations. I. Energy inequalities*, J. Math. Kyoto Univ. **10** (1970), 349–373. MR283400
- [Sak70b] R. Sakamoto, *Mixed problems for hyperbolic equations. II. Existence theorems with zero initial data and energy inequalities with initial data*, J. Math. Kyoto Univ. **10** (1970), 403–417. MR283401
- [Tay11] M. E. Taylor, *Partial differential equations I. Basic theory*, Second, Applied Mathematical Sciences, vol. 115, Springer, New York, 2011. MR2744150
- [Wal84] R. M. Wald, *General relativity*, University of Chicago Press, Chicago, IL, 1984. MR757180

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ, JYVÄSKYLÄ, FINLAND
Email address: shiqi.s.ma@jyu.fi