

REGULARITY THEORY FOR MIXED LOCAL AND NONLOCAL PARABOLIC p -LAPLACE EQUATIONS

BIN SHANG, YUZHOU FANG AND CHAO ZHANG*

ABSTRACT. We investigate the mixed local and nonlocal parabolic p -Laplace equation

$$\partial_t u(x, t) - \Delta_p u(x, t) + \mathcal{L}u(x, t) = 0,$$

where Δ_p is the local p -Laplace operator and \mathcal{L} is the nonlocal p -Laplace operator. Based on the combination of suitable Caccioppoli-type inequality and Logarithmic Lemma with a De Giorgi-Nash-Moser iteration, we establish the local boundedness and Hölder continuity of weak solutions for such equations.

1. INTRODUCTION

In this paper, we are concerned with the local behaviour of weak solutions to the following mixed problem

$$\partial_t u(x, t) - \Delta_p u(x, t) + \mathcal{L}u(x, t) = 0 \quad \text{in } Q_T, \quad 1 < p < \infty, \quad (1.1)$$

where $Q_T := \Omega \times (0, T)$ with $T > 0$ and Ω is a bounded domain in \mathbb{R}^N . This kind of evolution equations arises from the Lévy process, image processing etc; see [14] and references therein. The local p -Laplace operator Δ_p is defined as

$$\Delta_p u := \operatorname{div}(|\nabla u|^{p-2} \nabla u),$$

and \mathcal{L} is a nonlocal p -Laplace operator given by

$$\mathcal{L}u(x, t) = \text{P.V.} \int_{\mathbb{R}^N} K(x, y, t) |u(x, t) - u(y, t)|^{p-2} (u(x, t) - u(y, t)) dy, \quad (1.2)$$

where the symbol P.V. stands for the Cauchy principle value. Here K is a symmetric kernel fulfilling

$$K(x, y, t) = K(y, x, t)$$

and

$$\frac{\Lambda^{-1}}{|x - y|^{N+sp}} \leq K(x, y, t) \leq \frac{\Lambda}{|x - y|^{N+sp}} \quad (1.3)$$

with $\Lambda \geq 1$ and $0 < s < 1$ for all $x, y \in \mathbb{R}^N$ and $t \in (0, T)$.

Before stating our main results, let us mention some known results. For the nonlocal parabolic equations of p -Laplacian type,

$$\partial_t u(x, t) + \mathcal{L}u(x, t) = 0, \quad (1.4)$$

the existence and uniqueness of strong solutions were verified by Vázquez [23], where the author studied the long time behaviours as well. Mazón-Rossi-Toledo [21] established the

2010 *Mathematics Subject Classification.* 35B45, 35B65, 35D30, 35K55, 35R11.

Key words and phrases. Local boundedness; Hölder continuity; mixed local and nonlocal parabolic p -Laplace equation.

*Corresponding author.

well-posedness of solutions to Eq. (1.4) together with the asymptotic property. When it comes to regularity theory for this equation, Strömqvist [22] obtained the existence and local boundedness of weak solutions provided $p \geq 2$. Hölder regularity with specific exponents in the case $p \geq 2$ was proved by Brasco-Lindgren-Strömqvist [4]. Furthermore, Ding-Zhang-Zhou [10] showed the local boundedness and Hölder continuity of weak solutions to the nonhomogeneous case under the conditions that $1 < p < \infty$ and $2 < p < \infty$ respectively. We refer the readers to [5, 16, 20, 24, 25] and references therein for more results.

In the mixed local and nonlocal setting, for the case $p = 2$,

$$-\Delta u + (-\Delta)^s u = 0, \quad (1.5)$$

Foondun [17] have derived Harnack inequality and interior Hölder estimates for nonnegative solutions, see also [7] for a diverse approach. In addition, the Harnack inequality regarding the parabolic version of (1.5) was established in [1, 6], where however the authors only proved such inequality for globally nonnegative solutions. Very recently, Garain-Kinnunen [19] proved a weak Harnack inequality with a tail term for sign changing solutions to the parabolic problem of (1.5). For what concerns maximum principles, interior sobolev regularity along with symmetry results among many other quantitative and qualitative properties for solutions to (1.5), one can see for instance [2, 3, 11–13]. In the nonlinear framework (i.e., $p \neq 2$), Garain-Kinnunen [18] developed the local regularity theory for

$$-\Delta_p u + \text{P.V.} \int_{\mathbb{R}^N} K(x, y) |u(x) - u(y)|^{p-2} (u(x) - u(y)) dy = 0$$

with $K(x, y) \simeq |x - y|^{-(N+sp)}$, involving boundedness, Hölder continuity, Harnack inequality as well as lower/upper semicontinuity of weak supersolutions/subsolutions. Nonetheless, to the best of our knowledge, there are little results concerning on the mixed local and nonlocal nonlinear parabolic problems. To this end, we aim to establish the interior Hölder regularity and boundedness of weak slutions to Eq. (1.1), which could be regarded as an analogue of [10, 18]. It is noteworthy that our results are new even for the case $p = 2$.

Before giving the notion of weak solutions to (1.1), let us recall the tail space

$$L_\alpha^q(\mathbb{R}^N) := \left\{ v \in L_{\text{loc}}^q(\mathbb{R}^N) : \int_{\mathbb{R}^N} \frac{|v(x)|^q}{1 + |x|^{N+\alpha}} dx < +\infty \right\}, \quad q > 0 \text{ and } \alpha > 0.$$

Then we define the tail appearing in estimates throughout this article,

$$\begin{aligned} \text{Tail}_\infty(v; x_0, r, I) &= \text{Tail}_\infty(v; x_0, r, t_0 - T_1, t_0 + T_2) \\ &:= \text{ess sup}_{t \in I} \left(r^p \int_{\mathbb{R}^N \setminus B_r(x_0)} \frac{|v(x, t)|^{p-1}}{|x - x_0|^{N+sp}} dx \right)^{\frac{1}{p-1}}, \end{aligned} \quad (1.6)$$

where $(x_0, t_0) \in \mathbb{R}^N \times (0, T)$ and the interval $I = [t_0 - T_1, t_0 + T_2] \subseteq (0, T)$. This is a parabolic counterpart to the tail introduced in [9]. It is easy to check that $\text{Tail}_\infty(v; x_0, r, I)$ is well-defined for any $v \in L^\infty(I; L_{sp}^{p-1}(\mathbb{R}^N))$.

Definition 1.1. *A function $u \in L^p(I; W_{\text{loc}}^{s,p}(\Omega)) \cap C(I; L_{\text{loc}}^2(\Omega)) \cap L^\infty(I; L_{sp}^{p-1}(\mathbb{R}^N))$ is a local weak subsolution (super-) to (1.1) if for any closed interval $I := [t_1, t_2] \subseteq (0, T)$, there holds that*

$$\int_{\Omega} u(x, t_2) \varphi(x, t_2) dx - \int_{\Omega} u(x, t_1) \varphi(x, t_1) dx - \int_{t_1}^{t_2} \int_{\Omega} u(x, t) \partial_t \varphi(x, t) dx dt$$

$$+ \int_{t_1}^{t_2} \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \varphi \, dx dt + \int_{t_1}^{t_2} \mathcal{E}(u, \varphi, t) \, dt \leq (\geq) 0, \quad (1.7)$$

for every nonnegative test function $\varphi \in L^p(I; W^{s,p}(\Omega)) \cap W^{1,2}(I; L^2(\Omega))$ with the property that φ has spatial support compactly contained in Ω , where

$$\mathcal{E}(u, \varphi, t) := \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \left[|u(x, t) - u(y, t)|^{p-2} (u(x, t) - u(y, t)) \right. \\ \left. (\varphi(x, t) - \varphi(y, t)) K(x, y, t) \right] dx dy.$$

A function u is a local weak solution to (1.1) if and only if u is a local weak subsolution and supersolution.

We now are in a position to state the main contribution of this work. First, we provide the local boundedness of weak solutions in the cases that $p > \frac{2N}{N+2}$ and $1 < p \leq \frac{2N}{N+2}$. For two real number, set

$$a \vee b := \max\{a, b\}, \quad a_+ := \max\{a, 0\}, \quad a_- := -\min\{a, 0\}.$$

Theorem 1.2 (Local boundedness). *Let $p > 2N/(2+N)$ and $q := \max\{p, 2\}$. Assume that u is a local weak subsolution to (1.1). Let $(x_0, t_0) \in Q_T$, $R \in (0, 1)$ and $Q_{\bar{R}}^- \equiv B_R(x_0) \times (t_0 - R^p, t_0)$ such that $\bar{B}_R(x_0) \subseteq \Omega$ and $[t_0 - R^p, t_0] \subseteq (0, T)$. Then it holds that*

$$\operatorname{ess\,sup}_{Q_{\bar{R}/2}^-} u \leq \operatorname{Tail}_{\infty}(u_+; x_0, R/2, t_0 - R^p, t_0) + C \left(\int_{Q_{\bar{R}}^-} u_+^q \, dx dt \right)^{\frac{p}{N(p\kappa - q)}} \vee 1,$$

where $\kappa := 1 + 2/N$ and $C > 0$ only depends on N, p, s and Λ .

In the scenario that $1 < p \leq 2N/(N+2)$, assuming that the weak subsolution have the following constructions: for $m > \max\left\{2, \frac{N(2-p)}{p}\right\}$, there exists a sequence of $\{u_k\}_{k \in \mathbb{N}}$ whose components are bounded subsolutions of (1.1) such that

$$\|u_k\|_{L_{\text{loc}}^{\infty}(0, T; L_{sp}^{p-1}(\mathbb{R}^N))} \leq C \quad (1.8)$$

and

$$u_k \rightarrow u \quad \text{in } L_{\text{loc}}^m(Q_T) \text{ as } k \rightarrow \infty. \quad (1.9)$$

Theorem 1.3 (Local boundedness). *Let $1 < p \leq 2N/(N+2)$, $\kappa = 1 + 2/N$ and $m > \max\left\{2, \frac{N(2-p)}{p}\right\}$. Suppose that $u \in L_{\text{loc}}^m(Q_T)$ with the properties (1.8) and (1.9) is a local weak subsolution of (1.1). Let $(x_0, t_0) \in Q_T$, $R \in (0, 1)$ and $Q_{\bar{R}}^- \equiv B_R(x_0) \times (t_0 - R^p, t_0)$ such that $\bar{B}_R(x_0) \subseteq \Omega$ and $[t_0 - R^p, t_0] \subseteq (0, T)$. Then it holds that*

$$\operatorname{ess\,sup}_{Q_{\bar{R}/2}^-} u \leq \operatorname{Tail}_{\infty}(u_+; x_0, R/2, t_0 - R^p, t_0) \\ + C \left(\int_{Q_{\bar{R}}^-} u_+^m \, dx dt \right)^{\frac{p}{(N+p)(m-2-\beta)}} \vee \left(\int_{Q_{\bar{R}}^-} u_+^m \, dx dt \right)^{\frac{p}{(N+p)(m-p-\beta)}},$$

where $\beta = N(m - p\kappa)/(p + N)$ and $C > 0$ only depends on N, p, s, m and Λ .

Based on the boundedness result (Theorem 1.2), we are able to deduce that the weak solutions are locally Hölder continuous for $p > 2$.

Theorem 1.4 (Hölder continuity). *Let $p > 2$. Assume that u is a local weak solution of (1.1). Let $(x_0, t_0) \in Q_T$, $R \in (0, 1)$ and $Q_R \equiv B_R(x_0) \times (t_0 - R^p, t_0 + R^p)$ such that $\overline{Q_R} \subseteq Q_T$. Moreover, there exists a constant $\alpha \in (0, p/(p-1))$ such that for every $\rho \in (0, R/2)$,*

$$\operatorname{ess\,osc}_{Q_{\rho, d\rho^p}} u < C \left(\frac{\rho}{R} \right)^\alpha \left[\operatorname{Tail}_\infty(u; x_0, R/2, t_0 - R^p, t_0 + R^p) + \left(\int_{Q_R} |u|^p dx dt \right)^{\frac{1}{2}} \vee 1 \right]$$

with $d \in (0, 1)$ and $C \geq 1$ depending on N, p, s and Λ .

The paper is organized as follows. In Section 2, we gather some notations and auxiliary inequalities. Necessary energy estimates are showed in Section 3. Sections 4 and 5 are devoted to proving the local boundedness and Hölder regularity of weak solutions, respectively.

2. PRELIMINARIES

In this section, we first give some notations for clarity and then provide some important inequalities to be used later.

2.1. Notation. Let $B_\rho(x)$ be the open ball with radius ρ and centered at $x \in \mathbb{R}^N$. We denote the parabolic cylinders by $Q_{\rho, r}(x, t) := B_\rho(x) \times (t - r, t + r)$, $Q_\rho(x, t) := Q_{\rho, \rho^p}(x, t) = B_\rho(x) \times (t - \rho^p, t + \rho^p)$ and $Q_\rho^-(x, t) := Q_{\rho, \rho^p}^-(x, t) = B_\rho(x) \times (t - \rho^p, t)$ with $r, \rho > 0$ and $(x, t) \in \mathbb{R}^N \times (0, T)$. If not important, or clear from the context, we simply write these symbols by $B_\rho = B_\rho(x)$, $Q_{\rho, r} = Q_{\rho, r}(x, t)$, $Q_\rho = Q_\rho(x, t)$ and $Q_\rho^- = Q_\rho^-(x, t)$. Moreover for $g \in L^1(V)$, we denote the integral average of g by

$$(g)_V := \int_V g(x) dx := \frac{1}{|V|} \int_V g(x) dx.$$

Define

$$J_p(a, b) = |a - b|^{p-2}(a - b)$$

for any $a, b \in \mathbb{R}$. We also use the notation

$$d\mu = d\mu(x, y, t) = K(x, y, t) dx dy.$$

It is worth mentioning that the constant C represents a general positive constant which may differ from each other.

Next, we will show several fundamental but very useful Sobolev inequalities.

2.2. Sobolev inequalities.

Lemma 2.1. *Let $1 \leq p, \ell \leq q < \infty$ satisfy $\frac{N}{p} - \frac{N}{q} \leq 1$ and*

$$\theta \left(1 - \frac{N}{p} + \frac{N}{q} \right) + (1 - \theta) \left(\frac{N}{q} - \frac{N}{\ell} \right) = 0$$

with $\theta \in (0, 1)$. Then there exists a constant $C > 0$ only depending on N, p, q, ℓ such that

$$\|u\|_{L^q(B_1)} \leq C \|Du\|_{L^p(B_1)}^\theta \|u\|_{L^\ell(B_1)}^{1-\theta} \quad (2.1)$$

for all $u \in W^{1,p}(B_1) \cap L^\ell(B_1)$.

Lemma 2.2. *Let $1 < p < N$. Then for every $u \in W^{1,p}(B_1)$, there holds that*

$$\|u\|_{L^{\frac{Np}{N-p}}(B_1)} \leq C \|u\|_{W^{1,p}(B_1)},$$

where $C > 0$ only depends on N and p .

Lemma 2.3. *Let $0 < t_1 < t_2$ and $p \in (1, \infty)$. Then for every*

$$u \in L^p(t_1, t_2; W^{1,p}(B_r)) \cap L^\infty(t_1, t_2; L^2(B_r)),$$

it holds that

$$\begin{aligned} & \int_{t_1}^{t_2} \int_{B_r} |u(x, t)|^{p(1+\frac{2}{N})} dx dt \\ & \leq C \left(r^p \int_{t_1}^{t_2} \int_{B_r} |\nabla u(x, t)|^p dx dt + \int_{t_1}^{t_2} \int_{B_r} |u(x, t)|^p dx dt \right) \\ & \quad \times \left(\operatorname{ess\,sup}_{t_1 < t < t_2} \int_{B_r} |u(x, t)|^2 dx \right)^{\frac{p}{N}}, \end{aligned} \tag{2.2}$$

where $C > 0$ only depends on p and N .

Proof. We divide this proof into two cases.

Case 1: $1 < p < N$. Applying Lemma 2.2 and Hölder inequality, we infer that

$$\begin{aligned} & \int_{t_1}^{t_2} \int_{B_1} |u(x, t)|^{p(1+\frac{2}{N})} dx dt \\ & = \int_{t_1}^{t_2} \int_{B_1} |u(x, t)|^{\frac{2p}{N}} |u(x, t)|^p dx dt \\ & \leq \int_{t_1}^{t_2} \left(\int_{B_1} |u(x, t)|^2 dx \right)^{\frac{p}{N}} \left(\int_{B_1} |u(x, t)|^{\frac{pN}{N-p}} dx \right)^{\frac{N-p}{N}} dt \\ & \leq C \left(\operatorname{ess\,sup}_{t_1 < t < t_2} \int_{B_1} |u(x, t)|^2 dx \right)^{\frac{p}{N}} \int_{t_1}^{t_2} \int_{B_1} |u(x, t)|^p + |\nabla u(x, t)|^p dx dt. \end{aligned}$$

By scaling argument, we get

$$\begin{aligned} & \int_{t_1}^{t_2} \int_{B_r} |u(x, t)|^{p(1+\frac{2}{N})} dx dt \\ & \leq C \left(r^p \int_{t_1}^{t_2} \int_{B_r} |\nabla u(x, t)|^p dx dt + \int_{t_1}^{t_2} \int_{B_r} |u(x, t)|^p dx dt \right) \\ & \quad \times \left(\operatorname{ess\,sup}_{t_1 < t < t_2} \int_{B_r} |u(x, t)|^2 dx \right)^{\frac{p}{N}}. \end{aligned}$$

Case 2: $p \geq N$. We can easily find that

$$1 \geq \frac{N}{p} - \frac{N}{p(1+\frac{2}{N})}, \quad p \left(1 + \frac{2}{N} \right) > 2$$

and

$$\theta \left(1 - \frac{N}{p} + \frac{N}{p(1+\frac{2}{N})} \right) + (1-\theta) \left(\frac{N}{p(1+\frac{2}{N})} - \frac{N}{2} \right) = 0$$

with $\theta = \frac{N}{N+2}$. Thus by Lemma 2.1, it follows that

$$\|u\|_{L^{p(1+\frac{2}{N})}(B_1)} \leq C \|Du\|_{L^p(B_1)}^{\frac{N}{N+2}} \|u\|_{L^2(B_1)}^{\frac{2}{N+2}}$$

for any $t \in (t_1, t_2)$. Using scaling argument, we have

$$\int_{B_r} |u(x, t)|^{p(1+\frac{2}{N})} dx \leq Cr^p \int_{B_r} |\nabla u(x, t)|^p dx \times \left(\int_{B_r} |u(x, t)|^2 dx \right)^{\frac{p}{N}}$$

for any $t \in (t_1, t_2)$. Integrating the above inequality over (t_1, t_2) , we get

$$\begin{aligned} & \int_{t_1}^{t_2} \int_{B_r} |u(x, t)|^{p(1+\frac{2}{N})} dx dt \\ & \leq Cr^p \int_{t_1}^{t_2} \int_{B_r} |\nabla u(x, t)|^p dx \times \left(\int_{B_r} |u(x, t)|^2 dx \right)^{\frac{p}{N}} dt \\ & \leq Cr^p \int_{t_1}^{t_2} \int_{B_r} |\nabla u(x, t)|^p dx dt \times \left(\operatorname{ess\,sup}_{t_1 < t < t_2} \int_{B_r} |u(x, t)|^2 dx \right)^{\frac{p}{N}}. \end{aligned}$$

□

Lemma 2.4. *Let $0 < t_1 < t_2$ and $p \in (1, \infty)$. Then for every*

$$u \in L^p(t_1, t_2; W^{1,p}(B_r)) \cap L^\infty(t_1, t_2; L^p(B_r)),$$

it holds that

$$\begin{aligned} & \int_{t_1}^{t_2} \int_{B_r} |u(x, t)|^{p(1+\frac{p}{N})} dx dt \\ & \leq C \left(r^p \int_{t_1}^{t_2} \int_{B_r} |\nabla u(x, t)|^p dx dt + \int_{t_1}^{t_2} \int_{B_r} |u(x, t)|^p dx dt \right) \\ & \quad \times \left(\operatorname{ess\,sup}_{t_1 < t < t_2} \int_{B_r} |u(x, t)|^p dx \right)^{\frac{p}{N}}, \end{aligned} \tag{2.3}$$

where $C > 0$ only depends on p and N .

Proof. It follows from Lemma 2.1 that

$$\|u\|_{L^{p(1+\frac{p}{N})}(B_1)} \leq C \|Du\|_{L^p(B_1)}^{\frac{N}{N+p}} \|u\|_{L^p(B_1)}^{\frac{p}{N+p}}$$

for all $t \in (t_1, t_2)$, where $C > 0$ only depends on p and N . Using scaling argument, we have

$$\int_{B_r} |u(x, t)|^{p(1+\frac{p}{N})} dx \leq Cr^p \int_{B_r} |\nabla u(x, t)|^p dx \times \left(\int_{B_r} |u(x, t)|^p dx \right)^{\frac{p}{N}}.$$

Integrating the above inequality over (t_1, t_2) , we get

$$\begin{aligned} & \int_{t_1}^{t_2} \int_{B_r} |u(x, t)|^{p(1+\frac{p}{N})} dx dt \\ & \leq Cr^p \int_{t_1}^{t_2} \int_{B_r} |\nabla u(x, t)|^p dx \times \left(\int_{B_r} |u(x, t)|^p dx \right)^{\frac{p}{N}} dt \\ & \leq Cr^p \int_{t_1}^{t_2} \int_{B_r} |\nabla u(x, t)|^p dx dt \times \left(\operatorname{ess\,sup}_{t_1 < t < t_2} \int_{B_r} |u(x, t)|^p dx \right)^{\frac{p}{N}}. \end{aligned}$$

□

3. ENERGY ESTIMATES

In this section, we will establish the Caccioppoli inequality and Logarithmic form inequality for mixed local and nonlocal type equation (1.1). The first step of the proof should be the regularization procedure with respect to the time variable, which can be performed by straightforward adaptation of standard reasonings as used in [4, 10]. We omit this step here.

Lemma 3.1 (Caccioppoli-type inequality). *Let $p > 1$ and u be a local subsolution of (1.1). Let $B_r \equiv B_r(x_0)$ satisfy $\overline{B_r} \subseteq \Omega$ and $0 < \tau_1 < \tau_2$, $\ell > 0$ satisfy $[\tau_1 - \ell, \tau_2] \subseteq (0, T)$. For any nonnegative functions $\psi \in C_0^\infty(B_r)$ and $\eta \in C^\infty(\mathbb{R})$ such that $\eta(t) \equiv 0$ if $t \leq \tau_1 - \ell$ and $\eta(t) \equiv 1$ if $t \geq \tau_1$, it holds that*

$$\begin{aligned}
& \int_{\tau_1 - \ell}^{\tau_2} \int_{B_r} |\nabla w_+(x, t)|^p \psi^p(x) \eta^2(t) \, dx dt + \operatorname{ess\,sup}_{\tau_1 < t < \tau_2} \int_{B_r} w_+^2(x, t) \psi^p(x) \, dx \\
& + \int_{\tau_1 - \ell}^{\tau_2} \int_{B_r} \int_{B_r} |w_+(x, t) \psi(x) - w_+(y, t) \psi(y)|^p \eta^2(t) \, d\mu dt \\
\leq & C \int_{\tau_1 - \ell}^{\tau_2} \int_{B_r} |\nabla \psi(x)|^p w_+^p(x, t) \eta^2(t) \, dx dt \\
& + C \int_{\tau_1 - \ell}^{\tau_2} \int_{B_r} \int_{B_r} (\max\{w_+(x, t), w_+(y, t)\})^p |\psi(x) - \psi(y)|^p \eta^2(t) \, d\mu dt \\
& + C \operatorname{ess\,sup}_{\substack{\tau_1 - \ell < t < \tau_2 \\ x \in \operatorname{supp} \psi}} \int_{\mathbb{R}^N \setminus B_r} \frac{w_+^{p-1}(y, t)}{|x - y|^{N+sp}} \, dy \int_{\tau_1 - \ell}^{\tau_2} \int_{B_r} w_+(x, t) \psi^p(x) \eta^2(t) \, dx dt \\
& + C \int_{\tau_1 - \ell}^{\tau_2} \int_{B_r} w_+^2(x, t) \psi^p(x) \eta(t) |\partial_t \eta(t)| \, dx dt, \tag{3.1}
\end{aligned}$$

where $w_+(x, t) := (u(x, t) - k)_+$ and $C > 0$ depends on N, p, s, Λ .

Proof. Taking $\varphi(x, t) := w_+(x, t) \psi^p(x) \eta^2(t) = (u(x, t) - k)_+ \psi^p(x) \eta^2(t)$ to test the weak formulation of (1.1). For $s \in [\tau_1, \tau_2]$, we can obtain

$$\begin{aligned}
0 & \geq \int_{\tau_1 - \ell}^s \int_{B_r} \partial_t u(x, t) w_+(x, t) \psi^p(x) \eta^2(t) \, dx dt \\
& + \int_{\tau_1 - \ell}^s \int_{B_r} |\nabla u(x, t)|^{p-2} \nabla u(x, t) \cdot \nabla (w_+(x, t) \psi^p(x) \eta^2(t)) \, dx dt \\
& + \int_{\tau_1 - \ell}^s \int_{B_r} \int_{B_r} J_p(w(x, t), w(y, t)) (w_+(x, t) \psi^p(x) \eta^2(t) - w_+(y, t) \psi^p(y) \eta^2(t)) \, d\mu dt \\
& + 2 \int_{\tau_1 - \ell}^s \int_{\mathbb{R}^N \setminus B_r} \int_{B_r} J_p(w(x, t), w(y, t)) w_+(x, t) \psi^p(x) \eta^2(t) \, d\mu dt \\
& =: I_1 + I_2 + I_3 + I_4.
\end{aligned}$$

Then we are going to estimate I_1, I_2, I_3 and I_4 . First, we evaluate

$$\begin{aligned}
I_1 & = \frac{1}{2} \int_{\tau_1 - \ell}^s \int_{B_r} \partial_t w_+^2(x, t) \psi^p(x) \eta^2(t) \, dx dt \\
& = \frac{1}{2} \int_{B_r} w_+^2(x, t) \psi^p(x) \eta^2(t) \, dx \Big|_{\tau_1 - \ell}^s - \int_{\tau_1 - \ell}^s \int_{B_r} w_+^2(x, t) \psi^p(x) \eta(t) \partial_t \eta(t) \, dx dt
\end{aligned}$$

$$= \frac{1}{2} \int_{B_r} w_+^2(x, s) \psi^p(x) dx - \int_{\tau_1 - \ell}^s \int_{B_r} w_+^2(x, t) \psi^p(x) \eta(t) \partial_t \eta(t) dx dt, \quad (3.2)$$

where in the last line we note that $\eta(\tau_1 - \ell) = 0$ and $\eta(s) = 1$ when $s \geq \tau_1$. We produce with estimating I_2 . By Young's inequality with ε , it yields that

$$\begin{aligned} I_2 &= \int_{\tau_1 - \ell}^s \int_{B_r} \psi^p \eta^2 |\nabla u|^{p-2} \nabla u \cdot \nabla w_+ + p \psi^{p-1} \eta^2 w_+ |\nabla u|^{p-2} \nabla u \cdot \nabla \psi dx dt \\ &\geq \int_{\tau_1 - \ell}^s \int_{B_r} |\nabla w_+|^p \psi^p \eta^2 dx dt - \int_{\tau_1 - \ell}^s \int_{B_r} p \psi^{p-1} \eta^2 w_+ |\nabla u|^{p-1} |\nabla \psi| dx dt \\ &\geq \int_{\tau_1 - \ell}^s \int_{B_r} |\nabla w_+|^p \psi^p \eta^2 dx dt - \varepsilon \int_{\tau_1 - \ell}^s \int_{B_r} \psi^p \eta^2 |\nabla w_+|^p dx dt \\ &\quad - C(p, \varepsilon) \int_{\tau_1 - \ell}^s \int_{B_r} |\nabla \psi|^p \eta^2 w_+^p dx dt \\ &= \frac{1}{2} \int_{\tau_1 - \ell}^s \int_{B_r} |\nabla w_+|^p \psi^p \eta^2 dx dt - C(p) \int_{\tau_1 - \ell}^s \int_{B_r} |\nabla \psi|^p \eta^2 w_+^p dx dt, \end{aligned} \quad (3.3)$$

with $\varepsilon = \frac{1}{2}$. As the same proof of Lemma 3.3 in [10], we have

$$\begin{aligned} I_3 &\geq \frac{1}{2^{p+1}} \int_{\tau_1 - \ell}^s \int_{B_r} \int_{B_r} |w_+(x, t) \psi(x) - w_+(y, t) \psi(y)|^p \eta^2(t) d\mu dt \\ &\quad - C \int_{\tau_1 - \ell}^s \int_{B_r} \int_{B_r} (\max\{w_+(x, t), w_+(y, t)\})^p |\psi(x) - \psi(y)|^p \eta^2(t) d\mu dt \end{aligned} \quad (3.4)$$

and

$$I_4 \geq -C \operatorname{ess\,sup}_{\substack{\tau_1 - \ell < t < s \\ x \in \operatorname{supp} \psi}} \int_{\mathbb{R}^N \setminus B_r} \frac{w_+^{p-1}(y, t)}{|x - y|^{N+sp}} dy \int_{\tau_1 - \ell}^s \int_{B_r} w_+(x, t) \psi^p(x) \eta^2(t) dx dt. \quad (3.5)$$

Merging the estimates on (3.2)–(3.5), we get

$$\begin{aligned} &\int_{\tau_1 - \ell}^s \int_{B_r} |\nabla w_+(x, t)|^p \psi^p(x) \eta^2(t) dx dt + \int_{B_r} w_+^2(x, s) \psi^p(x) dx \\ &\quad + \int_{\tau_1 - \ell}^s \int_{B_r} \int_{B_r} |w_+(x, t) \psi(x) - w_+(y, t) \psi(y)|^p \eta^2(t) d\mu dt \\ &\leq C \int_{\tau_1 - \ell}^{\tau_2} \int_{B_r} |\nabla \psi(x)|^p w_+^p(x, t) \eta^2(t) dx dt \\ &\quad + C \int_{\tau_1 - \ell}^{\tau_2} \int_{B_r} \int_{B_r} \max\{w_+(x, t), w_+(y, t)\}^p |\psi(x) - \psi(y)|^p \eta^2(t) d\mu dt \\ &\quad + C \operatorname{ess\,sup}_{\substack{\tau_1 - \ell < t < \tau_2 \\ x \in \operatorname{supp} \psi}} \int_{\mathbb{R}^N \setminus B_r} \frac{w_+^{p-1}(y, t)}{|x - y|^{N+sp}} dy \int_{\tau_1 - \ell}^{\tau_2} \int_{B_r} w_+(x, t) \psi^p(x) \eta^2(t) dx dt \\ &\quad + C \int_{\tau_1 - \ell}^{\tau_2} \int_{B_r} w_+^2(x, t) \psi^p(x) \eta(t) |\partial_t \eta(t)| dx dt, \end{aligned}$$

which leads to the desired result. \square

From the forthcoming lemma, one can interpret the reason why we only derive the Hölder continuity of weak solutions in the case $p > 2$. For the subquadratic case, we at present can not show a similar logarithmic estimate.

Lemma 3.2 (Logarithmic estimates). *Let $p > 2$ and u be a local solution to (1.1). Let $B_r \equiv B_r(x_0)$ and $(x_0, t_0) \in Q_T, T_0 > 0, 0 < r \leq R/2$. We also note that $\tilde{Q} \equiv B_R(x_0) \times (t_0 - 2T_0, t_0 + 2T_0)$ such that $\overline{B}_R(x_0) \subseteq \Omega$ and $[t_0 - 2T_0, t_0 + 2T_0] \subseteq (0, T)$. If $u \in L^\infty(\tilde{Q})$ and $u \geq 0$ in \tilde{Q} . For any $d > 0$, it holds that*

$$\begin{aligned} & \int_{t_0-T_0}^{t_0+T_0} \int_{B_r} |\nabla \log(u(x, t) + d)|^p dxdt + \int_{t_0-T_0}^{t_0+T_0} \int_{B_r} \int_{B_r} \left| \log \left(\frac{u(x, t) + d}{u(y, t) + d} \right) \right|^p d\mu dt \\ & \leq CT_0 r^N d^{1-p} R^{-p} [\text{Tail}_\infty(u; x_0, R, t_0 - 2T_0, t_0 + 2T_0)]^{p-1} \\ & \quad + Cr^N d^{2-p} + CT_0 r^{N-sp} + CT_0 r^{N-p}, \end{aligned} \quad (3.6)$$

where $C > 0$ depends on N, p, s and Λ .

Proof. Let $\psi \in C_0^\infty(B_{3r/2})$ and $\eta \in C_0^\infty(t_0 - 2T_0, t_0 + 2T_0)$ fulfill

$$0 \leq \psi \leq 1, \quad |\nabla \psi| < Cr^{-1} \text{ in } B_{2r}, \quad \psi \equiv 1 \text{ in } B_r,$$

and

$$0 \leq \eta \leq 1, \quad |\partial_t \eta| < CT_0^{-1} \text{ in } (t_0 - 2T_0, t_0 + 2T_0), \quad \eta \equiv 1 \text{ in } (t_0 - T_0, t_0 + T_0).$$

Choosing $\varphi(x, t) := (u(x, t) + d)^{1-p} \psi^p(x) \eta^2(t)$ to test the weak formulation of (1.1), we get

$$\begin{aligned} 0 &= - \int_{t_0-2T_0}^{t_0+2T_0} \int_{B_{2r}} \partial_t ((u(x, t) + d)^{1-p} \psi^p(x) \eta^2(t)) u(x, t) dxdt \\ & \quad + \int_{t_0-2T_0}^{t_0+2T_0} \int_{B_{2r}} |\nabla u|^{p-2} \nabla u \cdot \nabla ((u(x, t) + d)^{1-p} \psi^p(x) \eta^2(t)) dxdt \\ & \quad + \int_{t_0-2T_0}^{t_0+2T_0} \int_{B_{2r}} \int_{B_{2r}} J_p(u(x, t), u(y, t)) \left[\frac{\psi^p(x)}{(u(x, t) + d)^{p-1}} - \frac{\psi^p(y)}{(u(y, t) + d)^{p-1}} \right] \eta^2(t) d\mu dt \\ & \quad + 2 \int_{t_0-2T_0}^{t_0+2T_0} \int_{\mathbb{R}^N \setminus B_{2r}} \int_{B_{2r}} J_p(u(x, t), u(y, t)) \frac{\psi^p(x)}{(u(x, t) + d)^{p-1}} \eta^2(t) d\mu dt \\ & =: I_1 + I_2 + I_3 + I_4. \end{aligned}$$

It follows from the proof of [10, Lemma 3.5] that

$$\begin{aligned} I_1 &= \int_{t_0-2T_0}^{t_0+2T_0} \int_{B_{2r}} ((u(x, t) + d)^{1-p} \psi^p(x) \eta^2(t)) \partial_t u(x, t) dxdt \\ & \leq Cr^N d^{2-p}, \end{aligned} \quad (3.7)$$

$$I_3 \leq -C \int_{t_0-T_0}^{t_0+T_0} \int_{B_r} \int_{B_r} \left| \log \left(\frac{u(x, t) + d}{u(y, t) + d} \right) \right|^p dx dy dt + CT_0 r^{N-sp}, \quad (3.8)$$

and

$$I_4 \leq CT_0 r^{N-sp} + CT_0 r^N R^{-p} d^{1-p} [\text{Tail}_\infty(u; x_0, R, t_0 - 2T_0, t_0 + 2T_0)]^{p-1}. \quad (3.9)$$

By Young's inequality with ε , we estimate the integral I_2 as

$$I_2 = -(p-1) \int_{t_0-2T_0}^{t_0+2T_0} \int_{B_{2r}} (u(x, t) + d)^{-p} \psi^p(x) |\nabla u(x, t)|^p \eta^2(t) dxdt$$

$$\begin{aligned}
& + p \int_{t_0-2T_0}^{t_0+2T_0} \int_{B_{2r}} (u(x, t) + d)^{1-p} \psi^{p-1}(x) |\nabla u(x, t)|^{p-2} \nabla u(x, t) \cdot \nabla \psi(x) \eta^2(t) \, dx dt \\
& \leq -(p-1) \int_{t_0-2T_0}^{t_0+2T_0} \int_{B_{2r}} (u(x, t) + d)^{-p} \psi^p(x) |\nabla(u(x, t) + d)|^p \eta^2(t) \, dx dt \\
& \quad + \varepsilon \int_{t_0-2T_0}^{t_0+2T_0} \int_{B_{2r}} (u(x, t) + d)^{-p} \psi^p(x) |\nabla(u(x, t) + d)|^p \eta^2(t) \, dx dt \\
& \quad + C(\varepsilon) \int_{t_0-2T_0}^{t_0+2T_0} \int_{B_{2r}} |\nabla \psi(x)|^p \eta^2(t) \, dx dt \\
& \leq -C(p) \int_{t_0-2T_0}^{t_0+2T_0} \int_{B_r} (u(x, t) + d)^{-p} |\nabla(u(x, t) + d)|^p \eta^2(t) \, dx dt + C(N, p) T_0 r^{N-p} \\
& \leq -C(p) \int_{t_0-T_0}^{t_0+T_0} \int_{B_r} |\nabla \log(u(x, t) + d)|^p \, dx dt + C(N, p) T_0 r^{N-p} \tag{3.10}
\end{aligned}$$

with ε satisfying $\varepsilon < p - 1$. Combining with (3.7)–(3.10), we can obtain the Logarithmic estimates. \square

Next, we will give a corollary of Lemma 3.2, which plays a crucial role in obtaining the Hölder continuity.

Corollary 3.3. *Let $p > 2$ and u be a local solution to (1.1). Let $B_r \equiv B_r(x_0)$ and $(x_0, t_0) \in Q_T$, $T_0 > 0$, $0 < r \leq R/2$. We also note that $\tilde{Q} \equiv B_R(x_0) \times (t_0 - 2T_0, t_0 + 2T_0)$ such that $\overline{B_R}(x_0) \subseteq \Omega$ and $[t_0 - 2T_0, t_0 + 2T_0] \subseteq (0, T)$. Suppose that $u \in L^\infty(\tilde{Q})$ and $u \geq 0$ in \tilde{Q} . Let $a, d > 0$, $b > 1$ and define*

$$v := \min \{ (\log(a + d) - \log(u + d))_+, \log b \}.$$

Then it holds that

$$\begin{aligned}
& \int_{t_0-T_0}^{t_0+T_0} \int_{B_r} |v(x, t) - (v)_{B_r}(t)|^p \, dx dt \\
& \leq CT_0 d^{1-p} \left(\frac{r}{R} \right)^p [\text{Tail}_\infty(u; x_0, R, t_0 - 2T_0, t_0 + 2T_0)]^{p-1} \\
& \quad + CT_0 + Cd^{2-p} r^p + CT_0 r^{p-sp}, \tag{3.11}
\end{aligned}$$

where $C > 0$ depends on N, p, s and Λ .

Proof. By the Poincaré inequality from Theorem 2 in [15], it yields that

$$\int_{B_r} |v(x, t) - (v)_{B_r}(t)|^p \, dx \leq Cr^{p-N} \int_{B_r} |\nabla v(x, t)|^p \, dx$$

for any $t \in (t_0 - T_0, t_0 + T_0)$. Integrating the above inequality over $(t_0 - T_0, t_0 + T_0)$ leads to

$$\int_{t_0-T_0}^{t_0+T_0} \int_{B_r} |v(x, t) - (v)_{B_r}(t)|^p \, dx dt \leq Cr^{p-N} \int_{t_0-T_0}^{t_0+T_0} \int_{B_r} |\nabla v(x, t)|^p \, dx dt, \tag{3.12}$$

where $C > 0$ depends only on N, p . Observing that v is a truncation function of the sum of a constant and $\log(u + d)$, which gives that

$$\int_{t_0-T_0}^{t_0+T_0} \int_{B_r} |\nabla v(x, t)|^p \, dx dt \leq \int_{t_0-T_0}^{t_0+T_0} \int_{B_r} |\nabla \log(u(x, t) + d)|^p \, dx dt. \tag{3.13}$$

We can get the results from Lemma 3.2 along with (3.12) and (3.13). \square

4. LOCAL BOUNDEDNESS

In this part, we are ready to study the local boundedness of weak solutions. To this end, we first introduce some notations. For $\sigma \in [1/2, 1)$, set

$$r_0 := r, \quad r_j := \sigma r + 2^{-j}(1 - \sigma)r, \quad \tilde{r}_j := \frac{r_j + r_{j+1}}{2}, \quad j = 0, 1, 2, \dots$$

and

$$Q_j^- := B_j \times \Gamma_j := B_{r_j}(x_0) \times (t_0 - r_j^p, t_0), \quad j = 0, 1, 2, \dots,$$

$$\tilde{Q}_j^- := \tilde{B}_j \times \tilde{\Gamma}_j := B_{\tilde{r}_j}(x_0) \times (t_0 - \tilde{r}_j^p, t_0), \quad j = 0, 1, 2, \dots$$

Denote

$$k_j := (1 - 2^{-j})\tilde{k}, \quad \tilde{k}_j := \frac{k_{j+1} + k_j}{2}, \quad j = 0, 1, 2, \dots$$

with

$$\tilde{k} \geq \frac{\text{Tail}_\infty(u_+; x_0, \sigma r, t_0 - r^p, t_0)}{2}.$$

Let

$$w_j := (u - k_j)_+, \quad \tilde{w}_j := (u - \tilde{k}_j)_+, \quad j = 0, 1, 2, \dots$$

We now provide a Caccioppoli type inequality in a special cylinder, which leads to the recursive inequalities.

Lemma 4.1. *Let $p > 1$ and u be a subsolution to (1.1). Let $(x_0, t_0) \in Q_T$, $0 < r < 1$ and $Q_r^- = B_r(x_0) \times (t_0 - r^p, t_0)$ such that $\overline{B_r}(x_0) \subseteq \Omega$ and $[t_0 - r^p, t_0] \subseteq (0, T)$. Suppose q is a parameter satisfying $q \geq \max\{p, 2\}$, it holds that*

$$\begin{aligned} & \int_{\Gamma_{j+1}} \int_{B_{j+1}} |\nabla \tilde{w}_j(x, t)|^p dx dt + \text{ess sup}_{t \in \Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^2(x, t) dx \\ & + \int_{\Gamma_{j+1}} \int_{B_{j+1}} \int_{B_{j+1}} \frac{|\tilde{w}_j(x, t) - \tilde{w}_j(y, t)|^p}{|x - y|^{N+sp}} dx dy dt \\ & \leq \frac{C}{r^p} \left(\frac{1}{\sigma^p(1 - \sigma)^{N+sp}} + \frac{1}{(1 - \sigma)^p} \right) \cdot \left(\frac{2^{(p+q-2)j}}{\tilde{k}^{q-2}} + \frac{2^{(N+sp+q-1)j}}{\tilde{k}^{q-p}} \right) \int_{\Gamma_j} \int_{B_j} w_j^q(x, t) dx dt, \end{aligned} \quad (4.1)$$

where $C > 0$ only depends on N, p, s, Λ and q .

Proof. First we give a trivial but very useful inequality

$$\tilde{w}_j^\tau(x, t) \leq \frac{C 2^{(q-\tau)j}}{\tilde{k}^{q-\tau}} w_j^q(x, t) \quad \text{in } Q_T, \quad (4.2)$$

where $0 \leq \tau < q$. We take the cut-off functions $\psi_j \in C_0^\infty(\tilde{B}_j)$ and $\eta_j \in C_0^\infty(\tilde{\Gamma}_j)$ such that

$$0 \leq \psi_j \leq 1, \quad |\nabla \psi_j| \leq \frac{C 2^j}{(1 - \sigma)r} \text{ in } \tilde{B}_j, \quad \psi_j \equiv 1 \text{ in } B_{j+1}$$

and

$$0 \leq \eta_j \leq 1, \quad |\partial_t \eta_j| \leq \frac{C2^{pj}}{(1-\sigma)^p r^p} \text{ in } \tilde{\Gamma}_j, \quad \eta_j \equiv 1 \text{ in } \Gamma_{j+1}.$$

Let $r = r_j$, $\tau_2 = t_0$, $\tau_1 = t_0 - r_{j+1}^p$ and $\ell = \tilde{r}_j^p - r_{j+1}^p$ in Lemma 3.1. Then we arrive at

$$\begin{aligned} & \int_{\tilde{\Gamma}_j} \int_{B_j} |\nabla \tilde{w}_j(x, t)|^p \psi_j^p(x) \eta_j^2(t) \, dx dt + \operatorname{ess\,sup}_{t \in \Gamma_{j+1}} \int_{B_j} \tilde{w}_j^2(x, t) \psi_j^p(x) \, dx \\ & + \int_{\tilde{\Gamma}_j} \int_{B_j} \int_{B_j} |\tilde{w}_j(x, t) \psi_j(x) - \tilde{w}_j(y, t) \psi_j(y)|^p \eta_j^2(t) \, d\mu dt \\ & \leq C \int_{\tilde{\Gamma}_j} \int_{B_j} |\nabla \psi_j(x)|^p \tilde{w}_j^p(x, t) \eta_j^2(t) \, dx dt \\ & + C \int_{\tilde{\Gamma}_j} \int_{B_j} \int_{B_j} (\max\{\tilde{w}_j(x, t), \tilde{w}_j(y, t)\})^p |\psi_j(x) - \psi_j(y)|^p \eta_j^2(t) \, d\mu dt \\ & + C \operatorname{ess\,sup}_{\substack{t \in \tilde{\Gamma}_j \\ x \in \operatorname{supp} \psi_j}} \int_{\mathbb{R}^N \setminus B_j} \frac{\tilde{w}_j^{p-1}(y, t)}{|x-y|^{N+sp}} \, dy \int_{\tilde{\Gamma}_j} \int_{B_j} \tilde{w}_j(x, t) \psi_j^p(x) \eta_j^2(t) \, dx dt \\ & + C \int_{\tilde{\Gamma}_j} \int_{B_j} \tilde{w}_j^2(x, t) \psi_j^p(x) \eta_j(t) |\partial_t \eta_j(t)| \, dx dt \\ & =: I_1 + I_2 + I_3 + I_4. \end{aligned} \tag{4.3}$$

Using inequality (4.2) and the definition of ψ_j , we estimate I_1 as

$$\begin{aligned} I_1 & \leq \frac{C2^{pj}}{(1-\sigma)^p r^p} \int_{\Gamma_j} \int_{B_j} \tilde{w}_j^p(x, t) \, dx dt \\ & \leq \frac{C2^{qj}}{\tilde{k}^{q-p}(1-\sigma)^p r^p} \int_{\Gamma_j} \int_{B_j} w_j^q(x, t) \, dx dt. \end{aligned} \tag{4.4}$$

Analogous to the proof of Lemma 4.1 in [10], we have

$$I_2 \leq \frac{C2^{qj}}{\tilde{k}^{q-p}(1-\sigma)^p r^p} \int_{\Gamma_j} \int_{B_j} w_j^q(x, t) \, dx dt, \tag{4.5}$$

$$I_3 \leq \frac{C2^{(N+sp+q-1)j}}{\tilde{k}^{q-p}\sigma^p(1-\sigma)^{N+sp} r^p} \int_{\Gamma_j} \int_{B_j} w_j^q(x, t) \, dx dt \tag{4.6}$$

and

$$I_4 \leq \frac{C2^{(p+q-2)j}}{\tilde{k}^{q-2}(1-\sigma)^p r^p} \int_{\Gamma_j} \int_{B_j} w_j^q(x, t) \, dx dt. \tag{4.7}$$

By virtue of the facts that $\psi_j \equiv 1$ in B_{j+1} , $\eta_j \equiv 1$ in Γ_{j+1} and (1.3), merging inequalities (4.3)–(4.7), we get

$$\begin{aligned} & \int_{\Gamma_{j+1}} \int_{B_{j+1}} |\nabla \tilde{w}_j(x, t)|^p \, dx dt + \operatorname{ess\,sup}_{t \in \Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^2(x, t) \, dx \\ & + \int_{\Gamma_{j+1}} \int_{B_{j+1}} \int_{B_{j+1}} \frac{|\tilde{w}_j(x, t) - \tilde{w}_j(y, t)|^p}{|x-y|^{N+sp}} \, dx dy dt \end{aligned}$$

$$\begin{aligned} &\leq \frac{C}{r^p(1-\sigma)^p} \left(\frac{2^{qj}}{r^{(s-1)p}\tilde{k}^{q-p}} + \frac{2^{(N+sp+q-1)j}}{\sigma^p(1-\sigma)^{N+p(s-1)}\tilde{k}^{q-p}} + \frac{2^{qj}}{\tilde{k}^{q-p}} + \frac{2^{(p+q-2)j}}{\tilde{k}^{q-2}} \right) \\ &\quad \times \int_{\Gamma_j} \int_{B_j} w_j^q(x, t) \, dxdt. \end{aligned}$$

After rearrangement, we get the desired result. \square

The following two lemmas are the consequences of Lemmas 2.3 and 4.1.

Lemma 4.2. *Let $p > 2N/(N+2)$ and $\max\{p, 2\} \leq q < p(N+2)/N$. Let $(x_0, t_0) \in Q_T$, $0 < r < 1$ and $Q_r^- = B_r(x_0) \times (t_0 - r^p, t_0)$ such that $\overline{B_r}(x_0) \subseteq \Omega$ and $[t_0 - r^p, t_0] \subseteq (0, T)$. Then for a local subsolution u to (1.1), we infer that*

$$\begin{aligned} &\int_{\Gamma_{j+1}} \int_{B_{j+1}} w_{j+1}^q(x, t) \, dxdt \\ &\leq \frac{C2^{bj}}{r^{\frac{2p}{\kappa N}}} \left(\frac{1}{\sigma^{\frac{q(N+p)}{\kappa N}}(1-\sigma)^{\frac{q(N+p)(N+sp)}{p\kappa N}}} + \frac{1}{(1-\sigma)^{\frac{q(N+p)}{\kappa N}}} \right) \\ &\quad \times \left(\frac{1}{\tilde{k}^{\frac{q}{\kappa}(\frac{q}{N}+1-\frac{2}{p})}} + \frac{1}{\tilde{k}^{\frac{q}{\kappa}(\frac{q}{N}+\frac{2}{N}-\frac{p}{N})}} \right) \left(\int_{\Gamma_j} \int_{B_j} w_j^q(x, t) \, dxdt \right)^{1+\frac{q}{\kappa N}} \end{aligned} \quad (4.8)$$

for $j \in \mathbb{N}$, where $b := (1+p/N)(N+p+q)$, $\kappa := 1+2/N$ and $C > 0$ only depends on N, p, s, Λ and q .

Proof. Since $q < p\kappa$, it follows from Hölder inequality that

$$\begin{aligned} &\int_{\Gamma_{j+1}} \int_{B_{j+1}} w_{j+1}^q(x, t) \, dxdt \\ &\leq \int_{\Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^q(x, t) \, dxdt \\ &\leq \left(\int_{\Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^{p\kappa}(x, t) \, dxdt \right)^{\frac{q}{p\kappa}} \left(\int_{\Gamma_{j+1}} \int_{B_{j+1}} \chi_{\{u \geq \tilde{k}_j\}}(x, t) \, dxdt \right)^{1-\frac{q}{p\kappa}}. \end{aligned} \quad (4.9)$$

From (4.2), we can get

$$\int_{\Gamma_{j+1}} \int_{B_{j+1}} \chi_{\{u \geq \tilde{k}_j\}}(x, t) \, dxdt \leq \frac{C2^{qj}}{\tilde{k}^{q-p}} \int_{\Gamma_j} \int_{B_j} w_j^q(x, t) \, dxdt \quad (4.10)$$

and

$$\int_{\Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^p(x, t) \, dxdt \leq \frac{C2^{(q-p)j}}{\tilde{k}^{q-p}} \int_{\Gamma_j} \int_{B_j} w_j^q(x, t) \, dxdt. \quad (4.11)$$

Now by (4.11), Lemma 2.3 and Lemma 4.1 we estimate

$$\begin{aligned} &\int_{\Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^{p\kappa}(x, t) \, dxdt \\ &\leq C \left(r^p \int_{\Gamma_{j+1}} \int_{B_{j+1}} |\nabla \tilde{w}_j(x, t)|^p \, dxdt + \int_{\Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^p(x, t) \, dxdt \right) \end{aligned}$$

$$\begin{aligned}
& \times \left(\operatorname{ess\,sup}_{t \in \Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^2(x, t) \, dx \right)^{\frac{p}{N}} \\
& \leq Cr^{-\frac{p^2}{N}} \left(\frac{1}{\sigma^p(1-\sigma)^{N+sp}} + \frac{1}{(1-\sigma)^p} \right)^{\frac{p}{N}} \left(\frac{2^{(p+q-2)j}}{\tilde{k}^{q-2}} + \frac{2^{(N+sp+q-1)j}}{\tilde{k}^{q-p}} \right)^{\frac{p}{N}} \\
& \quad \times \left[\left(\frac{1}{\sigma^p(1-\sigma)^{N+sp}} + \frac{1}{(1-\sigma)^p} \right) \left(\frac{2^{(p+q-2)j}}{\tilde{k}^{q-2}} + \frac{2^{(N+sp+q-1)j}}{\tilde{k}^{q-p}} \right) + \frac{2^{(q-p)j}}{\tilde{k}^{q-p}} \right] \\
& \quad \times \left(\int_{\Gamma_j} \int_{B_j} w_j^q(x, t) \, dx dt \right)^{1+\frac{p}{N}} \\
& \leq \frac{C2^{bj}}{r^{\frac{p^2}{N}}} \left(\frac{1}{\sigma^{\frac{p(N+p)}{N}}(1-\sigma)^{\frac{(N+p)(N+sp)}{N}}} + \frac{1}{(1-\sigma)^{\frac{p(N+p)}{N}}} \right) \\
& \quad \times \left(\frac{1}{\tilde{k}^{q-2}} + \frac{1}{\tilde{k}^{q-p}} \right)^{1+\frac{p}{N}} \left(\int_{\Gamma_j} \int_{B_j} w_j^q(x, t) \, dx dt \right)^{1+\frac{p}{N}} \tag{4.12}
\end{aligned}$$

with $b = (1 + p/N)(N + p + q)$. Combining (4.9), (4.10) and (4.12), we get the desired result. \square

Lemma 4.3. *Let $1 < p \leq 2N/(N + 2)$ and $m > \max\left\{2, \frac{N(2-p)}{p}\right\}$. Let $(x_0, t_0) \in Q_T$, $0 < r < 1$ and $Q_r^- = B_r(x_0) \times (t_0 - r^p, t_0)$ such that $\overline{B_r}(x_0) \subseteq \Omega$ and $[t_0 - r^p, t_0] \subseteq (0, T)$. Suppose that $u \in L_{\text{loc}}^\infty(Q_T)$ is a weak subsolution to (1.1). Then for any $j \in \mathbb{N}$, we have*

$$\begin{aligned}
& \int_{\Gamma_{j+1}} \int_{B_{j+1}} w_{j+1}^m(x, t) \, dx dt \\
& \leq \frac{C2^{bj}}{r^{\frac{p^2}{N}}} \left(\frac{1}{\sigma^{\frac{p(N+p)}{N}}(1-\sigma)^{\frac{(N+p)(N+sp)}{N}}} + \frac{1}{(1-\sigma)^{\frac{p(N+p)}{N}}} \right) \left(\frac{1}{\tilde{k}^{m-p}} + \frac{1}{\tilde{k}^{m-2}} \right)^{1+\frac{p}{N}} \\
& \quad \times \|\tilde{w}_j\|_{L^\infty(Q_{j+1}^-)}^{m-p\kappa} \left(\int_{\Gamma_j} \int_{B_j} w_j^m(x, t) \, dx dt \right)^{1+\frac{p}{N}}, \tag{4.13}
\end{aligned}$$

where $b := (1 + p/N)(N + p + m)$, $\kappa := 1 + 2/N$ and $C > 0$ only depends on N, p, s, m and Λ .

Proof. Based on the assumptions, we have

$$\begin{aligned}
\int_{\Gamma_{j+1}} \int_{B_{j+1}} w_{j+1}^m(x, t) \, dx dt & \leq \int_{\Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^m(x, t) \, dx dt \\
& \leq \|\tilde{w}_j\|_{L^\infty(Q_{j+1}^-)}^{m-p\kappa} \int_{\Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^{p\kappa}(x, t) \, dx dt. \tag{4.14}
\end{aligned}$$

By utilizing Lemma 2.3, Lemma 4.1 with $q = m$ and inequality (4.11) with $q = m$, it yields that

$$\begin{aligned}
& \int_{\Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^{p\kappa}(x, t) \, dx dt \\
& \leq C \left(r^p \int_{\Gamma_{j+1}} \int_{B_{j+1}} |\nabla \tilde{w}_j(x, t)|^p \, dx dt + \int_{\Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^p(x, t) \, dx dt \right)
\end{aligned}$$

$$\begin{aligned}
& \times \left(\operatorname{ess\,sup}_{t \in \Gamma_{j+1}} \int_{B_{j+1}} \tilde{w}_j^2(x, t) \, dx \right)^{\frac{p}{N}} \\
& \leq \frac{C2^{bj}}{r^{\frac{p^2}{N}}} \left[\frac{1}{\sigma^{\frac{p(N+p)}{N}} (1-\sigma)^{\frac{(N+p)(N+sp)}{N}}} + \frac{1}{(1-\sigma)^{\frac{p(N+p)}{N}}} \right] \left(\frac{1}{\tilde{k}^{m-p}} + \frac{1}{\tilde{k}^{m-2}} \right)^{1+\frac{p}{N}} \\
& \quad \times \left(\int_{\Gamma_j} \int_{B_j} w_j^m(x, t) \, dx dt \right)^{1+\frac{p}{N}} \tag{4.15}
\end{aligned}$$

with $b := (1 + p/N)(N + p + m)$. Thus combining (4.14) and (4.15), we get the desired inequality (4.13). \square

Remark 4.4. In Lemma 4.3, the quantity $\sigma^{\frac{p(N+p)}{N}}$ can be removed, since $\sigma \in [1/2, 1)$. In addition,

$$\max \left\{ (1-\sigma)^{\frac{(N+p)(N+sp)}{N}}, (1-\sigma)^{\frac{p(N+p)}{N}} \right\} \geq (1-\sigma)^{\frac{(N+p)^2}{N}}.$$

Thus, we can get

$$\begin{aligned}
& \int_{\Gamma_{j+1}} \int_{B_{j+1}} w_{j+1}^m(x, t) \, dx dt \\
& \leq \frac{C2^{bj}}{r^{\frac{p^2}{N}}} \cdot \frac{1}{(1-\sigma)^{\frac{(N+p)^2}{N}}} \left(\frac{1}{\tilde{k}^{m-p}} + \frac{1}{\tilde{k}^{m-2}} \right)^{1+\frac{p}{N}} \\
& \quad \times \|\tilde{w}_j\|_{L^\infty(Q_{j+1}^-)}^{m-p\kappa} \left(\int_{\Gamma_j} \int_{B_j} w_j^m(x, t) \, dx dt \right)^{1+\frac{p}{N}}.
\end{aligned}$$

Next, we introduce an analysis lemma which will be used later.

Lemma 4.5. ([8, Lemma 4.1]) Let $\{Y_j\}_{j=0}^\infty$ be a sequence of positive numbers such that

$$Y_0 \leq K^{-\frac{1}{\delta}} b^{-\frac{1}{\delta^2}} \quad \text{and} \quad Y_{j+1} \leq Kb^j Y_j^{1+\delta}, \quad j = 0, 1, 2, \dots$$

for some constants $K, b > 1$ and $\delta > 0$. Then we have $\lim_{j \rightarrow \infty} Y_j = 0$.

Finally, we end this section by proving the results of local boundedness.

Proof of Theorem 1.2. Let $r = R$, $\sigma = \frac{1}{2}$, then $r_j = \frac{R}{2} + 2^{-j-1}R$. Fix $q = \max\{2, p\}$. We denote

$$Y_j = \int_{\Gamma_j} \int_{B_j} (u - k_j)_+^q \, dx dt, \quad j = 0, 1, 2, \dots$$

Supposing $\tilde{k} \geq 1$ and recalling $r < 1$, we derive from Lemma 4.2 that

$$\begin{aligned}
\frac{Y_{j+1}}{r^p} & \leq \frac{C2^{bj} Y_j^{1+\frac{q}{N\kappa}}}{r^{p(1+\frac{q}{N\kappa})} \tilde{k}^{\frac{q}{\kappa}(\frac{q}{N} + \frac{2}{N} - \frac{p}{N})}} + \frac{C2^{bj} Y_j^{1+\frac{q}{N\kappa}}}{r^{p(1+\frac{q}{N\kappa})} \tilde{k}^{\frac{q}{\kappa}(\frac{q}{N} + 1 - \frac{2}{p})}} \\
& \leq \frac{C2^{bj}}{\tilde{k}^{q(1-\frac{q}{p\kappa})}} \left(\frac{Y_j}{r^p} \right)^{1+\frac{q}{N\kappa}}, \tag{4.16}
\end{aligned}$$

where $b := (1 + p/N)(N + p + q)$, $\kappa := 1 + 2/N$ and $C > 0$ only depends on N, p, s, Λ . For any $j \in \mathbb{N}$, define $W_j = Y_j/r^p$. Thus we get

$$W_{j+1} \leq \frac{C2^{bj}}{\tilde{k}^q(1-\frac{q}{p\kappa})} W_j^{1+\frac{q}{N\kappa}},$$

where \tilde{k} is such that

$$\tilde{k} \geq \max \left\{ \text{Tail}_\infty(u_+; x_0, R/2, t_0 - R^p, t_0), C \left(\int_{t_0 - R^p}^{t_0} \int_{B_R} u_+^q dx dt \right)^{\frac{p}{N(p\kappa - q)}} \vee 1 \right\}$$

with C only depending on N, p, s and Λ . Thus along with Lemma 4.5 we can deduce that $\lim_{j \rightarrow \infty} W_j = 0$. Consequently, we get the desired result

$$\text{ess sup}_{Q_{R/2}^-} u \leq \text{Tail}_\infty(u_+; x_0, R/2, t_0 - R^p, t_0) + C \left(\int_{Q_R^-} u_+^q dx dt \right)^{\frac{p}{N(p\kappa - q)}} \vee 1.$$

We now finish the proof. \square

Proof of Theorem 1.3. Let $R_0 = R/2$ and $R_n = R/2 + \sum_{i=1}^n 2^{-i-1}R$ for $n \in \mathbb{N}^+$, and $Q_n^- = B_{R_n}(x_0) \times (t_0 - R_n^p, t_0)$. Set

$$M_n = \text{ess sup}_{Q_n^-} u_+, \quad n = 0, 1, 2, 3, \dots$$

Choosing $r = R_{n+1}$ and $\sigma r = R_n$, then

$$\sigma = \frac{1/2 + \sum_{i=1}^n 2^{-i-1}}{1/2 + \sum_{i=1}^{n+1} 2^{-i-1}} \geq \frac{1}{2}.$$

We denote

$$Y_j = \int_{\Gamma_j} \int_{B_j} (u - k_j)_+^m dx dt, \quad j = 0, 1, 2, \dots$$

Due to Lemma 4.3, we obtain

$$\begin{aligned} Y_{j+1} &\leq \frac{C2^{bj}}{R_{n+1}^{\frac{p^2}{N}}} \|u_+\|_{L^\infty(Q_{n+1}^-)}^{m-p\kappa} \left(\frac{1}{(1-\sigma)^{\frac{(N+p)^2}{N}}} + \frac{1}{(1-\sigma)^{\frac{p(N+p)}{N}}} \right) \\ &\quad \times \left(\frac{1}{\tilde{k}^{m-p}} + \frac{1}{\tilde{k}^{m-2}} \right)^{1+\frac{p}{N}} Y_j^{1+\frac{p}{N}} \\ &\leq \frac{C2^{bj+dn}}{R_{n+1}^{\frac{p^2}{N}}} M_{n+1}^{m-p\kappa} \left(\frac{1}{\tilde{k}^{m-p}} + \frac{1}{\tilde{k}^{m-2}} \right)^{1+\frac{p}{N}} Y_j^{1+\frac{p}{N}} \end{aligned} \quad (4.17)$$

with $b := (1 + p/N)(N + p + m)$ and $d = (N + p)^2/N$. For any $j \in \mathbb{N}$, define $W_j = Y_j/R_n^p$. Then we get

$$W_{j+1} \leq C2^{bj+dn} M_{n+1}^{m-p\kappa} \left(\frac{1}{\tilde{k}^{m-p}} + \frac{1}{\tilde{k}^{m-2}} \right)^{1+\frac{p}{N}} W_j^{1+\frac{p}{N}}.$$

According to Lemma 4.5, we can see that

$$\lim_{j \rightarrow \infty} Y_j = 0$$

if

$$W_0 \leq C 2^{-\frac{dnN}{p} - \frac{bN^2}{p^2}} M_{n+1}^{-\frac{N(m-p\kappa)}{p}} \left(\frac{1}{\tilde{k}^{m-p}} + \frac{1}{\tilde{k}^{m-2}} \right)^{-\frac{p+N}{p}}. \quad (4.18)$$

To ensure the above inequality, we need choose \tilde{k} properly large. Indeed,

$$\begin{aligned} W_0 &= \frac{Y_0}{R_n^p} = R_n^{-p} \int_{t_0 - R_{n+1}^p}^{t_0} \int_{B_{R_{n+1}}} u_+^m dx dt \\ &\leq 2^p \int_{t_0 - R_{n+1}^p}^{t_0} \int_{B_{R_{n+1}}} u_+^m dx dt. \end{aligned}$$

Namely,

$$2^p \int_{t_0 - R_{n+1}^p}^{t_0} \int_{B_{R_{n+1}}} u_+^m dx dt \leq C 2^{-\frac{dnN}{p} - \frac{bN^2}{p^2}} M_{n+1}^{-\frac{N(m-p\kappa)}{p}} \left(\frac{1}{\tilde{k}^{m-p}} + \frac{1}{\tilde{k}^{m-2}} \right)^{-\frac{p+N}{p}},$$

which implies that

$$C 2^{\frac{dnN}{N+p}} M_{n+1}^{\frac{N(m-p\kappa)}{N+p}} \left(\int_{t_0 - R_{n+1}^p}^{t_0} \int_{B_{R_{n+1}}} u_+^m dx dt \right)^{\frac{p}{N+p}} \leq \left(\frac{1}{\tilde{k}^{m-p}} + \frac{1}{\tilde{k}^{m-2}} \right)^{-1}.$$

Therefore, we take

$$\begin{aligned} \tilde{k} &= C 2^{\frac{dnN}{(N+p)(m-p)}} M_{n+1}^{\frac{N(m-p\kappa)}{(N+p)(m-p)}} \left(\int_{t_0 - R_{n+1}^p}^{t_0} \int_{B_{R_{n+1}}} u_+^m dx dt \right)^{\frac{p}{(N+p)(m-p)}} \\ &\quad + C 2^{\frac{dnN}{(N+p)(m-2)}} M_{n+1}^{\frac{N(m-p\kappa)}{(N+p)(m-2)}} \left(\int_{t_0 - R_{n+1}^p}^{t_0} \int_{B_{R_{n+1}}} u_+^m dx dt \right)^{\frac{p}{(N+p)(m-2)}} \\ &\quad + \frac{\text{Tail}_\infty(u_+; x_0, R_n, t_0 - R_{n+1}^p, t_0)}{2}, \end{aligned}$$

which makes inequality (4.18) hold true. Here the constant C only depends on N, p, s, m and Λ . Under this choice, it follows from Lemma 4.5 that

$$\begin{aligned} M_n = \text{ess sup}_{Q_{R_n}^-} u^+ &\leq C 2^{\frac{dnN}{(N+p)(m-p)}} M_{n+1}^{\frac{N(m-p\kappa)}{(N+p)(m-p)}} \left(\int_{t_0 - R_{n+1}^p}^{t_0} \int_{B_{R_{n+1}}} u_+^m dx dt \right)^{\frac{p}{(N+p)(m-p)}} \\ &\quad + C 2^{\frac{dnN}{(N+p)(m-2)}} M_{n+1}^{\frac{N(m-p\kappa)}{(N+p)(m-2)}} \left(\int_{t_0 - R_{n+1}^p}^{t_0} \int_{B_{R_{n+1}}} u_+^m dx dt \right)^{\frac{p}{(N+p)(m-2)}} \\ &\quad + \frac{\text{Tail}_\infty(u_+; x_0, R_n, t_0 - R_{n+1}^p, t_0)}{2}. \end{aligned} \quad (4.19)$$

Obverse $m > \max\{2, \frac{N(2-p)}{p}\}$ and $\kappa = 1 + 2/N$, which indicates that

$$0 < \frac{N(m-p\kappa)}{(N+p)(m-p)}, \frac{N(m-p\kappa)}{(N+p)(m-2)} < 1.$$

Now we apply the Young's inequality with ε to (4.19), arriving at

$$M_n \leq \varepsilon M_{n+1} + C 2^{\frac{dnN}{(N+p)(m-p-\beta)}} \varepsilon^{-\frac{\beta}{m-p-\beta}} \left(\int_{t_0 - R^p}^{t_0} \int_{B_R} u_+^m dx dt \right)^{\frac{p}{(N+p)(m-p-\beta)}}$$

$$\begin{aligned}
& + C 2^{\frac{d_n N}{(N+p)(m-2-\beta)}} \varepsilon^{-\frac{\beta}{m-2-\beta}} \left(\int_{t_0-R^p}^{t_0} \int_{B_R} u_+^m dx dt \right)^{\frac{p}{(N+p)(m-2-\beta)}} \\
& + \frac{\text{Tail}_\infty(u_+; x_0, R/2, t_0 - R^p, t_0)}{2}
\end{aligned}$$

with $\beta = (m - p\kappa)N/(p + N)$, where we used the fact $R/2 \leq R_n < R$. Via the induction argument, we can derive

$$\begin{aligned}
M_0 & \leq \varepsilon^{n+1} M_{n+1} \\
& + C \varepsilon^{-\frac{\beta}{m-p-\beta}} \left(\int_{t_0-R^p}^{t_0} \int_{B_R} u_+^m dx dt \right)^{\frac{p}{(N+p)(m-p-\beta)}} \sum_{i=0}^n \left(2 \frac{dN}{(N+p)(m-p-\beta)} \varepsilon \right)^i \\
& + C \varepsilon^{-\frac{\beta}{m-2-\beta}} \left(\int_{t_0-R^p}^{t_0} \int_{B_R} u_+^m dx dt \right)^{\frac{p}{(N+p)(m-2-\beta)}} \sum_{i=0}^n \left(2 \frac{dN}{(N+p)(m-2-\beta)} \varepsilon \right)^i \\
& + \frac{\text{Tail}_\infty(u_+; x_0, R/2, t_0 - R^p, t_0)}{2} \sum_{i=0}^n \varepsilon^i, \quad n = 0, 1, 2, \dots
\end{aligned}$$

It is easy to see that the sum on the right-hand side could be revised by a convergent series, provided that we take

$$\varepsilon = 2^{-[\frac{dN}{(N+p)(m-2-\beta)} + 1]}.$$

Finally, letting $n \rightarrow \infty$, we deduce that

$$\begin{aligned}
\text{ess sup}_{Q_{R/2}^-} u & \leq \text{Tail}_\infty(u_+; x_0, R/2, t_0 - R^p, t_0) \\
& + C \left(\int_{Q_R^-} u_+^m dx dt \right)^{\frac{p}{(N+p)(m-2-\beta)}} \vee \left(\int_{Q_R^-} u_+^m dx dt \right)^{\frac{p}{(N+p)(m-p-\beta)}},
\end{aligned}$$

where $C > 0$ depends only on N, p, s, m and Λ . \square

5. LOCAL HÖLDER CONTINUITY

In this section we aim at establishing the Hölder continuity of weak solutions to (1.1) in the case that $p > 2$, based on the local boundedness results. Before verifying this conclusion, we introduce some notations.

Let

$$0 < \alpha < \frac{p}{p-1}$$

be a constant to be determined later. Set

$$r_j := \frac{\sigma^j r}{2}, \quad \omega(r_0) = \omega(r/2) := M, \quad \omega(r_j) := \left(\frac{r_j}{r_0} \right)^\alpha \omega(r_0), \quad j = 0, 1, 2, 3 \dots \quad (5.1)$$

and

$$M := C \left[\text{Tail}_\infty(u; \bar{x}_0, r/2, \bar{t}_0 - r^p, \bar{t}_0 + r^p) + \left(\int_{Q_r} |u|^p dx dt \right)^{\frac{1}{2}} \vee 1 \right] \quad (5.2)$$

with C depending on N, p, s, Λ . Define

$$d_j := \begin{cases} [\varepsilon \sigma^{(j-1)\alpha} M]^{2-p} & \text{if } j \geq 1, \\ 1 & \text{if } j = 0, \end{cases}$$

where

$$\varepsilon = \sigma^{\frac{p}{p-1}-\alpha}.$$

Thus it is easy to obtain

$$\frac{1}{d_{j+1}} = [\varepsilon \omega(r_j)]^{p-2} \text{ for all } j \geq 0. \quad (5.3)$$

Denote

$$B_j := B_{r_j}(\bar{x}_0) \text{ and } t_j := d_j r_j^p$$

and

$$Q_j := Q_{r_j, t_j}(\bar{x}_0, \bar{t}_0) = B_j \times (\bar{t}_0 - t_j, \bar{t}_0 + t_j).$$

Hence, for $j \geq 1$, we have

$$4 \left(\sigma^{\frac{p}{p-1}-\alpha} \right)^{2-p} r_1^p \leq r_0^p \quad \text{and} \quad 4 \sigma^{\alpha(2-p)} r_{j+1}^p \leq r_j^p.$$

The above inequalities combine with the definitions of d_j and t_j gives that

$$4t_{j+1} \leq t_j \text{ for all } j \geq 0. \quad (5.4)$$

Now we are going to deduce an oscillation reduction on weak solutions.

Lemma 5.1. *Let $p > 2$ and u be a local solution to (1.1). Let $(\bar{x}_0, \bar{t}_0) \in Q_T$, $r \in (0, R]$ for some $R \in (0, 1)$ and $Q_R \equiv B(\bar{x}_0) \times (\bar{t}_0 - R^p, \bar{t}_0 + R^p)$ such that $\bar{Q}_R \subseteq Q_T$. Then*

$$\text{ess osc}_{Q_j} u \leq \omega(r_j) \text{ for all } j = 0, 1, 2, \dots \quad (5.5)$$

Proof. We prove this Lemma by induction argument. It follows from Theorem 1.2 and the definition of $\omega(r_0)$ that (5.5) holds true for $j = 0$. We may assume that (5.5) is valid for $i \in \{0, \dots, j\}$ with some $j \geq 0$. Then we devoted to proving it holds for $i = j + 1$. It is obvious that either

$$\frac{|2Q_{j+1} \cap \{u \geq \text{ess inf}_{Q_j} u + \omega(r_j)/2\}|}{|2Q_{j+1}|} \geq \frac{1}{2} \quad (5.6)$$

or

$$\frac{|2Q_{j+1} \cap \{u \leq \text{ess inf}_{Q_j} u + \omega(r_j)/2\}|}{|2Q_{j+1}|} \geq \frac{1}{2} \quad (5.7)$$

must hold. In the case of (5.6), we set $u_j := u - \text{ess inf}_{Q_j} u$. In the case of (5.7), we set $u_j := \omega(r_j) - (u - \text{ess inf}_{Q_j} u)$. In all cases, we have

$$\frac{|2Q_{j+1} \cap \{u_j \geq \omega(r_j)/2\}|}{|2Q_{j+1}|} \geq \frac{1}{2} \quad (5.8)$$

and

$$0 \leq \text{ess inf}_{Q_i} u_j \leq \text{ess sup}_{Q_i} u_j \leq 2\omega(r_i) \text{ for } i = 0, \dots, j. \quad (5.9)$$

Now we provide an important estimate to be used later,

$$[\text{Tail}_\infty (u_j; \bar{x}_0, r_j, \bar{t}_0 - t_j, \bar{t}_0 + t_j)]^{p-1} \leq C \sigma^{-\alpha(p-1)} [\omega(r_j)]^{p-1} \text{ for } j = 0, 1, 2, \dots, \quad (5.10)$$

where C only depends on N, p, s , the difference of $p/(p-1)$ and α . Indeed, it is easy to see the claim is true when $j = 0$. For $j \geq 1$, we have

$$\begin{aligned} & [\text{Tail}_\infty (u_j; \bar{x}_0, r_j, \bar{t}_0 - t_j, \bar{t}_0 + t_j)]^{p-1} \\ &= r_j^p \text{ess sup}_{t \in (\bar{t}_0 - t_j, \bar{t}_0 + t_j)} \sum_{i=1}^j \int_{B_{i-1} \setminus B_i} \frac{|u_j(x, t)|^{p-1}}{|x - \bar{x}_0|^{N+sp}} dx \\ & \quad + r_j^p \text{ess sup}_{t \in (\bar{t}_0 - t_j, \bar{t}_0 + t_j)} \int_{\mathbb{R}^N \setminus B_0} \frac{|u_j(x, t)|^{p-1}}{|x - \bar{x}_0|^{N+sp}} dx \\ &\leq r_j^p \sum_{i=1}^j \left(\text{ess sup}_{Q_{i-1}} u_j \right)^{p-1} \int_{\mathbb{R}^N \setminus B_i} \frac{1}{|x - \bar{x}_0|^{N+sp}} dx \\ & \quad + r_j^p \text{ess sup}_{t \in (\bar{t}_0 - t_j, \bar{t}_0 + t_j)} \int_{\mathbb{R}^N \setminus B_0} \frac{|u_j(x, t)|^{p-1}}{|x - \bar{x}_0|^{N+sp}} dx \\ &\leq C \sum_{i=1}^j \left(\frac{r_j}{r_i} \right)^p [\omega(r_{i-1})]^{p-1}, \end{aligned}$$

where in the last line we used (5.9) and the definition of u_j . Since $\sigma \leq 1/4$ and $\alpha < p/(p-1)$, we estimate the right hand side as

$$\begin{aligned} & \sum_{i=1}^j \left(\frac{r_j}{r_i} \right)^p [\omega(r_{i-1})]^{p-1} \\ &= [\omega(r_0)]^{p-1} \left(\frac{r_j}{r_0} \right)^{\alpha(p-1)} \sum_{i=1}^j \left(\frac{r_{i-1}}{r_i} \right)^{\alpha(p-1)} \left(\frac{r_j}{r_i} \right)^{p-\alpha(p-1)} \\ &= [\omega(r_j)]^{p-1} \sigma^{-\alpha(p-1)} \sum_{i=0}^{j-1} \sigma^{i(p-\alpha(p-1))} \\ &\leq [\omega(r_j)]^{p-1} \frac{\sigma^{-\alpha(p-1)}}{1 - \sigma^{p-\alpha(p-1)}} \\ &\leq \frac{4^{p-\alpha(p-1)}}{(p-\alpha(p-1)) \log 4} \sigma^{-\alpha(p-1)} [\omega(r_j)]^{p-1}. \end{aligned}$$

Thus, we have proved (5.10) with the constant C depending on N, p, s , the difference of $p/(p-1)$ and α . Next, we define

$$v := \min \left\{ \left[\log \left(\frac{\omega(r_j)/2 + d}{u_j + d} \right) \right]_+, k \right\} \text{ for } k > 0. \quad (5.11)$$

It follows from Corollary 3.3 with $a \equiv \omega(r_j)/2$ and $b \equiv \exp(k)$ that

$$\int_{\bar{t}_0 - 2t_{j+1}}^{\bar{t}_0 + 2t_{j+1}} \int_{2B_{j+1}} |v(x, t) - (v)_{2B_{j+1}}(t)|^p dx dt$$

$$\begin{aligned}
&\leq Ct_{j+1}d^{1-p} \left(\frac{r_{j+1}}{r_j} \right)^p [\text{Tail}_\infty(u_j; \bar{x}_0, r_j, \bar{t}_0 - 4t_{j+1}, \bar{t}_0 + 4t_{j+1})]^{p-1} \\
&\quad + Ct_{j+1} + Cd^{2-p}r_{j+1}^p + Ct_{j+1}r_{j+1}^{p-sp} \\
&\leq Ct_{j+1}d^{1-p} [\varepsilon\omega(r_j)]^{p-1} + Ct_{j+1} + Cd^{2-p}r_{j+1}^p,
\end{aligned}$$

where in the last line we used the fact $4t_{j+1} \leq t_j$ and (5.10). Choosing $d = \varepsilon\omega(r_j)$ in (5.11), we get

$$d^{2-p} = d_{j+1}.$$

Since $\alpha < p/(p-1)$, we can verify

$$d^{1-p} \leq r_{j+1}^{-p}.$$

Thus, we arrive at

$$\int_{\bar{t}_0 - 2t_{j+1}}^{\bar{t}_0 + 2t_{j+1}} \int_{2B_{j+1}} |v(x, t) - (v)_{2B_{j+1}}(t)|^p dxdt \leq Ct_{j+1},$$

where C depends on N, p, s, Λ , the difference of $p/(p-1)$ and α . Following the calculation in Pages 38-39 in [10], there holds that

$$\frac{|2Q_{j+1} \cap \{u_j \leq 2\varepsilon\omega(r_j)\}|}{|2Q_{j+1}|} \leq \frac{\bar{C}}{\log\left(\frac{1}{\sigma}\right)}, \quad (5.12)$$

where $\bar{C} > 0$ depends on N, p, s, Λ , the difference of $p/(p-1)$ and α .

In what follows, we will proceed by a suitable iteration to infer the desired oscillation decay over the domain Q_{j+1} . For any $i = 0, 1, 2, \dots$, we define

$$\begin{aligned}
\varrho_i &:= r_{j+1} + 2^{-i}r_{j+1}, \quad \tilde{\varrho}_i := \frac{\varrho_i + \varrho_{i+1}}{2}, \\
\theta_i &:= t_{j+1} + 2^{-i}t_{j+1}, \quad \tilde{\theta}_i := \frac{\theta_i + \theta_{i+1}}{2}, \\
Q^i &:= B^i \times \Gamma_i := B_{\varrho_i}(\bar{x}_0) \times (\bar{t}_0 - \theta_i, \bar{t}_0 + \theta_i), \\
\tilde{Q}^i &:= \tilde{B}^i \times \tilde{\varrho}_i := B_{\tilde{\varrho}_i}(\bar{x}_0) \times (\bar{t}_0 - \tilde{\theta}_i, \bar{t}_0 + \tilde{\theta}_i).
\end{aligned}$$

Then we take the cut-off function $\psi_i \in C_0^\infty(\tilde{B}^i)$ and $\eta_i \in C_0^\infty(\tilde{\Gamma}_i)$ such that

$$0 \leq \psi_i \leq 1, \quad |\nabla\psi_i| \leq C2^i r_{j+1}^{-1} \text{ in } \tilde{B}^i, \quad \psi_i \equiv 1 \text{ in } B^{i+1}$$

and

$$0 \leq \eta_i \leq 1, \quad |\partial_t \eta_i| \leq C2^i t_{j+1}^{-1} \text{ in } \tilde{\Gamma}_i, \quad \eta_i \equiv 1 \text{ in } \Gamma_{i+1}.$$

Define

$$k_i := (1 + 2^{-i})\varepsilon\omega(r_j), \quad v_i := (k_i - u_j)_+.$$

Taking $\ell = \theta_i - \theta_{i+1}$, $\tau_1 = \bar{t}_0 - \theta_{i+1}$ and $\tau_2 = \bar{t}_0 + \theta_{i+1}$ in Lemma 3.1 to get

$$\begin{aligned}
&\int_{\Gamma_{i+1}} \int_{B^i} |\nabla v_i(x, t)|^p \psi_i^p(x) dxdt + \text{ess sup}_{t \in \Gamma_{i+1}} \int_{B^i} v_i^2(x, t) \psi_i^p(x) dx \\
&\quad + \int_{\Gamma_{i+1}} \int_{B^i} \int_{B^i} \frac{|v_i(x, t)\psi_i(x) - v_i(y, t)\psi_i(y)|^p}{|x - y|^{N+sp}} dx dy dt \\
&\leq C \int_{\Gamma_i} \int_{B^i} |\nabla\psi_i(x)|^p v_i^p(x, t) \eta_i^2(t) dxdt
\end{aligned}$$

$$\begin{aligned}
& + C \int_{\Gamma_i} \int_{B^i} \int_{B^i} \max\{v_i(x, t), v_i(y, t)\}^p |\psi_i(x) - \psi_i(y)|^p \eta_i^2(t) \, d\mu dt \\
& + C \operatorname{ess\,sup}_{\substack{t \in \Gamma_i \\ x \in \operatorname{supp} \psi_i}} \int_{\mathbb{R}^N \setminus B^i} \frac{v_i^{p-1}(y, t)}{|x - y|^{N+sp}} \, dy \int_{\Gamma_i} \int_{B^i} v_i(x, t) \psi_i^p(x) \eta_i^2(t) \, dx dt \\
& + C \int_{\Gamma_i} \int_{B^i} v_i^2(x, t) \psi_i^p(x) \eta_i(t) |\partial_t \eta_i(t)| \, dx dt \\
& =: I_1 + I_2 + I_3 + I_4.
\end{aligned}$$

We estimate I_1 as

$$\begin{aligned}
I_1 & \leq C k_i^p 2^{pi} r_{j+1}^{-p} \int_{\Gamma_i} \int_{B^i} \chi_{\{u_j \leq k_i\}}(x, t) \, dx dt \\
& \leq C 2^{pi} r_{j+1}^{-p} [\varepsilon \omega(r_j)]^p \int_{\Gamma_i} \int_{B^i} \chi_{\{u_j \leq k_i\}}(x, t) \, dx dt,
\end{aligned}$$

where we used the properties of ψ_i . As the computations in Pages 39-40 in [10], we derive

$$\begin{aligned}
I_2 & \leq C 2^{pi} r_{j+1}^{-p} [\varepsilon \omega(r_j)]^p \int_{\Gamma_i} \int_{B^i} \chi_{\{u_j \leq k_i\}}(x, t) \, dx dt, \\
I_3 & \leq C 2^{(N+sp)i} r_{j+1}^{-p} [\varepsilon \omega(r_j)]^p \int_{\Gamma_i} \int_{B^i} \chi_{\{u_j \leq k_i\}}(x, t) \, dx dt
\end{aligned}$$

and

$$I_4 \leq C 2^{sp_i} r_{j+1}^{-p} [\varepsilon \omega(r_j)]^p \int_{\Gamma_i} \int_{B^i} \chi_{\{u_j \leq k_i\}}(x, t) \, dx dt.$$

Since $\psi_i \equiv 1$ in B^{i+1} , we can deduce that

$$\begin{aligned}
& \int_{\Gamma_{i+1}} \int_{B^{i+1}} |\nabla v_i(x, t)|^p \, dx dt + \int_{\Gamma_{i+1}} \int_{B^{i+1}} \int_{B^{i+1}} \frac{|v_i(x, t) - v_i(y, t)|^p}{|x - y|^{N+sp}} \, dx dy dt \\
& + \operatorname{ess\,sup}_{t \in \Gamma_{i+1}} \int_{B^{i+1}} v_i^2(x, t) \, dx \\
& \leq C 2^{(N+p)i} r_{j+1}^{-p} [\varepsilon \omega(r_j)]^p \int_{\Gamma_i} \int_{B^i} \chi_{\{u_j \leq k_i\}}(x, t) \, dx dt.
\end{aligned} \tag{5.13}$$

From (5.3), we get

$$\begin{aligned}
\operatorname{ess\,sup}_{t \in \Gamma_{i+1}} \int_{B^{i+1}} v_i^p(x, t) \, dx & \leq k_i^{p-2} \operatorname{ess\,sup}_{t \in \Gamma_{i+1}} \int_{B^{i+1}} v_i^2(x, t) \, dx \\
& \leq C d_{j+1}^{-1} \operatorname{ess\,sup}_{t \in \Gamma_{i+1}} \int_{B^{i+1}} v_i^2(x, t) \, dx.
\end{aligned} \tag{5.14}$$

Combining (5.13) with (5.14) gives that

$$\begin{aligned}
& r_{j+1}^p \int_{\Gamma_{i+1}} \int_{B^{i+1}} |\nabla v_i(x, t)|^p \, dx dt + \operatorname{ess\,sup}_{t \in \Gamma_{i+1}} \int_{B^{i+1}} v_i^p(x, t) \, dx \\
& \leq d_{j+1}^{-1} \int_{\Gamma_{i+1}} \int_{B^{i+1}} |\nabla v_i(x, t)|^p \, dx dt + d_{j+1}^{-1} \operatorname{ess\,sup}_{t \in \Gamma_{i+1}} \int_{B^{i+1}} v_i^2(x, t) \, dx \\
& \leq C 2^{(N+p)i} r_{j+1}^{-p} d_{j+1}^{-1} [\varepsilon \omega(r_j)]^p \int_{\Gamma_i} \int_{B^i} \chi_{\{u_j \leq k_i\}}(x, t) \, dx dt
\end{aligned}$$

$$\leq C2^{(N+p)i}[\varepsilon\omega(r_j)]^p A_i, \quad (5.15)$$

where A_i is denoted by

$$A_i := \frac{|Q_i \cap \{u_j \leq k_i\}|}{|Q_i|}.$$

In view of Lemma 2.4, we can see

$$\begin{aligned} \int_{\Gamma_{i+1}} \int_{B^{i+1}} v_i^{p(1+\frac{p}{N})}(x, t) dx dt &\leq Cr_{j+1}^p \int_{\Gamma_{i+1}} \int_{B^{i+1}} |\nabla v_i(x, t)|^p dx dt \\ &\quad \times \left(\operatorname{ess\,sup}_{t \in \Gamma_{i+1}} \int_{B^{i+1}} v_i^p(x, t) dx \right)^{\frac{p}{N}}. \end{aligned} \quad (5.16)$$

Merging (5.15) and (5.16) leads to

$$\begin{aligned} A_{i+1} (k_i - k_{i+1})^{p(1+\frac{p}{N})} &\leq \int_{\Gamma_{i+1}} \int_{B^{i+1} \cap \{u_j \leq k_{i+1}\}} v_i^{p(1+\frac{p}{N})}(x, t) dx dt \\ &\leq Cr_{j+1}^p \int_{\Gamma_{i+1}} \int_{B^{i+1}} |\nabla v_i(x, t)|^p dx dt \times \left(\operatorname{ess\,sup}_{t \in \Gamma_{i+1}} \int_{B^{i+1}} v_i^p(x, t) dx \right)^{\frac{p}{N}} \\ &\leq C \left[2^{(N+p)i} (\varepsilon\omega(r_j))^p A_i \right]^{1+\frac{p}{N}}. \end{aligned}$$

We can readily get the recursive inequality

$$A_{i+1} \leq \tilde{C} 2^{(N+p)(1+\frac{p}{N})i} A_i^{1+\frac{p}{N}},$$

where \tilde{C} depends on N, p, s, Λ , the difference of $p/(p-1)$ and α . Set

$$\nu^* := \tilde{C}^{-\frac{N}{p}} 2^{\frac{-N(N+p)^2}{p^2}}.$$

Then we take

$$\sigma = \min \left\{ \frac{1}{4}, \exp \left(-\frac{\tilde{C}}{\nu^*} \right) \right\}.$$

Utilizing the definition of A_i , we obtain

$$A_0 = \frac{|2Q_{j+1} \cap \{u_j \leq 2\varepsilon\omega(r_j)\}|}{|2Q_{j+1}|} \leq \nu^*.$$

It follows from Lemma 4.5 that

$$\lim_{i \rightarrow \infty} A_i = 0,$$

which implies that

$$u_j(x, t) \geq \varepsilon\omega(r_j) \text{ in } Q_{j+1}.$$

Recalling the definition of u_j , we have

$$\operatorname{ess\,osc}_{Q_{j+1}} u \leq (1-\varepsilon)\omega(r_j) = (1-\varepsilon)\sigma^{-\alpha}\omega(r_{j+1}). \quad (5.17)$$

Now, we pick $\alpha \in (0, p/(p-1))$ such that

$$\sigma^\alpha \geq 1 - \varepsilon = 1 - \sigma^{\frac{p}{p-1}-\alpha},$$

which together with (5.17) ensures that

$$\operatorname{ess\,osc}_{Q_{j+1}} u \leq \omega(r_{j+1}).$$

Now we finish the proof. \square

Proof of Theorem 1.4. Let $p > 2$. Assume that u be a local weak solution of (1.1). Let $(x_0, t_0) \in Q_T$, $R \in (0, 1)$ and $Q_R \equiv B_R(x_0) \times (t_0 - R^p, t_0 + R^p)$ such that $\overline{Q_R} \subseteq Q_T$. Taking $r = R$ in Lemma 5.1 we can get

$$\operatorname{ess\,osc}_{Q_j} u \leq C \left(\frac{r_j}{R}\right)^\alpha \omega\left(\frac{R}{2}\right) \text{ for all } j \in \mathbb{N} \quad (5.18)$$

with $\alpha < p/(p-1)$, $\sigma < 1/4$, where $C \geq 1$ depends on N, p, s, Λ , and

$$\omega\left(\frac{R}{2}\right) = \operatorname{Tail}_\infty(u; x_0, R/2, t_0 - R^p, t_0 + R^p) + \left(\int_{Q_R} |u|^p dxdt\right)^{\frac{1}{2}} \vee 1. \quad (5.19)$$

For every $\rho \in (0, R/2]$, we have $\rho \in (r_{j_0+1}, r_{j_0}]$ for $j_0 \in \mathbb{N}$. Choosing $d = [C\omega(R/2)]^{2-p}$, it follows that $Q_{\rho, d\rho^p} \subseteq Q_{j_0}$. By applying (5.18), we get

$$\operatorname{ess\,osc}_{Q_{\rho, d\rho^p}} u \leq \operatorname{ess\,osc}_{Q_{j_0}} u \leq C\sigma^{-\alpha} \left(\frac{r_{j_0+1}}{R}\right)^\alpha \omega\left(\frac{R}{2}\right) \leq C\sigma^{-\alpha} \left(\frac{\rho}{R}\right)^\alpha \omega\left(\frac{R}{2}\right).$$

We can obtain the Hölder continuity from the above inequality along with (5.19). \square

ACKNOWLEDGMENT

This work was supported by the National Natural Science Foundation of China (No. 12071098).

REFERENCES

- [1] M. T. Barlow, R. F. Bass, Z. Chen and M. Kassmann, Non-local Dirichlet forms and symmetric jump processes, *Trans. Amer. Math. Soc.* 361 (4) (2009) 1963–1999.
- [2] S. Biagi, S. Dipierro, E. Valdinoci and E. Vecchi, Mixed local and nonlocal elliptic operators: regularity and maximum principles, arXiv:2005.06907.
- [3] S. Biagi, S. Dipierro, E. Valdinoci and E. Vecchi, Semilinear elliptic equations involving mixed local and nonlocal operators, *Proc. Roy. Soc. Edinburgh Sect. A*, <https://doi.org/10.1017/prm.2020.75>.
- [4] L. Brasco, E. Lindgren and M. Strömqvist, Continuity of solutions to a nonlinear fractional diffusion equation, *J. Evol. Equ.* (2021), <https://doi.org/10.1007/s00028-021-00721-2>.
- [5] L. Caffarelli, C. Chan and A. Vasseur, Regularity theory for parabolic nonlinear integral operators, *J. Amer. Math. Soc.* 24 (3) (2011) 849–869.
- [6] Z. Chen and T. Kumagai, A priori Hölder estimate, parabolic Harnack principle and heat kernel estimates for diffusions with jumps, *Rev. Mat. Iberoam.* 26 (2) (2010) 551–589.
- [7] Z. Chen, P. Kim, R. Song and Zoran Vondraček, Boundary Harnack principle for $\Delta + \Delta^{\alpha/2}$, *Trans. Amer. Math. Soc.* 364 (8) (2012) 4169–4205.
- [8] E. DiBenedetto, *Degenerate parabolic equations*, In: Universitext. Springer, New York (1993).
- [9] A. Di Castro, T. Kuusi and G. Palatucci, Local behavior of fractional p -minimizers, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 33 (2016) 1279–1299.
- [10] M. Ding, C. Zhang and S. Zhou, Local boundedness and Hölder continuity for the parabolic fractional p -Laplace equations, *Calc. Var. Partial Differential Equations*, 60 (2021), Art. 38.
- [11] S. Dipierro, E. Proietti Lippi and E. Valdinoci, Linear theory for a mixed operator with Neumann conditions, arXiv:2006.03850.
- [12] S. Dipierro, E. Proietti Lippi and E. Valdinoci, (Non)local logistic equations with Neumann conditions, arXiv:2101.02315.
- [13] S. Dipierro, X. Ros-Oton, J. Serra and E. Valdinoci, Non-symmetric stable operators: regularity theory and integration by parts, arXiv:2012.04833.

- [14] S. Dipierro and E. Valdinoci, Description of an ecological niche for a mixed local/nonlocal dispersal: an evolution equation and a new Neumann condition arising from the superposition of Brownian and Lévy processes, *Phys. A*, 575 (2021), 126052, 20 pp.
- [15] L. C. Evans, *Partial Differential Equations*, Graduate Studies in Mathematics, vol. 19. American Mathematical Society, Providence (1998).
- [16] M. Felsinger and M. Kassmann, Local regularity for parabolic nonlocal operators, *Comm. Partial Differential Equations* 38 (9) (2013) 1539–1573.
- [17] M. Foondun, Heat kernel estimates and Harnack inequalities for some Dirichlet forms with non-local part. *Electron. J. Probab.* 14 (11) (2009) 314–340.
- [18] P. Garain and J. Kinnunen, On the regularity theory for mixed local and nonlocal quasilinear elliptic equations, arXiv:2102.13365.
- [19] P. Garain and J. Kinnunen, Weak Harnack inequality for a mixed local and nonlocal parabolic equation, arXiv:2105.15016.
- [20] Y. Kim, Nonlocal Harnack inequalities for nonlocal heat equations, *J. Differential Equations*, 267 (11) (2019) 6691–6757.
- [21] J. M. Mazón, J. D. Rossi and J. Toledo, Fractional p -Laplacian evolution equations, *J. Math. Pures Appl.* 105 (6) (2016) 810–844.
- [22] M. Strömqvist, Local boundedness of solutions to nonlocal parabolic equations modeled on the fractional p -Laplacian, *J. Differential Equations*, 266 (2019) 7948–7979.
- [23] J. L. Vázquez, The Dirichlet problem for the fractional p -Laplacian evolution equation, *J. Differential Equations* 260 (7) (2016) 6038–6056.
- [24] J. L. Vázquez, The evolution fractional p -Laplacian equation in \mathbb{R}^N . Fundamental solution and asymptotic behaviour, *Nonlinear Anal.* 199 (2020), 112034, 32 pp.
- [25] J. L. Vázquez, The fractional p -Laplacian evolution equation in \mathbb{R}^N in the sublinear case, *Calc. Var. Partial Differential Equations*, 60 (2021), Art. 140.

BIN SHANG

SCHOOL OF MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN 150001, P.R. CHINA

Email address: shangbin0521@163.com

YUZHOU FANG

SCHOOL OF MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN 150001, P.R. CHINA

Email address: 18b912036@hit.edu.cn

CHAO ZHANG

SCHOOL OF MATHEMATICS AND INSTITUTE FOR ADVANCED STUDY IN MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN 150001, P.R. CHINA

Email address: czhangmath@hit.edu.cn