

# FACTORING POLYNOMIALS OVER FUNCTION FIELDS

JOSÉ FELIPE VOLOCH

**ABSTRACT.** If  $K/k$  is a function field in one variable of positive characteristic, we describe a general algorithm to factor one-variable polynomials with coefficients in  $K$ . The algorithm is flexible enough to find factors subject to additional restrictions, e.g., to find all roots that belong to a given finite dimensional  $k$ -subspace of  $K$ , more efficiently. For bounded characteristic, it runs in polynomial time, relative to factorizations over the constant field  $k$  and also provides a deterministic polynomial time irreducibility test. We also discuss applications to places of reducible reduction, when  $k$  is a global field, and to list decoding of Reed-Solomon codes.

## 1. INTRODUCTION

Let  $K/k$  be a function field in one variable, that is, a finitely generated extension of transcendence degree one with  $k$  algebraically closed in  $K$ . Let  $G(T)$  be a polynomial in one variable over  $K$ . The algorithmic problem of finding the irreducible factors of  $G(T)$  in  $K[T]$  and, in particular its roots in  $K$ , is a much-studied problem with many applications, see e.g. [vzGK85, Poh05, BvHKS09] and the references therein. A noteworthy special case is the case where  $K = k(x)$ , the rational function field, and the coefficients of  $G(T)$  are in  $k[x]$ . This case corresponds to factoring polynomials in two variables with coefficients in  $k$ . Some recent papers representing the state of the art of this special case are [Lec10, Wei17].

Many of the applications of the above problem actually require the solution of a more restricted problem. For instance, given  $G(T)$  and a finite dimensional  $k$ -subspace  $V$  of  $K$ , find the roots of  $G(T)$  in  $V$ . An example of an application where this restricted problem suffices is the Guruswami–Sudan list-decoding algorithm. See [GS00] for a comprehensive discussion and [BLQ13, NRS17] for other approaches to this problem. We discuss an instance of this in subsection 5.2 below.

Throughout this paper  $k$  has characteristic  $p > 0$  and we make the assumption that the polynomial  $G(T)$  to be factored is separable. The reduction to this case is a standard first step in all algorithms and is presented in the above cited papers. We also assume  $G(0) \neq 0$ . This paper describes an algorithm that solves the general problem of factoring such a  $G(T)$ . Additionally, the algorithm has improved performance when applied to the more restricted problems described above. Indeed, we will describe an algorithm that finds a factor of  $G(T)$  of prescribed degree whose coefficients are on prescribed finite dimensional  $k$ -subspaces of  $K$ . We prove that, assuming the characteristic is bounded, our algorithm provides a deterministic polynomial time absolute irreducibility test and, up to factorization of polynomials in  $k[T]$ , the algorithm runs in deterministic polynomial time. See Theorem 3.2, Remark 3.3 and Section 4 below for precise statements and discussion. We also discuss some other applications of the ideas in this paper in Section 5. The approach is novel and

---

2020 *Mathematics Subject Classification.* Primary: 12-08 ; Secondary: 11R09.

*Key words and phrases.* Polynomial factorization, function fields, irreducibility test, list decoding.

relies on casting the problem of finding a factor of  $G(T)$  as finding a  $k$ -linear dependence relation among some elements of a quotient ring of  $K[T]/(G(T))$  and applying a linear independence criterion over  $k$  involving Wronskian matrices.

Our algorithm, like most standard algorithms, needs at the beginning, a place of the function field  $K$  with a few additional properties. In our presentation of the main part of the algorithm, in Section 3, we just assume its existence. In the subsequent Section 4 we discuss how to find such a place. This seems to, unavoidably, require a search.

Most of the standard algorithms then proceed to compute a complete factorization of the image of the polynomial when specialized to residue field of the place just discussed (see the description of a generic factorization algorithm in [Poh05]). This is known as the lifting and recombination strategy. The factorization step in the residue field is often easy in practice but can be difficult in certain circumstances (see the beginning of Section 2). These algorithms also often have a bottleneck reconstructing global factorizations from local ones. In contrast, our algorithm does not need to compute this factorization in the residue field at the beginning, nor tries to reconstruct global factorizations from local ones. Instead, it may do a partial factorization during intermediate steps using easy gcd computations. At the end, our algorithm may need to further factor some of these partial factors. If the objective is an irreducibility criterion, it does not require finding such a factorization at all.

There are two algorithms in the literature that use differential operators to factor polynomials and don't follow the lifting and recombination strategy.

The first is the algorithm of [Rup99, Gao03] that uses a certain first order partial differential equation and shares some of the advantages of our approach. It only applies however when  $K$  is a rational function field. One application of this approach [Rup99] is to bound the size of the largest prime  $p$  for which an irreducible polynomial in  $\mathbb{Z}[x, y]$  factors modulo  $p$ . Our approach allows us to obtain similar bounds in full generality when  $k$  itself is a global field.

The second is the algorithm of [CSTU02] which uses the linear differential operator annihilating all roots of  $G(T)$  (and also uses Wronskians). This algorithm is restricted to characteristic zero as the crucial [CSTU02, Proposition 4.2] does not extend to positive characteristic. Indeed, the proposition there is about the  $k$ -dimension of the set of solutions in  $K$  of the linear differential operator annihilating roots of  $G(T)$  (assumed irreducible). In characteristic  $p$ , the set of solutions of a linear differential operator of order  $< p$  in a field  $E/K$  is a  $E^p$ -vector space. But even considering  $E^p$ -dimension, the set of solutions in  $K$  has the same  $K^p$ -dimension as the  $E^p$ -dimension of the set of solutions in the splitting field  $E$  of  $G(T)$ , as follows from [Hon81, Lemma 1], as opposed to having dimension 1, which would be the analogue of the characteristic zero result.

## 2. PRELIMINARIES

Let  $K/k$  be a function field in one variable, that is, a finitely generated extension of transcendence degree one with  $k$  algebraically closed in  $K$ . Factoring polynomials in  $K[T]$  includes factoring polynomials in  $k[T]$  and the latter could be hard, depending on  $k$ . For example, if  $k$  is a finite field or a number field, the existence of a polynomial time deterministic factoring algorithm for  $k[T]$  is an open problem, although there are polynomial time probabilistic algorithms that also perform well in practice. If  $k$  is the field of real numbers, there may be issues with precision. Our goal, therefore, is to present an algorithm to factor polynomials in  $K[T]$  relative to factoring polynomials in  $k[T]$ .

Also, in discussing the running time of the algorithm, the estimates will be in terms of the number of field operations in  $k$  or finite extensions thereof.

**2.1. The basic rings.** Let  $v$  be a place of  $K$  with ring of integers  $\mathcal{O}$  and maximal ideal  $\mathfrak{m}$ . If  $q$  is a power of the characteristic  $p$  of  $K$ , we let  $R = \mathcal{O}/\mathfrak{m}^q$ . We have that the completion of  $\mathcal{O}$  is isomorphic to  $\ell[[x]]$ , where  $\ell = \mathcal{O}/\mathfrak{m}$  and  $x$  is a uniformiser of  $v$ , that is, an element of  $\mathfrak{m} \setminus \mathfrak{m}^2$ . It follows that  $R$  is isomorphic to  $\ell[[x]]/(x^q)$ .

Let  $G[T] \in \mathcal{O}[T]$  be a monic polynomial such that  $G \pmod{v}$  is separable.

The ring  $R_1 = (\mathcal{O}/\mathfrak{m}^q)[T]/(G(T))$  is an Artinian ring and, thus, a direct sum of Artinian local rings. These summands correspond to irreducible factors of  $G(T)$  which, in turn, correspond to irreducible factors of  $G(T) \in (\mathcal{O}/\mathfrak{m})[T]$ . The standard factorization version of Hensel's lemma gives an algorithmic way to go from the latter to the former.

**2.2. Hasse derivatives and Wronskians.** We begin with presenting some concepts and results from [Sch39, SV86]. See also [GV87] which proves stronger versions of the main results of [Sch39] and may be more accessible, as well as [Hes02, Section 6.1] which presents relevant algorithms (particularly algorithms 26 and 28 there). Let  $K/k$  be a function field. The usual higher derivatives do not work well in small characteristics. A suitable replacement for higher derivatives that work in general are the Hasse derivatives. We denote by  $D^{(i)}, i = 0, 1, \dots$ , the Hasse derivatives with respect to some separating variable  $x$  on  $K$ . These are  $k$ -linear operators on  $K$  with  $i!D^{(i)} = (d/dx)^i$  and satisfying:

$$\begin{aligned} D^{(i)} \circ D^{(j)} &= \binom{i+j}{j} D^{(i+j)}, \\ D^{(i)}(uv) &= \sum_{j=0}^i D^{(j)}(u) D^{(i-j)}(v). \end{aligned} \tag{2.1}$$

The second formula is an extension of the Leibniz rule. They are defined first in  $k[x]$  by setting  $D^{(i)}x^n = \binom{n}{i}x^{n-i}$  and extending  $k$ -linearly. These operators then extend uniquely to  $k(x)$  and any separable extension thereof by requiring that they satisfy the formulas 2.1 above.

By [Sch39, Satz 2],  $f_0, \dots, f_m \in K$  are linearly independent over  $k$  if and only if the Wronskian matrix  $(D^{(i)}(f_j))$  has maximal rank  $m+1$ . In this case, there is a minimal list of integers  $0 = \varepsilon_0 < \dots < \varepsilon_m$  such that the matrix  $(D^{(\varepsilon_i)}(f_j))$  has maximal rank  $m+1$ . We have  $\varepsilon_i = i$  if the characteristic is zero or large enough but that is not going to be the situation in this paper. Also, to a place  $v$  of  $K$ , we can associate a minimal list of integers  $0 = j_0 < \dots < j_m$  (which depend on  $v$ ) such that the matrix  $(D^{(j_i)}(f_j) \pmod{v})$  has maximal rank  $m+1$ . If  $K$  is the function field of an algebraic curve  $Y$ , then the morphism  $(f_0 : \dots : f_m) : Y \rightarrow \mathbb{P}^m$  has some degree  $\Delta$ , then we can take  $\varepsilon_i \leq j_i \leq \Delta$  when the  $f_i$  are linearly independent ([SV86, Section 2], [Hes02, Prop. 13]). On the other hand, if the Wronskian matrix has rank  $m$  and  $a_0, \dots, a_m \in K$  satisfy  $\sum_{j=0}^m a_j D^{(j)}(f_j) = 0, i = 0, 1, 2, \dots$  and  $a_0 = 1$ , then  $a_j \in k, j = 0, 1, \dots, m$  as follows from the proof of [Sch39, Satz 1].

Consider a monic, separable polynomial

$$G(T) = \sum_{j=0}^s a_j T^j, a_j \in K, a_s = 1. \tag{2.2}$$

Let  $R = K[T]/G(T)$  and  $t$  the image of  $T$  in  $R$ . We extend the operators  $D^{(i)}, i \geq 0$  to  $R$ . We need an expression for  $D^{(i)}(t)$ . We have that

$$0 = D^{(i)}(G(t)) = \sum_{j_1+2j_2+\dots+ij_i \leq i} A_{j_1,\dots,j_i} (D^{(1)}(t))^{j_1} \dots (D^{(i)}(t))^{j_i} \quad (2.3)$$

where the  $A_{j_1,\dots,j_i}$  are polynomials in  $t$  and, in particular,  $A_{0,\dots,0,1} = G'(t)$  which is invertible in  $R$  and this determines  $D^{(i)}(t)$  uniquely, by induction. They can be computed more efficiently by the algorithms of [Hes02].

If  $v$  is a place of  $K$  with ring of integers  $\mathcal{O}$  and maximal ideal  $\mathfrak{m} = (x)$  then, for any  $q$  power of  $p$ , the operators  $D^{(i)}, i < q$ , defined using  $x$  as the separating variable, induce operators on  $\mathcal{O}/\mathfrak{m}^q$ . Indeed,  $D^{(i)}(x^q) = 0, i < q$ , so  $D^{(i)}(x^q y) = x^q D^{(i)}(y), i < q$  for any  $y \in \mathcal{O}$  and it follows that  $D^{(i)}, i < q$  preserve  $\mathfrak{m}^q$ . In our main algorithm, we will choose  $q$  large enough so that the non-zero elements of  $K$  involved in the algorithm have zeros of multiplicity less than  $q$ , so, if such an element is in  $\mathfrak{m}^q$ , it will vanish.

Now, if  $G(T) \in \mathcal{O}[T]$  is as above and is such that  $G \pmod{v}$  is separable, then the operators  $D^{(i)}, i < q$  induce operators on  $R_1 = (\mathcal{O}/\mathfrak{m}^q)[T]/(G(T))$  by the same formulas. As mentioned before,  $R_1$  is an Artinian ring and, thus, a direct sum of Artinian local rings. For each irreducible factor  $H_0$  of  $G \pmod{\mathfrak{m}}$ , there is a corresponding factor  $H$  of  $G \pmod{\mathfrak{m}^q}$  and a corresponding summand  $(\mathcal{O}/\mathfrak{m}^q)[T]/(H(T))$  of  $R_1$ . If  $\mathfrak{m} = (x)$ , then  $\mathcal{O}/\mathfrak{m}^q$  is isomorphic to  $(\mathcal{O}/\mathfrak{m})[X]/(X^q)$  and  $(\mathcal{O}/\mathfrak{m}^q)[T]/(H(T))$  is isomorphic to  $((\mathcal{O}/\mathfrak{m})[T]/(H_0(T)))[X]/(X^q)$  and we refer to the subring  $(\mathcal{O}/\mathfrak{m})[T]/(H_0(T))$  of this latter ring as its constants.

**Lemma 2.1.** *If  $u \in R_1$  satisfies  $D^{(i)}(u) = 0, 0 < i < q$ , then in each local summand of  $R_1$ ,  $u$  is constant (in the above sense).*

*Proof.* Each summand is isomorphic to  $\ell[X]/(X^q)$  for some field  $\ell$  and  $D^{(i)}(x^r) = \binom{r}{i} x^{r-i}$ , where  $x$  is the image of  $X$ , so it is clear that  $D^{(i)}(u) = 0, 0 < i < q$  if and only if  $u \in \ell$ .  $\square$

**Remark 2.2.** *In the process of algorithm 1 below, we will decompose  $R_1$  as a direct sum but we may not go all the way to the full decomposition as a sum of Artinian local rings.*

### 3. THE MAIN ALGORITHM

To obtain a full factorization algorithm, the following lemma (a variant of [Poh05, Lemma 4.1]) provides bounds for the valuations of the coefficients of the potential factors of  $G(T)$  and these bounds can be used to define suitable  $V_i$  as Riemann-Roch spaces.

**Lemma 3.1.** *Let  $G(T) = \sum_{i=0}^s a_i T^i \in K[T]$  with  $a_s = 1, a_0 \neq 0$  and let  $t$  be a root of  $G(T)$  in some finite extension  $L/K$  and  $H(T) = \sum_{i=0}^r b_i T^i$  be its monic minimal polynomial over  $K$ . Then, for any place  $v$  of  $L$  extending a place of  $K$ ,*

$$v(t) \geq \min_{0,1,\dots,s-1} v(a_i)/(s-i)$$

*and, for any place  $v$  of  $K$ ,*

$$v(b_j) \geq (r-j) \min_{0,1,\dots,s-1} v(a_i)/(s-i).$$

*Proof.* Recall that we assume throughout that  $G(0) \neq 0$ , so  $t \neq 0$ . If  $iv(t) + v(a_i) > sv(t)$  for all  $i < s$ , then  $\infty = v(G(t)) = \min\{iv(t) + v(a_i)\} = sv(t)$ , contradiction. This gives the first part of the lemma.

---

**Algorithm 1** Find factor of  $G(T)$  with restricted coefficients

---

**Input**

A function field  $K/k$

A polynomial  $G(T) \in K[T]$  monic, separable, of degree  $s$  with discriminant  $f \neq 0$ .

A place  $v$  of  $K/k$ , with ring of integers  $\mathcal{O}$  and maximal ideal  $\mathfrak{m}$  with  $G(T) \in \mathcal{O}[T]$  and  $v(f) = 0$  and an uniformizer of  $v$  used to define the  $D^{(i)}$ .

Finite dimensional  $k$ -vector spaces  $V_i \subset K, i = 0, \dots, r-1$ , with  $1 \in V_0$ , together with a  $k$ -basis  $\{h_{ij}\}$  for each  $V_i$ , where  $r < s$ . Put  $h_{01} = h_{r1} = 1, m = \sum \dim V_i$ .

**Output**

Either a proof that  $G(T)$  has no monic factor of the form  $H(T) = \sum_{i=0}^r b_i T^i, b_i \in V_i, i < r, b_r = 1$ .

Or a direct summand  $S$  of the ring  $R_1 = (\mathcal{O}/\mathfrak{m}^q)[T]/(G(T))$  and elements  $b_i, i = 0, \dots, r$  of  $S$  such that, for each local summand of  $S$ , a factor of  $G(T)$  of the required form can be obtained (see Theorem 3.2 and Remark 3.3).

- 1:  $R = K[T]/(G(T))$ .
  - 2:  $\Phi \in R^{m+1}$  the row vector with entries (in some order)  $h_{ij}t^i \in R, i = 0, \dots, r$  and for each  $i < r, j = 1, \dots, \dim V_i$  and  $j = 1$  for  $i = r$ .
  - 3: Compute an integer  $\Delta$  which bounds  $j_i, \varepsilon_i$  for any map to  $\mathbb{P}^m$  obtained by viewing  $\Phi$  in a quotient field of  $R$  in which its entries are linearly independent over  $k$  (use 3.1 if no better one is available).
  - 4: Compute  $q$ , smallest power of  $p$  with  $q > \max\{m, \Delta\}$  (so  $q \leq p \max\{m, \Delta\}$ ).
  - 5:  $R_1 = (\mathcal{O}/\mathfrak{m}^q)[T]/(G(T)), R_0 = (\mathcal{O}/\mathfrak{m})[T]/(G(T))$ .
  - 6: Compute matrix  $M$  with rows  $D^{(i)}(\Phi), i = 0, \dots, q-1$ , working in  $R_1$ .
  - 7: Attempt to do Gaussian elimination on  $M$ , working in  $R_1$ .
  - 8: **if** Some pivot  $P(T) \in R_1$  is not invertible **then**
  - 9:     Compute  $H_0(T) = \gcd(G(T), P(T))$  and  $E_0(T) = G(T)/H_0(T)$  in  $(\mathcal{O}/\mathfrak{m})[T]$ .
  - 10:    Lift factorization  $G(T) = H_0(T)E_0(T)$  in  $R_0$  to factorization  $G(T) = H(T)E(T)$  in  $R_1$  and replace  $G(T)$  by  $H(T)$  and  $E(T)$  in step 5.
  - 11: **end if**
  - 12: **if**  $M$  has full rank  $m+1$  **then**
  - 13:     **return**  $G(T)$  has no factor of the form required for the output.
  - 14: **else if**  $M$  has rank  $m$  **then**
  - 15:     Compute solution  $u_{ij} \in R_1$  of  $\sum_{ij} u_{ij} D^{(\ell)}(h_{ij}t^i) = 0, u_{r1} = 1, \ell = 0, 1, \dots, q-1$ .
  - 16:     **return**  $b_i = \sum_j u_{ij} h_{ij}, i = 0, \dots, r$ .
  - 17: **else**
  - 18:     Go back to 2 and remove the last entry from  $\Phi$ .
  - 19: **end if**
- 

We have that  $b_j$  is the  $(r-j)$ -th symmetric function on the conjugates of  $t$  so the second part follows from the first by extending  $v$  to a valuation of the splitting field of  $H$ .  $\square$

It follows from Lemma 3.1 that, without any further restriction on the valuations of the elements of  $V_i$ , that we can take

$$\Delta = -r \sum_v \min\{0, \min_{0,1,\dots,s-1} v(a_i)/(s-i)\} \quad (3.1)$$

as a bound for the  $j_i, \varepsilon_i$ . Such a bound can be improved if additional restrictions are put in the  $V_i$ .

We begin by proving that Algorithm 1 performs as described and runs in polynomial time. We defer a detailed estimate of the running time and a discussion of how to compute the auxiliary place to Section 4.

**Theorem 3.2.** *Given the above input, Algorithm 1 runs in deterministic polynomial time in  $p, s, \Delta$  (measured in number of operations in the field  $\mathcal{O}/\mathfrak{m}$ ) and outputs either a certificate that  $G(T)$  has no factor of the required form or a decomposition of  $R_1$  as a direct sum of at most  $s$  rings. Moreover, for each such summand ( $R'$ , say), the algorithm outputs elements  $u_{ij}$  of  $R'$  that are constant in each summand of the decomposition of  $R'$  into local rings and from which a factor of  $G(T)$  of the required form can be constructed or a certificate that this summand does not yield such a factor. In particular, the algorithm provides a deterministic polynomial time absolute irreducibility test in characteristic  $p$  for  $p$  polynomially bounded in  $s, \Delta$ .*

*Proof.* By induction on  $m$ . Assume  $m = 1$ . Since  $1 \in V_0, \Phi = (1, t^r)$  and  $M$  has rank 1 or 2. If the rank is 2 in  $R'$ , it is clear there is no factor of the required form. If the rank is 1 in  $R'$ , this means that  $D^{(i)}(t^r) = 0, i = 1, \dots, q-1$ , so  $t^r$  is constant in any local summand of  $R'$ , the solution of the linear system is  $u_{01} = -t, u_{r1} = 1$  and  $H(T) = T - t$  is a factor of  $G(T)$  of the required form.

As mentioned above, step 3 is dealt with in general by equation 3.1 unless there is a better bound available.

As mentioned above, the operators  $D^{(i)}, i < q$  act on  $R_1$  and the computation of the matrix  $M$  in step 6 is polynomial in operations in  $R_1$  but  $\dim_{\mathcal{O}/\mathfrak{m}} R_1 \leq sq$  giving a bound in terms of the number of operations in  $\mathcal{O}/\mathfrak{m}$ .

In the process of Gaussian elimination (whose running time is polynomial in the size of the matrix), for each candidate pivot in the ring  $R_1$ , the factorization from steps 9 and 10 allows us to write  $R_1$  as a direct sum of two rings for which the pivot is invertible in one factor and zero in the other. In the first factor, we use the pivot as in the usual Gaussian elimination and, in the second factor, we look for a new pivot. The work done before splitting  $R_1$  is reused. At the end of steps 1-11, we arrive at a decomposition of  $R$  as a sum of at most  $s$  rings and the image of  $M$  in each of these is put in row echelon form by the Gaussian elimination process.

Those summands of  $R_1$  where  $M$  has full rank  $m+1$  yield no factor of  $G(T)$  of the required form, since such a factor is a linear relation among the entries of  $\Phi$  with constant coefficients and, applying  $D^{(i)}, i < q$  to this relation shows that the coefficients are in the kernel of  $M$ . For other factors  $R'$  where  $M$  has rank  $m$ , we compute the  $u_{ij}$  as described in step 15. It follows from the proof of [GV87, Theorem 1] that  $D^{(r)}(u_{ij}) = 0, 0 < r < q$  and thus, from Lemma 2.1 the  $u_{ij}$  are constant in each local factor of the decomposition of  $R'$ . Let  $S$  be one such factor. It determines a factor  $H(T)$  of  $G(T)$  and a place  $w|v$  in the function field  $L$  obtained by adjoining a root  $t$  of  $H(T)$  to  $K$ . Moreover, if  $\hat{u}_{ij}$  are the constants in  $S$



corresponding to the  $u_{ij}$ , the fact that  $\sum u_{ij}h_{ij}t^i = 0$  in  $S$ , shows that  $\sum \hat{u}_{ij}h_{ij}t^i$  vanishes to order at least  $q$  at  $w$ . But  $q > \Delta$  and  $\Delta$  is an upper bound for the degree of the morphism associated to  $\Phi$ , therefore an upper bound for the degree of  $\sum \hat{u}_{ij}h_{ij}t^i$  and this implies that  $\sum \hat{u}_{ij}h_{ij}t^i = 0$  in  $L$ . Hence  $\sum \hat{u}_{ij}h_{ij}T^i$  yields a factor of  $G(T)$  of the required form. For the factors  $R'$  where  $M$  has rank smaller than  $m$ , we decrement  $m$ , drop an entry of  $\Phi$  as described, rerun the algorithm and are done by induction.  $\square$

**Remark 3.3.** *What Theorem 3.2 does not do is identify the  $u_{ij}$  with specific elements of an algebraic extension of  $k$ , necessarily. For that, we need to further factor the factor of  $G(T)$  corresponding to the ring  $R'$  as a product of irreducibles to obtain the full decomposition of  $R'$  as a sum of local rings and identify the  $u_{ij}$  with constants in each summand.*

*As mentioned in the introduction, the first step in most standard factoring algorithms for  $K[T]$  is to fully factor  $G(T) \in (\mathcal{O}/\mathfrak{m})[T]$  and how it is performed depends on the nature of the field  $k$ . In our algorithm, this step may not be required at all (if  $G(T)$  has no factors, or a single irreducible factor, of the required form) or it may only be needed for a proper factor of  $G(T)$ .*

*On the other hand, it is not claimed that the algorithm, as described, will output all factors of the required form, only that it will output at least one such factor, in case it exists. This can be easily remedied by dividing by the output factors and rerunning the algorithm with the quotient as input.*

We will discuss an example but, beforehand, here is a non-example. If  $G(T) \in k[T]$ , that is, has constant coefficients, then  $D^{(i)}(t) = 0, i > 0$ , the matrix  $M$  has always rank one and the algorithm unravels to the base case  $m = 1$ . The polynomial  $T - t$  is a factor of  $G(T)$  for  $G(t) = 0$  and we are left with the task of factoring  $G(T)$  over the constant field  $k$ .

For a more representative example consider

$$G(T) = T^4 + (x+1)T^3 + (x^2+1)T^2 + (x^3+x^2+1)T + (x^2+x) \in \mathbb{F}_2(x)[T].$$

We look for factors of  $G(T)$  of the form  $T+ax+b$ , so  $r = 1$ ,  $V_0$  is spanned by  $1, x, \Phi = (1, x, t)$  where  $t$  is the image of  $T$  in  $(k[x]/(x^4)[T]/(G(T)))$  and  $m = 2$ . Modulo the ideal  $(x+1)$ ,  $G(T)$  reduces to  $T^4 + T$ . We find, using 2.3 that  $D^{(2)}(t) = J(t)/G'(t)^3$ , where

$$J(T) = (xT^5 + (x^2+x)T^4 + x^5T + (x^6+x^5)).$$

The gcd of  $G(T)$  and  $J(T)$  is  $H(T) = T^2 + T + x^2 + x$  and, switching to the ring  $R_1 = (k[x]/(x+1)^4)[T]/(H(T))$ , we let  $t$  be the image of  $T$ , forcing  $D^{(2)}(t) = 0$ . Using 2.3 now with  $H$ , we find that  $D(t) = 1$ . The matrix  $M$  is therefore

$$M = \begin{pmatrix} 1 & x & t \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

Now, we solve the system  $t + ax + b = D(t) + a = 0$ , so  $a = D(t) = 1, b = t + x$ . Now, modulo the ideal  $(x+1)$ , we have  $H(T) = T^2 + T = T(T+1)$  and we lift this factorization to  $k[x]/(x+1)^4$  and find  $H(T) = (T+x)(T+x+1)$ . Now,  $R_1$  is a direct sum of two rings from this factorization and  $b = 0, 1$  respectively in each of the factors. Consequently,  $G(T)$  has the factors  $T+x, T+x+1$  of the required form.

**Remark 3.4.** *If the characteristic is zero,  $R_1$  is not defined and if the characteristic is large,  $R_1$  is too big. An extension of the algorithm of this paper to those cases would require*

a replacement for  $R_1$ . For theoretical results, such as Theorem 5.1, we can work with  $R = \mathcal{O}[T]/(G(T))$  but for algorithmic purposes we need a ring finite dimensional over  $k$ . We could work with the ring  $(\mathcal{O}/\mathfrak{m}^n)[T]/(G(T))$  for some appropriate choice of  $n$ , but the higher derivatives are not typically well-defined as operators on this ring. Instead, it makes sense to look at  $D^{(i)} : (\mathcal{O}/\mathfrak{m}^n)[T]/(G(T)) \rightarrow (\mathcal{O}/\mathfrak{m}^{n-i})[T]/(G(T))$ . We have not worked out the full details of this possibility.

#### 4. RUNNING TIME ESTIMATES AND FINDING A SUITABLE PLACE

We retain the notation of the previous section and especially of Algorithm 1.

**Theorem 4.1.** *Given a place  $v$  as in the input of Algorithm 1 with  $[\mathcal{O}/\mathfrak{m} : k] = d$ , the algorithm runs in  $O(ds(p \max\{m, \Delta\})^4)$  field operations in  $k$ .*

*Proof.* The ring  $R_1$  is initialized in the algorithm and during its execution it is replaced by its direct summands, of which there are at most  $s$ . From [Hes02, Section 6.1], the computation of the matrix  $M$  takes  $O(q^2)$  operations in each ring  $R_1$  and Gaussian elimination of the matrix  $M$  takes  $O(q^3)$  operations in each ring  $R_1$ . The latter dominates the former. The initial ring  $R_1$  is a vector space over  $\mathcal{O}/\mathfrak{m}$  of dimension  $sq$ , so we can bound the running time of these calculations as  $O(sq^4)$  operations in  $\mathcal{O}/\mathfrak{m}$ . Using now that  $q \leq p \max\{m, \Delta\}$  we arrive at a running time of  $O(s(p \max\{m, \Delta\})^4)$  operations in  $\mathcal{O}/\mathfrak{m}$ . If we let  $d = [\mathcal{O}/\mathfrak{m} : k]$ , we finally arrive at  $O(ds(p \max\{m, \Delta\})^4)$  operations in  $k$ .  $\square$

**Remark 4.2.** *There are faster algorithms in the special cases of most interest, e.g.  $k$  finite,  $K = k(x)$  [Lec10, Wei17] or finding a root of  $G(T)$ , again for  $k$  finite [GS00]. There does not seem to be an algorithm with the same generality and flexibility with a stated running time for direct comparison.*

It remains to make explicit how to find a suitable place  $v$  and compute  $d$ . We are given a function field  $K/k$  and  $f \in K, f \neq 0$ , we need to construct a place  $v$  of  $K$  such that  $v(f) = 0$ , as well as a uniformizer for this place. Without loss of generality, we assume that  $k$  is finitely generated over its prime field. We can also assume that  $f \notin k$ , for otherwise the condition  $v(f) = 0$  is automatic and we can replace  $f$  by an element of  $K \setminus k$  in order to produce the place  $v$ . Under this additional hypothesis, there are only at most  $2[K : k(f)]$  places of  $K$  with  $v(f) \neq 0$  as  $[K : k(f)] = \sum_v \max\{0, v(f)\} = -\sum_v \min\{0, v(f)\}$ . Hence, to find  $v$  it suffices to generate enough places of  $K$  until a place with  $v(f) = 0$  is found. It seems that such a search is, at the moment, unavoidable. While, in some cases (e.g.  $k$  finite,  $K = k(x)$ ) this is easy, it can be difficult in the generality we work with and some of the literature seems to gloss over this point. We assume  $K/k(f)$  separable. For a general discussion of how to reduce to this case, see [Ste05]. Then  $K = k(f, g)$  for some  $g \in K$  and there is  $P(x, y) \in k[x, y], P(f, g) = 0$ . We then find  $\alpha \neq 0$  in  $k$  or in an extension field such that  $P(\alpha, y)$  is separable. There is a place of  $K$  corresponding to each irreducible factor of  $P(\alpha, y)$  over  $k(\alpha)$  and, if  $m(x)$  is the minimal polynomial of  $\alpha$  over  $k$ , then  $m(f)$  is a uniformizer for any such place, as  $m(f)$  is a uniformizer for the corresponding place of  $k(f)$  which, by construction, is unramified in  $K$ . An easy way of ensuring that  $P(\alpha, y)$  is separable is to require that it be irreducible in  $k(\alpha)[y]$ . This has the advantage that irreducibility tests are easier than factorization and that  $m(x) = x - \alpha$  in this case. On the other hand, this makes the degree of the place bigger.



Assume first that  $k$  is a finite field. This case is discussed in [GS00, Algorithm 3.2] with some additional assumptions which are relevant there but not here, so we just give a simplified discussion. Basically, one just searches for  $\alpha$  in  $k$  or an extension thereof until a suitable element is found. While this procedure is straightforward enough, it is worth pointing out that we may not have a suitable place with residue field  $k$  or even an extension of  $k$  of small degree as we can see by considering a simple example such as  $G(T) = T^p - (x^{p^n} - x)T + 1 \in \mathbb{F}_p(x)[T]$  (which has  $f = \pm(x^{p^n} - x)^p$ ). It is possible to ensure that a suitable place with residue field  $k$  exists if  $\#k$  is large enough in terms of  $[K : k(f)]$  and the genus of  $K$ .

The case where  $k$  is not algebraic over its prime field can be tackled as follows. Let  $k_0$  be the algebraic closure in  $k$  of its prime field. Then  $K$  and  $k$  are respectively the function field of varieties  $X, Y$  over  $k_0$  with a map  $X \rightarrow Y$  of relative dimension 1. We realize  $X$  as a hypersurface of projective space and intersect  $X$  with random hypersurfaces. As long as these hypersurfaces intersect  $X$  in an irreducible subset that is not a component of the divisor of  $f$  and is transversal to the generic fiber of the map  $X \rightarrow Y$ , such a hypersurface will define a place of  $K/k$  satisfying our requirements and the equation of the hypersurface is the uniformizer we need. The version of Bertini's theorem over finite fields from [CP16] guarantees that, for high enough degree, most hypersurfaces satisfy our conditions. Checking irreducibility can be done using the main algorithm of this paper (applied to a field of smaller transcendence degree) and the other conditions can be checked directly. Alternatively, one can use the explicit version of Hilbert's irreducibility theorem for function fields from [BSE21] to estimate how far one might need to search.

## 5. OTHER APPLICATIONS

**5.1. Arithmetic properties.** As mentioned in the introduction, [Rup99] has a bound on the size of the largest prime  $p$  for which an irreducible polynomial in  $\mathbb{Z}[x, y]$  factors modulo  $p$ . We substantially extend this result. We will consider polynomials in  $K[T]$  where  $K/k$  is a function field and  $k$  itself is a global field. We impose no restriction on the characteristic of  $k$ . We can associate a height to elements of  $k$  in the usual way ([Lan83, Chapter 3, §3]), namely, for  $a \in k$ , the height of  $a$  is  $-\sum_v \min\{0, v(a)\}$ , where  $v$  runs through the places of  $k$ . If we represent  $K$  as a finite extension of  $k(x)$  for some transcendental element  $x$  of  $K$ , for an element  $a \in K$ , there is a polynomial  $P(x, y) \in k[x, y]$ ,  $P(x, a) = 0$  of minimal degree. We call the coefficients (in  $k$ ) of  $P$  simply the coefficients of  $a$  and use the maximum of their heights as a measure of the complexity of  $a$ . Given  $G(T) \in K[T]$ , for all but finitely primes  $\mathfrak{p}$  of  $k$ , we can consider the reduction of  $G(T)$  modulo  $\mathfrak{p}$ .

**Theorem 5.1.** *Let  $k$  be a global field,  $K/k$  a function field and  $G(T) \in K[T]$  an irreducible polynomial, as in equation 2.2 and define  $\Delta$  as in equation 3.1. Let  $H$  be the maximum height of the coefficients of the  $a_i$ . The norm  $N(\mathfrak{p})$  of the primes  $\mathfrak{p}$  of  $k$  for which the reduction of  $G(T)$  modulo  $\mathfrak{p}$  is either undefined or reducible satisfies  $N(\mathfrak{p}) = O(H^{(\Delta+1)^3})$ , where the implied constant depends on  $s, \Delta$ .*

*Proof.* Since  $G(T)$  is assumed irreducible, the matrix  $M$  in Algorithm 1 has maximal rank and so a maximal minor has non-zero determinant. Note that  $\Delta + 1$  is a bound for the number of columns (hence also of rows) of  $M$ . We now compute  $M$  with entries in  $R$  (with no restriction on the characteristic of  $k$ ). It follows by induction from equation 2.3 that the height of the coefficients of  $D^{(i)}(t^j)$  is  $O(H^{ij}) = O(H^{(\Delta+1)^2})$ , hence the height of the

coefficients of the (non-zero) determinant of a maximal  $(\Delta + 1) \times (\Delta + 1)$ -minor of  $M$  is  $O(H^{(\Delta+1)^3})$ . But for the reduction of  $G(T)$  modulo  $\mathfrak{p}$  to be either undefined or reducible,  $\mathfrak{p}$  needs to divide the determinant of this minor and the result follows.  $\square$

In the case where  $k = \mathbb{F}_q(Z)$ , the approach of the previous theorem can be used to prove a proximity gap statement (in the sense of [BSCI<sup>+</sup>20, BN20]) for Algebraic Geometry codes.

**5.2. List decoding.** A different application concerns the the Guruswami–Sudan list-decoding algorithm for Reed–Solomon or Algebraic Geometry codes [GS00]. If one has a fixed code, there are fixed finite dimensional  $k$ -vector spaces  $V_0, W_0, \dots, W_{s-1} \subset K$  and, for each received message, a polynomial  $G(T) \in K[T]$  is constructed with coefficients  $a_i \in W_i$  (the precise construction is irrelevant at the moment), for which we want to know its roots in  $V_0$ . That is precisely the problem we dealt with above, with  $r = 1$ .

For an alternative approach to improving the factorization step of the Guruswami–Sudan algorithm, see [AP00]. Additionally, [AP00, Theorem 5] provides an alternative way of finding a suitable place (in the sense of Section 4) to be used in either approach to factorization in the context of the Guruswami–Sudan algorithm.

For a prime power  $q$  and an integer  $k, 1 \leq k \leq q$ , the Reed–Solomon code  $RS_k(q)$  is the subspace of the  $q$ -dimensional space of functions  $\mathbb{F}_q \rightarrow \mathbb{F}_q$  corresponding to the functions represented by a polynomial  $f(x) \in \mathbb{F}_q[x], \deg f < k$ . Define, as usual for  $g_1, g_2 : \mathbb{F}_q \rightarrow \mathbb{F}_q$ , the Hamming distance  $d(g_1, g_2) = \#\{\alpha \in \mathbb{F}_q \mid g_1(\alpha) \neq g_2(\alpha)\}$ .

McEliece [RJM03] has shown that, for a fixed Reed–Solomon code, most invocations of the Guruswami–Sudan algorithm output a list of codewords of size 0 or 1. We will strengthen this result and show that, for a certain range of Reed–Solomon codes, for most invocations of the Guruswami–Sudan algorithm, the polynomial  $G(T) \in K[T]$  has at most one root in  $V_0$ , so we can take full advantage of our factoring algorithm’s enhanced performance in this situation. Our result is meant to be illustrative and one might expect a similar result in much greater generality.

The following proposition sets the stage for our result and is a special case of Sudan’s original precursor [Sud97] to the Guruswami–Sudan algorithm.

**Proposition 5.2.** *Assume  $(q+5)/10 < k \leq q/8$  and that  $g : \mathbb{F}_q \rightarrow \mathbb{F}_q$  is any function. Then there exists a non-zero polynomial  $G(x, y) = a_0(x) + a_1(x)y + a_2(x)y^2 + a_3(x)y^3 + a_4(x)y^4 \in \mathbb{F}_q[x, y]$  with  $\deg a_i \leq (4-i)(k-1), i = 0, \dots, 4$  such that  $\forall \alpha \in \mathbb{F}_q, G(\alpha, g(\alpha)) = 0$ . Moreover, if  $f(x) \in RS_k(q)$  satisfies  $d(f, g) \leq [q/2]$  then  $G(x, f(x))$  is identically zero.*

*Proof.* Each condition  $G(\alpha, g(\alpha)) = 0$  is a linear equation on the coefficients of the  $a_i$ . The total number of coefficients is  $\sum_{i=0}^4 ((4-i)(k-1)+1) = 10k-5 > q$ , by hypothesis. So there is a  $G$ , as prescribed. If  $f(x) \in RS_k(q)$  satisfies  $d(f, g) \leq [q/2]$  then  $G(x, f(x))$  is a polynomial with at least  $q/2$  zeros but, since  $\deg f \leq k-1$ , we have  $\deg G(x, f(x)) \leq 4(k-1) < q/2$  by hypothesis, hence  $G(x, f(x))$  is identically zero.  $\square$

So, as mentioned above, given such  $G$ , we need to find its factors of the form  $y - f(x), \deg f < k$  to find the elements  $f(x) \in RS_k(q), d(f, g) \leq [q/2]$  and, by construction, there are at most 4 such. This is the “list” in the list-decoding algorithm. It is easy to see that, for most choices of  $g$ , the list is empty. Indeed, there are  $q^q$  possible  $g$ ’s and at most  $q^k \sum_{i=0}^{[q/2]} \binom{q}{i} (q-1)^i$  of those satisfy  $d(f, g) \leq [q/2]$  for some  $f \in RS_k(q)$  and the sum is asymptotic to  $q^{k+q/2+o(q)}$ , hence bounded above by  $q^{5q/8+o(q)}$ , much smaller than  $q^q$ .

More interestingly, McEliece [RJM03] has shown that, for most choices of  $g$  such that there exists some  $f \in RS_k(q)$  with  $d(f, g) \leq [q/2]$ , the  $f$  is unique. It follows that  $q^{k+q/2+o(q)}$  is the correct order of magnitude for the number of  $g$  with  $d(f, g) \leq [q/2]$  for some  $f \in RS_k(q)$ . The following result strengthens this result by showing that, for most choices of  $g$  such that there exists some  $f \in RS_k(q)$  with  $d(f, g) \leq [q/2]$ , any polynomial  $G$ , output of Proposition 5.2, has only one factor of the form  $y - f(x)$ ,  $\deg f < k$ . This justifies our claim that a factoring algorithm that performs better when there is only one factor of prescribed type improves the running time of the list decoding algorithm.

**Theorem 5.3.** *Assume  $(q + 5)/10 < k < q/8$ . Among those  $g : \mathbb{F}_q \rightarrow \mathbb{F}_q$  such that there exists some  $f \in RS_k(q)$  with  $d(f, g) \leq q/2$ , the number of those for which a polynomial  $G$ , output of Proposition 5.2, has more than one factor of the form  $y - f(x)$ ,  $\deg f < k$  is at most  $q^{5k+o(q)}$ .*

*Proof.* We assume that  $d(f_0, g) = \delta \leq [q/2]$ , for some  $f_0$  as above. Replacing  $g$  by  $g - f_0$ , where  $f_0$  is one choice, we can assume  $f_0 = 0$ . By Proposition 5.2,  $G(x, y) = yH(x, y)$ , for some  $H$ . Moreover,  $H(\alpha, g(\alpha)) = 0$  for  $\delta$  values of  $\alpha$ . We split the proof in two cases, depending on whether or not there exists  $\alpha \in \mathbb{F}_q$  with  $H(\alpha, y)$  identically zero.

Suppose first that there is no such  $\alpha$ . The number of  $g$ 's for a given choice of  $H$  is at most  $\binom{q}{\delta} 3^\delta$ . Indeed, there are  $\binom{q}{\delta}$  choices for a subset of size  $\delta$  of  $\mathbb{F}_q$  and, for each  $\alpha$  in this subset, at most 3 choices for a solution of  $H(\alpha, y) = 0$  as  $H$  is cubic in  $y$  (for the  $\alpha$  outside of the subset we set  $g(\alpha) = 0$ ). If, in turn,  $H(x, y) = (y - f(x))S(x, y)$  with  $f \in RS_k(q)$  and

$$S(x, y) = b_0(x) + b_1(x)y + b_2(x)y^2, \deg b_i \leq (2 - i)(k - 1), i = 0, 1, 2,$$

we can count the number of coefficients, namely  $k$  for  $f$  and  $3k$  for the  $b_i$ 's. Hence at most  $q^{4k}$  possible such  $H$ 's and at most  $\sum_{\delta \leq [q/2]} \binom{q}{\delta} 3^\delta q^{4k}$  possible such  $g$ 's. If we now allow the initial  $f_0$  to vary (as opposed to be set to  $f_0 = 0$ ) we end up with a total of at most  $\sum_{\delta \leq [q/2]} \binom{q}{\delta} 3^\delta q^{5k}$  for the count of the theorem and this is at most  $q^{5k+o(q)}$ , as desired.

Let's now assume that there are  $\alpha \in \mathbb{F}_q$  with  $H(\alpha, y)$  identically zero and let  $c(x)$  be the monic polynomial that has these  $\alpha$ 's as roots. Since the coefficient of  $y^3$  in  $H(x, y)$  as a polynomial in  $y$  is constant, we conclude in this case that it is zero. We can then write  $H$  (in the case that  $G$  has a second root  $f \in RS_k(q)$ ) as

$$H(x, y) = c(x)(y - f(x))(b_0(x)/c(x) + (b_1(x)/c(x))y).$$

If we fix  $\deg c = m$ , the number of possible  $H$ 's is at most  $q^m q^k q^{2k-m} q^{k-m} = q^{4k-m}$  by counting the number of possible  $c, f, b_0/c, b_1/c$  respectively. The number of  $g$  given  $H$  is bounded above by  $3^{q-m} q^m$  by counting the options for  $g(\alpha)$  when  $c(\alpha) \neq 0$  or  $c(\alpha) = 0$ , respectively. So, the count in this case is at most  $q^{4k+o(q)}$  and again, allowing  $m$  and the initial  $f_0$  to vary (as opposed to be set to  $f_0 = 0$ ) we end up with a total of at most  $q^{5k+o(q)}$   $g$ 's, completing the proof.  $\square$

We note that  $5k < k + q/2$  if  $k < q/8$  which justifies the claim made immediately before the theorem. We also note that there are functions, as in the statement of the theorem, for which  $G$  has more than one factor of the required form. For example, if  $q$  is even, we can take any  $g$  that takes only values 0, 1 each  $q/2$  times. Then  $G = y(y + 1)$  is an output of Proposition 5.2 for this  $g$ .

Finally, we note that Algorithm 1 vastly simplifies in the present context. In this case  $K = \mathbb{F}_q(x), T = y, r = 1$ . The vector  $\Phi$  is  $(1, x, x^2, \dots, x^{k-1}, y)$  and the matrix  $M$  is already in echelon form with rows  $(0, 0, 0, \dots, 0, D^{(\epsilon)}(y))$  after the first  $k$  rows. The only real computation is to express the pivots  $D^{(\epsilon)}(y)$  as elements of  $R_1$  and do the computations of step 9 of the algorithm.

## ACKNOWLEDGEMENTS

This work was supported by MBIE. I would also like to thank M. Esgin, V. Kuchta, S. Ruj, A. Sakzad and R. Steinfeld of the Trans-Tasman ZK group for questions that motivated part of this research.

## REFERENCES

- [AP00] Daniel Augot and Lancelot Pecquet, *A Hensel lifting to replace factorization in list-decoding of algebraic-geometric and Reed-Solomon codes*, IEEE Trans. Inform. Theory **46** (2000), no. 7, 2605–2614. [↑10](#)
- [BSE21] Lior Bary-Soroker and Alexei Entin, *Explicit Hilbert’s irreducibility theorem in function fields*, Abelian varieties and number theory, Contemp. Math., vol. 767, Amer. Math. Soc., Providence, RI, 2021, pp. 125–134. [↑9](#)
- [BvHKS09] Karim Belabas, Mark van Hoeij, Jürgen Klüners, and Allan Steel, *Factoring polynomials over global fields*, J. Théor. Nombres Bordeaux **21** (2009), no. 1, 15–39. [↑1](#)
- [BSCI<sup>+</sup>20] Eli Ben-Sasson, Dan Carmon, Yuval Ishai, Swastik Kopparty, and Shubhangi Saraf, *Proximity Gaps for Reed-Solomon Codes* (2020). Cryptology ePrint Archive, Report 2020/654. [↑10](#)
- [BLQ13] Jérémy Berthomieu, Grégoire Lecerf, and Guillaume Quintin, *Polynomial root finding over local rings and application to error correcting codes*, Appl. Algebra Engrg. Comm. Comput. **24** (2013), no. 6, 413–443. [↑1](#)
- [BN20] Sarah Bordage and Jade Nardi, *Interactive Oracle Proofs of Proximity to Algebraic Geometry Codes* (2020). arXiv:2011.04295. [↑10](#)
- [CP16] François Charles and Bjorn Poonen, *Bertini irreducibility theorems over finite fields*, J. Amer. Math. Soc. **29** (2016), no. 1, 81–94. [↑9](#)
- [CSTU02] Olivier Cormier, Michael F. Singer, Barry M. Trager, and Felix Ulmer, *Linear differential operators for polynomial equations*, J. Symbolic Comput. **34** (2002), no. 5, 355–398. [↑2](#)
- [GS00] Shuhong Gao and M. Amin Shokrollahi, *Computing roots of polynomials over function fields of curves*, Coding theory and cryptography (Annapolis, MD, 1998), Springer, Berlin, 2000, pp. 214–228. [↑1](#), [8](#), [9](#), [10](#)
- [Gao03] Shuhong Gao, *Factoring multivariate polynomials via partial differential equations*, Math. Comp. **72** (2003), no. 242, 801–822. [↑2](#)
- [GV87] Arnaldo García and José Felipe Voloch, *Wronskians and linear independence in fields of prime characteristic*, Manuscripta Math. **59** (1987), no. 4, 457–469. [↑3](#), [6](#)
- [vzGK85] J. von zur Gathen and E. Kaltofen, *Factorization of multivariate polynomials over finite fields*, Math. Comp. **45** (1985), no. 171, 251–261. [↑1](#)
- [Hes02] Florian Hess, *An algorithm for computing Weierstrass points*, Algorithmic number theory (Sydney, 2002), Lecture Notes in Comput. Sci., vol. 2369, Springer, Berlin, 2002, pp. 357–371. [↑3](#), [4](#), [8](#)
- [Hon81] Taira Honda, *Algebraic differential equations*, Symposia Mathematica, Vol. XXIV (Sympos., INDAM, Rome, 1979), Academic Press, London-New York, 1981, pp. 169–204. [↑2](#)
- [Lan83] Serge Lang, *Fundamentals of Diophantine geometry*, Springer-Verlag, New York, 1983. MR715605 [↑9](#)
- [Lec10] Grégoire Lecerf, *New recombination algorithms for bivariate polynomial factorization based on Hensel lifting*, Appl. Algebra Engrg. Comm. Comput. **21** (2010), no. 2, 151–176. [↑1](#), [8](#)

- [RJM03] R. J. McEliece, *On the Average List Size for the Guruswami-Sudan Decoder*, 7th International Symposium on Communications Theory and Applications (ISCTA), 2003. [↑10](#), [11](#)
- [NRS17] Vincent Neiger, Johan Rosenkilde, and Éric Schost, *Fast computation of the roots of polynomials over the ring of power series*, ISSAC'17—Proceedings of the 2017 ACM International Symposium on Symbolic and Algebraic Computation, ACM, New York, 2017, pp. 349–356. [↑1](#)
- [Poh05] Michael E. Pohst, *Factoring polynomials over global fields. I*, J. Symbolic Comput. **39** (2005), no. 6, 617–630. [↑1](#), [2](#), [4](#)
- [Rup99] Wolfgang M. Ruppert, *Reducibility of polynomials  $f(x, y)$  modulo  $p$* , J. Number Theory **77** (1999), no. 1, 62–70. [↑2](#), [9](#)
- [Sch39] Friedrich Karl Schmidt, *Die Wronskische Determinante in beliebigen differenzierbaren Funktionskörpern*, Mathematische Zeitschrift **45** (1939), no. 1, 62–74. [↑3](#)
- [Ste05] Allan Steel, *Conquering inseparability: primary decomposition and multivariate factorization over algebraic function fields of positive characteristic*, J. Symbolic Comput. **3** (2005), 1053–1075. [↑8](#)
- [SV86] Karl-Otto Stöhr and José Felipe Voloch, *Weierstrass points and curves over finite fields*, Proc. London Math. Soc. (3) **52** (1986), no. 1, 1–19. [↑3](#)
- [Sud97] Madhu Sudan, *Decoding of Reed Solomon codes beyond the error-correction bound*, J. Complexity **13** (1997), no. 1, 180–193. [↑10](#)
- [Wei17] Martin Weimann, *Bivariate factorization using a critical fiber*, Found. Comput. Math. **17** (2017), no. 5, 1219–1263. [↑1](#), [8](#)

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF CANTERBURY, PRIVATE BAG 4800,  
CHRISTCHURCH 8140, NEW ZEALAND

*Email address:* felipe.voloch@canterbury.ac.nz

*URL:* <http://www.math.canterbury.ac.nz/~f.voloch>