

# On Liouville type theorems in the stationary non-Newtonian fluids

Dongho Chae\*, Junha Kim† and Jörg Wolf‡

Department of Mathematics,  
Chung-Ang University  
Seoul 06974, Republic of Korea

## Abstract

In this paper we prove a Liouville type theorem for the stationary equations of a non-Newtonian fluid in  $\mathbb{R}^3$  with the viscous part of the stress tensor  $\mathbf{A}_p(u) = \operatorname{div}(|\mathbf{D}(u)|^{p-2}\mathbf{D}(u))$ , where  $\mathbf{D}(u) = \frac{1}{2}(\nabla u + (\nabla u)^\top)$  and  $\frac{9}{5} < p < 3$ . We consider a weak solution  $u \in W_{loc}^{1,p}(\mathbb{R}^3)$  and its potential function  $\mathbf{V} = (V_{ij}) \in W_{loc}^{2,p}(\mathbb{R}^3)$ , i.e.  $\nabla \cdot \mathbf{V} = u$ . We show that there exists a constant  $s_0 = s_0(p)$  such that if the  $L^s$  mean oscillation of  $\mathbf{V}$  for  $s > s_0$  satisfies a certain growth condition at infinity, then the velocity field vanishes. Our result includes the previous results [5, 6] as particular cases.

**AMS Subject Classification Number:** 35Q35, 35Q30, 76A05, 76D05, 76D03  
**keywords:** non-Newtonian fluid equations, Liouville type theorem

## 1 Introduction

We consider a power law model of non-Newtonian fluid in  $\mathbb{R}^3$

$$\begin{cases} -\mathbf{A}_p(u) + (u \cdot \nabla)u = -\nabla\pi & \text{in } \mathbb{R}^3, \\ \operatorname{div} u = 0, \end{cases} \quad (1.1)$$

where  $u = (u_1(x), u_2(x), u_3(x))$  is the velocity field,  $\pi = \pi(x)$  is the pressure field. The diffusion term is represented by

$$\mathbf{A}_p(u) = \operatorname{div}(|\mathbf{D}(u)|^{p-2}\mathbf{D}(u)), \quad 1 < p < +\infty$$

---

\*E-mail: dchae@cau.ac.kr

†E-mail: jha02@cau.ac.kr

‡E-mail: jwolf2603@gmail.com

and the deviatoric stress tensor is interpreted as  $|\mathbf{D}|^{p-2}\mathbf{D} = \boldsymbol{\sigma}(\mathbf{D})$ , where  $\mathbf{D}(u) = \frac{1}{2}(\nabla u + (\nabla u)^\top)$  is the symmetric gradient. If  $2 < p < +\infty$ , the equations describe shear thickening fluids, of which viscosity increases along with shear rate  $|\mathbf{D}(u)|$ . If  $1 < p < 2$ , shear thinning fluids satisfy them. In the case of  $p = 2$ , (1.1) corresponds to the usual stationary Navier-Stokes equations which represent Newtonian fluids. We refer to Wilkinson [1] for continuum mechanical background of the above system.

The Liouville problem for the stationary Navier-Stokes equations (Galdi [4], Remark X. 9.4, pp. 729) has attracted considerable attention in the mathematical fluid mechanics. Though it is still open, there are positive answers under additional conditions (see [7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17]). And as a generalization, Liouville type theorems for non-Newtonian fluids have been investigated (see [18, 5]).

Let  $u \in L^1_{loc}(\mathbb{R}^3)$  be a vector field, and let  $\mathbf{V} = (V_{ij}) \in L^1_{loc}(\mathbb{R}^3; \mathbb{R}^3 \times \mathbb{R}^3)$  be a matrix valued function satisfying  $\operatorname{div} \mathbf{V} = u$  in the distributional sense. In [6] Chae and Wolf proved Liouville type theorem for the stationary Navier-Stokes equations when the following is assumed

$$\left( \int_{B(r)} |\mathbf{V} - \mathbf{V}_{B(r)}|^s dx \right)^{\frac{1}{s}} \leq Cr^{\min\{\frac{1}{3} - \frac{1}{s}, \frac{1}{6}\}} \quad \forall 1 < r < +\infty$$

for some  $3 < s < +\infty$ . They also considered it in (1.1) when  $\frac{9}{5} < p < 3$  but only for  $s = \frac{3p}{2p-3}$  in [5]. We generalize these results.

As is well known, weak solutions are actually smooth for  $p = 2$ . Otherwise there is only partial regularity of weak solutions [2, 3]. In this paper, we consider weak solutions, which is defined as follows :

**Definition 1.1.** Let  $\frac{9}{5} < p < 3$ . A function  $u \in W^{1,p}_{loc}(\mathbb{R}^3)$  is called a weak solution to (1.1) if

$$\int_{\mathbb{R}^3} (|\mathbf{D}(u)|^{p-2}\mathbf{D}(u) - u \otimes u) : \mathbf{D}(\varphi) dx = 0 \quad (1.2)$$

is fulfilled for all vector fields  $\varphi \in C_c^\infty(\mathbb{R}^3)$  with  $\operatorname{div} \varphi = 0$ .

**Remark 1.2.** Let  $u$  be a weak solution. Then, there exists  $\pi \in L^{\frac{p}{p-1}}_{loc}(\mathbb{R}^3)$  satisfying

$$\int_{B(r)} |\pi - \pi_{B(r)}|^s dx \leq C \int_{B(r)} ||\mathbf{D}(u)|^{p-2}\mathbf{D}(u) - u \otimes u|^s dx, \quad \forall 0 < r < +\infty \quad (1.3)$$

for all  $\frac{3}{2} \leq s \leq \frac{p}{p-1}$ . And  $(u, \pi)$  holds

$$\int_{\mathbb{R}^3} (|\mathbf{D}(u)|^{p-2}\mathbf{D}(u) - u \otimes u) : \mathbf{D}(\varphi) dx = \int_{\mathbb{R}^3} \pi \operatorname{div} \varphi dx \quad (1.4)$$

for any vector field  $\varphi \in W^{1,p}(\mathbb{R}^3)$  with compact support. Hence, (1.4) replaces (1.2). We refer to [5] for a brief explanation.

**Remark 1.3.** Let  $(u, \pi)$  be a weak solution and  $\phi \in C_c^\infty(\mathbb{R}^3)$ . If we take  $\varphi = u\phi$ , then (1.4) with  $\varphi \in W^{1,p}(\mathbb{R}^3)$  yields the local energy equality

$$\int_{\mathbb{R}^3} |\mathbf{D}(u)|^p \phi dx = - \int_{\mathbb{R}^3} |\mathbf{D}(u)|^{p-2} \mathbf{D}(u) : u \otimes \nabla \phi dx + \int_{\mathbb{R}^3} \left( \frac{1}{2} |u|^2 + \pi \right) u \cdot \nabla \phi dx. \quad (1.5)$$

**Theorem 1.4.** Let  $\frac{9}{5} < p < 3$ , and  $\frac{3}{2} < s < +\infty$  satisfy

$$s \geq \frac{3p}{2(2p-3)}, \quad s > \frac{9-3p}{2p-3}.$$

Let  $(u, \pi) \in W_{loc}^{1,p}(\mathbb{R}^3) \times L_{loc}^{\frac{p}{p-1}}(\mathbb{R}^3)$  be a weak solution to (1.1). We set

$$\alpha(p, s) := \min \left\{ \frac{1}{3} - \frac{5p-9}{s(2p-3)}, \frac{3}{p} - \frac{4}{3} \right\}.$$

If we assume there exists a potential  $\mathbf{V} \in W_{loc}^{2,p}(\mathbb{R}^3; \mathbb{R}^{3 \times 3})$  such that

$$\left( \int_{B(r)} |\mathbf{V} - \mathbf{V}_{B(r)}|^s dx \right)^{\frac{1}{s}} \leq Cr^{\alpha(p,s)} \quad \forall 1 < r < +\infty, \quad (1.6)$$

then  $u \equiv 0$ .

**Remark 1.5.** If  $s = \frac{3p}{2p-3}$ , then we obtain  $\alpha = \frac{9-4p}{3p}$ , which consistent with Theorem 1.3 (ii) in [5].

**Remark 1.6.** In the case of  $p = 2$ , we have  $3 < s < +\infty$  and  $\alpha = \min\{\frac{s}{3} - 1, \frac{s}{6}\}$ , corresponding to Theorem 1.1 in [6].

We denote by  $C(p, s) = C$  a generic constant that may vary from line to line.

## 2 Caccioppoli type inequalities

To prove Theorem 1.4, we need the following two lemmas.

**Lemma 2.1.** Let  $\frac{9}{5} < p < 3$ . Let  $u \in W_{loc}^{1,p}(\mathbb{R}^3)$  and  $1 \leq R < +\infty$ . For  $0 < \rho < R$ , we let  $\psi \in C_c^\infty(B(R))$  satisfy  $0 \leq \psi \leq 1$  and  $|\nabla \psi| \leq C(R-\rho)^{-1}$ . If we assume there exists  $\mathbf{V} \in W_{loc}^{2,p}(\mathbb{R}^3)$  such that  $\operatorname{div} \mathbf{V} = u$  and (1.6) for some  $\frac{3p}{2(2p-3)} \leq s \leq \frac{3p}{2p-3}$ , then for  $s \geq p$

$$\int_{B(R)} |\psi^p u|^p dx \leq CR^{\frac{3}{2} + \frac{p}{6} - \frac{p(5p-9)}{2s(2p-3)}} \|\psi \nabla u\|_{L^p}^{\frac{p}{2}} + C(R-\rho)^{-p} R^{3 + \frac{p}{3} - \frac{p(5p-9)}{s(2p-3)}}, \quad (2.7)$$

and for  $s < p$

$$\begin{aligned} \int_{B(R)} |\psi^p u|^p dx &\leq CR^{\frac{p^2}{3(2p-3)}} \|\psi \nabla u\|_{L^p}^{\frac{p(sp-3s+3p)}{2sp-3s+3p}} + C(R-\rho)^{-\frac{3p}{2s} + \frac{3}{2}} R^{\frac{p}{6} + \frac{p^2}{2s(2p-3)}} \|\psi \nabla u\|_{L^p}^{\frac{p}{2}} \\ &+ C(R-\rho)^{-\frac{sp^2}{sp+3p-3s}} R^{\frac{p^2(2sp+3p-3s)}{3(2p-3)(sp+3p-3s)}} \|\psi \nabla u\|_{L^p}^{\frac{p(3p-3s)}{sp-3s+3p}} + C(R-\rho)^{-p - \frac{3p}{s} + 3} R^{\frac{p}{3} + \frac{p^2}{s(2p-3)}}. \end{aligned} \quad (2.8)$$

*Proof.* We first let  $\frac{9}{5} < p < 2$ . Beacuse  $s \geq \frac{3p}{2(2p-3)} \geq p$ , we show (2.7) in this case. Hölder's inequality implies that

$$\int_{B(R)} |\psi^p u|^p dx \leq |B(R)|^{1-\frac{p}{2}} \left( \int_{B(R)} |\psi^p u|^2 dx \right)^{\frac{p}{2}} = CR^{3-\frac{3p}{2}} \left( \int_{B(R)} |u|^2 \psi^{2p} dx \right)^{\frac{p}{2}}. \quad (2.9)$$

Recalling  $u = \operatorname{div} \mathbf{V}$  and using integration by parts, we have

$$\begin{aligned} \int_{B(R)} |u|^2 \psi^{2p} dx &= \int_{B(R)} \partial_i (\mathbf{V}_{ij} - (\mathbf{V}_{ij})_{B(R)}) u_j \psi^{2p} dx \\ &= - \int_{B(R)} (\mathbf{V}_{ij} - (\mathbf{V}_{ij})_{B(R)}) \partial_i u_j \psi^{2p} dx - 2p \int_{B(R)} (\mathbf{V}_{ij} - (\mathbf{V}_{ij})_{B(R)}) u_j \psi^{2p-1} \partial_i \psi dx \\ &\leq \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}| |\psi \nabla u| \psi^{2p-1} dx + 2p \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}| |\psi^p u| |\nabla \psi| \psi^{p-1} dx \\ &\leq \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}| |\psi \nabla u| dx + C(R - \rho)^{-1} \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}| |\psi^p u| dx. \end{aligned}$$

Note that  $\frac{3p}{2(2p-3)} \leq s \leq \frac{3p}{2p-3}$  implies

$$\frac{1}{s} + \frac{1}{p} < 1$$

and

$$\alpha = \frac{1}{3} - \frac{5p-9}{s(2p-3)}.$$

Using Hölder's inequality and (1.6), we obtain

$$\begin{aligned} \int_{B(R)} |u|^2 \psi^{2p} dx &\leq \left( \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}|^s dx \right)^{\frac{1}{s}} \|\psi \nabla u\|_{L^p} |B(R)|^{1-\frac{1}{p}} \\ &\quad + C(R - \rho)^{-1} \left( \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}|^s dx \right)^{\frac{1}{s}} \|\psi^p u\|_{L^p} |B(R)|^{1-\frac{1}{p}} \\ &\leq CR^{\frac{10}{3} - \frac{5p-9}{s(2p-3)} - \frac{3}{p}} \|\psi \nabla u\|_{L^p} + C(R - \rho)^{-1} R^{\frac{10}{3} - \frac{5p-9}{s(2p-3)} - \frac{3}{p}} \|\psi^p u\|_{L^p}. \end{aligned}$$

Inserting this inequality into (2.9) and applying Young's inequality, we have

$$\begin{aligned} \int_{B(R)} |\psi^p u|^p dx &\leq CR^{\frac{3}{2} + \frac{p}{6} - \frac{p(5p-9)}{2s(2p-3)}} \|\psi \nabla u\|_{L^p}^{\frac{p}{2}} + C(R - \rho)^{-\frac{p}{2}} R^{\frac{3}{2} + \frac{p}{6} - \frac{p(5p-9)}{2s(2p-3)}} \|\psi^p u\|_{L^p}^{\frac{p}{2}} \\ &\leq CR^{\frac{3}{2} + \frac{p}{6} - \frac{p(5p-9)}{2s(2p-3)}} \|\psi \nabla u\|_{L^p}^{\frac{p}{2}} + C(R - \rho)^{-p} R^{3 + \frac{p}{3} - \frac{p(5p-9)}{s(2p-3)}} + \frac{1}{2} \int_{B(R)} |\psi^p u|^p dx, \end{aligned}$$

which implies (2.7).

Now, we let  $2 \leq p < 3$ . Using  $u = \operatorname{div} \mathbf{V}$ , integration by parts and Hölder's inequality, we find

$$\begin{aligned}
& \int_{B(R)} |\psi^p u|^p dx = \int_{B(R)} \partial_i (\mathbf{V}_{ij} - (\mathbf{V}_{ij})_{B(R)}) u_j |u|^{p-2} \psi^{p^2} dx \\
& = - \int_{B(R)} (\mathbf{V}_{ij} - (\mathbf{V}_{ij})_{B(R)}) (\partial_i u_j |u|^{p-2} + (p-2) u_j u_k \partial_i u_k |u|^{p-4}) \psi^{p^2} dx \\
& \quad - p^2 \int_{B(R)} (\mathbf{V}_{ij} - (\mathbf{V}_{ij})_{B(R)}) u_j |u|^{p-2} \psi^{p^2-1} \partial_i \psi dx \\
& \leq C \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}| |\psi \nabla u| |\psi^{p+1} u|^{p-2} dx \\
& \quad + C(R-\rho)^{-1} \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}| |\psi^{p+1} u|^{p-1} dx \\
& \leq C \|\mathbf{V} - \mathbf{V}_{B(R)}\|_{L^s(B(R))} \|\psi \nabla u\|_{L^p} \|\psi^{p+1} u\|_{L^{\frac{sp}{sp-s-p}}}^{p-2} \\
& \quad + C(R-\rho)^{-1} \|\mathbf{V} - \mathbf{V}_{B(R)}\|_{L^s(B(R))} \|\psi^{p+1} u\|_{L^{\frac{s}{s-1}}}^{p-1} = I + II.
\end{aligned}$$

We assume  $s \geq p$  first. Since

$$1 \leq \frac{sp}{sp-s-p} \leq \frac{p}{p-2}$$

by applying Hölder's inequality to  $I$  we have

$$I \leq C \|\mathbf{V} - \mathbf{V}_{B(R)}\|_{L^s(B(R))} \|\psi \nabla u\|_{L^p} \|\psi^p u\|_{L^p}^{p-2} |B(R)|^{\frac{1}{p} - \frac{1}{s}}.$$

Subsequently, we use (1.6) and Young's inequality. This yields

$$\begin{aligned}
I & \leq CR^{\frac{3}{p}} \left( \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}|^s dx \right)^{\frac{1}{s}} \|\psi \nabla u\|_{L^p} \|\psi^p u\|_{L^p}^{p-2} \\
& \leq CR^{\frac{3}{p} + \frac{1}{3} - \frac{5p-9}{s(2p-3)}} \|\psi \nabla u\|_{L^p} \|\psi^p u\|_{L^p}^{p-2} \\
& \leq CR^{\frac{3}{2} + \frac{p}{6} - \frac{p(5p-9)}{2s(2p-3)}} \|\psi \nabla u\|_{L^p}^{\frac{p}{2}} + \frac{1}{4} \int |\psi^p u|^p dx. \tag{2.10}
\end{aligned}$$

Arguing similarly to the above, having

$$1 < \frac{s}{s-1} \leq \frac{p}{p-1},$$

we can calculate  $II$  as follows

$$\begin{aligned}
II & \leq C(R-\rho)^{-1} \|\mathbf{V} - \mathbf{V}_{B(R)}\|_{L^s(B(R))} \|\psi^{p+1} u\|_{L^p}^{p-1} |B(R)|^{\frac{1}{p} - \frac{1}{s}} \\
& \leq C(R-\rho)^{-1} R^{\frac{3}{p}} \left( \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}|^s dx \right)^{\frac{1}{s}} \|\psi^p u\|_{L^p}^{p-1}
\end{aligned}$$

$$\begin{aligned}
&\leq C(R - \rho)^{-1} R^{\frac{3}{s} + \frac{1}{3} - \frac{5p-9}{s(2p-3)}} \|\psi^p u\|_{L^p}^{p-1} \\
&\leq C(R - \rho)^{-p} R^{3 + \frac{p}{s} - \frac{p(5p-9)}{s(2p-3)}} + \frac{1}{4} \int |\psi^p u|^p dx. \tag{2.11}
\end{aligned}$$

(2.10) and (2.11) show (2.7).

Now we assume  $s < p$ . Since it holds

$$\frac{p}{p-2} < \frac{sp}{sp-s-p} \leq \frac{3p}{3-p} < \frac{3p}{(3-p)(p-2)},$$

the standard interpolation inequality implies that

$$\begin{aligned}
\|\psi^{p+1} u\|_{L^{\frac{sp}{sp-s-p}}}^{p-2} &\leq \|\psi^{p+1} u\|_{L^{\frac{3p}{(3-p)(p-2)}}}^{p-2} \|\psi^{p+1} u\|_{L^{\frac{sp(p-2)-3p+3s}{sp(p-2)}}}^{\frac{3p-3s}{sp(p-2)}} \\
&= \|\psi^{p+1} u\|_{L^{\frac{3p}{3-p}}}^{\frac{3p-3s}{sp}} \|\psi^{p+1} u\|_{L^p}^{\frac{sp(p-2)-3p+3s}{sp}}.
\end{aligned}$$

Thus, by using Sobolev inequality here, we arrive at

$$\begin{aligned}
\|\psi^{p+1} u\|_{L^{\frac{sp}{sp-s-p}}}^{p-2} &\leq C \left( \|\psi^{p+1} \nabla u\|_{L^p} + (R - \rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{3p-3s}{sp}} \|\psi^{p+1} u\|_{L^p}^{\frac{sp(p-2)-3p+3s}{sp}} \\
&\leq C \|\psi \nabla u\|_{L^p}^{\frac{3p-3s}{sp}} \|\psi^p u\|_{L^p}^{\frac{sp(p-2)-3p+3s}{sp}} + C(R - \rho)^{-\frac{3p-3s}{sp}} \|\psi^p u\|_{L^p}^{p-2}.
\end{aligned}$$

Inserting the inequality we just obtained into  $I$  and applying (1.6) along with Young's inequality, we find

$$\begin{aligned}
I &\leq CR^{\frac{3}{s}} \left( \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}|^s dx \right)^{\frac{1}{s}} \|\psi \nabla u\|_{L^p}^{\frac{sp+3p-3s}{sp}} \|\psi^p u\|_{L^p}^{\frac{sp(p-2)-3p+3s}{sp}} \\
&\quad + C(R - \rho)^{-\frac{3p-3s}{sp}} R^{\frac{3}{s}} \left( \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}|^s dx \right)^{\frac{1}{s}} \|\psi \nabla u\|_{L^p} \|\psi^p u\|_{L^p}^{p-2} \\
&\leq CR^{\frac{3}{s} + \frac{1}{3} - \frac{5p-9}{s(2p-3)}} \|\psi \nabla u\|_{L^p}^{\frac{sp+3p-3s}{sp}} \|\psi^p u\|_{L^p}^{\frac{sp(p-2)-3p+3s}{sp}} \\
&\quad + C(R - \rho)^{-\frac{3p-3s}{sp}} R^{\frac{3}{s} + \frac{1}{3} - \frac{5p-9}{s(2p-3)}} \|\psi \nabla u\|_{L^p} \|\psi^p u\|_{L^p}^{p-2} \\
&\leq CR^{\frac{p^2}{3(2p-3)}} \|\psi \nabla u\|_{L^p}^{\frac{p(sp+3p-3s)}{2sp+3p-3s}} + C(R - \rho)^{-\frac{3p}{2s} + \frac{3}{2}} R^{\frac{3p}{2s} - \frac{3}{2} + (\frac{3}{2} + \frac{p}{6} - \frac{p(5p-9)}{2s(2p-3)})} \|\psi \nabla u\|_{L^p}^{\frac{p}{2}} \\
&\quad + \frac{1}{4} \int |\psi^p u|^p dx. \tag{2.12}
\end{aligned}$$

We note that  $s < p$  implies

$$\frac{p}{p-1} < \frac{s}{s-1} < \frac{3p}{2(3-p)} < \frac{3p}{(3-p)(p-1)}.$$

Thus, using Interpolation inequality and Sobolev inequality, we infer

$$\|\psi^{p+1} u\|_{L^{\frac{s}{s-1}}}^{p-1} \leq \|\psi^{p+1} u\|_{L^{\frac{3p}{(3-p)(p-1)}}}^{p-1} \|\psi^{p+1} u\|_{L^{\frac{p}{p-1}}}^{\frac{sp(p-1)-3p+3s}{sp(p-1)}}$$

$$\begin{aligned}
&= \|\psi^{p+1}u\|_{L^{\frac{3p-3s}{3p}}}^{\frac{3p-3s}{sp}} \|\psi^{p+1}u\|_{L^p}^{\frac{sp(p-1)-3p+3s}{sp}} \\
&\leq C \left( \|\psi^{p+1}\nabla u\|_{L^p} + (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{3p-3s}{sp}} \|\psi^{p+1}u\|_{L^p}^{\frac{sp(p-1)-3p+3s}{sp}} \\
&\leq C \|\psi\nabla u\|_{L^p}^{\frac{3p-3s}{sp}} \|\psi^p u\|_{L^p}^{\frac{sp(p-1)-3p+3s}{sp}} + C(R-\rho)^{-\frac{3p-3s}{sp}} \|\psi^p u\|_{L^p}^{p-1}.
\end{aligned}$$

Inserting this inequality into  $II$  and using (1.6) and Young's inequality, it follows

$$\begin{aligned}
II &\leq C(R-\rho)^{-1} R^{\frac{3}{s}} \left( \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}|^s dx \right)^{\frac{1}{s}} \|\psi\nabla u\|_{L^p}^{\frac{3p-3s}{sp}} \|\psi^p u\|_{L^p}^{\frac{sp(p-1)-3p+3s}{sp}} \\
&\quad + C(R-\rho)^{-1-\frac{3p-3s}{sp}} R^{\frac{3}{s}} \left( \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}|^s dx \right)^{\frac{1}{s}} \|\psi^p u\|_{L^p}^{p-1} \\
&\leq C(R-\rho)^{-1} R^{\frac{3}{s} + \frac{1}{3} - \frac{5p-9}{s(2p-3)}} \|\psi\nabla u\|_{L^p}^{\frac{3p-3s}{sp}} \|\psi^p u\|_{L^p}^{\frac{sp(p-1)-3p+3s}{sp}} \\
&\quad + C(R-\rho)^{-1-\frac{3p-3s}{sp}} R^{\frac{3}{s} + \frac{1}{3} - \frac{5p-9}{s(2p-3)}} \|\psi^p u\|_{L^p}^{p-1} \\
&\leq C(R-\rho)^{-\frac{sp^2}{sp+3p-3s}} R^{\frac{p^2(2sp+3p-3s)}{3(2p-3)(sp+3p-3s)}} \|\psi\nabla u\|_{L^p}^{\frac{p(3p-3s)}{sp+3p-3s}} \\
&\quad + C(R-\rho)^{-p-\frac{3p}{s}+3} R^{\frac{3p}{s} + \frac{p}{3} - \frac{p(5p-9)}{s(2p-3)}} + \frac{1}{4} \int |\psi^p u|^p dx. \tag{2.13}
\end{aligned}$$

By (2.12) and (2.13) we complete the proof.  $\square$

**Lemma 2.2.** *Let  $\frac{9}{5} < p < 3$ . Let  $u \in W_{loc}^{1,p}(\mathbb{R}^3)$  and  $1 \leq R < +\infty$ . For  $0 < \rho < R$ , we let  $\psi \in C_c^\infty(B(R))$  satisfy  $0 \leq \psi \leq 1$  and  $|\nabla\psi| \leq C(R-\rho)^{-1}$ . If we assume there exists  $\mathbf{V} \in W_{loc}^{2,p}(\mathbb{R}^3)$  such that  $\operatorname{div} \mathbf{V} = u$  and (1.6) for some  $\frac{3p}{2(2p-3)} \leq s \leq \frac{3p}{2p-3}$ , then for  $s \geq 3$*

$$\begin{aligned}
\int_{B(R)} |\psi^3 u|^3 dx &\leq CR \|\psi\nabla u\|_{L^p}^{\frac{9p}{2sp+3p-3s}} + CR \left( (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{9p}{2sp+3p-3s}} \\
&\quad + C(R-\rho)^{-3} R^{4-\frac{3(5p-9)}{s(2p-3)}}, \tag{2.14}
\end{aligned}$$

and for  $s < 3$

$$\begin{aligned}
\int_{B(R)} |\psi^3 u|^3 dx &\leq CR \|\psi\nabla u\|_{L^p}^{\frac{9p}{2sp+3p-3s}} + CR \left( (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{9p}{2sp+3p-3s}} \\
&\quad + C(R-\rho)^{-\frac{3s(2p-3)}{sp+3p-3s}} R^{\frac{2sp+3p-3s}{sp+3p-3s}} \|\psi\nabla u\|_{L^p}^{\frac{3p(3-s)}{sp+3p-3s}} \\
&\quad + C(R-\rho)^{-\frac{3s(2p-3)}{sp+3p-3s}} R^{\frac{2sp+3p-3s}{sp+3p-3s}} \left( (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{3p(3-s)}{sp+3p-3s}}. \tag{2.15}
\end{aligned}$$

*Proof.* Arguing similarly to the proof of Lemma 2.1, we get

$$\begin{aligned} \int_{B(R)} |\psi^3 u|^3 dx &\leq C \|\mathbf{V} - \mathbf{V}_{B(R)}\|_{L^s(B(R))} \|\psi \nabla u\|_{L^p} \|\psi^4 u\|_{L^{\frac{sp}{sp-s-p}}} \\ &\quad + C(R-\rho)^{-1} \|\mathbf{V} - \mathbf{V}_{B(R)}\|_{L^s(B(R))} \|\psi^4 u\|_{L^{\frac{s}{s-1}}}^2 = I + II. \end{aligned}$$

Firstly, we estimate  $I$ . Since

$$3 \leq \frac{sp}{sp-s-p} \leq \frac{3p}{3-p},$$

Gagliardo–Nirenberg interpolation inequality and Hölder’s inequality imply that

$$\begin{aligned} \|\psi^4 u\|_{L^{\frac{sp}{sp-s-p}}} &\leq C \left( \|\psi^4 \nabla u\|_{L^p} + (R-\rho)^{-1} \|\psi^3 u\|_{L^p} \right)^{\frac{3p+3s-2sp}{s(2p-3)}} \|\psi^4 u\|_{L^3}^{\frac{4sp-3p-6s}{s(2p-3)}} \\ &\leq C \|\psi \nabla u\|_{L^p}^{\frac{3p+3s-2sp}{s(2p-3)}} \|\psi^3 u\|_{L^3}^{\frac{4sp-3p-6s}{s(2p-3)}} + C \left( (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{3p+3s-2sp}{s(2p-3)}} \|\psi^3 u\|_{L^3}^{\frac{4sp-3p-6s}{s(2p-3)}}. \end{aligned}$$

We insert it into  $I$  and use (1.6), and then we apply Young’s inequality twice. This yields

$$\begin{aligned} I &\leq CR^{\frac{3}{s} + \frac{1}{3} - \frac{5p-9}{s(2p-3)}} \|\psi \nabla u\|_{L^p}^{\frac{3p}{s(2p-3)}} \|\psi^3 u\|_{L^3}^{\frac{4sp-3p-6s}{s(2p-3)}} \\ &\quad + CR^{\frac{3}{s} + \frac{1}{3} - \frac{5p-9}{s(2p-3)}} \|\psi \nabla u\|_{L^p} \left( (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{3p+3s-2sp}{s(2p-3)}} \|\psi^3 u\|_{L^3}^{\frac{4sp-3p-6s}{s(2p-3)}} \\ &\leq CR \|\psi \nabla u\|_{L^p}^{\frac{9p}{2sp+3p-3s}} + CR \|\psi \nabla u\|_{L^p}^{\frac{3s(2p-3)}{2sp+3p-3s}} \left( (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{9p+9s-6sp}{2sp+3p-3s}} \\ &\quad + \frac{1}{4} \int |\psi^3 u|^3 dx \\ &\leq CR \|\psi \nabla u\|_{L^p}^{\frac{9p}{2sp+3p-3s}} + CR \left( (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{9p}{2sp+3p-3s}} + \frac{1}{4} \int |\psi^3 u|^3 dx. \quad (2.16) \end{aligned}$$

Secondly, we estimate  $II$ . First, let  $s \geq 3$ . If we apply Hölder’s inequality to  $II$  and use (1.6) along with Young’s inequality, it follows

$$\begin{aligned} II &\leq C(R-\rho)^{-1} R \left( \int_{B(R)} |\mathbf{V} - \mathbf{V}_{B(R)}|^s dx \right)^{\frac{1}{s}} \|\psi^3 u\|_{L^3}^2 \\ &\leq C(R-\rho)^{-3} R^4 \frac{3(5p-9)}{s(2p-3)} + \frac{1}{4} \int |\psi^3 u|^3 dx. \quad (2.17) \end{aligned}$$

(2.16) and (2.17) shows (2.14). Now we let  $s < 3$ . Since

$$\frac{3}{2} < \frac{s}{s-1} \leq \frac{3p}{6-p} < \frac{3p}{2(3-p)},$$

the interpolation inequality and Sobolev inequality imply that

$$\|\psi^4 u\|_{L^{\frac{s}{s-1}}}^2 \leq \|\psi^4 u\|_{L^{\frac{p(3-s)}{2s(2p-3)}}}^2 \|\psi^4 u\|_{L^{\frac{3p}{2(3-p)}}}^{\frac{5sp-3p-6s}{2s(2p-3)}} \|\psi^4 u\|_{L^{\frac{3}{2}}}^{\frac{3}{2}}$$



$$\begin{aligned}
&\leq \|\psi^4 u\|_{L^{\frac{3p}{3-p}}}^{\frac{p(3-s)}{s(2p-3)}} \|\psi^4 u\|_{L^3}^{\frac{5sp-3p-6s}{s(2p-3)}} \\
&\leq C \left( \|\psi^4 \nabla u\|_{L^p} + (R-\rho)^{-1} \|\psi^3 u\|_{L^p} \right)^{\frac{p(3-s)}{s(2p-3)}} \|\psi^4 u\|_{L^3}^{\frac{5sp-3p-6s}{s(2p-3)}} \\
&\leq C \|\psi \nabla u\|_{L^p}^{\frac{p(3-s)}{s(2p-3)}} \|\psi^3 u\|_{L^3}^{\frac{5sp-3p-6s}{s(2p-3)}} + C \left( (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{p(3-s)}{s(2p-3)}} \|\psi^3 u\|_{L^3}^{\frac{5sp-3p-6s}{s(2p-3)}}.
\end{aligned}$$

Similarly as above, we obtain

$$\begin{aligned}
II &\leq C(R-\rho)^{-1} R^{\frac{3}{s} + \frac{1}{3} - \frac{5p-9}{s(2p-3)}} \|\psi \nabla u\|_{L^p}^{\frac{p(3-s)}{s(2p-3)}} \|\psi^3 u\|_{L^3}^{\frac{5sp-3p-6s}{s(2p-3)}} \\
&+ C(R-\rho)^{-1} R^{\frac{3}{s} + \frac{1}{3} - \frac{5p-9}{s(2p-3)}} \left( (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{p(3-s)}{s(2p-3)}} \|\psi^3 u\|_{L^3}^{\frac{5sp-3p-6s}{s(2p-3)}} \\
&\leq C(R-\rho)^{-\frac{3s(2p-3)}{sp+3p-3s}} R^{\frac{2sp+3p-3s}{sp+3p-3s}} \|\psi \nabla u\|_{L^p}^{\frac{3p(3-s)}{sp+3p-3s}} \\
&+ C(R-\rho)^{-\frac{3s(2p-3)}{sp+3p-3s}} R^{\frac{2sp+3p-3s}{sp+3p-3s}} \left( (R-\rho)^{-1} \|\psi^p u\|_{L^p} \right)^{\frac{3p(3-s)}{sp+3p-3s}} + \frac{1}{4} \int |\psi^3 u|^3 dx. \quad (2.18)
\end{aligned}$$

By (2.16) and (2.18) we complete the proof.  $\square$

### 3 Proof of Theorem 1.4

We assume all conditions for Theorem 1.4 are fulfilled. Note that in the case of  $\frac{3p}{2p-3} < s < +\infty$ , we have by Jensen's inequality and (1.6)

$$\left( \frac{1}{|B(r)|} \int_{B(r)} |\mathbf{V} - \mathbf{V}_{B(r)}|^{\frac{3p}{2p-3}} dx \right)^{\frac{2p-3}{3p}} \leq \left( \frac{1}{|B(r)|} \int_{B(r)} |\mathbf{V} - \mathbf{V}_{B(r)}|^s dx \right)^{\frac{1}{s}} \leq C r^{\frac{3}{p} - \frac{4}{3}}.$$

This shows that we can reduce (1.6) to that case of  $s = \frac{3p}{2p-3}$ . Hence, in general we may restrict the range of  $s$  to

$$\frac{3p}{2(2p-3)} \leq s \leq \frac{3p}{2p-3}, \quad s > \frac{9-3p}{2p-3}.$$

Let  $1 < r < +\infty$  be arbitrarily chosen. We set  $r \leq \rho < R \leq 4r$  and  $\bar{R} = \frac{R+\rho}{2}$ . The first claim is that

$$\int_{\mathbb{R}^3} |\nabla u|^p dx < +\infty. \quad (3.19)$$

Let  $\zeta \in C_c^\infty(B(\bar{R}))$  be a radially non-increasing function such that  $0 \leq \zeta \leq 1$ ,  $\zeta = 1$  on  $B(\rho)$  and  $|\nabla \zeta| \leq C(R-\rho)^{-1}$  for some  $C > 0$ . If we insert  $\phi = \zeta^p$  into (1.5), then we have

$$\begin{aligned}
&\int_{B(\bar{R})} |\mathbf{D}(u)|^p \zeta^p dx = - \int_{B(\bar{R})} |\mathbf{D}(u)|^{p-2} \mathbf{D}(u) : u \otimes \nabla \zeta^p dx \\
&+ \frac{1}{2} \int_{B(\bar{R})} |u|^2 u \cdot \nabla \zeta^p dx + \int_{B(\bar{R})} (\pi - \pi_{B(\bar{R})}) u \cdot \nabla \zeta^p dx.
\end{aligned}$$

Hölder's inequality and Young's inequality imply that

$$\begin{aligned} \int_{B(\bar{R})} |\mathbf{D}(u)|^p \zeta^p dx &\leq C \int_{B(\bar{R})} |u|^p |\nabla \zeta|^p dx \\ + C \int_{B(\bar{R})} |u|^3 |\nabla \zeta| |\zeta|^{p-1} dx &+ C \int_{B(\bar{R})} |\pi - \pi_{B(\bar{R})}|^{\frac{3}{2}} |\nabla \zeta| |\zeta|^{p-1} dx. \end{aligned}$$

Employing Calderón-Zygmund's inequality, we obtain

$$\int |\nabla(u\zeta)|^p dx \leq C \int |\mathbf{D}(u)|^p \zeta^p dx + C \int |u|^p |\nabla \zeta|^p dx. \quad (3.20)$$

Using (3.20) along with (1.3), it follows

$$\begin{aligned} \int_{B(\rho)} |\nabla u|^p dx &\leq C(R - \rho)^{-p} \int_{B(\bar{R})} |u|^p dx + C(R - \rho)^{-1} \int_{B(\bar{R})} |u|^3 dx \\ &+ C(R - \rho)^{-1} R^{\frac{3}{2}(\frac{3}{p}-1)} \left( \int_{B(R)} |\nabla u|^p dx \right)^{\frac{3}{2}(1-\frac{1}{p})}. \end{aligned}$$

We consider  $\psi \in C_c^\infty(B(R))$  a radially non-increasing function satisfying  $0 \leq \psi \leq 1$ ,  $\psi = 1$  on  $B(\bar{R})$  and  $|\nabla \psi| \leq C(R - \rho)^{-1}$  for some  $C > 0$ . By the properties of  $\psi$  we have that

$$\begin{aligned} \int_{B(\rho)} |\nabla u|^p dx &\leq C(R - \rho)^{-p} \int_{B(R)} |\psi^p u|^p dx + C(R - \rho)^{-1} \int_{B(R)} |\psi^3 u|^3 dx \\ &+ C(R - \rho)^{-1} R^{\frac{3}{2}(\frac{3}{p}-1)} \left( \int_{B(R)} |\nabla u|^p dx \right)^{\frac{3}{2}(1-\frac{1}{p})} = I + II + III. \end{aligned} \quad (3.21)$$

Let  $\epsilon > 0$  be an arbitrary real number. Before calculating  $II$  first, we note that  $\psi$  satisfies the assumptions for Lemma 2.1 and Lemma 2.2. Observing that for  $s > \frac{9-3p}{2p-3}$

$$\frac{9p}{2sp + 3p - 3s} < p,$$

we may apply Young's inequality to (2.14) for  $s \geq 3$ . This yields

$$\begin{aligned} II &\leq C(\epsilon)(R - \rho)^{-\frac{2sp+3p-3s}{2sp-3s+3p-9}} R^{\frac{2sp+3p-3s}{2sp-3s+3p-9}} + I \\ &+ C(R - \rho)^{-4} R^{4-\frac{3(5p-9)}{s(2p-3)}} + \epsilon \int_{B(R)} |\psi \nabla u|^p dx. \end{aligned}$$

We continue estimating  $I$ . Since

$$4 - \frac{3(5p-9)}{s(2p-3)} < 4,$$

we get for  $R > 1$

$$II \leq C(\epsilon)(R - \rho)^{-\frac{2sp+3p-3s}{2sp-3s+3p-9}} R^{\frac{2sp+3p-3s}{2sp-3s+3p-9}} + C(R - \rho)^{-4} R^4 + \epsilon \int_{B(R)} |\psi \nabla u|^p dx.$$

In the case of  $s < 3$ , we see that for  $s > \frac{9-3p}{2p-3}$  it holds

$$\frac{3p(3-s)}{sp-3s+3p} < \frac{3p(3-s)}{s(2p-3)-3s+3p} < p.$$

Thus (2.15) and Young's inequality give

$$\begin{aligned} II &\leq C(\epsilon)(R - \rho)^{-\frac{2sp+3p-3s}{2sp-3s+3p-9}} R^{\frac{2sp+3p-3s}{2sp-3s+3p-9}} + I \\ &+ C(\epsilon)(R - \rho)^{-\frac{7sp+3p-12s}{sp+3p-9}} R^{\frac{2sp+3p-3s}{sp+3p-9}} + \epsilon \int_{B(R)} |\psi \nabla u|^p dx. \end{aligned}$$

Since

$$\frac{2sp+3p-3s}{sp+3p-9} < \frac{7sp+3p-12s}{sp+3p-9},$$

$R > 1$  shows that

$$\begin{aligned} II &\leq C(\epsilon)(R - \rho)^{-\frac{2sp+3p-3s}{2sp-3s+3p-9}} R^{\frac{2sp+3p-3s}{2sp-3s+3p-9}} + C(\epsilon)(R - \rho)^{-\frac{7sp+3p-12s}{sp+3p-9}} R^{\frac{7sp+3p-12s}{sp+3p-9}} \\ &+ \epsilon \int_{B(R)} |\psi \nabla u|^p dx. \end{aligned}$$

Hence, in both cases we obtain the following estimate

$$\begin{aligned} II &\leq C(\epsilon)(R - \rho)^{-\frac{7sp+3p-12s}{sp+3p-9}} R^{\frac{7sp+3p-12s}{sp+3p-9}} + C(\epsilon)(R - \rho)^{-\frac{2sp+3p-3s}{2sp-3s+3p-9}} R^{\frac{2sp+3p-3s}{2sp-3s+3p-9}} \\ &+ \epsilon \int_{B(R)} |\psi \nabla u|^p dx. \end{aligned} \quad (3.22)$$

Now we estimate  $I$ . If  $s \geq p$ , using (2.7) and Young's inequality, we see that

$$I \leq C(\epsilon)(R - \rho)^{-2p} R^{3+\frac{p}{3}-\frac{p(5p-9)}{s(2p-3)}} + \epsilon \int_{B(R)} |\psi \nabla u|^p dx.$$

Since

$$3 + \frac{p}{3} - \frac{p(5p-9)}{s(2p-3)} \leq 6 - \frac{4p}{3}$$

for  $s \leq \frac{3p}{2p-3}$ ,  $R > 1$  implies

$$\begin{aligned} I &\leq C(\epsilon)(R - \rho)^{-2p} R^{6-\frac{4p}{3}} + \epsilon \int_{B(R)} |\psi \nabla u|^p dx \\ &\leq C(\epsilon)(R - \rho)^{-2p} R^{2p} + \epsilon \int_{B(R)} |\psi \nabla u|^p dx. \end{aligned} \quad (3.23)$$

Notice that for  $s < p$ ,

$$0 < \frac{p(sp - 3s + 3p)}{2sp - 3s + 3p} < p$$

and

$$0 < \frac{p(3p - 3s)}{sp - 3s + 3p} < p.$$

By applying Young's inequality to (2.8) we get

$$I \leq C(\epsilon)(R - \rho)^{-2p+3-\frac{3p}{s}} R^{\frac{p}{3}+\frac{p^2}{s(2p-3)}} + \epsilon \int_{B(R)} |\psi \nabla u|^p dx. \quad (3.24)$$

Since

$$\frac{p}{3} + \frac{p^2}{s(2p-3)} < 2p - 3 + \frac{3p}{s} \leq 6p - 9 \quad (3.25)$$

for  $s \geq \frac{3p}{2(2p-3)}$ ,  $R > 1$  and  $R(R - \rho)^{-1} > 1$  imply that

$$I \leq C(\epsilon)(R - \rho)^{-(6p-9)} R^{6p-9} + \epsilon \int_{B(R)} |\psi \nabla u|^p dx.$$

Therefore, in each case we obtain

$$I \leq C(\epsilon)(R - \rho)^{-2p} R^{2p} + C(\epsilon)(R - \rho)^{-(6p-9)} R^{6p-9} + \epsilon \int_{B(R)} |\psi \nabla u|^p dx. \quad (3.26)$$

Applying Young's inequality to  $III$ , we find

$$III \leq C(\epsilon)(R - \rho)^{-\frac{2p}{3-p}} R^3 + \epsilon \int_{B(R)} |\psi \nabla u|^p dx.$$

By  $3 < \frac{2p}{3-p}$  and  $R > 1$ , it follows that

$$III \leq C(\epsilon)(R - \rho)^{-\frac{2p}{3-p}} R^{\frac{2p}{3-p}} + \epsilon \int_{B(R)} |\psi \nabla u|^p dx. \quad (3.27)$$

We define

$$\gamma := \max \left\{ \frac{7sp + 3p - 12s}{sp + 3p - 9}, \frac{2sp + 3p - 3s}{2sp - 3s + 3p - 9}, 2p, 6p - 9, \frac{2p}{3-p} \right\}.$$

From (3.22), (3.26), (3.27) and  $R(R - \rho)^{-1} > 1$  we deduce that

$$I + II + III \leq C(\epsilon)(R - \rho)^{-\gamma} R^\gamma + \epsilon \int_{B(R)} |\nabla u|^p dx.$$

Inserting this estimate into (3.21) and applying the iteration Lemma in [19, Lemma 3.1] for sufficiently small  $\epsilon > 0$ , we are led to

$$\int_{B(\rho)} |\nabla u|^p dx \leq C(R - \rho)^{-\gamma} R^\gamma.$$

By taking  $R = 2r$ ,  $\rho = r$  and passing  $r \rightarrow +\infty$ , we obtain (3.19).

Secondly we claim that

$$r^{-p} \int_{B(2r) \setminus B(r)} |u|^p dx = o(1) \quad \text{as} \quad r \rightarrow +\infty. \quad (3.28)$$

We consider a cut-off function  $\psi \in C_c^\infty(B(4r) \setminus B(\frac{r}{2}))$  satisfying  $0 \leq \psi \leq 1$ ,  $\psi = 1$  on  $B(2r) \setminus B(r)$  and  $|\nabla \psi| \leq Cr^{-1}$ . Then  $\psi$  satisfies the assumptions for Lemma 2.1 when  $R = 4r$  and  $\rho = r$ . Hence, in the case of  $s \geq p$  we use (3.23) to obtain

$$\begin{aligned} r^{-p} \int_{B(4r)} |\psi^p u|^p dx &\leq Cr^{6-\frac{10p}{3}} + C \int_{B(4r)} |\psi \nabla u|^p dx \\ &\leq Cr^{6-\frac{10p}{3}} + C \int_{B(4r) \setminus B(\frac{r}{2})} |\nabla u|^p dx. \end{aligned}$$

If  $s < p$ , using (3.24), we have that

$$r^{-p} \int_{B(4r)} |\psi^p u|^p dx \leq Cr^{\frac{p}{3} + \frac{p^2}{s(2p-3)} - (2p-3 + \frac{3p}{s})} + C \int_{B(4r) \setminus B(\frac{r}{2})} |\nabla u|^p dx.$$

Thus, observing (3.25) and (3.19), we obtain

$$r^{-p} \int_{B(4r)} |\psi^p u|^p dx = o(1) \quad \text{as} \quad r \rightarrow +\infty \quad (3.29)$$

which implies (3.28).

Next, we claim

$$r^{-1} \int_{B(2r) \setminus B(r)} |u|^3 dx = o(1) \quad \text{as} \quad r \rightarrow +\infty \quad (3.30)$$

and

$$r^{-1} \int_{B(r)} |u|^3 dx = O(1) \quad \text{as} \quad r \rightarrow +\infty. \quad (3.31)$$

We set the same function  $\psi \in C_c^\infty(B(4r) \setminus B(\frac{r}{2}))$  with  $R = 4r$  and  $\rho = r$ . For  $s \geq 3$  we can use (2.14) to infer

$$\begin{aligned} &r^{-1} \int_{B(4r)} |\psi^3 u|^3 dx \\ &\leq C \left( \int_{B(4r) \setminus B(\frac{r}{2})} |\nabla u|^p dx \right)^{\frac{9}{2sp+3p-3s}} + C \left( r^{-p} \int_{B(4r)} |\psi^p u|^p dx \right)^{\frac{9}{2sp+3p-3s}} + Cr^{-\frac{3(5p-9)}{s(2p-3)}}. \end{aligned}$$

In case  $s < 3$ , (2.15) gives that

$$r^{-1} \int_{B(4r)} |\psi^3 u|^3 dx$$

$$\begin{aligned}
&\leq C \left( \int_{B(4r) \setminus B(\frac{r}{2})} |\nabla u|^p dx \right)^{\frac{9}{2sp+3p-3s}} + C \left( r^{-p} \int_{B(4r)} |\psi^p u|^p dx \right)^{\frac{9}{2sp+3p-3s}} \\
&+ Cr^{\frac{s(9-5p)}{sp+3p-3s}} \left( \int_{B(4r) \setminus B(\frac{r}{2})} |\nabla u|^p dx \right)^{\frac{3(3-s)}{sp+3p-3s}} + Cr^{\frac{s(9-5p)}{sp+3p-3s}} \left( r^{-p} \int_{B(4r)} |\psi^p u|^p dx \right)^{\frac{3(3-s)}{sp+3p-3s}}.
\end{aligned}$$

In each case (3.19) and (3.29) imply (3.30).

To verify (3.31) we choose  $r_0 > 1$  arbitrarily and let  $r > r_0$ . Then for  $j \in \mathbb{N}$  satisfying  $r < 2^j r_0$ , we have

$$\begin{aligned}
r^{-1} \int_{B(r)} |u|^3 dx &\leq r^{-1} \int_{B(2r) \setminus B(r)} |u|^3 dx + \frac{1}{2} \left( \left( \frac{r}{2} \right)^{-1} \int_{B(r) \setminus B(\frac{r}{2})} |u|^3 dx \right) + \dots \\
&+ \frac{1}{2^j} \left( \left( \frac{r}{2^j} \right)^{-1} \int_{B(\frac{r}{2^{j-1}}) \setminus B(\frac{r}{2^j})} |u|^3 dx \right) + r^{-1} \int_{B(r_0)} |u|^3 dx \\
&\leq 2 \sup_{r_0 \leq \tilde{r} < +\infty} \left\{ \tilde{r}^{-1} \int_{B(2\tilde{r}) \setminus B(\tilde{r})} |u|^3 dx \right\} + r^{-1} \int_{B(r_0)} |u|^3 dx.
\end{aligned}$$

Then for sufficiently large  $r_0 > 1$ , (3.31) is obtained by (3.30) and  $u \in W^{1,p}(B(r_0)) \hookrightarrow L^3(B(r_0))$ .

**Proof of Theorem 1.4** Let  $\psi \in C_c^\infty(B(2r))$  satisfy  $0 \leq \psi \leq 1$ ,  $\psi = 1$  on  $B(r)$  and  $|\nabla \psi| \leq Cr^{-1}$ . We observe (1.5) with  $\psi^2 = \phi$ , and apply Hölder's inequality, Young's inequality and Calderón-Zygmund's inequality to get

$$\begin{aligned}
\int_{B(r)} |\nabla u|^p dx &\leq C \int_{B(2r)} |u|^p |\nabla \phi|^p dx + C \int_{B(2r)} |u|^3 |\nabla \phi| dx \\
&+ C \int_{B(2r)} |\pi - \pi_{B(2r)}| |u| |\nabla \phi| dx \\
&\leq Cr^{-p} \int_{B(2r) \setminus B(r)} |u|^p dx + Cr^{-1} \int_{B(2r) \setminus B(r)} |u|^3 dx \\
&+ Cr^{-1} \int_{B(2r) \setminus B(r)} |\pi - \pi_{B(2r)}| |u| dx = IV + V + VI.
\end{aligned}$$

The properties (3.28) and (3.30) directly shows that

$$IV + V \rightarrow 0 \quad \text{as} \quad r \rightarrow +\infty.$$

To estimate  $VI$  we use Hölder's inequality and (1.3) when  $s = \frac{3}{2}$ . This yields

$$VI \leq C \left( r^{-1} \int_{B(2r)} |\pi - \pi_{B(2r)}|^{\frac{3}{2}} dx \right)^{\frac{2}{3}} \left( r^{-1} \int_{B(2r) \setminus B(r)} |u|^3 dx \right)^{\frac{1}{3}}$$

$$\leq C \left( r^{-1} \int_{B(2r)} |\nabla u|^{\frac{3(p-1)}{2}} dx + r^{-1} \int_{B(2r)} |u|^3 dx \right)^{\frac{2}{3}} \left( r^{-1} \int_{B(2r) \setminus B(r)} |u|^3 dx \right)^{\frac{1}{3}}.$$

According to (3.31) it is sufficient to show that

$$r^{-1} \int_{B(2r)} |\nabla u|^{\frac{3(p-1)}{2}} dx = O(1) \quad \text{as} \quad r \rightarrow +\infty.$$

On the other hand, Hölder's inequality with  $\frac{3(p-1)}{2} < p$  implies that

$$\begin{aligned} r^{-1} \int_{B(2r)} |\nabla u|^{\frac{3(p-1)}{2}} dx &\leq r^{-1} |B(2r)|^{\frac{3}{2p} - \frac{1}{2}} \left( \int_{B(2r)} |\nabla u|^p dx \right)^{\frac{3(p-1)}{2p}} \\ &= r^{\frac{9}{2p} - \frac{5}{2}} \left( \int_{B(2r)} |\nabla u|^p dx \right)^{\frac{3(p-1)}{2p}}. \end{aligned}$$

This implies

$$VI \rightarrow 0 \quad \text{as} \quad r \rightarrow +\infty$$

and

$$\int_{B(r)} |\nabla u|^p = o(1) \quad \text{as} \quad r \rightarrow +\infty.$$

Accordingly,  $u \equiv \text{const}$  and by means of (3.30), we have that  $u \equiv 0$ . □

### Acknowledgements

Chae's research was partially supported by NRF grants 2021R1A2C1003234, and by the Chung-Ang University research grant in 2019. Wolf has been supported by NRF grants 2017R1E1A1A01074536. The authors declare that they have no conflict of interest.

## References

- [1] Wilkinson, W.L. *Non-Newtonian Fluids. Fluid Mechanics, Mixing and Heat Transfe*, Pergamon Press, London (1960).
- [2] Pokorný, M. *Cauchy problem for the non-Newtonian viscous incompressible fluid*, Appl. Math. **41** (3), 169–201 (1996).
- [3] Frehse, J., Málek, J., Steinhauer, M. *On analysis of steady flows of fluids with shear-dependent viscosity based on the Lipschitz truncation method*, SIAM J. Math. Anal. **34**(5), 1064–1083 (2003).

- [4] Galdi, G.P.: *An Introduction to the Mathematical Theory of the Navier–Stokes Equations: Steady-State Problems*, Springer Monographs in Mathematics, 2nd edn. Springer, New York (2011).
- [5] Chae, D., Wolf, J. *On Liouville Type Theorem for Stationary Non-Newtonian Fluid Equations*, J. Nonlinear Sci, **30**, no. 4, 1503-1517(2020).
- [6] Chae, D., Wolf, J. *On Liouville type theorem for the stationary Navier-Stokes equations*, Cal. Var. PDE **58**(3), 111(11pp) (2019).
- [7] Chae, D., Yoneda, T. *On the Liouville theorem for the stationary Navier-Stokes equations in a critical space*, J. Math. Anal. Appl. **405** (2), 706–710 (2013).
- [8] Chae, D. *Liouville-type theorem for the forced Euler equations and the Navier-Stokes equations*, Commun. Math. Phys. **326**, 37-48 (2014).
- [9] Chae, D., Wolf, J. *On Liouville type theorems for the steady Navier-Stokes equations in  $\mathbb{R}^3$* , J. Differ. Equ. **261** , 5541-5560 (2016).
- [10] Chamorro, D., Jarrin, O., Lemarié-Rieusset, P.-G. *Some Liouville theorems for stationary Navier-Stokes equations in Lebesgue and Morrey spaces*, C.R. Math. Acad. Sci. Paris. **357**(2), 175-179 (2019).
- [11] Gilbarg, D., Weinberger, H.F., *Asymptotic properties of steady plane solutions of the Navier-Stokes equations with bounded Dirichlet integral*, Ann. Sc. Norm. Super. Pisa **4**(5), 381-404 (1978).
- [12] Koch, G., Nadirashvili, N., Seregin, G., Šverák, V. *Liouville theorems for the Navier-Stokes equations and applications*, Acta Math. **203**, 83-105 (2009).
- [13] Korobkov, M., Pileckas, K., Russo, R. *The Liouville theorem for the steady-state Navier-Stokes problem for axially symmetric 3D solutions in absence of swirl*, J. Math. Fluid Mech. **17** (2), 287-293 (2015).
- [14] Kozono, H., Terasawa, Y., Wakasugi, Y. *A remark on Liouville-type theorems for the stationary Navier-Stokes equations in three space dimensions*, J. Funct. Anal. **272**, 804-818 (2017).
- [15] Seregin, G. *Liouville type theorem for stationary Navier–Stokes equations*, Non-linearity **29**, 2191-2195 (2016).
- [16] Seregin, G. *Remarks on Liouville type theorems for steady-state Navier-Stokes equations* Algebra Anal. **30** (2), 238-248 (2018).



- [17] Seregin, G., Wang, W.: Sufficient conditions on Liouville type theorems for the 3D steady Navier-Stokes equations. *Algebra i Analiz.* **31**(2), 269-278 (2019).
- [18] Jin, B.J., Kang, K. *Liouville Theorem for the Steady-State Non-Newtonian Navier-Stokes Equations in Two Dimensions*, *J. Math. Fluid Mech.* **16**, 275-292 (2014).
- [19] Giaquinta, M. *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*, *Annals of Mathematics Studies* No. 105. Princeton University Press, Princeton (1983).