

HOLOMORPHIC FUNCTIONS OF FINITE ORDER GENERATED BY DIRICHLET SERIES

ANDREAS DEFANT AND INGO SCHOOLMANN

ABSTRACT. Given a frequency $\lambda = (\lambda_n)$ and $\ell \geq 0$, we introduce the scale of Banach spaces $H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$ of holomorphic functions f on the open right half-plane $[\operatorname{Re} > 0]$, which satisfy (A) the growth condition $|f(s)| = O((1 + |s|)^{\ell})$, and (B) have a Riesz germ, i.e. on some open subset and for some $m \geq 0$ the function f coincides with the pointwise limit (as $x \rightarrow \infty$) of the so-called (λ, m) -Riesz means $\sum_{\lambda_n < x} a_n e^{-\lambda_n s} (1 - \frac{\lambda_n}{x})^m$, $x > 0$ of some λ -Dirichlet series $\sum a_n e^{-\lambda_n s}$. Reformulated in our terminology, an important result of M. Riesz shows that in this case the function f for every $k > \ell$ is the pointwise limit of the (λ, k) -Riesz means of D on $[\operatorname{Re} > 0]$.

Our main contribution is an extension – showing that ‘after translation’ every bounded set in $H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$ is uniformly approximable by all its (λ, k) -Riesz means of order $k > \ell$. This follows from an appropriate maximal theorem, which in fact turns out to be at the very heart of a seemingly interesting structure theory of the Banach spaces $H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$. One of the many consequences is that $H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$ basically consists of those holomorphic functions on $[\operatorname{Re} > 0]$, which have a Riesz germ and are of finite uniform order ℓ on $[\operatorname{Re} > 0]$.

To establish all this and more, we need to reorganize (and to improve) various aspects and keystones of the classical theory of Riesz summability of general Dirichlet series as invented by Hardy and M. Riesz.

CONTENTS

1. Introduction	2
2. Riesz summation revisited	7
2.1. Ordinary summation	8
2.2. Riesz means	11
2.3. Integral representation	13
2.4. Perron formula	16
2.5. Bohr-Cahen formula	18
3. A new scale of Banach spaces	18
3.1. Ordinary and power case	19
3.2. Perron formula – a variant	21
3.3. A maximal inequality for Riesz means	22
3.4. Uniform Riesz approximation	23
3.5. Theorem 41 of M. Riesz revisited	24
3.6. The bounded case $\ell = 0$	25
3.7. Boundedness of coefficient functionals	25
3.8. A Montel theorem	26

3.9.	Completeness	27
3.10.	Behavior far left	28
3.11.	Behavior far right	29
3.12.	Finite order	30
3.13.	Equivalence	31
4.	Appendix	36
4.1.	Integral forms	36
4.2.	Changing orders	38
4.3.	Estimates	42
4.4.	The technical heart	45
4.5.	Proof of Theorem 2.9	49
4.6.	Proof of Theorem 2.16	50
4.7.	Proof of Theorem 2.7	50
4.8.	Proof of Theorem 2.8	54
	References	58

1. Introduction

Recently, the theory of ordinary Dirichlet series $\sum a_n n^{-s} = \sum a_n e^{-s \log n}$ and that of general Dirichlet series $\sum a_n e^{-\lambda_n s}$, where the frequency $\lambda = (\lambda_n)$ is a strictly increasing, non-negative real sequence, saw a remarkable renaissance.

General Dirichlet series $\sum a_n e^{-\lambda_n s}$ converge on maximal half-planes $[\operatorname{Re} > \sigma]$ in \mathbb{C} , where they define holomorphic functions f , and even though in concrete situations these half-planes might be small, the holomorphic functions f often extend to holomorphic functions on larger half-planes.

The problem of determining whether a holomorphic f , defined on some half-plane, is or is not representable in the form of a λ -Dirichlet series $\sum a_n e^{-\lambda_n s}$, is in general difficult.

An important criterion is that the limit function of a λ -Dirichlet series has finite order in each closed half-plane of its domain of convergence. Moreover, if this limit function can be holomorphically extended to a larger half-plane, it may happen that its continuation still has finite order in a larger domain.

Recall that, given a holomorphic function $f : [\operatorname{Re} > \sigma_0] \rightarrow \mathbb{C}$, $\sigma_0 \in \mathbb{R}$, its order $\mu_f(\sigma) \in [\sigma_0, \infty]$ on the line $[\operatorname{Re} = \sigma]$, $\sigma > \sigma_0$ is the infimum over all $\ell \in [0, \infty]$ such that

$$(1) \quad \exists C, t_0 > 0 \forall |t| > t_0: |f(\sigma + it)| \leq C|t|^\ell.$$

2010 *Mathematics Subject Classification*: Primary 43A17, Secondary 30B50, 43A50

Key words and phrases: general Dirichlet series, finite order, Riesz summability, almost everywhere convergence, Hardy spaces.

The function f is said to be of finite order on $[\operatorname{Re} > \sigma_0]$ whenever $\mu_f < \infty$ for all $\sigma > \sigma_0$, and in this case $\mu_f :]\sigma_0, \infty[\rightarrow \mathbb{R}_{\geq 0}$ is continuous and convex.

Similarly, the uniform order $\nu_f \in [0, \infty]$ of f on the half-plane $[\operatorname{Re} > \sigma_0]$ is given by the infimum over all $\ell \in [0, \infty]$ such that the above condition holds uniformly, i.e.,

$$(2) \quad \exists C, t_0 > 0 \forall \sigma > \sigma_0, |t| > t_0: |f(\sigma + it)| \leq C|t|^\ell.$$

Moreover, the function f is said to have finite uniform order on $[\operatorname{Re} > \sigma_0]$ whenever $\nu_f < \infty$.

To see a famous example, denote by $\zeta : \mathbb{C} \setminus \{1\} \rightarrow \mathbb{C}$ the zeta-function, which is holomorphic with a simple pole in $s = 1$, and which on $[\operatorname{Re} > 1]$ is the pointwise limit of the zeta-Dirichlet series $\sum n^{-s}$. The famous Lindelöf conjecture asks whether the order of ζ on the critical line $[\operatorname{Re} = \frac{1}{2}]$ equals 0. Equivalently, consider the entire function

$$\eta : \mathbb{C} \rightarrow \mathbb{C}, \quad \eta(s) = (1 - 2^{1-s})\zeta(s),$$

which on $[\operatorname{Re} > 0]$ is nothing else than the pointwise limit of the η -Dirichlet series $\sum (-1)^{n+1} n^{-s}$. It is known that $\mu_\eta(\sigma) = \frac{1}{2} - \sigma$ for $\sigma < 0$ and $\mu_\eta(\sigma) = 0$ for $\sigma > 1$. Hence Lindelöf's conjecture is equivalent to the question whether or not we have that

$$\mu_\eta(\sigma) = \begin{cases} \frac{1}{2} - \sigma & 0 \leq \sigma < \frac{1}{2} \\ 0 & \frac{1}{2} < \sigma \leq 1. \end{cases}$$

The main aim here is to study holomorphic functions of finite uniform order generated by Dirichlet series from the point of view of functional analysis – and our main inspiration comes from the classical monograph [13] of Hardy and Riesz on so-called Riesz summation of general Dirichlet series.

Given a frequency λ and a holomorphic function $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$, we say that the λ -Dirichlet series $D = \sum a_n(D)e^{-\lambda_n s}$ is a λ -Riesz germ of f whenever D for some $m \geq 0$ and some $\sigma \in \mathbb{R}$ satisfies that

$$f(s) = \lim_{x \rightarrow \infty} \sum_{x < \lambda_n} a_n(D)e^{-\lambda_n s} \left(1 - \frac{\lambda_n}{x}\right)^m \quad \text{for all } s \in [\operatorname{Re} > \sigma].$$

The trigonometric polynomials

$$R_x^{\lambda, m}(f)(s) = \sum_{\lambda_n < x} a_n(D)e^{-\lambda_n s} \left(1 - \frac{\lambda_n}{x}\right)^m, \quad x > 0$$

are said to be the (λ, m) -Riesz means of f of order m .

An important fact will be that such λ -Riesz germs D of f , whenever they exist (!), are unique (Corollary 2.15) – and so in this situation the coefficients $a_n(f) := a_n(D)$ uniquely determine the function f .

We study a graduated scale of Banach spaces that consist of holomorphic functions on the right half-plane, which are of finite uniform order on $[\operatorname{Re} > 0]$ and

are generated by λ -Riesz germs. More precisely, given a frequency $\lambda = (\lambda_n)$ and $\ell \geq 0$, the linear space

$$(3) \quad H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$$

consists of all holomorphic functions $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$, which are generated by a λ -Riesz germ and satisfy the growth condition

$$(4) \quad \|f\|_{\infty, \ell} = \sup_{\operatorname{Re} s > 0} \frac{|f(s)|}{(1 + |s|)^{\ell}} < \infty.$$

This defines a scale of Banach spaces (a non-trivial fact from Theorem 3.16),

$$(H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0], \|\cdot\|_{\infty, \ell})_{\ell \geq 0},$$

which is intimately connected with the notion of finite uniform order (Theorem 3.21):

- Each $f \in H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$ has finite uniform order $\nu_f \leq \ell$ on $[\operatorname{Re} > 0]$, and conversely,
- if f is a holomorphic function on $[\operatorname{Re} > 0]$, which is generated by a λ -Riesz germ and is of finite uniform order ν_f on $[\operatorname{Re} > 0]$, then for every $k > \nu_f$ all translations $f_{\sigma} = f(\sigma + \cdot) \in H_{\infty, k}^{\lambda}[\operatorname{Re} > 0]$, $\sigma > 0$.

More background. Mainly inspired by the classical monograph [13] of Hardy and M. Riesz from 1915, the recent works [2], [6], [8], [9], [10], [17], and [20] suggest a modern study of general Dirichlet series $D = \sum a_n e^{-\lambda_n s}$. Whereas the dominant tool of the early days of this theory was complex analysis, the idea now is to implement modern techniques like functional analysis, Fourier analysis, or abstract harmonic analysis on compact abelian groups.

An important object of study are Hardy spaces $\mathcal{H}_p(\lambda)$, $1 \leq p \leq \infty$, which may be defined as follows:

Let G be a compact abelian group, and $\beta : (\mathbb{R}, +) \rightarrow G$ a continuous homomorphism with dense range. Then (G, β) is said to be λ -Dirichlet group, provided, for all $n \in \mathbb{N}$, there exists $h_{\lambda_n} \in \widehat{G}$ such that $h_{\lambda_n} \circ \beta = e^{-i\lambda_n}$. The Banach space $\mathcal{H}_p(\lambda)$ then consists of all Dirichlet series $D = \sum a_n e^{-\lambda_n s}$ for which there is $f \in L_p(G)$ such that $\operatorname{supp} \widehat{f} \subset \{h_{\lambda_n} : n \in \mathbb{N}\}$ and $\widehat{f}(h_{\lambda_n}) = a_n$ for all $n \in \mathbb{N}$, endowed with the norm $\|D\|_p = \|f\|_p$. It is important to note that for every frequency λ there always exist a λ -Dirichlet group, and that $\mathcal{H}_p(\lambda)$ does not depend on the choice of this group. For ordinary Dirichlet series these spaces have been introduced in [1] and [14], and this has in fact caused a fruitful renaissance of the analysis of such series. Within general Dirichlet series various aspects of $\mathcal{H}_p(\lambda)$ have been studied in [2], [6], [8], [9], [10], and [20].

The particular case $p = \infty$ is of special interest, since then $\mathcal{H}_{\infty}(\lambda)$ may be described in terms of holomorphic functions on the right half-plane, and in fact we are only interested trying to develop this case further here.

Recall that $H_{\infty}^{\lambda}[\operatorname{Re} > 0]$ (as defined in [9]) denotes the linear space of all holomorphic and bounded functions $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$, which are uniformly almost

periodic function on all vertical lines $[\operatorname{Re} = \sigma]$ (or equivalently, some line $[\operatorname{Re} = \sigma]$) with Bohr coefficients

$$a_x(f) = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T f(\sigma + it) e^{-(\sigma + it)x} dt, \quad x > 0.$$

supported in $\{\lambda_n \mid n \in \mathbb{N}\}$. Note that here the integrals are independent of the choice of σ . Together with the sup norm on the right half-plane, $H_\infty^\lambda[\operatorname{Re} > 0]$ forms a Banach space, and by [9, Theorem 2.16] there is a coefficient preserving isometric linear bijection identifying $\mathcal{H}_\infty(\lambda)$ and $H_\infty^\lambda[\operatorname{Re} > 0]$,

$$(5) \quad \mathcal{H}_\infty(\lambda) = H_\infty^\lambda[\operatorname{Re} > 0].$$

The most fundamental theorem of the theory of almost periodic functions on \mathbb{R} is due to Bohr, and states that a function $f : \mathbb{R} \rightarrow \mathbb{C}$ is uniformly almost periodic if and only if it is uniformly approximable by trigonometric polynomials of the form $p(t) = \sum_{k=1}^n \alpha_k e^{-i\alpha_k t}$ with $\alpha_k \in \mathbb{R}$, $1 \leq k \leq n$. We refer to [3] for all needed information on (uniformly) almost periodic functions.

For bounded, holomorphic functions on half-planes the following result from [9, Corollary 2.19] (see also [11, Lemma 3.3] for a may be more elementary proof) should be viewed as an analogue of Bohr's approximation theorem: If $f \in H_\infty^\lambda[\operatorname{Re} > 0]$, then for every $k > 0$ and $s \in [\operatorname{Re} > 0]$,

$$(6) \quad f(s) = \lim_{x \rightarrow \infty} \sum_{\lambda_n < x} a_{\lambda_n}(f) e^{-\lambda_n s} \left(1 - \frac{\lambda_n}{x}\right)^k,$$

and the convergence is even uniform on every half-plane $[\operatorname{Re} > \sigma]$, $\sigma > 0$.

In Corollary 3.11 we characterize $H_\infty^\lambda[\operatorname{Re} > 0]$ in terms of our new setting showing that the canonical embedding from $H_{\infty,0}^\lambda[\operatorname{Re} > 0]$ into $H_\infty^\lambda[\operatorname{Re} > 0]$ is in fact isometric (obvious), onto and coefficient preserving,

$$(7) \quad H_\infty^\lambda[\operatorname{Re} > 0] = H_{\infty,0}^\lambda[\operatorname{Re} > 0].$$

But then in view of Bohr's approximation theorem for uniformly almost periodic functions on \mathbb{R} , and in view of the approximation theorem from (6) for almost periodic holomorphic functions on $[\operatorname{Re} > 0]$, the following question arises:

Given $\ell > 0$, to which extend is it possible to approximate any function $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ by (λ, k) -Riesz means

$$R_x^{\lambda,k}(D)(s) = \sum_{\lambda_n < x} a_n(f) e^{-\lambda_n s} \left(1 - \frac{\lambda_n}{x}\right)^k, \quad x > 0$$

pointwise on $[\operatorname{Re} > 0]$ or even in the norm of $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$? More precisely, is there some $k > 0$, and if yes for which set of k 's, is it possible to approximate any $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ pointwise on $[\operatorname{Re} > 0]$ or in the norm of $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ by their (λ, k) -Riesz means?

A first answer to this question is given by an important theorem of M. Riesz, first published in [13, Theorem 41], which inspired the definition from (3) as well as the whole article. It basically shows that in view of (7) the validity of (6)

exceeds the class of functions in $H_\infty^\lambda[\operatorname{Re} > 0]$ considerably, and in our terminology it reads as follows:

Theorem 1.1. *Let λ be any frequency and $\ell \geq 0$. Then every $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ is (λ, k) -Riesz summable on $[\operatorname{Re} > 0]$ for every $k > \ell$, i.e.*

$$f(s) = \lim_{x \rightarrow \infty} R_x^{\lambda,k}(f)(s), \quad s \in [\operatorname{Re} > 0].$$

We reprove this result in Corollary 3.9, where we also carefully compare our formulation with the original formulation given in [13, Theorem 41].

Main results. Theorem 1.1 is a pointwise approximation theorem of holomorphic functions on half-planes by Riesz means, and we are going to extend various of its different aspects – including equivalent reformulations and sufficient conditions on λ for the validity of Theorem 1.1 when $k = \ell > 0$ (proved in Theorem 3.26). But in general, it fails for $k = \ell = 0$; see the forthcoming Theorems 3.23 and 3.24.

Since we recover this result of Riesz in terms of the Banach spaces $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$, we in Theorem 3.6 are able to isolate the corresponding maximal inequality:

Given a frequency λ and $k > \ell \geq 0$, there is a constant $C = C(k, \ell, \lambda)$ such that for all $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$

$$\sup_{x > 0} \|R_x^{\lambda,k}(f)\|_{\infty,\ell} \leq C \|f\|_{\infty,\ell}.$$

This then in Theorem 3.7 leads to a uniform approximation theorem of bounded sets in $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ under Riesz summation:

For every bounded set $M \subset H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$, $k > \ell$ and $\varepsilon, u > 0$ there exists $x_0 > 0$ such that

$$\sup_{x > x_0} \sup_{f \in M} \|f(u + \cdot) - R_x^{\lambda,k}(f)(u + \cdot)\|_{\infty,\ell} \leq \varepsilon.$$

Indeed, the preceding two facts form the main body of this article, since they turn out to be the driving forces for most of the structure theory on the Banach spaces $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ we intend to present.

This applies both, to the results on $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ that we have already mentioned here (like e.g. the facts that these spaces are complete and intimately linked with the notion of 'finite order'), but also to a few others to come (like e.g. a Montel theorem for $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ from Theorem 3.15).

Studying problems for general Dirichlet series $\sum a_n e^{-\lambda_n s}$, the very first orientation usually comes from

- the frequency $\lambda = (n)$, which after the substitution $e^{-s} = z$ generates power series $\sum a_n z^n$,
- and the frequency $\lambda = (\log n)$ generating ordinary Dirichlet series $\sum a_n n^{-s}$.

In Proposition 3.3 we show that the scale $H_{\infty,\ell}^{(n)}[\operatorname{Re} > 0]$, $\ell \geq 0$, collapses:

$$H_{\infty}^{(n)}[\operatorname{Re} > 0] = H_{\infty,\ell}^{(n)}[\operatorname{Re} > 0].$$

But, as shown in Proposition 3.4 the situation for the frequency $\lambda = (\log n)$ is very different. For example $\eta \notin H_{\infty,0}^{(\log n)}[\operatorname{Re} > 0]$, but $\eta \in H_{\infty,\ell}^{(\log n)}[\operatorname{Re} > 0]$ for all

$\ell > \frac{1}{2}$. In Proposition 3.4 we show that $f \in H_{\infty,1}^{(\log n)}[\operatorname{Re} > 0]$ if and only if the growth condition (4) holds, and moreover there exists an ordinary Dirichlet series which on $[\operatorname{Re} > 0]$ is Cesàro summable with limit f .

Twofold interest. The proof of our main results and all its consequences for the structure theory of our new scale of Banach spaces $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ is very much inspired by the theory of Riesz summation from [13].

Hardy and Riesz write in the preface of their book ‘...*The task of condensing any account of so extensive a theory into the compass of one of these tracts has proved an exceedingly difficult one. Many important theorems are stated without proof, and many details are left to the reader.*’

In fact, at several occasions we even need improvements of the results from [13], and this has prompted us to add an appendix devoted to a detailed, self contained, differently organized, and in parts more modern presentation of several key stones of the theory of Hardy and Riesz. We hope that apart from our new results on Banach spaces of holomorphic functions of finite order generated by Dirichlet series, this creates an additional benefit of our work.

Outlook. The classical Carleson-Hunt result implies that the Taylor series of bounded holomorphic functions on the open unit disk converge almost everywhere on the boundary. In contrast, there exists an ordinary Dirichlet series, which on the open right half-plane converges to a bounded, holomorphic function – but diverges at each point of the imaginary line, although its limit function extends continuously to the closed right half plane. As a continuation of the present article we in the forthcoming work [12] study Riesz summability of functions in the spaces $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$, $\ell \geq 0$, on the imaginary axis. Again our inspiration comes from an important, surprisingly deep result of M. Riesz published (without proof) in [13, Theorem 42].

Structure of the article. In the Sections 2 and 4 we develop parts of the summation theory on Riesz means, which provides us with an important integral formula of the limit function of λ -Dirichlet series under Riesz summation, which then in a second step by ‘Laplace inversion’ leads to the so-called Perron formulas for the summatory function of these series. Then Section 3 covers the main results of this article including the basic properties of our new scale of Banach spaces $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$, $\ell \geq 0$.

2. Riesz summation revisited

The aim of this first section is to recall the most crucial ingredients of the theory of Riesz summation of general Dirichlet series. We isolate three fundamental pillars on which the theory of Riesz-summability of general Dirichlet series $D = \sum a_n e^{-\lambda_n s}$ essentially rests : the so-called ‘theorems of consistency’ of such series as presented in Theorem 2.7, their integral representation from Theorem 2.9, and the Bohr-Cahen formula for their abscissas of convergence formulated in Theorem 2.16. We offer full self contained proof of these results in our appendix from Section 4.

All important consequences, as e.g. Perron's formula from Theorem 2.13 are already given in this introductory section. Fixing a frequency λ , for simplicity the collection of all such λ -Dirichlet series $\sum a_n e^{-\lambda n s}$ is denoted by $\mathcal{D}(\lambda)$.

2.1. Ordinary summation. Every presentation of the classical convergence theory of general Dirichlet series necessarily starts with the following theorem, and for the sake of completeness we repeat the standard proof based on Abel summation (see e.g. [13, Theorem 1]).

Theorem 2.1. *Let $D = \sum a_n e^{-\lambda n s}$ be a λ -Dirichlet series which converges at $s_0 \in [\operatorname{Re} \geq 0]$. Then D converges uniformly on each cone $|\arg(s - s_0)| \leq \gamma < \frac{\pi}{2}$.*

Proof. Assume without loss of generality that $s_0 = 0$, and fix some $\gamma < \frac{\pi}{2}$ and $0 \neq s \in \mathbb{C}$ with $|\arg(s)| \leq \gamma$. Then by standard Abel summation for all $m < n$

$$(8) \quad \sum_{\nu=m}^n a_\nu e^{-\lambda \nu s} = \sum_{\nu=m}^{n-1} \left(\sum_{\ell=m}^{\nu} a_\ell \right) (e^{-\lambda \nu s} - e^{-\lambda(\nu+1)s}) + \left(\sum_{\nu=m}^n a_\nu \right) e^{-\lambda n s}.$$

But with $\operatorname{Re} s = \sigma$

$$|e^{-\lambda \nu s} - e^{-\lambda(\nu+1)s}| \leq \frac{|s|}{\operatorname{Re} s} (e^{-\lambda \nu \sigma} - e^{-\lambda(\nu+1)\sigma}) \leq \sec \gamma (e^{-\lambda \nu \sigma} - e^{-\lambda(\nu+1)\sigma}),$$

and hence, for m and n large, we have that

$$\left| \sum_{\nu=n}^m a_\nu e^{-\lambda \nu s} \right| \leq \varepsilon$$

uniformly on the cone $|\arg(s)| \leq \gamma$. □

Theorem 2.1 suggests the following basic definition. For each λ -Dirichlet series $D = \sum a_n e^{-\lambda n s}$ the number

$$\sigma_c^\lambda(D) = \inf \{ \sigma \in \mathbb{R} : D \text{ converges at } \sigma \} \in \mathbb{R} \cup \{\pm\infty\}$$

is called abscissa of convergence of D . Obviously, D converges on $[\operatorname{Re} > \sigma_c^\lambda(D)]$ and diverges on $[\operatorname{Re} < \sigma_c^\lambda(D)]$.

The following integral formula describes the limit function f of a λ -Dirichlet series $D = \sum a_n e^{-\lambda n s}$ on the half-plane $[\operatorname{Re} > \sigma_c^\lambda(D)]$ of convergence in terms of the Laplace transform of the so-called summatory function which for $s \in \mathbb{C}$ and $t \geq 0$ is given by

$$S_t^\lambda(D)(s) = \sum_{\lambda_n < t} a_n e^{-\lambda n s}.$$

Our proof is inspired by Helson [17, (2.3)], and serves as a model for the proof of the much more involved variant Theorem 2.9 on Riesz summation.

Theorem 2.2. *Let $D = \sum a_n e^{-\lambda n s}$ be convergent at $s_0 \in [\operatorname{Re} \geq 0]$. Then D converges on $[\operatorname{Re} > \operatorname{Re} s_0]$, and the limit function*

$$(9) \quad f : [\operatorname{Re} > \operatorname{Re} s_0] \rightarrow \mathbb{C}, \quad s \mapsto \lim_{x \rightarrow \infty} \sum_{\lambda_n < x} a_n e^{-\lambda n s}$$

is holomorphic and satisfies for all $s \in [\operatorname{Re} > \operatorname{Re} s_0]$

$$\frac{f(s)}{s} = \int_0^\infty e^{-st} \sum_{\lambda_n < t} a_n dt.$$

Moreover, the convergence in (9) is uniform on each cone $|\arg(s - s_0)| \leq \gamma < \frac{\pi}{2}$.

Note that the preceding theorem shows that the function $s \mapsto f(s)/s$ is nothing else than the Laplace transform of the summatory function $t \mapsto S_t(D)(0)$.

We prepare the proof with two simple lemmas.

Lemma 2.3. *Let $D \in \mathcal{D}(\lambda)$. Then for all $s, w \in \mathbb{C}$ and all $x \geq 0$*

$$S_x^\lambda(D)(s+w) = S_x^\lambda(D)(w)e^{-sx} - \int_0^x S_t^\lambda(D)(w)se^{-st}dt.$$

Proof. Since $S_x^\lambda(D)(s+w) = S_x^\lambda(D_w)(s)$, we concentrate on the case $w = 0$; recall that here D_w stands for the translation of $D = \sum a_n e^{-\lambda_n s}$ about w , i.e. $D_w = \sum a_n e^{-\lambda_n w} e^{-\lambda_n s}$. Then

$$S_x^\lambda(D)(s) = \int_0^x e^{-ts} dS_t^\lambda(D)(0) = \int_0^x e^{-ts} (S_\bullet^\lambda(D)(0))'(t)dt,$$

where the first integral is a Stieltjes integral. Obviously, summatory functions are of bounded variation, hence by partial integration (see e.g. Helson [16, Appendix])

$$S_x^\lambda(D)(s) = e^{-xs} S_x^\lambda(D)(0) - \int_0^x se^{-ts} S_t^\lambda(D)(0)dt. \quad \square$$

Lemma 2.4. *Let $D \in \mathcal{D}(\lambda)$, and $s_0, s \in \mathbb{C}$. Then for all $x > 0$*

$$|S_x^\lambda(D)(s)| \leq (1 + |s_0|)e^{x \operatorname{Re} s_0} \sup_{y \leq x} |S_y^\lambda(D)(s_0 + s)|.$$

Proof. Since $S_x^\lambda(D)(s) = S_x^\lambda(D_s)(0)$, we may assume that $s = 0$. By Lemma 2.3

$$(10) \quad S_x^\lambda(D)(0) = S_x^\lambda(D)(-s_0 + s_0) = e^{s_0 x} S_x^\lambda(D)(s_0) - \int_0^x S_t^\lambda(D)(s_0)(-s_0)e^{s_0 t} dt.$$

Then for the first summand we have

$$|e^{s_0 x} S_x^\lambda(D)(s_0)| \leq e^{x \operatorname{Re} s_0} \sup_{y \leq x} |S_y^\lambda(D)(s_0)|,$$

and for the second

$$\left| \int_0^x S_t^\lambda(D)(s_0) s_0 e^{s_0 t} dt \right| \leq |s_0| e^{x \operatorname{Re} s_0} \sup_{y \leq x} |S_y^\lambda(D)(s_0)|. \quad \square$$

Proof of Theorem 2.2. Deduce first by Lemma 2.4 (with $s = 0$) and the assumption (D converges in s_0) that there is a constant $C = C(s_0) > 0$ such that for all $x > 0$

$$(11) \quad |S_x^\lambda(D)(0)| \leq C e^{x \operatorname{Re} s_0}.$$

Now fix some $s \in [\operatorname{Re} > \operatorname{Re} s_0]$. Using Lemma 2.3, for all $x > 0$

$$(12) \quad S_x^\lambda(D)(s) = S_x^\lambda(D)(0)e^{-sx} - \int_0^x S_t^\lambda(D)(0)se^{-st} dt.$$

By (11) for all $x > 0$

$$e^{-x \operatorname{Re} s} |S_x^\lambda(D)(0)| \leq C e^{x(\operatorname{Re} s_0 - \operatorname{Re} s)},$$

which converges to 0 whenever $x \rightarrow \infty$. Hence it remains to show that

$$\lim_{x \rightarrow \infty} \int_0^x S_t^\lambda(D)(0)e^{-st} dt = \int_0^\infty S_t^\lambda(D)(0)e^{-st} dt.$$

But this follows from the dominated convergence theorem – indeed, for $x > 0$ we conclude from another application of (11) that for all $t > 0$

$$|S_t^\lambda(D)(0)e^{-st} \chi_{[0,x]}(t)| \leq C e^{t \operatorname{Re} s_0} e^{-t \operatorname{Re} s} \chi_{[0,x]}(t) \leq C e^{t(\operatorname{Re} s_0 - \operatorname{Re} s)}.$$

The last statement of the theorem is then a consequence of Theorem 2.1. \square

The following formula, which we for historical reasons call Bohr-Cahen formula, is taken from [13, Theorem 7].

Corollary 2.5. *Let $D \in \mathcal{D}(\lambda)$. Then*

$$\sigma_c^\lambda(D) \leq \limsup_{x \rightarrow \infty} \frac{\log(|S_x^\lambda(D)(0)|)}{x},$$

with equality whenever $\sigma_c^\lambda(D)$ is non-negative.

Proof. We denote the limes superior by L , assume that it is finite, and choose some $\sigma_0 > L$. Then there is a constant $C > 0$ such that for all $x > 0$ we have that

$$|S_x^\lambda(D)(0)| \leq C e^{\sigma_0 x}.$$

Now, starting as in (12), we show that D converges on $[\operatorname{Re} > \sigma_0]$, so $\sigma_c^\lambda(D) \leq L$. To finish, assume that $\sigma_c^\lambda(D) \geq 0$. Let $\varepsilon > 0$, and define $\sigma_0 = \sigma_c^\lambda(D) + \varepsilon$. Then D converges at σ_0 (Theorem 2.1), and so by Lemma 2.4 for all $x > 0$

$$|S_x^\lambda(D)(0)| \leq C(\sigma_0) e^{\sigma_0 x},$$

implying $L \leq \sigma_0 = \sigma_c^\lambda(D) + \varepsilon$ for all $\varepsilon > 0$. \square

Given a frequency λ , we also need the abscissa of uniform convergence of a λ -Dirichlet series $D = \sum a_n e^{-\lambda n s}$ determined by the number

$$\sigma_u^\lambda(D) = \inf \{ \sigma \in \mathbb{R} : D \text{ converges uniformly on } [\operatorname{Re} > \sigma] \}.$$

Clearly, D for each $\varepsilon > 0$ converges uniformly on $[\operatorname{Re} > \sigma_u^\lambda(D) + \varepsilon]$, whereas it does not converge uniformly on $[\operatorname{Re} > \sigma_u^\lambda(D) - \varepsilon]$.

We finish with the Bohr-Cahen formula for this abscissa.

Corollary 2.6. *Let $D \in \mathcal{D}(\lambda)$. Then*

$$\sigma_u^\lambda(D) \leq \limsup_{x \rightarrow \infty} \frac{\log \left(\sup_{t \in \mathbb{R}} |S_x^\lambda(D)(it)| \right)}{x},$$

with equality whenever $\sigma_u^\lambda(D)$ is non-negative.

In fact this is a relatively simple consequence of Corollary 2.5, once one realizes that all we proved so far, after proper modifications, also holds for λ -Dirichlet series $E = \sum x_n e^{-\lambda_n s}$ with coefficients x_n in a Banach space X .

Proof. Define $X = H_\infty[\operatorname{Re} > 0]$ (the Banach space of all holomorphic and bounded functions on the right half-plane) and the X -valued λ -Dirichlet series

$$E = \sum (a_n e^{-\lambda_n s}) e^{-\lambda_n w}$$

with the coefficients $x_n = a_n e^{-\lambda_n s} \in X$. We easily see that

$$\sigma_u^\lambda(D) = \sigma_c^\lambda(E),$$

and that by the Hahn-Banach theorem and the maximum modulus theorem we, for all $x > 0$, have

$$\|S_x^\lambda(E)(0)\|_X = \sup_{\operatorname{Re} s > 0} |S_x^\lambda(D)(s)| = \sup_{t \in \mathbb{R}} |S_x^\lambda(D)(it)|.$$

Then by the vector-valued extension of Corollary 2.5 we immediately obtain

$$\begin{aligned} \sigma_u^\lambda(D) = \sigma_c^\lambda(E) &\leq \limsup_{x \rightarrow \infty} \frac{\log(\|S_x^\lambda(D)(0)\|_X)}{x} \\ &= \limsup_{x \rightarrow \infty} \frac{\log \left(\sup_{t \in \mathbb{R}} |S_x^\lambda(D)(it)| \right)}{x}, \end{aligned}$$

with equality whenever $\sigma_u^\lambda(D) = \sigma_c^\lambda(E) \geq 0$. □

2.2. Riesz means. All definitions following are inspired by [13] (see also [9]). Let λ be a frequency, $k \geq 0$, and $C = \sum a_n$ a series in a normed space X . Then C is said to be (λ, k) -Riesz summable whenever the limit

$$\lim_{x \rightarrow \infty} \sum_{\lambda_n < x} \left(1 - \frac{\lambda_n}{x}\right)^k a_n$$

exists, and the finite sums

$$R_x^{\lambda, k}(C) := \sum_{\lambda_n < x} \left(1 - \frac{\lambda_n}{x}\right)^k a_n, \quad x > 0$$

are called (λ, k) -Riesz means of $C = \sum a_n$ of first kind. More generally, we consider functions

$$r : [0, \infty[\times [0, \infty[\rightarrow \mathbb{R}_{\geq 0}, \quad (x, t) \mapsto r(x, t),$$

which are C^∞ in t and satisfy $r(x, x) = 0$ for all $x \geq 0$, and call them Riesz weights. This definition helps to unify some of our coming proofs – although we mainly focus on the two special examples:

$$u_k(x, t) = (x - t)^k \quad \text{and} \quad v_k(x, t) = (e^x - e^t)^k.$$

Given a frequency λ , a series $C = \sum a_n$ (in a normed space), and a Riesz weight r , we define the summatory function by

$$S_x^{\lambda, r}(C) = \sum_{\lambda_n < x} a_n r(x, \lambda_n), \quad x \geq 0,$$

and for the most important cases u_k and v_k we abbreviate

$$S_x^{\lambda, k}(C) = S_x^{\lambda, u_k}(C),$$

and

$$U_x^{\lambda, k}(C) = S_x^{\lambda, v_k}(C).$$

Note that, if $C = \sum a_n$ is (e^λ, k) -Riesz summable, then

$$\lim_{x \rightarrow \infty} R_x^{e^\lambda, k}(C) = \lim_{x \rightarrow \infty} R_{e^x}^{e^\lambda, k}(C) = \lim_{x \rightarrow \infty} T_x^{\lambda, k}(C),$$

where

$$(13) \quad T_x^{\lambda, k}(C) = \sum_{\lambda_n < x} \left(1 - \frac{e^{\lambda_n}}{e^x}\right)^k a_n = e^{-kx} U_x^{\lambda, k}(C)$$

are the (λ, k) -Riesz means of second kind.

Hardy and Riesz in [13, Theorem 16, 17] collected the following two basic properties of Riesz summability (in the order given). Since we try to keep our article as self contained as possible, we are going to give proofs with full details in our appendix from Section 4 (in particular, the original arguments for the second statement are, according to [13, p.30], somewhat 'intricate').

Theorem 2.7. *Let λ be a frequency, $k \geq 0$, and $D = \sum a_n$ a series in a normed space X .*

- (i) *If D is (λ, k) -Riesz summable, then it is (λ, ℓ) -Riesz summable for any $\ell \geq k$, and the associated limits coincide.*
- (ii) *If D is (e^λ, k) -Riesz summable, then D is (λ, k) -Riesz summable, and the associated limits coincide.*

Let us come back to λ -Dirichlet series $D = \sum a_n e^{-\lambda_n s}$. For such series and $k \geq 0$, the Dirichlet polynomials

$$R_x^{\lambda, k}(D)(s) = \sum_{\lambda_n < x} a_n e^{-\lambda_n s} \left(1 - \frac{\lambda_n}{x}\right)^k, \quad x \geq 0, s \in \mathbb{C}$$

are called (λ, k) -Riesz means of D , and D is said to be (λ, k) -Riesz summable at $s_0 \in \mathbb{C}$, whenever the limit

$$\lim_{x \rightarrow \infty} R_x^{\lambda, k}(D)(s_0)$$

exists. Clearly, by Theorem 2.7,(ii) we know that $D = \sum a_n e^{-\lambda_n s}$ is (e^λ, k) -Riesz summable at $s_0 \in \mathbb{C}$, if

$$\lim_{x \rightarrow \infty} T_x^{\lambda, k}(D)(s_0)$$

exists.

A basic property states that the (λ, k) -Riesz summability of D at s_0 implies the $(\lambda, k + \varepsilon)$ -Riesz summability of D at s_0 for every $\varepsilon > 0$ (see Theorem 2.7,(i)). As done for $k = 0$ above, we define

$$\sigma_c^{\lambda, k}(D) = \inf \{ \sigma \in \mathbb{R} : D \text{ } (\lambda, k)\text{-Riesz summable in } \sigma \},$$

which characterizes the largest possible open half-plane on which D is pointwise (λ, k) -Riesz summable. Similarly, we put

$$\sigma_u^{\lambda, k}(D) = \inf \{ \sigma \in \mathbb{R} : D \text{ uniformly } (\lambda, k)\text{-Riesz summable in } \sigma \},$$

defining the largest possible open half-plane on which D is uniformly (λ, k) -Riesz summable.

Let us compare summability with respect to Riesz means of first and second kind. Note first that, given a λ -Dirichlet series D , we by Theorem 2.7,(ii) always have $\sigma_c^{\lambda, k}(D) \leq \sigma_c^{e^\lambda, k}(D)$, where

$$\sigma_c^{e^\lambda, k}(D)$$

now characterizes the largest possible open half-plane on which D is (e^λ, k) -Riesz summable. Hardy and Riesz in [13, Theorem 30] even prove equality.

Theorem 2.8. *For every λ -Dirichlet series D and every $k > 0$*

$$\sigma_c^{\lambda, k}(D) = \sigma_c^{e^\lambda, k}(D).$$

The proof of this result is only sketched in [13]. As a by-product of what we intend to do, we in our appendix from Section 4 will give a full self contained proof.

2.3. Integral representation. The following analog of Theorem 2.2 rules the theory of Riesz summation of general Dirichlet series. The result is taken from [13, Theorem 23]. In fact, (16) is an improvement of the original form from [13], which we repeat in Remark 2.11,(i).

Given $k \geq 0$ and a frequency λ , the theory of (λ, k) -Riesz summability of λ -Dirichlet series $\sum a_n e^{-\lambda s}$ is ruled by the so-called summatory functions of order k , which for $s \in \mathbb{C}$ and $x > 0$ are defined by

$$(14) \quad S_x^{\lambda, k}(D)(s) = \sum_{\lambda_n < x} a_n e^{-\lambda_n s} (x - \lambda_n)^k = x^k R_x^{\lambda, k}(D)(s).$$

Theorem 2.9. *Let $k \geq 0$, and $D = \sum a_n e^{-\lambda_n s} \in \mathcal{D}(\lambda)$ be (λ, k) -summable at $s_0 \in [\operatorname{Re} \geq 0]$. Then D is (λ, k) -summable on $[\operatorname{Re} > \operatorname{Re} s_0]$ and the limit function*

$$(15) \quad f: [\operatorname{Re} > \operatorname{Re} s_0] \rightarrow \mathbb{C}, \quad s \mapsto \lim_{x \rightarrow \infty} \sum_{\lambda_n < x} a_n \left(1 - \frac{\lambda_n}{x}\right)^k e^{-\lambda_n s},$$

is holomorphic and satisfies for all $s \in [\operatorname{Re} > \operatorname{Re} s_0]$

$$(16) \quad \Gamma(1+k) \frac{f(s)}{s^{1+k}} = \int_0^\infty e^{-st} S_t^{\lambda,k}(D)(0) dt,$$

where as usual Γ denotes the Gamma function. Moreover, the convergence in (15) is uniform on each cone $|\arg(s - s_0)| \leq \gamma < \frac{\pi}{2}$.

The proof, although far more complex, is analog to that for the case $k = 0$ stated in Theorem 2.2. A proof of the two cases $k \in \mathbb{N}_0$ and $0 < k < 1$ is given in [13], whereas the proof of the general case is omitted. Very much inspired by the ideas used in [13], we give a self contained proof with full details in our appendix from Section 4 which incorporates all different cases simultaneously.

For the rest of this section we collect a few crucial consequences of the preceding theorem.

Remark 2.10. *As above, we like to mention that the integral formula from (16) may be read as follows: The function $s \mapsto \Gamma(1+k) \frac{f(s)}{s^{1+k}}$ is the Laplace transform of $t \mapsto S_t^{\lambda,k}(D)(0)$, in short*

$$\mathcal{L}(t \mapsto S_t^{\lambda,k}(D)(0))(s) = \Gamma(1+k) \frac{f(s)}{s^{1+k}}.$$

The case $f = 1$ of (16) (assuming $\lambda_1 = 0$) reproves that

$$\mathcal{L}(t^k)(s) = \frac{\Gamma(1+k)}{s^{1+k}}, \quad s \in [\operatorname{Re} > 0].$$

The following remark collects two useful reformulations of Theorem 2.9 in terms of the abscissa of (λ, k) -Riesz summability of $D = \sum a_n e^{-\lambda_n s}$.

Remark 2.11. *Let $k \geq 0$, and $D = \sum a_n e^{-\lambda_n s} \in \mathcal{D}(\lambda)$. Assume that $\sigma_c^{\lambda,k}(D) \in \mathbb{R}$, and let $f : [\operatorname{Re} > \sigma_c^{\lambda,k}(D)] \rightarrow \mathbb{C}$ be the limit function of D .*

(i) *For every $s_0 \in [\operatorname{Re} \geq \sigma_c^{\lambda,k}(D)]$ and $s \in [\operatorname{Re} > 0]$*

$$f(s + s_0) \frac{\Gamma(1+k)}{s^{1+k}} = \int_0^\infty e^{-st} S_t^{\lambda,k}(D)(s_0) dt.$$

(ii) *For every $s_0 \in [\operatorname{Re} > \sigma_c^{\lambda,k}(D)]$ and $s \in [\operatorname{Re} > 0]$*

$$f(s + s_0) = f(s_0) + \frac{s^{1+k}}{\Gamma(1+k)} \int_0^\infty e^{-st} (S_t^{\lambda,k}(D)(s_0) - f(s_0)) dt.$$

Proof. In order to prove (i) assume first that $\operatorname{Re} s_0 > \sigma_c^{\lambda,k}(D)$, so D is (λ, k) -summable in s_0 . Clearly, we obtain (i), if we apply Theorem 2.9 to the translation $D_{s_0} = \sum a_n e^{-\lambda_n s_0} e^{-\lambda_n s}$. If $\operatorname{Re} s_0 = \sigma_c^{\lambda,k}(D)$, then take $s = \sigma + i\tau$ and $\varepsilon > 0$ such that $2\varepsilon < \sigma$. Then we conclude from the first case that

$$f(s + s_0 + \varepsilon) \frac{\Gamma(1+k)}{s^{1+k}} = \int_0^\infty e^{-st} S_t^{\lambda,k}(D)(s_0 + \varepsilon) dt,$$

and hence by continuity of f the claim follows once we prove that

$$\lim_{\varepsilon \rightarrow 0} \int_0^\infty e^{-st} S_t^{\lambda,k}(D)(s_0 + \varepsilon) dt = \int_0^\infty e^{-st} S_t^{\lambda,k}(D)(s_0) dt.$$

Indeed, by (the forthcoming) Lemma 4.7 we have

$$|R_t^{\lambda,k}(D)(s_0 + \varepsilon)| \leq e^{(\sigma - \varepsilon)t} \sup_{0 < y < t} |R_y^{\lambda,k}(D)(s_0 + \varepsilon + (\sigma - \varepsilon))| \leq C(\sigma) e^{\frac{\sigma}{2}t},$$

and hence the dominated convergence theorem does the job. Finally, in order to see (ii), apply (i) to the function $g(s) = f(s) - f(s_0)$, assuming without loss of generality that $\lambda_1 = 0$. \square

Corollary 2.12. *Let k, D and f be as in Theorem 2.9. Then*

$$\forall \varepsilon, \delta > 0 \exists t_0 > 0 \forall |t| \geq t_0, \sigma > \delta + \sigma_c^{\lambda,k}(D): |f(\sigma + it)| \leq \varepsilon |t|^{k+1}.$$

In particular, the function f for every $\delta > 0$ has finite uniform order $\leq k + 1$ on $[\text{Re} > \delta + \sigma_c^{\lambda,k}(D)]$.

The result is taken from [13, Theorem 38].

Proof. Without loss of generality we assume that D is (λ, k) -Riesz summable in 0. We choose $\varepsilon, \delta > 0$, an arbitrary $0 < \gamma \leq \frac{\pi}{2}$, and put

$$M = \sup_{x > 0} |R_x^{\lambda,k}(D)(0)| < \infty.$$

In a first step we consider f on the complement of $[|\arg \bullet| \leq \gamma]$ in $[\text{Re} > 0]$ intersected with $[\text{Re} > \delta]$, i.e. all $s = \sigma + it \in [\text{Re} > \delta]$ such that $\frac{|s|}{|t|} \leq \frac{1}{\sin \gamma}$. Fix such s , and choose $x_0 > 0$ such that

$$\int_{x_0}^\infty y^k e^{-\delta y} dy < \frac{\Gamma(1+k) (\sin \gamma)^{1+k}}{M} \varepsilon.$$

Applying again Theorem 2.9, we see that

$$\begin{aligned} f(s) &= \frac{s^{k+1}}{\Gamma(1+k)} \int_0^{x_0} S_y^{\lambda,k}(D)(0) e^{-sy} dy \\ &\quad + \frac{s^{k+1}}{\Gamma(1+k)} \int_{x_0}^\infty S_y^{\lambda,k}(D)(0) e^{-sy} dy = J_1(s) + J_2(s). \end{aligned}$$

Then

$$J_2(s) \leq |s|^{k+1} \frac{M}{\Gamma(1+k)} \int_{x_0}^\infty y^k e^{-\delta y} dy \leq \varepsilon |t|^{k+1}.$$

Moreover, by partial integration we have

$$J_1(s) = \frac{s^k}{\Gamma(1+k)} \left(-S_{x_0}^{\lambda,k}(D)(0) e^{-sx_0} + \int_0^{x_0} (S_\bullet^{\lambda,k}(D)(0))'(y) e^{-sy} dy \right),$$

and then, since $|e^{-sy}| \leq 1$ for all $0 \leq y \leq x_0$, there is $C = C(x_0, k) > 0$ such that

$$|J_1(s)| \leq |s|^k C \leq \frac{C}{\sin \gamma} |t|^k \leq \varepsilon |t|^{k+1},$$

whenever $\frac{C}{\varepsilon \sin \gamma} \leq |t|$. All in all this shows that $|f(s)| \leq \varepsilon |t|^{k+1}$ for all $s = \sigma + it$ in the complement of $[|\arg \bullet| \leq \gamma]$ in $[\operatorname{Re} > 0]$ intersected with $[\operatorname{Re} > \delta]$ provided $\frac{C}{\varepsilon \sin \gamma} \leq |t|$. Using again Theorem 2.9, we know that f is bounded on the cone $[|\arg \bullet| \leq \gamma]$, which then clearly leads to the desired conclusion. \square

2.4. Perron formula. We prove Perron's formula – an integral formula for the summatory function of a λ -Dirichlet series in terms of its limit function. The result was first presented in [13, Theorem 39]. In fact, it turns out that it follows by Fourier inversion from the integral formula in Theorem 2.9. The original proof of Perron's formula for general Dirichlet series as given in [13] is mainly a consequence of Cauchy's integral theorem. Our approach, deeply inspired by ideas used in [17] to cover the case $k = 0$, differs considerable.

Theorem 2.13. *For $k \geq 0$ let $D = \sum a_n e^{-\lambda_n s}$ be somewhere (λ, k) -Riesz summable and $f : [\operatorname{Re} > \sigma_c^{\lambda, k}(D)] \rightarrow \mathbb{C}$ its limit function. Then for each $x > 0$ and $c > \max\{\sigma_c^{\lambda, k}(D), 0\}$ we have*

$$S_x^{\lambda, k}(D)(0) = \frac{\Gamma(1+k)}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{f(s)}{s^{1+k}} e^{xs} ds.$$

Proof. Let us denote by $\mathcal{F}_{L_1(\mathbb{R})}$ the Fourier transform on $L_1(\mathbb{R})$ (with respect to the ordinary Lebesgue measure). Then for all $y \in \mathbb{R}$ by Theorem 2.9

$$\begin{aligned} \frac{f(c+iy)}{(c+iy)^{1+k}} &= \frac{1}{\Gamma(1+k)} \int_0^\infty [S_t^{\lambda, k}(D)(0) e^{-ct}] e^{-iyt} dt \\ &= \frac{1}{\Gamma(1+k)} \mathcal{F}_{L_1(\mathbb{R})}(t \mapsto S_t^{\lambda, k}(D)(0) e^{-ct})(y). \end{aligned}$$

Since the function $g(t) = S_t^{\lambda, k}(D)(0) e^{-ct}$ satisfies a Lipschitz condition at all $t \notin \{\lambda_n \mid n \in \mathbb{N}\}$, we by [15, Corollary on p.9] for all such t obtain

$$S_t^{\lambda, k}(D)(0) e^{-ct} = \lim_{A \rightarrow \infty} \frac{1}{2\pi} \int_{-A}^A \Gamma(1+k) \frac{f(c+iy)}{(c+iy)^{1+k}} e^{iyt} dy.$$

Hence,

$$S_t^{\lambda, k}(D)(0) = \frac{\Gamma(1+k)}{2\pi} \lim_{A \rightarrow \infty} \int_{-A}^A \frac{f(c+iy)}{(c+iy)^{1+k}} e^{t(c+iy)} dy,$$

which is the equality we intended to prove. \square

It is important for our later purposes that the preceding integral formula for the summatory function of a given λ -Dirichlet series D , can be extended considerably, whenever the limit function of f extends holomorphically to the full right half-plane – still satisfying a 'finite order type-condition' (a result sketched in [13, p.51]).

Corollary 2.14. For $k \geq 0$ let $D = \sum a_n e^{-\lambda_n s}$ be (λ, k) -Riesz summable for some $s_0 \in [\text{Re} > 0]$. Assume that the limit function f of D extends holomorphically to $[\text{Re} > 0]$ such that

$$\forall \varepsilon, \delta > 0 \exists t_0 > 0 \forall |t| \geq t_0, \sigma > \delta: |f(\sigma + it)| \leq \varepsilon |\sigma + it|^{1+k}.$$

Then for each $x > 0$ and $c > 0$

$$S_x^{\lambda, k}(D)(0) = \frac{\Gamma(1+k)}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{f(s)}{s^{1+k}} e^{xs} ds.$$

Proof. By Theorem 2.13 we know that for $x > 0$ and $\sigma > \text{Re } s_0$

$$S_x^{\lambda, k}(D)(0) = \frac{\Gamma(1+k)}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \frac{f(s)}{s^{1+k}} e^{xs} ds.$$

We claim that for each $x > 0$ and $c > 0$

$$(17) \quad \int_{c-i\infty}^{c+i\infty} \frac{f(s)}{s^{1+k}} e^{xs} ds = \int_{\sigma-i\infty}^{\sigma+i\infty} \frac{f(s)}{s^{1+k}} e^{xs} ds.$$

Indeed, an application of Cauchy's integral theorem shows that (17) follows, once we prove that

$$(18) \quad \lim_{T \rightarrow \infty} \int_c^\sigma \frac{f(y \pm iT)}{(y \pm iT)^{1+k}} e^{x(y \pm iT)} dy = 0.$$

To see this, choose, given $\varepsilon > 0$ and $\delta = c$, some $t_0 > 0$ according to the assumption. Then for all $T > t_0$

$$\left| \int_c^\sigma \frac{f(y \pm iT)}{(y \pm iT)^{1+k}} e^{x(y \pm iT)} dy \right| \leq \varepsilon \int_c^\sigma e^{xy} dy = \varepsilon (e^{x\sigma} - e^{xc}) x^{-1},$$

which implies (18). □

Finally, we show that the coefficients of a λ -Dirichlet series are uniquely determined by the values of its limit function on some abscissa - another crucial point for our coming purposes.

Corollary 2.15. Let $D = \sum a_n e^{-\lambda_n s}$ be (λ, k) -Riesz summable at $s_0 = \sigma_0 + i\tau_0$ for some $k \geq 0$. If the limit function f of D vanishes on some vertical line $[\text{Re} = \sigma]$, where $\sigma > \sigma_0$, then $a_n = 0$ for all n .

Proof. Let first $\lambda_1 < x < \lambda_2$. Then by Theorem 2.13 and our assumption

$$a_1 e^{-\lambda_1 s_0} \left(1 - \frac{\lambda_1}{x}\right)^k = S_x^{\lambda, k}(D)(0) = \frac{\Gamma(1+k)}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \frac{f(s) e^{xs}}{s^{1+k}} ds = 0,$$

which implies $a_1 = 0$. Proceeding successively by considering a sequence (x_n) with $\lambda_n < x_n < \lambda_{n+1}$, we obtain that $a_n = 0$ for all n . □

2.5. Bohr-Cahen formula. The following formula for the abscissa of convergence is again due to Hardy and Riesz [13, Theorem 31].

Theorem 2.16. *Let $D = \sum a_n e^{-\lambda_n s}$ and $k \geq 0$. Then*

$$\sigma_c^{\lambda,k}(D) \leq \limsup_{x \rightarrow \infty} \frac{\log(|R_x^{\lambda,k}(D)(0)|)}{x},$$

with equality whenever $\sigma_c^{\lambda,k}(D)$ is non-negative.

The proof will be given in Section 4, mainly as a consequence of Theorem 2.9. Note that

$$\limsup_{x \rightarrow \infty} \frac{\log(|R_x^{\lambda,k}(D)(0)|)}{x} = \limsup_{x \rightarrow \infty} \frac{\log(|S_x^{\lambda,k}(D)(0)|)}{x}.$$

Recall that Theorem 2.16 for the special case $k = 0$ was already presented in Corollary 2.5, and that we in Corollary 2.6 formulated a uniform counterpart.

Again a simple analysis shows that the proof of Theorem 2.16 also works for vector-valued Dirichlet series. Hence we exactly as in Section 2.1 obtain the following uniform variant of Theorem 2.16 (note that for $0 < k < 1$ a direct, but technically more involved, argument was given in [20, Lemma 3.8]).

Corollary 2.17. *Let $D = \sum a_n e^{-\lambda_n s}$ and $k \geq 0$. Then*

$$\sigma_u^{\lambda,k}(D) \leq \limsup_{x \rightarrow \infty} \frac{\log(\sup_{t \in \mathbb{R}} |R_x^{\lambda,k}(D)(it)|)}{x},$$

with equality whenever $\sigma_u^{\lambda,k}(D)$ is non-negative.

3. A new scale of Banach spaces

We start repeating some of our basic definitions from the introduction. Given a frequency $\lambda = (\lambda_n)$, we call a λ -Dirichlet series $D = \sum a_n e^{-\lambda_n s}$ a λ -Riesz germ of the holomorphic function $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$, whenever $\sigma_c^{\lambda,m}(D) < \infty$ for some $m \geq 0$ and f is the holomorphic extension of the limit function of D to all of $[\operatorname{Re} > 0]$.

Two remarks are in order.

Remark 3.1. *Let $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$ be a holomorphic function generated by the λ -Riesz germ $D = \sum a_n e^{-\lambda_n s}$. Then D by Corollary 2.15 is unique, and as a consequence we may assign to every such f the unique sequence $(a_n(f))_n = (a_n)_n$, which we (like for Dirichlet series) again call the 'sequence of Bohr coefficients of f '.*

Remark 3.2. *Let $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$ be holomorphic. Then f has a λ -Riesz germ if and only if f on some half-plane and for some m equals the limit function of some λ -Dirichlet series under (e^λ, m) -Riesz summation. This is an immediate consequence of Theorem 2.8.*

Moreover, given a holomorphic function $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$ generated by the λ -Riesz germ D , the x th Riesz mean of order $k \geq 0$ of f in $s \in \mathbb{C}$ is given by

$$R_x^{\lambda,k}(f)(s) = \sum_{\lambda < x} a_n(f) e^{-\lambda_n s} \left(1 - \frac{\lambda_n}{x}\right)^k,$$

and its summatory function in $s \in \mathbb{C}$ by

$$S_x^{\lambda,k}(f)(s) = \sum_{\lambda < x} a_n(f) e^{-\lambda_n s} (x - \lambda_n)^k, \quad x > 0.$$

As explained above the spaces

$$H_{\infty,\ell}^{\lambda}[\operatorname{Re} > 0], \quad \ell \geq 0,$$

accumulate all holomorphic functions $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$, which are generated by a λ -Riesz germ and satisfy the growth condition

$$(19) \quad \|f\|_{\infty,\ell} = \sup_{\operatorname{Re} s > 0} \frac{|f(s)|}{(1 + |s|)^{\ell}} < \infty.$$

For each such f

$$\sup_{\operatorname{Re} s > 0} \frac{|f(s)|}{(1 + |s|)^{\ell}} \leq \sup_{\operatorname{Re} s > 0} \frac{|f(s)|}{|(1 + s)^{\ell}|} \leq 2^{\ell} \sup_{\operatorname{Re} s > 0} \frac{|f(s)|}{(1 + |s|)^{\ell}},$$

since $|1 + s| \leq 1 + |s| \leq 2|1 + s|$ for all $s \in [\operatorname{Re} > 0]$. Obviously, the pairs

$$(H_{\infty,\ell}^{\lambda}[\operatorname{Re} > 0], \|\cdot\|_{\infty,\ell})$$

form an increasing scale of normed spaces. That all these spaces in fact are complete, so Banach spaces, is a non-trivial fact which is given in Theorem 3.16.

Finally, we remark that a holomorphic function satisfying the growth condition (19), by Remark 3.2 (see also (13)) belongs to $H_{\infty,\ell}^{\lambda}[\operatorname{Re} > 0]$ if and only if there is a λ -Dirichlet series $D = \sum a_n e^{-\lambda_n s}$ such that for some $m, \sigma \geq 0$ and for every $s \in [\operatorname{Re} > \sigma]$

$$(20) \quad f(s) = \lim_{x \rightarrow \infty} R_x^{e^{\lambda},m}(D) = \lim_{x \rightarrow \infty} T_x^{\lambda,m}(D) = \lim_{x \rightarrow \infty} \sum_{\lambda_n < x} a_n e^{-\lambda_n s} \left(1 - \frac{e^{\lambda_n}}{e^x}\right)^m.$$

3.1. Ordinary and power case. We start considering the power case $\lambda = (n)$. Then, as already announced in the introduction, the scale $H_{\infty,\ell}^{(n)}[\operatorname{Re} > 0]$, $\ell \geq 0$, collapses to the case $\ell = 0$.

Proposition 3.3. *For every $\ell \geq 0$*

$$H_{\infty}^{(n)}[\operatorname{Re} > 0] = H_{\infty,\ell}^{(n)}[\operatorname{Re} > 0].$$

Proof. Take $f \in H_{\infty,\ell}^{(n)}[\operatorname{Re} > 0]$, let $D = \sum a_n e^{-ns}$ be its associated Dirichlet series, and choose any $m \in \mathbb{N}$ such that $\ell < m$. Then we deduce from Corollary 3.9 (proved below) and Theorem 2.8 (for which we give a complete proof in our

appendix from Section 4) that D is $((e^n), m)$ -Riesz summable on $[\operatorname{Re} > 0]$ with limit f . Then by [13, Theorem 21]

$$\lim_{N \rightarrow \infty} \left(\frac{e^{N+1} - e^N}{e^{N+1}} \right)^m \left(\sum_{n=1}^N a_n - \sum_{n=1}^{\infty} a_n \right) = 0,$$

i.e. D on $[\operatorname{Re} > 0]$ converges pointwise to f .

We remark that [13, Theorem 21] is a result on arbitrary frequencies λ and arbitrary orders $k > 0$, and we here only use this result for the integer case $k = m \in \mathbb{N}$, which according to [13, p.36] has an 'extremely simple' proof.

Continuing the proof, we see that D converges uniformly on $[\operatorname{Re} > \varepsilon]$ for every $\varepsilon > 0$ (indeed, for the frequency $\lambda = (n)$ the abscissas of convergence and absolute convergence coincide – see [4, 12, §3, Hilfssatz 2,3]). So in particular f is bounded on $[\operatorname{Re} > 1]$. Moreover, $f(\sigma + i \cdot) : \mathbb{R} \rightarrow \mathbb{C}$ is 2π -periodic for every $\sigma > 0$, which implies that

$$\begin{aligned} \sup_{\substack{0 < \sigma < 2 \\ t \in \mathbb{R}}} |f(\sigma + it)| &= \sup_{\substack{0 < \sigma < 2 \\ t \in [0, 2\pi]}} |f(\sigma + it)| \\ &\leq \|f\|_{\infty, \ell} \sup_{\substack{0 < \sigma < 2 \\ t \in [0, 2\pi]}} (1 + |\sigma + it|)^\ell \leq \|f\|_{\infty, \ell} (1 + \sqrt{4 + 4\pi^2})^\ell. \end{aligned}$$

All in all we conclude that $f \in H_\infty^{(n)}[\operatorname{Re} > 0]$. \square

Next we consider the frequency $\lambda = (\log n)$, which generates ordinary Dirichlet series $\sum a_n n^{-s}$. Looking at (20) (with $x = \log y$), we see that a holomorphic function $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$ belongs to the Banach space $\mathcal{H}_{\infty, 1}^{(\log n)}[\operatorname{Re} > 0]$ if and only if the growth condition (19) holds, and there exists an ordinary Dirichlet series $D = \sum a_n n^{-s}$ such that for some $m, \sigma \geq 0$

$$f(s) = \lim_{y \rightarrow \infty} \sum_{n < y} a_n n^{-s} \left(1 - \frac{n}{y}\right)^m, \quad s \in [\operatorname{Re} > \sigma].$$

Actually, we will see in Corollary 3.26 that Theorem 3.10 in the special case $\lambda = (\log n)$ is valid for $k = \ell$, and therefore in the particular case $\ell = 1$

$$f(s) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \sum_{k=1}^n a_k k^{-s}, \quad s \in [\operatorname{Re} > 0],$$

so that eventually we arrive at the following result.

Proposition 3.4. *Let $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$ be holomorphic. Then the growth condition (19) holds and there exists an ordinary Dirichlet series which on $[\operatorname{Re} > 0]$ is Cesàro summable with limit f , if and only if $f \in H_{\infty, 1}^{(\log n)}[\operatorname{Re} > 0]$.*

Note that the space $H_{\infty, \ell}^{(\log n)}[\operatorname{Re} > 0]$ is strictly larger than $H_\infty^{(\log n)}[\operatorname{Re} > 0]$. To see an example, we again look at the eta-function η (defined in the introduction), for which

$$\ell > \frac{1}{2} \quad \Rightarrow \quad \eta \in H_{\infty, \ell}^{(\log n)}[\operatorname{Re} > 0] \quad \Rightarrow \quad \ell \geq \frac{1}{2}.$$

In particular, $\eta \notin H_\infty^{(\log n)}[\operatorname{Re} > 0]$. Alternatively, this may also be seen as a consequence of Bohr's inequality (see [7, Corollary 4.3] or [21, Theorem 4.4.1]): Assuming $\eta \in \mathcal{H}_\infty((\log n))$ implies that $N = \sum_{n=1}^N |(-1)^{p_n}| \leq \|D\|_\infty < \infty$ for all N , which clearly is a contradiction.

This example in particular shows that, although we in general have

$$\mathcal{H}_{\infty, \ell_1}^\lambda[\operatorname{Re} > 0] \subset \mathcal{H}_{\infty, \ell_2}^\lambda[\operatorname{Re} > 0] \quad \text{for all } 0 \leq \ell_1 < \ell_2 ,$$

this inclusion for certain λ 's may be strict whereas for other λ 's it actually may be an equality (of sets).

3.2. Perron formula – a variant. We now for the new scale of Banach spaces prove an important variant of Perron's formula from Theorem 2.13.

Theorem 3.5. *Let $f \in H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$ and $k > \ell \geq 0$. Then for all $s_0 \in [\operatorname{Re} \geq 0]$, $x > 0$ and $c > 0$*

$$S_x^{\lambda, k}(f)(s_0) = \frac{\Gamma(1+k)}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{f(s+s_0)}{s^{1+k}} e^{xs} ds.$$

Proof. After translation we may assume that $s_0 = 0$. By assumption the Dirichlet series $D = \sum a_n e^{-\lambda_n s}$ is (λ, m) -Riesz summable at some $s_1 \in [\operatorname{Re} > 0]$ for some $m \geq 0$. Hence D is $(\lambda, m+k)$ -Riesz summable at s_1 (Theorem 2.7,(i)), and for every $\varepsilon, \sigma > 0$ and $|t| \geq t_0 > 0$

$$\frac{|f(\sigma + it)|}{|\sigma + it|^{1+m+k}} \leq \|f\|_{\infty, \ell} |\sigma + it|^{-(1+m+k-\ell)} \leq t_0^{-(1+m+k-\ell)} \|f\|_{\infty, \ell} \leq \varepsilon ,$$

whenever t_0 is large enough. Corollary 2.14 then implies that for all $c, x > 0$

$$\sum_{\lambda_n < x} a_n (x - \lambda_n)^{m+k} = \frac{\Gamma(1+m+k)}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{f(s) e^{xs}}{s^{1+m+k}} ds.$$

Now we differentiate m -times obtaining

$$\frac{d^m}{dx} \int_{c-i\infty}^{c+i\infty} \frac{f(s) e^{xs}}{s^{1+k+m}} ds = \int_{c-i\infty}^{c+i\infty} \frac{f(s) e^{xs}}{s^{1+k}} ds$$

and

$$\frac{d^m}{dx} \sum_{\lambda_n < x} a_n (x - \lambda_n)^{m+k} = \prod_{j=0}^{m-1} (m+k-j) \sum_{\lambda_n < x} a_n (x - \lambda_n)^k ;$$

for the second equality see (33), and for the first note that the differentiation under the integral is legit, since for every $j = 0, \dots, m-1$

$$\int_{c-i\infty}^{c+i\infty} \left| \frac{f(s) e^{xs}}{s^{1+k+j}} \right| ds \leq e^{xc} \|f\|_{\infty, \ell} \int_{c-i\infty}^{c+i\infty} |s|^{-(1+k+j-\ell)} ds < \infty.$$

Additionally by the functional equation of the Gamma function we have

$$\Gamma(1+k)\Gamma(1+m+k) = \prod_{j=0}^{m-1} (m+k-j),$$

so that altogether we obtain

$$S_x^{\lambda,k}(f)(0) = \sum_{\lambda_n < x} a_n(x - \lambda_n)^k = \frac{\Gamma(1+k)}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{f(s)e^{xs}}{s^{1+k}} ds. \quad \square$$

3.3. A maximal inequality for Riesz means. The following maximal inequality is going to be our major tool to perform a substantial structure theory of the spaces $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$. It shows that for $k > \ell$ the linear projections

$$R_x^{\lambda,k} : H_{\infty,\ell}^\lambda[\operatorname{Re} > 0] \rightarrow H_{\infty,\ell}^\lambda[\operatorname{Re} > 0], \quad f \rightarrow R_x^{\lambda,k}f, \quad x > 0,$$

are uniformly bounded.

Theorem 3.6. *Let $k > \ell \geq 0$ and λ a frequency. Then there is a constant $C = C(k, \ell, \lambda) > 0$ such that for all $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$*

$$\sup_{x>0} \|R_x^{\lambda,k}(f)\|_{\infty,\ell} \leq C\|f\|_{\infty,\ell}.$$

In particular, for every $\sigma \geq 0$ there is a constant $C = C(\ell, k, \lambda, \sigma) > 0$ such that for every $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ and $t \in \mathbb{R}$

$$\sup_{x>0} |R_x^{\lambda,k}(f)(\sigma + it)| \leq C(1 + |t|)^\ell \|f\|_{\infty,\ell}.$$

Proof. We assume first that $\lambda_1 > 0$, and take $s = \sigma + it \in [\operatorname{Re} > 0]$ and $y \geq \lambda_1$. Then by Theorem 3.5 with $c = y^{-1}$

$$|R_y^{\lambda,k}(D)(s)| \leq \frac{e\Gamma(1+k)}{2\pi} y^{-k} \int_{\mathbb{R}} \frac{|f(s + \frac{1}{y} + ix)|}{|\frac{1}{y} + ix|^{1+k}} dx,$$

and

$$\begin{aligned} \|f\|_{\infty,\ell}^{-1} y^{-k} \int_{\mathbb{R}} \frac{|f(s + \frac{1}{y} + ix)|}{|\frac{1}{y} + ix|^{1+k}} dx &\leq \int_{\mathbb{R}} \frac{y|1 + s + \frac{1}{y} + ix|^\ell}{|1 + ixy|^{1+k}} dx \\ &\leq 2^{\max\{0,\ell-1\}} \int_{\mathbb{R}} \frac{y(|1 + s|^\ell + |\frac{1}{y} + ix|^\ell)}{|1 + ixy|^{1+k}} dx \\ &\leq 2^{\max\{0,\ell-1\}} \left((1 + |s|)^\ell \int_{\mathbb{R}} \frac{y}{|1 + ixy|^{1+k}} dx + y^{-\ell} \int_{\mathbb{R}} \frac{y}{|1 + ixy|^{1+k-\ell}} dx \right) \\ &\leq 2^{\max\{0,\ell-1\}} \left((1 + |s|)^\ell \int_{\mathbb{R}} \frac{1}{|1 + iv|^{1+k}} dv + \lambda_1^{-\ell} \int_{\mathbb{R}} \frac{1}{|1 + iv|^{1+k-\ell}} dv \right) \\ &\leq 2^{\max\{0,\ell-1\}} \left((1 + |s|)^\ell \left(1 + \frac{1}{k}\right) + \lambda_1^{-\ell} \left(1 + \frac{1}{k-\ell}\right) \right) \\ &\leq 2^{\max\{0,\ell-1\}} \left(\left(1 + \frac{1}{k}\right) + \lambda_1^{-\ell} \left(1 + \frac{1}{k-\ell}\right) \right) (1 + |s|)^\ell. \end{aligned}$$

Assume now that $\lambda_1 = 0$ and choose $0 < y < \lambda_2$. Then by the previous calculation

$$|a_0| \leq \|f\|_{\infty,\ell} 2^{\max\{0,\ell-1\}} \left(\left(1 + \frac{1}{k}\right) + y^{-\ell} \left(1 + \frac{1}{k-\ell}\right) \right) (1 + |s|)^\ell.$$

Tending $y \rightarrow \lambda_2$ we obtain

$$|a_0| \leq C(\lambda, k, \ell)(1 + |s|)^\ell \|f\|_{\infty, \ell}.$$

The 'in particular part' is then a straight forward consequence, since

$$1 + |s| \leq (1 + \sigma)\left(1 + \frac{|t|}{1 + \sigma}\right) \leq (1 + \sigma)(1 + |t|). \quad \square$$

3.4. Uniform Riesz approximation. Next we give the main application of the preceding maximal theorem. We prove that for $k > \ell$ all functions in a bounded subset of $H_{\infty, \ell}^\lambda[\text{Re} > 0]$, after translation by some arbitrary $u > 0$, are uniformly (λ, k) -Riesz approximable in the norm of $H_{\infty, \ell}^\lambda[\text{Re} > 0]$.

Theorem 3.7. *Let λ be a frequency and $k > \ell \geq 0$. Then for every bounded set $M \subset H_{\infty, \ell}^\lambda[\text{Re} > 0]$ and every choice of $\varepsilon, u > 0$ there is $x_0 > 0$ such that*

$$\sup_{x > x_0} \sup_{f \in M} \|f(u + \cdot) - R_x^{\lambda, k}(f)(u + \cdot)\|_{\infty, \ell} \leq \varepsilon.$$

Proof. Let $m \in N_0$ such that $m < k \leq m + 1$, and $C = \sup_{f \in M} \|f\|_{\infty, \ell} < \infty$. Fix $f \in M$ and $u > 0$. Then by Lemma 4.11 and Theorem 3.6 we for every $s \in [\text{Re} > 0]$ have

$$\begin{aligned} & \left| \frac{x^{-k}}{\Gamma(m+2)} \int_0^x \frac{S_t^{\lambda, m+1}(f)(s)}{(1+|s|)^\ell} u^{m+2} (x-t)^k e^{-ut} dt - \frac{R_x^{\lambda, k}(f)(s+u)}{(1+|s|)^\ell} \right| \\ & \leq e^{-ux} \frac{|R_x^{\lambda, k}(f)(s)|}{(1+|s|)^\ell} + C_1(m, k) x^{-k} \int_0^x \frac{|S_y^{\lambda, k}(f)(s)|}{(1+|s|)^\ell} \sum_{j=1}^{m+1} u^j e^{-uy} (x-y)^{j-1} dy \\ & \leq e^{-ux} C + C_2(m, k, u) C \sum_{j=1}^{m+1} x^{-k} \int_0^x y^k (x-y)^{j-1} e^{-uy} dy, \end{aligned}$$

where

$$\begin{aligned} x^{-k} \int_0^x y^k (x-y)^{j-1} e^{-uy} dy &= x^j \int_0^x x^{-1} \left(\frac{y}{x}\right)^k \left(1 - \frac{y}{x}\right)^{j-1} e^{-uy} dy \\ &= x^j e^{-ux} \int_0^1 \alpha^k (1-\alpha)^{j-1} d\alpha = x^j e^{-ux} \frac{\Gamma(k+1)\Gamma(j)}{\Gamma(k+1-j)}. \end{aligned}$$

Hence for $x \geq 1$ and $f \in M$

$$\begin{aligned} & \sup_{s \in [\text{Re} > 0]} \left| \frac{x^{-k}}{\Gamma(m+2)} \int_0^x \frac{S_t^{\lambda, m+1}(f)(s)}{(1+|s|)^\ell} u^{m+2} (x-t)^k e^{-ut} dt - \frac{R_x^{\lambda, k}(D)(s+u)}{(1+|s|)^\ell} \right| \\ & \leq C e^{-ux} (1 + C_3(m, k, u) x^{m+1}), \end{aligned}$$

which vanishes uniformly for all $f \in M$ as $x \rightarrow \infty$. Additionally, following the proof of Lemma 4.14 and Lemma 4.5 the limit

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{x^{-k}}{\Gamma(m+2)} \int_0^x \frac{S_t^{\lambda, m+1}(f)(s)}{(1+|s|)^\ell} u^{m+2} (x-t)^k e^{-ut} dt \\ = \frac{1}{\Gamma(1+k)} \int_0^\infty \frac{S_t^{\lambda, k}(f)(s)}{(1+|s|)^\ell} u^{k+1} e^{-ut} dt = \frac{f(s+u)}{(1+|s|)^\ell}, \end{aligned}$$

exists uniformly on $[\operatorname{Re} > 0]$ and $f \in M$, where the last equality holds by Theorem 2.9 (or more precisely Remark 2.11). This proves the claim. \square

The following corollary is an immediate consequence considering singleton sets.

Corollary 3.8. *Let $f \in H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$ and $k > \ell \geq 0$. Then for every $u > 0$*

$$\lim_{x \rightarrow \infty} R_x^{\lambda, k}(f)(u + \cdot) = f(u + \cdot) \quad \text{in } H_{\infty, \ell}^\lambda[\operatorname{Re} > 0].$$

In particular, for every $u > 0$

$$\lim_{x \rightarrow \infty} \sup_{s \in [\operatorname{Re} > u]} \left| \frac{f(s) - R_x^{\lambda, k}(f)(s)}{(1+s)^\ell} \right| = 0.$$

To check the 'in particular' part fix $u > 0$ and let $s = w + u$, where $w \in [\operatorname{Re} > 0]$. Then for large x by the first part

$$\begin{aligned} \frac{|f(s) - R_x^{\lambda, k}(f)(s)|}{|1+s|^\ell} \\ = \frac{|f(u+w) - R_x^{\lambda, k}(f)(u+w)|}{|1+w|^\ell} \frac{|1+w|^\ell}{|1+u+w|^\ell} \leq \varepsilon \frac{|1+u+w|^\ell}{|1+u+w|^\ell} = \varepsilon. \end{aligned}$$

3.5. Theorem 41 of M. Riesz revisited. Corollary 3.8 immediately implies the following reformulation of an important result of M. Riesz published in [13, Theorem 41]. Actually, this result motivated much of the content of this article.

Corollary 3.9. *If $f \in H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$, then for every $k > \ell \geq 0$ and $s \in [\operatorname{Re} > 0]$*

$$f(s) = \lim_{x \rightarrow \infty} R_x^{\lambda, k}(f)(s).$$

Let us point out that the mentioned result of Riesz actually is formulated in terms of (e^λ, k) -Riesz summability (recall the definition of (λ, k) -Riesz means of second kind from (13)). In its original form as given in [13] it reads as follows (using our terminology of having a Riesz germ).

Theorem 3.10. *Let $f: [\operatorname{Re} > 0] \rightarrow \mathbb{C}$ be holomorphic with a λ -Riesz germ $D = \sum a_n e^{-\lambda_n s}$. Assume that there is $\ell \geq 0$ such that*

$$(21) \quad \forall \varepsilon > 0 \exists C > 0: |f(s)| \leq C|s|^\ell, \quad s \in [\operatorname{Re} > \varepsilon].$$

Then for every $k > \ell$ and $s \in [\operatorname{Re} > 0]$

$$f(s) = \lim_{x \rightarrow \infty} T_x^{\lambda, k}(D)(s) = \lim_{x \rightarrow \infty} \sum_{\lambda_n < x} a_n (1 - e^{\lambda_n - x})^k e^{-\lambda_n s}.$$

Let us explain why Theorem 3.10 and Corollary 3.9 in fact are equivalent. Recall first, that by Theorem 2.8 every λ -Dirichlet series is (λ, k) -Riesz summable on all half-planes $[\operatorname{Re} > \sigma]$, $\sigma > 0$ if and only if this series is (e^λ, k) -Riesz summable on all such half-planes. So it is obvious that Theorem 3.10 is applicable to every $f \in H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$; indeed, given $\varepsilon > 0$, we have

$$\sup_{\operatorname{Re} s > \varepsilon} \frac{|f(s)|}{|s|^\ell} \leq \|f\|_{\infty, \ell} \sup_{\operatorname{Re} s > \varepsilon} \frac{(1 + |s|)^\ell}{|s|^\ell} < \infty.$$

Conversely, if a holomorphic function f on $[\operatorname{Re} > 0]$ has a λ -Riesz germ and satisfies (21), then $f_\varepsilon = f(\varepsilon + \cdot) \in H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$ for every $\varepsilon > 0$, and consequently we may apply Corollary 3.9.

The decision on why we focus on (λ, k) -Riesz means instead of (e^λ, k) -Riesz means is based on the integral representation

$$R_x^{\lambda, k}(f)(s_0) = \frac{\Gamma(1+k)}{2\pi i} x^{-k} \int_{c-i\infty}^{c+i\infty} f(s+s_0) \frac{e^{xs}}{s^{1+k}} ds$$

from Theorem 2.13 of the x th Riesz means of first kind of a Dirichlet series $D = \sum a_n e^{-\lambda_n s}$ in terms of its limit function f , whereas by [13, Theorem 40, p. 51] for Riesz means of second kind we have

$$T_x^{\lambda, k}(D)(s_0) = \frac{\Gamma(1+k)}{2\pi i} \int_{c-i\infty}^{c+i\infty} f(s+s_0) \frac{\Gamma(s)e^{xs}}{\Gamma(1+k+s)} ds.$$

In fact, we believe that the handling of the kernels e^{xs}/s^{1+k} is more simple than that of the kernels $\Gamma(s)e^{xs}/\Gamma(1+k+s)$.

3.6. The bounded case $\ell = 0$. For $\ell = 0$ and $f \in H_{\infty, 0}^\lambda[\operatorname{Re} > 0]$, Corollary 3.8 shows that for every $\sigma > 0$

$$f = \lim_{x \rightarrow \infty} R_x^{\lambda, k}(f) \quad \text{uniformly on } [\operatorname{Re} > \sigma].$$

Hence, as the uniform limit of uniformly almost periodic function on \mathbb{R} is again uniformly almost periodic, the function $t \mapsto f(\sigma + it)$ is uniformly almost periodic for every $\sigma > 0$, this has the following consequence.

Corollary 3.11. *For every frequency λ*

$$H_{\infty, 0}^\lambda[\operatorname{Re} > 0] = H_\infty^\lambda[\operatorname{Re} > 0].$$

3.7. Boundedness of coefficient functionals. We explained above that every function in $H_{\infty, \ell}^\lambda[\operatorname{Re} > 0] \rightarrow \mathbb{C}$ is uniquely determined by its Bohr coefficients $a_N(f)$, $N \in \mathbb{N}$ (Remark 3.1). The following result shows that this assignment in fact is continuous.

Theorem 3.12. *Let $k > \ell \geq 0$ and $\lambda = (\lambda_n)$ a frequency. Then for each $N \in \mathbb{N}$ the coefficient mapping*

$$\mathcal{C}_N: H_{\infty, \ell}^\lambda[\operatorname{Re} > 0] \rightarrow \mathbb{C}, \quad f \mapsto a_N(f)$$

is bounded. More precisely, with $\lambda_{-1} := 0$

$$\|\mathcal{C}_N\| \leq C(k, \ell, \lambda_1) \left(\left(\frac{\lambda_{N+1}}{\lambda_{N+1} - \lambda_N} \right)^k + \left(\frac{\lambda_N}{\lambda_N - \lambda_{N-1}} \right)^k \right), \quad N \in \mathbb{N}.$$

Since $a_N(f) = \sum_{n=1}^N a_n(f) - \sum_{n=1}^{N-1} a_n(f)$, the proof follows from the following independently interesting lemma.

Lemma 3.13. *Let $k > \ell \geq 0$. Then for every $f \in H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$ and $M \in \mathbb{N}$*

$$\left| \sum_{n=1}^M a_n(f) \right| \leq C(k, \ell, \lambda) \left(\frac{\lambda_{M+1}}{\lambda_{M+1} - \lambda_M} \right)^k \|f\|_{\infty, \ell}.$$

Proof. From [20, Lemma 3.5] we know that for every $M \in \mathbb{N}$

$$\left| \sum_{n=1}^M a_{\lambda_n}(f) \right| \leq 3 \left(\frac{\lambda_{M+1}}{\lambda_{M+1} - \lambda_M} \right)^k \sup_{\lambda_1 < x < \lambda_{M+1}} \left| \sum_{\lambda_n < x} a_{\lambda_n}(f) \left(1 - \frac{\lambda_n}{x}\right)^k \right|.$$

Then the conclusion follows from Theorem 3.6. \square

3.8. A Montel theorem. According to [11, Theorem 3.4], we for any frequency λ have a Montel-type theorem in $H_\infty^\lambda[\operatorname{Re} > 0]$. The aim now is to extend this result from $H_\infty^\lambda[\operatorname{Re} > 0]$ to $H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$, $\ell > 0$.

Recall from (5) that, given a frequency λ , there is an isometric and coefficient preserving identity $H_\infty^\lambda[\operatorname{Re} > 0] = \mathcal{H}_\infty(\lambda)$, so Montel's theorem holds in $\mathcal{H}_\infty(\lambda)$, i.e. for each bounded sequence (D_N) in $\mathcal{H}_\infty(\lambda)$ there is a subsequence (D_{N_k}) and $D \in \mathcal{H}_\infty(\lambda)$ such that for every $\varepsilon > 0$ the sequence of all translations $D_{N_k}^\varepsilon = \sum a_n e^{-\lambda_n \varepsilon} e^{-\lambda_n s}$ converges to $D \in \mathcal{H}_\infty(\lambda)$.

The important special case $\lambda = (\log n)$ was first proved by Bayart in [1], and within the setting of Hardy space of general Dirichlet series it was substantially extended in [8], [9], and [11]. See also the discussion in Section 3.13.

Theorem 3.14. *Let λ be an arbitrary frequency, and (f_N) a bounded sequence in $H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$, $\ell \geq 0$. Then there is a subsequence $(f_{N_j})_j$ of (f_N) and $f \in H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$ such that for every $u > 0$*

$$f_{N_j}(u + \cdot) \rightarrow f(u + \cdot) \quad \text{in } H_{\infty, \ell}^\lambda[\operatorname{Re} > 0].$$

In fact we are going to deduce this result from the following more general version.

Theorem 3.15. *Let (f_N) be a bounded sequence in $H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$. Assume that for every n the limit*

$$a_n = \lim_{N \rightarrow \infty} a_n(f_N)$$

exists. Then there exists $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ such that $a_n(f) = a_n$ for all n , and for every $u > 0$

$$f_N(u + \cdot) \rightarrow f(u + \cdot) \quad \text{in } H_{\infty,\ell}^\lambda[\operatorname{Re} > 0].$$

Proof. We define formally the Dirichlet series $E = \sum a_n e^{-\lambda n s}$. Then Theorem 3.6 implies that

$$(22) \quad |R_x^{\lambda,k}(E)(0)| = \lim_{N \rightarrow \infty} |R_x^{\lambda,k}(f_N)(0)| \leq C \sup_{N \in \mathbb{N}} \|f_N\|_{\infty,\ell}.$$

Hence, applying Theorem 2.16, we see that E is (λ, k) -Riesz summable on $[\operatorname{Re} > 0]$. Moreover, denoting by f its limit function, we by (22) have

$$\frac{|f(s)|}{(1 + |s|)^\ell} = \lim_{x \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{|R_x^{\lambda,k}(f_N)(s)|}{(1 + |s|)^\ell} \leq C \sup_{N \in \mathbb{N}} \|f_N\|_{\infty,\ell},$$

and so $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$. It remains to show that $f_N(u + \cdot) \rightarrow f(u + \cdot)$ in $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$. Fixing $u > 0$ and applying Theorem 3.7 to the bounded set $M = \{f_N \mid N \in \mathbb{N}\} \cup \{f\}$, we for every $\varepsilon > 0$ obtain some $x > 0$ such that

$$\begin{aligned} \|f(u + \cdot) - f_N(u + \cdot)\|_{\infty,\ell} &\leq \|f(u + \cdot) - R_x^{\lambda,k}(f)(u + \cdot)\|_{\infty,\ell} \\ &+ \|R_x^{\lambda,k}(f - f_N)(u + \cdot)\|_{\infty,\ell} + \|f(u + \cdot) - R_x^{\lambda,k}(f_N)(u + \cdot)\|_{\infty,\ell} \\ &\leq 2\varepsilon + \|R_x^{\lambda,k}(f - f_N)(u + \cdot)\|_{\infty,\ell}. \end{aligned}$$

Now tending $N \rightarrow \infty$ gives the claim. \square

Proof of Theorem 3.14. From Theorem 3.12 we know that $|a_n(g)| \leq C(n)\|g\|_{\infty,\ell}$ for every $g \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$. This allows to apply a diagonal process to find a subsequence (N_j) such that the limit

$$a_n = \lim_{j \rightarrow \infty} a_n(f_{N_j})$$

exists for every n . Then Theorem 3.15 finishes the proof. \square

3.9. Completeness. As we have already announced several times, all spaces $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$, $\ell \geq 0$ are indeed Banach spaces, which after all seems non-trivial.

Theorem 3.16. For each $\ell \geq 0$

$$H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$$

together with the norm $\|\cdot\|_{\infty,\ell}$ forms a Banach space.

Proof. Given a Cauchy sequence (f_N) , Theorem 3.14 proves the existence of a subsequence (N_j) and $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$, such that for every $u > 0$ we have

$$\lim_{j \rightarrow \infty} \|f_{N_j}(u + \cdot) - f(u + \cdot)\|_{\infty,\ell} = 0,$$

which in particular shows that (f_{N_j}) converges pointwise to f on $[\operatorname{Re} > 0]$. Hence for large N and $j = j(s)$ we obtain

$$\left| \frac{f(s) - f_N(s)}{(1 + s)^\ell} \right| \leq \left| \frac{f(s) - f_{N_j}(s)}{(1 + s)^\ell} \right| + \|f_{N_j} - f_N\|_{\infty,\ell} \leq |f(s) - f_{N_j}(s)| + \varepsilon,$$

and so $\|f - f_N\|_{\infty, \ell} \leq 2\varepsilon$ tending $j \rightarrow \infty$. \square

3.10. Behavior far left. The following result shows that the norm of any $f \in H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$ determines the growth of f on vertical abscissas close to the imaginary axis.

Theorem 3.17. *Let $f \in H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$ with $\ell \geq 0$. Then*

$$\|f\|_{\infty, \ell} = \lim_{\sigma \rightarrow 0} \sup_{t \in \mathbb{R}} \left| \frac{f(\sigma + it)}{(1 + (\sigma + it))^{\ell}} \right|.$$

The proof needs a well-known Phragmen-Lindelöf type lemma (see e.g. [3, p. 138]).

Lemma 3.18. *Let g be a bounded function on the strip $[\sigma_1 \leq \operatorname{Re} \leq \sigma_1]$, which is holomorphic on $[\sigma_1 < \operatorname{Re} < \sigma_1]$. Then the function*

$$L_g(\sigma) = \sup_{t \in \mathbb{R}} |g(\sigma + it)|, \quad \sigma \in [\sigma_1, \sigma_2].$$

is logarithmically convex on $[\sigma_1, \sigma_2]$, i.e. $\log L_g(\sigma)$ is convex on $[\sigma_1, \sigma_2]$.

Proof of Theorem 3.17. Define the function

$$g(s) = \frac{f(s + \varepsilon)}{(1 + s)^{\ell}},$$

and for each x

$$g_x(s) = \frac{\sum_{\lambda_n < x} a_n \left(1 - \frac{\lambda_n}{x}\right)^k e^{-s\lambda_n}}{(1 + s)^{\ell}} = \frac{R_x^{\lambda, k}(f)(s)}{(1 + s)^{\ell}},$$

which are both obviously holomorphic and bounded on $[\operatorname{Re} > 0]$. Note now that $L_{g_x}(\sigma)$ for each $x > 0$ is non-increasing for $\sigma > 0$; indeed, by Lemma 3.18 we know that $\log L_{g_x}(\sigma)$ is convex and moreover $\log L_{g_x}(\infty) = -\infty$. But since Corollary 3.8 implies that for every $u > 0$

$$g_x \rightarrow g \quad \text{uniformly on } [\operatorname{Re} > u] \text{ as } x \rightarrow \infty,$$

we see that $L_g(\sigma)$ is non-increasing for $\sigma > 0$. This gives

$$\|f\|_{\infty, \ell} = \sup_{\sigma > 0} |L_g(\sigma)| = \lim_{\sigma \rightarrow 0} |L_g(\sigma)|,$$

which is exactly what we were looking for. \square

3.11. Behavior far right. In contrast to what we did in Section 3.10 we now prove in which sense the growth of functions $f \in H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$ subsides on abscissas far right.

Theorem 3.19. *Let $\ell \geq 0$ and $\lambda = (\lambda_n)$ a frequency with $\lambda_1 > 0$. Then for every $f \in H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$ we have*

$$\lim_{\sigma \rightarrow \infty} \sup_{s \in [\operatorname{Re} > \sigma]} \frac{|f(s)|}{(1 + |s|)^{\ell}} = \lim_{\sigma \rightarrow \infty} \sup_{\substack{\delta \geq \sigma \\ t \in \mathbb{R}}} \frac{|f(\delta + it)|}{(1 + |t|)^{\ell}} = 0.$$

Proof. Observe, that it suffices to check the equality for the second limit. Moreover recall that by Corollary 3.9 the Dirichlet series $D = \sum a_n e^{-\lambda_n s}$ associated to f is (λ, k) -Riesz summable on $[\operatorname{Re} > 0]$ for every $k > \ell$. Hence, fixing $k > \ell$, $t \in \mathbb{R}$ and $\sigma > 0$, we apply Theorem 2.9 to the translation $\sum a_n e^{-\lambda_n it} e^{-\lambda_n s}$ and obtain for $\delta > \sigma$

$$f(\delta + it) = \frac{\delta^{1+k}}{\Gamma(1+k)} \int_{\lambda_1}^{\infty} e^{-\delta y} S_y^{\lambda, k}(D)(it) dy.$$

Hence with Theorem 3.6

$$\begin{aligned} \Gamma(1+k)|f(\delta + it)| &\leq \delta^{1+k} \int_{\lambda_1}^{\infty} e^{-\delta y} y^k \left| R_y^{\lambda, k}(D)(it) \right| dy \\ &\leq \|f\|_{\infty, \ell} (1 + |t|)^{\ell} \delta^{1+k} \int_{\lambda_1}^{\infty} e^{-\delta y} y^k dy \\ &\leq \|f\|_{\infty, \ell} (1 + |t|)^{\ell} \delta \int_{\lambda_1}^{\infty} e^{-\delta y} (\delta y)^k dy \\ &\leq \|f\|_{\infty, \ell} (1 + |t|)^{\ell} \int_{\delta \lambda_1}^{\infty} e^{-y} y^k dy \leq \|f\|_{\infty, \ell} (1 + |t|)^{\ell} \int_{\sigma \lambda_1}^{\infty} e^{-y} y^k dy, \end{aligned}$$

and so

$$\lim_{\sigma \rightarrow \infty} \sup_{\substack{\delta \geq \sigma \\ t \in \mathbb{R}}} \frac{|f(\delta + it)|}{(1 + |t|)^{\ell}} \leq \frac{\|f\|_{\infty, \ell}}{\Gamma(1+k)} \lim_{\sigma \rightarrow \infty} \int_{\sigma \lambda_1}^{\infty} e^{-y} y^k dy = 0. \quad \square$$

With a similar technique we obtain the following lemma – needed for the proof of Theorem 3.21 below.

Lemma 3.20. *Given a frequency λ with $\lambda_1 = 0$, let f be the limit function of $D = \sum a_n e^{-\lambda_n s}$ with $\sigma_c^{\lambda, m}(D) \in \mathbb{R}$ for some $m \geq 0$. Then for every $\gamma > 0$*

$$\lim_{\substack{\sigma \rightarrow \infty \\ \sigma > \sigma_c^{\lambda, m}(D)}} \sup_{\substack{s \in [\operatorname{Re} \geq \sigma] \\ |Im(s)| \leq \gamma}} |f(s) - a_1| = 0.$$

Proof. We may assume that $a_1 = 0$. Fix some $\sigma_0 > \sigma_c^{\lambda, m}(D)$ and $\gamma > 0$. Then by Theorem 2.9 (second part) the (λ, m) -Riesz means of D converge uniformly on $\sigma_0 + i[-\gamma, \gamma]$, hence

$$\sup_{|t| \leq \gamma} |R_y^{\lambda, m}(D)(\sigma_0 + it)| = C(\sigma_0, \gamma) < \infty.$$

Given $|t| > \gamma$, we apply (the integral formula from) Theorem 2.9 to the translation D_{σ_0+it} : For all $\sigma > 0$

$$f(\sigma_0 + \sigma + it) = \frac{\sigma^{1+m}}{\Gamma(1+m)} \int_{\lambda_2}^{\infty} e^{-\delta y} y^{1+m} R_y^{\lambda, m}(D_{\sigma_0})(it) dy.$$

Now we follow the estimates of the proof of Theorem 3.19. Replacing the expression $\|f\|_{\infty, \ell}(1+|t|)^{\ell}$ by $C(\sigma_0, \gamma)$, we conclude

$$\lim_{\sigma \rightarrow \infty} \sup_{\substack{s \in [\operatorname{Re} \geq \sigma] \\ |\operatorname{Im}(s)| \leq \gamma}} |f(s)| \leq \frac{C(\sigma_0, \gamma)}{\Gamma(1+m)} \lim_{\sigma \rightarrow \infty} \int_{\lambda_2 \sigma}^{\infty} e^{-y} y^m dy = 0. \quad \square$$

3.12. Finite order. In (2) we defined what it means that a holomorphic function $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$ has finite uniform order ν_f (see also [13, p.14]) The following result shows that our new scale of Banach spaces is intimately linked with this notion.

Theorem 3.21. *Let λ be a frequency, $\ell \geq 0$, and $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$ a holomorphic function which has a Riesz germ.*

- (1) *If $f \in H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$, then there is $C > 0$ such that for all $\sigma > 0$ and all $|t| > 1$*

$$|f(\sigma + it)| \leq C|t|^{\ell}.$$

In particular, if $f \in H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$, then f has finite uniform order $\nu_f \leq \ell$ on $[\operatorname{Re} > 0]$.

- (2) *Conversely, $f_{\mu} = f(\mu + \cdot) \in H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$ for all $\mu > 0$ whenever there are $C, t_0 > 0$ such that for all $\sigma > 0$ and all $|t| > t_0$*

$$|f(\sigma + it)| \leq C|t|^{\ell},$$

so in particular if f has finite uniform order $\nu_f < \ell$ on $[\operatorname{Re} > 0]$.

The proof relies on Lemma 3.20 and the following lemma.

Lemma 3.22. *Let $\ell \geq 0$ and λ an arbitrary frequency. Then for every $f \in H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$ we have*

$$\sup_{\substack{\sigma > 0 \\ t \in \mathbb{R}}} \frac{|f(\sigma + it)|}{(1+|t|)^{\ell}} < \infty.$$

Proof. Since $|a_1| \leq C\|f\|_{\infty, \ell}$ by Theorem 3.12, we may assume that $\lambda_1 > 0$. Note first that by Theorem 3.19 there is some $\sigma_0 > 0$ such that

$$\sup_{\substack{\sigma > \sigma_0 \\ t \in \mathbb{R}}} \frac{|f(\sigma + it)|}{(1+|t|)^{\ell}} \leq 1.$$

On the other hand, for $\sigma \leq \sigma_0$ and $t \in \mathbb{R}$ we have that

$$\frac{|f(\sigma + it)|}{(1+|t|)^{\ell}} = \frac{|f(\sigma + it)|}{(1+|\sigma + it|)^{\ell}} \frac{(1+|\sigma + it|)^{\ell}}{(1+|t|)^{\ell}} \leq \frac{|f(\sigma + it)|}{(1+|\sigma + it|)^{\ell}} (1+\sigma_0)^{\ell}. \quad \square$$

Proof of Theorem 3.21. (1) Let $f \in H_{\infty, \ell}^{\lambda}[\operatorname{Re} > 0]$. Then by Lemma 3.22 we know that

$$\exists R > 0 \forall \sigma > 0, t \in \mathbb{R}: |f(\sigma + it)| \leq R(1 + |t|)^{\ell}.$$

Then f satisfies (2) with $t_0 = 1$ and $C = R2^{\ell}$, since $(1 + |t|)^{\ell} \leq 2^{\ell}|t|^{\ell}$ for all $|t| \geq 1$.

(2) We assume that there are $C, t_0 > 0$ such that for all $\sigma > 0$ and all $|t| > |t_0|$

$$|f(\sigma + it)| \leq C|t|^{\ell} \leq C(1 + |t|)^{\ell}.$$

Fix some $\mu > 0$, and assume without loss of generality that $\lambda_1 = 0$. Then we know from Lemma 3.20 (with $\gamma = t_0$) that there is some $\sigma_0 > \mu$ with $|f(\sigma + it)| \leq C$ for all $\sigma > \sigma_0$ and $|t| \leq t_0$. Since the continuous function f_{μ} is bounded on the rectangle $[\mu, \sigma_0] \times [-t_0, t_0]$, we have that

$$\|f_{\mu}\|_{\infty, \ell} \leq \sup_{\substack{\sigma > 0 \\ |t| \geq t_0}} \frac{|f(s)|}{|(1+s)^{\ell}|} + \sup_{\substack{\sigma_0 > \sigma > \mu \\ |t| \leq t_0}} \frac{|f(s)|}{|(1+s)^{\ell}|} + \sup_{\substack{\sigma > \sigma_0 \\ |t| \leq t_0}} \frac{|f(s)|}{|(1+s)^{\ell}|},$$

which then is finite. \square

3.13. Equivalence. Recall that λ -Dirichlet series $D = \sum a_n e^{-\lambda_n s}$ converge on half-planes in the complex plane, where they define holomorphic functions, and in general the largest possible half-planes of convergence, uniform convergence, or absolute convergence differ.

In the early days of the theory, a very prominent research problem was to characterize the class of those frequencies λ for which boundedness of the limit function of D on $[\operatorname{Re} > 0]$ implies uniform convergence of D on every smaller half plane $[\operatorname{Re} > \varepsilon]$. In other words, if $\mathcal{D}_{\infty}(\lambda)$ stands for all λ -Dirichlet series $D = \sum a_n e^{-\lambda_n s}$ that on $[\operatorname{Re} > 0]$ converges to a bounded (holomorphic) function, then the question is whether all these series even converges uniformly on all smaller half-planes $[\operatorname{Re} > \varepsilon]$.

Bohr was the first who in [5] managed to isolate a prominent class of such λ 's, namely those for which there is some $\beta > 0$ such that

$$(BC) \quad \frac{1}{\lambda_{n+1} - \lambda_n} = O(e^{\beta \lambda_n});$$

slightly more precise, we in this case say that λ satisfies (BC) with constant β .

We say that 'Bohr's theorem holds for λ ', whenever every $D \in \mathcal{D}_{\infty}(\lambda)$ converges uniformly on $[\operatorname{Re} > \varepsilon]$ for every $\varepsilon > 0$. In particular, frequencies λ with (BC) satisfy Bohr's theorem. Later this result was extended by Landau in [18] assuming the less restrictive assumption

$$(LC) \quad \forall \delta > 0 \exists C > 0 \forall n: \lambda_{n+1} - \lambda_n \geq C e^{-e^{\delta \lambda_n}},$$

and very recently, Bayart in [2] added a further interesting condition, which provides a nontrivial extension of (LC): A frequency λ satisfies (NC) if

$$(NC) \quad \forall \delta > 0 \exists C > 0 \forall m > n: \log \left(\frac{\lambda_m + \lambda_n}{\lambda_m - \lambda_n} \right) + (m - n) \leq C e^{\delta \lambda_n}.$$

Examples show that any concrete λ may or may not satisfy Bohr's theorem, and we refer to [2] and [20] for detailed information on all this.

In this context, an important recent achievement is that the property ' λ satisfies Bohr's theorem ' in fact is equivalent to a number of facts which seem absolutely unavoidable for a reasonable abstract theory of general Dirichlet series. The following so-called 'equivalence theorem' is taken from [6, Theorem 4.6]) (in fact, the main part of this result was proved earlier in [9, Theorem 2.16] and [10, Theorem 5.1]).

Theorem 3.23. *Let λ be a frequency. Then the following are equivalent:*

- (1) λ satisfies Bohr's theorem.
- (2) $\mathcal{D}_\infty(\lambda)$ is a Banach space.
- (3) $\mathcal{D}_\infty(\lambda) = \mathcal{H}_\infty(\lambda)$ isometrically and coefficient preserving.
- (4) $\mathcal{D}_\infty(\lambda) = H_\infty^\lambda[\operatorname{Re} > 0]$ isometrically and coefficient preserving.
- (5) λ satisfies Montel's theorem in $\mathcal{D}_\infty(\lambda)$.

Let us take some time to explain this result in more detail. Note first that $\mathcal{D}_\infty(\lambda)$ endowed with the sup norm $\|D\|_\infty = \sup_{\operatorname{Re} > 0} |f(s)|$, where f denotes the limit function of D , forms a normed space. But unfortunately this space in general is not complete, and so the theorem ensures that this only holds true if λ satisfies Bohr's theorem – in particular under each of the conditions of Bayart, Bohr, or Landau.

To see a non-trivial example, $(\sqrt{\log n})$ doesn't satisfy (BC) but (LC). Hence, Bohr's theorem holds for $(\sqrt{\log n})$, and so $\mathcal{D}_\infty((\sqrt{\log n}))$ forms a Banach space – a seemingly non-trivial fact.

An important step is to understand that the Banach space $H_\infty^\lambda[\operatorname{Re} > 0]$ always contains the (in general non-complete) space $\mathcal{D}_\infty(\lambda)$ isometrically (see [20, Proposition 3.4]). Hence both spaces in general differ, but in view of the theorem they coincide iff Bohr's theorem holds true for λ .

Moreover, as already mentioned in Section 3.8, Bayart proved a 'Montel type theorem' for $\mathcal{D}_\infty((\log n)) = \mathcal{H}_\infty((\log n))$, which turned out to be a corner stone of the ordinary theory. The equivalence theorem shows that the analog result holds true for $\mathcal{D}_\infty(\lambda)$ if and only if λ satisfies Bohr's theorem.

The idea now is to extend parts of Theorem 3.23 to the new setting. Given a frequency λ and $\ell \geq 0$, let

$$\mathcal{D}_{\infty,\ell}(\lambda)$$

denote the space of all λ -Dirichlet series $D = \sum a_n e^{-\lambda_n s}$ that are (λ, ℓ) -Riesz summable on all of $[\operatorname{Re} > 0]$ and have limit functions $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$. Together with the norm $\|D\|_{\infty,\ell} = \|f\|_{\infty,\ell}$, this leads to another scale of normed spaces of λ -Dirichlet series.

Recall again from Corollary 3.9 that the λ -Riesz germ of each $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ is (λ, k) -Riesz summable for each $k > \ell$, and so one of the questions in the following will be, under with additional assumptions on λ , Corollary 3.9 even holds for $k = \ell$.

As mentioned above, $\mathcal{D}_\infty(\lambda)$ is always an isometric subspace of $H_\infty^\lambda[\operatorname{Re} > 0]$, hence in the particular case $\ell = 0$ we by definition have the isometric equality

$$(23) \quad \mathcal{D}_{\infty,0}(\lambda) = \mathcal{D}_\infty(\lambda).$$

Together with Corollary 3.11 we see that Theorem 3.23 settles the case $\ell = 0$.

For $\ell > 0$ the following result partly serves as a sort of substitute of Theorem 3.23.

Theorem 3.24. *Let λ be a frequency and $\ell \geq 0$. Then the following are equivalent:*

- (1) $\mathcal{D}_{\infty,\ell}(\lambda)$ satisfies Montel's theorem.
- (2) $\mathcal{D}_{\infty,\ell}(\lambda)$ is a Banach space.
- (3) The λ -Riesz germ of every $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ is (λ, ℓ) -Riesz summable on $[\operatorname{Re} > 0]$.
- (4) $\mathcal{D}_{\infty,\ell}(\lambda) = H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ holds isometrically and coefficient preserving.

Proof. If $\mathcal{D}_{\infty,\ell}(\lambda)$ satisfies Montel's theorem, then the same proof as for Theorem 3.16 shows that $\mathcal{D}_{\infty,\ell}(\lambda)$ is complete, so we have that (1) \Rightarrow (2). Let us show that (2) \Rightarrow (4), and fix some $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ with associated λ -Riesz germ D . By Corollary 3.8 we know that for every $u > 0$ and every $k > \ell$

$$\lim_{x \rightarrow \infty} R_x^{\lambda,k}(f)(u + \cdot) = f(u + \cdot) \text{ in } H_{\infty,\ell}^\lambda[\operatorname{Re} > 0].$$

But since

$$R_x^{\lambda,k}(D_u) = R_x^{\lambda,k}(f)(u + \cdot) \in \mathcal{D}_{\infty,\ell}(\lambda),$$

where $D_u = \sum a_n e^{-\lambda_n u} e^{-\lambda_n s} \in \mathcal{D}_{\infty,\ell}(\lambda)$ as usual denotes the translation of D about $u > 0$, we eventually see that $D \in \mathcal{D}_{\infty,\ell}(\lambda)$. Hence $\mathcal{D}_{\infty,\ell}(\lambda) = H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$, the claim from (4). The equivalence (3) \Leftrightarrow (4) is obvious, so that it finally suffices to check that (4) \Rightarrow (1). But if $\mathcal{D}_{\infty,\ell}(\lambda) = H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ holds true, then trivially by Theorem 3.14 Montel's theorem holds in $\mathcal{D}_{\infty,\ell}(\lambda)$. \square

The case $\ell = 0$ from the preceding theorem allows another interesting remark on Theorem 3.23.

Remark 3.25. *Bohr's theorem is valid for λ if and only if every Dirichlet series which for some $m \geq 0$ is somewhere (λ, m) -Riesz summable and has a limit function that extends holomorphically to a bounded function on $[\operatorname{Re} > 0]$, converges uniformly on $[\operatorname{Re} > \sigma]$ for every $\sigma > 0$.*

For concrete frequencies λ , in particular $\lambda = (\log n)$, the following result is our main application of Theorem 3.24.

Theorem 3.26. *Let $\ell > 0$. If (BC) holds for λ , then the λ -Riesz germ D of every $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ is (λ, ℓ) -Riesz summable on $[\operatorname{Re} > 0]$. In particular, all equivalent statements from Theorem 3.24 hold.*

It seems very interesting to decide whether or not the preceding result even holds under the condition (LC) or more generally (NC).

Divided into three lemmas of independent interest, the proof of Theorem 3.26 is given at the very end of this section. The first lemma gives a necessary condition

under which $H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$ and $\mathcal{D}_{\infty,\ell}(\lambda)$ coincide, and is based on a Phragmen-Lindelöf type argument again borrowed from [13].

Lemma 3.27. *Let $\ell > 0$ and $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$. Assume that there exist $\sigma_0 > 0$ and $0 \leq \ell' < \ell$ such that $f_{\sigma_0} = f(\sigma_0 + \cdot) \in H_{\infty,\ell'}^\lambda[\operatorname{Re} > 0]$. Then the λ -Riesz germ D of f is (λ, ℓ) -Riesz summable on $[\operatorname{Re} > 0]$, i.e. $D \in \mathcal{D}_{\infty,\ell}(\lambda)$.*

Proof. Fix $\varepsilon > 0$. By Theorem 3.21, part (1) there is $D_0 > 0$ such that for all $\sigma > 0$ and $|t| > 1$

$$(24) \quad |f(\sigma + it)| \leq D_0 |t|^\ell,$$

and some $D_1 > 0$ such that for all $\sigma > \sigma_0$ and $|t| > 1$

$$(25) \quad |f(\sigma + it)| \leq D_1 |t|^{\ell'}.$$

Combining (24) and (25) by [13, Theorem 14] (another Phragmen-Lindelöf principle), there are $C, t_0 > 0$ such that for every $\varepsilon \leq \sigma \leq \sigma_0 + 1$ and every $|t| \geq t_0$

$$(26) \quad |f(\sigma + it)| \leq C |t|^{\kappa(\sigma)},$$

where $\kappa : [\varepsilon, \sigma_0 + 1] \rightarrow [0, \infty[$ is the affine linear function linking the points (ε, ℓ) and $(\sigma_0 + 1, \ell')$. We claim that

$$f(\varepsilon + \sigma + \cdot) \in H_{\infty,\kappa(\beta)}^\lambda[\operatorname{Re} > 0] \text{ for every } \sigma > 0,$$

and we do this with Theorem 3.21, part (2) showing for all $\sigma > 0$ and $|t| \geq \max\{1, t_0\}$

$$(27) \quad |f(\varepsilon + \sigma + it)| \leq \max\{C, D_1\} |t|^{\kappa(\sigma)}.$$

Indeed, since $\kappa \geq \ell'$ on $[\varepsilon, \sigma_0 + 1]$, this is immediate from (25) and (26).

Since $\kappa < \ell$ on $[\varepsilon, \sigma_0 + 1]$ (recall that $\ell' < \ell$), we by Corollary 3.9 obtain from (3.13) that $\sigma_c^{\lambda,\ell}(D_\varepsilon) \leq 0$ for every $\varepsilon > 0$, or equivalently $\sigma_c^{\lambda,\ell}(D) \leq \varepsilon$ for every $\varepsilon > 0$. \square

The next tool, based on the preceding one, is a reformulation of [13, Theorem 44]. For the sake of completeness we include an argument within our new setting.

Lemma 3.28. *Let $\ell > 0$, and $f \in H_{\infty,\ell}^\lambda[\operatorname{Re} > 0]$. Then the λ -Riesz germ D of f is (λ, ℓ) -Riesz summable on $[\operatorname{Re} > 0]$, i.e. $D \in \mathcal{D}_{\infty,\ell}(\lambda)$, provided $\sigma_a(D) < \infty$.*

Proof. Since $\sigma_a(D) < \infty$, we have that

$$f(s) = \sum_{n=1}^{\infty} a_n(f) e^{-\lambda_n s}, \quad s \in [\operatorname{Re} > \sigma_a(D)],$$

and so we see that f is bounded on $[\operatorname{Re} > \sigma_a(D) + 1]$. Hence Lemma 3.27 with $\sigma_0 = \sigma_a(D) + 1$ and $\ell' = 0$ implies $D \in \mathcal{D}_{\infty,\ell}(\lambda)$. \square

For the last lemma we need further notation and information. Given a frequency λ , define

$$L(\lambda) := \sup_{D \in \mathcal{D}(\lambda)} \sigma_a(D) - \sigma_c(D),$$

where we recall that $\sigma_c(D)$ and $\sigma_a(D)$ define the abscissas of convergence and absolute convergence (with respect to ordinary summation). Then straightforward arguments show that

$$(28) \quad L(\lambda) = \sigma_c \left(\sum e^{-\lambda_n s} \right) = \sigma_a \left(\sum e^{-\lambda_n s} \right).$$

With a less obvious arguments, Bohr in [4, §3, Hilfssatz 4] showed that his condition (BC) implies $L(\lambda) < \infty$. In passing we remark (although this is not needed for what's coming) that by [4, §3, Hilfssatz 4]

$$(29) \quad L(\lambda) = \limsup_{N \rightarrow \infty} \frac{\log(N)}{\lambda_N}.$$

It is evident that $L((n)) = 0$ and $L((\log n)) = 1$.

Lemma 3.29. *Let $\ell > 0$, and $f \in H_{\infty, \ell}^\lambda[\operatorname{Re} > 0]$ with its associated λ -Riesz germ D . Then $\sigma_a(D) \leq L(\lambda) + \beta\ell < \infty$, whenever λ satisfies (BC) with exponent β .*

To see an example, we note that for $\lambda = (n)$ we get the conclusion $\sigma_a(D) \leq 0$, whereas for $\lambda = (\log n)$ we obtain that $\sigma_a(D) \leq 1 + \ell$.

Proof. Condition (BC) for λ with exponent β means that there is some $C > 0$ such that for all n

$$\frac{1}{\lambda_{n+1} - \lambda_n} \leq C e^{\lambda_n \beta}.$$

Adding more elements to λ , we assume without loss of generality that $\lambda_{n+1} - \lambda_n \leq 1$ for all n (see the first part of the proof of [20, Theorem 4.2] for details). We now claim that the Bohr coefficients $(a_n(D)) = (a_n(f))$ of D for every $\delta > 0$ and $k > \ell$ satisfy

$$(30) \quad A = \sup_n |a_n(f) e^{-(\beta k + \delta)\lambda_n}| < \infty.$$

Indeed, given $\delta > 0$ and $k > \ell$, by Theorem 3.12 we conclude that for every n and some constant $C = C(k, \ell, \lambda_1)$

$$\begin{aligned} |a_n(f)| &\leq C \left(\left(\frac{\lambda_{n+1}}{\lambda_{n+1} - \lambda_n} \right)^k + \left(\frac{\lambda_n}{\lambda_n - \lambda_{n-1}} \right)^k \right) \|f\|_{\infty, \ell} \\ &\leq 2C \left(\lambda_{n+1}^k e^{\beta k \lambda_n} + \lambda_n^k e^{\beta k \lambda_{n-1}} \right) \|f\|_{\infty, \ell} \\ &\leq 2C \left((\lambda_{n+1})^k e^{\beta k \lambda_n} \right) \|f\|_{\infty, \ell}. \end{aligned}$$

Hence, choosing $D = D(\delta, k, \lambda) > 0$ such that $(1 + \lambda_n)^k \leq D e^{\delta \lambda_n}$ for all n , we finally obtain (30), namely that for all n

$$|a_n(f)| \leq 2CD e^{(\delta + \beta k)\lambda_n} \|f\|_{\infty, \ell}.$$

Then by (30) and by the fact that under Bohr's condition $L(\lambda) < \infty$, we see that for every $\varepsilon > 0$

$$\begin{aligned} \sum |a_n| e^{-((\beta k + \delta) + (L(\lambda) + \varepsilon))\lambda_n} &= \sum |a_n e^{-(\beta k + \delta)\lambda_n}| e^{-(L(\lambda) + \varepsilon)\lambda_n} \\ &\leq A \sum e^{-(L(\lambda) + \varepsilon)\lambda_n} < \infty. \end{aligned}$$

Consequently, $\sigma_a(D) \leq (\beta k + \delta) + (L(\lambda) + \varepsilon)$, which is the conclusion (whenever $k \rightarrow \ell$, $\delta \rightarrow 0$, and $\varepsilon \rightarrow 0$). \square

Proof of Theorem 3.26. Applying first Lemma 3.29 and then second Lemma 3.28, we finally obtain as desired Theorem 3.26. \square

4. Appendix

This section is devoted to the proofs of Theorem 2.7, Theorem 2.8, Theorem 2.9, and Theorem 2.16. These are those results from [13], and in a few cases even improvements of them, which were of central importance for what we tried to explain in the preceding sections. In fact, most of the proofs from [13] are not given in full detail and only sketches – together with this appendix our article is indeed self contained.

We need a couple of technical lemmas to properly prepare these proofs. To do so, recall from Section 2.2 the definition of summatory functions $S_x^{\lambda,r}(D)(s)$ with respect to a frequency λ , a Riesz weight r , and a λ -Dirichlet series D .

4.1. Integral forms. Given a frequency λ and a λ -Dirichlet series D , we in Section 2.2 defined the summatory function $S_x^{\lambda,k}(D)$. We start with the following integral form of this function, which, simple as it is, seems to be the seed for most of the coming arguments.

Lemma 4.1. *Let $D \in \mathcal{D}(\lambda)$, r a Riesz weight, and $x \geq 0$. Then for all $s, w \in \mathbb{C}$*

$$(31) \quad S_x^{\lambda,r}(D)(s+w) = - \int_0^x S_t^{\lambda,0}(D)(s) \left(e^{-w \bullet} r(x, \bullet) \right)'(t) dt,$$

and in particular,

$$(32) \quad S_x^{\lambda,r}(D)(s) = - \int_0^x S_t^{\lambda,0}(D)(s) r(x, \bullet)'(t) dt.$$

Moreover, if $C = \sum a_n$ is a series and $k \geq 0$, then for $r(x, t) = (x - t)^{k+1}$

$$(33) \quad \frac{d}{dt} S_t^{\lambda,k+1}(C)|_{t=x} = (k+1) S_x^{\lambda,k}(C).$$

Proof. It suffices to do the proof for $s = 0$. We write $S_x^{\lambda,r}(D)(w)$ as a Stieltjes integral,

$$S_x^{\lambda,r}(D)(w) = \sum_{\lambda_n < x} a_n e^{-\lambda_n w} r(x, \lambda_n) = \int_0^x e^{-wt} r(x, t) dS_t^{\lambda,0}(D)(0),$$

and then (31) follows by partial integration. Clearly, (32) is the special case $w = 0$ in (31). The proof of (33) follows applying (32) two times:

$$\begin{aligned} \frac{d}{dt} S_t^{\lambda, k+1}(D) &= \frac{d}{dt} \int_0^t S_u^{\lambda, 0}(D)(k+1)(t-u)^k du \\ &= (k+1)k \int_0^t S_u^{\lambda, 0}(D)(t-u)^{k-1} du = (k+1)S_t^{\lambda, k}(D). \quad \square \end{aligned}$$

Recall that, the first statement of Theorem 2.9 claims (λ, k) -Riesz summability of the λ -Dirichlet series on the half plane $[\text{Re} > \text{Re } s_0]$, given (λ, k) -Riesz summability at s_0 . Roughly speaking, the strategy for proving this claim is to substitute the expression $S_t^{\lambda, 0}(D)(s)$ of (31) by $S_t^{\lambda, k}(D)(s)$. In other terms, we want to increase the order of the summatory function of f on the right hand side of (31) from zero to k .

The following lemma is the first step in this direction – partial integration allows to increase the order zero to the order $m+1$, where $m \in \mathbb{N}$ with $m < k \leq m+1$. Then, for the case $k \notin \mathbb{N}$, we utilize the subsequent Lemma 4.4 to drop the order to k as desired.

Lemma 4.2. *Let $D \in \mathcal{D}(\lambda)$, r a Riesz weight, and $m \in \mathbb{N}_0$. Then for all $s, w \in \mathbb{C}$ and all $x > 0$*

$$S_x^{\lambda, r}(D)(s+w) = -e^{-wx} S_x^{\lambda, r}(D)(s) + \frac{(-1)^m}{(m+1)!} \int_0^x S_t^{\lambda, m+1}(D)(s) \partial^{m+2} h_w(t) dt,$$

where $h_w(t) = (e^{-wt} - e^{-wx})r(x, t)$. In particular,

$$\begin{aligned} (34) \quad S_x^{\lambda, r}(D)(s+w) &- \frac{1}{\Gamma(m+2)} \int_0^x S_t^{\lambda, m+1}(D)(s) w^{m+2} e^{-wt} r(x, t) dt \\ &= -e^{-wt} S_x^{\lambda, r}(D)(s) + \int_0^x S_t^{\lambda, m+1}(D)(s) (g_1(t) + g_2(t)) dt, \end{aligned}$$

where

$$\begin{aligned} g_1(t) &= \sum_{j=1}^{m+1} \binom{m+2}{j} (-w)^j e^{-wt} r(x, \bullet)^{(m+2-j)}(t) \\ g_2(t) &= (e^{-wt} - e^{-wx}) r(x, \bullet)^{(m+2)}(t). \end{aligned}$$

Proof. By partial integration and Lemma 4.1 we have

$$\begin{aligned} S_x^{\lambda, r}(D)(s+w) &= - \int_0^x S_t^{\lambda, 0}(D)(s) \left(e^{-w\bullet} r(x, \bullet) \right)'(t) dt \\ &= -e^{-wx} \int_0^x S_t^{\lambda, 0}(D)(s) r(x, \bullet)'(t) dt - \int_0^x S_t^{\lambda, 0}(D)(s) \left((e^{-w\bullet} - e^{-wx}) r(x, \bullet) \right)'(t) dt. \end{aligned}$$

Using (32) we know

$$e^{-wx} \int_0^x S_t^{\lambda, 0}(D)(s) r(x, \bullet)'(t) dt = e^{-wx} S_x^{\lambda, r}(D)(s),$$

and get

$$S_x^{\lambda,r}(D)(s+w) = -e^{-wx} S_x^{\lambda,r}(D)(s) - \int_0^x S_t^{\lambda,0}(D)(s) ((e^{-w\bullet} - e^{-wx}) r(x, \bullet))'(t) dt.$$

Finally, combining partial integration and (33), gives the first conclusion. For the proof of the second claim note that

$$\partial^{m+2} h_w(t) = (-w)^{m+2} e^{-wt} r(x, t) + g_1(t) + g_2(t),$$

which finishes the proof. \square

We go on analyzing the integral on the right side of the preceding lemma. We split it into two summands – one which will lead us to the integral describing the (λ, k) -Riesz sum in Theorem 2.9, and an other one which we in fact control whenever x increases to infinity.

Lemma 4.3. *Let $D \in \mathcal{D}(\lambda)$, r a Riesz weight, $k > 0$ and $m \in \mathbb{N}_0$ with $m < k < m + 1$. Then for all $s, w \in \mathbb{C}$ and all $x > 0$*

$$\begin{aligned} S_x^{\lambda,r}(D)(s+w) &= \frac{1}{\Gamma(m+2)} \int_0^x S_t^{\lambda,m+1}(D)(s) w^{m+2} e^{-wt} r(x, t) dt \\ &= -e^{-wt} S_x^{\lambda,r}(D)(s) \\ &\quad + C(k) \int_0^x S_y^{\lambda,k}(D)(s) \int_y^x (t-y)^{m-k} (g_1(t) + g_2(t)) dt dy, \end{aligned}$$

where g_1 and g_2 are as in (4.2) and $C(k) = (-1)^m \frac{1}{(m+1)\Gamma(1+k)\Gamma(1+m-k)}$.

The proof of this lemma needs further independently useful preparation, and we postpone it to the next section.

4.2. Changing orders. Given a frequency λ and a series $C = \sum a_n$, we present two devices which allow to increase or decrease the order κ of a given summatory function $S_x^{\lambda,\kappa}(C)$. Both lemmas are indispensable technical tools for the proofs of Theorem 2.9, Theorem 2.16, and Theorem 2.8.

We begin collecting a few basic fact about the classical Gamma function given by

$$\Gamma(z) = \int_0^\infty x^{z-1} e^{-x} dx, \quad \operatorname{Re} z > 0.$$

Recall that for every $z \in [\operatorname{Re} > 0]$

$$(35) \quad \Gamma(1+z) = \Gamma(z)z.$$

Moreover, the Beta function is given by

$$B(p, q) = \int_0^1 y^{p-1} (1-y)^{q-1} dy, \quad p, q \in [\operatorname{Re} > 0],$$

which in terms of Gamma functions reads

$$(36) \quad B(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}.$$

This shows that for $p, q > -1$ with $q > p$ and $x > 0$

$$(37) \quad \int_0^x y^p(x-y)^{q-p-1}dy = x^q \frac{\Gamma(p+1)\Gamma(q-p)}{\Gamma(q+1)}.$$

which indeed follows by a simple substitution:

$$\begin{aligned} \int_0^x y^p(x-y)^{q-p-1}dy &= x^q \int_0^1 x^{-1}\left(\frac{y}{x}\right)^p \left(1-\frac{y}{x}\right)^{q-p-1}dy \\ &= x^q \int_0^1 \alpha^p(1-\alpha)^{q-p-1}d\alpha = x^q \frac{\Gamma(p+1)\Gamma(q-p)}{\Gamma(q+1)}. \end{aligned}$$

We several times need (37) in combination with the fact that for arbitrary $c, d \in \mathbb{R}$, $w \in \mathbb{C}$ and $y < x$

$$(38) \quad \int_y^x e^{-wt}(t-y)^c(x-t)^d dt = (x-y)^{c+d+1} e^{-wy} \int_0^1 e^{-w\beta(x-y)} \beta^c(1-\beta)^d d\beta;$$

indeed,

$$\begin{aligned} \int_y^x e^{-wt}(t-y)^c(x-t)^d dt &= \int_0^{x-y} e^{-w(y+\alpha)} \alpha^c(x-y-\alpha)^d d\alpha \\ &= (x-y)^{c+d+1} \int_0^1 e^{-w(y+\beta(x-y))} \beta^c(1-\beta)^d d\beta \\ &= (x-y)^{c+d+1} e^{-wy} \int_0^1 e^{-w\beta(x-y)} \beta^c(1-\beta)^d d\beta. \end{aligned}$$

We are ready for the first lemma taken from [13, Lemma 6]. Following the idea of our article we for the sake of completeness repeat its proof.

Lemma 4.4. *Let $C = \sum a_n$ be a series, λ a frequency, and $\kappa, \mu > 0$. Then for all $x > 0$*

$$S_x^{\lambda, \kappa + \mu}(C) = \frac{\Gamma(\kappa + \mu + 1)}{\Gamma(\kappa + 1)\Gamma(\mu)} \int_0^x S_u^{\lambda, \kappa}(C)(x-u)^{\mu-1} du.$$

Moreover, if $\kappa > 0$, $\mu < 1$ and $\mu \leq \kappa$, then for all $x > 0$

$$S_x^{\lambda, \kappa - \mu}(C) = \frac{\Gamma(\kappa - \mu + 1)}{\Gamma(\kappa + 1)\Gamma(1 - \mu)} \int_0^x \frac{d}{du} S_u^{\lambda, \kappa}(C)(x-u)^{-\mu} du.$$

Proof. For the proof of the first formula we start with the integral on the left side, and use (32), Fubini's theorem and (37) to obtain

$$\begin{aligned}
& \int_0^x S_u^{\lambda,k}(C)(x-u)^{\mu-1} du \\
&= \int_0^x k \int_t^x S_t^{\lambda,0}(C)(u-t)^{k-1}(x-u)^{\mu-1} dudt \\
&= k \int_0^x S_t^{\lambda,0}(C) \int_t^x (u-t)^{k-1}(x-u)^{\mu-1} dudt \\
&= \frac{\Gamma(k)\Gamma(\mu)}{\Gamma(k+\mu)} k \int_0^x S_t^{\lambda,0}(C)(x-t)^{\mu+k-1} dt = \frac{\Gamma(k)\Gamma(\mu)}{\Gamma(k+\mu)} \frac{k}{\mu+k} S_t^{\lambda,\mu+k}(C),
\end{aligned}$$

the conclusion. To prove the second formula we use (33) and the first formula to see that for all $x > 0$

$$\begin{aligned}
S_x^{\lambda,\kappa-\mu}(C) &= \frac{1}{\kappa-\mu+1} \frac{d}{dx} S_x^{\lambda,\kappa+(1-\mu)}(C) \\
&= \frac{\Gamma(\kappa-\mu+1)}{\Gamma(\kappa+1)\Gamma(1-\mu)} \frac{d}{dx} \int_0^x S_u^{\lambda,\kappa}(C)(x-u)^{-\mu} du.
\end{aligned}$$

On the other hand, integrating by parts we have

$$\int_0^x S_u^{\lambda,\kappa}(C)(x-u)^{-\mu} du = \frac{1}{1-\mu} \int_0^x \frac{d}{du} S_u^{\lambda,\kappa}(C)(x-u)^{1-\mu} du.$$

For the desired formula we differentiate the right side with respect to x . \square

As announced we use the preceding lemma to add the still missing proof of Lemma 4.3.

Proof of Lemma 4.3. By Lemma 4.2 for every $s, w \in \mathbb{C}$

$$S_x^{\lambda,r}(D)(s+w) = -e^{-wx} S_x^{\lambda,r}(D)(s) + \frac{(-1)^m}{\Gamma(m+2)} \int_0^x S_t^{\lambda,m+1}(D)(s) \partial^{m+2} h_w(t) dt,$$

where $h_w(t) = (e^{-wt} - e^{-wx})r(x, t)$. We calculate

$$\partial^{m+2} h_w(t) = (-w)^{m+2} e^{-wt} r(x, t) + g_1(t) + g_2(t).$$

The aim is to replace the summatory function of order $m+1$ by the summatory function of order k . If $k = m+1$, then we are fine – but if $k < m+1$, then we need Lemma 4.4 (with the choices $\mu = m-k+1$ and $\kappa = k$):

$$S_t^{\lambda,m+1}(D)(s) = C(m, k) \int_0^t S_y^{\lambda,k}(D)(s)(t-y)^{m-k} dy,$$

where

$$C(k) = \frac{\Gamma(1+m)}{\Gamma(1+k)\Gamma(1+m-k)}.$$

For the rest of the proof we assume that $k < m + 1$; in fact, it is easier to handle the case $k = m + 1$ following basically the same lines. We apply Fubini's theorem and obtain for $i = 1, 2$

$$\begin{aligned} C(k)^{-1} \int_0^x S_t^{\lambda, m+1}(D)(s) g_i(t) dt &= \int_0^x \int_0^t S_y^{\lambda, k}(D)(s) (t-y)^{m-k} dy g_i(t) dt \\ &= \int_0^\infty S_y^{\lambda, k}(D)(s) \int_0^x \chi_{(0,t)}(y) (t-y)^{m-k} g_i(t) dt dy \\ &= \int_0^x S_y^{\lambda, k}(D)(s) \int_y^x (t-y)^{m-k} g_i(t) dt dy. \end{aligned}$$

Combining all this gives the conclusion. \square

Recall that Theorem 2.9 provides us with an integral representation for (λ, k) -Riesz limits. In fact this representation comes from the integral on the left side of the equality given in Lemma 4.3. As discussed earlier we have to switch from the order $m + 1$ to the order k which will be done using the following general device.

Lemma 4.5. *Let $D \in \mathcal{D}(\lambda)$ and $0 < p < q$. Then for all $w, s \in \mathbb{C}$ and all $x > 0$*

$$\int_0^\infty S_x^{\lambda, q}(D)(s) w^{q+1} e^{-wx} dx = \frac{\Gamma(q+1)}{\Gamma(p+1)} \int_0^\infty S_u^{\lambda, p}(D)(s) w^{p+1} e^{-wu} du.$$

Proof. Again we assume that $s = 0$. We first use Lemma 4.4 (applied to $k = p$ and $\mu = q - p$) to show that

$$\begin{aligned} &\int_0^\infty S_x^{\lambda, q}(D)(0) e^{-wx} dx \\ &= \frac{\Gamma(q+1)}{\Gamma(p+1)\Gamma(q-p)} \int_0^\infty \int_0^x S_u^{\lambda, p}(D)(0) (x-u)^{q-p-1} du e^{-wx} dx \\ &= \frac{\Gamma(q+1)}{\Gamma(p+1)\Gamma(q-p)} \int_0^\infty S_u^{\lambda, p}(D)(0) \int_u^\infty (x-u)^{q-p-1} e^{-wx} dx du. \end{aligned}$$

Now we claim that

$$\int_u^\infty (x-u)^{q-p-1} e^{-wx} dx = w^{p-q} e^{-wu} \Gamma(q-p).$$

Indeed, since both sides define holomorphic functions in w , it suffices to check for all $w = \sigma > 0$. In this case we obtain by substitution that

$$\begin{aligned} \int_u^\infty (x-u)^{q-p-1} e^{-wx} dx &= \int_0^\infty \alpha^{q-p-1} e^{-\sigma(\alpha+u)} d\alpha \\ &= e^{-\sigma u} \sigma^{p-q} \int_0^\infty \beta^{q-p-1} e^{-\beta} d\beta = e^{-\sigma u} \sigma^{p-q} \Gamma(q-p). \end{aligned}$$

Altogether

$$w^{q+1} \int_0^\infty S_x^{\lambda, q}(D)(0) e^{-wx} dx = w^{p+1} \frac{\Gamma(q+1)}{\Gamma(p+1)} \int_0^\infty S_u^{\lambda, p}(D)(0) e^{-wu} du. \quad \square$$

4.3. Estimates. We need several estimates for Riesz means which also play a crucial role for the proofs of Theorem 2.9, Theorem 2.16, and Theorem 2.8.

The first one allows to estimate Riesz means of higher order by Riesz means of smaller orders.

Lemma 4.6. *Given $q > p \geq 0$, $s \in \mathbb{C}$ and $D \in \mathcal{D}(\lambda)$, we for all $x > 0$ have*

$$|R_x^{\lambda,q}(D)(s)| \leq \sup_{0 < y < x} |R_y^{\lambda,p}(D)(s)|.$$

Proof. From Lemma 4.4 (with $k = p$ and $\mu = q - p$) we get

$$R_x^{\lambda,q}(D)(s) = x^{-q} \frac{\Gamma(q+1)}{\Gamma(1+p)\Gamma(q-p)} \int_0^x R_y^{\lambda,p}(D)(s) y^p (x-y)^{q-p-1} dy,$$

and then we conclude from (37) that

$$|R_x^{\lambda,q}(D)(s)| \leq \sup_{0 < y < x} |R_x^{\lambda,p}(D)(s)|. \quad \square$$

The second device is an analog of Lemma 2.4 for Riesz means of arbitrary orders.

Lemma 4.7. *Let $D \in \mathcal{D}(\lambda)$ and $k > 0$. Then for $s_0 = \sigma + it \in [\text{Re} > 0]$, $s \in \mathbb{C}$ and $x > 0$*

$$|R_x^{\lambda,k}(D)(s)| \leq C(s_0, k) e^{\sigma x} \sup_{0 < y < x} |R_y^{\lambda,k}(D)(s + s_0)|.$$

Proof. Using translation, we may assume without loss of generality that $s = 0$. Then the choice $s = s_0$ and $w = -s_0$ in Lemma 4.2 leads to

$$\begin{aligned} R_x^{\lambda,k}(D)(0) &= R_x^{\lambda,k}(D)(s_0 - s_0) \\ &= e^{s_0 x} R_x^{\lambda,k}(D)(s_0) + \frac{(-1)^m}{(m+1)!} x^{-k} \int_0^x S_t^{\lambda,m+1}(D)(s_0) \partial^{m+2} h_{-s_0}(t) dt. \end{aligned}$$

where

$$h_{-s_0}(t) = (e^{s_0 t} - e^{s_0 x})(x-t)^k.$$

By Lemma 4.6 we have

$$|R_t^{\lambda,m+1}(s_0)| \leq \sup_{0 < y < t} |R_y^{\lambda,k}(s_0)|,$$

and so

$$\left| x^{-k} \int_0^x S_t^{\lambda,m+1}(s_0) \partial^{m+2} h_{-s_0}(t) dt \right| \leq \sup_{0 < y < x} |R_y^{\lambda,k}(s_0)| x^{-k} \int_0^x t^{m+1} \partial^{m+2} h_{-s_0}(t) dt,$$

where

$$\begin{aligned} (39) \quad \partial^{m+2} h_{-s_0}(t) &= s_0^{m+2} e^{s_0 t} (x-t)^k + \sum_{j=1}^{m+1} s_0^j e^{s_0 t} c_{k,j} (x-t)^{k-(m+2-j)}(t) \\ &\quad + (e^{s_0 t} - e^{s_0 x}) c_{k,m+1} (x-t)^{k-(m+2)}. \end{aligned}$$

We now take into account that

$$(40) \quad |e^{s_0 t} - e^{s_0 x}| \leq |e^{\sigma t} - e^{\sigma x}| \frac{|s_0|}{\sigma} \leq (x-t) \sigma e^{\sigma t} \frac{|s_0|}{\sigma} = (x-t) e^{\sigma t} |s_0|,$$

and hence

$$\begin{aligned} \left| x^{-k} \int_0^x t^{m+1} \partial^{m+2} h_{-s_0}(t) dt \right| &\leq C(s_0, k) e^{\sigma x} x^{-k} \sum_{j=1}^{m+2} \int_0^x t^{m+1} (x-t)^{k-(m+2-j)} dt \\ &\leq C(s_0, k) e^{\sigma x} \sum_{j=1}^{m+2} x^{m+2+j} \int_0^x t^{m+1} \left(1 - \frac{t}{x}\right)^{k-(m+2-j)} dt. \end{aligned}$$

Finally, we calculate

$$\begin{aligned} \int_0^x t^{m+1} \left(1 - \frac{t}{x}\right)^{k-(m+2-j)} dt &= x^{m+2} \int_0^x \left(\frac{t}{x}\right)^{m+1} \left(1 - \frac{t}{x}\right)^{k-(m+2-j)} x^{-1} dt \\ &= x^{m+2} \int_0^1 y^{m+1} (1-y)^{k-(m+2-j)} dy, \end{aligned}$$

and by (37) with $x = 1$

$$\int_0^1 y^{m+1} (1-y)^{k-(m+2-j)} dy = \frac{\Gamma(m+2)\Gamma(k+j-m-1)}{\Gamma(k+j+1)}.$$

This finishes the proof. \square

The third tool we present, is in fact one of the decisive ingredients for the proof of Theorem 2.8. See Section 2.2 for the definition of the summatory function U of the second kind.

Lemma 4.8. *Let $C = \sum a_n$ be a series, λ a frequency, and $k > 0$. Then there is $C = C(k) > 0$ such that for all $x > 0$*

$$|U_x^{\lambda, k}(C)| \leq C e^{kx} \sup_{0 < y < x} |S_y^{\lambda, k}(C)|.$$

The proof of Lemma 4.8 needs another estimate taken from [13, Lemma 8], and (as above) we for the sake of completeness again repeat its argument.

Lemma 4.9. *Let $C = \sum a_n$ be a real series, λ a frequency, $\mu \geq 0$, and $0 < \kappa \leq 1$. Then for all $0 < \xi < x$*

$$\frac{\Gamma(\kappa + \mu + 1)}{\Gamma(\kappa)\Gamma(\mu + 1)} \left| \int_0^\xi S_t^{\lambda, \mu}(C) (x-t)^{\kappa-1} dt \right| \leq \sup_{0 < t < \xi} |S_y^{\lambda, \kappa+\mu}(C)|.$$

Proof. We put

$$c(\mu, \kappa) = \frac{\Gamma(\mu + 1)}{\Gamma(\mu + \kappa + 1)\Gamma(1 - \kappa)},$$

and show with the second formula from Lemma 4.4 and Fubini's theorem that

$$\begin{aligned}
& c(\mu, \kappa)^{-1} \frac{1}{\Gamma(1-\kappa)\Gamma(\kappa)} \int_0^\xi S_t^{\lambda, \mu}(C)(x-t)^{\kappa-1} dt \\
&= c(\mu, \kappa)^{-1} \frac{1}{\Gamma(1-\kappa)\Gamma(\kappa)} \int_0^\xi S_t^{\lambda, (\mu+\kappa)-\kappa}(C)(x-t)^{\kappa-1} dt \\
&= \frac{1}{\Gamma(1-\kappa)\Gamma(\kappa)} \int_0^\xi \int_0^t \frac{d}{du} S_u^{\lambda, \mu+\kappa}(C)(t-u)^{-\kappa} du (x-t)^{\kappa-1} dt \\
&= \int_0^\xi \frac{d}{du} S_u^{\lambda, \mu+\kappa}(C) \frac{1}{\Gamma(1-\kappa)\Gamma(\kappa)} \int_u^\xi (t-u)^{-\kappa} (x-t)^{\kappa-1} dt du \\
&= \int_0^\xi \frac{d}{du} S_u^{\lambda, \mu+\kappa}(C) \left(1 - \frac{1}{\Gamma(1-\kappa)\Gamma(\kappa)} \int_\xi^x (t-u)^{-\kappa} (x-t)^{\kappa-1} dt\right) du.
\end{aligned}$$

Now define on the interval $[0, \xi]$ the function

$$h(u) = \frac{1}{\Gamma(1-\kappa)\Gamma(\kappa)} \int_u^\xi (t-u)^{-\kappa} (x-t)^{\kappa-1} dt.$$

Since

$$\int_u^x (t-u)^{-\kappa} (x-t)^{\kappa-1} dt = \Gamma(1-\kappa)\Gamma(\kappa),$$

we see that

$$h(u) = 1 - \frac{1}{\Gamma(1-\kappa)\Gamma(\kappa)} \int_\xi^x (t-u)^{-\kappa} (x-t)^{\kappa-1} dt.$$

Moreover, on $[0, \xi]$ the function $(t-u)^{-\kappa}$ increases in u , implying that h on $[0, \xi]$ defines a positive decreasing function, always less than 1. By the second mean value theorem there is $0 \leq \eta \leq \xi$ such that

$$\begin{aligned}
& \frac{\Gamma(\kappa + \mu + 1)}{\Gamma(\kappa)\Gamma(\mu + 1)} \int_0^\xi S_t^{\lambda, \mu}(C)(x-t)^{\kappa-1} dt \\
&= h(0) \int_0^\eta \partial_u S_u^{\lambda, \mu+\kappa}(C) du = h(0) S_\eta^{\lambda, \mu+\kappa}(C),
\end{aligned}$$

which completes the argument. \square

We still need one more tool for the proof of Lemma 4.8.

Lemma 4.10. *Let $C = \sum a_n$ be a real series, λ a frequency, and $k > 0$. Then there $c = c(k)$ such that for all $0 < \xi < x$*

$$\left| \int_0^\xi S_t^{\lambda, 0}(C)(x-t)^{k-1} dt \right| \leq c \sup_{0 < t < \xi} |S_t^{\lambda, k}(C)|.$$

Proof. Choose $m \in \mathbb{N}_0$ such that $0 < k - m \leq 1$. By (33) and partial integration we have

$$\begin{aligned} \int_0^\xi S_t^{\lambda,0}(C)(x-t)^{k-1} dt &= c(k) \int_0^\xi \partial_t^m [S_t^{\lambda,m}(C)](x-t)^{k-1} dt \\ &= c(k) \int_0^\xi S_t^{\lambda,m}(C)(x-t)^{(k-1)-m} dt. \end{aligned}$$

On the other hand we deduce from Lemma 4.9 that

$$\left| \int_0^\xi S_t^{\lambda,m}(C)(x-t)^{(k-m)-1} dt \right| \leq c(k) \sup_{0 < t < \xi} |S_t^{\lambda,k}(C)|. \quad \square$$

Finally, we are ready for the proof of Lemma 4.8.

Proof of Lemma 4.8. Without loss of generality we assume that $C = \sum a_n$ is a real series. By (32) we have

$$\begin{aligned} U_x^{\lambda,k}(D) &= k \int_0^x S_t^{\lambda,0}(C)(e^x - e^t)^{k-1} e^t dt \\ &= k \int_0^x S_t^{\lambda,0}(C)(x-t)^{k-1} \left(\frac{e^x - e^t}{x-t} \right)^{k-1} e^t dt. \end{aligned}$$

The function $\left(\frac{e^x - e^\bullet}{x - \bullet} \right)^{k-1} e^\bullet$ is positive and increasing on $[0, x]$ (for $k \geq 1$ this is obvious since then $\frac{e^x - e^\bullet}{x - \bullet}$ increases, and for $0 < k < 1$ differentiate), and the limit as t tends to x equals e^{kx} . Then by the second mean value theorem there is $0 < \xi < x$

$$U_x^{\lambda,k}(C) = k e^{kx} \int_\xi^x S_t^{\lambda,0}(C)(x-t)^{k-1} dt = k e^{kx} \left(\int_0^x - \int_0^\xi \right) S_t^{\lambda,0}(C)(x-t)^{k-1} dt.$$

Hence the conclusion follows from Lemma 4.10. \square

4.4. The technical heart. The following two lemmas finish our preparation of the proofs of Theorem 2.13 and Theorem 2.16. We believe that they in a sense form the 'technical heart' of much of the theory of Riesz summation as created in [13]. In fact, they will be used at various places of this work (see e.g. the proof of one of our main contributions, Theorem 3.7).

Basically, we execute the indicated strategy explained in the previous sections, by carefully analyzing the formula given in Lemma 4.3.

Lemma 4.11. *Let $D \in \mathcal{D}(\lambda)$, $k > 0$ and $m \in \mathbb{N}_0$ such that $m < k \leq m + 1$. Then there is a constant $C = C(m, k)$ such that for all $w = \sigma + i\tau \in [\text{Re} > 0]$, all $s \in \mathbb{C}$ and all $x > 0$*

$$\begin{aligned} &\left| \Gamma(m+2)^{-1} \int_0^x S_t^{\lambda,m+1}(D)(s) w^{m+2} (x-t)^k e^{-wt} dt - S_x^{\lambda,k}(D)(s+w) \right| \\ &\leq e^{-\sigma x} |S_x^{\lambda,k}(D)(s)| + C \sum_{j=1}^{m+1} |w|^j \int_0^x |S_y^{\lambda,k}(D)(s)| e^{-\sigma y} (x-y)^{j-1} dy. \end{aligned}$$

In particular, there is a constant $L = L(m, k)$ such that for all $w = \sigma + it \in [\operatorname{Re} > 0]$ with $|\arg(w)| \leq \gamma$ and all $x > 0$

$$\begin{aligned} & \left| \Gamma(m+2)^{-1} \int_0^x S_t^{\lambda, m+1}(D)(0) w^{m+2} (x-t)^k e^{-wt} dt - S_x^{\lambda, k}(D)(w) \right| \\ & \leq e^{-\sigma x} |S_x^{\lambda, k}(D)(0)| + L \sum_{j=1}^{m+1} |\sec(\gamma)|^j \int_0^x |S_y^{\lambda, k}(D)(0)| y^{-j} (x-y)^{j-1} dy. \end{aligned}$$

Proof. We first handel the more complicated case $k < m + 1$, and at the end of this proof we comment on the easier case $k = m + 1$.

If $m < k < m + 1$, then we know from Lemma 4.3 that for all $s, w \in \mathbb{C}$ and all $x > 0$

$$\begin{aligned} S_x^{\lambda, k}(D)(s+w) - \Gamma(m+2)^{-1} \int_0^x S_t^{\lambda, m+1}(D)(s) w^{m+2} e^{-wt} (x-t)^k dt \\ = -e^{-wt} S_x^{\lambda, k}(D)(s) \\ + C(k) \int_0^x S_y^{\lambda, k}(D)(s) \int_y^x (t-y)^{m-k} (g_1(t) + g_2(t)) dt dy, \end{aligned}$$

where

$$\begin{aligned} g_1(t) &= \sum_{j=1}^{m+1} \binom{m+2}{j} (-w)^j e^{-wt} [(x-\bullet)^k]^{(m+2-j)}(t) \\ g_2(t) &= (e^{-wt} - e^{-wx}) [(x-\bullet)^k]^{(m+2)}(t). \end{aligned}$$

We calculate (as in (39))

$$(41) \quad g_1(t) + g_2(t) = \sum_{j=1}^{m+1} (-w)^j e^{-wt} c_{k,j} (x-t)^{k-(m+2-j)}$$

$$(42) \quad + (e^{-wt} - e^{-wx}) c_{k, m+1} (x-t)^{k-(m+2)}.$$

Then it remains to control the integral

$$C(k) \int_0^x S_y^{\lambda, k}(D)(s) \int_y^x (t-y)^{m-k} (g_1(t) + g_2(t)) dt dy.$$

For $i = 1$ we use (37) (with $x = 1$) and (38) to obtain with $w = \sigma + i\tau \in [\text{Re} > 0]$

$$\begin{aligned}
& \left| \int_y^x (t-y)^{m-k} g_2(t) dt \right| \\
&= \left| \sum_{j=1}^{m+1} (-1)^j w^j c_{k,j} (x-y)^{j-1} e^{-wy} \int_0^1 e^{-w\beta(x-y)} \beta^{m-k} (1-\beta)^{k-(m+2-j)} d\beta \right| \\
&\leq C_1(m, k) \sum_{j=1}^{m+1} |w|^j (x-y)^{j-1} e^{-\sigma y} \int_0^1 \beta^{m-k} (1-\beta)^{k-(m+2-j)} d\beta \\
&= C_1(m, k) \sum_{j=1}^{m+1} \frac{\Gamma(1+m-k)\Gamma(k-m-1+j)}{\Gamma(j)} |w|^j (x-y)^{j-1} e^{-\sigma y}.
\end{aligned}$$

For $i = 2$ we claim that

$$\left| \int_y^x (t-y)^{m-k} g_3(t) dt \right| \leq C(m, k) |w| e^{-\sigma y}.$$

Indeed, by the mean value theorem (as in (40)) for $0 \leq t \leq x$ and $w \in [\text{Re} > 0]$

$$(43) \quad |e^{-wt} - e^{-wx}| \leq |e^{-\sigma t} - e^{-\sigma x}| \frac{|w|}{\sigma} \leq (x-t) \sigma e^{-\sigma t} \frac{|w|}{\sigma} = (x-t) e^{-\sigma t} |w|.$$

Then (again using (37) with $x = 1$)

$$\begin{aligned}
& \left| \int_y^x (t-y)^{m-k} g_3(t) dt \right| \\
&= \left| c_{k,m+1} \int_y^x (t-y)^{m-k} (e^{-wt} - e^{-wx}) (x-t)^{k-(m+2)} dt \right| \\
&\leq c_{k,m+1} \int_y^x (t-y)^{m-k} |e^{-\sigma t} - e^{-\sigma x}| \frac{|w|}{\sigma} (x-t)^{k-(m+2)} dt \\
&\leq e^{-\sigma y} |w| c_{k,m+1} \int_y^x (t-y)^{m-k} (x-t)^{k-m-1} dt \\
&= e^{-\sigma y} |w| c_{k,m+1} (x-y)^{m-k+(k-m-1)+1} \int_0^1 \beta^{m-k} (1-\beta)^{k-m-1} d\beta \\
&= C_2(m, k) \frac{\Gamma(1+m-k)\Gamma(k-m)}{\Gamma(1)} e^{-\sigma y} |w|.
\end{aligned}$$

Putting everything together completes the argument of the first part. For the second, take $s = 0$, and observe that $e^{\sigma y} \leq \frac{(\sigma y)^j}{j!}$ and $\frac{|w|}{\sigma} \leq \sec(\gamma)$. Finally, we as announced comment on the simpler case $m < k = m + 1$. In this case we know from Lemma 4.2 that

$$\begin{aligned}
S_x^{\lambda,r}(D)(s+w) &- \frac{1}{\Gamma(m+2)} \int_0^x S_t^{\lambda,m+1}(D)(s) w^{m+2} e^{-wt} r(x,t) dt \\
&= -e^{-wt} S_x^{\lambda,r}(D)(s) + \int_0^x S_t^{\lambda,m+1}(D)(s) (g_1(t) + g_2(t)) dt.
\end{aligned}$$

As above, the claim is an immediate consequence of (41) and (43). \square

Roughly speaking, the integral representation (16) (from Theorem 2.9) for (λ, k) -Riesz limits is determined by the first summand of the left-hand side of the first inequality of Lemma 4.11, tending $x \rightarrow \infty$. The next lemma proves existence of this limit – given a growth condition on the summatory function.

Lemma 4.12. *Given $D \in \mathcal{D}(\lambda)$, let $s \in \mathbb{C}$ and $k \geq 0$, $q > 0$. Then for all $w \in \mathbb{C}$ and $\varepsilon > 0$ such that $0 < \varepsilon < \operatorname{Re} w$ and*

$$|S_t^{\lambda, q}(D)(s)| \leq C(q, \varepsilon)t^q e^{\varepsilon t}, \quad t > 0,$$

we have

$$(44) \quad \lim_{x \rightarrow \infty} \int_0^x S_t^{\lambda, q}(D)(s) \left(1 - \frac{t}{x}\right)^k w^{q+1} e^{-wt} dt = \int_0^\infty S_y^{\lambda, q}(D)(s) w^{q+1} e^{-wy} dy.$$

Moreover, the convergence in (56) is uniform on each cone $|\arg(w - s)| \leq \gamma < \frac{\pi}{2}$, whenever

$$|S_t^{\lambda, q}(D)(s)| \leq C(q)t^q, \quad t > 0.$$

Proof. We assume without loss of generality that $s = 0$. In order to prove (56), we fix w and ε , and put $\sigma = \operatorname{Re} w - \varepsilon$. Then for all x, t

$$\left| S_t^{\lambda, q}(D)(0) \left(1 - \frac{t}{x}\right)^k w^{q+1} e^{-wt} \chi_{[0, x]}(t) \right| \leq C(q, \varepsilon) |w|^{q+1} e^{-\sigma t} t^q,$$

and hence (56) is an immediate consequence of the dominated convergence theorem.

The proof of the 'moreover-part' is similar: For each w with $|\arg(w)| \leq \gamma < \frac{\pi}{2}$ we abbreviate $\sigma_w = \operatorname{Re} w$. Then for all x, t, w

$$\left| S_t^{\lambda, q}(D)(0) \left(1 - \frac{t}{x}\right)^k w^{q+1} e^{-wt} \chi_{[0, x]}(t) \right| \leq C(q) \sec(\gamma) t^q \sigma_w^{q+1} e^{-\sigma_w t},$$

and hence by the substitution $u = \sigma_w t$ we get for all x, u, w

$$\left| S_{\frac{u}{\sigma_w}}^{\lambda, q}(D)(0) \left(1 - \frac{\frac{u}{\sigma_w}}{x}\right)^k w^{q+1} e^{-w \frac{u}{\sigma_w}} \chi_{[0, x]} \left(\frac{u}{\sigma_w}\right) \right| \frac{1}{\sigma_w} \leq C(q) \sec(\gamma) u^q e^{-u}.$$

Hence again by the dominated convergence theorem (uniform variant)

$$\int_0^{x\sigma_w} S_{\frac{u}{\sigma_w}}^{\lambda, q}(D)(0) \left(1 - \frac{\frac{u}{\sigma_w}}{x}\right)^k w^{q+1} e^{-w \frac{u}{\sigma_w}} \frac{1}{\sigma_w} du \rightarrow \int_0^\infty S_{\frac{u}{\sigma_w}}^{\lambda, q}(D)(0) w^{q+1} e^{-w \frac{u}{\sigma_w}} \frac{1}{\sigma_w} du$$

uniformly in w as $x \rightarrow \infty$. Substituting back, finishes the proof. \square

4.5. Proof of Theorem 2.9.

Proof of Theorem 2.9. Let $k > 0$ and $m \in \mathbb{N}_0$ such that $m < k \leq m + 1$. If D is (λ, k) -summable at $s_0 = \sigma_0 + i\tau_0$, then for some constant $C = C(s_0, k)$ we by Lemma 4.7 have that for all $x > 0$

$$(45) \quad |R_x^{\lambda, k}(D)(0)| \leq C e^{\sigma_0 x}.$$

We fix $s = \sigma + i\tau \in [\operatorname{Re} > \sigma_0]$, and deduce from Lemma 4.14 and Lemma 4.5 that

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{x^{-k}}{\Gamma(m+2)} \int_0^x S_t^{\lambda, m+1}(D)(0) s^{m+2} (x-t)^k e^{-st} dt \\ = \Gamma(1+k)^{-1} s^{k+1} \int_0^\infty S_y^{\lambda, k}(D)(0) e^{-sy} dy. \end{aligned}$$

According to Lemma 4.11 we have to show that the term

$$e^{-\sigma x} |R_x^{\lambda, k}(D)(0)| + C \sum_{j=1}^{m+1} x^{-k} |s|^j \int_0^x |S_y^{\lambda, k}(D)(0)| e^{-\sigma y} (x-y)^{j-1} dy$$

vanishes as $x \rightarrow \infty$. Observe that (45) implies the claim for the first summand. Writing $\varepsilon = \sigma - \sigma_0 > 0$ we use again (45) to see that for $j = 1, \dots, m+1$

$$\begin{aligned} x^{-k} \int_0^x |S_y^{\lambda, k}(D)(0)| e^{-\sigma y} (x-y)^{j-1} dy &\leq C x^{-k} \int_0^x e^{-\varepsilon y} y^k (x-y)^{j-1} dy \\ &\leq C x^{m-k} \int_0^\infty e^{-\varepsilon y} y^k dy, \end{aligned}$$

which vanishes as $x \rightarrow \infty$, since $k > m$.

It remains to prove the 'moreover-part'. We may assume without loss of generality that $\lambda_1 = 0$, $s_0 = 0$ and $\lim_{x \rightarrow \infty} R_x^{\lambda, k}(D)(0) = 0$; if this particular case is settled and s_0 is arbitrary, then we consider the modified Dirichlet series $\sum b_n e^{-s_0 \lambda_n} e^{-s \lambda_n}$ with $b_1 = a_1 - \lim_{x \rightarrow \infty} R_x^{\lambda, k}(D)(s_0)$ and $b_n = a_n$, $n \geq 2$, which then converges uniformly on each cone $|\arg(s)| \leq \gamma < \frac{\pi}{2}$, and so implies the conclusion.

So let $s_0 = 0$. According to Lemma 4.11 and Lemma 4.14, we have to check that uniformly for all $s \in [\operatorname{Re} > 0]$ with $\arg(s) \leq \gamma < \frac{\pi}{2}$

$$(46) \quad \lim_{x \rightarrow \infty} \sum_{j=1}^{m+1} |\sec(\gamma)|^j x^{-k} \int_0^x |S_y^{\lambda, k}(D)(0)| y^{-j} (x-y)^{j-1} dy = 0.$$

Therefore, choose $\varepsilon > 0$ and $x_0 > 1$ such that $|R_x^{\lambda, k}(D)(0)| \leq \varepsilon$ for all $x > x_0$. We split the integral

$$\begin{aligned} x^{-k} \int_0^x |S_y^{\lambda, k}(D)(0)| y^{-j} (x-y)^{j-1} dy \\ = x^{-k} \int_0^{x_0} |S_y^{\lambda, k}(D)(0)| y^{-j} (x-y)^{j-1} dy + x^{-k} \int_{x_0}^x |S_y^{\lambda, k}(D)(0)| y^{-j} (x-y)^{j-1} dy. \end{aligned}$$

Then note that for $1 \leq j \leq m+1$ and $x \geq 2x_0$

$$\begin{aligned} x^{-k} \int_0^{x_0} |S_y^{\lambda,k}(D)(0)| y^{-j} (x-y)^{j-1} dy &\leq C(D, x_0) x^{-k} \int_0^{x_0} y^{k-j} (x-y)^{j-1} dy \\ &\leq C(D, x_0) x^{m-k} \int_0^{x_0} y^{k-j} dy \leq C(D, x_0) x^{m-k} x_0^{k+1-j}, \end{aligned}$$

which vanishes as $x \rightarrow \infty$, since $k > m$, and moreover for $x \geq 2x_0$

$$\begin{aligned} x^{-k} \int_{x_0}^x |S_y^{\lambda,k}(D)(0)| y^{-j} (x-y)^{j-1} dy &\leq x^{-k} \varepsilon \int_{x_0}^x y^{k-j} (x-y)^{j-1} dy \\ &\leq \varepsilon x^{-k} x^{k-j} \int_{x_0}^x (x-y)^{j-1} dy = \varepsilon x^{-j} j^{-1} (x-x_0)^j \leq \varepsilon \left(1 - \frac{x_0}{x}\right)^j \leq \varepsilon. \end{aligned}$$

This finishes the proof of (46). □

4.6. Proof of Theorem 2.16.

Proof of Theorem 2.16. Assume that

$$\sigma_0 > \limsup_{x \rightarrow \infty} x^{-1} \log(|R_x^{\lambda,k}(D)(0)|) =: L.$$

Then (45) is satisfied, and from the previous proof of Theorem 2.9 (using again Lemma 4.14, Lemma 4.5 and Lemma 4.11) we see that D is (λ, k) -summable on $[\operatorname{Re} > \sigma_0]$. This proves that $L \leq \sigma_c^{\lambda,k}(D)$.

Assume conversely that $\sigma_c^{\lambda,k}(D) \geq 0$, and let $\varepsilon > 0$. We define $\sigma_0 = \sigma_c^{\lambda,k}(D) + \varepsilon$. Then D is (λ, k) -summable at σ_0 and so by Lemma 4.7 we have

$$|R_x^{\lambda,k}(D)(0)| \leq C(\sigma_0) e^{\sigma_0 x}, \quad x > 0.$$

Consequently, $L \leq \sigma_0 = \sigma_c^{\lambda,k}(D) + \varepsilon$, and hence as desired $L = \sigma_c^{\lambda,k}(D)$. □

4.7. Proof of Theorem 2.7.

Proof of Theorem 2.7,(i). Without loss of generality we may assume that the (λ, k) -Riesz sum of D is 0, i.e. $S_x^{\lambda,k}(D) = o(x^k)$. Since by Lemma 4.4 for all x

$$S_x^{\lambda,\ell}(D) = \frac{\Gamma(\ell+1)}{\Gamma(k+1)\Gamma(\ell-k)} \int_0^x S_u^{\lambda,k}(D) (x-u)^{\ell-k-1} du,$$

we check that

$$\int_0^x S_u^{\lambda,k}(D) (x-u)^{\ell-k-1} du = o(x^\ell).$$

Fix some $\varepsilon > 0$, and choose x_0 such that for all $x > x_0$ we have $S_x^{\lambda,k}(D) < \varepsilon x^k$. Then, using (37), for all $x > x_0$

$$\begin{aligned} & \int_0^x S_u^{\lambda,k}(D)(x-u)^{\ell-k-1} du \\ &= \int_0^{x_0} S_u^{\lambda,k}(D)(x-u)^{\ell-k-1} du + \int_{x_0}^x S_u^{\lambda,k}(D)(x-u)^{\ell-k-1} du \\ &\leq \sup_{u < x_0} S_u^{\lambda,k}(D) \int_0^{x_0} (x-u)^{\ell-k-1} du + \varepsilon \int_{x_0}^x u^k (x-u)^{\ell-k-1} du \\ &\leq \sup_{u < x_0} S_u^{\lambda,k}(D) \int_0^{x_0} (x-u)^{\ell-k-1} du + \varepsilon \int_0^x u^k (x-u)^{\ell-k-1} du \\ &\leq \sup_{u < x_0} S_u^{\lambda,k}(D) x_0^{\ell-k-1} + \varepsilon x^\ell \frac{\Gamma(\ell+1)\Gamma(\ell-k)}{\Gamma(k+1)}, \end{aligned}$$

the conclusion. \square

Proof of Theorem 2.7,(ii). Without loss of generality we may assume that the (e^λ, k) -Riesz sum of D is zero. Hence we know by assumption from Lemma 4.1, (32) that

$$(47) \quad U_x^{\lambda,k}(D) = k \int_0^x S_t^{\lambda,0}(D) (e^x - e^t)^{k-1} e^t dt = o(e^{kx}),$$

and the job is to show that

$$(48) \quad S_y^{\lambda,k}(D) = k \int_0^y S_s^{\lambda,0}(D) (y-s)^{k-1} ds = o(y^k).$$

Substitution with $s = \log t$ and $y = \log x$ gives that (48) in fact is equivalent to

$$(49) \quad S_{\log x}^{\lambda,k}(D) = k \int_1^x S_t^{e^\lambda,0}(D) \frac{(\log x - \log t)^{k-1}}{t} dt = o(\log^k x).$$

Indeed, the proof in [13, p. 32] proves this for the two cases $k \in \mathbb{N}$ and $0 < k < 1$ separately, and sketches an argument for the case $k > 1, k \notin \mathbb{N}$. We, in a first step, for the case $0 < k < 1$ follow the proof from [13], and modify it in a second step to verify the general case $k \geq 1$.

The case $0 < k < 1$: By Theorem 2.7,(i) we know that $S_x^{e^\lambda,1}(D) = o(x)$, hence, given $\varepsilon > 0$, there is $x_0 > 1$ such that for all $x > x_0$

$$(50) \quad S_x^{e^\lambda,1}(D) \leq \varepsilon x.$$

For $x > 3x_0$ we split the integral from (49) into three pieces,

$$(51) \quad \int_1^x S_t^{e^\lambda,0}(D) \frac{(\log x - \log t)^{k-1}}{t} dt = \int_1^{x_0} + \int_{x_0}^{x/3} + \int_{x/3}^x = J_1 + J_2 + J_3,$$

and estimate each integral separately.

Integral J_1 : Clearly,

$$J_1 \leq \left[\sup_{t \leq x_0} S_t^{e^\lambda,0}(D) \right] x_0 \log^{k-1} \left(\frac{x}{x_0} \right) = o(\log^k(x)),$$

where we use that here $\log^{k-1}\left(\frac{x}{t}\right) \leq \log^{k-1}\left(\frac{x}{x_0}\right)$, since $t \leq x_0 \leq x$ and $k-1 < 0$.

Integral J_2 : Using (33) and integrating by parts we have

$$\begin{aligned} J_2 &= \log^{k-1}(3) \frac{3}{x} S_t^{e^\lambda, 1}(D) - \log^{k-1}\left(\frac{x}{x_0}\right) \frac{1}{x_0} S_{x_0}^{e^\lambda, 1}(D) \\ &\quad + \int_{x_0}^{x/3} S_t^{e^\lambda, 1}(D) \left[(k-1) \log^{k-2}\left(\frac{x}{t}\right) + \log^{k-1}\left(\frac{x}{t}\right) \right] \frac{dt}{t^2}. \end{aligned}$$

In absolute value the first two terms are less than an absolute constant times ε (we use (50), and the fact that $\log(x/x_0) > \log 3 > 1$ since $x_0 < x/3$). The integral in the preceding equality we estimate from above by

$$\varepsilon k \int_{x_0}^{x/3} \log^{k-1}\left(\frac{x}{t}\right) \frac{dt}{t} = \varepsilon k \left(\frac{1}{k} \log^k(3) - \frac{1}{k} \log^k\left(\frac{x}{x_0}\right) \right),$$

using again (50) together with the fact that $\log\left(\frac{x}{t}\right) > \log\left(\frac{3t}{t}\right) = \log(3) > 1$ for $x_0 < t < x/3$, implying $\log^{k-2}\left(\frac{x}{t}\right) < \log^{k-1}\left(\frac{x}{t}\right)$. All together we have proved that $J_2 = o(\log^k(x))$.

Integral J_3 : By the second mean value theorem there is $x/3 \leq \xi \leq x$ such that

$$\begin{aligned} J_3 &= \int_{x/3}^x S_t^{e^\lambda, 0}(D) \log^{k-1}\left(\frac{x}{t}\right) \frac{1}{t} dt \\ &= \frac{3}{x} \int_{x/3}^{\xi} S_t^{e^\lambda, 0}(D) \log^{k-1}\left(\frac{x}{t}\right) dt \\ &= \frac{3}{x} \int_{x/3}^{\xi} S_t^{e^\lambda, 0}(D) (x-t)^{k-1} \left(\frac{\log x - \log t}{x-t} \right)^{k-1} dt. \end{aligned}$$

But the function $\left(\frac{\log x - \log \bullet}{x - \bullet} \right)^{k-1}$ is increasing on $[1, x]$ with limit x^{1-k} as t tends to x (the quotient is decreasing and $k-1 < 0$). Hence another application of the second mean value theorem shows that there is $x/3 \leq \xi_1 \leq \xi \leq x$ such that

$$\begin{aligned} |J_3| &= \frac{3}{x} \left(\frac{\log x - \log \xi_1}{x - \xi_1} \right)^{k-1} \left| \int_{x/3}^{\xi} S_t^{e^\lambda, 0}(D) (x-t)^{k-1} dt \right| \\ &\leq 3x^{-k} \left| \int_{\xi_1}^{\xi} S_t^{e^\lambda, 0}(D) (x-t)^{k-1} dt \right|. \end{aligned}$$

Now by Lemma 4.10 (write $\int_{\xi_1}^{\xi} = \int_0^{\xi} - \int_0^{\xi_1}$) there is some constant $c = c(k) > 0$ for which

$$\left| \int_{\xi_1}^{\xi} S_t^{e^\lambda, 0}(D) (x-t)^{k-1} dt \right| \leq c \sup_{x/3 < t < x} |S_t^{e^\lambda, k}(D)|.$$

Since by assumption $S_x^{e^\lambda, k}(D) = o(x^k)$, we see that $J_3 = o(1) = o(\log^k(x))$. Collecting all estimates we got for the three integrals J_1 , J_2 , and J_3 , we in fact

proved (what we were aiming at in (49))

$$\int_1^x S_t^{e^\lambda, 0}(D) \frac{(\log x - \log t)^{k-1}}{t} dt = o(\log^k x).$$

It remains to consider the case $k \geq 1$, which is going to follow from a refinement of the preceding arguments: Choose some $m \in \mathbb{N}$ such that $m < k \leq m + 1$, and note again that by Theorem 2.7,(i) we have $S_x^{e^\lambda, m+1}(D) = o(x)$. Hence, given $\varepsilon > 0$, there is $x_0 > 1$ such that for all $x > x_0$

$$(52) \quad S_x^{e^\lambda, m+1}(D) \leq \varepsilon x.$$

For $x > 3x_0$ we again consider the three integrals J_1, J_2 , and J_3 from (51), and handel each piece separately in order to show (49).

Integral J_1 : Clearly,

$$J_2 \leq \left[\sup_{t \leq x_0} S_t^{e^\lambda, 0}(D) \right] x_0 \log^{k-1}(x) = o(\log^k(x)),$$

since now $k - 1 \geq 0$.

Integral J_2 : We have that for all $1 \leq \ell \leq m + 1$

$$\frac{d^\ell (\log x - \log t)^{k-1}}{dt} = \frac{1}{t^{\ell+1}} \sum_{j=0}^{\ell} c_j(k) (\log x - \log t)^{k-1-j}.$$

Integrating $m + 1$ times by parts (again using (33)) we have

$$\begin{aligned} J_2 &= S_t^{e^\lambda, m+1}(D) \frac{1}{t^{m+1}} \sum_{j=0}^m c_j(k) \log^{k-1-j} \left(\frac{x}{t} \right) \Big|_{x_0}^{x/3} \\ &\quad + \int_{x_0}^{x/3} S_t^{e^\lambda, m+1}(D) \frac{1}{t^{m+2}} \sum_{j=0}^{m+1} c_j(k) \log^{k-1-j} \left(\frac{x}{t} \right) dt. \end{aligned}$$

As above we have that $\log \left(\frac{x}{t} \right) > \log(3) > 1$ for $x_0 \leq t \leq x/3$, and hence

$$\begin{aligned} \log^{k-1} \left(\frac{x}{t} \right) &\geq \dots \geq \log^{(k-1)-(m-1)} \left(\frac{x}{t} \right) \\ &\geq \log^{(k-1)-m} \left(\frac{x}{t} \right) \geq \log^{(k-1)-(m+1)} \left(\frac{x}{t} \right). \end{aligned}$$

Together with (52) this shows that the first term in the above formula for J_2 is less than a constant (only depending on k) times $\varepsilon \log^{k-1}(x)$, and the second term is less than a constant (only depending on k) times

$$\varepsilon \int_{x_0}^{x/3} \log^{k-1} \left(\frac{x}{t} \right) \frac{dt}{t} \leq \log^{k-1}(x) \int_{x_0}^{x/3} \frac{1}{t} dt = o(\log^k(x))$$

(again taking into account that $k - 1 \geq 0$). All together we get $J_2 = o(\log^k(x))$.

Integral J_3 : We write

$$J_3 = \int_{x/3}^x S_t^{e^\lambda, 0}(D)(x-t)^{k-1} \left(\frac{\log x - \log t}{x-t} \right)^{k-1} \frac{1}{t} dt.$$

Since $k \geq 1$, the function $\left(\frac{\log x - \log \bullet}{x - \bullet} \right)^{k-1} \frac{1}{\bullet}$ is decreasing on $[1, x]$ (look at the graph of $\log \bullet$), and

$$\left(\frac{3 \log 3}{2} \right)^{k-1} x^{1-k} \frac{3}{x} = \lim_{t \rightarrow x/3} \left(\frac{\log x - \log t}{x-t} \right)^{k-1} \frac{1}{t}.$$

Then the second mean value theorem yields some $x/3 < \xi < x$ such that

$$J_3 = 3 \left(\frac{3 \log 3}{2} \right)^{k-1} x^{-k} \int_{x/3}^{\xi} S_t^{e^\lambda, 0}(D)(x-t)^{k-1} dt = \dots \int_0^{\xi} - \int_0^{\frac{x}{3}} \dots$$

Now by Lemma 4.10 there is some constant $c = c(k) > 0$ such that

$$\left| \int_0^{\xi} S_t^{e^\lambda, 0}(D)(x-t)^{k-1} dt \right| \leq c \sup_{0 < t < x} |S_t^{e^\lambda, k}(D)|,$$

and hence by (52) we finally get $J_3 = o(1) = o(\log^k(x))$. As in the first case this proves the claim. \square

4.8. Proof of Theorem 2.8. Let us finally turn to the proof of Theorem 2.8. Our task is to prove that, given a λ -Dirichlet series $D = \sum a_n e^{-\lambda_n s} \in \mathcal{D}(\lambda)$ and $k > 0$,

$$\sigma_c^{\lambda, k}(D) \leq \sigma_c^{e^\lambda, k}(D).$$

It suffices to check that D is (e^λ, k) -summable on $[\operatorname{Re} > \operatorname{Re} s_0]$, provided D is (λ, k) -summable at some $s_0 \in [\operatorname{Re} \geq 0]$. As a by-product our arguments again show that then the limit function $f : [\operatorname{Re} > 0] \rightarrow \mathbb{C}$ is given by

$$(53) \quad \Gamma(1+k) \frac{f(s)}{s^{1+k}} = \int_0^\infty e^{-st} S_t^{\lambda, k}(D)(0) dt,$$

as it should be according to Theorem 2.7 and Theorem 2.9. In fact the proof we give is a modification of the proof given for Theorem 2.9, and so we start modifying Lemma 4.11.

Lemma 4.13. *Let $D \in \mathcal{D}(\lambda)$, $k > 0$, and $m \in \mathbb{N}_0$ with $m < k \leq m+1$. Then there is a constant $C = C(k)$ such that for all $w = \sigma + i\tau \in [\operatorname{Re} > 0]$, all $s \in \mathbb{C}$,*

all $0 \leq \varepsilon < \frac{1}{m+2} \min\{k, \sigma\}$, and all $x > 0$

$$\begin{aligned} & \left| \Gamma(m+2)^{-1} \int_0^x S_t^{\lambda, m+1}(D)(s) w^{m+2} (e^x - e^t)^k e^{-wt} dt - U_x^{\lambda, k}(D)(s+w) \right|, \\ & \leq e^{-\sigma x} |U_x^{\lambda, k}(D)(s)| \\ & \quad + C(k) \left[\sum_{j=1}^{m+1} |w^j| \sum_{\ell=1}^{m+2-j} e^{(k-\varepsilon\ell)x} x^{m-\ell+1} \int_0^x |S_y^{\lambda, k}(D)(s)| e^{-(\sigma-\varepsilon\ell)y} dy \right. \\ & \quad \left. + |w| \sum_{\ell=1}^{m+2} e^{(k-\varepsilon\ell)x} x^{m+2-\ell} \int_0^x |S_y^{\lambda, k}(D)(s)| e^{-(\sigma-\varepsilon\ell)y} dy \right] \end{aligned}$$

Proof. Again we first deal with the case $k < m+1$, and at the end we consider the more simple case $k = m+1$. By Lemma 4.3 for all $s, w \in \mathbb{C}$ and all $x > 0$

$$\begin{aligned} (54) \quad & U_x^{\lambda, k}(D)(s+w) - \frac{1}{\Gamma(m+2)} \int_0^x S_t^{\lambda, m+1}(D)(s) w^{m+2} e^{-wt} (e^x - e^t)^k dt \\ & = -e^{-wt} U_x^{\lambda, k}(D)(s) \\ & \quad + C(k) \int_0^x S_y^{\lambda, k}(D)(s) \int_y^x (t-y)^{m-k} (g_1(t) + g_2(t)) dt dy, \end{aligned}$$

where

$$\begin{aligned} (55) \quad & g_1(t) = \sum_{j=1}^{m+1} \binom{m+2}{j} (-w)^j e^{-wt} \partial_t^{(m+2-j)} (e^x - e^t)^k \\ & g_2(t) = (e^{-wt} - e^{-wx}) \partial_t^{(m+2)} (e^x - e^t)^k. \end{aligned}$$

A straight forward calculation shows that for every $N \in \mathbb{N}$ we have

$$\partial_t^N (e^x - e^t)^k = \sum_{\ell=1}^N c_\ell(k) (e^x - e^t)^{k-\ell} e^{\ell t},$$

with constants $c_\ell(k) > 0$. Then by the mean value theorem and the equations (37), (38) we have

$$\begin{aligned}
& \left| \int_y^x (t-y)^{m-k} g_1(t) dt \right| \\
&= \left| \int_y^x (t-y)^{m-k} \sum_{j=1}^{m+1} \binom{m+2}{j} (-w)^j e^{-wt} \sum_{\ell=1}^{m+2-j} c_\ell(k) (e^x - e^t)^{k-\ell} e^{\ell t} dt \right| \\
&\leq \sum_{j=1}^{m+1} \binom{m+2}{j} |w|^j \sum_{\ell=1}^{m+2-j} c_\ell(k) \int_y^x (t-y)^{m-k} (x-t)^{k-\ell} e^{(k-\ell)x} e^{-\sigma t} e^{\ell t} dt \\
&= \sum_{j=1}^{m+1} \binom{m+2}{j} |w|^j \sum_{\ell=1}^{m+2-j} c_\ell(k) \int_y^x (t-y)^{m-k} (x-t)^{k-\ell} e^{(k-\ell)x} e^{-\sigma t} e^{(1-\varepsilon)\ell t} e^{\varepsilon \ell t} dt \\
&\leq \sum_{j=1}^{m+1} \binom{m+2}{j} |w|^j \sum_{\ell=1}^{m+2-j} c_\ell(k) e^{(k-\varepsilon\ell)x} e^{-(\sigma-\varepsilon\ell)y} (x-y)^{m-\ell+1} c(k),
\end{aligned}$$

where

$$c(k) = \int_0^1 \beta^{m-k} (1-\beta)^{m-\ell+1} d\beta = \frac{\Gamma(m-k+1)\Gamma(m-\ell+2)}{\Gamma(2(m+1)-k-\ell)}.$$

And similarly, using (43), we have

$$\begin{aligned}
& \left| \int_y^x (t-y)^{m-k} g_2(t) dt \right| \\
&\leq \int_y^x (t-y)^{m-k} (x-t) e^{-\sigma t} |w| \sum_{\ell=1}^{m+2} (x-t)^{k-\ell} e^{(k-\ell)x} e^{\ell t} dt \\
&= |w| \sum_{\ell=1}^{m+2} \int_y^x (t-y)^{m-k} (x-t)^{k-\ell+1} e^{(k-\ell)x} e^{-\sigma t} e^{\ell t} dt \\
&\leq |w| \sum_{\ell=1}^{m+2} e^{(k-\varepsilon\ell)x} \int_y^x (t-y)^{m-k} (x-t)^{k-\ell+1} e^{-(\sigma-\varepsilon\ell)t} dt \\
&\leq |w| \sum_{\ell=1}^{m+2} e^{(k-\varepsilon\ell)x} (x-y)^{m+2-\ell} e^{-(\sigma-\varepsilon\ell)y} c(k).
\end{aligned}$$

Implementing the preceding two estimates into (54), gives exactly what we were aiming at.

Let us finally look at the case $k = m + 1$. This case follows similarly – but it is more simple. We start replacing (54) by (34), implement the derivatives of g_1 and g_2 from (55), and finish as above (using again the mean value theorem, the estimate (43), and the splitting $e^{\ell t} = e^{(1-\varepsilon)\ell t} e^{\varepsilon \ell t}$). \square

Next we need a simple modification of Lemma 4.14.

Lemma 4.14. *Let $D \in \mathcal{D}(\lambda)$, $k, q > 0$, and $m \in \mathbb{N}_0$ with $m < k \leq m + 1$. Then for all $w \in \mathbb{C}$ and $\varepsilon > 0$ such that $0 < \varepsilon < \operatorname{Re} w$ and*

$$|S_t^{\lambda, q}(D)(s)| \leq C(q, \varepsilon) t^q e^{\varepsilon t}, \quad t > 0,$$

we have

$$(56) \quad \lim_{x \rightarrow \infty} \int_0^x S_t^{\lambda, q}(D)(s) \left(1 - \frac{e^t}{e^x}\right)^k w^{q+1} e^{-wt} dt = \int_0^\infty S_y^{\lambda, q}(D)(s) w^{q+1} e^{-wy} dy.$$

Proof. We assume without loss of generality that $s = 0$. In order to prove (56), we fix w and ε , and put $\sigma = \operatorname{Re} w - \varepsilon$. Then for all x, t

$$\left| S_t^{\lambda, q}(D)(0) \left(1 - \frac{e^t}{e^x}\right)^k w^{q+1} e^{-wt} \chi_{[0, x]}(t) \right| \leq C(q, \varepsilon) |w|^{q+1} e^{-\sigma t} t^q,$$

and hence (56) is an immediate consequence of the dominated convergence theorem. \square

The final argument for Theorem 2.8 is now very similar to that of Theorem 2.9.

Proof of Theorem 2.8. Let $k > 0$ and $m \in \mathbb{N}_0$ such that $m < k \leq m + 1$. If D is (λ, k) -summable at $s_0 = \sigma_0 + i\tau_0$, then for some constant $C = C(s_0, k)$ we by Lemma 4.7 have that for all $x > 0$

$$(57) \quad |R_x^{\lambda, k}(D)(0)| \leq C e^{\sigma_0 x}.$$

We fix $s = \sigma + i\tau \in [\operatorname{Re} > \sigma_0]$, and deduce from Lemma 4.14 and Lemma 4.5 that

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{e^{-kx}}{\Gamma(m+2)} \int_0^x S_t^{\lambda, m+1}(D)(0) s^{m+2} (e^x - e^t)^k e^{-st} dt \\ = \Gamma(1+k)^{-1} s^{k+1} \int_0^\infty S_y^{\lambda, k}(D)(0) e^{-st} dy. \end{aligned}$$

We choose some $0 < \varepsilon < \frac{1}{m+2} \min\{k, \sigma - \sigma_0\}$. According to Lemma 4.13 we have to show that the term

$$(58) \quad \begin{aligned} & e^{-\sigma x} |T_x^{\lambda, k}(D)(s)| \\ & + C(k) \left[\sum_{j=1}^{m+1} |w^j| \sum_{\ell=1}^{m+2-j} \frac{e^{(k-\varepsilon\ell)x}}{e^{kx}} e^{(k-\varepsilon\ell)x} x^{m-\ell+1} \int_0^x |S_y^{\lambda, k}(D)(s)| e^{-(\sigma-\varepsilon\ell)y} dy \right. \\ & \left. + |w| \sum_{\ell=1}^{m+2} \frac{e^{(k-\varepsilon\ell)x}}{e^{kx}} x^{m+2-\ell} \int_0^x |S_y^{\lambda, k}(D)(s)| e^{-(\sigma-\varepsilon\ell)y} dy \right] \end{aligned}$$

vanishes as $x \rightarrow \infty$. From (57) we know that for all $y > 0$

$$|S_y^{\lambda, k}(D)(0)| \leq C y^k e^{\sigma_0 y}.$$

Hence for each of the integrals in (58) we have

$$\int_0^x |S_y^{\lambda, k}(D)(0)| e^{-(\sigma-\varepsilon\ell)y} dy \leq C \int_0^\infty y^k e^{-(\sigma-\sigma_0-\varepsilon\ell)y} dy < \infty,$$

and it remains to check that

$$0 = \lim_{x \rightarrow \infty} e^{-\sigma x} |T_x^{\lambda, k}(D)(0)|.$$

To do this, we note first that by Lemma 4.8 for some constant $C > 0$ and all $x > 0$

$$|U_x^{\lambda, k}(D)(0)| \leq C e^{kx} \sup_{y < x} |S_x^{\lambda, k}(D)(0)|,$$

and hence by (57)

$$e^{-\sigma x} |T_x^{\lambda, k}(D)(0)| \leq C e^{-\sigma x} \sup_{y < x} |y^k R_x^{\lambda, k}(D)(0)| \leq C e^{(\sigma_0 - \sigma)x} x^k,$$

implying the conclusion. \square

REFERENCES

- [1] F. Bayart. Hardy spaces of Dirichlet series and their composition operators. *Monatsh. Math.* 136 (2002) 203-236
- [2] F. Bayart. Convergence and almost sure properties in Hardy spaces of Dirichlet series. To appear in *Math. Ann.* 2021, arXiv:2101.02990v1.
- [3] A. S. Besicovitch. *Almost periodic functions*. Dover Publications, Inc., New York, 1955.
- [4] H. Bohr. Einige Bemerkung über das Konvergenzproblem Dirichletscher Reihen. *Rendiconti del circolo Matematica di Palermo* 37 (1913) 1-16.
- [5] H. Bohr. Über die gleichmäßige Konvergenz Dirichletscher Reihen *J. Reine Angew. Math.* 143 (1913) 203-211.
- [6] D. Carando, A. Defant, F. Marceca, and I. Schoolmann. Vector-valued general Dirichlet series. *Studia Mathematica* 258 (2021) 269-316.
- [7] A. Defant, D. García, M. Maestre, and P. Sevilla-Peris. *Dirichlet Series and Holomorphic Functions in High Dimensions*. volume 37 of *New Mathematical Monographs*, Cambridge University Press, Cambridge (2019).
- [8] A. Defant and I. Schoolmann. \mathcal{H}_p -theory of general Dirichlet series. *J. Fourier Anal. Appl.*, 25(6)(2019) 3220–3258.
- [9] A. Defant and I. Schoolmann. Riesz means in Hardy spaces on Dirichlet groups. *Math. Ann.*, 378(1-2)(2020) 57–96.
- [10] A. Defant and I. Schoolmann. Variants of a theorem of Helson on general Dirichlet series. *J. Funct. Anal.*, 279 (5) (2020).
- [11] A. Defant, T. Fernández-Vidal, I. Schoolmann, and P. Sevilla-Peris. Fréchet spaces of general Dirichlet series. *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas* volume 115, Article number: 138 (2021).
- [12] A. Defant and I. Schoolmann. Riesz summability on boundary lines of holomorphic functions generated by Dirichlet series. Preprint 2021.
- [13] G. H. Hardy and M. Riesz. *The general theory of Dirichlet series*. Cambridge Tracts in Mathematics and Mathematical Physics 18 (1915).
- [14] H. Hedenmalm, P. Lindqvist, and K. Seip. A Hilbert space of Dirichlet series and systems of dilated function in $L^2(0, 1)$. *Duke Math. J.* 86 (1997) 1-37.
- [15] H. Helson. *Harmonic analysis*, Springer (2010).
- [16] H. Helson. Compact groups and Dirichlet series. *Ark. Mat.* 8 (1969) 139-143.
- [17] H. Helson. *Dirichlet series*. Regent Press (2005).
- [18] E. Landau. Über die gleichmäßige Konvergenz Dirichletscher Reihen. *J. Reine Angew. Math.* 143 (1921) 203-211.
- [19] I. Schoolmann. *Hardy spaces of general Dirichlet series and their maximal inequalities*. PhD thesis, Carl von Ossietzky University of Oldenburg, 2020

- [20] I. Schoolmann. On Bohr's theorem for general Dirichlet series. *Math. Nachr.* 293 (8) (2020) 1591–1612.
- [21] H. Queffélec and M. Queffélec. Diophantine approximation and Dirichlet series. Volume 2 of Harish-Chandra Research Institute Lecture Notes. Hindustan Book Agency (2013).

ANDREAS DEFANT
INSTITUT FÜR MATHEMATIK,
CARL VON OSSIETZKY UNIVERSITÄT,
26111 OLDENBURG, GERMANY.
Email address: `defant@mathematik.uni-oldenburg.de`

INGO SCHOOLMANN
INSTITUT FÜR MATHEMATIK,
CARL VON OSSIETZKY UNIVERSITÄT,
26111 OLDENBURG, GERMANY.
Email address: `ingo.schoolmann@uni-oldenburg.de`