

# ON COMPLEX-TIME HEAT KERNELS OF FRACTIONAL SCHRÖDINGER OPERATORS VIA PHRAGMÉN–LINDELÖF PRINCIPLE

KONSTANTIN MERZ

ABSTRACT. We consider fractional Schrödinger operators with possibly singular potentials and derive certain spatially averaged estimates for its complex-time heat kernel. The main tool is a Phragmén–Lindelöf theorem for polynomially bounded functions on the right complex half-plane. The interpolation leads to possibly nonoptimal off-diagonal bounds.

## 1. INTRODUCTION

Phragmén–Lindelöf theorems (cf. [2, 31, 40, 43]) are powerful complex analysis tools that extend the maximum modulus principle to certain unbounded domains. They are often used in the presence of exponential bounds, which are, e.g., available in the analysis of heat kernels of elliptic second-order differential operators. Suppose, e.g., that  $e^{-tH}$  is a symmetric Markov semigroup on  $L^2(\mathbb{R}^d)$  whose integral kernel satisfies

$$0 \leq e^{-tH}(x, y) \leq ct^{-\frac{d}{2}} \exp\left[-\frac{b|x-y|^2}{t}\right], \quad t > 0, x, y \in \mathbb{R}^d, \quad (1.1)$$

where  $b, c$  are positive constants. In [20, Theorem 3.4.8] and [21, Lemma 9, Theorem 10] Davies used the Phragmén–Lindelöf principle to show that for all  $\varepsilon > 0$  there is  $c_\varepsilon > 0$  such that

$$|e^{-zH}(x, y)| \leq c_\varepsilon (\operatorname{Re} z)^{-\frac{d}{2}} \exp\left[-\operatorname{Re}\left(\frac{b|x-y|^2}{(1+\varepsilon)z}\right)\right], \quad x, y \in \mathbb{R}^d \quad (1.2)$$

holds for all complex times  $z \in \mathbb{C}_+ := \{z \in \mathbb{C} : \operatorname{Re}(z) > 0\}$ , whenever (1.1) holds.

Coulhon and Sikora [19, Section 4] reversed Davies' ideas and used the Phragmén–Lindelöf principle to derive gaussian heat kernel estimates like (1.2) from suitably weighted off-diagonal estimates – so-called Davies–Gaffney estimates (cf. [19, (3.2)] or (3.20) for  $\theta = 0$  and  $\alpha = 2$ ) – and on-diagonal estimates like  $e^{-tH}(x, x) \leq ct^{-\frac{d}{2}}$ . The latter can often be derived from Sobolev inequalities corresponding to  $H$  by Nash's method, cf. Davies [20, Section 2.4] and Milman and Semenov [38].

Complex-time heat kernel estimates like (1.2) are of paramount importance in many problems in harmonic analysis and partial differential equations, e.g., in proving  $L^p$  boundedness of spectral multipliers and convergence of Riesz means, and investigating maximal regularity properties of the Schrödinger evolution  $e^{itH}$  for operators  $H$  whose heat kernels satisfy (sub-)gaussian estimates, see, e.g., [4, 9, 7, 5, 6, 8, 12, 13, 15, 14, 16, 17, 18, 23, 24, 29, 42].

*Date:* July 29, 2021.

*2010 Mathematics Subject Classification.* Primary: 35K08; Secondary: 35J10.

*Key words and phrases.* Complex-time heat kernel, Phragmén–Lindelöf principle, fractional Laplace, fractional Schrödinger operator, Davies–Gaffney estimate.

On the other hand, there has been recent interest in the fractional Laplace operator  $(-\Delta)^{\alpha/2}$  in  $L^2(\mathbb{R}^d)$  for  $\alpha > 0$  and its generated holomorphic semigroup  $e^{-z(-\Delta)^{\alpha/2}}$  for  $z \in \mathbb{C}_+$ . For  $\alpha > 0$  Blumenthal and Gettoor [3] proved

$$|e^{-t(-\Delta)^{\alpha/2}}(x)| \leq c_{d,\alpha} \frac{t}{(t^{1/\alpha} + |x|)^{d+\alpha}}, \quad t > 0, x \in \mathbb{R}^d, \quad (1.3)$$

whereas for  $\alpha = 1$  one has the representation via Poisson's kernel

$$e^{-t(-\Delta)^{1/2}}(x) = c_d \frac{t}{(t^2 + |x|^2)^{(d+1)/2}}$$

for an explicit constant  $c_d > 0$ . The slow decay of these kernels and their extension to  $\mathbb{C}_+$  – that we discuss momentarily – complicates many arguments in problems where heat kernel estimates are the central element of the analysis like [26, 37, 11]. In particular, they impede the generalization of the above mentioned works on spectral multipliers for operators whose heat kernels only decay algebraically. As (1.2) indicates, complex-time heat kernels satisfy worse bounds as  $|\arg(z)|$  increases. Thus, the derivation of sharp estimates becomes even more crucial in this scenario.

Zhao and Zheng [46, Theorem 1.3] recently proved uniform complex-time heat kernel estimates for  $(-\Delta)^{\alpha/2}$  and all  $\alpha > 0$  using the stationary phase method. While for  $\alpha = 1$  one has the explicit formula

$$e^{-z(-\Delta)^{1/2}}(x) = c_d \frac{z}{(z^2 + |x|^2)^{(d+1)/2}}, \quad z \in \mathbb{C}_+, x \in \mathbb{R}^d,$$

the derivation of such estimates for  $\alpha \neq 1$  is rather intricate. Pointwise estimates in the case  $|\arg(z)| = \pi/2$  are however well known, see, e.g., Miyachi [39, Proposition 5.1], Wainger [44, pp. 41-52], Huang et al [33] and [46, pp. 2-3].

Using perturbation theory, Zhao and Zheng extended their results and derived uniform complex-time heat kernel estimates for fractional Schrödinger operators

$$H_\alpha := (-\Delta)^{\alpha/2} + V \quad \text{in } L^2(\mathbb{R}^d) \quad (1.4)$$

when  $V \in L^1_{\text{loc}}(\mathbb{R}^d; \mathbb{R})$  belongs to the higher order Kato class  $K_\alpha(\mathbb{R}^d)$  (cf. [22, 28, 47] for a precise definition). In this case  $V$  is infinitesimally form bounded with respect to  $(-\Delta)^{\alpha/2}$  and  $H_\alpha$  can be defined as the self-adjoint Friedrichs extension of the corresponding quadratic form with form core  $C_c^\infty(\mathbb{R}^d)$ . Their estimates for the kernel of the holomorphic extension of  $e^{-tH_\alpha}$  to  $\mathbb{C}_+$  read as follows.

**Theorem 1.1** ([46, Theorem 1.5]). *Let  $\alpha > 0$ ,  $z = |z|e^{i\theta} \in \mathbb{C}_+$ , and  $V \in K_\alpha(\mathbb{R}^d)$ . Then for any  $0 < \varepsilon \ll 1$  there are  $c_{d,\alpha} > 0$  and  $\mu_{\varepsilon,V,d,\alpha} > 0$  such that*

$$|e^{-zH_\alpha}(x, y)| \leq c_{d,\alpha} e^{\mu_{\varepsilon,V,d,\alpha}|z|} (\cos \theta)^{-d(\frac{1}{\alpha} - \frac{1}{2})\mathbf{1}_{\{\alpha < 1\}} - (\frac{d}{2} + \alpha - 1)\mathbf{1}_{\{\alpha \geq 1\}}} \frac{|z|}{(|z|^{\frac{1}{\alpha}} + |x - y|)^{d+\alpha}} \quad (1.5)$$

holds for all  $x, y \in \mathbb{R}^d$ .

Estimate (1.5) reflects the best possible off-diagonal pointwise decay for  $|x - y| \gg |z|^{1/\alpha}$  and all  $|\theta| < \pi/2$  (compare with (1.3)) and is particularly useful for  $|z| \lesssim 1$ . (See [46, p. 22] for the precise definition of  $\mu_{\varepsilon,V,d,\alpha}$ .)

In this note we derive a Phragmén–Lindelöf principle for polynomially bounded functions and apply it to obtain suitably weighted and averaged estimates for  $e^{-zH_\alpha}$  that are not uniform in  $\theta$  but do not deteriorate for  $|z| \gg 1$ . Moreover, we can

allow for critically singular potentials  $V$ , like the Hardy potential  $|x|^{-\alpha}$ , which do not belong to  $K_\alpha(\mathbb{R}^d)$ . In this case  $e^{-tH_\alpha}$  is generally not  $L^1 \rightarrow L^\infty$  bounded. On the downside, it is not clear whether the off-diagonal decay of our estimates is optimal, especially when  $|\pi/2 - |\theta|| \ll 1$ .

**Organization and notation.** In Section 2 we state and prove a Phragmén–Lindelöf theorem for polynomially bounded functions. We apply this theorem in Section 3 to derive estimates for the complex-time heat kernel of fractional Schrödinger operators with non-negative potentials (Corollary 3.1 and Corollary 3.2 for a pointwise estimate), and potentials with a non-vanishing, possibly singular, negative part (Theorem 3.7 and Corollaries 3.11 and 3.12). In Section 4 we collect some properties of the estimates – so-called dyadic Davies–Gaffney estimates (Definitions 3.3 and 3.9) – that we discuss when  $V$  has a negative part.

We write  $A \lesssim B$  for two non-negative quantities  $A, B \geq 0$  to indicate that there is a constant  $C > 0$  such that  $A \leq CB$ . If  $C = C_\tau$  depends on a parameter  $\tau$ , we write  $A \lesssim_\tau B$ . The dependence on fixed parameters like  $d$  and  $\alpha$  is sometimes omitted. The notation  $A \sim B$  means  $A \lesssim B \lesssim A$ . All constants are denoted by  $c$  or  $C$  and are allowed to change from line to line. We abbreviate  $A \wedge B := \min\{A, B\}$  and  $A \vee B := \max\{A, B\}$ . The Heaviside function is denoted by  $\theta(x)$ . We use the convention  $\theta(0) = 1$ . The indicator function and the Lebesgue measure of a set  $\Omega \subseteq \mathbb{R}^d$  are denoted by  $\mathbf{1}_\Omega$  and  $|\Omega|$ , respectively. The euclidean distance between two sets  $\Omega_1, \Omega_2 \subseteq \mathbb{R}^d$  is denoted by  $d(\Omega_1, \Omega_2) := \inf_{x \in \Omega_1, y \in \Omega_2} |x - y|$ . If  $T : L^p(\mathbb{R}^d) \rightarrow L^q(\mathbb{R}^d)$  is a bounded linear operator, we write  $T \in \mathcal{B}(L^p \rightarrow L^q)$  and denote its operator norm by  $\|T\|_{p \rightarrow q}$ . For  $1 \leq p \leq \infty$  we write  $p' = (1 - 1/p)^{-1}$ .

## 2. PHRAGMÉN–LINDELÖF PRINCIPLE WITH POLYNOMIAL BOUNDS

**Theorem 2.1.** *Let  $X$  be a Banach space equipped with a norm  $\|\cdot\|$  and  $F : \mathbb{C}_+ = \{z \in \mathbb{C} : \operatorname{Re}(z) > 0\} \rightarrow X$  be a holomorphic function satisfying*

$$\|F(|z|e^{i\theta})\| \leq a_1(|z|\cos\theta)^{-\beta_1} \quad \text{and} \quad (2.1)$$

$$\|F(|z|)\| \leq a_1|z|^{-\beta_1} \left(\frac{a_2}{|z|}\right)^{-\beta_2} \cdot \left(\frac{a_3}{|z|}\right)^{\beta_3} \quad (2.2)$$

for some  $a_1, a_2, a_3 > 0$ ,  $\beta_1, \beta_2, \beta_3 \geq 0$ , all  $|z| > 0$ , and all  $|\theta| < \pi/2$ . Then for all  $\varepsilon \in (0, 1)$  one has

$$\|F(|z|e^{i\theta})\| \leq a_1(|z|\cos\theta)^{-\beta_1} \left[ 1 \wedge \varepsilon^{-\beta_1} \left( \left(\frac{a_2}{|z|}\right)^{-\beta_2} \cdot \left(\frac{a_3}{|z|}\right)^{\beta_3} \right)^{1-|\theta|/\gamma(\varepsilon, \theta)} \right] \quad (2.3)$$

for all  $|\theta| < \pi/2$  and  $|z| > 0$ , where  $\gamma(\varepsilon, \theta) := \varepsilon|\theta| + (1 - \varepsilon)\pi/2$ .

*Proof.* For  $\gamma \in (0, \pi/2)$  and  $z = |z|e^{i\theta}$  let

$$G(z) := z^{-\beta_1} F(z^{-1}) \cdot H_2(z) \cdot H_3(z) \quad (2.4)$$

with

$$H_2(z) := \exp\left(\beta_2 \log(a_2 z) \left(1 + \frac{i}{2\gamma} \log(a_2 z)\right)\right) = (a_2 z)^{\beta_2(1 + \frac{i}{2\gamma} \log(a_2 z))} \quad (2.5)$$

and

$$H_3(z) := \exp\left(-\beta_3 \log(a_3 z) \left(1 + \frac{i}{2\gamma} \log(a_3 z)\right)\right) = (a_3 z)^{-\beta_3(1 + \frac{i}{2\gamma} \log(a_3 z))}. \quad (2.6)$$

Here  $\log(|z|e^{i\theta}) := \log(|z|) + i\theta$  with  $|\theta| < \pi$  is the principal branch of the logarithm. By the assumptions on  $\|F(z)\|$  for  $\theta = 0$  and  $\theta = \gamma$ , and the bounds

$$|H_2(|z|)| \leq (a_2|z|)^{\beta_2}$$

and

$$|H_2(|z|e^{i\gamma})| = \left| \exp\left(\frac{i\beta_2}{2}((\log|a_2z|)^2/\gamma + \gamma)\right) \right| = 1,$$

and

$$|H_3(|z|)| \leq (a_3|z|)^{-\beta_3} \quad \text{and} \quad |H_3(|z|e^{i\gamma})| = 1,$$

respectively, we have

$$\|G(|z|)\| \leq |z|^{-\beta_1} \cdot a_1|z|^{\beta_1} (a_2|z|)^{-\beta_2} (a_2|z|)^{\beta_2} \cdot (a_3|z|)^{\beta_3} (a_3|z|)^{-\beta_3} \leq a_1 \quad (2.7)$$

and

$$\|G(|z|e^{i\gamma})\| \leq |z|^{-\beta_1} \cdot a_1|z|^{\beta_1} (\cos \gamma)^{-\beta_1} = a_1 (\cos \gamma)^{-\beta_1}. \quad (2.8)$$

Combining (2.7) and (2.8) using the three lines lemma shows

$$\|G(|z|e^{i\theta})\| \leq a_1 (\cos \gamma)^{-\beta_1 \theta / \gamma} \leq a_1 (\cos \gamma)^{-\beta_1}$$

for all  $0 \leq \theta \leq \gamma$  and  $|z| > 0$ . Plugging this estimate and the identities

$$\begin{aligned} |H_2(z^{-1})^{-1}| &= |(a_2 z^{-1})^{-\beta_2(1 + \frac{i}{2\gamma} \log(a_2 z^{-1}))}| = (a_2 |z|^{-1})^{-\beta_2(1 + \theta/\gamma)} \\ |H_3(z^{-1})^{-1}| &= (a_3 |z|^{-1})^{\beta_3(1 + \theta/\gamma)} \end{aligned}$$

into the expression for  $F(z)$  in (2.4), i.e.,

$$F(|z|e^{i\theta}) = z^{-\beta_1} G(z^{-1}) H_2(z^{-1})^{-1} \cdot H_3(z^{-1})^{-1},$$

implies for  $-\gamma \leq \theta < 0$  and  $|z| > 0$ ,

$$\|F(|z|e^{i\theta})\| \leq a_1 |z|^{-\beta_1} (\cos \gamma)^{-\beta_1} (a_2 |z|^{-1})^{-\beta_2(1 + \theta/\gamma)} \cdot (a_3 |z|^{-1})^{\beta_3(1 + \theta/\gamma)}. \quad (2.9)$$

By a reflection along the real axis we conclude the corresponding bound with  $\theta$  replaced by  $-\theta \in [-\gamma, 0)$ . Choosing  $\gamma = \gamma(\varepsilon, \theta) = \varepsilon|\theta| + (1 - \varepsilon)\pi/2$  for any  $0 < \varepsilon < 1$  (which ensures  $|\theta| < \gamma < \pi/2$ ) lets us estimate  $(\cos \gamma)^{-\beta_1} \leq \varepsilon^{-\beta_1} (\cos \theta)^{-\beta_1}$  and conclude the proof of (2.3), upon taking the minimum between (2.9) and (2.1).  $\square$

*Remarks 2.2.* (1) For  $\beta_3 = 0$  one observes that the decay of  $|F(z)|$  in the regime  $a_2 \gg |z|$  becomes weaker as  $|\theta|$  increases.

(2) The choice  $\gamma = \varepsilon|\theta| + (1 - \varepsilon)\pi/2$  suppresses the effect of large  $|\theta|$  for  $\varepsilon \ll 1$ , but does not prevent the vanishing of decay as  $|\theta| \rightarrow \pi/2$ .

### 3. APPLICATION - SEMIGROUPS OF FRACTIONAL SCHRÖDINGER OPERATORS

Let  $d \in \mathbb{N}$ ,  $\alpha > 0$ ,  $V \in L^1_{\text{loc}}(\mathbb{R}^d : \mathbb{R})$ , and recall the notation (1.4) for the fractional Schrödinger operator  $H_\alpha = (-\Delta)^{\alpha/2} + V$  in  $L^2(\mathbb{R}^d)$ . We assume that  $V$  is such that the quadratic form of  $H_\alpha$  is non-negative on  $C_c^\infty(\mathbb{R}^d)$  so that it gives rise to a self-adjoint operator by Friedrichs' theorem. In particular,  $e^{-zH_\alpha}$  is a bounded, holomorphic semigroup on  $L^2(\mathbb{R}^d)$  whenever  $\text{Re}(z) > 0$  [34, p. 493]. In the following we derive certain weighted and averaged estimates for the extension of the semigroup  $e^{-tH_\alpha}$  to complex times  $z \in \mathbb{C}_+$  and distinguish between the cases where  $V \geq 0$  and where  $V$  has a non-vanishing negative part, respectively.

**3.1. Non-negative potentials.** If  $\alpha \in (0, 2)$ , then  $e^{-t(-\Delta)^{\alpha/2}}(x) \sim_{d,\alpha} t^{-d/\alpha}(1 + |x|/t^{1/\alpha})^{-d/\alpha}$ , i.e., the kernel is in particular positive [3]. Therefore, Trotter's formula implies for  $V \geq 0$  the upper bound

$$e^{-tH_\alpha}(x, y) \lesssim_{d,\alpha} t^{-\frac{d}{\alpha}} \left(1 + \frac{|x-y|}{t^{1/\alpha}}\right)^{-d-\alpha}. \quad (3.1)$$

In turn, (3.1) implies the following weighted  $L^2 \rightarrow L^\infty$  estimate for any  $r, t > 0$ ,

$$\sup_{y \in \mathbb{R}^d} \int_{\mathbb{R}^d \setminus B_y(r)} |\exp(-tH_\alpha)(x, y)|^2 dx \lesssim t^{-\frac{d}{\alpha}} \left(1 + \frac{r}{t^{1/\alpha}}\right)^{-d-2\alpha}. \quad (3.2)$$

The uniform complex-time heat kernel estimates (1.5) by Zhao and Zheng yield the following extension to  $z \in \mathbb{C}_+$ , i.e.,

$$\begin{aligned} & \sup_{y \in \mathbb{R}^d} \int_{\mathbb{R}^d \setminus B_y(r)} |\exp(-zH_\alpha)(x, y)|^2 dx \\ & \lesssim e^{2\mu_\varepsilon, \nu|z|} \cdot (\cos \theta)^{-2d(\frac{1}{\alpha} - \frac{1}{2})\mathbf{1}_{\{\alpha < 1\}} - 2(\frac{d}{2} + \alpha - 1)\mathbf{1}_{\{\alpha \geq 1\}}} |z|^{-\frac{d}{\alpha}} \left(1 + \frac{r}{|z|^{1/\alpha}}\right)^{-d-2\alpha}. \end{aligned} \quad (3.3)$$

In some applications one is merely in possession of averaged and possibly weighted analogs of (3.1), such as (3.2). This is typically the case when  $V$  has a singular negative part, which will be discussed in more detail in Section 3.2. The following corollary illustrates how the Phragmén–Lindelöf principle can be used to extend weighted  $L^2 \rightarrow L^\infty$  estimates like (3.2) to complex times.

**Corollary 3.1.** *Let  $\alpha \in (0, 2)$  and  $V \geq 0$ . Let further  $z = |z|e^{i\theta}$  with  $|\theta| \in [0, \pi/2)$ ,  $r > 0$ ,  $\varepsilon \in (0, 1)$ , and*

$$\beta_{d,\alpha,\varepsilon}(\theta) := (d + 2\alpha) \left(1 - \frac{|\theta|}{\varepsilon|\theta| + (1-\varepsilon)\pi/2}\right) \geq 0. \quad (3.4)$$

Then

$$\sup_{y \in \mathbb{R}^d} \int_{\mathbb{R}^d \setminus B_y(r)} |\exp(-zH_\alpha)(x, y)|^2 dx \lesssim_{d,\alpha,\varepsilon} (|z| \cos \theta)^{-\frac{d}{\alpha}} \left(1 + \frac{r}{|z|^{1/\alpha}}\right)^{-\beta_{d,\alpha,\varepsilon}(\theta)}. \quad (3.5)$$

Although the singularity of  $(\cos \theta)^{-d/\alpha}$  in (3.5) is less severe than in (3.3), the decay in the region  $r \gg |z|^{1/\alpha}$  becomes weaker as  $|\theta|$  increases.

*Proof.* For  $y \in \mathbb{R}^d$  define the analytic function  $F_y : \mathbb{C}_+ \rightarrow \mathbb{C}$  by

$$F_y(z) = \left( \int_{\mathbb{R}^d} e^{-zH_\alpha}(x, y) f(x) dx \right)^2$$

for any normalized  $f \in L^2(\mathbb{R}^d)$  with  $\text{supp } f \subseteq \mathbb{R}^d \setminus B_y(r)$ . For  $|\theta| < \frac{\pi}{2}$  and  $z = |z|e^{i\theta}$ , we have, using the unitarity of  $e^{-itH_\alpha}$  and (3.1) with  $t$  replaced by  $|z| \cos \theta$ ,

$$\begin{aligned} \sup_{y \in \mathbb{R}^d} |F_y(z)| & \leq \|e^{-zH_\alpha}\|_{2 \rightarrow \infty}^2 \leq \|e^{-|z| \cos \theta H_\alpha}\|_{2 \rightarrow \infty}^2 \|e^{-i|z| \sin \theta H_\alpha}\|_{2 \rightarrow 2}^2 \\ & \leq C_{1,d,\alpha} (|z| \cos \theta)^{-\frac{d}{\alpha}}. \end{aligned} \quad (3.6)$$

On the other hand we have for  $\theta = 0$ ,

$$\begin{aligned} \sup_{y \in \mathbb{R}^d} |F_y(|z|)| &\leq \int_{\mathbb{R}^d \setminus B_y(r)} \left| e^{-|z|H_\alpha(x, y)} \right|^2 dx \leq C_{2,d,\alpha} |z|^{-d/\alpha} \left( 1 + \frac{r}{|z|^{1/\alpha}} \right)^{-d-2\alpha} \\ &\leq C_{2,d,\alpha} |z|^{-d/\alpha} \left( \frac{r^\alpha}{|z|} \right)^{-\frac{d}{\alpha}-2} \end{aligned}$$

by (3.2). Thus, we may apply Theorem 2.1 with  $X = \mathbb{C}$ ,  $a_1 = \max\{C_{1,d,\alpha}, C_{2,d,\alpha}\}$ ,  $a_2 = r^\alpha$ ,  $\beta_1 = d/\alpha$ ,  $\beta_2 = (d + 2\alpha)/\alpha$ , and  $\beta_3 = 0$ , and obtain

$$|F_y(z)| \lesssim_{d,\alpha,\varepsilon} (|z| \cos \theta)^{-\frac{d}{\alpha}} \left[ 1 \wedge \left( \frac{r^\alpha}{|z|} \right)^{-\frac{d+2\alpha}{\alpha} \left( 1 - \frac{|\theta|}{\varepsilon|\theta| + (1-\varepsilon)\pi/2} \right)} \right], \quad (3.7)$$

which shows (3.5).  $\square$

Similarly, one can use Theorem 2.1 to derive pointwise estimates.

**Corollary 3.2.** *Let  $\alpha \in (0, 2)$  and  $V \geq 0$ . Let further  $z = |z|e^{i\theta}$  with  $|\theta| \in [0, \pi/2)$ ,  $x, y \in \mathbb{R}^d$ ,  $\varepsilon \in (0, 1)$ , and*

$$\beta_{d,\alpha,\varepsilon}(\theta) := (d + \alpha) \left( 1 - \frac{|\theta|}{\varepsilon|\theta| + (1-\varepsilon)\pi/2} \right) \geq 0. \quad (3.8)$$

Then

$$|e^{-zH_\alpha}(x, y)| \lesssim_{d,\alpha,\varepsilon} (|z| \cos \theta)^{-\frac{d}{\alpha}} \left( 1 + \frac{|x-y|}{|z|^{1/\alpha}} \right)^{-\beta_{d,\alpha,\varepsilon}(\theta)}. \quad (3.9)$$

*Proof.* For  $z = |z|e^{i\theta}$  with  $|\theta| \in [0, \pi/2)$  we use  $\|e^{-i\operatorname{Im}(z)H_\alpha}\|_{2 \rightarrow 2} = 1$  and estimate

$$\|e^{-zH_\alpha}\|_{1 \rightarrow \infty} \leq \|e^{-\operatorname{Re}(z)H_\alpha/2}\|_{1 \rightarrow 2} \|e^{-\operatorname{Re}(z)H_\alpha/2}\|_{2 \rightarrow \infty} \leq C_{1,d,\alpha} (|z| \cos \theta)^{-\frac{d}{\alpha}},$$

which shows

$$|e^{-zH_\alpha}(x, y)| \leq C_{1,d,\alpha} (|z| \cos \theta)^{-\frac{d}{\alpha}}, \quad x, y \in \mathbb{R}^d. \quad (3.10)$$

On the other hand (3.1) implies

$$|e^{-|z|H_\alpha}(x, y)| \leq C_{2,d,\alpha} |z|^{-\frac{d}{\alpha}} \left( \frac{|x-y|^\alpha}{|z|} \right)^{-\frac{d+\alpha}{\alpha}}, \quad x, y \in \mathbb{R}^d. \quad (3.11)$$

Thus, by Theorem 2.1 with  $X = \mathbb{C}$ ,  $a_1 = \max\{C_{1,d,\alpha}, C_{2,d,\alpha}\}$ ,  $a_2 = |x-y|^\alpha$ ,  $\beta_1 = d/\alpha$ ,  $\beta_2 = (d + \alpha)/\alpha$ , and  $\beta_3 = 0$ , we obtain

$$|e^{-zH_\alpha}(x, y)| \lesssim_{d,\alpha,\varepsilon} (|z| \cos \theta)^{-\frac{d}{\alpha}} \left[ 1 \wedge \left( \frac{|x-y|^\alpha}{|z|} \right)^{-\frac{d+\alpha}{\alpha} \left( 1 - \frac{|\theta|}{\varepsilon|\theta| + (1-\varepsilon)\pi/2} \right)} \right] \quad (3.12)$$

for all  $x, y \in \mathbb{R}^d$ . This shows (3.9).  $\square$

**3.2. Potentials with a negative part.** We now consider the situation where the semigroup  $e^{-tH_\alpha}$  is not necessarily  $L^2 \rightarrow L^\infty$  bounded anymore. This typically occurs when  $V$  has a non-vanishing, singular negative part [36, 38]. In these cases  $e^{-tH_\alpha}$  may nevertheless satisfy certain weighted  $L^p \rightarrow L^q$  estimates with  $1 < p \leq q < \infty$ . To introduce the estimates that we discuss here, we denote by

$$A_2(x, r, k) := B_x(2^k r) \setminus B_x(2^{k-1} r) \quad \text{with} \quad A_2(x, r, 0) := B_x(r) \quad \text{and} \quad k \in \mathbb{N}_0$$

dyadic annuli around  $x \in \mathbb{R}^d$ .

**Definition 3.3.** Let  $r > 0$ ,  $(T_r)_{r>0}$  be a family of linear bounded operators on  $L^2(\mathbb{R}^d)$ ,  $1 \leq p \leq q \leq \infty$ ,  $\beta, \sigma > 0$ , and  $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  satisfy  $g(\lambda) \sim_\beta (1 + \lambda)^{-\beta}$ . Then  $T_r$  is said to satisfy the *dyadic*  $(p, q, \sigma)$  *Davies–Gaffney estimate* if there is a finite constant  $C_{\text{DG}} = C_{\text{DG}}(d, p, q, \beta, \sigma) > 0$  such that

$$\|\mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x,r,k)}\|_{p \rightarrow q} \leq C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{q})} g(2^k) 2^{\frac{kd}{\sigma}}, \quad x \in \mathbb{R}^d, k \in \mathbb{N}_0, \quad (3.13)$$

and if  $\beta > d(1/\sigma + 1/q)$ .

This definition is inspired by Davies [21] and the notion of generalized gaussian estimates introduced by Blunck and Kunstmann, cf. [9, p. 920]. Heuristically, the projection onto  $B_x(r)$  captures the singularity of  $T_r$  at  $x$ , whereas the projection onto  $A_2(x, r, k)$  controls its decay at distance  $2^k r$ . We use dyadic annuli instead of annuli with constant thickness, since they allow to exploit decay more effectively. Similar estimates were, e.g., used by Schreieck and Voigt [41] on  $L^p$  independence of the spectrum of Schrödinger operators with form small potentials, and later systematically studied and exploited in a series of works by Blunck and Kunstmann [4, 9, 7, 5, 6, 8] on spectral multiplier theorems for operators whose semigroups need not have a bounded kernel but obey the mentioned generalized gaussian estimates.

Estimate (3.13) with  $p = 1$ ,  $q = \infty$ , and the relaxed assumption  $\beta \geq d/\sigma$  is equivalent to a pointwise estimate for the kernel of  $T_r$ .

**Proposition 3.4.** *Suppose  $r, \beta, \sigma > 0$ ,  $\beta \geq d/\sigma$ , and  $(T_r)_{r>0}$  is a family of linear operators in  $\mathcal{B}(L^1 \rightarrow L^\infty)$  with integral kernel  $T_r(x, y)$ . Then the following statements are equivalent.*

(1)  $T_r$  satisfies (3.13) with  $p = 1$ ,  $q = \infty$ , and  $C_{\text{DG}}$  replaced by  $c_{\beta, d, \sigma} C_{\text{DG}}$  for some  $c_{\beta, d, \sigma} > 0$ .

(2) One has  $|T_r(x, y)| \lesssim_{\beta, d, \sigma} C_{\text{DG}} r^{-d} (1 + |x - y|/r)^{-(\beta - d/\sigma)}$  for all  $x, y \in \mathbb{R}^d$ .

*Proof.* (1)  $\Rightarrow$  (2): In this case (3.13) asserts

$$\begin{aligned} \sup_{y, z \in \mathbb{R}^d} \mathbf{1}_{B_x(r)}(y) |T_r(y, z)| \mathbf{1}_{A_2(x, r, k)}(z) &\leq c_{\beta, d, \sigma} C_{\text{DG}} r^{-d} g(2^k) 2^{kd/\sigma} \\ &\lesssim_{\beta, d, \sigma} C_{\text{DG}} r^{-d} 2^{-k(\beta - d/\sigma)} \end{aligned}$$

for all  $x \in \mathbb{R}^d$  and  $k \in \mathbb{N}_0$ . Choosing  $y = x$  and  $k \in \mathbb{N}_0$  such that  $2^k = \max\{1, |x - z|/r\}$  yields  $|T_r(x, z)| \lesssim_{\beta, d, \sigma} C_{\text{DG}} r^{-d} (1 + |x - z|/r)^{-(\beta - d/\sigma)}$  for all  $x, z \in \mathbb{R}^d$ .

(2)  $\Rightarrow$  (1): We estimate

$$\begin{aligned} \|\mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x, r, k)}\|_{1 \rightarrow \infty} &= \sup_{y, z \in \mathbb{R}^d} \mathbf{1}_{B_x(r)}(y) |T_r(y, z)| \mathbf{1}_{A_2(x, r, k)}(z) \\ &\leq \sup_{y \in B_x(r)} \sup_{z \in \mathbb{R}^d \setminus B_x(2^{k-1}r)} |T_r(y, z)| \\ &\lesssim_{\beta, d, \sigma} C_{\text{DG}} r^{-d} (1 + 2^k \theta(k-2))^{-(\beta - d/\sigma)}, \end{aligned}$$

which concludes the proof.  $\square$

Further consequences of Definition 3.3 will be discussed in Section 4. They play an important role in the subsequent analysis.

The example of the semigroup of  $H_\alpha$  with the Hardy potential  $V = a|x|^{-\alpha}$  illustrates that (3.13) is a reasonable assumption. Note that  $|x|^{-\alpha} \notin K_\alpha(\mathbb{R}^d)$ . The resulting operator

$$\mathcal{L}_{a, \alpha} := (-\Delta)^{\alpha/2} + \frac{a}{|x|^\alpha} \quad \text{in } L^2(\mathbb{R}^d) \quad (3.14)$$

for  $\alpha \in (0, 2 \wedge d)$  and  $a \geq a_* \equiv a_*(\alpha, d) > -\infty$  is sometimes called generalized or fractional Hardy operator. By the sharp Hardy–Kato–Herbst inequality,  $\mathcal{L}_{a,\alpha}$  is bounded from below and non-negative in the sense of quadratic forms if and only if  $a \geq a_*$ . We refer to Kato [34, Chapter 5, Equation (5.33)] (for  $\alpha = 1$  and  $d = 3$ ) and Herbst [30, Equation (2.6)] and also [25, 27, 35, 45] for proofs of this fact and the explicit expression of  $a_*$ .

For  $a \in [a_*, 0]$  Bogdan et al [10] proved pointwise heat kernel bounds, namely

$$e^{-t\mathcal{L}_{a,\alpha}}(x, y) \sim_{d,\alpha,a} \left(1 \vee \frac{t^{\frac{1}{\alpha}}}{|x|}\right)^\delta \left(1 \vee \frac{t^{\frac{1}{\alpha}}}{|y|}\right)^\delta \frac{t}{(t^{\frac{1}{\alpha}} + |x - y|)^{d+\alpha}}, \quad t > 0, x, y \in \mathbb{R}^d \setminus \{0\}, \quad (3.15)$$

where  $\delta = \delta(a, d, \alpha) \in [0, (d - \alpha)/2]$  satisfies  $\delta(0, d, \alpha) = 0$ ,  $\delta(a_*, d, \alpha) = (d - \alpha)/2$ , and increases monotonously as  $a$  decreases. An explicit formula for  $\delta(a, d, \alpha)$  is, e.g., contained in [10] or Frank et al [25]. In particular,  $e^{-t\mathcal{L}_{a,\alpha}}$  is  $L^p \rightarrow L^p$  bounded for  $a < 0$  if and only if  $p \in (d/(d - \delta), d/\delta)$ . This follows from (3.15), duality, and  $(e^{-\mathcal{L}_{a,\alpha}} \mathbf{1}_{B_0(1)})(x) \gtrsim |x|^{-\delta}$ . Moreover,  $e^{-t\mathcal{L}_{a,\alpha}}$  satisfies the  $L^2 \rightarrow L^\infty$  estimates in (3.2) if and only if  $a \geq 0$ .

**Example 3.5.** Let  $\alpha \in (0, 2 \wedge d)$ ,  $t > 0$ , and  $r_t := t^{1/\alpha}$ .

(1) If  $a \in [a_*, 0)$  and  $p \in (d/(d - \delta), 2]$ , then  $T_{r_t} := e^{-t\mathcal{L}_{a,\alpha}}$  satisfies the dyadic  $(p, p', p')$ ,  $(p, 2, p')$ , and  $(2, p', 2)$  Davies–Gaffney estimate (3.13) with  $g(\lambda) \sim_{d,\alpha} (1 + \lambda)^{-d-\alpha}$ .

(2) If  $a \geq 0$  and  $1 \leq p \leq 2 \leq q \leq \infty$ , then  $T_{r_t} := e^{-t\mathcal{L}_{a,\alpha}}$  satisfies the dyadic  $(p, q, p')$  Davies–Gaffney estimate (3.13) with  $g(\lambda) \sim_{d,\alpha} (1 + \lambda)^{-d-\alpha}$ .

*Proof.* (1) By Hölder’s inequality and (3.15), one obtains for  $x \in \mathbb{R}^d$  and  $k \in \mathbb{N} \setminus \{1\}$

$$\begin{aligned} & \sup_{f \in L^p: \|f\|_p=1} \|\mathbf{1}_{B_x(r_t)} e^{-t\mathcal{L}_{a,\alpha}} \mathbf{1}_{A_2(x, r_t, k)} f\|_{p'}^{p'} \\ & \leq \int_{\frac{|z-x|}{r_t} \leq 1} dz \left(1 \vee \frac{r_t}{|z|}\right)^{\delta p'} \int_{\frac{|y-x|}{r_t} \in [2^{k-1}, 2^k]} dy \left(1 \vee \frac{r_t}{|y|}\right)^{\delta p'} \frac{r_t^{\alpha p'}}{(r_t + |z - y|)^{p'(d+\alpha)}} \\ & = r_t^{-d(p'-2)} (1 + 2^k)^{-(d+\alpha)p'} \cdot 2^{kd} \end{aligned}$$

where we used  $|z - y| \geq |y - x| - r_t \geq (2^{k-1} - 1)r_t$ . For  $k \in \{0, 1\}$  it suffices to integrate  $y$  over  $|y - x| \leq 2r_t$ , in which case the left side is bounded by a constant times  $r_t^{-d(p'-2)} \lesssim r_t^{-d(p'-2)} (1 + 2^k)^{-(d+\alpha)p'} \cdot 2^{kd}$  as well. Taking the  $p'$ -th root yields the dyadic  $(p, p', p')$  Davies–Gaffney estimate (3.13). The  $(p, 2, p')$  and  $(2, p', 2)$  Davies–Gaffney estimates follow similarly. For  $x \in \mathbb{R}^d$  and  $k \in \mathbb{N}_0$ , one has

$$\begin{aligned} & \sup_{f \in L^2: \|f\|_2=1} \|\mathbf{1}_{B_x(r_t)} e^{-t\mathcal{L}_{a,\alpha}} \mathbf{1}_{A_2(x, r_t, k)} f\|_{p'}^{p'} \\ & \leq \int_{\frac{|z-x|}{r_t} \leq 1} dz \left(1 \vee \frac{r_t}{|z|}\right)^{\delta p'} \left[ \int_{\frac{|y-x|}{r_t} \in [2^{k-1}, 2^k]} dy \left(1 \vee \frac{r_t}{|y|}\right)^{2\delta} \frac{r_t^{2\alpha}}{(r_t + |z - y|)^{2(d+\alpha)}} \right]^{\frac{p'}{2}} \\ & \lesssim r_t^{-d(p'-2)/2} (1 + 2^k)^{-(d+\alpha)p'} \cdot 2^{kdp'/2} \end{aligned}$$

which shows the  $(2, p', 2)$  estimate. The  $(p, 2, p')$  estimate is shown analogously.



(2) By (3.1) (Trotter's formula) and a similar computation, one obtains

$$\|\mathbf{1}_{B_x(r_t)} e^{-t\mathcal{L}_{a,\alpha}} \mathbf{1}_{A_2(x,r_t,k)}\|_{p \rightarrow q} \lesssim r_t^{-d(\frac{1}{p}-\frac{1}{q})} (1+2^k)^{-(d+\alpha)} \cdot 2^{\frac{kd}{p'}}$$

for all  $x \in \mathbb{R}^d$  and  $k \in \mathbb{N}_0$ .  $\square$

*Remark 3.6.* In fact, (3.15) allows to derive pointwise estimates for  $e^{-z\mathcal{L}_{a,\alpha}}$  using Davies' method, cf. [38, Theorem A] and Bui and D'Ancona [11, Proposition 3.4] when  $|\arg(z)| < \pi/4$ .

We now use Theorem 2.1 to extend dyadic Davies–Gaffney estimates (3.13) for  $e^{-tH_\alpha}$  and  $t > 0$  to complex times  $z \in \mathbb{C}_+$ .

**Theorem 3.7.** *Let  $\alpha > 0$ ,  $\beta > 0$ ,  $1 \leq p \leq 2 \leq q \leq \infty$ , and  $\sigma, t > 0$ . Suppose  $e^{-tH_\alpha}$  satisfies the dyadic  $(p, q, \sigma)$  Davies–Gaffney estimate (3.13) (Definition 3.3) with  $r \equiv r_t := t^{1/\alpha}$ , i.e., there is a constant  $C_{\text{DG}} > 0$  such that*

$$\|\mathbf{1}_{B_x(r_t)} e^{-tH_\alpha} \mathbf{1}_{A_2(x,r_t,k)}\|_{p \rightarrow q} \lesssim_\beta C_{\text{DG}} r_t^{-d(\frac{1}{p}-\frac{1}{q})} 2^{-k(\beta-\frac{d}{\sigma})}, \quad x \in \mathbb{R}^d, k \in \mathbb{N}_0. \quad (3.16)$$

*In case  $p \in [1, 2)$  and  $q \in (2, \infty]$ , and  $q \neq p'$ , assume additionally the bounds  $\max\{\|e^{-tH_\alpha}\|_{p \rightarrow p}, \|e^{-tH_\alpha}\|_{q \rightarrow q}\} \lesssim_{d,p,q,\alpha,\beta,\sigma} 1$ . Then, for  $z = |z|e^{i\theta} \in \mathbb{C}_+$ ,  $\zeta \geq 0$ ,  $r_z := |z|^{1/\alpha}(\cos\theta)^{-\zeta}$ ,  $\varepsilon \in (0, 1)$ , and*

$$\tilde{\beta} \equiv \tilde{\beta}_{d,\beta,\sigma,\varepsilon}(\theta) := \left(\beta - \frac{d}{\sigma}\right) \cdot \left(1 - \frac{|\theta|}{\varepsilon|\theta| + (1-\varepsilon)\pi/2}\right) \geq 0, \quad (3.17)$$

one has

$$\begin{aligned} & \|\mathbf{1}_{B_x(r_z)} e^{-zH_\alpha} \mathbf{1}_{A_2(x,r_z,k)}\|_{p \rightarrow q} \\ & \lesssim_{d,\alpha,\beta,\sigma,p,q,\varepsilon} C_{\text{DG}} (|z| \cos\theta)^{-\frac{d}{\alpha}(\frac{1}{p}-\frac{1}{q})} \cdot (\cos\theta)^{-\frac{d\zeta}{\alpha}} \cdot 2^{-k\tilde{\beta}} \\ & = C_{\text{DG}} r_z^{-d(\frac{1}{p}-\frac{1}{q})} (\cos\theta)^{-d(\zeta+\frac{1}{\alpha})(\frac{1}{p}-\frac{1}{q})-\frac{d\zeta}{\alpha}} \cdot 2^{-k\tilde{\beta}} \end{aligned} \quad (3.18)$$

for all  $x \in \mathbb{R}^d$  and  $k \in \mathbb{N}_0$ . Moreover,

$$\|e^{-zH_\alpha}\|_{p \rightarrow q} \lesssim_{d,\alpha,p,q,\beta,\sigma} C_{\text{DG}} (|z| \cos\theta)^{-\frac{d}{\alpha}(\frac{1}{p}-\frac{1}{q})}. \quad (3.19)$$

*Remarks 3.8.* (1) Example 3.5 and (1.5) indicate that one will have  $\beta = d + \alpha$  in many scenarios. Nevertheless, we prefer to keep  $\beta$  as a free parameter here and in the following to illustrate that certain estimates for  $e^{-tH_\alpha}$  can be extended to complex times under less severe decay conditions.

(2) We make some remarks on the choices  $r_t = t^{\frac{1}{\alpha}}$  and  $r_z = |z|^{\frac{1}{\alpha}}(\cos\theta)^{-\zeta}$ . The power  $\alpha^{-1}$  reflects the scaling relation between time and space in  $e^{-zH_\alpha}$  and is dictated by the order of the principal symbol of  $(-\Delta)^{\frac{\alpha}{2}} + \min\{V, 0\}$ . For  $t > 0$  this is seen in (1.3) for  $V = 0$ , in (3.1) for  $\alpha \in (0, 2)$  and  $V \geq 0$ , in (3.15) for  $V = a|x|^{-\alpha}$  with  $a \geq a_*$ , and in Huang et al [32, Theorem 1.3] when  $V \in K_\alpha(\mathbb{R}^d)$  is a perturbation. For complex times the power is expected to be  $\alpha^{-1}$ , too. This is confirmed by Corollaries 3.1 and 3.2 for  $V \geq 0$ , (1.5) for  $V \in K_\alpha(\mathbb{R}^d)$ , and estimates for  $e^{-zH_\alpha}$  when  $\alpha \in 2\mathbb{N}$ , see (3.20) below. On the other hand, a natural choice for  $\zeta$  is not obvious due to the complicated relation between  $\theta$  and  $|z|$  in (3.18). For  $\alpha \in 2\mathbb{N}$ ,  $V \geq 0$ ,  $x, y \in \mathbb{R}^d$ ,  $r, s > 0$ , and  $|x - y| > r + s$ , Davies [21, Theorem 10] showed

$$\|\mathbf{1}_{B_x(r)} e^{-|z|e^{i\theta}H_\alpha} \mathbf{1}_{B_y(s)}\|_{2 \rightarrow 2} \leq \exp\left(-c \left(\frac{d(B_x(r), B_y(s))}{|z|^{\frac{1}{\alpha}}(\cos\theta)^{-\frac{\alpha-1}{\alpha}}}\right)^{\frac{\alpha}{\alpha-1}}\right) \quad (3.20)$$

with similar estimates being available for more general (even singular)  $V$ , cf. [8, pp. 154-156]. This shows that  $r_z = |z|^{1/\alpha}(\cos \theta)^{-\zeta}$  with  $\zeta \equiv (\alpha - 1)/\alpha > 0$  is a natural choice when  $\alpha \in 2\mathbb{N}$ . In fact,  $\zeta > 0$  is necessary for (3.20) to hold on all of  $\mathbb{C}_+$ , which can be seen by fixing  $x \neq y$  and letting  $|\theta| \rightarrow \pi/2$ .

(3) Estimate (3.20) and many related variants, such as [12, Proposition 4.1] by Carron et al and [8, Theorem 2.1] by Blunck, were proved using Phragmén–Lindelöf principles, see Davies [21, Lemma 9] for the original version. The presence of exponential bounds for  $\alpha \in 2\mathbb{N}$  and  $t > 0$  is essential for the derivation of the clean dependence of (3.20) on  $\cos \theta$ .

The proof of Theorem 3.7 is inspired by Blunck [8, Theorem 2.1] and uses two consequences (Propositions 4.3 and 4.5) of the dyadic Davies–Gaffney estimates that are contained in Section 4.

*Proof of Theorem 3.7.* For  $\mu = 1/2 - 1/q$  and  $\nu = 1/p - 1/2$  we have

$$r_t^{d\mu} \|e^{-tH_\alpha}\|_{2 \rightarrow q} \lesssim C_{\text{DG}}^{\mu/(\mu+\nu)} \quad \text{and} \quad r_t^{d\nu} \|e^{-tH_\alpha}\|_{p \rightarrow 2} \lesssim C_{\text{DG}}^{\nu/(\mu+\nu)}, \quad t > 0 \quad (3.21)$$

by Proposition 4.3, whenever  $p = 2$  and  $q \in [2, \infty)$ , or  $p \in [1, 2]$  and  $q = 2$ , or  $q = p'$ . In all other cases (3.21) follows from Riesz–Thorin interpolation between (4.3) in Proposition 4.3 and the  $L^p \rightarrow L^p$  and  $L^q \rightarrow L^q$  boundedness of  $e^{-tH_\alpha}$ . Thus, we obtain for  $z = |z|e^{i\theta} \in \mathbb{C}_+$  and  $t \in (0, \text{Re}(z)/2)$ ,

$$\|e^{-zH_\alpha}\|_{p \rightarrow q} \leq \|e^{-tH_\alpha}\|_{2 \rightarrow q} \|e^{-(z-2t)H_\alpha}\|_{2 \rightarrow 2} \|e^{-tH_\alpha}\|_{p \rightarrow 2} \lesssim C_{\text{DG}} r_t^{-d(\frac{1}{p} - \frac{1}{q})}. \quad (3.22)$$

Combining (3.22) with  $\lim_{t \nearrow \text{Re } z/2} r_t^{-1} = 2^{1/\alpha}(|z| \cos \theta)^{-\frac{1}{\alpha}}$  yields

$$\|e^{-zH_\alpha}\|_{p \rightarrow q} \lesssim C_{\text{DG}} (|z| \cos \theta)^{-\frac{d}{\alpha}(\frac{1}{p} - \frac{1}{q})}, \quad (3.23)$$

which proves (3.19). Moreover, for any ball  $B \equiv B_{x_0}(r_0)$  and its  $k$ -th dyadic annulus  $A_2^{(k)} \equiv A_2(x_0, r_0, k)$  centered around  $B$  (with  $k \in \mathbb{N}_0$  and the convention  $A_2^{(0)} = B$ ), we obtain

$$\|\mathbf{1}_B e^{-zH_\alpha} \mathbf{1}_{A_2^{(k)}}\|_{p \rightarrow q} \leq c_{d,\alpha,\beta,p,q,\sigma} C_{\text{DG}} (|z| \cos \theta)^{-\frac{d}{\alpha}(\frac{1}{p} - \frac{1}{q})}. \quad (3.24)$$

On the other hand, (4.5) in Proposition 4.5 implies for  $\theta = 0$ ,

$$\begin{aligned} \|\mathbf{1}_B e^{-tH_\alpha} \mathbf{1}_{A_2^{(k)}}\|_{p \rightarrow q} &\leq c_{d,\alpha,\beta,p,q,\sigma} C_{\text{DG}} t^{-\frac{d}{\alpha}(\frac{1}{p} - \frac{1}{q})} \cdot \left(1 + \frac{|B|^{\frac{\alpha}{d}}}{t}\right)^{\frac{d}{\alpha q}} \\ &\quad \times \left[ \left(\frac{d(B, A_2^{(k)})}{r_t}\right)^{-\beta} \cdot \left(\frac{|A_2^{(k)}|}{r_t^d}\right)^{\frac{1}{\sigma}} \theta(k-2) + \theta(1-k) \right]. \end{aligned} \quad (3.25)$$

Define the analytic function  $F : \mathbb{C}_+ \rightarrow \mathcal{B}(L^p \rightarrow L^q)$  by

$$F(z) := (c_{d,\alpha,\beta,p,q,\sigma} C_{\text{DG}})^{-1} \left(1 + \frac{|B|^{\frac{\alpha}{d}}}{z}\right)^{-\frac{d}{\alpha q}} \mathbf{1}_B e^{-zH_\alpha} \mathbf{1}_{A_2^{(k)}}. \quad (3.26)$$

By (3.24) and (3.25) we have (using  $|1 + |B|^{\alpha/d}/z| \geq 1$  for  $z \in \mathbb{C}_+$ )

$$\|F(|z|e^{i\theta})\|_{p \rightarrow q} \leq (|z| \cos \theta)^{-\frac{d}{\alpha}(\frac{1}{p} - \frac{1}{q})}, \quad (3.27)$$

$$\|F(|z|)\|_{p \rightarrow q} \leq |z|^{-\frac{d}{\alpha}(\frac{1}{p}-\frac{1}{q})} \left[ \left( \frac{d(B, A_2^{(k)})}{|z|^{1/\alpha}} \right)^{-\beta} \cdot \left( \frac{|A_2^{(k)}|^{d/\alpha}}{|z|} \right)^{\frac{d}{\alpha\sigma}} \theta(k-2) + \theta(1-k) \right]. \quad (3.28)$$

We now apply Theorem 2.1 with  $a_1 = 1$ ,  $\beta_1 = d(1/p-1/q)/\alpha$ , and  $a_2 = d(B, A_2^{(k)})^\alpha$ ,  $\beta_2 = \beta/\alpha$ ,  $a_3 = |A_2^{(k)}|^{d/\alpha}$ ,  $\beta_3 = \frac{d}{\alpha\sigma}$  if  $k \in \mathbb{N} \setminus \{1\}$ , and  $a_2 = a_3 = 1$  and  $\beta_2 = \beta_3 = 0$  if  $k \in \{0, 1\}$ . Abbreviating  $\gamma_\varepsilon = \varepsilon|\theta| + (1-\varepsilon)\pi/2$ , we obtain

$$\begin{aligned} \|F(|z|e^{i\theta})\|_{p \rightarrow q} &\lesssim_\varepsilon (|z| \cos \theta)^{-\frac{d}{\alpha}(\frac{1}{p}-\frac{1}{q})} \\ &\times \left\{ \theta(1-k) + \theta(k-2) \left[ 1 \wedge \left( \frac{d(B, A_2^{(k)})^\alpha}{|z|} \right)^{-\frac{\beta}{\alpha}} \cdot \left( \frac{|A_2^{(k)}|^{d/\alpha}}{|z|} \right)^{\frac{d}{\alpha\sigma}} \right]^{1-\frac{|\theta|}{\gamma_\varepsilon}} \right\} \end{aligned} \quad (3.29)$$

for all  $|z| > 0$  and  $|\theta| < \pi/2$ . By the definition (3.26) and

$$\left| 1 + \frac{|B|^{\alpha/d}}{z} \right|^{\frac{d}{\alpha q}} \lesssim \left( 1 + \frac{|B|}{|z|^{d/\alpha}} \right)^{1/q},$$

this implies

$$\begin{aligned} \|\mathbf{1}_B e^{-zH_\alpha} \mathbf{1}_{A_2^{(k)}}\|_{p \rightarrow q} &\lesssim C_{\text{DG}} (|z| \cos \theta)^{-\frac{d}{\alpha}(\frac{1}{p}-\frac{1}{q})} \cdot \left( 1 + \frac{|B|}{|z|^{d/\alpha}} \right)^{\frac{1}{q}} \\ &\times \left\{ \theta(1-k) + \theta(k-2) \left[ 1 \wedge \left( \frac{d(B, A_2^{(k)})^\alpha}{|z|} \right)^{-\frac{\beta}{\alpha}} \cdot \left( \frac{|A_2^{(k)}|^{d/\alpha}}{|z|} \right)^{\frac{d}{\alpha\sigma}} \right]^{1-\frac{|\theta|}{\gamma_\varepsilon}} \right\}. \end{aligned}$$

Choosing  $B \equiv B_x(r_z)$  and  $A_2^{(k)} \equiv A_2(x, r_z, k)$ , and recalling  $r_z = |z|^{1/\alpha}(\cos \theta)^{-\zeta}$  and  $r_{|z|} = |z|^{1/\alpha}$  yields

$$\begin{aligned} \|\mathbf{1}_{B_x(r_z)} e^{-zH_\alpha} \mathbf{1}_{A_2(x, r_z, k)}\|_{p \rightarrow q} &\lesssim C_{\text{DG}} (|z| \cos \theta)^{-\frac{d}{\alpha}(\frac{1}{p}-\frac{1}{q})} \cdot \left( 1 + \frac{r_z}{r_{|z|}} \right)^{\frac{d}{q}} \\ &\times \left\{ \theta(1-k) + \theta(k-2) \left( \frac{2^k r_z}{r_{|z|}} \right)^{-(\beta-\frac{d}{\sigma}) \cdot (1-\frac{|\theta|}{\gamma_\varepsilon})} \right\} \quad (3.30) \\ &\lesssim C_{\text{DG}} (|z| \cos \theta)^{-\frac{d}{\alpha}(\frac{1}{p}-\frac{1}{q})} \cdot (\cos \theta)^{-\frac{d\zeta}{q}} \cdot 2^{-k(\beta-\frac{d}{\sigma})(1-\frac{|\theta|}{\varepsilon|\theta|+(1-\varepsilon)\pi/2})} \end{aligned}$$

for all  $z \in \mathbb{C}_+$ . Here we used  $\beta \geq d/\sigma$  and  $r_z \geq r_{|z|}$  since  $\zeta \geq 0$ . This shows (3.18) and concludes the proof of Theorem 3.7.  $\square$

For  $p \in [1, 2]$  Theorem 3.7 can be used to extend dyadic  $(2, p', \sigma)$  and  $(p, 2, \tilde{\sigma})$  Davies–Gaffney estimates for  $e^{-tH_\alpha}$  to complex times. In Proposition 4.6 we show that for  $t > 0$ ,  $\sigma = 2$ , and  $\tilde{\sigma} = p'$  they can be inferred from  $(p, p', p')$  estimates under a slightly stronger decay condition.

**Definition 3.9.** Let  $r > 0$ ,  $(T_r)_{r>0}$  be a family of linear bounded operators on  $L^2(\mathbb{R}^d)$ ,  $1 \leq p \leq q \leq \infty$ ,  $\beta, \sigma > 0$ , and  $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  satisfy  $g(\lambda) \sim_\beta (1+\lambda)^{-\beta}$ . Then  $T_r$  is said to satisfy the

- (1) *restricted dyadic*  $(p, q, \sigma)$  *Davies–Gaffney estimate* if there is a finite constant  $C_{\text{DG}} = C_{\text{DG}}(d, p, q, \beta, \sigma) > 0$  such that (3.13) holds, and if  $\beta > d(1/p + 1/\sigma)$ .
- (2) *dual dyadic*  $(p, p', p')$  *Davies–Gaffney estimate* if  $p \in [1, 2]$ , if there is a finite constant  $C_{\text{DG}} = C_{\text{DG}}(d, p, \beta, \sigma) > 0$  such that (3.13) holds, and if  $\beta > d(1/2 + 1/p')$ .

*Remarks 3.10.* (1) If  $p \in [1, 2]$ , then the restricted dyadic  $(p, p', p')$  estimate implies the dual dyadic  $(p, p', p')$  and the dyadic  $(p, p', p')$  Davies–Gaffney estimate.

(2) In Proposition 4.6 we show that dual dyadic  $(p, p', p')$  estimates imply dyadic  $(2, p', 2)$  and  $(p, 2, p')$  estimates, whereas restricted dyadic  $(p, p', p')$  estimates imply restricted dyadic  $(2, p', 2)$  and  $(p, 2, p')$  Davies–Gaffney estimates.

(3) The notions of dyadic  $(p, p, \sigma)$  and restricted dyadic  $(p, p, \sigma)$  Davies–Gaffney estimates coincide.

(4) The semigroup  $e^{-t\mathcal{L}^{a, \alpha}}$  in Example 3.5 satisfies the restricted dyadic  $(p, p', p')$  estimates whenever  $a \in [-a_*, 0)$  and  $p \in (d/(d - \delta), 2]$ . Moreover, if  $V \geq 0$ , then  $e^{-tH_\alpha}$  satisfies the restricted dyadic  $(p, p', p')$  Davies–Gaffney estimates for all  $p \in [1, 2]$ .

The following corollary shows that dual dyadic  $(p, p', p')$  Davies–Gaffney estimates can be used to derive complex-time  $(2, p', 2)$  and  $(p, 2, p')$  estimates.

**Corollary 3.11.** *Let  $\alpha > 0$ ,  $\beta > 0$ ,  $1 \leq p \leq 2$ , and  $t > 0$ . Suppose  $e^{-tH_\alpha}$  satisfies the dual dyadic  $(p, p', p')$  Davies–Gaffney estimate (Definition 3.9) with  $r \equiv r_t := t^{1/\alpha}$  and  $g(\lambda) \sim_\beta (1 + \lambda)^{-\beta}$ . Then, for  $z = |z|e^{i\theta} \in \mathbb{C}_+$ ,  $\zeta \geq 0$ ,  $r_z := |z|^{1/\alpha}(\cos \theta)^{-\zeta}$ ,  $\varepsilon \in (0, 1)$ , and*

$$\begin{aligned} \tilde{\beta}^{(1)} &\equiv \tilde{\beta}_{d, \beta, \varepsilon}^{(1)}(\theta) := \left( \beta - \frac{d}{2} \right) \left( 1 - \frac{|\theta|}{\varepsilon|\theta| + (1 - \varepsilon)\pi/2} \right) \geq 0, \\ \tilde{\beta}^{(2)} &\equiv \tilde{\beta}_{d, \beta, p, \varepsilon}^{(2)}(\theta) := \left( \beta - \frac{d}{p'} \right) \left( 1 - \frac{|\theta|}{\varepsilon|\theta| + (1 - \varepsilon)\pi/2} \right) \geq 0, \end{aligned} \quad (3.31)$$

one has

$$\begin{aligned} &\| \mathbf{1}_{B_x(r_z)} e^{-zH_\alpha} \mathbf{1}_{A_2(x, r_z, k)} \|_{2 \rightarrow p'} \\ &\lesssim_{d, \alpha, p, \beta, \varepsilon} C_{\text{DG}} (|z| \cos \theta)^{-\frac{d}{\alpha} \left( \frac{1}{2} - \frac{1}{p'} \right)} (\cos \theta)^{-\frac{d\zeta}{p'}} \cdot 2^{-k\tilde{\beta}^{(1)}} \\ &= C_{\text{DG}} r_z^{-d \left( \frac{1}{2} - \frac{1}{p'} \right)} (\cos \theta)^{-d \left( \zeta + \frac{1}{\alpha} \right) \left( \frac{1}{2} - \frac{1}{p'} \right) - \frac{d\zeta}{p'}} \cdot 2^{-k\tilde{\beta}^{(1)}} \end{aligned} \quad (3.32)$$

and

$$\begin{aligned} &\| \mathbf{1}_{B_x(r_z)} e^{-zH_\alpha} \mathbf{1}_{A_2(x, r_z, k)} \|_{p \rightarrow 2} \\ &\lesssim_{d, \alpha, p, \beta, \varepsilon} C_{\text{DG}} (|z| \cos \theta)^{-\frac{d}{\alpha} \left( \frac{1}{2} - \frac{1}{p'} \right)} (\cos \theta)^{-\frac{d\zeta}{2}} \cdot 2^{-k\tilde{\beta}^{(2)}} \\ &= C_{\text{DG}} r_z^{-d \left( \frac{1}{2} - \frac{1}{p'} \right)} (\cos \theta)^{-d \left( \zeta + \frac{1}{\alpha} \right) \left( \frac{1}{2} - \frac{1}{p'} \right) - \frac{d\zeta}{2}} \cdot 2^{-k\tilde{\beta}^{(2)}} \end{aligned} \quad (3.33)$$

for all  $x \in \mathbb{R}^d$  and  $k \in \mathbb{N}_0$ . Moreover,

$$\| e^{-zH_\alpha} \|_{2 \rightarrow p'} = \| e^{-\bar{z}H_\alpha} \|_{p \rightarrow 2} \lesssim_{d, \alpha, p, \beta} C_{\text{DG}} (|z| \cos \theta)^{-\frac{d}{\alpha} \left( \frac{1}{p} - \frac{1}{2} \right)}. \quad (3.34)$$

*Proof.* By Proposition 4.6 the dual dyadic  $(p, p', p')$  Davies–Gaffney estimate (3.13) implies the dyadic  $(2, p', 2)$  and  $(p, 2, p')$  Davies–Gaffney estimates. Thus, assumption (3.16) in Theorem 3.7 with  $(p, q, \sigma) = (2, p', 2)$  or  $(p, q, \sigma) = (p, 2, p')$  there is satisfied. The proof is concluded by an application of Theorem 3.7.  $\square$

While the  $L^p \rightarrow L^q$  estimates (3.19) and  $L^2 \rightarrow L^{p'}$  and  $L^p \rightarrow L^2$  estimates (3.34) could be proved rather directly, one could have also obtained them by combining Proposition 4.3 with (3.18), (3.32), and (3.33). However, this argument requires a smallness assumption on  $|\theta|$  and produces another non-positive power of  $\cos \theta$ . On the other hand, it seems difficult to extend estimates for  $\|e^{-tH_\alpha}\|_{p \rightarrow p}$  (cf. Corollary 4.8) to complex times without any restrictions on  $|\theta|$  or  $V$ .

**Corollary 3.12.** *Let  $\alpha > 0$ ,  $\beta > 0$ ,  $1 \leq p \leq 2$ , and  $t > 0$ .*

(1) *Suppose  $e^{-tH_\alpha}$  satisfies the dyadic  $(p, 2, p')$  Davies–Gaffney estimate (Definition 3.3) or the dual dyadic  $(p, p', p')$  Davies–Gaffney estimate (Definition 3.9) with  $r \equiv r_t := t^{1/\alpha}$  and  $g(\lambda) \sim_\beta (1 + \lambda)^{-\beta}$ . Then, for  $z = |z|e^{i\theta} \in \mathbb{C}_+$ ,  $\zeta \geq 0$ ,  $r_z := |z|^{1/\alpha}(\cos \theta)^{-\zeta}$ ,  $\varepsilon \in (0, 1)$ , and*

$$\tilde{\beta}^{(2)} \equiv \tilde{\beta}_{d,\beta,p,\varepsilon}^{(2)}(\theta) := \left( \beta - \frac{d}{p'} \right) \left( 1 - \frac{|\theta|}{\varepsilon|\theta| + (1-\varepsilon)\pi/2} \right) \geq 0,$$

one has

$$\|\mathbf{1}_{B_x(r_z)} e^{-zH_\alpha} \mathbf{1}_{A_2(x,r_z,k)} f\|_{p \rightarrow p} \lesssim_{d,\alpha,p,\beta,\varepsilon} C_{\text{DG}} (\cos \theta)^{-d(\zeta + \frac{1}{\alpha})(\frac{1}{2} - \frac{1}{p'}) - \frac{d\zeta}{2}} 2^{-k\tilde{\beta}^{(2)}} \quad (3.35)$$

for all  $x \in \mathbb{R}^d$  and  $k \in \mathbb{N}_0$ .

(2) *If  $e^{-tH_\alpha}$  satisfies the restricted dyadic  $(p, 2, p')$  or  $(p, p', p')$  Davies–Gaffney estimate (Definition 3.9) and  $|\theta| < \pi/2$  satisfies  $\tilde{\beta}^{(2)} > d/p$ , i.e.,*

$$\frac{|\theta|}{\varepsilon|\theta| + (1-\varepsilon)\pi/2} < 1 - \frac{d}{p} \left( \beta - \frac{d}{p'} \right)^{-1}, \quad (3.36)$$

then

$$\|e^{-zH_\alpha}\|_{p \rightarrow p} = \|e^{-\bar{z}H_\alpha}\|_{p' \rightarrow p'} \lesssim_{d,\alpha,p,\beta,\varepsilon} C_{\text{DG}} (\cos \theta)^{-d(\zeta + \frac{1}{\alpha})(\frac{1}{2} - \frac{1}{p'}) - \frac{d\zeta}{2}}. \quad (3.37)$$

(3) *Suppose  $V \in K_\alpha(\mathbb{R}^d)$  and  $\mu_{\varepsilon,V,d,\alpha} > 0$  is the constant appearing in (1.5). Then for all  $z \in \mathbb{C}_+$  and  $0 < \varepsilon \ll 1$ ,*

$$\|e^{-zH_\alpha}\|_{p \rightarrow p} \lesssim_{d,\alpha,p} e^{\mu_{\varepsilon,V,d,\alpha}|z|} (\cos \theta)^{-d(\frac{1}{\alpha} - \frac{1}{2})\mathbf{1}_{\{\alpha < 1\}} - (\frac{d}{2} + \alpha - 1)\mathbf{1}_{\{\alpha \geq 1\}}}. \quad (3.38)$$

*Remarks 3.13.* (1) The power of  $\cos \theta$  in (3.37) could not be correct if the estimate held for all  $z \in \mathbb{C}_+$ . This can be seen by considering  $H_\alpha = -\Delta$ , since  $\|e^{z\Delta}\|_{1 \rightarrow 1} \lesssim (\cos \theta)^{-\frac{d}{2}}$  and  $\|e^{z\Delta}\|_{2 \rightarrow 2} \leq 1$  imply  $\|e^{z\Delta}\|_{p \rightarrow p} \lesssim (\cos \theta)^{-d(\frac{1}{2} - \frac{1}{p'})}$  for all  $z \in \mathbb{C}_+$ . Moreover, this upper bound is sharp as there is a matching lower bound, cf. Arendt et al [1, Lemma 2.2].

(2) Since  $\beta > d$  in the case of restricted dyadic  $(p, 2, p')$  or  $(p, p', p')$  Davies–Gaffney estimates, there exist  $|\theta| \in [0, \pi/2)$  for which  $\tilde{\beta}^{(2)} > d/p$  is satisfied.

*Proof.* To prove (3.35) it suffices to assume that  $e^{-tH_\alpha}$  satisfies the dyadic  $(p, 2, p')$  Davies–Gaffney estimate by Proposition 4.6. By Hölder’s inequality and (3.18) in Theorem 3.7 (which reduces to (3.33) in this case) we have

$$\begin{aligned} \|\mathbf{1}_{B_x(r_z)} e^{-zH_\alpha} \mathbf{1}_{A_2(x,r_z,k)} f\|_p &\leq |B_x(r_z)|^{\frac{2-p}{2p}} \|\mathbf{1}_{B_x(r_z)} e^{-zH_\alpha} \mathbf{1}_{A_2(x,r_z,k)} f\|_2 \\ &\lesssim_{d,\alpha,p,\beta,\varepsilon} C_{\text{DG}} r_z^{d\frac{2-p}{2p} - d(\frac{1}{p} - \frac{1}{2})} (\cos \theta)^{-d(\zeta + \frac{1}{\alpha})(\frac{1}{2} - \frac{1}{p'}) - \frac{d\zeta}{2}} 2^{-k\tilde{\beta}^{(2)}} \|f\|_p \end{aligned}$$

for any  $f \in L^p$ ,  $x \in \mathbb{R}^d$ , and  $k \in \mathbb{N}_0$ . This proves (3.35).

To prove (3.37) it suffices to assume that  $e^{-tH_\alpha}$  satisfies the restricted dyadic  $(p, 2, p')$  Davies–Gaffney estimate by Proposition 4.6. If  $\tilde{\beta}^{(2)} > d/p$ , then (3.35)

shows that  $e^{-zH_\alpha}$  satisfies the (restricted) dyadic  $(p, p, p' \cdot (1 - |\theta|/\gamma_\varepsilon)^{-1})$  Davies–Gaffney estimate with  $c_{d,\alpha,p,\beta,\varepsilon} C_{\text{DG}} (\cos \theta)^{-d(\zeta + \frac{1}{\alpha})(\frac{1}{2} - \frac{1}{p'}) - \frac{d\zeta}{2}}$  instead of  $C_{\text{DG}}$ , and  $\beta$  replaced by  $\beta(1 - |\theta|/\gamma_\varepsilon)$ , where  $\gamma_\varepsilon = \varepsilon|\theta| + (1 - \varepsilon)\pi/2$ . Thus, (3.37) follows from (4.3) in Proposition 4.3. Estimate (3.38) follows from (1.5).  $\square$

#### 4. CONSEQUENCES OF DYADIC DAVIES–GAFFNEY ESTIMATES

We collect some consequences of the dyadic Davies–Gaffney estimate (3.13) (Definitions 3.3 and 3.9). The dyadic partition  $1 = \sum_{k \geq 0} \mathbf{1}_{A_2(x,r,k)}$  for any  $x \in \mathbb{R}^d$  and  $r > 0$ , and the triangle inequality yield the following preliminary estimate.

**Corollary 4.1.** *Let  $r > 0$ ,  $(T_r)_{r>0}$  be a family of linear operators that satisfy the dyadic  $(p, q, \sigma)$  Davies–Gaffney estimate (3.13) (Definition 3.3). Then*

$$\|\mathbf{1}_{B_x(r)} T_r\|_{p \rightarrow q} \lesssim_{d,\beta,\sigma} C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{q})}, \quad x \in \mathbb{R}^d. \quad (4.1)$$

In fact, for (4.1) to hold, one only needs  $\sum_{k \geq 0} g(k) 2^{kd/\sigma} < \infty$ . We will upgrade this estimate in Proposition 4.3 with the help of the ball averages

$$(N_{p,r} f)(x) := r^{-\frac{d}{p}} \|\mathbf{1}_{B_x(r)} f\|_{L^p(\mathbb{R}^d)} \quad \text{and} \quad (N_{p,q,r} f)(x) := r^{-\frac{d}{q}} \|\mathbf{1}_{B_x(r)} f\|_{L^p(\mathbb{R}^d)} \quad (4.2)$$

for  $p, q \in [1, \infty)$ . The following lemma by Blunck and Kunstmann summarizes useful estimates involving these averaging operators.

**Lemma 4.2** ([6, Lemma 3.3]). *Let  $1 \leq p \leq q \leq \infty$  and  $r > 0$ . Then*

- (1)  $(N_{p,r} f)(x) \lesssim_{p,q} (N_{q,r} f)(x)$  for all  $x \in \mathbb{R}^d$ ,
- (2)  $\|f\|_{L^p(\mathbb{R}^d)} \sim_p \|N_{p,r} f\|_{L^p(\mathbb{R}^d)}$ , and
- (3)  $\|N_{p,q,r} f\|_{L^q(\mathbb{R}^d)} \lesssim_{p,q} \|f\|_{L^p(\mathbb{R}^d)}$ .

**Proposition 4.3.** *Let  $r > 0$  and  $(T_r)_{r>0}$  be a family of linear operators that satisfy the dyadic  $(p, q, \sigma)$  Davies–Gaffney estimate (3.13) (Definition 3.3). Then*

$$\|T_r\|_{p \rightarrow q} \lesssim_{d,\beta,\sigma,p,q} C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{q})}. \quad (4.3)$$

If  $q = p'$  (so in particular  $1 \leq p \leq 2$ ) and  $T_r$  is in addition self-adjoint in  $L^2(\mathbb{R}^d)$  and satisfies  $T_{r+s} = T_r T_s$  for all  $r, s > 0$ , then

$$\|T_r\|_{p \rightarrow 2} = \|T_r\|_{2 \rightarrow p'} \lesssim_{d,\beta,\sigma,p} C_{\text{DG}}^{1/2} r^{-d(\frac{1}{2} - \frac{1}{p'})}. \quad (4.4)$$

*Proof.* By  $\|f\|_q \sim \|N_{q,r} f\|_q$ , a dyadic partition of unity, assumption (3.13), and  $\|N_{p,q,r} f\|_q \lesssim \|f\|_p$  for  $1 \leq p \leq q \leq \infty$ , we have

$$\begin{aligned} \|T_r f\|_q &\lesssim_q \|N_{q,r} T_r f\|_q = \left( \int_{\mathbb{R}^d} \left( r^{-d/q} \|\mathbf{1}_{B_x(r)} T_r \sum_{k=0}^{\infty} \mathbf{1}_{A_2(x,r,k)} f\|_q \right)^q dx \right)^{1/q} \\ &\leq C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{q})} \left( \int_{\mathbb{R}^d} \left( \sum_{k=0}^{\infty} g(2^k) 2^{kd(\frac{1}{\sigma} + \frac{1}{q})} (2^k r)^{-\frac{d}{q}} \|\mathbf{1}_{B_x(2^k r)} f\|_p \right)^q dx \right)^{1/q} \\ &\leq C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{q})} \sum_{k=0}^{\infty} g(2^k) 2^{kd(\frac{1}{\sigma} + \frac{1}{q})} \|N_{p,q,2^k r} f\|_q \\ &\lesssim_{p,q} C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{q})} \sum_{k=0}^{\infty} g(2^k) 2^{kd(\frac{1}{\sigma} + \frac{1}{q})} \|f\|_p \lesssim_{d,\sigma,\beta,q} C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{q})} \|f\|_p. \end{aligned}$$

This shows (4.3). Formula (4.4) follows from  $\|T_{2r}\|_{p \rightarrow p'} = \|T_r\|_{p \rightarrow 2}^2 = \|T_r\|_{2 \rightarrow p'}^2$ .  $\square$

We need one other consequence of (3.13) in the proof of Theorem 3.7. To that end we record the following geometric observations.

**Lemma 4.4.** *Let  $x, z \in \mathbb{R}^d$ ,  $r, r_0 > 0$ , and  $|x - z| \leq r + r_0$ .*

- (1) *If  $j, k \in \mathbb{N} \setminus \{1\}$  and  $A_2(z, r, j) \cap A_2(x, r_0, k) \neq \emptyset$ , then  $2^{k-3}r_0 \leq 2^j r \leq 2^{k+3}r_0$ .*
- (2) *If  $j \in \mathbb{N}$  and  $A_2(z, r, j) \cap B_x(r_0) \neq \emptyset$ , then  $(2^{j-1} - 1)r \leq 2r_0$ , i.e.,  $j \leq 1 + \log_2(1 + 2r_0/r)$ .*

*Proof.* By a translation we can assume  $z = 0$  without loss of generality. In that case  $|x| \leq r + r_0$ . Let  $y \in \mathbb{R}^d$  denote those points belonging to the intersection of the geometric objects in claims (1) and (2). We use that, if  $a \leq b$  and  $c \leq d$  and, if  $[a, b] \cap [c, d] \neq \emptyset$ , then either  $a \leq c \leq b \leq d$ , or  $c \leq a \leq d \leq b$ , or  $a \leq c \leq d \leq b$ , or  $c \leq a \leq b \leq d$ . If in addition  $a = 0$  and  $b, c, d \geq 0$ , then  $c \leq b$ .

- (1) If the annuli intersect, then  $|y| \in [2^{j-1}r, 2^j r]$  and  $|x - y| \in [2^{k-1}r_0, 2^k r_0]$ . By the triangle inequality and the bounds on  $|x|$  and  $|y|$ , we also have

$$\begin{aligned} |x - y| &\leq |x| + |y| \leq r_0 + (2^j + 1)r \quad \text{and} \\ |x - y| &\geq \max\{|y| - |x|, 0\} \geq \max\{(2^{j-1} - 1)r - r_0, 0\}. \end{aligned}$$

Thus,  $|x - y| \in [2^{k-1}r_0, 2^k r_0] \cap [\max\{(2^{j-1} - 1)r - r_0, 0\}, (2^j + 1)r + r_0]$ .

This shows that, if  $(2^{j-1} - 1)r > r_0$ , then either

- (a)  $2^{k-1}r_0 \leq (2^{j-1} - 1)r - r_0 \leq 2^k r_0 \leq (2^j + 1)r + r_0$ , or
- (b)  $(2^{j-1} - 1)r - r_0 \leq 2^{k-1}r_0 \leq (2^j + 1)r + r_0 \leq 2^k r_0$ , or
- (c)  $2^{k-1}r_0 \leq (2^{j-1} - 1)r - r_0 \leq (2^j + 1)r + r_0 \leq 2^k r_0$ , or
- (d)  $(2^{j-1} - 1)r - r_0 \leq 2^{k-1}r_0 \leq 2^k r_0 \leq (2^j + 1)r + r_0$ .

In particular,  $2^{k-3}r_0 \leq 2^j r \leq 2^{k+3}r_0$ . If instead  $(2^{j-1} - 1)r \leq r_0$ , then  $(2^{k-1} - 1)r_0 \leq (2^j + 1)r$ , so in particular  $2^{k-3}r_0 \leq 2^j r \leq 2^{k+3}r_0$ .

- (2) If  $y \in A_2(0, r, j) \cap B_x(r_0) \neq \emptyset$ , then  $|y| \in [2^{j-1}r, 2^j r]$  and  $|x - y| \leq r_0$ . The triangle inequality and the bounds on  $|x - y|$  and  $|x|$  imply  $|y| = |y - x + x| \leq 2r_0 + r$  and therefore  $2^{j-1}r \leq 2r_0 + r$ .

$\square$

**Proposition 4.5.** *Let  $r > 0$  and  $(T_r)_{r>0}$  be a family of linear operators that satisfy the dyadic  $(p, q, \sigma)$  Davies–Gaffney estimate (3.13) (Definition 3.3). Then*

$$\begin{aligned} &\|\mathbf{1}_{B_x(r_0)} T_r \mathbf{1}_{A_2(x, r_0, k)}\|_{p \rightarrow q} \\ &\lesssim_{d, \sigma, \beta, p, q} C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{q})} \left(1 + \frac{|B_x(r_0)|}{r^d}\right)^{1/q} \\ &\quad \times \left[ \left(\frac{d(B_x(r_0), A_2(x, r_0, k))}{r}\right)^{-\beta} \cdot \left(\frac{|A_2(x, r_0, k)|}{r^d}\right)^{\frac{1}{\sigma}} \theta(k-2) + \theta(1-k) \right] \end{aligned} \quad (4.5)$$

holds for all  $x \in \mathbb{R}^d$ ,  $r_0 > 0$ , and  $k \in \mathbb{N}_0$ . In fact, the factor  $(1 + |B_x(r_0)|/r^d)^{1/q}$  can be removed for  $k \in \{0, 1\}$ , i.e.,

$$\begin{aligned} & \| \mathbf{1}_{B_x(r_0)} T_r \mathbf{1}_{A_2(x, r_0, k)} \|_{p \rightarrow q} \\ & \lesssim_{d, \sigma, \beta, p, q} C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{q})} \left[ \left( \frac{d(B_x(r_0), A_2(x, r_0, k))}{r} \right)^{-\beta} \cdot \left( \frac{|A_2(x, r_0, k)|}{r^d} \right)^{\frac{1}{\sigma}} \right. \\ & \quad \left. \times \left( 1 + \frac{|B_x(r_0)|}{r^d} \right)^{\frac{1}{q}} \theta(k-2) + \theta(1-k) \right] \end{aligned} \quad (4.6)$$

holds for all  $x \in \mathbb{R}^d$ ,  $r_0 > 0$ , and  $k \in \mathbb{N}_0$ .

Recall that (3.13) only involved one radius. Here the radii  $r$  appearing in  $T_r$ , and  $r_0$  appearing in  $B_x(r_0)$  and  $A_2(x, r_0, k)$  in formula (4.5) are independent of each other. Note also that the proof of (4.5) only needs  $\beta > d/\sigma$ .

*Proof.* To prove (4.5) we use  $\|f\|_q \sim \|N_{q,r} f\|_q$ , decompose dyadically, apply (3.13), and obtain

$$\begin{aligned} & \| \mathbf{1}_{B_x(r_0)} T_r \mathbf{1}_{A_2(x, r_0, k)} f \|_q \lesssim_q \| N_{q,r} (\mathbf{1}_{B_x(r_0)} T_r \mathbf{1}_{A_2(x, r_0, k)} f) \|_q \\ & = \left( \int_{\mathbb{R}^d} \left( r^{-\frac{d}{q}} \| \mathbf{1}_{B_z(r)} \mathbf{1}_{B_x(r_0)} T_r \sum_{j=0}^{\infty} \mathbf{1}_{A_2(z, r, j)} \mathbf{1}_{A_2(x, r_0, k)} f \|_q \right)^q dz \right)^{1/q} \\ & \leq C_{\text{DG}} r^{-\frac{d}{p}} \sum_{j=0}^{\infty} g(2^j) 2^{\frac{jd}{\sigma}} \left( \int_{|z-x| \leq r+r_0} \| \mathbf{1}_{A_2(z, r, j)} \mathbf{1}_{A_2(x, r_0, k)} f \|_p^q dz \right)^{1/q}. \end{aligned} \quad (4.7)$$

To estimate  $(\dots)^{1/q}$  on the right side we use Lemma 4.4 (with  $|x-z| \leq r+r_0$ ) and distinguish between the following cases.

- (1) If  $j, k \in \mathbb{N} \setminus \{1\}$ , and  $A_2(z, r, j) \cap A_2(x, r_0, k) \neq \emptyset$ , then  $2^{k-3}r_0 \leq 2^j r \leq 2^{k+3}r_0$  by (1) in Lemma 4.4.
- (2) If  $k \in \mathbb{N} \setminus \{1\}$ ,  $j \in \{0, 1\}$ , and  $A_2(z, r, j) \cap A_2(x, r_0, k) \neq \emptyset$ , then  $B_z(2^j r) \cap A_2(x, r_0, k) \neq \emptyset$  and so  $2^j r \geq 2^{k-3}r_0$  by (2) in Lemma 4.4.
- (3) If  $k \in \{0, 1\}$ , we put no restriction on  $j \in \mathbb{N}_0$ .

Thus,

$$\begin{aligned} & \| \mathbf{1}_{A_2(z, r, j)} \mathbf{1}_{A_2(x, r_0, k)} f \|_p \\ & \lesssim \|f\|_p \left[ \mathbf{1}_{\{2^j r \geq 2^{k-3}r_0\}} \theta(k-2) \left( \mathbf{1}_{\{2^j r \leq 2^{k+3}r_0\}} \theta(j-2) + \theta(1-j) \right) + \theta(1-k) \right]. \end{aligned}$$

If  $k \in \{0, 1\}$ , we simply sum over all  $j \in \mathbb{N}_0$ . If  $k \in \mathbb{N} \setminus \{1\}$  and  $j \in \mathbb{N} \setminus \{1\}$ , we use  $g(2^j) 2^{jd/\sigma} \lesssim_{\beta} 2^{-j(\beta-d/\sigma)}$  and that we are summing over dyadic numbers. If  $k \in \mathbb{N} \setminus \{1\}$  and  $j \in \{0, 1\}$ , we use  $\beta \geq d/\sigma$  and  $2^j r \geq 2^{k-3}r_0$  to estimate



$g(2^j) \cdot 2^{\frac{j d}{\sigma}} \lesssim_{d,\beta,\sigma} (2^k r_0/r)^{-\beta+d/\sigma}$ . Thus, (4.7) can be estimated by

$$\begin{aligned} & \|\mathbf{1}_{B_x(r_0)} T_r \mathbf{1}_{A_2(x,r_0,k)} f\|_q \\ & \lesssim_q C_{\text{DG}} r^{-\frac{d}{p}} \left( \int_{|z-x| \leq r+r_0} dz \right)^{1/q} \|f\|_p \\ & \times \sum_{j=0}^{\infty} \left[ 2^{j(\frac{d}{\sigma}-\beta)} (\mathbf{1}_{\{2^j r \geq 2^{k-3} r_0\}} \theta(k-2) (\mathbf{1}_{\{2^j r \leq 2^{k+3} r_0\}} \theta(j-2) + \theta(1-j)) + \theta(1-k)) \right] \\ & \lesssim_{d,\sigma,\beta,q} C_{\text{DG}} r^{-d(\frac{1}{p}-\frac{1}{q})} \cdot \left(1 + \frac{r_0}{r}\right)^{\frac{d}{q}} \left[ \left(\frac{2^k r_0}{r}\right)^{-\beta+d/\sigma} \theta(k-2) + \theta(1-k) \right] \|f\|_p. \end{aligned}$$

Since  $2^k r_0 \sim d(B_x(r_0), A_2(x, r_0, k)) \sim |A_2(x, r_0, k)|^{\frac{1}{d}}$  for  $k \geq 2$ , this proves (4.5). The improved estimate (4.6) for  $k \in \{0, 1\}$  follows from  $\|\mathbf{1}_{B_x(r_0)} T_r \mathbf{1}_{A_2(x,r_0,k)}\|_{p \rightarrow q} \leq \|T_r\|_{p \rightarrow q}$  and Proposition 4.3. This concludes the proof of Proposition 4.5.  $\square$

The proof of Corollary 3.11 relies on the following proposition which says that  $(p, p', p')$  estimates imply  $(2, p', 2)$  and  $(p, 2, p')$  estimates, if one assumes additionally  $\sum_{k \geq 0} g(2^k) 2^{kd(\frac{1}{p'} + \frac{1}{2})} < \infty$ . This assumption is contained in the notion of dual dyadic Davies–Gaffney estimates.

**Proposition 4.6.** *Let  $p \in [1, 2]$  and  $r > 0$ .*

(1) *If  $(T_r)_{r>0}$  is a family of linear operators that satisfy the dual dyadic  $(p, p', p')$  Davies–Gaffney estimate (3.13) (Definition 3.9), then  $T_r$  satisfies the dyadic  $(2, p', 2)$  Davies–Gaffney estimate (Definition 3.3), i.e.,*

$$\|\mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x,r,k)}\|_{2 \rightarrow p'} \lesssim_{d,\beta,p} C_{\text{DG}} r^{-d(\frac{1}{2}-\frac{1}{p'})} g(2^k) 2^{\frac{k d}{2}}, \quad x \in \mathbb{R}^d, k \in \mathbb{N}_0 \quad (4.8a)$$

and the dyadic  $(p, 2, p')$  Davies–Gaffney estimate, i.e.,

$$\|\mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x,r,k)}\|_{p \rightarrow 2} \lesssim_{d,\beta,p} C_{\text{DG}} r^{-d(\frac{1}{p}-\frac{1}{2})} g(2^k) 2^{\frac{k d}{p'}}, \quad x \in \mathbb{R}^d, k \in \mathbb{N}_0. \quad (4.8b)$$

(2) *If  $(T_r)_{r>0}$  satisfies the restricted dyadic  $(p, p', p')$  Davies–Gaffney estimate (3.13) (Definition 3.9), then it satisfies the restricted dyadic  $(2, p', 2)$  and  $(p, 2, p')$  Davies–Gaffney estimates (4.8a) and (4.8b).*

*Proof.* Since  $\beta > d(1/2 + 1/p')$  for the dual dyadic  $(p, p', p')$  estimates and  $\beta > d$  for the restricted dyadic  $(p, p', p')$  estimates, it suffices to show (4.8a) and (4.8b).

To prove (4.8a) we use  $\|f\|_{p'} \sim_p \|N_{p',r} f\|_{p'}$  and obtain for  $f \in L^2$ ,

$$\begin{aligned} & \|\mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x,r,k)} f\|_{p'} \lesssim_p \|N_{p',r} \mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x,r,k)} f\|_{p'} \\ & = \left( \int_{\mathbb{R}^d} dz \left( r^{-\frac{d}{p'}} \|\mathbf{1}_{B_z(r)} \mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x,r,k)} \sum_{j \geq 0} \mathbf{1}_{A_2(z,r,j)} f\|_{p'} \right)^{p'} \right)^{1/p'} \\ & \leq C_{\text{DG}} r^{-\frac{d}{p}} \sum_{j \geq 0} g(2^j) 2^{\frac{j d}{p'}} \left( \int_{|x-z| \leq 2r} dz \|\mathbf{1}_{A_2(x,r,k)} \mathbf{1}_{A_2(z,r,j)} f\|_p^{p'} \right)^{1/p'}. \end{aligned}$$

We use Lemma 4.4 (with  $|x-z| \leq 2r$ ) and distinguish between the following cases.

- (1) If  $k, j \in \mathbb{N} \setminus \{1\}$  and  $A_2(x, r, k) \cap A_2(z, r, j) \neq \emptyset$ , then  $2^{k-3} \leq 2^j \leq 2^{k+3}$  by (1) in Lemma 4.4.

- (2) If  $k \in \mathbb{N} \setminus \{1\}$ ,  $j \in \{0, 1\}$ , and  $A_2(x, r, k) \cap A_2(z, r, j) \neq \emptyset$ , then  $A_2(x, r, k) \cap B_z(2^j r) \neq \emptyset$  and so  $k \leq 4$  by (2) in Lemma 4.4.
- (3) If  $k \in \{0, 1\}$ ,  $j \in \mathbb{N}_0$ , and  $A_2(x, r, k) \cap A_2(z, r, j) \neq \emptyset$ , then  $B_x(2^k r) \cap A_2(z, r, j) \neq \emptyset$  and so  $j \leq 4$  by (2) in Lemma 4.4.

Using this observation and Hölder's inequality, we can estimate

$$\|\mathbf{1}_{A_2(x, r, k)} \mathbf{1}_{A_2(z, r, j)} f\|_p \lesssim_{d, p, \beta} \mathbf{1}_{\{|j-k| \leq 4\}} (2^k r)^{d(\frac{1}{p} - \frac{1}{2})} \|f\|_2, \quad |x - z| \leq 2r$$

and deduce (using  $g(2^j) \sim_\beta 2^{-j\beta}$ )

$$\begin{aligned} & \|\mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x, r, k)} f\|_{p'} \\ & \lesssim C_{\text{DG}} r^{-\frac{d}{2}} \sum_{j \geq 0} g(2^j) 2^{\frac{jd}{p'}} 2^{kd(\frac{1}{p} - \frac{1}{2})} \mathbf{1}_{\{|j-k| \leq 4\}} \left( \int_{|x-z| \leq 2r} dz \right)^{\frac{1}{p'}} \|f\|_2 \\ & \lesssim_{p, d, \beta} C_{\text{DG}} r^{-d(\frac{1}{2} - \frac{1}{p'})} g(2^k) 2^{kd/2} \|f\|_2. \end{aligned}$$

This concludes the proof of (4.8a).

The proof of (4.8b) is similar but uses also (1) of Lemma 4.2, i.e.,  $(N_{2, r} f)(x) \lesssim_p (N_{p', r} f)(x)$ . By the above reasoning and  $\|\mathbf{1}_{A_2(x, r, k)} \mathbf{1}_{A_2(z, r, j)} f\|_p \leq \mathbf{1}_{\{|j-k| \leq 4\}} \|f\|_p$  for  $|x - z| < 2r$ , we obtain

$$\begin{aligned} & \|\mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x, r, k)} f\|_2 \lesssim \|N_{2, r} \mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x, r, k)} f\|_2 \lesssim_p \|N_{p', r} \mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x, r, k)} f\|_2 \\ & = \left( \int_{\mathbb{R}^d} dz \left( r^{-\frac{d}{p'}} \|\mathbf{1}_{B_z(r)} \mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x, r, k)} \sum_{j \geq 0} \mathbf{1}_{A_2(z, r, j)} f\|_{p'} \right)^2 \right)^{1/2} \\ & \leq C_{\text{DG}} r^{-\frac{d}{p}} \sum_{j \geq 0} g(2^j) 2^{\frac{jd}{p'}} \left( \int_{|x-z| \leq 2r} dz \|\mathbf{1}_{A_2(x, r, k)} \mathbf{1}_{A_2(z, r, j)} f\|_p^2 \right)^{1/2} \\ & \lesssim C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{2})} \|f\|_p \sum_{j \geq 0} g(2^j) 2^{\frac{jd}{p'}} \mathbf{1}_{\{|j-k| \leq 4\}} \lesssim C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{2})} g(2^k) 2^{\frac{kd}{p'}} \|f\|_p. \end{aligned}$$

This concludes the proof of Proposition 4.6.  $\square$

Combining Proposition 4.6 with Proposition 4.3 yields the following variant of (4.4), which does not need that  $T_r$  has the semigroup property.

**Corollary 4.7.** *Let  $p \in [1, 2]$ ,  $r > 0$ , and  $(T_r)_{r>0}$  be a family of linear operators that satisfy the dual dyadic  $(p, p', p')$  Davies–Gaffney estimate (3.13) (Definition 3.9). Then*

$$\|T_r\|_{2 \rightarrow p'} \lesssim_{d, \beta, p} C_{\text{DG}} r^{-d(\frac{1}{2} - \frac{1}{p'})} \quad (4.9a)$$

and

$$\|T_r\|_{p \rightarrow 2} \lesssim_{d, \beta, p} C_{\text{DG}} r^{-d(\frac{1}{p} - \frac{1}{2})} \quad (4.9b)$$

for all  $x \in \mathbb{R}^d$ .

We now show that restricted dyadic  $(p, 2, p')$  and  $(p, p', p')$  estimates imply (restricted) dyadic  $(p, p, p')$  estimates and  $L^p$  boundedness of  $T_r$ .

**Corollary 4.8.** *Let  $p \in [1, 2]$ ,  $r > 0$ , and  $(T_r)_{r>0}$  be a family of linear operators that satisfy the restricted dyadic  $(p, 2, p')$  or  $(p, p', p')$  Davies–Gaffney estimate (3.13) (Definition 3.9). Then  $T_r$  satisfies the restricted dyadic  $(p, p, p')$  Davies–Gaffney estimate*

$$\|\mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x,r,k)} f\|_{p \rightarrow p} \lesssim_{d,\beta,p} C_{\text{DG}} 2^{\frac{kd}{p'}} g(2^k), \quad x \in \mathbb{R}^d, k \in \mathbb{N}_0. \quad (4.10)$$

Moreover,

$$\|T_r\|_{p \rightarrow p} = \|(T_r)^*\|_{p' \rightarrow p'} \lesssim_{d,\beta,p} C_{\text{DG}}. \quad (4.11)$$

*Proof.* By Proposition 4.6 it suffices to assume that  $(T_r)_{r>0}$  satisfies the restricted  $(p, 2, p')$  Davies–Gaffney estimates. By Hölder’s inequality and (4.8b) we have

$$\begin{aligned} \|\mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x,r,k)} f\|_p &\leq |B_x(r)|^{\frac{2-p}{2p}} \|\mathbf{1}_{B_x(r)} T_r \mathbf{1}_{A_2(x,r,k)} f\|_2 \\ &\lesssim_{d,\beta,p} C_{\text{DG}} r^{d\frac{2-p}{2p} - d(\frac{1}{p} - \frac{1}{2})} 2^{\frac{kd}{p'}} g(2^k) \|f\|_p \end{aligned}$$

for any  $f \in L^p$ ,  $x \in \mathbb{R}^d$ , and  $k \in \mathbb{N}_0$ . This proves (4.10). Formula (4.11) follows from (4.10) and Proposition 4.3. This concludes the proof.  $\square$

#### ACKNOWLEDGMENTS

I wish to thank Volker Bach and Jean–Claude Cuenin for valuable discussions.

#### REFERENCES

- [1] Wolfgang Arendt, Omar El-Mennaoui, and Matthias Hieber. Boundary values of holomorphic semigroups. *Proc. Amer. Math. Soc.*, 125(3):635–647, 1997.
- [2] Joseph Bak and Donald J. Newman. *Complex Analysis*. Undergraduate Texts in Mathematics. Springer, New York, third edition, 2010.
- [3] R. M. Blumenthal and R. K. Gettoor. Some theorems on stable processes. *Trans. Amer. Math. Soc.*, 95:263–273, 1960.
- [4] S. Blunck and P. C. Kunstmann. Weighted norm estimates and maximal regularity. *Adv. Differential Equations*, 7(12):1513–1532, 2002.
- [5] S. Blunck and P. C. Kunstmann. Weak type  $(p, p)$  estimates for Riesz transforms. *Math. Z.*, 247(1):137–148, 2004.
- [6] S. Blunck and P. C. Kunstmann. Generalized Gaussian estimates and the Legendre transform. *J. Operator Theory*, 53(2):351–365, 2005.
- [7] Sönke Blunck. A Hörmander-type spectral multiplier theorem for operators without heat kernel. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 2(3):449–459, 2003.
- [8] Sönke Blunck. Generalized Gaussian estimates and Riesz means of Schrödinger groups. *J. Aust. Math. Soc.*, 82(2):149–162, 2007.
- [9] Sönke Blunck and Peer Christian Kunstmann. Calderón-Zygmund theory for non-integral operators and the  $H^\infty$  functional calculus. *Rev. Mat. Iberoamericana*, 19(3):919–942, 2003.
- [10] Krzysztof Bogdan, Tomasz Grzywny, Tomasz Jakubowski, and Dominika Pilarczyk. Fractional Laplacian with Hardy potential. *Communications in Partial Differential Equations*, 44(1):20–50, 2019.
- [11] The Anh Bui and Piero D’Ancona. Generalized Hardy operators. *Preprint*, pages 1–24, April 2021.
- [12] Gilles Carron, Thierry Coulhon, and El-Maati Ouhabaz. Gaussian estimates and  $L^p$ -boundedness of Riesz means. *J. Evol. Equ.*, 2(3):299–317, 2002.
- [13] Peng Chen, El Maati Ouhabaz, Adam Sikora, and Lixin Yan. Restriction estimates, sharp spectral multipliers and endpoint estimates for Bochner-Riesz means. *J. Anal. Math.*, 129:219–283, 2016.
- [14] Peng Chen, Xuan Thinh Duong, Zhijie Fan, Ji Li, and Lixin Yan. The Schrödinger equation in  $L^p$  spaces for operators with heat kernel satisfying Poisson type bounds. *arXiv e-prints*, page arXiv:2007.01469, July 2020.

- [15] Peng Chen, Xuan Thinh Duong, Ji Li, and Lixin Yan. Sharp endpoint  $L^p$  estimates for Schrödinger groups. *arXiv e-prints*, page arXiv:1811.03326, November 2018.
- [16] Thierry Coulhon and Xuan Thinh Duong. Riesz transforms for  $1 \leq p \leq 2$ . *Trans. Amer. Math. Soc.*, 351(3):1151–1169, 1999.
- [17] Thierry Coulhon and Xuan Thinh Duong. Maximal regularity and kernel bounds: observations on a theorem by Hieber and Prüss. *Adv. Differential Equations*, 5(1-3):343–368, 2000.
- [18] Thierry Coulhon and Xuan Thinh Duong. Riesz transforms for  $p > 2$ . *C. R. Acad. Sci. Paris Sér. I Math.*, 332(11):975–980, 2001.
- [19] Thierry Coulhon and Adam Sikora. Gaussian heat kernel upper bounds via the Phragmén-Lindelöf theorem. *Proc. Lond. Math. Soc. (3)*, 96(2):507–544, 2008.
- [20] E. B. Davies. *Heat Kernels and Spectral Theory*, volume 92 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1990.
- [21] E. B. Davies. Uniformly elliptic operators with measurable coefficients. *J. Funct. Anal.*, 132(1):141–169, 1995.
- [22] E. B. Davies and A. M. Hinz. Kato class potentials for higher order elliptic operators. *J. London Math. Soc. (2)*, 58(3):669–678, 1998.
- [23] Xuan T. Duong and Derek W. Robinson. Semigroup kernels, Poisson bounds, and holomorphic functional calculus. *J. Funct. Anal.*, 142(1):89–128, 1996.
- [24] Xuan Thinh Duong, El Maati Ouhabaz, and Adam Sikora. Plancherel-type estimates and sharp spectral multipliers. *J. Funct. Anal.*, 196(2):443–485, 2002.
- [25] Rupert L. Frank, Elliott H. Lieb, and Robert Seiringer. Hardy-Lieb-Thirring inequalities for fractional Schrödinger operators. *J. Amer. Math. Soc.*, 21(4):925–950, 2008.
- [26] Rupert L. Frank, Konstantin Merz, and Heinz Siedentop. Equivalence of Sobolev norms involving generalized Hardy operators. *International Mathematics Research Notices*, 2021(3):2284–2303, February 2021.
- [27] Rupert L. Frank and Robert Seiringer. Non-linear ground state representations and sharp Hardy inequalities. *J. Funct. Anal.*, 255(12):3407–3430, 2008.
- [28] A. Gulisashvili. On the Kato classes of distributions and the BMO-classes. In *Differential equations and control theory (Athens, OH, 2000)*, volume 225 of *Lecture Notes in Pure and Appl. Math.*, pages 159–176. Dekker, New York, 2002.
- [29] Waldemar Hebisch. A multiplier theorem for Schrödinger operators. *Colloq. Math.*, 60/61(2):659–664, 1990.
- [30] Ira W. Herbst. Spectral theory of the operator  $(p^2 + m^2)^{1/2} - Ze^2/r$ . *Comm. Math. Phys.*, 53:285–294, 1977.
- [31] Einar Hille. *Analytic Function Theory. Vol. II*. Introductions to Higher Mathematics. Ginn and Co., Boston, Mass.-New York-Toronto, Ont., 1962.
- [32] Shanlin Huang, Ming Wang, Quan Zheng, and Zhiwen Duan.  $L^p$  estimates for fractional Schrödinger operators with Kato class potentials. *J. Differential Equations*, 265(9):4181–4212, 2018.
- [33] Tianxiao Huang, Shanlin Huang, and Quan Zheng. Inhomogeneous oscillatory integrals and global smoothing effects for dispersive equations. *J. Differential Equations*, 263(12):8606–8629, 2017.
- [34] Tosio Kato. *Perturbation Theory for Linear Operators*. Springer-Verlag, Berlin-New York, second edition, 1976. Grundlehren der Mathematischen Wissenschaften, Band 132.
- [35] V. F. Kovalenko, M. A. Perelmuter, and Ya. A. Semenov. Schrödinger operators with  $L_w^{1/2}(\mathbb{R}^l)$ -potentials. *J. Math. Phys.*, 22:1033–1044, 1981.
- [36] Vitali Liskevich, Zeev Sobol, and Hendrik Vogt. On the  $L_p$ -theory of  $C_0$ -semigroups associated with second-order elliptic operators. II. *J. Funct. Anal.*, 193(1):55–76, 2002.
- [37] Konstantin Merz. On scales of Sobolev spaces associated to generalized Hardy operators. *Math. Z.*, 2021. To appear in print. Electronically published on January 09, 2021. Available at <https://doi.org/10.1007/s00209-020-02651-0>.
- [38] Pierre D. Milman and Yu. A. Semenov. Global heat kernel bounds via desingularizing weights. *J. Funct. Anal.*, 212(2):373–398, 2004.
- [39] Akihiko Miyachi. On some singular Fourier multipliers. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 28(2):267–315, 1981.
- [40] Walter Rudin. *Real and Complex Analysis*. McGraw-Hill Book Co., New York, third edition, 1987.

- [41] Gabriele Schreieck and Jürgen Voigt. Stability of the  $L_p$ -spectrum of Schrödinger operators with form-small negative part of the potential. In *Functional Analysis (Essen, 1991)*, volume 150 of *Lecture Notes in Pure and Appl. Math.*, pages 95–105. Dekker, New York, 1994.
- [42] Adam Sikora, Lixin Yan, and Xiaohua Yao. Sharp spectral multipliers for operators satisfying generalized Gaussian estimates. *J. Funct. Anal.*, 266(1):368–409, 2014.
- [43] Elias M. Stein and Rami Shakarchi. *Complex Analysis*, volume 2 of *Princeton Lectures in Analysis*. Princeton University Press, Princeton, NJ, 2003.
- [44] Stephen Wainger. Special trigonometric series in  $k$ -dimensions. *Mem. Amer. Math. Soc.*, 59:102, 1965.
- [45] D. Yafaev. Sharp constants in the Hardy-Rellich inequalities. *Journ. Functional Analysis*, 168(1):212–144, October 1999.
- [46] Shiliang Zhao and Quan Zheng. Uniform complex time heat kernel estimates without Gaussian bounds. *arXiv e-prints*, page arXiv:2012.08763, December 2020.
- [47] Quan Zheng and Xiaohua Yao. Higher-order Kato class potentials for Schrödinger operators. *Bull. Lond. Math. Soc.*, 41(2):293–301, 2009.

(Konstantin Merz) INSTITUT FÜR ANALYSIS UND ALGEBRA, TECHNISCHE UNIVERSITÄT BRAUNSCHWEIG, UNIVERSITÄTSPLATZ 2, 38106 BRAUNSCHWEIG, GERMANY

*Email address:* `k.merz@tu-bs.de`