

# Large time behavior of global strong solutions to the 2-D compressible FENE dumbbell model

Zhaonan Luo<sup>1,\*</sup>, Wei Luo<sup>1†</sup> and Zhaoyang Yin<sup>1,2‡</sup>

<sup>1</sup>Department of Mathematics, Sun Yat-sen University, Guangzhou 510275, China

<sup>2</sup>Faculty of Information Technology,  
Macau University of Science and Technology, Macau, China

## Abstract

In this paper, we mainly study large time behavior of the strong solutions to the 2-D compressible finite extensible nonlinear elastic (FENE) dumbbell model. The Fourier splitting method yields that the  $L^2$  decay rate is  $\ln^{-l}(e+t)$  for any  $l \in \mathbb{N}$ . By virtue of the time weighted energy estimate, we can improve the decay rate to  $(1+t)^{-\frac{1}{4}}$ . Under the low-frequency condition and by the Littlewood-Paley theory, we show that the solutions belong to some Besov space with negative index and obtain the optimal  $L^2$  decay rate.

*2010 Mathematics Subject Classification:* 35Q30, 76B03, 76D05, 76D99.

*Keywords:* The compressible FENE dumbbell model; Besov spaces; Time decay rate.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Short reviews for the incompressible FENE model . . . . .	3
1.2	Short reviews for the CNS equations . . . . .	3
1.3	Main results . . . . .	4
<b>2</b>	<b>Preliminaries</b>	<b>5</b>
<b>3</b>	<b>The <math>L^2</math> decay rate</b>	<b>7</b>
	<b>References</b>	<b>18</b>

\*email: 1411919168@qq.com

†E-mail: luowei23@mail2.sysu.edu.cn

‡E-mail: mcsyzy@mail.sysu.edu.cn

# 1 Introduction

In this paper we are concerned with the compressible finite extensible nonlinear elastic (FENE) dumbbell model [2, 14]:

$$(1.1) \quad \begin{cases} \varrho_t + \operatorname{div}(\varrho u) = 0, \\ (\varrho u)_t + \operatorname{div}(\varrho u \otimes u) - \operatorname{div}\Sigma(u) + \nabla_x P(\varrho) = \operatorname{div} \tau, \\ \psi_t + u \cdot \nabla \psi = \operatorname{div}_R[-\sigma(u) \cdot R\psi + \nabla_R \psi + \nabla_R \mathcal{U}\psi], \\ \tau_{ij} = \int_B (R_i \nabla_j \mathcal{U}) \psi dR, \\ \varrho|_{t=0} = \varrho_0, \quad u|_{t=0} = u_0, \quad \psi|_{t=0} = \psi_0, \\ (\nabla_R \psi + \nabla_R \mathcal{U}\psi) \cdot n = 0 \quad \text{on} \quad \partial B(0, R_0). \end{cases}$$

In (1.1),  $\varrho(t, x)$  stands for the density of the solvent,  $u(t, x)$  denotes the velocity of the polymeric liquid and  $\psi(t, x, R)$  represents the distribution function for the internal configuration. The notation  $\Sigma(u) = \mu(\nabla u + \nabla^T u) + \mu' \operatorname{div} u \cdot Id$  stands for the stress tensor, with the viscosity coefficients  $\mu$  and  $\mu'$  satisfying  $\mu > 0$  and  $2\mu + \mu' > 0$ . The pressure satisfies  $P(\varrho) = \varrho^\gamma$  with  $\gamma \geq 1$ . Moreover,  $\tau$  stands for an additional stress tensor.  $\sigma(u)$  is the drag term. In the general case,  $\sigma(u) = \nabla u$ .

The compressible FENE dumbbell model describes the system coupling fluids and polymers. A polymer is described as an "elastic dumbbell" consisting of two "beads" joined by a spring that can be modeled by  $R$ . The finite extensibility of the polymers means that  $R$  is bounded in ball  $B = B(0, R_0)$ . Let  $\mathcal{U}(R) = -k \log(1 - (\frac{|R|}{R_0})^2)$  represents the potential with  $k > 0$ . The polymer particles are described by  $\psi(t, x, R)$  satisfies  $\int_B \psi(t, x, R) dR = 1$ . At the level of liquid, the system couples compressible Navier-Stokes (CNS) equations with a Fokker-Planck equation. This is a micro-macro model (For more details, one can refer to [31] and [32]).

In this paper, we take  $R_0 = 1$  and  $x \in \mathbb{R}^2$ . Notice that  $(\varrho, u, \psi)$  with  $\varrho = 1$ ,  $u = 0$  and

$$\psi_\infty(R) = \frac{e^{-\mathcal{U}(R)}}{\int_B e^{-\mathcal{U}(R)} dR} = \frac{(1 - |R|^2)^k}{\int_B (1 - |R|^2)^k dR},$$

is a trivial solution of (1.1). Taking the perturbations near the global equilibrium:

$$\rho = \varrho - 1, \quad u = u, \quad g = \frac{\psi - \psi_\infty}{\psi_\infty},$$

then we can rewrite (1.1) as the following system:

$$(1.2) \quad \begin{cases} \rho_t + \operatorname{div} u(1 + \rho) = -u \cdot \nabla \rho, \\ u_t - \frac{1}{1+\rho} \operatorname{div}\Sigma(u) + \frac{P'(1+\rho)}{1+\rho} \nabla \rho = -u \cdot \nabla u + \frac{1}{1+\rho} \operatorname{div} \tau, \\ g_t + \mathcal{L}g = -u \cdot \nabla g - \frac{1}{\psi_\infty} \nabla_R \cdot (\nabla u R g \psi_\infty) - \operatorname{div} u - \nabla u R \nabla_R \mathcal{U}, \\ \tau_{ij}(g) = \int_B (R_i \nabla_j \mathcal{U}) g \psi_\infty dR, \\ \rho|_{t=0} = \rho_0, \quad u|_{t=0} = u_0, \quad g|_{t=0} = g_0, \\ \psi_\infty \nabla_R g \cdot n = 0 \quad \text{on} \quad \partial B(0, 1), \end{cases}$$

where  $\mathcal{L}g = -\frac{1}{\psi_\infty} \nabla_R \cdot (\psi_\infty \nabla_R g)$ .

**Remark.** As in the reference [32], one can deduce that  $\psi = 0$  on  $\partial B(0, 1)$ .

### 1.1. Short reviews for the incompressible FENE model

Let us review some mathematical results about the incompressible FENE model. At first, M. Renardy [38] established the local well-posedness in Sobolev spaces with potential  $\mathcal{U}(R) = (1 - |R|^2)^{1-\sigma}$  for  $\sigma > 1$ . Later, B. Jourdain, T. Lelièvre, and C. Le Bris [22] proved local existence of a stochastic differential equation with potential  $\mathcal{U}(R) = -k \log(1 - |R|^2)$  in the case  $k > 3$  for a Couette flow. H. Zhang and P. Zhang [47] proved local well-posedness of (1.4) with  $d = 3$  in weighted Sobolev spaces. For the co-rotation case, F. Lin, P. Zhang, and Z. Zhang [26] obtain a global existence results with  $d = 2$  and  $k > 6$ . If the initial data is perturbation around equilibrium, N. Masmoudi [31] proved global well-posedness of (1.4) for  $k > 0$ . In the co-rotation case with  $d = 2$ , he [31] obtained a global result for  $k > 0$  without any small conditions. A. V. Busuioc, I. S. Ciuperca, D. Iftimie and L. I. Palade [5] obtain a global existence result with only the small condition on  $\psi_0$ . The global existence of weak solutions in  $L^2$  was proved recently by N. Masmoudi [32] under some entropy conditions. Recently, M. Schonbek [40] studied the  $L^2$  decay of the velocity for the co-rotation FENE dumbbell model, and obtained the decay rate  $(1+t)^{-\frac{d}{4}+\frac{1}{2}}$ ,  $d \geq 2$  with  $u_0 \in L^1$ . Moreover, she conjectured that the sharp decay rate should be  $(1+t)^{-\frac{d}{4}}$ ,  $d \geq 2$ . However, she failed to get it because she could not use the bootstrap argument as in [39] due to the additional stress tensor. Recently, W. Luo and Z. Yin [27] improved Schonbek's result and showed that the decay rate is  $(1+t)^{-\frac{d}{4}}$  with  $d \geq 3$  and  $\ln^{-l}(1+t)$  with  $d = 2$  for any  $l \in \mathbb{N}^+$ . This result shows that M. Schonbek's conjecture is true when  $d \geq 3$ . More recently, W. Luo and Z. Yin [28] improved the decay rate to  $(1+t)^{-\frac{d}{4}}$  with  $d = 2$ . In [19], Y. Jia and H. Ye also prove the optimal time decay rate via a different approach.

### 1.2. Short reviews for the CNS equations

Taking  $\psi \equiv 0$ , the system (1.1) reduce to the CNS equations. In order to study about the (1.1), we cite some reference about the CNS equations. The first local existence and uniqueness results were obtained by J. Nash [36] for smooth initial data without vacuum. Later on, A. Matsumura and T. Nishida [34] proved the global well-posedness for smooth data close to equilibrium with  $d = 3$ . In [23], A. V. Kazhikhov and V. V. Shelukhin established the first global existence result with large data in one dimensional space under some suitable condition on  $\mu$  and  $\lambda$ . If  $\mu$  is constant and  $\lambda(\rho) = b\rho^\beta$ , X. Huang and J. Li [17] obtained a global existence and uniqueness result for large initial data in two dimensional space(See also [21]). In [18], X. Huang, J. Li, and Z. Xin proved the global well-posedness with vacuum. The Blow-up phenomenons were studied by Z. Xin et al in [43, 24, 45]. Concerning the global existence of weak solutions for the large initial data, we may refer to [3, 4, 35, 42].

To catch the scaling invariance property of the CNS equations. R. Danchin introduced the "critical spaces" in his series papers [8, 9, 10, 11, 13] and obtained several important existence and uniqueness results. Recently, Q. Chen, C. Miao and Z. Zhang [6] proved the local existence and uniqueness in critical homogeneous Besov spaces. The ill-posedness results was obtained in [7]. In [15], L. He, J. Huang and C. Wang proved the global stability with  $d = 3$  i.e. for any perturbed solutions will remain close to the reference solutions if initially they are close to another one.

The large time behaviour of the global solutions  $(\rho, u)$  to the CNS equations with  $d = 3$  was firstly proved by A. Matsumura and T. Nishida in [34]. Recently, H. Li and T. Zhang [25] obtain the optimal time decay rate for the CNS equations with  $d = 3$  by spectrum analysis in Sobolev spaces. R. Danchin and J. Xu [12] studied about the large time behaviour in the critical Besov space with  $d \geq 2$ . J. Xu

[46] obtained the optimal time decay rate with a small low-frequency assumption in some Besov spaces with negative index. More recently, Z. Xin and J. Xu [44] studied about the large time behaviour and removed the smallness of low frequencies.

### 1.3. Main results

The well-posedness for the system (1.1) was established by J. Ning, Y. Liu and T. Zhang [20]. They proved the global well-posedness for (1.1) if the initial data is close to the equilibrium and  $R \in \mathbb{R}^3$ . For the compressible FENE system, Z. Luo, W. Luo and Z. Yin [30] proved the global well-posedness results with  $d \geq 2$  and studied large time behaviour with  $d \geq 3$ .

Recently, N. Masmoudi [33] is concerning with the long time behavior for polymeric models. Large time behaviour of the compressible FENE system with  $d \geq 3$  has been studied in [29] and [30]. To our best knowledge, for  $d = 2$ , large time behaviour for the compressible FENE system (1.2) has not been studied yet. This problem is interesting and more difficult than the case with  $d \geq 3$ . In this paper, we will study about the optimal time decay rate for global strong solutions in  $L^2$ . The proof is based on the Littlewood-Paley decomposition theory and the Fourier splitting method. Motivated by [16] and [48], we deal with the linear term  $\operatorname{div} \tau$  by virtue of the cancellation relation in Fourier space. Firstly we can prove the logarithmic decay rate for the velocity via the method in [27]. By virtue of the time weighted energy estimate and the logarithmic decay rate, then we improve the time decay rate to  $(1+t)^{-\frac{1}{4}}$  for the velocity in  $L^2$ . The main difficult to get optimal time decay rate is that we can not get any information of  $u$  in  $L^1$  from (1.2). Fortunately, similar to [41], we can prove  $\|u\|_{L^\infty(0,\infty;\dot{B}_{2,\infty}^{-\frac{1}{2}})} \leq C$  from (1.2) by using the time decay rate  $(1+t)^{-\frac{1}{4}}$ . Then we improve the time decay rate to  $(1+t)^{-\frac{5}{16}}$  for the velocity in  $L^2$  by the Littlewood-Paley decomposition theory and the Fourier splitting method. We obtain a slightly weaker conclusion  $\|u\|_{L^\infty(0,\infty;\dot{B}_{2,\infty}^{-1})} \leq C$  from (1.2) by using the time decay rate  $(1+t)^{-\frac{5}{16}}$ . Finally, we obtain optimal time decay rate for the velocity in  $L^2$  by the Littlewood-Paley decomposition theory and the standard Fourier splitting method.

Our main result can be stated as follows.

**Theorem 1.1** (Large time behaviour). *Let  $d = 2$ . Let  $(\rho, u, g)$  be a global strong solution of (1.2) with the initial data  $(\rho_0, u_0, g_0)$  under the condition in Theorem 2.6. In addition, if  $(\rho_0, u_0) \in \dot{B}_{2,\infty}^{-1} \times \dot{B}_{2,\infty}^{-1}$  and  $g_0 \in \dot{B}_{2,\infty}^{-1}(\mathcal{L}^2)$ , then there exists a constant  $C$  such that*

$$(1.3) \quad \|\rho\|_{L^2} + \|u\|_{L^2} \leq C(1+t)^{-\frac{1}{2}}$$

and

$$(1.4) \quad \|g\|_{L^2(\mathcal{L}^2)} \leq C(1+t)^{-1}.$$

**Remark 1.2.** *Taking  $\psi \equiv 0$  and combining with the result in [46], we can see that the  $L^2$  decay rate for  $(\rho, u)$  obtained in Theorem 1.1 is optimal.*

**Remark 1.3.** *In previous papers, researchers usually add the condition  $(\rho_0, u_0) \in L^1 \times L^1$  to obtain the optimal time decay rate. Since  $L^1 \hookrightarrow \dot{B}_{2,\infty}^{-1}$ , it follows that our condition is weaker and the results still hold true for  $(\rho_0, u_0) \in L^1 \times L^1$ . Moreover, the assumption can be replaced with a weaker assumption  $\sup_{j \leq j_0} 2^{-j} \|\dot{\Delta}_j(\rho_0, u_0, g_0)\|_{L^2 \times L^2 \times L^2(\mathcal{L}^2)} < \infty$ , for any  $j_0 \in \mathbb{Z}$ .*

The paper is organized as follows. In Section 2 we introduce some notations and give some preliminaries which will be used in the sequel. In Section 3 we prove the  $L^2$  decay of solutions to the 2-D compressible FENE model by using the Fourier splitting method, the Littlewood-Paley decomposition theory and the bootstrap argument.

## 2 Preliminaries

In this section we introduce some notations and useful lemmas which will be used in the sequel.

For  $p \geq 1$ , we denote by  $\mathcal{L}^p$  the space

$$\mathcal{L}^p = \{f \mid \|f\|_{\mathcal{L}^p}^p = \int_B \psi_\infty |f|^p dR < \infty\}.$$

We use the notation  $L_x^p(\mathcal{L}^q)$  to denote  $L^p[\mathbb{R}^2; \mathcal{L}^q]$ :

$$L_x^p(\mathcal{L}^q) = \{f \mid \|f\|_{L_x^p(\mathcal{L}^q)} = \left( \int_{\mathbb{R}^2} \left( \int_B \psi_\infty |f|^q dR \right)^{\frac{2}{q}} dx \right)^{\frac{1}{p}} < \infty\}.$$

The symbol  $\widehat{f} = \mathcal{F}(f)$  represents the Fourier transform of  $f$ . Let  $\Lambda^s f = \mathcal{F}^{-1}(|\xi|^s \widehat{f})$ . If  $s \geq 0$ , we denote by  $H^s(\mathcal{L}^2)$  the space

$$H^s(\mathcal{L}^2) = \{f \mid \|f\|_{H^s(\mathcal{L}^2)}^2 = \int_{\mathbb{R}^2} \int_B (|f|^2 + |\Lambda^s f|^2) \psi_\infty dR dx < \infty\}.$$

Sometimes we write  $f \lesssim g$  instead of  $f \leq Cg$ , where  $C$  is a constant. We agree that  $\nabla$  stands for  $\nabla_x$  and  $div$  stands for  $div_x$ .

For the convenience of readers, we recall the Littlewood-Paley decomposition theory and Besov spaces.

**Proposition 2.1.** [1] *Let  $\mathcal{C}$  be the annulus  $\{\xi \in \mathbb{R}^2 : \frac{3}{4} \leq |\xi| \leq \frac{8}{3}\}$ . There exist radial function  $\varphi$ , valued in the interval  $[0, 1]$ , belonging respectively to  $\mathcal{D}(\mathcal{C})$ , and such that*

$$\forall \xi \in \mathbb{R}^2 \setminus \{0\}, \quad \sum_{j \in \mathbb{Z}} \varphi(2^{-j} \xi) = 1,$$

$$|j - j'| \geq 2 \Rightarrow \text{Supp } \varphi(2^{-j} \cdot) \cap \text{Supp } \varphi(2^{-j'} \cdot) = \emptyset.$$

Further, we have

$$\forall \xi \in \mathbb{R}^2 \setminus \{0\}, \quad \frac{1}{2} \leq \sum_{j \in \mathbb{Z}} \varphi^2(2^{-j} \xi) \leq 1.$$

Let  $u$  be a tempered distribution in  $\mathcal{S}'_h(\mathbb{R}^2)$ . For all  $j \in \mathbb{Z}$ , define

$$\dot{\Delta}_j u = \mathcal{F}^{-1}(\varphi(2^{-j} \cdot) \mathcal{F}u).$$

Then the Littlewood-Paley decomposition is given as follows:

$$u = \sum_{j \in \mathbb{Z}} \dot{\Delta}_j u \quad \text{in } \mathcal{S}'(\mathbb{R}^2).$$

Let  $s \in \mathbb{R}$ ,  $1 \leq p, r \leq \infty$ . The homogeneous Besov space  $\dot{B}_{p,r}^s$  and  $\dot{B}_{p,r}^s(\mathcal{L}^q)$  is defined by

$$\dot{B}_{p,r}^s = \{u \in \mathcal{S}'_h : \|u\|_{\dot{B}_{p,r}^s} = \left\| (2^{js} \|\dot{\Delta}_j u\|_{L^p})_j \right\|_{l^r(\mathbb{Z})} < \infty\},$$

$$\dot{B}_{p,r}^s(\mathcal{L}^q) = \{\phi \in \mathcal{S}'_h : \|\phi\|_{\dot{B}_{p,r}^s(\mathcal{L}^q)} = \left\| (2^{js} \|\dot{\Delta}_j \phi\|_{L_x^p(\mathcal{L}^q)})_j \right\|_{l^r(\mathbb{Z})} < \infty\}.$$

We introduce the following results about inclusions between Lebesgue and Besov spaces

**Lemma 2.2.** [1] *Let  $1 \leq p \leq 2$  and  $d = 2$ . Then we have*

$$L^p \hookrightarrow \dot{B}_{2,\infty}^{1-\frac{2}{p}}.$$

The following lemma is the Gagliardo-Nirenberg inequality of Sobolev type.

**Lemma 2.3.** [37] *Let  $d \geq 2$ ,  $p \in [2, +\infty)$  and  $0 \leq s, s_1 \leq s_2$ , then there exists a constant  $C$  such that*

$$\|\Lambda^s f\|_{L^p} \leq C \|\Lambda^{s_1} f\|_{L^2}^{1-\theta} \|\Lambda^{s_2} f\|_{L^2}^\theta,$$

where  $0 \leq \theta \leq 1$  and  $\theta$  satisfy

$$s + d\left(\frac{1}{2} - \frac{1}{p}\right) = s_1(1 - \theta) + \theta s_2.$$

Note that we require that  $0 < \theta < 1$ ,  $0 \leq s_1 \leq s$ , when  $p = \infty$ .

The following lemma allows us to estimate the extra stress tensor  $\tau$ .

**Lemma 2.4.** [31] *If  $\int_B g \psi_\infty dR = 0$ , then there exists a constant  $C$  such that*

$$\|g\|_{\mathcal{L}^2} \leq C \|\nabla_R g\|_{\mathcal{L}^2}.$$

**Lemma 2.5.** [31] *For all  $\delta > 0$ , there exists a constant  $C_\delta$  such that*

$$|\tau(g)|^2 \leq \delta \|\nabla_R g\|_{\mathcal{L}^2}^2 + C_\delta \|g\|_{\mathcal{L}^2}^2.$$

If  $(p-1)k > 1$ , then

$$|\tau(g)| \leq C \|g\|_{\mathcal{L}^p}.$$

Denote the energy and energy dissipation functionals for  $(\rho, u, g)$  as follows:

$$E(t) = \|\rho\|_{H^s}^2 + \|u\|_{H^s}^2 + \|g\|_{H^s(\mathcal{L}^2)}^2,$$

and

$$D(t) = \|\nabla \rho\|_{H^{s-1}}^2 + \mu \|\nabla u\|_{H^s}^2 + (\mu + \mu') \|\operatorname{div} u\|_{H^s}^2 + \|\nabla_R g\|_{H^s(\mathcal{L}^2)}^2.$$

To study the  $L^2$  decay rate, we recall the following global existence of strong solutions for (1.2).

**Theorem 2.6.** [30] *Let  $d = 2$  and  $s > 2$ . Let  $(\rho, u, g)$  be a strong solution of (1.2) with the initial data  $(\rho_0, u_0, g_0)$  satisfying the conditions  $\int_B g_0 \psi_\infty dR = 0$  and  $1 + g_0 > 0$ . Then, there exists some sufficiently small constant  $\epsilon_0$  such that if*

$$(2.1) \quad E(0) = \|\rho_0\|_{H^s}^2 + \|u_0\|_{H^s}^2 + \|g_0\|_{H^s(\mathcal{L}^2)}^2 \leq \epsilon_0,$$

then (1.2) admits a unique global strong solution  $(\rho, u, g)$  with  $\int_B g \psi_\infty dR = 0$  and  $1 + g > 0$ , and we have

$$(2.2) \quad \sup_{t \in [0, +\infty)} E(t) + \int_0^\infty D(t) dt \leq C_0 E(0),$$

where  $C_0 > 1$  is a constant.

### 3 The $L^2$ decay rate

This section is devoted to investigating the long time behaviour for the compressible FENE dumbbell model with  $d = 2$ . Since the additional stress tensor  $\tau$  does not decay fast enough, we failed to use the bootstrap argument as in [39, 28]. To deal with this term, we consider the coupling effect between  $\rho$ ,  $u$  and  $g$ . Motivated by [16], [27] and [48], we obtain the initial  $L^2$  decay rate by taking Fourier transform in (1.2) and using the Fourier splitting method in following Proposition.

**Proposition 3.1.** *Let  $d = 2$ . Under the condition in Theorem 1.1, for any  $l \in \mathbb{N}^+$ , then there exists a constant  $C$  such that*

$$(3.1) \quad E_\eta(t) \leq C \ln^{-l}(e + t)$$

*Proof.* Consider that

$$E_\eta(t) = \sum_{n=0,s} ( \|h(\rho)^{\frac{1}{2}} \Lambda^n \rho\|_{L^2}^2 + \|(1 + \rho)^{\frac{1}{2}} \Lambda^n u\|_{L^2}^2 ) + \lambda \|g\|_{H^s(\mathcal{L}^2)}^2 + 2\eta \sum_{m=0,s-1} \int_{\mathbb{R}^2} \Lambda^m u \nabla \Lambda^m \rho dx,$$

and

$$D_\eta(t) = \eta \gamma \|\nabla \rho\|_{H^{s-1}}^2 + \mu \|\nabla u\|_{H^s}^2 + (\mu + \mu') \|\operatorname{div} u\|_{H^s}^2 + \lambda \|\nabla_R g\|_{H^s(\mathcal{L}^2)}^2.$$

For some sufficiently small constant  $\eta > 0$ , we obtain  $E(t) \sim E_\eta(t)$  and  $D(t) \sim D_\eta(t)$ .

In the proof of Theorem 2.6, see [30], we have the following global energy estimation:

$$(3.2) \quad \frac{d}{dt} E_\eta + D_\eta \leq 0.$$

Define  $S_0(t) = \{\xi : |\xi|^2 \leq 2C_2 \frac{f'(t)}{f(t)}\}$  with  $f(t) = \ln^3(e + t)$  and  $C_2$  large enough. According to Schonbek's strategy, we obtain

$$(3.3) \quad \begin{aligned} & \frac{d}{dt} [f(t) E_\eta(t)] + C_2 f'(t) (\mu \|u\|_{H^s}^2 + \eta \gamma \|\rho\|_{H^{s-1}}^2) + f(t) \|\nabla_R g\|_{H^s(\mathcal{L}^2)}^2 \\ & \leq C f'(t) \int_{S_0(t)} |\hat{u}(\xi)|^2 + |\hat{\rho}(\xi)|^2 d\xi + f'(t) \|\Lambda^s \rho\|_{L^2}^2. \end{aligned}$$

We now focus on the  $L^2$  estimate to the low frequency part of  $\rho$  and  $u$ . Applying Fourier transform to (1.2), we obtain

$$(3.4) \quad \begin{cases} \hat{\rho}_t + i\xi_k \hat{u}^k = \hat{F}, \\ \hat{u}_t^j + \mu |\xi|^2 \hat{u}^j + (\mu + \mu') \xi_j \xi_k \hat{u}^k + i\xi_j \gamma \hat{\rho} - i\xi_k \hat{\tau}^{jk} = \hat{G}^j, \\ \hat{g}_t + \mathcal{L} \hat{g} - i\xi_k \hat{u}^j R_j \partial_{R_k} \mathcal{U} + i\xi_k \hat{u}^k = \hat{H}, \end{cases}$$

where  $F = -\operatorname{div}(\rho u)$ ,  $G = -u \cdot \nabla u + [i(\rho) - 1](\operatorname{div} \Sigma(u) + \operatorname{div} \tau) + [\gamma - h(\rho)] \nabla \rho$  and  $H = -u \cdot \nabla g - \frac{1}{\psi_\infty} \nabla_R \cdot (\nabla u R g \psi_\infty)$ .

From (3.4), using the fact  $\int_B i\xi_k \hat{u}^k \bar{\hat{g}} \psi_\infty dR = 0$ , we deduce that

$$(3.5) \quad \begin{cases} \frac{1}{2} \frac{d}{dt} |\hat{\rho}|^2 + \operatorname{Re}[i\xi \cdot \hat{u} \bar{\hat{\rho}}] = \operatorname{Re}[\hat{F} \bar{\hat{\rho}}], \\ \frac{1}{2} \frac{d}{dt} |\hat{u}|^2 + \operatorname{Re}[\gamma \hat{\rho} i\xi \cdot \hat{u}] + \mu |\xi|^2 |\hat{u}|^2 + (\mu + \mu') |\xi \cdot \hat{u}|^2 - \operatorname{Re}[i\xi \otimes \hat{u}(t, \xi) : \hat{\tau}] = \operatorname{Re}[\hat{G} \cdot \hat{u}], \\ \frac{1}{2} \frac{d}{dt} \|\hat{g}\|_{\mathcal{L}^2}^2 + \|\nabla_R \hat{g}\|_{\mathcal{L}^2}^2 - \operatorname{Re}[i\xi \otimes \hat{u} : \hat{\tau}] = \operatorname{Re}[\int_B \hat{H} \bar{\hat{g}} \psi_\infty dR]. \end{cases}$$

One can verify that

$$\mathcal{R}e[i\xi \cdot \hat{u}\bar{\rho}] + \mathcal{R}e[\hat{\rho}i\xi \cdot \hat{u}] = \mathcal{R}e[i\xi \otimes \hat{u}(t, \xi) : \hat{\tau}] + \mathcal{R}e[i\xi \otimes \hat{u} : \hat{\tau}] = 0,$$

which implies that

$$(3.6) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} (\gamma |\hat{\rho}|^2 + |\hat{u}|^2 + \|\hat{g}\|_{\mathcal{L}^2}^2) + \mu |\xi|^2 |\hat{u}|^2 + (\mu + \mu') |\xi \cdot \hat{u}|^2 + \|\nabla_{R\hat{g}}\|_{\mathcal{L}^2}^2 \\ & = \mathcal{R}e[\gamma \hat{F}\bar{\rho}] + \mathcal{R}e[\hat{G} \cdot \hat{u}] + \mathcal{R}e\left[\int_B \hat{H}\hat{g}\psi_\infty dR\right]. \end{aligned}$$

Multiplying  $i\xi \cdot \hat{u}$  to the first equation of (3.4), multiplying  $-i\xi_j \bar{\rho}$  with  $1 \leq j \leq 2$  to the second equation of (3.4) and taking the real part, we can deduce that

$$(3.7) \quad \mathcal{R}e[\hat{\rho}_t i\xi \cdot \hat{u}] - |\xi \cdot \hat{u}|^2 = \mathcal{R}e[\hat{F}i\xi \cdot \hat{u}],$$

and

$$(3.8) \quad \mathcal{R}e[\hat{\rho}i\xi \cdot \hat{u}_t] + \gamma |\xi|^2 |\hat{\rho}|^2 + (2\mu + \mu') |\xi|^2 \mathcal{R}e[\hat{\rho}i\xi \cdot \hat{u}] - \mathcal{R}e[\hat{\rho}\xi \otimes \xi : \hat{\tau}] = \mathcal{R}e[\hat{G} \cdot i\xi \hat{\rho}].$$

It follows from (3.6)–(3.8) that

$$(3.9) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} (\gamma |\hat{\rho}|^2 + |\hat{u}|^2 + \|\hat{g}\|_{\mathcal{L}^2}^2 + 2(\mu + \mu') \mathcal{R}e[\hat{\rho}i\xi \cdot \hat{u}]) + \mu |\xi|^2 |\hat{u}|^2 + (\mu + \mu') \gamma |\xi|^2 |\hat{\rho}|^2 + \|\nabla_{R\hat{g}}\|_{\mathcal{L}^2}^2 \\ & = -(\mu + \mu')(2\mu + \mu') |\xi|^2 \mathcal{R}e[\hat{\rho}i\xi \cdot \hat{u}] + (\mu + \mu') \mathcal{R}e[\hat{\rho}\xi \otimes \xi : \hat{\tau}] + (\mu + \mu') \mathcal{R}e[\hat{F}i\xi \cdot \hat{u}] \\ & + (\mu + \mu') \mathcal{R}e[\hat{G} \cdot i\xi \hat{\rho}] + \mathcal{R}e[\gamma \hat{F}\bar{\rho}] + \mathcal{R}e[\hat{G} \cdot \hat{u}] + \mathcal{R}e\left[\int_B \hat{H}\hat{g}\psi_\infty dR\right]. \end{aligned}$$

Consider  $\xi \in S_0(t)$  and sufficiently large  $t$ , by Lemma 2.4 and 2.5, we deduce that

$$(3.10) \quad \begin{aligned} |\hat{\rho}|^2 + |\hat{u}|^2 + \|\hat{g}\|_{\mathcal{L}^2}^2 & \leq C(|\hat{\rho}_0|^2 + |\hat{u}_0|^2 + \|\hat{g}_0\|_{\mathcal{L}^2}^2) + C \int_0^t |\hat{G} \cdot \hat{u}| + |\widehat{\rho u}|^2 + |\hat{G}|^2 ds \\ & + C_\delta \int_0^t \int_B \psi_\infty |\mathcal{F}(u \cdot \nabla g)|^2 + \psi_\infty |\mathcal{F}(\nabla u \cdot Rg)|^2 dR ds. \end{aligned}$$

Integrating over  $S_0(t)$  with  $\xi$ , then we have the following estimation to (3.4):

$$(3.11) \quad \begin{aligned} & \int_{S_0(t)} (|\hat{\rho}|^2 + |\hat{u}|^2 + \|\hat{g}\|_{\mathcal{L}^2}^2) d\xi \leq C \int_{S_0(t)} (|\hat{\rho}_0|^2 + |\hat{u}_0|^2 + \|\hat{g}_0\|_{\mathcal{L}^2}^2) d\xi + C \int_{S_0(t)} \int_0^t |\hat{G} \cdot \hat{u}| + |\widehat{\rho u}|^2 + |\hat{G}|^2 ds d\xi \\ & + C_\delta \int_{S_0(t)} \int_0^t \int_B \psi_\infty |\mathcal{F}(u \cdot \nabla g)|^2 + \psi_\infty |\mathcal{F}(\nabla u \cdot Rg)|^2 dR ds d\xi. \end{aligned}$$

If  $E(0) < \infty$  and  $(\rho_0, u_0, g_0) \in \dot{B}_{2,\infty}^{-1} \times \dot{B}_{2,\infty}^{-1} \times \dot{B}_{2,\infty}^{-1}(\mathcal{L}^2)$ , applying Proposition 2.1, we have

$$\begin{aligned} \int_{S_0(t)} (|\hat{\rho}_0|^2 + |\hat{u}_0|^2 + \|\hat{g}_0\|_{\mathcal{L}^2}^2) d\xi & \leq \sum_{j \leq \log_2[\frac{4}{3}C_2^{\frac{1}{2}} \sqrt{\frac{f'(t)}{f(t)}}]} \int_{\mathbb{R}^d} 2\varphi^2(2^{-j}\xi) (|\hat{\rho}_0|^2 + |\hat{u}_0|^2 + \|\hat{g}_0\|_{\mathcal{L}^2}^2) d\xi \\ & \leq \sum_{j \leq \log_2[\frac{4}{3}C_2^{\frac{1}{2}} \sqrt{\frac{f'(t)}{f(t)}}]} (\|\dot{\Delta}_j u_0\|_{L^2}^2 + \|\dot{\Delta}_j \rho_0\|_{L^2}^2 + \|\dot{\Delta}_j g_0\|_{L^2(\mathcal{L}^2)}^2) \end{aligned}$$



$$\begin{aligned}
 &\leq \sum_{j \leq \log_2 [\frac{4}{3} C_2^{\frac{1}{2}} \sqrt{\frac{f'(t)}{f(t)}}]} 2^{2j} (\|u_0\|_{\dot{B}_{2,\infty}^{-1}}^2 + \|\rho_0\|_{\dot{B}_{2,\infty}^{-1}}^2 + \|g_0\|_{\dot{B}_{2,\infty}^{-1}(\mathcal{L}^2)}^2) \\
 &\leq C \frac{f'(t)}{f(t)} (\|u_0\|_{\dot{B}_{2,\infty}^{-1}}^2 + \|\rho_0\|_{\dot{B}_{2,\infty}^{-1}}^2 + \|g_0\|_{\dot{B}_{2,\infty}^{-1}(\mathcal{L}^2)}^2).
 \end{aligned}$$

Thanks to Minkowski's inequality and Theorem 2.6, we get

$$\begin{aligned}
 (3.12) \quad \int_{S_0(t)} \int_0^t |\hat{G}|^2 ds d\xi &\leq C \int_{S_0(t)} d\xi \int_0^t \|\hat{G}\|_{L^\infty}^2 ds \\
 &\leq C \frac{f'(t)}{f(t)}.
 \end{aligned}$$

Using Theorem 2.6 and Lemma 2.4, we get

$$\begin{aligned}
 (3.13) \quad \int_{S_0(t)} \int_0^t \int_B \psi_\infty |\mathcal{F}(u \cdot \nabla g)|^2 + \psi_\infty |\mathcal{F}(\nabla u \cdot Rg)|^2 dR ds d\xi \\
 \leq C \frac{f'(t)}{f(t)} \int_0^t (\|u\|_{L^2}^2 \|\nabla g\|_{L^2(\mathcal{L}^2)}^2 + \|\nabla u\|_{L^2}^2 \|g\|_{L^2(\mathcal{L}^2)}^2) ds \\
 \leq C \frac{f'(t)}{f(t)}.
 \end{aligned}$$

Similarly, we have

$$\begin{aligned}
 (3.14) \quad \int_{S_0(t)} \int_0^t |\hat{G} \cdot \bar{u}| + |\widehat{\rho u}|^2 ds d\xi &= \int_0^t \int_{S_0(t)} |\hat{G} \cdot \bar{u}| + |\widehat{\rho u}|^2 d\xi ds \\
 &\leq C \sqrt{\frac{f'(t)}{f(t)}} \int_0^t (\|u\|_{L^2}^2 + \|\rho\|_{L^2}^2) D(s)^{\frac{1}{2}} ds + C \frac{f'(t)}{f(t)} \int_0^t \|u\|_{L^2}^2 \|\rho\|_{L^2}^2 ds \\
 &\leq C \sqrt{\frac{f'(t)}{f(t)}} (1+t)^{\frac{1}{2}} + C \frac{f'(t)}{f(t)} (1+t).
 \end{aligned}$$

Plugging the above estimates into (3.11), we obtain

$$(3.15) \quad \int_{S_0(t)} |\hat{\rho}|^2 + |\hat{u}|^2 d\xi \leq C \ln^{-\frac{1}{2}}(e+t).$$

According to (3.3) and (3.15), we deduce that

$$\begin{aligned}
 (3.16) \quad \frac{d}{dt} [f(t) E_\eta(t)] + C_2 f'(t) (\mu \|u\|_{H^s}^2 + \eta \gamma \|\rho\|_{H^{s-1}}^2) + f(t) \|\nabla Rg\|_{H^s(\mathcal{L}^2)}^2 \\
 \leq C f'(t) \ln^{-\frac{1}{2}}(e+t) + 2f'(t) \|\Lambda^s \rho\|_{L^2}^2,
 \end{aligned}$$

which implies that

$$f(t) E_\eta(t) \leq C \int_0^t f'(s) \ln^{-\frac{1}{2}}(e+s) ds + C \int_0^t 2f'(s) \|\Lambda^s \rho\|_{L^2}^2 ds \leq C \ln^{\frac{5}{2}}(e+t).$$

We thus get

$$(3.17) \quad E_\eta(t) \leq C \ln^{-\frac{1}{2}}(e+t).$$

We improve the  $L^2$  decay rate in (3.17) by using the bootstrap argument. According to (3.14) and (3.17), we have

$$\begin{aligned}
 (3.18) \quad & \int_{S_0(t)} \int_0^t |\hat{G} \cdot \bar{u}| + |\widehat{\rho u}|^2 ds d\xi \\
 & \leq C \sqrt{\frac{f'(t)}{f(t)}} \int_0^t (\|u\|_{L^2}^2 + \|\rho\|_{L^2}^2) D(s)^{\frac{1}{2}} ds + C \frac{f'(t)}{f(t)} \int_0^t \|u\|_{L^2}^2 \|\rho\|_{L^2}^2 ds \\
 & \leq C \sqrt{\frac{f'(t)}{f(t)}} (1+t)^{\frac{1}{2}} \ln^{-\frac{1}{2}}(e+t) + C \frac{f'(t)}{f(t)} (1+t) \ln^{-1}(e+t),
 \end{aligned}$$

where in the last inequality we have used the fact that

$$\lim_{t \rightarrow \infty} \frac{\int_0^t \ln^{-1}(e+s) ds}{(1+t) \ln^{-1}(e+t)} = \lim_{t \rightarrow \infty} \frac{\ln^{-1}(e+t)}{\ln^{-1}(1+t) - \ln^{-2}(e+t)} = 1.$$

Then the proof of (3.15) implies that

$$(3.19) \quad \int_{S_0(t)} |\hat{\rho}|^2 + |\hat{u}|^2 d\xi \leq C \ln^{-1}(e+t).$$

According to (3.3) and (3.19), we deduce that

$$E_\eta \leq C \ln^{-1}(e+t).$$

Using the bootstrap argument, for any  $l \in N^+$ , we obtain

$$(3.20) \quad E_\eta \leq C \ln^{-l}(e+t).$$

We thus complete the proof of Proposition 3.1 □

In order to improve the decay rate, we estimate the following high order energy:

$$\begin{aligned}
 E_\eta^1(t) &= \sum_{n=1, s} (\|h(\rho)^{\frac{1}{2}} \Lambda^n \rho\|_{L^2}^2 + \|(1+\rho)^{\frac{1}{2}} \Lambda^n u\|_{L^2}^2) \\
 &\quad + \|\Lambda^1 g\|_{H^{s-1}(\mathcal{L}^2)}^2 + 2\eta \sum_{m=1, s-1} \int_{\mathbb{R}^2} \Lambda^m u \nabla \Lambda^m \rho dx,
 \end{aligned}$$

and

$$D_\eta^1(t) = \eta \gamma \|\nabla \Lambda^1 \rho\|_{H^{s-2}}^2 + \mu \|\nabla \Lambda^1 u\|_{H^{s-1}}^2 + (\mu + \mu') \|\operatorname{div} \Lambda^1 u\|_{H^{s-1}}^2 + \|\Lambda^1 \nabla_R g\|_{H^{s-1}(\mathcal{L}^2)}^2.$$

The following Lemma can be proved by the standard energy method. Thus we omit the proof here.

**Lemma 3.2.** *Let  $(\rho, u, g) \in L^\infty([0, +\infty); H^s \times H^s \times H^s(\mathcal{L}^2))$  be global strong solutions constructed in Theorem 2.6. If  $t \in (0, +\infty)$ , then we have*

$$(3.21) \quad \frac{d}{dt} E_\eta^1(t) + D_\eta^1(t) \leq 0.$$

The following proposition is about the high order energy estimate.

**Proposition 3.3.** *Let  $d = 2$ . Under the condition in Theorem 1.1, for any  $l \in \mathbb{N}^+$ , then there exists a constant  $C$  such that*

$$(3.22) \quad E_\eta^1(t) \leq C(1+t)^{-1} \ln^{-l}(e+t),$$

and

$$(3.23) \quad \int_0^t (1+s)f(s) \|\Lambda^1 \nabla_{RG}\|_{H^{s-1}(\mathcal{L}^2)}^2 ds \leq C.$$

*Proof.* Applying Lemma 3.2, we have

$$(3.24) \quad \frac{d}{dt} E_\eta^1 + D_\eta^1 \leq 0,$$

which implies that

$$(3.25) \quad \begin{aligned} & \frac{d}{dt} [f(t)E_\eta^1] + C_2 f'(t) (\mu \|\Lambda^1 u\|_{H^{s-1}}^2 + \eta \gamma \|\Lambda^1 \rho\|_{H^{s-2}}^2) + f(t) \|\Lambda^1 \nabla_{RG}\|_{H^{s-1}(\mathcal{L}^2)}^2 \\ & \leq C f'(t) \int_{S_0(t)} |\xi|^2 (|\hat{u}(\xi)|^2 + |\hat{\rho}(\xi)|^2) d\xi + f'(t) \|\Lambda^s \rho\|_{L^2}^2. \end{aligned}$$

According to (3.20), we have

$$f'(t) \int_{S_0(t)} |\xi|^2 (|\hat{u}(\xi)|^2 + |\hat{\rho}(\xi)|^2) d\xi \leq C(1+t)^{-2} \ln^{-l+1}(e+t).$$

This together with (3.20), (3.25) and (3.2) ensures that

$$\begin{aligned} (1+t) \ln^{l+1}(e+t) E_\eta^1 & \leq C + C \ln(e+t) + C \int_0^t \ln^l(e+s) \|\Lambda^s \rho\|_{L^2}^2 ds + C \int_0^t \ln^{l+1}(e+s) E_\eta^1 ds \\ & \leq C \ln(e+t) + C \int_0^t \ln^{l+1}(e+s) D_\eta ds \\ & \leq C \ln(e+t) + C \int_0^t (1+s)^{-1} \ln^l(e+s) E_\eta ds \\ & \leq C \ln(e+t), \end{aligned}$$

which implies that

$$(3.26) \quad E_\eta^1 \leq C(1+t)^{-1} \ln^{-l}(e+t).$$

Using (3.25) and (3.26) with  $l \geq 5$ , we have the following estimate for  $\nabla_{RG}$ :

$$(3.27) \quad \begin{aligned} & \int_0^t (1+s)f(s) \|\Lambda^1 \nabla_{RG}\|_{H^{s-1}(\mathcal{L}^2)}^2 ds \\ & \leq C + \int_0^t (1+s)f'(s) \|\Lambda^s \rho\|_{L^2}^2 ds + C \int_0^t f(s) E_\eta^1 ds \leq C. \end{aligned}$$

We thus complete the proof of Proposition 3.3.  $\square$

By virtue of the standard method, one can not obtain the optimal decay rate. However, we can obtain a weak result as follow.

**Proposition 3.4.** *Under the condition in Theorem 1.1, then there exists a constant  $C$  such that*

$$(3.28) \quad E_\eta(t) \leq C(1+t)^{-\frac{1}{2}},$$

and

$$(3.29) \quad E_\eta^1(t) \leq C(1+t)^{-\frac{3}{2}}.$$

*Proof.* Define  $S(t) = \{\xi : |\xi|^2 \leq C_2(1+t)^{-1}\}$  where the constant  $C_2$  will be chosen later on. Using Schonbek's strategy, we split the phase space into two time-dependent domain:

$$\|\nabla u\|_{H^s}^2 = \int_{S(t)} (1 + |\xi|^{2s}) |\xi|^2 |\hat{u}(\xi)|^2 d\xi + \int_{S(t)^c} (1 + |\xi|^{2s}) |\xi|^2 |\hat{u}(\xi)|^2 d\xi.$$

Then we can easily deduce that

$$\frac{C_2}{1+t} \int_{S(t)^c} (1 + |\xi|^{2s}) |\hat{u}(\xi)|^2 d\xi \leq \|\nabla u\|_{H^s}^2,$$

and

$$\frac{C_2}{1+t} \int_{S(t)^c} (1 + |\xi|^{2s-2}) |\hat{\rho}(\xi)|^2 d\xi \leq \|\nabla \rho\|_{H^{s-1}}^2.$$

According to (3.2), we have

$$(3.30) \quad \frac{d}{dt} E_\eta(t) + \frac{\mu C_2}{1+t} \|u\|_{H^s}^2 + \frac{\eta \gamma C_2}{1+t} \|\rho\|_{H^{s-1}}^2 + \|\nabla Rg\|_{H^s(\mathcal{L}^2)}^2 \leq \frac{CC_2}{1+t} \int_{S(t)} |\hat{u}(\xi)|^2 + |\hat{\rho}(\xi)|^2 d\xi.$$

Integrating (3.10) over  $S(t)$  with  $\xi$ , then we have

$$(3.31) \quad \int_{S(t)} |\hat{\rho}|^2 + |\hat{u}|^2 + \|\hat{g}\|_{\mathcal{L}^2}^2 d\xi \leq C \int_{S(t)} (|\hat{\rho}_0|^2 + |\hat{u}_0|^2 + \|\hat{g}_0\|_{\mathcal{L}^2}^2) d\xi + C \int_{S(t)} \int_0^t |\hat{G} \cdot \hat{u}| + |\widehat{\rho u}|^2 + |\hat{G}|^2 ds d\xi \\ + C_\delta \int_{S(t)} \int_0^t \int_B \psi_\infty |\mathcal{F}(u \cdot \nabla g)|^2 + \psi_\infty |\mathcal{F}(\nabla u \cdot Rg)|^2 dR ds d\xi.$$

If  $E(0) < \infty$  and  $(\rho_0, u_0, g_0) \in \dot{B}_{2,\infty}^{-1} \times \dot{B}_{2,\infty}^{-1} \times \dot{B}_{2,\infty}^{-1}(\mathcal{L}^2)$ , applying Proposition 2.1, we have

$$\int_{S(t)} (|\hat{\rho}_0|^2 + |\hat{u}_0|^2 + \|\hat{g}_0\|_{\mathcal{L}^2}^2) d\xi \leq \sum_{j \leq \log_2[\frac{4}{3}C_2^{\frac{1}{2}}(1+t)^{-\frac{1}{2}}]} \int_{\mathbb{R}^d} 2\varphi^2(2^{-j}\xi) (|\hat{\rho}_0|^2 + |\hat{u}_0|^2 + \|\hat{g}_0\|_{\mathcal{L}^2}^2) d\xi \\ \leq \sum_{j \leq \log_2[\frac{4}{3}C_2^{\frac{1}{2}}(1+t)^{-\frac{1}{2}}]} (\|\dot{\Delta}_j u_0\|_{L^2}^2 + \|\dot{\Delta}_j \rho_0\|_{L^2}^2 + \|\dot{\Delta}_j g_0\|_{L^2(\mathcal{L}^2)}^2) \\ \leq \sum_{j \leq \log_2[\frac{4}{3}C_2^{\frac{1}{2}}(1+t)^{-\frac{1}{2}}]} 2^{2j} (\|u_0\|_{\dot{B}_{2,\infty}^{-1}}^2 + \|\rho_0\|_{\dot{B}_{2,\infty}^{-1}}^2 + \|g_0\|_{\dot{B}_{2,\infty}^{-1}(\mathcal{L}^2)}^2) \\ \leq C(1+t)^{-1} (\|u_0\|_{\dot{B}_{2,\infty}^{-1}}^2 + \|\rho_0\|_{\dot{B}_{2,\infty}^{-1}}^2 + \|g_0\|_{\dot{B}_{2,\infty}^{-1}(\mathcal{L}^2)}^2).$$

Thanks to Minkowski's inequality, we get

$$\int_{S(t)} \int_0^t |\widehat{\rho u}|^2 ds d\xi = \int_0^t \int_{S(t)} |\widehat{\rho u}|^2 d\xi ds$$

$$\begin{aligned} &\leq C \int_{S(t)} d\xi \int_0^t \|\widehat{\rho u}\|^2_{L^\infty} ds \\ &\leq C(1+t)^{-1} \int_0^t \|\rho\|_{L^2}^2 \|u\|_{L^2}^2 ds, \end{aligned}$$

and

$$\begin{aligned} \int_{S(t)} \int_0^t |\hat{G}|^2 ds d\xi &\leq C \int_{S(t)} d\xi \int_0^t \|\hat{G}\|^2_{L^\infty} ds \\ &\leq C(1+t)^{-1}. \end{aligned}$$

Thanks to Lemma 2.4 and Lemma 2.5, we have

$$\begin{aligned} (3.32) \quad \int_{S(t)} \int_0^t |\hat{G} \cdot \bar{u}| ds d\xi &\leq C \left( \int_{S(t)} d\xi \right)^{\frac{1}{2}} \int_0^t \|\hat{G} \cdot \bar{u}\|_{L^2} ds \\ &\leq C(1+t)^{-\frac{1}{2}} \int_0^t (\|u\|_{L^2}^2 + \|\rho\|_{L^2}^2) (\|\nabla u\|_{H^1} + \|\nabla \rho\|_{L^2} + \|\nabla \nabla_{Rg}\|_{L^2(\mathcal{L}^2)}) ds. \end{aligned}$$

Using Theorem 2.6 and Lemma 2.4, we get

$$\begin{aligned} (3.33) \quad \int_{S(t)} \int_0^t \int_B \psi_\infty |\mathcal{F}(u \cdot \nabla g)|^2 + \psi_\infty |\mathcal{F}(\nabla u \cdot Rg)|^2 dR ds d\xi \\ \leq C(1+t)^{-1} \int_0^t \|u\|_{L^2}^2 \|\nabla g\|_{L^2(\mathcal{L}^2)}^2 + \|\nabla u\|_{L^2}^2 \|g\|_{L^2(\mathcal{L}^2)}^2 ds \\ \leq C(1+t)^{-1}. \end{aligned}$$

Plugging the above estimates into (3.31), we obtain

$$\begin{aligned} (3.34) \quad \int_{S(t)} |\hat{\rho}(t, \xi)|^2 + |\hat{u}(t, \xi)|^2 d\xi &\leq C(1+t)^{-1} + C(1+t)^{-1} \int_0^t \|\rho\|_{L^2}^2 \|u\|_{L^2}^2 ds \\ &+ C(1+t)^{-\frac{1}{2}} \int_0^t (\|u\|_{L^2}^2 + \|\rho\|_{L^2}^2) (\|\nabla u\|_{H^1} + \|\nabla \rho\|_{L^2} + \|\nabla \nabla_{Rg}\|_{L^2(\mathcal{L}^2)}) ds. \end{aligned}$$

According to (3.30) and (3.34), we deduce that

$$\begin{aligned} &\frac{d}{dt} E_\eta(t) + \frac{\mu C_2}{1+t} \|u\|_{H^s}^2 + \frac{\eta \gamma C_2}{1+t} \|\rho\|_{H^{s-1}}^2 + \|\nabla_{Rg}\|_{H^s(\mathcal{L}^2)}^2 \\ &\leq \frac{CC_2}{1+t} [(1+t)^{-1} + (1+t)^{-1} \int_0^t \|\rho\|_{L^2}^2 \|u\|_{L^2}^2 ds \\ &+ (1+t)^{-\frac{1}{2}} \int_0^t (\|u\|_{L^2}^2 + \|\rho\|_{L^2}^2) (\|\nabla u\|_{H^1} + \|\nabla \rho\|_{L^2} + \|\nabla \nabla_{Rg}\|_{L^2(\mathcal{L}^2)}) ds.] \end{aligned}$$

If  $C_2$  large enough, then we have

$$\begin{aligned} (3.35) \quad (1+t)^{\frac{3}{2}} E_\eta(t) &\leq C \int_0^t \|\Lambda^s \rho\|_{L^2}^2 (1+s)^{\frac{1}{2}} ds + C(1+t)^{\frac{1}{2}} + C \int_0^t (1+s)^{-\frac{1}{2}} \int_0^s \|\rho\|_{L^2}^2 \|u\|_{L^2}^2 ds' ds \\ &+ C(1+t) \int_0^t (\|u\|_{L^2}^2 + \|\rho\|_{L^2}^2) (\|\nabla u\|_{H^1} + \|\nabla \rho\|_{L^2} + \|\nabla \nabla_{Rg}\|_{L^2(\mathcal{L}^2)}) ds \\ &\leq C(1+t)^{\frac{1}{2}} + C \int_0^t (1+s)^{-\frac{1}{2}} \int_0^s \|\rho\|_{L^2}^2 \|u\|_{L^2}^2 ds' ds \end{aligned}$$

$$\begin{aligned}
 & + C(1+t) \int_0^t (\|u\|_{L^2}^2 + \|\rho\|_{L^2}^2) (\|\nabla u\|_{H^1} + \|\nabla \rho\|_{L^2} + \|\nabla \nabla_{Rg}\|_{L^2(\mathcal{L}^2)}) ds \\
 & \leq C(1+t)^{\frac{1}{2}} + C(1+t)^{\frac{1}{2}} \int_0^t \|\rho\|_{L^2}^2 \|u\|_{L^2}^2 ds \\
 & + C(1+t) \int_0^t (\|u\|_{L^2}^2 + \|\rho\|_{L^2}^2) (\|\nabla u\|_{H^1} + \|\nabla \rho\|_{L^2} + \|\nabla \nabla_{Rg}\|_{L^2(\mathcal{L}^2)}) ds.
 \end{aligned}$$

Define  $N(t) = \sup_{0 \leq s \leq t} (1+s)^{\frac{1}{2}} E_\eta(s)$ . According to (3.35), we get

$$\begin{aligned}
 (3.36) \quad N(t) & \leq C + C \sup_{0 \leq s \leq t} (1+s)^{-\frac{1}{2}} \int_0^s \|\rho\|_{L^2}^2 \|u\|_{L^2}^2 ds' \\
 & + C \int_0^t (\|u\|_{L^2}^2 + \|\rho\|_{L^2}^2) (\|\nabla u\|_{H^1} + \|\nabla \rho\|_{L^2} + \|\nabla \nabla_{Rg}\|_{L^2(\mathcal{L}^2)}) ds \\
 & \leq C + C \sup_{0 \leq s \leq t} \int_0^s \|\rho\|_{L^2}^2 \|u\|_{L^2}^2 (1+s')^{-\frac{1}{2}} ds' \\
 & + C \int_0^t N(s) (1+s)^{-\frac{1}{2}} (\|\nabla u\|_{H^1} + \|\nabla \rho\|_{L^2} + \|\nabla \nabla_{Rg}\|_{L^2(\mathcal{L}^2)}) ds \\
 & \leq C + C \int_0^t \|\rho\|_{L^2}^2 N(s) (1+s)^{-1} ds \\
 & + C \int_0^t N(s) (1+s)^{-\frac{1}{2}} (\|\nabla u\|_{H^1} + \|\nabla \rho\|_{L^2} + \|\nabla \nabla_{Rg}\|_{L^2(\mathcal{L}^2)}) ds.
 \end{aligned}$$

Applying Gronwall's inequality, Propositions 3.1 and 3.3, we obtain  $N(t) \leq C$ , which implies that

$$(3.37) \quad E_\eta(t) \leq C(1+t)^{-\frac{1}{2}}.$$

Applying Lemma 3.2, we have

$$(3.38) \quad \frac{d}{dt} E_\eta^1 + D_\eta^1 \leq 0,$$

which implies that

$$\begin{aligned}
 (3.39) \quad \frac{d}{dt} E_\eta^1 + \frac{C_2}{1+t} (\mu \|\Lambda^1 u\|_{H^{s-1}}^2 + \eta \gamma \|\Lambda^1 \rho\|_{H^{s-2}}^2) + \|\Lambda^1 \nabla_{Rg}\|_{H^{s-1}(\mathcal{L}^2)}^2 \\
 \leq \frac{CC_2}{1+t} \int_{S(t)} |\xi|^2 (|\hat{u}(\xi)|^2 + |\hat{\rho}(\xi)|^2) d\xi.
 \end{aligned}$$

According to (3.37), we have

$$\frac{CC_2}{1+t} \int_{S(t)} |\xi|^2 (|\hat{u}(\xi)|^2 + |\hat{\rho}(\xi)|^2) d\xi \leq CC_2^2 (1+t)^{-2} (\|\rho\|_{L^2}^2 + \|u\|_{L^2}^2) \leq C(1+t)^{-\frac{5}{2}}.$$

This together with (3.2), (3.37) and (3.39) ensure that

$$\begin{aligned}
 (1+t)^{\frac{5}{2}} E_\eta^1 & \leq C(1+t) + C \int_0^t \|\Lambda^s \rho\|_{L^2}^2 (1+s)^{\frac{3}{2}} ds \\
 & \leq C(1+t) + C \int_0^t E_\eta(s) (1+s)^{\frac{1}{2}} ds \\
 & \leq C(1+t),
 \end{aligned}$$

which implies that

$$(3.40) \quad E_\eta^1 \leq C(1+t)^{-\frac{3}{2}}.$$

Therefore, we complete the proof of Proposition 3.4.  $\square$

**Remark 3.5.** *The proposition 3.4 indicates that*

$$\|\rho\|_{L^2} + \|u\|_{L^2} \leq C(1+t)^{-\frac{1}{4}}.$$

*Combining with the incompressible FENE model and CNS system, one can see that this is not the optimal time decay.*

In order to obtain optimal  $L^2$  decay rate, we have to use the estimate of lower frequency. The following lemma show that the decay rate implies that the solution will belong to some Besov space with negative index.

**Lemma 3.6.** *Let  $0 < \alpha, \sigma \leq 1$  and  $\sigma < 2\alpha$ . Assume that  $(\rho_0, u_0, g_0)$  satisfy the condition in Theorem 1.1. For any  $t \in [0, +\infty)$ , if*

$$(3.41) \quad E_\eta(t) \leq C(1+t)^{-\alpha}, \quad E_\eta^1(t) \leq C(1+t)^{-\alpha-1} \quad \text{and} \quad \int_0^t (1+s)^\alpha \|\Lambda^1 \nabla_R g\|_{L^2(\mathcal{L}^2)}^2 ds \leq C,$$

then we have

$$(3.42) \quad (\rho, u, g) \in L^\infty(0, \infty; \dot{B}_{2,\infty}^{-\sigma}) \times L^\infty(0, \infty; \dot{B}_{2,\infty}^{-\sigma}) \times L^\infty(0, \infty; \dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)).$$

*Proof.* Applying  $\dot{\Delta}_j$  to (1.2), we get

$$(3.43) \quad \begin{cases} \dot{\Delta}_j \rho_t + \operatorname{div} \dot{\Delta}_j u = \dot{\Delta}_j F, \\ \dot{\Delta}_j u_t - \operatorname{div} \Sigma(\dot{\Delta}_j u) + \gamma \nabla \dot{\Delta}_j \rho - \operatorname{div} \dot{\Delta}_j \tau = \dot{\Delta}_j G, \\ \dot{\Delta}_j g_t + \mathcal{L} \dot{\Delta}_j g - \nabla \dot{\Delta}_j u R_j \partial_{R_k} \mathcal{U} + \operatorname{div} \dot{\Delta}_j u = \dot{\Delta}_j H, \end{cases}$$

where  $F = -\operatorname{div}(\rho u)$ ,  $G = -u \cdot \nabla u + [i(\rho) - 1](\operatorname{div} \Sigma(u) + \operatorname{div} \tau) + [\gamma - h(\rho)] \nabla \rho$  and  $H = -u \cdot \nabla g - \frac{1}{\psi_\infty} \nabla_R \cdot (\nabla u R g \psi_\infty)$ .

Using  $\int_B \dot{\Delta}_j g \psi_\infty dR = 0$  and integrating by parts, then we have

$$(3.44) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} (\gamma \|\dot{\Delta}_j \rho\|_{L^2}^2 + \|\dot{\Delta}_j u\|_{L^2}^2 + \|\dot{\Delta}_j g\|_{L^2(\mathcal{L}^2)}^2) \\ & + \mu \|\nabla \dot{\Delta}_j u\|_{L^2}^2 + (\mu + \mu') \|\operatorname{div} \dot{\Delta}_j u\|_{L^2}^2 + \|\nabla_R \dot{\Delta}_j g\|_{L^2(\mathcal{L}^2)}^2 \\ & = \int_{\mathbb{R}^2} \gamma \dot{\Delta}_j F \dot{\Delta}_j \rho dx + \int_{\mathbb{R}^2} \dot{\Delta}_j G \dot{\Delta}_j u dx + \int_{\mathbb{R}^2} \int_B \dot{\Delta}_j H \dot{\Delta}_j g \psi_\infty dx dR \\ & \leq C(\|\dot{\Delta}_j F\|_{L^2} \|\dot{\Delta}_j \rho\|_{L^2} + \|\dot{\Delta}_j G\|_{L^2} \|\dot{\Delta}_j u\|_{L^2}) \\ & + C(\|\dot{\Delta}_j(u \nabla g)\|_{L^2(\mathcal{L}^2)}^2 + \|\dot{\Delta}_j(\nabla u R g)\|_{L^2(\mathcal{L}^2)}^2) + \frac{1}{2} \|\nabla_R \dot{\Delta}_j g\|_{L^2(\mathcal{L}^2)}^2. \end{aligned}$$

Multiplying both sides of (3.44) by  $2^{-2j\sigma}$  and taking  $l^\infty$  norm, we get

$$(3.45) \quad \frac{d}{dt} (\gamma \|\rho\|_{\dot{B}_{2,\infty}^{-\sigma}}^2 + \|u\|_{\dot{B}_{2,\infty}^{-\sigma}}^2 + \|g\|_{\dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)}^2)$$

$$\leq C(\|F\|_{\dot{B}_{2,\infty}^{-\sigma}} \|\rho\|_{\dot{B}_{2,\infty}^{-\sigma}} + \|G\|_{\dot{B}_{2,\infty}^{-\sigma}} \|u\|_{\dot{B}_{2,\infty}^{-\sigma}} + \|\nabla u Rg\|_{\dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)}^2 + \|u \nabla g\|_{\dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)}^2).$$

Define  $M(t) = \sum_{0 \leq s \leq t} \|\rho\|_{\dot{B}_{2,\infty}^{-\sigma}} + \|u\|_{\dot{B}_{2,\infty}^{-\sigma}} + \|g\|_{\dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)}$ . According to (3.45), we deduce that

$$(3.46) \quad \begin{aligned} M^2(t) &\leq M^2(0) + CM(t) \int_0^t \|F\|_{\dot{B}_{2,\infty}^{-\sigma}} + \|G\|_{\dot{B}_{2,\infty}^{-\sigma}} ds \\ &\quad + C \int_0^t \|\nabla u Rg\|_{\dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)}^2 + \|u \nabla g\|_{\dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)}^2 ds. \end{aligned}$$

Using Lemmas 2.2, 2.3 and (3.41), we obtain

$$\begin{aligned} \int_0^t \|\nabla u Rg\|_{\dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)}^2 + \|u \nabla g\|_{\dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)}^2 ds &\leq C \int_0^t \|\nabla u Rg\|_{L^{\frac{2}{\sigma+1}}(\mathcal{L}^2)}^2 + \|u \nabla g\|_{L^{\frac{2}{\sigma+1}}(\mathcal{L}^2)}^2 ds \\ &\leq C \int_0^t \|\nabla u\|_{L^2}^2 \|g\|_{L^{\frac{2}{\sigma}}(\mathcal{L}^2)}^2 + \|\nabla g\|_{L^2(\mathcal{L}^2)}^2 \|u\|_{L^{\frac{2}{\sigma}}}^2 ds \leq C, \end{aligned}$$

and

$$\begin{aligned} \int_0^t \|F\|_{\dot{B}_{2,\infty}^{-\sigma}} ds &\leq C \int_0^t \|F\|_{L^{\frac{2}{\sigma+1}}} ds \\ &\leq C \int_0^t \|u\|_{L^{\frac{2}{\sigma}}} \|\nabla \rho\|_{L^2} + \|\operatorname{div} u\|_{L^2} \|\rho\|_{L^{\frac{2}{\sigma}}} ds \\ &\leq C \int_0^t \|u\|_{L^2}^\sigma \|\nabla u\|_{L^2}^{1-\sigma} \|\nabla \rho\|_{L^2} + \|\operatorname{div} u\|_{L^2} \|\rho\|_{L^2}^\sigma \|\nabla \rho\|_{L^2}^{1-\sigma} ds \\ &\leq C \int_0^t (1+s)^{-(1+\alpha-\frac{\sigma}{2})} ds \leq C. \end{aligned}$$

By virtue of (3.41), we similarly get

$$(3.47) \quad \begin{aligned} \int_0^t \|G\|_{\dot{B}_{2,\infty}^{-\sigma}} ds &\leq C \int_0^t (1+s)^{-(1+\alpha-\frac{\sigma}{2})} ds + C \int_0^t \|\operatorname{div} \tau\|_{L^2} \|\rho\|_{L^{\frac{2}{\sigma}}} ds \\ &\leq C + C \int_0^t \|\rho\|_{L^2}^\sigma \|\nabla \rho\|_{L^2}^{1-\sigma} \|\nabla \nabla Rg\|_{L^2(\mathcal{L}^2)} ds \\ &\leq C + C \left( \int_0^t \|\rho\|_{L^2}^{2\sigma} \|\nabla \rho\|_{L^2}^{2-2\sigma} (1+s)^{-\alpha} ds \right)^{\frac{1}{2}} \left( \int_0^t (1+s)^\alpha \|\nabla \nabla Rg\|_{L^2(\mathcal{L}^2)}^2 ds \right)^{\frac{1}{2}} \\ &\leq C + C \left( \int_0^t (1+s)^{-(1+2\alpha-\sigma)} ds \right)^{\frac{1}{2}} \leq C. \end{aligned}$$

From (3.46), we have  $M^2(t) \leq CM^2(0) + M(t)C + C$ . By virtue of interpolation theory, we can deduce that  $(\rho_0, u_0, g_0) \in \dot{B}_{2,\infty}^{-\sigma} \times \dot{B}_{2,\infty}^{-\sigma} \times \dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)$  with  $0 < \sigma \leq 1$ . Then  $M^2(0) \leq C$  implies that  $M(t) \leq C$ .  $\square$

Using the above lemma one can prove that the solution belongs to some Besov space. The following proposition indicates that if the solution belongs to some Besov space with negative index, then the decay rate can be improved.

**Proposition 3.7.** *Let  $0 < \beta, \sigma \leq 1$  and  $\frac{1}{2} \leq \alpha$ . Assume that  $(\rho_0, u_0, g_0)$  satisfy the condition in Theorem 1.1. For any  $t \in [0, +\infty)$ , if*

$$(3.48) \quad E_\eta(t) \leq C(1+t)^{-\alpha}, \quad E_\eta^1(t) \leq C(1+t)^{-\alpha-1} \quad \text{and} \quad \int_0^t (1+s)^{\frac{3}{4}} \|\Lambda^1 \nabla Rg\|_{L^2(\mathcal{L}^2)}^2 ds \leq C,$$



and

$$(3.49) \quad (\rho, u, g) \in L^\infty(0, \infty; \dot{B}_{2,\infty}^{-\sigma}) \times L^\infty(0, \infty; \dot{B}_{2,\infty}^{-\sigma}) \times L^\infty(0, \infty; \dot{B}_{2,\infty}^{-\sigma}(\mathcal{L}^2)),$$

then there exists a constant  $C$  such that

$$(3.50) \quad E_\eta(t) \leq C(1+t)^{-\beta} \quad \text{and} \quad E_\eta^1(t) \leq C(1+t)^{-\beta-1}$$

where  $\beta < \frac{\sigma+1}{2}$  for  $\alpha = \frac{1}{2}$  and  $\beta = \frac{\sigma+1}{2}$  for  $\alpha > \frac{1}{2}$ .

*Proof.* According to (3.48), we obtain

$$\int_{S(t)} \int_0^t |\widehat{\rho u}|^2 ds d\xi \leq C(1+t)^{-1} \int_0^t \|\rho\|_{L^2}^2 \|u\|_{L^2}^2 ds \leq C(1+t)^{-1} \int_0^t (1+s)^{-2\alpha} ds \leq C(1+t)^{-\beta}.$$

By virtue of (3.48) and (3.49), we have

$$\begin{aligned} \int_{S(t)} \int_0^t |\hat{G} \cdot \hat{u}| ds d\xi &\leq C \int_0^t \|G\|_{L^1} \int_{S(t)} |\hat{u}| d\xi ds \\ &\leq C(1+t)^{-\frac{1}{2}} \int_0^t \|G\|_{L^1} \left( \int_{S(t)} |\hat{u}|^2 d\xi \right)^{\frac{1}{2}} ds \\ &\leq C(1+t)^{-\frac{\sigma+1}{2}} M(t) \int_0^t \|G\|_{L^1} ds \\ &\leq C(1+t)^{-\frac{\sigma+1}{2}} \int_0^t (1+s)^{-\alpha-\frac{1}{2}} + \|\rho\|_{L^2} \|\nabla \nabla_{Rg}\|_{L^2(\mathcal{L}^2)} ds \\ &\leq C(1+t)^{-\beta} + C(1+t)^{-\frac{\sigma+1}{2}} \left( \int_0^t \|\rho\|_{L^2}^2 (1+s)^{-\frac{3}{4}} ds \right)^{\frac{1}{2}} \\ &\leq C(1+t)^{-\beta}. \end{aligned}$$

Similar to the proof of Proposition 3.4, we obtain

$$(3.51) \quad \frac{d}{dt} E_\eta(t) + \frac{\mu C_2}{1+t} \|u\|_{H^s}^2 + \frac{\eta \gamma C_2}{1+t} \|\rho\|_{H^{s-1}}^2 + \|\nabla_{Rg}\|_{H^s(\mathcal{L}^2)}^2 \leq \frac{CC_2}{1+t} (1+t)^{-\beta}.$$

Then the proof of (3.40) implies that

$$(3.52) \quad E_\eta(t) \leq C(1+t)^{-\beta}.$$

Using Lemma 3.2, we have

$$(3.53) \quad \begin{aligned} \frac{d}{dt} E_\eta^1 + \frac{C_1}{1+t} (\mu \|\Lambda^1 u\|_{H^{s-1}}^2 + \eta \gamma \|\Lambda^1 \rho\|_{H^{s-2}}^2) + \|\Lambda^1 \nabla_{Rg}\|_{H^{s-1}(\mathcal{L}^2)}^2 \\ \leq \frac{CC_1}{1+t} \int_{S(t)} |\xi|^2 (|\hat{u}(\xi)|^2 + |\hat{\rho}(\xi)|^2) d\xi. \end{aligned}$$

According to (3.52), we have

$$(3.54) \quad \frac{CC_2}{1+t} \int_{S(t)} |\xi|^2 (|\hat{u}(\xi)|^2 + |\hat{\rho}(\xi)|^2) d\xi \leq CC_2^2 (1+t)^{-2} (\|\rho\|_{L^2}^2 + \|u\|_{L^2}^2) \leq C(1+t)^{-2-\beta}.$$

Then the proof of (3.40) implies that  $E_\eta^1 \leq C(1+t)^{-1-\beta}$ . We thus finish the proof of Proposition 3.7.  $\square$

**The proof of Theorem 1.1:**

We now improve the decay rate in Proposition 3.4. According to Propositions 3.3, 3.4 and Lemma 3.6 with  $\sigma = \alpha = \frac{1}{2}$ , we obtain

$$(\rho, u, g) \in L^\infty(0, \infty; \dot{B}_{2, \infty}^{-\frac{1}{2}}) \times L^\infty(0, \infty; \dot{B}_{2, \infty}^{-\frac{1}{2}}) \times L^\infty(0, \infty; \dot{B}_{2, \infty}^{-\frac{1}{2}}(\mathcal{L}^2)).$$

Taking advantage of Propositions 3.7 with  $\alpha = \sigma = \frac{1}{2}$  and  $\beta = \frac{5}{8}$ , we deduce that

$$E_\eta(t) \leq C(1+t)^{-\frac{5}{8}} \quad \text{and} \quad E_\eta^1(t) \leq C(1+t)^{-\frac{5}{8}-1}.$$

According to Proposition 3.3 and Lemma 3.6 with  $\sigma = 1$  and  $\alpha = \frac{5}{8}$ , we obtain

$$(\rho, u, g) \in L^\infty(0, \infty; \dot{B}_{2, \infty}^{-1}) \times L^\infty(0, \infty; \dot{B}_{2, \infty}^{-1}) \times L^\infty(0, \infty; \dot{B}_{2, \infty}^{-1}(\mathcal{L}^2)).$$

Using Propositions 3.7 again with  $\alpha = \frac{5}{8}$  and  $\sigma = \beta = 1$ , we verify that

$$E_\eta(t) \leq C(1+t)^{-1} \quad \text{and} \quad E_\eta^1(t) \leq C(1+t)^{-2}.$$

To get the faster decay rate for  $g$  in  $L^2$ , we need the following standard energy estimation to (1.2):

$$\frac{1}{2} \frac{d}{dt} \|g\|_{L^2(\mathcal{L}^2)}^2 + \|\nabla_R g\|_{L^2(\mathcal{L}^2)}^2 \lesssim - \int_{\mathbb{R}^d} \nabla u : \tau dx + \|\nabla u\|_{L^\infty} \|g\|_{L^2(\mathcal{L}^2)} \|\nabla_R g\|_{L^2(\mathcal{L}^2)}.$$

By Lemma 2.4, Lemma 2.5, we have

$$\int_{\mathbb{R}^d} \nabla u : \tau dx \leq \delta \|\nabla_R g\|_{L^2(\mathcal{L}^2)}^2 + C_\delta \|\nabla u\|_{L^2}^2.$$

Using Lemma 2.4 and Theorem 2.6, we deduce that

$$\frac{d}{dt} \|g\|_{L^2(\mathcal{L}^2)}^2 + \|g\|_{L^2(\mathcal{L}^2)}^2 \leq C_\delta \|\nabla u\|_{L^2}^2,$$

which implies that

$$\begin{aligned} \|g\|_{L^2(\mathcal{L}^2)}^2 &\leq \|g_0\|_{L^2(\mathcal{L}^2)}^2 e^{-t} + C \int_0^t e^{-(t-s)} \|\nabla u\|_{L^2}^2 ds \\ &\leq C(e^{-t} + \int_0^t e^{-(t-s)} (1+s)^{-2} ds) \\ &\leq C(1+t)^{-2}. \end{aligned}$$

We thus complete the proof of Theorem 1.1. □

**Remark 3.8.** *One can see that the decay rate for the  $\dot{H}^1$ -norm obtained in Propositions 3.4 is not optimal. However, in the proof of Theorem 1.1, we improve the decay rate to  $(1+t)^{-1}$ .*

**Remark 3.9.** *In Theorem 1.1, we only obtain the optimal decay rate with  $d = 2$ . The decay rate for  $d = 1$  is an interesting problem. However, the technique in this paper fail to obtain the optimal decay rate when  $d = 1$ . We are going to study about this problem in the future.*

**Acknowledgments** This work was partially supported by the National Natural Science Foundation of China (No.11671407 and No.11701586), the Macao Science and Technology Development Fund (No. 098/2013/A3), and Guangdong Province of China Special Support Program (No. 8-2015), and the key project of the Natural Science Foundation of Guangdong province (No. 2016A030311004).

## References

- [1] H. Bahouri, J.-Y. Chemin, and R. Danchin. *Fourier analysis and nonlinear partial differential equations*, volume 343 of *Grundlehren der Mathematischen Wissenschaften*. Springer, Heidelberg, 2011.
- [2] R. B. Bird, R. C. Armstrong, and O. Hassager. *Dynamics of Polymeric Liquids*, volume 1. Wiley, New York, 1977.
- [3] D. Bresch and B. Desjardins. Existence of global weak solutions for a 2d viscous shallow water equations and convergence to the quasi-geostrophic model. *Communications in Mathematical Physics.*, 238(1-2):211–233, 2003.
- [4] D. Bresch, B. Desjardins, and C. K. Lin. On some compressible fluid models: Korteweg, lubrication, and shallow water systems. *Communications in Partial Differential Equations.*, 28(3-4):843–868, 2003.
- [5] A. V. Busuioc, I. S. Ciuperca, D. Iftimie, and L. I. Palade. The FENE dumbbell polymer model: existence and uniqueness of solutions for the momentum balance equation. *J. Dynam. Differential Equations*, 26(2):217–241, 2014.
- [6] Q. Chen, C. Miao, and Z. Zhang. Well-posedness in critical spaces for the compressible Navier-Stokes equations with density dependent viscosities. *Revista Matemática Iberoamericana.*, 26(3):915–946, 2010.
- [7] Q. Chen, C. Miao, and Z. Zhang. On the ill-posedness of the compressible Navier-Stokes equations in the critical besov spaces. *Revista Matemática Iberoamericana.*, 31(4):165–186, 2011.
- [8] R. Danchin. Global existence in critical spaces for compressible Navier-Stokes equations. *Inventiones Mathematicae.*, 141(3):579–614, 2000.
- [9] R. Danchin. Local theory in critical spaces for compressible viscous and heat-conductive gases. *Comm Partial Differential Equations.*, 141(26):1183–1233, 2001.
- [10] R. Danchin. On the uniqueness in critical spaces for compressible Navier-Stokes equations. *Nonlinear Differential Equations Applications Nodda.*, 12(1):111–128, 2005.
- [11] R. Danchin. Well-posedness in critical spaces for barotropic viscous fluids with truly not constant density. *Communications in Partial Differential Equations.*, 32(9):111–128, 2007.
- [12] R. Danchin and X. Jiang. Optimal time-decay estimates for the compressible Navier-Stokes equations in the critical  $L^p$  framework. *Archive for Rational Mechanics and Analysis*, 2016.
- [13] R. Danchin and P. B. Mucha. A lagrangian approach for the incompressible Navier-Stokes equations with variable density. *Communications on Pure and Applied Mathematics.*, 65(10), 2012.
- [14] M. Doi and S. F. Edwards. *The Theory of Polymer Dynamics*. Oxford University Press, Oxford, 1988.

- 
- [15] L. He, J. Huang, and C. Wang. Global stability of large solutions to the 3d compressible Navier-Stokes equations. *Archive for Rational Mechanics and Analysis.*, 234(3):1167–1222, 2019.
- [16] L. He and P. Zhang.  $L^2$  Decay of Solutions to a Micro-Macro Model for Polymeric Fluids Near Equilibrium. *Siam Journal on Mathematical Analysis*, 40(5):1905–1922, 2009.
- [17] X. Huang and J. Li. Existence and blowup behavior of global strong solutions to the two-dimensional barotropic compressible Navier-Stokes system with vacuum and large initial data. *J. Math. Pures Appl. (9)*, 106(1):123–154, 2016.
- [18] X. Huang, J. Li, and Z. Xin. Global well-posedness of classical solutions with large oscillations and vacuum to the three-dimensional isentropic compressible Navier-Stokes equations. *Communications on Pure Applied Mathematics.*, 234, 2011.
- [19] Y. Jia and H. Ye. Optimal decay rates of weak solutions for the 2D co-rotation FENE dumbbell model of polymeric flows. *Appl. Math. Lett.*, 113:106860, 8, 2021.
- [20] N. Jiang, Y. Liu, and T.-F. Zhang. Global classical solutions to a compressible model for micro-macro polymeric fluids near equilibrium. *SIAM J. Math. Anal.*, 50(4):4149–4179, 2018.
- [21] Q. Jiu, Y. Wang, and Z. Xin. Global well-posedness of 2d compressible Navier-Stokes equations with large data and vacuum. *Journal of Mathematical Fluid Mechanics.*, 16(3):483–521, 2014.
- [22] B. Jourdain, T. Lelièvre, and C. Le Bris. Existence of solution for a micro-macro model of polymeric fluid: the FENE model. *J. Funct. Anal.*, 209(1):162–193, 2004.
- [23] A. V. Kazhikhov and V. V. Shelukhin. Unique global solution with respect to time of initial-boundary value problems for one-dimensional equations of a viscous gas. *J. Appl. Math. Mech.*, 41(2):273–282., 1977.
- [24] H. Li, J. Li, and Z. Xin. Vanishing of vacuum states and blow-up phenomena of the compressible Navier-Stokes equations. *Communications in Mathematical Physics.*, 281(2):401, 2008.
- [25] H.-L. Li and T. Zhang. Large time behavior of isentropic compressible Navier-Stokes system in  $\mathbb{R}^3$ . *Math. Methods Appl. Sci.*, 34(6):670–682, 2011.
- [26] F. Lin, P. Zhang, and Z. Zhang. On the global existence of smooth solution to the 2-D FENE dumbbell model. *Comm. Math. Phys.*, 277(2):531–553, 2008.
- [27] W. Luo and Z. Yin. The Liouville Theorem and the  $L^2$  Decay for the FENE Dumbbell Model of Polymeric Flows. *Arch. Ration. Mech. Anal.*, 224(1):209–231, 2017.
- [28] W. Luo and Z. Yin. The  $L^2$  decay for the 2D co-rotation FENE dumbbell model of polymeric flows. *Adv. Math.*, 343:522–537, 2019.
- [29] Z. Luo, W. Luo, and Z. Yin. Global strong solutions and large time behavior to the compressible co-rotation FENE dumbbell model of polymeric flows near equilibrium. arXiv:2104.10844.
- [30] Z. Luo, W. Luo, and Z. Yin. Global strong solutions and optimal  $L^2$  decay to the compressible fene dumbbell model. arXiv:2107.08646.

- 
- [31] N. Masmoudi. Well-posedness for the FENE dumbbell model of polymeric flows. *Comm. Pure Appl. Math.*, 61(12):1685–1714, 2008.
- [32] N. Masmoudi. Global existence of weak solutions to the FENE dumbbell model of polymeric flows. *Invent. Math.*, 191(2):427–500, 2013.
- [33] N. Masmoudi. *Equations for Polymeric Materials*. Handbook of Mathematical Analysis in Mechanics of Viscous Fluids, 2016.
- [34] A. Matsumura and T. Nishida. The initial value problem for the equations of motion of compressible viscous and heat-conductive fluids. *Proc. Japan Acad. Ser. A Math. Sci.*, 55:337–342, 1979.
- [35] A. Mellet and A. Vasseur. On the barotropic compressible Navier-Stokes equations. *Proc. Japan Acad. Ser. A Math. Sci.*, 32(3):431–452, 2007.
- [36] J. Nash. Le probleme de cauchy pour les equations differentielles d’un fluide general. *Bull.soc.math.france.*, 90(4):487–497, 1962.
- [37] L. Nirenberg. On elliptic partial differential equations. *Ann.scuola Norm.sup.pisa*, 1959.
- [38] M. Renardy. An existence theorem for model equations resulting from kinetic theories of polymer solutions. *SIAM J. Math. Anal.*, 22(2):313–327, 1991.
- [39] M. E. Schonbek.  $L^2$  decay for weak solutions of the Navier-Stokes equations. *Arch. Rational Mech. Anal.*, 88(3):209–222, 1985.
- [40] M. E. Schonbek. Existence and decay of polymeric flows. *SIAM J. Math. Anal.*, 41(2):564–587, 2009.
- [41] L. Tong, Z. Tan, and Y. Wang. The asymptotic behavior of globally smooth solutions to the compressible magnetohydrodynamic equations with Coulomb force. *Analysis and Applications*, 2017.
- [42] A. F. Vasseur and C. Yu. Existence of global weak solutions for 3D degenerate compressible Navier-Stokes equations. *Invent. Math.*, 206(3):935–974, 2016.
- [43] Z. Xin. Blowup of smooth solutions to the compressible Navier-Stokes equation with compact density. *Communications on Pure Applied Mathematics.*, 1998.
- [44] Z. Xin and J. Xu. Optimal decay for the compressible Navier-Stokes equations without additional smallness assumptions. 2018.
- [45] Z. Xin and W. Yan. On blow-up of classical solutions to the compressible Navier-Stokes equations. *Communications in Mathematical Physics.*, 321(2):529–541, 2013.
- [46] J. Xu. A low-frequency assumption for optimal time-decay estimates to the compressible Navier-Stokes equations. *Comm. Math. Phys.*, 371(2):525–560, 2019.
- [47] H. Zhang and P. Zhang. Local existence for the FENE-dumbbell model of polymeric fluids. *Arch. Ration. Mech. Anal.*, 181(2):373–400, 2006.

- [48] Z. Zhou, C. Zhu, and R. Zi. Global well-posedness and decay rates for the three dimensional compressible Oldroyd-B model. *Journal of Differential Equations*, 265(4), 2018.