IDEAL STRUCTURE OF C^* -ALGEBRAS OF SINGLY GENERATED DYNAMICAL SYSTEMS

TAKESHI KATSURA

ABSTRACT. In this paper, we show that the set of all ideals of the C^* -algebras of a singly generated dynamical system corresponds bijectively to the set of all subsets of the product of the space of the system and the circle satisfying three conditions.

0. Introduction

In the theory of C^* -algebras, it is very important yet very difficult to list all ideals of a given C^* -algebra. In this paper, we list all ideals of a C^* -algebra of a singly generated dynamical system in terms of subsets of a certain space.

In [R], Renault introduces the notion of a singly generated dynamical system (SGDS) and associates a C^* -algebra with it. In [K1], the author introduces the notion of a topological graph and associates a C^* -algebra with it. In [K2], the author shows that an SGDS can be considered as a topological graph and their C^* -algebras coincide. Conversely in [K4] every C^* -algebra of a topological graph is a C^* -algebra of some SGDS (see also [Y] and [KL]). Thus the class of C^* -algebras of SGDSs is fairly large. Although in [R] spaces in SGDSs are assumed to be second countable, there is no such restriction in [K1]. In this paper, spaces are not assumed to be second countable, and hence the associated C^* -algebras are not necessarily separable.

The purpose of this paper is to investigate the ideal structure of C^* -algebras of SGDSs. Since we do not assume second countability, it is almost impossible to list primitive ideals of the associated C^* -algebra in terms of points of the space. Instead in this paper, we list all ideals directly in terms of subsets of the product of the space and the circle (Theorem 7.8). The investigation here may be useful to describe the topology of primitive ideal space when the space is second countable (cf. [SW]).

This paper is organized as follows. In Section 1, we introduce SGDSs and their C^* -algebras. To define a C^* -algebra of an SGDS, we do not use groupoids as in [R], but use a definition similar to [K1]. We examine the structure of a C^* -algebra of an SGDS which we need later. In Section 2, we construct lots of irreducible representations of a C^* -algebra of an SGDS. In Section 3, we associate a subset of a space to each ideal, and an ideal to each subset. Then, we list all gauge-invariant ideals in terms of invariant subsets (Proposition 3.11). In Section 4, we list all prime ideals, and show that the set of primitive ideals constructed in Section 2 is sufficiently large (Corollary 4.19). In Section 5, we associate a subset of the product of a space and the circle to each ideal, and an ideal to each subset. We show that ideals of the C^* -algebra of an SGDS correspond injectively to a subset of the product of a

space and the circle (Proposition 5.5). In Section 6, we show that this subset is an admissible set which is defined in Definition 5.7 (Proposition 6.8). In Section 7, we show that the set of all ideals of the C^* -algebra of an SGDS corresponds bijectively to the set of all admissible sets (Theorem 7.8).

We try to make this paper self-contained as well as possible. However, in Section 4 we need two kinds of uniqueness theorems for C^* -algebras of topological graphs which are quoted in Appendix A.

Acknowledgments. This work was supported by JSPS KAKENHI Grant Number JP18K03345.

1. Singly generated dynamical systems

Definition 1.1. A pair $\Sigma = (X, \sigma)$ is a singly generated dynamical system (in short SGDS) if X is a locally compact space and σ is a local homeomorphism from an open subset U of X to X.

It should be emphasized that X is not necessarily second countable. Throughout this paper, $\Sigma = (X, \sigma)$ means an SGDS. We denote by $\mathbb{N} := \{0, 1, 2, \ldots\}$ the set of natural numbers.

Definition 1.2. We define a decreasing sequence $\{U_n\}_{n\in\mathbb{N}}$ of open subsets of X by $U_0 = X$, $U_1 = U$ and $U_{n+1} = \{x \in U_n \mid \sigma^n(x) \in U\}$.

The open set U_n is the domain of σ^n , and $\sigma^n : U_n \to X$ is a local homeomorphism.

Definition 1.3. For $\xi, \eta \in C_c(U)$, we define $\langle \xi, \eta \rangle \in C_c(X)$ by

$$\langle \xi, \eta \rangle(x) = \sum_{\sigma(y)=x} \overline{\xi}(y)\eta(y)$$

for $x \in X$, where $\overline{\xi} \in C_c(U)$ is defined by $\overline{\xi}(y) = \overline{\xi(y)}$ for $y \in U$.

In the expression above, we implicitly assume that y is in U. We use this convention. Namely if we write $\sigma^n(x) = \cdots$, we assume $x \in U_n$. The fact that $\langle \xi, \eta \rangle$ is in $C_c(X)$ can be shown by using the fact that σ is a local homeomorphism (see [K1, Lemma 1.5]).

Definition 1.4. The C^* -algebra $C^*(\Sigma)$ of an SGDS $\Sigma = (X, \sigma)$ is the universal C^* -algebra generated by the images of a *-homomorphism $t^0: C_0(X) \to C^*(\Sigma)$ and a linear map $t^1: C_c(U) \to C^*(\Sigma)$ satisfying

- (i) $t^1(\xi)^*t^1(\eta) = t^0(\langle \xi, \eta \rangle)$ for $\xi, \eta \in C_c(U)$,
- (ii) $t^1(\xi)t^1(\eta)^* = t^0(\xi\overline{\eta})$ for $\xi, \eta \in C_c(V)$ where $V \subset U$ is an open subset on which σ is injective.

We call the pair (t^0, t^1) in Definition 1.4 the *universal* pair for $C^*(\Sigma)$. Hereafter, we investigate properties of the universal pair (t^0, t^1) for $C^*(\Sigma)$. For $f \in C_0(X)$ and $\xi \in C_c(U)$, we have $t^1(\xi)t^0(f) = t^1(\xi(f \circ \sigma))$ because one can see $d^*d = 0$ for $d = t^1(\xi)t^0(f) - t^1(\xi(f \circ \sigma))$ by the computation

$$t^{1}(\eta)^{*} \left(t^{1}(\xi) t^{0}(f) - t^{1}(\xi(f \circ \sigma)) \right) = t^{0} (\langle \eta, \xi \rangle) t^{0}(f) - t^{0} (\langle \eta, \xi(f \circ \sigma) \rangle)$$
$$= t^{0} \left(\langle \eta, \xi \rangle f - \langle \eta, \xi(f \circ \sigma) \rangle \right) = 0$$

for arbitrary $\eta \in C_c(U)$. For $f \in C_0(X)$ and $\xi \in C_c(U)$, we have $t^0(f)t^1(\xi) = t^1(f\xi)$. To see this, first we may assume $\xi \in C_c(V)$ where $V \subset U$ is an open subset on which σ is injective by the partition of unity because the fact that σ is locally homeomorphic implies that the set of such V covers U. Now for arbitrary $\eta \in C_c(V)$, we have

$$(t^{0}(f)t^{1}(\xi) - t^{1}(f\xi))t^{1}(\eta)^{*} = t^{0}(f)t^{0}(\xi\overline{\eta}) - t^{0}(f\xi\overline{\eta}) = 0.$$

Therefore we get $dd^* = 0$ for $d = t^0(f)t^1(\xi) - t^1(f\xi)$. We have shown that $t^0(f)t^1(\xi) = t^1(f\xi)$ for all $f \in C_0(X)$ and $\xi \in C_c(U)$. Finally we see that for $f \in C_c(U)$ there exist $\xi_1, \ldots, \xi_n, \eta_1, \ldots, \eta_n \in C_c(U)$ such that

$$t^{0}(f) = \sum_{k=1}^{n} t^{1}(\xi_{k})t^{1}(\eta_{k})^{*}.$$

By the partition of unity, there exist $f_k \in C_c(V_k)$ for k = 1, ..., n such that $f = \sum_{k=1}^n f_k$ where $V_k \subset U$ is an open subset on which σ is injective for k = 1, ..., n. Then for k = 1, ..., n one can find $\xi_k, \eta_k \in C_c(V_k)$ such that $\xi_k \overline{\eta_k} = f_k$ (for example one can take $\eta_k = |f_k|^{1/2}$). Then we have

$$t^{0}(f) = \sum_{k=1}^{n} t^{0}(f_{k}) = \sum_{k=1}^{n} t^{1}(\xi_{k})t^{1}(\eta_{k})^{*}.$$

From these computation, one can see that the C^* -algebra $C^*(\Sigma)$ is the C^* -algebra of the topological graph $E = (X, U, \sigma, \iota)$, where $\iota \colon U \to X$ is the embedding, as defined in [K1, Definition 2.10]. Thus by [K2, Proposition 10.9], the C^* -algebra $C^*(\Sigma)$ is isomorphic to the C^* -algebra defined in [R] when X is second countable.

We try to make this paper self-contained as well as possible. We quote results in [K1] which is used to prove the main theorem (Theorem 7.8) only in Appendix A. We know that both t^0 and t^1 are injective by [K1, Proposition 3.7]. This fact is reproved in Lemma 2.8.

Definition 1.5. Let $n \in \mathbb{N}$. For $\xi, \eta \in C_c(U_n)$, we define $\langle \xi, \eta \rangle_n \in C_c(X)$ by

$$\langle \xi, \eta \rangle_n(x) = \sum_{\sigma^n(y)=x} \overline{\xi}(y)\eta(y)$$

for $x \in X$.

Note that $\langle \xi, \eta \rangle_0 = \overline{\xi} \eta$ for $\xi, \eta \in C_c(X)$, and $\langle \xi, \eta \rangle_1 = \langle \xi, \eta \rangle$ for $\xi, \eta \in C_c(U)$.

Lemma 1.6. Let $n, m \in \mathbb{N}$. For $\xi_1, \eta_1 \in C_c(U_n)$ and $\xi_2, \eta_2 \in C_c(U_m)$, define $\xi, \eta \in C_c(U_{n+m})$ by $\xi(x) = \xi_1(x)\xi_2(\sigma^n(x))$ and $\eta(x) = \eta_1(x)\eta_2(\sigma^n(x))$ for $x \in U_{n+m}$. Then we have

$$\langle \xi_2, \langle \xi_1, \eta_1 \rangle_n \eta_2 \rangle_m = \langle \xi, \eta \rangle_{n+m}$$

Proof. For $x \in X$, we have

$$\langle \xi_{2}, \langle \xi_{1}, \eta_{1} \rangle_{n} \eta_{2} \rangle_{m}(x) = \sum_{\sigma^{m}(y)=x} \overline{\xi_{2}}(y) \langle \xi_{1}, \eta_{1} \rangle_{n}(y) \eta_{2}(y)$$

$$= \sum_{\sigma^{m}(y)=x} \overline{\xi_{2}}(y) \Big(\sum_{\sigma^{n}(z)=y} \overline{\xi_{1}}(z) \eta_{1}(z) \Big) \eta_{2}(y)$$

$$= \sum_{\sigma^{m}(y)=x} \sum_{\sigma^{n}(z)=y} \overline{\xi_{2}}(\sigma^{n}(z)) \overline{\xi_{1}}(z) \eta_{1}(z) \eta_{2}(\sigma^{n}(z))$$

$$= \sum_{\sigma^{n+m}(z)=x} \overline{\xi}(z) \eta(z)$$

$$= \langle \xi, \eta \rangle_{n+m}(x).$$

Lemma 1.7. Let $n \in \mathbb{N}$. For $\xi_1, \ldots, \xi_n, \eta_1, \ldots, \eta_n \in C_c(U)$, define $\xi, \eta \in C_c(U_n)$ by

$$\xi(x) = \xi_1(x)\xi_2(\sigma(x))\cdots\xi_n(\sigma^{n-1}(x))$$

$$\eta(x) = \eta_1(x)\eta_2(\sigma(x))\cdots\eta_n(\sigma^{n-1}(x))$$

for $x \in U_n$. Then we have

$$\langle \xi_n, \langle \xi_{n-1}, \cdots, \langle \xi_2, \langle \xi_1, \eta_1 \rangle \eta_2 \rangle \cdots \eta_{n-1} \rangle \eta_n \rangle = \langle \xi, \eta \rangle_n$$

Proof. Apply Lemma 1.6 (n-1) times.

Lemma 1.8. Let $n \in \mathbb{N}$. For $\xi_1, \ldots, \xi_n, \eta_1, \ldots, \eta_n \in C_c(U)$, we have

$$(t^{1}(\xi_{1})t^{1}(\xi_{2})\cdots t^{1}(\xi_{n}))^{*}(t^{1}(\eta_{1})t^{1}(\eta_{2})\cdots t^{1}(\eta_{n}))$$

$$= t^{0}(\langle \xi_{n}, \langle \xi_{n-1}, \cdots, \langle \xi_{2}, \langle \xi_{1}, \eta_{1} \rangle \eta_{2} \rangle \cdots \eta_{n-1} \rangle \eta_{n} \rangle)$$

Proof. This follows from direct computation.

Lemma 1.9. Let $n, m \in \mathbb{N}$. For $\xi \in C_c(U_{n+m})$, there exist $\xi_1 \in C_c(U_n)$ and $\xi_2 \in C_c(U_m)$ such that $\xi(x) = \xi_1(x)\xi_2(\sigma^n(x))$ for $x \in U_{n+m}$.

Proof. Take $\xi_1 = \xi$, and choose $\xi_2 \in C_c(U_m)$ which is 1 on the compact set $\sigma^n(C) \subset U_m$ where $C \subset U_{n+m}$ is the compact support of ξ .

Lemma 1.10. Let $n \in \mathbb{N}$. For $\xi \in C_c(U_n)$, there exist $\xi_1, \xi_2, \dots, \xi_n \in C_c(U)$ such that $\xi(x) = \xi_1(x)\xi_2(\sigma(x))\cdots\xi_n(\sigma^{n-1}(x))$ for $x \in U_n$.

Proof. Apply Lemma 1.9 (n-1) times.

For each integer $n \geq 2$, we define a linear map $t^n : C_c(U_n) \to C^*(\Sigma)$. Take $\xi \in C_c(U_n)$. Take $\xi_1, \ldots, \xi_n \in C_c(U)$ such that

$$\xi(x) = \xi_1(x)\xi_2(\sigma(x))\cdots\xi_n(\sigma^{n-1}(x))$$

for all $x \in U_n$. Such functions exist by Lemma 1.10. We set $t^n(\xi) = t^1(\xi_1)t^1(\xi_2)\cdots t^1(\xi_n)$. This definition is well-defined because Lemma 1.7 and Lemma 1.8 imply $d^*d = 0$ for the difference d of two definitions. By the same reason, we see that t^n is linear. We also see the following.

Lemma 1.11. For $\xi, \eta \in C_c(U_n)$, we have $t^n(\xi)^*t^n(\eta) = t^0(\langle \xi, \eta \rangle_n)$.

Proof. This follows from Lemma 1.7 and Lemma 1.8.

Lemma 1.12. Let $n, m \in \mathbb{N}$. For $\xi_1 \in C_c(U_n)$ and $\xi_2 \in C_c(U_m)$, define $\xi \in C_c(U_{n+m})$ by $\xi(x) = \xi_1(x)\xi_2(\sigma^n(x))$ for $x \in U_{n+m}$. Then we have

$$t^{n}(\xi_{1})t^{m}(\xi_{2}) = t^{n+m}(\xi)$$

Proof. When $n \ge 1$ and $m \ge 1$, this follows from the definition of t^n . When n = 0 or m = 0, this follows from a similar computation as the one after Definition 1.4. \square

Lemma 1.13. Let $n, m \in \mathbb{N}$. For $\xi \in C_c(U_n)$ and $\eta \in C_c(U_{n+m}) \subset C_c(U_n)$, we have $t^n(\xi)^*t^{n+m}(\eta) = t^m(\langle \xi, \eta \rangle_n)$.

Proof. By Lemma 1.9, choose $\eta_1 \in C_c(U_n)$ and $\eta_2 \in C_c(U_m)$ such that $\eta(x) = \eta_1(x)\eta_2(\sigma^n(x))$ for $x \in U_{n+m}$. By Lemma 1.12 and Lemma 1.11, we have

$$t^{n}(\xi)^{*}t^{n+m}(\eta) = t^{n}(\xi)^{*}t^{n}(\eta_{1})t^{m}(\eta_{2}) = t^{0}(\langle \xi, \eta_{1} \rangle_{n})t^{m}(\eta_{2})$$
$$= t^{m}(\langle \xi, \eta_{1} \rangle_{n}\eta_{2}) = t^{m}(\langle \xi, \eta_{1} (\eta_{2} \circ \sigma^{n}) \rangle_{n}) = t^{m}(\langle \xi, \eta \rangle_{n}). \qquad \Box$$

Lemma 1.14. The C^* -algebra $C^*(\Sigma)$ is the closure of the linear span of the set $\{t^n(\xi)t^m(\eta)^* \mid n, m \in \mathbb{N}, \xi \in C_c(U_n), \eta \in C_c(U_m)\}.$

Proof. By Lemma 1.12 and Lemma 1.13, the set above is closed under multiplication. This set contains the images of t^0 and t^1 . Hence the closure of the linear span of the set is $C^*(\Sigma)$.

2. Irreducible representations of $C^*(\Sigma)$

Take an SGDS $\Sigma = (X, \sigma)$. In this section, we construct irreducible representations $\pi_{x_0,\gamma}$ of the C^* -algebra $C^*(\Sigma)$.

Definition 2.1. For $x_0 \in X$, we define the *orbit* of x_0 by

$$Orb(x_0) = \{ x \in X \mid \sigma^n(x) = \sigma^m(x_0) \text{ for some } n, m \in \mathbb{N} \}.$$

Note that for $x \in \text{Orb}(x_0)$ we have $\text{Orb}(x) = \text{Orb}(x_0)$. Hence two orbits Orb(x) and Orb(y) are either same or disjoint.

Definition 2.2. For $x_0 \in X$, let H_{x_0} be the Hilbert space whose complete orthonormal system is given by $\{\delta_x\}_{x \in \text{Orb}(x_0)}$. The inner product of H_{x_0} is denoted by $\langle \cdot, \cdot \rangle_{x_0}$ which is linear in the second variable.

Definition 2.3. For $(x_0, \gamma_0) \in X \times \mathbb{T}$, we define two maps $t^0_{(x_0, \gamma_0)} : C_0(X) \to B(H_{x_0})$ and $t^1_{(x_0, \gamma_0)} : C_c(U) \to B(H_{x_0})$ by

$$t^{0}_{(x_{0},\gamma_{0})}(f)\delta_{x} = f(x)\delta_{x},$$
 $t^{1}_{(x_{0},\gamma_{0})}(\xi)\delta_{x} = \gamma_{0}\sum_{\sigma(y)=x}\xi(y)\delta_{y},$

for $x \in Orb(x_0)$.

Take $(x_0, \gamma_0) \in X \times \mathbb{T}$ and fix it for a while. It is routine to check that $t^0_{(x_0, \gamma_0)}$ is a well-defined *-homomorphism and $t^1_{(x_0, \gamma_0)}$ is a well-defined linear map.

Lemma 2.4. For $\xi \in C_c(U)$ and $y \in Orb(x_0)$, we have

$$t_{(x_0,\gamma_0)}^1(\xi)^*\delta_y = \begin{cases} \gamma_0^{-1}\overline{\xi}(y)\delta_{\sigma(y)} & if \ y \in U\\ 0 & otherwise. \end{cases}$$

Proof. For $x, y \in Orb(x_0)$, we have

$$\begin{split} \langle \delta_x, t^1_{(x_0, \gamma_0)}(\xi)^* \delta_y \rangle_{x_0} &= \langle t^1_{(x_0, \gamma_0)}(\xi) \delta_x, \delta_y \rangle_{x_0} \\ &= \left\langle \gamma_0 \sum_{\sigma(y') = x} \xi(y') \delta_{y'}, \delta_y \right\rangle_{x_0} \\ &= \sum_{\sigma(y') = x} \gamma_0^{-1} \overline{\xi(y')} \langle \delta_{y'}, \delta_y \rangle_{x_0} \\ &= \begin{cases} \gamma_0^{-1} \overline{\xi}(y) & \text{if } x = \sigma(y), \\ 0 & \text{otherwise.} \end{cases} \end{split}$$

This completes the proof.

Proposition 2.5. The pair $(t^0_{(x_0,\gamma_0)}, t^1_{(x_0,\gamma_0)})$ satisfies (i) and (ii) in Definition 1.4.

Proof. Take $\xi, \eta \in C_c(U)$. For $x \in Orb(x_0)$, we have

$$t_{(x_0,\gamma_0)}^1(\xi)^* t_{(x_0,\gamma_0)}^1(\eta) \delta_x = t_{(x_0,\gamma_0)}^1(\xi)^* \left(\gamma_0 \sum_{\sigma(y)=x} \eta(y) \delta_y \right)$$

$$= \gamma_0 \sum_{\sigma(y)=x} \eta(y) t_{(x_0,\gamma_0)}^1(\xi)^* \delta_y$$

$$= \gamma_0 \sum_{\sigma(y)=x} \eta(y) \gamma_0^{-1} \overline{\xi}(y) \delta_{\sigma(y)}$$

$$= \sum_{\sigma(y)=x} \overline{\xi}(y) \eta(y) \delta_x$$

$$= \langle \xi, \eta \rangle(x) \delta_x$$

by Lemma 2.4. This shows $t^1_{(x_0,\gamma_0)}(\xi)^*t^1_{(x_0,\gamma_0)}(\eta) = t^0_{(x_0,\gamma_0)}(\langle \xi, \eta \rangle)$. Now take an open subset $V \subset U$ on which σ is injective, and take $\xi, \eta \in C_c(V)$. For $y \in \mathrm{Orb}(x_0) \cap V$, we have

$$t_{(x_0,\gamma_0)}^1(\xi)t_{(x_0,\gamma_0)}^1(\eta)^*\delta_y = t_{(x_0,\gamma_0)}^1(\xi)\left(\gamma_0^{-1}\overline{\eta}(y)\delta_{\sigma(y)}\right)$$
$$= \gamma_0^{-1}\overline{\eta}(y)\gamma_0 \sum_{\sigma(y')=\sigma(y)} \xi(y')\delta_{y'}$$
$$= \overline{\eta}(y)\xi(y)\delta_y$$

by Lemma 2.4. For $y \in \text{Orb}(x_0) \setminus V$, the same equation holds because the both sides become 0. Hence we get $t^1_{(x_0,\gamma_0)}(\xi)t^1_{(x_0,\gamma_0)}(\eta)^* = t^0_{(x_0,\gamma_0)}(\xi\overline{\eta})$.

Definition 2.6. We denote by $\pi_{(x_0,\gamma_0)} \colon C^*(\Sigma) \to B(H_{x_0})$ the *-homomorphism induced by the pair $(t^0_{(x_0,\gamma_0)}, t^1_{(x_0,\gamma_0)})$.

Definition 2.7. For $(x_0, \gamma_0) \in X \times \mathbb{T}$, we set $P_{(x_0, \gamma_0)} = \ker \pi_{(x_0, \gamma_0)}$

Lemma 2.8. Let (t^0, t^1) be the universal pair for $C^*(\Sigma)$. Then both t^0 and t^1 are injective.

Proof. For each $x \in X$, we have $\ker t^0_{(x,1)} = C_0(X \setminus \overline{\operatorname{Orb}(x)})$. Since $t^0_{(x,1)} = \pi_{(x,1)} \circ t^0$, we have $\ker t^0 \subset \ker t^0_{(x,1)}$. Hence we have

$$\ker t^0 \subset \bigcap_{x \in X} \ker t^0_{(x,1)} = \bigcap_{x \in X} C_0(X \setminus \overline{\operatorname{Orb}(x)}) = 0.$$

This shows that t^0 is injective. Take $\xi \in C_c(U)$ with $t^1(\xi) = 0$. Then we have $t^0(\langle \xi, \xi \rangle) = t^1(\xi)^* t^1(\xi) = 0$. Since t^0 is injective, we have $\langle \xi, \xi \rangle = 0$. This shows $\xi = 0$. Therefore t^1 is injective.

We are going to see that the representation $\pi_{(x_0,\gamma_0)}$ is irreducible, and hence $P_{(x_0,\gamma_0)}$ is a primitive ideal of $C^*(\Sigma)$.

Lemma 2.9. For $n \in \mathbb{N}$, $\xi \in C_0(U_n)$ and $x \in Orb(x_0)$, we have

$$\pi_{(x_0,\gamma_0)}(t^n(\xi))\delta_x = \gamma_0^n \sum_{\sigma^n(y)=x} \xi(y)\delta_y$$

$$\pi_{(x_0,\gamma_0)}(t^n(\xi))^*\delta_x = \begin{cases} \gamma_0^{-n}\overline{\xi}(x)\delta_{\sigma^n(x)} & \text{if } x \in U_n, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. By Lemma 1.10, we can choose $\xi_1, \ldots, \xi_n \in C_c(U)$ such that

$$\xi(x) = \xi_1(x)\xi_2(\sigma(x))\cdots\xi_n(\sigma^{n-1}(x))$$

for all $x \in U_n$. Then we have

$$\pi_{(x_{0},\gamma_{0})}(t^{n}(\xi))\delta_{x} = \pi_{(x_{0},\gamma_{0})}(t^{1}(\xi_{1})t^{1}(\xi_{2})\dots t^{1}(\xi_{n-1})t^{1}(\xi_{n}))\delta_{x}$$

$$= \pi_{(x_{0},\gamma_{0})}(t^{1}(\xi_{1})t^{1}(\xi_{2})\dots t^{1}(\xi_{n-1}))\gamma_{0}\sum_{\sigma(y_{1})=x}\xi_{n}(y_{1})\delta_{y_{1}}$$

$$= \pi_{(x_{0},\gamma_{0})}(t^{1}(\xi_{1})\dots t^{1}(\xi_{n-2}))\gamma_{0}^{2}\sum_{\sigma(y_{1})=x}\xi_{n}(y_{1})\sum_{\sigma(y_{2})=y_{1}}\xi_{n-1}(y_{2})\delta_{y_{2}}$$

$$= \pi_{(x_{0},\gamma_{0})}(t^{1}(\xi_{1})\dots t^{1}(\xi_{n-2}))\gamma_{0}^{2}\sum_{\sigma^{2}(y_{2})=x}\xi_{n-1}(y_{2})\xi_{n}(\sigma(y_{2}))\delta_{y_{2}}$$

$$\vdots$$

$$= \pi_{(x_{0},\gamma_{0})}(t^{1}(\xi_{1}))\gamma_{0}^{n-1}\sum_{\sigma^{n-1}(y_{n-1})=x}\xi_{2}(y_{n-1})\xi_{3}(\sigma(y_{n-1}))\cdots$$

$$\cdots \xi_{n-1}(\sigma^{n-3}(y_{n-1}))\xi_{n}(\sigma^{n-2}(y_{n-1}))\delta_{y_{n-1}}$$

$$= \gamma_{0}^{n}\sum_{\sigma^{n}(y)=x}\xi(y)\delta_{y}.$$

From this equation, we can compute $\pi_{(x_0,\gamma_0)}(t^n(\xi))^*\delta_x$ in a similar way to the proof of Lemma 2.4.

Proposition 2.10. For $(x_0, \gamma_0) \in X \times \mathbb{T}$, the representation $\pi_{(x_0, \gamma_0)} \colon C^*(\Sigma) \to B(H_{x_0})$ is irreducible.

Proof. Let $\{e_{x,y}\}_{x,y\in Orb(x_0)}$ be the matrix units of $B(H_{x_0})$. Namely $e_{x,y}\in B(H_{x_0})$ satisfies

$$e_{x,y}\delta_z = \begin{cases} \delta_x & \text{if } z = y, \\ 0 & \text{otherwise} \end{cases}$$

for $z \in \operatorname{Orb}(x_0)$. For all $x \in \operatorname{Orb}(x_0)$, it is standard to see that $e_{x,x}$ is in the weak closure of $\pi_{(x_0,\gamma_0)}(t^0(C_0(X))) = t^0_{(x_0,\gamma_0)}(C_0(X)) \subset B(H_{x_0})$. Take $x \in \operatorname{Orb}(x_0)$ with $x \in U$. Take $\xi \in C_0(U)$ with $\xi(x) = \gamma_0^{-1}$. Then we have $e_{x,x}t^1_{(x_0,\gamma_0)}(\xi) = e_{x,\sigma(x)}$. Hence $e_{x,\sigma(x)}$ is in the weak closure of $\pi_{(x_0,\gamma_0)}(C^*(\Sigma))$ for all $x \in \operatorname{Orb}(x_0)$ with $x \in U$. Take $x, y \in \operatorname{Orb}(x_0)$. Then there exist $n, m \in \mathbb{N}$ with $\sigma^n(x) = \sigma^m(y)$. We have that

$$e_{x,y} = e_{x,\sigma^{n}(x)}(e_{y,\sigma^{m}(y)})^{*}$$

$$= e_{x,x}e_{x,\sigma(x)}e_{\sigma(x),\sigma^{2}(x)}\dots e_{\sigma^{n-1}(x),\sigma^{n}(x)}(e_{y,y}e_{y,\sigma(y)}e_{\sigma(y),\sigma^{2}(y)}\dots e_{\sigma^{m-1}(y),\sigma^{m}(y)})^{*}$$

is in the weak closure of $\pi_{(x_0,\gamma_0)}(C^*(\Sigma))$. Hence the weak closure of $\pi_{(x_0,\gamma_0)}(C^*(\Sigma))$. is whole $B(H_{x_0})$.

Corollary 2.11. For $(x_0, \gamma_0) \in X \times \mathbb{T}$, the ideal $P_{(x_0, \gamma_0)}$ is primitive.

Lemma 2.12. For $x \in U$ and $\gamma \in \mathbb{T}$, we have $P_{(x,\gamma)} = P_{(\sigma(x),\gamma)}$.

Proof. Since
$$Orb(x) = Orb(\sigma(x))$$
, we have $\pi_{(x,\gamma)} = \pi_{(\sigma(x),\gamma)}$. Hence $P_{(x,\gamma)} = P_{(\sigma(x),\gamma)}$.

Definition 2.13. Let $x \in X$. If there exist $k, n \in \mathbb{N}$ with $n \geq 1$ such that $\sigma^{k+n}(x) = \sigma^k(x)$, then we say that x is *periodic*, and define its *period* p(x) to be the smallest positive integer n satisfying $\sigma^{k+n}(x) = \sigma^k(x)$ for some k. We also denote l(x) be the smallest natural number k satisfying $\sigma^{k+p(x)}(x) = \sigma^k(x)$. A point x which is not periodic is said to be *aperiodic*. We set $p(x) = l(x) = \infty$ for an aperiodic point x.

It is fairly easy to see, but worth remarking, that we have $\sigma^k(x) = \sigma^l(x)$ for $k, l \in \mathbb{N}$ with k > l if and only if $l \geq l(x)$ and $k - l \in p(x)\mathbb{N}$.

Lemma 2.14. For an aperiodic point $x_0 \in X$, we have $P_{(x_0,\gamma)} = P_{(x_0,1)}$ for all $\gamma \in \mathbb{T}$.

Proof. If $x_0 \in X$ is aperiodic, we can define a map c: $\operatorname{Orb}(x_0) \to \mathbb{Z}$ such that $c(x_0) = 0$ and $c(\sigma(x)) = c(x) - 1$ for $x \in U$. We define a unitary $u_{\gamma} \in B(H_{x_0})$ by $u_{\gamma}\delta_x = \gamma^{c(x)}\delta_x$ for $x \in \operatorname{Orb}(x_0)$. It is not difficult to check that

$$u_{\gamma}t^{0}_{(x_{0},1)}(f)u^{*}_{\gamma} = t^{0}_{(x_{0},1)}(f) = t^{0}_{(x_{0},\gamma)}(f),$$

$$u_{\gamma}t^{1}_{(x_{0},1)}(\xi)u^{*}_{\gamma} = \gamma t^{1}_{(x_{0},1)}(\xi) = t^{1}_{(x_{0},\gamma)}(\xi)$$

for $f \in C_0(X)$ and $\xi \in C_c(U)$. Hence two representation $\pi_{(x_0,\gamma)}$ and $\pi_{(x_0,1)}$ are unitarily equivalent. This shows $P_{(x_0,\gamma)} = P_{(x_0,1)}$.

We denote the elements of $\mathbb{Z}/n\mathbb{Z}$ by $\{0,1,\ldots,n-1\}$, and sometimes consider them as elements in \mathbb{Z} .

Lemma 2.15. For a periodic point $x_0 \in X$ with period n, we have $P_{(x_0,\gamma)} = P_{(x_0,\mu)}$ if $\gamma^n = \mu^n$.

Proof. Similarly as in the proof of Lemma 2.14, we can define a map $c: \operatorname{Orb}(x_0) \to \mathbb{Z}/n\mathbb{Z}$ such that $c(x_0) = 0$ and $c(\sigma(x)) = c(x) - 1$ for $x \in U$. Set $\lambda := \gamma \overline{\mu} \in \mathbb{T}$. We have $\lambda^n = 1$. Hence we can define $u_{\lambda} \in B(H_{x_0})$ by $u_{\lambda} \delta_x = \lambda^{c(x)} \delta_x$ for $x \in \operatorname{Orb}(x_0)$.

Similarly as in the proof of Lemma 2.14, two representations $\pi_{(x_0,\gamma)}$ and $\pi_{(x_0,\mu)}$ are unitarily equivalent by the unitary u_{λ} . Hence we obtain $P_{(x_0,\gamma)} = P_{(x_0,\mu)}$.

Definition 2.16. For an integer $n \geq 2$, we denote by $\zeta_n = e^{2\pi i/n}$ the *n*-th root of unity.

Lemma 2.17. Let $x_0 \in X$ be a periodic point with period n. If x_0 is isolated in $Orb(x_0)$, then for $\gamma, \mu \in \mathbb{T}$, $P_{(x_0,\gamma)} = P_{(x_0,\mu)}$ if and only if $\gamma^n = \mu^n$.

Proof. Let $\Lambda = \{\sigma^{l(x_0)}(x_0), \ldots, \sigma^{l(x_0)+n-1}(x_0)\}$. Since σ is a local homeomorphism, if x_0 is isolated in $\operatorname{Orb}(x_0)$ we can find an open subset V of X such that $V \cap \operatorname{Orb}(x_0) = \Lambda$. Hence there exists $\xi \in C_c(U)$ such that $\xi(x) = 1$ for $x \in \Lambda$ and $\xi(x) = 0$ for $x \in \operatorname{Orb}(x_0) \setminus \Lambda$. For each $\gamma \in \mathbb{T}$, non-zero elements of the spectrum of $t^1_{(x_0,\gamma)}(\xi) \in B(H_{x_0})$ are $\gamma, \gamma\zeta_n, \ldots, \gamma\zeta_n^{n-1}$. Hence those of the image of $t^1(\xi)$ via the natural surjection $C^*(\Sigma) \to C^*(\Sigma)/P_{(x_0,\gamma)}$ are also $\gamma, \gamma\zeta_n, \ldots, \gamma\zeta_n^{n-1}$. This shows that if $P_{(x_0,\gamma)} = P_{(x_0,\mu)}$ then $\gamma^n = \mu^n$. The converse follows from Lemma 2.15. \square

Remark 2.18. We will see in Proposition 4.10 that for a periodic point $x_0 \in X$ such that x_0 is not isolated in $Orb(x_0)$, we have $P_{(x_0,\gamma)} = P_{(x_0,1)}$ for all $\gamma \in \mathbb{T}$.

Remark 2.19. We will see in Remark 4.17 that if X is second countable $P_{(x,\gamma)}$'s are whole primitive ideal. (See [SW] for the case U=X.) In general, there is a primitive ideal which is not in the form $P_{(x,\gamma)}$ (see [K3, Example 13.2]).

It is probably impossible to list all primitive ideals in terms of elements of X and \mathbb{T} . However it is possible to list all prime ideals in terms of subsets of X and \mathbb{T} . From next section, we will do this. This will show that the set $\{P_{(x,\gamma)}\}$ is sufficiently large (Corollary 4.19).

3. Gauge invariant ideals and σ -invariant sets

Definition 3.1. A subset X' of X is said to be σ -invariant when $x \in X'$ if and only if $\sigma(x) \in X'$ for all $x \in U$.

An ideal I of $C^*(\Sigma)$ is said to be gauge-invariant if $\beta_z(I) = I$ for all $z \in \mathbb{T}$ where β is the gauge action defined in Appendix A. We will see in Proposition 3.11 that the set of gauge-invariant ideals corresponds bijectively to the set of closed σ -invariant subsets of X.

Definition 3.2. For an ideal I of $C^*(\Sigma)$, we define a closed set X_I of X by $t^0(C_0(X \setminus X_I)) = t^0(C_0(X)) \cap I$.

The following lemma is easy to see from the definition.

Lemma 3.3. For two ideals I_1, I_2 of $C^*(\Sigma)$, we have $X_{I_1 \cap I_2} = X_{I_1} \cup X_{I_2}$. If $I_1 \subset I_2$, then $X_{I_1} \supset X_{I_2}$.

Proposition 3.4. For an ideal I of $C^*(\Sigma)$, the closed set X_I is σ -invariant.

Proof. Take $x \in U$. Suppose $x \notin X_I$. Choose an open subset V such that $x \in V \subset U$, $V \cap X_I = \emptyset$ and σ is injective on V. Take $\xi \in C_c(V)$ with $\xi(x) = 1$. Then $\xi \overline{\xi} \in C_0(X \setminus X_I)$. Hence $t^1(\xi)t^1(\xi)^* = t^0(\xi \overline{\xi}) \in I$. This implies $t^0(\langle \xi, \xi \rangle) = t^1(\xi)^*t^1(\xi) \in I$. Hence $\langle \xi, \xi \rangle \in C_0(X \setminus X_I)$. Since $\langle \xi, \xi \rangle (\sigma(x)) = |\xi(x)|^2 = 1$, we

have $\sigma(x) \notin X_I$. Now suppose $\sigma(x) \notin X_I$. Choose an open subset V such that $x \in V \subset U$, $V \cap \sigma^{-1}(X_I) = \emptyset$ and σ is injective on V. Take $\xi \in C_c(V)$ with $\xi(x) = 1$. Then $\langle \xi, \xi \rangle \in C_0(X \setminus X_I)$. Hence $t^1(\xi)^*t^1(\xi) = t^0(\langle \xi, \xi \rangle) \in I$. This implies $t^0(\xi\overline{\xi}) = t^1(\xi)t^1(\xi)^* \in I$. Hence $\xi\overline{\xi} \in C_0(X \setminus X_I)$. Since $\xi\overline{\xi}(x) = |\xi(x)|^2 = 1$, we have $x \notin X_I$. Thus we have shown that $x \in X_I$ if and only if $\sigma(x) \in X_I$.

Take a closed σ -invariant subset X' of X. We define an SGDS $\Sigma' = (X', \sigma')$ where σ' is a restriction of σ on $U' := X' \cap U$. Let (t'^0, t'^1) be the universal pair for $C^*(\Sigma')$. We have the following.

Lemma 3.5. There exists a surjection $\Phi: C^*(\Sigma) \to C^*(\Sigma')$ such that $\Phi(t^0(f)) = t'^0(f|_{X'})$ and $\Phi(t^1(\xi)) = t'^1(\xi|_{U'})$.

Proof. It suffices to see that the pair of maps $C_0(X) \ni f \mapsto t'^0(f|_{X'}) \in C^*(\Sigma')$ and $C_c(U) \ni \xi \mapsto t'^1(\xi|_{U'}) \in C^*(\Sigma')$ satisfy (i) and (ii) in Definition 1.4 for Σ . This follows from the fact that t'^0 and t'^1 satisfy (i) and (ii) in Definition 1.4 for Σ' . \square

Definition 3.6. For a closed σ -invariant subset X' of X, let $I_{X'}$ be the kernel of the surjection $\Phi: C^*(\Sigma) \to C^*(\Sigma')$ in Lemma 3.5.

Proposition 3.7. For a closed σ -invariant subset X' of X, we have $X_{I_{X'}} = X'$.

Proof. This follows from the fact that the kernel of $\Phi \circ t^0$ in Lemma 3.5 is $C_0(X \setminus X')$.

Lemma 3.8. For a closed σ -invariant subset X' of X, the ideal $I_{X'}$ is gauge-invariant.

Proof. This follows from the fact that the surjection $\Phi: C^*(\Sigma) \to C^*(\Sigma')$ in Lemma 3.5 commutes with the gauge actions.

Lemma 3.9. For an ideal I of $C^*(\Sigma)$, we have $I_{X_I} \subset I$.

Proof. Takes an deal I of $C^*(\Sigma)$, and denote the natural surjection by $\pi\colon C^*(\Sigma)\to C^*(\Sigma)/I$. We set $\Sigma_I:=(X_I,\sigma_I)$ where σ_I is the restriction of σ to $U_I:=U\cap X_I$. By the definition of X_I , we can define a *-homomorphism $t'^0\colon C_0(X_I)\to C^*(\Sigma)/I$ such that $t'^0(f|_{X_I})=\pi(t^0(f))$ for all $f\in C_0(X)$. We can also define a linear map $t'^1\colon C_c(U_I)\to C^*(\Sigma)/I$ such that $t'^1(\xi|_{U_I})=\pi(t^1(\xi))$ for all $\xi\in C_c(U)$ because for $\eta\in C_c(U)$ with $\eta|_{U_I}=0$ we have $\langle \eta,\eta\rangle\in C_0(X\setminus X_I)$ and hence $t^1(\eta)\in I$. The pair (t'^0,t'^1) satisfies (i) and (ii) in Definition 1.4 for Σ_I . Hence we get a *-homomorphism $\Psi\colon C^*(\Sigma_I)\to C^*(\Sigma)/I$ with $\pi=\Psi\circ \Phi_I$ where $\Phi_I\colon C^*(\Sigma)\to C^*(\Sigma_I)$ is the surjection whose kernel is I_{X_I} . Hence we get $I_{X_I}\subset I$.

Proposition 3.10. For an ideal I of $C^*(\Sigma)$, we have $I_{X_I} = I$ if and only if I is gauge-invariant.

Proof. The "only if" part follows from Lemma 3.8. Let I be a gauge-invariant ideal I of $C^*(\Sigma)$. Then we can define an action of \mathbb{T} on $C^*(\Sigma)/I$ so that the natural surjection $\pi \colon C^*(\Sigma) \to C^*(\Sigma)/I$ becomes equivariant. Then the map $\Psi \colon C^*(\Sigma_I) \to C^*(\Sigma)/I$ in the proof of Lemma 3.9 becomes also equivariant. The map Ψ is injective on $t^0(C_0(X_I))$ by the definition of X_I . Hence by Proposition A.1 Ψ is injective. Therefore we have $I_{X_I} = I$.

Proposition 3.11. Through the maps $I \mapsto X_I$ and $X' \mapsto I_{X'}$, the set of gauge-invariant ideals corresponds bijectively to the set of closed σ -invariant subsets of X.

Proof. This follows from Proposition 3.7 and Proposition 3.10. \Box

Now we have the following.

Lemma 3.12. For closed σ -invariant subsets X_1 and X_2 of X, we have $I_{X_1 \cup X_2} = I_{X_1} \cap I_{X_2}$. If $X_1 \subset X_2$, then $I_{X_1} \supset I_{X_2}$.

Proof. By Lemma 3.3 and Proposition 3.7, we have

$$X_{I_{X_1}\cap I_{X_2}}=X_{I_{X_1}}\cup X_{I_{X_2}}=X_1\cup X_2.$$

Since $I_{X_1} \cap I_{X_2}$ is gauge-invariant, we have

$$I_{X_1} \cap I_{X_2} = I_{X_{I_{X_1} \cap I_{X_2}}} = I_{X_1 \cup X_2}$$

by Proposition 3.10. If $X_1 \subset X_2$, then $X_2 = X_1 \cup X_2$. Hence we have $I_{X_2} = I_{X_1 \cup X_2} = I_{X_1} \cap I_{X_2}$. This shows $I_{X_1} \supset I_{X_2}$.

Definition 3.13. For a closed σ -invariant set X', we say X' is essentially free if the SGDS $\Sigma' = (X', \sigma|_{U \cap X'})$ is essentially free as defined in Definition A.2.

Proposition 3.14. For an ideal I of $C^*(\Sigma)$, if X_I is essentially free then we have $I_{X_I} = I$.

Proof. The proof goes similarly as in Proposition 3.10 using Proposition A.3 instead of Proposition A.1. \Box

4. Prime ideals

An ideal I of a C^* -algebra A is said to be *prime* if $I \neq A$ and for two ideals I_1, I_2 of a C^* -algebra A with $I_1 \cap I_2 \subset I$ we have either $I_1 \subset I$ or $I_2 \subset I$. A primitive ideal is prime [B, II.6.1.11]. The converse is true when X is second countable (cf. [B, II.6.5.15]), but not true in general (see [K3, Example 13.3] and [K4, Theorem 5.4]).

Definition 4.1. A closed σ -invariant set X is said to be *irreducible* if it is non-empty and we have either $X = X_1$ or $X = X_2$ for two closed σ -invariant sets X_1, X_2 satisfying $X = X_1 \cup X_2$.

Proposition 4.2. If an ideal I of $C^*(\Sigma)$ is prime, then X_I is irreducible.

Proof. If $X_I = \emptyset$, then I = A which contradicts that I is prime. Hence $X_I \neq \emptyset$. Take two closed σ -invariant sets X_1, X_2 satisfying $X_I = X_1 \cup X_2$. Then

$$I_{X_1} \cap I_{X_2} = I_{X_1 \cup X_2} = I_{X_I} \subset I$$

by Lemma 3.12 and Lemma 3.9. Since I is prime, we have either $I_{X_1} \subset I$ or $I_{X_2} \subset I$. When $I_{X_1} \subset I$, we have $X_I \subset X_{I_{X_1}} = X_1 \subset X_I$ by Lemma 3.3 and Proposition 3.7, and hence $X_1 = X_I$. Similarly we have $X_2 = X_I$ when $I_{X_2} \subset I$. Thus X_I is irreducible.

Proposition 4.3. For $(x, \gamma) \in X \times \mathbb{T}$, we have $X_{P_{(x,\gamma)}} = \overline{\operatorname{Orb}(x)}$.

Proof. This follows from $\ker t_{(x,\gamma)}^0 = C_0(X \setminus \overline{\operatorname{Orb}(x)}).$

Proposition 4.4. For $x \in X$, $\overline{Orb(x)}$ is an irreducible closed σ -invariant set.

Proof. This follows from Proposition 4.2 and Proposition 4.3. One can also show this directly as follows. Since $\operatorname{Orb}(x)$ is σ -invariant and since σ is locally homeomorphic, $\overline{\operatorname{Orb}(x)}$ is closed and σ -invariant. Take two closed σ -invariant sets X_1, X_2 satisfying $\overline{\operatorname{Orb}(x)} = X_1 \cup X_2$. Then either $x \in X_1$ or $x \in X_2$. When $x \in X_1$ we have $X_1 = \overline{\operatorname{Orb}(x)}$, whereas when $x \in X_2$ we have $X_2 = \overline{\operatorname{Orb}(x)}$. Thus $\overline{\operatorname{Orb}(x)}$ is irreducible.

Proposition 4.5. Let X' be an irreducible closed σ -invariant set. If X' is essentially free, then $I_{X'}$ is prime.

Proof. Since $X' \neq \emptyset$, we have $I_{X'} \neq A$. Take ideals I_1, I_2 of $C^*(\Sigma)$ such that $I_1 \cap I_2 \subset I_{X'}$. Then we have

$$X_{I_1+I_{X'}} \cup X_{I_2+I_{X'}} = X_{(I_1+I_{X'})\cap(I_2+I_{X'})} = X_{I_{X'}} = X'$$

by Lemma 3.3 and Proposition 3.7. Since X' is irreducible, either we have $X_{I_1+I_{X'}}=X'$ or $X_{I_2+I_{X'}}=X'$. When $X_{I_1+I_{X'}}=X'$ we have $I_1+I_{X'}=I_{X'}$ by Proposition 3.14 because X' is essentially free. This shows $I_1\subset I_{X'}$. Similarly, when $X_{I_2+I_{X'}}=X'$ we have $I_2\subset I_{X'}$. Thus $I_{X'}$ is prime.

From now on, we investigate which irreducible closed σ -invariant set X' becomes essentially free.

Definition 4.6. We set

$$P(\Sigma) := \{ x \in X \mid l(x) = 0, x \text{ is isolated in } Orb(x) \}.$$

Definition 4.7. We denote by $\mathcal{A}(\Sigma)$ the set of all irreducible closed σ -invariant set which is not in the form $\overline{\mathrm{Orb}(x)}$ for $x \in \mathrm{P}(\Sigma)$.

Lemma 4.8. For $x \in P(\Sigma)$, an irreducible closed σ -invariant $\overline{Orb(x)}$ is not essentially free.

Proof. In the SGDS $\Sigma' = (X', \sigma|_{U \cap X'})$ where $X' = \operatorname{Orb}(x)$, $\{x\}$ is an open subset and l(x) = 0. Hence Σ' is not essentially free.

Proposition 4.9 (cf. [K3, Proposition 11.3]). For an irreducible closed σ -invariant set X', X' is essentially free if and only if $X' \in \mathcal{A}(\Sigma)$

Proof. Take an irreducible closed σ -invariant set X'. By Lemma 4.8, if X' is essentially free then $X' \in \mathcal{A}(\Sigma)$. Suppose X' is not essentially free. Then there exists a non-empty open subset V' of X' such that l(x) = 0 for all $x \in V'$. By Baire's category theorem (see, for example, [T1, Proposition 2.2]), there exist a non-empty open subset $V \subset V'$ of X' and $n \in \mathbb{N}$ such that p(x) = n for all $x \in V$. Take $x \in V$ arbitrarily. We will show that $V = \{x, \sigma(x), \dots, \sigma^{n-1}(x)\}$. To the contrary, suppose there exists $y \in V \setminus \{x, \sigma(x), \dots, \sigma^{n-1}(x)\}$. Choose open subsets W_1, W_2 of V such that $x \in W_1$, $y \in W_2$ and $W_2 \cap \bigcup_{k=0}^{n-1} \sigma^k(W_1) = \emptyset$. Then $\bigcup_{k=0}^{\infty} (\sigma^k)^{-1}(W_1)$ is an open σ -invariant subset of X' because for $l = 1, 2, \dots$ we have $\sigma^l(W_1) \subset (\sigma^k)^{-1}(W_1)$ for $k \in \mathbb{N}$ with $k + l \in n\mathbb{N}$. Hence $X_1 := X' \setminus \bigcup_{k=0}^{\infty} (\sigma^k)^{-1}(W_1)$ is a closed σ -invariant subset of X' with $X_1 \neq X'$. Similarly, $X_2 := X' \setminus \bigcup_{k=0}^{\infty} (\sigma^k)^{-1}(W_2)$ is a closed σ -invariant subset of X' with $X_2 \neq X'$. Since X' is irreducible, we have $X_1 \cup X_2 \neq X'$.

Hence we have $(\sigma^{k_0})^{-1}(W_1) \cap (\sigma^{l_0})^{-1}(W_2) \neq \emptyset$ for some $k_0, l_0 \in \mathbb{N}$. Thus we get $z \in X'$ such that $\sigma^{k_0}(z) \in W_1$ and $\sigma^{l_0}(z) \in W_2$. Then for $m \in \mathbb{N}$ with $l_0 + mn \geq k_0$, $\sigma^{l_0+mn}(z)$ is in $W_2 \cap \bigcup_{k=0}^{n-1} \sigma^k(W_1)$ which is empty. This is a contradiction. Hence we have shown $V = \{x, \sigma(x), \dots, \sigma^{n-1}(x)\}$. Since V is open in X', $\{x\}$ is open in X'. Hence x is isolated in $\operatorname{Orb}(x) \subset X'$. This shows $x \in P(\Sigma)$. Finally we show that $X' = \overline{\operatorname{Orb}(x)}$. To do so, take $y \in X'$. Take a neighborhood W of y arbitrarily. Then

$$X_1 := X' \setminus \bigcup_{k=0}^{\infty} (\sigma^k)^{-1} \Big(\bigcup_{l=0}^{\infty} \sigma^l(W)\Big)$$

is a closed σ -invariant subset of X' with $X_1 \neq X'$. Since $\{x\}$ is open in X', $X_2 = X' \setminus \operatorname{Orb}(x)$ is also a closed σ -invariant subset of X' with $X_2 \neq X'$. Since X' is irreducible, we have $X_1 \cup X_2 \neq X'$. Hence we have

$$\operatorname{Orb}(x) \cap \bigcup_{k=0}^{\infty} (\sigma^k)^{-1} \Big(\bigcup_{l=0}^{\infty} \sigma^l(W)\Big) \neq \emptyset.$$

Therefore there exists $x' \in \operatorname{Orb}(x)$ and $k_0, l_0 \in \mathbb{N}$ such that $\sigma^{k_0}(x') \in \sigma^{l_0}(W)$. Hence $W \cap \operatorname{Orb}(x) \neq \emptyset$. Since W is arbitrary, we have $y \in \overline{\operatorname{Orb}(x)}$. Hence $X' = \overline{\operatorname{Orb}(x)}$. This shows $X' \notin \mathcal{A}(\Sigma)$.

Proposition 4.10. For a periodic point $x_0 \in X$ such that x_0 is not isolated in $Orb(x_0)$, we have $P_{(x_0,\gamma)} = P_{(x_0,1)}$ for all $\gamma \in \mathbb{T}$.

Proof. If x_0 is not isolated in $\operatorname{Orb}(x_0)$, then x is not isolated in $\operatorname{Orb}(x_0)$ for all $x \in \operatorname{Orb}(x_0)$ because σ is locally homeomorphic. Hence $X' := \operatorname{Orb}(x_0)$ has no isolated point. Thus we have $X' \in \mathcal{A}(\Sigma)$. By Proposition 4.9, X' is essentially free. By Proposition 3.14 and Proposition 4.3, we have $P_{(x_0,\gamma)} = I_{X'} = P_{(x_0,1)}$ for all $\gamma \in \mathbb{T}$.

Take $x_0 \in P(\Sigma)$. We set $n_0 := p(x_0)$. We are going to show that prime ideals I with $X_I = \overline{\operatorname{Orb}(x_0)}$ is parametrized as $P_{(x_0,e^{2\pi it})}$ for $0 \le t < 1/n_0$. Let us set $X' := \overline{\operatorname{Orb}(x_0)}$ and $X'' := \overline{\operatorname{Orb}(x_0)} \setminus \operatorname{Orb}(x_0)$ which is a closed σ -invariant subset of X'.

Lemma 4.11 (cf. [K3, Lemma 11.10]). For an ideal I of $C^*(\Sigma)$, we have $X_I = X'$ if and only if $I_{X'} \subset I$ and $I_{X''} \not\subset I$.

Proof. Take an ideal I of $C^*(\Sigma)$ with $X_I = X'$. Then $I_{X'} = I_{X_I} \subset I$ by Lemma 3.9. If $I_{X''} \subset I$ then we have $X_I \subset X_{I_{X''}} = X''$ by Proposition 3.7. This contradicts $X_I = X'$. Thus we get $I_{X''} \not\subset I$.

Conversely, take an ideal I of $C^*(\Sigma)$ with $I_{X'} \subset I$ and $I_{X''} \not\subset I$. By Lemma 3.3 and Proposition 3.7, we have $X_I \subset X_{I_{X'}} = X'$. Suppose $x_0 \notin X_I$. Then $X_I \subset X''$ since X_I is σ -invariant. This shows $I_{X''} \subset I_{X_I} \subset I$ by Lemma 3.12 and Lemma 3.9. This contradicts $I_{X''} \not\subset I$. Hence we have $x_0 \in X_I$. This shows $X' \subset X_I$. Therefore we get $X_I = X'$.

Let us set $\Sigma' = (X', \sigma|_{U \cap X'})$. Then we have $C^*(\Sigma') = C^*(\Sigma)/I_{X'}$ by the definition of $I_{X'}$. Let (t^0, t^1) be the universal pair for $C^*(\Sigma')$, and define t^n as in Section 1.

Let us define $f_0 \in C_c(U_{n_0})$ by

$$f_0(x) = \begin{cases} 1 & x = x_0 \\ 0 & x \neq x_0 \end{cases}.$$

We set $p_0 = t^0(f_0) \in C^*(\Sigma')$ and $u_0 = t^{n_0}(f_0) \in C^*(\Sigma')$. Note that we have $u_0^* u_0 = u_0 u_0^* = p_0$.

Lemma 4.12 (cf. [K3, Lemma 11.12]). The corner $p_0C^*(\Sigma')p_0$ is the C^* -subalgebra $C^*(u_0)$ generated by u_0 .

Proof. A corner $p_0C^*(\Sigma')p_0$ is a closure of a linear span of the set

$$\{p_0 t^n(\xi) t^m(\eta)^* p_0 \mid n, m \in \mathbb{N}, \xi \in C_c(U_n), \eta \in C_c(U_m)\}.$$

by Lemma 1.14. For $n, m \in \mathbb{N}, \xi \in C_c(U_n), \eta \in C_c(U_m)$, the element $p_0 t^n(\xi) t^m(\eta)^* p_0$ is $\alpha t^n(f_0) t^m(f_0)^*$ for some $\alpha \in \mathbb{C}$. We also have

$$t^{n}(f_{0})t^{m}(f_{0})^{*} = \begin{cases} p_{0} & \text{if } n = m \\ u_{0}^{k} & \text{if } n - m = n_{0}k \text{ for some } k = 1, 2, \dots \\ (u_{0}^{k})^{*} & \text{if } m - n = n_{0}k \text{ for some } k = 1, 2, \dots \\ 0 & \text{otherwise.} \end{cases}$$

Hence the corner $p_0C^*(\Sigma')p_0$ is $C^*(u_0)$.

Lemma 4.13. The corner $p_0C^*(\Sigma')p_0 = C^*(u_0)$ is full in the ideal $I_{X''}/I_{X'} \subset C^*(\Sigma)/I_{X'} = C^*(\Sigma')$.

Proof. Let I be the ideal of $C^*(\Sigma')$ generated by p_0 . Then I is gauge-invariant. Consider $X_I \subset X'$. Since X_I does not contain x_0 . we have $X_I \subset X''$. Therefore $I_{X''} \subset I_{X_I} = I$ by Lemma 3.12 and Proposition 3.10. Since $p_0 \in I_{X''}$, we have $I \subset I_{X''}$. Hence we get $I = I_{X''}$.

The ideal $I_{X''}$ of $C^*(\Sigma')$ is equal to $I_{X''}/I_{X'} \subset C^*(\Sigma)/I_{X'}$ via the identification $C^*(\Sigma)/I_{X'} = C^*(\Sigma')$. This completes the proof.

The following lemma should be known, but the author could not find it in the literature. A similar result on primitive ideals can be found, for example, in [B, II.6.5.4]

Lemma 4.14. Let A be a C^* -algebra, and I', I'' be ideals of A with $I' \subset I''$. Then $I \mapsto (I \cap I'')/I'$ is a bijection from the set of prime ideals I of A with $I' \subset I$ and $I'' \not\subset I$ to the set of prime ideals of I''/I'.

Proof. We first show that $I \mapsto I \cap I''$ is a bijection from the set of prime ideals I of A with $I'' \not\subset I$ to the set of prime ideals of I''. Take a prime ideal I of A with $I'' \not\subset I$. We show that $I \cap I''$ is a prime ideal of I'' Since $I'' \not\subset I$, we have $I \cap I'' \neq I''$. Take ideals I_1, I_2 of I'' with $I_1 \cap I_2 \subset I \cap I''$. Then I_1, I_2 are ideals of A with $I_1 \cap I_2 \subset I$. Since I is prime, either $I_1 \subset I$ or $I_2 \subset I$ holds. Thus either $I_1 \subset I \cap I''$ or $I_2 \subset I \cap I''$ holds. This shows $I \cap I''$ is a prime ideal of I''. Next take two prime ideals I_1, I_2 of A with $I'' \not\subset I_1$ and $I'' \not\subset I_2$. Suppose $I_1 \cap I'' = I_2 \cap I''$ and we will show $I_1 = I_2$. We have $I_1 \supset I_1 \cap I'' = I_2 \cap I''$. Since I_1 is prime and $I'' \not\subset I_1$, we get $I_2 \subset I_1$. Similarly

we get $I_1 \subset I_2$. Thus $I_1 = I_2$. This shows that the map $I \mapsto I \cap I''$ is injective. Now take a prime ideal P of I''. Define $I \subset A$ by

$$I := \{ a \in A \mid ab \in P \text{ for all } b \in I'' \}.$$

Then I is an ideal of A satisfying $I \cap I'' = P$. We will show I is prime. Take two ideals I_1, I_2 of A with $I_1 \cap I_2 \subset I$. Then two ideals $I_1 \cap I'', I_2 \cap I''$ of I'' satisfy $(I_1 \cap I'') \cap (I_2 \cap I'') \subset I \cap I'' = P$. Since P is a prime ideal of I'', we have either $I_1 \cap I'' \subset P$ or $I_2 \cap I'' \subset P$. When $I_1 \cap I'' \subset P$, we have $I_1 \subset I$ by the definition of I. Similarly we have $I_2 \subset I$ when $I_2 \cap I'' \subset P$. Thus we have shown that I is prime. This shows that the map $I \mapsto I \cap I''$ is surjective. Therefore the map $I \mapsto I \cap I''$ is bijective. Hence the map $I \mapsto I \cap I''$ is a bijection from the set of prime ideals I of A with $I' \subset I$ and $I'' \not\subset I$ to the set of prime ideals P of I'' with $I' \subset P$.

It is well-known and clear that $P \mapsto P/I'$ is a bijection from the set of prime ideals P of I'' with $I' \subset P$ to the set of prime ideals of I''/I'. This completes the proof.

Proposition 4.15. The prime ideals I with $X_I = \overline{\text{Orb}(x_0)}$ is parametrized as $P_{(x_0,e^{2\pi it})}$ for $0 \le t < 1/n_0$.

Proof. Since $C^*(u_0)$ is hereditary and full in the ideal $I_{X''}/I_{X'}$, the map $I \mapsto I \cap C^*(u_0)$ induces the bijection from the set of all ideals of $I_{X''}/I_{X'}$ to the one of $C^*(u_0)$ preserving the intersections and inclusion. Hence it induces the bijection between the set of prime ideals. Combine this with Lemma 4.11 and Lemma 4.14, we see that $I \mapsto I \cap C^*(u_0)$ induces the bijection from the set of prime ideals I with $X_I = \overline{\operatorname{Orb}(x_0)}$ to the set of all prime ideals of $C^*(u_0)$. The prime ideals of $C^*(u_0)$ is parametrized as Q_z for $z \in \mathbb{T}$, where Q_z is the ideal generated by $p_0 - zu_0$. Since we have $\pi_{(x_0,\gamma)}(u_0) = \gamma^{n_0}\pi_{(x_0,\gamma)}(p_0)$, we get $P_{(x_0,\gamma)} \cap C^*(u_0) = Q_{\gamma^{-n_0}}$. Hence we see that $P_{(x_0,e^{2\pi it})}$ for $0 \le t < 1/n_0$ is the parametrization of the prime ideals I with $X_I = \overline{\operatorname{Orb}(x_0)}$.

Note that for $x_0 \in P(\Sigma)$ with $n_0 = p(x_0)$, we have $P_{(x,\gamma)} = P_{(x,\mu)}$ if and only if $\gamma^{n_0} = \mu^{n_0}$ by Lemma 2.17.

Combine the discussion above, we get the following.

Theorem 4.16. The set of all prime ideals of $C^*(\Sigma)$ is

$$\{I_{X'} \mid X' \in \mathcal{A}(\Sigma)\} \cup \{P_{(x,e^{2\pi it})} \mid x \in P(\Sigma), 0 \le t < 1/p(x)\}.$$

Remark 4.17. One can show that when X is second countable for $X' \in \mathcal{A}(\Sigma)$ there exists $x \in X$ such that $X' = \overline{\operatorname{Orb}(x)}$ (cf. [K3, Proposition 4.14]). Therefore when X is second countable, $\{P_{(x,\gamma)}\}$ are all prime (primitive) ideals of $C^*(\Sigma)$.

Proposition 4.18. Every prime ideals of $C^*(\Sigma)$ is an intersection of primitive ideals in the form $P_{(x,\gamma)}$.

Proof. Take $X' \in \mathcal{A}(\Sigma)$. We set $I = \bigcap_{x \in X', \gamma \in \mathbb{T}} P_{(x,\gamma)}$. Then I is gauge-invariant. For $x \in X'$ and $\gamma \in \mathbb{T}$, we have $X_{P_{(x,\gamma)}} = \overline{\operatorname{Orb}(x)} \subset X'$. Hence $I_{X'} \subset I_{X_{P_{(x,\gamma)}}} \subset P_{(x,\gamma)}$ by Lemma 3.12 and Lemma 3.9. Therefore we get $I_{X'} \subset I$. For $x \in X'$ and $\gamma \in \mathbb{T}$, $I \subset P_{(x,\gamma)}$ implies $X_I \supset X_{P_{(x,\gamma)}} = \overline{\operatorname{Orb}(x)}$. Hence we get $X_I \supset X'$. By Proposition 3.10 and Lemma 3.12, we get $I = I_{X_I} \subset I_{X'}$. Thus we have $I_{X'} = I$. \square

Corollary 4.19. Every primitive ideals of $C^*(\Sigma)$ is an intersection of primitive ideals in the form $P_{(x,\gamma)}$.

5.
$$Y_I$$
, I_Y AND $I_{Y_I} = I$

Definition 5.1. For an ideal I of $C^*(\Sigma)$, we define $Y_I \subset X \times \mathbb{T}$ by

$$Y_I = \{(x, \gamma) \in X \times \mathbb{T} \mid I \subset P_{(x, \gamma)}\}.$$

Lemma 5.2. For two ideals I_1, I_2 of $C^*(\Sigma)$, we have $Y_{I_1 \cap I_2} = Y_{I_1} \cup Y_{I_2}$. If $I_1 \subset I_2$, we have $Y_{I_1} \supset Y_{I_2}$.

Proof. To show $Y_{I_1 \cap I_2} = Y_{I_1} \cup Y_{I_2}$, it suffices to show for $(x, \gamma) \in X \times \mathbb{T}$, $I_1 \cap I_2 \subset P_{(x,\gamma)}$ if and only if $I_1 \subset P_{(x,\gamma)}$ or $I_2 \subset P_{(x,\gamma)}$. This follows from the fact that $P_{(x,\gamma)}$ is prime. The latter assertion is clear by definition.

Definition 5.3. For a subset Y of $X \times \mathbb{T}$, we define an ideal I_Y of $C^*(\Sigma)$ by

$$I_Y = \bigcap_{(x,\gamma)\in Y} P_{(x,\gamma)}.$$

Lemma 5.4. For two subsets Y_1, Y_2 of $X \times \mathbb{T}$, we have $I_{Y_1 \cup Y_2} = I_{Y_1} \cap I_{Y_2}$. If $Y_1 \subset Y_2$, we have $I_{Y_1} \supset I_{Y_2}$.

Proof. Clear by definition.

Proposition 5.5. For an ideal I of $C^*(\Sigma)$, we have $I_{Y_I} = I$.

Proof. It is known that I is the intersection of all primitive ideals of $C^*(\Sigma)$ containing I ([B, II.6.5.3]). By Corollary 4.19, all primitive ideals of $C^*(\Sigma)$ are the intersection of ideals in the form $P_{(x,\gamma)}$. We are done.

Thus $I \mapsto Y_I$ is an injective map from the set of ideals of $C^*(\Sigma)$ to the set of subsets of $X \times \mathbb{T}$. We will determine the image of this map in order to get the complete description of ideals of $C^*(\Sigma)$.

Definition 5.6. Let Y be a subset of $X \times \mathbb{T}$. For each $x \in X$, we set

$$Y_x = \{ \gamma \in \mathbb{T} \mid (x, \gamma) \in Y \} \subset \mathbb{T}.$$

Definition 5.7. We say a subset Y of $X \times \mathbb{T}$ is said to be admissible if

- (i) Y is a closed subset of $X \times \mathbb{T}$ with respect to the product topology,
- (ii) $Y_x = Y_{\sigma(x)}$ for all $x \in U$,
- (iii) $Y_{x_0} \neq \emptyset$, \mathbb{T} implies that x_0 is periodic, $\zeta_{p(x_0)}Y_{x_0} = Y_{x_0}$, and there exists a neighborhood V of x_0 such that all $x \in V$ with $l(x) \neq l(x_0)$ satisfies $Y_x = \emptyset$.

6. Y_I is admissible

We will show that Y_I is admissible for all ideal I of $C^*(\Sigma)$ (Proposition 6.8).

For a periodic point $x_0 \in X$ with $p := p(x_0)$ and a positive integer n, we define a map $\sigma_n \colon \operatorname{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z} \to \operatorname{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z}$ as follows.

We set $y_0 = \sigma^{l(x_0)+p-1}(x_0)$. We define $d: \operatorname{Orb}(x_0) \to \{0,1\}$ by d(x) = 0 for $x \in \operatorname{Orb}(x_0) \setminus \{y_0\}$ and $d(y_0) = 1$, and a map $\sigma_n: \operatorname{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z} \to \operatorname{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z}$ by $\sigma_n(x,j) = (\sigma(x),j+d(x))$. Let $H_{x_0}^{(n)}$ be the Hilbert space whose complete orthonormal system is given by $\{\delta_{(x,j)}\}_{(x,j)\in\operatorname{Orb}(x_0)\times\mathbb{Z}/n\mathbb{Z}}$. In a similar way to define

representations $\pi_{(x_0,\gamma_0)}$, we can define a representation $\pi_{(x_0,\gamma_0)}^{(n)} \colon C^*(\Sigma) \to B(H_{x_0}^{(n)})$ such that

$$\pi_{(x_0,\gamma_0)}^{(n)}(t^0(f))\delta_{(x,j)} = f(x)\delta_{(x,j)}$$

$$\pi_{(x_0,\gamma_0)}^{(n)}(t^1(\xi))\delta_{(x,j)} = \gamma_0 \sum_{\sigma_n((y,j'))=(x,j)} \xi(y)\delta_{(y,j')} = \gamma_0 \sum_{\sigma(y)=x} \xi(y)\delta_{(y,j-d(y))}.$$

By construction, we have $\pi_{(x_0,\gamma_0)}^{(1)}=\pi_{(x_0,\gamma_0)}$. We will see that $\pi_{(x_0,\gamma_0)}^{(n)}$ is not irreducible for $n\geq 2$ in Lemma 6.2. For $x\in \mathrm{Orb}(x_0)$, let $c(x)\in \mathbb{N}$ be the smallest natural number satisfying $\sigma^{c(x)}(x)=y_0$. We have $c(\sigma(x))=c(x)-1$ for $x\in \mathrm{Orb}(x_0)\setminus \{y_0\}$, and $c(\sigma(y_0))=p-1=c(y_0)+p-1$. Hence we get $d(y)p=c(\sigma(y))-c(y)+1$ for all $y\in \mathrm{Orb}(x_0)$.

Lemma 6.1. For $k = 0, 1, \ldots, n-1$, the map $\iota_k \colon H_{x_0} \to H_{x_0}^{(n)}$ defined by

$$\iota_k(\delta_x) = \frac{1}{\sqrt{n}} \sum_{j=0}^{n-1} \zeta_{np}^{(jp-c(x))k} \delta_{(x,j)},$$

is an isometry satisfying $\iota_k \circ \pi_{(x_0,\zeta_{np}^k\gamma_0)}(a) = \pi_{(x_0,\gamma_0)}^{(n)}(a) \circ \iota_k$ for $a \in C^*(\Sigma)$.

Proof. Clearly ι_k is isometric. For every $f \in C_0(X)$ and $x \in Orb(x_0)$, we have

$$\iota_k(\pi_{(x_0,\zeta_{np}^k\gamma_0)}(t^0(f))\delta_x) = \iota_k(f(x)\delta_x) = f(x)\frac{1}{\sqrt{n}}\sum_{j=0}^{n-1}\zeta_{np}^{(jp-c(x))k}\delta_{(x,j)},$$

and

$$\pi_{(x_0,\gamma_0)}^{(n)}(t^0(f))(\iota_k(\delta_x)) = \pi_{(x_0,\gamma_0)}^{(n)}(t^0(f))\left(\frac{1}{\sqrt{n}}\sum_{j=0}^{n-1}\zeta_{np}^{(jp-c(x))k}\delta_{(x,j)}\right)$$
$$= \frac{1}{\sqrt{n}}\sum_{j=0}^{n-1}\zeta_{np}^{(jp-c(x))k}f(x)\delta_{(x,j)}.$$

Hence $\iota_k \circ \pi_{(x_0,\zeta_{np}^k\gamma_0)}(a) = \pi_{(x_0,\gamma_0)}^{(n)}(a) \circ \iota_k$ for $a \in t^0(C_0(X))$. For every $\xi \in C_c(U)$ and $x \in \operatorname{Orb}(x_0)$, we have

$$\iota_{k}\left(\pi_{(x_{0},\zeta_{np}^{k}\gamma_{0})}(t^{1}(\xi))\delta_{x}\right) = \iota_{k}\left(\zeta_{np}^{k}\gamma_{0}\sum_{\sigma(y)=x}\xi(y)\delta_{y}\right)$$

$$= \zeta_{np}^{k}\gamma_{0}\sum_{\sigma(y)=x}\xi(y)\left(\frac{1}{\sqrt{n}}\sum_{j=0}^{n-1}\zeta_{np}^{(jp-c(y))k}\delta_{(y,j)}\right)$$

$$= \frac{1}{\sqrt{n}}\gamma_{0}\sum_{\sigma(y)=x}\xi(y)\sum_{j=0}^{n-1}\zeta_{np}^{(jp-c(y)+1)k}\delta_{(y,j)},$$

and

$$\pi_{(x_{0},\gamma_{0})}^{(n)}(t^{1}(\xi))(\iota_{k}(\delta_{x})) = \pi_{(x_{0},\gamma_{0})}^{(n)}(t^{1}(\xi))\left(\frac{1}{\sqrt{n}}\sum_{j=0}^{n-1}\zeta_{np}^{(jp-c(x))k}\delta_{(x,j)}\right)$$

$$= \frac{1}{\sqrt{n}}\sum_{j=0}^{n-1}\zeta_{np}^{(jp-c(x))k}\left(\gamma_{0}\sum_{\sigma(y)=x}\xi(y)\delta_{(y,j-d(y))}\right)$$

$$= \frac{1}{\sqrt{n}}\gamma_{0}\sum_{\sigma(y)=x}\xi(y)\sum_{j=0}^{n-1}\zeta_{np}^{(jp-c(x))k}\delta_{(y,j-d(y))}$$

$$= \frac{1}{\sqrt{n}}\gamma_{0}\sum_{\sigma(y)=x}\xi(y)\sum_{j=0}^{n-1}\zeta_{np}^{((j+d(y))p-c(x))k}\delta_{(y,j)}$$

$$= \frac{1}{\sqrt{n}}\gamma_{0}\sum_{\sigma(y)=x}\xi(y)\sum_{j=0}^{n-1}\zeta_{np}^{((jp-c(y)+1)k}\delta_{(y,j)}$$

where in the last equality we use the fact that d(y)p = c(x) - c(y) + 1 for $y \in Orb(x_0)$ with $\sigma(y) = x$. Hence $\iota_k \circ \pi_{(x_0, \zeta_{np}^k \gamma_0)}(a) = \pi_{(x_0, \gamma_0)}^{(n)}(a) \circ \iota_k$ for $a \in t^1(C_c(U))$. Since $C^*(\Sigma)$ is generated by $t^0(C_0(X)) \cup t^1(C_c(U))$, we have $\iota_k \circ \pi_{(x_0,\zeta_{np}^k\gamma_0)}(a) = \pi_{(x_0,\gamma_0)}^{(n)}(a) \circ \iota_k$ for every $a \in C^*(\Sigma)$.

Lemma 6.2. The representation $\pi_{(x_0,\gamma_0)}^{(n)}$ is unitarily equivalent to $\bigoplus_{k=0}^{n-1} \pi_{(x_0,\zeta_{np}^k\gamma_0)}$. Hence we have $\ker \pi_{(x_0,\gamma_0)}^{(n)} = \bigcap_{k=0}^{n-1} P_{(x_0,\zeta_{nx}^k\gamma_0)}$.

Proof. For $k=0,1,\ldots,n-1$, let ι_k be the isometies in Lemma 6.1. Then for each $x\in \mathrm{Orb}(x_0),\ \{\iota_k(\delta_x)\}_{k=0}^{n-1}$ are complete orthonormal system of the linear space spanned by $\{\delta_{(x,j)} \mid j \in \mathbb{Z}/n\mathbb{Z}\}$. Hence $\bigoplus_{k=0}^{n-1} \iota_k$ is a unitary from $\bigoplus_{k=0}^{n-1} H_{x_0}$ to $H_{x_0}^{(n)}$ which intertwines $\bigoplus_{k=0}^{n-1} \pi_{(x_0,\zeta_{np}^k\gamma_0)}$ and $\pi_{(x_0,\gamma_0)}^{(n)}$. This completes the proof.

Note that Lemma 2.15 implies $P_{(x_0,\zeta_{np}^k\gamma_0)} = P_{(x_0,\zeta_{np}^{k+n}\gamma_0)}$. Hence we get $\ker \pi_{(x_0,\gamma_0)}^{(n)} =$ $\bigcap_{k\in\mathbb{Z}} P_{(x_0,\zeta_{n_p}^k\gamma_0)}.$

Lemma 6.3. Let $l, m \in \mathbb{N}$ with $m \geq 1$. Let $\{(x_{\lambda}, \gamma_{\lambda})\}_{{\lambda} \in \Lambda}$ be a net in $X \times \mathbb{T}$ converges to (x_0, γ_0) such that $l(x_\lambda) = l$ and $p(x_\lambda) = m$ for all $\lambda \in \Lambda$. Then we have the following.

- (i) $l(x_0) = l$ and $p(x_0) = m/n$ for some $n \in \mathbb{N}$.
- (ii) for each $(x,j) \in \text{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z}$, there exist $\lambda_{(x,j)} \in \Lambda$ and $\varphi_{\lambda}(x,j) \in \mathbb{Z}$ $\operatorname{Orb}(x_{\lambda}) \text{ for } \lambda \succeq \lambda_{(x,j)} \text{ satisfying }$
 - $\lim_{\lambda} \varphi_{\lambda}(x,j) = x$ for all (x,j),
 - $\sigma(\varphi_{\lambda}(x,j)) = \varphi_{\lambda}(\sigma_{n}(x,j))$ for all (x,j) and all λ with $\lambda \succeq \lambda_{(x,j)}$ and
- for two distinct elements $(x,j),(x',j') \in \operatorname{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z}$, we have $\varphi_{\lambda}(x,j) \neq \varphi_{\lambda}(x',j')$ for sufficiently large λ . (iii) $\bigcap_{\lambda} P_{(x_{\lambda},\gamma_{\lambda})} \subset \ker \pi_{(x_{0},\gamma_{0})}^{(n)}$.

Proof. (i) For each λ , we have $\sigma^{l+m}(x_{\lambda}) = \sigma^{l}(x_{\lambda})$. Hence we get $\sigma^{l+m}(x_{0}) = \sigma^{l}(x_{0})$. This implies that x_0 is a periodic point whose period $p(x_0)$ divides m. Thus we can find $n \in \mathbb{N}$ with $p(x_0) = m/n$. We clearly have $l(x_0) \leq l$. To drive a contradiction, assume $l(x_0) = k < l$. Then we have $\sigma^{k+m}(x_0) = \sigma^k(x_0)$. Since σ^{l-k} is locally homeomorphic, there exists a neighborhood V of $\sigma^{k+m}(x_0) = \sigma^k(x_0)$ such that σ^{l-k} is injective on V. We can find λ such that $\sigma^{k+m}(x_{\lambda}), \sigma^{k}(x_{\lambda}) \in V$. Since k < l, we have $\sigma^{k+m}(x_{\lambda}) \neq \sigma^{k}(x_{\lambda})$. However, we have

$$\sigma^{l-k}(\sigma^{k+m}(x_{\lambda})) = \sigma^{l+m}(x_{\lambda}) = \sigma^{l}(x_{\lambda}) = \sigma^{l-k}(\sigma^{k}(x_{\lambda})).$$

This is a contradiction. Hence $l(x_0) = l$.

(ii) Let us set $p = p(x_0)$, and $x^{(i)} = \sigma^{l+i}(x_0)$ for i = 0, ..., p-1. Take $(x, j) \in$ $\operatorname{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z}$. Let k := l(x) which is the smallest natural number with $\sigma^{k+p}(x) =$ $\sigma^k(x)$. Choose $i \in \{0, 1, \dots, p-1\}$ such that $\sigma^k(x) = x^{(i)}$. Choose a neighborhood V of x such that σ^k is injective on V. Since $\sigma^{l+i+jp}(x_0) = x^{(i)}$, there exists $\lambda_{(x,j)} \in \Lambda$ such that we have $\sigma^{l+i+jp}(x_{\lambda}) \in \sigma^{k}(V)$ for all $\lambda \succeq \lambda_{(x,j)}$. For $\lambda \succeq \lambda_{(x,j)}$, define $\varphi_{\lambda}(x,j)$ to be the element in V with $\sigma^{k}(\varphi_{\lambda}(x,j)) = \sigma^{l+i+jp}(x_{\lambda})$. Since the restriction of σ^k to V is a homeomorphism and $\lim_{\lambda} \sigma^{l+i+jp}(x_{\lambda}) = \sigma^{l+i+jp}(x_0) = x^{(i)}$, we get $\lim_{\lambda} \varphi_{\lambda}(x,j) = x$. By construction, it is easy to see that $\sigma(\varphi_{\lambda}(x,j)) = \varphi_{\lambda}(\sigma_{n}(x,j))$ for $(x,j) \in (\mathrm{Orb}(x_0) \setminus \{x^{(p-1)}\}) \times \mathbb{Z}/n\mathbb{Z}$ and $\lambda \in \Lambda$ with $\lambda \succeq \lambda_{(x,j)}$ and $\lambda \succeq \lambda_{(x,j)}$ $\lambda_{\sigma_n(x,j)}$. For $(x^{(p-1)},j) \in \operatorname{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z}$, we have $\sigma_n(x^{(p-1)},j) = (x^{(0)},j+1)$. We get $\varphi_{\lambda}(x^{(p-1)},j) = \sigma^{l+p-1+jp}(x_{\lambda})$ and $\varphi_{\lambda}(x^{(0)},j+1) = \sigma^{l+0+(j+1)p}(x_{\lambda})$. Hence the equation $\sigma(\varphi_{\lambda}(x^{(p-1)},j)) = \varphi_{\lambda}(\sigma_{n}(x^{(p-1)},j))$ is easy to see when $j \neq n-1$, and follows from the fact that $\sigma^{l+np}(x_{\lambda}) = \sigma^{l}(x_{\lambda})$ when j = n-1. Take two distinct elements $(x,j), (x',j') \in \text{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z}$. If $x \neq x'$, then $\varphi_{\lambda}(x,j) \neq \varphi_{\lambda}(x',j')$ for sufficiently large λ because $\lim_{\lambda} \varphi_{\lambda}(x,j) = x$ and $\lim_{\lambda} \varphi_{\lambda}(x',j') = x'$. If x = x', then $j \neq j'$. Take k := l(x) and $i \in \{0, 1, \dots, p-1\}$ with $\sigma^k(x) = x^{(i)}$. For every λ we have $\sigma^{l+i+jp}(x_{\lambda}) \neq \sigma^{l+i+j'p}(x_{\lambda})$ because 0 < |jp-j'p| < m. Hence $\varphi_{\lambda}(x,j) \neq \varphi_{\lambda}(x',j')$ when they are defined because $\sigma^{k}(\varphi_{\lambda}(x,j)) = \sigma^{l+i+jp}(x_{\lambda})$ and $\sigma^k(\varphi_\lambda(x',j')) = \sigma^{l+i+j'p}(x_\lambda).$

(iii) Take $(x,j), (x',j') \in \text{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z}$. We will show that for all $a \in C^*(\Sigma)$ we have

$$\lim_{\lambda} \langle \delta_{\varphi_{\lambda}(x',j')}, \pi_{(x_{\lambda},\gamma_{\lambda})}(a) \delta_{\varphi_{\lambda}(x,j)} \rangle_{x_{\lambda}} = \langle \delta_{(x',j')}, \pi_{(x_{0},\gamma_{0})}^{(n)}(a) \delta_{(x,j)} \rangle_{x_{0}}^{(n)},$$

where $\langle \cdot, \cdot \rangle_{x_0}^{(n)}$ is the inner product of $H_{x_0}^{(n)}$ which is linear in the second variable. To do so, it suffices to assume that $a = \sum_{k=1}^K t^{n_k}(\xi_k)t^{m_k}(\eta_k)^* \in C^*(\Sigma)$ for some $K, n_k, m_k \in \mathbb{N}, \, \xi_k \in C_c(U_{n_k})$ and $\eta_k \in C_c(U_{m_k})$ for $k = 1, \ldots, K$ by Lemma 1.14. We can find $\lambda_0 \in \Lambda$ such that

- $\lambda_0 \succeq \lambda_{(x,j)}$ and $\lambda_0 \succeq \lambda_{(x',j')}$,
- $\lambda_0 \succeq \lambda_{\sigma_n^{m_k}(x,j)}$ and $\lambda_0 \succeq \lambda_{\sigma_n^{n_k}(x',j')}$ for $k = 1, \ldots, K$, φ_{λ} is injective on $\{\sigma_n^{m_k}(x,j), \sigma_n^{n_k}(x',j') \mid k = 1, \ldots, K\}$ for all $\lambda \succeq \lambda_0$.

For $\lambda \succeq \lambda_0$, we have

$$\pi_{(x_{\lambda},\gamma_{\lambda})}(a)\delta_{\varphi_{\lambda(x,j)}} = \sum_{k=1}^{K} \pi_{(x_{\lambda},\gamma_{\lambda})} \left(t^{n_{k}}(\xi_{k}) t^{m_{k}}(\eta_{k})^{*} \right) \delta_{\varphi_{\lambda}(x,j)}$$

$$= \sum_{k=1}^{K} \pi_{(x_{\lambda},\gamma_{\lambda})} \left(t^{n_{k}}(\xi_{k}) \right) \left(\gamma_{\lambda}^{-m_{k}} \overline{\eta_{k}}(\varphi_{\lambda}(x,j)) \delta_{\sigma^{m_{k}}(\varphi_{\lambda}(x,j))} \right)$$

$$= \sum_{k=1}^{K} \sum_{\sigma^{n_{k}}(y) = \sigma^{m_{k}}(\varphi_{\lambda}(x,j))} \gamma_{\lambda}^{n_{k}} \xi_{k}(y) \gamma_{\lambda}^{-m_{k}} \overline{\eta_{k}}(\varphi_{\lambda}(x,j)) \delta_{y}.$$

Hence we get

$$\langle \delta_{\varphi_{\lambda}(x',j')}, \pi_{(x_{\lambda},\gamma_{\lambda})}(a) \delta_{\varphi_{\lambda}(x,j)} \rangle_{x_{\lambda}} = \sum_{k \in N_{\lambda}} \gamma_{\lambda}^{n_{k}-m_{k}} \xi_{k}(\varphi_{\lambda}(x',j')) \overline{\eta_{k}}(\varphi_{\lambda}(x,j)),$$

where

$$N_{\lambda} = \{k \in \{1, \dots, K\} \mid \sigma^{n_k}(\varphi_{\lambda}(x', j')) = \sigma^{m_k}(\varphi_{\lambda}(x, j))\}.$$

Similarly, we get

$$\langle \delta_{(x',j')}, \pi_{(x_0,\gamma_0)}^{(n)}(a)\delta_{(x,j)}\rangle_{x_0}^{(n)} = \sum_{k\in N_0} \gamma_0^{n_k-m_k} \xi_k(x')\overline{\eta_k}(x),$$

where

$$N_0 = \{k \in \{1, \dots, K\} \mid \sigma_n^{n_k}(x', j') = \sigma_n^{m_k}(x, j)\}.$$

Since

$$\sigma^{n_k}(\varphi_{\lambda}(x',j')) = \varphi_{\lambda}(\sigma^{n_k}_n(x',j')), \quad \sigma^{m_k}(\varphi_{\lambda}(x,j)) = \varphi_{\lambda}(\sigma^{m_k}_n(x,j))$$

and φ_{λ} is injective on $\{\sigma_{n}^{m_{k}}(x,j), \sigma_{n}^{n_{k}}(x',j') \mid k=1,\ldots,K\}$, we have $N_{\lambda}=N_{0}$ for $\lambda \succeq \lambda_{0}$. Since $\lim_{\lambda} \varphi_{\lambda}(x,j)=x$ and $\lim_{\lambda} \varphi_{\lambda}(x',j')=x'$, we have

$$\lim_{\lambda} \langle \delta_{\varphi_{\lambda}(x',j')}, \pi_{(x_{\lambda},\gamma_{\lambda})}(a) \delta_{\varphi_{\lambda}(x,j)} \rangle_{x_{\lambda}} = \langle \delta_{(x',j')}, \pi_{(x_{0},\gamma_{0})}^{(n)}(a) \delta_{(x,j)} \rangle_{x_{0}}^{(n)}.$$

Thus we get this equality for all $(x, j), (x', j') \in \text{Orb}(x_0) \times \mathbb{Z}/n\mathbb{Z}$ and all $a \in C^*(\Sigma)$. Hence for $a \in \bigcap_{\lambda} P_{(x_{\lambda}, \gamma_{\lambda})}$ we have $\pi_{(x_0, \gamma_0)}^{(n)}(a) = 0$. We are done.

Corollary 6.4. Under the condition in Lemma 6.3, we have

$$\bigcap_{\lambda} P_{(x_{\lambda}, \gamma_{\lambda})} \subset P_{(x_{0}, \gamma_{0})}.$$

Proof. This follows from Lemma 6.3 (iii) and Lemma 6.2.

For a periodic point $x_0 \in X$ with $p := p(x_0)$, we define a representation $\pi_{(x_0,\gamma_0)}^{(\infty)} \colon C^*(\Sigma) \to B(H_{x_0}^{(\infty)})$ in a similar way to the definition of $\pi_{(x_0,\gamma_0)}^{(n)}$ where $H_{x_0}^{(\infty)}$ is the Hilbert space whose complete orthonormal system is given by $\{\delta_{(x,j)}\}_{(x,j)\in Orb(x_0)\times\mathbb{Z}}$ using the map $\sigma_{\infty} \colon Orb(x_0) \times \mathbb{Z} \to Orb(x_0) \times \mathbb{Z}$ defined by $\sigma_{\infty}(x,j) = (\sigma(x),j+d(x))$. Let us define a probability space

$$[1, \zeta_p) = \{e^{2\pi i\theta/p} \mid 0 \le \theta < 1\}$$

on which we consider the measure $d\gamma$ defined from the Lebesgue measure via the bijection $[0,1) \ni \theta \mapsto e^{2\pi i\theta/p} \in [1,\zeta_p)$. The Hilbert space $L^2([1,\zeta_p),H_{x_0})$ of the all

 H_{x_0} -valued second power integrable functions on $[1, \zeta_p)$ has spanned by the elements in the form $\eta \delta_x$ with $\eta \in L^2([1, \zeta_p))$ and $x \in \text{Orb}(x_0)$. We define a representation $\pi_{(x_0, [\gamma_0, \zeta_p \gamma_0))} : C^*(\Sigma) \to B(L^2([1, \zeta_p), H_{x_0}))$ by

$$(\pi_{(x_0,[\gamma_0,\zeta_p\gamma_0))}(a)\Psi)(\gamma) = \pi_{(x_0,\gamma\gamma_0)}(a)\Psi(\gamma) \in H_{x_0}$$

for $\Psi \in L^2([1,\zeta_p), H_{x_0})$ and $\gamma \in [1,\zeta_p)$.

Lemma 6.5. The map $\iota: L^2([1,\zeta_p),H_{x_0}) \to H_{x_0}^{(\infty)}$ defined by

$$\iota(\eta \delta_x) = \sum_{j \in \mathbb{Z}} \left(\int_{[1,\zeta_p)} \gamma^{jp-c(x)} \eta(\gamma) d\gamma \right) \delta_{(x,j)},$$

for $\eta \in L^2([1,\zeta_p))$ and $x \in \operatorname{Orb}(x_0)$ is a unitary satisfying $\iota \circ \pi_{(x_0,[\gamma_0,\zeta_p\gamma_0))}(a) = \pi_{(x_0,\gamma_0)}^{(\infty)}(a) \circ \iota$ for $a \in C^*(\Sigma)$.

Proof. For each $x \in \operatorname{Orb}(x_0)$, the map $L^2([1,\zeta_p)) \ni \eta(\cdot) \mapsto (\cdot)^{-c(x)}\eta(\cdot) \in L^2([1,\zeta_p))$ is a unitary. Since $[1,\zeta_p) \ni \gamma \to \gamma^p \in \mathbb{T}$ is an isomorphism between probability spaces, the natural isomorphism $L^2(\mathbb{T}) \cong \ell^2(\mathbb{Z})$ of the Fourier transform shows that the map

$$L^{2}([1,\zeta_{p})) \ni \eta \mapsto \sum_{j\in\mathbb{Z}} \left(\int_{[1,\zeta_{p})} \gamma^{jp-c(x)} \eta(\gamma) d\gamma \right) \delta_{(x,j)}$$

is an isomorphism onto the Hilbert subspace spanned by $\{\delta_{(x,j)}\}_{j\in\mathbb{Z}}$. Thus ι is a unitary.

Take $\xi \in C_c(U)$, $\eta \in L^2([1,\zeta_p))$ and $x \in Orb(x_0)$. For each $\gamma \in [1,\zeta_p)$, we obtain

$$\left(\pi_{(x_0,[\gamma_0,\zeta_p\gamma_0))}(t^1(\xi))\eta\delta_x\right)(\gamma) = \pi_{(x_0,\gamma\gamma_0)}(t^1(\xi))(\eta(\gamma)\delta_x)$$

$$= \gamma\gamma_0 \sum_{\sigma(y)=x} \xi(y)\eta(\gamma)\delta_y.$$

If we denote by $z \in L^2([1,\zeta_p))$ the identity function $\gamma \mapsto \gamma$, then we have

$$\pi_{(x_0,[\gamma_0,\zeta_p\gamma_0))}(t^1(\xi))\eta\delta_x = \gamma_0 \sum_{\sigma(y)=x} \xi(y)\eta z\delta_y.$$

Hence

$$\iota\left(\pi_{(x_0,[\gamma_0,\zeta_p\gamma_0))}(t^1(\xi))\eta\delta_x\right) = \iota\left(\gamma_0\sum_{\sigma(y)=x}\xi(y)\eta z\delta_y\right)$$

$$= \gamma_0\sum_{\sigma(y)=x}\xi(y)\sum_{j\in\mathbb{Z}}\left(\int_{[1,\zeta_p)}\gamma^{jp-c(y)}\eta(\gamma)\gamma d\gamma\right)\delta_{(y,j)}$$

$$= \gamma_0\sum_{\sigma(y)=x}\sum_{j\in\mathbb{Z}}\left(\int_{[1,\zeta_p)}\gamma^{jp-c(y)+1}\eta(\gamma)d\gamma\right)\xi(y)\delta_{(y,j)}.$$

On the other hand, we get

$$\pi_{(x_{0},\gamma_{0})}^{(\infty)}(t^{1}(\xi))(\iota(\eta\delta_{x})) = \pi_{(x_{0},\gamma_{0})}^{(\infty)}(t^{1}(\xi))\left(\sum_{j\in\mathbb{Z}}\left(\int_{[1,\zeta_{p})}\gamma^{jp-c(x)}\eta(\gamma)d\gamma\right)\delta_{(x,j)}\right)$$

$$= \sum_{j\in\mathbb{Z}}\left(\int_{[1,\zeta_{p})}\gamma^{jp-c(x)}\eta(\gamma)d\gamma\right)\gamma_{0}\sum_{\sigma(y)=x}\xi(y)\delta_{(y,j-d(y))}$$

$$= \gamma_{0}\sum_{\sigma(y)=x}\sum_{j\in\mathbb{Z}}\left(\int_{[1,\zeta_{p})}\gamma^{jp-c(x)}\eta(\gamma)d\gamma\right)\xi(y)\delta_{(y,j-d(y))}$$

$$= \gamma_{0}\sum_{\sigma(y)=x}\sum_{j\in\mathbb{Z}}\left(\int_{[1,\zeta_{p})}\gamma^{jp-c(y)+1}\eta(\gamma)d\gamma\right)\xi(y)\delta_{(y,j)}$$

$$= \gamma_{0}\sum_{\sigma(y)=x}\sum_{j\in\mathbb{Z}}\left(\int_{[1,\zeta_{p})}\gamma^{jp-c(y)+1}\eta(\gamma)d\gamma\right)\xi(y)\delta_{(y,j)}.$$

Hence we have $\iota \circ \pi_{(x_0,[\gamma_0,\zeta_p\gamma_0))}(a) = \pi_{(x_0,\gamma_0)}^{(\infty)}(a) \circ \iota$ for all $a \in t^1(C_c(U))$. The proof of this equality for $a \in t^0(C_0(X))$ is much easier. Therefore we get the equality for all $a \in C^*(\Sigma)$.

Corollary 6.6. We have $\ker \pi_{(x_0,\gamma_0)}^{(\infty)} = \bigcap_{\gamma \in \mathbb{T}} P_{(x_0,\gamma)}$.

Proof. By Lemma 6.5, we have

$$\ker \pi_{(x_0,\gamma_0)}^{(\infty)} = \ker \pi_{(x_0,[\gamma_0,\zeta_p\gamma_0))}.$$

Since $\gamma \mapsto \pi_{(x_0,\gamma)}$ is pointwise norm continuous, we have

$$\ker \pi_{(x_0,[\gamma_0,\zeta_p\gamma_0))} = \bigcap_{\gamma \in [\gamma_0,\zeta_p\gamma_0)} P_{(x_0,\gamma)}.$$

By Lemma 2.15, we get

$$\bigcap_{\gamma \in [\gamma_0, \zeta_p \gamma_0)} P_{(x_0, \gamma)} = \bigcap_{\gamma \in \mathbb{T}} P_{(x_0, \gamma)}.$$

This completes the proof.

As usual, we consider $\mathbb{N} \cup \{\infty\}$ as the one-point compactification of the discrete set \mathbb{N} .

Lemma 6.7. Let $\{(x_{\lambda}, \gamma_{\lambda})\}_{{\lambda} \in \Lambda}$ be a net in $X \times \mathbb{T}$ converges to (x_0, γ_0) such that $l(x_{\lambda}) + p(x_{\lambda})$ converges to ∞ . Then we have $\bigcap_{\lambda} P_{(x_{\lambda}, \gamma_{\lambda})} \subset P_{(x_0, \gamma)}$ for all $\gamma \in \mathbb{T}$.

Proof. We divide the proof into two cases; the case that x_0 is periodic and the case that x_0 is aperiodic. In the both cases, the proofs go in a similar way to the proof of Lemma 6.3, and so we just sketch the proofs.

First consider the case that x_0 is periodic with period $p(x_0) = p$. We first show that for each $(x, j) \in \text{Orb}(x_0) \times \mathbb{Z}$, there exist $\lambda_{(x,j)} \in \Lambda$ and $\varphi_{\lambda}(x,j) \in \text{Orb}(x_{\lambda})$ for $\lambda \succeq \lambda_{(x,j)}$ satisfying

- $\lim_{\lambda} \varphi_{\lambda}(x,j) = x$,
- $\sigma(\varphi_{\lambda}(x,j)) = \varphi_{\lambda}(\sigma_{\infty}(x,j)).$

• for two distinct elements $(x, j), (x', j') \in \text{Orb}(x_0) \times \mathbb{Z}$, we have $\varphi_{\lambda}(x, j) \neq \varphi_{\lambda}(x', j')$ for sufficiently large λ .

We set $x'_0 := \sigma^{l(x_0)}(x_0)$. For $(x'_0, 0) \in \text{Orb}(x_0) \times \mathbb{Z}$, we set $\varphi_{\lambda}(x'_0, 0) = \sigma^{l(x_0)}(x_{\lambda})$ for all λ . There exists $\lambda_{(x'_0,1)} \in \Lambda$ such that $\varphi_{\lambda}(x'_0,0) \in U$ for all $\lambda \succeq \lambda_{(x'_0,0)}$. Set $\varphi_{\lambda}(\sigma(x'_0,0)) = \sigma(\varphi_{\lambda}(x'_0,0))$ for all $\lambda \succeq \lambda_{(x'_0,0)}$. Similarly for $k=2,3,\ldots,p-1$ $\varphi_{\lambda}(\sigma^{k}(x'_{0}),0) = \sigma(\varphi_{\lambda}(\sigma^{k-1}(x'_{0}),0))$ for sufficiently large λ . We also set for sufficiently large λ , $\varphi_{\lambda}(x'_0, 1) = \sigma(\varphi_{\lambda}(\sigma^{p-1}(x'_0), 0))$ and $\varphi_{\lambda}(\sigma^k(x'_0), 1) = \sigma(\varphi_{\lambda}(\sigma^{k-1}(x'_0), 1))$ for $k=1,2,\ldots,p-1$. For $j=2,3,\ldots$, we set for sufficiently large $\lambda,\ \varphi_{\lambda}(x'_0,j)=$ $\sigma(\varphi_{\lambda}(\sigma^{p-1}(x'_0), j-1))$ and $\varphi_{\lambda}(\sigma^k(x'_0), j) = \sigma(\varphi_{\lambda}(\sigma^{k-1}(x'_0), j))$ for $k = 1, 2, \dots, p-1$. Next we choose a neighborhood V of x'_0 such that σ^p is injective on V. We set $\lambda_{(x_0',-1)} \in \Lambda$ such that $\varphi_{\lambda}(x_0',0) \in \sigma^p(V)$ for all $\lambda \succeq \lambda_{(x_0',-1)}$. We set $\varphi_{\lambda}(x_0',-1) \in V$ such that $\sigma^p(\varphi_\lambda(x_0',-1)) = \varphi_\lambda(x_0',0)$ and set $\varphi_\lambda(\sigma^k(x_0'),-1) = \sigma(\varphi_\lambda(\sigma^{k-1}(x_0'),-1))$ for k = 1, 2, ..., p - 1. We set $\lambda_{(x'_0, -2)} \in \Lambda$ such that $\varphi_{\lambda}(x'_0, -1) \in \sigma^p(V)$ for all $\lambda \succeq \lambda_{(x'_0,-2)}$. We set $\varphi_{\lambda}(x'_0,-2) \in V$ such that $\sigma^p(\varphi_{\lambda}(x'_0,-2)) = \varphi_{\lambda}(x'_0,-1)$ and set $\varphi_{\lambda}(\sigma^{k}(x'_{0}), -2) = \sigma(\varphi_{\lambda}(\sigma^{k-1}(x'_{0}), -2))$ for k = 1, 2, ..., p-1. For j = 3, 4, ...,we set $\lambda_{(x'_0,-j)} \in \Lambda$ such that $\varphi_{\lambda}(x'_0,-j+1) \in \sigma^p(V)$ for all $\lambda \succeq \lambda_{(x'_0,-j)}$. We set $\varphi_{\lambda}(x'_0,-j) \in V$ such that $\sigma^p(\varphi_{\lambda}(x'_0,-j)) = \varphi_{\lambda}(x'_0,-j+1)$ and set $\varphi_{\lambda}(\sigma^k(x'_0),-j) = \varphi_{\lambda}(x'_0,-j+1)$ $\sigma(\varphi_{\lambda}(\sigma^{k-1}(x'_0), -j))$ for $k = 1, 2, \dots, p-1$. We have defined $\varphi_{\lambda}(x, j)$ for all $(x, j) \in$ $\operatorname{Orb}(x_0) \times \mathbb{Z}$ with l(x) = 0. For $(x,j) \in \operatorname{Orb}(x_0) \times \mathbb{Z}$ with $l(x) \geq 1$, we choose a neighborhood V of x such that σ^k is injective on V where k := l(x). We set $\lambda_{(x,j)} \in \Lambda$ such that $\varphi_{\lambda}(\sigma^k(x),j) \in \sigma^k(V)$ for all $\lambda \succeq \lambda_{(x,j)}$. We set $\varphi_{\lambda}(x,j) \in V$ such that $\sigma^k(\varphi_\lambda(x,j)) = \varphi_\lambda(\sigma^k(x),j)$. It is very similar as in the proof of Lemma 6.3 (ii) to check that $\varphi_{\lambda}(x,j)$'s satisfy the desired conditions except the proof of the statement that for $x \in \operatorname{Orb}(x_0)$ and $j, j' \in \mathbb{Z}$ with j < j', we have $\varphi_{\lambda}(x, j) \neq \varphi_{\lambda}(x, j')$ for sufficiently large λ . This follows from the fact that, we have

$$\sigma^{l(x)+N}(\varphi_{\lambda}(x,j)) = \sigma^{N'}(x_{\lambda}), \qquad \sigma^{l(x)+N}(\varphi_{\lambda}(x,j')) = \sigma^{N'+p(j'-j)}(x_{\lambda})$$

for sufficiently large N and some N'. Since $l(x_{\lambda}) + p(x_{\lambda})$ converges to ∞ we have $\sigma^{N'}(x_{\lambda}) \neq \sigma^{N'+p(j'-j)}(x_{\lambda})$ for sufficiently large λ . This shows $\varphi_{\lambda}(x,j) \neq \varphi_{\lambda}(x,j')$ for sufficiently large λ .

Now, in a similar to the proof of Lemma 6.3 (iii), we can show that for all $(x, j), (x', j') \in \text{Orb}(x_0) \times \mathbb{Z}$ and all $a \in C^*(\Sigma)$, we get

$$\lim_{\lambda} \langle \delta_{\varphi_{\lambda}(x',j')}, \pi_{(x_{\lambda},\gamma_{\lambda})}(a) \delta_{\varphi_{\lambda}(x,j)} \rangle_{x_{\lambda}} = \langle \delta_{(x',j')}, \pi_{(x_{0},\gamma_{0})}^{(\infty)}(a) \delta_{(x,j)} \rangle_{x_{0}}^{(\infty)}$$

where $\langle \cdot, \cdot \rangle_{x_0}^{(\infty)}$ is the inner product of $H_{x_0}^{(\infty)}$. Hence we have $\bigcap_{\lambda} P_{(x_{\lambda}, \gamma_{\lambda})} \subset \ker \pi_{(x_0, \gamma_0)}^{(\infty)}$. We finish the proof for the case that x_0 is periodic by Corollary 6.6.

For the case that x_0 is aperiodic, we can similarly construct $\varphi_{\lambda}(x) \in \text{Orb}(x_{\lambda})$ for $x \in \text{Orb}(x_0)$ and sufficiently large $\lambda \in \Lambda$, so that we get

$$\lim_{\lambda} \langle \delta_{\varphi_{\lambda}(x')}, \pi_{(x_{\lambda}, \gamma_{\lambda})}(a) \delta_{\varphi_{\lambda}(x)} \rangle_{x_{\lambda}} = \langle \delta_{x'}, \pi_{(x_{0}, \gamma_{0})}(a) \delta_{x} \rangle_{x_{0}},$$

for all $x, x' \in \text{Orb}(x_0)$ and all $a \in C^*(\Sigma)$. Hence we have $\bigcap_{\lambda} P_{(x_{\lambda}, \gamma_{\lambda})} \subset P_{(x_0, \gamma_0)}$. Since $P_{(x_0, \gamma_0)} = P_{(x_0, \gamma_0)}$ for all $\gamma \in \mathbb{T}$ by Lemma 2.14, we finish the case that x_0 is aperiodic.

Proposition 6.8. For an ideal I, the set Y_I is admissible.

Proof. Take a net $\{(x_{\lambda}, \gamma_{\lambda})\}_{{\lambda} \in \Lambda}$ in Y_I converges to $(x_0, \gamma_0) \in X \times \mathbb{T}$. By replacing the net to a subnet if necessary, we may assume that either $l(x_{\lambda})$ and $p(x_{\lambda})$ are finite and constant or $l(x_{\lambda}) + p(x_{\lambda})$ converges ∞ . When $l(x_{\lambda})$ and $p(x_{\lambda})$ are finite and constant, we have

$$I \subset \bigcap_{\lambda} P_{(x_{\lambda}, \gamma_{\lambda})} \subset P_{(x_{0}, \gamma_{0})}$$

by Corollary 6.4. When $l(x_{\lambda}) + p(x_{\lambda})$ converges ∞ , we have

$$I \subset \bigcap_{\lambda} P_{(x_{\lambda}, \gamma_{\lambda})} \subset P_{(x_{0}, \gamma_{0})}$$

by Lemma 6.7. Hence we get $(x_0, \gamma_0) \in Y_I$. This shows that Y_I is closed. By Lemma 2.12, Y_I satisfies the condition (ii) in Definition 5.7.

Take $x_0 \in X$ with $(Y_I)_{x_0} \neq \emptyset$, \mathbb{T} . By Lemma 2.14, we see that x_0 is periodic, and by Lemma 2.15 we see that $\zeta_{p(x_0)}(Y_I)_{x_0} = (Y_I)_{x_0}$. To the contrary, suppose that for any neighborhood V of x_0 we can find $x \in V$ with $l(x) \neq l(x_0)$ satisfying $(Y_I)_x \neq \emptyset$. Then we can find a net $\{(x_\lambda, \gamma_\lambda)\}_{\lambda \in \Lambda}$ in Y_I with $l(x_\lambda) \neq l(x_0)$ which converges (x_0, γ_0) for some $\gamma_0 \in \mathbb{T}$. By taking a subnet if necessary, we may assume that either $l(x_\lambda)$ and $p(x_\lambda)$ are finite and constant or $l(x_\lambda) + p(x_\lambda)$ converges to ∞ . If $l(x_\lambda)$ and $p(x_\lambda)$ are finite and constant, then $l(x_0) = l(x_\lambda)$ by Lemma 6.3 (i). This is a contradiction. If $l(x_\lambda) + p(x_\lambda)$ converges to ∞ then we have $(Y_I)_{x_0} = \mathbb{T}$ by Lemma 6.7. This is also a contradiction. Therefore we can find a neighborhood V of x_0 such that all $x \in V$ with $l(x) \neq l(x_0)$ satisfies $(Y_I)_x = \emptyset$. We have proved that Y_I is admissible.

7. A proof of
$$Y_{I_Y} = Y$$

Take an admissible subset Y of $X \times \mathbb{T}$. In this section we will prove that $Y_{I_Y} = Y$. By definition, we have $Y_{I_Y} \supset Y$. Thus all we have to do is to prove the other inclusion $Y_{I_Y} \subset Y$. We set $X' := \{x \in X \mid Y_x \neq \emptyset\}$.

Lemma 7.1. The set X' is closed and σ -invariant.

Proof. The set X' is closed because Y is closed and \mathbb{T} is compact. By the condition (ii) in Definition 5.7, X' is σ -invariant.

Take $x_0 \notin X'$ and $\gamma_0 \in \mathbb{T}$. We can find $f \in C_0(X)$ such that $f(x_0) = 1$ and f(x) = 0 for all $x \in X'$. Then we have $t^0(f) \in I_Y$ because $\pi_{(x,\gamma)}(t^0(f)) = 0$ for all $x \in X'$ and all $\gamma \in \mathbb{T}$. However $t^0(f) \notin P_{(x_0,\gamma_0)}$ because $\pi_{(x_0,\gamma_0)}(t^0(f))\delta_{x_0} = \delta_{x_0}$. This implies $(x_0,\gamma_0) \notin Y_{I_Y}$. Therefore we have $Y_{I_Y} \subset X' \times \mathbb{T}$.

Take $(x_0, \gamma_0) \in (X' \times \mathbb{T}) \setminus Y$, and we will show that $(x_0, \gamma_0) \notin Y_{I_Y}$ which is equivalent to $I_Y \not\subset P_{(x_0, \gamma_0)}$. To do so, it suffices to construct $a \in C^*(\Sigma)$ such that $\pi_{(x_0, \gamma_0)}(a) \neq 0$ and $\pi_{(x, \gamma)}(a) = 0$ for all $(x, \gamma) \in Y$. Note that the representation $\pi_{(x_0, \gamma_0)}$ factors through the natural surjection $C^*(\Sigma) \to C^*(\Sigma')$ where $\Sigma' = (X', \sigma|_{U \cap X'})$. The induced representation of $C^*(\Sigma')$ can be denoted by the same notation $\pi_{(x_0, \gamma_0)}$ considered as $x_0 \in X'$. Similarly for each $(x, \gamma) \in Y$, the representation $\pi_{(x, \gamma)} : C^*(\Sigma) \to B(H_{x_0})$ factors through the representation $\pi_{(x, \gamma)} : C^*(\Sigma') \to B(H_{x_0})$ considered as $x \in X'$. Hence to finish the proof it suffices to construct $a \in C^*(\Sigma')$ such that $\pi_{(x_0, \gamma_0)}(a) \neq 0$ and $\pi_{(x, \gamma)}(a) = 0$ for all $(x, \gamma) \in Y$.

We have $Y_{x_0} \neq \emptyset$, \mathbb{T} . Hence by the condition (iii) in Definition 5.7, there exists an open neighborhood V of x_0 such that $l(x) = l(x_0)$ for all $x \in X' \cap V$. We set $x'_0 = \sigma^{l(x_0)}(x_0)$ which satisfies $l(x'_0) = 0$. Set $V' = \sigma^{l(x_0)}(V) \cap X'$ which is an open neighborhood of $x'_0 \in X'$ such that all $x' \in V'$ satisfies l(x') = 0. We have $(x'_0, \gamma_0) \notin Y$ because $(x_0, \gamma_0) \notin Y$. Since Y is closed, we can find an open neighborhood W' of $x'_0 \in X'$ and an open neighborhood Z of $\gamma_0 \in \mathbb{T}$ such that $(W' \times Z) \cap Y = \emptyset$. We may assume that $W' \subset V'$. For each $x \in W'$, we have $Y_x \cap Z = \emptyset$ and $\zeta_{p(x)}Y_x = Y_x$. Hence $\sup_{x \in W'} p(x) < \infty$. Thus we can find $N \in \mathbb{N}$ such that $\sigma^N(x) = x$ for all $x \in W'$ (recall that l(x) = 0 for $x \in W'$). Let us set $W = \bigcup_{k=0}^{N-1} \sigma^k(W')$ which is an open subset of X' satisfying $\sigma(W) = W$, and we denote by $\sigma_W : W \to W$ the restriction of σ to W, which is a homeomorphism satisfying $\sigma_W^N = \mathrm{id}_W$. By the condition (ii) in Definition 5.7, we have $(W \times Z) \cap Y = \emptyset$. For $f \in C_0(W)$ and a negative integer n, we define $t^n(f) \in C^*(\Sigma)$ by $t^n(f) = t^{-n}(\overline{f} \circ \sigma_W^{-n})^*$.

Lemma 7.2. Let $f \in C_0(W)$. For $(x, \gamma) \in X' \times \mathbb{T}$, $y \in Orb(x)$ and $n \in \mathbb{Z}$, we have

$$\pi_{(x,\gamma)}(t^n(f))\delta_y = \begin{cases} \gamma_0^n f(\sigma_W^{-n}(y))\delta_{\sigma_W^{-n}(y)} & \text{if } y \in W\\ 0 & \text{otherwise.} \end{cases}$$

Proof. We first consider the case $n \geq 0$. By Lemma 2.9, we have

$$\pi_{(x,\gamma)}(t^n(f))\delta_y = \gamma_0^n \sum_{\sigma^n(z)=y} f(z)\delta_z.$$

We have $f(z) \neq 0$ only when $z \in W$. There exists $z \in W$ with $\sigma^n(z) = y$ only when $y \in W$, and in this case $z = \sigma_W^{-n}(y)$ is the only element in W satisfying $\sigma^n(z) = y$. This proves the case $n \geq 0$. Next, we consider the case n < 0. By Lemma 2.9, we have

$$\pi_{(x,\gamma)}\big(t^n(f)\big)\delta_y = \pi_{(x,\gamma)}\big(t^{-n}(\overline{f}\circ\sigma_W^{-n})^*\big)\delta_y = \begin{cases} \gamma_0^n(f\circ\sigma_W^{-n})(y)\delta_{\sigma^{-n}(y)} & \text{if } y\in U_n,\\ 0 & \text{otherwise.} \end{cases}$$

When $y \notin W$, we have $(f \circ \sigma_W^{-n})(y) = 0$. This proves the case n < 0.

Choose $f \in C_0(W)$ with $f \ge 0$ and $f(x'_0) \ne 0$ and set

$$f_0 = \frac{1}{n} \sum_{k=0}^{n-1} f \circ \sigma_W^k.$$

Then we have $f_0 \geq 0$, $f_0(x'_0) \neq 0$ and $f_0 \circ \sigma_W = f_0$.

Take $(x, \gamma) \in X' \times \mathbb{T}$ and $y \in \text{Orb}(x)$ with $y \in W$. For k = 0, 1, ..., p(y) - 1, we set $\xi_{y,k} \in H_x$ by

$$\xi_{y,k} := \frac{1}{\sqrt{p(y)}} \sum_{j=0}^{p(y)-1} \zeta_{p(y)}^{jk} \delta_{\sigma_W^j(y)}.$$

Then $\{\xi_{y,k}\}_{k=0}^{p(y)-1}$ is a basis of the span of $\{\delta_{\sigma_W^k(y)}\}_{k=0}^{p(y)-1}$. In these situation, we have the following.

Lemma 7.3. For k = 0, 1, ..., p(y) - 1 and $n \in \mathbb{Z}$, we have

$$\pi_{(x,\gamma)}(t^n(f_0))\xi_{y,k} = (\gamma \zeta_{p(y)}^k)^n f_0(y)\xi_{y,k}.$$

Proof. By Lemma 7.2, we have

$$\pi_{(x,\gamma)}(t^{n}(f_{0}))\xi_{y,k} = \frac{1}{\sqrt{p(y)}} \sum_{j=0}^{p(y)-1} \zeta_{p(y)}^{jk} \pi_{(x,\gamma)}(t^{n}(f_{0})) \delta_{\sigma_{W}^{j}(y)}$$

$$= \frac{1}{\sqrt{p(y)}} \sum_{j=0}^{p(y)-1} \zeta_{p(y)}^{jk} \gamma^{n} (f_{0} \circ \sigma_{W}^{-n}) (\sigma_{W}^{j}(y)) \delta_{\sigma_{W}^{-n}(\sigma_{W}^{j}(y))}$$

$$= \gamma^{n} \frac{1}{\sqrt{p(y)}} \sum_{j=0}^{p(y)-1} \zeta_{p(y)}^{jk} f_{0}(y) \delta_{\sigma_{W}^{j-n}(y)}$$

$$= \gamma^{n} f_{0}(y) \frac{1}{\sqrt{p(y)}} \sum_{j=0}^{p(y)-1} \zeta_{p(y)}^{(j+n)k} \delta_{\sigma_{W}^{j}(y)}$$

$$= (\gamma \zeta_{p(y)}^{k})^{n} f_{0}(y) \xi_{y,k}.$$

Lemma 7.4. For a trigonometric polynomial $q(z) = \sum_{n=-N}^{N} \alpha_n z^n$, we set $q(f_0) = \sum_{n=-N}^{N} \alpha_n t^n(f_0)$. Then for k = 0, 1, ..., p(y) - 1, we have

$$\pi_{(x,\gamma)}(q(f_0))\xi_{y,k} = q(\gamma\zeta_{p(y)}^k)f_0(y)\xi_{y,k}.$$

Proof. This follows from Lemma 7.3.

Take $g \in C_c(Z) \subset \mathbb{C}(\mathbb{T})$ with $g(\gamma_0) \neq 0$. Choose a sequence q_1, q_2, \ldots of trigonometric polynomials converging to g uniformly on \mathbb{T} . We have the following.

Lemma 7.5. The sequence $q_1(f_0), q_2(f_0), \ldots$ converges to an element $a \in C^*(\Sigma')$. We also have

$$\pi_{(x,\gamma)}(a)\xi_{y,k} = g(\gamma_0\zeta_{p(y)}^k)f_0(y)\xi_{y,k}.$$

for
$$(x, \gamma) \in X' \times \mathbb{T}$$
, $y \in Orb(x) \cap W$ and $k = 0, 1, \dots, p(y) - 1$.

Proof. Let $B = C^*(W, \sigma_W) = C_0(W) \rtimes_{\sigma_W} \mathbb{Z}$ be the universal C^* -algebra generated by the products of the elements in $C_0(W) \subset B$ and a unitary $u \in M(B)$ which satisfy $ufu^* = f \circ \sigma_W$ for $f \in C_0(W)$ where M(B) is the multiplier algebra of B. By the universality, there exists a *-homomorphism $\iota \colon B \to C^*(\Sigma')$ such that $\iota(f) = t^0(f)$ and $\iota(fu) = t^1(f)$. (We know that ι is injective by Proposition A.1.) For $n \in \mathbb{N}$, we have $\iota(f_0u^n) = t^n(f_0)$. To see this for $n \geq 0$, choose $\xi_1, \ldots, \xi_n \in C_c(U)$ such that

$$f_0(x) = \xi_1(x)\xi_2(\sigma(x))\cdots\xi_n(\sigma^{n-1}(x))$$

for all $x \in U_n$ by Lemma 1.10, and compute

$$\iota(f_0u^n) = \iota(\xi_1(\xi_2 \circ \sigma) \cdots (\xi_n \circ \sigma^{n-1})u^n)$$

$$= \iota(\xi_1u\xi_2u \cdots \xi_{n-1}u\xi_nu)$$

$$= \iota(\xi_1u)\iota(\xi_2u) \cdots \iota(\xi_{n-1}u)\iota(\xi_nu)$$

$$= t_1(\xi_1)t_1(\xi_2) \cdots t_1(\xi_{n-1})t_1(\xi_n)$$

$$= t^n(f_0).$$

For $n \leq -1$, we also have $\iota(f_0u^n) = t^n(f_0)$ because

$$\iota(f_0 u^n) = \iota(u^{-n} \overline{f_0})^* = \iota((\overline{f_0} \circ \sigma_W^{-n}) u^{-n})^*$$
$$= t^{-n} (\overline{f_0} \circ \sigma_W^{-n})^* = t^n (f_0)$$

Thus for a trigonometric polynomial q, we have $\iota(f_0q(u)) = q(f_0)$. Since $(f_0q_k(u))_k$ converges $f_0g(u)$ in B, $(q_k(f_0))_k$ converges $\iota(f_0g(u))$ in $C^*(\Sigma')$. Thus we get $a = \iota(f_0g(u))$. The last equation follows easily from Lemma 7.4.

Proposition 7.6. Let $a \in C^*(\Sigma')$ be as in Lemma 7.5. We have $\pi_{(x_0,\gamma_0)}(a) \neq 0$ and $\pi_{(x,\gamma)}(a) = 0$ for all $(x,\gamma) \in Y$.

Proof. We have $\pi_{(x_0,\gamma_0)}(a) \neq 0$ because $\pi_{(x_0,\gamma_0)}(a)\xi_{x_0,0} = g(\gamma_0)f_0(x_0)\xi_{x_0,0} \neq 0$ by Lemma 7.5.

Take $(x, \gamma) \in Y$. Take $y \in \operatorname{Orb}(x)$. If $y \notin W$, then we can see $\pi_{(x,\gamma)}(a)\delta_y = 0$ by Lemma 7.2. Suppose $y \in W$. By the condition (ii) in Definition 5.7, we have $(y, \gamma) \in Y$. By the condition (iii) in Definition 5.7, we have $(y, \zeta_{p(y)}^k \gamma) \in Y$ for $k = 0, 1, \ldots, p(y) - 1$. Hence $\zeta_{p(y)}^k \gamma \notin Z$ for $k = 0, 1, \ldots, p(y) - 1$. By Lemma 7.5 we have $\pi_{(x,\gamma)}(a)\xi_{y,k} = 0$ for $k = 0, 1, \ldots, p(y) - 1$. Hence we get $\pi_{(x,\gamma)}(a)\delta_y = 0$. These show that $\pi_{(x,\gamma)}(a) = 0$.

We have shown the following.

Proposition 7.7. For an admissible subset Y of $X \times \mathbb{T}$, we have $Y_{I_Y} = Y$.

Theorem 7.8. The set of ideals of $C^*(\Sigma)$ corresponds bijectively to the set of all admissible subsets of $X \times \mathbb{T}$ via the maps $I \mapsto Y_I$ and $Y \mapsto I_Y$.

Proof. This follows from Proposition 5.5, Proposition 6.8 and Proposition 7.7. \square

APPENDIX A. UNIQUENESS THEOREMS

Let $\Sigma = (X, \sigma)$ be an SGDS. By the universality of $C^*(\Sigma)$, we have an action β of the group $\mathbb{T} := \{z \in \mathbb{C} \mid |z| = 1\}$ on $C^*(\Sigma)$ so that for $z \in \mathbb{T}$ we have $\beta_z(t^0(f)) = t^0(f)$ for $f \in C_0(X)$ and $\beta_z(t^1(\xi)) = zt^1(\xi)$ for $\xi \in C_c(U)$. This action β is called the *gauge* action (see [K1, Section 4]).

Proposition A.1. Let A be a C^* -algebra. A *-homomorphism $\Phi: C^*(\Sigma) \to A$ is injective if $\Phi \circ t^0$ is injective and there exists an action of \mathbb{T} on A with which and the gauge action on $C^*(\Sigma)$, Φ is equivariant.

Proof. This follows from [K1, Theorem 4.5]. \Box

Definition A.2 (cf. [R, Definition 2.5]). An SGDS Σ is said to be essentially free if the interior of the set $\{x \in X \mid l(x) = 0\}$ is empty.

By Baire's category theorem, Σ is essentially free if and only if the interior of the set $\{x \in X \mid \sigma^n(x) = x\}$ is empty for every positive integer n. Thus this definition coincides with [R, Definition 2.5].

Proposition A.3. Suppose an SGDS Σ is essentially free. Let A be a C^* -algebra. A *-homomorphism $\Phi: C^*(\Sigma) \to A$ is injective if $\Phi \circ t^0$ is injective

Proof. An SGDS Σ is essentially free if and only if the associate topological graph $E = (X, U, \sigma, \iota)$ is topologically free in the sense of [K1, Definition 5.4]. Thus this proposition follows from [K1, Theorem 5.12].

References

- [B] Blackadar, B. Operator algebras. Theory of C*-algebras and von Neumann algebras. Encyclopaedia of Mathematical Sciences, 122. Operator Algebras and Non-commutative Geometry, III. Springer-Verlag, Berlin, 2006.
- [K1] Katsura, T. A class of C^* -algebras generalizing both graph algebras and homeomorphism C^* -algebras I, fundamental results. Trans. Amer. Math. Soc. **356** (2004), no. 11, 4287-4322.
- [K2] Katsura, T. A class of C*-algebras generalizing both graph algebras and homeomorphism C*-algebras II, examples. Internat. J. Math. 17 (2006), No. 7, 791–833.
- [K3] Katsura, T. A class of C*-algebras generalizing both graph algebras and homeomorphism C*-algebras III, ideal structures. Ergodic Theory Dynam. Systems 26 (2006), no. 6, 1805–1854.
- [K4] Katsura, T. Topological graphs and singly generated dynamical systems. Preprint 2021, arXiv:2107.01389, submitted.
- [KL] Kumjian, A.; Li, H. Twisted topological graph algebras are twisted groupoid C*-algebras. J. Operator Theory, 78 (2017), no. 1, 201–225.
- [R] Renault, J. Cuntz-like algebras. Operator theoretical methods, 371–386, Theta Found., Bucharest, 2000.
- [SW] Sims, A.; Williams, D. P. The primitive ideals of some étale groupoid C*-algebras. Algebr. Represent. Theory 19 (2016), no. 2, 255–276.
- [T1] Tomiyama, J. The interplay between topological dynamics and theory of C*-algebras. Lecture Notes Series, 2. Seoul National University, Research Institute of Mathematics, Global Analysis Research Center, Seoul, 1992.
- [W] Williams, D. P. The topology on the primitive ideal space of transformation group C*-algebras and C.C.R. transformation group C*-algebras. Trans. Amer. Math. Soc. **266** (1981), no. 2, 335–359.
- [Y] Yeend, T. Topological higher-rank graphs and the C*-algebras of topological 1-graphs. Operator theory, operator algebras, and applications, Amer. Math. Soc., Providence, RI, 2006, 231–244.

DEPARTMENT OF MATHEMATICS, KEIO UNIVERSITY, YOKOHAMA, 223-8522 JAPAN *Email address*: katsura@math.keio.ac.jp