VECTOR BUNDLES AND CONNECTIONS ON RIEMANN SURFACES WITH PROJECTIVE STRUCTURE

INDRANIL BISWAS, JACQUES HURTUBISE, AND VLADIMIR ROUBTSOV

ABSTRACT. Let $\mathcal{B}_g(r)$ be the moduli space of triples of the form $(X, K_X^{1/2}, F)$, where X is a compact connected Riemann surface of genus g, with $g \geq 2$, $K_X^{1/2}$ is a theta characteristic on X, and F is a stable vector bundle on X of rank r and degree zero. We construct a $T^*\mathcal{B}_g(r)$ -torsor $\mathcal{H}_g(r)$ over $\mathcal{B}_g(r)$. This generalizes on the one hand the torsor over the moduli space of stable vector bundles of rank r, on a fixed Riemann surface Y, given by the moduli space of holomorphic connections on the stable vector bundles of rank r on Y, and on the other hand the torsor over the moduli space of Riemann surfaces given by the moduli space of Riemann surfaces with a projective structure. It is shown that $\mathcal{H}_g(r)$ has a holomorphic symplectic structure compatible with the $T^*\mathcal{B}_g(r)$ -torsor structure. We also describe $\mathcal{H}_g(r)$ in terms of the second order matrix valued differential operators. It is shown that $\mathcal{H}_g(r)$ is identified with the $T^*\mathcal{B}_g(r)$ -torsor given by the sheaf of holomorphic connections on the theta line bundle over $\mathcal{B}_g(r)$.

CONTENTS

1. Introduction	2
2. Stable r -opers and their properties	4
2.1. Stable <i>r</i> -opers	4
2.2. An equivalence relation	9
3. The space of holomorphic connections	13
4. Infinitesimal deformations	15
5. Affine space structure	19
6. Differential operators, integral kernels and r -opers	22
6.1. Differential operators and r -opers	22
6.2. Integral kernels and differential operators	26
7. A canonical holomorphic section	28
8. A canonical isomorphism of torsors and a symplectic structure	31
8.1. A canonical isomorphism of torsors	31
8.2. A holomorphic symplectic form	36
9. Isomonodromy and symplectic form	37
9.1. Projective structure on a Riemann surface	37

²⁰¹⁰ Mathematics Subject Classification. 14H60, 16S32, 14D21, 53D30.

Key words and phrases. Projective structure, differential operator, holomorphic connection, oper, torsor.

9.2. Isomonodromy

References

1. INTRODUCTION

Let X be a compact connected Riemann surface of genus g, with $g \geq 2$, and $\mathcal{N}_X(r)$ the moduli space of stable vector bundles on X of rank r and degree zero. Let $\mathcal{C}_X(r)$ be the moduli space of rank r holomorphic connections (E, D) on X such that the underlying holomorphic vector bundle E is stable. It has a forgetful map to $\mathcal{N}_X(r)$ that sends any (E, D) to E, and $\mathcal{C}_X(r)$ is a torsor over $\mathcal{N}_X(r)$ for the holomorphic cotangent bundle $T^*\mathcal{N}_X(r)$. Moreover, $\mathcal{C}_X(r)$ has a holomorphic symplectic structure which is compatible with the $T^*\mathcal{N}_X(r)$ -torsor structure.

There is another $T^*\mathcal{N}_X(r)$ -torsor that one can define. To each bundle $E \in \mathcal{N}_X(r)$ we associate the Quillen determinant line for its $\overline{\partial}$ operator, and this defines a holomorphic line bundle L over $\mathcal{N}_X(r)$. Then consider the sheaf of pointwise connections $\operatorname{Conn}(L)$ over $\mathcal{N}_X(r)$ (i.e., the sheaf on $\mathcal{N}_X(r)$ whose sections are holomorphic connections on L). This is again a $T^*\mathcal{N}_X(r)$ -torsor over $\mathcal{N}_X(r)$.

The surprise, established in [BH1, BH2] is that there is a canonical isomorphism between these two torsors. The isomorphism is constructed by taking C^{∞} sections on both sides that at the first sight do not seem to have much to do with each other, but have the same data encoding the torsor. For $\mathcal{C}_X(r)$, we associate to each bundle its unitary connection given by the Narasimhan–Seshadri theorem; for Conn(L) the Chern connection for the Quillen metric on L. There is a holomorphic version of this equivalence by sections, expanded in [BH2], in which the line bundle L is restricted to the complement \mathcal{U} of the theta divisor in $\mathcal{N}_X(r)$ associated to a theta characteristic $K_X^{1/2}$ of X, so the canonical trivialization of L over \mathcal{U} produces an integrable holomorphic connection on $L|_{\mathcal{U}}$. On the other hand, a holomorphic connection on any bundle $E \in \mathcal{U}$ is obtained as the asymptotic data of a canonical section of $(E \otimes K_X^{1/2}) \boxtimes (E^* \otimes K_X^{1/2})$ over $X \times X$, with singularities on the diagonal. There is a holomorphic isomorphism between $\operatorname{Conn}(L)$ and $\mathcal{C}_X(r)$ that maps the section of $\operatorname{Conn}(L)|_{\mathcal{U}}$ given be the trivialization of $L|_{\mathcal{U}}$ to the section of $\mathcal{C}_X(r)|_{\mathcal{U}}$ given by the above canonical section. In fact, not only is there an isomorphism, but it is obtained by linking together rather special and hitherto unrelated objects on each side of the equivalence, in several different ways.

The holomorphic version allows us, of course to move not only the bundle, but the base curve as well, and the question we are examining in this paper is whether we can again have two a priori inequivalent holomorphic $T^*\mathcal{B}_q(r)$ torsors over the space $\mathcal{B}_q(r)$ of pairs (bundles, curves) actually turn out to be the same. The answer turns out to be yes. As a bonus, we have a hereto undefined symplectic structure for one of the torsors.

To explain with more details, let \mathcal{P}_g denote the moduli space of Riemann surfaces of genus q equipped with a projective structure. It has a natural map to the moduli space \mathcal{M}_q of Riemann the surfaces of genus g that simply forgets the projective structure, and \mathcal{P}_g is a holomorphic torsor over \mathcal{M}_g for the holomorphic cotangent bundle $T^*\mathcal{M}_g$. Also, \mathcal{P}_g has a holomorphic symplectic structure which is compatible with the $T^*\mathcal{M}_g$ -torsor structure.

To combine the torsors \mathcal{P}_g and $\mathcal{C}_X(r)$ into a single one, let $\mathcal{B}_g(r)$ be the moduli space of triples of the form $(X, K_X^{1/2}, F)$, where

- X is a compact connected Riemann surface of genus g, with $g \ge 2$,
- $K_X^{1/2}$ is a theta characteristic on X (the holomorphic cotangent bundle of X is denoted by K_X), and
- F is a stable vector bundle over X of rank r and degree zero.

Fix a theta characteristic $K_X^{1/2}$ on a compact Riemann surface X of genus g. Given a stable vector bundle F on X of rank r and degree zero, we construct a certain quotient space of the space of all holomorphic connections on the first order jet bundle $J^1(F \otimes (K_X^{1/2})^*)$; this quotient space is denoted by $\mathcal{D}(F)$ (see (2.29) and Corollary 3.2). Let $\mathcal{H}_g(r)$ denote the moduli space of objects of the form $(X, K_X^{1/2}, F, D)$, where $(X, K_X^{1/2}, F) \in \mathcal{B}_g(r)$ and $D \in \mathcal{D}(F)$. It has a natural map

$$\gamma : \mathcal{H}_g(r) \longrightarrow \mathcal{B}_g(r), \quad (X, K_X^{1/2}, F, D) \longmapsto (X, K_X^{1/2}, F).$$

We prove the following (see Theorem 5.3 and Corollary 8.2):

Theorem 1.1.

- (1) $\mathcal{H}_g(r)$ is a torsor over $\mathcal{B}_g(r)$ for the holomorphic cotangent bundle $T^*\mathcal{B}_q(r)$.
- (2) $\mathcal{H}_q(r)$ has an algebraic symplectic structure $\Omega_{\mathcal{H}_q(r)}$.
- (3) The symplectic form $\Omega_{\mathcal{H}_g(r)}$ on $\mathcal{H}_g(r)$ is compatible with the $T^*\mathcal{B}_g(r)$ -torsor structure of $\mathcal{H}_g(r)$.
- (4) There is a holomorphic line bundle \mathbf{L} on $\mathcal{H}_g(r)$, and a holomorphic connection $\nabla^{\mathbf{L}}$ on \mathbf{L} , such that the curvature of $\nabla^{\mathbf{L}}$ is the symplectic form $\Omega_{\mathcal{H}_g(r)}$.

Consider the standard subbundle $F \otimes (K_X^{1/2})^* \otimes K_X = F \otimes K_X^{1/2} \subset J^1(F \otimes (K_X^{1/2})^*)$. For any holomorphic connection \mathbb{D} on $J^1(F \otimes (K_X^{1/2})^*)$, the second fundamental form of $F \otimes K_X^{1/2}$ for \mathbb{D} is the identity map of F (see Corollary 3.2). This property is very similar to the defining property of opers. We recall that opers were introduced by Beilinson and Drinfeld [BD1], [BD2]. Their motivation came from the works of Drinfeld and Sokolov [DS1], [DS2]. We note that the study of opers within geometry and mathematical physics has received much attention in the recent times.

There is a natural divisor $\Theta \subset \mathcal{B}_g(r)$ consisting of all $(X, K_X^{1/2}, F)$ such that $H^0(X, F \otimes K_X^{1/2}) \neq 0$. The line bundle on $\mathcal{B}_g(r)$ defined by Θ will be denoted by \mathcal{L} . Let

$$\mathcal{C}(\mathcal{L}) \longrightarrow \mathcal{B}_q(r)$$

be the holomorphic fiber bundle defined by the sheaf of holomorphic connection on \mathcal{L} . So the space of holomorphic sections of $\mathcal{C}(\mathcal{L})$ over an open subset $U \subset \mathcal{B}_g(r)$ is the space of all holomorphic connections on $\mathcal{L}|_U \longrightarrow U$. This $\mathcal{C}(\mathcal{L})$ is an algebraic torsor over $\mathcal{B}_g(r)$ for the holomorphic cotangent bundle $T^*\mathcal{B}_g(r)$.

We prove the following (see Theorem 8.1):

Theorem 1.2. There is a canonical algebraic isomorphism of $T^*\mathcal{B}_q(r)$ -torsors

$$\mathcal{H}: \mathcal{H}_q(r) \longrightarrow \mathcal{C}(\mathcal{L})$$

Projective structures on a Riemann surface X are defined by giving a holomorphic coordinate atlas on X such that all the transition functions are Möbius transformations. Projective structures on a Riemann surface X are identified with holomorphic ordinary differential operators \mathcal{D} on X of order two such that

- the symbol of \mathcal{D} is the constant function 1 on X, and
- the sub-leading term of \mathcal{D} vanishes identically (equivalently, the $\operatorname{GL}(2, \mathbb{C})$ -local system of X defined by the sheaf of solutions of \mathcal{D} is actually a $\operatorname{SL}(2, \mathbb{C})$ -local system).

The above mentioned space $\mathcal{H}_g(r)$ admits a similar description in terms of the second order matrix valued differential operators. To explain this, for any $(X, K_X^{1/2}, F) \in \mathcal{B}_g(r)$, let

$$\widetilde{\mathcal{D}}'(X, K_X^{1/2}, F) \subset H^0(X, \operatorname{Diff}_X^2(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$$

be the locus of holomorphic differential operators whose symbol is $\mathrm{Id}_F \in H^0(X, \mathrm{End}(F))$. The vector space $H^0(X, \mathrm{End}(F) \otimes K_X^{\otimes 2})$ acts freely on $\widetilde{\mathcal{D}}'(X, K_X^{1/2}, F)$ (but this action is not transitive).

We prove the following (see Theorem 6.4):

Theorem 1.3. There is a canonical bijection between $\mathcal{D}(F)$ and the quotient space

$$\widetilde{\mathcal{D}}'(X, K_X^{1/2}, F) / H^0(X, \operatorname{ad}(F) \otimes K_X^{\otimes 2}).$$

There is a natural holomorphic projection $\mathcal{H}_g(r) \longrightarrow \mathcal{P}_g$ (see Proposition 9.1), using which we may pullback, to $\mathcal{H}_g(r)$, the symplectic 2-form on \mathcal{P}_g . On the other hand, using isomonodromic deformations, the symplectic form on $\mathcal{C}_X(r)$ produces a holomorphic 2form on $\mathcal{H}_g(r)$. It is natural to ask whether the symplectic form on $\mathcal{H}_g(r)$ in Theorem 1.1(2) is a combination of these two 2-forms (see Conjecture 9.3 for a precise formulation).

The result of this paper, as well as those of [BH1], [BH2], have a flavor of geometric quantization: a symplectic manifold projects to a manifold of half the dimension, equipped with a line bundle with a connection. The fibers of the projection are Lagrangians, and the curvature of the connection is the symplectic form. We are unaware, however, of any concrete link to geometric quantization.

2. Stable r-opers and their properties

2.1. Stable *r*-opers. Let X be a compact connected Riemann surface; its holomorphic cotangent (respectively, tangent) bundle will be denoted by K_X (respectively, TX). We shall assume throughout that

$$\operatorname{genus}(X) = g \ge 2$$
.

A holomorphic connection on a holomorphic vector bundle E over X is a first order holomorphic differential operator

$$D: E \longrightarrow E \otimes K_X$$

such that $D(fs) = fD(s) + s \otimes df$ for all locally defined holomorphic functions f on X and all locally defined holomorphic sections s of E; see [At]. A holomorphic connection on E is automatically flat because $\Omega_X^{2,0} = 0$.

If D is a holomorphic connection on E, and $S \subset E$ is a holomorphic subbundle of E, then consider the composition of homomorphisms

$$S \hookrightarrow E \xrightarrow{D} E \otimes K_X \xrightarrow{q \otimes \operatorname{Id}_{K_X}} (E/S) \otimes K_X,$$

where

$$q: E \longrightarrow E/S \tag{2.1}$$

is the quotient map. This composition of homomorphisms is \mathcal{O}_X -linear, and hence it corresponds to a holomorphic section

$$D_S \in H^0(X, \operatorname{Hom}(S, E/S) \otimes K_X).$$
 (2.2)

This homomorphism D_S is called the *second fundamental form* of S for the connection D.

A holomorphic vector bundle V on X is called stable if

$$\frac{\operatorname{degree}(W)}{\operatorname{rank}(W)} < \frac{\operatorname{degree}(V)}{\operatorname{rank}(V)}$$

for every holomorphic subbundle $0 \neq W \subsetneq V$.

Definition 2.1. For a positive integer r, a r-oper on X is a triple (E, S, D), where

- E is a holomorphic vector bundle on X of rank 2r,
- D is a holomorphic connection on E, and
- $S \subset E$ is a holomorphic subbundle of rank r,

such that the second fundamental form D_S defined in (2.2) is an isomorphism.

A r-oper (E, S, D) is called *stable* if the holomorphic vector bundle S of rank r is stable.

Using the isomorphism $D_S : S \longrightarrow (E/S) \otimes K_X$ in Definition 2.1, we may identify E/S with $S \otimes TX$. Invoking this isomorphism D_S is made into the identity map of S. Henceforth, for any r-oper we shall always execute the option of using this isomorphism.

Fix a holomorphic line bundle $K_X^{1/2}$ on X such that $K_X^{1/2} \otimes K_X^{1/2} = K_X$; also fix a holomorphic isomorphism of $K_X^{1/2} \otimes K_X^{1/2}$ with K_X . Such a line bundle is called a *theta* characteristic on X. The dual line bundle $(K_X^{1/2})^*$ will be denoted by $K_X^{-1/2}$; also $(K_X^{1/2})^{\otimes n}$ (respectively, $(K_X^{1/2})^{-\otimes n}$) will be denoted by $K_X^{n/2}$ (respectively, $K_X^{-n/2}$) for every $n \geq 1$.

Take a r-oper (E, S, D) on X. Define the vector bundle $F := S \otimes K_X^{-1/2}$, so we have

$$S = F \otimes K_X^{1/2}. (2.3)$$

Then from the isomorphism D_S we have

$$E/S = S \otimes (K_X)^* = F \otimes K_X^{1/2} \otimes (K_X)^* = F \otimes K_X^{-1/2}.$$
 (2.4)

Since E admits a holomorphic connection, we have

$$\operatorname{degree}(E) = 0 \tag{2.5}$$

[At, p. 202, Proposition 18(i)]. Using (2.5), (2.4) and (2.3) it follows that

0 = degree(E) = degree(S) + degree(E/S)

 $= \text{degree}(F \otimes K_X^{1/2}) + \text{degree}(F \otimes K_X^{-1/2}) = 2 \cdot \text{degree}(F) \,.$ Hence we have degree(F) = 0.

For any holomorphic vector bundle W on X, the first jet bundle $J^1(W)$ fits into the following short exact sequence of holomorphic vector bundles on X:

$$0 \longrightarrow W \otimes K_X \xrightarrow{\iota} J^1(W) \longrightarrow J^0(W) = W \longrightarrow 0.$$
 (2.6)

Lemma 2.2. Let (E, S, D) be a r-oper on X. Then the first jet bundle $J^1(E/S)$ is canonically identified with the holomorphic vector bundle E. Also, E is holomorphically identified with $J^1(S \otimes TX)$.

Proof. Using the flat holomorphic connection D we shall construct a homomorphism

$$\varphi: E \longrightarrow J^1(E/S). \tag{2.7}$$

For this, take any point $x \in X$ and any element $v \in E_x$ in the fiber over x. Let \hat{v} be the unique flat section of E (for the connection D), defined on a simply connected open neighborhood $U \subset X$ of x, such that $\hat{v}(x) = v$. Now restrict the section $q(\hat{v}) \in H^0(U, E/S)$, where q is the quotient map in (2.1), to the first order infinitesimal neighborhood of x; let $\hat{v}' \in J^1(E/S)_x$ be the element obtained this way from $q(\hat{v})$. The map φ in (2.7) sends any $v \in E_x$, $x \in X$, to $\hat{v}' \in J^1(E/S)_x$ constructed above from it.

The map φ in (2.7) fits in the following commutative diagram of homomorphisms

where φ' is the restriction of φ to the subbundle S, and the exact sequence at the bottom of (2.8) is the one in (2.6) for W = E/S. It is straightforward to check that the homomorphism φ' in (2.8) actually coincides with the second fundamental form D_S defined in (2.2). Since D_S is an isomorphism, from the commutativity of (2.8) it follows immediately that φ constructed in (2.7) is an isomorphism.

Using the isomorphism $E/S \xrightarrow{\sim} S \otimes TX$ in (2.4), the isomorphism φ identifies E with $J^1(S \otimes TX).$

Lemma 2.3. Let (E, S, D) be a r-oper on X. Let $D^1 := (\varphi^{-1})^*D$ be the connection on $J^1(E/S)$, where φ is the isomorphism in (2.8). Then the second fundamental form of $(E/S) \otimes K_X \hookrightarrow J^1(E/S)$ for the connection D^1 is the identity map of $(E/S) \otimes K_X$.

Proof. This follows immediately from the construction of φ , the commutativity of the diagram in (2.8) and the fact that φ' coincides with the second fundamental form D_S defined in (2.2).

Proposition 2.4. Let F be a stable vector bundle on X of rank r and degree zero. Then there is a holomorphic vector bundle E on X of rank 2r, and a holomorphic connection Don E, such that

- (1) $S := F \otimes K_X^{1/2}$ is a holomorphic subbundle of E, and (2) the triple (E, S, D) is a stable r-oper. In particular, $E = J^1(S \otimes TX)$ by Lemma 2.2.

Proof. The stable vector bundle F of degree zero admits a holomorphic connection [At, p. 203, Proposition 19], [We] (in fact F admits a unique holomorphic connection whose monodromy representation is unitary [NS]). Fix a holomorphic connection D^F on F. Now consider the holomorphic vector bundle

$$E := F \otimes J^1(K_X^{-1/2}).$$

Since $J^1(K_X^{-1/2})$ is indecomposable of degree zero, it admits a holomorphic connection [At, p. 203, Proposition 19], [We]. In fact, a projective structure on X produces a holomorphic connection on $J^1(K_X^{-1/2})$ [Gu]; the definition of projective structure is recalled in Section 9.1. Let D^J be a holomorphic connection on $J^1(K_X^{-1/2})$. Consider the natural inclusion map

$$K_X^{1/2} = K_X^{-1/2} \otimes K_X \stackrel{\iota}{\hookrightarrow} J^1(K_X^{-1/2})$$
(2.9)

(see (2.6)). Let

$$D^{J}(K_{X}^{1/2}) : K_{X}^{1/2} \longrightarrow K_{X}^{-1/2} \otimes K_{X} = K_{X}^{1/2}$$
 (2.10)

be the second fundamental form of this subbundle $K_X^{1/2} \hookrightarrow J^1(K_X^{-1/2})$ for the above connection D^J on $J^1(K_X^{-1/2})$. This homomorphism $D^J(K_X^{1/2})$ in (2.10) is a nonzero constant scalar multiple of the identity map of $K_X^{1/2}$; the scalar is nonzero because $K_X^{1/2}$ does not admit any holomorphic connection as its degree is nonzero.

The holomorphic connections D^F and D^J together produce the holomorphic connection

$$D := D^F \otimes \mathrm{Id}_{J^1(K_X^{-1/2})} + \mathrm{Id}_F \otimes D^J$$
(2.11)

on $F \otimes J^1(K_X^{-1/2}) = E$. The inclusion map ι in (2.9) produces an inclusion map

$$S := F \otimes K_X^{1/2} \xrightarrow{\operatorname{Id}_F \otimes \iota} F \otimes J^1(K_X^{-1/2}) = E.$$

The second fundamental form D_S (see (2.2)) of S for the connection D in (2.11) coincides with

$$\operatorname{Id}_F \otimes D^J(K_X^{1/2}) : F \otimes K_X^{1/2} \longrightarrow F \otimes (K_X^{-1/2} \otimes K_X) = F \otimes K_X^{1/2},$$

where $D^J(K_X^{1/2})$ is the homomorphism in (2.10). Since $D^J(K_X^{1/2})$ is a nonzero scalar multiple of the identity map of $K_X^{1/2}$, we now conclude that D_S is a nonzero scalar multiple of the identity map of $F \otimes K_X^{1/2}$, in particular, D_S is an isomorphism. Therefore, the triple (E, S, D) is a stable *r*-oper on *X*.

Proposition 2.5. Let F be a stable vector bundle on X of degree zero, and let V be any holomorphic vector bundle on X. Then there is a holomorphic isomorphism

 $h \, : \, F \otimes J^1(V) \, \longrightarrow \, J^1(F \otimes V)$

such that the exact sequence in (2.6) for $W = F \otimes V$ coincides with the exact sequence in (2.6) for W = V tensored with F.

Proof. As in the proof of Proposition 2.4, fix a holomorphic connection D^F on F. For any point $x \in X$, and any $v \in F_x$, let \hat{v} be the unique flat section of F, defined on a simply connected open neighborhood of $x \in X$, such that $\hat{v}(x) = x$ (see the proof of Lemma 2.2). Take any element $w \in J^1(V)_x$, so w is a section of V defined on the first order infinitesimal neighborhood of x. Therefore, $\hat{v}' \otimes w$ is a section of $F \otimes V$ defined on the first order infinitesimal neighborhood of x, where \hat{v}' is the restriction of \hat{v} to the first order infinitesimal neighborhood of x. Now we have a map

$$h(x) : F_x \otimes J^1(V)_x \longrightarrow J^1(F \otimes V)_x$$

that sends any $v \otimes w \in F_x \otimes J^1(V)_x$ to the element $\hat{v}' \otimes w$ of $J^1(F \otimes V)_x$ constructed above. It is straightforward to check that h(x) is an isomorphism. Moreover, we get a holomorphic isomorphism

$$h: F \otimes J^1(V) \longrightarrow J^1(F \otimes V)$$

which coincides with h(x) for any $x \in X$.

From the above construction of h we conclude that the following diagram of homomorphisms is commutative

where the top one is the exact sequence in (2.6) for W = V tensored with F and the bottom one is the exact sequence in (2.6) for $W = F \otimes V$. This completes the proof. \Box

For F as in Proposition 2.5, consider the short exact sequence

$$0 \longrightarrow F \otimes K_X^{-1/2} \otimes K_X = F \otimes K_X^{1/2} \longrightarrow J^1(F \otimes K_X^{-1/2}) \longrightarrow F \otimes K_X^{-1/2} \longrightarrow 0$$

in (2.6) for $W = F \otimes K_X^{-1/2}$. It corresponds to an extension class

$$c(F) \in H^1(X, \operatorname{Hom}(F \otimes K_X^{-1/2}, F \otimes K_X^{1/2})) = H^1(X, \operatorname{End}(F) \otimes K_X).$$
(2.13)

Since $H^1(X, K_X) = H^0(X, \mathcal{O}_X)^* = \mathbb{C}^* = \mathbb{C}$, we have

$$\mathrm{Id}_F \otimes 1 \in H^0(X, \mathrm{End}(F)) \otimes H^1(X, K_X) \hookrightarrow H^1(X, \mathrm{End}(F) \otimes K_X); \qquad (2.14)$$

the above map $H^0(X, \operatorname{End}(F)) \otimes H^1(X, K_X) \longrightarrow H^1(X, \operatorname{End}(F) \otimes K_X)$ is injective because $H^0(X, \operatorname{End}(F)) = \mathbb{C}$.

Corollary 2.6. Let F be a stable vector bundle on X of degree zero. Then the cohomology class c(F) in (2.13) coincides with $Id_F \otimes 1$ in (2.14).

Proof. Consider the short exact sequence

$$0 \longrightarrow K_X^{-1/2} \otimes K_X = K_X^{1/2} \longrightarrow J^1(K_X^{-1/2}) \longrightarrow K_X^{-1/2} \longrightarrow 0$$

in (2.6) for $W = K_X^{-1/2}$. The corresponding extension class coincides with

$$1 \in H^1(X, \operatorname{Hom}(K_X^{-1/2}, K_X^{1/2})) = H^1(X, K_X) = \mathbb{C}$$

[Gu]. Hence the extension class of the tensor product of the above exact sequence with F

$$0 \longrightarrow F \otimes K_X^{1/2} \longrightarrow F \otimes J^1(K_X^{-1/2}) \longrightarrow F \otimes K_X^{-1/2} \longrightarrow 0$$

is $\mathrm{Id}_F \otimes 1 \in H^1(X, \mathrm{End}(F) \otimes K_X) = H^1(X, \mathrm{Hom}(F \otimes K_X^{-1/2}, F \otimes K_X^{1/2}))$ in (2.14). Therefore, the proof is completed using the isomorphism h in (2.12) together with the commutativity of the diagram in (2.12). 2.2. An equivalence relation. Take a holomorphic vector bundle F on X of rank r and degree zero; it need not be stable. Assume that the jet bundle $J^1(F \otimes K_X^{-1/2})$ has a holomorphic connection D_1 . Consider the subbundle

$$F \otimes K_X^{1/2} = F \otimes K_X^{-1/2} \otimes K_X \subset J^1(F \otimes K_X^{-1/2})$$

(see (2.6)). The second fundamental form of this subbundle for the connection D_1 is a homomorphism

$$\mathcal{S}(F, D_1) : F \otimes K_X^{1/2} \longrightarrow (J^1(F \otimes K_X^{-1/2})/(F \otimes K_X^{1/2})) \otimes K_X = F \otimes K_X^{-1/2} \otimes K_X = F \otimes K_X^{1/2}$$
(2.15)

(see (2.2) and (2.6)).

Now assume that $J^1(F \otimes K_X^{-1/2})$ admits a holomorphic connection D_1 for which the second fundamental form $\mathcal{S}(F, D_1)$ in (2.15) is the identity map of $F \otimes K_X^{1/2}$. Note that for such a holomorphic connection D_1 , the triple $(J^1(F \otimes K_X^{-1/2}), F \otimes K_X^{1/2}, D_1)$ is a *r*-oper (see Definition 2.1, (2.3) and (2.4)).

Let

$$\operatorname{Conn}(J^1(F \otimes K_X^{-1/2})) \tag{2.16}$$

denote the space of all holomorphic connection on the vector bundle $J^1(F \otimes K_X^{-1/2})$.

Definition 2.7. Let

$$\widetilde{\mathcal{C}}(F) \subset \operatorname{Conn}(J^1(F \otimes K_X^{-1/2}))$$

be the space of all holomorphic connections D on $J^1(F \otimes K_X^{-1/2})$ with the property that the corresponding second fundamental form $\mathcal{S}(F, D)$ in (2.15) is the identity map of $F \otimes K_X^{1/2}$.

We note that $\widetilde{\mathcal{C}}(F)$ is nonempty by the assumption on F.

So $(J^1(F \otimes K_X^{-1/2}), F \otimes K_X^{1/2}, D)$ is a *r*-oper for every $D \in \widetilde{\mathcal{C}}(F)$.

We shall see in Corollary 3.2 that $\widetilde{\mathcal{C}}(F) = \operatorname{Conn}(J^1(F \otimes K_X^{-1/2}))$ when the vector bundle F is stable.

For notational convenience, define

$$S := F \otimes K_X^{1/2} \quad \text{and} \quad E := J^1(F \otimes K_X^{-1/2}) \tag{2.17}$$

(see (2.3), (2.4) and Lemma 2.2), so S is a holomorphic subbundle of E by (2.6) (see (2.18) below). Note that we have $\operatorname{End}(S) = \operatorname{End}(F) \otimes \operatorname{End}(K_X^{1/2}) = \operatorname{End}(F)$.

Setting $W = F \otimes K_X^{-1/2}$ in (2.6), we get a short exact sequence

$$0 \longrightarrow F \otimes K_X^{-1/2} \otimes K_X = S \xrightarrow{\iota} J^1(F \otimes K_X^{-1/2}) = E \xrightarrow{q_0} F \otimes K_X^{-1/2} = S \otimes TX \longrightarrow 0$$
(2.18)

(see (2.17)). Consequently, there is a natural inclusion map $\psi : H^0(X, \operatorname{End}(S) \otimes K_X^{\otimes 2}) = H^0(X, \operatorname{Hom}(S \otimes TX, S \otimes K_X)) \hookrightarrow H^0(X, \operatorname{End}(E) \otimes K_X)$ (2.19)

that sends any homomorphism $\alpha : S \otimes TX \longrightarrow S \otimes K_X$ to the following composition of homomorphisms

$$E \xrightarrow{q_0} S \otimes TX \xrightarrow{\alpha} S \otimes K_X \xrightarrow{\iota \otimes \mathrm{Id}_{K_X}} E \otimes K_X,$$

where q_0 and ι are the homomorphisms in (2.18).

The space of holomorphic connections on E is an affine space modeled on the vector space $H^0(X, \operatorname{End}(E) \otimes K_X)$; recall that E is assumed to admit a holomorphic connection. In view of the homomorphism ψ in (2.19), for any holomorphic connection D on E and any $\alpha \in H^0(X, \operatorname{End}(S) \otimes K_X^{\otimes 2})$, we get a holomorphic connection $D + \psi(\alpha)$ on E. If $D \in \widetilde{\mathcal{C}}(F)$ (see Definition 2.7), then it can be shown that

$$D + \psi(\alpha) \in \widetilde{\mathcal{C}}(F)$$
. (2.20)

Indeed, the restrictions of D and $D + \psi(\alpha)$ to the subbundle S in (2.18) coincide. Therefore, the second fundamental form of S for the connection D coincides with the second fundamental form of S for the connection $D + \psi(\alpha)$. Hence (2.20) holds. Consequently, the vector space $H^0(X, \operatorname{End}(S) \otimes K_X^{\otimes 2})$ acts on $\widetilde{\mathcal{C}}(F)$. Let

$$\widetilde{\mathcal{C}}(F) \times H^0(X, \operatorname{End}(S) \otimes K_X^{\otimes 2}) \longrightarrow \widetilde{\mathcal{C}}(F)$$
 (2.21)

be this action. The action in (2.21) is evidently free; however, the action is not transitive (the dimension of $H^0(X, \operatorname{End}(S) \otimes K_X^{\otimes 2})$ is smaller than that of $\widetilde{\mathcal{C}}(F)$).

Let $\operatorname{ad}(S) \subset \operatorname{End}(S)$ be the subbundle of rank $r^2 - 1$ given by the sheaf of endomorphisms of S of trace zero. Note that we have

$$\operatorname{End}(S) = \operatorname{ad}(S) \oplus \mathcal{O}_X;$$
 (2.22)

the inclusion map $\mathcal{O}_X \hookrightarrow \operatorname{End}(S)$ sends a locally defined holomorphic function f on X to the locally defined endomorphism of S that maps any locally defined section s of E to $f \cdot s$. It is evident that $\operatorname{ad}(S) = \operatorname{ad}(F)$ (see (2.17)). The decomposition of $\operatorname{End}(S)$ in (2.22) produces a decomposition

$$H^{0}(X, \operatorname{End}(S) \otimes K_{X}^{\otimes 2}) = H^{0}(X, \operatorname{ad}(S) \otimes K_{X}^{\otimes 2}) \oplus H^{0}(X, K_{X}^{\otimes 2}).$$
(2.23)

Consider the action of $H^0(X, \operatorname{End}(S) \otimes K_X^{\otimes 2})$ on $\widetilde{\mathcal{C}}(F)$ in (2.21). In view of (2.23), from this action we obtain an action of $H^0(X, \operatorname{ad}(S) \otimes K_X^{\otimes 2})$ on $\widetilde{\mathcal{C}}(F)$.

Definition 2.8. Define $\mathcal{C}(F)$ to be the quotient space $\widetilde{\mathcal{C}}(F)/H^0(X, \operatorname{ad}(S) \otimes K_X^{\otimes 2})$ for the above action of $H^0(X, \operatorname{ad}(S) \otimes K_X^{\otimes 2})$ on the space $\widetilde{\mathcal{C}}(F)$ in Definition 2.7.

From (2.23) it follows immediately that $H^0(X, K_X^{\otimes 2})$ acts freely on $\mathcal{C}(F)$. It is evident that

$$\mathcal{C}(F)/H^0(X, K_X^{\otimes 2}) = \widetilde{\mathcal{C}}(F)/H^0(X, \operatorname{End}(S) \otimes K_X^{\otimes 2}).$$

We have a homomorphism

$$\varpi : \operatorname{End}(F) \otimes K_X = \operatorname{Hom}(F \otimes K_X^{-1/2}, F \otimes K_X^{1/2}) \longrightarrow \operatorname{End}(J^1(F \otimes K_X^{-1/2})) \quad (2.24)$$

that sends any locally defined homomorphism $\eta : F \otimes K_X^{-1/2} \longrightarrow F \otimes K_X^{1/2}$ to the following composition of (locally defined) homomorphisms:

$$J^{1}(F \otimes K_{X}^{-1/2}) \xrightarrow{q_{0}} F \otimes K_{X}^{-1/2} \xrightarrow{\eta} F \otimes K_{X}^{1/2} \xrightarrow{\iota} J^{1}(F \otimes K_{X}^{-1/2}),$$

where ι and q_0 are the homomorphisms in (2.18).

Lemma 2.9. Let F be a stable vector bundle on X of rank r and degree zero. The homomorphism

$$\widehat{\varpi} : H^0(X, \operatorname{End}(F) \otimes K_X) \oplus \mathbb{C} \longrightarrow H^0(X, \operatorname{End}(J^1(F \otimes K_X^{-1/2}))),$$

that sends any $(v, c) \in H^0(X, \operatorname{End}(F) \otimes K_X) \oplus \mathbb{C}$ to $\varpi(v) + c \cdot \operatorname{Id}_{J^1(F \otimes K_X^{-1/2})}$, where ϖ is the homomorphism in (2.24), is an isomorphism.

Proof. The holomorphic vector bundle $F \otimes K_X^{-1/2}$ does not admit any holomorphic connection because degree $(F \otimes K_X^{-1/2}) = r(1-g) \neq 0$ [At, p. 202, Proposition 18(i)] (recall that $g \geq 2$). The statement that $F \otimes K_X^{-1/2}$ does not admit any holomorphic connection is equivalent to the statement that the short exact sequence in (2.18) does not split holomorphically [At]. Indeed, a holomorphic splitting homomorphism

$$D_1: J^1(F \otimes K_X^{-1/2}) \longrightarrow (F \otimes K_X^{-1/2}) \otimes K_X$$

for (2.18) such that $D_1 \circ \iota = \text{Id}$, where ι is the homomorphism in (2.18), defines a holomorphic differential operator of order one $\widetilde{D}_1 : F \otimes K_X^{-1/2} \longrightarrow (F \otimes K_X^{-1/2}) \otimes K_X$ which satisfies the Leibniz identity, in other words, \widetilde{D}_1 is a holomorphic connection on $F \otimes K_X^{-1/2}$; conversely, the homomorphism $D_2 : J^1(F \otimes K_X^{-1/2}) \longrightarrow (F \otimes K_X^{-1/2}) \otimes K_X$ corresponding to any holomorphic connection on $F \otimes K_X^{-1/2}$ produces a holomorphic splitting of (2.18).

Take any endomorphism $\rho : J^1(F \otimes K_X^{-1/2}) \longrightarrow J^1(F \otimes K_X^{-1/2})$ over X. Consider the composition of homomorphisms

$$F \otimes K_X^{1/2} \xrightarrow{\iota} J^1(F \otimes K_X^{-1/2}) \xrightarrow{\rho} J^1(F \otimes K_X^{-1/2}) \xrightarrow{q_0} F \otimes K_X^{-1/2}, \qquad (2.25)$$

where ι and q_0 are the homomorphisms in (2.18). Both the vector bundles $F \otimes K_X^{1/2}$ and $F \otimes K_X^{-1/2}$ are stable because F is so. We have

$$\frac{\operatorname{degree}(F \otimes K_X^{1/2})}{\operatorname{rank}(F \otimes K_X^{1/2})} > \frac{\operatorname{degree}(F \otimes K_X^{-1/2})}{\operatorname{rank}(F \otimes K_X^{-1/2})}$$

since $g \ge 2$. Therefore, there is no nonzero homomorphism from $F \otimes K_X^{1/2}$ to $F \otimes K_X^{-1/2}$. In particular, the composition of homomorphisms in (2.25) vanishes identically.

Now let $\rho' := \rho |_{F \otimes K_X^{1/2}} : F \otimes K_X^{1/2} \longrightarrow F \otimes K_X^{1/2}$ be the restriction of ρ to the subbundle $F \otimes K_X^{1/2}$ in (2.18). Since $F \otimes K_X^{1/2}$ is stable, there is a $c \in \mathbb{C}$ such that $\rho' = c \cdot \mathrm{Id}_{F \otimes K_X^{1/2}}$. Define the endomorphism over X

$$\rho_1 := \rho - c \cdot \mathrm{Id}_{J^1(F \otimes K_X^{-1/2})} : J^1(F \otimes K_X^{-1/2}) \longrightarrow J^1(F \otimes K_X^{-1/2}).$$
(2.26)

So we have $\rho_1(F \otimes K_X^{1/2}) = 0$. This implies that there is a homomorphism

$$\rho_2' : J^1(F \otimes K_X^{-1/2}) / (F \otimes K_X^{1/2}) = F \otimes K_X^{-1/2} \longrightarrow J^1(F \otimes K_X^{-1/2})$$

such that

$$\rho_1 = \rho_2' \circ q_0 \,, \tag{2.27}$$

where q_0 and ρ_1 are the homomorphisms in (2.18) and (2.26) respectively. Now $q_0 \circ \rho'_2 \in H^0(X, \operatorname{End}(F \otimes K_X^{-1/2}))$ coincides with $c' \cdot \operatorname{Id}$ for some $c' \in \mathbb{C}$, because $F \otimes K_X^{-1/2}$ is stable. Next we note that if $c' \neq 0$, then $\frac{1}{c'}\rho'_2 : F \otimes K_X^{-1/2} \longrightarrow J^1(F \otimes K_X^{-1/2})$ is a holomorphic splitting of the short exact sequence in (2.18). Since the short exact sequence in (2.18) does not admit a holomorphic splitting (this was shown earlier), we conclude that c' = 0.

Given that c' = 0, it is deduced that there is a homomorphism

$$\rho_2 : F \otimes K_X^{-1/2} \longrightarrow F \otimes K_X^{1/2}$$

such that $\rho'_2 = \iota \circ \rho_2$, where ι and ρ'_2 are the homomorphisms in (2.18) and (2.27) respectively. Therefore, from (2.27) and the definition of ϖ in (2.24) it follows that

$$\rho_1 = \iota \circ \rho_2 \circ q_0 = \varpi(\rho_2).$$

So from (2.26) we know that $\rho = \varpi(\rho_2) + c \cdot \operatorname{Id}_{J^1(F \otimes K_X^{-1/2})}$.

As in Lemma 2.9, F is a stable vector bundle on X of rank r and degree zero. Let

$$\operatorname{Aut}(J^1(F \otimes K_X^{-1/2}))$$

be the group of all holomorphic automorphisms of $J^1(F \otimes K_X^{-1/2})$; it is a complex affine algebraic group, in fact, it is a Zariski open subset of the affine space $H^0(X, \operatorname{End}(J^1(F \otimes K_X^{-1/2})))$.

From Lemma 2.9 it follows immediately that

$$\operatorname{Aut}(J^{1}(F \otimes K_{X}^{-1/2})) = H^{0}(X, \operatorname{End}(F) \otimes K_{X}) \times \mathbb{C}^{\star}, \qquad (2.28)$$

where $\mathbb{C}^{\star} = \mathbb{C} \setminus \{0\}$ is the multiplicative group.

The group $\operatorname{Aut}(J^1(F \otimes K_X^{-1/2}))$ has a natural action on $\operatorname{Conn}(J^1(F \otimes K_X^{-1/2}))$ defined in (2.16). The action of any $T \in \operatorname{Aut}(J^1(F \otimes K_X^{-1/2}))$ sends any holomorphic connection D to the holomorphic connection given by the composition $(T \otimes \operatorname{Id}_{K_X}) \circ D \circ T^{-1}$ of operators.

The isomorphism in (2.28) has the following consequence:

Corollary 2.10. Let F be a stable vector bundle on X of rank r and degree zero. The action of $\operatorname{Aut}(J^1(F \otimes K_X^{-1/2}))$ on $\operatorname{Conn}(J^1(F \otimes K_X^{-1/2}))$ preserves the subset $\widetilde{\mathcal{C}}(F)$ in Definition 2.7.

The action of $\operatorname{Aut}(J^1(F \otimes K_X^{-1/2}))$ on $\widetilde{\mathcal{C}}(F)$ descends to an action of $\operatorname{Aut}(J^1(F \otimes K_X^{-1/2}))$ on the quotient space $\mathcal{C}(F)$ in Definition 2.8.

Proof. The first statement is straightforward. Take any $D \in \text{Conn}(J^1(F \otimes K_X^{-1/2}))$ and $T \in \text{Aut}(J^1(F \otimes K_X^{-1/2}))$. From (2.28) it follows immediately that the second fundamental forms of the subbundle $F \otimes K_X^{1/2} \subset J^1(F \otimes K_X^{-1/2})$ (see (2.18)) for the two connections D and $(T \otimes \text{Id}_{K_X}) \circ D \circ T^{-1}$ coincide; see also Corollary 3.2. This implies that the action of $\text{Aut}(J^1(F \otimes K_X^{-1/2}))$ on $\text{Conn}(J^1(F \otimes K_X^{-1/2}))$ preserves $\widetilde{\mathcal{C}}(F)$.

Take any $D \in \operatorname{Conn}(J^1(F \otimes K_X^{-1/2})), T \in \operatorname{Aut}(J^1(F \otimes K_X^{-1/2}))$ and $B \in H^0(X, \operatorname{ad}(S) \otimes K_X^{\otimes 2}).$

From (2.28) it follows that

$$(T \otimes \mathrm{Id}_{K_X}) \circ (D+B) \circ T^{-1} = (T \otimes \mathrm{Id}_{K_X}) \circ D \circ T^{-1} + B$$

Therefore, the translation action of $H^0(X, \operatorname{ad}(S) \otimes K_X^{\otimes 2})$ on $\widetilde{\mathcal{C}}(F)$ and the action of $\operatorname{Aut}(J^1(F \otimes K_X^{-1/2}))$ on $\widetilde{\mathcal{C}}(F)$ commute. The second statement of the corollary follows from this.

In the next section we will put structures on the quotient space obtained from Corollary 2.10

$$\mathcal{D}(F) := \mathcal{C}(F) / \operatorname{Aut}(J^1(F \otimes K_X^{-1/2})).$$
(2.29)

3. The space of holomorphic connections

Take a stable holomorphic vector bundle F on X of rank r and degree zero.

As before, $K_X^{1/2}$ is a theta characteristic on X. From Proposition 2.4 and Lemma 2.3 we know that $J^1(F \otimes K_X^{-1/2})$ admits a holomorphic connection D_1 for which the second fundamental form $\mathcal{S}(F, D_1)$ in (2.15) is the identity map of $F \otimes K_X^{1/2}$.

Recall that the space $\operatorname{Conn}(J^1(F \otimes K_X^{-1/2}))$ in (2.16) is an affine space modeled on the vector space $H^0(X, \operatorname{End}(J^1(F \otimes K_X^{-1/2})) \otimes K_X)$. Let

$$0 \longrightarrow F \otimes K_X^{3/2} = F \otimes K_X^{1/2} \otimes K_X \xrightarrow{\iota \otimes \operatorname{Id}_{K_X}} J^1(F \otimes K_X^{-1/2}) \otimes K_X \qquad (3.1)$$
$$\xrightarrow{q_0 \otimes \operatorname{Id}_{K_X}} F \otimes K_X^{-1/2} \otimes K_X = F \otimes K_X^{1/2} \longrightarrow 0$$

be the short exact sequence of vector bundles obtained by tensoring the short exact sequence in (2.18) by K_X .

Lemma 3.1. Take any

$$\Psi \in H^0(X, \operatorname{Hom}(J^1(F \otimes K_X^{-1/2}), J^1(F \otimes K_X^{-1/2}) \otimes K_X))$$

Then

$$\Psi(F\otimes K_X^{1/2}) \subset F\otimes K_X^{3/2},$$

where $F \otimes K_X^{1/2}$ (respectively, $F \otimes K_X^{3/2}$) is the subbundle of $J^1(F \otimes K_X^{-1/2})$ (respectively, $J^1(F \otimes K_X^{-1/2}) \otimes K_X$) in (2.18) (respectively, (3.1)).

Proof. Take any holomorphic homomorphism

$$\Psi : J^1(F \otimes K_X^{-1/2}) \longrightarrow J^1(F \otimes K_X^{-1/2}) \otimes K_X.$$

Consider the composition of homomorphisms

$$F \otimes K_X^{1/2} \hookrightarrow J^1(F \otimes K_X^{-1/2}) \xrightarrow{\Psi} J^1(F \otimes K_X^{-1/2}) \otimes K_X$$

$$\xrightarrow{q_0 \otimes \operatorname{Id}_{K_X}} F \otimes K_X^{-1/2} \otimes K_X = F \otimes K_X^{1/2},$$
(3.2)

where $q_0 \otimes \operatorname{Id}_{K_X}$ is the homomorphism in (3.1). Since the vector bundle F, and hence $F \otimes K_X^{1/2}$, is stable, any nonzero holomorphic endomorphism of $F \otimes K_X^{1/2}$ is an isomorphism.

We will now show that the composition of homomorphisms in (3.2) is the zero homomorphism.

To prove this by contradiction, assume that the composition of homomorphisms in (3.2) is nonzero. As observed above, this implies that the composition of homomorphisms in (3.2) is an isomorphism. Consequently, $\Psi(F \otimes K_X^{1/2})$ is a subbundle of $J^1(F \otimes K_X^{-1/2}) \otimes K_X$. Now consider the subbundle

$$\Psi(F \otimes K_X^{1/2}) \otimes TX \subset J^1(F \otimes K_X^{-1/2}) \otimes K_X \otimes TX = J^1(F \otimes K_X^{-1/2}).$$
(3.3)

The condition that the composition of homomorphisms in (3.2) is an isomorphism implies that the homomorphism

$$\Psi(F \otimes K_X^{1/2}) \otimes TX \xrightarrow{q'_0} F \otimes K_X^{-1/2}$$

where q'_0 is the restriction of the projection q_0 in (2.18) (see (3.3)), is an isomorphism. Consequently, the subbundle $\Psi(F \otimes K_X^{1/2}) \otimes TX \subset J^1(F \otimes K_X^{-1/2})$ in (3.3) produces a holomorphic splitting of the short exact sequence in (2.18). But it was observed in the proof of Lemma 2.9 that the short exact sequence in (2.18) does not split holomorphically.

In view of the above contradiction we conclude that the composition of homomorphisms in (3.2) is the zero homomorphism. This proves the lemma.

The following two results are deduced using Lemma 3.1.

Corollary 3.2. As in Lemma 3.1, F is a stable vector bundle on X of degree zero. The space $\widetilde{\mathcal{C}}(F)$ in Definition 2.7 actually coincides with $\operatorname{Conn}(J^1(F \otimes K_X^{-1/2}))$ in (2.16).

Proof. Take any holomorphic connection D on $J^1(F \otimes K_X^{-1/2})$ and any

$$\Psi \in H^0(X, \operatorname{Hom}(J^1(F \otimes K_X^{-1/2}), J^1(F \otimes K_X^{-1/2}) \otimes K_X)).$$

So $D + \Psi$ is a holomorphic connection on $J^1(F \otimes K_X^{-1/2})$. Consider the second fundamental forms of the subbundle $F \otimes K_X^{1/2} \subset J^1(F \otimes K_X^{-1/2})$ for the two connections D and $D + \Psi$. From Lemma 3.1 it follows immediately that these two second fundamental forms actually coincide. Since $J^1(F \otimes K_X^{-1/2})$ admits a holomorphic connection D_1 for which the second fundamental form $\mathcal{S}(F, D_1)$ in (2.15) is the identity map of $F \otimes K_X^{1/2}$ (see Proposition 2.4), the proof is complete.

Lemma 3.3. The vector space $H^0(X, \operatorname{Hom}(J^1(F \otimes K_X^{-1/2}), J^1(F \otimes K_X^{-1/2}) \otimes K_X))$ fits in the following short exact sequence:

$$0 \longrightarrow H^{0}(X, \operatorname{End}(F) \otimes K_{X}^{\otimes 2}) \longrightarrow H^{0}(X, \operatorname{Hom}(J^{1}(F \otimes K_{X}^{-1/2}), J^{1}(F \otimes K_{X}^{-1/2}) \otimes K_{X}))$$
$$\longrightarrow H^{0}(X, \operatorname{End}(F) \otimes K_{X})^{\oplus 2} \longrightarrow 0.$$

Proof. Let

$$\operatorname{Hom}_{P}(J^{1}(F \otimes K_{X}^{-1/2}), J^{1}(F \otimes K_{X}^{-1/2}) \otimes K_{X}) \subset \operatorname{Hom}(J^{1}(F \otimes K_{X}^{-1/2}), J^{1}(F \otimes K_{X}^{-1/2}) \otimes K_{X})$$

be the subbundle defined by the sheaf of homomorphisms from $J^{1}(F \otimes K_{X}^{-1/2})$ to $J^{1}(F \otimes K_{X}^{-1/2}) \otimes K_{X}$ that take the subbundle $F \otimes K_{X}^{1/2}$ in (2.18) to the subbundle $F \otimes K_{X}^{3/2}$ in (3.1). Let

$$\operatorname{Hom}(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}) = \operatorname{End}(F) \otimes K_X^{\otimes 2}$$

$$\xrightarrow{\mathbf{j}} \operatorname{Hom}_P(J^1(F \otimes K_X^{-1/2}), J^1(F \otimes K_X^{-1/2}) \otimes K_X)$$
(3.4)

be the homomorphism that sends any locally defined homomorphism

$$\eta \,:\, F \otimes K_X^{-1/2} \,\longrightarrow\, F \otimes K_X^{3/2}$$

to the following composition of (locally defined) homomorphisms:

$$J^{1}(F \otimes K_{X}^{-1/2}) \xrightarrow{q_{0}} F \otimes K_{X}^{-1/2} \xrightarrow{\eta} F \otimes K_{X}^{3/2} \xrightarrow{\iota \otimes \operatorname{Id}_{K_{X}}} J^{1}(F \otimes K_{X}^{-1/2}) \otimes K_{X},$$

where q_0 and $\iota \otimes \mathrm{Id}_{K_X}$ are the homomorphisms in (2.18) and (3.1) respectively.

We have a natural surjective homomorphism

$$\varpi_1 : \operatorname{Hom}_P(J^1(F \otimes K_X^{-1/2}), J^1(F \otimes K_X^{-1/2}) \otimes K_X) \\ \longrightarrow \operatorname{Hom}(F \otimes K_X^{1/2}, F \otimes K_X^{3/2}) \oplus \operatorname{Hom}(F \otimes K_X^{-1/2}, F \otimes K_X^{1/2}) = (\operatorname{End}(F) \otimes K_X)^{\oplus 2}$$

that sends any homomorphism to the induced homomorphism from the subbundle (respectively, quotient bundle) in (2.18) to the subbundle (respectively, quotient bundle) in (3.1). Consequently, we have a short exact sequence of holomorphic vector bundles on X

$$0 \longrightarrow \operatorname{End}(F) \otimes K_X^{\otimes 2} \xrightarrow{\mathbf{j}} \operatorname{Hom}_P(J^1(F \otimes K_X^{-1/2}), J^1(F \otimes K_X^{-1/2}) \otimes K_X)$$
$$\xrightarrow{\varpi_1} (\operatorname{End}(F) \otimes K_X)^{\oplus 2} \longrightarrow 0, \qquad (3.5)$$

where \mathbf{j} is the homomorphism in (3.4).

Let

$$0 \longrightarrow H^{0}(X, \operatorname{End}(F) \otimes K_{X}^{\otimes 2}) \longrightarrow H^{0}(X, \operatorname{Hom}_{P}(J^{1}(F \otimes K_{X}^{-1/2}), J^{1}(F \otimes K_{X}^{-1/2}) \otimes K_{X}))$$
$$\longrightarrow H^{0}(X, (\operatorname{End}(K) \otimes K_{Y})^{\oplus 2}) \longrightarrow H^{1}(X, \operatorname{End}(F) \otimes K^{\otimes 2})$$
(3.6)

be the long exact sequence of cohomologies corresponding to the exact sequence in (3.5). By Serre duality,

$$H^{1}(X, \operatorname{End}(F) \otimes K_{X}^{\otimes 2}) = H^{0}(X, \operatorname{End}(F) \otimes TX)^{*} = 0, \qquad (3.7)$$

because $\operatorname{End}(F)$ is semistable of degree zero (recall that F is stable) and degree(TX) < 0 (recall that $g \geq 2$); the unitary flat connection on F, [NS], induces a unitary flat connection on $\operatorname{End}(F)$ and hence $\operatorname{End}(F)$ is polystable, in particular, $\operatorname{End}(F)$ is semistable. From Lemma 3.1 we have

$$H^{0}\left(X, \operatorname{Hom}_{P}(J^{1}(F \otimes K_{X}^{-1/2}), J^{1}(F \otimes K_{X}^{-1/2}) \otimes K_{X})\right)$$

= $H^{0}\left(X, \operatorname{Hom}(J^{1}(F \otimes K_{X}^{-1/2}), J^{1}(F \otimes K_{X}^{-1/2}) \otimes K_{X})\right).$

Consequently, the lemma follows from (3.6) and (3.7).

4. INFINITESIMAL DEFORMATIONS

The space of all infinitesimal deformations of a compact Riemann surface X is identified with $H^1(X, TX)$. By Serre duality,

$$H^{1}(X, TX)^{*} = H^{0}(X, K_{X}^{\otimes 2})$$

For a holomorphic vector bundle V on X, the infinitesimal deformations of V, keeping X fixed, are parametrized by $H^1(X, \operatorname{End}(V))$. By Serre duality,

$$H^1(X, \operatorname{End}(V))^* = H^0(X, \operatorname{End}(V) \otimes K_X).$$

Consider the holomorphic vector bundle

$$\operatorname{Diff}_{X}^{i}(V, V) := \operatorname{Hom}(J^{i}(V), V)$$

$$(4.1)$$

on X given by the sheaf of all holomorphic differential operators of order *i* from V to itself. Take the dual of the exact sequence in (2.6) for W = V

$$0 \longrightarrow V^* \stackrel{\iota}{\longrightarrow} J^1(V)^* \longrightarrow V^* \otimes TX \longrightarrow 0.$$

Tensoring it with V the following short exact sequence of holomorphic vector bundles on X is obtained

$$0 \longrightarrow \operatorname{Diff}_{X}^{0}(V, V) = \operatorname{End}(V) \longrightarrow \operatorname{Diff}_{X}^{1}(V, V) \xrightarrow{\sigma_{0}} \operatorname{End}(V) \otimes TX \longrightarrow 0; \quad (4.2)$$

the above projection σ_0 coincides with the symbol map. Using the natural inclusion $\mathcal{O}_X \subset \operatorname{End}(V)$ (see (2.22)), we have $TX \subset \operatorname{End}(V) \otimes TX$. Now define the Atiyah bundle for V

$$\operatorname{At}(V) := \sigma_0^{-1}(TX) \subset \operatorname{Diff}_X^1(V, V), \qquad (4.3)$$

where σ_0 is the projection in (4.2) [At]. The exact sequence in (4.2) produces the short exact sequence of holomorphic vector bundles on X

$$0 \longrightarrow \operatorname{Diff}_{X}^{0}(V, V) = \operatorname{End}(V) \longrightarrow \operatorname{At}(V) \xrightarrow{\sigma} TX \longrightarrow 0, \qquad (4.4)$$

where σ is the restriction of σ_0 (in (4.2)) to the subbundle At(V); this exact sequence is known as the Atiyah exact sequence for V (see [At]).

The space of all infinitesimal deformations of the pair (X, V) is known to be identified with $H^1(X, \operatorname{At}(V))$ (see [Ch, p. 1413, Proposition 4.3], [BHH, p. 127, (2.12)]). The homomorphism

$$H^1(X, \operatorname{At}(V)) \longrightarrow H^1(X, TX)$$
 (4.5)

produced by the projection σ in (4.4) coincides with the forgetful map that sends an infinitesimal deformation of (X, V) to the infinitesimal deformation of X obtained from it by simply forgetting the vector bundle. The homomorphism

$$H^1(X, \operatorname{End}(V)) \longrightarrow H^1(X, \operatorname{At}(V))$$

induced by the homomorphism $\operatorname{End}(V) \longrightarrow \operatorname{At}(V)$ in (4.4) coincides with the map that sends an infinitesimal deformation of V to the infinitesimal deformation of (X, V) associated to it that keeps the Riemann surface fixed.

From the construction of $\operatorname{At}(V)$ in (4.3) it follows immediately that the dual vector bundle $\operatorname{At}(V)^*$ is a quotient of $\operatorname{Diff}^1_X(V, V)^* = J^1(V) \otimes V^*$ (see (4.1)). To describe this quotient, consider the subbundle $V \otimes K_X \subset J^1(V)$ in (2.6). Using it, we have

$$\operatorname{ad}(V) \otimes K_X \subset \operatorname{End}(V) \otimes K_X = (V \otimes K_X) \otimes V^* \subset J^1(V) \otimes V^*,$$
 (4.6)

where ad(V) as before is the sheaf of trace zero endomorphisms of V. From (4.3) it is deduced that

$$\operatorname{At}(V)^* = (J^1(V) \otimes V^*) / (\operatorname{ad}(V) \otimes K_X) = \operatorname{Diff}_X^1(V, V)^* / (\operatorname{ad}(V) \otimes K_X), \qquad (4.7)$$

where the quotient is by the subbundle in (4.6); note that $\operatorname{Diff}_X^1(V, V)^* = J^1(V) \otimes V^*$. In view of (4.7), by Serre duality,

$$H^{1}(X, \operatorname{At}(V))^{*} = H^{0}\left(X, \frac{J^{1}(V) \otimes V^{*}}{\operatorname{ad}(V) \otimes K_{X}} \otimes K_{X}\right).$$

$$(4.8)$$

As in Section 2.1, let $K_X^{1/2}$ be a theta characteristic on X. Since the collection of theta characteristics on X is a discrete set, there is unique way to move the theta characteristic when X moves over a family of Riemann surfaces parametrized by a simply connected space. A consequence of this observation will be explained now.

Consider the Atiyah exact sequence

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \operatorname{At}(K_X^{1/2}) \longrightarrow TX \longrightarrow 0$$

in (4.4) for $V = K_X^{1/2}$. Let $0 = H^0(X, TX) \longrightarrow H^1(X, \mathcal{O}_X) \xrightarrow{h} H^1(X, \operatorname{At}(K_X^{1/2})) \xrightarrow{\xi} H^1(X, TX) \longrightarrow H^2(X, \mathcal{O}_X) = 0$ (4.9)

be the long exact sequence of cohomologies associated to it. The above observation implies that there is a canonical homomorphism

$$\phi : H^1(X, TX) \longrightarrow H^1(X, \operatorname{At}(K_X^{1/2}))$$
(4.10)

that sends an infinitesimal deformation of X to the corresponding infinitesimal deformation of the pair $(X, K_X^{1/2})$. In particular, we have $\xi \circ \phi = \mathrm{Id}_{H^1(X, TX)}$, where ξ is the homomorphism in (4.9).

An alternative description of the homomorphism ϕ in (4.10) is the following. Let $\{U_i\}_{i \in I}$ be a covering of X by open subsets, and let

$$\lambda_{i,j} : U_{ij} = U_i \cap U_j \longrightarrow TU_{ij}, \quad i, j \in I,$$

be a 1-cocycle giving an element of $H^1(X, TX)$. Then $\lambda_{i,j}$ acts on $H^0(U_{ij}, K_X^{1/2})$ by Lie derivation. To define the operation Lie derivation, take a locally defined holomorphic vector field v and a locally defined holomorphic section s of $K_X^{1/2}$; so $s \otimes s$ is a locally defined holomorphic 1-from on X. Now define $L_v s$ by the equation

$$s \otimes L_v s = \frac{1}{2} L_v(s \otimes s) = \frac{1}{2} \left(i_v d(s \otimes s) + d((s \otimes s)(v)) \right).$$

$$(4.11)$$

Using this action by Lie derivation, $\{\lambda_{i,j}\}$ is considered as a 1-cocycle with values in $\operatorname{At}(K_X^{1/2})$. The corresponding element of $H^1(X, \operatorname{At}(K_X^{1/2}))$ is the image, under the homomorphism ϕ in (4.10), of the cohomology class of $\{\lambda_{i,j}\} \in H^1(X, TX)$.

Lemma 4.1. For any holomorphic vector bundle V on X, there is a natural isomorphism

$$\zeta : H^1(X, \operatorname{At}(V)) \xrightarrow{\sim} H^1(X, \operatorname{At}(V \otimes K_X^{-1/2})).$$

Proof. Fix X, V and a theta characteristic $K_X^{1/2}$ on X. Take any infinitesimal deformation $v \in H^1(X, \operatorname{At}(V))$ of (X, V). It corresponds to a family of curves $\mathcal{X} \longrightarrow \operatorname{Spec} \mathbb{C}[\epsilon]/\epsilon^2$ together with a holomorphic vector bundle $\mathcal{V} \longrightarrow \mathcal{X}$. The fiber of \mathcal{X} over $\operatorname{Spec} \mathbb{C}$ is X. Let $v' \in H^1(X, TX)$ be the image of v under the homomorphism $H^1(X, \operatorname{At}(V)) \longrightarrow H^1(X, TX)$ in (4.5). The image of v' under the homomorphism ϕ in (4.10) gives a relative theta line bundle $\mathcal{K}^{1/2} \longrightarrow \mathcal{X}$, whose restriction to X is the chosen theta characteristic $K_X^{1/2}$. Now the pair $(\mathcal{X}, \mathcal{V} \otimes (\mathcal{K}^{1/2})^*)$ corresponds to an element $v'' \in H^1(X, \operatorname{At}(V \otimes K_X^{-1/2}))$. The map

$$\zeta : H^1(X, \operatorname{At}(V)) \longrightarrow H^1(X, \operatorname{At}(V \otimes K_X^{-1/2})), v \longmapsto v''$$

constructed above is evidently an isomorphism.

The above homomorphism ζ can be described in terms of the cocycles as follows. Let $\{U_i\}_{i\in I}$ be a covering of X by open subsets, and let

$$\delta_{i,j} : U_{ij} = U_i \cap U_j \longrightarrow \operatorname{At}(V)|_{U_{ij}}, \quad i, j \in I,$$

be a 1-cocycle giving an element $c \in H^1(X, \operatorname{At}(V))$. So

$$\sigma \circ \delta_{i,j} : U_{ij} = U_i \cap U_j \longrightarrow TU_{ij}, \quad i, j \in I,$$

is a 1-cocycle with values in TX, where σ is the projection in (4.4). Its cohomology class in $H^1(X, TX)$ coincides with the image of c by the homomorphism in (4.5). For any holomorphic sections

$$v \in H^0\left(U_{ij}, V|_{U_{ij}}\right), \ \kappa \in H^0\left(U_{ij}, K_X^{-1/2}|_{U_{ij}}\right),$$

define $\Delta_{ij}(v \otimes \kappa) = \delta_{i,j}(v) \otimes \kappa + v \otimes L_{\sigma \circ \delta_{i,j}} \kappa$, where $L_{\sigma \circ \delta_{i,j}}$ is a the Lie derivation by the vector field $\sigma \circ \delta_{i,j}$ (Lie derivation is defined in (4.11)); recall that $\delta_{i,j}$ is a first order holomorphic differential operator $V|_{U_{ij}} \longrightarrow V|_{U_{ij}}$. It is straightforward to check that

$$\Delta_{ij}((fv)\otimes\kappa) = \Delta_{ij}(v\otimes(f\kappa))$$

for any holomorphic function f on U_{ij} . Consequently, we have

$$\Delta_{ij} \in H^0(U_{ij}, \operatorname{At}(V \otimes K_X^{-1/2})).$$

The homomorphism ζ takes the cohomology class c of $\{\delta_{i,j}\}$ to the cohomology class of $\{\Delta_{ij}\}$ in $H^1(X, \operatorname{At}(V \otimes K_X^{-1/2}))$.

Now take a triple (X, V, D), where X is a compact Riemann surface of genus g, V is a holomorphic vector bundle on X, and D is a holomorphic connection on V. We recall that holomorphic connections on V are precisely the holomorphic splittings of the Atiyah exact sequence in (4.4) [At]. The connection D on V produces a holomorphic differential operator of order one

$$D : \operatorname{At}(V) \longrightarrow \operatorname{End}(V) \otimes K_X$$
 (4.12)

which is constructed as follows. Let $p_1 : \operatorname{At}(V) \longrightarrow \operatorname{End}(V)$ be the holomorphic projection defined by the holomorphic splitting of (4.4) given by D. Let

D': End $(V) \longrightarrow$ End $(V) \otimes K_X$

be the holomorphic connection on End(V) induced by the connection D on V. Then

$$\widetilde{D} = D' \circ p_1$$

Note that \widetilde{D} is not \mathcal{O}_X -linear; it is a differential operator of order one.

Let

$$\mathcal{A}_{\bullet} : \mathcal{A}_{0} := \operatorname{At}(V) \xrightarrow{\widetilde{D}} \mathcal{A}_{1} := \operatorname{End}(V) \otimes K_{X}$$

$$(4.13)$$

be the two term complex of sheaves on X, where D is the differential operator in (4.12), and \mathcal{A}_i is at the *i*-th position. The infinitesimal deformations of the triple (X, V, D) are parametrized by the first hypercohomology

$$\mathbb{H}^{1}(\mathcal{A}_{\bullet}), \qquad (4.14)$$

where \mathcal{A}_{\bullet} is the complex constructed in (4.13) [Ch, p. 1415, Proposition 4.4] (see also [BHH]).

Consider the homomorphisms of complexes

It induces homomorphisms

$$\mathbb{H}^1(\mathcal{A}_{\bullet}) \longrightarrow H^1(X, \operatorname{At}(V)) \text{ and } H^0(X, \operatorname{End}(V) \otimes K_X) \longrightarrow \mathbb{H}^1(\mathcal{A}_{\bullet}).$$

The first homomorphism corresponds to the forgetful map that sends an infinitesimal deformation of the triple (X, V, D) to the infinitesimal deformation of the pair (X, V) obtained from it by simply forgetting the connection. The second homomorphism corresponds to the map that sends an infinitesimal deformation of the connection D to the infinitesimal deformation of the triple (X, V, D) produced by it by keeping the pair (X, V) fixed.

5. Affine space structure

As before, F is a stable vector bundle on X of rank r and degree zero. Set $V = F \otimes K_X^{-1/2}$, and consider the vector space

$$H^0\left(X, \quad \frac{J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{1/2}}{\mathrm{ad}(F \otimes K_X^{-1/2}) \otimes K_X} \otimes K_X\right)$$

constructed in (4.8). Recall the quotient space $\mathcal{D}(F)$ in (2.29).

Proposition 5.1. The space $\mathcal{D}(F)$ in (2.29) is an affine space modeled on the above vector space

$$H^{0}\left(X, \quad \frac{J^{1}(F \otimes K_{X}^{-1/2}) \otimes F^{*} \otimes K_{X}^{1/2}}{\operatorname{ad}(F \otimes K_{X}^{-1/2}) \otimes K_{X}} \otimes K_{X}\right).$$
(5.1)

Proof. Using the natural inclusion map ι in (2.18), we have

$$\operatorname{ad}(F \otimes K_X^{-1/2}) \otimes K_X^{\otimes 2} \hookrightarrow \operatorname{End}(F \otimes K_X^{-1/2}) \otimes K_X^{\otimes 2} = \operatorname{End}(F) \otimes K_X^{\otimes 2}$$
$$= (F \otimes K_X^{1/2}) \otimes (F^* \otimes K_X^{3/2}) \xrightarrow{\iota \otimes \operatorname{Id}_{F^* \otimes K_X^{3/2}}} J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2}$$

(see also (2.22)). It produces a short exact sequence of vector bundles on X

$$0 \longrightarrow \operatorname{ad}(F \otimes K_X^{-1/2}) \otimes K_X^{\otimes 2} \longrightarrow J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2}$$
(5.2)
$$\longrightarrow \frac{J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{1/2}}{\operatorname{ad}(F \otimes K_X^{-1/2}) \otimes K_X} \otimes K_X \longrightarrow 0.$$

Since F is stable, we have

$$H^{1}(X, \operatorname{ad}(F \otimes K_{X}^{-1/2}) \otimes K_{X}^{\otimes 2}) = H^{0}(X, \operatorname{ad}(F) \otimes TX)^{*} = 0$$

because $\operatorname{ad}(F) \otimes TX$ is polystable of negative degree (recall that $g \geq 2$). Therefore, the long exact sequence of cohomologies associated to this short exact sequence in (5.2) gives a short exact sequence

$$0 \longrightarrow H^{0}(X, \operatorname{ad}(F \otimes K_{X}^{-1/2}) \otimes K_{X}^{\otimes 2}) \longrightarrow H^{0}(X, J^{1}(F \otimes K_{X}^{-1/2}) \otimes F^{*} \otimes K_{X}^{3/2}) \quad (5.3)$$
$$\longrightarrow H^{0}\left(X, \ \frac{J^{1}(F \otimes K_{X}^{-1/2}) \otimes F^{*} \otimes K_{X}^{1/2}}{\operatorname{ad}(F \otimes K_{X}^{-1/2}) \otimes K_{X}} \otimes K_{X}\right) \longrightarrow 0.$$

Consequently, the vector space in (5.1) coincides with the quotient space

$$\mathcal{Q}(F) =: \frac{H^0(X, J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2})}{H^0(X, \operatorname{ad}(F \otimes K_X^{-1/2}) \otimes K_X^{\otimes 2})}$$
$$= \frac{H^0(X, J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2})}{H^0(X, \operatorname{ad}(F) \otimes K_X^{\otimes 2})}.$$
(5.4)

Now, we have a natural inclusion map

$$H^{0}(X, J^{1}(F \otimes K_{X}^{-1/2}) \otimes F^{*} \otimes K_{X}^{3/2}) = H^{0}(X, \operatorname{Hom}(F \otimes K_{X}^{-/2}, J^{1}(F \otimes K_{X}^{-1/2}) \otimes K_{X}))$$

$$\hookrightarrow H^{0}(X, \operatorname{Hom}(J^{1}(F \otimes K_{X}^{-1/2}), J^{1}(F \otimes K_{X}^{-1/2}) \otimes K_{X}))$$
(5.5)

that sends any homomorphism $\eta : F \otimes K_X^{-1/2} \longrightarrow J^1(F \otimes K_X^{-1/2}) \otimes K_X$ over X to

$$\eta \circ q_0 : J^1(F \otimes K_X^{-1/2}) \longrightarrow J^1(F \otimes K_X^{-1/2}) \otimes K_X$$

where q_0 is the projection in (2.18). Using the inclusion map in (5.5), the natural action of the vector space $H^0(X, \operatorname{Hom}(J^1(F \otimes K_X^{-1/2}), J^1(F \otimes K_X^{-1/2}) \otimes K_X))$ on the affine space $\operatorname{Conn}(J^1(F \otimes K_X^{-1/2}))$ (defined in (2.16)) restricts to an action of $H^0(X, J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2})$ on $\operatorname{Conn}(J^1(F \otimes K_X^{-1/2}))$.

Recall that Corollary 3.2 says that $\operatorname{Conn}(J^1(F \otimes K_X^{-1/2})) = \widetilde{\mathcal{C}}(F)$. Therefore, the vector space $H^0(X, J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2})$ acts on $\widetilde{\mathcal{C}}(F)$.

It can be shown that the above action of $H^0(X, J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2})$ on $\widetilde{\mathcal{C}}(F)$ produces an action of $H^0(X, J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2})$ on the quotient space $\mathcal{C}(F)$ in Definition 2.8. Indeed, the action of $H^0(X, J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2})$ on $\widetilde{\mathcal{C}}(F)$, and also the quotient map $\widetilde{\mathcal{C}}(F) \longrightarrow \mathcal{C}(F)$, are both constructed using the action of $H^0(X, \operatorname{End}(J^1(F \otimes K_X^{-1/2})) \otimes K_X)$ on $\widetilde{\mathcal{C}}(F)$; as $S = F \otimes K_X^{1/2}$ (see (2.17)), we have $\operatorname{End}(S) = \operatorname{End}(F) = \operatorname{End}(F \otimes K_X^{-1/2})$. Since the group $H^0(X, \operatorname{End}(J^1(F \otimes K_X^{-1/2})) \otimes K_X)$ is abelian, the two actions on $\widetilde{\mathcal{C}}(F)$ commute, and hence we get an action of $H^0(X, J^1(F \otimes K_X^{-1/2})) \otimes F^* \otimes K_X^{3/2}$) on $\mathcal{C}(F)$.

For the above action of $H^0(X, J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2})$ on $\mathcal{C}(F)$, it is evident that the subspace

$$H^{0}(X, \operatorname{ad}(F) \otimes K_{X}^{\otimes 2}) = H^{0}(X, \operatorname{ad}(F \otimes K_{X}^{-1/2}) \otimes K_{X}^{\otimes 2})$$

$$\subset H^{0}(X, J^{1}(F \otimes K_{X}^{-1/2}) \otimes F^{*} \otimes K_{X}^{3/2})$$

in (5.3) acts trivially. Indeed, this follows from the fact that $\mathcal{C}(F)$ is the quotient of $\widetilde{\mathcal{C}}(F)$ by the action of $H^0(X, \operatorname{ad}(F) \otimes K_X^{\otimes 2})$. Thus we have an action on $\mathcal{C}(F)$ of the quotient space $\mathcal{Q}(F)$ in (5.4). This action of $\mathcal{Q}(F)$ on $\mathcal{C}(F)$ is free, because

- the action of $H^0(X, \operatorname{Hom}(J^1(F \otimes K_X^{-1/2}), J^1(F \otimes K_X^{-1/2}) \otimes K_X))$ on $\operatorname{Conn}(J^1(F \otimes K_X^{-1/2}))$ is free, and
- $\mathcal{Q}(F)$ and $\mathcal{C}(F)$ are constructed by quotienting with the same subspace, namely

$$H^0(X, \operatorname{ad}(F) \otimes K_X^{\otimes 2}) \subset H^0(X, J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{3/2})$$

(see (5.4) and Definition 2.8).

Finally from the structure of $\operatorname{Aut}(J^1(F \otimes K_X^{-1/2}))$ shown in (2.28) it follows that the above action of $\mathcal{Q}(F)$ on $\mathcal{C}(F)$ produces an action of $\mathcal{Q}(F)$ on the quotient space $\mathcal{D}(F)$ in (2.29).

Using Lemma 3.3 it follows that the above action of $\mathcal{Q}(F)$ on $\mathcal{D}(F)$ is transitive. This action is also free. Indeed, this follows from (2.28) and the above observation that the action of $\mathcal{Q}(F)$ on $\mathcal{C}(F)$ is free.

We noted earlier that $\mathcal{Q}(F)$ is identified with the vector space in (5.1). Therefore, the above action of $\mathcal{Q}(F)$ on $\mathcal{D}(F)$ produces an action, on $\mathcal{D}(F)$, of the vector space in (5.1). Now the proposition follows from the above properties of the action of $\mathcal{Q}(F)$ on $\mathcal{D}(F)$. \Box

Proposition 5.1 and Lemma 4.1 together give the following:

Corollary 5.2. The space $\mathcal{D}(F)$ is an affine space modeled on the complex vector space $H^{1}(X, \operatorname{At}(F))^{*}$.

Proof. From Lemma 4.1 we know that $H^1(X, \operatorname{At}(F))^* = H^1(X, \operatorname{At}(F \otimes K_X^{-1/2}))^*$. From (4.8) it follows that

 $H^{1}(X, \operatorname{At}(F \otimes K_{X}^{-1/2}))^{*}$

$$= H^0\left(X, \left((J^1(F \otimes K_X^{-1/2}) \otimes F^* \otimes K_X^{1/2}) / (\operatorname{ad}(F \otimes K_X^{-1/2}) \otimes K_X) \right) \otimes K_X \right).$$
Proposition 5.1 completes the proof.

Now Proposition 5.1 completes the proof.

Let \mathcal{M}^{θ}_{a} denote the moduli space of irreducible smooth complex projective curves of genus g, with $g \geq 2$, equipped with a theta characteristic. It is a smooth orbifold of complex dimension 3g - 3. We note that \mathcal{M}_g^{θ} is not connected; the loci of curves with an odd theta characteristic and curves with an even theta characteristic are disconnected. For any fixed $r \geq 1$, let

$$\beta : \mathcal{B}_g(r) \longrightarrow \mathcal{M}_g^\theta \tag{5.6}$$

be the moduli space of triples of the form $(X, K_X^{1/2}, F)$, where

- X is a compact connected Riemann surface of genus g,
- K_X^{1/2} is a theta characteristic on X, and
 F is a stable vector bundle on X of rank r and degree zero.

The map β in (5.6) sends any $(X, K_X^{1/2}, F)$ to $(X, K_X^{1/2})$ by forgetting F. The moduli space $\mathcal{B}_g(r)$ is a smooth orbifold of complex dimension $3g + r^2(g-1) - 2$. Note that $\mathcal{B}_g(r)$ is not connected as \mathcal{M}_q^{θ} is not connected. Let

$$\gamma : \mathcal{H}_g(r) \longrightarrow \mathcal{B}_g(r) \tag{5.7}$$

be the moduli space of quadruples of the form $(X, K_X^{1/2}, F, D)$, where

- X is a compact connected Riemann surface of genus g,
- K_X^{1/2} is a theta characteristic on X,
 F is a stable vector bundle on X of rank r and degree zero, and
- $D \in \mathcal{D}(F)$ (see (2.29)).

The projection γ in (5.7) sends any $(X, K_X^{1/2}, F, D)$ to $(X, K_X^{1/2}, F)$ by simply forgetting D. The moduli space $\mathcal{H}_g(r)$ is a smooth orbifold of complex dimension $2(3g+r^2(g-1)-2)$.

Theorem 5.3. The algebraic fiber bundle $\gamma : \mathcal{H}_g(r) \longrightarrow \mathcal{B}_g(r)$ in (5.7) is an algebraic affine bundle modeled on the holomorphic cotangent bundle $T^*\mathcal{B}_g(r)$. In other words, $\mathcal{H}_g(r)$ is an algebraic torsor over $\mathcal{B}_g(r)$ for $T^*\mathcal{B}(r)$.

Proof. From Proposition 2.4 and Corollary 3.2 we know that γ is surjective.

Since the space of infinitesimal deformations of a pair $(X, K_X^{1/2}, F) \in \mathcal{B}_g(r)$ are parametrized by $H^1(X, \operatorname{At}(F))$ (see (4.5) and (4.10)), we conclude that

$$T^*_{(X,K^{1/2}_X,F)}\mathcal{B}_g(r) = H^1(X, \operatorname{At}(F))^*$$

Therefore, the theorem follows from Corollary 5.2. It should be clarified that the canonical nature of the action of $H^1(X, \operatorname{At}(F))^*$ on $\mathcal{D}(F)$ in Corollary 5.2 ensures that it extends to any given family of Riemann surfaces equipped with a theta characteristic and a stable vector bundle of rank r and degree zero.

On $\mathcal{B}_q(r)$ there is a natural reduced divisor Θ which is defined as follows:

$$\Theta := \{ (X, K_X^{1/2}, F) \in \mathcal{B}_g(r) \mid H^0(X, F \otimes K_X^{1/2}) \neq 0 \} \subset \mathcal{B}_g(r).$$
 (5.8)

Since $\chi(X, F \otimes K_X^{1/2}) = 0$ (Riemann–Roch theorem), it follows that $H^0(X, F \otimes K_X^{1/2}) \neq 0$ if and only if $H^1(X, F \otimes K_X^{1/2}) \neq 0$. For any $(X, K_X^{1/2}, F) \in \Theta$, we have

$$(X, K_X^{1/2}, F^*) \in \Theta,$$
 (5.9)

because $H^0(X, F^* \otimes K_X^{1/2}) = H^1(X, F \otimes K_X^{1/2})^* \neq 0$ by Serre duality.

6. Differential operators, integral kernels and r-opers

6.1. Differential operators and *r*-opers. As before, take any $(X, K_X^{1/2}, F) \in \mathcal{B}_g(r)$ (see (5.6)). Let *D* be a holomorphic connection on $E := J^1(F \otimes K_X^{-1/2})$; it exists by Proposition 2.4. We know that the triple $(E, F \otimes K_X^{1/2}, D)$ is a *r*-oper (see Corollary 3.2 and Proposition 2.4), where $F \otimes K_X^{1/2}$ is considered as a subbundle of *E* using ι in (2.18). Consider the automorphism

$$\varphi : E = J^1(F \otimes K_X^{-1/2}) \xrightarrow{\sim} J^1(F \otimes K_X^{-1/2}) = E$$
(6.1)

constructed in (2.7). It should be clarified that the construction of φ uses D in an essential way; so φ need not be the identity map of E. Imitating the construction of φ in (2.7) we will construct a homomorphism

$$\varphi_2 : E = J^1(F \otimes K_X^{-1/2}) \longrightarrow J^2(F \otimes K_X^{-1/2}).$$
(6.2)

To construct φ_2 , take any $x \in X$ and any $v \in E_x$. As in the proof of Lemma 2.2, let \hat{v} be the unique flat section of E for the connection D, defined on a simply connected open neighborhood $U \subset X$ of x, such that $\hat{v}(x) = v$. So $q_0(\hat{v}) \in H^0(U, F \otimes K_X^{-1/2})$, where q_0 is the projection in (2.18). Now restrict $q_0(\hat{v})$ to the second order infinitesimal neighborhood of x; let $\hat{v}'_2 \in J^2(F \otimes K_X^{-1/2})_x$ be the element obtained this way from $q_0(\hat{v})$. The homomorphism φ_2 sends any $v \in E_x$, $x \in X$, to $\hat{v}'_2 \in J^2(F \otimes K_X^{-1/2})_x$ constructed above from it.

Let

$$0 \longrightarrow F \otimes K_X^{-1/2} \otimes K_X^2 = F \otimes K_X^{3/2} \xrightarrow{\iota_2} J^2(F \otimes K_X^{-1/2}) \xrightarrow{q_2} J^1(F \otimes K_X^{-1/2}) \longrightarrow 0$$
(6.3)

be the canonical short exact sequence of jet bundles. The homomorphism

$$\varphi_2 \circ \varphi^{-1} : J^1(F \otimes K_X^{-1/2}) \longrightarrow J^2(F \otimes K_X^{-1/2}),$$

where φ_2 and φ are the homomorphisms in (6.2) and (6.1) respectively, satisfies the equation

$$q_2 \circ (\varphi_2 \circ \varphi^{-1}) = \operatorname{Id}_{J^1(F \otimes K_X^{-1/2})}, \qquad (6.4)$$

where q_2 is the projection in (6.3). In other words, $\varphi_2 \circ \varphi^{-1}$ gives a holomorphic splitting of the short exact sequence in (6.3). Therefore, there is a unique holomorphic homomorphism

$$\Delta_D : J^2(F \otimes K_X^{-1/2}) \longrightarrow F \otimes K_X^{3/2}$$
(6.5)

such that

- kernel (Δ_D) = image $(\varphi_2 \circ \varphi^{-1})$ = image (φ_2) , and
- $\Delta_D \circ \iota_2 = \mathrm{Id}_{F \otimes K_{\mathbf{v}}^{3/2}}$, where ι_2 is the homomorphism in (6.3).

We note that the homomorphism Δ_D in (6.5) defines a holomorphic differential operator of order two

$$\Delta_D \in H^0(X, \operatorname{Diff}_X^2(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$$
(6.6)

from $F \otimes K_X^{-1/2}$ to $F \otimes K_X^{3/2}$. The symbol of any holomorphic differential operator of order two from $F \otimes K_X^{-1/2}$ to $F \otimes K_X^{3/2}$ is a holomorphic section of

$$\operatorname{Hom}(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}) \otimes (TX)^{\otimes 2} = \operatorname{End}(F).$$

From the above equation $\Delta_D \circ \iota_2 = \operatorname{Id}_{F \otimes K_X^{3/2}}$ it follows immediately that the symbol of the differential operator Δ_D in (6.6) is actually $\operatorname{Id}_F \in H^0(X, \operatorname{End}(F))$.

For any
$$A \in H^0(X, \operatorname{Diff}_X^2(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$$
 and
 $B \in H^0(X, \operatorname{Diff}_X^0(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$
 $= H^0(X, \operatorname{Hom}(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2})) = H^0(X, \operatorname{End}(F) \otimes K_X^2),$

we have

$$A + B \in H^0(X, \operatorname{Diff}_X^2(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$$

Also, the symbol of A + B coincides with the symbol of A, because B is a lower order differential operator.

Two holomorphic differential operators $A_1, A_2 \in H^0(X, \operatorname{Diff}_X^2(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$ will be called *equivalent* if

$$A_1 - A_2 \in H^0(X, \operatorname{ad}(F) \otimes K_X^2) \subset H^0(X, \operatorname{End}(F) \otimes K_X^2).$$
(6.7)

Definition 6.1. The space of all equivalence classes of differential operators

$$A \in H^0(X, \operatorname{Diff}_X^2(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$$

such that the symbol of A is $\mathrm{Id}_F \in H^0(X, \mathrm{End}(F))$ will be denoted by $\widetilde{\mathcal{D}}(X, K_X^{1/2}, F)$.

The above construction of Δ_D in (6.6) from D gives the following:

Lemma 6.2. For any $(X, K_X^{1/2}, F) \in \mathcal{B}_g(r)$, there is a natural map

$$\Psi : \mathcal{D}(F) \longrightarrow \mathcal{D}(X, K_X^{1/2}, F),$$

where $\mathcal{D}(F)$ and $\widetilde{\mathcal{D}}(X, K_X^{1/2}, F)$ are constructed in (2.29) and Definition 6.1 respectively.

Proof. Let $\widetilde{\mathcal{D}}'(X, K_X^{1/2}, F)$ denote the space of all holomorphic differential operators $A \in H^0(X, \operatorname{Diff}^2_X(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$ such that the symbol of A is $\operatorname{Id}_F \in H^0(X, \operatorname{End}(F))$. The construction of Δ_D in (6.6) from D clearly gives a map

$$\Psi'\,:\,\widetilde{\mathcal{C}}(F)\,\longrightarrow\,\widetilde{\mathcal{D}}'(X,\,K^{1/2}_X,\,F)$$

(see Definition 2.7). Now consider the quotient space $\mathcal{C}(F)$ of $\widetilde{\mathcal{C}}(F)$ in Definition 2.8. Take two element $D_1, D_2 \in \widetilde{\mathcal{C}}(F)$ that give the same element of $\mathcal{C}(F)$. It is straightforward to check that the differential operators $\Psi'(D_1)$ and $\Psi'(D_2)$ are equivalent. Therefore, Ψ' gives a map

$$\Psi'' : \mathcal{C}(F) \longrightarrow \widetilde{\mathcal{D}}(X, K_X^{1/2}, F).$$

The map Ψ'' clearly factors through the quotient $\mathcal{D}(F)$ of $\mathcal{C}(F)$ in (2.29). Hence Ψ'' produces a map Ψ as in the statement of the lemma.

We will construct an inverse of the map Ψ in Lemma 6.2.

For a holomorphic vector bundle W on X, there is a canonical commutative diagram of homomorphisms

where the rows and columns are exact; the top (respectively, bottom) row is the one as in (6.3) (respectively, (2.6)), while the left column is (2.6) tensored with K_X . The map **b** is tautological; it follows from the definition of jet bundles. Now set

$$W = F \otimes K_X^{-1/2}$$

in (6.8), where F is a stable vector bundle of rank r and degree zero on X, and $K_X^{1/2}$ is a theta characteristic on X. Let

$$\Delta_0 : J^2(F \otimes K_X^{-1/2}) \longrightarrow F \otimes K_X^{-1/2} \otimes K_X^2 = F \otimes K_X^{3/2}$$

be a holomorphic homomorphism such that

$$\Delta_0 \circ \iota_2 = \operatorname{Id}_{F \otimes K_X^{3/2}},\tag{6.9}$$

where ι_2 is the homomorphism in (6.8). In other words,

$$\Delta_0 \in H^0(X, \operatorname{Diff}^2_X(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$$

and the symbol of Δ_0 is $\mathrm{Id}_F \in H^0(X, \mathrm{End}(F))$ (it is equivalent to the equation in (6.9)). From (6.9) it follows that Δ_0 gives a holomorphic splitting of the top row in (6.8). Consequently, there is a unique holomorphic homomorphism

$$D': J^{1}(F \otimes K_{X}^{-1/2}) \longrightarrow J^{2}(F \otimes K_{X}^{-1/2})$$

and
$$q_{2} \circ D' = \operatorname{Id}_{J^{1}(F \otimes K_{X}^{-1/2})}, \qquad (6.10)$$

such that $\Delta_0 \circ D' = 0$ and

where q_2 is the projection in (6.8).

Now consider the composition of homomorphisms

$$D := \mathbf{b} \circ D' : J^1(F \otimes K_X^{-1/2}) \longrightarrow J^1(J^1(F \otimes K_X^{-1/2})), \qquad (6.11)$$

where **b** is the homomorphism in (6.8). Since the diagram in (6.8) is commutative, from (6.10) it follows that

$$q' \circ D = q' \circ \mathbf{b} \circ D' = \mathrm{Id}_{J^{1}(F \otimes K_{X}^{-1/2})} \circ q_{2} \circ D' = \mathrm{Id}_{J^{1}(F \otimes K_{X}^{-1/2})},$$

where q' is the projection in (6.8). In other words, D in (6.11) gives a holomorphic connection on the vector bundle $J^1(F \otimes K_X^{-1/2})$. The differential operator $J^1(F \otimes K_X^{-1/2}) \longrightarrow J^1(F \otimes K_X^{-1/2}) \otimes K_X$ for the connection D is the unique \mathcal{O}_X -linear homomorphism

$$\widehat{D} : J^1(J^1(F \otimes K_X^{-1/2})) \longrightarrow J^1(F \otimes K_X^{-1/2}) \otimes K_X$$

that satisfies the following conditions:

- kernel (\widehat{D}) = image(D), and
- $\widehat{D} \circ \iota' = \operatorname{Id}_{J^1(F \otimes K_Y^{-1/2}) \otimes K_X}$, where ι' is the homomorphism in (6.8).

From Corollary 3.2 we know that any holomorphic connection on $J^1(F \otimes K_X^{-1/2})$ gives an element of $\mathcal{C}(F)$ (see Definition 2.8). Hence the holomorphic connection D in (6.11) gives an element

$$D_{\Delta_0} \in \mathcal{D}(F), \tag{6.12}$$

where $\mathcal{D}(F)$ is the quotient of $\mathcal{C}(F)$ defined in (2.29).

Lemma 6.3. For any $(X, K_X^{1/2}, F) \in \mathcal{B}_g(r)$ (see (5.6)), there is a natural map $\Phi : \widetilde{\mathcal{D}}(X, K_X^{1/2}, F) \longrightarrow \mathcal{D}(F),$

where $\widetilde{\mathcal{D}}(X, K_X^{1/2}, F)$ and $\mathcal{D}(F)$ are constructed in Definition 6.1 and (2.29) respectively.

Proof. The above construction of D_{Δ_0} (in (6.12)) from Δ_0 produces a map

$$\Phi': \widetilde{\mathcal{D}}'(X, K_X^{1/2}, F) \longrightarrow \mathcal{D}(F), \tag{6.13}$$

where $\widetilde{\mathcal{D}}'(X, K_X^{1/2}, F)$ is defined in the proof of Lemma 6.2.

Take $\Delta_1, \Delta_2 \in \widetilde{\mathcal{D}}'(X, K_X^{1/2}, F)$ such that Δ_1 is equivalent to Δ_2 (see (6.7)). Let D_1 (respectively, D_2) be the holomorphic connection on $J^1(F \otimes K_X^{-1/2})$ corresponding to Δ_1 (respectively, Δ_2); see (6.11). Since

$$\Delta_1 - \Delta_2 \in H^0(X, \operatorname{ad}(F) \otimes K_X^2),$$

we have

$$D_1 - D_2 = \Delta_1 - \Delta_2 \in H^0(X, \operatorname{ad}(F) \otimes K_X^2).$$

Hence D_1 and D_2 give the same element of $\mathcal{C}(F)$ defined in Definition 2.8. This implies that Φ' in (6.13) produces a map Φ as in the statement of the lemma.

Theorem 6.4. The two maps Ψ and Φ , constructed in Lemma 6.2 and Lemma 6.3 respectively, are inverses of each other.

Proof. In view of the explicit nature of the maps Ψ and Φ , this is a verification by straightforward computations. We omit the details.

6.2. Integral kernels and differential operators. For i = 1, 2, let

 $q_i : X \times X \longrightarrow X$

be the projection to the *i*-th factor. For holomorphic vector bundles A, B on X, the holomorphic vector bundle $(q_1^*A) \otimes (q_2^*B)$ on $X \times X$ will be denoted by $A \boxtimes B$. Let

$$\mathbf{\Delta} := \{ (x, x) \mid x \in X \} \subset X \times X$$

be the reduced diagonal divisor.

For holomorphic vector bundles A, B on X, and a nonnegative integer d, we will construct a torsion sheaf on $X \times X$ supported on the divisor $(d+1)\Delta \subset X \times X$. Let r_A and r_B be the ranks of A and B respectively.

Consider the holomorphic vector bundles

$$A \boxtimes (B^* \otimes K_X)$$
 and $A \boxtimes (B^* \otimes K_X) \otimes \mathcal{O}_{X \times X}((d+1)\Delta)$

on $X \times X$. We note that $A \boxtimes (B^* \otimes K_X)$ is a subsheaf of $A \boxtimes (B^* \otimes K_X) \otimes \mathcal{O}_{X \times X}((d+1)\Delta)$ because $(d+1)\Delta$ is an effective divisor on $X \times X$. So we have a short exact sequence of coherent sheaves on $X \times X$

$$0 \longrightarrow A \boxtimes (B^* \otimes K_X) \longrightarrow A \boxtimes (B^* \otimes K_X) \otimes \mathcal{O}_{X \times X}((d+1)\mathbf{\Delta})$$
$$\longrightarrow \mathcal{Q}_d(A, B) := \frac{A \boxtimes (B^* \otimes K_X) \otimes \mathcal{O}_{X \times X}((d+1)\mathbf{\Delta})}{A \boxtimes (B^* \otimes K_X)} \longrightarrow 0;$$
(6.14)

the support of the above quotient sheaf $\mathcal{Q}_d(A, B)$ in (6.14) is $(d+1)\Delta$. The direct image

$$\mathcal{K}_d(A, B) := q_{1*}\mathcal{Q}_d(A, B) \tag{6.15}$$

is a holomorphic vector bundle on X of rank $r_A r_B(d+1)$. It is known that

$$\mathcal{K}_d(A, B) = \operatorname{Hom}(J^d(B), A) = \operatorname{Diff}_X^d(B, A), \qquad (6.16)$$

where $\mathcal{K}_d(A, B)$ is constructed in (6.15) (see [BS, Section 2.1], [Bi, p. 25, (5.1)], [BB, Section 3.1, p. 1314]).

For $d \geq 1$, the sheaf $\mathcal{Q}_d(A, B)$ in (6.14) fits in the following short exact sequence of sheaves on $X \times X$

$$0 \longrightarrow \mathcal{Q}_{d-1}(A, B) := \frac{A \boxtimes (B^* \otimes K_X) \otimes \mathcal{O}_{X \times X}(d\Delta)}{A \boxtimes (B^* \otimes K_X)} \longrightarrow \mathcal{Q}_d(A, B)$$
(6.17)

$$:=\frac{A\boxtimes (B^*\otimes K_X)\otimes \mathcal{O}_{X\times X}((d+1)\mathbf{\Delta})}{A\boxtimes (B^*\otimes K_X)}\longrightarrow \frac{A\boxtimes (B^*\otimes K_X)\otimes \mathcal{O}_{X\times X}((d+1)\mathbf{\Delta})}{A\boxtimes (B^*\otimes K_X)\otimes \mathcal{O}_{X\times X}(d\mathbf{\Delta})}\longrightarrow 0.$$

The above sheaf $\frac{A\boxtimes(B^*\otimes K_X)\otimes \mathcal{O}_{X\times X}((d+1)\Delta)}{A\boxtimes(B^*\otimes K_X)\otimes \mathcal{O}_{X\times X}(d\Delta)}$ is supported on the reduced divisor Δ . Taking direct image of the short exact sequence in (6.17) by the projection q_1 we get the following short exact sequence holomorphic vector bundles on X

$$0 \longrightarrow q_{1*}\mathcal{Q}_{d-1}(A, B) = \mathcal{K}_{d-1}(A, B) \longrightarrow q_{1*}\mathcal{Q}_d(A, B) = \mathcal{K}_d(A, B)$$
$$\xrightarrow{p_0} q_{1*}\left(\frac{A \boxtimes (B^* \otimes K_X) \otimes \mathcal{O}_{X \times X}((d+1)\mathbf{\Delta})}{A \boxtimes (B^* \otimes K_X) \otimes \mathcal{O}_{X \times X}(d\mathbf{\Delta})}\right) \longrightarrow 0.$$
(6.18)

Poincaré adjunction formula says that $\mathcal{O}_{X \times X}(\Delta)|_{\Delta}$ is the normal bundle of Δ [GH, p. 146]. So $\mathcal{O}_{X \times X}(\Delta)|_{\Delta} = TX$, using the identification of Δ with X defined by $x \mapsto (x, x)$. Therefore, we have

$$q_{1*}\left(\frac{A\boxtimes (B^*\otimes K_X)\otimes \mathcal{O}_{X\times X}((d+1)\mathbf{\Delta})}{A\boxtimes (B^*\otimes K_X)\otimes \mathcal{O}_{X\times X}(d\mathbf{\Delta})}\right) = \operatorname{Hom}(B, A)\otimes (TX)^{\otimes d}.$$

Hence the isomorphism in (6.16) and the projection p_0 in (6.18) together produce a homomorphism

$$\operatorname{Diff}_X^d(B, A) \longrightarrow \operatorname{Hom}(B, A) \otimes (TX)^{\otimes d}$$

The homomorphism of global sections corresponding to it

$$H^0(X, \operatorname{Diff}^d_X(B, A)) \longrightarrow H^0(X, \operatorname{Hom}(B, A) \otimes (TX)^{\otimes d})$$
 (6.19)

is the symbol map on the global differential operators.

Let F be a stable vector bundle on X of rank r and degree zero. Using the isomorphism in (6.16), the space of holomorphic differential operators $H^0(X, \operatorname{Diff}_X^2(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$ has the following isomorphism:

$$H^{0}(X, \operatorname{Diff}_{X}^{2}(F \otimes K_{X}^{-1/2}, F \otimes K_{X}^{3/2}))$$

$$\xrightarrow{\sim} H^{0}(3\Delta, ((F \otimes K_{X}^{3/2}) \boxtimes (F^{*} \otimes K_{X}^{3/2}) \otimes \mathcal{O}_{X \times X}(3\Delta))|_{3\Delta}).$$

$$(6.20)$$

Now, for any

$$s \in H^0(3\Delta, ((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\Delta))|_{3\Delta}),$$

let

$$s_0 := s \big|_{\mathbf{\Delta}} \in H^0\big(\mathbf{\Delta}, (F \boxtimes F)\big|_{\mathbf{\Delta}}\big) = H^0(X, \operatorname{End}(F))$$

$$(6.21)$$

be the restriction of it to $\Delta \subset 3\Delta$; note that $((K_X^{3/2} \boxtimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\Delta))|_{\Delta}$ is canonically trivialized, because $\mathcal{O}_{X \times X}(\Delta)|_{\Delta} = TX$ (after identifying Δ with X). Then s_0 coincides with the symbol of the differential operator

 $D_s \in H^0(X, \operatorname{Diff}^2_X(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$

corresponding to the above section s (see (6.19) and (6.20)).

Consider the natural short exact sequence

$$0 \longrightarrow K_{\mathbf{\Delta}}^{\otimes 2} = K_X^{\otimes 2} \longrightarrow \mathcal{O}_{3\mathbf{\Delta}} \longrightarrow \mathcal{O}_{2\mathbf{\Delta}} \longrightarrow 0.$$

Tensoring it with $(F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\Delta)$, and then taking global sections, we see that the vector space

$$H^{0}(\boldsymbol{\Delta}, \left((F \otimes K_{X}^{3/2}) \boxtimes (F^{*} \otimes K_{X}^{3/2}) \otimes \mathcal{O}_{X \times X}(3\boldsymbol{\Delta}) \right) \big|_{\boldsymbol{\Delta}} \otimes K_{\boldsymbol{\Delta}}^{\otimes 2} \right) = H^{0}(X, \operatorname{End}(F) \otimes K_{X}^{\otimes 2})$$

is a subspace of $H^0(3\Delta, ((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\Delta))|_{3\Delta})$. Take two sections $s, t \in H^0(3\Delta, ((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\Delta))|_{3\Delta}).$

Then the corresponding differential operators (see (6.20))

 $D_s, D_t \in H^0(X, \operatorname{Diff}_X^2(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}))$

are equivalent (see (6.7)) if and only if

$$s - t \in H^0(X, \operatorname{ad}(F) \otimes K_X^{\otimes 2}) \subset H^0(X, \operatorname{End}(F) \otimes K_X^{\otimes 2}).$$

Consequently, we obtain the following description of $\widetilde{\mathcal{D}}(X, K_X^{1/2}, F)$ (see Definition 6.1) in terms of integral kernels: $\widetilde{\mathcal{D}}(X, K_X^{1/2}, F)$ is identified with the quotient of

$$\{s \in H^0(3\mathbf{\Delta}, \left((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\mathbf{\Delta})\right)|_{3\mathbf{\Delta}}) |s|_{\mathbf{\Delta}} = H^0(X, \operatorname{End}(F))\}$$

by the subspace $H^0(X, \operatorname{ad}(F) \otimes K_X^{\otimes 2})$ of it. Using this description of $\widetilde{\mathcal{D}}(X, K_X^{1/2}, F)$, Theorem 6.4 gives the following description of $\mathcal{H}_g(r)$ in terms of the integral kernels.

Corollary 6.5. The moduli space $\mathcal{H}_g(r)$ in (5.7) is identified with the space of quadruples of the form $(X, K_X^{1/2}, F, s)$, where $(X, K_X^{1/2}, F) \in \mathcal{B}_g(r)$ (see (5.6)) and

$$s \in \frac{H^0\left(3\boldsymbol{\Delta}, \left((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\boldsymbol{\Delta})\right)\big|_{3\boldsymbol{\Delta}}\right)}{H^0(X, \operatorname{ad}(F) \otimes K_X^{\otimes 2})}$$

such that $s_0 := s \big|_{\mathbf{\Delta}} = \mathrm{Id}_F$ (see (6.21)).

7. A CANONICAL HOLOMORPHIC SECTION

Recall the projection γ in (5.7) and the effective divisor Θ in (5.8). In this section we we will construct a holomorphic map on the complement of the theta divisor

$$\mathbb{S}: \mathcal{B}_g(r) \setminus \Theta \longrightarrow \mathcal{H}_g(r)$$

such that $\gamma \circ \mathbb{S} = \mathrm{Id}_{\mathcal{B}_g(r) \setminus \Theta}$.

Take any
$$(X, K_X^{1/2}, F) \in \mathcal{B}_g(r) \setminus \Theta$$
. Since
 $H^0(X, F \otimes K_X^{1/2}) = 0 = H^1(X, F \otimes K_X^{1/2}),$

it follows that

$$H^{0}(X \times X, (F \otimes K_{X}^{1/2}) \boxtimes (F^{*} \otimes K_{X}^{1/2}) \otimes \mathcal{O}_{X \times X}(\mathbf{\Delta})) = H^{0}(X, \operatorname{End}(F \otimes K_{X}^{1/2})) = \operatorname{End}(F)$$

$$(7.1)$$

(see [BH1, Remark 2.4], [BH2, p. 8]). As in [BH2, (3.6)], let

$$\beta_F \in H^0(X \times X, (F \otimes K_X^{1/2}) \boxtimes (F^* \otimes K_X^{1/2}) \otimes \mathcal{O}_{X \times X}(\mathbf{\Delta}))$$
(7.2)

be the section that corresponds to $\mathrm{Id}_F \in H^0(X, \mathrm{End}(F))$ by the isomorphism in (7.1). The line bundle over 2Δ

$$K_X^{1/2} \boxtimes K_X^{1/2} \otimes \mathcal{O}_{X \times X}(\mathbf{\Delta}) \Big|_{2\mathbf{\Delta}} \longrightarrow 2\mathbf{\Delta}$$

has a natural trivialization given by a section

$$\sigma_0 \in H^0(2\mathbf{\Delta}, K_X^{1/2} \boxtimes K_X^{1/2} \otimes \mathcal{O}_{X \times X}(\mathbf{\Delta})\big|_{2\mathbf{\Delta}})$$
(7.3)

[BH2, (3.8)], [BR, p. 688, Theorem 2.2]. Let

$$\widehat{\beta}_F \in H^0(2\Delta, (F \boxtimes F^*)|_{2\Delta})$$
(7.4)

be the section defined by the equation $\beta_F|_{2\Delta} = \hat{\beta}_F \otimes \sigma_0$, where β_F is the section in (7.2) [BH2, (3.9)]. The restriction of $\hat{\beta}_F$ to $\Delta \subset 2\Delta$ is Id_F [BH2], and hence $\hat{\beta}_F$ defines a holomorphic connection on F [BH2, (3.10)]; the holomorphic connection on F given by $\hat{\beta}_F$ will be denoted by $\hat{\beta}'_F$.

Any holomorphic connection on X is integrable, because $\Omega_X^{2,0} = 0$. Therefore, using the above holomorphic connection $\hat{\beta}'_F$, for any simply connected open subset $U \subset X$, the restriction $F|_U$ is canonically identified with the trivialized holomorphic vector bundle $F_{x_0} \times U \longrightarrow U$, for any point $x_0 \in U$. More precisely, this identification of vector bundles is constructed by taking parallel translations of F_{x_0} , for the integrable connection $\hat{\beta}'_F$, along paths originating from x_0 . Consequently, the two holomorphic vector bundles q_1^*F and q_2^*F are holomorphically identified over an analytic neighborhood of $\Delta \subset X \times X$. The restriction of this isomorphism to 2Δ coincides with the isomorphism

$$(q_1^*F)\Big|_{2\mathbf{\Delta}} \xrightarrow{\sim} (q_2^*F)\Big|_{2\mathbf{\Delta}}$$
 (7.5)

given by $\widehat{\beta}_F$ in (7.4). Therefore, we have an extension of the isomorphism in (7.5) to an isomorphism

$$(q_1^*F)\Big|_{m\Delta} \xrightarrow{\sim} (q_2^*F)\Big|_{m\Delta}$$

for every $m \geq 2$, which is given by $\widehat{\beta}'_F$. In particular, we get an isomorphism

$$f_3 : (q_1^*F)\big|_{3\mathbf{\Delta}} \xrightarrow{\sim} (q_2^*F)\big|_{3\mathbf{\Delta}}$$
 (7.6)

extending the isomorphism in (7.5).

Using the isomorphism f_3 in (7.6), the section $\beta_F|_{3\Lambda}$ in (7.2) becomes a section

$$\beta_{3,F} := \beta_F \big|_{3\mathbf{\Delta}} \in H^0 \left((3\mathbf{\Delta}, \left(\left((F \otimes F^* \otimes K_X^{1/2}) \boxtimes K_X^{1/2} \right) \otimes \mathcal{O}_{X \times X}(\mathbf{\Delta}) \right) \big|_{3\mathbf{\Delta}} \right) \\ = H^0 \left(3\mathbf{\Delta}, \left((K_X^{1/2} \boxtimes K_X^{1/2}) \otimes q_1^* \operatorname{End}(F) \otimes \mathcal{O}_{X \times X}(\mathbf{\Delta}) \right) \big|_{3\mathbf{\Delta}} \right).$$

Now using the trace homomorphism

$$q_1^* \operatorname{End}(F) \longrightarrow q_1^* \mathcal{O}_X = \mathcal{O}_{X \times X}, \ w \longmapsto \frac{1}{r} \operatorname{trace}(w)$$

the above section $\beta_{3,F}$ produces a section

$$\sigma_1 \in H^0(3\Delta, \left(K_X^{1/2} \boxtimes K_X^{1/2} \otimes \mathcal{O}_{X \times X}(\Delta)\right)\big|_{3\Delta});$$
(7.7)

the restriction of σ_1 to $2\Delta \subset 3\Delta$ coincides with the section σ_0 in (7.3).

Now define

$$\gamma_F := (\beta_F)\big|_{3\mathbf{\Delta}} \otimes (\sigma_1)^{\otimes 2} \in H^0\left(3\mathbf{\Delta}, \left(\left((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2})\right) \otimes \mathcal{O}_{X \times X}(3\mathbf{\Delta})\right)\big|_{3\mathbf{\Delta}}\right) \\ = H^0\left(X, q_{1*}\left(\left(\left((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2})\right) \otimes \mathcal{O}_{X \times X}(3\mathbf{\Delta})\right)\big|_{3\mathbf{\Delta}}\right)\right),$$
(7.8)

where β_F and σ_1 are the sections constructed in (7.2) and (7.7) respectively; here the restriction of the projection q_1 to $3\Delta \subset X \times X$ is also denoted by q_1 .

We have

$$\left(\left((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2})\right) \otimes \mathcal{O}_{X \times X}(3\mathbf{\Delta})\right)\Big|_{3\mathbf{\Delta}} = \mathcal{Q}_2(F \otimes K_X^{3/2}, F \otimes K_X^{-1/2})$$

(see (6.14)), which implies that

$$q_{1*}\left(\left(\left((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2})\right) \otimes \mathcal{O}_{X \times X}(3\Delta)\right)\Big|_{3\Delta}\right) = \mathcal{K}_2(F \otimes K_X^{3/2}, F \otimes K_X^{-1/2})$$

(see (6.15)). Therefore, from (6.16) it follows that

$$H^{0}\left(X, q_{1*}\left(\left(\left(F \otimes K_{X}^{3/2}\right) \boxtimes \left(F^{*} \otimes K_{X}^{3/2}\right)\right) \otimes \mathcal{O}_{X \times X}(3\boldsymbol{\Delta})\right)\Big|_{3\boldsymbol{\Delta}}\right)\right)$$

= $H^{0}(X, \mathcal{K}_{2}(F \otimes K_{X}^{3/2}, F \otimes K_{X}^{-1/2})) = H^{0}(X, \operatorname{Diff}_{X}^{2}(F \otimes K_{X}^{-1/2}, F \otimes K_{X}^{3/2})).$

Using this isomorphism, the section γ_F constructed in (7.8) gives a holomorphic differential operator

$$\widehat{\gamma}_F \in H^0(X, \operatorname{Diff}_X^2(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2})).$$
(7.9)

From the symbol homomorphism constructed in (6.19) it can be shown that the symbol of the differential operator $\hat{\gamma}_F$ in (7.9) is

$$\mathrm{Id}_F \in H^0(X, \operatorname{Hom}(F \otimes K_X^{-1/2}, F \otimes K_X^{3/2}) \otimes (TX)^{\otimes 2}) = H^0(X, \operatorname{End}(F)).$$

Indeed, in view of the construction of γ_F in (7.8), this is a consequence the following two facts:

- The restriction of the section $\widehat{\beta}_F$ in (7.4) to $\Delta \subset 2\Delta$ is Id_F, and
- the restriction of the section σ_1 in (7.7) to $\Delta \subset 3\Delta$ is the constant function 1 on Δ (recall that the restriction of σ_1 to $2\Delta \subset 3\Delta$ coincides with σ_0 in (7.3)).

Recall the space of differential operators $\widetilde{\mathcal{D}}'(X, K_X^{1/2}, F)$ defined in the proof of Lemma 6.2. Since the symbol of the differential operator $\widehat{\gamma}_F$ in (7.9) is Id_F, it is an element of $\widetilde{\mathcal{D}}'(X, K_X^{1/2}, F)$. Therefore, $\widehat{\gamma}_F$ gives an element of the quotient space $\widetilde{\mathcal{D}}(X, K_X^{1/2}, F)$ of $\widetilde{\mathcal{D}}'(X, K_X^{1/2}, F)$ (see Definition 6.1). Let

$$\widehat{\gamma}'_F \in \widetilde{\mathcal{D}}(X, \, K_X^{1/2}, \, F) \tag{7.10}$$

be the element given by $\widehat{\gamma}_{F}$.

The above construction is summarized in the following lemma.

Lemma 7.1. There is a natural holomorphic map

$$\mathbb{S}: \mathcal{B}_g(r) \setminus \Theta \longrightarrow \mathcal{H}_g(r)$$

such that $\gamma \circ \mathbb{S} = \mathrm{Id}_{\mathcal{B}_g(r) \setminus \Theta}$, where γ is the projection in (5.7).

Proof. Take any $(X, K_X^{1/2}, F) \in \mathcal{B}_g(r) \setminus \Theta$. From Theorem 6.4 we know that $(X, K_X^{1/2}, F, \widehat{\gamma}'_F) \in \mathcal{H}_a(r),$

$$X, K_X^{1/2}, F, \widehat{\gamma}'_F) \in \mathcal{H}_g(r),$$

where $\widehat{\gamma}'_F$ is constructed from $(X, K_X^{1/2}, F)$ in (7.10). Let

$$\mathbb{S} : \mathcal{B}_g(r) \setminus \Theta \longrightarrow \mathcal{H}_g(r)$$

be the map that sends any $(X, K_X^{1/2}, F) \in \mathcal{B}_g(r) \setminus \Theta$ to $(X, K_X^{1/2}, F, \widehat{\gamma}'_F)$. It is evident that $\gamma \circ \mathbb{S} = \mathrm{Id}_{\mathcal{B}_q(r) \setminus \Theta}$.

The above map S can also be described using Corollary 6.5.

8. A CANONICAL ISOMORPHISM OF TORSORS AND A SYMPLECTIC STRUCTURE

8.1. A canonical isomorphism of torsors. Let

$$\mathcal{L} := \mathcal{O}_{\mathcal{B}_g(r)}(\Theta) \longrightarrow \mathcal{B}_g(r) \tag{8.1}$$

be the line bundle corresponding to the reduced divisor Θ in (5.8). Consider the short exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathcal{B}_g(r)} = \operatorname{Diff}^0_X(\mathcal{L}, \mathcal{L}) \longrightarrow \operatorname{At}(\mathcal{L}) := \operatorname{Diff}^1_X(\mathcal{L}, \mathcal{L}) \xrightarrow{\zeta} T\mathcal{B}_g(r) \longrightarrow 0, \quad (8.2)$$

where ζ is the symbol map. We note that (8.2) coincides with the Atiyah exact sequence for the line bundle \mathcal{L} in (8.1) (see (4.4)). Let

$$0 \longrightarrow T^* \mathcal{B}_g(r) \longrightarrow \operatorname{At}(\mathcal{L})^* \xrightarrow{\eta} \mathcal{O}_{\mathcal{B}_g(r)} \longrightarrow 0$$
(8.3)

be the dual of the sequence in (8.2). Define

$$\mathcal{C}(\mathcal{L}) := \eta^{-1}(\operatorname{image}(\mathbf{1})) \subset \operatorname{At}(\mathcal{L})^*, \qquad (8.4)$$

where η is the projection in (8.3), and **1** is the section of $\mathcal{O}_{\mathcal{B}_g(r)}$ given by the constant function 1 on $\mathcal{B}_q(r)$. Let

$$\rho : \mathcal{C}(\mathcal{L}) \longrightarrow \mathcal{B}_g(r) \tag{8.5}$$

be the restriction of the natural projection $\operatorname{At}(\mathcal{L})^* \longrightarrow \mathcal{B}_g(r)$. From (8.3) it follows that $\mathcal{C}(\mathcal{L})$ is an algebraic torsor over $\mathcal{B}_g(r)$ for the holomorphic cotangent bundle $T^*\mathcal{B}_g(r)$. Giving a holomorphic section of $\mathcal{C}(\mathcal{L})$ over an open subset $U \subset \mathcal{B}_g(r)$ is equivalent to giving a holomorphic connection on the line bundle $\mathcal{L}|_U \longrightarrow U$.

Recall from Theorem 5.3 the algebraic torsor $\gamma : \mathcal{H}_g(r) \longrightarrow \mathcal{B}_g(r)$ for the holomorphic cotangent bundle $T^*\mathcal{B}_g(r)$. Using Lemma 7.1 it can be deduced that the two torsors $\mathcal{C}(\mathcal{L})$ (constructed in (8.5)) and $\mathcal{H}_g(r)$, over $\mathcal{B}_g(r)$ for the holomorphic cotangent bundle $T^*\mathcal{B}_g(r)$, are naturally identified when restricted to the open subset $\mathcal{B}_g(r) \setminus \Theta \subset \mathcal{B}_g(r)$, where Θ is the divisor in (5.8). To explain this, define

$$\mathcal{H}_{g}^{0}(r) := \gamma^{-1}(\mathcal{B}_{g}(r) \setminus \Theta) \subset \mathcal{H}_{g}(r)$$
(8.6)

$$\mathcal{C}(\mathcal{L})^0 := \rho^{-1}(\mathcal{B}_g(r) \setminus \Theta) \subset \mathcal{C}(\mathcal{L})$$
(8.7)

where γ and ρ are the projections in (5.7) and (8.5) respectively. Since the restriction of \mathcal{L} to $\mathcal{B}_g(r) \setminus \Theta \subset \mathcal{B}_g(r)$ is the trivial bundle $\mathcal{O}_{\mathcal{B}_g(r)\setminus\Theta}$, there is a natural integrable algebraic connection ∇^0 on the restriction of \mathcal{L} to $\mathcal{B}_g(r) \setminus \Theta$ which is given by the de Rham differential d on $\mathcal{O}_{\mathcal{B}_g(r)\setminus\Theta}$. This connection ∇^0 produces an algebraic splitting of the Atiyah exact sequence for $\mathcal{L}|_{\mathcal{B}_g(r)\setminus\Theta}$ (see (8.2)), which in turn produces an algebraic splitting, over $\mathcal{B}_g(r) \setminus \Theta$, of the short exact sequence in (8.3). Hence we get an algebraic section

$$G : \mathcal{B}_q(r) \setminus \Theta \longrightarrow \mathcal{C}(\mathcal{L})^0 \tag{8.8}$$

of the bundle $\rho : \mathcal{C}(\mathcal{L})^0 \longrightarrow \mathcal{B}_g(r) \setminus \Theta$ in (8.7). To explain the construction of G, if

$$s_0 : \mathcal{O}_{\mathcal{B}_g(r)\setminus\Theta} \longrightarrow \operatorname{At}(\mathcal{L})^* \big|_{\mathcal{B}_g(r)\setminus\Theta}$$

is the splitting homomorphism for (8.3) over $\mathcal{B}_g(r) \setminus \Theta$, then $G(z) = s_0(\mathbf{1}(z))$ for all $z \in \mathcal{B}_g(r) \setminus \Theta$, where **1** as before is the section of $\mathcal{O}_{\mathcal{B}_g(r) \setminus \Theta}$ given by the constant function 1 on $\mathcal{B}_g(r) \setminus \Theta$.

Consider $\mathcal{H}_q^0(r)$ and $\mathcal{C}(\mathcal{L})^0$ constructed in (8.6) and (8.7) respectively. We have a map

$$H : \mathcal{H}_g^0(r) \longrightarrow \mathcal{C}(\mathcal{L})^0 \tag{8.9}$$

that sends any $v \in \mathcal{H}_{g}^{0}(r)$ to

$$G(\gamma(v)) + (v - \mathbb{S}(\gamma(v))), \qquad (8.10)$$

where \mathbb{S} , G and γ are constructed in Lemma 7.1, (8.8) and (5.7) respectively; note that $v - \mathbb{S}(\gamma(v))$ in (8.10) is an element of $T^*_{\gamma(v)}\mathcal{B}_g(r)$ (recall that $\mathcal{H}_g(r)$ is a torsor for $T^*\mathcal{B}_g(r)$ by Theorem 5.3), so the sum in (8.10) is an element of $\mathcal{C}(\mathcal{L})^0$ because $\mathcal{C}(\mathcal{L})^0$ is a torsor over $\mathcal{B}_g(r) \setminus \Theta$ for $T^*(\mathcal{B}_g(r) \setminus \Theta)$ (see (8.4) and (8.7)).

It is evident that H in (8.9) is an algebraic isomorphism of torsors over $\mathcal{B}_g(r) \setminus \Theta$ for $T^*(\mathcal{B}_g(r) \setminus \Theta)$.

Theorem 8.1. The algebraic isomorphism H in (8.9), of torsors over $\mathcal{B}_g(r) \setminus \Theta$ for $T^*(\mathcal{B}_g(r) \setminus \Theta)$, extends to an algebraic isomorphism of $T^*\mathcal{B}_g(r)$ -torsors

$$\mathcal{H} : \mathcal{H}_g(r) \longrightarrow \mathcal{C}(\mathcal{L})$$

over entire $\mathcal{B}_g(r)$.

Proof. Consider the submersion β in (5.6). The kernel of the differential of β

$$d\beta : T\mathcal{B}_g(r) \longrightarrow \beta^* T\mathcal{M}_g^{\theta}$$

will be denoted by T_{β} ; in other words, T_{β} is the relative tangent bundle for the projection β . So we have the short exact sequence of vector bundles

$$0 \longrightarrow \beta^* T^* \mathcal{M}_g^{\theta} \xrightarrow{(d\beta)^*} T^* \mathcal{B}_g(r) \longrightarrow T_{\beta}^* := (T_{\beta})^* \longrightarrow 0$$

over $\mathcal{B}_g(r)$. Let $\mathcal{W} \longrightarrow \mathcal{B}_g(r)$ be an algebraic torsor over $\mathcal{B}_g(r)$ for $T^*\mathcal{B}_g(r)$. Then

$$\mathcal{W}/(\beta^* T^* \mathcal{M}_g^{\theta}) \longrightarrow \mathcal{B}_g(r)$$
 (8.11)

is an algebraic torsor over $\mathcal{B}_q(r)$ for the vector bundle T^*_{β} .

Substitute the two $T^*\mathcal{B}_g(r)$ -torsors $\mathcal{H}_g(r)$ and $\mathcal{C}(\mathcal{L})$ in place of the above $T^*\mathcal{B}_g(r)$ -torsor \mathcal{W} . The construction in (8.11) produces two T^*_{β} -torsors over $\mathcal{B}_g(r)$ from $\mathcal{H}_g(r)$ and $\mathcal{C}(\mathcal{L})$; these two T^*_{β} -torsors will be denoted by $\widehat{\mathcal{H}}_g(r)$ and $\widehat{\mathcal{C}}(\mathcal{L})$ respectively. The restrictions of $\widehat{\mathcal{H}}_g(r)$ and $\widehat{\mathcal{C}}(\mathcal{L})$ to the Zariski open subset

$$\mathcal{B}_g(r) \setminus \Theta \subset \mathcal{B}_g(r)$$

will be denoted by $\widehat{\mathcal{H}}_{g}^{0}(r)$ and $\widehat{\mathcal{C}}(\mathcal{L})^{0}$ respectively.

The isomorphism H in (8.9) produces an algebraic isomorphism of T^*_{β} -torsors over $\mathcal{B}_g(r) \setminus \Theta$

$$\widehat{H} : \widehat{\mathcal{H}}_g^0(r) \longrightarrow \widehat{\mathcal{C}}(\mathcal{L})^0.$$
 (8.12)

For each point $z \in \mathcal{M}_{g}^{\theta}$, consider the restriction of the isomorphism \widehat{H} in (8.12) to the complement $\beta^{-1}(z) \setminus (\beta^{-1}(z) \cap \Theta)$, where β is the projection in (5.6). This restriction coincides with the isomorphism constructed in [BH2, Lemma 3.1]. Therefore, from [BH2, Corollary 4.5] we know that \widehat{H} in (8.12) extends to an algebraic isomorphism of T_{β}^{*} -torsors

$$\widehat{H}' : \widehat{\mathcal{H}}_g(r) \longrightarrow \widehat{\mathcal{C}}(\mathcal{L}).$$
 (8.13)

over entire $\mathcal{B}_q(r)$. Let

$$\mathbb{G} := \{ (y, \widehat{H}'(y)) \mid y \in \widehat{\mathcal{H}}_g(r) \} \subset \widehat{\mathcal{H}}_g(r) \times_{\mathcal{B}_g(r)} \widehat{\mathcal{C}}(\mathcal{L})$$
(8.14)

be the graph of the map \widehat{H}' in (8.13).

Consider the natural quotient map

 $\mathcal{H}_g(r) \times_{\mathcal{B}_g(r)} \mathcal{C}(\mathcal{L}) \xrightarrow{\mathbf{q}_1} (\mathcal{H}_g(r)/(\beta^* T^* \mathcal{M}_g^{\theta})) \times_{\mathcal{B}_g(r)} (\mathcal{C}(\mathcal{L})/(\beta^* T^* \mathcal{M}_g^{\theta})) = \widehat{\mathcal{H}}_g(r) \times_{\mathcal{B}_g(r)} \widehat{\mathcal{C}}(\mathcal{L}).$ The inverse image

$$\mathcal{G}_1 := \mathbf{q}_1^{-1}(\mathbb{G}) \subset \mathcal{H}_g(r) \times_{\mathcal{B}_g(r)} \mathcal{C}(\mathcal{L}),$$

where \mathbb{G} is constructed in (8.14), is an algebraic torsor over $\mathcal{B}_g(r)$ for the vector bundle $(\beta^*T^*\mathcal{M}_g^\theta)^{\oplus 2}$. Let

$$\mathcal{G} := \mathcal{G}_1/(\beta^* T^* \mathcal{M}_g^\theta) \xrightarrow{\mathbf{q}} \mathcal{B}_g(r)$$
(8.15)

be the quotient for the diagonal action of $\beta^* T^* \mathcal{M}_g^{\theta}$; so for any $z \in \mathcal{B}_g(r)$, two elements (x, y) and (x', y') of the fiber $(\mathcal{G}_1)_z$ give the same element of the fiber \mathcal{G}_z if and only if there is an element $v \in (\beta^* T^* \mathcal{M}_g^{\theta})_z = T^*_{\beta(z)} \mathcal{M}_g^{\theta}$ such that x' = x + v and y' = y + v. Therefore, we have a map

$$\nu : \mathcal{H}_g(r) \times_{\mathcal{B}_g(r)} \mathcal{G} \longrightarrow \mathcal{C}(\mathcal{L})$$

that sends any (h, (x, y)), where $h \in \mathcal{H}_g(r)_z$, $(x, y) \in \mathcal{G}_z$ and $z \in \mathcal{B}_g(r)$, to

$$y + (h - x) \in \mathcal{C}(\mathcal{L})_{z}$$

(note that $h - x \in T_z^* \mathcal{B}_g(r)$). Using this map ν , the space of holomorphic sections of \mathcal{G} over any open subset $U \subset \mathcal{B}_g(r)$ is identified with the space of holomorphic isomorphisms $\mathcal{H}_g(r)|_U \longrightarrow \mathcal{C}(\mathcal{L})|_U$, of the torsor over U for T^*U , that induce the isomorphism $\widehat{H}'|_U$ in (8.13) of $T_\beta^*|_U$ -torsors.

We note that \mathcal{G} is an algebraic torsor over $\mathcal{B}_g(r)$ for the vector bundle $\beta^* T^* \mathcal{M}_g^{\theta}$, where β is the projection in (5.6), as follows: For any $z \in \mathcal{B}_g(r)$, take $(x, y) \in \mathcal{G}_z$ and $v \in (\beta^* T^* \mathcal{M}_g^{\theta})_z = T^*_{\beta(z)} \mathcal{M}_g^{\theta}$; then we have

$$(x, y) + v = (x + v, y - v).$$

The moduli space $\mathcal{B}_q(r)$ has an algebraic involution

$$\mathcal{I}_B : \mathcal{B}_g(r) \longrightarrow \mathcal{B}_g(r), \quad (X, K_X^{1/2}, F) \longmapsto (X, K_X^{1/2}, F^*).$$
(8.16)

This involution \mathcal{I}_B preserves the divisor Θ in (5.8) (see (5.9)). Hence the action $\mathbb{Z}/2\mathbb{Z}$ on $\mathcal{B}_g(r)$ given by \mathcal{I}_B lifts to the line bundle \mathcal{L} in (8.1). Consequently, the involution \mathcal{I}_B lifts to an involution

$$\mathcal{I}_C : \mathcal{C}(\mathcal{L}) \longrightarrow \mathcal{C}(\mathcal{L})$$
 (8.17)

of $\mathcal{C}(\mathcal{L})$.

We will now describe the relationship between \mathcal{I}_C and the $T^*\mathcal{B}_g(r)$ -torsor structure of $\mathcal{C}(\mathcal{L})$. Take any $y \in \mathcal{B}_g(r), z \in \mathcal{C}(\mathcal{L})_y$ and $w \in T^*_y\mathcal{B}_g(r)$. Then we have

$$\mathcal{I}_C(z+w) = \mathcal{I}_C(z) - (d\mathcal{I}_B)^*_{\mathcal{I}_B(y)}(w), \qquad (8.18)$$

where $(d\mathcal{I}_B)^*_{\mathcal{I}_B(y)} : T^*_y \mathcal{B}_g(r) \longrightarrow T^*_{\mathcal{I}_B(y)} \mathcal{B}_g(r)$ is the dual of the differential $(d\mathcal{I}_B)_{\mathcal{I}_B(y)} : T_{\mathcal{I}_B(y)} \mathcal{B}_g(r) \longrightarrow T_y \mathcal{B}_g(r)$ of the map \mathcal{I}_B at the point $\mathcal{I}_B(y)$.

Since $\bigwedge^2 J^1(K_X^{-1/2}) = K_X \otimes (K_X^{-1/2})^{\otimes 2} = \mathcal{O}_X$ (see (2.6)), it follows that $J^1(K_X^{-1/2}) = J^1(K_X^{-1/2})^*$. Let *F* be a stable vector bundle on *X* of rank *r* and degree zero. Since *F* admits a holomorphic connection (see the proof of Proposition 2.4), from Proposition 2.5 we conclude that

$$J^{1}(F^{*} \otimes K_{X}^{-1/2}) = F^{*} \otimes J^{1}(K_{X}^{-1/2}) = F^{*} \otimes J^{1}(K_{X}^{-1/2})^{*} = J^{1}(F \otimes K_{X}^{-1/2})^{*}$$

Fixing an isomorphism of $J^1(F^* \otimes K_X^{-1/2})$ with $J^1(F \otimes K_X^{-1/2})^*$ we conclude that any holomorphic connection on $J^1(F \otimes K_X^{-1/2})$ produces a holomorphic connection on $J^1(F \otimes K_X^{-1/2})^* = J^1(F^* \otimes K_X^{-1/2})$. It is straightforward to check that this produces a bijection

$$M_F : \mathcal{D}(F) \longrightarrow \mathcal{D}(F^*),$$

where $\mathcal{D}(F)$ is constructed in (2.29). This map M_F does not depend on the choice of the isomorphism of $J^1(F^* \otimes K_X^{-1/2})$ with $J^1(F \otimes K_X^{-1/2})^*$; this is because in the construction of $\mathcal{D}(F)$ quotienting by $\operatorname{Aut}(F \otimes K_X^{-1/2})$ was executed. Now we have an algebraic involution

$$\mathcal{I}_H : \mathcal{H}_g(r) \longrightarrow \mathcal{H}_g(r), \quad (X, K_X^{1/2}, F, D) \longmapsto (X, K_X^{1/2}, F^*, M_F(D)), \quad (8.19)$$

where M_F is the map constructed above. It is evident that

$$\gamma \circ \mathcal{I}_H = \mathcal{I}_B \circ \gamma \,,$$

where γ and \mathcal{I}_B are the maps in (5.7) and (8.16) respectively.

We will describe the relationship between \mathcal{I}_H and the $T^*\mathcal{B}_g(r)$ -torsor structure of $\mathcal{H}_g(r)$. Take any $y \in \mathcal{B}_g(r), z \in \mathcal{H}_g(r)_y$ and $w \in T^*_y\mathcal{B}_g(r)$. Then we have

$$\mathcal{I}_H(z+w) = \mathcal{I}_H(z) - (d\mathcal{I}_B)^*_{\mathcal{I}_B(y)}(w), \qquad (8.20)$$

where $(d\mathcal{I}_B)^*_{\mathcal{I}_B(y)}$ is the homomorphism in (8.18).

We note that the projection β in (5.6) satisfies the equation

$$\beta \circ \mathcal{I}_B = \beta$$

In view of this, from (8.18) and (8.20) we conclude the following:

(1) The involution \mathcal{I}_B lifts to an involution

$$\mathcal{I}_G : \mathcal{G} \longrightarrow \mathcal{G} \tag{8.21}$$

of \mathcal{G} constructed in (8.15). For any $z \in \mathcal{B}_g(r)$, take $(x, y) \in \mathcal{G}_z$; the map \mathcal{I}_G sends (x, y) to $(\mathcal{I}_H(x), \mathcal{I}_C(y))$, where \mathcal{I}_H and \mathcal{I}_C are the involutions in (8.19) and (8.17) respectively.

(2) For any $v \in (\beta^* T^* \mathcal{M}_g^\theta)_z = T^*_{\beta(z)} \mathcal{M}_g^\theta$,

$$\mathcal{I}_G((x,y)+v) = \mathcal{I}_G((x,y))+v;$$
 (8.22)

recall that \mathcal{G} is a torsor for $\beta^* T^* \mathcal{M}_q^{\theta}$.

 \mathcal{G}^t

Take any point

Let

$$t := (X, K_X^{1/2}) \in \mathcal{M}_g^{\theta}.$$

:= $\mathcal{G}|_{\beta^{-1}(t)} \xrightarrow{\mathbf{q}^t} \mathcal{B}^t := \beta^{-1}(t)$ (8.23)

34

be the restriction, where \mathbf{q}^t is the restriction of the projection \mathbf{q} in (8.15), and β is the projection in (5.6). So \mathcal{G}^t is an algebraic torsor over \mathcal{B}^t for the trivial vector bundle

$$\mathcal{V}^t := \mathcal{B}^t \times T_t^* \mathcal{M}_g^\theta \longrightarrow \mathcal{B}^t$$

over \mathcal{B}^t with fiber $T^*_t \mathcal{M}^{\theta}_g$; this is because \mathcal{G} is a torsor for $\beta^* T^* \mathcal{M}^{\theta}_g$.

Let

$$\mathcal{I}_B^t := \mathcal{I}_B \big|_{\mathcal{B}^t} : \mathcal{B}^t \longrightarrow \mathcal{B}^t$$

be the restriction of \mathcal{I}_B in (8.16) to the subvariety \mathcal{B}^t in (8.23).

We note that the isomorphism classes of algebraic \mathcal{V}^t -torsors over the variety \mathcal{B}^t in (8.23) are parametrized by

$$H^1(\mathcal{B}^t, \mathcal{V}^t) = H^1(\mathcal{B}^t, \mathcal{O}_{\mathcal{B}^t}) \otimes T^*_t \mathcal{M}^{\theta}_g.$$

Let

$$\mathbf{c} \in H^1(\mathcal{B}^t, \mathcal{O}_{\mathcal{B}^t}) \otimes T^*_t \mathcal{M}^{\theta}_g$$

$$(8.24)$$

be the class of the \mathcal{V}^t -torsor \mathcal{G}^t in (8.23). From (8.22) it follows immediately that

$$(\mathcal{I}_B^t)^* \mathbf{c} = \mathbf{c} \,. \tag{8.25}$$

Consider the determinant map

$$\widetilde{\delta} : \mathcal{B}^t \longrightarrow \operatorname{Pic}^0(X), \quad F \longmapsto \bigwedge^r F.$$

The corresponding homomorphism

$$\widetilde{\delta}^* : H^1(\operatorname{Pic}^0(X), \mathcal{O}_{\operatorname{Pic}^0(X)}) \longrightarrow H^1(\mathcal{B}^t, \mathcal{O}_{\mathcal{B}^t})$$
(8.26)

is an isomorphism. Indeed, $\operatorname{Pic}(\mathcal{B}^t) = \operatorname{Pic}(\operatorname{Pic}^0(X)) \oplus \mathbb{Z}$ [DN, p. 57, Theorem D] when $r \geq 2$ and $(g, r) \neq (2, 2)$ (when g = 2 = r, the quotient $\operatorname{Pic}(\mathcal{B}^t)/\operatorname{Pic}(\operatorname{Pic}^0(X))$) is a finite group), so using the exact sequence of cohomologies

$$H^1(Y, 2\pi\sqrt{-1} \cdot \mathbb{Z}) \longrightarrow H^1(Y, \mathcal{O}_Y) \longrightarrow H^1(Y, \mathcal{O}_Y^*) \longrightarrow H^2(Y, 2\pi\sqrt{-1} \cdot \mathbb{Z})$$

associated to the exponential sequence

$$0 \longrightarrow 2\pi\sqrt{-1} \cdot \mathbb{Z} \longrightarrow \mathcal{O}_Y \xrightarrow{\exp} \mathcal{O}_Y^* \longrightarrow 0$$

on a complex variety Y we conclude that $\tilde{\delta}^*$ in (8.26) is an isomorphism. We also note that the involution

$$\operatorname{Pic}^{0}(X) \longrightarrow \operatorname{Pic}^{0}(X), \quad \xi \longmapsto \xi^{*}$$

acts on $H^1(\operatorname{Pic}^0(X), \mathcal{O}_{\operatorname{Pic}^0(X)})$ as multiplication by -1. In other words, no nonzero element of $H^1(\operatorname{Pic}^0(X), \mathcal{O}_{\operatorname{Pic}^0(X)})$ is fixed by this involution. In view of these, from (8.25) we conclude that

 $\mathbf{c} = 0.$

So from (8.24) we conclude that the \mathcal{V}^t -torsor \mathcal{G}^t in (8.23) is the trivial \mathcal{V}^t -torsor

$$\mathcal{V}^t = \mathcal{B}^t \times T^*_t \mathcal{M}^{\theta}_g \longrightarrow \mathcal{B}^t.$$

Restrict H (constructed in (8.9)) to $\mathcal{B}^t \subset \mathcal{B}_g(r)$; denote this restriction by H^t . From the above isomorphism of \mathcal{G}^t with the trivial \mathcal{V}^t -torsor \mathcal{V}^t it follows that H^t is a meromorphic function on \mathcal{B}^t with values in the vector space $T^*_t \mathcal{M}^\theta_g$ (recall that $\mathcal{V}^t := \mathcal{B}^t \times T^*_t \mathcal{M}^\theta_g$). This meromorphic function is evidently regular on the complement $\mathcal{B}^t \setminus (\mathcal{B}^t \cap \Theta)$, where Θ is constructed in (5.8). From the construction of H it is straightforward to deduce that H^t has a pole of order at most one on the divisor $\mathcal{B}^t \cap \Theta \subset \mathcal{B}^t$. On the other hand, we know that

$$H^0(\mathcal{B}^t, \mathcal{O}_{\mathcal{B}^t}(\mathcal{B}^t \cap \Theta)) = H^0(\mathcal{B}^t, \mathcal{O}_{\mathcal{B}^t}) = \mathbb{C}$$

[BNR, p. 169, Theorem 2]. Consequently, the section H^t over $\mathcal{B}^t \setminus (\mathcal{B}^t \cap \Theta)$ extends to entire \mathcal{B}^t as a regular section. From this it follows immediately that the isomorphism H of torsors over $\mathcal{B}_g(r) \setminus \Theta$ for $T^*(\mathcal{B}_g(r) \setminus \Theta)$, extends to an algebraic isomorphism of $T^*\mathcal{B}_g(r)$ -torsors $\mathcal{H}_g(r) \longrightarrow \mathcal{C}(\mathcal{L})$ over entire $\mathcal{B}_g(r)$. \Box

8.2. A holomorphic symplectic form. Recall the holomorphic line bundle $\mathcal{L} \longrightarrow \mathcal{B}_g(r)$ in (8.1). The space $\mathcal{C}(\mathcal{L})$ in (8.5) has a canonical holomorphic symplectic structure. We will briefly recall the construction of this symplectic form on $\mathcal{C}(\mathcal{L})$.

From the construction of $\mathcal{C}(\mathcal{L})$ in (8.4) it follows immediately that there is a tautological holomorphic splitting

$$\rho^* \operatorname{At}(\mathcal{L}) = \rho^* \mathcal{O}_{\mathcal{B}_g(r)} \oplus \rho^* T \mathcal{B}_g(r) = \mathcal{O}_{\mathcal{C}(\mathcal{L})} \oplus \rho^* T \mathcal{B}_g(r),$$

where ρ is constructed in (8.5). This decomposition of $\rho^* \operatorname{At}(\mathcal{L})$ gives a holomorphic projection

$$f_0: \rho^* \operatorname{At}(\mathcal{L}) \longrightarrow \mathcal{O}_{\mathcal{C}(\mathcal{L})}.$$
 (8.27)

Let

$$0 \longrightarrow \mathcal{O}_{\mathcal{C}(\mathcal{L})} \longrightarrow \operatorname{At}(\rho^* \mathcal{L}) \xrightarrow{\zeta'} T\operatorname{Conn}(\mathcal{L}) \longrightarrow 0$$
(8.28)

be the Atiyah exact sequence for $\rho^* \mathcal{L}$ (see (4.4)). We also have a tautological projection

 $h_0 : \operatorname{At}(\rho^* \mathcal{L}) \longrightarrow \rho^* \operatorname{At}(\mathcal{L})$

such that the diagram

$$\begin{array}{ccc} \operatorname{At}(\rho^{*}\mathcal{L}) & \stackrel{h_{0}}{\longrightarrow} & \rho^{*}\operatorname{At}(\mathcal{L}) \\ & & & \downarrow \zeta' & & \downarrow \rho^{*}\zeta \\ T\operatorname{Conn}(\mathcal{L}) & \stackrel{d\rho}{\longrightarrow} & \rho^{*}T\mathcal{B}_{g}(r) \end{array}$$

is commutative, where ζ and ζ' are the projections in (8.2) and (8.28) respectively, and $d\rho$ is the differential of the projection ρ in (8.5); see [BHS, (3.9)] [BH1, Section 3] for the construction of h_0 . The composition of homomorphisms

$$f_0 \circ h_0 : \operatorname{At}(\rho^* \mathcal{L}) \longrightarrow \mathcal{O}_{\mathcal{C}(\mathcal{L})},$$
 (8.29)

where f_0 is constructed in (8.27), gives a holomorphic splitting of the Atiyah exact sequence in (8.28). Hence $f_0 \circ h_0$ defines a holomorphic connection on $\rho^* \mathcal{L}$; see [BHS, Proposition 3.3].

The curvature $\Omega_{\mathcal{L}}$ of the holomorphic connection $f_0 \circ h_0$ on $\rho^* \mathcal{L}$ in (8.29) is a closed algebraic 2-form on Conn(\mathcal{L}). This algebraic 2-form $\Omega_{\mathcal{L}}$ is symplectic. (See [BH1, Section 3], [BHS].)

Recall the holomorphic $T^*\mathcal{B}_g(r)$ -torsor structure of $\operatorname{Conn}(\mathcal{L})$. The above symplectic form $\Omega_{\mathcal{L}}$ on $\operatorname{Conn}(\mathcal{L})$ is compatible with the $T^*\mathcal{B}_g(r)$ -torsor structure. This means that for any locally defined holomorphic section of the projection ρ in (8.5)

$$\mathcal{B}_g(r) \supset U \xrightarrow{s} \operatorname{Conn}(\mathcal{L})$$

and any holomorphic 1-form $\omega \in H^0(U, T^*U)$, we have

$$s^*\Omega_{\mathcal{L}} + d\omega = (s+\omega)^*\Omega_{\mathcal{L}}; \qquad (8.30)$$

note that $y \mapsto s(y) + \omega(y)$ is a holomorphic section, over U, of the projection ρ in (8.5).

Corollary 8.2.

- (1) The moduli space $\mathcal{H}_g(r)$ in (5.7) has a canonical algebraic symplectic structure $\Omega_{\mathcal{H}_g(r)}$.
- (2) The symplectic form $\Omega_{\mathcal{H}_g(r)}$ on $\mathcal{H}_g(r)$ is compatible with the $T^*\mathcal{B}_g(r)$ -torsor structure of $\mathcal{H}_g(r)$ obtained in Theorem 5.3.
- (3) There is a holomorphic line bundle \mathbf{L} on $\mathcal{H}_g(r)$ and a holomorphic connection $\nabla^{\mathbf{L}}$ on \mathbf{L} such that the curvature of $\nabla^{\mathbf{L}}$ is the symplectic form $\Omega_{\mathcal{H}_g(r)}$.

Proof. Using the isomorphism \mathcal{H} in Theorem 8.1, the above algebraic symplectic form $\Omega_{\mathcal{L}}$ on Conn(\mathcal{L}) (see (8.30)) produces an algebraic symplectic form

$$\Omega_{\mathcal{H}_q(r)} := \mathcal{H}^* \Omega_{\mathcal{L}}$$

on $\mathcal{H}_g(r)$.

The compatibility condition in the statement (2) says that for any locally defined holomorphic section of the projection γ in (5.7)

$$\mathcal{B}_q(r) \supset U \xrightarrow{s} \mathcal{H}_q(r)$$

and any holomorphic 1-form $\omega \in H^0(U, T^*U)$, the equality

$$s^* \Omega_{\mathcal{H}_q(r)} + d\omega = (s + \omega)^* \Omega_{\mathcal{H}_q(r)}$$
(8.31)

holds; note that $y \mapsto s(y) + \omega(y)$ is a holomorphic section, over U, of the projection γ in (5.7). Now, (8.31) follows immediately from (8.30).

We recall that $\Omega_{\mathcal{L}}$ is the curvature of the holomorphic connection $f_0 \circ h_0$ (see (8.29)) on the holomorphic line bundle $\rho^* \mathcal{L}$. Therefore, $\Omega_{\mathcal{H}_g(r)}$ is the curvature of the holomorphic connection

$$\nabla^{\mathbf{L}} := \mathcal{H}^*(f_0 \circ h_0)$$

on the holomorphic line bundle $\mathbf{L} := \mathcal{H}^* \rho^* \mathcal{L}$.

9. ISOMONODROMY AND SYMPLECTIC FORM

9.1. **Projective structure on a Riemann surface.** Let Y be a compact connected Riemann surface. A holomorphic coordinate chart on Y is a pair of the form (U, φ) , where $U \subset Y$ is an open subset and $\varphi : U \longrightarrow \mathbb{CP}^1$ is a holomorphic embedding. A holomorphic coordinate atlas on Y is a collection of holomorphic coordinate charts $\{(U_i, \varphi_i)\}_{i \in J}$ such that $Y = \bigcup_{i \in J} U_i$. A projective structure on Y is given by a holomorphic coordinate atlas $\{(U_i, \varphi_i)\}_{i \in J}$ satisfying the following condition: For every $i, j \in J \times J$ with $U_i \cap U_j \neq \emptyset$, and every connected component $V_c \subset U_i \cap U_j$, there is a $\tau_{j,i}^c \in PSL(2, \mathbb{C})$ such that the map $(\varphi_j \circ \varphi_i^{-1})|_{\varphi_i(V_c)}$ is the restriction, to $\varphi_i(V_c)$, of the automorphism of \mathbb{CP}^1 given by $\tau_{i,i}$. Recall that $PSL(2, \mathbb{C}) = Aut(\mathbb{CP}^1)$.

Two holomorphic coordinate atlases $\{(U_i, \varphi_i)\}_{i \in J}$ and $\{(U_i, \varphi_i)\}_{i \in J'}$ satisfying the above condition are called *equivalent* if their union $\{(U_i, \varphi_i)\}_{i \in J \cup J'}$ also satisfies the above condition. A *projective structure* on Y is an equivalence class of holomorphic coordinate atlases satisfying the above condition.

Giving a projective structure on Y is equivalent to giving a holomorphic Cartan geometry on Y for the pair of groups (PSL(2, \mathbb{C}), B), where $B \subset PSL(2, \mathbb{C})$ is the Borel subgroup

$$B := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{PSL}(2, \mathbb{C}) \mid c = 0 \right\};$$

see [Sh] for Cartan geometry. Let $\widetilde{B} \subset SL(2,\mathbb{C})$ be the Borel subgroup that projects to B defined above. Giving a holomorphic Cartan geometry on Y for the pair of groups $(SL(2,\mathbb{C}), B)$ is equivalent to giving a projective structure on Y together with a theta characteristic on Y [Gu].

For $g \geq 2$, let

$$\delta : \mathcal{P}_g \longrightarrow \mathcal{M}_g^\theta \tag{9.1}$$

be the moduli space of triples of the form $(X, K_X^{1/2}, P)$, where

- X is a compact connected Riemann surface of genus g,
- K_X^{1/2} is a theta characteristic on X, and
 P is a projective structure on X.

The map δ in (9.1) sends any $(X, K_X^{1/2}, P)$ to $(X, K_X^{1/2})$.

Proposition 9.1. The moduli space $\mathcal{H}_g(r)$ in (5.7) admits a natural projection

$$f : \mathcal{H}_g(r) \longrightarrow \mathcal{P}_g,$$

where \mathcal{P}_q is defined in (9.1).

Proof. Take any
$$(X, K_X^{1/2}, F, D) \in \mathcal{H}_g(r)$$
. Let

$$s \in \frac{H^0\left(3\Delta, \left((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\Delta)\right)\Big|_{3\Delta}\right)}{H^0(X, \operatorname{ad}(F) \otimes K_X^{\otimes 2})}$$
(9.2)

be the element corresponding to it given by Corollary 6.5. Restricting s to $2\Delta \subset 3\Delta$, we get a section

$$s_{1} \in H^{0}\left(2\boldsymbol{\Delta}, \left((F \otimes K_{X}^{3/2}) \boxtimes (F^{*} \otimes K_{X}^{3/2}) \otimes \mathcal{O}_{X \times X}(3\boldsymbol{\Delta})\right)\big|_{2\boldsymbol{\Delta}}\right)$$
$$= H^{0}\left(2\boldsymbol{\Delta}, \left((F \boxtimes F^{*}) \otimes ((K_{X}^{3/2} \boxtimes K_{X}^{3/2}) \otimes \mathcal{O}_{X \times X}(3\boldsymbol{\Delta}))\right)\big|_{2\boldsymbol{\Delta}}\right).$$

Since $((K_X^{3/2} \boxtimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\Delta))|_{2\Delta}$ is canonically trivialized [BR, p. 688, Theorem 2.2] (this was noted in (7.3)), there is a unique section

$$s'_1 \in H^0(2\mathbf{\Delta}, (F \boxtimes F^*)|_{2\mathbf{\Delta}})$$

$$(9.3)$$

such that $s_1 = s'_1 \otimes \mathbf{t}$, where \mathbf{t} is the section of $((K_X^{3/2} \boxtimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\Delta))|_{2\Delta}$ that trivializes it. On the other hand, $s_1|_{\Delta} = \mathrm{Id}_F$ (see Corollary 6.5), which implies that $s'_1|_{\mathbf{\Delta}} = \mathrm{Id}_F$, because the restriction of t to $\mathbf{\Delta}$ is the constant function 1 on X using the identification of Δ with X given by $x \mapsto (x, x)$. Consequently, s'_1 defines a holomorphic connection on F. This holomorphic connection in turn provides an extension of the section s'_1 to a section

$$s_2 \in H^0\left(3\Delta, (F \boxtimes F^*)\Big|_{3\Delta}\right);$$

we note that using parallel translations, for the integrable connection on F defined by s'_1 , we get a holomorphic isomorphism between q_1^*F and q_2^*F over an analytic neighborhood of $\Delta \subset X \times X$ (recall that q_i is the projection of $X \times X$ to the *i*-th factor). Now, invoking the isomorphism

$$s_2: (q_1^*F)\big|_{3\mathbf{\Delta}} \longrightarrow (q_2^*F)\big|_{3\mathbf{\Delta}},$$

the element s in (9.2) becomes

$$s \in \frac{H^0\left(3\boldsymbol{\Delta}, \left((F \otimes F^* \otimes K_X^{3/2}) \boxtimes (K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\boldsymbol{\Delta})\right)\big|_{3\boldsymbol{\Delta}}\right)}{H^0(X, \operatorname{ad}(F) \otimes K_X^{\otimes 2})}.$$

Composing with the trace homomorphism $F \otimes F^* = \operatorname{End}(F) \longrightarrow \mathcal{O}_X$ defined by $A \mapsto \frac{1}{r}\operatorname{trace}(A)$, the above element s gives a section

$$\phi_2 \in H^0\left(3\mathbf{\Delta}, \left((K_X^{3/2}) \boxtimes K_X^{3/2}\right) \otimes \mathcal{O}_{X \times X}(3\mathbf{\Delta})\right)\Big|_{3\mathbf{\Delta}}\right).$$

This section ϕ_2 defines a projective structure on X [BR, p. 688, Theorem 2.2]. Therefore, we have a map

$$f: \mathcal{H}_g(r) \longrightarrow \mathcal{P}_g$$

that sends any $(X, K_X^{1/2}, F, D) \in \mathcal{H}_g(r)$ to $(X, K_X^{1/2}, \phi_2) \in \mathcal{P}_g$, where ϕ_2 is constructed above from D.

9.2. **Isomonodromy.** For any compact Riemann surface X of genus g, with $g \ge 2$, let $\mathcal{D}_X(r)$ denote the moduli space of pairs of the form (F, D), where F is a stable vector bundle on X of rank r and degree zero, and D is a holomorphic connection on F.

Proposition 9.2. For the map f in Proposition 9.1, the fiber over any $(X, K_X^{1/2}, P) \in \mathcal{P}_g$ is canonically identified with the moduli space $\mathcal{D}_X(r)$.

Proof. Take any $(X, K_X^{1/2}, F, D) \in \mathcal{H}_g(r)$ in the fiber of f over $(X, K_X^{1/2}, P) \in \mathcal{P}_g$. As seen before, s'_1 in (9.3) defines a holomorphic connection on F. So we get a map from the fiber of f over $(X, K_X^{1/2}, P)$ to $\mathcal{D}_X(r)$ that sends any $(X, K_X^{1/2}, F, D)$ to (F, s'_1) .

Conversely, take any $(F, D) \in \mathcal{D}_X(r)$. Then D gives a section

$$\widehat{s}_D \in H^0(3\mathbf{\Delta}, (F \boxtimes F^*)|_{3\mathbf{\Delta}}).$$

On the other hand, the projective structure P gives a section

$$P' \in H^0(3\Delta, ((K_X^{3/2} \boxtimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\Delta))|_{3\Delta})$$

[BR, p. 688, Theorem 2.2]. So we have

$$\widehat{s}_D \otimes P' \in H^0\left(3\mathbf{\Delta}, \left((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\mathbf{\Delta})\right)\Big|_{3\mathbf{\Delta}}\right).$$

Consider the image of $\hat{s}_D \otimes P'$

$$\widetilde{\hat{s}_D \otimes P'} \in \frac{H^0\left(3\boldsymbol{\Delta}, \left((F \otimes K_X^{3/2}) \boxtimes (F^* \otimes K_X^{3/2}) \otimes \mathcal{O}_{X \times X}(3\boldsymbol{\Delta})\right)\big|_{3\boldsymbol{\Delta}}\right)}{H^0(X, \operatorname{ad}(F) \otimes K_X^{\otimes 2})}$$

in the quotient space. Now $(X, K_X^{1/2}, F, \widehat{s_D \otimes P'})$ gives an element of $\mathcal{H}_g(r)$ in the fiber of f over $(X, K_X^{1/2}, P) \in \mathcal{P}_g$ (see Corollary 6.5).

In view of Proposition 9.2, the isomonodromy condition for integrable holomorphic connections defines a holomorphic foliation

$$\mathcal{F} \subset T\mathcal{H}_a(r) \tag{9.4}$$

which gives the following decomposition:

$$T\mathcal{H}_g(r) = \mathcal{F} \oplus \operatorname{kernel}(df),$$
(9.5)

where $df : T\mathcal{H}_g(r) \longrightarrow f^*T\mathcal{P}_g$ is the differential of the projection f in Proposition 9.1. Consequently, the differential df identifies \mathcal{F} with $f^*T\mathcal{P}_g$.

We note that $\mathcal{D}_X(r)$ has a natural holomorphic symplectic form [Go], [AB]. Also, \mathcal{P}_g has a holomorphic symplectic form which is constructed using the monodromy representation associated to any projective structure [Go], [AB], [He]. Therefore, using the decomposition in (9.5) we obtain two closed holomorphic 2-forms on $\mathcal{H}_g(r)$: one is given by the symplectic form on $\mathcal{D}_X(r)$ and the other is given by the symplectic form on \mathcal{P}_g .

We end with the following conjecture:

Conjecture 9.3. The holomorphic symplectic form on $\mathcal{H}_g(r)$ in Corollary 8.2(1) is a constant linear combination of the above two holomorphic 2-forms on $\mathcal{H}_g(r)$.

References

- [At] M. F. Atiyah, Complex analytic connections in fibre bundles, *Trans. Amer. Math. Soc.* 85 (1957), 181–207.
- [AB] M. F. Atiyah and R. Bott, The Yang-Mills equations over Riemann surfaces, *Philos. Trans. Roy. Soc. London* 308 (1983), 523–615.
- [BNR] A. Beauville, M. S. Narasimhan and S. Ramanan, Spectral curves and the generalised theta divisor, *J. Reine Angew. Math.* **398** (1989), 169–179.
- [BD1] A. Beilinson and V. G. Drinfeld, Opers, arXiv math/0501398 (1993).
- [BD2] A. Beilinson and V. G. Drinfeld, Quantization of Hitchin's integrable system and Hecke eigensheaves, (1991).
- [BS] A. A. Beilinson and V. V. Schechtman, Determinant bundles and Virasoro algebras, Comm. Math. Phys. 118 (1988), 651–701.
- [BB] D. Ben-Zvi and I. Biswas, Theta functions and Szegő kernels, *Int. Math. Res. Not.* (2003), no. 24, 1305–1340.
- [Bi] I. Biswas, Coupled connections on a compact Riemann surface, *Jour. Math. Pures Appl.* 82 (2003), 1–42.
- [BH1] I. Biswas and J. Hurtubise, Meromorphic connections, determinant line bundles and the Tyurin parametrization, arXiv:1907.00133, Asian Jour. Math. (to appear).
- [BH2] I. Biswas and J. Hurtubise, A canonical connection on bundles on Riemann surfaces and Quillen connection on the theta bundle, *Adv. Math.* **389** (2021), Article 107918, arXiv:2102.00624.
- [BHH] I. Biswas, V. Heu and J. Hurtubise, Isomonodromic deformations of logarithmic connections and stability, *Math. Ann.* **366** (2016), 121–140.
- [BHS] I. Biswas, J. Hurtubise and J. Stasheff, A construction of a universal connection, Forum Math. 24 (2012), 365–378.
- [BR] I. Biswas and A. K. Raina, Projective structures on a Riemann surface. II, Internat. Math. Res. Not. (1999), no. 13, 685–716.
- [Ch] T. Chen, The associated map of the nonabelian Gauss-Manin connection, Cent. Eur. Jour. Math. 10 (2012), 1407–1421.
- [DN] J.-M. Drézet and M. S. Narasimhan, Groupe de Picard des variétés de modules de fibrés semistables sur les courbes algébriques, *Invent. Math.* 97 (1989), 53–94.
- [DS1] V. G. Drinfeld and V. V. Sokolov, Lie algebras and equations of Korteweg-de Vries type, Current problems in mathematics, Vol. 24, 81–180, Itogi Nauki i Tekhniki, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1984.

- [DS2] V. G. Drinfeld and V. V. Sokolov, Equations of Korteweg-de Vries type, and simple Lie algebras, Dokl. Akad. Nauk SSSR 258 (1981), 11–16.
- [Go] W. M. Goldman, The symplectic nature of fundamental groups of surfaces, Adv. in Math. 54 (1984), 200–225.
- [GH] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Pure and Applied Mathematics, Wiley-Interscience, New York, 1978.
- [Gu] R. C. Gunning, *Lectures on Riemann Surfaces*, Mathematical Notes 2, Princeton University Press, Princeton, NJ, 1966.
- [He] D. A. Hejhal, Monodromy groups and linearly polymorphic functions, *Acta Math.* **135** (1975), 1–55.
- [NS] M. S. Narasimhan and C. S. Seshadri, Stable and unitary vector bundles on a compact Riemann surface, *Ann. of Math.* 82 (1965), 540–567.
- [Sh] R. Sharpe, *Differential Geometry: Cartan's generalization of Klein's Erlangen program*, Graduate Texts in Mathematics, 166. Springer-Verlag, New York, 1997.
- [We] A. Weil, Généralisation des fonctions abéliennes, Jour. Math. Pures Appl. 17 (1938), 47–87.

School of Mathematics, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005

Email address: indranil@math.tifr.res.in

DEPARTMENT OF MATHEMATICS, MCGILL UNIVERSITY, BURNSIDE HALL, 805 SHERBROOKE ST. W., MONTREAL, QUE. H3A 2K6, CANADA

Email address: jacques.hurtubise@mcgill.ca

UNAM, LAREMA UMR 6093 DU CNRS, UNIVERSITÉ D'ANGERS, 49045 CEDEX 01, ANGERS, FRANCE

Email address: vladimir.roubtsov@univ-angers.fr