

# Characterization of the boundedness of generalized fractional integral and maximal operators on Orlicz-Morrey and weak Orlicz-Morrey spaces

Ryota Kawasumi, Eiichi Nakai and Minglei Shi

## Abstract

We give necessary and sufficient conditions for the boundedness of generalized fractional integral and maximal operators on Orlicz-Morrey and weak Orlicz-Morrey spaces. To do this we prove the weak-weak type modular inequality of the Hardy-Littlewood maximal operator with respect to the Young function. Orlicz-Morrey spaces contain  $L^p$  spaces ( $1 \leq p \leq \infty$ ), Orlicz spaces and generalized Morrey spaces as special cases. Hence we get necessary and sufficient conditions on these function spaces as corollaries.

## 1 Introduction

In this paper we consider the generalized fractional integral operator  $I_\rho$  and the generalized fractional maximal operator  $M_\rho$ . We give necessary and sufficient conditions for the boundedness of  $I_\rho$  and  $M_\rho$  on Orlicz-Morrey spaces  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and weak Orlicz-Morrey spaces  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$ . More precisely, we investigate the following boundedness of the operator  $T = I_\rho$  or  $M_\rho$ :

$$\begin{aligned} \|Tf\|_{wL^{(\Psi, \varphi)}} &\leq C\|f\|_{L^{(\Phi, \varphi)}}, \\ \|Tf\|_{L^{(\Psi, \varphi)}} &\leq C\|f\|_{L^{(\Phi, \varphi)}}, \\ \|Tf\|_{wL^{(\Psi, \varphi)}} &\leq C\|f\|_{wL^{(\Phi, \varphi)}}. \end{aligned}$$

We treat a wide class of Young functions as  $\Phi : [0, \infty] \rightarrow [0, \infty]$ . Orlicz-Morrey spaces contain  $L^p$  spaces ( $1 \leq p \leq \infty$ ), Orlicz spaces and generalized Morrey spaces as special cases. Hence we get necessary and sufficient conditions for the boundedness of  $I_\rho$  and  $M_\rho$  on these function spaces as corollaries.

Now, we recall the definitions of the Orlicz-Morrey and weak Orlicz-Morrey spaces. For a measurable set  $G \subset \mathbb{R}^n$ , we denote by  $|G|$  its Lebesgue measure. We

---

2010 *Mathematics Subject Classification.* 42B35, 46E30, 42B20, 42B25.

*Key words and phrases.* Fractional integral, fractional maximal operator, Orlicz-Morrey space, weak Orlicz-Morrey space, modular inequality.

denote by  $B(a, r)$  the open ball centered at  $a \in \mathbb{R}^n$  and of radius  $r$ . For a function  $f \in L^1_{\text{loc}}(\mathbb{R}^n)$  and a ball  $B$ , let

$$f_B = \int_B f = \int_B f(y) dy = \frac{1}{|B|} \int_B f(y) dy.$$

For a measurable set  $G \subset \mathbb{R}^n$ , a measurable function  $f$  and  $t \geq 0$ , let

$$m(G, f, t) = |\{x \in G : |f(x)| > t\}|.$$

In the case  $G = \mathbb{R}^n$ , we briefly denote it by  $m(f, t)$ .

**Definition 1.1.** For a Young function  $\Phi : [0, \infty] \rightarrow [0, \infty]$ , a function  $\varphi : (0, \infty) \rightarrow (0, \infty)$  and a ball  $B = B(a, r)$ , let

$$\|f\|_{\Phi, \varphi, B} = \inf \left\{ \lambda > 0 : \frac{1}{\varphi(r)} \int_B \Phi \left( \frac{|f(x)|}{\lambda} \right) dx \leq 1 \right\},$$

$$\|f\|_{\Phi, \varphi, B, \text{weak}} = \inf \left\{ \lambda > 0 : \sup_{t \in (0, \infty)} \frac{\Phi(t) m(B, f/\lambda, t)}{|B| \varphi(r)} \leq 1 \right\}.$$

Let  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  be the set of all measurable functions  $f$  on  $\mathbb{R}^n$  such that the following functionals are finite, respectively:

$$\|f\|_{L^{(\Phi, \varphi)}} = \sup_B \|f\|_{\Phi, \varphi, B},$$

$$\|f\|_{wL^{(\Phi, \varphi)}} = \sup_B \|f\|_{\Phi, \varphi, B, \text{weak}},$$

where the suprema are taken over all balls  $B$  in  $\mathbb{R}^n$ . (For the definition of the Young function, see Definition 2.2 below.)

Then  $\|\cdot\|_{L^{(\Phi, \varphi)}(\mathbb{R}^n)}$  is a norm and thereby  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  is a Banach space. On the other hand  $\|\cdot\|_{wL^{(\Phi, \varphi)}(\mathbb{R}^n)}$  is a quasi norm with the inequality

$$\|f + g\|_{wL^{(\Phi, \varphi)}} \leq 2(\|f\|_{wL^{(\Phi, \varphi)}} + \|g\|_{wL^{(\Phi, \varphi)}}),$$

and thereby  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  is a quasi Banach space.

If  $\varphi(r) = 1/r^n$ , then  $L^{(\Phi, \varphi)}(\mathbb{R}^n) = L^\Phi(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n) = wL^\Phi(\mathbb{R}^n)$ , which are the usual Orlicz and weak Orlicz spaces, respectively. In this paper we treat a wide class of Young functions such as

$$(1.1) \quad \Phi_1(t) = \begin{cases} t, & 0 \leq t \leq 1, \\ \infty, & t > 1, \end{cases} \quad \Phi_2(t) = \max(0, t^2 - 4), \quad t \geq 0.$$

We also treat generalized Young functions such as

$$(1.2) \quad \Phi_3(t) = \begin{cases} e^{1-1/t^p}, & 0 \leq t \leq 1, \\ e^{t^p-1}, & t > 1, \end{cases} \quad 0 < p < \infty,$$

which is not convex near  $t = 1$ .

If  $\Phi(t) = t^p$  ( $1 \leq p < \infty$ ), then we denote  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  by  $L^{(p, \varphi)}(\mathbb{R}^n)$  and  $wL^{(p, \varphi)}(\mathbb{R}^n)$ , which are the generalized Morrey and weak Morrey spaces, respectively. If  $\varphi_\lambda(r) = r^\lambda$ ,  $-n \leq \lambda < 0$ , then  $L^{(p, \varphi_\lambda)}(\mathbb{R}^n)$  is the classical Morrey space. In particular, if  $\lambda = -n$ , then it is the Lebesgue space  $L^p(\mathbb{R}^n)$ .

Orlicz spaces were introduced by [46, 47]. For the theory of Orlicz spaces, see [25, 26, 28, 32, 51] for example. Weak Orlicz spaces were studied in [21, 31, 40], etc. Morrey spaces were introduced by [35]. For their generalization, see [33, 36, 48, 49], etc. Weak Morrey spaces were studied in [13, 14, 53, 58, 63], etc. The Orlicz-Morrey space  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  was first studied in [41]. The spaces  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  were investigated in [17, 42, 43, 52], etc. For other kinds of Orlicz-Morrey spaces, see [4, 5, 8, 11, 55], etc. For the study related to weak Orlicz and weak Morrey spaces, see [7, 10, 15, 16, 19, 20, 29, 30, 64], etc.

Next we recall the generalized fractional integral operator  $I_\rho$ . For a function  $\rho : (0, \infty) \rightarrow (0, \infty)$ , the operator  $I_\rho$  is defined by

$$(1.3) \quad I_\rho f(x) = \int_{\mathbb{R}^n} \frac{\rho(|x-y|)}{|x-y|^n} f(y) dy, \quad x \in \mathbb{R}^n,$$

where we always assume that

$$(1.4) \quad \int_0^1 \frac{\rho(t)}{t} dt < \infty.$$

If  $\rho(r) = r^\alpha$ ,  $0 < \alpha < n$ , then  $I_\rho$  is the usual fractional integral operator  $I_\alpha$ . The condition (1.4) is needed for the integral in (1.3) to converge for bounded functions  $f$  with compact support. In this paper we also assume that there exist positive constants  $C$ ,  $K_1$  and  $K_2$  with  $K_1 < K_2$  such that, for all  $r > 0$ ,

$$(1.5) \quad \sup_{r \leq t \leq 2r} \rho(t) \leq C \int_{K_1 r}^{K_2 r} \frac{\rho(t)}{t} dt.$$

The condition (1.5) was considered in [50]. If  $\rho$  satisfies the doubling condition (2.12) below, then  $\rho$  satisfies (1.5). Let  $\rho(r) = \min(r^\alpha, e^{-r/2})$  with  $0 < \alpha < n$ , which controls the Bessel potential (see [59]). Then  $\rho$  also satisfies (1.5). The operator  $I_\rho$  was introduced in [38] to extend the Hardy-Littlewood-Sobolev theorem to Orlicz spaces whose partial results were announced in [37]. For example, the generalized fractional integral  $I_\rho$  is bounded from  $\exp L^p(\mathbb{R}^n)$  to  $\exp L^q(\mathbb{R}^n)$ , where

$$(1.6) \quad \rho(r) = \begin{cases} 1/(\log(1/r))^{\alpha+1} & \text{for small } r, \\ (\log r)^{\alpha-1} & \text{for large } r, \end{cases} \quad \alpha > 0,$$

$p, q \in (0, \infty)$ ,  $-1/p + \alpha = -1/q$  and  $\exp L^p(\mathbb{R}^n)$  is the Orlicz space  $L^\Phi(\mathbb{R}^n)$  with

$$(1.7) \quad \Phi(t) = \begin{cases} 1/\exp(1/t^p) & \text{for small } t, \\ \exp(t^p) & \text{for large } t. \end{cases}$$

See also [39, 40, 41, 42, 44].

We also consider the generalized fractional maximal operator  $M_\rho$  and compare its boundedness with  $I_\rho$ . For a function  $\rho : (0, \infty) \rightarrow (0, \infty)$ , the operator  $M_\rho$  is defined by

$$(1.8) \quad M_\rho f(x) = \sup_{B(a,r) \ni x} \rho(r) \int_{B(a,r)} |f(y)| dy, \quad x \in \mathbb{R}^n,$$

where the supremum is taken over all balls  $B(a, r)$  containing  $x$ . We need neither the condition (1.4) nor (1.5) on the definition of  $M_\rho$ . The operator  $M_\rho$  was studied in [54] on generalized Morrey spaces. See also [18]. If  $\rho(r) = |B(0, r)|^{\alpha/n}$ , then  $M_\rho$  is the usual fractional maximal operator  $M_\alpha$ . If  $\rho \equiv 1$ , then  $M_\rho$  is the Hardy-Littlewood maximal operator  $M$ . It is known that the usual fractional maximal operator  $M_\alpha$  is dominated pointwise by the fractional integral operator  $I_\alpha$ , that is,  $M_\alpha f(x) \leq CI_\alpha |f|(x)$  for all  $x \in \mathbb{R}^n$ . Then the boundedness of  $M_\alpha$  follows from one of  $I_\alpha$ . However, we have a better estimate of  $M_\rho$  than  $I_\rho$ .

To prove the boundedness of  $I_\rho$  and  $M_\rho$  on  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  we show the pointwise estimate by the Hardy-Littlewood maximal operator  $M$  and use the modular inequality of  $M$  with respect to Young functions  $\Phi$ . The strong and weak type modular inequalities are known. In this paper we show the weak-weak type modular inequality. In general, the modular inequality is stronger than the norm inequality. For the boundedness on generalized Morrey spaces  $L^{(p, \varphi)}(\mathbb{R}^n)$  we only need the  $L^p$ -norm inequality of  $M$ . However, for  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  we need the modular inequality.

The organization of this paper is as follows: In the next section we state precise definitions of the functions  $\Phi$  and  $\varphi$  by which we define  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$ . Then we state the main results in Section 3. To prove them in the final section, we give properties of Young functions, Orlicz-Morrey and weak Orlicz-Morrey spaces in Section 4.

At the end of this section, we make some conventions. Throughout this paper, we always use  $C$  to denote a positive constant that is independent of the main parameters involved but whose value may differ from line to line. Constants with subscripts, such as  $C_p$ , are dependent on the subscripts. If  $f \leq Cg$ , we then write  $f \lesssim g$  or  $g \gtrsim f$ ; and if  $f \lesssim g \lesssim f$ , we then write  $f \sim g$ .

## 2 On the functions $\Phi$ and $\varphi$

In this section we state on the functions  $\Phi$  and  $\varphi$  by which we define  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$ . We first recall the Young function and its generalization.

For an increasing (i.e. nondecreasing) function  $\Phi : [0, \infty] \rightarrow [0, \infty]$ , let

$$(2.1) \quad a(\Phi) = \sup\{t \geq 0 : \Phi(t) = 0\}, \quad b(\Phi) = \inf\{t \geq 0 : \Phi(t) = \infty\},$$

with convention  $\sup \emptyset = 0$  and  $\inf \emptyset = \infty$ . Then  $0 \leq a(\Phi) \leq b(\Phi) \leq \infty$ .

Let  $\overline{\Phi}$  be the set of all increasing functions  $\Phi : [0, \infty] \rightarrow [0, \infty]$  such that

$$(2.2) \quad 0 \leq a(\Phi) < \infty, \quad 0 < b(\Phi) \leq \infty,$$

$$(2.3) \quad \lim_{t \rightarrow +0} \Phi(t) = \Phi(0) = 0,$$

$$(2.4) \quad \Phi \text{ is left continuous on } [0, b(\Phi)),$$

$$(2.5) \quad \text{if } b(\Phi) = \infty, \text{ then } \lim_{t \rightarrow \infty} \Phi(t) = \Phi(\infty) = \infty,$$

$$(2.6) \quad \text{if } b(\Phi) < \infty, \text{ then } \lim_{t \rightarrow b(\Phi)-0} \Phi(t) = \Phi(b(\Phi)) (\leq \infty).$$

In what follows, if an increasing and left continuous function  $\Phi : [0, \infty) \rightarrow [0, \infty)$  satisfies (2.3) and  $\lim_{t \rightarrow \infty} \Phi(t) = \infty$ , then we always regard that  $\Phi(\infty) = \infty$  and that  $\Phi \in \overline{\Phi}$ .

For  $\Phi \in \overline{\Phi}$ , we recall the generalized inverse of  $\Phi$  in the sense of O'Neil [45, Definition 1.2].

**Definition 2.1.** For  $\Phi \in \overline{\Phi}$  and  $u \in [0, \infty]$ , let

$$(2.7) \quad \Phi^{-1}(u) = \begin{cases} \inf\{t \geq 0 : \Phi(t) > u\}, & u \in [0, \infty), \\ \infty, & u = \infty. \end{cases}$$

Let  $\Phi \in \overline{\Phi}$ . Then  $\Phi^{-1}$  is finite, increasing and right continuous on  $[0, \infty)$  and positive on  $(0, \infty)$ . If  $\Phi$  is bijective from  $[0, \infty]$  to itself, then  $\Phi^{-1}$  is the usual inverse function of  $\Phi$ . Moreover, if  $\Phi \in \overline{\Phi}$ , then

$$(2.8) \quad \Phi(\Phi^{-1}(u)) \leq u \leq \Phi^{-1}(\Phi(u)) \quad \text{for all } u \in [0, \infty],$$

which is a generalization of Property 1.3 in [45]. For its proof see [56, Proposition 2.2].

For  $\Phi, \Psi \in \overline{\Phi}$ , we write  $\Phi \approx \Psi$  if there exists a positive constant  $C$  such that

$$\Phi(C^{-1}t) \leq \Psi(t) \leq \Phi(Ct) \quad \text{for all } t \in [0, \infty].$$

For functions  $P, Q : [0, \infty] \rightarrow [0, \infty]$ , we write  $P \sim Q$  if there exists a positive constant  $C$  such that

$$C^{-1}P(t) \leq Q(t) \leq CP(t) \quad \text{for all } t \in [0, \infty].$$

Then, for  $\Phi, \Psi \in \overline{\Phi}$ ,

$$(2.9) \quad \Phi \approx \Psi \quad \Leftrightarrow \quad \Phi^{-1} \sim \Psi^{-1},$$

see [56, Lemma 2.3].

Now we recall the definition of the Young function and give its generalization.

**Definition 2.2.** A function  $\Phi \in \overline{\Phi}$  is called a Young function (or sometimes also called an Orlicz function) if  $\Phi$  is convex on  $[0, b(\Phi))$ . Let  $\Phi_Y$  be the set of all Young functions. Let  $\overline{\Phi}_Y$  be the set of all  $\Phi \in \overline{\Phi}$  such that  $\Phi \approx \Psi$  for some  $\Psi \in \Phi_Y$ .

For example,  $\Phi_1$  and  $\Phi_2$  defined by (1.1) are in  $\Phi_Y$ , and  $\Phi_3$  defined by (1.2) is in  $\overline{\Phi}_Y \setminus \Phi_Y$ .

Similar to Definition 1.1 we also define  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  by using generalized Young functions  $\Phi \in \overline{\Phi}_Y$  together with  $\|\cdot\|_{\Phi, \varphi, B}$  and  $\|\cdot\|_{\Phi, \varphi, B, \text{weak}}$ , respectively. Then  $\|\cdot\|_{\Phi, \varphi, B}$  and  $\|\cdot\|_{\Phi, \varphi, B, \text{weak}}$  are quasi norms and thereby  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  are quasi Banach spaces. Note that, for  $\Phi, \Psi \in \overline{\Phi}_Y$ , if  $\Phi \approx \Psi$  and  $\varphi \sim \psi$ , then  $L^{(\Phi, \varphi)}(\mathbb{R}^n) = L^{(\Psi, \psi)}(\mathbb{R}^n)$  and  $wL^{(\Phi, \varphi)}(\mathbb{R}^n) = wL^{(\Psi, \psi)}(\mathbb{R}^n)$  with equivalent quasi norms.

**Definition 2.3.** (i) A function  $\Phi \in \overline{\Phi}$  is said to satisfy the  $\Delta_2$ -condition, denoted by  $\Phi \in \overline{\Delta}_2$ , if there exists a constant  $C > 0$  such that

$$(2.10) \quad \Phi(2t) \leq C\Phi(t) \quad \text{for all } t > 0.$$

(ii) A function  $\Phi \in \overline{\Phi}$  is said to satisfy the  $\nabla_2$ -condition, denoted by  $\Phi \in \overline{\nabla}_2$ , if there exists a constant  $k > 1$  such that

$$(2.11) \quad \Phi(t) \leq \frac{1}{2k}\Phi(kt) \quad \text{for all } t > 0.$$

(iii) Let  $\Delta_2 = \Phi_Y \cap \overline{\Delta}_2$  and  $\nabla_2 = \Phi_Y \cap \overline{\nabla}_2$ .

Next, we say that a function  $\theta : (0, \infty) \rightarrow (0, \infty)$  satisfies the doubling condition if there exists a positive constant  $C$  such that, for all  $r, s \in (0, \infty)$ ,

$$(2.12) \quad \frac{1}{C} \leq \frac{\theta(r)}{\theta(s)} \leq C, \quad \text{if } \frac{1}{2} \leq \frac{r}{s} \leq 2.$$

We say that  $\theta$  is almost increasing (resp. almost decreasing) if there exists a positive constant  $C$  such that, for all  $r, s \in (0, \infty)$ ,

$$(2.13) \quad \theta(r) \leq C\theta(s) \quad (\text{resp. } \theta(s) \leq C\theta(r)), \quad \text{if } r < s.$$

In this paper we consider the following class of  $\varphi : (0, \infty) \rightarrow (0, \infty)$ .

**Definition 2.4.** Let  $\mathcal{G}^{\text{dec}}$  be the set of all functions  $\varphi : (0, \infty) \rightarrow (0, \infty)$  such that  $\varphi$  is almost decreasing and that  $r \mapsto \varphi(r)r^n$  is almost increasing. That is, there exists a positive constant  $C$  such that, for all  $r, s \in (0, \infty)$ ,

$$C\varphi(r) \geq \varphi(s), \quad \varphi(r)r^n \leq C\varphi(s)s^n, \quad \text{if } r < s.$$

If  $\varphi \in \mathcal{G}^{\text{dec}}$ , then  $\varphi$  satisfies doubling condition. Let  $\psi : (0, \infty) \rightarrow (0, \infty)$ . If  $\psi \sim \varphi$  for some  $\varphi \in \mathcal{G}^{\text{dec}}$ , then  $\psi \in \mathcal{G}^{\text{dec}}$ .

**Remark 2.5.** Let  $\varphi \in \mathcal{G}^{\text{dec}}$ . Then there exists  $\tilde{\varphi} \in \mathcal{G}^{\text{dec}}$  such that  $\varphi \sim \tilde{\varphi}$  and that  $\tilde{\varphi}$  is continuous and strictly decreasing, see [42, Proposition 3.4]. Moreover, if

$$(2.14) \quad \lim_{r \rightarrow 0} \varphi(r) = \infty, \quad \lim_{r \rightarrow \infty} \varphi(r) = 0,$$

then  $\tilde{\varphi}$  is bijective from  $(0, \infty)$  to itself.

At the end of this section, we note that, for  $\Phi \in \Phi_Y$  and a ball  $B$ , the following relation holds

$$(2.15) \quad \sup_{t \in (0, \infty)} \Phi(t)m(B, f, t) = \sup_{t \in (0, \infty)} t m(B, f, \Phi^{-1}(t)) = \sup_{t \in (0, \infty)} t m(B, \Phi(|f|), t).$$

See [22] for the proof of (2.15). Hence, the norm inequality  $\|Tf\|_{\text{w}L(\Psi, \varphi)} \leq C\|f\|_{L(\Phi, \varphi)}$  holds if and only if

$$t m \left( B, \Psi \left( \frac{|Tf|}{C\|f\|_{L(\Phi, \varphi)}} \right), t \right) \leq |B|\varphi(r)$$

holds for all balls  $B = B(a, r)$  and  $t \in (0, \infty)$ .

### 3 Main results

For a measurable function  $f$  and  $t \geq 0$ , recall that  $m(f, t) = |\{x \in \mathbb{R}^n : |f(x)| > t\}|$ . First we state known modular inequalities for the Hardy-Littlewood maximal operator  $M$ . For their proofs, see [26, Theorem 1.2.1 and Lemma 1.2.4] for example.

**Theorem 3.1** ([26, 62]). Let  $\Phi \in \overline{\Phi}_Y$ . Then there exists a positive constant  $C_\Phi$  such that

$$(3.1) \quad \sup_{t \in (0, \infty)} \Phi(t)m(Mf, t) \leq \int_{\mathbb{R}^n} \Phi(C_\Phi|f(x)|) dx.$$

Moreover, if  $\Phi \in \overline{\nabla}_2$ , then

$$(3.2) \quad \int_{\mathbb{R}^n} \Phi(Mf(x)) dx \leq \int_{\mathbb{R}^n} \Phi(C_\Phi|f(x)|) dx.$$

If  $\Phi \in \overline{\nabla}_2$  and  $\varphi \in \mathcal{G}^{\text{dec}}$ , then  $\text{w}L^{(\Phi, \varphi)}(\mathbb{R}^n) \subset L_{\text{loc}}^1(\mathbb{R}^n)$ , see Lemma 4.7 below. Hence,  $Mf$  is well defined for  $f \in \text{w}L^{(\Phi, \varphi)}(\mathbb{R}^n)$ , in particular, for  $f \in \text{w}L^\Phi(\mathbb{R}^n)$ .

Our first result is the following modular inequality.

**Theorem 3.2.** If  $\Phi \in \overline{\nabla}_2$ , then there exists a positive constant  $C_\Phi$  such that

$$(3.3) \quad \sup_{t \in (0, \infty)} \Phi(t)m(Mf, t) \leq \sup_{t \in (0, \infty)} \Phi(t)m(C_\Phi f, t).$$

Liu and Wang [31] proved the norm inequality for  $wL^\Phi(\mathbb{R}^n)$  with  $\Phi \in \Delta_2 \cap \nabla_2$ . Theorem 3.2 is its extension.

By Theorems 3.1 and 3.2 we will prove the following boundedness, which is an extension of [42, Theorem 6.1].

**Theorem 3.3.** *Let  $\Phi \in \overline{\Phi}_Y$  and  $\varphi \in \mathcal{G}^{\text{dec}}$ . Then the Hardy-Littlewood maximal operator  $M$  is bounded from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$ . Moreover, if  $\Phi \in \overline{\nabla}_2$ , then the operator  $M$  is bounded from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to itself and from  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  to itself.*

Next we state the boundedness of generalized fractional integral operators  $I_\rho$ .

**Theorem 3.4.** *Let  $\Phi, \Psi \in \overline{\Phi}_Y$ ,  $\varphi \in \mathcal{G}^{\text{dec}}$  and  $\rho : (0, \infty) \rightarrow (0, \infty)$ . Assume that  $\rho$  satisfies (1.4) and (1.5).*

- (i) *Assume that  $\varphi$  satisfies (2.14) and that there exists a positive constant  $A$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.4) \quad \int_0^r \frac{\rho(t)}{t} dt \Phi^{-1}(\varphi(r)) + \int_r^\infty \frac{\rho(t) \Phi^{-1}(\varphi(t))}{t} dt \leq A \Psi^{-1}(\varphi(r)).$$

*Then, for any positive constant  $C_0$ , there exists a positive constant  $C_1$  such that, for all  $f \in L^{(\Phi, \varphi)}(\mathbb{R}^n)$  with  $f \not\equiv 0$ ,*

$$(3.5) \quad \Psi \left( \frac{|I_\rho f(x)|}{C_1 \|f\|_{L^{(\Phi, \varphi)}}} \right) \leq \Phi \left( \frac{Mf(x)}{C_0 \|f\|_{L^{(\Phi, \varphi)}}} \right), \quad x \in \mathbb{R}^n.$$

*Consequently,  $I_\rho$  is bounded from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $wL^{(\Psi, \varphi)}(\mathbb{R}^n)$ . Moreover, if  $\Phi \in \overline{\nabla}_2$ , then, for all  $f \in wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  with  $f \not\equiv 0$ ,*

$$(3.6) \quad \Psi \left( \frac{|I_\rho f(x)|}{C_1 \|f\|_{wL^{(\Phi, \varphi)}}} \right) \leq \Phi \left( \frac{Mf(x)}{C_0 \|f\|_{wL^{(\Phi, \varphi)}}} \right), \quad x \in \mathbb{R}^n.$$

*Consequently, if  $\Phi \in \overline{\nabla}_2$ , then  $I_\rho$  is bounded from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $L^{(\Psi, \varphi)}(\mathbb{R}^n)$  by (3.5) and from  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $wL^{(\Psi, \varphi)}(\mathbb{R}^n)$  by (3.6).*

- (ii) *If  $I_\rho$  is bounded from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $wL^{(\Psi, \varphi)}(\mathbb{R}^n)$ , then there exists a positive constant  $A'$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.7) \quad \int_0^r \frac{\rho(t)}{t} dt \Phi^{-1}(\varphi(r)) \leq A' \Psi^{-1}(\varphi(r)).$$

*Moreover, under the assumption that there exists a positive constant  $C$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.8) \quad \int_0^r \varphi(t) t^{n-1} dt \leq C \varphi(r) r^n,$$

*if  $I_\rho$  is bounded from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $wL^{(\Psi, \varphi)}(\mathbb{R}^n)$ , then (3.4) holds for some  $A \in (0, \infty)$  and for all  $r \in (0, \infty)$ .*



(iii) In the part (ii) the boundedness from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $wL^{(\Psi, \varphi)}(\mathbb{R}^n)$  can be replaced by the boundedness from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $L^{(1, \Psi^{-1}(\varphi))}(\mathbb{R}^n)$ .

The part (i) in the above theorem is an extension of [42, Theorem 7.3]. In the part (iii), if  $\Psi \in \overline{\nabla}_2$ , then  $wL^{(\Psi, \varphi)}(\mathbb{R}^n) \subset L^{(1, \Psi^{-1}(\varphi))}(\mathbb{R}^n)$ , see Lemma 4.7 below. In this case the part (iii) implies the part (ii).

Thirdly, we state the boundedness of the generalized fractional maximal operators  $M_\rho$ . In this case we need neither the assumption (1.4) nor (1.5) on the function  $\rho : (0, \infty) \rightarrow (0, \infty)$ .

**Theorem 3.5.** *Let  $\Phi, \Psi \in \overline{\Phi}_Y$ ,  $\varphi \in \mathcal{G}^{\text{dec}}$  and  $\rho : (0, \infty) \rightarrow (0, \infty)$ .*

(i) *Assume that  $\lim_{r \rightarrow \infty} \varphi(r) = 0$  or that  $\Psi^{-1}(t)/\Phi^{-1}(t)$  is almost decreasing on  $(0, \infty)$ . If there exists a positive constant  $A$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.9) \quad \left( \sup_{0 < t \leq r} \rho(t) \right) \Phi^{-1}(\varphi(r)) \leq A \Psi^{-1}(\varphi(r)),$$

*then, for any positive constant  $C_0$ , there exists a positive constant  $C_1$  such that, for all  $f \in L^{(\Phi, \varphi)}(\mathbb{R}^n)$  with  $f \not\equiv 0$ ,*

$$(3.10) \quad \Psi \left( \frac{M_\rho f(x)}{C_1 \|f\|_{L^{(\Phi, \varphi)}}} \right) \leq \Phi \left( \frac{Mf(x)}{C_0 \|f\|_{L^{(\Phi, \varphi)}}} \right), \quad x \in \mathbb{R}^n.$$

*Consequently,  $M_\rho$  is bounded from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $wL^{(\Psi, \varphi)}(\mathbb{R}^n)$ . Moreover, if  $\Phi \in \overline{\nabla}_2$ , then, for all  $f \in wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  with  $f \not\equiv 0$ ,*

$$(3.11) \quad \Psi \left( \frac{M_\rho f(x)}{C_1 \|f\|_{wL^{(\Phi, \varphi)}}} \right) \leq \Phi \left( \frac{Mf(x)}{C_0 \|f\|_{wL^{(\Phi, \varphi)}}} \right), \quad x \in \mathbb{R}^n.$$

*Consequently, if  $\Phi \in \overline{\nabla}_2$ , then  $M_\rho$  is bounded from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $L^{(\Psi, \varphi)}(\mathbb{R}^n)$  by (3.10) and from  $wL^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $wL^{(\Psi, \varphi)}(\mathbb{R}^n)$  by (3.11).*

(ii) *If  $M_\rho$  is bounded from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $wL^{(\Psi, \varphi)}(\mathbb{R}^n)$ , then (3.9) holds for some  $A \in (0, \infty)$  and for all  $r \in (0, \infty)$ .*

(iii) *In the part (ii) the boundedness from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $wL^{(\Psi, \varphi)}(\mathbb{R}^n)$  can be replaced by the boundedness from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $L^{(1, \Psi^{-1}(\varphi))}(\mathbb{R}^n)$ .*

The part (i) in the above theorem is an extension of [57, Theorem 5.1]. If  $\Psi \in \overline{\nabla}_2$ , then the part (iii) implies the part (ii) as same as Theorem 3.4.

**Remark 3.6.** *From (1.5) and (3.4) it follows that*

$$\left( \sup_{0 < t \leq r} \rho(t) \right) \Phi^{-1}(\varphi(r)) \lesssim \int_0^{K_2 r} \frac{\rho(t)}{t} dt \Phi^{-1}(\varphi(r)) \lesssim \Psi^{-1}(\varphi(r)),$$

*which is the condition (3.9). If  $\rho(r) = (\log(1/r))^{-\alpha}$  for small  $r > 0$  or  $\rho(r) = (\log r)^\alpha$  for large  $r > 0$  with  $\alpha \geq 0$ , then the condition (3.9) is strictly weaker than (3.4).*

For the case  $\varphi(r) = 1/r^n$ , we have the following corollaries.

**Corollary 3.7** ([9, 27, 26, 31]). *Let  $\Phi \in \overline{\Phi}_Y$ . Then the Hardy-Littlewood maximal operator  $M$  is bounded from  $L^\Phi(\mathbb{R}^n)$  to  $wL^\Phi(\mathbb{R}^n)$ . Moreover, if  $\Phi \in \overline{\nabla}_2$ , then the operator  $M$  is bounded from  $L^\Phi(\mathbb{R}^n)$  to itself and from  $wL^\Phi(\mathbb{R}^n)$  to itself.*

For the boundedness of the operator  $M$  on Orlicz spaces, see also Cianchi [3] and Kita [23, 24]. Cianchi [3] also investigated the operators  $M_\alpha$  and  $I_\alpha$  on Orlicz spaces.

**Corollary 3.8.** *Let  $\Phi, \Psi \in \overline{\Phi}_Y$  and  $\rho : (0, \infty) \rightarrow (0, \infty)$ . Assume that  $\rho$  satisfies (1.4) and (1.5).*

(i) *Assume that there exists a positive constant  $A$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.12) \quad \int_0^r \frac{\rho(t)}{t} dt \Phi^{-1}(1/r^n) + \int_r^\infty \frac{\rho(t) \Phi^{-1}(1/t^n)}{t} dt \leq A \Psi^{-1}(1/r^n).$$

*Then, for any positive constant  $C_0$ , there exists a positive constant  $C_1$  such that, for all  $f \in L^\Phi(\mathbb{R}^n)$  with  $f \not\equiv 0$ ,*

$$(3.13) \quad \Psi \left( \frac{|I_\rho f(x)|}{C_1 \|f\|_{L^\Phi}} \right) \leq \Phi \left( \frac{Mf(x)}{C_0 \|f\|_{L^\Phi}} \right), \quad x \in \mathbb{R}^n.$$

*Consequently,  $I_\rho$  is bounded from  $L^\Phi(\mathbb{R}^n)$  to  $wL^\Psi(\mathbb{R}^n)$ . Moreover, if  $\Phi \in \overline{\nabla}_2$ , then, for all  $f \in wL^\Phi(\mathbb{R}^n)$  with  $f \not\equiv 0$ ,*

$$(3.14) \quad \Psi \left( \frac{|I_\rho f(x)|}{C_1 \|f\|_{wL^\Phi}} \right) \leq \Phi \left( \frac{Mf(x)}{C_0 \|f\|_{wL^\Phi}} \right), \quad x \in \mathbb{R}^n.$$

*Consequently, if  $\Phi \in \overline{\nabla}_2$ , then  $I_\rho$  is bounded from  $L^\Phi(\mathbb{R}^n)$  to  $L^\Psi(\mathbb{R}^n)$  and from  $wL^\Phi(\mathbb{R}^n)$  to  $wL^\Psi(\mathbb{R}^n)$ .*

(ii) *If  $I_\rho$  is bounded from  $L^\Phi(\mathbb{R}^n)$  to  $wL^\Psi(\mathbb{R}^n)$ , then there exists a positive constant  $A'$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.15) \quad \int_0^r \frac{\rho(t)}{t} dt \Phi^{-1}(1/r^n) \leq A' \Psi^{-1}(1/r^n).$$

*Moreover, under the assumption that there exists a positive constant  $A''$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.16) \quad \int_r^\infty \frac{\rho(t) \Phi^{-1}(1/t^n)}{t} dt \leq A'' \Psi^{-1}(1/r^n),$$

*if  $I_\rho$  is bounded from  $L^\Phi(\mathbb{R}^n)$  to  $wL^\Psi(\mathbb{R}^n)$ , then (3.12) holds for some  $A \in (0, \infty)$  and for all  $r \in (0, \infty)$ .*

(iii) In the part (ii) the boundedness from  $L^\Phi(\mathbb{R}^n)$  to  $wL^\Psi(\mathbb{R}^n)$  can be replaced by the boundedness from  $L^\Phi(\mathbb{R}^n)$  to  $L^{(1,\psi)}(\mathbb{R}^n)$  with  $\psi(r) = \Psi^{-1}(1/r^n)$ .

The above corollary is an extension of [4, Theorem 3]. See [39], for examples of  $\Phi, \Psi \in \overline{\Phi}_Y$  which satisfy the assumption in Corollary 3.8. The boundedness of the usual fractional integral operator on Orlicz spaces was given by O'Neil [45]. See also [34] for the boundedness of  $I_\rho$  on Orlicz space  $L^\Phi(\Omega)$  with bounded domain  $\Omega \subset \mathbb{R}^n$ .

**Corollary 3.9.** *Let  $\Phi, \Psi \in \overline{\Phi}_Y$  and  $\rho : (0, \infty) \rightarrow (0, \infty)$ .*

(i) *If there exists a positive constant  $A$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.17) \quad \left( \sup_{0 < t \leq r} \rho(t) \right) \Phi^{-1}(1/r^n) \leq A \Psi^{-1}(1/r^n),$$

*then, for any positive constant  $C_0$ , there exists a positive constant  $C_1$  such that, for all  $f \in L^\Phi(\mathbb{R}^n)$  with  $f \not\equiv 0$ ,*

$$(3.18) \quad \Psi \left( \frac{M_\rho f(x)}{C_1 \|f\|_{L^\Phi}} \right) \leq \Phi \left( \frac{Mf(x)}{C_0 \|f\|_{L^\Phi}} \right), \quad x \in \mathbb{R}^n.$$

*Consequently,  $M_\rho$  is bounded from  $L^\Phi(\mathbb{R}^n)$  to  $wL^\Psi(\mathbb{R}^n)$ . Moreover, if  $\Phi \in \overline{\nabla}_2$ , then, for all  $f \in wL^\Phi(\mathbb{R}^n)$  with  $f \not\equiv 0$ ,*

$$(3.19) \quad \Psi \left( \frac{M_\rho f(x)}{C_1 \|f\|_{wL^\Phi}} \right) \leq \Phi \left( \frac{Mf(x)}{C_0 \|f\|_{wL^\Phi}} \right), \quad x \in \mathbb{R}^n.$$

*Consequently, if  $\Phi \in \overline{\nabla}_2$ , then  $M_\rho$  is bounded from  $L^\Phi(\mathbb{R}^n)$  to  $L^\Psi(\mathbb{R}^n)$  and from  $wL^\Phi(\mathbb{R}^n)$  to  $wL^\Psi(\mathbb{R}^n)$ .*

(ii) *If  $M_\rho$  is bounded from  $L^\Phi(\mathbb{R}^n)$  to  $wL^\Psi(\mathbb{R}^n)$ , then (3.17) holds for some  $A \in (0, \infty)$  and for all  $r \in (0, \infty)$ .*

(iii) *In the part (ii) the boundedness from  $L^\Phi(\mathbb{R}^n)$  to  $wL^\Psi(\mathbb{R}^n)$  can be replaced by the boundedness from  $L^\Phi(\mathbb{R}^n)$  to  $L^{(1,\psi)}(\mathbb{R}^n)$  with  $\psi(r) = \Psi^{-1}(1/r^n)$ .*

The above corollary is an extension of [56, Theorem 3.8].

For the case  $\Phi(t) = t^p$ , we have the following corollaries.

**Corollary 3.10.** *Let  $1 \leq p < \infty$  and  $\varphi \in \mathcal{G}^{\text{dec}}$ . Then the Hardy-Littlewood maximal operator  $M$  is bounded from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $wL^{(p,\varphi)}(\mathbb{R}^n)$ . Moreover, if  $1 < p < \infty$ , then the operator  $M$  is bounded from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to itself and from  $wL^{(p,\varphi)}(\mathbb{R}^n)$  to itself.*

The above corollary is an extension of [36, Theorem 1] and [42, Corollary 6.2]. The boundedness of the Hardy-Littlewood maximal operator on the classical Morrey space was proven by Chiarenza and Frasca [2].

**Corollary 3.11.** *Let  $1 \leq p < q < \infty$ ,  $\varphi \in \mathcal{G}^{\text{dec}}$  and  $\rho : (0, \infty) \rightarrow (0, \infty)$ . Assume that  $\rho$  satisfies (1.4) and (1.5).*

- (i) *Assume that  $\varphi$  satisfies (2.14) and that there exists a positive constant  $A$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.20) \quad \int_0^r \frac{\rho(t)}{t} dt \varphi(r)^{1/p} + \int_r^\infty \frac{\rho(t)\varphi(t)^{1/p}}{t} dt \leq A\varphi(r)^{1/q}.$$

*Then there exists a positive constant  $C$  such that, for all  $f \in L^{(p,\varphi)}(\mathbb{R}^n)$ ,*

$$(3.21) \quad |I_\rho f(x)| \leq CMf(x)^{p/q} (\|f\|_{L^{(p,\varphi)}})^{1-p/q}, \quad x \in \mathbb{R}^n.$$

*Consequently,  $I_\rho$  is bounded from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $wL^{(q,\varphi)}(\mathbb{R}^n)$ . Moreover, if  $p > 1$ , then, for all  $f \in wL^{(p,\varphi)}(\mathbb{R}^n)$ ,*

$$(3.22) \quad |I_\rho f(x)| \leq CMf(x)^{p/q} (\|f\|_{wL^{(p,\varphi)}})^{1-p/q}, \quad x \in \mathbb{R}^n.$$

*Consequently, if  $p > 1$ , then  $I_\rho$  is bounded from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $L^{(q,\varphi)}(\mathbb{R}^n)$  by (3.21) and from  $wL^{(p,\varphi)}(\mathbb{R}^n)$  to  $wL^{(q,\varphi)}(\mathbb{R}^n)$  by (3.22).*

- (ii) *If  $I_\rho$  is bounded from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $wL^{(q,\varphi)}(\mathbb{R}^n)$ , then there exists a positive constant  $A'$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.23) \quad \int_0^r \frac{\rho(t)}{t} dt \varphi(r)^{1/p} \leq A'\varphi(r)^{1/q}.$$

*Moreover, under the assumption (3.8), if  $I_\rho$  is bounded from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $wL^{(q,\varphi)}(\mathbb{R}^n)$ , then (3.20) holds for some  $A \in (0, \infty)$  and for all  $r \in (0, \infty)$ .*

- (iii) *In the part (ii) the boundedness from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $wL^{(q,\varphi)}(\mathbb{R}^n)$  can be replaced by the boundedness from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $L^{(1,\varphi^{1/q})}(\mathbb{R}^n)$ .*

The above corollary is an extension of [6, Theorems 1.1, 1.2, 1.8 and 1.9]. The condition (3.20) was first given by Gunawan [12]. The boundedness of the usual fractional integral operator on the classical Morrey space was proven by Adams [1].

**Corollary 3.12.** *Let  $1 \leq p \leq q < \infty$ ,  $\varphi \in \mathcal{G}^{\text{dec}}$  and  $\rho : (0, \infty) \rightarrow (0, \infty)$ .*

- (i) *If there exists a positive constant  $A$  such that, for all  $r \in (0, \infty)$ ,*

$$(3.24) \quad \left( \sup_{0 < t \leq r} \rho(t) \right) \varphi(r)^{1/p} \leq A\varphi(r)^{1/q},$$

*then there exists a positive constant  $C$  such that, for all  $f \in L^{(p,\varphi)}(\mathbb{R}^n)$ ,*

$$(3.25) \quad M_\rho f(x) \leq CMf(x)^{p/q} (\|f\|_{L^{(p,\varphi)}})^{1-p/q}, \quad x \in \mathbb{R}^n.$$

Consequently,  $M_\rho$  is bounded from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $wL^{(q,\varphi)}(\mathbb{R}^n)$ . Moreover, if  $p > 1$ , then, for all  $f \in wL^{(p,\varphi)}(\mathbb{R}^n)$ ,

$$(3.26) \quad M_\rho f(x) \leq CMf(x)^{p/q} (\|f\|_{wL^{(p,\varphi)}})^{1-p/q}, \quad x \in \mathbb{R}^n.$$

Consequently, if  $p > 1$ , then  $M_\rho$  is bounded from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $L^{(q,\varphi)}(\mathbb{R}^n)$  by (3.25) and from  $wL^{(p,\varphi)}(\mathbb{R}^n)$  to  $wL^{(q,\varphi)}(\mathbb{R}^n)$  by (3.26).

- (ii) If  $M_\rho$  is bounded from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $wL^{(q,\varphi)}(\mathbb{R}^n)$ , then (3.24) holds for some  $A \in (0, \infty)$  and for all  $r \in (0, \infty)$ .
- (iii) In the part (ii) the boundedness from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $wL^{(q,\varphi)}(\mathbb{R}^n)$  can be replaced by the boundedness from  $L^{(p,\varphi)}(\mathbb{R}^n)$  to  $L^{(1,\varphi^{1/q})}(\mathbb{R}^n)$ .

## 4 Properties of Young functions and Orlicz-Morrey spaces

In this section we state the properties of Young functions, Orlicz-Morrey and weak Orlicz Morrey spaces. By the convexity, any Young function  $\Phi$  is continuous on  $[0, b(\Phi))$  and strictly increasing on  $[a(\Phi), b(\Phi)]$ . Hence  $\Phi$  is bijective from  $[a(\Phi), b(\Phi)]$  to  $[0, \Phi(b(\Phi))]$ .

**Definition 4.1.** *Let*

$$\begin{aligned} \mathcal{Y}^{(1)} &= \{\Phi \in \Phi_Y : b(\Phi) = \infty\}, \\ \mathcal{Y}^{(2)} &= \{\Phi \in \Phi_Y : b(\Phi) < \infty, \Phi(b(\Phi)) = \infty\}, \\ \mathcal{Y}^{(3)} &= \{\Phi \in \Phi_Y : b(\Phi) < \infty, \Phi(b(\Phi)) < \infty\}. \end{aligned}$$

**Remark 4.2.** (i) If  $\Phi \in \mathcal{Y}^{(1)} \cup \mathcal{Y}^{(2)}$ , then  $\Phi(\Phi^{-1}(u)) = u$  for all  $u \in [0, \infty]$ .

- (ii) If  $\Phi \in \mathcal{Y}^{(3)}$  and  $0 < \delta < 1$ , then there exists a Young function  $\Psi \in \mathcal{Y}^{(2)}$  such that  $b(\Phi) = b(\Psi)$  and

$$\Psi(\delta t) \leq \Phi(t) \leq \Psi(t) \quad \text{for all } t \in [0, \infty).$$

To see this we only set  $\Psi = \Phi + \Theta$ , where we choose  $\Theta \in \mathcal{Y}^{(2)}$  such that  $a(\Theta) = \delta b(\Phi)$  and  $b(\Theta) = b(\Phi)$ .

- (iii)  $\overline{\nabla}_2 \subset \overline{\Phi}_Y$  ([26, Lemma 1.2.3]).
- (iv) Let  $\Phi \in \overline{\Phi}_Y$ . Then  $\Phi \in \overline{\Delta}_2$  if and only if  $\Phi \approx \Psi$  for some  $\Psi \in \Delta_2$ , and,  $\Phi \in \overline{\nabla}_2$  if and only if  $\Phi \approx \Psi$  for some  $\Psi \in \nabla_2$ .
- (v) If  $\Phi \in \Phi_Y$ , then  $\Phi^{-1}$  satisfies the doubling condition by its concavity.

(vi) Let  $\Phi \in \Phi_Y$ . Then  $\Phi \in \nabla_2$  if and only if  $t \mapsto \frac{\Phi(t)}{t^p}$  is almost increasing for some  $p \in (1, \infty)$ .

For a Young function  $\Phi$ , its complementary function is defined by

$$\tilde{\Phi}(t) = \begin{cases} \sup\{tu - \Phi(u) : u \in [0, \infty)\}, & t \in [0, \infty), \\ \infty, & t = \infty. \end{cases}$$

Then  $\tilde{\Phi}$  is also a Young function, and  $(\Phi, \tilde{\Phi})$  is called a complementary pair. For example, if  $\Phi(t) = t^p/p$ , then  $\tilde{\Phi}(t) = t^{p'}/p'$  for  $p, p' \in (1, \infty)$  and  $1/p + 1/p' = 1$ . If  $\Phi(t) = t$ , then

$$\tilde{\Phi}(t) = \begin{cases} 0, & t \in [0, 1], \\ \infty, & t \in (1, \infty]. \end{cases}$$

Let  $(\Phi, \tilde{\Phi})$  be a complementary pair of functions in  $\Phi_Y$ . Then the following inequality holds:

$$(4.1) \quad t \leq \Phi^{-1}(t)\tilde{\Phi}^{-1}(t) \leq 2t \quad \text{for } t \in [0, \infty],$$

which is (1.3) in [60].

Orlicz and weak Orlicz spaces on a measure space  $(\Omega, \mu)$  defined by the following: For  $\Phi \in \overline{\Phi}_Y$ , let  $L^\Phi(\Omega, \mu)$  and  $wL^\Phi(\Omega, \mu)$  be the set of all measurable functions  $f$  such that the following functionals are finite, respectively:

$$\|f\|_{L^\Phi(\Omega, \mu)} = \inf \left\{ \lambda > 0 : \int_{\Omega} \Phi \left( \frac{|f(x)|}{\lambda} \right) d\mu(x) \leq 1 \right\},$$

$$\|f\|_{wL^\Phi(\Omega, \mu)} = \inf \left\{ \lambda > 0 : \sup_{t \in (0, \infty)} \Phi(t) \mu \left( \frac{f}{\lambda}, t \right) \leq 1 \right\},$$

where

$$\mu(f, t) = \mu(\{x \in \Omega : |f(x)| > t\}).$$

Then  $\|\cdot\|_{L^\Phi(\Omega, \mu)}$  and  $\|\cdot\|_{wL^\Phi(\Omega, \mu)}$  are quasi norms and thereby  $L^\Phi(\Omega, \mu)$  and  $wL^\Phi(\Omega, \mu)$  are quasi Banach spaces. If  $\Phi \in \Phi_Y$ , then  $\|\cdot\|_{L^\Phi(\Omega, \mu)}$  is a norm and thereby  $L^\Phi(\Omega, \mu)$  is a Banach space. For  $\Phi, \Psi \in \overline{\Phi}_Y$ , if  $\Phi \approx \Psi$ , then  $L^\Phi(\Omega, \mu) = L^\Psi(\Omega, \mu)$  and  $wL^\Phi(\Omega, \mu) = wL^\Psi(\Omega, \mu)$  with equivalent quasi norms, respectively.

For a measurable set  $G \subset \Omega$ , let  $\chi_G$  be its characteristic function. For  $\Phi \in \Phi_Y$ , it is known that

$$(4.2) \quad \|\chi_G\|_{wL^\Phi(\Omega, \mu)} = \|\chi_G\|_{L^\Phi(\Omega, \mu)} = \frac{1}{\Phi^{-1}(1/\mu(G))}, \quad \text{if } \mu(G) > 0.$$

Let  $(\Phi, \tilde{\Phi})$  be a complementary pair of functions in  $\Phi_Y$ . Then the following generalized Hölder's inequality holds:

$$(4.3) \quad \int_{\Omega} |f(x)g(x)| d\mu(x) \leq 2\|f\|_{L^\Phi(\Omega, \mu)}\|g\|_{L^{\tilde{\Phi}}(\Omega, \mu)}.$$

For a ball  $B = B(a, r) \subset \mathbb{R}^n$ , let  $\mu_B = dx/(|B|\varphi(r))$ . Then we have the following relations.

$$(4.4) \quad \|f\|_{\Phi, \varphi, B} = \|f\|_{L^\Phi(B, \mu_B)}, \quad \|f\|_{\Phi, \varphi, B, \text{weak}} = \|f\|_{\text{w}L^\Phi(B, \mu_B)}.$$

By the relations (4.2) and (4.4), we have

$$(4.5) \quad \|\chi_B\|_{\Phi, \varphi, B, \text{weak}} = \|\chi_B\|_{\Phi, \varphi, B} = \frac{1}{\Phi^{-1}(1/\mu_B(B))} = \frac{1}{\Phi^{-1}(\varphi(r))}.$$

For functions  $f, g$  on  $\mathbb{R}^n$ , by (4.3) and (4.4), we have

$$(4.6) \quad \frac{1}{|B|\varphi(r)} \int_B |f(x)g(x)| dx \leq 2\|f\|_{\Phi, \varphi, B} \|g\|_{\tilde{\Phi}, \varphi, B}.$$

**Lemma 4.3.** *Let  $\Phi \in \Phi_Y$  and  $\varphi \in \mathcal{G}^{\text{dec}}$ . Then there exists a constant  $C \geq 1$  such that, for any ball  $B = B(a, r)$ ,*

$$(4.7) \quad \frac{1}{\Phi^{-1}(\varphi(r))} \leq \|\chi_B\|_{\text{w}L^{(\Phi, \varphi)}} \leq \|\chi_B\|_{L^{(\Phi, \varphi)}} \leq \frac{C}{\Phi^{-1}(\varphi(r))}.$$

*Proof.* By (4.5) we have

$$\frac{1}{\Phi^{-1}(\varphi(r))} = \|\chi_B\|_{\Phi, \varphi, B, \text{weak}} \leq \|\chi_B\|_{\text{w}L^{(\Phi, \varphi)}} \leq \|\chi_B\|_{L^{(\Phi, \varphi)}}.$$

The last inequality in (4.7) has been proven in [57, Lemma 4.1].  $\square$

**Corollary 4.4.** *Let  $\varphi \in \mathcal{G}^{\text{dec}}$ . Then there exists a constant  $C \geq 1$  such that, for any ball  $B = B(a, r)$ ,*

$$\frac{1}{\varphi(r)} \leq \|\chi_B\|_{\text{w}L^{(1, \varphi)}} \leq \|\chi_B\|_{L^{(1, \varphi)}} \leq \frac{C}{\varphi(r)}.$$

The following relation between Orlicz-Morrey spaces and generalized Morrey spaces is known:

**Lemma 4.5** ([42, Corollary 4.7]). *Let  $\Phi \in \Phi_Y$  and  $\varphi : (0, \infty) \rightarrow (0, \infty)$ . Then there exists a positive constant  $C$  such that, for all  $f \in L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and for all balls  $B = B(a, r)$ ,*

$$(4.8) \quad \int_B |f(x)| dx \leq 2\Phi^{-1}(\varphi(r))\|f\|_{\Phi, \varphi, B}.$$

*Consequently,  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  continuously embeds in  $L^{(1, \Phi^{-1}(\varphi))}(\mathbb{R}^n)$ .*

In the above lemma we do not need to assume  $\varphi \in \mathcal{G}^{\text{dec}}$ , while it was assumed in [42, Corollary 4.7]. Actually, we can prove (4.8) by (4.1), (4.5) and (4.6).

**Remark 4.6.** If  $\Phi \in \Phi_Y$  and  $\varphi \in \mathcal{G}^{\text{dec}}$ , then  $\Phi^{-1}(\varphi) \in \mathcal{G}^{\text{dec}}$ , see [42, Corollary 4.7].

The following lemma shows that any weak Orlicz-Morrey space continuously embeds in a generalized Morrey space, whenever  $\Phi \in \overline{\nabla}_2$ , and it is an extension of [40, Theorem 3.4] which treats weak Orlicz spaces. Note that if  $\Phi \in \overline{\nabla}_2$ , then  $\Phi \approx \Phi_1$  for some  $\Phi_1 \in \nabla_2$ , see Remark 4.2 (iii) and (iv). In this case  $\|f\|_{\Phi, \varphi, B, \text{weak}} \sim \|f\|_{\Phi_1, \varphi, B, \text{weak}}$  and  $\Phi^{-1} \sim \Phi_1^{-1}$ .

**Lemma 4.7.** Let  $\Phi \in \nabla_2$  and  $\varphi : (0, \infty) \rightarrow (0, \infty)$ . Then there exists a positive constant  $C$  such that, for all  $f \in \text{wL}^{(\Phi, \varphi)}(\mathbb{R}^n)$  and for all balls  $B = B(a, r)$ ,

$$\int_B |f(x)| dx \leq C \Phi^{-1}(\varphi(r)) \|f\|_{\Phi, \varphi, B, \text{weak}}.$$

Consequently,  $\text{wL}^{(\Phi, \varphi)}(\mathbb{R}^n)$  continuously embeds in  $L^{(1, \Phi^{-1}(\varphi))}(\mathbb{R}^n)$ .

*Proof.* Case 1:  $\Phi \in \mathcal{Y}^{(1)} \cup \mathcal{Y}^{(2)}$ . We may assume that  $\|f\|_{\Phi, \varphi, B, \text{weak}} = 1$ . If  $b(\Phi) < \infty$  and  $t \in [b(\Phi), \infty)$ , then  $m(f, t) = 0$ . For any ball  $B$  of radius  $r$ , let  $t_0 = \Phi^{-1}(\varphi(r))$ . Then  $\Phi(t_0) = \varphi(r) \in (0, \infty)$  by Remark 4.2 (i). That is,  $t_0 \in (a(\Phi), b(\Phi))$ . Take  $p \in (1, \infty)$  such that  $\Phi(t)/t^p$  is almost increasing, see Remark 4.2 (vi). Then

$$\begin{aligned} \int_B |f(x)| dx &= \int_0^{t_0} m(B, f, t) dt + \int_{t_0}^{b(\Phi)} m(B, f, t) dt \\ &\leq t_0 |B| + \int_{t_0}^{b(\Phi)} \frac{\varphi(r) |B|}{\Phi(t)} dt \\ &= t_0 |B| + \varphi(r) |B| \int_{t_0}^{b(\Phi)} \frac{t^p}{\Phi(t)} t^{-p} dt \\ &\lesssim t_0 |B| + \varphi(r) |B| \frac{t_0^p}{\Phi(t_0)} \int_{t_0}^{b(\Phi)} t^{-p} dt \\ &= t_0 |B| + \frac{t_0 |B| \varphi(r)}{(p-1) \Phi(t_0)} \\ &= t_0 |B| + \frac{t_0 |B|}{(p-1)}. \end{aligned}$$

This shows the conclusion.

Case 2:  $\Phi \in \mathcal{Y}^{(3)}$ . In this case, for any  $\delta \in (0, 1)$ , there exists  $\Phi_1 \in \mathcal{Y}^{(2)}$  such that

$$\Phi_1(\delta t) \leq \Phi(t) \leq \Phi_1(t), \quad t \in [0, \infty],$$

see Remark 4.2 (ii). It follows that

$$\delta \Phi^{-1}(u) \leq \Phi_1^{-1}(u) \leq \Phi^{-1}(u) \quad \text{and} \quad \delta \|f\|_{\text{wL}^{(\Phi_1, \varphi)}} \leq \|f\|_{\text{wL}^{(\Phi, \varphi)}} \leq \|f\|_{\text{wL}^{(\Phi_1, \varphi)}}$$



By Case 1 we have

$$\int_B |f(x)| dx \leq C\Phi_1^{-1}(\varphi(r))\|f\|_{\text{w}L^{(\Phi_1, \varphi)}} \leq C\Phi^{-1}(\varphi(r))\|f\|_{\text{w}L^{(\Phi, \varphi)}}/\delta.$$

Letting  $\delta \rightarrow 1$ , we have the conclusion.  $\square$

**Lemma 4.8** ([42, Lemma 9.4]). *Let  $\Phi \in \Phi_Y$  and  $\varphi \in \mathcal{G}^{\text{dec}}$ , and let  $f \in L^{(\Phi, \varphi)}(\mathbb{R}^n)$ . For a ball  $B = B(a, r)$ , if  $\text{supp } f \cap 2B = \emptyset$ , then*

$$Mf(x) \leq C\Phi^{-1}(\varphi(r))\|f\|_{L^{(\Phi, \varphi)}} \quad \text{for } x \in B,$$

where the constant  $C$  depends only on  $\Phi$  and  $\varphi$ .

**Lemma 4.9.** *Let  $\Phi \in \nabla_2$  and  $\varphi \in \mathcal{G}^{\text{dec}}$ , and let  $f \in L^{(\Phi, \varphi)}(\mathbb{R}^n)$ . For a ball  $B = B(a, r)$ , if  $\text{supp } f \cap 2B = \emptyset$ , then*

$$Mf(x) \leq C\Phi^{-1}(\varphi(r))\|f\|_{\text{w}L^{(\Phi, \varphi)}} \quad \text{for } x \in B,$$

where the constant  $C$  depends only on  $\Phi$  and  $\varphi$ .

*Proof.* For any ball  $B' \ni x$  whose radius is  $s$ , if  $s \leq r/2$ , then  $\int_{B'} |f(x)| dx = 0$ , and, if  $s > r/2$ , then, using Lemma 4.7, we have

$$\begin{aligned} \int_{B'} |f(x)| dx &\lesssim \Phi^{-1}(\varphi(s))\|f\|_{\text{w}L^{(\Phi, \varphi)}} \\ &\lesssim \Phi^{-1}(\varphi(r))\|f\|_{\text{w}L^{(\Phi, \varphi)}}, \end{aligned}$$

since  $r \mapsto \Phi^{-1}(\varphi(r))$  is almost decreasing and satisfies the doubling condition.  $\square$

## 5 Proofs

We first note that, to prove the theorems, we may assume that  $\Phi, \Psi \in \Phi_Y$  instead of  $\Phi, \Psi \in \overline{\Phi}_Y$ . For example, if  $\Phi$  and  $\Psi$  satisfy (3.4) and  $\Phi \approx \Phi_1$ ,  $\Psi \approx \Psi_1$ , then  $\Phi_1$  and  $\Psi_1$  also satisfy (3.4) by the relation (2.9). Moreover,

$$L^{(\Phi, \varphi)}(\mathbb{R}^n) = L^{(\Phi_1, \varphi)}(\mathbb{R}^n), \quad L^{(\Psi, \varphi)}(\mathbb{R}^n) = L^{(\Psi_1, \varphi)}(\mathbb{R}^n),$$

and

$$\text{w}L^{(\Phi, \varphi)}(\mathbb{R}^n) = \text{w}L^{(\Phi_1, \varphi)}(\mathbb{R}^n), \quad \text{w}L^{(\Psi, \varphi)}(\mathbb{R}^n) = \text{w}L^{(\Psi_1, \varphi)}(\mathbb{R}^n),$$

with equivalent quasi norms. Similarly, by Remark 4.2 (iv) we may assume that  $\Phi \in \nabla_2$  instead of  $\Phi \in \overline{\nabla}_2$ . By Remark 2.5 we may also assume that  $\varphi$  is continuous and strictly decreasing.

## 5.1 Proof of Theorem 3.2

To prove Theorem 3.2 we use the following lemma.

**Lemma 5.1** ([61, page 92]). *If  $f \in L^1(\mathbb{R}^n)$ , then*

$$m(Mf, t) \leq \frac{C}{t} \int_{|f|>t/2} |f(x)| dx, \quad t \in (0, \infty).$$

**Proof of Theorem 3.2.** We may assume that  $\Phi \in \nabla_2$  and  $f \geq 0$ . By Remark 4.2 (vi) we can take  $p > 1$  such that  $\Phi(t)/t^p$  is almost increasing. Then,

$$\frac{\Phi(t)}{t^p} \leq C_p \frac{\Phi(s)}{s^p}, \quad t < s.$$

Let

$$N(f) = \sup_{t \in (0, \infty)} \Phi(t)m(2f, t) < \infty \quad \text{and} \quad G_t = \{2f > t\},$$

and let  $g_t(x) = (f(x) - t/2)\chi_{G_t}(x)$  and  $h_t(x) = f(x) - g_t(x)$ . Then  $Mh_t(x) \leq t/2$ , which implies

$$m(Mf, t) \leq m(Mg_t, t/2) + m(Mh_t, t/2) = m(Mg_t, t/2).$$

We consider three cases  $t \in (0, a(\Phi)]$ ,  $t \in (a(\Phi), b(\Phi))$  and  $t \in [b(\Phi), \infty)$ .

Case 1: If  $a(\Phi) > 0$  and  $t \in (0, a(\Phi)]$ , then  $\Phi(t)m(Mf, t) = 0$ .

Case 2: If  $0 \leq a(\Phi) < b(\Phi) \leq \infty$  and  $t \in (a(\Phi), b(\Phi))$ , then  $|G_t| = m(2f, t) < \infty$ . Hence

$$\begin{aligned} 2 \int_{\mathbb{R}^n} g_t(x) dx &= \int_{G_t} (2f(x) - t) dx \\ &= \int_0^t m(G_t, 2f, s) ds + \int_t^{b(\Phi)} m(G_t, 2f, s) ds - t|G_t| \\ &\leq \int_t^{b(\Phi)} m(2f, s) ds = \frac{1}{\Phi(t)} \int_t^{b(\Phi)} \frac{\Phi(t)}{\Phi(s)} \Phi(s)m(2f, s) ds \\ &\leq \frac{C_p N(f)}{\Phi(t)} \int_t^{b(\Phi)} \left(\frac{t}{s}\right)^p ds \leq \frac{C_p N(f)t}{(p-1)\Phi(t)} < \infty. \end{aligned}$$

That is,  $g_t \in L^1(\mathbb{R}^n)$  for  $t \in (a(\Phi), b(\Phi))$ . By Lemma 5.1 we have

$$\Phi(t)m(Mf, t) \leq \Phi(t)m(Mg_t, t/2) \leq \frac{2C\Phi(t)}{t} \int_{\mathbb{R}^n} g_t(x) dx \leq \frac{CC_p}{p-1} N(f).$$

Case 3: If  $b(\Phi) < \infty$  and  $t \in [b(\Phi), \infty)$ , then  $|G_t| = m(2f, t) = 0$  and  $g_t = 0$  a.e. Hence  $Mg_t = 0$  and

$$\Phi(t)m(Mf, t) \leq \Phi(t)m(Mg_t, t/2) = 0.$$

In conclusion, for some  $C' \geq 1$  and all  $t \in (0, \infty)$ ,

$$\begin{aligned} \Phi(t)m(Mf, t) &\leq C'N(f) \\ &\leq \sup_{s \in (0, \infty)} \Phi(C's)m(2f, s) \\ &= \sup_{s \in (0, \infty)} \Phi(s)m(2C'f, s). \end{aligned}$$

This shows the conclusion.  $\square$

## 5.2 Proof of Theorem 3.3

In this subsection we prove Theorem 3.3 by using Theorems 3.1 and 3.2.

**Proof of Theorem 3.3.** We may assume that  $\Phi \in \Phi_Y$ . Let  $f \in L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $\|f\|_{L^{(\Phi, \varphi)}} = 1$ . To prove the norm inequality  $\|Mf\|_{wL^{(\Phi, \varphi)}} \leq C_0$ , it is enough to prove that, for any ball  $B = B(a, r)$ ,

$$(5.1) \quad \|Mf\|_{\Phi, \varphi, B, \text{weak}} \leq C_0.$$

Let  $f = f_1 + f_2$ ,  $f_1 = f\chi_{2B}$ . Then we have

$$(5.2) \quad \int_{\mathbb{R}^n} \Phi(|f_1(x)|) dx \leq \int_{2B} \Phi\left(\frac{|f(x)|}{\|f\|_{\Phi, \varphi, 2B}}\right) dx \leq |2B|\varphi(2r) \leq C_{n, \varphi}|B|\varphi(r),$$

for some constant  $C_{n, \varphi} \geq 1$  by the doubling condition of  $\varphi$ . Hence, (3.1) yields

$$\begin{aligned} \sup_{t \in (0, \infty)} \Phi(t) m\left(B, \frac{Mf_1}{C_{n, \varphi} C_\Phi}, t\right) &\leq \frac{1}{C_{n, \varphi}} \sup_{t \in (0, \infty)} \Phi(t) m\left(\frac{Mf_1}{C_\Phi}, t\right) \\ &\leq \frac{1}{C_{n, \varphi}} \int_{\mathbb{R}^n} \Phi(|f_1(x)|) dx \leq |B|\varphi(r). \end{aligned}$$

Next, since  $\text{supp } f_2 \cap 2B = \emptyset$ , by Lemma 4.8 we have  $Mf_2(x) \leq C\Phi^{-1}(\varphi(r))$  for  $x \in B$ . Then

$$\sup_{t \in (0, \infty)} \Phi(t) m\left(B, \frac{Mf_2}{C}, t\right) \leq \int_B \Phi\left(\frac{Mf_2(x)}{C}\right) dx \leq \int_B \varphi(r) dx = |B|\varphi(r).$$

In the above we use (2.8) for the second inequality. Therefore,

$$\|Mf\|_{\Phi, \varphi, B, \text{weak}} \leq 2(\|Mf_1\|_{\Phi, \varphi, B, \text{weak}} + \|Mf_2\|_{\Phi, \varphi, B, \text{weak}}) \leq 2(C_{n, \varphi} C_\Phi + C),$$

which shows (5.1). If  $\Phi \in \nabla_2$ , then by the same way we have the norm inequality  $\|Mf\|_{L^{(\Phi, \varphi)}} \leq C_0$ , using (3.2) instead of (3.1).

Next, let  $f \in \text{w}L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $\|f\|_{\text{w}L^{(\Phi, \varphi)}} = 1$ . To prove the norm inequality  $\|Mf\|_{\text{w}L^{(\Phi, \varphi)}} \leq C_0$ , it is enough to prove (5.1) for any ball  $B = B(a, r)$ . Let  $f = f_1 + f_2$ ,  $f_1 = f\chi_{2B}$ . Then we have

$$\begin{aligned} \sup_{t \in (0, \infty)} \Phi(t)m(f_1, t) &\leq \sup_{t \in (0, \infty)} \Phi(t)m\left(2B, \frac{f}{\|f\|_{\Phi, \varphi, 2B, \text{weak}}}, t\right) \\ &\leq |2B|\varphi(2r) \leq C_{n, \varphi}|B|\varphi(r), \end{aligned}$$

instead of (5.2). Hence, (3.3) yields

$$\begin{aligned} \sup_{t \in (0, \infty)} \Phi(t) m\left(B, \frac{Mf_1}{C_{n, \varphi}C_\Phi}, t\right) &\leq \frac{1}{C_{n, \varphi}} \sup_{t \in (0, \infty)} \Phi(t) m\left(\frac{Mf_1}{C_\Phi}, t\right) \\ &\leq \frac{1}{C_{n, \varphi}} \sup_{t \in (0, \infty)} \Phi(t)m(f_1, t) \leq |B|\varphi(r). \end{aligned}$$

We also have  $Mf_2(x) \leq C\Phi^{-1}(\varphi(r))$  for  $x \in B$  by Lemma 4.9. Then

$$\sup_{t \in (0, \infty)} \Phi(t) m\left(B, \frac{Mf_2}{C}, t\right) \leq \sup_{t \in (0, \infty)} \Phi(t) m\left(B, \Phi^{-1}(\varphi(r)), t\right) \leq |B|\varphi(r).$$

Therefore, we have the conclusion.  $\square$

### 5.3 Proof of Theorem 3.4

We may assume that  $\Phi, \Psi \in \Phi_Y$  and that  $\varphi$  is continuous and strictly decreasing. First we state a lemma.

**Lemma 5.2** ([4, Proposition 1]). *Let  $\rho, \tau : (0, \infty) \rightarrow (0, \infty)$ . Assume that  $\rho$  satisfies (1.5) and that  $\tau$  satisfies the doubling condition (2.12). Define*

$$(5.3) \quad \tilde{\rho}(r) = \int_{K_1 r}^{K_2 r} \frac{\rho(t)}{t} dt, \quad r \in (0, \infty).$$

*Then there exists a positive constant  $C$  such that, for all  $r \in (0, \infty)$ ,*

$$(5.4) \quad \sum_{j=-\infty}^{-1} \tilde{\rho}(2^j r) \lesssim \int_0^{K_2 r} \frac{\rho(t)}{t} dt,$$

$$(5.5) \quad \sum_{j=0}^{\infty} \tilde{\rho}(2^j r)\tau(2^j r) \lesssim \int_{K_1 r}^{\infty} \frac{\rho(t)\tau(t)}{t} dt.$$

**Proof of Theorem 3.4** (i). By the assumption (2.14) we may assume that  $\varphi$  is bijective from  $(0, \infty)$  to itself. If  $\Phi^{-1}(0) > 0$ , then, from (3.4) it follows that

$$0 < \int_0^{\infty} \frac{\rho(t)}{t} dt \Phi^{-1}(0) \lesssim \Psi^{-1}(0),$$

since  $\lim_{r \rightarrow \infty} \varphi(r) = 0$ . Let  $f \in L^{(\Phi, \varphi)}(\mathbb{R}^n)$  and  $\|f\|_{L^{(\Phi, \varphi)}} = 1$ , and let  $x \in \mathbb{R}^n$ . We may assume that

$$0 < \frac{Mf(x)}{C_0} < \infty \quad \text{and} \quad 0 \leq \Phi\left(\frac{Mf(x)}{C_0}\right) < \infty,$$

otherwise there is nothing to prove.

If  $\Phi(Mf(x)/C_0) = 0$ , then, by (2.8) we have

$$\frac{Mf(x)}{C_0} \leq \Phi^{-1}(0) = \sup\{u \geq 0 : \Phi(u) = 0\}.$$

Hence, using (5.4), we have

$$\begin{aligned} |I_\rho f(x)| &\leq \sum_{j=-\infty}^{\infty} \int_{2^j \leq |x-y| < 2^{j+1}} \frac{\rho(|x-y|)}{|x-y|^n} |f(x)| dx \\ &\lesssim \sum_{j=-\infty}^{\infty} \frac{\tilde{\rho}(2^j)}{2^{jn}} \int_{|x-y| < 2^{j+1}} |f(y)| dy \lesssim \int_0^\infty \frac{\rho(t)}{t} dt Mf(x) \\ &\leq C_0 \int_0^\infty \frac{\rho(t)}{t} dt \Phi^{-1}(0) \lesssim \Psi^{-1}(0) \lesssim \Psi^{-1}\left(\Phi\left(\frac{Mf(x)}{C_0}\right)\right), \end{aligned}$$

which shows (3.5).

If  $\Phi(Mf(x)/C_0) > 0$ , choose  $r \in (0, \infty)$  such that

$$(5.6) \quad \varphi(r) = \Phi\left(\frac{Mf(x)}{C_0}\right),$$

and let

$$\begin{aligned} J_1 &= \sum_{j=-\infty}^{-1} \frac{\tilde{\rho}(2^j r)}{(2^j r)^n} \int_{|x-y| < 2^{j+1} r} |f(y)| dy, \\ J_2 &= \sum_{j=0}^{\infty} \frac{\tilde{\rho}(2^j r)}{(2^j r)^n} \int_{|x-y| < 2^{j+1} r} |f(y)| dy. \end{aligned}$$

Then

$$|I_\rho f(x)| \lesssim J_1 + J_2.$$

By (5.6) and (2.8) we have  $Mf(x) \leq C_0 \Phi^{-1}(\varphi(r))$ . Then, using (5.4), we have

$$J_1 \lesssim \int_0^{K_2 r} \frac{\rho(t)}{t} dt Mf(x) \lesssim \int_0^{K_2 r} \frac{\rho(t)}{t} dt \Phi^{-1}(\varphi(r)).$$

Next, by Lemma 4.5,  $\|f\|_{L^{(\Phi, \varphi)}} \leq 1$  and (5.5) we have

$$J_2 \lesssim \sum_{j=0}^{\infty} \tilde{\rho}(2^j r) \Phi^{-1}(\varphi(2^{j+1} r)) \lesssim \int_{K_1 r}^{\infty} \frac{\rho(t) \Phi^{-1}(\varphi(t))}{t} dt.$$

Then by (3.4), the doubling condition of  $\Phi^{-1}(\varphi(r))$  and  $\Psi^{-1}(\varphi(r))$ , and (5.6), we have

$$\begin{aligned} J_1 + J_2 &\lesssim \int_0^{K_2 r} \frac{\rho(t)}{t} dt \Phi^{-1}(\varphi(r)) + \int_{K_1 r}^{\infty} \frac{\rho(t)\Phi^{-1}(\varphi(t))}{t} dt \\ &\lesssim \frac{\Psi^{-1}(\varphi(K_2 r))}{\Phi^{-1}(\varphi(K_2 r))} \Phi^{-1}(\varphi(r)) + \Psi^{-1}(\varphi(K_1 r)) \\ &\sim \Psi^{-1}(\varphi(r)) = \Psi^{-1}\left(\Phi\left(\frac{Mf(x)}{C_0}\right)\right). \end{aligned}$$

Combining this inequality with (2.8), we have (3.5).

The proof of (3.6) is almost the same as one of (3.5). The only difference is that we use Lemma 4.7 instead of Lemma 4.5 to estimate  $J_2$ .  $\square$

To prove Theorem 3.4 (ii) we state three lemmas.

**Lemma 5.3** ([6, Lemma 2.1]). *There exists a positive constant  $C$  such that, for all  $R > 0$ ,*

$$\int_0^{R/2} \frac{\rho(t)}{t} dt \chi_{B(0, R/2)}(x) \leq CI_\rho \chi_{B(0, R)}(x), \quad x \in \mathbb{R}^n.$$

The following lemma is an extension of [6, Lemma 2.4] and gives a typical element in  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$ .

**Lemma 5.4.** *For  $\Phi \in \Phi_Y$  and  $\varphi \in \mathcal{G}^{\text{dec}}$ , let  $g(x) = \Phi^{-1}(\varphi(|x|))$ . If  $\varphi$  satisfies (3.8), then  $g \in L^{(\Phi, \varphi)}(\mathbb{R}^n)$ .*

*Proof.* By Remark 2.5 we may assume that  $\varphi$  is decreasing. In this case  $x \mapsto \varphi(|x|)$  is radial decreasing, so that, for any ball  $B(a, r)$ ,

$$\int_{B(a, r)} \varphi(|x|) dx \leq \int_{B(0, r)} \varphi(|x|) dx \sim \frac{1}{r^n} \int_0^r \varphi(t) t^{n-1} dt \lesssim \varphi(r).$$

In the above we used (3.8) for the last inequality. Then, taking a suitable constant  $C_g \geq 1$ , and using the convexity of  $\Phi$  and (2.8), we have

$$\begin{aligned} \frac{1}{\varphi(r)} \int_{B(a, r)} \Phi\left(\frac{\Phi^{-1}(\varphi(|x|))}{C_g}\right) dx &\leq \frac{1}{C_g \varphi(r)} \int_{B(a, r)} \Phi(\Phi^{-1}(\varphi(|x|))) dx \\ &\leq \frac{1}{C_g \varphi(r)} \int_{B(a, r)} \varphi(|x|) dx \leq 1. \end{aligned}$$

This shows  $g \in L^{(\Phi, \varphi)}(\mathbb{R}^n)$  with  $\|g\|_{L^{(\Phi, \varphi)}} \leq C_g$ .  $\square$

**Lemma 5.5.** Let  $\Phi \in \Phi_Y$  and  $\varphi \in \mathcal{G}^{\text{dec}}$ . For  $R > 0$ , let

$$g_R(x) = \Phi^{-1}(\varphi(|x|))\chi_{\mathbb{R}^n \setminus B(0,R)}(x).$$

Then there exists a positive constant  $C$  such that, for all  $R > 0$ ,

$$\int_{2R}^{\infty} \frac{\rho(t)\Phi^{-1}(\varphi(t))}{t} dt \chi_{B(0,R)}(x) \leq CI_{\rho}g_R(x).$$

*Proof.* Let  $x \in B(0, R)$ . Then  $B(0, R) \subset B(x, 2R)$  and  $|x - y| \sim |y|$  for all  $y \notin B(0, 2R)$ . Hence

$$\begin{aligned} I_{\rho}g_R(x) &= \int_{\mathbb{R}^n \setminus B(0,R)} \frac{\rho(|x-y|)\Phi^{-1}(\varphi(|y|))}{|x-y|^n} dy \\ &\geq \int_{\mathbb{R}^n \setminus B(x,2R)} \frac{\rho(|x-y|)\Phi^{-1}(\varphi(|y|))}{|x-y|^n} dy \\ &= \int_{\mathbb{R}^n \setminus B(0,2R)} \frac{\rho(|y|)\Phi^{-1}(\varphi(|x-y|))}{|y|^n} dy \\ &\sim \int_{2R}^{\infty} \frac{\rho(t)\Phi^{-1}(\varphi(t))}{t} dt. \end{aligned}$$

This shows the conclusion.  $\square$

**Proof of Theorem 3.4 (ii).** Firstly, by Lemma 5.3 and the boundedness of  $I_{\rho}$ , we have

$$\int_0^r \frac{\rho(t)}{t} dt \|\chi_{B(0,r)}\|_{\text{w}L(\Psi,\varphi)} \lesssim \|I_{\rho}\chi_{B(0,2r)}\|_{\text{w}L(\Psi,\varphi)} \lesssim \|\chi_{B(0,2r)}\|_{L(\Phi,\varphi)}.$$

By Lemma 4.3 and the doubling condition of  $\Phi^{-1}(\varphi(r))$  we have

$$\int_0^r \frac{\rho(t)}{t} dt \Phi^{-1}(\varphi(r)) \lesssim \Psi^{-1}(\varphi(r)).$$

Secondly, under the assumption (3.8), let  $g$  and  $g_R$  be functions as in Lemmas 5.4 and 5.5, respectively. Then by the boundedness of  $I_{\rho}$  we obtain

$$\int_r^{\infty} \frac{\rho(t)\Phi^{-1}(\varphi(t))}{t} dt \|\chi_{B(0,r/2)}\|_{\text{w}L(\Psi,\varphi)} \lesssim \|I_{\rho}g_{r/2}\|_{\text{w}L(\Psi,\varphi)} \lesssim \|g_{r/2}\|_{L(\Phi,\varphi)} \lesssim \|g\|_{L(\Phi,\varphi)}.$$

By Lemma 4.3 and the doubling condition of  $\Psi^{-1}(\varphi(r))$  we have

$$\int_r^{\infty} \frac{\rho(t)\Phi^{-1}(\varphi(t))}{t} dt \lesssim \Psi^{-1}(\varphi(r)).$$

Thus, we obtain the conclusion.  $\square$

**Proof of Theorem 3.4 (iii).** By Lemma 4.3, Corollary 4.4 and Remark 4.6 we have

$$\|\chi_{B(0,r)}\|_{\text{w}L(\Psi,\varphi)} \sim \|\chi_{B(0,r)}\|_{L(1,\Psi^{-1}(\varphi))} \sim \frac{1}{\Psi^{-1}(\varphi(r))}.$$

Then we can replace  $\|\chi_{B(0,r)}\|_{\text{w}L(\Psi,\varphi)}$  by  $\|\chi_{B(0,r)}\|_{L(1,\Psi^{-1}(\varphi))}$  in the proof of (ii).  $\square$

## 5.4 Proof of Theorem 3.5

We may assume that  $\Phi, \Psi \in \Phi_Y$  and that  $\varphi$  is continuous and strictly decreasing.

**Proof of Theorem 3.5 (i).** The pointwise estimate (3.10) was already proven in [57, Theorem 5.1]. The pointwise estimate (3.11) can be proven by almost the same way as (3.10). To prove (3.11) we use

$$\int_B |f(x)| dx \leq C\Phi^{-1}(\varphi(r))\|f\|_{\text{w}L^{(\Phi, \varphi)}}$$

by Lemma 4.7 instead of

$$\int_B |f(x)| dx \leq C\Phi^{-1}(\varphi(r))\|f\|_{L^{(\Phi, \varphi)}}.$$

Other parts are the same as the proof of (3.10). □

For the proof of Theorem 3.5 (ii) we use the following lemma.

**Lemma 5.6** ([56, Lemma 5.1]). *Let  $\rho : (0, \infty) \rightarrow (0, \infty)$ . Then, for all  $r \in (0, \infty)$ ,*

$$(5.7) \quad \left( \sup_{0 < t \leq r} \rho(t) \right) \chi_{B(0,r)}(x) \leq (M_\rho \chi_{B(0,r)})(x), \quad x \in \mathbb{R}^n.$$

**Proof of Theorem 3.5 (ii).** By Lemma 5.6 and the boundedness of  $M_\rho$  from  $L^{(\Phi, \varphi)}(\mathbb{R}^n)$  to  $\text{w}L^{(\Psi, \varphi)}(\mathbb{R}^n)$  we have

$$\left( \sup_{0 < t \leq r} \rho(t) \right) \|\chi_{B(0,r)}\|_{\text{w}L^{(\Psi, \varphi)}} \leq \|M_\rho \chi_{B(0,r)}\|_{\text{w}L^{(\Psi, \varphi)}} \lesssim \|\chi_{B(0,r)}\|_{L^{(\Phi, \varphi)}}.$$

Then, by Lemma 4.3 we have the conclusion. □

**Proof of Theorem 3.5 (iii).** The same as Proof of Theorem 3.4 (iii). □

## Acknowledgement

The second author was supported by Grant-in-Aid for Scientific Research (B), No. 15H03621, Japan Society for the Promotion of Science.

## References

- [1] D. R. Adams, A note on Riesz potentials, Duke Math. J. 42 (1975), No. 4, 765–778.



- [2] F. Chiarenza and M. Frasca, Morrey spaces and Hardy-Littlewood maximal function, *Rend. Mat. Appl.* (7) 7 (1987), No. 3-4, 273–279.
- [3] A. Cianchi, Strong and weak type inequalities for some classical operators in Orlicz spaces, *J. London Math. Soc.* (2) 60 (1999), No. 1, 187–202.
- [4] F. Deringoz, V.S. Guliyev, E. Nakai, Y. Sawano and M. Shi, Generalized fractional maximal and integral operators on Orlicz and generalized Orlicz-Morrey spaces of the third kind, *Positivity* 23 (2019), No. 3, 727–757.
- [5] F. Deringoz, V.S. Guliyev and S. Samko, Boundedness of the maximal and singular operators on generalized Orlicz-Morrey spaces. *Operator theory, operator algebras and applications*, 139–158, *Oper. Theory Adv. Appl.*, 242, Birkhäuser/Springer, Basel, 2014.
- [6] Eridani, H. Gunawan, E. Nakai and Y. Sawano, Characterizations for the generalized fractional integral operators on Morrey spaces, *Math. Inequal. Appl.* 17 (2014), No. 2, 761–777.
- [7] L. C. F. Ferreira, On a bilinear estimate in weak-Morrey spaces and uniqueness for Navier-Stokes equations, *J. Math. Pures Appl.* (9) 105 (2016), No. 2, 228–247.
- [8] S. Gala, Y. Sawano and H. Tanaka, A remark on two generalized Orlicz-Morrey spaces, *J. Approx. Theory* 198 (2015), 1–9.
- [9] D. Gallardo, Orlicz spaces for which the Hardy-Littlewood maximal operator is bounded, *Publ. Mat.* 32 (1988), No. 2, 261–266.
- [10] A. Gogatishvili, R. Mustafayev and M. Agcayazi, Weak-type estimates in Morrey spaces for maximal commutator and commutator of maximal function, *Tokyo J. Math.* 41 (2018), No. 1, 193–218.
- [11] V. S. Guliyev, S. G. Hasanov, Y. Sawano and T. Noi, Non-smooth atomic decompositions for generalized Orlicz-Morrey spaces of the third kind, *Acta Appl. Math.* 145 (2016), No. 1, 133–174.
- [12] H. Gunawan, A note on the generalized fractional integral operators, *J. Indones. Math. Soc.* 9 (2003), No. 1, 39–43.
- [13] H. Gunawan, D. I. Hakim, K. M. Limanta and A. A. Masta, Inclusion properties of generalized Morrey spaces, *Math. Nachr.* 290 (2017), No. 2-3, 332–340.
- [14] H. Gunawan, D. I. Hakim, E. Nakai and Y. Sawano. On inclusion relation between weak Morrey spaces and Morrey spaces, *Nonlinear Anal.* 168 (2018), 27–31.

- [15] D. I. Hakim and Y. Sawano, Interpolation of generalized Morrey spaces, *Rev. Mat. Complut.* 29 (2016), No. 2, 295–340.
- [16] N. Hatano, Boundedness of the Calderón-Zygmund operators and their commutators on Morrey-Lorentz spaces, preprint.
- [17] K.-P. Ho, Vector-valued maximal inequalities on weighted Orlicz-Morrey spaces, *Tokyo J. Math.* 36 (2013), No. 2, 499–512.
- [18] K.-P. Ho, Vector-valued operators with singular kernel and Triebel-Lizorkin block spaces with variable exponents, *Kyoto J. Math.* 56 (2016), No. 1, 97–124.
- [19] K.-P. Ho, Atomic decompositions and Hardy’s inequality on weak Hardy-Morrey spaces, *Sci. China Math.* 60 (2017), No. 3, 449–468.
- [20] K.-P. Ho, Weak type estimates of singular integral operators on Morrey-Banach spaces, *Integral Equations Operator Theory* 91 (2019), No. 3, Paper No. 20.
- [21] Y. Jiao, L. H. Peng and P. D. Liu, Interpolation for weak Orlicz spaces with  $M_\Delta$  condition. *Sci. China Ser. A* 51 (2008), No. 11, 2072–2080.
- [22] R. Kawasumi and E. Nakai, Pointwise multipliers on weak Orlicz spaces, *Hiroshima Math. J.* 50 (2020), No. 2, 169–184.
- [23] H. Kita, On maximal functions in Orlicz spaces, *Proc. Amer. Math. Soc.* 124 (1996), No. 10, 3019–3025.
- [24] H. Kita, On Hardy-Littlewood maximal functions in Orlicz spaces, *Math. Nachr.* 183 (1997), 135–155.
- [25] H. Kita, Orlicz spaces and their applications (Japanese), Iwanami Shoten, Publishers. Tokyo, 2009.
- [26] V. Kokilashvili and M. Krbeč, Weighted inequalities in Lorentz and Orlicz spaces, World Scientific Publishing Co., Inc., River Edge, NJ, 1991.
- [27] S. G. Kreĭn, Yu. Ī. Petunĭn and E. M. Semĕnov, Interpolation of linear operators, Translated from the Russian by J. Szűcs, Translations of Mathematical Monographs, 54, American Mathematical Society, Providence, R.I., 1982.
- [28] M. A. Krasnosel’skiĭ and Ja. B. Rutickiĭ, Convex functions and Orlicz spaces, Translated from the first Russian edition by Leo F. Boron, P. Noordhoff Ltd., Groningen 1961.
- [29] H. Li, Hardy-type inequalities on strong and weak Orlicz-Lorentz spaces, *Sci. China Math.* 55 (2012), No. 12, 2493–2505.

- [30] Y. Liang, D. Yang and R. Jiang, Weak Musielak-Orlicz Hardy spaces and applications, *Math. Nachr.* 289 (2016), No. 5-6, 634–677.
- [31] P. Liu and M. Wang, Weak Orlicz spaces: Some basic properties and their applications to harmonic analysis, *Sci. China Math.* 56 (2013), 789–802.
- [32] L. Maligranda, Orlicz spaces and interpolation, *Seminars in mathematics* 5, Departamento de Matemática, Universidade Estadual de Campinas, Brasil, 1989.
- [33] T. Mizuhara, Boundedness of some classical operators on generalized Morrey spaces, *Harmonic analysis (Sendai, 1990)*, 183–189, ICM-90 Satell. Conf. Proc., Springer, Tokyo, 1991.
- [34] Y. Mizuta, E. Nakai, T. Ohno and T. Shimomura, Boundedness of fractional integral operators on Morrey spaces and Sobolev embeddings for generalized Riesz potentials, *J. Math. Soc. Japan* 62 (2010), No. 3, 707–744.
- [35] C. B. Morrey, On the solutions of quasi-linear elliptic partial differential equations, *Trans. Amer. Math. Soc.* 43 (1938), No. 1, 126–166.
- [36] E. Nakai, Hardy-Littlewood maximal operator, singular integral operators and the Riesz potentials on generalized Morrey spaces, *Math. Nachr.* 166 (1994), 95–103.
- [37] E. Nakai, On generalized fractional integrals in the Orlicz spaces, *Proceedings of the Second ISAAC Congress, Vol. 1 (Fukuoka, 1999)*, 75–81, *Int. Soc. Anal. Appl. Comput.*, 7, Kluwer Acad. Publ., Dordrecht, 2000.
- [38] E. Nakai, On generalized fractional integrals, *Taiwanese J. Math.* 5 (2001), 587–602.
- [39] E. Nakai, On generalized fractional integrals in the Orlicz spaces on spaces of homogeneous type, *Sci. Math. Jpn.* 54 (2001), No. 3, 473–487.
- [40] E. Nakai, On generalized fractional integrals on the weak Orlicz spaces,  $BMO_\phi$ , the Morrey spaces and the Campanato spaces, *Function spaces, interpolation theory and related topics (Lund, 2000)*, 389–401, de Gruyter, Berlin, 2002.
- [41] E. Nakai, Generalized fractional integrals on Orlicz-Morrey spaces, *Banach and Function Spaces*, 323–333, Yokohama Publ., Yokohama, 2004.
- [42] E. Nakai, Orlicz-Morrey spaces and the Hardy-Littlewood maximal function, *Studia Math.* 188 (2008), No 3, 193–221.

- [43] E. Nakai, Calderón-Zygmund operators on Orlicz-Morrey spaces and modular inequalities, Banach and function spaces II, 393–410, Yokohama Publ., Yokohama, 2008.
- [44] E. Nakai and H. Sumitomo, On generalized Riesz potentials and spaces of some smooth functions, Sci. Math. Jpn. 54 (2001), No. 3, 463–472.
- [45] R. O’Neil, Fractional integration in Orlicz spaces. I., Trans. Amer. Math. Soc. 115 (1965), 300–328.
- [46] W. Orlicz, Über eine gewisse Klasse von Räumen vom Typus B, Bull. Acad. Polonaise A (1932), 207–220; reprinted in his Collected Papers, PWN, Warszawa 1988, 217–230.
- [47] W. Orlicz, Über Räume ( $L^M$ ), Bull. Acad. Polonaise A (1936), 93–107; reprinted in his Collected Papers, PWN, Warszawa 1988, 345–359.
- [48] J. Peetre, On convolution operators leaving  $L^{p,\lambda}$  spaces invariant, Ann. Mat. Pura Appl. (4) 72 (1966), 295–304.
- [49] J. Peetre, On the theory of  $\mathcal{L}_{p,\lambda}$  spaces, J. Funct. Anal. 4 (1969), 71–87.
- [50] C. Pérez, Two weighted inequalities for potential and fractional type maximal operators, Indiana Univ. Math. J. 43 (1994), No. 2, 663–683.
- [51] M. M. Rao and Z. D. Ren, Theory of Orlicz Spaces, Marcel Dekker, Inc., New York, Basel and Hong Kong, 1991.
- [52] Y. Sawano, Singular integral operators acting on Orlicz-Morrey spaces of the first kind, Nonlinear Stud. 26 (2019), No. 4, 895–910.
- [53] Y. Sawano and S. R. El-Shabrawy, Weak Morrey spaces with applications, Math. Nachr. 291 (2018), No. 1, 178–186.
- [54] Y. Sawano, S. Sugano and H. Tanaka, Generalized fractional integral operators and fractional maximal operators in the framework of Morrey spaces, Trans. Amer. Math. Soc. 363 (2011), No. 12, 6481–6503.
- [55] Y. Sawano, S. Sugano and H. Tanaka, Orlicz-Morrey spaces and fractional operators, Potential Anal. 36 (2012), No. 4, 517–556.
- [56] M. Shi, R. Arai and E. Nakai, Generalized fractional integral operators and their commutators with functions in generalized Campanato spaces on Orlicz spaces, Taiwanese J. Math. 23 (2019), No. 6, 1339–1364.
- [57] M. Shi, R. Arai and E. Nakai, Commutators of integral operators with functions in Campanato spaces on Orlicz-Morrey spaces, Banach J. Math. Anal. 15 (2021), Paper No. 22, 41 pp.

- [58] I. Sihwaningrum and Y. Sawano, Weak and strong type estimates for fractional integral operators on Morrey spaces over metric measure spaces, *Eurasian Math. J.* 4 (2013), No. 1, 76–81.
- [59] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, No. 30, Princeton University Press, Princeton, N.J. 1970.
- [60] A. Torchinsky, Interpolation of operations and Orlicz classes, *Studia Math.* 59 (1976), No. 2, 177–207.
- [61] A. Torchinsky, *Real-variable methods in harmonic analysis*, Academic Press, New York, 1986.
- [62] O. D. Tsereteli (Cereteli), The interpolation of operators by the truncation method, (Russian) *Sakharth. SSR Mecn. Akad. Math. Inst. Šrom.* 36 (1969), 111–122.
- [63] N. K. Tumalun and H. Gunawan, Morrey spaces are embedded between weak Morrey spaces and Stummel classes, *J. Indones. Math. Soc.* 25 (2019), No. 3, 203–209.
- [64] G. Xie and D. Yang, Atomic characterizations of weak martingale Musielak-Orlicz Hardy spaces and their applications, *Banach J. Math. Anal.* 13 (2019), No. 4, 884–917.

Ryota Kawasumi  
 Minohara 1-6-3 (B-2), Misawa, Aomori 033-0033, Japan  
 rykawasumi@gmail.com

Eiichi Nakai  
 Department of Mathematics  
 Ibaraki University  
 Mito, Ibaraki 310-8512, Japan  
 eiichi.nakai.math@vc.ibaraki.ac.jp

Minglei Shi  
 Jiangxi University of Engineering  
 Xinyu, Jiangxi 338000, China  
 shimingleiyy@163.com