

INTERPOLATION PROPERTIES OF CERTAIN CLASSES OF NET SPACES

A. H. KALIDOLDAY, E. D. NURSULTANOV

ABSTRACT. The paper studies the interpolation properties of net spaces $N_{p,q}(M)$, when M is the set of dyadic cubes in \mathbb{R}^n , and also when M is the family of all cubes with parallel faces to the coordinate axes in \mathbb{R}^n . It is shown that, in the case when M is the set of dyadic cubes the scale of spaces is closed with respect to the real interpolation method. In the case, when M is the set of all cubes with parallel faces to the coordinate axes, an analogue of the Marcinkiewicz-Calderon theorem on cones of non-negative functions is given.

1. INTRODUCTION

Let in \mathbb{R}^n is given n -dimensional Lebesgue measure μ , M is the set of all cubes in \mathbb{R}^n . Further M will be called as a "net". For function $f(x)$, defined and integrable on each e from M , we define the function

$$\bar{f}(t, M) = \sup_{\substack{e \in M \\ |e| \geq t}} \frac{1}{|e|} \left| \int_e f(x) dx \right|, \quad t > 0,$$

where the supremum is taken over all $e \in M$, whose measure is $|e| \stackrel{\text{def}}{=} \mu e \geq t$. In the case when $\sup\{|e|: e \in M\} = \alpha < \infty$ and $t > \alpha$ assuming that $\bar{f}(t, M) = 0$.

Let p, q parameters satisfy the conditions $0 < p \leq \infty$, $0 < q \leq \infty$. Let's define the net spaces $N_{p,q}(M)$, as a set of all functions f , such that for $q < \infty$

$$\|f\|_{N_{p,q}(M)} = \left(\int_0^\infty \left(t^{\frac{1}{p}} \bar{f}(t, M) \right)^q \frac{dt}{t} \right)^{\frac{1}{q}} < \infty,$$

and for $q = \infty$

$$\|f\|_{N_{p,\infty}(M)} = \sup_{t>0} t^{\frac{1}{p}} \bar{f}(t, M) < \infty.$$

These spaces were introduced in the work [N1].

Net spaces have found important applications in various problems of harmonic analysis, operator theory and the theory of stochastic processes [AR1, AR2, ARN, NK, NT1, N2, N3, NT, NA].

In this paper, we study the interpolation properties of these spaces. It should be noted here, that net spaces are in a sense close to the Morrey space:

$$M_p^\alpha = \left\{ f : \sup_{y \in \mathbb{R}^n, t > 0} t^{-\lambda} \left(\int_{|x+y| \leq t} |f(x)|^p dx \right)^{\frac{1}{p}} < \infty \right\}.$$

In the case when $f(x) \geq 0$, for $\frac{1}{p} = 1 - \frac{\lambda}{n}$

$$\|f\|_{N_{p,\infty}(M)} \asymp \|f\|_{M_1^\lambda}.$$

2020 *Mathematics Subject Classification*: 46B70.

Key words and phrases: Net spaces, Interpolation properties of net spaces, Marcinkiewicz-Calderon type interpolation theorem.

This research was supported by the Ministry of Education and Science of the Republic of Kazakhstan (projects no. AP08956157 and AP08856479).

The question of interpolation of Morrey spaces was considered in the works [St, CM, P, RV, BRV, LR2]. It follows from the results of [P] that

$$(M_p^{\lambda_0}, M_p^{\lambda_1})_{\theta, \infty} \hookrightarrow M_p^\lambda,$$

where $\lambda = (1 - \theta)\lambda_0 + \theta\lambda_1$. In the works [RV, BRV] it was established that this inclusion is strict.

For net spaces $N_{p,q}(M)$, where M is an arbitrary system of measurable sets from \mathbb{R}^2 , we also have an embedding (see [N1, Theorem 1])

$$(1) \quad (N_{p_0, q_0}(M), N_{p_1, q_1}(M))_{\theta, q} \hookrightarrow N_{p, q}(M),$$

where $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$, $0 < \theta < 1$, $0 < q \leq \infty$.

From (1) it follows that if the linear operator T bounded from A_i to $N_{p_i, \infty}(M)$, $i = 0, 1$, then the operator T bounded from $A_{\theta, q}$ to $N_{p, q}(M)$, where $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$.

The question arises whether the following equality will take place

$$(2) \quad (N_{p_0, q_0}(M), N_{p_1, q_1}(M))_{\theta, q} = N_{p, q}(M).$$

Here, in contrast to the Morrey spaces in the one-dimensional case, when M is the set of all segments, the answer is positive [NT].

In this paper we show that, if M is the set of dyadic cubes in \mathbb{R}^n , then the relation (2) holds. In case, if M is the set of all cubes, an analogue of the Marcinkiewicz-Calderon theorem on the cones of non-negative functions is obtained.

Given functions F and G , in this paper $F \lesssim G$ means that $F \leq CG$, where C is a positive number, depending only on numerical parameters, that may be different on different occasions. Moreover, $F \asymp G$ means that $F \lesssim G$ and $G \lesssim F$.

2. MAIN RESULTS

Let (A_0, A_1) – compatible pair of Banach spaces [BL].

$$K(t, a; A_0, A_1) = \inf_{a=a_0+a_1} (\|a_0\|_{A_0} + t\|a_1\|_{A_1}), \quad a \in A_0 + A_1,$$

– functional Petre. For $0 < q < \infty$, $0 < \theta < 1$

$$(A_0, A_1)_{\theta, q} = \left\{ a \in A_0 + A_1 : \|a\|_{(A_0, A_1)_{\theta, q}} = \left(\int_0^\infty (t^{-\theta} K(t, a))^q \frac{dt}{t} \right)^{1/q} < \infty \right\},$$

and for $q = \infty$

$$(A_0, A_1)_{\theta, q} = \left\{ a \in A_0 + A_1 : \|a\|_{(A_0, A_1)_{\theta, q}} = \sup_{0 < t < \infty} t^{-\theta} K(t, a) < \infty \right\}.$$

A family of sets from \mathbb{R}^n of the form

$$Q_k^m = \left[\frac{k_1}{2^m}, \frac{k_1 + 1}{2^m} \right) \times \dots \times \left[\frac{k_n}{2^m}, \frac{k_n + 1}{2^m} \right),$$

where $k \in \mathbb{Z}^n$, $m \in \mathbb{Z}$, is called the family of dyadic cubes and denoted by M .

Note that for arbitrary $m \in \mathbb{Z}$ the space \mathbb{R}^n can be represented in the form

$$\mathbb{R}^n = \bigcup_{k \in \mathbb{Z}^n} Q_k^m,$$

and the measure of intersection

$$|Q_k^m \cap Q_r^m| = \begin{cases} 2^{nm}, & k_i = r_i, \quad i = \overline{1, n} \\ 0, & \text{in other cases.} \end{cases}$$

Let $m \in \mathbb{Z}$, cubes Q_k^m , $k \in \mathbb{Z}^n$ are called cubes of m order.

Note also that if $n \geq m$, then each cube of n order is partitioned into 4^{n-m} cubes of m order.

Theorem 1. *Let $0 < p_0 < p_1 < \infty$ and $0 < q_0, q_1, q \leq \infty$. Let M be the family of dyadic cubes. Then*

$$(N_{p_0, q_0}(M), N_{p_1, q_1}(M))_{\theta, q} = N_{p, q}(M),$$

where $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$, $\theta \in (0, 1)$.

The following statement is an attempt to answer the question about the interpolation of net spaces, when M – is the family of all cubes with parallel faces to the coordinate axes in \mathbb{R}^n .

Theorem 2. *Let $n \leq p_0 < p_1 < \infty$, $0 < q \leq \infty$, M be the family of all cubes with parallel faces to the coordinate axes in \mathbb{R}^n . Let $G = \{f : f(x) \geq 0\}$, then for any $f \in G \cap N_{p, q}(M)$ it is true*

$$(3) \quad \|f\|_{(N_{p_0, q_0}(M), N_{p_1, q_1}(M))_{\theta, q}} \asymp \|f\|_{N_{p, q}(M)},$$

where the corresponding constants depend only on $p_i, q_i, \theta, q, i = 0, 1$.

The following corollary holds from Theorem 2.

Corollary 3. *Let $n \leq p_0 < p_1 < \infty$, $1 \leq q_0, q_1 < \infty$, $q_0 \neq q_1$, $0 < \tau, \sigma < \infty$. M and G – sets from Theorem 2. If the following inequalities hold for a quasilinear operator*

$$(4) \quad \|Tf\|_{N_{q_0, \infty}(M)} \leq F_0 \|f\|_{N_{p_0, \sigma}(M)}, \quad f \in N_{p_0, \sigma}(M),$$

$$(5) \quad \|Tf\|_{N_{q_1, \infty}(M)} \leq F_1 \|f\|_{N_{p_1, \sigma}(M)}, \quad f \in N_{p_1, \sigma}(M),$$

then for any $f \in G \cap N_{p, \tau}$ we have

$$(6) \quad \|Tf\|_{N_{q, \tau}(M)} \leq c F_0^{1-\theta} F_1^\theta \|f\|_{N_{p, \tau}(M)},$$

where $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$, $\theta \in (0, 1)$ and the corresponding constant depends only on $p_i, q_i, \sigma, i = 0, 1$.

3. PROOF OF THEOREM 1

Proof. Let M – be the set of dyadic cubes, $1 < p_0 < \infty$. Let us prove first

$$(7) \quad (N_{1, \infty}(M), N_{\infty, \infty}(M))_{\theta, q} = N_{p, q}(M),$$

where $\frac{1}{p} = 1 - \theta$, $\theta \in (0, 1)$.

Let $m \in \mathbb{Z}$, then the Euclidean space \mathbb{R}^n is partitioned into disjoint cubes of order m from M

$$\mathbb{R}^n = \bigcup_{k \in \mathbb{Z}^n} Q_k^m.$$

Let $f \in N_{p, q}(M)$, define the function

$$\varphi_0(x) = \frac{1}{|Q_k^m|} \int_{Q_k^m} f(x) dx, \quad x \in Q_k^m, \quad k \in \mathbb{Z}^n.$$

Then taking into account that the measure $|Q_k^m| = 2^{nm}$ we have

$$(8) \quad |\varphi_0(x)| \leq \bar{f}(2^{nm}, M), \quad x \in \mathbb{R}^n,$$

and

$$\int_{Q_k^m} (f(x) - \varphi_0(x)) dx = 0.$$

For the Petre functional, we have the following

$$\begin{aligned} K(t, f; N_{p_0, \infty}(M), N_{\infty, \infty}(M)) &= \inf_{f=f_0+f_1} (\|f_0\|_{N_{p_0, \infty}(M)} + t \|f_1\|_{N_{\infty, \infty}(M)}) \leq \\ &\leq \sup_{s>0} s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s, M) + t \sup_{s>0} \bar{\varphi}_0(s, M) = \sup_{s>0} s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s, M) + t \bar{f}(2^{nm}, M). \end{aligned}$$

Consider the first term

$$\sup_{s>0} s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s, M) \asymp \sup_{2^{nm} \geq s > 0} s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s, M) + \sup_{s \geq 2^{nm}} s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s, M).$$

Let I be an arbitrary cube from M such that $|I| \geq 2^{nm}$. Hence, I is some cube of order n , where $n \geq m$. Taking into account that each dyadic cube of n order is partitioned into mutually disjoint cubes of m order, we obtain

$$\left| \int_I (f - \varphi_0)(x) dx \right| = \left| \sum_{Q_k^m \subset I} \int_{Q_k^m} (f(x) - \varphi_0(x)) dx \right| = 0.$$

Hence,

$$\begin{aligned} \sup_{s>0} s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s, M) &= \sup_{2^{nm} \geq s > 0} s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s, M) \\ &\leq \sup_{2^{nm} \geq s > 0} s^{\frac{1}{p_0}} \bar{f}(s, M) + \sup_{2^{nm} \geq s > 0} s^{\frac{1}{p_0}} \bar{\varphi}_0(s, M) \leq \\ &\leq \sup_{2^{nm} \geq s > 0} \frac{1}{p_0} \int_0^s t^{\frac{1}{p_0}} \bar{f}(t, M) \frac{dt}{t} + \sup_{2^{nm} \geq s > 0} s^{\frac{1}{p_0}} \bar{f}(2^{nm}, M) \\ &\leq \sup_{2^{nm} \geq s > 0} \frac{1}{p_0} \int_0^s t^{\frac{1}{p_0}} \bar{f}(t, M) \frac{dt}{t} + 2^{\frac{nm}{p_0}} \bar{f}(2^{nm}, M) = \\ &= \frac{1}{p_0} \int_0^{2^{nm}} t^{\frac{1}{p_0}} \bar{f}(t, M) \frac{dt}{t} + \frac{1}{p_0} \int_0^{2^{nm}} t^{\frac{1}{p_0}} \bar{f}(t, M) \frac{dt}{t} = \frac{2}{p_0} \int_0^{2^{nm}} \bar{f}(y, M) y^{\frac{1}{p_0}-1} dy. \end{aligned}$$

In this way,

$$K(a^m, f; N_{p_0, \infty}(M), N_{\infty, \infty}(M)) \leq c \int_0^{2^{nm}} y^{\frac{1}{p_0}} \bar{f}(y, M) \frac{dy}{y} + a^m \bar{f}(2^{nm}, M),$$

where $a = 2^{\frac{n}{p_0}} > 1$. Then we have

$$\begin{aligned} \|f\|_{(N_{p_0, \infty}(M), N_{\infty, \infty}(M))_{\theta, q}} &\asymp \left(\sum_{m \in \mathbb{Z}} (a^{-\theta m} K(a^m, f))^q \right)^{\frac{1}{q}} \\ &\leq \left(\sum_{m \in \mathbb{Z}} \left(a^{-\theta m} \left(c \int_0^{4^m} y^{\frac{1}{p_0}} \bar{f}(y, M) \frac{dy}{y} + a^m \bar{f}(2^{nm}, M) \right) \right)^q \right)^{\frac{1}{q}}. \end{aligned}$$

Applying the Minkowski inequality, we obtain the following

$$\begin{aligned} \|f\|_{(N_{p_0, \infty}(M), N_{\infty, \infty}(M))_{\theta, q}} &\leq \left(\sum_{m \in \mathbb{Z}} \left(a^{-\theta m} c \int_0^{4^m} y^{\frac{1}{p_0}} \bar{f}(y, M) \frac{dy}{y} \right)^q \right)^{\frac{1}{q}} \\ &\quad + \left(\sum_{m \in \mathbb{Z}} (a^{m(1-\theta)} \bar{f}(4^m, M))^q \right)^{\frac{1}{q}}. \end{aligned}$$

Further, taking into account that $a = 2^{\frac{n}{p_0}}$ and applying Hardy's inequality for the first term, we obtain

$$\begin{aligned} \|f\|_{(N_{p_0, \infty}(M), N_{\infty, \infty}(M))_{\theta, q}} &\leq \left(\sum_{m \in \mathbb{Z}} \left(a^{-\theta m} \sum_{k=-\infty}^m 2^{\frac{nk}{p_0}} \bar{f}(2^{nk}, M) \right)^q \right)^{\frac{1}{q}} \\ &\quad + \left(\sum_{m \in \mathbb{Z}} (a^{m(1-\theta)} \bar{f}(2^{nm}, M))^q \right)^{\frac{1}{q}} \end{aligned}$$

$$\begin{aligned} &\leq c \left(\sum_{m \in \mathbb{Z}} \left(2^{\left((-\theta) \frac{1}{p_0} + \frac{1}{p_0} \right) nm} \bar{f}(2^{nm}, M) \right)^q \right)^{\frac{1}{q}} + \left(\sum_{m \in \mathbb{Z}} \left(2^{\frac{nm(1-\theta)}{p_0}} \bar{f}(2^{nm}, M) \right)^q \right)^{\frac{1}{q}} \\ &= c \left(\sum_{m \in \mathbb{Z}} \left(2^{\frac{nm}{p}} \bar{f}(2^{nm}, M) \right)^q \right)^{\frac{1}{q}} \asymp \|f\|_{N_{p,q}}, \end{aligned}$$

where $\frac{1}{p} = \frac{1-\theta}{p_0}$, $\theta \in (0, 1)$.

So we got the embedding

$$N_{p,q} \hookrightarrow (N_{1,\infty}(M), N_{\infty,\infty}(M))_{\theta,q},$$

where $\frac{1}{p} = 1 - \theta$, $\theta \in (0, 1)$. The reverse embedding follows from (1). Hence the relation (7) holds. To prove the general case, we use the reiteration theorem [BL, Theorem 3.5.3].

Let $1 < p_0 < p_1 < \infty$. From (7) it follows that there are $\theta_0, \theta_1 \in (0, 1)$ such that

$$(9) \quad \begin{aligned} (N_{1,\infty}(M), N_{\infty,\infty}(M))_{\theta_0,q_0} &= N_{p_0,q_0}(M) \\ (N_{1,\infty}(M), N_{\infty,\infty}(M))_{\theta_1,q_1} &= N_{p_1,q_1}(M), \end{aligned}$$

then by the reiteration theorem it follows that

$$(N_{p_0,q_0}(M), N_{p_1,q_1}(M))_{\theta,q} = (N_{1,\infty}(M), N_{\infty,\infty}(M))_{\eta,q} = N_{p,q}(M).$$

In the last equality, we took into account that $\eta = (1 - \theta)\theta_0 + \theta\theta_1$.

□

4. PROOF OF THEOREM 2

Proof. Let $\tau > 0$, euclidean space $\mathbb{R}^n = \bigcup_{k=1}^{\infty} I_k$ partitions into non-intersecting half-open cubes $\{I_k\}_{k=1}^{\infty}$ with faces parallel to the coordinate axes and such that $|I_k| = \tau$. Let $f \in G \cap N_{p,q}(M)$, define the function

$$\varphi_0(x) = \frac{1}{|I_k|} \int_{I_k} f(x) dx, \quad x \in I_k, \quad k \in \mathbb{N}.$$

Then it is obvious that

$$(10) \quad \varphi_0(x) \leq \bar{f}(\tau), \quad x \in \mathbb{R}^n,$$

and

$$\int_{I_k} (f(x) - \varphi_0(x)) dx = 0.$$

Let $0 < \sigma < \min\{q_0, q_1, q\}$, then

$$\begin{aligned} K(t, f; N_{p_0,\sigma}(M), N_{p_1,\sigma}(M)) &= \inf_{f=f_0+f_1} (\|f_0\|_{N_{p_0,\sigma}(M)} + t\|f_1\|_{N_{p_1,\sigma}(M)}) \\ &\leq \left(\int_0^\infty \left(s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} + t \left(\int_0^\infty \left(s^{\frac{1}{p_1}} \bar{\varphi}_0(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}}. \end{aligned}$$

Estimate the first term, taking into account the inequality (10), we have

$$\begin{aligned} &\left(\int_0^\infty \left(s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \asymp \left(\int_0^\tau \left(s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \\ &+ \left(\int_\tau^\infty \left(s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} = \left(\int_0^\tau \left(s^{\frac{1}{p_0}} \bar{f}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \\ &+ \left(\int_0^\tau \left(s^{\frac{1}{p_0}} \bar{\varphi}_0(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} + \left(\int_\tau^\infty \left(s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \\ &\leq \left(\int_0^\tau \left(s^{\frac{1}{p_0}} \bar{f}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} + \bar{f}(\tau) \left(\int_0^\tau s^{\frac{\sigma}{p_0}-1} ds \right)^{\frac{1}{\sigma}} \end{aligned}$$

$$\begin{aligned}
& + \left(\int_{\tau}^{\infty} \left(s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s) \right)^{\sigma} \frac{ds}{s} \right)^{\frac{1}{\sigma}} = 2 \left(\int_0^{\tau} \left(s^{\frac{1}{p_0}} \bar{f}(s) \right)^{\sigma} \frac{ds}{s} \right)^{\frac{1}{\sigma}} \\
& \quad + \left(\int_{\tau}^{\infty} \left(s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s) \right)^{\sigma} \frac{ds}{s} \right)^{\frac{1}{\sigma}}.
\end{aligned}$$

Let I be an arbitrary cube from M , ∂I – its boundary, then

$$\begin{aligned}
& \left| \int_I (f - \varphi_0)(x) dx \right| = \\
& = \left| \sum_{I_k \subset I} \int_{I_k} (f(x) - \varphi_0(x)) dx + \sum_{|I_k \cap \partial I| \neq 0} \int_{I_k \cap I} (f(x) - \varphi_0(x)) dx \right|.
\end{aligned}$$

Note that the first sum is equal to zero, and the second contains at most $2^n \left(\left(\frac{|I|}{\tau} \right)^{\frac{1}{n}} + 1 \right)^{n-1}$ terms. Moreover, taking into account the non-negativity of the function f , the second sum is estimated as follows

$$\begin{aligned}
& \left| \sum_{|I_k \cap \partial I| \neq 0} \int_{I_k \cap I} (f(x) - \varphi_0(x)) dx \right| \\
& \leq \sum_{|I_k \cap \partial I| \neq 0} \int_{I_k \cap I} f(x) dx + \sum_{|I_k \cap \partial I| \neq 0} |I_k \cap I| \frac{1}{|I_k|} \int_{I_k} f(x) dx \\
& \leq 2 \sum_{|I_k \cap \partial I| \neq 0} \int_{I_k} f(x) dx \leq 2 \sum_{|I_k \cap \partial I| \neq 0} \tau \bar{f}(\tau) \leq 4^n |I|^{\frac{n-1}{n}} \tau^{\frac{1}{n}} \bar{f}(\tau).
\end{aligned}$$

Hence,

$$\begin{aligned}
& \sup_{|I| \geq s} \frac{1}{|I|} \left| \int_I (f - \varphi_0)(x) dx \right| \leq \sup_{|I| \geq s} \frac{1}{|I|} 4^n |I|^{\frac{n-1}{n}} \tau^{\frac{1}{n}} \bar{f}(\tau) \\
& = 4^n \tau^{\frac{1}{n}} \bar{f}(\tau) \sup_{|I| \geq s} \frac{1}{|I|^{\frac{1}{n}}} \leq 4^n \cdot \frac{\tau^{\frac{1}{n}}}{s^{\frac{1}{n}}} \bar{f}(\tau).
\end{aligned}$$

Further, taking into account that $p_0 \geq n$ we have

$$\begin{aligned}
& \left(\int_{\tau}^{\infty} \left(s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s) \right)^{\sigma} \frac{ds}{s} \right)^{\frac{1}{\sigma}} = \left(\int_{\tau}^{\infty} \left(s^{\frac{1}{p_0}} \sup_{|I| \geq s} \frac{1}{|I|} \left| \int_I (f - \varphi_0)(x) dx \right| \right)^{\sigma} \frac{ds}{s} \right)^{\frac{1}{\sigma}} \\
& \leq 4^n \left(\int_{\tau}^{\infty} \left(s^{\frac{1}{p_0}} \frac{\tau^{\frac{1}{n}}}{s^{\frac{1}{n}}} \bar{f}(\tau) \right)^{\sigma} \frac{ds}{s} \right)^{\frac{1}{\sigma}} = 4^n \tau^{\frac{1}{n}} \bar{f}(\tau) \left(\int_{\tau}^{\infty} \left(s^{\frac{1}{p_0} - \frac{1}{n}} \right)^{\sigma} \frac{ds}{s} \right)^{\frac{1}{\sigma}} \\
& \asymp \tau^{\frac{1}{n}} \bar{f}(\tau) \tau^{\frac{1}{p_0} - \frac{1}{n}} = \tau^{\frac{1}{p_0}} \bar{f}(\tau) = c \left(\int_0^{\tau} \left(s^{\frac{1}{p_0}} \bar{f}(s) \right)^{\sigma} \frac{ds}{s} \right)^{\frac{1}{\sigma}}.
\end{aligned}$$

Hence,

$$\left(\int_0^{\infty} \left(s^{\frac{1}{p_0}} \overline{(f - \varphi_0)}(s) \right)^{\sigma} \frac{ds}{s} \right)^{\frac{1}{\sigma}} \leq c \left(\int_0^{\tau} \left(s^{\frac{1}{p_0}} \bar{f}(s) \right)^{\sigma} \frac{ds}{s} \right)^{\frac{1}{\sigma}},$$

where c depends only on parameters p_0, σ .

To estimate the second term, we first show that,

$$(11) \quad \bar{\varphi}_0(s) \leq \begin{cases} \bar{f}(\tau), & \text{at } s \leq \tau, \\ 4\bar{f}(s), & \text{at } s > \tau. \end{cases}$$

For $s \leq \tau$ from (10) we have $\bar{\varphi}_0(s) \leq \bar{f}(\tau)$. Let $s > \tau$, $Q \in M$, $|Q| = s$ and by Q' we denote a cube, whose center coincides with the center of Q and has an edge twice as large, then we have

$$\begin{aligned} \bar{\varphi}_0(s) &= \sup_{\substack{|Q| \geq s \\ Q \in M}} \frac{1}{|Q|} \sum_{I_k \cap Q \neq \emptyset} \int_{I_k \cap Q} \frac{1}{|I_k|} \int_{I_k} f(x) dx dy \\ &\leq \sup_{\substack{|Q| \geq s \\ Q \in M}} \frac{1}{|Q|} \sum_{I_k \cap Q \neq \emptyset} \int_{I_k} f(x) dx \frac{|I_k \cap Q|}{|I_k|} \leq \sup_{\substack{|Q| \geq s \\ Q \in M}} \frac{1}{|Q|} \sum_{I_k \cap Q \neq \emptyset} \int_{I_k} f(x) dx \\ &\leq \sup_{\substack{|Q| \geq s \\ Q \in M}} \frac{1}{|Q|} \int_{Q'} f(x) dx \leq 4\bar{f}(s). \end{aligned}$$

Further, using the relation (11) we obtain

$$\begin{aligned} &\left(\int_0^\infty \left(s^{\frac{1}{p_1}} \bar{\varphi}_0(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} = \left(\int_0^\tau \left(s^{\frac{1}{p_1}} \bar{\varphi}_0(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \\ &+ \left(\int_\tau^\infty \left(s^{\frac{1}{p_1}} \bar{\varphi}_0(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \leq \bar{f}(\tau) \left(\int_0^\tau \left(s^{\frac{1}{p_1}} \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \\ &+ 4 \left(\int_\tau^\infty \left(s^{\frac{1}{p_1}} \bar{f}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} = c\tau^{\frac{1}{p_1}} \bar{f}(\tau) + 4 \left(\int_\tau^\infty \left(s^{\frac{1}{p_1}} \bar{f}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}}, \end{aligned}$$

where c depends only on parameters p_1, σ .

Hence,

$$\begin{aligned} K(t, f; N_{p_0, \sigma}(M), N_{p_1, \sigma}(M)) &\leq c \left(\int_0^\tau \left(s^{\frac{1}{p_0}} \bar{f}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \\ &+ t \left(c\tau^{\frac{1}{p_1}} \bar{f}(\tau) + 4 \left(\int_\tau^\infty \left(s^{\frac{1}{p_1}} \bar{f}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \right). \end{aligned}$$

Taking into account the monotonicity of $\bar{f}(s)$, for $\tau = t^{\left(\frac{1}{p_0} - \frac{1}{p_1}\right)^{-1}}$ we obtain

$$\begin{aligned} \|f\|_{(N_{p_0, \sigma}(M), N_{p_1, \sigma}(M))_{\theta, q}} &\leq \left(\int_0^\infty \left[t^{-\theta} \left(\int_0^{t^{\left(\frac{1}{p_0} - \frac{1}{p_1}\right)^{-1}}} \left(s^{\frac{1}{p_0}} \bar{f}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \right. \right. \\ &\left. \left. + t \left(ct^{\left(\frac{1}{p_0} - \frac{1}{p_1}\right)^{-1} \frac{1}{p_1}} \bar{f}(t^{\left(\frac{1}{p_0} - \frac{1}{p_1}\right)^{-1}}) + 4 \left(\int_{t^{\left(\frac{1}{p_0} - \frac{1}{p_1}\right)^{-1}}^\infty \left(s^{\frac{1}{p_1}} \bar{f}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \right) \right]^q \frac{dt}{t} \right)^{\frac{1}{q}}. \end{aligned}$$

Making the replacement $\gamma = t^{\left(\frac{1}{p_0} - \frac{1}{p_1}\right)^{-1}}$ and applying the Minkowski inequality, we arrive at the following

$$\begin{aligned} \|f\|_{(N_{p_0, \sigma}(M), N_{p_1, \sigma}(M))_{\theta, q}} &\leq \left(\int_0^\infty \left(\gamma^{-\theta \left(\frac{1}{p_0} - \frac{1}{p_1}\right)} \left(\int_0^\gamma \left(s^{\frac{1}{p_0}} \bar{f}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \right)^q \frac{d\gamma}{\gamma} \right)^{\frac{1}{q}} \\ &+ \left(\int_0^\infty \left(\gamma^{\frac{1}{p}} \bar{f}(\gamma) \right)^q \frac{d\gamma}{\gamma} \right)^{\frac{1}{q}} + \left(\int_0^\infty \left(\gamma^{(1-\theta) \left(\frac{1}{p_0} - \frac{1}{p_1}\right)} \left(\int_\gamma^\infty \left(s^{\frac{1}{p_1}} \bar{f}(s) \right)^\sigma \frac{ds}{s} \right)^{\frac{1}{\sigma}} \right)^q \frac{d\gamma}{\gamma} \right)^{\frac{1}{q}}. \end{aligned}$$

Further, for the first and third terms, we apply the following variants of Hardy inequalities: if $\mu > 0, -\infty < \nu < \infty$ and $0 < \sigma, \tau \leq \infty$, then

$$\left(\int_0^\infty \left(y^{-\mu} \left(\int_0^y (r^{-\nu} |g(r)|)^\sigma \frac{dr}{r} \right)^{\frac{1}{\sigma}} \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}} \leq (\mu\sigma)^{-\frac{1}{\sigma}} \left(\int_0^\infty (y^{-\mu-\nu} |g(y)|)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}}$$

and

$$\left(\int_0^\infty \left(y^\mu \left(\int_0^y \left(r^{-\nu} |g(r)| \right)^\sigma \frac{dr}{r} \right)^{\frac{1}{\sigma}} \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}} \leq (\mu\sigma)^{-\frac{1}{\sigma}} \left(\int_0^\infty \left(y^{\mu-\nu} |g(y)| \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}}.$$

According to these inequalities we have

$$\|f\|_{(N_{p_0,\sigma}(M), N_{p_1,\sigma}(M))_{\theta,q}} \lesssim \left(\int_0^\infty \left(\gamma^{\frac{1}{p}} \bar{f}(\gamma) \right)^q \frac{d\gamma}{\gamma} \right)^{\frac{1}{q}} = \|f\|_{N_{p,q}},$$

and hence,

$$\|f\|_{(N_{p_0,q_0}(M), N_{p_1,q_1}(M))_{\theta,q}} \lesssim \|f\|_{(N_{p_0,\sigma}(M), N_{p_1,\sigma}(M))_{\theta,q}} \lesssim \|f\|_{N_{p,q}(M)},$$

where $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$. From the theorem in [N1, Theorem 1] we know that the following embedding holds

$$(N_{p_0,q_0}(M), N_{p_1,q_1}(M))_{\theta,q} \hookrightarrow N_{p,q}(M),$$

i.e.

$$\|f\|_{N_{p,q}(M)} \lesssim \|f\|_{(N_{p_0,q_0}(M), N_{p_1,q_1}(M))_{\theta,q}}.$$

This proves the equivalence (2). □

5. PROOF OF COROLLARY 1

Proof. According to the real interpolation method [BL, Theorem 3.1.2] and the inequalities (4) and (5) it follows

$$\|Tf\|_{(N_{q_0,\infty}(M), N_{q_1,\infty}(M))_{\theta,\tau}} \leq F_0^{1-\theta} F_1^\theta \|f\|_{(N_{p_0,\sigma}(M), N_{p_1,\sigma}(M))_{\theta,\tau}}.$$

From the relation (1) we have

$$\|Tf\|_{N_{q,\tau}(M)} \leq c \|Tf\|_{(N_{q_0,\infty}(M), N_{q_1,\infty}(M))_{\theta,\tau}}.$$

From Theorem 2, taking into account, that $f \geq 0$, we obtain

$$\|f\|_{N_{p,\tau}(M)} \asymp \|f\|_{(N_{p_0,\sigma}(M), N_{p_1,\sigma}(M))_{\theta,q}}.$$
□

REFERENCES

- [AR1] R. Akylzhanov, M. Ruzhansky *L_p – L_q multipliers on locally compact groups* // J. Fun. Anal. — 2020, Vol. 278, Issue 3.
- [AR2] R. Akylzhanov, M. Ruzhansky *Net spaces on lattices, Hardy-Littlewood type inequalities, and their converses.* // Eurasian Math. J. — 2017, Vol. 8 Issue 3, — P. 10-27.
- [ARN] R. Akylzhanov, M. Ruzhansky, E.D. Nursultanov *Hardy-Littlewood, Hausdorff-Young-Paley inequalities, and L_p – L_q Fourier multipliers on compact homogeneous manifolds* // J. Math. Anal. Appl. — 2019, Vol. 479 Issue 2, — P. 1519-1548.
- [BL] J. Bergh, J. Löfström, *Interpolation Spaces. An Introduction.* Springer, Berlin, 1976.
- [BRV] O. Blasco, A. Ruiz, L. Vega *Non interpolation in Morrey-Campanato and block spaces.* // Ann. Scuola Norm. Sup. Pisa Cl. — 1999, Sci. 4, — P. 31-40.
- [CM] A. Campanato, M.K.V. Murthy, *Una generalizzazione del teorema di Riesz-Thorin.* Ann. Scuola Norm. Super. Pisa 19 (1965), 87-100.
- [LR2] P.G. Lemarié-Rieusset, *Multipliers and Morrey spaces.* Potential Analysis, Springer 38 (2013), no. 3, 741-752
- [N1] E. D. Nursultanov *Net spaces and inequalities of Hardy-Littlewood type* // Sb. Math., — 1998. — Vol. 189, — C. 399-419.
- [N2] E.D. Nursultanov *On the coefficients of multiple Fourier series in L_p - spaces* // Izv. Math. — 2000, Vol. 64, N:1, — P. 93-120.
- [N3] E.D. Nursultanov *Application of Interpolation Methods to the Study of Properties of Functions of Several Variables.* // Mathematical Notes 75, 341–351 (2004).
- [NA] E.D. Nursultanov, T.U. Aubakirov *Interpolation methods for stochastic processes spaces* // Abstr. Appl. Anal. — 2013, Vol. 2013, — P. 1-12.

- [NK] E.D. Nursultanov, A.G. Kostyuchenko *Theory of control of "catastrophes"* // Russ. Math. Surv.— 1998, Vol. 53, N:3, — P. 628-629.
- [NTI] E.D. Nursultanov, N.T. Tleukhanova *Lower and upper bounds for the norm of multipliers of multiple trigonometric Fourier series in Lebesgue spaces* // Func. Anal. Appl. — 2000, Vol. 34, N:2, — P. 151-153.
- [NT] E.D. Nursultanov, S. Tikhonov *Net spaces and boundedness of integral operators* // J. Geom. Anal. — 2011, Vol. 21, — P. 950-981.
- [P] J. Peetre, *On the theory of $\mathcal{L}_{p,\lambda}$ spaces*. J. Func. Anal. 4 (1969), 71-87.
- [RV] A. Ruiz, L. Vega *Corrigenda to "Unique continuation for Schrodinger operators" and a remark on interpolation of Morrey spaces*. // Publicacions Matemàtiques. — 1995, Vol. 39, — P. 405-411.
- [St] G. Stampacchia, *$\mathcal{L}_{p,\lambda}$ -spaces and interpolation*. Comm. Pure Appl. Math. 17 (1964), 293-306.

L.N.GUMILYOV EURASIAN NATIONAL UNIVERSITY, NUR-SULTAN, KAZAKHSTAN

M.V.LOMONOSOV MOSCOW STATE UNIVERSITY (KAZAKHSTAN BRANCH), NUR-SULTAN, KAZAKHSTAN
Email address: aitolkynnur@gmail.com, er-nurs@yandex.ru

This figure "universe.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/2107.10633v1>