

# AN ELEMENTARY PROOF OF THE 3 DIMENSIONAL SIMPLEX MEAN WIDTH CONJECTURE

AARON GOLDSMITH

ABSTRACT. After a Hessian computation, we quickly prove the 3D simplex mean width conjecture using classical methods in section 2. Then, we generalize some components to  $d$  dimensions in section 4.

## 1. INTRODUCTION

Let  $B_2^d \subset \mathbb{R}^d$  be the standard Euclidean ball, with basis  $(e_1, \dots, e_d)$ . Denote  $\mathbb{S}^{d-1} := \partial B_2^d$ . Let  $\delta(X, Y)$  be the arclength on  $\mathbb{S}^{d-1}$  and  $\mu$  be the uniform probability measure on  $\mathbb{S}^{d-1}$ .

The support function of a convex body  $K \subset B_2^d$  is

$$h_K(u) := \max_{x \in K} u \cdot x$$

and the mean width

$$w(K) := 2 \int_{\mathbb{S}^{d-1}} h_K(u) d\mu(u)$$

**Conjecture 1.1.** (*Simplex Mean Width Conjecture*) *Of all simplices contained in  $B_2^d$ , the inscribed regular simplex has the maximum mean width, and is unique up to isometry.*

Conjecture 1.1 was mentioned in the survey by Gritzmann and Klee [4] and by Klee several times in his talks. Litvak surveyed the problem more recently in arXiv:math/0606350 [math.DG]. It's related to the problem of recovering transmissions from a noisy signal and has been assumed to be true by information theorists [2, 3]. The  $d + 1$  dimensional Gaussian Random Vector Maximum Conjecture is equivalent and was proved in 4 dimensions arXiv:2008.04827v2 [math.PR]. In section 2, we give a simpler and more classical proof. The following claim will be useful throughout.

**Claim 1.2.** *In order that  $\Delta := \text{conv}(v_0, \dots, v_d)$  maximizes  $w(\cdot)$  over all simplices contained in  $B_2^d$ , it must be that*

- (a)  $B_2^d$  is the smallest ball containing  $\Delta$ .
- (b)  $v_i \in \mathbb{S}^{d-1}$  for  $0 \leq i \leq n$ , i.e.  $\mathbb{S}^{d-1}$  is the circumsphere of  $\Delta$ .

---

*Date:* December 11, 2021.

(c) The closed hemispheres centered at  $v_i$  cover  $\mathbb{S}^{d-1}$ .

*Proof.* (a) If  $\Delta \subset Q + rB_2^d$  with  $r < 1$ , then  $(\Delta - Q)/r$  is a simplex contained in  $B_2^d$  and

$$w((\Delta - Q)/r) = w(\Delta/r) = w(\Delta)/r > w(\Delta)$$

(b) Suppose that  $|v_0| < 1$ . Extend the edge  $v_1v_0$  past  $v_0$  until it meets  $\mathbb{S}^{d-1}$  at a point  $v_{n+1}$ . The simplex  $\tilde{\Delta} := \text{conv}\{v_1, \dots, v_{n+1}\} \subset B_2^d$  contains  $v_0$  as a convex combination of  $v_{n+1}$  and  $v_1$ , so it strictly contains  $\Delta$ . Since  $w$  is strictly increasing under set inclusion,  $w(\Delta) < w(\tilde{\Delta})$ . Therefore, if  $|v_i| < 1$  for any  $i$ , then  $w(\Delta)$  is not maximum.

(c) Suppose  $v \in \mathbb{S}^{d-1}$  is such that  $\max v \cdot v_i = m < 0$  for all  $i$ . It follows that each  $v_i$  is contained in the spherical cap  $\{v \cdot x \leq m\}$ , which has diameter less than 1. Part (a) implies there is no such  $v$ . □

The support function of the simplex  $\Delta = \text{conv}\{v_0, \dots, v_d\}$  reduces to a maximum over the vertex set:

$$\max_{x \in \Delta} u \cdot x = \max_{0 \leq i \leq d} u \cdot v_i$$

As such, for each  $i$  define the  $i$ th Voronoi cell on the sphere to be

$$S_i := \{u \in \mathbb{S}^{d-1} : u \cdot v_i = \max_{x \in \Delta} u \cdot x\}$$

The mean width of a simplex is then

$$\begin{aligned} w(\Delta) &= 2 \int_{\mathbb{S}^{d-1}} \max_{0 \leq i \leq d} X \cdot v_i d\mu(X) \\ &= 2 \sum_{i=0}^d \int_{S_i} X \cdot v_i d\mu(X) \end{aligned} \tag{1.1}$$

Now define

$$M_A(R) := \int_R X \cdot A d\mu(X) \tag{1.2}$$

## 2. SMWC DIMENSION 3

Capital Roman letters  $A, B, C$  represent either vertices of a spherical triangle or their angles, depending on context; small Roman letters  $a, b, c$ , are used for either the opposite legs or their arclength. Next, we present some classical results about trigonometry in  $\mathbb{S}^2$  (see [6]).

**Theorem 2.1** (Spherical Law of Sines). *For a general triangle  $\triangle ABC \subset \mathbb{S}^2$ ,*

$$\frac{\sin a}{\sin A} = \frac{\sin b}{\sin B} = \frac{\sin c}{\sin C}$$

**Theorem 2.2** (Napier). *Suppose  $\triangle ABC$  is a right spherical triangle. Then,*

$$(2.1) \quad \cos c = \cos a \cos b$$

$$(2.2) \quad \cos A = \cos a \sin B$$

$$(2.3) \quad \cos A = \tan b \cot c$$

$$(2.4) \quad \tan a = \tan A \sin b$$

*and, of course, the same if  $(A, a)$  and  $(B, b)$  are interchanged.*

Claim 2.3 gives a parametrization of an arc as measured from the opposite vertex and Theorem 2.4 a simple formula reminiscent of the area of a Euclidean triangle.

**Claim 2.3.** *Let  $\triangle ABC \subset \mathbb{S}^2$  be a right spherical triangle with  $A = (1, 0, 0)$  and the right angled vertex  $C = (\cos b, \sin b, 0)$ . Side  $a$  is parametrized by  $(\cos \Phi(\theta), \cos \theta \sin \Phi(\theta), \sin \theta \sin \Phi(\theta))$  as  $\theta \in [0, A]$  where*

$$\sin^2 \Phi(\theta) = \frac{\tan^2 b}{\tan^2 b + \cos^2 \theta}$$

*Proof.* Napier's rule 2.3 says  $\cos \theta = \tan b \cot \Phi(\theta)$ , so

$$\sin^2 \Phi(\theta) = \frac{1}{1 + \cot^2 \Phi(\theta)} = \frac{\tan^2 b}{\tan^2 b + \cos^2 \theta}$$

□

**Theorem 2.4.** *Let  $\triangle ABC$  be a right spherical triangle with  $C = \pi/2$ . Then,*

$$4\pi\sigma(\triangle ABC)M_A(\triangle ABC) = \frac{1}{2}a \sin b$$

*Proof.* Set  $A = (1, 0, 0)$ . Uniform measure on  $\mathbb{S}^2$  is  $d\mu = (4\pi)^{-1} \sin \varphi d\varphi d\theta$ , giving

$$\begin{aligned}
8\pi \int_{\Delta ABC} X \cdot Ad\mu(X) &= \int_0^A \int_0^{\Phi(\theta)} \cos \varphi \sin \varphi d\varphi d\theta \\
&= \int_0^A \sin^2 \Phi(\theta) d\theta \\
&= \int_0^A \tan^2 b / (\tan^2 b + \cos^2 \theta) d\theta \\
&= \tan^2 b \int_0^A \sec^2 \theta / (1 + \tan^2 b + \tan^2 b \tan^2 \theta) d\theta \\
&= \tan^2 b \int_0^{\tan A} 1 / (\sec^2 b + u^2 \tan^2 b) du \\
&= \sin b \int_0^{\tan A \sin b} 1 / (1 + u^2) du \\
&= \sin b (\tan^{-1}(\sin b \tan A)) \\
&= a \sin b
\end{aligned}$$

Where Claim 2.3 was used in the third line and the last line follows from Napier's rule 2.4. □

**Remark 2.5.** Denote the area of  $\Delta ABC$  as  $[ABC]$ . The centroid  $G$  satisfies

$$[ABC]G = \int_{\Delta ABC} X d\mu(X)$$

Let  $d$  be the length of the altitude from angle  $A$  to side  $a$ . Either  $\Delta ABC$  is the union of two right triangles with common side  $d$ , or the difference. Either way, use the previous claim for the second line and Theorem 2.1 in the third of the following,

$$\begin{aligned}
2[ABC](G \cdot A) &= 2M_A(\Delta ABC) \\
&= a \sin d \\
&= a \sin b \sin C
\end{aligned}$$

Since these quantities are independent of coordinates, the two other cases ( $A \rightarrow B \rightarrow C$ ) follow similarly. If  $A, B, C$  have no common great circle,  $G$  is the unique point satisfying these equations. The formula

$$(2.5) \quad 2[ABC]G \det(A, B, C) = \sum_{\text{cyc}} (B \times C) a \sin b \sin C$$

is verified by taking dot products with each of  $A, B, C$  separately. Recall the triple product  $\det(A, B, C) = A \times B \cdot C$ . Now, arbitrarily putting  $A = (1, 0, 0)$  and  $C_2 = 0$  and  $\sigma(\Delta ABC) = 1$ , we see that  $C \times B \cdot A = C_1 B_2 =$

$\sin b(\sin c \sin A)$ , the volume of the zonotope generated by  $A, B, C$ , is the ubiquitous quantity  $n$  associated to a spherical triangle in [6]. Dividing the equation 2.5 thru by this quantity becomes Brock's formula [7],

$$2[ABC]G = \sum_{\text{cyc}} (B \times C) \frac{a}{\sin a}$$

Consider a right spherical triangle  $\Delta$  with  $C = \pi/2$ . We use equation 2.2 to convert the formula from Theorem 2.4 in terms of  $A$  and  $B$ .

**Theorem 2.6.** *The function defined by*

$$f(A, B) := a \sin b = \left( \cos^{-1} \frac{\cos B}{\sin A} \right) \left( 1 - \frac{\cos^2 A}{\sin^2 B} \right)^{1/2}$$

is negative definite (i.e.  $-f$  is convex) in the region

$$R := \{-\pi/2 < A, B < \pi/2\} \cap \{\cos^2 A + \cos^2 B < 1\}$$

*Proof.* See section 5 □

**Theorem 2.7.** *The Simplex Mean Width Conjecture is true for  $d = 3$*

*Proof.* Let the orientation of a spherical triangle be  $\sigma(\Delta ABC) = +1$  for a counterclockwise ordering of  $A, B, C$  and  $-1$  for clockwise. If  $\sigma(\Delta ABC) = -1$ , measure both the angles and edges as negative.

Suppose  $\Delta = \text{conv}(v_0, \dots, v_3)$  is any simplex with  $|v_i| = 1$  for each  $i$  (by Claim 1.2). The Voronoi cells

$$S_i(\Delta) := \{u \in \mathbb{S}^2 : i \in \text{argmax}_i u \cdot v_i\}$$

are each the intersection of three hemispheres and so are spherical triangles tiling  $\mathbb{S}^2$ . For each  $i$ , label the vertices of  $S_i$  as  $A, B, C$  (suppressing subscript  $i$ ) in the counterclockwise orientation. Draw arcs from  $v_i$  to  $A, B, C$  and drop altitudes from  $v_i$  to the edges  $a, b, c$  bounding  $S_i$  (they may leave the interior of  $S_i$ ) with feet  $D, E, F$  on  $a, b, c$  (resp.). Form the collection  $\mathcal{T}$  of right triangles  $\Delta v_i XY$  such that  $X, Y$  come from the list  $A, F, B, D, C, E$  and  $Y$  is immediately to the right of  $X$ . Since

$$\sigma(\Delta XYZ) \mathbb{1}_{\Delta XYZ} + \sigma(\Delta XZW) \mathbb{1}_{\Delta XZW} = \sigma(\Delta XYW) \mathbb{1}_{\Delta XYW}$$

we can write

$$M_{v_i}(S_i) = \sum_{T \in \mathcal{T}} \sigma(T) M_{v_i}(T)$$

Also, the signed sum of the six angles meeting at  $v_i$  is  $2\pi$  while the signed sum of the other six non-right angles equals the sum of the angles of  $S_i$ , which exceeds the area of  $S_i$  by  $\pi$  (Girard's Theorem). It follows that the

whole of  $\mathbb{S}^2$  may be divided into 24 right triangles where the signed sum of the 24 angles measured at  $\{v_0, \dots, v_3\}$  is  $4 * 2\pi$  and the sum of the other 24 non-right angles is  $4\pi + 4\pi$ . To see that none of these 24 right triangles contains an angle exceeding  $\pi/2$ , note that the spherical Pythagorean (2.1) implies there would be two such edges from the same triangle, in particular one meeting at a vertex  $v_i$ . But, from (Claim. 1.2(c)) every point in  $\mathbb{S}^2$  is within  $\pi/2$  from a vertex  $v_i$ . As no edge exceeds  $\pi/2$ , neither does any angle. Theorems 2.4 and 2.6 imply that the maximum mean width occurs when all these angles are equal, i.e. a regular tetrahedron.  $\square$

### 3. INTEGRATION ON SPHERES

To generalize Theorem 2.4 we need to generalize a right triangle to a right angled simplex, called a path simplex, on higher dimensional spheres. Then, we find the marginal mean of an RV uniform over the simplex. Let us start with the measure of spherical caps. Denote a spherical cap of geodesic radius  $r$ , centered at  $e_1$ , by  $B_{\mathbb{S}}^{d-1}(r) \subset \mathbb{S}^{d-1}$ . Define the incomplete Wallis integral by

$$\mathcal{W}^d(x) = \int_0^x \sin^d t dt$$

and denote  $W^d := \mathcal{W}^d(\pi)$ .

**Theorem 3.1.** *The uniform measure of a spherical cap is*

$$(3.1) \quad \mu(B_{\mathbb{S}}^{d-1}(r)) = \frac{\mathcal{W}^{d-2}(r)}{W^{d-2}}$$

where

$$(3.2) \quad \mathcal{W}^d(r) = \frac{d-1}{d} \mathcal{W}^{d-2}(r) - \frac{1}{d} \cos r \sin^{d-1} r$$

*Proof.* Cut the sphere into thin slices perpendicular to the radius at the center of the cap. The cross section at geodesic radius  $\varphi$  is  $\sin \varphi \cdot \mathbb{S}^{d-2}$ , so the measure of a thin slice is proportional to  $\sin^{d-2} \varphi d\varphi$ , and equation 3.1 follows.

For the recursion formula, consider  $(\sin^d \varphi)''$  for  $d \geq 2$ :

$$\begin{aligned} (d \cos \varphi \sin^{d-1} \varphi)' &= d(d-1) \cos^2 \varphi \sin^{d-2} \varphi - d \sin^d \varphi \\ &= d(d-1) \sin^{d-2} \varphi - d^2 \sin^d \varphi \end{aligned}$$

Integrating gives 3.2.  $\square$

**Remark 3.2.** *Equation 3.2 may be used to find the trigonometric series for  $\mathcal{W}^d(r)$ . Note that  $\mathcal{W}^0(r) = r$  and  $\mathcal{W}^1(r) = 1 - \cos r$ . The denominator of 3.1 is the well known Wallis Integral. When  $r = \pi$ , we can multiply equation 3.2 through by  $W^{d-1}$  to get*

$$d \cdot W^d W^{d-1} = (d-1)W^{d-1}W^{d-2}$$

which shows

$$d \cdot W^d W^{d-1} = W^1 W^0 = 2\pi$$

for all  $d$ . Since  $W^d$  is decreasing with  $d$ , we come upon the bound (see also [9]),

$$(3.3) \quad \sqrt{\frac{2\pi}{d+1}} < W^d < \sqrt{\frac{2\pi}{d}}$$

**Corollary 3.3.** *Averaging the marginal of an RV uniform over a cap,*

$$\begin{aligned} M_{e_1}(B_{\mathbb{S}}^{d-1}(r)) &= \int_0^r \cos t \, d\mu(B_{\mathbb{S}}^{d-1}(t)) \\ &= \frac{\int_0^r \cos t \sin^{d-2} t \, dt}{W^{d-2}} \\ &= \frac{\sin^{d-1} r}{(d-1)W^{d-2}} \end{aligned}$$

Now, fix an axis  $A \in \mathbb{S}^{d-1}$  and parametrize  $\mathbb{S}^{d-1}$  by  $X = (r, \theta) \in [0, \pi] \times \mathbb{S}^{d-2}$  where  $\cos r = X \cdot A$  and

$$\theta = \frac{X - A \cos r}{|X - A \cos r|}$$

**Corollary 3.4.** *Let a region  $R \subset \mathbb{S}^{d-1}$  be defined by  $0 \leq r \leq \varphi(\theta)$  under the above parameterization for some measurable  $\varphi$ . Then*

$$M_A(R) = \int_{\mathbb{S}^{d-2}} \frac{\sin^{d-1} \varphi(\theta)}{(d-1)W^{d-2}} d\mu(\theta)$$

When this region is a spherical simplex, say  $T \subset \mathbb{S}^{d-1}$ , it is the intersection of  $d$  hemispheres. Arrange the vertices as columns in the matrix  $V$ . We describe  $T$  with matrix  $H \in \mathbb{R}^{d \times d}$ , whose rows are the centers of the  $d$  hemispheres, ordered so that the  $i$ th vertex is in the interior of the  $i$ th hemisphere. Then  $x/|x| \in T$  iff

$$Hx \geq \mathbf{0}$$

Since each vertex  $v_i$  is on the boundary of the other  $d-1$  hemispheres,  $H$  is a diagonal multiple of  $V^{-1}$ . Further, if  $e_1$  is a vertex, we may assume the first column of  $H$  is  $(1, 0, 0, \dots, 0)^t$ . Also,  $T$  is a union of arcs originating at  $e_1$ , and  $T$  has a parameterization  $0 \leq r \leq \varphi(\theta)$  as in Corollary 3. To find  $\varphi(\theta)$ , note that the first row of  $H$  describes the facet opposite  $e_1$ .

$$\cos \varphi(\theta) + \theta \cdot (H_{12}, \dots, H_{1d}) \sin \varphi(\theta) = 0$$

giving

$$\tan \varphi(\theta) = -\frac{1}{\theta \cdot (H_{12}, \dots, H_{1d})}$$

or

$$\sin^2 \varphi(\theta) = \frac{1}{1 + (\theta \cdot (H_{12}, \dots, H_{1d}))^2}$$

The support  $\tilde{T}$  of  $\varphi$  is the simplex described by omitting the first row and column of  $H$ . Applying Corollary 3, we get

$$(3.4) \quad M_{e_1}(T) = \frac{1}{(d-2)W^{d-2}} \int_{\tilde{T}} (1 + (\theta \cdot (H_{12}, \dots, H_{1d}))^2)^{(1-d)/2} d\mu(\theta)$$

#### 4. SMWC FOR $d > 3$

In this section, we assume the set of points  $V = \{v_0, \dots, v_d\} \subset \mathbb{S}^{d-1}$  is in general position so that the feet of the altitudes to a face never land in the boundary of that face.

**Definition 4.1.** A path simplex in  $\mathbb{S}^{d-1}$  is a spherical simplex with  $d$  edges forming a path such that any pair of them determine great circles meeting at right angles. The endpoints of the path are called end vertices.

**Definition 4.2.** Let  $T$  be a spherical simplex with vertices  $\{p_1, \dots, p_d\}$  and inward facing normals  $\{g_1, \dots, g_d\}$  corresponding to the opposite facets. Let  $G_T = [g_1 \dots g_d]$  and the (angle) Gram matrix be  $G_T^t G_T$ .

Note, many authors use  $2I_{d \times d} - G_T$  as the Gram matrix. It is known that  $G$  determines  $T$  up to isometry, e.g. [8].

**Claim 4.3.** If  $T$  is a path simplex in  $\mathbb{S}^{d-1}$ , it has a tridiagonal Gram matrix.

*Proof.* Say the vertices along the path are  $p_1, \dots, p_d$ , in order. The  $i$ th row of  $P^{-1} = [p_1, \dots, p_d]^{-1}$  is normal to  $F_i$ , so there is a diagonal matrix  $D$  with

$$(4.1) \quad G = (DP^{-1})(DP^{-1})^t = D(P^t P)^{-1} D$$

The path simplex condition may be restated as the arc  $p_i p_{i+1}$  must be orthogonal to  $\text{span}\{p_{i+1}, \dots, p_d\}$  for  $1 \leq i \leq d-1$ . That is, the vector  $p_i - (p_i \cdot p_{i+1})p_{i+1}$  tangent to the arc at  $p_{i+1}$  is orthogonal to each of  $p_{i+1}, \dots, p_d$ . It follows that



$$P^t P \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ p_1 \cdot p_2 & 1 & 0 & \dots & 0 \\ 0 & p_2 \cdot p_3 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix}$$

is upper triangular. Substitute this into equation 4.1. Since  $G$  is symmetric,  $G$  must be tridiagonal.  $\square$

**Claim 4.4.** *Let  $\Delta$  be spherical simplex in  $\mathbb{S}^{d-1}$ . For almost every  $O \in \mathbb{S}^{d-1}$ , there is a collection  $\mathcal{T}$  of  $d!$  path simplices with end vertex  $O$  and a sign  $\sigma : \mathcal{T} \rightarrow \{-1, 1\}$  satisfying*

$$(4.2) \quad \mathbb{1}_\Delta = \sum_{T \in \mathcal{T}} \sigma(T) \mathbb{1}_T \quad (a.e.)$$

*Proof.* Proceed by induction;  $d = 2$  is trivial. Suppose the lemma is true at level  $d$ , and we will show it is also true at the level  $d + 1$ .

Let  $\Delta \subset \mathbb{S}^d$  be a simplex. Drop an altitude from  $O \in \mathbb{S}^{d-1}$  to any facet  $F$  of  $\Delta$  with foot  $O_F \notin \partial F$  (excluding a measure 0 case). Both  $F$  and  $O_F$  lie in a common great sphere  $S_F$  of dimension  $d - 1$ . By induction, we have a collection  $\mathcal{T}_F$  of  $d!$  path simplices with paths beginning at  $O_F$  and satisfying

$$\mathbb{1}_F = \sum_{T_F \in \mathcal{T}_F} \sigma(T_F) \mathbb{1}_{T_F}$$

Since  $OO_F$  is perpendicular to  $S_F$ , each of these paths may append  $OO_F$  to the beginning to represent the path simplices  $T^F := \text{conv}(O, T_F)$ . Form the collection  $\mathcal{T}^F := \{T^F : T_F \in \mathcal{T}_F\}$ , and the simplex  $\Delta^F := \text{conv}(O, F)$ . Now, identify points  $X \in S_F$  with segments  $OX$ , so that

$$(4.3) \quad \mathbb{1}_{\Delta^F} = \sum_{T^F \in \mathcal{T}^F} \sigma(T^F) \mathbb{1}_{T^F}$$

$$\mathbb{1}_\Delta = \sum_F \left\{ \begin{array}{ll} -1 & \text{if } \mu(\Delta^F \cap \Delta) = 0 \\ +1 & \text{o.w.} \end{array} \right\} \mathbb{1}_{\Delta^F}$$

Since there are  $d + 1$  faces  $F$  (in the second line above), the total sum involves  $(d + 1)!$  path simplices.  $\square$

**Claim 4.5.** *The Voronoi cells generated by  $V$  are simplices. The set  $\mathcal{C}$  of faces across all  $d + 1$  cells corresponds to the nonempty subsets of  $V$  in that each face is the set of points equidistant from the points in its corresponding subset. Also, the intersection of any two of these faces is again a face ( $\mathcal{C}$  is a simplicial complex).*

*Proof.* The Voronoi cells are bounded by perpendicular bisectors between each pair of vertices  $v_i, v_j$ . That's  $d$  facets for each cell, i.e. a simplex. Each face of a simplex corresponds to an intersection of facets, or in this case, perpendicular bisectors. Thus, each facet is the set of points equidistant from some subset of  $V$ . It follows that the intersection of two faces corresponds to the union of the two corresponding subsets.  $\square$

For a set  $V = \{v_0, \dots, v_d\} \subset \mathbb{S}^{d-1}$  and a maximal chain  $\tau$  in the power set of  $V$ , denote by  $S_\tau$  the path simplex starting at the vertex in the singleton set of  $\tau$  and landing successively in the faces determined by the subsets in  $\tau$ , increasing by inclusion.

**Claim 4.6.** *Let  $\mathcal{C}$  be as in the previous claim. To each face  $F \in \mathcal{C}$  (and hence every subset  $Q \subset V$ ) call  $p(F) \in \text{span}(F)$  the point that minimizes the distance from the points in  $Q$ . Then,  $p(F)$  is a vertex of  $S_\tau$  iff  $Q \in \tau$ .*

*Proof.*  $\text{span}(F)$  is the set of points equidistant from each point in  $Q$ . Since the composition of projections onto subspaces is the projection onto the intersection of the subspaces, each point in  $Q$  projects onto  $\text{span}(F)$  with foot  $p(F)$ . Finally,  $p(F)$  is a vertex of  $S_\tau$  if  $p(F)$  is equidistant from points in  $Q$ , but no other points in  $V$ .  $\square$

The following is the main result.

**Theorem 4.7.** *Let  $T \subset \mathbb{S}^{d-1}$  be the path simplex with  $\theta_{ij}$  the angle between facets  $F_i$  and  $F_j$ . If  $A$  is an end vertex of  $T$  and the Hessian of  $M_A(T)$  from equation 3.4 as a function of  $\theta_{12}, \dots, \theta_{d,d+1}$  is negative definite, then the simplex mean width conjecture is true.*

*Proof.* Let  $\Delta$  be a simplex in  $\mathbb{R}^d$  with vertices  $V = \{v_0, \dots, v_d\} \subset \mathbb{S}^{d-1}$  (Claim 1.2(b)). The altitudes from  $v_i$  and  $v_j$  to the perpendicular bisector between them meet at the same point. The rest of the triangulations of each cell, as in Claim 4.4, coincide along facets, forming a  $d+1$  times larger simplicial complex  $\mathcal{C}^\dagger$  where every face is the intersection of cells.

By Claim 4.6 the vertices (or facets or acute dihedral angles) of any path simplex  $S_\tau$  may be ordered as  $\tau$  is ordered. Given three consecutive subsets  $Q, Q \cup \{x\}, Q \cup \{x, y\}$  in a maximal chain  $\tau$  of the subsets of  $V$ , the only replacement subset for  $Q \cup \{x\}$  to keep  $\tau$  a chain is  $Q \cup \{y\}$ . It follows that if a dihedral angle at a  $d-3$  dimensional face in a path simplex  $S_\tau$  is between two nonadjacent facets, there are  $2 \times 2 = 4$  path simplices meeting at that face, at right angles as we know from Claim 4.3. Similarly, a dihedral angle between two adjacent facets has  $3 \times 2 = 6$  path simplices meeting at that face.

$\mathcal{C}^\dagger$  has a total of  $(d+1)!$  path simplices, each with  $d$  acute dihedral angles meeting together at the  $d-3$  dimensional faces, according to their position

in the  $\tau$  ordering. So, for each  $i$ , the sum of the angles  $\theta_{i,i+1}$  over all path simplices is  $2\pi(d+1)!/6$ . If  $M_A(T)$  is negative definite as a function of these adjacent dihedral angles, the mean width from equation 1.1 will be maximum when all the adjacent dihedral angles are equal to  $\pi/3$ , that is when  $\triangle$  is regular.  $\square$

## 5. PROOF OF THEOREM 2.3

Use Napier's formula 2.2 to phrase quantities in terms of 'intermediate variables'  $a, b$ . Differentiating this formula with respect to  $A$  and using the spherical law of sines 2.1,

$$\sin A = \sin a \sin B \frac{da}{dA} = \sin A \sin b \frac{da}{dA}$$

Now with respect to  $B$ , and using again 2.1 and 2.2 (cycled  $A \rightarrow B$ ),

$$\begin{aligned} 0 &= \sin a \sin B \frac{da}{dB} - \cos a \cos B \\ &= \sin A \sin b \frac{da}{dB} - \cos a \cos b \sin A \end{aligned}$$

The same holds if  $(A, a)$  and  $(B, b)$  are interchanged, yielding

$$\begin{array}{ll} \frac{\partial a}{\partial A} = \frac{1}{\sin b} & \frac{\partial a}{\partial B} = \frac{\cos a \cos b}{\sin b} \\ \frac{\partial b}{\partial A} = \frac{\cos a \cos b}{\sin a} & \frac{\partial b}{\partial B} = \frac{1}{\sin a} \end{array}$$

The first order partials are then

$$\begin{aligned} f_A &= \frac{da}{dA} \sin b + a \cos b \frac{db}{dA} = 1 + \frac{a \cos a \cos^2 b}{\sin a} \\ f_B &= \frac{da}{dB} \sin b + a \cos b \frac{db}{dB} = \cos b \left( \cos a + \frac{a}{\sin a} \right) \end{aligned}$$

For second order partials,

$$\begin{aligned} \frac{\sin a}{\cos a \cos^2 b} f_{AA} &= \frac{1}{\sin b} - \left( 2 \sin b \cos a + \frac{1}{\sin b \cos a} \right) \frac{a}{\sin a} \\ \frac{\sin a}{\cos a} f_{BB} &= -\sin b + \frac{\cos^2 a \cos^2 b}{\sin b} - \left( \frac{\sin b}{\cos a} + \frac{\cos^2 b}{\sin b} \cos a \right) \frac{a}{\sin a} \\ \frac{\sin a}{\cos a \cos b} f_{AB} &= \frac{\cos^2 b}{\sin b} \cos a - \left( \frac{1 + \sin^2 b}{\sin b} \right) \frac{a}{\sin a} \end{aligned}$$

Uniquely represent

$$\begin{aligned}\frac{\sin^2 a}{\cos^2 a \cos^2 b} \det \text{Hess}(f) &= \frac{\sin^2 a}{\cos^2 a \cos^2 b} (f_{AA} f_{BB} - f_{AB}^2) \\ &= P(\sin a, \sin b) + Q(\sin a, \sin b) \frac{a \cos a}{\sin a} + R(\sin a, \sin b) \frac{a^2}{\sin^2 a}\end{aligned}$$

for some rational functions  $P, Q, R$ . Now we will compute  $P, Q, R$ .

$$\begin{aligned}P(x, y) &= \left(\frac{1}{y}\right) \left(-y + \frac{(1-x^2)(1-y^2)}{y}\right) - \left(\frac{(1-y^2)^2}{y^2}(1-x^2)\right) \\ &= -1 + \frac{(1-x^2)(1-y^2)(1-(1-y^2))}{y^2} \\ &= (1-x^2)(1-y^2) - 1 \\ Q(x, y) &= -\frac{1}{y} \left(\frac{y}{1-x^2} + \frac{1-y^2}{y}\right) \\ &\quad - \left(2y + \frac{1}{y(1-x^2)}\right) \left(-y + \frac{(1-x^2)(1-y^2)}{y}\right) \\ &\quad + 2 \left(\frac{1-y^2}{y}\right) \left(\frac{1+y^2}{y}\right) \\ &= 2 \left(y^2 - \frac{1-y^2}{y^2} - (1-x^2)(1-y^2) + \frac{1-y^4}{y^2}\right) \\ &= 2(1 - (1-x^2)(1-y^2)) \\ R(x, y) &= \left(2y + \frac{1}{y(1-x^2)}\right) \left(y + \frac{(1-y^2)(1-x^2)}{y}\right) - \left(y + \frac{1}{y}\right)^2 \\ &= 2y^2 + \frac{1}{1-x^2} + 2(1-x^2)(1-y^2) + \frac{1}{y^2} - 1 - \left(y + \frac{1}{y}\right)^2 \\ &= -y^2 - 1 - 2x^2 + 2x^2y^2 + \frac{1}{1-x^2} \\ &= -2 + (1-2x^2)(1-y^2) + \frac{1}{1-x^2} \\ &= (2x^2 - 1) \left(y^2 - 1 + \frac{1}{1-x^2}\right) \\ &= \left(\frac{x^2}{1-x^2} - \frac{1-x^2}{x^2}\right) x^2 (1 - (1-x^2)(1-y^2))\end{aligned}$$

From the spherical law of sines 2.1 and spherical Pythagorean 2.1,

$$\frac{\sin^2 a}{\sin^2 A} = \sin^2 c = 1 - \cos^2 c = 1 - \cos^2 a \cos^2 b$$

and so we reduce the Hessian to

$$\begin{aligned}\frac{\sin^2 A}{\cos^2 a \cos^2 b} \det \text{Hess} f &= (\tan^2 a - \cot^2 a) a^2 + (2 \cot a) a - 1 \\ &= a^2 \tan^2 a - (1 - a \cot a)^2\end{aligned}$$

This can be seen to be positive by following the chain of inequalities backward:

$$\begin{aligned}0 &> (1 - a \cot a - a \tan a)(1 - a \cot a + a \tan a) \\ &= \left(1 - \frac{a}{\sin a \cos a}\right) \left(1 - a \frac{\cos^2 a - \sin^2 a}{\sin a \cos a}\right) \\ &= \left(1 - \frac{2a}{\sin 2a}\right) \left(1 - \frac{2a}{\tan 2a}\right)\end{aligned}$$

for  $|a| < \pi/2$ , which follows since  $x/\sin x$  (resp.  $x/\tan x$ ) is tangent to  $y = 1$  and increasing (resp. decreasing) away from 0 for  $-\pi < x < \pi$ .

Thanks to Joshua Horowitz and the use of his copy of Mathematica, which cleared the way for this proof.

#### REFERENCES

- [1] T. Kwok-keung Au, F. Luo, R. Stong, Comparing Corresponding Dihedral Angles on Classical Geometric Simplices, In: Asian J. Math. 12 (2) 203 - 212, June 2008. arXiv:math/0606350 [math.DG]
- [2] A.V. Balakrishnan, Research Problem No. 9, In: Geometry, Bull. Amer. Math. Soc. 69 (1963), 737738.
- [3] A.V. Balakrishnan, Signal selection for space communication channels, In: Advances in Communication Systems, (ed. by A.V. Balakrishnan), Academic Press, New York, 1965, 131. 373466, 1994.
- [4] P. Gritzmann, V. Klee, On the complexity of some basic problems in computational convexity II: Volume and mixed volumes, Bisztriczky, T. (ed.) et al., Polytopes: abstract, convex and computational. Proceedings of the NATO Advanced Study Institute. Kluwer Academic Publishers. NATO ASI Ser., Ser. C, Math. Phys. Sci. 440,
- [5] Wei Sun and Ze-Chun Hu and Guolie Lan, The 4-D Gaussian Random Vector Maximum Conjecture and the 3-D Simplex Mean Width Conjecture (2020), eprint arXiv:2008.04827v2 [math.PR]
- [6] I. Todhunter, Spherical Trigonometry: For the Use of Colleges and Schools, In: Macmillan and Co. (1886), (open source)
- [7] J. Brock, The centroid and inertia tensor for a spherical triangle, In: Calhoun (1974)
- [8] R. Díaz, A Characterization of Gram Matrices of Polytopes, In: Discrete & Computational Geometry Vol. 21 pg 581-601 (1999).
- [9] S. Taschuk, Integral of Powers of Sine (2013), (open source)

AARON GOLDSMITH, UNAFFILIATED  
*Email address:* aaron.goldsmith163@gmail.com