

REMARKS ON PARTIALLY ABELIAN EXACT CATEGORIES

THEO BÜHLER

ABSTRACT. The purpose of this short and elementary note is to identify some classes of exact categories introduced in L. Previdi's thesis. Among other things we show:

- (i) An exact category is partially abelian exact if and only if it is abelian.
- (ii) An exact category satisfies the axioms (AIC) and (AIC)^o if and only if it is quasi-abelian in the sense of J.-P. Schneiders.
- (iii) An exact category satisfies (AIC) if and only if it is an additive category of the type considered by G. Laumon in his work on derived categories of filtered \mathcal{D} -modules.

In all of the above classes all morphisms have kernels and coimages and the exact structure must be given by all kernel-cokernel pairs.

1. PRELIMINARY OBSERVATIONS

For every morphism $f: A \rightarrow B$ in an additive category there is an associated automorphism $\begin{bmatrix} 1 & 0 \\ f & 1 \end{bmatrix}$ of $A \oplus B$. It yields an isomorphism of short sequences

$$\begin{array}{ccccc} A & \xrightarrow{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} & A \oplus B & \xrightarrow{\begin{bmatrix} 0 & 1 \end{bmatrix}} & B \\ \parallel & & \cong \downarrow \begin{bmatrix} 1 & 0 \\ f & 1 \end{bmatrix} & & \parallel \\ A & \xrightarrow{\begin{bmatrix} 1 \\ f \end{bmatrix}} & A \oplus B & \xrightarrow{\begin{bmatrix} -f & 1 \end{bmatrix}} & B. \end{array}$$

The morphism $\begin{bmatrix} 1 \\ f \end{bmatrix}$ can be interpreted as the graph of f . It has a left inverse given by the projection onto the first coordinate, while composition with the projection onto the second coordinate yields the morphism $f = \begin{bmatrix} 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ f \end{bmatrix}$. In particular, every morphism in an additive category is the composition of a split monic followed by a split epic.

The following lemma simply states that the intersection of the graph of f with the graph of the zero morphism coincides with the kernel of f . The verification is straightforward.

LEMMA. *Let $f: A \rightarrow B$ be a morphism in an additive category. Consider the diagram of solid arrows below*

$$\begin{array}{ccccc} K & \xrightarrow{k} & A & \xrightarrow{f} & B \\ \downarrow k & \text{PB} & \downarrow \begin{bmatrix} 1 \\ f \end{bmatrix} & & \parallel \\ A & \xrightarrow{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} & A \oplus B & \xrightarrow{\begin{bmatrix} 0 & 1 \end{bmatrix}} & B. \end{array}$$

The pull-back square on the left exists if and only if f has a kernel $k: K \rightarrow A$. \square

COROLLARY. *An additive category admits pull-backs of all pairs of split monics if and only if it has kernels (if and only if it is finitely complete). \square*

Date: February 25, 2013.
2010 Mathematics Subject Classification. 18E10, 18G25.

2. INTRINSIC CHARACTERIZATION OF EXACT CATEGORIES SATISFYING (AIC)

Previdi's additional axiom (AIC) for exact categories [Pre10, 3.17] states that pull-backs of admissible monics along admissible monics exist and are admissible monics. Since split monics with cokernels are always admissible, we infer from (AIC) and the previous section that the underlying additive category must have kernels. Moreover, (AIC) and the lemma imply that all kernels are admissible monics.

Suppose f has a cokernel c . Then $\ker c$ is an admissible monic and $c = \text{coker } \ker c$ implies that c is an admissible epic. Therefore the class of cokernels is equal to the class of admissible epics.

This discussion establishes the necessity part of the following characterization of exact categories satisfying (AIC):

PROPOSITION. $(\mathcal{A}, \mathcal{E})$ is an exact category satisfying (AIC) if and only if:

- (i) Every morphism in \mathcal{A} has a kernel.
- (ii) The push-out of a kernel along an arbitrary morphism exists and is a kernel.
- (iii) Cokernels are stable under pull-backs along arbitrary morphisms.
- (iv) All kernels are admissible monics and all cokernels are admissible epics and \mathcal{E} is the class of all kernel-cokernel pairs in \mathcal{A} .

In particular, if \mathcal{A} admits an exact structure satisfying (AIC), it is the unique maximal exact structure on \mathcal{A} . Moreover, kernels are stable under pull-backs along arbitrary morphisms.

PROOF. Suppose $(\mathcal{A}, \mathcal{E})$ satisfies (i), (ii), (iii) and (iv). Note that (ii) implies that every kernel has a cokernel and with (i) this shows that every morphism has a coimage (form the push-out of the kernel along zero).

In order to check that \mathcal{E} is an exact structure, Keller [Kel90, A.1] shows that it is enough to verify that the composition of kernels is again a kernel.

To this end, let (i, p) and (j, q) be short exact such that ji is defined. We need to prove that ji is a kernel. Form the push-out under j and p to obtain the diagram

$$\begin{array}{ccccc}
 A' & \xrightarrow{i} & A & \xrightarrow{p} & A'' \\
 \parallel & & \downarrow j & \text{PO} & \downarrow j'' \\
 A' & \xrightarrow{ji} & B & \xrightarrow{r} & B'' \\
 & & \downarrow q & & \downarrow q'' \\
 & & C & \xlongequal{\quad} & C
 \end{array}$$

where q'' is uniquely determined by $q''r = q$ and $q''j'' = 0$. Since $q'' = \text{coker } j''$ we have $(j'', q'') \in \mathcal{E}$. Using the push-out property one checks that $r = \text{coker}(ji)$ and finally one checks that $ji = \ker r$ so that $(ji, r) \in \mathcal{E}$.

It remains to verify (AIC). Consider the following diagram in which $(b, b') \in \mathcal{E}$ and f is arbitrary:

$$\begin{array}{ccccc}
 A' & \xrightarrow{a} & A & \xrightarrow{a'} & A'' \\
 \downarrow f' & & \downarrow f & & \downarrow f'' \\
 B' & \xrightarrow{b} & B & \xrightarrow{b'} & B''
 \end{array}$$

Define $(a, a') = (\ker(b'f), \text{coim}(b'f))$. Since $b'fa = 0$, there are unique f' and f'' making the diagram commutative. If $\alpha: X \rightarrow A$ and $\beta': X \rightarrow B'$ are such that $f\alpha = b'\beta'$ then $b'f\alpha = b'b\beta' = 0$ and since $a = \ker(b'f)$, there is a unique morphism $\alpha': X \rightarrow A'$ such that $a\alpha' = \alpha$. Because b is monic we conclude $\beta' = f'\alpha'$. Thus, the left hand square is a pull-back and (AIC) follows. \square

REMARK. Exact categories satisfying (AIC) are studied in [Lau83, (1.3.0), p.160ff].

REMARK. Since admissible monics have cokernels, every morphism has in addition a coimage.

$$\begin{array}{ccc}
 & A & \xrightarrow{f} B \\
 \text{ker } f \nearrow & & \searrow \\
 \text{Ker } f & & \text{Coim } f \\
 & p = \text{coim } f & \nearrow m
 \end{array}
 \qquad
 \begin{array}{ccc}
 Y & \xrightarrow{y} & A \\
 \downarrow q & \text{PB} & \downarrow p \\
 X & \xrightarrow{x} & \text{Coim } f
 \end{array}$$

The morphism $m: \text{Coim } f \rightarrow B$ is monic: if $mx = 0$, form the pull-back over x and p , notice that $fy = 0$ so that y factors through $\text{Ker } f$. Therefore $0 = py = xq$, so $x = 0$ since q is epic. [A variant of this argument can be used to give a direct proof of [Pre10, Lemma 3.18].]

3. EXACT CATEGORIES SATISFYING (AIC) AND (AIC) $^\circ$

The characterization of exact categories satisfying (AIC) in the previous section is very nearly self-dual. The only missing piece is the existence of cokernels. Thus:

PROPOSITION. $(\mathcal{A}, \mathcal{E})$ is an exact category with (AIC) and (AIC) $^\circ$ if and only if:

- (i) Every morphism in \mathcal{A} has a kernel and a cokernel.
- (ii) Kernels are stable under push-outs along arbitrary morphisms.
- (iii) Cokernels are stable under pull-backs along arbitrary morphisms.
- (iv) \mathcal{E} is the class of all kernel-cokernel pairs in \mathcal{A} .

These are precisely the quasi-abelian categories of Schneiders [Sch99] and the almost abelian categories of Rump [Rum01]. \square

4. PARTIALLY ABELIAN EXACT CATEGORIES ARE ABELIAN

An exact category is called *partially abelian exact* if every morphism which is the composition of an admissible monic followed by an admissible epic has a factorization as an admissible epic followed by an admissible monic:

$$\begin{array}{ccc}
 & C & \\
 m \nearrow & & \searrow e \\
 A & \xrightarrow{f} & B \\
 \exists p \searrow & & \nearrow \exists i \\
 & I &
 \end{array}$$

that is, $f = em$ with m admissible monic and e admissible epic implies $f = ip$ with p admissible epic and i admissible monic. This factorization shows that $\ker f = \ker p$ and $\text{coker } f = \text{coker } i$ and thus I is both coimage and image of f .

From the first section we know that every morphism $f: A \rightarrow B$ in an additive category can be written as the composition of a split monic followed by a split epic $f = \begin{bmatrix} 0 & 1 \\ \hline & f \end{bmatrix}$. Thus, in a partially abelian exact category every morphism f factors as $f = ip$ via admissible epic p followed by an admissible monic i and hence every f has a kernel and a cokernel.

Consider the factorization $f = ip$ of a monic f into an admissible epic followed by an admissible monic. Then p is monic and since monic cokernels in an additive category are isomorphisms, p is an isomorphism from f to the admissible monic i , so f is an admissible monic, and hence it is a kernel. Dually, every epic is an admissible epic and hence it is a cokernel.

PROPOSITION. A category is partially abelian exact if and only if it is abelian. \square

REFERENCES

- [Kel90] Bernhard Keller, *Chain complexes and stable categories*, Manuscripta Math. **67** (1990), no. 4, 379–417. MR1052551 (91h:18006)
- [Lau83] G. Laumon, *Sur la catégorie dérivée des \mathcal{D} -modules filtrés*, Algebraic geometry (Tokyo/Kyoto, 1982), Lecture Notes in Math., vol. 1016, Springer, Berlin, 1983, pp. 151–237. MR726427 (85d:32022)
- [Pre10] Luigi Previdi, *Sato Grassmannians for generalized Tate spaces*, preprint arXiv:1002.4863 (2010), 1–57.
- [Rum01] Wolfgang Rump, *Almost abelian categories*, Cahiers Topologie Géom. Différentielle Catég. **42** (2001), no. 3, 163–225. MR1856638 (2002m:18008)
- [Sch99] Jean-Pierre Schneiders, *Quasi-abelian categories and sheaves*, Mém. Soc. Math. Fr. (N.S.) (1999), no. 76, vi+134. MR1779315 (2001i:18023)

POSTSCRIPTUM

This is an unmodified version of a note sent to Braunling, Groechenig and Wolfson. It was written late 2012 or early 2013 when I answered some questions that arose in their work on Tate objects in exact categories. At that point I also informed Previdi of these results. While others have since rediscovered variants of these ideas, there is still no citable reference for this material, so I made this publicly available on Braunling's request.

Email address: `math@theobuehler.org`