

DO LAW-INVARIANT LINEAR FUNCTIONALS COLLAPSE TO THE MEAN?

SHENGZHONG CHEN, NIUSHAN GAO, DENNY H. LEUNG, AND LEI LI

ABSTRACT. In this note, we show that, on a wide range of rearrangement-invariant spaces, a law-invariant bounded linear functional is a scalar multiple of the expectation. We also construct a rearrangement-invariant space on which this property fails.

1. INTRODUCTION AND NOTATION

In the past few years, law-invariant risk measures have been of intense research interest in Financial Mathematics; see, e.g., [2, 3, 5, 7, 8, 10, 12, 13, 14, 15, 17, 19, 20, 21, 22]. Of particular interest to us are two recent papers [3, 11] that investigate when a (convex) law-invariant risk measure collapses to the mean, i.e., being a scalar multiple of expectation. We are motivated to study the problem for linear functionals. Precisely, we investigate whether law-invariant bounded linear functionals on rearrangement-invariant spaces automatically collapse to the mean.

In Section 2 of this paper, we collect some preliminary facts on automatically collapsing to the mean of law-invariant bounded linear functionals. In Section 3, we establish a sufficient condition for automatically collapsing to the mean. It turns out that nearly all classical rearrangement-invariant spaces, including Lebesgue spaces L^p ($1 \leq p < \infty$), Lorentz spaces $L^{p,q}$ ($1 < p < \infty, 1 \leq q \leq \infty$) and Orlicz spaces, satisfy this condition and thus law-invariant bounded linear functionals on these spaces all collapse to the mean. In Section 4, we construct a rearrangement-invariant space on which some law-invariant bounded linear functionals fail to collapse to the mean.

We introduce the terminology and notation for this paper. Throughout the paper, $(\Omega, \mathcal{F}, \mathbb{P})$ stands for a non-atomic probability space. Given a random variable X' on another probability space $(\Omega', \mathcal{F}', \mathbb{P}')$, there exists a random variable X on Ω having the same distribution as X' , i.e. $\mathbb{P}(X \leq c) = \mathbb{P}'(X' \leq c)$ for any $c \in \mathbb{R}$; in this case,

Date: November 15, 2021.

2010 Mathematics Subject Classification. 46E30, 60E05.

Key words and phrases. Law invariance, linear functional, collapse to the mean, rearrangement-invariant space.

The first author is supported by an NSERC fellowship. The second author acknowledges support of an NSERC Discovery Grant.

we write $X \sim X'$ (cf., e.g., [9, Appendix 3]). For a random variable X on Ω , define its decreasing rearrangement by

$$X^*(t) = \inf \{ \lambda > 0 : \mathbb{P}(|X| > \lambda) \leq t \}, \quad t \in (0, 1).$$

Clearly, if $X \sim Y$, then $X^* \equiv Y^*$. Moreover, if $(0, 1)$ is endowed with the Lebesgue measure, then $X^* \sim |X|$. These notions and facts extend in a plain manner to finite measure spaces of the same measure. In particular, we will work on a set $A \in \mathcal{F}$ endowed with the probability structure restricted from $(\Omega, \mathcal{F}, \mathbb{P})$. Given a (measurable) partition $\pi = (A_i)_{i \in I}$ of Ω , where I is at most countable, we can define random variables on Ω by specifying its values on each A_i and then gluing the pieces together. A useful fact is as follows. Let $\pi' = (A'_i)_{i \in I}$ be a partition of Ω' for another probability space $(\Omega', \mathcal{F}', \mathbb{P}')$, such that $\mathbb{P}(A_i) = \mathbb{P}'(A'_i)$ for each $i \in I$. If X and X' are random variables on Ω and Ω' , respectively, such that $X|_{A_i} \sim X'|_{A'_i}$ for any $i \in I$, then $X \sim X'$.

Let L^0 be the space of all random variables (to be precise, equivalent classes of random variables modulo a.s. equality) on $(\Omega, \mathcal{F}, \mathbb{P})$. Throughout the paper, \mathcal{X} stands for a rearrangement-invariant (abbr., r.i.) space over $(\Omega, \mathcal{F}, \mathbb{P})$. That is, \mathcal{X} is a Banach space of functions in L^0 such that for any $X \in \mathcal{X}$, (1) if $Y \in L^0$ and $|Y| \leq |X|$ a.s. then $Y \in \mathcal{X}$ and $\|Y\| \leq \|X\|$, and (2) if $Z \in L^0$ and $Z \sim X$ then $Z \in \mathcal{X}$ and $\|Z\| = \|X\|$. We only consider $\mathcal{X} \neq \{0\}$. In this case, it is well known (see, e.g., [4, Ch. 2]) that $L^\infty \subset \mathcal{X} \subset L^1$ and there exist constants $C, C' > 0$ such that

$$(1.1) \quad \|X\|_1 \leq C\|X\| \text{ for every } X \in \mathcal{X}, \text{ and } \|X\| \leq C'\|X\|_\infty \text{ for every } X \in L^\infty.$$

Given an r.i. space \mathcal{X} , its *associate space* \mathcal{X}' is the space of all $Y \in L^0$ such that $\mathbb{E}[|XY|] < \infty$ for every $X \in \mathcal{X}$. \mathcal{X}' itself is also an r.i. space. In the Banach lattice language, \mathcal{X}' corresponds to the order continuous dual \mathcal{X}'_n of \mathcal{X} , which consists of all linear functionals ρ on \mathcal{X} such that $\rho(X_n) \rightarrow \rho(X)$ whenever $(X_n) \subset \mathcal{X}$ converges a.s. to $X \in \mathcal{X}$ and there exists $X_0 \in \mathcal{X}$ such that $|X_n| \leq X_0$ for all $n \in \mathbb{N}$. The *order continuous part* \mathcal{X}'_a of \mathcal{X}' is the set of all functions in \mathcal{X}' that are order continuous (in other words, have absolutely continuous norm), i.e., all $X \in \mathcal{X}'$ such that $\lim_{\mathbb{P}(A) \rightarrow 0} \|X \mathbf{1}_A\| = 0$. It is known that when $\mathcal{X} \neq L^\infty$, \mathcal{X}'_a is the norm closure of L^∞ in \mathcal{X}' . Moreover, in this case, $X \in \mathcal{X}'_a \iff \lim_n \|X \mathbf{1}_{\{|X| \geq n\}}\| = 0$.

An r.i. space \mathcal{X} is said to be *order continuous* if $\mathcal{X}'_a = \mathcal{X}'$, or equivalently, if $\mathcal{X}' = \mathcal{X}^*$ ([18, Theorem 2.4.2]), where \mathcal{X}^* is the norm continuous dual of \mathcal{X} . \mathcal{X} is said to be *monotonically complete*¹ if the pointwise supremum of any increasing, norm bounded sequence in \mathcal{X}_+ belongs to \mathcal{X} , or equivalently, if \mathcal{X} is isomorphic to $(\mathcal{X}')'$ via the evaluation mapping ([18, Theorem 2.4.22]). In this case, there exists a constant $M > 0$

¹In the literature, monotonic completeness bears many other names such as *maximal* and *the weak Fatou property*. It is sometimes included in the definition of function spaces.

such that

$$(1.2) \quad \|X\| \leq M \sup_{Y \in \mathcal{X}', \|Y\| \leq 1} |\mathbb{E}[XY]| \quad \text{for all } X \in \mathcal{X},$$

where the norm of $Y \in \mathcal{X}'$ is the dual norm on \mathcal{X} .

We refer the reader to [1, 18] for unexplained facts and terminology on Banach lattices and order structures and to [4] for facts and results on general r.i. spaces.

2. PRELIMINARY OBSERVATIONS

The functionals considered in [3, 11] are assumed to be $\sigma(\mathcal{X}, \mathcal{X}')$ lower semicontinuous. In general, weak continuities are rather restrictive assumptions to impose. For linear functionals, it is straightforward to see that under law invariance, $\sigma(\mathcal{X}, \mathcal{X}')$ lower semicontinuity is equivalent to collapsing to the mean.

Lemma 2.1. *Let \mathcal{X} be an r.i. space over a non-atomic probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let ρ be a law-invariant linear functional on \mathcal{X} .*

- (1) ρ is $\sigma(\mathcal{X}, \mathcal{X}')$ lower semicontinuous iff $\rho(X) = \rho(\mathbf{1})\mathbb{E}[X]$ for any $X \in \mathcal{X}$.
- (2) If ρ is norm continuous, then $\rho(X) = \rho(\mathbf{1})\mathbb{E}[X]$ for all $X \in \mathcal{X}_a$.

Proof. (1) Recall that a linear functional ρ on \mathcal{X} is $\sigma(\mathcal{X}, \mathcal{X}')$ lower semicontinuous iff it is $\sigma(\mathcal{X}, \mathcal{X}')$ continuous iff it lies in \mathcal{X}' , i.e., there exists $Y \in \mathcal{X}'$ such that

$$(2.1) \quad \rho(X) = \mathbb{E}[XY] \quad \text{for any } X \in \mathcal{X}.$$

Thus the “if” direction follows by taking $Y = \rho(\mathbf{1})\mathbf{1} \in \mathcal{X}'$. For the reverse direction, suppose that ρ is given by (2.1). If Y is not constant, then there exist two sets $A, B \in \mathcal{F}$ with positive probabilities and $\alpha \in \mathbb{R}$ such that $Y|_A > \alpha > Y|_B$. By using non-atomicity and shrinking to smaller sets, we may assume that $\mathbb{P}(A) = \mathbb{P}(B)$. Then $\mathbf{1}_A \sim \mathbf{1}_B$ but $\rho(\mathbf{1}_A) = \mathbb{E}[Y\mathbf{1}_A] > \alpha\mathbb{P}(A) = \alpha\mathbb{P}(B) > \mathbb{E}[Y\mathbf{1}_B] = \rho(\mathbf{1}_B)$, contradicting law-invariance of ρ . Thus $Y \equiv c$ for some constant $c \in \mathbb{R}$. Taking $X = \mathbf{1}$ in (2.1), we get $c = \rho(\mathbf{1})$. Putting this back into (2.1), we get $\rho(X) = \rho(\mathbf{1})\mathbb{E}[X]$ for any $X \in \mathcal{X}$.

(2) If $\mathcal{X} = L^\infty$, then $\mathcal{X}_a = \{0\}$ so that there is nothing to prove. Assume that $\mathcal{X} \neq L^\infty$. Then \mathcal{X}_a is an r.i. space over Ω and by order continuity of \mathcal{X}_a , $(\mathcal{X}_a)^* = (\mathcal{X}_a)'$. Thus being norm continuous, $\rho|_{\mathcal{X}_a}$ is $\sigma(\mathcal{X}_a, (\mathcal{X}_a)^*) = \sigma(\mathcal{X}_a, (\mathcal{X}_a)')$ continuous. The desired result follows from (1). \square

Recall that for a linear functional, $\sigma(\mathcal{X}, \mathcal{X}')$ lower semicontinuity implies norm continuity (boundedness). Thus following Lemma 2.1, it is natural to consider what happens if $\sigma(\mathcal{X}, \mathcal{X}')$ lower semicontinuity is weakened to plain norm boundedness. It turns out that collapsing to the mean no longer always holds for law-invariant norm bounded linear functionals. It, however, does hold on r.i. spaces satisfying (3.1) below.

Lemma 2.3 below reduces the problem to positive functionals. For the proof, we first recall [6, Lemma 2.3], which will be used later in the paper as well.

Lemma 2.2 ([6]). *Let X_1, X_2 be random variables on a non-atomic probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and X' be a random variable on a non-atomic probability space $(\Omega', \mathcal{F}', \mathbb{P}')$. If $X' \sim X_1 \geq X_2$, then there exists a random variable X'_2 on Ω' such that*

$$X' \geq X'_2 \sim X_2.$$

The conclusion still holds if both “ \geq ” are replaced by “ \leq ”.

Lemma 2.3. *Let \mathcal{X} be an r.i. space over a non-atomic probability space. Let ρ be a law-invariant bounded linear functional on \mathcal{X} . Then ρ^\pm is also law invariant.*

Proof. Take any $X \in \mathcal{X}_+$. Recall from the Riesz-Kantorovich formula ([1, Theorem 1.18]) that $\rho^+(X) = \sup\{\rho(Y) : 0 \leq Y \leq X\}$. Take any $X', Y \in \mathcal{X}$ such that $X' \sim X \geq Y \geq 0$. By Lemma 2.2, there exists a random variable Z such that $X' \geq Z \sim Y$. Since $Y \geq 0, Z \geq 0$. Thus by law invariance of ρ , $\rho(Y) = \rho(Z) \leq \rho^+(Z) \leq \rho^+(X')$. Taking supremum over Y , we have $\rho^+(X) \leq \rho^+(X')$. Hence, $\rho^+(X) = \rho^+(X')$ by symmetry. For general random variables $X, X' \in \mathcal{X}$ with $X \sim X', X^\pm \sim (X')^\pm$. Thus $\rho^+(X) = \rho^+(X^+) - \rho^+(X^-) = \rho^+((X')^+) - \rho^+((X')^-) = \rho^+(X')$. Therefore, ρ^+ is law invariant and hence so is $\rho^- = \rho^+ - \rho$. \square

A simple, but key, observation is that for a bounded linear functional ρ , collapsing to the mean and being law invariant can be characterized in terms of its kernel. Let \mathcal{X} be an r.i. space over a non-atomic probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Define

$$\mathcal{S} = \text{Span}\{X - Y : X, Y \in \mathcal{X}, X \sim Y\}, \text{ and } \mathcal{M} = \{X \in \mathcal{X} : \mathbb{E}[X] = 0\}.$$

Clearly, \mathcal{M} is a closed subspace of \mathcal{X} and

$$\overline{\mathcal{S}} \subset \mathcal{M} = \{X - \mathbb{E}[X] \cdot \mathbf{1} : X \in \mathcal{X}\},$$

where the closure is taken in \mathcal{X} with respect to the norm topology. The following lemma is immediate; we skip the proof.

Lemma 2.4. *Let \mathcal{X} be an r.i. space over a non-atomic probability space. Let ρ be a bounded linear functional on \mathcal{X} . Consider the following statements.*

- (1) $\rho(X) = \rho(\mathbf{1})\mathbb{E}[X]$ for all $X \in \mathcal{X}$.
- (2) $\mathcal{M} \subset \ker \rho$.
- (3) ρ is law invariant.
- (4) $\overline{\mathcal{S}} \subset \ker \rho$.

Then (1) \iff (2) \implies (3) \iff (4).

The following proposition reveals some elements in $\overline{\mathcal{S}}$.

Proposition 2.5. *Let \mathcal{X} be an r.i. space over a non-atomic probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Then $X - \mathbb{E}[X] \cdot \mathbf{1} \in \overline{\mathcal{S}}$ for any $X \in L^\infty$. The same holds for any $X \in \mathcal{X}_a$.*

Proof. First, let A be a measurable set such that $\mathbb{P}(A) \leq \frac{1}{m}$ for some $m \in \mathbb{N}$. By non-atomicity, there exist disjoint measurable sets $(A_i)_{i=1}^m$ of Ω such that $A_1 = A$ and $\mathbb{P}(A_i) = \mathbb{P}(A)$ for all $i = 1, \dots, m$. Then

$$(2.2) \quad \mathbf{1}_A - \mathbb{P}(A) \cdot \mathbf{1} = \frac{1}{m} \sum_{i=1}^m (\mathbf{1}_A - \mathbf{1}_{A_i}) + \frac{1}{m} \mathbf{1}_{\bigcup_{i=1}^m A_i} - \mathbb{P}(A) \cdot \mathbf{1}.$$

In particular, if $\mathbb{P}(A) = \frac{1}{m}$, then $\mathbb{P}(\bigcup_{i=1}^m A_i) = 1$, so that $\mathbf{1}_{\bigcup_{i=1}^m A_i} = \mathbf{1}$. Hence $\mathbf{1}_A - \mathbb{P}(A) \cdot \mathbf{1} = \frac{1}{m} \sum_{i=1}^m (\mathbf{1}_A - \mathbf{1}_{A_i}) \in \mathcal{S}$.

Next, let A be a measurable set such that $\mathbb{P}(A) = \frac{n}{m}$ for some $m, n \in \mathbb{N}$ with $1 \leq n < m$. Partition A into measurable sets B_1, \dots, B_n , where $\mathbb{P}(B_i) = \frac{1}{m}$ for $i = 1, \dots, n$. By the above, $\mathbb{P}(B_i) - \frac{1}{m} \cdot \mathbf{1} \in \mathcal{S}$ for all $i = 1, \dots, n$. Hence

$$\mathbf{1}_A - \frac{n}{m} \cdot \mathbf{1} = \sum_{i=1}^n \left(\mathbf{1}_{B_i} - \frac{1}{m} \cdot \mathbf{1} \right) \in \mathcal{S}.$$

Now, let A be any measurable set with $\mathbb{P}(A) \in (0, 1)$. For any $m \geq 3$, there exist disjoint subsets B, C of A such that $A = B \cup C$, $\mathbb{P}(B) = \frac{n-1}{m}$ and $0 \leq \mathbb{P}(C) < \frac{1}{m}$, where $n \in [1, m] \cap \mathbb{N}$. Clearly, $\mathbf{1}_A - \mathbb{P}(A) \cdot \mathbf{1} = \mathbf{1}_B - \mathbb{P}(B) \cdot \mathbf{1} + \mathbf{1}_C - \mathbb{P}(C) \cdot \mathbf{1}$. By the second case above, $\mathbf{1}_B - \mathbb{P}(B) \cdot \mathbf{1} \in \mathcal{S}$. By (2.2), $d(\mathbf{1}_C - \mathbb{P}(C) \cdot \mathbf{1}, \mathcal{S}) \leq \frac{2}{m} \|\mathbf{1}\|$. Therefore,

$$d(\mathbf{1}_A - \mathbb{P}(A) \cdot \mathbf{1}, \mathcal{S}) \leq \frac{2}{m} \|\mathbf{1}\|.$$

Since $m \geq 3$ is arbitrary, we obtain $\mathbf{1}_A - \mathbb{P}(A) \cdot \mathbf{1} \in \overline{\mathcal{S}}$.

From the above, it follows immediately that $X - \mathbb{E}[X] \cdot \mathbf{1} \in \overline{\mathcal{S}}$ for all simple functions X . If $X \in L^\infty$, take a sequence of simple functions (X_n) such that $\|X_n - X\|_\infty \rightarrow 0$. Since $X_n - \mathbb{E}[X_n] \cdot \mathbf{1} \in \overline{\mathcal{S}}$ for all $n \in \mathbb{N}$ and the sequence converges to $X - \mathbb{E}[X] \cdot \mathbf{1}$ in the norm of \mathcal{X} by (1.1), $X - \mathbb{E}[X] \cdot \mathbf{1} \in \overline{\mathcal{S}}$. This proves the first assertion.

The second assertion follows from the fact that, if $\mathcal{X}_a \neq \{0\}$, then \mathcal{X}_a is the norm closure of L^∞ in \mathcal{X} . \square

Remark 2.6. (1) Taking $\mathcal{X} = L^\infty$ in Proposition 2.5 and applying the implication (3) \implies (4) in Lemma 2.4, we obtain that if ρ is a law-invariant bounded linear functional on L^∞ , then it collapses to the mean. This fact, which may be known in the literature, can also be proved using the Radon-Nikydome theorem, by setting $\mu(A) = \rho(\mathbf{1}_A)$ for $A \in \mathcal{F}$. We skip the details.

(2) Similarly, the assertion for \mathcal{X}_a in Proposition 2.5 implies Lemma 2.1(2). The proof in Lemma 2.1(2) also uses the Radon-Nikydome theorem, although implicitly, when realizing $(\mathcal{X}_a)^*$ as $(\mathcal{X}_a)'$. We point out that Lemma 2.1(2) implies

that law-invariant bounded linear functionals on order continuous r.i. spaces collapse to the mean, which is also likely known in the literature.

Below is an interesting consequence of Proposition 2.5/Remark 2.6(1).

Corollary 2.7. *Let \mathcal{X} be an r.i. space over a non-atomic probability space and ρ be a law-invariant positive linear functional on \mathcal{X} . Then $\rho(X) \geq \rho(\mathbf{1})\mathbb{E}[X]$ for any $X \in \mathcal{X}_+$.*

Proof. Let $X \in \mathcal{X}_+$. By Remark 2.6(1), $\rho|_{L^\infty}$ collapses to the mean. Thus for any $n \geq 1$, since $X\mathbf{1}_{\{X \leq n\}} \in L^\infty$, $\rho(X) \geq \rho(X\mathbf{1}_{\{X \leq n\}}) = \rho(\mathbf{1})\mathbb{E}[X\mathbf{1}_{\{X \leq n\}}]$. Letting $n \rightarrow \infty$, we get $\rho(X) \geq \rho(\mathbf{1})\mathbb{E}[X]$. \square

We now present some equivalent conditions for collapsing to the mean.

Proposition 2.8. *Let \mathcal{X} be an r.i. space over a non-atomic probability space. The following are equivalent.*

- (1) *Every law-invariant positive linear functional ρ on \mathcal{X} has the form $\rho(X) = \rho(\mathbf{1})\mathbb{E}[X]$ for all $X \in \mathcal{X}$.*
- (2) *Every law-invariant bounded linear functional ρ on \mathcal{X} has the form $\rho(X) = \rho(\mathbf{1})\mathbb{E}[X]$ for all $X \in \mathcal{X}$.*
- (3) $\overline{\mathcal{S}} = \mathcal{M}$.
- (4) $\mathcal{M} \subseteq \overline{\mathcal{S} - \mathcal{X}_+}$.

Proof. As remarked earlier, (1) \iff (2) follows from Lemma 2.3. If (2) holds, then by Lemma 2.4, any bounded linear functional on \mathcal{X} that vanishes on $\overline{\mathcal{S}}$ also vanishes on \mathcal{M} . Hence, $\mathcal{M} \subseteq \overline{\mathcal{S}}$. The reverse inclusion is trivial. Thus (3) holds. (3) \implies (4) is clear. Assume that (4) holds. Then

$$\mathcal{X} = \mathbb{R} \cdot \mathbf{1} + \mathcal{M} \subseteq \mathbb{R} \cdot \mathbf{1} + \overline{\mathcal{S} - \mathcal{X}_+}.$$

Hence, $\mathbb{R} \cdot \mathbf{1} + \mathcal{S} - \mathcal{X}_+$ is dense in \mathcal{X} . Let ρ be a law-invariant positive functional on \mathcal{X} . Then the bounded linear functional $\tau(X) := \rho(X) - \rho(\mathbf{1})\mathbb{E}[X]$ vanishes on $\mathbb{R} \cdot \mathbf{1}$ and \mathcal{S} and $\tau \leq 0$ on $-\mathcal{X}_+$ by Corollary 2.7. It follows that $\tau(X) \leq 0$ for all $X \in \mathbb{R} \cdot \mathbf{1} + \mathcal{S} - \mathcal{X}_+$. Hence $\tau(X) \leq 0$ for all $X \in \mathcal{X}$. Thus $\tau = 0$, i.e., (1) holds. \square

3. A SUFFICIENT CONDITION

The following property was shown by the authors in [6] to play a pivotal role in automatic continuity of law-invariant convex functionals:

$$(3.1) \quad d(\mathcal{CL}(X), \mathcal{X}_a) = 0 \text{ for any } X \in \mathcal{X}_+, \text{ where } \mathcal{CL}(X) = \text{co}(\{Y : Y \sim X\}).$$

It was termed there as the *Almost Order Continuous Equidistributional Average (AOCEA)* property. Here, we show that the AOCEA property is also sufficient to ensure that all law-invariant bounded linear functionals collapse to the mean.

Proposition 3.1. *Let \mathcal{X} be an r.i. space over a non-atomic probability space other than L^∞ that satisfies the AOCEA property. Then $\overline{\mathcal{S}} = \mathcal{M}$. Hence every law-invariant bounded linear functional ρ on \mathcal{X} has the form $\rho(X) = \rho(\mathbf{1})\mathbb{E}[X]$ for all $X \in \mathcal{X}$.*

Proof. By (1.1), there is a finite constant C such that $\|X\|_1 \leq C\|X\|$ for all $X \in \mathcal{X}$. We may also assume that $\|\mathbf{1}\| = 1$. Let $X \in \mathcal{X}_+$ and let $\varepsilon > 0$ be given. By (3.1), choose $X' \in \mathcal{CL}(X)$ and $V \in \mathcal{X}_a$ such that $\|X' - V\| < \varepsilon$. By Proposition 2.5,

$$V - \mathbb{E}[V] \cdot \mathbf{1} \in \overline{\mathcal{S}}.$$

From the equation $X' - \mathbb{E}[X'] \cdot \mathbf{1} = (X' - V) - \mathbb{E}[X' - V] \cdot \mathbf{1} + V - \mathbb{E}[V] \cdot \mathbf{1}$, one sees that

$$d(X' - \mathbb{E}[X'] \cdot \mathbf{1}, \overline{\mathcal{S}}) \leq \|X' - V\| + \|X' - V\|_1 \|\mathbf{1}\| < (C + 1)\varepsilon.$$

Since $X - X' \in \mathcal{S}$ and $\mathbb{E}[X] = \mathbb{E}[X']$, it follows from $X - \mathbb{E}[X] \cdot \mathbf{1} = (X - X') + X' - \mathbb{E}[X'] \cdot \mathbf{1}$ that

$$d(X - \mathbb{E}[X] \cdot \mathbf{1}, \overline{\mathcal{S}}) \leq (C + 1)\varepsilon.$$

By arbitrariness of ε , $X - \mathbb{E}[X] \cdot \mathbf{1} \in \overline{\mathcal{S}}$ for all $X \in \mathcal{X}_+$. It follows easily that $X - \mathbb{E}[X] \cdot \mathbf{1} \in \overline{\mathcal{S}}$ for all $X \in \mathcal{X}$. Therefore, $\mathcal{M} \subseteq \overline{\mathcal{S}}$. The reverse inclusion is clear. This proves that $\mathcal{M} = \overline{\mathcal{S}}$. The second assertion follows from Proposition 2.8. \square

An alternative proof of this proposition in the spirit of [6] is included in the appendix.

It was also proved in [6] that when $\mathcal{X} \neq L^\infty$, (3.1) is equivalent to the following condition:

(3.2) For any $X \in \mathcal{X}_+$, any $\varepsilon > 0$, and any sequence of measurable sets $(A_n)_{n=1}^\infty$ with $A_n \downarrow \emptyset$, there exist natural numbers $(n_i)_{i=1}^k$, random variables $(Z_i)_{i=1}^k$, and a convex combination $\sum_{i=1}^k \lambda_i Z_i$ such that $Z_i \sim X \mathbf{1}_{A_{n_i}}$ for $i = 1, \dots, k$ and $\left\| \sum_{i=1}^k \lambda_i Z_i \right\| < \varepsilon$.

We compare it with the following more verifiable condition that uses disjoint Z_i 's:

(3.3) For any $X \in \mathcal{X}$, any $\varepsilon > 0$, any sequence of measurable sets $(A_n)_{n=1}^\infty$ with $A_n \downarrow \emptyset$ and $\sum_{n=1}^\infty \mathbb{P}(A_n) \leq 1$, and any disjoint sequence of measurable functions $(Z_n)_{n=1}^\infty$ such that $Z_n \sim X \mathbf{1}_{A_n}$ for all $n \geq 1$, there exist $n_1, \dots, n_k \in \mathbb{N}$ such that $\left\| \frac{1}{k} \sum_{i=1}^k Z_{n_i} \right\| < \varepsilon$.

Proposition 3.2. *Let \mathcal{X} be an r.i. space over a non-atomic probability space $(\Omega, \mathcal{F}, \mathbb{P})$ other than L^∞ . Then (3.3) \implies (3.2) (equivalently, (3.1)). The reverse is true if \mathcal{X} is either monotonically complete or order continuous.*

Proof of Proposition 3.2; the first assertion. Assume that (3.3) holds. To verify (3.2), take any $X \in \mathcal{X}_+$, $A_n \downarrow \emptyset$, and $\varepsilon > 0$. By passing to a subsequence of (A_n) , we may assume that $\sum_{n=1}^\infty \mathbb{P}(A_n) \leq 1$. By non-atomicity, there exists a disjoint sequence of measurable sets $(B_n)_{n=1}^\infty$ such that $\mathbb{P}(A_n) = \mathbb{P}(B_n)$ for all $n \geq 1$. For each $n \in \mathbb{N}$, since $\mathbb{P}(X\mathbf{1}_{A_n} \neq 0) \leq \mathbb{P}(A_n) = \mathbb{P}(B_n)$, by non-atomicity again, there exists a random variable Z_n , supported in B_n , such that $Z_n \sim X\mathbf{1}_{A_n}$. In particular, Z_n 's are disjoint. Applying (3.3) to (Z_n) , we obtain the desired Z_i 's and convex combination. \square

For the proof of (3.2) \implies (3.3) in the second assertion, we need some technical lemmas. For random variables X_1, \dots, X_k such that $\sum_{i=1}^k \mathbb{P}(X_i \neq 0) \leq 1$, let

$$\bigoplus_{i=1}^k X_i = \sum_{i=1}^k Y_i,$$

where Y_i 's are disjoint and $Y_i \sim X_i$ for each i . Clearly,

$$\left| \bigoplus_{i=1}^k X_i \right| = \left| \sum_{i=1}^k Y_i \right| = \sum_{i=1}^k |Y_i| = \bigoplus_{i=1}^k |X_i|.$$

Although $\bigoplus_{i=1}^k X_i$ is not uniquely defined as a random variable, its distribution and hence its norm are uniquely determined by X_i 's and that are what we use.

Lemma 3.3. *Let \mathcal{X} be an r.i. space over a non-atomic probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and $X \in \mathcal{X}$. Let $A \in \mathcal{F}$ be such that $\mathbb{P}(A) \leq \frac{1}{k}$ for some $k \in \mathbb{N}$ and that*

$$(3.4) \quad \alpha := \inf\{|X(\omega)| : \omega \in A\} \geq \|X\mathbf{1}_{A^c}\|_\infty.$$

Let $A_i \in \mathcal{F}$ be such that $\mathbb{P}(A_i) \leq \mathbb{P}(A)$, $i = 1, \dots, k$. Then $\|\bigoplus_{i=1}^k X\mathbf{1}_{A_i}\| \leq \|\bigoplus_{i=1}^k X\mathbf{1}_A\|$.

Proof. By replacing X with $|X|$, we assume that $X \geq 0$. For any $i = 1, \dots, k$, since $\mathbb{P}(A_i \cap A^c) = \mathbb{P}(A_i) - \mathbb{P}(A_i \cap A) \leq \mathbb{P}(A) - \mathbb{P}(A_i \cap A) = \mathbb{P}(A \cap A_i^c)$, take $B_i \subset A \cap A_i^c$ such that $\mathbb{P}(B_i) = \mathbb{P}(A_i \cap A^c)$. By non-atomicity, find $Z_i|_{B_i} \sim X|_{A_i \cap A^c}$. Then by (3.4), $Z_i|_{B_i} \leq \alpha \leq X|_{B_i}$. Furthermore, set $Z_i = X$ on $A_i \cap A$ and $Z_i = 0$ off $B_i \cup (A_i \cap A)$. Then $Z_i \sim X\mathbf{1}_{A_i}$ and $Z_i \leq X\mathbf{1}_A$. Since $\mathbb{P}(A) \leq \frac{1}{k}$, we can take disjoint random variables U_i 's such that $U_i \sim X\mathbf{1}_A$, $i = 1, \dots, k$. By Lemma 2.2 applied to the chain $U_i \sim X\mathbf{1}_A \geq Z_i$, there exist random variables V_i 's such that $U_i \geq V_i \sim Z_i$, $i = 1, \dots, k$. In particular, V_i 's are disjoint. Then $0 \leq \bigoplus_{i=1}^k X\mathbf{1}_{A_i} = \sum_{i=1}^k V_i \leq \sum_{i=1}^k U_i = \bigoplus_{i=1}^k X\mathbf{1}_A$ so that $\|\bigoplus_{i=1}^k X\mathbf{1}_{A_i}\| \leq \|\bigoplus_{i=1}^k X\mathbf{1}_A\|$. \square

For two random variables X, Y , write

$$Y \prec X \quad \text{if} \quad \int_0^s Y^* dt \leq \int_0^s X^* dt \quad \text{for all } s \in [0, 1].$$

Slightly modifying the proof of [4, Ch. 2, Corollary 4.7], one obtains that if \mathcal{X} is monotonically complete and two random variables X, Y satisfy that $X \in \mathcal{X}$ and $Y \prec X$, then $Y \in \mathcal{X}$ and $\|Y\| \leq M\|X\|$, where M is the constant in (1.2).

Lemma 3.4. *Let X, X_1, \dots, X_k be non-negative random variables over a non-atomic probability space $(\Omega, \mathcal{F}, \mathbb{P})$ such that $\mathbb{P}(X \neq 0) \leq \frac{1}{k}$ and $X_i \sim X$ for $i = 1, \dots, k$. Let $\lambda_i \geq 0$, $i = 1, \dots, k$, be such that $\sum_{i=1}^k \lambda_i = 1$. Then*

$$\frac{1}{k} \bigoplus_{i=1}^k X \prec \sum_{i=1}^k \lambda_i X_i.$$

If in addition X lies in a monotonically complete r.i. space \mathcal{X} over Ω , then $\|\frac{1}{k} \bigoplus_{i=1}^k X\| \leq M\|\sum_{i=1}^k \lambda_i X_i\|$, where M is the constant in (1.2) for \mathcal{X} .

Proof. For any $s \in [0, 1]$, by [4, Ch. 2, Lemma 2.5], there exists a measurable set A_i with $\mathbb{P}(A_i) = \frac{s}{k}$ such that

$$\int_{A_i} X_i = \int_0^{\frac{s}{k}} X_i^* dt.$$

Let $A = \bigcup_{i=1}^k A_i$. Then $\mathbb{P}(A) \leq s$. Hence by [4, Ch. 2, Lemma 2.1],

$$\begin{aligned} \int_0^s \left(\sum_{i=1}^k \lambda_i X_i \right)^* dt &\geq \int_A \sum_{i=1}^k \lambda_i X_i \geq \sum_{i=1}^k \lambda_i \int_{A_i} X_i \\ &= \sum_{i=1}^k \lambda_i \int_0^{\frac{s}{k}} X_i^* dt = \int_0^{\frac{s}{k}} X^* dt \\ &= \int_0^s \left(\frac{1}{k} \bigoplus_{i=1}^k X \right)^* dt, \end{aligned}$$

where the last equality follows from the fact $(\frac{1}{k} \bigoplus_{i=1}^k X)^*(t) = \frac{1}{k} X^*(\frac{t}{k})$. This proves the first assertion.

The additional assertion follows from the remark preceding the lemma. \square

Proof of Proposition 3.2; the second assertion. Assume that (3.2) holds. Let $X \in \mathcal{X}$, $\varepsilon > 0$, and $A_n \downarrow \emptyset$ with $\sum_{n=1}^{\infty} \mathbb{P}(A_n) \leq 1$ be given. Note that (3.3) is equivalent to finding n_1, \dots, n_k such that $\|\frac{1}{k} \bigoplus_{i=1}^k X \mathbf{1}_{A_{n_i}}\| < \varepsilon$. If $X \in \mathcal{X}_a$ (in particular, if \mathcal{X} is order continuous), then $\lim_{\mathbb{P}(A) \rightarrow 0} \|X \mathbf{1}_A\| = 0$. Thus it is easy to find the desired n_i 's. For the following, we assume that $X \notin \mathcal{X}_a$ and \mathcal{X} is monotonically complete.

Take a strictly increasing sequence of positive numbers $(\alpha_m)_{m=1}^{\infty}$ such that $\sum_{m=1}^{\infty} \mathbb{P}(|X| \geq \alpha_m) \leq 1$. By (3.2) applied to $|X|$ and the sequence of measurable sets $(\{|X| \geq$

$\alpha_m\}_{m=1}^\infty$, we obtain natural numbers $m_1 < \dots < m_k$, random variables $(Z_i)_{i=1}^k$ and a convex combination $\sum_{i=1}^k \lambda_i Z_i$ such that

$$Z_i \sim |X| \mathbf{1}_{\{|X| \geq \alpha_{m_i}\}} \text{ for } i = 1, \dots, k \quad \text{and} \quad \left\| \sum_{i=1}^k \lambda_i Z_i \right\| < \frac{\varepsilon}{M},$$

where M is the constant in (1.2) for \mathcal{X} . For $i = 1, \dots, k$, since $Z_i \sim |X| \mathbf{1}_{\{|X| \geq \alpha_{m_i}\}} \geq |X| \mathbf{1}_{\{|X| \geq \alpha_{m_k}\}}$, by Lemma 2.2, there exists a random variable Z'_i such that $Z_i \geq Z'_i$ and

$$Z'_i \sim |X| \mathbf{1}_{\{|X| \geq \alpha_{m_k}\}}.$$

Since $0 \leq Z'_i \leq Z_i$ for each i ,

$$\left\| \sum_{i=1}^k \lambda_i Z'_i \right\| < \frac{\varepsilon}{M}.$$

Clearly, $k\mathbb{P}(|X| \geq \alpha_{m_k}) \leq \sum_{i=1}^k \mathbb{P}(|X| \geq \alpha_{m_i}) \leq 1$ so that $\mathbb{P}(|X| \geq \alpha_{m_k}) \leq \frac{1}{k}$. By Lemma 3.4 applied to $|X| \mathbf{1}_{\{|X| \geq \alpha_{m_k}\}}$ and Z'_i 's,

$$\left\| \frac{1}{k} \bigoplus_{i=1}^k |X| \mathbf{1}_{\{|X| \geq \alpha_{m_k}\}} \right\| < \varepsilon.$$

Since $X \notin \mathcal{X}_a$, $X \notin L^\infty$, consequently $\mathbb{P}(|X| \geq \alpha_{m_k}) > 0$. Thus we can take $n_1, \dots, n_k \in \mathbb{N}$ such that $\mathbb{P}(A_{n_i}) \leq \mathbb{P}(|X| \geq \alpha_{m_k})$ for $i = 1, \dots, k$. By Lemma 3.3 applied to $A = \{|X| \geq \alpha_{m_k}\}$, we obtain $\left\| \frac{1}{k} \bigoplus_{i=1}^k X \mathbf{1}_{A_{n_i}} \right\| < \varepsilon$, as desired. \square

Remark 3.5. It was shown in [6] that the AOCEA property (3.1) is satisfied by nearly all classical r.i. spaces other than L^∞ , including Lebesgue spaces L^p ($1 \leq p < \infty$), Lorentz spaces $L^{p,q}$ ($1 < p < \infty, 1 \leq q \leq \infty$) and Orlicz spaces. Proposition 3.2 gives another class of r.i. spaces with the AOCEA property: Let \mathcal{X} be an r.i. space over a non-atomic probability space other than L^∞ . If \mathcal{X} satisfies an upper p -estimate for some $p \in (1, \infty)$ (see, e.g., [16, Definition 1.f.4]), then it clearly satisfies (3.3) and hence (3.2) and (3.1). By Proposition 3.1, law-invariant bounded linear functionals on all of these spaces collapse to the mean.

The AOCEA property (3.1), however, is not universally satisfied by all r.i. spaces. In the next section, we construct an r.i. space on which there is a positive law-invariant linear functional that does not collapse to the mean and thus, in particular, (3.1) fails in this space.

4. A COUNTEREXAMPLE

For simplicity, we work on $[0, 1]$ endowed with the Lebesgue measure. Let \mathcal{X} be the space of all random variables X on $[0, 1]$ such that

$$\|X\| := \sup_{n \in \mathbb{N}} \left(n 2^n \int_0^{\frac{1}{2^n \cdot n!}} X^* dt \right) < \infty.$$

Then $(\mathcal{X}, \|\cdot\|)$ is an r.i. space over $[0, 1]$. Indeed, law-invariance of \mathcal{X} and $\|\cdot\|$ is obvious. Other defining properties of an r.i. space can be easily verified using results in [4, Ch. 2], in particular, Proposition 1.7 there. In fact, $\|\cdot\|$ is obviously positive homogeneous. To verify the triangle inequality, recall from [4, P. 54] that for any $s \in [0, 1]$ and any random variables X, Y ,

$$\int_0^s (X + Y)^* dt \leq \int_0^s X^* dt + \int_0^s Y^* dt.$$

It is then immediate that if $X, Y \in \mathcal{X}$ then $X + Y \in \mathcal{X}$ and $\|X + Y\| \leq \|X\| + \|Y\|$. Next, we check that \mathcal{X} is complete. Since X^* is decreasing and $X^* \sim |X|$,

$$\|X\| \geq 2 \int_0^{\frac{1}{2}} X^* dt \geq \int_0^1 X^* dt = \|X\|_1.$$

Thus if X_n 's are random variables such that $\sup_{n \in \mathbb{N}} \|X_n\| < \infty$ and $0 \leq X_n \uparrow$ a.s., then $\sup_{n \in \mathbb{N}} \|X_n\|_1 < \infty$, so that there exists a random variable $X \in L^1$ such that $X_n \uparrow X$ a.s. By [4, Ch. 2, Proposition 1.7], $X_n^*(t) \uparrow X^*(t)$ for any $t \in (0, 1)$, so that $\int_0^s X_n^* dt \uparrow \int_0^s X^* dt$ for any $s \in [0, 1]$. It follows that $\|X_n\| \uparrow \|X\|$. To sum up, we have

$$(4.1) \quad \text{if } \sup_{n \in \mathbb{N}} \|X_n\| < \infty \text{ and } 0 \leq X_n \uparrow, \text{ then } X := \lim_n X_n \in \mathcal{X} \text{ and } \|X_n\| \uparrow \|X\|.$$

This in turn implies the Riesz-Fisher property, i.e, if (X_n) is a sequence in \mathcal{X} such that $\sum_{n=1}^{\infty} \|X_n\| < \infty$, then $\sum_{n=1}^{\infty} X_n$ converges in \mathcal{X} in norm (also almost surely). It is well known that the Riesz-Fisher property implies completeness of the norm. Furthermore, (4.1) actually says that $(\mathcal{X}, \|\cdot\|)$ is monotonically complete.

Following the notation in Section 2, set $\mathcal{S} = \text{Span}\{U - V : U, V \in \mathcal{X}, U \sim V\}$ and $\mathcal{M} = \{X \in \mathcal{X} : \mathbb{E}[X] = 0\}$. We show that \mathcal{S} is not dense in \mathcal{M} . This, together with Proposition 2.8, implies that some law-invariant positive linear functional on \mathcal{X} does not collapse to the mean. Consequently, \mathcal{X} fails (3.1) and (3.2) by Proposition 3.1 and also fails (3.3) by Proposition 3.2.

We begin with the following lemma.

Lemma 4.1. *Let $m, n \in \mathbb{N}$ and $\varepsilon > 0$ be given. Suppose that Y, Z, U_i, V_i ($1 \leq i \leq m$) are random variables in \mathcal{X}_+ such that $\|Z\| < \varepsilon$, $U_i \sim V_i$, $i = 1, \dots, m$, and that*

$$U := \sum_{i=1}^m U_i \geq Y - Z + \sum_{i=1}^m V_i.$$

Then for any measurable set A with $\mathbb{P}(A) \leq \frac{1}{(m+1)2^{n!}}$, there exists a measurable set A' containing A such that $\mathbb{P}(A') \leq (m+1)\mathbb{P}(A)$ and that

$$\mathbb{E}[U\mathbf{1}_{A'}] - \mathbb{E}[U\mathbf{1}_A] \geq \mathbb{E}[Y\mathbf{1}_A] - \frac{\varepsilon}{n2^n}.$$

Proof. For $1 \leq i \leq m$, by Lemma 2.2, there exists a random variable W_i such that $U_i \mathbf{1}_A \sim W_i \leq V_i$. Set $A' = A \cup \bigcup_{i=1}^m \{W_i \neq 0\}$. Then $\mathbb{P}(A') \leq (m+1)\mathbb{P}(A)$. Since $V_i \mathbf{1}_{A'} \geq W_i \mathbf{1}_{A'} = W_i$, we have

$$\begin{aligned} \mathbb{E}[U \mathbf{1}_{A'}] &\geq \mathbb{E}[Y \mathbf{1}_{A'}] + \sum_{i=1}^m \mathbb{E}[V_i \mathbf{1}_{A'}] - \mathbb{E}[Z \mathbf{1}_{A'}] \\ &\geq \mathbb{E}[Y \mathbf{1}_{A'}] + \sum_{i=1}^m \mathbb{E}[W_i] - \mathbb{E}[Z \mathbf{1}_{A'}] \\ &= \mathbb{E}[Y \mathbf{1}_{A'}] + \sum_{i=1}^m \mathbb{E}[U_i \mathbf{1}_A] - \mathbb{E}[Z \mathbf{1}_{A'}] \\ &\geq \mathbb{E}[Y \mathbf{1}_A] + \mathbb{E}[U \mathbf{1}_A] - \mathbb{E}[Z \mathbf{1}_{A'}]. \end{aligned}$$

Since $\mathbb{P}(A') \leq (m+1)\mathbb{P}(A) \leq \frac{1}{2^{n n!}}$, we have, by definition of $\|\cdot\|$,

$$\mathbb{E}[Z \mathbf{1}_{A'}] \leq \int_0^{\frac{1}{2^{n n!}}} Z^* dt \leq \frac{1}{n 2^n} \cdot \|Z\| \leq \frac{\varepsilon}{n 2^n},$$

where the first inequality is due to [4, Ch. 2, Lemma 2.1]. This proves the lemma. \square

Iterating the lemma k times gives the following lemma.

Lemma 4.2. *Let $m, n, k \in \mathbb{N}$ and $\varepsilon > 0$ be given. Suppose that Y, Z, U_i, V_i ($1 \leq i \leq m$) are random variables in \mathcal{X}_+ such that $\|Z\| < \varepsilon$, $U_i \sim V_i$, $i = 1, \dots, m$, and that*

$$U := \sum_{i=1}^m U_i \geq Y - Z + \sum_{i=1}^m V_i.$$

For any measurable set A with $\mathbb{P}(A) \leq \frac{1}{(m+1)^k 2^{n n!}}$, there exists a measurable set A' containing A such that $\mathbb{P}(A') \leq (m+1)^k \mathbb{P}(A)$ and that

$$\mathbb{E}[U \mathbf{1}_{A'}] \geq k \mathbb{E}[Y \mathbf{1}_A] + \mathbb{E}[U \mathbf{1}_A] - \frac{k\varepsilon}{n 2^n}.$$

We are ready to complete the proof of that $\overline{\mathcal{S}} \subsetneq \mathcal{M}$.

Proof of $\overline{\mathcal{S}} \subsetneq \mathcal{M}$. Set

$$c_n = \frac{1}{2^n \cdot (n+1)!} \quad \text{for any } n \in \mathbb{N},$$

and

$$Y = \sum_{n=3}^{\infty} n! \mathbf{1}_{[c_{n+1}, c_n]}.$$

We claim that $Y \in \mathcal{X}$. Indeed, since Y is non-negative and decreasing on $(0, 1)$, $Y^* = Y$ a.s. For any $l \geq 4$, since $c_l < \frac{1}{2^{l \cdot l}} < c_{l-1}$,

$$\int_0^{\frac{1}{2^{l \cdot l}}} Y^* dt \leq \frac{(l-1)!}{2^l \cdot l!} + \sum_{n=l}^{\infty} n! c_n = \frac{1}{l2^l} + \sum_{n=l}^{\infty} \frac{1}{2^n(n+1)} \leq \frac{3}{l2^l}.$$

It follows that $Y \in \mathcal{X}$, as claimed. Clearly, $X := Y - \mathbb{E}[Y] \cdot \mathbf{1} \in \mathcal{M}$.

We claim that $X \notin \overline{\mathcal{S}}$. Assume on the contrary that $X \in \overline{\mathcal{S}}$. By splitting a random variable into the positive and negative parts, it is easy to see that $\mathcal{S} = \text{Span}\{U - V : U, V \in \mathcal{X}, U \sim V, U, V \geq 0\}$. Thus there are sequences $(U_i)_{i=1}^m, (V_i)_{i=1}^m$ in \mathcal{X}_+ and $W \in \mathcal{X}$ with $\|W\| < \frac{1}{8}$ such that $U_i \sim V_i$ for $i = 1, \dots, m$ and $X = \sum_{i=1}^m (U_i - V_i) + W$. Hence

$$U := \sum_{i=1}^m U_i \geq Y - Z + \sum_{i=1}^m V_i,$$

where $Z = \mathbb{E}[Y] \cdot \mathbf{1} + W^+$. Observe that $\mathbb{E}[Y] = \sum_{n=3}^{\infty} n!(c_n - c_{n+1}) \leq \sum_{n=3}^{\infty} \frac{1}{2^n(n+1)} < \frac{1}{8}$ and that $\|\mathbf{1}\| = 1$. Hence

$$\|Z\| \leq \|\mathbb{E}[Y] \cdot \mathbf{1}\| + \|W\| < \frac{1}{4}.$$

For any $k \in \mathbb{N}$, let $n \geq 6$ be such that

$$c_n - c_{n+1} = \frac{n + \frac{3}{2}}{2^n \cdot (n+2)!} \leq \frac{1}{(m+1)^k 2^n \cdot n!}.$$

Taking $A = (c_{n+1}, c_n)$ and $\varepsilon = \frac{1}{4}$ and applying Lemma 4.2, we obtain a measurable set A' such that $\mathbb{P}(A') \leq (m+1)^k \mathbb{P}(A) \leq \frac{1}{2^n \cdot n!}$ and that

$$\mathbb{E}[U \mathbf{1}_{A'}] \geq k \mathbb{E}[Y \mathbf{1}_A] - \frac{k}{4n2^n} = \frac{k}{2^n} \left(\frac{n + \frac{3}{2}}{(n+1)(n+2)} - \frac{1}{4n} \right) > \frac{k}{2^{n+2} n}.$$

Therefore,

$$\|U\| \geq n2^n \int_0^{\frac{1}{2^n \cdot n!}} U^* dt \geq n2^n \mathbb{E}[U \mathbf{1}_{A'}] \geq \frac{k}{4}.$$

This cannot hold for all $k \in \mathbb{N}$ and concludes the proof. \square

APPENDIX A. AN ALTERNATIVE PROOF OF PROPOSITION 3.1

Alternative proof. Let ρ be a positive law-invariant linear functional on \mathcal{X} . Recall from Lemma 2.1 that $\rho(X) = \rho(\mathbf{1})\mathbb{E}(X)$ for all $X \in \mathcal{X}_a$. Let $X \in \mathcal{X}_+$. By [6, Theorem 2.2], $-\rho$ has the Fatou property at 0, i.e., $-\rho(X) \leq \liminf_n (-\rho(X_n))$ whenever $X_n \xrightarrow{a.s.} 0$ and there exists $X_0 \in \mathcal{X}$ such that $|X_n| \leq X_0$ for all $n \geq 1$. In particular, take $X_n = X \mathbf{1}_{\{X \geq n\}}$ for $n \geq 1$. Since the sequence $(-\rho(X \mathbf{1}_{\{X \geq n\}}))$ is increasing,

$$0 = -\rho(0) \leq \liminf_n -\rho(X \mathbf{1}_{\{X \geq n\}}) = -\lim_n \rho(X \mathbf{1}_{\{X \geq n\}}) \leq 0.$$

Thus $\lim_n \rho(X\mathbf{1}_{\{X \geq n\}}) = 0$. Therefore, $\rho(X) = \lim_n [\rho(X\mathbf{1}_{\{X < n\}}) + \rho(X\mathbf{1}_{\{X \geq n\}})] = \lim_n \rho(X\mathbf{1}_{\{X < n\}})$. Since $X\mathbf{1}_{\{X < n\}} \in L^\infty \subset \mathcal{X}_a$,

$$\rho(X) = \lim_n \rho(X\mathbf{1}_{\{|X| < n\}}) = \lim_n \rho(\mathbf{1})\mathbb{E}(X\mathbf{1}_{\{|X| < n\}}) = \rho(\mathbf{1})\mathbb{E}(X).$$

Since $\rho(X) = \rho(\mathbf{1})\mathbb{E}(X)$ for all $X \in \mathcal{X}_+$, the same equation holds for all $X \in \mathcal{X}$ by linearity of ρ . For a general ρ , invoke Lemma 2.3 to consider ρ^\pm . \square

REFERENCES

- [1] Aliprantis, C.D., Burkinshaw, O.: *Positive Operators*, Springer, Dordrecht (2006)
- [2] Bellini, F., Koch-Medina, P., Munari, C., Svindland, G.: Law-invariant functionals on general spaces of random variables, *SIAM Journal on Financial Mathematics*, 12(1), 318-341 (2021)
- [3] Bellini, F., Koch-Medina, P., Munari, C., Svindland, G.: Law-invariant functionals that collapse to the mean, *Insurance: Mathematics and Economics*, 98, 83-91 (2021)
- [4] Bennet, C., Sharpley, R.: *Interpolation of Operators*, Academic Press, Inc., Boston (1988)
- [5] Chen, S., Gao, N., Xanthos, F.: On the strong Fatou property of risk measures, *Dependence Modeling*, 6(1), 183-196 (2018)
- [6] Chen, S., Gao, N., Leung, D., Lei, L: Automatic Fatou property of law-invariant risk measures. Preprint available at *arXiv:2107.08109*
- [7] Filipović, D., Svindland, G.: Optimal capital and risk allocations for law- and cash-invariant convex functions, *Finance and Stochastics*, 12(3), 423-439 (2008)
- [8] Filipović, D., Svindland, G.: The canonical model space for law-invariant convex risk measures is L^1 , *Mathematical Finance*, 22(3), 585-589 (2012).
- [9] Föllmer, H., Schied, A.: *Stochastic Finance An Introduction in Discrete Time*, 4th Edition, de Gruyter, Berlin/Boston (2016)
- [10] Gao, N., Leung, D., Munari, C., Xanthos, F.: Fatou property, representations, and extensions of law-invariant risk measures on general Orlicz spaces, *Finance and Stochastics*, 22, 395-415 (2018)
- [11] Liebrich, F., Munari, C.: Law-invariant functionals that collapse to the mean: beyond convexity. Preprint available at *arXiv:2106.01281*
- [12] Jouini, E., Schachermayer, W., Touzi, N.: Law invariant risk measures have the Fatou property, *Adv. Math. Econ.*, 9, 49-71 (2006)
- [13] Jouini E., Schachermayer W., Touzi N.: Optimal risk sharing for law invariant monetary utility functions, *Mathematical Finance*, 18(2), 269-292 (2008)
- [14] Krättschmer, V, Schied, A., Zähle, H.: Comparative and qualitative robustness for law-invariant risk measures, *Finance and Stochastics*, 18, 271-295 (2014)
- [15] Kusuoka, S.: On law invariant coherent risk measures, *Advances in Mathematical Economics*, 3, 83-95 (2001)
- [16] Lindenstrauss, J., Tzafriri, L.: *Classical Banach Spaces II*, Springer-Verlag, Berlin (1979)
- [17] Liu, P., Wang, R., Wei, L.: Is the inf-convolution of law-invariant preferences law-invariant? *Insurance: Mathematics and Economics*, 91, 144-154 (2020)
- [18] Meyer-Nieberg, P.: *Banach Lattices*, Springer-Verlag, Berlin (1991)
- [19] Svindland, G.: Continuity properties of law-invariant (quasi-)convex risk functions on L^∞ , *Mathematics and Financial Economics*, 3(1), 39-43 (2010)

- [20] Tantrawan, M., Leung, D.: On closedness of law-invariant convex sets in rearrangement invariant spaces, *Archiv der Mathematik*, 114(2), 175-183 (2020)
- [21] Wang, R., Zitikis, R.: An axiomatic foundation for the Expected Shortfall, *Management Science*, 67(3), 1413-1429 (2021)
- [22] Weber S.: Distribution-invariant risk measures, information, and dynamic consistency, *Mathematical Finance*, 16, 419-441 (2006)

DEPARTMENT OF MATHEMATICS, RYERSON UNIVERSITY, 350 VICTORIA STREET, TORONTO, CANADA M5B2K3

Email address: sz.chen@ryerson.ca

DEPARTMENT OF MATHEMATICS, RYERSON UNIVERSITY, 350 VICTORIA STREET, TORONTO, CANADA M5B2K3

Email address: niushan@ryerson.ca

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 117543

Email address: dennyhl@u.nus.edu

SCHOOL OF MATHEMATICAL SCIENCES AND LPMC, NANKAI UNIVERSITY, TIANJIN 300071, CHINA

Email address: leilee@nankai.edu.cn