

ROUGH CAUCHY SEQUENCES IN A CONE METRIC SPACE

RAHUL MONDAL

ABSTRACT. Here we have introduced the idea of rough Cauchyness of sequences in a cone metric space. Also here we have discussed several basic properties of rough Cauchy sequences in a cone metric space using the idea of Phu [8].

1. Introduction

Metric spaces have been generalized in many ways. Huang and Zhang [5] introduced the idea of cone metric spaces where the idea of distance have been generalized to a vector through the idea of a cone defined in a ordered Banach space.

Phu [8] introduced the idea of rough convergence of sequences as a generalization of ordinary convergence of sequences in a normed linear space in 2001. There he also introduced the idea of rough Cauchyness of sequences as a generalization of Cauchyness of sequences. Phu [8] discussed about the usefulness of rough convergence and some of its basic properties. Phu [9] also extended the idea of rough convergence of sequences in an infinite dimensional normed linear space in 2003 and also in 2008 Ayter [1] introduced the idea of rough statistical convergence. Recently Banerjee and Mondal [3] studied the idea of rough convergence of sequences in a cone metric space. Many works have been done by many authors [2, 6, 7] using the idea of Phu. Here we have discussed the rough Cauchyness of sequences in a cone metric space.

2. Preliminaries

Definition 2.1. [8] *Let $\{x_n\}$ be a sequence in a normed linear space $(X, \|\cdot\|)$, and r be a nonnegative real number. Then $\{x_n\}$ is said to be r -convergent to x if for any $\epsilon > 0$, there exists a natural number k such that $\|x_n - x\| < r + \epsilon$ for all $n \geq k$.*

Definition 2.2. [5] *Let E be a real Banach space and P be a subset of E . Then P is called a cone if and only if (i) P is closed nonempty, and $P \neq \{0\}$*

(ii) $a, b \in \mathbb{R}$, $a, b \geq 0$, $x, y \in P$ implies $ax + by \in P$.

(iii) $x \in P$ and $-x \in P$ implies $x = 0$.

⁰2010 AMS Subject Classification: 40A05, 40A99.

Key words and phrases: Rough convergence, rough Cauchy sequence, cone, cone metric space.

Let E be a real Banach space and P be a cone in E . Let us use the partial ordering [5] with respect to P by $x \leq y$ if and only if $y - x \in P$. We shall write $x < y$ to indicate that $x \leq y$ but $x \neq y$.

Also by $x \ll y$, we mean $y - x \in \text{int}P$, the interior of P . The cone P is called normal if there is a number $K > 0$ such that for all $x, y \in E$, $0 \leq x \leq y$ implies $\|x\| \leq K\|y\|$.

Definition 2.3. [5] *Let X be a non empty set. If the mapping $d : X \times X \rightarrow E$ satisfies the following three conditions*

(d1) $0 \leq d(x, y)$ for all $x, y \in X$ and $d(x, y) = 0$ if and only if $x = y$;

(d2) $d(x, y) = d(y, x)$ for all $x, y \in X$;

(d3) $d(x, y) \leq d(x, z) + d(z, y)$ for all $x, y, z \in X$; then d is called a cone metric on X , and (X, d) is called a cone metric space.

It is clear that a cone metric space is a generalization of metric spaces. Throughout (X, d) or simply X stands for a cone metric space which is associated with a real Banach space E with a cone P , \mathbb{R} for the set of all real numbers, \mathbb{N} for the set of all natural numbers, sets are always subsets of X unless otherwise stated.

Definition 2.4. [5] *Let (X, d) be a cone metric space. A sequence $\{x_n\}$ in X is said to be convergent to $x \in X$ if for every $c \in E$ with $0 \ll c$ there is $k \in \mathbb{N}$ such that $d(x_n, x) \ll c$, whenever for all $n > k$.*

We know that in a real Banach space E with cone P . If $x_0 \in \text{int}P$ and $c(> 0) \in \mathbb{R}$ then $cx_0 \in \text{int}P$ and if $x_0 \in P$ and $y_0 \in \text{int}P$ then $x_0 + y_0 \in \text{int}P$. Hence we can also say that if $x_0, y_0 \in \text{int}P$ then $x_0 + y_0 \in \text{int}P$. Also it has been discussed in [3] that a real normed linear space is always connected and if E be a real Banach space with cone P then $0 \notin \text{int}P$.

Definition 2.5. [8, 3] *Let (X, d) be a cone metric space. A sequence $\{x_n\}$ in X is said to be r -convergent to x for some $r \in E$ with $0 \ll r$ or $r = 0$ if for every ϵ with $0 \ll \epsilon$ there exists a $k \in \mathbb{N}$ such that $d(x_n, x) \ll r + \epsilon$ for all $n \geq k$.*

Let (X, d) be a cone metric space with normal cone P and normal constant k . Then from [3] we can say for every $\epsilon > 0$, we can choose $c \in E$ with $c \in \text{int}P$ and $k\|c\| < \epsilon$ and also for each $c \in E$ with $0 \ll c$, there is a $\delta > 0$, such that $\|x\| < \delta$ implies $c - x \in \text{int}P$. We will use this idea in the next section of our work.

3. r -Cauchy sequences in a cone metric space

Definition 3.1. Let (X, d) be a cone metric space. A sequence $\{x_n\}$ in X is said to be a Cauchy sequence in X if for every $(0 \ll) \epsilon$ there exists a natural number m such that $d(x_i, x_j) \ll \epsilon$ for all $i, j \geq m$.

Definition 3.2. A sequence $\{x_n\}$ in a cone metric space (X, d) is said to be a r -Cauchy sequence for some $(0 \ll) r$ or $r = 0$ in this space if for every $(0 \ll) \epsilon$ there exists a natural number m such that $d(x_i, x_j) \ll r + \epsilon$ for all $i, j \geq m$.

It should be noted that when $r = 0$ then rough Cauchyness coincides with the Cauchyness.

Theorem 3.1. Every $\frac{r}{2}$ -convergent sequence in a cone metric space (X, d) is r -Cauchy for every r as defined above.

Proof. Let $\{x_n\}$ be a $\frac{r}{2}$ -convergent sequence in a cone metric space (X, d) and converges to x in X and consider a arbitrary $\epsilon \in \text{int}P$. Now for the $(0 \ll) \epsilon$ there exists a natural number m such that $d(x_n, x) \ll \frac{r}{2} + \frac{\epsilon}{2}$ for all $n \geq m$, so $[\frac{r}{2} + \frac{\epsilon}{2}] - d(x_n, x) \in \text{int}P$ for all $n \geq m$. Now if $r \in \text{int}P$ then for any two natural numbers i, j we have $d(x_i, x_j) \leq d(x_i, x) + d(x_j, x) \rightarrow (i)$. So $[d(x_i, x) + d(x_j, x)] - d(x_i, x_j) \in P$. Also for $i \geq m$ and $j \geq m$ we have $[\frac{r}{2} + \frac{\epsilon}{2}] - d(x_i, x) \in \text{int}P$ and $[\frac{r}{2} + \frac{\epsilon}{2}] - d(x_j, x) \in \text{int}P$. Hence $[r + \epsilon] - [d(x_i, x) + d(x_j, x)] \in \text{int}P \rightarrow (ii)$ Therefore using (i) and (ii) we have $d(x_i, x_j) \ll r + \epsilon$. \square

The idea of boundedness of a sequence in cone metric spaces is similar as in metric spaces and this has been discussed in [3]. We recall that a sequence $\{x_n\}$ in a cone metric space (X, d) is bounded if there is a $g \in \text{int}P$ such that $d(x_m, x_n) \ll g$ for all $m \in \mathbb{N}$ and $n \in \mathbb{N}$.

Theorem 3.2. A bounded sequence in a cone metric space is always r -Cauchy for some r as defined above.

Proof. Let $\{x_n\}$ be a bounded sequence in a cone metric space (X, d) . So there exists a $(0 \ll) s$ such that $d(x_n, x_m) \ll s$ for all $m, n \in \mathbb{N}$. Therefore $[s - d(x_n, x_m)] \in \text{int}P$ for all $m, n \in \mathbb{N}$. Hence for any $(0 \ll) \epsilon$ we have $[s - d(x_n, x_m)] + \epsilon \in \text{int}P$ for all $m, n \in \mathbb{N}$. Therefore $d(x_n, x_m) \ll s + \epsilon$ for all $m, n \in \mathbb{N}$. So $\{x_n\}$ is a s -Cauchy sequence in X . \square

Theorem 3.3. Let (X, d) be a cone metric space with normal cone P and normal constant k and with the given condition $*$ as given bellow. If $\{x_n\}$ and $\{y_n\}$ be two $\frac{r}{2k^2}$ -Cauchy sequence $(0 \ll r)$ in X then the sequence $\{d(x_n, y_n)\}$ is $\|r\|$ -Cauchy.

(*) If $p \leq c$ and $-p \leq c$ with $c \in P$ then $\|p\| \leq k\|c\|$, where $p, c \in E$.

Proof. Let $\epsilon > 0$ be preassigned real number. So by using the property of a normal cone there exists a $c \in \text{int}P$ such that $k\|c\| < \epsilon$. Since $c \in \text{int}P$, we have $\frac{c}{2} \in \text{int}P$ and also $\frac{r}{2k^2} \in \text{int}P$. Now there exists two positive integers k_1 and k_2 such that $d(x_i, x_j) \ll \frac{r}{2k^2} + \frac{c}{2}$ for all $i, j \geq k_1$ and $d(y_i, y_j) \ll \frac{r}{2k^2} + \frac{c}{2}$ for all $i, j \geq k_2$. If k be the maximum of k_1 and k_2 then by using the property of normal cone we have $\|d(x_i, x_j)\| \leq k\|\frac{r}{2k^2} + \frac{c}{2}\|$ and $\|d(y_i, y_j)\| \leq k\|\frac{r}{2k^2} + \frac{c}{2}\|$ for all $i, j \geq k$.

Now $d(x_i, y_i) - d(x_j, y_j) \leq d(x_i, x_j) + d(y_i, y_j)$ and $d(x_j, y_j) - d(x_i, y_i) \leq d(x_i, x_j) + d(y_i, y_j)$. Since $[d(x_i, x_j) + d(y_i, y_j)] \in P$, by the given condition we have $\|d(x_i, y_i) - d(x_j, y_j)\| \leq k\|d(x_i, x_j) + d(y_i, y_j)\| \leq k\|d(x_i, x_j)\| + k\|d(y_i, y_j)\|$. Hence for all $i, j \geq k$ we have $\|d(x_i, y_i) - d(x_j, y_j)\| \leq \|r\| + k\|c\| < \|r\| + \epsilon$. \square

Theorem 3.4. Let $\{x_n\}$ and $\{y_n\}$ be two sequences in a cone metric space (X, d) such that $d(x_n, y_n) \rightarrow 0$ as $n \rightarrow \infty$. Then $\{x_n\}$ is r -Cauchy if and only if $\{y_n\}$ is r -Cauchy.

Proof. Let $\{x_n\}$ be a r -Cauchy sequence in (X, d) . Then for $0 \ll \epsilon$ there exists two positive integers k_1 and k_2 such that $d(x_i, x_j) \ll r + \frac{\epsilon}{3}$ for all $i, j \geq k_1$ and $d(x_i, y_i) \ll \frac{\epsilon}{3}$ for all $i \geq k_2$. Now $d(y_i, y_j) \leq d(x_i, y_i) + d(x_i, y_j)$ and also $d(x_i, y_j) \leq d(x_i, x_j) + d(x_j, y_j)$. Hence $d(x_j, y_j) \leq d(x_i, y_i) + d(x_i, x_j) + d(x_j, y_j)$. If $k = \max(k_1, k_2)$, then for all $i, j \geq k$ we have $d(x_i, y_j) \ll r + \epsilon$ and hence $\{y_n\}$ is r -Cauchy.

Conversely if $\{y_n\}$ is r -Cauchy then similarly we can show that $\{x_n\}$ is r -Cauchy. \square

Theorem 3.5. Let (X, d) be a cone metric space with normal cone P and normal constant k . If a sequence $\{x_n\}$ in (X, d) is r -Cauchy and also converges to x in X then the sequence $\{d(x_n, x) - r\}$ is converges to 0 in E , provided that $\{d(x_n, x) - r\}$ is a sequence in P .

Proof. Let $\{x_n\}$ be a r -Cauchy sequence and converges to x in X . Let $\epsilon > 0$ be preassigned. Then we have an element $c \in E$ with $0 \ll c$ and $k\|c\| < \epsilon$. Now for $0 \ll c$ there exists $k_1, k_2 \in \mathbb{N}$ such that $d(x_n, x_m) \ll r + \frac{c}{2}$ for all $n, m \geq k_1$ and $d(x_m, x) \ll \frac{c}{2}$ for all $m \geq k_2$. Hence $0 \leq d(x_n, x) - r \ll c$ for all $n \geq k$, k is the maximum of k_1 and k_2 . Now by using the property of a normal cone we have $\|d(x_n, x) - r\| \leq k\|c\| < \epsilon$ for all $n \geq k$. Therefore $\{d(x_n, x) - r\}$ converges to 0 in E . \square

REFERENCES

- [1] S. Aytar, Rough statistical convergence, Numer. Funct. Anal. and Optimiz, 29(3-4) (2008), 291-303.
- [2] S. Aytar, The rough limit set and the core of a real sequence, Numer. Funct. Anal. and Optimiz, 29(3-4) (2008), 283-290.

- [3] A. K. Banerjee, R. Mondal, Rough convergence of sequences in a cone metric space, J Anal (2019), <https://doi.org/10.1007/s41478-019-00168-2>.
- [4] Kieu Phuong Chi, Tran Van An; Dugundji's theorem for cone metric spaces, Applied Mathematics Letters, 24(2011) 387-390.
- [5] Huang Long-Gung, Zhang X: Cone metric spaces and fixed point theorems of contractive mappings, Journal of Mathematical Analysis and Applications, 332(2) (2007), 1468-1476.
- [6] P. Malik, and M. Maity, M. On rough convergence of double sequence in normed linear spaces, Bull. Allah. Math. Soc. 28(1), 89-99, 2013.
- [7] P. Malik, and M. Maity, On rough statistical convergence of double sequences in normed linear spaces, Afr. Mat. 27, 141-148, 2016.
- [8] H. X. Phu, Rough convergence in normed linear spaces, Numer. Funct. Anal. and Optimiz, 22: (2001), 199-222.
- [9] H. X. Phu, Rough convergence in infinite dimensional normed spaces, Numer. Funct. Anal. and Optimiz, 24: (2003), 285-301.

DEPARTMENT OF MATHEMATICS, VIVEKANANDA SATAVARSHIKI MAHAVIDYALAYA, MANIKPARA, JHAR-GRAM -721513, WEST BENGAL, INDIA.

Email address: imondalrahul@gmail.com, rahul04051987@gmail.com