

The Möbius Function on Affine Grassmannian elements

Michael Lugo and Mark Shimozono

December 23, 2021

Abstract

To any saturated chain in the affine Weyl group whose translation parts are sufficiently regular, we associate a near path and a far path in the quantum Bruhat graph. Using this, working in the Bruhat order on the minimal-length representatives of the cosets in the affine Weyl group with respect to the finite Weyl group, we characterize the pairs of elements for which the Möbius function is nonzero. This is applied to obtain explicit expansions in the K -theory of affine Grassmannians, of the basis of ideal sheaves into the basis of structure sheaves of Schubert varieties.

1 Introduction

1.1 Möbius Function on Affine Grassmannian elements

Let W be the Weyl group of a root system of untwisted affine type or the dual of untwisted affine type. For ease of exposition we shall work with untwisted affine type, leaving the statements of the dual case to §4. The group W is generated by simple reflections s_i for i in the affine Dynkin node set I , with distinguished affine node $0 \in I$. Say that $w \in W$ is *affine Grassmannian* (denoted $w \in W^0$) if $ws_i > w$ for all $i \in I_0 = I \setminus \{0\}$. Let $\tilde{\mu}$ be the Möbius function of the Bruhat order on W^0 . The following is the affine Grassmannian case of Deodhar's general formula [3] for the Möbius function of a parabolic quotient of the Bruhat order. For $u, v \in W$ with $u \leq v$, denote by $[u, v] = \{z \in W \mid u \leq z \leq v\}$ the interval between u and v in the Bruhat order on W .

Theorem 1 ([3, Theorem 1.2]). *For $u, v \in W^0$ with $u \leq v$,*

$$\tilde{\mu}(u, v) = \begin{cases} (-1)^{\ell(v) - \ell(u)} & \text{if } [u, v] \subset W^0 \\ 0 & \text{otherwise.} \end{cases} \quad (1)$$

Moreover the first case holds if and only if there is no $i \in I_0$ such that $us_i \leq v$.

For $y \in W^0$ let $B_y = \{x \in W^0 \mid [x, y] \subset W^0\}$. Our main theorem (Theorem 3) is an explicit description of the subposet B_y of W^0 in the case that $y \in W^0$ is

superregular, that is, its translation part is sufficiently regular. For superregular y , we show that B_y bijects with the classical Weyl group W_0 . Moreover, all of the possible poset structures on W_0 which arise this way, come from a single directed graph structure Γ on W_0 . This directed graph, first studied by Brenti, Fomin, and Postnikov [2], has become known as the quantum Bruhat graph due to its connection with quantum cohomology of flag varieties; it indicates which Schubert classes appear in the quantum product of a Schubert class with a Schubert divisor.

1.2 Quantum Bruhat Graph

For a positive classical root $\alpha \in \Phi_0^+$ let $r_\alpha \in W_0$ be the associated reflection and $\alpha^\vee$ the associated coroot. Let $\rho = (1/2) \sum_{\alpha \in \Phi_0^+} \alpha$.

The quantum Bruhat graph (QBG) [2] is the directed graph Γ with vertex set W_0 and a directed edge $w \rightarrow wr_\alpha$ for each pair $(w, \alpha) \in W_0 \times \Phi_0^+$ such that one of the following holds.

1. $\ell(wr_\alpha) = \ell(w) + 1$. Such edges are called *Bruhat* edges.
2. $\ell(wr_\alpha) = \ell(w) - \langle \alpha^\vee, 2\rho \rangle + 1$. Such edges are called *quantum* edges.

The *label* of the edge $w \rightarrow wr_\alpha$ is $\alpha \in \Phi_0^+$. The *weight* of the edge $w \rightarrow wr_\alpha$ in Γ is $\alpha^\vee$ if the edge is quantum and 0 if it is Bruhat. The *weight* of a path¹ P , denoted $\text{wt}(P)$, is the sum of the weights of its edges. For any $u, v \in W_0$, let $M(u, v)$ be the weight of any shortest path in Γ from u to v ; this is well defined by [8, Lemma 1].

Example 2. The quantum Bruhat graph of type A_2 is pictured in Figure 1. Let $w_0 = s_1 s_2 s_1$ and $\alpha_{12}^\vee = \alpha_1^\vee + \alpha_2^\vee$. The quantum edges are given in red and their weights are indicated. We have $M(w_0, s_1) = \alpha_1^\vee + \alpha_2^\vee$, which is realized by either of the shortest paths $w_0 \rightarrow \text{id} \rightarrow s_1$ or $w_0 \rightarrow s_1 s_2 \rightarrow s_1$.

1.3 Main Theorem

An element λ of the classical coroot lattice $Q^\vee$ is *superregular* if $|\langle \lambda, \alpha_i \rangle| \gg 0$ ² for all $i \in I_0$. Any element $x \in W$ can be written uniquely in the form $x = wt_\lambda$ with $w \in W_0$ and $\lambda \in Q^\vee$ where $t_\lambda \in W$ is the translation element. For $w \in W_0$ we define the affine Weyl group element wt_λ to be superregular if λ is.

Theorem 3. *Let $x, y \in W^0$ with $x = w't_{\lambda'}$, $y = wt_\lambda$ with $w, w' \in W_0$, $\lambda, \lambda' \in Q^\vee$ and λ superregular. Then*

$$\tilde{\mu}(x, y) = \begin{cases} (-1)^{\ell(y) - \ell(x)} & \text{if } \lambda' = \lambda + M(w, w') \\ 0 & \text{else} \end{cases}$$

where M is the weight of a shortest path in Γ from w to w' .

¹All our paths are directed.

²Lower bounds for superregularity are discussed in §3.




Figure 1: Quantum Bruhat Graph of type A_2

Corollary 4. *Let $y = wt_\lambda \in W^0$ be superregular. Then the set*

$$B_y = \{x \in W^0 \mid x \leq y \text{ and } \tilde{\mu}(x, y) \neq 0\}$$

is given by

$$\{ut_{\lambda+M(w,u)} \mid u \in W_0\}. \quad (2)$$

Remark 5. The poset structure on B_y can be constructed from the shortest paths from w to all the other vertices in Γ . This poset structure is given by the w -tilted Bruhat order $\preceq_w$ of [2], which is defined by $u \preceq_w v$ if u is on some shortest path in Γ from w to v .

1.4 Application to the K -theory of the affine Grassmannian

Let $\mathcal{O} = \mathbb{C}[[t]]$ be the ring of formal power series with coefficients in $\mathbb{C}$, $\mathcal{K} = \mathbb{C}((t))$ the field of formal Laurent series, G a simple Lie group over $\mathbb{C}$, $\text{Gr}_G = G(\mathcal{K})/G(\mathcal{O})$ the affine Grassmannian, $\mathcal{I}$ the Iwahori subgroup of the affine Kac-Moody group. For $x \in W^0$ let $C_x = \mathcal{I}xG(\mathcal{O})/G(\mathcal{O}) \subset \text{Gr}_G$ be the affine Grassmannian Schubert cell and $X_x = \overline{C_x} \subset \text{Gr}_G$ the Schubert variety. For $x \in W^0$ let O_x and I_x be the elements of $K(\text{Gr}_G)$ given by the K -classes of the structure sheaf of X_x and the ideal sheaf of the boundary ∂X_x .

The following is due to Kumar [5], who produced the proof (for Kac-Moody partial flag varieties) upon being asked about it by the second author.

Proposition 6 ([5]). *For any $y \in W^0$*

$$O_y = \sum_{\substack{x \in W^0 \\ x \leq y}} I_x. \quad (3)$$

By the definition of the Möbius function and Theorem 3 we deduce:

Corollary 7. *Let $y = wt_\lambda \in W^0$ be superregular. Then*

$$I_y = \sum_{u \in W_0} (-1)^{\ell(w,u)} O_{ut_{\lambda+M(w,u)}}$$

where $\ell(w, u)$ is the length of a shortest path from w to u in Γ .

2 Near and far paths in Γ associated to superregular saturated chains in W

2.1 Sufficiently Regular Bruhat covers

Say that $\lambda \in Q^\vee$ is *sufficiently regular* if $|\langle \lambda, \alpha_i \rangle| \gg 0$ for all $i \in I_0$ and that $y = wt_\lambda \in W$ is sufficiently regular if λ is³. The question of sufficient regularity to qualify will be addressed in §3. Say that $\lambda \in Q^\vee$ is *antidominant* if $\langle \lambda, \alpha_i \rangle \leq 0$ for all $i \in I_0$. Let $\tilde{Q}$ be the set of antidominant elements in $Q^\vee$.

Proposition 8 ([6, Proposition 4.1]). *Let $\lambda \in \tilde{Q}$ be sufficiently regular and $y = wt_{v\lambda}$ where $w, v \in W_0$. Then $x = yr_{v\alpha+n\delta} \leq y$ for some $\alpha \in \Phi_0^+$ and $n \in \mathbb{Z}$ if and only if one of the following conditions hold:*

1. $\ell(wv) = \ell(wvr_\alpha) - 1$ and $n = \langle \lambda, \alpha \rangle$, giving $x = wr_{v\alpha}t_{v\lambda}$.
2. $\ell(wv) = \ell(wvr_\alpha) + \langle \alpha^\vee, 2\rho \rangle - 1$ and $n = \langle \lambda, \alpha \rangle + 1$, giving $x = wr_{v\alpha}t_{v(\lambda+\alpha^\vee)}$.
3. $\ell(v) = \ell(vr_\alpha) + 1$ and $n = 0$, giving $x = wr_{v\alpha}t_{vr_\alpha(\lambda)}$.
4. $\ell(v) = \ell(vr_\alpha) - \langle \alpha^\vee, 2\rho \rangle + 1$ and $n = -1$, giving $x = wr_{v\alpha}t_{vr_\alpha(\lambda+\alpha^\vee)}$.

That is, every sufficiently regular cover in the Bruhat order on W has a corresponding edge in Γ . In the terminology introduced in [6], the first two cases are called *near*, since the chamber of the translation component remains fixed, and the last two cases *far*.

Corollary 9. *The four possibilities in the previous Proposition may be replaced by the following two:*

1. $wv \rightarrow wvr_\alpha$ is an edge E in Γ and $x = wr_{v\alpha}t_{v(\lambda+\text{wt}(E))}$. We call this a near covering.
2. $vr_\alpha \rightarrow v$ is an edge E in Γ and $x = wr_{v\alpha}t_{vr_\alpha(\lambda+\text{wt}(E))}$. We call this a far covering.

³This definition follows the definition of superregular given in [6]. In this paper, superregularity has a different requirement.

2.2 Superregular saturated Bruhat chains

We extend this idea to saturated chains in the Bruhat order. We say that $y = wt_\lambda \in W$ is *superregular* with respect to x if for every saturated chain $x \leq z_{k-1} \leq \dots \leq z_1 \leq y$ it is true that y and z_i for $i = 1, \dots, k-1$ are all sufficiently regular. We say a saturated chain C is superregular if its top element is superregular with respect to its bottom element. Suppose that

$$x \xrightarrow{\beta_k} z_{k-1} \xrightarrow{\beta_{k-1}} \dots \xrightarrow{\beta_2} z_1 \xrightarrow{\beta_1} y \quad (4)$$

is a superregular saturated chain in W , where $\beta_i \in \Phi_0^+$ is the root corresponding to α in Proposition 8 which labels the corresponding edge in Γ . Let $1 \leq f_1 < f_2 < \dots < f_i \leq k$ be the ordered subindices relating to far coverings and $1 \leq n_1 < n_2 < \dots < n_j \leq k$ be the ordered subindices relating to near coverings. Define the *far product* $r_f = r_{\beta_{f_1}} r_{\beta_{f_2}} \dots r_{\beta_{f_i}}$ and *near product* $r_n = r_{\beta_{n_1}} r_{\beta_{n_2}} \dots r_{\beta_{n_j}}$.

Lemma 10. *Suppose $x < y = wt_{v\lambda}$ with $\lambda \in \tilde{Q}$ and superregular chain as in Equation (4). Let r_f and r_n be defined as above. Then there exist well defined paths $P_f : vr_f \rightarrow v$ and $P_n : wv \rightarrow wvr_n$ such that*

$$x = wvr_n(vr_f)^{-1} t_{vr_f(\lambda + \text{wt}(P_n) + \text{wt}(P_f))}$$

Example 11. Let 1 denote the identity element of W_0 . Consider the following superregular saturated chain in type $A_2^{(1)}$.

$$\begin{aligned} x &= s_2 t_{s_1 s_2 s_1} (-2\alpha_1^\vee - 3\alpha_2^\vee) \xrightarrow{\alpha_1 + \alpha_2} \\ z_2 &= s_1 s_2 t_{-3\alpha_1^\vee - 4\alpha_2^\vee} \xrightarrow{\alpha_1} \\ z_1 &= s_1 s_2 s_1 t_{-4\alpha_1^\vee - 4\alpha_2^\vee} \xrightarrow{\alpha_1} \\ y &= s_1 s_2 t_{-4\alpha_1^\vee - 4\alpha_2^\vee} \end{aligned}$$

We observe that $x \leq z_2$ is a far covering that bijects with the quantum edge $s_1 s_2 s_1 \rightarrow 1$ with path weight $\alpha_1^\vee + \alpha_2^\vee$. $z_2 \leq z_1$ is a near covering that bijects with the quantum edge $s_1 s_2 s_1 \rightarrow s_1 s_2$ with path weight $\alpha_1^\vee$. $z_1 \leq y$ is a near covering that bijects with the Bruhat edge $s_1 s_2 \rightarrow s_1 s_2 s_1$ with path weight 0.

In terms of Lemma 10, this implies that

- $w = s_1 s_2$, $v = 1$, and $\lambda = -4\alpha_1^\vee - 4\alpha_2^\vee$
- $r_f = s_1 s_2 s_1$ and $r_n = (s_1)(s_1) = 1$
- $P_f : s_1 s_2 s_1 \rightarrow 1$ with $\text{wt}(P_f) = \alpha_1^\vee + \alpha_2^\vee$
- $P_n : s_1 s_2 \rightarrow s_1 s_2$ with $\text{wt}(P_n) = \alpha_1^\vee$

We find x can be expressed in these terms.

$$\begin{aligned}
x &= wvr_n(vr_f)^{-1}t_{vr_f(\lambda+\text{wt}(P_n)+\text{wt}(P_f))} \\
&= (s_1s_2)(1)(1)[(1)(s_1s_2s_1)]^{-1}t_{(1)(s_1s_2s_1)(-4\alpha_1^\vee-4\alpha_2^\vee+\alpha_1^\vee+\alpha_1^\vee+\alpha_2^\vee)} \\
&= s_1s_2s_1s_2s_1t_{s_1s_2s_1(-2\alpha_1^\vee-3\alpha_2^\vee)} \\
&= s_2t_{s_1s_2s_1(-2\alpha_1^\vee-3\alpha_2^\vee)}
\end{aligned}$$

Proof. We proceed by induction. The base case for $k = 1$ is given explicitly by Corollary 9: the near case yields P_f as trivial while P_n is the edge described, and the far case yields P_n as trivial while P_f is the edge described. Suppose the result is true for any fixed length difference $k \geq 1$ and that $\ell(y) - \ell(x) = k + 1$. By induction, there are elements r'_n, r'_f and associated paths $P'_n : wv \rightarrow wvr'_n, P'_f : vr'_f \rightarrow v$ such that $z_k = wvr'_n(vr'_f)^{-1}t_{vr'_f(\lambda+\text{wt}(P'_n)+\text{wt}(P'_f))}$. Let $\beta = \beta_{k+1}$. We will perform cases based on whether the covering $x \leq z_k$ is a near or far covering. Note that in either case, we observe the following is the resulting classical Weyl component of x :

$$wvr'_n(vr'_f)^{-1}r_{vr'_f(\beta)} = wvr'_n(vr'_f)^{-1}(vr'_f)r_\beta(vr'_f)^{-1} = wvr'_nr_\beta(vr'_f)^{-1}$$

If $x \leq z_k$ by a near covering, then $r'_f = r_f, r'_nr_\beta = r_n$, and $P'_f = P_f$. The covering yields an edge E from $wvr'_n(vr'_f)^{-1}(vr'_f) = wvr'_n \rightarrow wvr'_nr_\beta = wvr_n$ and shows that $P'_n + E = P_n$ and that $\text{wt}(P'_n) + \text{wt}(E) = \text{wt}(P_n)$. Therefore

$$\begin{aligned}
x &= wvr'_nr_\beta(vr'_f)^{-1}t_{vr'_f(\lambda+\text{wt}(P'_n)+\text{wt}(P'_f)+\text{wt}(E))} \\
&= wvr'_nr_\beta(vr_f)^{-1}t_{vr_f(\lambda+\text{wt}(P'_n)+\text{wt}(P_f)+\text{wt}(E))} \\
&= wvr_n(vr_f)^{-1}t_{vr_f(\lambda+\text{wt}(P_n)+\text{wt}(P_f))}
\end{aligned}$$

If $x \leq z_k$ by a far edge, then $r'_n = r_n, r'_fr_\beta = r_f$, and $P'_n = P_n$. The covering yields an edge E from $vr'_fr_\beta = vr_f \rightarrow vr'_f$ and shows that $E + P'_f = P_f$ and $\text{wt}(P'_f) + \text{wt}(E) = \text{wt}(P_f)$. Therefore

$$\begin{aligned}
x &= wvr'_nr_\beta(vr'_f)^{-1}t_{vr'_fr_\beta(\lambda+\text{wt}(P'_n)+\text{wt}(P'_f)+\text{wt}(E))} \\
&= wvr_n r_\beta r'_f{}^{-1} v^{-1} t_{vr'_fr_\beta(\lambda+\text{wt}(P_n)+\text{wt}(P'_f)+\text{wt}(E))} \\
&= wvr_n r_f{}^{-1} v^{-1} t_{vr_f(\lambda+\text{wt}(P_n)+\text{wt}(P_f))} \\
&= wvr_n(vr_f)^{-1}t_{vr_f(\lambda+\text{wt}(P_n)+\text{wt}(P_f))}
\end{aligned}$$

So by induction, the result is true for all superregular saturated chains. $\square$

For a superregular saturated chain (4), we call P_f the associated *far path* and P_n the associated *near path*.

The following Corollary is the specialization of Lemma 10 to W^0 .

Corollary 12. *Suppose $x, y \in W^0$, $x < y = wt_\lambda$ and let (4) be a superregular saturated chain where $\beta_i, r_f, r_n, P_f, P_n$ are defined as in Lemma 10. Then P_f goes from 1 to 1, P_n goes from w to wr_n , and*

$$x = wr_n t_{\lambda+\text{wt}(P_n)+\text{wt}(P_f)}$$

Proof. Since $y \in W^0$, $v = 1$ in Lemma 10. Similarly, since $x \in W^0$, $vr_f = r_f = 1$. Therefore P_f is a closed loop at the identity. The proof is complete. $\square$

2.3 Loops of length 2 in Γ

We note two facts about the Bruhat order.

Theorem 13 ([1, Chain Property]). *If $u < w$ and $u, w \in W^0$, then there exists a saturated chain $u = w_0 < w_1 < w_2 < \dots < w_k = w$ with $w_i \in W^0$.*

Lemma 14 ([1, Lemma 2.7.3]). *If $x < y$ in W with $\ell(y) = \ell(x) + 2$, then $[x, y] = \{x, u, z, y\}$ consists of four distinct elements with $x < u < y$ and $x < z < y$.*

We observe that any loop of length 2 in Γ has a simple root as the label for both of its edges; apply both quantum and Bruhat edge conditions to α .

Lemma 15. *Let $x, y \in W^0$ with y superregular, $\ell(y) = \ell(x) + 2$ and $x < y$. Let $z \in W^0$ be such that $x < z < y$ is the chain guaranteed to exist by Theorem 13. Then $[x, y] \not\subset W^0$ if and only if the near path associated to $x < z < y$ is a 2-loop.*

Proof. Let $y = wt_\lambda$. Suppose $u \notin W^0$. Then by Corollary 12, $x < u < y$ associates to a trivial near path, a far path that is a loop of length 2 at the identity with weight $\alpha^\vee$ for some $\alpha \in \Delta_0$, $u = ws_\alpha t_{s_\alpha(\lambda)}$, and $x = wt_{\lambda+\alpha^\vee}$. Thus by Corollary 12, the associated near path for $x < z < y$ is a loop of length 2 at w . Now suppose the near path associated to $x < z < y$ is a loop $P_n : w \rightarrow ws_\alpha \rightarrow w$ ($\alpha \in \Delta_0$). Then $x = wt_{\lambda+\alpha^\vee}$ by Corollary 12. If $u \in W^0$, this and Lemma 12 would imply a different $\beta^\vee$ such that

$$wt_{\lambda+\beta^\vee} = x = wt_{\lambda+\alpha^\vee} \Rightarrow \alpha = \beta$$

But this contradicts $u \neq z$. $\square$

Example 16. In Example 11, we find that $z_1 < z_2 < y$ corresponds with a near path that is a 2-loop at $s_1 s_2$ and $z_1, z_2, y \in W^0$. In this case, $u = s_1 s_2 s_1 t_{s_1(-3\alpha_1^\vee - 4\alpha_2^\vee)}$ satisfies $u \notin W^0$ and $u \in [z_1, y]$.

Remark 17. It is always the case that if the near path 2-loop from Lemma 15 utilizes the simple root α , then the associated far path which induces u is the far path is the loop $1 \rightarrow r_\alpha \rightarrow 1$.

2.4 Interval equivalent paths

We now discuss and extend properties of paths in Γ uncovered by Postnikov [8]. Following Dyer [4], a reflection ordering is equivalent to a total order on Φ_0^+ such that for all $\alpha, \beta \in \Delta_0$ with $\alpha < \beta$ and $\alpha + \beta \in \Phi_0^+$, then $\alpha < \alpha + \beta < \beta$. Given a reflection ordering on Φ_0^+ , then the following is true.

Lemma 18 (Lemma 6.7 [2]). *If $a \xrightarrow{\alpha_1} x \xrightarrow{\beta_1} c$ is a path in Γ with $\alpha_1 > \beta_1$, then there exists a unique $y \in W_0$ such that $a \xrightarrow{\alpha_2} y \xrightarrow{\beta_2} c$ where $\beta_1 < \beta_2 > \alpha_2 < \alpha_1$.*

In essence, any descent in the label sequence of a path can be uniquely exchanged for an ascent without changing the endpoints of the path. Note that the reversed ordering of a reflection ordering ($\alpha < \beta$ in the reversed ordering if and only if $\beta < \alpha$ in the original reflection ordering) is itself a reflection ordering. By applying Lemma 18 to the reversed ordering, we find that we may also uniquely exchange any descent in the label sequence of a path with an ascent.

Given a path P , we can apply these swaps to sort the label sequence of a path. Pairing this information with [2, Theorem 6.4], we obtain two key facts.

Corollary 19 ([8, proof of Lemma 1], [2]). *Suppose $P : w \rightarrow w'$ is a non-minimal path in Γ .*

1. *P is interval equivalent to a path containing a loop of length 2.*
2. *$\text{wt}(P) - M(w, w')$ is a positive sum of coroots.*

We say that two paths in Γ are *interval equivalent* if the paths can be obtained from each other by a series of exchanges using Lemma 18. In Postnikov's proof of [8, Lemma 1], he notes Lemma 18 also preserves the weight of the path.

Lemma 20. *Let C be a superregular saturated chain from x to y which utilizes only near (resp. far) coverings, P be the associated near (resp. far) path, and P' be interval equivalent to P . Then there is a unique superregular saturated chain C' from x to y which utilizes only near (resp. far) coverings with corresponding near (resp. far) path P' .*

Proof. By Proposition 8, each edge of P bijects to a covering in C . By Lemma 10, the exact element x can be found using the weight and endpoint of P . If P' is interval equivalent, then P' shares the same endpoint, length, and weight of P . Thus, if we biject the edges of P' with near (resp. far) coverings then Lemma 10 states that the lower element from the resulting saturated chain C' must be x as well. $\square$

This previous lemma is sufficient to prove Theorem 3. However, it can be strengthened by the following proposition.

Proposition 21. *Let C be a superregular saturated chain from x to y , P_n be the associated near path, and P_f be the associated far path. If P_n is interval equivalent to P'_n , and P_f is interval equivalent to P'_f , then*

1. *there exists a saturated chain C' from x to y with associated near and far paths P'_n and P'_f .*
2. *any saturated chain C' with upper element y , associated near path P'_n , and associated far path P'_f has lower element x .*

Proof. Note that the proof by induction of Lemma 10 shows that each near covering only modifies r_n and P_n while leaving the far components fixed. Similarly, each far covering only modifies r_f and P_f . Thus, having a near covering then

far covering results in the same lower element as when we take the same far covering first and then the near covering. The only restriction is one cannot change the order of near coverings with respect to other near coverings and similarly for far coverings.

To show (1) and (2), we may take the saturated chain C_1 given by Lemma 20 from z to y generated by P'_n , then append the saturated chain C_2 from x to z given by P'_f . $\square$

Example 22. Consider the sub-chain $x \xrightarrow{\alpha_1+\alpha_2} z_2 \xrightarrow{\alpha_1} z_1$ from Example 11. The left covering was a far covering corresponding to the quantum edge $s_1 s_2 s_1 \rightarrow 1$ while the right covering was a near covering corresponding to the quantum edge $s_1 s_2 s_1 \rightarrow s_1 s_2$. If instead we used the quantum edge $s_1 s_2 s_1 \rightarrow 1$ of a far covering with upper element z_1 , we generate the element $z_3 = t_{s_1 s_2 s_1}(-3\alpha_1^\vee - 3\alpha_2^\vee)$. We also observe that $x \xrightarrow{\alpha_1} z_3$ by a near covering. This demonstrates that changing the order of near and far covers with respect to each other results in a chain still contained in the interval: $z_3 \in [x, z_1]$.

2.5 Proof of Theorem 3

Note that proving Theorem 3 is equivalent to showing that for $x = w't_{\lambda'}$, $y = wt_{\lambda} \in W^0$ and y superregular with respect to x , then $[x, y] \subset W^0$ if and only if $\lambda' = \lambda + M(w, w')$.

Suppose $[x, y] \subset W^0$. Theorem 13 implies there exists a saturated chain in W^0 from x to y , which by Corollary 12 implies $x = w't_{\lambda+\text{wt}(P_n)}$ for the associated near path. If P_n has non-minimal length, then by Corollary 19 part 1, there exists a P'_n which is interval equivalent to P_n but contains a loop of length 2. However, Lemma 15 implies the induced saturated chain in $[x, y]$ is not contained in W^0 , which is a contradiction. Thus P_n has minimal length, i.e. $\text{wt}(P_n) = M(w, w')$.

Suppose $\lambda' = \lambda + M(w, w')$. If $[x, y] \not\subset W^0$, then there exists a saturated chain from x to y which utilizes far coverings. Corollary 12 supplies that $\lambda' = \lambda + \text{wt}(P_n) + \text{wt}(P_f)$ for the associated near and far paths. P_f must be a non-empty loop at the identity and $P_n : w \rightarrow w'$. But this implies $\text{wt}(P_n) - M(w, w') = -\text{wt}(P_f)$ with $\text{wt}(P_f) \neq 0$ since P_f is a non-empty loop, which is a contradiction to Corollary 19 part 2. $\square$

3 Bounds on Regularity

Initially, the definition of sufficiently regular in [6] left a large bound of $|\langle \lambda, \alpha \rangle| > 2|W_0| + 2$ for all $\alpha \in \Delta_0$. Later, in [7, Proposition 4.2], it was found that one only requires $|\langle \lambda, \alpha \rangle| \geq 2\ell(w_0)$ if $W_0 \neq G_2$ (change to $3\ell(w_0)$ in the case of G_2) for all such $\alpha \in \Delta_0$. Recently in [9], it was shown that Proposition 8 holds when W_0 is simply laced and $|\langle \lambda, \alpha \rangle| \geq 3$ for all $\alpha \in \Delta_0$. Furthermore, this hints at a similar, smaller bound for the non-simply laced cases using a similar proof to

[7]. For the following discussion, suppose the regularity bound for Proposition 8 is k .

We can say that y is superregular with respect to x if we are able to use Proposition 8 for every covering between x and y in a saturated chain. If $\ell(y) - \ell(x) = m$, then we must use Proposition 8 $(m - 1)$ -times. We note that each use of Proposition 8 can modify the translation by adding $\alpha^\vee$. Thus we can decrease the regularity by at most the maximum of $\langle \beta, \alpha^\vee \rangle$ over $\beta \in \Delta_0$. If j is the mentioned maximum, then one functioning bound for λ is $k + (m - 1)j$.

Theorem 3 requires the use of Proposition 8 up to the length of the longest shortest path starting at w . Note that $|W_0|$ is an upper bound on the longest shortest path. One functioning lower bound on the regularity of λ in Theorem 3 is $k + |W_0|j$.

4 Dual untwisted affine root systems

Consider the dual of an untwisted affine root system. There is an associated quantum Bruhat graph defined in the same way as for the untwisted affine root systems, except that the roles of roots and coroots are exchanged. For dual untwisted type $s_0 = t_\varphi s_\varphi$ where φ is the short dominant root.

Let $\rho^\vee = (1/2) \sum_{\alpha^\vee \in \Phi_0^+} \alpha^\vee$.

The quantum Bruhat graph (QBG) is the directed graph Γ with vertex set W_0 and a directed edge $w \rightarrow wr_\alpha$ for each pair $(w, \alpha) \in W_0 \times \Phi_0^+$ such that one of the following holds.

1. (Bruhat edge) $\ell(wr_\alpha) = \ell(w) + 1$.
2. (Quantum edge) $\ell(wr_\alpha) = \ell(w) - \langle 2\rho^\vee, \alpha \rangle + 1$.

The *weight* of the edge $w \rightarrow wr_\alpha$ in Γ is α if the edge is quantum and 0 if it is Bruhat. For any $u, v \in W_0$, let $M(u, v)$ be the weight of any shortest path in Γ from u to v .

With these definitions in place, analogues of the statements for the untwisted affine root systems hold for the dual root systems with essentially the same proofs.

Example 23. 1. Consider the root system $A_{2,3-1}^{(2)}$, which is dual to $B_3^{(1)}$.

The classical subroot system has type C_3 realized in $\mathbb{R}^3$ with simple classical roots $\alpha_1 = (1, -1, 0)$, $\alpha_2 = (0, 1, -1)$, $\alpha_3 = (0, 0, 2)$, simple coroots $\alpha_i^\vee = \alpha_i$ for $i \in \{1, 2\}$ and $\alpha_3^\vee = (0, 0, 1)$, and $\varphi = (1, 1, 0)$. We have $2\rho^\vee = (5, 3, 1)$, $\langle 2\rho^\vee, \varphi \rangle = 8$, $s_\varphi = s_2 s_1 s_3 s_2 s_1 s_3 s_2$ and $s_0 = t_\varphi s_\varphi$. Γ has a quantum edge from s_φ to 1 since $\ell(s_\varphi) = 7 = \langle 2\rho^\vee, \varphi \rangle - 1$.

2. Consider the root system $D_{3+1}^{(2)}$, which is dual to $C_3^{(1)}$. The classical subroot system is type B_3 realized in $\mathbb{R}^3$ by $\alpha_1 = (1, -1, 0)$, $\alpha_2 = (0, 1, -1)$, and $\alpha_3 = (0, 0, 1)$, $\alpha_i^\vee = \alpha_i$ for $i = 1, 2$, $\alpha_3^\vee = (0, 0, 2)$, $2\rho^\vee = (6, 4, 2)$, $\varphi = (1, 0, 0)$, $\langle 2\rho^\vee, \varphi \rangle = 6$, and $s_\varphi = s_1 s_2 s_3 s_2 s_1$. Γ has a quantum edge from s_φ to 1 since $\ell(s_\varphi) = 5 = \langle 2\rho^\vee, \varphi \rangle - 1$.




Figure 2: Some dual untwisted affine Dynkin diagrams

References

- [1] Anders Björner and Francesco Brenti. Combinatorics of Coxeter groups. Graduate Texts in Mathematics, 231. Springer, New York, 2005. [13](#), [14](#)
- [2] Francesco Brenti, Sergey Fomin, and Alexander Postnikov. Mixed Bruhat operators and Yang-Baxter equations for Weyl groups. Internat. Math. Res. Notices 1999, no. 8, 419–441. [1.1](#), [1.2](#), [5](#), [18](#), [2.4](#), [19](#)
- [3] Vinay Deodhar. Some characterizations of Bruhat ordering on a Coxeter group and determination of the relative Möbius function. Invent. Math. 39 (1977), no. 2, 187–198. [1.1](#), [1](#)
- [4] M. J. Dyer. Hecke algebras and shellings of Bruhat intervals. Compositio Math. 89 (1993), no. 1, 91–115. [2.4](#)
- [5] Shrawan Kumar. Personal correspondence, 2019. [1.4](#), [6](#)
- [6] Thomas Lam and Mark Shimozono. Quantum cohomology of G/P and homology of affine Grassmannian. Acta Math. 204 (2010), no. 1, 49–90. [8](#), [2.1](#), [3](#), [3](#)
- [7] Elizabeth Milićević. Maximal Newton polygons via the quantum Bruhat graph. 24th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2012), 899–910, Discrete Math. Theor. Comput. Sci. Proc., AR, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2012. [3](#)
- [8] Alexander Postnikov. Quantum Bruhat graph and Schubert polynomials. Proc. Amer. Math. Soc. 133 (2005), no. 3, 699–709. [1.2](#), [2.4](#), [19](#), [2.4](#)
- [9] Amanda Welch. Double Affine Bruhat Order. PhD thesis, Virginia Tech, May 2019. [3](#)