

HYPER-REINHARDT FREE SPECTRAHEDRA

SCOTT MCCULLOUGH

ABSTRACT. The free automorphisms of a class of Reinhardt free spectrahedra are trivial.

1. INTRODUCTION

Fix a positive integer g . For positive integers n , let $M_n(\mathbb{C})$ denote the set of $n \times n$ matrices and $M_n(\mathbb{C})^g$ the set of g -tuples $X = (X_1, \dots, X_g)$ with entries X_j from $M_n(\mathbb{C})$. Let $M(\mathbb{C})^g$ denote the sequence $(M_n(\mathbb{C})^g)_n$. Given a positive integer d and a tuple $B \in M_d(\mathbb{C})^g$, let $\Lambda_B(x)$ denote the linear (matrix) polynomial,

$$\Lambda_B(x) = \sum_{j=1}^g B_j x_j.$$

The polynomial Λ_B naturally *evaluates* at an $X \in M(\mathbb{C})^g$ using the (Kronecker) tensor product as

$$\Lambda_B(X) = \sum_{j=1}^g B_j \otimes X_j.$$

Define L_B by

$$L_B(X) = I_d \otimes I + \Lambda_B(X) + \Lambda_B(X)^*.$$

For a square matrix T the notation $T \succ 0$ (resp. $T \succeq 0$) indicates T is positive definite (resp. positive semidefinite). With this notation, let, for positive integers n ,

$$\mathfrak{P}_B[n] = \{X \in M_n(\mathbb{C})^g : L_B(X) \succ 0\}.$$

The set $\mathfrak{P}_B[1]$ is a *spectrahedron* and the sequence of sets $\mathfrak{P}_B = (\mathfrak{P}_B[n])_{n=1}^\infty$ is a *free spectrahedron*. Alternately,

$$\mathfrak{P}_B = \{X \in M(\mathbb{C})^g : L_B(X) \succ 0\}.$$

Spectrahedra and free spectrahedra occur in a number of areas of mathematics and its applications. For instance, spectrahedra are basic objects in semidefinite programming and convex optimization; free spectrahedra are connected to operator systems and spaces and

2010 *Mathematics Subject Classification.* 47L25, 32H02 (Primary); 52A05, 46L07 (Secondary).

Key words and phrases. bianalytic map, birational map, free spectrahedron, free analysis.

Research Supported by NSF grant DMS-1764231.

complete positivity. Spectrahedra, both free and not, arise in certain engineering applications. For further details see [HKMV20, BPT, dOH06, dOHMP, WSV12, SIG97] and the references therein.

Free spectrahedra are *free sets* in the following sense. If $X \in \mathfrak{P}_B[n]$ and $Y \in \mathfrak{P}_B[m]$, then

$$X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in \mathfrak{P}[n+m],$$

where

$$X_j \oplus Y_j = \begin{pmatrix} X_j & 0 \\ 0 & Y_j \end{pmatrix};$$

and if $X \in \mathfrak{P}_B[n]$ and $U \in M_n(\mathbb{C})$ is unitary, then $U^*XU = (U^*X_1U, \dots, U^*X_gU) \in \mathfrak{P}_B[n]$.

A *free map* $f : \mathfrak{P}_B \rightarrow \mathfrak{P}_C$ between free spectrahedra is a sequence $f = (f[n])$, where $f[n] : \mathfrak{P}_B[n] \rightarrow \mathfrak{P}_C[n]$, that satisfies the axioms

(i) if $X \in \mathfrak{P}_B[n]$ and $Y \in \mathfrak{P}_B[m]$, then

$$f[n+m](X \oplus Y) = f[n](X) \oplus f[m](Y);$$

(ii) if $X \in \mathfrak{P}_B[n]$ and U is an $n \times n$ unitary matrix, then

$$f[n](U^*XU) = U^*f[n](X)U.$$

We typically write f in place of $f[n]$. The free map f is *analytic* if each $f[n]$ is analytic; and f is *free bianalytic* or *free biholomorphic* if f is analytic and has a free analytic inverse. In the case $C = B$, a free bianalytic map is a *free automorphism*. We will often just say f is bianalytic or an automorphism, dropping the adjective free (and also analytic). Motivations for studying free biholomorphic maps between free spectrahedra stem from connections with matrix inequalities that arise in systems engineering [dOHMP, dOH06] and by analogy with rigidity theory in several complex variables. Free biholomorphisms can also be viewed as nonlinear completely positive maps. For a comprehensive treatment of free maps see [KVV14].

We now introduce the central object of this paper. Fix positive integers d_1, \dots, d_{g+1} with $d = \sum d_j$ and norm one matrices C_1, \dots, C_g , where C_j has size $d_j \times d_{j+1}$. Let A_j denote the $(g+1) \times (g+1)$ block matrix with $(j, j+1)$ entry C_j and all other entries 0. Thus each A_j is of size $d \times d$ and the (k, ℓ) block of A_j has size $d_k \times d_\ell$. Such a tuple A is now fixed for the remainder of this paper. The case $g = 2$ was treated in the article [MT+].

Let \mathbb{T} denote the unit circle in the complex plane \mathbb{C} . Given $\gamma \in \mathbb{T}^g$, the mapping

$$\varphi(x) = \gamma \cdot x := (\gamma_1 x_1, \dots, \gamma_g x_g)$$

evidently restricts to an automorphism of \mathfrak{P}_A . We refer to these automorphisms as *trivial automorphisms* and \mathfrak{P}_A as a *hyper-Reinhardt* free spectrahedron.

The *free polydisc* in \mathfrak{g} -variables is the free set

$$\mathcal{F}^{\mathfrak{g}} = \{X \in M(\mathbb{C})^{\mathfrak{g}} : \|X_j\| < 1\}.$$

It is easily seen to be a free spectrahedron (see Section 1.1). Because the C_j have norm one, $\mathfrak{P}_A \subseteq \mathcal{F}^{\mathfrak{g}}$.

Given $\mathfrak{J} \subseteq \{1, 2, \dots, \mathfrak{g}\}$, define $\pi_{\mathfrak{J}} : M(\mathbb{C})^{\mathfrak{g}} \rightarrow M(\mathbb{C})^{\mathfrak{g}}$ by

$$\pi_{\mathfrak{J}}(T)_j = \begin{cases} T_j & \text{if } j \notin \mathfrak{J} \\ 0 & \text{if } j \in \mathfrak{J}. \end{cases}$$

It is immediate that if $T \in \mathfrak{P}_A$, then $\pi_{\mathfrak{J}}(T) \in \mathfrak{P}_A$. As a definition, \mathfrak{P}_A *contains a polydisc as a distinguished summand* if there exists $\emptyset \neq \mathfrak{J} \subseteq \{1, 2, \dots, \mathfrak{g}\}$ such that, for $T \in M(\mathbb{C})^{\mathfrak{g}}$, if $\|T_j\| < 1$ for $j \in \mathfrak{J}$ and $\pi_{\mathfrak{J}}(T) \in \mathfrak{P}_A$, then $T \in \mathfrak{P}_A$. Similarly, \mathfrak{P}_A is a *coordinate direct sum* if there exists a $1 \leq \nu < \mathfrak{g}$ such that $T \in M(\mathbb{C})^{\mathfrak{g}}$ is an element of \mathfrak{P}_A if and only both $\pi_{\mathfrak{J}}(T) \in \mathfrak{P}_A$ and $\pi_{\tilde{\mathfrak{J}}}(T) \in \mathfrak{P}_A$, where $\mathfrak{J} = \{1, 2, \dots, \nu\}$ and $\tilde{\mathfrak{J}}$ is the complement of \mathfrak{J} in $\{1, 2, \dots, \mathfrak{g}\}$. Theorem 1.1 below is the main result of this article.

Theorem 1.1. *Suppose φ is an automorphism of \mathfrak{P}_A .*

- (i) *If \mathfrak{P}_A does not contain a polydisc as a distinguished summand, then $\varphi(0) = 0$;*
- (ii) *If $\varphi(0) = 0$, then there exists a $\gamma \in \mathbb{T}^{\mathfrak{g}}$ and a permutation π of $\{1, 2, \dots, \mathfrak{g}\}$ such that $\varphi_j(x) = \gamma_j x_{\pi(j)}$;*
- (iii) *If $\varphi(0) = 0$ and \mathfrak{P}_A is not a coordinate direct sum, then φ is trivial.*

The proof of Theorem 1.1 occupies the body of this paper.

Corollary 1.2. *If \mathfrak{P}_A is neither a polydisc nor a direct sum of free spectrahedra, then its automorphisms are trivial.*

The case of $\mathfrak{g} = 2$ of Corollary 1.2 is one of the principle results in [MT+], though the proof here diverges materially from that in [MT+].

Proof. If \mathfrak{P}_A is not a direct sum, then it does not contain a polydisc as a proper distinguished summand. By assumption it is not a polydisc. Hence \mathfrak{P}_A does not contain a polydisc as a distinguished summand and therefore, by item (i) its automorphisms send 0 to 0. An application of item (iii) now says the automorphisms of \mathfrak{P}_A are trivial, since, by assumption, \mathfrak{P}_A does not a coordinate direct sum. ■

1.1. Context and prior results. Given a tuple $E = (E_1, \dots, E_{\mathfrak{g}})$ of $k \times \ell$ (not necessarily square) matrices, the spectrahedron \mathfrak{P}_B , where $B \in M_{k+\ell}(\mathbb{C})^{\mathfrak{g}}$ is given by

$$B_j = \begin{pmatrix} 0 & E_j \\ 0 & 0 \end{pmatrix},$$

is a *spectraball*. It is routine to see that a tuple $X \in M(\mathbb{C})^{\mathfrak{g}}$ is in \mathfrak{P}_B if and only if $\|\Lambda_E(X)\| < 1$. We mention two special cases. One is the free polydisc $\mathcal{F}^{\mathfrak{g}}$

$$\mathcal{F}^{\mathfrak{g}} = \{X \in M(\mathbb{C})^{\mathfrak{g}} : \|X_j\| < 1\}$$

in \mathfrak{g} variables and the other is the free $\mathfrak{g} \times \mathfrak{h}$ matrix ball,

$$\mathcal{M}^{\mathfrak{g} \times \mathfrak{h}} = \{X = (X_{j,k})_{j,k=1}^{\mathfrak{g},\mathfrak{h}} : X_{j,k} \in M(\mathbb{C}), \|X\| < 1\}.$$

In each case, the automorphisms at the scalar level - those of $\mathcal{F}^{\mathfrak{g}}[1]$ and $\mathcal{M}^{\mathfrak{g} \times \mathfrak{h}}[1]$ - naturally extend to free automorphisms. See for instance [Pop10, MT16]. In particular, automorphisms of $\mathcal{F}^{\mathfrak{g}}$ are composites of permutation of the coordinates with mobius maps of the unit disc in each coordinate.

More generally, the results of [AHKM, HKMV20] characterize bianalytic maps between free spectrahedra \mathfrak{P}_B and \mathfrak{P}_C under certain generic irreducibility hypotheses on the tuples B and C . They also characterize bianalytic maps between spectraballs absent any additional hypotheses. A canonical class of free spectrahedra not covered by these results are those with circular symmetry.

A free spectrahedron \mathfrak{P}_B is *circular* if $X \in \mathfrak{P}_B$ and $\gamma \in \mathbb{T}$ implies $\gamma X = (\gamma X_1, \dots, \gamma X_{\mathfrak{g}}) \in \mathfrak{P}_B$. From [HKM12], circular free spectrahedra that are bianalytic are linearly equivalent. Thus, in this case, there is an automorphism that sends 0 to 0. As a conjecture, an automorphism of a circular free spectrahedron that does not contain a spectraball as a direct summand, must send 0 to 0.

2. CARATHEODORY INTERPOLATION PRELIMINARIES

In this section consequence of, and results related to, Caratheodory interpolation needed in the sequel are collected for easy reference. We begin with the following well known lemma.

Lemma 2.1. *Given $c_0, c_1 \in \mathbb{C}$, let*

$$X = \begin{pmatrix} c_0 & c_1 \\ 0 & c_0 \end{pmatrix}.$$

- (i) $\|X\| = 1$ if and only if $|c_0| \leq 1$ and $|c_1| = 1 - |c_0|^2$; (and hence there is a $\theta \in \mathbb{R}$ such that $c_1 = e^{i\theta}(|c_0|^2 - 1)$);
- (ii) $\|X\| < 1$ if and only if $|c_0| < 1$ and $|c_1| < 1 - |c_0|^2$;
- (iii) $\|X\| > 1$ if and only if $|c_0| > 1$ or $|c_0| \leq 1$ and $|c_1| > 1 - |c_0|^2$.

Further, in the first case, the kernel of $I - XX^*$ is spanned by the vector

$$v = \begin{pmatrix} 1 \\ -e^{-i\theta}c_0 \end{pmatrix}.$$

Proof. The matrix $I - XX^*$ has norm at most 1 (resp. equal 1) if and only if $|c_0| < 1$ (resp. $|c_0| \leq 1$) and its determinant, $(1 - |c_0|^2)^2 - |c_1|^2$, is non-negative (resp. equal 0), proving the first two items. A similar argument proves the third item. Direct computation verifies that v is in the kernel of $I - XX^*$ in the case that X has norm 1. ■

Lemma 2.2. *Suppose*

- (i) $n \geq 2$;
- (ii) $c_0, \dots, c_n \in \mathbb{C}$ and $|c_0| < 1$;
- (iii) $\theta \in \mathbb{R}$ and $c_1 = e^{i\theta}(|c_0|^2 - 1)$.

Let $\lambda_1 = 1$, fix $\lambda_2, \dots, \lambda_n \in \mathbb{C} \setminus \{0\}$ and let $S = (S_{j,k})_{j,k=1}^{n+1}$ denote the $(n+1) \times (n+1)$ matrix with $S_{j,j+1} = \lambda_j$ for $1 \leq j \leq n$ and $S_{j,k} = 0$ otherwise. Thus the nonzero entries of S are exactly on the first super diagonal. The matrix

$$T = \sum_{j=0}^n c_j S^j$$

is a contraction if and only if $c_j = e^{i\theta}(e^{i\theta}c_0^*)^{j-1}(|c_0|^2 - 1)$ and $|\lambda_j| \leq 1$ for $j \geq 1$ if and only if it has norm 1.

Moreover, in this case, letting P denote the $(n+1) \times (n+1)$ matrix with $P_{1,n+1} = 1$ and $P_{j,k} = 0$ otherwise, if $\mu \in \mathbb{C}$ and $T + \mu P$ is a contraction, then $\mu = 0$.

Let \mathbb{D} denote the open unit disc in \mathbb{C} .

Proof. First note, for each $1 \leq j \leq n$ that

$$X_j := \begin{pmatrix} T_{j,j} & T_{j,j+1} \\ T_{j+1,j} & T_{j+1,j+1} \end{pmatrix} = \begin{pmatrix} c_0 & \lambda_j c_1 \\ 0 & c_0 \end{pmatrix}.$$

Observe that X_1 is the matrix X from Lemma 2.1 and in particular has norm 1. Hence T has norm at least one and is thus a contraction if and only if it has norm one. Further, if T is a contraction, then each X_j is a contraction and hence, by Lemma 2.1, $|\lambda_j| \leq 1$.

Suppose now the norm of T is one and write T as

$$T = \begin{pmatrix} X & Y \\ 0 & Z \end{pmatrix},$$

where $X = X_1$. Using Lemma 2.1 and its notation, since T is a contraction and $\|X^*v\| = \|v\|$,

$$\|v\|^2 \geq \|T^* \begin{pmatrix} v \\ 0 \end{pmatrix}\|^2 = \left\| \begin{pmatrix} X^*v \\ Y^*v \end{pmatrix} \right\|^2 = \|v\|^2 + \|Y^*v\|^2.$$

Thus the columns of Y are orthogonal to v . On the other hand, y_k , the k -th column of Y is given by

$$y_k = \lambda_2 \cdots \lambda_{k+1} \begin{pmatrix} c_{k+1} \\ c_k \end{pmatrix},$$

for $1 \leq k \leq n-1$. Using the assumption that the λ_j are non-zero,

$$\begin{pmatrix} c_{k+1} \\ c_k \end{pmatrix} = c_k \begin{pmatrix} e^{i\theta} c_0^* \\ 1 \end{pmatrix}.$$

Thus, $c_{k+1} = e^{i\theta} c_0^* c_k$. In particular, $c_2 = e^{2i\theta} c_0^* (|c_0|^2 - 1)$ and by induction, $c_k = e^{ik\theta} (c_0^*)^{k-1} (|c_0|^2 - 1)$.

To prove the converse, suppose $|\lambda_j| \leq 1$ and $c_j = e^{ij\theta} c_0^{*(j-1)} (|c_0|^2 - 1)$. In particular, S is a contraction and

$$f(z) = \frac{c_0 - e^{i\theta} z}{1 - c_0^* e^{i\theta} z} = \sum_{j=0}^{\infty} c_j z^j$$

is an automorphism of \mathbb{D} . Hence $T = f(S)$ is a contraction.

Finally suppose $T = f(S)$ and $T + \mu P$ is a contraction. The argument above with, $T + \mu P$ in place of T , shows $y_{n-1} + \mu w$ is orthogonal to the vector v from Lemma 2.1, where $w \in \mathbb{C}^2$ is the vector with first entry 1 and second entry 0. Since y_{n-1} is orthogonal to v , it follows that μw is orthogonal to v . Hence $\mu = 0$ and the proof is complete. \blacksquare

3. THE AFFINE LINEAR TERMS

Suppose $\varphi : \mathfrak{P}_A \rightarrow \mathfrak{P}_A$ is bianalytic,

$$\varphi = \begin{pmatrix} \varphi_1 & \cdots & \varphi_g \end{pmatrix}.$$

Let $b_j = \varphi_j(0)$.

The coordinate functions φ_j of φ have power series expansions,

$$\varphi_j = b_j + \sum_{k=1}^g \ell_{j,k} x_k + h_j(x),$$

that converge in some neighborhood of 0 and for all nilpotent tuples, where $h_j(x)$ consists of terms of order two and higher. For notational purposes, let

$$(3.1) \quad \mathcal{L} = \left(\ell_{j,k} \right)_{j,k=1}^g.$$

Since φ is bianalytic, \mathcal{L} is invertible.

For integers $N \geq 2$, let \mathscr{W}_N denote the words in $\{x_1, \dots, x_{\mathbf{g}}\}$ of length between 2 and N . Let \mathscr{W}_N^j denote the words in \mathscr{W}_N that are **not** of the form x_j^n (for $2 \leq n \leq N$). Let \mathscr{W}^j denote the union (over $N \geq 2$) of the \mathscr{W}_N^j and let \mathfrak{H}^j denote power series of the form

$$(3.2) \quad \mathfrak{h}(x) = \sum_{\alpha \in \mathscr{W}^j} a_\alpha x^\alpha.$$

Let $\{\delta_1, \dots, \delta_{\mathbf{g}}\}$ denote the standard orthonormal basis for $\mathbb{C}^{\mathbf{g}}$. Given $p \in \mathbb{D}$, let

$$\mathfrak{m}_p(z) = \frac{p + z}{1 + p^* z}.$$

Proposition 3.1 below is the main results of this section.

Proposition 3.1. *There exists a permutation π of $\{1, 2, \dots, \mathbf{g}\}$ and a tuple $\theta = (\theta_1, \dots, \theta_{\mathbf{g}}) \in \mathbb{R}$ such that, for each $1 \leq j \leq \mathbf{g}$,*

$$\varphi_j(\delta_{\pi(j)} z) = \mathfrak{m}_{b_j}(e^{i\theta_j} z) = \frac{b_j + e^{i\theta_j} z}{1 + b_j^* e^{i\theta_j} z} = e^{i\theta_j} \mathfrak{m}_{e^{-i\theta_j} b_j}(z).$$

More generally, with $k = \pi(j)$, there exists an $\mathfrak{h}_k \in \mathfrak{H}^k$ such that

$$\varphi_j(x) = \mathfrak{m}_{b_j}(e^{i\theta_j} x_k) + \mathfrak{h}_k(x).$$

In particular, there exists $a_\alpha^k \in \mathbb{C}$ such that if T is nilpotent of order $N + 1$, then

$$(3.3) \quad \varphi_j(T) = \mathfrak{m}_{b_j}(e^{i\theta_j} T_k) + \mathfrak{h}_k(T) = \mathfrak{m}_{b_j}(e^{i\theta_j} T_k) + \sum_{\alpha \in \mathscr{W}_N^k} a_\alpha^k T^\alpha.$$

We refer to π as the *permutation associated to φ* .

Before proving Proposition 3.1, we pause to prove item (ii) of Theorem 1.1, stated as Proposition 3.2 below.

Proposition 3.2. *If $\varphi(0) = 0$, then there exists $\gamma \in \mathbb{T}^{\mathbf{g}}$ and permutation π such that $\varphi_j(x) = \gamma_j x_{\pi(j)}$.*

Proof. In this case φ is an automorphism of a circular domain that maps 0 to 0 and is thus linear by [HKM11, Theorem 4.4]. On the other hand, by Proposition 3.1, $\varphi_j(x) = e^{i\theta_j} x_k + \mathfrak{h}_k(x)$. Linearity forces $\mathfrak{h}_k = 0$ completing the proof. ■

The proof of Proposition 3.1 is broken down into several lemmas.

Lemma 3.3. *There is a permutation π of $\{1, \dots, \mathbf{g}\}$ and a tuple $\theta \in \mathbb{R}^{\mathbf{g}}$ such that, for tuples $T = (T_1, \dots, T_{\mathbf{g}})$ nilpotent of order 2,*

$$\varphi_j(T) = b_j + e^{i\theta_j} (|b_j|^2 - 1) T_{\pi(j)}$$

Before proving Lemma 3.3, we first establish Lemma 3.4 below. Recall equation (3.1).

Lemma 3.4. *There is a permutation π of $\{1, \dots, \mathbf{g}\}$ such that $\ell_{j,k} \neq 0$ if and only if $k = \pi(j)$. In particular, for tuples $T = (T_1, \dots, T_{\mathbf{g}})$ nilpotent of order 2,*

$$\varphi_j(T) = b_j + \ell_{j,\pi(j)} T_{\pi(j)}.$$

Thus $\ell_{j,k} \neq 0$ if and only if $k = \pi(j)$ and \mathcal{L} has exactly one nonzero entry in each row and column.

The proof of Lemma 3.4 in turn uses the following elementary linear algebra lemma included here for completeness.

Lemma 3.5. *Suppose R and Q are self-adjoint $n \times n$ matrices and $\gamma \in \mathbb{C}^n$. If $R \pm Q \succeq 0$ and $R\gamma = 0$, then $Q\gamma = 0$.*

Proof. It is immediate that $\langle Q\gamma, \gamma \rangle = 0$. For $\lambda \in \mathbb{C}$ and $\delta \in \mathbb{C}^n$, the inequality

$$0 \leq \langle (R + Q)(\gamma + \lambda\delta), \gamma + \lambda\delta \rangle = |\lambda|^2 \langle (R + Q)\delta, \delta \rangle + \lambda \langle Q\gamma, \delta \rangle + \lambda^* \langle Q\delta, \gamma \rangle$$

implies $\langle Q\gamma, \delta \rangle = 0$ and hence $Q\gamma = 0$. ■

Define $\mathbf{T} = (T_1, \dots, T_{\mathbf{g}}) = \mathbf{T}(t) = \mathbf{T}(t_1, \dots, t_{\mathbf{g}})$ by $T_j = t_j S$, where

$$(3.4) \quad S = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Recall the definition of \mathcal{L} from equation (3.1) and let $B = I + \Lambda(b) + \Lambda(b)^*$. Since

$$L_A(\varphi(\mathbf{T}(t))) = B + \Lambda_A(\mathbf{T}(\mathcal{L}t)) + \Lambda_A(\mathbf{T}(\mathcal{L}t))^*,$$

if $s, t \in \mathbb{C}^{\mathbf{g}}$ and $0 < c < 1$, then

$$L_A(\varphi(\mathbf{T}(t + s))) = L_A(\varphi(\mathbf{T}(t))) + \Lambda_A(\mathbf{T}(\mathcal{L}s)) + \Lambda_A(\mathbf{T}(\mathcal{L}s))^*.$$

Lemma 3.6. *If $t \in \mathbb{C}^{\mathbf{g}}$ and $|t_j| \leq 1$ for all j , then $\mathbf{T}(t)$ is in the closure of \mathfrak{P}_A . If in addition, there is an $1 \leq m \leq \mathbf{g}$ such that $|t_m| = 1$, then $\mathbf{T}(t)$ is in the boundary of \mathfrak{P}_A .*

Given $1 \leq m \leq \mathbf{g}$, if $s \in \mathbb{C}^{\mathbf{g}}$ satisfies $s_m = 0$ and $|s_j| \leq 1$ for all j , then $\varphi(\mathbf{T}(\delta_m))$ and $\varphi(\mathbf{T}(\delta_m + s))$ are in the boundary of \mathfrak{P}_A and thus each of $\mathbf{T}(\delta_m)$; and $\mathbf{T}(\delta_m \pm s)$ are in the boundary of \mathfrak{P}_A . Equivalently,

(a) $R := L_A(\varphi(\mathbf{T}(\delta_m)))$; and

(b) $L_A(\varphi(\mathbf{T}(\delta_m \pm s)))$

are positive semidefinite with non-trivial kernel. In particular, there is a vector $0 \neq \gamma$ such that $R\gamma = 0$.

Moreover,

$$L_A(\varphi(\mathbf{T}(\delta_m \pm s))) = R \pm Q(s),$$

where

$$Q(s) = \Lambda_A(\mathbb{T}(\mathcal{L}s)) + \Lambda_A(\mathbb{T}(\mathcal{L}s))^*.$$

Finally, $Q(s)\gamma = 0$.

Proof. The first statements about $\mathbb{T}(t)$ are evident.

By the first part of the lemma, $\mathbb{T}(\delta_m)$ and $\mathbb{T}(\delta_m \pm s)$ are in the boundary of \mathfrak{P}_A and hence so are there images under φ .

The moreover portion of the lemma is straightforward, since $\varphi(\mathbb{T}(t))$ is affine linear in t ,

$$\varphi(\mathbb{T}(t)) = b + \mathbb{T}(\mathcal{L}t).$$

The last statement follows from the moreover part of this lemma and Lemma 3.5. ■

Proof of Lemma 3.4. Recall, $\{\delta_1, \dots, \delta_{\mathfrak{g}}\}$ is the standard orthonormal basis for $\mathbb{C}^{\mathfrak{g}}$. Let H_j denote the orthogonal complement of δ_j and suppose, for each $1 \leq k \leq \mathfrak{g}$, there is a $1 \leq \pi(k) \leq \mathfrak{g}$ such that $\mathcal{L}H_k \subseteq H_{\pi(k)}$. Since \mathcal{L} is invertible, it follows that $\mathcal{L}H_k = H_{\pi(k)}$ and $\pi(k) \neq \pi(j)$ for $j \neq k$. In particular, π is a permutation. Moreover, for each $1 \leq j \leq \mathfrak{g}$, using \mathcal{L} is invertible gives

$$\mathcal{L}[\delta_j] = \mathcal{L} \cap_{k \neq j} H_k = \cap_{k \neq j} \mathcal{L}H_k = \cap_{k \neq j} H_{\pi(k)} = [\delta_{\pi(j)}],$$

where $[x]$ denotes the span of a vector $x \in \mathbb{C}^{\mathfrak{g}}$. Thus, for j fixed, $\ell_{j,k} \neq 0$ if and only if $k = \pi(j)$ and the conclusion of the lemma holds.

Now suppose there is an $1 \leq m \leq \mathfrak{g}$ such that $\mathcal{L}H_m \not\subseteq H_k$ for each k . It follows that $\dim[\mathcal{L}H_m] \cap H_k < \dim H_k = \mathfrak{g} - 1$ and consequently $\cup_{k=1}^{\mathfrak{g}} [\mathcal{L}H_m] \cap H_k \subsetneq \mathcal{L}H_m$. Thus there exists an $s \in H_m \subseteq \mathbb{C}^{\mathfrak{g}}$ such that each entry of $\mathcal{L}s$ is non-zero. By scaling s , we assume the entries of s have absolute value at most one. Applying Lemma 3.6 (and using its notation), fix a (non-zero) vector γ in the kernel of R so that $Q(s)\gamma = 0$.

Expressing $\gamma = \oplus_{j=1}^{\mathfrak{g}+1} \gamma_j$ and

$$\gamma_j = \begin{pmatrix} \gamma_{j,1} \\ \gamma_{j,2} \end{pmatrix} \in \begin{matrix} \mathbb{C}^{d_j} \\ \mathbb{C}^{d_j} \end{matrix},$$

and setting $\sigma = \mathcal{L}s$, the condition $Q(s)\gamma = 0$ implies $0 = \sigma_j C_j \gamma_{j+1,2}$ and $0 = \sigma_j^* C_j^* \gamma_{j,1}$ for $1 \leq j \leq \mathfrak{g}$. Thus,

$$(3.5) \quad \begin{aligned} C_j \gamma_{j+1,2} &= 0 \\ C_j^* \gamma_{j,1} &= 0, \end{aligned}$$

since $\sigma_j \neq 0$. In particular, $C_{\mathfrak{g}}^* \gamma_{\mathfrak{g},1} = 0$.

Since $R\gamma = 0$,

$$b_{\mathbf{g}}^* C_{\mathbf{g}}^* \gamma_{\mathbf{g},1} + \gamma_{\mathbf{g}+1,1} = 0$$

and it follows that $\gamma_{\mathbf{g}+1,1} = 0$.

Arguing by induction, suppose $\gamma_{\mathbf{g}+1,1}, \dots, \gamma_{k+1,1}$ are all 0 for some $2 \leq k \leq \mathbf{g}$. Since $R\gamma = 0$ implies

$$0 = b_{k-1}^* C_{k-1}^* \gamma_{k-1,1} + \gamma_{k,1} + b_k C_k \gamma_{k+1,1} + \ell_{k,m} C_k \gamma_{k+1,2},$$

it follows from equation (3.5) and the induction hypothesis that $\gamma_{k,1} = 0$. Hence $\gamma_{k,1} = 0$ for all $2 \leq k \leq \mathbf{g} + 1$. Finally, from $\gamma_{1,1} + b_1 C_1 \gamma_{2,1} = 0$ it follows that $\gamma_{1,1} = 0$. A similar argument starting from

$$0 = \gamma_{1,2} + b_1 C_1 \gamma_{2,2}$$

shows $\gamma_{k,2} = 0$ for each $1 \leq k \leq \mathbf{g} + 1$ too. We have now reached the contradiction $\gamma = 0$, completing the proof. \blacksquare

Proof of Lemma 3.3. Let π denote the permutation of Lemma 3.4. Thus, for each $1 \leq j \leq \mathbf{g}$ and each tuple T nilpotent of order two,

$$\varphi_j(T) = b_j + \ell_{j,\pi(j)} T_{\pi(j)}.$$

Fix $1 \leq m \leq \mathbf{g}$ and let $v = \pi^{-1}(m)$. In particular, $\pi(v) = m$. Let $s \in \mathbb{C}^{\mathbf{g}}$ denote the tuple with $s_a = 1$ for $a \neq m$ and $s_m = 0$. Using Lemma 3.6 and its notation, there is a non-zero vector γ such that $R\gamma = 0$ and $Q(s)\gamma = 0$. In this case,

$$Q(s) = \sum_{j \neq v} \ell_{j,\pi(j)} A_j \otimes S + \left[\sum_{j \neq v} \ell_{j,\pi(j)} A_j \otimes S \right]^*$$

where S is given in equation (3.4).

Expressing $\gamma = \bigoplus_{j=1}^{\mathbf{g}+1} \gamma_j$ and

$$\gamma_j = \begin{pmatrix} \gamma_{j,1} \\ \gamma_{j,2} \end{pmatrix} \in \begin{matrix} \mathbb{C}^{m_j} \\ \oplus \\ \mathbb{C}^{m_j} \end{matrix},$$

the condition $Q(s)\gamma = 0$ (and $\ell_{j,\pi(j)} \neq 0$) implies, for $j \neq v$,

$$(3.6) \quad \begin{aligned} 0 &= C_j \gamma_{j+1,2} \\ 0 &= C_j^* \gamma_{j,1}. \end{aligned}$$

As in the proof of Lemma 3.4, $R\gamma = 0$ implies $\gamma_{\mathbf{g}+1,1} + b_{\mathbf{g}}^* C_{\mathbf{g}}^* \gamma_{\mathbf{g},1} = 0$ and also $\gamma_{1,2} + b_1 C_1 \gamma_{2,2} = 0$. Assuming $v < \mathbf{g}$, it follows that $\gamma_{\mathbf{g}+1,1} = 0$. Another application of $R\gamma = 0$ gives

$$b_{\mathbf{g}-1}^* C_{\mathbf{g}-1}^* \gamma_{\mathbf{g}-1,1} + \gamma_{\mathbf{g},1} + b_{\mathbf{g}} C_{\mathbf{g}} \gamma_{\mathbf{g}+1,1} + \ell_{\mathbf{g},\pi(\mathbf{g})} C_{\mathbf{g}} \gamma_{\mathbf{g}+1,2} = 0.$$

Thus, if $v < \mathfrak{g} - 1$, then in view of equation (3.6) and using the already established $\gamma_{\mathfrak{g}+1,1} = 0$, it follows that $\gamma_{\mathfrak{g},1} = 0$. By induction, $\gamma_{j,1} = 0$ for $v+2 \leq j$. A similar argument shows $\gamma_{j,2} = 0$ for $j \leq v - 1$. On the other hand, $R\gamma = 0$ gives

$$(3.7) \quad \begin{aligned} 0 &= b_{v-1}^* C_{v-1}^* \gamma_{v-1,1} + \gamma_{v,1} + b_v C_v \gamma_{v+1,1} + \ell_{v,m} C_{v+1} \gamma_{v+1,2} \\ 0 &= \ell_{v-1,m} C_{v-1}^* \gamma_{v-1,1} + b_{v-1}^* C_{v-1}^* \gamma_{v-1,2} + \gamma_{v,2} + b_v C_v \gamma_{v+1,2} \\ 0 &= b_v^* C_v^* \gamma_{v,1} + \gamma_{v+1,1} + b_{v+1} C_{v+1} \gamma_{v+2,1} + \ell_{v+1,m} C_{v+1} \gamma_{v+2,2} \\ 0 &= \ell_{v,m}^* C_v^* \gamma_{v,1} + b_v^* C_v^* \gamma_{v,2} + \gamma_{v+1,2} + b_{v+1} C_{v+1} \gamma_{v+2,2}. \end{aligned}$$

Combining equations (3.6), (3.7) and $\gamma_{v-1,2} = 0 = \gamma_{v+2,1}$, it follows that

$$\begin{aligned} 0 &= \gamma_{v,1} + b_v C_v \gamma_{v+1,1} + \ell_{v,m} C_v \gamma_{v+1,2} \\ 0 &= \gamma_{v,2} + b_v C_v \gamma_{v+1,2} \\ 0 &= b_v^* C_v^* \gamma_{v,1} + \gamma_{v+1,1} \\ 0 &= \ell_{v,m}^* C_v^* \gamma_{v,1} + b_v^* C_v^* \gamma_{v,2} + \gamma_{v+1,2}. \end{aligned}$$

Thus,

$$0 = \begin{pmatrix} I & \varphi_v(T(\delta_m)) \otimes C_v \\ \varphi_v(T(\delta_m))^* \otimes C_v^* & I \end{pmatrix} \begin{pmatrix} \gamma_{v,1} \\ \gamma_{v,2} \\ \gamma_{v+1,1} \\ \gamma_{v+1,2} \end{pmatrix}$$

and hence

$$\varphi_v(T(\delta_m)) = \begin{pmatrix} b_v & \ell_{v,m} \\ 0 & b_v \end{pmatrix}$$

has norm one or $\gamma_{a,b} = 0$ for $a = v, v+1$ and $b = 1, 2$. In the second case, $L_A(cI_2)\gamma = 0$, where $c \in \mathbb{C}^{\mathfrak{s}}$ is the tuple with $c_j = b_j$ for $j \neq v$ and $c_v = 0$. Thus c is in the boundary of $\mathfrak{P}_A[1]$. On the other hand, since b is not in this boundary of $\mathfrak{P}_A[1]$, it is easy to see that neither is c , a contradiction. In the first case, from Lemma 2.1, $\ell_{v,m} = e^{i\theta_v}(|b_v|^2 - 1)$, for some $\theta_v \in \mathbb{R}$ and the proof is complete. \blacksquare

Proof of Proposition 3.1. By Lemma 3.3, there is a permutation π of $\{1, \dots, \mathfrak{g}\}$ such that, for tuples $T = (T_1, \dots, T_{\mathfrak{g}})$ nilpotent of order 2 and $1 \leq j \leq \mathfrak{g}$,

$$\varphi_j(T) = b_j + e^{i\theta_j}(|b_j|^2 - 1)T_{\pi(j)},$$

for some $\theta_j \in \mathbb{R}$. Hence, with $k = \pi(j)$, there exists a_{α}^k for words α of length at least two, such that for a tuple T nilpotent of order $N + 1 \geq 2$,

$$\varphi_j(T) = b_j + e^{i\theta_j}(|b_j|^2 - 1)T_{\pi(j)} + \sum_{\alpha \in \mathcal{W}_N} a_{\alpha}^k T^{\alpha}.$$

For $N \in \mathbb{N}^+$, let R denote the $(N+1) \times (N+1)$ matrix with $R_{j,j+1} = 1$ for $1 \leq j \leq N$ and $R_{j,k} = 0$ otherwise. Let $k = \pi(j)$. The tuple $Y = \delta_k R$ is nilpotent of order $N+1$ and is in the boundary of \mathfrak{P}_A and hence so is $\varphi(Y)$. In particular,

$$\varphi_j(Y) = b_j + e^{i\theta_j}(|b_j|^2 - 1)R + \sum_{n=2}^N a_{x_k^n}^k R^n.$$

is a contraction. By Lemma 2.2, $a_{x_k^n}^k = e^{in\theta_j}(b_j^*)^{n-1}(|b_j|^2 - 1)$. Hence equation (3.3) holds for T nilpotent of order $N+1$ and the remainder of the proposition follows. \blacksquare

4. HIGHER ORDER TERMS

We continue to work with a fixed automorphism φ of \mathfrak{P}_A , with $b = \varphi(0)$. In particular, there exists a permutation π and a tuple $\theta \in \mathbb{R}^{\mathfrak{g}}$ such that the conclusion of Proposition 3.1 holds. Let

$$(4.1) \quad \mathfrak{g}_j(z) = e^{i\theta_j} \mathfrak{m}_{e^{-i\theta_j} b_j}(z) = \mathfrak{m}_{b_j}(e^{i\theta_j} z) = b_j + e^{i\theta_j}(1 - |b_j|^2)z + \dots$$

Let $\mathfrak{Z}^+ \subseteq \{1, 2, \dots, \mathfrak{g}\}$ denote those indices k such that, for each $\epsilon > 0$, the matrix $L_A(\delta_k + \epsilon\delta_{k+1})$ is not positive semidefinite; let \mathfrak{Z}^- denote those indices k such that, for each $\epsilon > 0$, the matrix $L_A(\delta_k + \epsilon\delta_{k-1})$ is not positive semidefinite; and let $\mathfrak{Z} = \mathfrak{Z}^+ \cup \mathfrak{Z}^-$. Let \mathfrak{N} denote the complement of \mathfrak{Z} . It is straight forward consequence of convexity that \mathfrak{N} is the set of those indices j for which there exists an $\epsilon > 0$ such that $L_A(\epsilon\delta_{k-1} + \delta_k + \epsilon\delta_{k+1}) \succeq 0$, (in the case that $k = 1$ or $k = \mathfrak{g}$ we interpret this condition in the natural way).

The following alternate criteria for membership in the sets \mathfrak{Z}^\pm will often be used without comment.

Lemma 4.1. *An index j is in \mathfrak{Z}^+ if and only if, for each $\epsilon > 0$,*

$$C_j^* C_j + \epsilon C_{j+1} C_{j+1}^* \not\succeq I.$$

Likewise $j \in \mathfrak{Z}^-$ if and only if for each $\epsilon > 0$, $\epsilon C_{j-1}^ C_{j-1} + C_j C_j^* \not\succeq I$.*

Proof. Observe $L_A(\delta_j + \epsilon\delta_{j+1}) \succeq 0$ if and only if

$$Z = \begin{pmatrix} I & C_j & 0 \\ C_j^* & I & \epsilon C_{j+1} \\ 0 & \epsilon C_{j+1}^* & I \end{pmatrix} \succeq 0$$

if and only if (after a couple of Schur complements of $Z \succeq 0$) $C_j^* C_j + \epsilon^2 C_{j+1} C_{j+1}^* \preceq I$. \blacksquare

Recall the definition of \mathfrak{H}^k from equation (3.2). Let \mathfrak{H}_\pm^k denote the set of those $\mathfrak{h} \in \mathfrak{H}^k$ such that $\mathfrak{h}(T) = 0$ for tuples $T \in \mathfrak{P}_A$ with $T_{k-1} = T_{k+1} = 0$.

Lemma 4.2. Fix $1 \leq j \leq \mathfrak{g}$ and let $k = \pi(j)$.

(i) If $k \in \mathfrak{N}$, then

$$\varphi_j(x) = \mathfrak{g}_j(x_k).$$

(ii) In any case, there exists $\mathfrak{h}_k \in \mathfrak{H}_\pm^k$ such that

$$\varphi_j(T) = \mathfrak{g}_j(T_k) + \mathfrak{h}_k(T).$$

Lemma 4.2 is a preliminary version of Proposition 4.10, the main result of this section. Before proving Lemma 4.2, we collect a couple of lemmas related to the hyper-Reinhardt nature of \mathfrak{P}_A .

4.1. **The hyper-Reinhardt condition.** Given $T \in M(\mathbb{C})^\mathfrak{g}$, note

$$L_A(T) = \begin{pmatrix} I & C_1 \otimes T_1 & 0 & 0 & \dots & 0 & 0 \\ C_1^* \otimes T_1^* & I & C_2 \otimes T_2 & 0 & \dots & 0 & 0 \\ 0 & C_2^* \otimes T_2^* & I & C_3 \otimes T_3 & \dots & 0 & 0 \\ 0 & 0 & C_3^* \otimes T_3^* & I & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \\ 0 & 0 & 0 & 0 & \dots & I & C_g \otimes T_g \\ 0 & 0 & 0 & 0 & \dots & C_g^* \otimes T_g^* & I \end{pmatrix}.$$

Given tuples $T \in M_n(\mathbb{C})^\mathfrak{g}$ and $W = (W_0, \dots, W_g) \in M_n(\mathbb{C})^{\mathfrak{g}+1}$, let

$$(4.2) \quad W \circ T := (W_0^* T_1 W_1, W_1^* T_2 W_2, \dots, W_{g-1}^* T_g W_g).$$

Lemma 4.3. If $T \in \mathfrak{P}_A[n]$ and $W \in M_n(\mathbb{C})^{\mathfrak{g}+1}$ is a tuple of unitary matrices, then $W \circ T$ is also in \mathfrak{P}_A .

Remark 4.4. A natural question is whether a free spectrahedra $\mathfrak{P}_B \subseteq M(\mathbb{C})^\mathfrak{g}$ is hyper-Reinhardt if and only if for each n , each $X \in \mathfrak{P}_B[n]$ and tuple $W = (W_0, \dots, W_g) \in M_n(\mathbb{C})^{\mathfrak{g}+1}$ of unitary matrices

$$W \circ X = (W_0^* X_1 W_1, \dots, W_{g-1}^* X_g W_g) \in \mathfrak{P}_A.$$

Proof of Lemma 4.3. Letting D_W denote the block diagonal $(\mathfrak{g}+1) \times (\mathfrak{g}+1)$ with $(j+1, j+1)$ entry $I \otimes W_j$ for $0 \leq j \leq \mathfrak{g}$,

$$D_W^* L_A(T) D_W = L_A(W \circ T). \quad \blacksquare$$

Lemma 4.5. If $\lambda \in \mathfrak{P}_A[1]$, and $X \in M_n(\mathbb{C})^\mathfrak{g}$ is a tuple of contractions, then $\lambda \cdot X = (\lambda_1 X_1, \dots, \lambda_g X_g) \in \mathfrak{P}_A[n]$.

Proof. Given a tuple $U \in M_n(\mathbb{C})^{\mathfrak{g}}$ of unitary matrices, let $W_0 = I$ and $W_j = U_1 U_2 \dots U_j$ and let $\Lambda = \lambda I_n$. Observe that $\Lambda \circ W = \lambda \cdot U$. Thus

$$L_A(\Lambda \circ W) = L_A(\lambda \cdot U)$$

and consequently, by Lemma 4.3, $\lambda \cdot U \in \mathfrak{P}_A$.

By the Russo-Dye Theorem, there is an n and unitary tuples $U^j \in M_n(\mathbb{C})^{\mathfrak{g}}$ for $1 \leq j \leq n$ such that

$$X = \frac{1}{n} \sum_{\ell=1}^n U^\ell, \quad X_j = \frac{1}{n} \sum_{\ell=1}^n U_j^\ell.$$

Thus

$$L_A(\lambda \cdot X) = \frac{1}{n} \sum_{j=1}^n L_A(\lambda \cdot U^j) \succ 0$$

and it follows that $\lambda \cdot X \in \mathfrak{P}_A$. ■

We will also need the following variation of Lemma 4.5.

Lemma 4.6. *Fix $\mathcal{I} \subseteq \{1, 2, \dots, \mathfrak{g}\}$ and $\lambda_j \geq 0$ for $j \in \mathcal{I}$. If $Y \in \mathfrak{P}_A$ and $Y_j = 0$ for $j \in \mathcal{I}$ implies Y^λ defined by*

$$Y_j^\lambda = \begin{cases} \lambda_j & \text{if } j \in \mathcal{I} \\ Y_j & \text{if } j \notin \mathcal{I}, \end{cases}$$

is in \mathfrak{P}_A , then, if $Y \in \mathfrak{P}_A$ and $Y_j = 0$ for $j \in \mathcal{I}$ and $\|T_j\| \leq 1$ for $j \in \mathcal{I}$, then the tuple

$$Z = \begin{cases} \lambda_j T_j & \text{if } j \in \mathcal{I} \\ Y_j & \text{if } j \notin \mathcal{I} \end{cases}$$

is in \mathfrak{P}_A .

Proof. Given a tuple $W = (W_0, W_1, \dots, W_{\mathfrak{g}})$ of $n \times n$ unitary matrices, the tuple $W^* \circ Y \in \mathfrak{P}_A$ and $(W^* \circ Y)_j = 0$ for $j \in \mathcal{I}$. Hence the tuple $(W^* \circ Y)^\lambda$ is in \mathfrak{P}_A and therefore so is

$$W \circ (W^* \circ Y)^\lambda = \begin{cases} \lambda_j W_j^* W_{j+1} & \text{if } j \in \mathcal{I} \\ Y_j & \text{if } j \notin \mathcal{I}. \end{cases}$$

An application of the Russo-Dye lemma as in the proof of Lemma 4.3 completes the proof. ■

Lemma 4.7. *Fix $1 \leq k \leq \mathfrak{g}$. If there is an $\epsilon > 0$ such that both $L_A(\epsilon \delta_{k-1} + \delta_k) \succeq 0$ and $L_A(\delta_k + \epsilon \delta_{k+1}) \succeq 0$, then there is an $\eta > 0$ such that if $T \in M_n(\mathbb{C})^{\mathfrak{g}}$ satisfies $\|T_k\| \leq 1$ and $\|T_j\| \leq \eta$ for $j \neq k$, then T is in the closure of $\mathfrak{P}_A[n]$. If, in addition, $\|T_k\| = 1$, then T is in the boundary of $\mathfrak{P}_A[n]$.*

Similarly, if just $L_A(\delta_k + \epsilon \delta_{k+1}) \succeq 0$, then there is an $\eta > 0$ such that if $\|T_k\| \leq 1$, $T_{k-1} = 0$ and $\|T_j\| < \eta$ for $j \neq k$, then T is in the closure of $\mathfrak{P}_A[n]$.

Proof. Observe that there is a $\sigma > 0$ such that $L_A(\delta_k + \sigma \sum_{j \neq k, k+1, k-1} \delta_j) \succeq 0$. Equivalently $\delta_k + \sigma \sum_{j \neq k, k+1, k-1} \delta_j$ is in the closure of \mathfrak{P}_A . By convexity, the average of $\delta_k + \sigma \sum_{j \neq k, k+1, k-1} \delta_j$ and $\delta_k + \epsilon \delta_{k+1}$ and $\delta_k + \epsilon \delta_{k-1}$ is also in \mathfrak{P}_A . Thus, there is an $\eta > 0$ such that $L_A(\delta_k + \eta \sum_{j \neq k} \delta_j) \succeq 0$. From here an application of Lemma 4.5, with $X_k = T_k$ and $X_j = \frac{T_j}{\eta}$ for $j \neq k$, establishes the first part of the lemma.

To prove the second part of the lemma, note that there is an $\eta > 0$ such that $L_A(\delta_k + \eta \sum_{k-1 \neq j \neq k} \delta_j) \succeq 0$ and argue as above. \blacksquare

4.2. Proof of Lemma 4.2. Fix j and recall the set \mathscr{W}^j defined just below equation (3.1). For notational convenience, let $k = \pi(j)$. From Proposition 3.1, there exists $\mathfrak{h}_k \in \mathfrak{H}^j$ such that

$$\varphi_j(x) = \mathfrak{g}_j(x_k) + \mathfrak{h}_k(x),$$

where \mathfrak{g}_k is given in equation (4.1).

Now suppose $k \in \mathfrak{N}$. We argue by induction on N that the result of item (i) of Lemma 4.2 holds when evaluating φ_j on a tuple T that is nilpotent of order N . That is, $\mathfrak{h}_k(T) = 0$ if T is nilpotent of order N . Equivalently, when writing

$$\mathfrak{h}_k(x) = \sum_{\alpha \in \mathscr{W}^j} c_\alpha x^\alpha,$$

we have $c_\alpha = 0$ whenever $|\alpha| < N$. The case of $N = 2$ follows from Lemma 3.3. Now suppose the result is true for an $N \geq 2$. Let T_k denote the $(N+1) \times (N+1)$ matrix with $\lambda_{k,1} = 1$ in the $(1, 2)$ entry; $\lambda_{k,u} \in \mathbb{C} \setminus \{0\}$ in the $(u, u+1)$ entry for $u \geq 2$ and 0 elsewhere. Similarly, for $\ell \neq k$, let T_ℓ denote the $(N+1) \times (N+1)$ matrix with $\lambda_{\ell,u} \in \mathbb{C}$ in the $(u, u+1)$ entry for $1 \leq u \leq N$ and 0 elsewhere. The tuple $T = T(\lambda)$ thus depends (linearly) on the $\lambda_{\ell,u} \in \mathbb{C}$ for $1 \leq \ell \leq \mathfrak{g}$ and $1 \leq u \leq N$ and $(\ell, u) \neq (k, 1)$. Given such a λ and a word $\alpha = x_{j_1} x_{j_2} \cdots x_{j_N}$, let

$$\lambda^\alpha = \lambda_{j_1,1} \lambda_{j_2,2} \cdots \lambda_{j_N,N}$$

and note that any $\lambda_{j,u}$ appears at most once. Hence, if $c_\alpha \in \mathbb{C}$ and

$$\sum_{|\alpha|=N} c_\alpha \lambda^\alpha = 0,$$

for λ an open set of such λ (with $\lambda_{k,1} = 1$), then $c_\alpha = 0$ for all α .

In the case of item (i), there is an open neighborhood U of 0 in $\mathbb{C}^{N\mathfrak{g}-1}$ such that for $\lambda \in U$, the tuple $T(\lambda)$ is in the boundary of \mathfrak{P}_A by Lemma 4.7. Let $M(\lambda) = \varphi_j(T(\lambda))$. By the induction hypothesis, and using notation from equation (3.3),

$$M(\lambda) = \mathfrak{g}_j(T_k(\lambda)) + \sum_{|\alpha|=N, \alpha \neq x_k^N} a_\alpha^k T^\alpha(\lambda).$$

Since, for $|\alpha| = N$, the matrix T^α is 0 except for its upper right entry. When $\lambda_{k,j} \neq 0$, Lemma 2.2 implies

$$\sum_{|\alpha|=N, \alpha \neq x_k^N} a_\alpha^k T^\alpha = 0.$$

On the other hand, letting P denote the matrix with 1 in the upper right entry and 0 elsewhere,

$$\sum_{|\alpha|=N, \alpha \neq x_k^N} a_\alpha^k T^\alpha = \sum a_\alpha^k \lambda^\alpha P.$$

Hence $a_\alpha^k = 0$ for each $\alpha \in \mathcal{W}_N^k$. Thus $\varphi_j(T) = \mathfrak{g}_j(T)$, completing the induction argument in the setting of item (i).

We now drop the $k \in \mathfrak{N}$ assumption and proceeds as above, but with $\lambda_{k-1,u} = 0 = \lambda_{k+1,u}$ for each u . The conclusion in this setting becomes: if α is a word the does not include either x_{k-1} and x_{k+1} (and $\alpha \neq x_k^N$), then $c_\alpha = 0$. Hence if $T \in \mathfrak{P}_A$ and $T_{k-1} = 0 = T_{k+1}$, then $\mathfrak{h}_k(T) = 0$.

4.3. The sets \mathfrak{N} and \mathfrak{J} . This subsection develops criteria for membership in the sets \mathfrak{N} and \mathfrak{J} .

Lemma 4.8. *Fix $1 \leq j < \mathfrak{g}$ and let $k = \pi(j)$ and $\ell = \pi(j+1)$. If $L_A(\delta_j + \delta_{j+1})$ is not positive semidefinite, then either $\ell = k-1$ or $\ell = k+1$.*

Proof. Suppose $L_A(\delta_j + \delta_{j+1})$ is not positive definite. Arguing by contradiction, suppose $\ell \neq k+1$ and $\ell \neq k-1$. By Lemma 4.2 item (ii), $\varphi_j(x\delta_k + w\delta_\ell) = \mathfrak{g}_j(x)$ and $\varphi_{j+1}(x\delta_k + w\delta_\ell) = \mathfrak{g}_{j+1}(w)$, in the latter case since $\ell = \pi(j+1)$ and $k \neq \ell+1$ and $k \neq \ell-1$. On the other hand $t(\delta_k + \delta_\ell)$ is in \mathfrak{P}_A for $0 < t < 1$ and hence so is $\varphi(t(\delta_k + \delta_\ell))$. In particular,

$$(4.3) \quad \begin{pmatrix} I & \mathfrak{g}_{b_j}(t) \otimes C_j & 0 \\ \mathfrak{g}_{b_j}(t)^* \otimes C_j^* & I & \mathfrak{g}_{b_{j+1}}(t) \otimes C_{j+1} \\ 0 & \mathfrak{g}_{b_{j+1}}(t)^* \otimes C_{j+1}^* & I \end{pmatrix} \succeq 0.$$

Because \mathfrak{g}_{b_j} and $\mathfrak{g}_{b_{j+1}}$ extend continuously across the boundary of the unit disc, it follows that the inequality of equation (4.3) holds for $t = 1$. Since $\mathfrak{g}_{b_j}(1)$ and $\mathfrak{g}_{b_{j+1}}(1)$ both have modulus 1, it follows that $L_A(\delta_j + \delta_{j+1}) \succeq 0$, a contradiction. Thus either $\ell = k+1$ or $\ell = k-1$. \blacksquare

Lemma 4.9. *Fix $1 \leq j \leq \mathfrak{g}$ and let $k = \pi(j)$.*

(a) *If, $j \in \mathfrak{J}^+$, then*

- (i) $\pi(j+1) = k+1$
- (ii) $b_j = 0 = b_{j+1}$; and
- (iii) $k \in \mathfrak{J}^+$.

(b) If $j \in \mathfrak{Z}^-$, then

- (i) $\pi(j-1) = k-1$;
- (ii) $b_j = 0 = b_{j-1}$; and
- (iii) $k \in \mathfrak{Z}^-$.

Proof. Suppose $j \in \mathfrak{Z}^+$. Thus, for each $\epsilon > 0$, the matrix $L_A(\delta_j + \epsilon\delta_{j+1})$ is not positive semidefinite. In particular, if $\epsilon > 0$, then

$$C_j^*C_j + \epsilon^2C_{j+1}C_{j+1}^* \not\leq I,$$

by Lemma 4.1.

Choose

$$T_k = S = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

and $T_a = 0$ otherwise. Thus T is in the boundary of \mathfrak{P}_A , the matrix $\varphi_j(T)$ has norm one and there is a scalar e_{j+1} such that

$$\varphi_{j+1}(T) = b_{j+1}I_2 + e_{j+1}T_{\pi(j+1)} = b_{j+1}I.$$

On the other hand, $L_A(\varphi(T)) \succeq 0$ implies

$$\varphi_j(T)^*\varphi_j(T) \otimes C_j^*C_j + |b_{j+1}|^2C_{j+1}C_{j+1}^* \preceq I,$$

and thus $b_{j+1} = 0$ by Lemma 4.1, proving half of item (aii).

From Lemma 4.8, $\ell = k-1$ or $\ell = k+1$. Arguing by contradiction, suppose $\ell = k-1$. In this case, let $T_a = 0$ for $k-1 \neq a \neq k$ and

$$T_k = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad T_{k-1} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}.$$

Thus T is in the boundary of \mathfrak{P}_A and hence so is $\varphi(T)$. The only word in T of length two or longer with non-zero coefficient is T_kT_{k-1} . Hence, letting $e_j = e^{i\theta_j}(|b_j|^2 - 1)$ and $e_{j+1} = e^{i\theta_{j+1}}$,

$$\begin{aligned} \varphi_j(T) &= \mathfrak{g}_j(T_k) + \alpha T_k T_{k-1} = b_j + e_j T_k + \alpha T_k T_{k-1} = \begin{pmatrix} b_j & e_j & \alpha \\ 0 & b_j & 0 \\ 0 & 0 & b_j \end{pmatrix} \\ \varphi_{j+1}(T) &= \mathfrak{g}_{j+1}(T_{k-1}) + \beta T_k T_{k-1} = e_{j+1} T_{k-1} + \beta T_k T_{k-1} = \begin{pmatrix} 0 & 0 & \beta \\ 0 & 0 & e_{j+1} \\ 0 & 0 & 0 \end{pmatrix}, \end{aligned}$$

for some $\alpha, \beta \in \mathbb{C}$. From Lemma 2.2, $\alpha = 0 = \beta$. Hence there exists a unit vector $\gamma \in \mathbb{C}^3$ such that $\|\varphi_j(T)\gamma\| = 1$ and $\gamma_2 \neq 0$ but $\gamma_3 = 0$. Since $|e_{j+1}| = 1$ and $\gamma_2 \neq 0$, it follows that $\varphi_{j+1}(T)^*\gamma \neq 0$. Since $L_A(\varphi(T)) \succeq 0$,

$$\begin{pmatrix} I & \varphi_j(T) \otimes C_j & 0 \\ \varphi_j(T)^* \otimes C_j^* & I & \varphi_{j+1}(T) \otimes C_{j+1} \\ 0 & \varphi_{j+1}(T)^* \otimes C_{j+1}^* & I \end{pmatrix} \succeq 0$$

and hence

$$I \succeq \|\varphi_j(T)\gamma\|^2 C_j^* C_j + \|\varphi_{j+1}(T)^*\gamma\|^2 C_{j+1} C_{j+1}^* = C_j^* C_j + \|\varphi_{j+1}(T)^*\gamma\|^2 C_{j+1} C_{j+1}^*.$$

Lemma 4.1 now gives the contradiction $\varphi_{j+1}(T)^*\gamma = 0$. Thus $\ell = k + 1$, proving item (ai).

Next choose $T_k = S = T_{k+1}$ and $T_a = 0$ otherwise, where S is given in equation (3.4). Hence T is in the boundary of \mathfrak{P}_A and thus so is $\varphi(T)$. Letting $X = \varphi_j(T)$ and $Y = \varphi_{j+1}(T)$, it follows that

$$X^* X \otimes C_j^* C_j + Y Y^* \otimes C_{j+1} C_{j+1}^* \preceq I.$$

Letting γ denote a unit vector with $X^* X \gamma = \gamma$, we find $Y^* \gamma = 0$. Since $b_{j+1} = 0$,

$$Y = \begin{pmatrix} 0 & e_{j+1} \\ 0 & 0 \end{pmatrix},$$

where $|e_{j+1}| = 1$. It follows that the first component of γ is zero and thus $b_j = 0$, and the proof of item (aii) is complete.

To prove item (aiii), suppose, by way of contradiction, there is an $\epsilon > 0$ such that $L_A(\delta_k + \epsilon \delta_{k+1}) \succeq 0$. The tuple T defined by $T_a = 0$ for $k \neq a \neq k + 1$ and

$$T_k = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad T_{k+1} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & \epsilon \\ 0 & 0 & 0 \end{pmatrix}$$

is in the boundary of \mathfrak{P}_A . Thus $\varphi(T)$ is in the boundary of \mathfrak{P}_A . Since $b_j = 0$,

$$X = \varphi_j(T) = \begin{pmatrix} 0 & e^{i\theta_j} & y \\ 0 & 0 & e^{i\theta_j} \\ 0 & 0 & 0 \end{pmatrix},$$

for some y . Since $\varphi_j(T)$ is a contraction, $y = 0$. Similarly, since $b_{j+1} = 0$,

$$Y = \varphi_{j+1}(T) = \begin{pmatrix} 0 & e^{i\theta_{j+1}} & 0 \\ 0 & 0 & \epsilon e^{i\theta_{j+1}} \\ 0 & 0 & 0 \end{pmatrix}.$$

The condition

$$X^* X \otimes C_j^* C_j + Y Y^* \otimes C_{j+1} C_{j+1}^* \preceq I$$

now implies that

$$C_j^* C_j + \epsilon^2 C_j C_j^* \preceq I.$$

Equivalently $L_A(\delta_j + \epsilon\delta_{j+1}) \succeq 0$, a contradiction that completes the proof of item (a).

Item (b) is proved similarly. ■

4.4. Summary. The following proposition records the key findings from this section.

Proposition 4.10. *The sets \mathfrak{N} and \mathfrak{Z} partition $\{1, 2, \dots, \mathfrak{g}\}$.*

If φ is an automorphism of \mathfrak{P}_A , then there is a permutation π of $\{1, 2, \dots, \mathfrak{g}\}$ and a tuple $\theta \in \mathbb{R}^{\mathfrak{g}}$, such that

- (i) *the sets \mathfrak{N} , \mathfrak{Z} , \mathfrak{Z}^+ , \mathfrak{Z}^- are invariant under π ;*
- (ii) *if $j \in \mathfrak{Z}$ and $k = \pi(j)$, then*

$$(4.4) \quad \varphi_j(x) = e^{i\theta_j} x_{\pi(j)} + \mathfrak{h}_k(x),$$

for some $\mathfrak{h}_k \in \mathfrak{H}^k$;

- (iii) *if $j \in \mathfrak{N}$ and $k = \pi(j)$, then there is an automorphism \mathfrak{g}_j of \mathbb{D} such that*

$$(4.5) \quad \varphi_j(x) = \mathfrak{m}_{b_j}(e^{i\theta_j} x_k) = \mathfrak{g}_j(x_k) = \frac{b_j + e^{i\theta_j} x_k}{1 + b_j^* e^{i\theta_j} x_k},$$

where $b_j = \varphi_j(0)$.

Proof. By definition, \mathfrak{N} and \mathfrak{Z} partition.

Part of the conclusion of Lemma 4.9 is that \mathfrak{Z}^+ and \mathfrak{Z}^- are invariant under π , from which it follows that \mathfrak{Z} and \mathfrak{N} are also invariant.

If $j \in \mathfrak{Z}$ then, by Lemma 4.9, $\varphi_j(0) = 0$ and thus, from Lemma 4.2 item (ii), there is a $\theta_j \in \mathbb{R}$ and $\mathfrak{h}_k \in \mathfrak{H}^k$ such that equation (4.4) holds.

Finally, if $j \in \mathfrak{N}$, then equation (4.5) holds by Lemma 4.2 item (i). ■

5. NORMALIZED AUTOMORPHISMS AND COMPOSITIONS

Recall that $\mathfrak{Z}, \mathfrak{N} \subseteq \{1, 2, \dots, \mathfrak{g}\}$ depend only on \mathfrak{P}_A and are thus independent of any particular automorphism of \mathfrak{P}_A .

Suppose ψ is an automorphism of \mathfrak{P}_A with associated permutation κ . In particular the conclusions of Proposition 4.10 hold. We say that ψ is a *normalized automorphism* if

- (i) $c_j = \psi_j(0) \geq 0$; for each j ;
- (ii) for $j \in \mathfrak{Z}$ and $k = \kappa(j)$,

$$\psi_j(x) = \gamma_j x_k + \mathfrak{h}_k(x),$$

where $\mathfrak{h}_k \in \mathfrak{H}^k$ and $\gamma_j \in \mathbb{T}$;

(iii) for $j \in \mathfrak{N}$ and $k = \kappa(j)$,

$$\psi_j(x) = \mathbf{m}_{c_j}(x_k) := \frac{c_j + x_k}{1 + c_j x_k}.$$

Lemma 5.1. *If φ is an automorphism of \mathfrak{P}_A with associated permutation π , then there is a normalized automorphism ψ of \mathfrak{P}_A with associated permutation π such that $\psi_j(0) = |\varphi_j(0)|$ for each j .*

Proof. Let $b_j = \varphi_j(0)$. From Proposition 4.10, there is a permutation π and a tuple $\theta \in \mathbb{R}^{\mathfrak{g}}$ such that, for $j \in \mathfrak{J}$,

$$\varphi_j(x) = e^{i\theta_j} x_k + \mathfrak{h}_k(x);$$

and, for $j \in \mathfrak{N}$,

$$\varphi_j(x) = \mathbf{m}_{b_j}(e^{i\theta_j} x_{\pi(j)}).$$

Let $\gamma_j = 1$ if $b_j = 0$ and $\gamma_j = \frac{b_j^*}{|b_j|}$ otherwise. Let $\rho : \mathfrak{P}_A \rightarrow \mathfrak{P}_A$ denote the trivial automorphism

$$\rho(x) = \gamma \cdot x.$$

Let $\alpha_k = e^{-i\theta_{\pi^{-1}(k)}} \gamma_{\pi^{-1}(k)}^*$ and $\tau : \mathfrak{P}_A \rightarrow \mathfrak{P}_A$ denote the trivial automorphism defined by

$$\tau(x) = \alpha \cdot x.$$

Finally, let $\psi : \mathfrak{P}_A \rightarrow \mathfrak{P}_A$ denote the automorphism $\psi = \rho \circ \varphi \circ \tau$ and note the conclusion of the lemma is satisfied with $\kappa = \pi$. Indeed, with $k = \pi(j)$, if $j \in \mathfrak{N}$, then,

$$\begin{aligned} \psi_j(x) &= \gamma_j \varphi_j(\alpha \cdot x) = \gamma_j \mathbf{m}_{b_j}(e^{i\theta_j} (\alpha \cdot x)_k) \\ &= \gamma_j \mathbf{m}_{b_j}(\gamma_j^* x_k) = \gamma_j \frac{b_j + \gamma_j^* x_k}{1 + |b_j| x_k} \\ &= \frac{|b_j| + x_k}{1 + |b_j| x_k} = \mathbf{m}_{|b_j|}(x_k). \quad \blacksquare \end{aligned}$$

Hence item (iii) holds and moreover $\psi_j(0) = |b_j| \geq 0$.

If instead $j \in \mathfrak{J}$, then, since ψ is an automorphism, then item (ii) holds by item (ii) of Proposition 4.10. Further $\psi_j(0) = 0$ and hence item (i) holds.

Lemma 5.2. *Suppose φ and $\tilde{\varphi}$ are automorphisms of \mathfrak{P}_A with associated permutations π and $\tilde{\pi}$. Let ψ denote the automorphism $\tilde{\varphi} \circ \varphi$ and let τ be its associated permutation. If $j \in \mathfrak{N}$, then $\tau(j) = \pi(\tilde{\pi}(j))$ and moreover, with $\tilde{\varphi}_j(x) = \tilde{g}_j(x_{\tilde{\pi}(j)})$ and $\varphi_{\tilde{\pi}(j)}(x) = \mathfrak{g}_{\tilde{\pi}(j)}(x_{\pi(\tilde{\pi}(j))})$, we have $\psi_j(x) = \tilde{g}_j \circ \mathfrak{g}_{\tilde{\pi}(j)}(x_{\tau(j)})$.*

Proof. Let $\tilde{b} = \tilde{\varphi}(0)$ and $b = \varphi(0)$. Given $j \in \mathfrak{N}$, it follows that $\pi(j) \in \mathfrak{N}$ and $\pi(\tilde{\pi}(j)) \in \mathfrak{N}$ by Proposition 4.10. From item (i) of Lemma 4.2, there exists functions g_j and \tilde{g}_j as well as $\mathfrak{g}_{\tilde{\pi}(j)}$ of one variable such that $\tilde{\varphi}_j(x) = \tilde{g}_j(x_{\tilde{\pi}(j)})$ and $\psi_j(x) = g_j(x_{\tau(j)})$ as well as $\varphi_{\tilde{\pi}(j)}(x) = \mathfrak{g}_{\tilde{\pi}(j)}(x_{\pi(\tilde{\pi}(j))})$. Hence

$$g_j(x_{\tau(j)}) = \psi_j(x) = \tilde{\varphi}_j(\varphi(x)) = \tilde{g}_j(\varphi_{\tilde{\pi}(j)}(x)) = \tilde{g}_j(\mathfrak{g}_{\tilde{\pi}(j)}(x_{\pi(\tilde{\pi}(j))})) = \tilde{g}_j \circ \mathfrak{g}_{\tilde{\pi}(j)}(x_{\pi(\tilde{\pi}(j))}).$$

It follows that $\tau(j) = \pi(\tilde{\pi}(j))$ and $g_j(y) = \tilde{g}_j \circ \mathfrak{g}_{\tilde{\pi}(j)}(y)$. ■

Given an automorphism φ of \mathfrak{P}_A , let

$$\mathcal{F}_\varphi = \{j : \varphi_j(0) \neq 0\}$$

and note that $\mathcal{F}_\varphi \subseteq \mathfrak{N}$ by Proposition 4.10.

Lemma 5.3. *If φ and $\tilde{\varphi}$ are normalized automorphisms of \mathfrak{P}_A with associated permutations π and $\tilde{\pi}$ respectively, then $\psi = \tilde{\varphi} \circ \varphi$ is a normalized automorphism of \mathfrak{P}_A and moreover, $\mathcal{F}_\psi = \mathcal{F}_{\tilde{\varphi}} \cup \tilde{\pi}^{-1}(\mathcal{F}_\varphi)$.*

Proof. Let τ denote the permutation associated with ψ . By Lemma 5.2 $\tau = \pi \circ \tilde{\pi}$ on \mathfrak{N} . Let $\tilde{b} = \tilde{\varphi}(0)$ and $b = \varphi(0)$. By assumption $\tilde{b}_j, b_j \geq 0$.

Suppose $j \in \mathfrak{N}$. From Proposition 4.10, $\tau(j), \tilde{\pi}(j), \pi(j) \in \mathfrak{N}$ and

$$\psi_j(x) = \tilde{\varphi}_j(\varphi(x)) = \mathfrak{m}_{\tilde{b}_j}(\varphi_{\tilde{\pi}(j)}(x)) = \mathfrak{m}_{\tilde{b}_j}(\mathfrak{m}_{b_{\tilde{\pi}(j)}}(x_{\tau(j)})) = \mathfrak{m}_{c_j}(x_{\tau(j)}),$$

where $c_j = (\tilde{b}_j + b_{\tilde{\pi}(j)})(1 + \tilde{b}_j b_{\tilde{\pi}(j)})^{-1} \geq 0$. In particular, $\psi_j(0) \geq 0$. Moreover, $\psi_j(0) > 0$ if and only either $\tilde{b}_j > 0$ or $b_{\tilde{\pi}(j)} > 0$.

Hence, if $j \in \mathfrak{N}$, then $j \in \mathcal{F}_\psi$ if and only if either $j \in \mathcal{F}_{\tilde{\varphi}}$ or $\tilde{\pi}(j) \in \mathcal{F}_\varphi$. On the other hand, if $j \notin \mathfrak{N}$, then $\tilde{\pi}(j) \notin \mathfrak{N}$ and therefore $j \notin \mathcal{F}_\psi \cup \mathcal{F}_{\tilde{\varphi}}$ and $\tilde{\pi}(j) \notin \mathcal{F}_\varphi$. Hence, $\mathcal{F}_\psi = \mathcal{F}_{\tilde{\varphi}} \cup \tilde{\pi}^{-1}(\mathcal{F}_\varphi)$. ■

Given a self map f of a set X and a positive integer n , let $f^{(n)}$ denote the composition of f with itself n -times.

Lemma 5.4. *If φ is a normalized automorphism of \mathfrak{P}_A , then there is a positive integer n such that $\psi = \varphi^{(n)}$ is a normalized automorphism of \mathfrak{P}_A such that, for each positive integer m ,*

- (a) *the permutation associated to $\psi^{(m)}$ is the identity on \mathfrak{N} ; and*
- (b) $\mathcal{F}_\varphi \subseteq \mathcal{F}_\psi = \mathcal{F}_{\psi^{(m)}}$.

Proof. By Lemma 5.3, for each positive integer p we have $\varphi^{(p)}$ is a normalized automorphism of \mathfrak{P}_A and

$$\mathcal{F}_\varphi \subseteq \mathcal{F}_{\varphi^{(2)}} \subseteq \mathcal{F}_{\varphi^{(3)}} \subseteq \cdots$$

Hence, there is N such that $\mathcal{F}_{\varphi^{(p)}} = \mathcal{F}_{\varphi^{(N)}}$ for all $p \geq N$.

By Proposition 4.10, \mathfrak{N} is invariant under π . Let $\rho = \pi|_{\mathfrak{N}} : \mathfrak{N} \rightarrow \mathfrak{N}$. There is an $\ell \geq 1$ such that $\rho^{(\ell)}$ is the identity. Let $n = N\ell$. Thus $n \geq N$ and $\rho^{(n)}$ is the identity. The conclusion of this lemma now follows from Lemma 5.2. \blacksquare

6. THE CASE OF THE IDENTITY PERMUTATION

In this section, we fix an automorphism ψ of \mathfrak{P}_A with associated permutation π satisfying the conclusion of Proposition 5.4 and assume $\mathcal{F}_\psi \neq \emptyset$. (In particular, $\mathfrak{N} \neq \emptyset$ by Proposition 4.10.) Thus ψ is a normalized automorphism, π is the identity on \mathfrak{N} and $\mathcal{F}_{\psi^{(m)}} = \mathcal{F}_\psi \neq \emptyset$ for all positive integers m . For notational convenience, let $\mathcal{F} = \mathcal{F}_\psi$. Since $\mathcal{F} \subseteq \mathcal{N}$, the permutation π is the identity on \mathcal{F} . Moreover, the complement, $\tilde{\mathcal{F}} = \{1, 2, \dots, \mathfrak{g}\} \setminus \mathcal{F}$, of \mathcal{F} is invariant under π . Let $b = \psi(0)$. Thus $b_j = \psi_j(0) \geq 0$ for all j and $b_j > 0$ if and only if $j \in \mathcal{F}$.

6.1. **Two auxiliary Reinhardt spectrahedra.** Let

$$M_n(\mathbb{C})^{\mathcal{F}} = \{X = (X_j)_{j \in \mathcal{F}} : X_j \in M_n(\mathbb{C})\}.$$

Similarly, let

$$M_n(\mathbb{C})^{\tilde{\mathcal{F}}} = \{Y = (Y_j)_{j \in \tilde{\mathcal{F}}} : Y_j \in M_n(\mathbb{C})\}.$$

Given $X \in M_n(\mathbb{C})^{\mathcal{F}}$ and $Y \in M_n(\mathbb{C})^{\tilde{\mathcal{F}}}$, let $T = (X, Y) \in M_n(\mathbb{C})^{\mathfrak{g}}$ denote the tuple with $T_j = X_j$ for $j \in \mathcal{F}$ and $T_j = Y_j$ for $j \in \tilde{\mathcal{F}}$.

Let $\hat{b} = (b_j I_n)_{j \in \mathcal{F}} \in M_n(\mathbb{C})^{\mathcal{F}}$. (Thus the n is understood from context.) Similarly, let $\hat{0} = (0 I_n)_{j \in \tilde{\mathcal{F}}}$. With these notations, let

$$\begin{aligned} \mathfrak{P}_A^{\mathcal{F}} &= \{Y \in M(\mathbb{C})^{\tilde{\mathcal{F}}} : (\hat{0}, Y) \in \mathfrak{P}_A\} \\ \mathfrak{E}_A^{\mathcal{F}} &= \{Y \in M(\mathbb{C})^{\tilde{\mathcal{F}}} : (\hat{b}, Y) \in \mathfrak{P}_A\}. \end{aligned}$$

Since \mathfrak{P}_A is Reinhardt, it immediately follows that $\mathfrak{E}_A^{\mathcal{F}}$ is a subset of $\mathfrak{P}_A^{\mathcal{F}}$.

Lemma 6.1. *Both $\mathfrak{P}_A^{\mathcal{F}}$ and $\mathfrak{E}_A^{\mathcal{F}}$ are Reinhardt free spectrahedra in $\mathfrak{g} - |\mathcal{F}|$ variables.*

Proof. That both domains are Reinhardt is immediate. It is also evident that $(\hat{0}, Y) \in \mathfrak{P}_A$ if and only if

$$I + \sum_{j \in \tilde{\mathcal{F}}} A_j \otimes Y_j + \sum_{j \in \tilde{\mathcal{F}}} A_j^* \otimes Y_j^* \succ 0,$$

and hence $\mathfrak{P}_A^{\mathcal{F}}$ is a free spectrahedron.

Let

$$P = I + \sum_{j \in \mathcal{F}} A_j b_j + \sum_{j \in \mathcal{F}} A_j^* b_j^*,$$

and note $Y \in \mathfrak{E}_A^{\mathcal{F}}$ if and only if $(\hat{b}, Y) \in \mathfrak{P}_A$ if and only if

$$P \otimes I + \sum_{j \notin \mathcal{F}} A_j \otimes Y_j + \sum_{j \notin \mathcal{F}} A_j^* \otimes Y_j^* \succ 0.$$

For $j \in \tilde{\mathcal{F}}$, let $B_j = P^{-\frac{1}{2}} A_j P^{-\frac{1}{2}}$ and observe $Y \in \mathfrak{E}_A^{\mathcal{F}}$ if and only if

$$I + \sum_{j \notin \mathcal{F}} B_j \otimes Y_j + \sum_{j \notin \mathcal{F}} B_j^* \otimes Y_j^* \succ 0.$$

Hence $\mathfrak{E}_A^{\mathcal{F}}$ is a free spectrahedron too. ■

6.2. Two auxiliary automorphisms. Given $Y \in \mathfrak{P}_A^{\mathcal{F}}$, let $T = (\hat{0}, Y)$ and define

$$\psi^{\mathcal{F}}(Y) = (\psi_j(T))_{j \in \tilde{\mathcal{F}}}.$$

If $j \in \mathcal{F}$, then, since (1) π is the identity on \mathcal{F} , (2) the automorphism ψ is normalized and (3) $j \in \mathfrak{N}$,

$$\psi_j(T) = \mathfrak{m}_{b_j}(T_{\pi(j)}) = \mathfrak{m}_{b_j}(T_j) = \mathfrak{m}_{b_j}(0) = b_j.$$

Thus

$$\psi(T) = \psi((\hat{0}, Y)) = (\hat{b}, \psi^{\mathcal{F}}(Y)) \in \mathfrak{P}_A.$$

It follows that $\psi^{\mathcal{F}}(Y) \in \mathfrak{E}_A^{\mathcal{F}}$ and thus $\psi^{\mathcal{F}}$ determines a free mapping $\psi^{\mathcal{F}} : \mathfrak{P}_A^{\mathcal{F}} \rightarrow \mathfrak{E}_A^{\mathcal{F}}$.

Lemma 6.2. *The mapping $\psi^{\mathcal{F}} : \mathfrak{P}_A^{\mathcal{F}} \rightarrow \mathfrak{E}_A^{\mathcal{F}}$ is bianalytic.*

Proof. Evidently $\psi^{\mathcal{F}}$ is analytic and one-one. It remains to show it is onto. To this end, let $Z \in \mathfrak{E}_A^{\mathcal{F}}$ be given. Thus $(\hat{b}, Z) \in \mathfrak{P}_A$. Since ψ is onto, there is a $T \in \mathfrak{P}_A$ such that $\psi(T) = (\hat{b}, Z)$. For $j \in \mathcal{F} \subseteq \mathfrak{N}$,

$$b_j I = \psi_j(T) = \mathfrak{m}_{b_j}(T_j) = (b_j + T_j)(I + b_j T_j)^{-1}$$

and hence $T_j = 0$. Thus, setting $Y = T_j$ for $j \in \tilde{\mathcal{F}}$, it follows that $T = (\hat{0}, Y)$, so that $Y \in \mathfrak{P}_A^{\mathcal{F}}$, and $\psi((\hat{0}, Y)) = (\hat{b}, Z)$. Hence $Y \in \mathfrak{P}_A^{\mathcal{F}}$ and $\psi^{\mathcal{F}}(Y) = Z$. ■

Lemma 6.3. *For each $j \notin \mathcal{F}$ there exists $\gamma_j \in \mathbb{T}$ such that,*

$$\psi_j(\hat{0}, Y) = \gamma_j Y_{\pi(j)},$$

for all $Y \in \mathfrak{P}_A^{\mathcal{F}}$. In particular, for each $Y \in \mathfrak{P}_A^{\mathcal{F}}$,

$$\psi_j^{\mathcal{F}}(Y) = \gamma_j Y_{\pi(j)}.$$

Proof. The mapping $\psi^{\mathcal{F}} : \mathfrak{P}_A^{\mathcal{F}} \rightarrow \mathfrak{E}_A^{\mathcal{F}}$ is bianalytic by Lemma 6.2. Moreover, $\psi^{\mathcal{F}}(0) = 0$ since $b_k = 0$ for $k \in \tilde{\mathcal{F}}$. Since, both $\mathfrak{P}_A^{\mathcal{F}}$ and $\mathfrak{E}_A^{\mathcal{F}}$ are circular free spectrahedra (Lemma 6.1) and since $\psi^{\mathcal{F}}(0) = 0$, it follows from [HKM11, Theorem 4.4] that $\psi^{\mathcal{F}}$ is linear.

Now suppose $j \in \mathfrak{J}$. Since ψ is normalized, there exists $\gamma_j \in \mathbb{T}$ and $\mathfrak{h}_{\pi(j)} \in \mathfrak{H}^{\pi(j)}$ such that

$$\psi_j(x) = \gamma_j x_{\pi(j)} + \mathfrak{h}_{\pi(j)}(x).$$

Thus, for $Y \in \mathfrak{P}_A^{\mathcal{F}}$,

$$\psi_j^{\mathcal{F}}(Y) = \psi_j(\hat{0}, Y) = \gamma_j Y_{\pi(j)} + \mathfrak{h}_{\pi(j)}(\hat{0}, Y),$$

It follows that $\mathfrak{h}_{\pi(j)}(\hat{0}, Y) = 0$ for $Y \in \mathfrak{P}_A^{\mathcal{F}}$.

If $j \in \mathfrak{N} \cap \tilde{\mathcal{F}}$ and $Y \in \mathfrak{P}_A^{\mathcal{F}}$, then $\psi_j^{\mathcal{F}}(Y) = \mathfrak{m}_{b_j}(Y_j) = \mathfrak{m}_0(Y_j) = Y_j$, since π is the identity on \mathfrak{N} . \blacksquare

Let ι denote the inclusion of $\mathfrak{E}^{\mathcal{F}}$ into $\mathfrak{P}_A^{\mathcal{F}}$ and $\phi^{\mathcal{F}} = \iota \circ \psi^{\mathcal{F}} : \mathfrak{P}_A^{\mathcal{F}} \rightarrow \mathfrak{P}_A^{\mathcal{F}}$.

Lemma 6.4. *The mapping $\phi^{\mathcal{F}} : \mathfrak{P}_A^{\mathcal{F}} \rightarrow \mathfrak{P}_A^{\mathcal{F}}$ is bianalytic, and, for $j \notin \mathcal{F}$, there exists $\gamma_j \in \mathbb{T}$ such that*

$$\phi_j^{\mathcal{F}}(T) = \gamma_j T_{\pi(j)}.$$

Proof. For notational ease, let $\phi = \phi^{\mathcal{F}}$. Since both ι and $\psi^{\mathcal{F}}$ are free analytic and injective, so is ϕ . It remains to prove ϕ is onto.

For each $j \notin \mathcal{F}$, there exist $\gamma_j \in \mathbb{T}$ such that, if $Y \in \mathfrak{P}_A^{\mathcal{F}}$, then

$$\phi_j(Y) = \psi_j^{\mathcal{F}}(Y) = \gamma_j Y_{\pi(j)}$$

by Lemma 6.3. Since $\tilde{\mathcal{F}}$ is invariant for π , there is a positive integer n such that $\pi^{(n)}$ is the identity on $\tilde{\mathcal{F}}$ (π was already the identity on $\mathcal{N} \supseteq \mathcal{F}$). Thus, for $j \notin \mathcal{F}$, there exists $\delta_j \in \mathbb{T}$ such that

$$\phi_j^{(n)}(Y) = \delta_j Y_j.$$

Since the mapping $Y \mapsto \delta \cdot Y$, where $(\delta \cdot Y)_j = \delta_j Y_j$, is an automorphism of \mathfrak{P}_A (Lemma 6.1) it follows that $\phi^{(n)}$ is onto and thus so is ϕ . \blacksquare

6.3. How to spot a polydisc. Recall the notation $(X, Y) \in \mathfrak{P}_A$ from the first paragraph of subsection 6.1. Given $Y \in M_n(\mathbb{C})^{\mathcal{F}}$ and $\lambda \in \mathbb{C}$, let $Y^\lambda = (\mathfrak{J}^\lambda, Y) \in M_n(\mathbb{C})^{\mathfrak{g}}$, where $\mathfrak{J}_j^\lambda = \lambda I_n$ for $j \in \mathcal{F}$.

Lemma 6.5. *If $Y^\lambda \in \mathfrak{P}_A^{\mathcal{F}}$ for each $0 \leq \lambda < 1$ and $Y \in \mathfrak{P}_A^{\mathcal{F}}$, then \mathfrak{P}_A contains a polydisc as a distinguished summand.*

Proof. Let $T = (X, Y)$ such that $(0, Y) \in \mathfrak{P}_A$ and $\|X_j\| \leq 1$ for $j \in \mathcal{F}$ be given. Thus $Y \in \mathfrak{P}_A^{\mathcal{F}}$. Given $0 \leq \lambda < 1$, by assumption $Y^\lambda \in \mathfrak{P}_A$. An application of Lemma 4.6 implies $(\lambda X, Y) \in \mathfrak{P}_A$. \blacksquare

7. PROOF OF THEOREM 1.1

It remains to prove items (i) and (iii) of Theorem 1.1, since item (ii) was already established as Proposition 3.2.

7.1. Proof of item (i).

Proposition 7.1. *If \mathfrak{P}_A does not contain a polydisc as a distinguished summand and if φ is an automorphism of \mathfrak{P}_A , then $\varphi(0) = 0$.*

Proof. We may assume φ normalized. In particular, $\mathcal{F}_\varphi = \{j : \varphi_j(0) \neq 0\}$. Let $\mathcal{F} = \mathcal{F}_\varphi$. Arguing the contrapositive, suppose $\mathcal{F} \neq \emptyset$. By Proposition 5.4 we may (and do) assume that for each positive integer m , the automorphism $\varphi^{(m)}$ is a normalized automorphism whose associated permutation is the identity on \mathfrak{N} and $\mathcal{F}_{\varphi^{(m)}} = \mathcal{F}$. In particular, φ satisfies the assumptions (on ψ) at the outset of Section 6.

Fix $j \in \mathcal{F}$. Since (1) φ is normalized, (2) $j \in \mathfrak{N}$, and (3) π is the identity on \mathfrak{N} ,

$$(7.1) \quad \varphi_j(x) = \mathbf{m}_{b_j}(x_j) = \frac{b_j + x_j}{1 + b_j x_j}.$$

Now suppose m is a positive integer and $\varphi_j^{(m)}(0) = \mathbf{m}_{b_j}^{(m)}(0)$. Since the permutation associated to $\varphi^{(m)}$ is the identity on \mathfrak{N} , an application of Lemma 5.2 together with equation (7.1) implies

$$\varphi_j^{(m+1)}(0) = \varphi_j(\varphi^{(m)}(0)) = \mathbf{m}_{b_j}(\varphi_j^{(m)}(0)) = \mathbf{m}_{b_j}(\mathbf{m}_{b_j}^{(m)}(0)).$$

Thus, by induction, $\varphi_j^{(m)}(0) = \mathbf{m}_{b_j}^{(m)}(0)$ for all m .

It is easily seen that the sequence $(\mathbf{m}_{b_j}^{(m)}(0))_m$ converges to 1. (It is an increasing sequence from $(0, 1)$ that cannot converge to an $L < 1$.) Hence, given $0 < \lambda < 1$, there is an m such that

$$\varphi_j^{(m)}(0) \geq \lambda$$

for each $j \in \mathcal{F}$. Let $\psi = \varphi^{(m)}$. In particular, ψ is a normalized automorphism, the permutation κ associated to ψ is the identity on $\mathfrak{N} \supseteq \mathcal{F}$ and $\mathcal{F}_\psi = \mathcal{F}$.

By Lemma 6.4, given $Y \in \mathfrak{P}_A^{\mathcal{F}}$, there exists a $Z \in \mathfrak{P}_A^{\mathcal{F}}$ such that $\psi^{\mathcal{F}}(Z) = Y$. It follows that

$$\psi(\hat{0}, Z) = (\hat{c}, \psi^{\mathcal{F}}(Z)) = (\hat{c}, Y),$$

where $c_j = \psi_j(0) \geq \lambda$ for $j \in \mathcal{F}$. In particular, $(\mathfrak{J}^\lambda, Y) \in \mathfrak{P}_A$. Thus, by Lemma 6.5, \mathfrak{P}_A contains a polydisc as a distinguished summand. ■

7.2. **Proof of item (iii).** Item (iii) of Theorem 1.1 follows from Proposition 7.2 below.

Proposition 7.2. *If there exists non-identity permutation ρ of $\{1, 2, \dots, \mathfrak{g}\}$ such that the mapping $\psi_j(x) = x_{\rho(j)}$ is an automorphism of \mathfrak{P}_A , then \mathfrak{P}_A is a coordinate direct sum.*

Let $\{\mathbf{e}_1, \mathbf{e}_2\}$ denote the standard basis for \mathbb{C}^2 . Thus $\mathbf{e}_j \mathbf{e}_k^*$ are the usual matrix units for $M_2(\mathbb{C})$.

Lemma 7.3. *Fix $1 \leq \mu \leq \mathfrak{g} - 1$ and let $X \in M_n(\mathbb{C})^{\mathfrak{g}}$ be given. If $(X_1, \dots, X_\mu, 0, \dots, 0)$, and $(0, \dots, 0, X_{\mu+1}, \dots, X_{\mathfrak{g}})$ are both in \mathfrak{P}_A , then, for each choice of $\ell_j \in \{1, 2\}$ for $j > \mu + 1$, the tuple Z defined by*

$$(a) \ Z_j = \mathbf{e}_1 \mathbf{e}_1^* \otimes X_j \text{ for } 1 \leq j < \mu;$$

$$(b) \ Z_\mu = \mathbf{e}_1 \mathbf{e}_2^* \otimes X_\mu;$$

$$(c) \ Z_{\mu+1} = \mathbf{e}_1 \mathbf{e}_1^* \otimes X_{\mu+1};$$

$$(d) \ Z_j = \mathbf{e}_{\ell_j} \mathbf{e}_{\ell_j}^* \otimes X_j \text{ for } j > \mu + 1,$$

is in \mathfrak{P}_A .

Proof. Let

$$(a) \ Y_\mu = \mathbf{e}_1 \mathbf{e}_1^* \otimes X_\mu;$$

$$(b) \ Y_{\mu+1} = \mathbf{e}_2 \mathbf{e}_2^* \otimes X_{\mu+1}; \text{ and}$$

$$(c) \ Y_j = I_2 \otimes X_j \text{ for } \mu + 1 \neq j \neq \mu.$$

Since $L_A(Y)$ is a direct sum of $L_A((X_1, \dots, X_\mu, 0, \dots, 0))$ and $L_A((0, \dots, 0, X_{\mu+1}, \dots, X_{\mathfrak{g}}))$ with the identity, it follows that $Y \in \mathfrak{P}_A$.

Let

$$W_j = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$$

for $j \geq \mu$ and $W_j = I$ for $0 \leq j < \mu$ and observe that, in the notation of equation (4.2),

$$(i) \ (W \circ Y)_\mu = \mathbf{e}_1 \mathbf{e}_2^* \otimes X_\mu;$$

$$(ii) \ (W \circ Y)_{\mu+1} = \mathbf{e}_1 \mathbf{e}_1^* \otimes X_{\mu+1}; \text{ and}$$

$$(iii) \ (W \circ Y)_j = Y_j \text{ otherwise.}$$

By Lemma 4.3, $W \circ Y \in \mathfrak{P}_A$, from which it readily follows that $Z \in \mathfrak{P}_A$. ■

Lemma 7.4. *Fix $2 \leq \nu \leq \mathfrak{g}$. Given a tuple $T \in M(\mathbb{C})^{\mathfrak{g}}$, if the tuple Z_j defined by*

$$(i) \ Z_j = \mathbf{e}_1 \mathbf{e}_1^* \otimes T_j \text{ for } j < \nu;$$

$$(ii) \ Z_\nu = \mathbf{e}_1 \mathbf{e}_2^* \otimes T_\nu \text{ for } j = \nu; \text{ and}$$

(iii) $Z_j = \mathbf{e}_2 \mathbf{e}_2^* \otimes T_j$ for $j > \nu$

is in \mathfrak{P}_A , then $T \in \mathfrak{P}_A$.

Proof. Observe that $L_A(Z) \succ 0$ since $Z \in \mathfrak{P}_A$ and that $L_A(Z)$ is the direct sum of an identity matrix with $L_A(T)$. Hence $T \in \mathfrak{P}_A$. ■

Proof of Proposition 7.2. Let μ denote the first index μ such that $\rho^{-1}(\mu + 1) < \rho^{-1}(\mu)$. Such an index must exist since ρ is not the identity permutation. Hence $\rho^{-1}(\mu) > \rho^{-1}(\mu - 1) > \dots > \rho^{-1}(1)$; in particular $\rho^{-1}(j) < \rho^{-1}(\mu)$ for $j < \mu$.

Let $X \in M_n(\mathbb{C})^g$ be given and suppose $(X_1, \dots, X_\mu, 0, \dots, 0)$, and $(0, \dots, 0, X_{\mu+1}, \dots, X_g)$ are both in \mathfrak{P}_A . Define a tuple Z as follows:

$$Z_j = \begin{cases} \mathbf{e}_1 \mathbf{e}_2^* \otimes X_\mu & \text{for } j = \mu \\ \mathbf{e}_1 \mathbf{e}_1^* \otimes X_j & \text{for } \rho^{-1}(j) < \rho^{-1}(\mu) \\ \mathbf{e}_2 \mathbf{e}_2^* \otimes X_j & \text{for } \rho^{-1}(j) > \rho^{-1}(\mu). \end{cases}$$

Let $\nu = \rho^{-1}(\mu)$. Since $\rho^{-1}(j) < \nu$ for $j < \mu$ and $\rho^{-1}(\mu + 1) < \nu = \rho^{-1}(\mu)$, Lemma 7.3 implies $Z \in \mathfrak{P}_A$. Thus $S = \psi(Z) \in \mathfrak{P}_A$ and $S_k = Z_{\rho(k)}$. If $k < \nu$ and $k = \rho^{-1}(j)$, then $S_k = Z_j = \mathbf{e}_1 \mathbf{e}_1^* \otimes X_j$. If $\rho^{-1}(j) = k > \nu$, then $S_k = Z_j = \mathbf{e}_1 \mathbf{e}_2^* \otimes X_j$. Finally, if $k = \nu$, then $S_k = Z_{\rho(k)} = Z_\mu = \mathbf{e}_1 \mathbf{e}_2^* \otimes X_\mu$. An application of Lemma 7.4 implies the tuple T with $T_k = X_{\rho(k)}$ is in \mathfrak{P}_A ; that is $\psi(X) \in \mathfrak{P}_A$. Hence $\psi^{-1}(\psi(X)) = X \in \mathfrak{P}_A$. Thus \mathfrak{P}_A is a coordinate direct sum. ■

REFERENCES

- [AHKM] Meric Augat, Igor Klep, Bill Helton and Scott McCullough, *Bianalytic maps between free spectrahedra*, Math. Annalen 371 (2018), no. 1-2, 883–959. [4](#)
- [BPT] G. Blekherman, P.A. Parrilo, R.R. Thomas (editors), *Semidefinite optimization and convex algebraic geometry*, SIAM, 2013. [2](#)
- [dOH06] M.C. De Oliveira and J. W. Helton, *Computer algebra tailored to matrix inequalities in control* International Journal of Control, (2006) 79:11, 1382-1400, DOI:10.1080/00207170600725529 [2](#)
- [dOHMP] Mauricio C. de Oliveira, J. William Helton, Scott McCullough and Mihai Putinar, *Engineering systems and free semi-algebraic geometry* in Emerging applications of algebraic geometry, 17–61, IMA Vol. Math. Appl., 149, Springer, New York, 2009. [2](#)
- [HKM11] Bill Helton, Igor Klep and Scott McCullough, *Proper analytic free maps*, J. Functional Analysis 260 (2011), no. 5, 1476–1490. [7](#), [23](#)
- [HKM12] Bill Helton, Igor Klep and Scott McCullough, *Free analysis, convexity and LMI domains*. Mathematical methods in systems, optimization, and control, 195–219, Oper. Theory Adv. Appl., 222, Birkhäuser/Springer Basel AG, Basel, 2012. [4](#)

- [HKMV20] Bill Helton, Igor Klep, Scott McCullough and Juriij Volčič, *Bianalytic free maps between spectrahedra and spectraballs*, Journal of Functional Analysis, **278** no. 11, 15 June 2020, doi.org/10.1016/j.jfa.2020.108472. [2](#), [4](#)
- [KVV14] Dmitry Kalyuzhnyi-Verbovetskiĭ and Victor Vinnikov, *Foundations of free noncommutative function theory*, Mathematical Surveys and Monographs 199, Amer. Math. Soc., 2014. [2](#)
- [MT+] S. McCullough and N. Tuovila, *Reinhardt Free Spectrahedra*, Linear Algebra and its Applications (to appear) [arXiv:2012.02289](#) [2](#), [3](#)
- [MT16] J.E. McCarthy, R.M. Timoney: *Non-commutative automorphisms of bounded non-commutative domains*, Proc. Roy. Soc. Edinburgh Sect. A 146 (2016) 1037–1045. [4](#)
- [SIG97] R.E. Skelton, T. Iwasaki, K.M. Grigoriadis, *A Unified Algebraic Approach to Linear Control Design*, Taylor & Francis, 1997. [2](#)
- [Pop10] G. Popescu: *Free holomorphic automorphisms of the unit ball of $B(H)^n$* , J. reine angew. Math. 638 (2010) 119–168. [4](#)
- [WSV12] H. Wolkowicz, R. Saigal, L. Vandenberghe (eds.), *Handbook of semidefinite programming: theory, algorithms, and applications*, International Series in Operations Research & Management Science 27, Springer, 2012 [2](#)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FL

Email address: sam@uf1.edu

INDEX

- $L_B(X)$, 1
- $M_n(\mathbb{C})^{\mathcal{F}}$, 22
- $M_n(\mathbb{C})^{\tilde{\mathcal{F}}}$, 22
- $M_n(\mathbb{C})$, 1
- $M_n(\mathbb{C})^{\mathfrak{g}}$, 1
- $T \succ 0$, 1
- $T \succeq 0$, 1
- $W \circ T$, 13
- $\Lambda_B(x)$, 1
- \mathcal{F} , 22
- \mathcal{F}_φ , 21
- δ_k , 7
- \mathcal{L} , 6
- \mathfrak{H}_\pm^k , 12
- \mathfrak{J}^λ , 24
- $\mathfrak{P}_A^{\mathcal{F}}$, 22
- \mathfrak{P}_B , 1
- $\mathfrak{P}_B[n]$, 1
- \mathfrak{z} , 12
- \mathfrak{z}^+ , 12
- \mathfrak{z}^- , 12
- \mathfrak{g} , 12
- \mathfrak{m} , 20
- \mathfrak{m} , 7
- $\hat{0}$, 22
- $\hat{1}$, 22
- \mathbb{D} , 5
- \mathbb{T} , 2
- \mathfrak{H}^j , 7
- \mathfrak{N} , 12
- \mathcal{W}^j , 7
- \mathcal{W}_N^j , 7
- \mathcal{W}_N , 7
- $\mathfrak{E}_A^{\mathcal{F}}$, 22
- $\phi^{\mathcal{F}}$, 24
- $\pi_{\mathfrak{z}}$, 3
- $\psi^{\mathcal{F}}$, 23
- $\mathbf{T}(t)$, 8
- $f[n]$, 2
- $f^{(n)}$, 21
- $h_j(x)$, 6
- analytic, 2
- circular, 4
- contains a polydisc as a distinguished summand, 3
- coordinate direct sum, 3
- evaluates, 1
- free automorphism, 2
- free bianalytic, 2
- free biholomorphic, 2
- free map, 2
- free polydisc, 3
- free sets, 2
- free spectrahedron, 1
- hyper-Reinhardt, 2
- normalized automorphism, 19
- permutation associated to, 7
- spectraball, 4
- spectrahedron, 1
- trivial automorphisms, 2